Uncertainty quantification for intervals

Carlos García Meixide^{1,2}, Michael R. Kosorok³ Marcos Matabuena⁴

¹Instituto de Ciencias Matemáticas, CSIC

²Departamento de Matemáticas, Universidad Autónoma de Madrid

³University of North Carolina, Chapel Hill

⁴*Harvard University*

Abstract: Data following an interval structure are increasingly prevalent in many scientific applications. In medicine, clinical events are often monitored between two clinical visits, making the exact time of the event unknown and generating outcomes with a range format. As interest in automating healthcare decisions grows, uncertainty quantification via predictive regions becomes essential for developing reliable and trusworthy predictive algorithms. However, the statistical literature currently lacks a general methodology for interval targets, especially when these outcomes are incomplete due to censoring. We propose a uncertainty quantification algorithm and establish its theoretical properties using empirical process arguments based on a newly developed class of functions specifically designed for interval data structures. Although this paper primarily focuses on deriving predictive regions for interval-censored data, the approach can also be applied to other statistical modeling tasks, such as goodness-of-fit assessments. Finally, the applicability of the methods developed here is illustrated through various biomedical applications, including two clinical examples: i) sleep time and its link with cardiovasculuar diseases ii) survival time and physical activity values.

MSC2020 subject classifications: Primary 62F, 62G, 62N Keywords: Interval-censoring, Conformal inference, Empirical processes.

1. Introduction

Survival analysis is a core statistical methodology in biomedical research and precision medicine [40], providing methods to quantify the lifetime of patients under various conditions and treatments. In these fields, survival models are the primary tools for establishing the effectiveness of drugs and clinical interventions concerning the occurrence of specific events, as is the case in clinical trials [18].

Traditionally, practitioners have focused on right-censored outcomes, which constitute the conventional setting in survival analysis [38]. However, in other clinical applications, such as digital health, outcomes often possess an interval structure. For instance, with electronic records [34], the time to a clinical event may occur between two medical visits, as seen in the case of diabetes mellitus or other chronic diseases. Another example of an interval setup is found in clinical settings related to sleep studies. Here, patients typically record the time they go to bed and the time they wake up; however, the precise moment they actually fall asleep is not documented. This results in an interval of time during which the event (falling asleep) occurs, but the exact time within this interval remains unknown. Similarly, with wearables, devices are not worn continuously, and some defined clinical events may occur between two wear periods. Given that patient monitoring is often quasi-continuous, it is imperative to develop new methodologies for handling these interval-structured outcomes. A comprehensive illustration is depicted in Figure 1.

Quantifying the uncertainty of predictive outcomes is one of the most crucial modeling tasks in statistical sciences, particularly in clinical applications within personalized medicine [30]. In clinical studies, the response of patients over time exhibits individual and heterogeneous behavior. Therefore, when predicting clinical outcomes, there is considerable variability in patient responses. Reporting only point estimates, such as the conditional mean, provides partial and imperfect knowledge of the phenomenon under study and can lead to misleading clinical conclusions regarding the effects of drugs and interventions [5]. Consequently, measures of uncertainty, such as predictive regions, should be included in analyses. Although the construction of covariate-dependent predictive regions has received significant attention throughout the historical development of statistics [69, 70], the literature on survival analysis and censored outcomes remains more limited. With the proliferation of data-driven systems across various scientific and engineering fields, quantifying uncertainty has become a central research topic within the statistical and machine learning communities.

A particularly prolific framework that has emerged in recent years is conformal inference [69], which encompasses a general family of methods. Given a random sample $\mathcal{D}_N = \{(X_i, T_i) \in \mathcal{X} \times \mathcal{T} : 1 \leq i \leq N\}$, assumed to be at least exchangeable, and a new observation $(X_{N+1}, T_{N+1}) \in \mathcal{X} \times \mathcal{T}$ with





Fig 1: Illustration of a target variable with an interval structure. Each segment represents a patient. The borders of the black lines indicate medical checkpoints that bound the true, unknown disease onset. The light grey segments represent right-censored patients, while the dark grey segments represent left-censored patients.

the same exchangeability property relative to \mathcal{D}_N , we can define a predictive region $\widehat{\mathcal{C}}_{1-\alpha}(\cdot) \subset \mathcal{T}$ such that the following non-asymptotic property is satisfied:

$$\mathbb{P}(T_{N+1} \in \mathcal{C}_{1-\alpha}(X_{N+1})) \ge 1 - \alpha.$$

Here, \mathbb{P} represents the joint probability law over the entire dataset \mathcal{D}_N and the new data point (X_{N+1}, T_{N+1}) . Other methods in the data analysis literature that address similar scientific problems include Bayesian methods, which require the subjective specification of priors for automatic uncertainty quantification, asymptotic approximations based on the central limit theorem, and, more broadly, bootstrapping techniques [61].

The key characteristic of the various methods mentioned is that they operate on scalar, multivariate, and functional random variables $T \in \mathcal{T}$ (see review [49]). However, in several scientific applications, the only available information is a random interval (L, U), where $0 \leq L \leq U \leq \infty$ almost surely, and almost surely $T \in (L, U) \subset \mathcal{T}$. This setup encompasses right-censoring

when $U = \infty$ and left-censoring when L = 0.

General methodologies for providing predictive regions of an interval-censored target typically focus on estimating the conditional distribution [32], but they may result in poorly calibrated predictive intervals and a higher dependence on the chosen model. The non-parametric estimation of survival functions with interval-censored data is based on [66]'s algorithms, which can be derived using a self-consistency argument [16]. Efron also proved that this approach yields the nonparametric maximum likelihood estimator (NPMLE). However, significant issues highlighted in the literature include the lack of uniqueness when the log-likelihood is not strictly concave, and the fact that consistency results are typically available only under the assumption of a fixed number of inspection times [26].

In [50] the authors deduce that the NPMLE for the joint distribution function of an interval-censored survival time and a continuous mark variable is generally not consistent. Furthermore, the self-consistent approach may not be consistent even when the inspection times are limited to finitely many values, as demonstrated by a counterexample in [76]. Additionally, traditional methodologies in computational statistics, such as the bootstrap, are known to be inconsistent when using the NPMLE as the reference measure in the presence of interval censoring [57]. Due to these challenges, practitioners often resort to parametric and semi-parametric approaches for modeling interval-censored data, as these methods are numerically more stable and yield consistent solutions from both optimization and statistical perspectives [53].

To address this gap in the literature, the goal of this paper is to introduce a new uncertainty quantification algorithm, uncervals. This algorithm is theoretically grounded in a blend of conformal prediction and bootstrap methods [57], following the resampling ideas proposed by [55]. Our approach achieves significantly better calibration, as confirmed by our simulation studies, which are consistent with previous work (see, for example, [78]). The new method could be applied to develop novel algorithms for interval-censored data, including variable selection methods and goodness-of-fit tests. To derive the theoretical properties of uncervals, we introduce a new class of functions that leverage empirical process theory. This new uncertainty interval quantification framework is highly relevant in clinical applications for precision medicine and digital health, where outcomes often exhibit an interval structure.

The main contributions of this paper are summarized below:

- 1. **Practical implications:** We introduce a comprehensive uncertainty quantification algorithm, **uncervals**, which is compatible with off-the-shelf regression models tailored for interval-censored data.
- 2. Theoretical support: To derive the formal guarantees of uncervals, we introduce a new class of functions that inherently capture the statistical nature of interval structures. This enables the use of empirical process theory to establish consistency and convergence rates.
- 3. Biomedical applications: We illustrate the potential of uncervals through relevant cases involving interval-censored data. We compare our approach to naive adaptations of right-censored algorithms and the consideration of quantiles for conditional survival functions. For these purposes, we examine the performance of our proposal across different scenarios using parametric, semi-parametric, and non-parametric regression algorithms.

1.1. Paper Structure

The structure of the paper is defined as follows. Section 2 describes the key literature on uncertainty quantification and conformal prediction for rightcensored data, providing a natural reference context. Section 3 introduces the algorithms for uncertainty quantification of interval-censored data and their theoretical analysis (Section 3.1). Section 4 presents a simulation analysis to examine the theoretical properties of uncervals. Section 5 introduces two examples to illustrate the potential of the methodology in biomedical applications. Finally, Section 6 discusses the contributions, limitations and future research directions of our interval uncertainty quantification framework.

1.2. Notation

Operating mode (0: deterministic, 1: randomized)
Predictive region
Lower Predictive Bound
Unobserved underlying time
Lower and upper bounds of a time interval, respectively
p-dimensional covariates
True conditional distribution function
Estimator of the conditional distribution function

Φ_i	Conditional probability integral transform of T_i through \hat{F}_1
Λ_i, Υ_i	Evaluations of \hat{F}_1 on observable data (L_i, X_i) and (U_i, X_i) respectively
\mathbb{P}	Joint probability law over \mathcal{D}_N and (X_{N+1}, Y_{N+1})
Р	Marginal probability law of split 1 $\mathbf{P}(\cdot) = \mathbb{P}(\cdot \cap \mathcal{I}_1)$
P	Probability law conditional on split 1 $P(\cdot) = \mathbb{P}(\cdot \mathcal{I}_1)$
n	Last index in split 2
N	Last index in split 1 and in \mathcal{D}_N
G(t)	Population cdf of Φ_1, \ldots, Φ_n conditional on \hat{F}_1
$\mathbb{Q}_n(t)$	Interval measure, $\mathbb{Q}_n(t) := \frac{1}{n} \sum_{i=1}^n \frac{1}{\Upsilon_i - \Lambda_i} \mathbb{1}_{(\Lambda_i, \Upsilon_i)}(t)$
$\mathbb{I}_n(t)$	Interval distribution, $\mathbb{I}_n(t) = \int_0^t d\mathbb{Q}_n(t')$
\mathbb{H}_n	Abstract empirical process, $\sqrt{n}(P_n - P)$
\mathbb{H}_n^*	Bootstrap empirical process, $\sqrt{n}(P_n^* - P_n)$
G_n	Empirical distribution function of Φ_1, \ldots, Φ_n
$\psi(\cdot)$	$ \cdot -b , b \in [0, 1]$
V_i	Unobserved scores, $V_i := \psi(\hat{F}_1(T_i, X_i))$
r_{N-n}	Uniform rate of convergence for survival curves

2. Related work

2.1. Statistical literature on uncertainty quantification

In recent years, uncertainty quantification became an active research area [25, 55]. The impact of uncertainty quantification on data-driven systems has led to a remarkable surge of interest in both applied and theoretical domains. These works delve into the profound implications of uncertainty quantification in statistics and beyond, such as in the biomedical field [3, 30].

Geisser's pioneering book [25] develops a mathematical theory of predictive inference. Building upon Geisser's foundations, Politis presented a comprehensive methodology [55] that effectively harnesses resampling techniques. Additionally, the book of Vovk, Gammerman and Shafer [69] has been pivotal, see recent reviews [2, 20].

One of the most widely used and robust frameworks for quantifying uncertainty in statistical and machine learning models is conformal inference [59]. The central idea of conformal inference is rooted in the concept of exchangeability [41], even though we will assume that the observedd random elements \mathcal{D}_N are independent and identically distributed (i.i.d.). Now, we present a general overview of conformal inference methods for regression models with

 $\mathbf{6}$

scalar responses. Consider the sequence $\mathcal{D}_N = \{(X_i, T_i)\}_{i=1}^N$ of i.i.d. random variables. Given a new i.i.d. random pair (X, T) with respect to \mathcal{D}_N , conformal prediction provides a family of algorithms for constructing predictive intervals independently of the regression method used, as introduced by [69].

Fix any regression algorithm

$$\mathcal{A}: \ \cup_{N \ge 0} \left(\mathcal{X} \times \mathbb{R} \right)^N \ \to \ \{\text{measurable functions } \widetilde{m}: \mathcal{X} \to \mathbb{R} \}\,,$$

which maps a data set containing any number of pairs (X_i, T_i) , to a fitted regression function \tilde{m} . The algorithm \mathcal{A} is required to treat data points symmetrically, i.e.,

$$\mathcal{A}\big((x_{\pi(1)}, t_{\pi(1)}), \dots, (x_{\pi(N)}, t_{\pi(N)})\big) = \mathcal{A}\big((x_1, t_1), \dots, (x_N, t_N)\big)$$
(1)

for all $N \ge 1$, all permutations π on $[N] = \{1, \ldots, N\}$, and all $\{(x_i, t_i)\}_{i=1}^N$. Next, for each $t \in \mathbb{R}$, let

$$\widetilde{m}^t = \mathcal{A}\big((X_1, Y_1), \dots, (X_N, T_N), (X, t)\big)$$

denote the trained model, fitted to the training data together with the test covariate value X and a hypothesized test response t. Let

$$R_{i}^{t} = \begin{cases} |T_{i} - \widetilde{m}^{t}(X_{i})|, & i = 1, \dots, N \\ |t - \widetilde{m}^{t}(X)|, & i = N + 1. \end{cases}$$
(2)

The prediction interval for X given by *full conformal* is then defined as

$$\widehat{C}_{1-\alpha}(X) = \left\{ t \in \mathbb{R} : R_{N+1}^t \le (1-\alpha) \text{ -quantile of } \sum_{i=1}^{N+1} \frac{1}{N+1} \cdot \delta_{R_i^t} \right\}$$
(3)

The *full conformal* method is known to guarantee distribution-free finitesample predictive coverage at the target level $1 - \alpha$:

Theorem 2.1 (Full conformal prediction [69]). If the data points $\mathcal{D}_N \cup \{(X,T)\}$ are *i.i.d.* (or more generally, exchangeable), and the algorithm \mathcal{A} treats the input data points symmetrically as in (1), then the full conformal prediction set defined in (3) satisfies

$$\mathbb{P}(T \in \widehat{C}_{1-\alpha}(X)) \ge 1 - \alpha.$$

The same result holds for split conformal methods, which separates the fitting and ranking steps using sample splitting, and its computational cost is simply that of the fitting step [42].

Conformal inference techniques have been applied to various regression settings, including estimation of the conditional mean [42], conditional quantiles [58], and different functionals of conditional distributions [8, 60]. In recent years, multiple extensions of conformal inference techniques have emerged to handle counterfactual inference problems [7, 36, 74], heterogeneous policy effect [6], reinforcement learning [13], federated learning [48], outlier detection [4], hyphotesis testing [31], robust optimization [37], multilevel structures [15, 19], missing data [51, 77], and survival analysis problems [5, 63], as well as problems involving dependent data such as time series and spatial data [7, 62, 72, 73].

Asymptotic marginal- and in certain cases, conditional- guarantees for the coverage are provided in [11, 71, 79]. The key idea behind these approaches is the application of resampling techniques, originally proposed by [56], to residuals or other score measures. Notably, these methodologies are applicable regardless of the predictive algorithm being used, as emphasized by [55].

Bayesian methods are also an important framework to quantify uncertainty (see [9]), which can also be integrated with conformal inference methods [2, 19, 54]. Assuming Gaussian errors and linearity [46, 64] is a classical and popular approach. However, the latter techniques generally introduce stronger parametric assumptions in statistical modeling and include the limitation or difficulty in selecting the appropriate prior distribution in Bayesian modeling [23]. The theory of tolerance regions gives another connection with the problem studied here [21, 29], which was generalized for the multivariate case with the notion of depth bands (see [45]). However, a few conditional depth measures are available in the literature [22]. Depth band measures for statistical objects that take values in metric spaces have recently been proposed [14, 24, 47, 68] but only in the unconditional case.

We now discuss in detail the non-parametric literature on conformal prediction for censored outcomes, focusing on right-censoring, as presented in [5] and [28]

2.2. Non-parametric conformal prediction for right-censored outcomes

Survival analysis traditionally aims to infer the probability that a patient will survive beyond a specified time. This is often complicated by the presence of censored data, where for some patients, survival times are only known to exceed a certain value due to limitations such as the end of a study period. Machine learning methods are introduced as promising tools for handling such complex data without relying on strong modeling assumptions. However, these methods face challenges in quantifying uncertainty, which is critical for making reliable predictions in high-stakes situations. To overcome this problem, [5] present a fully non-parametric conformal methodology for right-censoring outcomes in survival analysis to construct prediction intervals for survival times. This new methodology is particularly relevant in modern healthcare applications, such as the COVID-19 pandemic, where accurate prediction of survival times is crucial for resource allocation and decisionmaking in public health crises.

The type of predictive region they propose is called the lower predictive bound (LPB), which can be seen as a one-sided tolerance region. The LPB serves as a conservative estimate of the survival time, offering a critical tool for high-stakes decision-making by ensuring that the true survival time falls above this bound with a specified probability. We say an LPB $\hat{L}_{1-\alpha}(\cdot)$ is calibrated if it satisfies the following coverage criterion:

$$\mathbb{P}(T_{N+1} \ge \widehat{L}_{1-\alpha}(X_{N+1})) \ge 1 - \alpha,$$

where α is a pre-specified level (e.g., $\alpha \in [0, 1]$), and the probability is computed over both \mathcal{D}_N and a future unit (X_{N+1}, T_{N+1}) that is independent of the training sample $\mathcal{D}_N = \{(X_1, T_1), \ldots, (X_N, T_N)\}$. This can be seen as a predictive interval whose upper border is $+\infty$, serving as a conservative assessment of survival.

Let $T_i = \min(T_i, C_i)$ be the censored survival times. Upon making the following elementary observation:

 $\tilde{T} \leq T \implies$ any calibrated LPB on \tilde{T} is also a calibrated LPB on T (4)

a naive approach to construct distribution-free LPBs in right-censored scenarios immediately arises: just apply conformal prediction over the censored target \tilde{T} in a one-sided fashion. However, this approach is overconservative in

the sense that it will lead to an excessively low $\hat{L}_{1-\alpha}(\cdot)$. [5] observe that this phenomenon is related to censoring times being overall smaller than survival times. This led them to apply conformal inference on subpopulations with larger censoring times and correcting for the covariate shift this involved by using a one-sided version of weighted conformal inference [65]. It can even be shown that if $(T, X) \perp C$, then the covariate shift vanishes and the technique reduces to applying conformal inference on $(X_i, T_i \wedge c_0)_{C_i \geq c_0}$. This has two main problems:

- In general, there is no access to censoring times in practical situations. This would involve, for instance, knowing when a person would quit from a clinical trial even though this individual has experienced the event of death.
- Their approach depends on a hyperparameter c_0 that selects a subpopulation with censoring times bigger than c_0 . Figure 2 serves as an illustration of how sensitive the approach by [5] is to the hyperparameter c_0 .
- In some clinical applications, such as oncology where some patients have a short expected survival, truncating the observations to $C \ge c_0$ does not make biological sense.



Fig 2: Evolution of empirical coverage with the threshold c_0 used in [5]. The desirable outcome would be a horizontal line at 0.9. Setup details in 4.3

To generalize the non-asymptotic guarantees for interval-censored data, where only (L, U) containing almost surely the true survival time T are observed, we base our approach on an observation analogous to (4):

 $L \leq T \implies$ any calibrated LPB on L is also a calibrated LPB on T (5)

The first immediate issue of this approach is that it would lead to even less tight LPBs. However, there is a clear benefit from this, which is the well-known finite sample size coverage guarantee inherent to conformal inference, as stated in Theorem 3.1. Another basic observation is that an interval whose lower and upper borders are respectively the $\frac{\alpha}{2}$ and $1 - \frac{\alpha}{2}$ quantiles of T given X = x is conditionally valid. This would be brought into practice by plugging consistent estimators of such quantiles, recoverable from the conditional survival function estimated by, for instance, interval-censored recursive forests [33] algorithm. However, this naive algorithm suffers from inherent problems due to the lack of calibration steps.

3. Predictive regions for interval-censored targets

Let T be an a.s. positive random variable representing true event times. In our setup, there will never be access to a sample from T but from L and U, which are two neighboring checkup points quantifying the interval where Tis known to lie within. Let X be real p-dimensional covariates. Our sample is

$$(L_1, U_1, X_1), (L_2, U_2, X_2), \dots, (L_N, U_N, X_N) \sim (L, U, X)$$
 i.i.d

where we assume the existence of underlying T_1, T_2, \ldots, T_N respecting $L_i \leq T_i \leq U_i$ almost surely for each $1 \leq i \leq N$. The formalism we propose includes interval-, right- and left-censoring observations as special cases. We encode with $U_i = \infty$ a right-censored observation and with $L_i = 0$ a left-censored observation. Otherwise, (L_i, U_i) represents an interval-censored observation.

Let $F(t, x) = P(T_i \leq t \mid X_i = x)$ denote the conditional cumulative distribution function (cdf) of T_i given $X_i = x$. Throughout the paper, we will split the whole data $\{1, \ldots, N\}$ into $\mathcal{I}_1 := \{n + 1, \ldots, N\}$ and $\mathcal{I}_2 := \{1, \ldots, n\}$. Because of being our data i.i.d, random splits are also possible.

$$\underbrace{1,\ldots,n-1,n}_{\text{split 2}},\underbrace{n+1,\ldots,N-1,N}_{\text{split 1}},\underbrace{N+1}_{\text{new}}$$

We assume we have obtained an estimator \hat{F}_1 of F based on the halfsample \mathcal{I}_1 . We denote by $\Lambda_i := \hat{F}_1(L_i, X_i), \, \Upsilon_i := \hat{F}_1(U_i, X_i)$ the evaluations of \hat{F}_1 on observable data coming \mathcal{I}_1 and indexed by $i = 1, \ldots, n$ according to how it was chosen. We denote by $\Phi_i := F_1(T_i, X_i)$ the evaluations of F_1 on the underlying survival times T_i . We do not have access to Φ_i but we need to postulate them. Upon fixing $\{(L_i, U_i, X_i)\}_{i \in \mathcal{I}_1}$ - and therefore also $\hat{F}_1(\cdot, \cdot)$ - then $\{(\Phi_i, \Lambda_i, \Upsilon_i)\}_{i \in \mathcal{I}_2}$ are i.i.d. because they are evaluations on an i.i.d. sample of a function that became deterministic after conditioning. We denote by P the conditional-on-split-1 joint law of $(\Phi, \Lambda, \Upsilon)$ and by G the population cdf of Φ_1, \ldots, Φ_n conditional on \hat{F}_1 .

Let $\psi(\cdot) = |\cdot - b|$. For now, $b = \frac{1}{2}$. When $F(\cdot, x)$ is a symmetric unimodal distribution with a well-defined conditional density, then b = 1/2 is optimal in the sense that the interval outputted by Algorithm 1 has minimum length asymptotically¹. An extension to choose b optimally to ensure efficiency as in [8] would be certainly possible, but in this article we are interested in LPBs as is natural in survival analysis applications, so that we set b = 1 unless explicitly stated.

Algorithm 1 uncervals

Require: $\{(L_i, U_i, X_i)\}_{i=1}^N, \alpha, X_{N+1}, e$

- 1: Split $\{1, \ldots, N\}$ into $\mathcal{I}_1 := \{n + 1, \ldots, N\}$ and $\mathcal{I}_2 := \{1, \ldots, n\}$
- 2: Obtain F_1 (1- conditional survival function) using an off-the-shelf semi-parametric model for interval-censored responses based on \mathcal{I}_1 .
- 3: for i = 1, ..., n do
- $j \leftarrow \texttt{sample}(\{1, \ldots, n\}, \texttt{replace} = \text{True})$ 4:
- if e = 0 then 5:
- $\Phi_i^* \leftarrow F_1(L_i, X_i)$ 6: 7: else
- $\Phi_i^* \leftarrow \hat{F}_1(L_j, X_j) + \texttt{runif}(0, 1)(\hat{F}_1(U_j, X_j) \hat{F}_1(L_j, X_j))$ 8:
- 9: end if
- $\hat{V}_i^* \leftarrow \psi\left(\Phi_i^*\right)$ 10:
- 11: end for

12: $\hat{Q}_{\mathcal{I}_2}^* \leftarrow (1-\alpha) (1+1/n)$ empirical quantile of $\left\{ \hat{V}_i^* \right\}$ Final $(1 - \alpha)$ prediction set $\widehat{\mathcal{C}}^{e}_{(1-\alpha)}(X_{N+1}) \leftarrow \left\{ t \ge 0 : \psi(\widehat{F}_{1}(t, X_{N+1})) \le \widehat{Q}^{*}_{\mathcal{I}_{2}} \right\}$

The user is free to choose between two operating modes indexed by e = 0, *.

¹The fact that b = 1/2 leads to result that are asymptotically similar to the oracle has nothing to do with optimal (smallest) length.

The main, general one e = * has desirable asymptotic coverage properties provided the base method employed for estimation of the conditional survival function is well behaved when sample size tends to infinite. The second operating mode e = 0 is a generalization of [5] which enjoys finite sample size calibration guarantees paying the price of overconservativeness; as the original approach suffers. All in all, our new approach does not rely in choosing any hyperparameter.

Observe that if we set b = 1 then

$$\widehat{\mathcal{C}}_{(1-\alpha)}^{e}\left(X_{N+1}\right) = \left\{t : \left|\widehat{F}_{1}\left(t, X_{N+1}\right) - 1\right| \le \widehat{Q}_{\mathcal{I}_{2}}^{*}\right\} \\
= \left\{t : 1 - \widehat{F}_{1}\left(t, X_{N+1}\right) \le \widehat{Q}_{\mathcal{I}_{2}}^{*}\right\} = \left\{t : \widehat{S}_{1}\left(t, X_{N+1}\right) \le \widehat{Q}_{\mathcal{I}_{2}}^{*}\right\}.$$

As (empirical) survival functions are monotonically decreasing, we have that the previous set is indeed given by an LPB upon setting b = 1:

$$\widehat{L}_{(1-\alpha)}(X_{N+1}) = \left(\inf\{t : \widehat{S}_1(t, X_{N+1}) \le \widehat{Q}_{\mathcal{I}_2}^*\}, +\infty\right)$$

Theorem 3.1 (Finite-sample validity of uncervals). Set e = 0, or equivalently, consider the evaluations of \hat{F}_1 on the left borders of the intervals. Suppose that the data are i.i.d. or exchangeable and that the estimator of the conditional survival function is invariant to permutations of the data. Then $\hat{C}^0_{(1-\alpha)}(\cdot)$ is an LPB and

$$\mathbb{P}\left(T_{n+1} \in \hat{\mathcal{C}}^{\theta}_{(1-\alpha)}\left(X_{n+1}\right)\right) \ge 1 - \alpha.$$

3.1. Statistical theory: the interval process

We have introduced uncervals, an algorithm that in practice is constituted by a symbiosis between conformal prediction and the bootstrap. However, from a theoretical perspective its foundations are rooted in a new class of functions whose properties capture the phenomenology of interval data and enable the establishment of performance guarantees through VC theory arguments.

Upon fixing \hat{F}_1 , we have transitioned to an auxiliary sample

$$(\Lambda_1, \Upsilon_1), (\Lambda_2, \Upsilon_2), \ldots, (\Lambda_n, \Upsilon_n)$$

which is i.i.d. (exchangeable provided the original sample was so) after having fixed \hat{F}_1 and respects the constraint $0 \leq \Lambda_i \leq \Phi_i \leq \Upsilon_i \leq 1$ P-a.s. for each $1 \leq i \leq n$. Classical split distributional conformal prediction relies on the $(1 - \alpha)(1 + 1/n)$ empirical quantile of $\psi(\Phi_1), \ldots, \psi(\Phi_2)$. In the presence of interval-censoring, we do not have access to Φ_1, \ldots, Φ_n and even less to the aforementioned empirical quantile. However, we will use them as auxiliary entities to support our theoretical arguments, even though they are never utilized by Algorithm 1.

We will use the following conceptual bridge to open a connection between conformal inference and empirical process theory. Suppose $\Phi_1, \ldots, \Phi_{n+1}$ are exchangeable random variables free from ties. The rank of the n + 1-th observation Φ_{n+1} among $\Phi_1, \ldots, \Phi_{n+1}$ is uniformly distributed over the set $\{1, \ldots, n+1\}$. What is to say, under exchangeability we have

$$\mathbb{P}\left\{\frac{\operatorname{rank}(\Phi_{n+1})}{n+1} > \epsilon\right\} \ge 1 - \epsilon, \text{ for all } \epsilon \in [0,1]$$

The key connection with empirical process theory is the following observation:

$$\frac{\operatorname{rank}(\Phi_{n+1})}{n+1} = \frac{\#\{i=1,\dots,n+1:\Phi_i \ge \Phi_{n+1}\}}{n+1}$$
$$= \sum_{i=1}^n \frac{1}{n+1} \mathbb{1}\{\Phi_i \ge \Phi_{n+1}\}$$
$$= \mathbb{1} - \operatorname{ecdf}_{\Phi_1,\dots,\Phi_{n+1}}(\Phi_{n+1})$$

In this way, the the problem reduces to the study of the asymptotic properties of the empirical distribution function of the Φ_1, \ldots, Φ_n . One of the core mechanisms of split conformal is the following fact: if original data was i.i.d. (resp. exchangeable), then conditional on split 1 scores are i.i.d (resp. exchangeable) even though model was not well specified. If we condition on the proper training set then the calibration residuals in split 2 and the test residual are all i.i.d. (resp. exchangeable). We will see that it is the theoretical mechanism inherent to split conformal of conditioning on a subset of data what makes bootstrap being valid. Under interval censoring, we lose access to the Φ_1, \ldots, Φ_n , but we will see that it is still possible to recover bootstrap replicates $\Phi_1^*, \ldots, \Phi_n^*$ from a very particular distribution coined by us the *interval measure*.

Definition 3.2. For $0 \le t \le 1$ given a sample $(\Lambda_1, \Upsilon_1), (\Lambda_2, \Upsilon_2), \ldots, (\Lambda_n, \Upsilon_n)$ satisfying $0 \le \Lambda_i \le \Upsilon_i \le 1$ a.s. for each $1 \le i \le n$, we define the interval

measure as

$$\mathbb{Q}_n(t) := rac{1}{n} \sum_{i=1}^n rac{1}{\Upsilon_i - \Lambda_i} \mathbb{1}_{(\Lambda_i, \Upsilon_i)}(t)$$

Having access to a hypothetical sample $\{(\Phi_i, \Lambda_i, \Upsilon_i)\}_{i=1}^n$ (which is not the case because we miss the Φ_i 's), the *empirical measure* is traditionally defined for $0 \leq t, \phi, \lambda, \upsilon \leq 1$ as $P_n(\phi, \lambda, \upsilon) = \frac{1}{n} \sum_{i=1}^n \delta_{\Phi_i, \Lambda_i, \Upsilon_i}(\phi, \lambda, \upsilon)$. The first observation to be made is that, in contrast with the empirical measure, the interval measure is not composed by atoms placed on the observations but by probability boxes.

It is easy to prove by basic integration of constant functions that the length of [0, t] with respect to the interval measure is given by

$$\int_0^t d\mathbb{Q}_n(t')$$

= $\frac{1}{n} \sum_{i=1}^n \frac{1}{\Upsilon_i - \Lambda_i} \left[(t - \Lambda_i) \mathbb{1}_{[0,t]}(\Lambda_i) + (\Upsilon_i - t) \mathbb{1}_{[0,t]}(\Upsilon_i) \right]$
=: $\mathbb{I}_n(t)$

We call $\mathbb{I}_n(t)$ the *interval distribution* for $0 \le t \le 1$. It is important not to miss the original sample picture:

$$\begin{array}{ccc} \mathcal{I}_1 & \longrightarrow & \widehat{F}_1 \\ & & & \downarrow \\ \mathcal{I}_2 & \longrightarrow & \mathbb{I}_n \end{array}$$

Now let $\mathcal{G} = \{g_t(l, u) : t \in [0, 1]\}$, where

$$g_t: (l, u) \mapsto \mathbf{1}_{\{0 \le l \le u \le 1\}} \frac{1}{u - l} \left[(t - l) \mathbf{1}_{[0,t]}(l) + (u - t) \mathbf{1}_{[0,t]}(u) \right].$$

Remark 3.1. Note that for l = u =: w with $0 \le w \le 1$ we have $g_t(w, w) = 1_{[0,t]}(w), 0 \le t \le 1$

Let $t \in [0, 1]$. Then $\mathbb{I}_n(t) = P_n g_t$. Provided that $\mathbb{E}g_t(\Lambda, \Upsilon) = G(t)$, where G is the true cumulative distribution function of the $\Phi'_i s$, the interval process $\sqrt{n} (\mathbb{I}_n - G)$ coincides with the abstract empirical process indexed by the class $\mathcal{G}: \{\sqrt{n}(P_n - P)g_t, g_t \in \mathcal{G}\}$. As \mathcal{G} has a finite envelope function, the map $\sqrt{n}(P_n - P) =: \mathbb{H}_n$ can be viewed as an element in $\ell^{\infty}(\mathcal{G})$ classically known as the *empirical process*.





Fig 3: Combinatorial ground of Theorem 3.3: an illustration of how the shuttering principle of VC theory acts at the core of the arguments encountered in its proof.

3.1.1. Donsker-universality

The main theoretical result of this paper is the proof that the abstract empirical process converges weakly in $\ell^{\infty}(\mathcal{G})$, meaning that \mathcal{G} is a Donsker class. This is particularly useful because the bootstrap method is always asymptotically valid for Donsker classes of functions.

Theorem 3.3. \mathcal{G} is Donsker for any distribution on (L, U) such that $0 \leq L \leq U \leq 1$ almost surely. Moreover, \mathcal{G} has a finite envelope function.

We depict in Figure 3 the main step in the proof of Theorem 3.3, which essentially involves demonstrating that the class $\{1_{\{l \le t < u\}}(t-l) : t \in [0,1]\}$ has a VC index ≤ 3 . To show it using a shattering principle, one needs to demonstrate that no set of 3 points can be shattered by the class. This involves proving that for any set of 3 points, there does not exist a subset of functions in the class that can realize all possible dichotomies (i.e., all possible ways to separate the points into two groups). In other words, there must be at least one dichotomy that cannot be represented by any function in the class for any set of 3 points. If this condition holds for every set of 3 points, then the VC index of the class is indeed 3.

3.1.2. Unbiasedness

In scenarios involving right-censored failure time data, it is typically assumed that the censoring time is independent of the survival time, either marginally

or conditionally when accounting for external covariates. However, this assumption does not readily extend to cases of interval censoring. In the context of the previously defined notation, where L and U denote the interval endpoints and T represents the survival time, there exists an inherent relationship expressed as $L < T \leq U$. For interval-censored data, the appropriate assumption to adopt is the *noninformative interval censoring assumption* [80], specified as follows.

Assumption 1.

 $\mathbb{P}(T \le t \mid L = l, U = u, L < T \le U, X = x) = \mathbb{P}(T \le t \mid l < T \le u, X = x).$

Assumption 1 essentially states that, except for the fact that T lies between l and u, which are the realizations of L and U, the interval (L, U) (or equivalently its endpoints L and U) does not provide any extra information for T. In other words, the probabilistic behavior of T remains the same except that the original sample space $T \ge 0$ is now reduced to $l = L < T \le U = u$.

Assumption 2. There exist $0 < \tau < \infty$ and r_{N-n} such that

$$\sup_{t \le \tau, x \in \mathbb{R}^p} \left| \hat{F}_1(t, x) - F(t, x) \right| = \mathcal{O}_{\mathbf{P}}(r_{N-n})$$

Have into account that N - n is the sample size of split 1, the data used to estimate \hat{F}_1 . For example, under certain assumptions, in [10] the authors derive the following uniform rate of convergence for single-draw-per-person random survival forest (technical details regarding n_{min} , α , φ and d_l can be found in that article)

$$r'_{N-n} = \max\left\{\sqrt{\frac{\log(N-n)\left\{\log(n_{\min}) + \log\log(N-n)\right\}}{n_{\min}}}, \left(\frac{n_{\min}}{N-n}\right)^{\frac{\log((1-\alpha)-1)}{\log(\alpha^{-1})}\frac{0.991\varphi}{\max_l d_l}}\right\}$$

Assumption 3.

 $F(T, X) \perp X$

Assumption 3 states that the model is well-specified and has as an immediate consequence the following result.

Proposition 3.1. Let Assumption 1 hold. If F is continuous on t for all x,

$$F(T,X) \mid L = l, U = u, L < T \le U, X = x \sim U(F(l,x), F(u,x))$$

Notice that if Φ, Λ, Υ were built oracle-wise; i.e. using the true conditional distribution function F instead of \hat{F}_1 , then if Assumption 1 is true and F is continuous on t for all x we have that Φ is uniformly distributed on $[\Lambda, \Upsilon]$ conditional on Λ, Υ .

Proposition 3.2. Let Assumption 1 hold and let F be continuous on its first argument for all x. Let $U(\hat{F}_1(l, x), \hat{F}_1(u, x))(\cdot)$ be the cdf of the uniform distribution over the interval $(\hat{F}_1(l, x), \hat{F}_1(u, x))$ and define $J(\cdot, l, u, x) := P\left(\hat{F}_1(T, X) \leq \cdot \mid l \leq T \leq u, X = x\right)$. Then,

$$J(\cdot, l, u, x) - U(\hat{F}_1(l, x), \hat{F}_1(u, x))(\cdot) = \mathcal{O}_{\mathbf{P}}(r_{N-n})$$

Note that the **P** involved in the stochastic order symbol $\mathcal{O}_{\mathbf{P}}$ represents the marginal probability distribution of split 1. We need the following auxiliary result to ensure that the interval distribution is an unbiased estimator of G.

For the sake of formal simplicity in the proofs, we will assume that the Φ_i are genuine probability integral transforms constructed using F, as justified by Proposition 3.2.

Lemma 3.4. Let Φ be uniformly distributed on $[\Lambda, \Upsilon]$ conditional on Λ, Υ . Then

$$E1\{\Phi \le \cdot \le \Upsilon\} = E\left[1\{\Lambda \le \cdot \le \Upsilon\}\frac{\cdot - \Lambda}{\Upsilon - \Lambda}\right],$$

where the expectation is taken jointly wrt to Φ, Λ, Υ .

The proof of the next Corollary is inmediate having into account the two previous results and writing $G(t) = P(\Phi \le t) = E1\{\Phi \le t\}$

Corollary 3.1. Suppose Assumption 1 is true and that F is continuous on t for all x. Then for any $t \in [0, 1]$, the following holds

$$Eg_t(\Lambda, \Upsilon) = G(t)$$

where the expectation is taken jointly wrt to Φ, Λ, Υ .

Corollary 3.2. The sequence of interval processes

$$\sqrt{n}\left(\mathbb{I}_n-G\right)$$

converges in distribution in the space D[0,1] to a tight random element \mathbb{H}_P , whose marginal distributions are zero-mean normal.

The limit process that shows up in Corollary 3.2, \mathbb{H}_P , is a Brownian bridge because of multivariate central limit theorem: given any finite set of measurable functions g_i with $Pg_i^2 < \infty$ then $(\mathbb{H}_n g_1, \ldots, \mathbb{H}_n g_k) \rightsquigarrow (\mathbb{H}_P(1), \ldots, \mathbb{H}_P(k))$, where the vector on the right possesses a multivariate-normal distribution with mean zero, provided that Assumption 1 is true and that F is continuous on t for all x.

Bootstrapping \mathbb{I}_n generates a sample $\Phi_1^*, \ldots, \Phi_n^*$ giving rise to the *bootstrap* empirical measure, which corresponds to

$$P_n^* = n^{-1} \sum_{i=1}^n \delta_{\Phi_i^* = \Lambda_i^* = \Upsilon_i^*}(\phi, \lambda, \upsilon)$$

and to the bootstrap empirical process

$$\mathbb{H}_n^* = \sqrt{n} \left(P_n^* - P_n \right) \in \ell^\infty(\mathcal{G})$$

Remark 3.2. Let $t \in [0, 1]$. Then, in virtue of Remark 3.1 we have $P_n^* g_t = n^{-1} \sum_{i=1}^n 1\{\Phi_i^* \le t\} := \mathbb{I}_n^*(t)$.

Corollary 3.3. Conditionally given $\Phi_1, \Lambda_1, \Upsilon_1, \Phi_2, \Lambda_2, \Upsilon_2...$ the sequence $\sqrt{n} (\mathbb{I}_n^* - \mathbb{I}_n)$ converges to \mathbb{H}_P .

Corollary 3.3, whose proof is inmediate by using Theorem 23.7 in [67], is the key formal step of our theoretical arguments, which can be graphically regarded as

or informally writing,

$$\boxed{\sqrt{n}\left(\mathbb{I}_{n}^{*}-\mathbb{I}_{n}\right)\mid\mathbb{I}_{n}\overset{d}{\approx}\sqrt{n}\left(\mathbb{I}_{n}-G\right)}$$

The next result constitutes the last auxiliary step to be taken towards deriving the asymptotic guarantees of uncervals.

Lemma 3.5. Let G_n be the empirical distribution function of Φ_1, \ldots, Φ_n (unknown). Then,

$$G_n(\phi) - \mathbb{I}_n(\phi) = o_P(1)$$

The proof is as follows: $G_n(\phi) - \mathbb{I}_n(\phi) = G_n(\phi) - G(\phi) + G(\phi) - \mathbb{I}_n(\phi) = o_P(1) + o_P(1) = o_P(1)$ because of the classical Glivenko-Cantelli Theorem and Theorem 3.2.

We now present two significant results regarding the consistency of **untervals**, which form a crucial part of this paper.

Theorem 3.6 (Asymptotic unconditional validity). Let Assumptions 1 and 2 hold. Then,

$$\mathbb{P}\left(T_{N+1}\in\widehat{\mathcal{C}}^*_{(1-\alpha)}\left(X_{N+1}\right)\right) = 1 - \alpha + o_{\mathbb{P}}(1).$$

Theorem 3.7 (Asymptotic conditional validity). Let Assumptions 1,2 and 3 hold. Then,

$$\mathbb{P}\left(T_{N+1}\in\widehat{\mathcal{C}}^*_{(1-\alpha)}\left(X_{N+1}\right)\mid X_{N+1}\right)=1-\alpha+o_{\mathbb{P}}(1).$$

Remark 3.3. Have into account that the probability \mathbb{P} involved in Theorems 3.6 and 3.7 (unconditional and conditional on X_{N+1} respectively) is the one from which the entire dataset \mathcal{D}_N and the new data point (X_{N+1}, Y_{N+1}) were originally sampled.

The proof of Theorem 3.7 is the same as that of Theorem 3.6 having into account that under correct specification, ranks are independent of the predictors unlike regression residuals. Theorems 3.6 and 3.7 establish the asymptotic validity of our procedure under weak and easy to verify conditions. Also, Theorem 3.7 implies that under its assumptions, Algorithm 1 outputs an interval that asymptotically coincides with the oracle when $b = \frac{1}{2}$.

3.1.3. Convergence rates

We start with a finite-sample bound for the supremum of the absolute difference between the empirical and theoretical cumulative distribution functions (cdfs) of the unobserved scores. For $i \in \mathcal{I}_2$ define $F_i = F(T_i, X_i)$ and let $\psi(F_i)$ be the oracle scores. Let $\tilde{G}(v) = \frac{1}{n} \sum_{i=1}^{n} 1 \{ \psi(F_i) < v \}$ be the empirical cdf of oracle scores and $\hat{G}(v) = n^{-1} \sum_{i=1}^{n} 1 \{ V_i < v \}$. Recall that $V_i = \hat{F}_1(T_i, X_i)$.

Lemma 3.8. Assume that G is such that $\sup_{x_1 \neq x_2} |G(x_1) - G(x_2)| / |x_1 - x_2| := W < \infty$, i.e. G is W-Lipschitz. Then we have for any $\delta > 0$

$$\begin{split} \sup_{v \in \mathbb{R}} \left| \widehat{G}(v) - G(b+v) + G(b-v) \right| &\leq \\ 2 \frac{\sup_{t \leq \tau, x \in \mathbb{R}^p} \left| \widehat{F}_1(t,x) - F(t,x) \right|^2}{\delta^2} + 2\delta W + 3 \sup_{v \in \mathbb{R}} \left| \widetilde{G}(v) - G(b+v) + G(b-v) \right| \quad \mathbb{P}-a.s. \end{split}$$

Observe that G(b+v) - G(b-v) is the theoretical cdf of the oracle scores $|F(T_i, X_i) - b|$.

The proof of Lemma 3.8 is an adaptation from that of Lemma 1 in [8] and having into account that

$$n^{-1} \sum_{i=1}^{n} \left(|V_i - \psi(F_i)| \right)^2 = n^{-1} \sum_{i=1}^{n} \left(\left| \hat{F}_1(T_i, X_i) - \frac{1}{2} \right| - \left| F(T_i, X_i) - \frac{1}{2} \right| \right)^2$$
$$\leq n^{-1} \sum_{i=1}^{n} \left(\left| \hat{F}_1(T_i, X_i) - F(T_i, X_i) \right| \right)^2$$

Next, we use the classical Glivenko-Cantelli Theorem for the part involving \tilde{G} , and for the rest, we choose δ optimally to tighten the bound in Lemma 3.8 as much as possible.

Proposition 3.3. Let Assumption 2 hold and let G be W-Lipschitz. Then,

$$\sup_{v \in \mathbb{R}} \left| \widehat{G}(x) - G(b+v) + G(b-v) \right| = \mathcal{O}_{\mathbf{P}} \left((Wr_{N-n})^{2/3} \right) + \mathcal{O}_{P} \left(n^{-1/2} \right)$$

Corollary 3.4 (Convergence rate of uncervals). Let Assumptions 1,2 and 3 hold. Also, let G be Lipschitz. Then,

$$\mathbb{P}\left(T_{N+1} \in \widehat{\mathcal{C}}_{(1-\alpha)}^{*}\left(X_{N+1}\right) \mid X_{N+1}\right) = 1 - \alpha + O_{\mathbb{P}}\left(\max\{r_{N-n}^{2/3}, n^{-1/2}\}\right).$$

3.2. Technical considerations of LPBs under right-censoring

Consider a covariate-free setup where only right censoring takes place. Consider survival functions with S(0) = 1 and restricted to $[0, \tau]$, where $\tau < \infty$ is such that neither the survival function of the true times nor the one of

the censoring times is zero coming from the left. Then \hat{S} is uniformly consistent for S over $[0, \tau]$, where \hat{S} is the Kaplan-Meier estimator defined as $\hat{S}(t) = \prod_{i:T_i \leq t} \left(1 - \frac{d_i}{n_i}\right)$, where T_i denote observed events, d_i denote number of deaths that happen at T_i and n_i denote number of individuals at risk just before T_i (units that have been censored are not considered to be at risk), see [38, 39]. The Kaplan-Meier estimator does not reach zero beyond the last observed event time if there are still individuals at risk who have not experienced the event (right-censored). Beyond the last event time, there are no further decreases in the estimator, leading to a tail that does not drop to zero. Using predictive regions of the form $\left\{t: b - \hat{Q}_{\mathcal{I}_2}^* \leq \hat{F}_1(t) \leq b + \hat{Q}_{\mathcal{I}_2}^*\right\}$ can lead to LPBs because there might not exist $0 < t < \tau$ such that $b - \hat{Q}_{\tau_2}^* > \hat{F}_1(t)$, resulting in a confidence interval with an infinite upper bound. This inconsistency arises when the desired regions' image under F_1 is centered around $b \in (0,1)$, such as the median when $b = \frac{1}{2}$. Recall that setting b = 1 when using uncervals produces predictive regions for survival time of the form $\left\{y: \hat{S}_1(y) \leq \hat{Q}^*_{\mathcal{I}_2}\right\}$, ensuring that the definition of the confidence region is not contradicted despite the tail behaviours of the Kaplan-Meier estimator.

4. Simulation studies

To simulate interval-censored responses, we modify the function $\mathtt{simIC_weib}^2$ from the R package icenReg [1] so that we can directly provide the regression surface evaluated on the sampled covariates. Consider the following model for the underlying observations:

$$p\log\left(sT\right) = -r(X) + H\tag{6}$$

where p > 0, s > 0, H follows the standard minimum extreme value type I (*Gumbel*) distribution. $r(\cdot)$ is a function of the covariates. The cdf of T conditional on X = x is

$$F(t, x) = P (T \le t \mid X = x) = P (p \log (sT) \le p \log (st) \mid X = x)$$

= $P (-r(x) + H \le p \log (st))$
= $P (H \le p \log (st) + r(x))$

And therefore we have that

²https://github.com/cran/icenReg/blob/master/R/user_utilities.R

$$S(t, x) = 1 - P (T \le t \mid X = x)$$

= 1 - (1 - exp(- exp(p log (st) + r(x))))
= exp(- exp(p log (st) + r(x)))

We use (4) to create a simulation setup ("oracle") where instead of estimating \hat{F}_1 with a distributional regression algorithm on split 1, we just evaluate 1 - S(t, x) on split 2. The cumulative hazard is

$$H(t,x) = -\log(S(t,x)) = \exp(p\log(st) + r(x)) = (st)^p e^{r(x)}$$

Therefore, Equation 6 hides a Cox model with baseline cumulative hazard $H_0(t) = (st)^p$, corresponding to a Weibull distribution of scale and shape parameters s and p respectively. We could have changed $p \log (sT)$ by any strictly monotonically increasing function (first, we just need to preserve inequalities inside probabilities and second, inverse of these functions is always guaranteed to exist).

4.1. Verfication of asympttic properties

We empirically corroborate what Theorem 3.6 implies: the probability that the new observation T_{N+1} falls within the prediction interval $\widehat{\mathcal{C}}^*_{(1-\alpha)}(X_{N+1})$ is approximately $1-\alpha$ as the sample size grows. The term $o_{\mathbb{P}}(1)$ denotes a small error term that converges to zero in probability as the sample size increases, signifying that the approximation becomes more accurate with larger sample sizes.

In this setup, uncervals is working under e = 1 operating mode, i.e. we uniformly randomize Φ^* over the interval whose upper and lower borders are \hat{F}_1 evaluations on the original intervals.

We prepare four different simulation scenarios, the setup of which is described in Table 1. The underlying true times are simulated according to model 6. These are then censored with a case II interval-censoring mechanism (see https://cran.r-project.org/web/packages/icenReg/index. html). Time between inspections is distributed as runif(min = 0, max = inspectLength). Then, we check the frequency with which the intervals outputted by our algorithm contain the true time-to-event over 100 test points. We average the results across 100 different training sets. This procedure is repeated for six different nominal coverages (0.5, 0.7, 0.8, 0.9, 0.95, 0.99) and four (100, 200, 500, 1000) different sample sizes. For each row in Table 1,

we choose different base estimators for \hat{F}_1 . In the "Linear PH" case, we use $ic_sp(Surv(L, U, type = 'interval2'), model = 'ph')$ from the icenReg package to fit \hat{F}_1 . In the "Non-linear PH" case, we use Interval Censored Recursive Forests [33] with ntree=20, nfold=2. The marginal case is the same as "Linear PH" but with no covariates specified. For the "Oracle" mode, the true survival function, as derived in 4, is directly passed to our approach. The results are visible in Figure 4.

	p	inspections	inspectLength	r(X)
Linear PH	2	5	0.2	$-0.1X_1$
Non-linear PH	2	5	0.5	$ 5X_2 - 0.5 $
No covariates	0	3	0.5	0
Oracle	5	5	0.2	$\sin(\pi X_1) + 2 X_2 - 0.5 + X_3^3$
			TABLE 1	

Note the increased difficulty for the "No covariates" case, where there are only 3 checkup points. Also, in the "Linear PH" case the covariates were simulated with correlation $\rho = 0.1$

4.2. Comparison of different LPBs

We consider three different sample sizes and numbers of covariates. For each sample size and number of covariates, we evaluate three different algorithms for uncertainty quantification. We propose $\alpha_1 = 0.8$, $\alpha_2 = 0.9$, $\alpha_3 = 0.95$. In all cases, we set b = 1 (therefore LPB).

- 1. Checking if $\hat{F}_1(t, X_{\text{new}})$ is greater than α : the LPB has as a lower bound the estimated α quantile (naive quantile).
- 2. Our approach with e = 0, this is, considering evaluations of \hat{F}_1 just on the left borders, thus the trivial extension of [5]. Instead of randomizing $\Phi_i^* \leftarrow \operatorname{runif}\left(\hat{F}_1(L_j, X_j), \hat{F}_1(U_j, X_j)\right)$ we set $\Phi_i^* \leftarrow \hat{F}_1(L_j, X_j)$ so that the finite sample coverage bound holds.
- 3. Our approach with e = * (randomizing between border evaluations).

The results are visible in Tables 2-10.

4.3. Evaluating Conformalized Survival Analysis

In the Conformalized Survival Analysis paper [5], the censoring mechanism is leveraged to mitigate the conservativeness of naive approaches driven by

Rest $quantile$	TABLE 2 ults for 1 and Lev	2 naive vel 1 – α ₁	L	Results e=0 at	TABLE : for unc nd Level	3 cervals, $1-\alpha_1$		Results e=* ar	TABLE 4 for unc nd Level	$\frac{1}{2}$ ervals, $1-lpha_1$	
p = 1	p=3	p = 5	p = 10	p = 1	p=3	p = 5	p = 10	p = 1	p = 3	p = 5	p = 10
20.79 20.26 20.18	$21.08 \\ 20.51 \\ 20.24$	22.1 20.79 20.49	$23.58 \\ 21.73 \\ 21.19$	38.34 38.08 38.05	38.23 38.74 38.63	$39.28 \\ 40.05 \\ 40.32$	$39.44 \\ 40.85 \\ 41.25$	20.67 20.23 20.2	20.64 20.12 19.95	20.81 20.2 20.06	$21.48 \\ 20.44 \\ 20.63$

Res [.] quantile	TABLE ults for a and Lea	5 naive vel 1 – α ₂	2	Results e=0 at	TABLE (for unc nd Level	5 cervals, $1-lpha_2$, .	Results e=* at	TABLE ' for unc nd Level	7 cervals, $1-lpha_2$	
p = 1	p = 3	p = 5	p = 10	p = 1	p=3	p = 5	p = 10	p = 1	p=3	p = 5	p = 10
11.85 10.9	$12.55 \\ 11.51$	$13.87 \\ 12.21$	$15.87 \\ 13.47$	26.06 25.81	$25.69 \\ 26.1$	$25.9 \\ 26.43$	$25.22 \\ 26.95$	$11.76 \\ 10.9$	$12.21 \\ 11.01$	$12.74 \\ 11.7$	$13.84 \\ 12.24$
10.71	10.94	11.69	12.82	25.8	26.07	27.02	27.42	10.58	10.68	11.29	12.31

_	TABLE 8	8			TABLE 9	9		r	Table 1	0	
Rest quantile	ults for a and Lev	naive vel $1 - \alpha_i$	3	e=0 an	for unc nd Level	cervals, $1 - \alpha_3$, .	Results e=* a	for unc nd Level	cervals: $1 - \alpha_3$,
$\frac{1}{p=1}$	p = 3	p = 5	p = 10	p = 1	p = 3	p=5	p = 10	p = 1	p = 3	p = 5	p = 10
7.99	8.79	9.98	11.92	18.35	18.04	17.68	17.33	8.11	8.64	9.21	10.36
$\begin{array}{c} 6.93 \\ 6.5 \end{array}$	$7.46 \\ 6.81$	$8.35 \\ 7.75$	$9.69 \\ 8.96$	$17.97 \\ 17.88$	$17.84 \\ 17.86$	$17.82 \\ 18.43$	$18.17 \\ 18.68$	$\begin{array}{c} 6.93 \\ 6.36 \end{array}$	$7.17 \\ 6.67$	$8.02 \\ 7.47$	$\begin{array}{c} 8.7\\ 8.6\end{array}$
6.5	6.81	7.75	8.96	17.88	17.80	18.43	18.68	0.30	0.07	(.4(8.0



Fig 4: Empirical vs nominal coverage for the setup in Section 4.1. It must be noted that throughout our simulations, uncervals was operating b = 0.5. This means that we are considering intervals whose image by $S(\cdot, x)$ is centered around 0.5.

small censoring times. By introducing a threshold c_0 and focusing on subpopulations where $C \geq c_0$, they address the distributional shift between subpopulations and the whole population. This shift arises because patients with larger censoring times tend to be healthier, leading to different joint and conditional distributions of the variables X, C, and T. The authors suggest using a secondary censored outcome $T \wedge c_0$ and highlight that while there is a covariate shift, this can be adjusted by reweighting the samples through weighted conformal inference [65], allowing for a calibrated lower prediction bound (LPB) on $T \wedge c_0$ and thus on T. This approach reduces the power loss due to censoring, but depends on choosing the threshold hy-

perparameter c_0 . We performed a sensitivity analysis of the **cfsurv** function from the **cfsurvival** package. Performance in terms of empirical coverage depends strongly on the chosen c_0 , as seen in Figure 2. We simulate as in Section 4 but now setting **par_shape=2**, **par_scale=1**, **par_inspections=5**, $n = 2000, X \sim U(0, 2)$ and r(X) = X without left censoring. Under this configuration, right-censoring amounts to approximately 30%. For a grid of c_0 ranging from 0.1 to 1 we launch conformalized survival analysis by setting \tilde{T} to the lower bounds, the event indicator to $1\{U = +\infty\}$ and the censoring times to $1\{U = +\infty\}L + 1\{U < +\infty\}U$. See also in Figure 2 how empirical coverage depends on the hyperparameter c_0 . α was set to 0.1. Also, recall that the methodology in [5] requires observing C even when survival time Tis available.

5. Evaluation and comparision of the models with empirical data

5.1. Systolic Blood Pressure Intervention Trial (SPRINT)

In order to assess the conservative non-asymptotic generalization for intervalcensored data of the right-censored algorithm provided in [5], in comparison with our algorithm, we the sleep time from the NIH's Systolic Blood Pressure Intervention Trial (SPRINT) study [27] as an outcome, in which we manage to recover the censoring time C > 0 as it is needed in [5].

SPRINT [27] is a clinical trial that was conducted with the goal to inform the new blood pressure medication guidelines in the US population by testing the effects that a lower blood pressure target has on reducing heart disease risk. Observational studies had shown that individuals with lower systolic blood pressure (SBP) levels had fewer complications and deaths due to cardiovascular disease (CVD). Building on this observation, SPRINT was designed to test the effects of a lower blood pressure target on reducing heart disease risk. Specifically, SPRINT aimed to compare treating high blood pressure to a target SBP goal of less than 120 mmHg against treating to a goal of less than 140 mmHg.

The "Ambulatory Blood Pressure Monitoring" (ABPM) study, an ancillary study within the SPRINT trial, investigates the effects of different hypertension management strategies on nighttime blood pressure. This research explores whether targeting a clinic systolic blood pressure (SBP) of less than 120 mmHg provided clinical advantages in reducing nighttime BP, a critical

predictor of cardiovascular events, over a standard target of less than 140 mmHg.

In our analysis, we propose examining the relationship between "Time that sleep ended" and "Morning Systolic BP". This relationship is pertinent because morning BP can influence wake-up times due to the physiological stress or discomfort associated with elevated BP levels. Understanding how morning systolic BP affects when individuals wake up can provide valuable insights into the interplay between BP management and sleep quality. By analyzing this connection, we aim to enhance hypertension treatment strategies, ensuring better BP control throughout the night and improving overall health outcomes for patients.

We build a couple of semi-synthetic responses: an interval-censored one employable by our algorithm and a right-censored one that can be fed to [5]. Assume that people in the study misunderstood their indications and did not write down the time when they woke up. However, we have access to the time that was set in their alarms and they remember if they woke up naturally or using the alarm. We craft the interval-censored (L, U) and the right-censored $(\tilde{T}, \text{Event indicator})$ following Table 11.

		\mathbf{L}	U	$ $ $\mathbf{\tilde{T}}$	Event indicator	\mathbf{C}	Т
Natural wake up	(zzzzz)!	(!	(1	!)
Alarm wake up	(zz!)	!	$+\infty$!	0	!)
TABLE 11							

Casuistic that has been followed in order to build the semi-synthetic data. The original interval-censored observations were given by two numbers (and) representing the time when sleep started and ended respectively. ! encodes an artificial independent censoring time that has been simulated according to rexp(ninitial, rate = 0.01), ensuring that the new target amounts to 10% right-censoring approximately.

The final dataset analyzed contains a total of 863 individuals, which we randomly split into 80% train and 20% test in two independent datasets.

On the one hand, we consider the approach in [5] using a Cox model to fit the conditional survival function and the corresponding quantile conformal score and we feed it with its semi-synthetic training data, which are samples from $(X, \tilde{T}, \text{Event indicator}, C)$ where X is MORNING_SYSTOLIC. Since $\tilde{T} \leq T$, any calibrated lower predictive bound on the censored survival time \tilde{T} is also a calibrated LPB on the uncensored survival time T. We try several values of c_0 and the coverage is always 1, a consequence of overconservative behavior.

On the other hand, we challenge our approach using as a base model

ic_sp(Surv(L, U, type = 'interval2'), model = 'ph') to fit \hat{F}_1 . We repeat the procedure B = 100 times by randomizing the train-test split and, within the training set, the split used in the two phases of conformal inference. We obtain a median coverage of 0.8576, indicating that the Cox model might be misspecified in this setting. Next, we try Interval Censored Recursive Forests another B = 100 times and observe a median coverage equal to 0.93.

5.2. NHANES Physical Activity Example

The National Health and Nutrition Examination Survey (NHANES) is a comprehensive program conducted by the National Center for Health Statistics (NCHS), with the primary objective of collecting health and nutrition data from the United States population. In this study, we utilized a subset of n = 2977 patients aged between 50 and 80 years, gathered from NHANES waves 2003-2006. This subset contains high-resolution physical activity information measured by means of accelerometer devices during 3-7 days with a time resolution of one physical activity measure per minute.

From a public health and epidemiological perspective, establishing a statistical association between physical activity and survival in the elderly population is of paramount importance. This analysis can serve as the foundation for promoting healthy habits across the population and designing targeted interventions to mitigate the adverse effects of inactivity and the natural decline in functional capacity that occurs with aging [43, 81]. Furthermore, given that physical inactivity is a modifiable risk factor for various chronic diseases, including heart disease, diabetes, and specific types of cancer, the implementation of effective physical activity policies has the potential to extend the lifespan of this at-risk population, enhance their quality of life, and significantly reduce healthcare costs while alleviating the burden on the medical healthcare systems.

There is a corpus of prior research examining the statistical association between survival and mortality in NHANES cohorts (see for example [35, 44, 44, 52]). However, none of these papers examine the uncertainty quantification problems and only focus on examining the statistical association, for example, in terms of measures of the ROC curve such as the area under the curve (AUC) with a pointwise estimation. A basic preliminary exploratory analysis of the NHANES dataset can be found in Figure 5.

Uncertainty quantification can be determinant in making informed public





Fig 5: Estimated survival functions for four different groups in the NHANES data. Each marginal Turnbull estimator is fed with data from one of the four subsets, created by partitioning the dataset based on whether individuals' age and TLAC are below or above the medians of these covariates. Note that older individuals who engage in sufficient physical activity exhibit survival probabilities comparable to those of younger individuals who lead sedentary lifestyles.

health decisions about the life expectancy of the population. This, in turn, impacts decision-making in multiple settings, such as social policies and telecare in residences—for instance, in rural areas.

Our off-the-shelf algorithm in this case is the following Cox model:

 $h(T \mid Age, TLAC) = h_0(T) \exp(-Age\beta_{Age} - TLAC\beta_{TLAC})$

where $h(\cdot)$ denotes the baseline hazard function, T is patient survival in

months, Age is measured in years and TLAC is a variable related to the physical activity levels of the individual (with higher physical activity levels corresponding to higher TLAC values). We use uncervals with e = *. Table 12 shows the intervals [left, ∞] for the 20th and 80th percentiles of the subsample considered for Age and TLAC, indicating the LPB with a probability of 95 percent. Younger individuals (Age=58) and more active (TLAC=3241) can live at least 124 months with high probability. Conversely, more elderly individuals (Age=73) and less active (TLAC=2263) have a threshold of 45.45 months, about one-third of the time. In intermediate cases, there is an increase of 50 percent for increased TLAC values and 100 percent for reducing the Age to 58 years, indicating a stronger influence of Age compared to TLAC. The results indicate that the uncertainty of survival depends on the patients' characteristics, and informed decisions must take into account public health criteria based on uncertainty. For example, inactive people aged 73 years lose 30 percent survival with respect to the active group.

 TABLE 12

 Intervals whose left boundaries are the estimated LPBs for different values of the covariates Age and TLAC

	$\mathbf{TLAC} = 2263$	$\mathbf{TLAC} = 3241$
Age = 58	$(87.89, +\infty)$	$(124.24, +\infty)$
Age = 73	$(45.45, +\infty)$	$(62.12, +\infty)$

5.2.1. Goodness of fit for conditional survival function for interval-censored data

The goal of this subsection is to illustrate how we can exploit uncervals to evaluate whether using a Cox model in the NHANES case application is well specified via a goodness-of-fit approach. The general idea of extending such a procedure is similar to [17], but in that research the authors focus solely on right-censored data.

Assuming that the random sample \mathcal{D}_N is distributed according to a theoretical conditional cdf F_0 , under the null hypothesis $H_0: F = F_0$ we have $F(X_i, T_i) \sim U(0, 1)$ in virtue of the probability integral transform theorem provided that F_0 is continuous. Therefore, testing the null hypothesis is equivalent to testing $H_0: F \circ F_0^{-1} = F_{\text{unif}}$, where F_{unif} denotes the uniform distribution function on (0, 1). Notice that since we have interval-censored

data, we do not observe the failure times T_i but instead observe (L_i, U_i) . Under non-informative interval-censoring (Assumption 1), the empirical distribution function $\frac{1}{n} \sum_{i=1}^{n} 1\{\Phi_i^* \leq \cdot\}$ converges in distribution to U(0, 1) under the null hypothesis. Recall that

$$\Phi_i^* = \hat{F}_1(L_i, X_i) + U(0, 1) \cdot (\hat{F}_1(U_i, X_i) - \hat{F}_1(L_i, X_i)).$$

A natural application in our NHANES application case arises, which is testing whether the Cox model is well-specified by evaluating the pseudo-random scores $\Phi_i^* \sim U(0, 1)$, $i = 1, \ldots, n$ via a graphical criterion (as shown in Figure 6). In this case, we can see that the method is well-calibrated and there is no evidence that the conditional survival function class is misspecified with a Cox model. A Kolmogorov-Smirnov test would have to be performed in order to conduct a formal hypothesis test.

Users might notice that using the constructed Φ_i^* , based on F_1 as Turnbull's estimator or the recently explained graphical criterion, also seems to be correct and might lead one to believe that it is acceptable to disregard the covariates. This is entirely expected, as Turnbull's estimator is fully nonparametric and, therefore, will naturally produce residuals with the correct distribution. However, what we are testing is whether the parametric specification formulated in the Cox model is correct for the two chosen covariates: Age and TLAC.

6. Discussion

The present paper proposes a new framework for uncertainty quantification in interval-censored data. We introduce novel algorithms along with new empirical process tools to examine the theoretical properties of the methods presented. In certain special cases, the methods introduced here are conformal algorithms that exhibit non-asymptotic guarantees of the form $\mathbb{P}(T_{N+1} \in \widehat{C}_{1-\alpha}(X_{N+1})) \geq 1 - \alpha$. These proposed methods are a natural progression of those proposed by [5] for right-censored data, extending them to intervalcensored data. In this sense, **untervals** can be seen as a trade-off between [5] and [55]. Diverse numerical results demonstrate clear advantages in terms of coverage approximation, especially when compared to [5] and the quantilebased approach using raw conditional survival models without calibration.

For the theoretical analysis, we introduce a new class of functions that allows to derive new asymptotic properties of the uncertainty quantification methods presented.





Fig 6: In blue, empirical cumulative distribution function of $\{\Phi_i^*\}_{i\in\mathcal{I}_2}$ using the NHANES database. The *ic_sp* algorithm (which uses a Cox model) was utilized as the base procedure to compute \hat{F}_1 For comparison, the black line illustrates the theoretical cumulative distribution function of a uniform distribution U(0, 1), which is the identity function.

We illustrate the advantages of our proposal over a naive adaptation for interval-censored data of [5] in a clinical problem related to sleep disorders. Additionally, we highlight the application and scientific interest of these methods in quantifying the impact of physical activity on patient survival. We also introduce the core steps to use the interval-censored methodology to do goodness of fit in this setting.

As future work, we suggest extending the proposed framework to handle truncated or double-truncated censored data (see [12]). Additionally, we propose examining the methods introduced here for multivariate interval data. We aim to develop tolerance regions through resampling techniques by exploiting the interval process and its corresponding universal-Donsker property. To the best of our knowledge, this would be the first general framework for tolerance regions [45, 75] for interval-censored data.

References

- [1] Clifford Anderson-Bergman. icenReg: Regression models for interval censored data in R. Journal of Statistical Software, 81(12):1–23, 2017.
- [2] Anastasios N. Angelopoulos and Stephen Bates. Conformal prediction: A gentle introduction. Foundations and Trends in Machine Learning, 16(4):494-591, 2023. ISSN 1935-8237. URL http://dx.doi.org/10. 1561/2200000101.
- [3] Christopher R. S. Banerji, Tapabrata Chakraborti, Chris Harbron, and Ben D. MacArthur. Clinical ai tools must convey predictive uncertainty for each individual patient. *Nature Medicine*, Oct 2023. ISSN 1546-170X. URL https://doi.org/10.1038/s41591-023-02562-7.
- [4] Stephen Bates, Emmanuel Candès, Lihua Lei, Yaniv Romano, and Matteo Sesia. Testing for outliers with conformal p-values. *The Annals of Statistics*, 51(1):149–178, 2023.
- [5] Emmanuel Candès, Lihua Lei, and Zhimei Ren. Conformalized survival analysis. Journal of the Royal Statistical Society Series B: Statistical Methodology, 85(1):24–45, 01 2023. ISSN 1369-7412. URL https:// doi.org/10.1093/jrsssb/qkac004.
- [6] Jiayi Cheng and Peilin Yang. Conformal inference for heterogeneous policy effect. 2022.
- [7] Victor Chernozhukov, Kaspar Wüthrich, and Yinchu Zhu. An exact and robust conformal inference method for counterfactual and synthetic controls. Journal of the American Statistical Association, 116(536):1849– 1864, 2021.
- [8] Victor Chernozhukov, Kaspar Wüthrich, and Yinchu Zhu. Distributional conformal prediction. *Proceedings of the National Academy of Sciences*, 118(48):e2107794118, 2021.
- [9] Raj S Chhikara and Irwin Guttman. Prediction limits for the inverse gaussian distribution. *Technometrics*, 24(4):319–324, 1982.
- [10] Hunyong Cho, Shannon T Holloway, David J Couper, and Michael R Kosorok. Multi-stage optimal dynamic treatment regimes for survival outcomes with dependent censoring. *Biometrika*, 110(2):395–410, 2023.
- [11] Srinjoy Das, Yiwen Zhang, and Dimitris N Politis. Model-based and model-free point prediction algorithms for locally stationary random fields. arXiv preprint arXiv:2212.03079, 2022.
- [12] Jacobo de Uña-Alvarez. Testing for an ignorable sampling bias under random double truncation. *Statistics in Medicine*, 42(20):3732–3744,

2023. URL https://onlinelibrary.wiley.com/doi/abs/10.1002/ sim.9828.

- [13] Tom Dietterich and Jesse Hostetler. Reinforcement learning prediction intervals with guaranteed fidelity.
- [14] Paromita Dubey, Yaqing Chen, and Hans-Georg Müller. Depth profiles and the geometric exploration of random objects through optimal transport. arXiv preprint arXiv:2202.06117, 2022.
- [15] Robin Dunn, Larry Wasserman, and Aaditya Ramdas. Distributionfree prediction sets for two-layer hierarchical models. *Journal of the American Statistical Association*, pages 1–12, 2022.
- [16] Bradley Efron. The two sample problem with censored data. In Proceedings of the fifth Berkeley symposium on mathematical statistics and probability, volume 4, pages 831–853, 1967.
- [17] Tamara Fernández and Arthur Gretton. A maximum-mean-discrepancy goodness-of-fit test for censored data. In *The 22nd International Conference on Artificial Intelligence and Statistics*, pages 2966–2975. PMLR, 2019.
- [18] Thomas R Fleming and DY Lin. Survival analysis in clinical trials: past developments and future directions. *Biometrics*, 56(4):971–983, 2000.
- [19] Edwin Fong and Chris C Holmes. Conformal bayesian computation. Advances in Neural Information Processing Systems, 34:18268–18279, 2021.
- [20] Matteo Fontana, Gianluca Zeni, and Simone Vantini. Conformal prediction: a unified review of theory and new challenges. *Bernoulli*, 29(1): 1–23, 2023.
- [21] Donald AS Fraser and Irwin Guttman. Tolerance regions. The Annals of Mathematical Statistics, 27(1):162–179, 1956.
- [22] Carlos García-Meixide and Marcos Matabuena. Causal survival embeddings: non-parametric counterfactual inference under censoring. arXiv preprint arXiv:2306.11704, 2023.
- [23] Jakob Gawlikowski, Cedrique Rovile Njieutcheu Tassi, Mohsin Ali, Jongseok Lee, Matthias Humt, Jianxiang Feng, Anna Kruspe, Rudolph Triebel, Peter Jung, Ribana Roscher, et al. A survey of uncertainty in deep neural networks. *Artificial Intelligence Review*, pages 1–77, 2023.
- [24] Gery Geenens, Alicia Nieto-Reyes, and Giacomo Francisci. Statistical depth in abstract metric spaces. *Statistics and Computing*, 33(2): 46, Feb 2023. ISSN 1573-1375. URL https://doi.org/10.1007/s11222-023-10216-4.

- [25] Seymour Geisser. *Predictive inference*. Chapman and Hall/CRC, 2017.
- [26] Robert Gentleman and Charles J. Geyer. Maximum likelihood for interval censored data: Consistency and computation. *Biometrika*, 81(3): 618-623, 09 1994. ISSN 0006-3444. URL https://doi.org/10.1093/biomet/81.3.618.
- [27] The SPRINT Research Group. A randomized trial of intensive versus standard blood-pressure control. New England Journal of Medicine, 373 (22):2103–2116, 2015.
- [28] Yu Gui, Rohan Hore, Zhimei Ren, and Rina Foygel Barber. Conformalized survival analysis with adaptive cut-offs. *Biometrika*, 111(2): 459–477, 2024.
- [29] Michael Hamada, Valen Johnson, Leslie M Moore, and Joanne Wendelberger. Bayesian prediction intervals and their relationship to tolerance intervals. *Technometrics*, 46(4):452–459, 2004.
- [30] Ziad Akram Ali Hammouri, Pablo Rodríguez Mier, Paulo Félix, Mohammad Ali Mansournia, Fernando Huelin, Martí Casals, and Marcos Matabuena. Uncertainty quantification in medicine science: The next big step. Archivos de Bronconeumología, 2023. ISSN 0300-2896. URL https://www.sciencedirect.com/science/article/ pii/S0300289623002338.
- [31] Xiaoyu Hu and Jing Lei. A two-sample conditional distribution test using conformal prediction and weighted rank sum. *Journal of the American Statistical Association*, pages 1–19, 2023.
- [32] Jing Huang. Efficient estimation for the cox model with interval censoring. Annals of Statistics, 24(2):540–568, 1996.
- [33] Nicholas P. Jewell Hunyong Cho and Michael R. Kosorok. Interval censored recursive forests. Journal of Computational and Graphical Statistics, 31(2):390-402, 2022. URL https://doi.org/10.1080/10618600. 2021.1987253.
- [34] Noorie Hyun, Li C Cheung, Qing Pan, Mark Schiffman, and Hormuzd A Katki. Flexible risk prediction models for left or interval-censored data from electronic health records. *The annals of applied statistics*, 11(2): 1063, 2017.
- [35] Ying Jin and Andrew Leroux. Comparing estimators of discriminative performance of time-to-event models. arXiv preprint arXiv:2406.04167, 2024.
- [36] Ying Jin, Zhimei Ren, and Emmanuel J Candès. Sensitivity analysis of individual treatment effects: A robust conformal inference approach.

Proceedings of the National Academy of Sciences, 120(6):e2214889120, 2023.

- [37] Chancellor Johnstone and Bruce Cox. Conformal uncertainty sets for robust optimization. In *Conformal and Probabilistic Prediction and Applications*, pages 72–90. PMLR, 2021.
- [38] Edward L Kaplan and Paul Meier. Nonparametric estimation from incomplete observations. Journal of the American statistical association, 53(282):457–481, 1958.
- [39] Michael R Kosorok. Introduction to empirical processes and semiparametric inference, volume 61. Springer, 2008.
- [40] Michael R Kosorok and Eric B Laber. Precision medicine. Annual review of statistics and its application, 6(1):263–286, 2019.
- [41] Arun Kumar Kuchibhotla. Exchangeability, conformal prediction, and rank tests. *arXiv preprint arXiv:2005.06095*, 2020.
- [42] Jing Lei, Max G'Sell, Alessandro Rinaldo, Ryan J Tibshirani, and Larry Wasserman. Distribution-free predictive inference for regression. *Journal* of the American Statistical Association, 113(523):1094–1111, 2018.
- [43] Michael F. Leitzmann, Yikyung Park, Aaron Blair, Rachel Ballard-Barbash, Traci Mouw, Albert R. Hollenbeck, and Arthur Schatzkin. Physical Activity Recommendations and Decreased Risk of Mortality. *Archives of Internal Medicine*, 167(22):2453–2460, 12 2007. ISSN 0003-9926.
- [44] Andrew Leroux, Erjia Cui, Ekaterina Smirnova, John Muschelli, Jennifer A. Schrack, and Ciprian M. Crainiceanu. Nhanes 2011-2014: Objective physical activity is the strongest predictor of all-cause mortality. *Medicine & Science in Sports & Exercise*, 9900. URL https://journals.lww.com/acsm-msse/fulltext/9900/ nhanes_2011_2014__objective_physical_activity_is.568.aspx.
- [45] Jun Li and Regina Y Liu. Multivariate spacings based on data depth:
 I. construction of nonparametric multivariate tolerance regions. The Annals of Statistics, 36(3):1299–1323, 2008.
- [46] Xun Li, Joyee Ghosh, and Gabriele Villarini. A comparison of bayesian multivariate versus univariate normal regression models for prediction. *The American Statistician*, 0(0):1–9, 2022. URL https://doi.org/10. 1080/00031305.2022.2087735.
- [47] Hang Liu, Xueqin Wang, and Jin Zhu. Quantiles, ranks and signs in metric spaces. arXiv preprint arXiv:2209.04090, 2022.
- [48] Charles Lu, Yaodong Yu, Sai Praneeth Karimireddy, Michael Jordan,

and Ramesh Raskar. Federated conformal predictors for distributed uncertainty quantification. In *International Conference on Machine Learning*, pages 22942–22964. PMLR, 2023.

- [49] Gábor Lugosi and Marcos Matabuena. Uncertainty quantification in metric spaces. arXiv preprint arXiv:2405.05110, 2024.
- [50] Marloes H Maathuis and Jon A Wellner. Inconsistency of the MLE for the joint distribution of interval-censored survival times and continuous marks. *Scand. Stat. Theory Appl.*, 35(1):83–103, March 2008.
- [51] Marcos Matabuena, Paulo Félix, Carlos García-Meixide, and Francisco Gude. Kernel machine learning methods to handle missing responses with complex predictors. application in modelling five-year glucose changes using distributional representations. *Computer Methods* and Programs in Biomedicine, page 106905, 2022.
- [52] Marcos Matabuena, Paulo Félix, Ziad Akram Ali Hammouri, Jorge Mota, and Borja del Pozo Cruz. Physical activity phenotypes and mortality in older adults: a novel distributional data analysis of accelerometry in the nhanes. Aging Clinical and Experimental Research, 34(12): 3107–3114, 2022.
- [53] Carlos García Meixide, Marcos Matabuena, Louis Abraham, and Michael R. Kosorok. Neural interval-censored survival regression with feature selection. *Statistical Analysis and Data Mining: The ASA Data Science Journal*, 17(4):e11704, 2024. URL https://onlinelibrary. wiley.com/doi/abs/10.1002/sam.11704.
- [54] Yash Patel, Declan McNamara, Jackson Loper, Jeffrey Regier, and Ambuj Tewari. Variational inference with coverage guarantees. arXiv preprint arXiv:2305.14275, 2023.
- [55] Dimitris N Politis. Model-free prediction in regression. Springer, 2015.
- [56] Dimitris N Politis and Joseph P Romano. Large sample confidence regions based on subsamples under minimal assumptions. *The Annals* of Statistics, pages 2031–2050, 1994.
- [57] Bodhisattva Sen and Gongjun Xu. Model based bootstrap methods for interval censored data. Computational Statistics and Data Analysis, 81: 121-129, 2015. ISSN 0167-9473. URL https://www.sciencedirect. com/science/article/pii/S0167947314002114.
- [58] Matteo Sesia and Emmanuel J. Candès. A comparison of some conformal quantile regression methods. *Stat*, 9(1):e261, 2020. URL https://onlinelibrary.wiley.com/doi/abs/10.1002/ sta4.261. e261 sta4.261.

- [59] Glenn Shafer and Vladimir Vovk. A tutorial on conformal prediction. Journal of Machine Learning Research, 9(3), 2008.
- [60] Sukrita Singh, Neeraj Sarna, Munich Re, Yuanyuan Li, Yang Lin, Agni Orfanoudaki, and Michael Berger. Distribution-free risk assessment of regression-based machine learning algorithms. arXiv preprint arXiv:2310.03545, 2023.
- [61] Robert A. Stine. Bootstrap prediction intervals for regression. Journal of the American Statistical Association, 80(392):1026–1031, 1985. URL http://doi.org/10.1080/01621459.1985.10478220.
- [62] Sophia Sun. Conformal methods for quantifying uncertainty in spatiotemporal data: A survey. arXiv preprint arXiv:2209.03580, 2022.
- [63] Jiaye Teng, Zeren Tan, and Yang Yuan. T-sci: A two-stage conformal inference algorithm with guaranteed coverage for cox-mlp. In *International Conference on Machine Learning*, pages 10203–10213. PMLR, 2021.
- [64] Lori A Thombs and William R Schucany. Bootstrap prediction intervals for autoregression. Journal of the American Statistical Association, 85 (410):486–492, 1990.
- [65] Ryan J Tibshirani, Rina Foygel Barber, Emmanuel Candes, and Aaditya Ramdas. Conformal prediction under covariate shift. In H. Wallach, H. Larochelle, A. Beygelzimer, F. d'Alché-Buc, E. Fox, and R. Garnett, editors, *Advances in Neural Information Processing Systems*, volume 32. Curran Associates, Inc., 2019. URL https://proceedings.neurips.cc/paper_files/paper/2019/ file/8fb21ee7a2207526da55a679f0332de2-Paper.pdf.
- [66] Bruce W Turnbull. The empirical distribution function with arbitrarily grouped, censored and truncated data. *Journal of the Royal Statistical Society: Series B (Methodological)*, 38(3):290–295, 1976.
- [67] A. W. van der Vaart. Asymptotic Statistics. Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press, 1998.
- [68] Joni Virta. Spatial depth for data in metric spaces. arXiv preprint arXiv:2306.09740, 2023.
- [69] Vladimir Vovk, Alex Gammerman, and Glenn Shafer. Algorithmic Learning in a Random World. Springer Science & Business Media, New York, 2005. ISBN 9780387001524.
- [70] Samuel S. Wilks. Determination of sample sizes for setting tolerance limits. The Annals of Mathematical Statistics, 12(1):91–96, 1941. URL http://doi.org/10.1214/aoms/1177731788.

- [71] Kejin Wu and Dimitris N Politis. Bootstrap prediction inference of nonlinear autoregressive models. arXiv preprint arXiv:2306.04126, 2023.
- [72] Chen Xu and Yao Xie. Conformal prediction interval for dynamic timeseries. In *International Conference on Machine Learning*, pages 11559– 11569. PMLR, 2021.
- [73] Chen Xu and Yao Xie. Conformal prediction for time series. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, 2023.
- [74] Mingzhang Yin, Claudia Shi, Yixin Wang, and David M Blei. Conformal sensitivity analysis for individual treatment effects. *Journal of the American Statistical Association*, pages 1–14, 2022.
- [75] Derek S Young and Thomas Mathew. Nonparametric hyperrectangular tolerance and prediction regions for setting multivariate reference regions in laboratory medicine. *Statistical Methods in Medical Research*, 29(12): 3569–3585, 2020.
- [76] Qiqing Yu, Linxiong Li, and George YC Wong. On consistency of the self-consistent estimator of survival functions with interval-censored data. *Scandinavian Journal of Statistics*, 27(1):35–44, 2000.
- [77] Margaux Zaffran, Aymeric Dieuleveut, Julie Josse, and Yaniv Romano. Conformal prediction with missing values. *arXiv preprint arXiv:2306.02732*, 2023.
- [78] Yunyi Zhang. Regression with complex data: regularization, prediction and bootstrap. University of California, San Diego, 2022.
- [79] Yunyi Zhang and Dimitris N Politis. Bootstrap prediction intervals with asymptotic conditional validity and unconditional guarantees. *Information and Inference: A Journal of the IMA*, 12(1):157–209, 06 2022. ISSN 2049-8772. URL https://doi.org/10.1093/imaiai/iaac017.
- [80] Zhigang Zhang and Jianguo Sun. Interval censoring. Statistical methods in medical research, 19(1):53–70, 2010.
- [81] Min Zhao, Sreenivas P Veeranki, Costan G Magnussen, and Bo Xi. Recommended physical activity and all cause and cause specific mortality in us adults: prospective cohort study. *BMJ*, 370, 2020. URL https://www.bmj.com/content/370/bmj.m2031.

Proof of Theorem 3

The elements in \mathcal{G} are well-defined. For $t \geq u$ the numerator is u - l, which yields the ratio 1. For t < u, the ratio is bounded by 1. First note that

$$g_t(l,u) = \mathbf{1}_{\{0 \le l \le u \le 1\}} \mathbf{1}_{\{u \le t\}} + \mathbf{1}_{\{l \le t < u\}} (t-l) \times \mathbf{1}_{\{0 \le l < u \le 1\}} (u-l)^{-1},$$

and thus $\mathcal{G} \subset \mathcal{F}_1 + \mathcal{F}_2 \cdot \tilde{g}(l, u)$, where $\mathcal{F}_1 = \{1_{\{0 \leq l \leq u \leq 1\}} 1_{\{u \leq t\}} : t \in [0, 1]\}$, $\mathcal{F}_2 = \{1_{\{l \leq t < u\}}(t-l) : t \in [0, 1]\}$, and $\tilde{g}(l, u) = 1_{\{0 \leq l < u \leq 1\}}(u-l)^{-1}$. Standard arguments yield that \mathcal{F}_1 is a Donsker class. We will next show that \mathcal{F}_2 is a VC class with VC index ≤ 3 . This then implies that $\mathcal{F}_3 \equiv \mathcal{F}_2 \cdot \tilde{g}$ is also VC with VC index ≤ 5 by part (vi) of Lemma 9.9 of [39]. We will then show that \mathcal{F}_3 is also pointwise measurable. This, combined with Proposition 8.11 (to establish sufficient measureability), Theorem 9.3 (to establish boundedness of the uniform entropy integral), and Theorem 8.19 of [39], yield that \mathcal{F}_3 is Donsker. Since the sum of two Donsker classes is also Donsker, the desired conclusion follows.

We next prove that \mathcal{F}_2 is VC with VC index ≤ 3 . Let $p_j = (l_j, u_j, c_j) \in \mathbb{R}^3$, for $j = 1, \ldots, 3$, be three distinct points. Also, define \mathcal{F}_2^* as \mathcal{F}_2 but with the permissible range for t expanded to all of \mathbb{R} . Then, since $\mathcal{F}_2 \subset \mathcal{F}_2^*$, showing that the new class is VC with VC index ≤ 3 will imply the same for \mathcal{F}_2 . Define $f_t(l, u) = 1_{\{l \leq t < u\}}(t - l)$, for all $(t, l, u) \in \mathbb{R}^3$. Let $T \subset \mathbb{R}$ be eight distinct points. Our task is to show that it is impossible to find such a T such that all 8 possible subsets of the collection $C = \{p_1, p_2, p_3\}$ can be obtained via sets of the form

$$C \cap \{(l, u, c) \in \mathbb{R}^3 : f_t(l, u) > c\},$$
(7)

Note that if $c_1 < 0$, then p_1 will never be excluded from sets of the form given in 7 since $f_{t_1}(l_1, u_1) \ge 0$ for any possible values of $(t_1, l_1, u_1) \in \mathbb{R}^3$. This is also true for c_2 and c_3 . Thus, we will assume going forward without loss of generality that $c_j \ge 0$ for $1 \le j \le 3$. Similarly, if $u_1 \le l_1$, then $f_{t_1}(l_1, u_1) = 0$ for all $t_1 \in \mathbb{R}$. Hence, we will also assume going forward that $l_j < u_j$ for $1 \le j \le 3$. Next, let $D_t^* = \{(l, u) : 1_{\{l < t < u\}} > 0\}, p_j^* = (l_j, u_j),$ for $1 \le j \le 3$, and $C^* = \{p_1^*, p_2^*, p_3^*\}$; and note that if we can't obtain all possible eight subsets of C^* from sets of the form $C^* \cap D_t^*$, as t ranges over T, then it will be impossible for the sets D_t to shatter C. To see this, note first that $f_t(l, u) > 0$ if and only if $1_{\{l < t < u\}} > 0$, and that $p_1 \in C \cap D_t$ only if $f_t(l_1, u_1) > 0$. Generalizing this, we deduce that $p_j \in C \cap D_t$ only if $p_j^* \in C^* \cap D_t^*, 1 \le j \le 3$. The implication does not go in the other way since

 c_j could be greater than 1, preventing p_j from being in $C \cap D_t$ even if p_j^* is in $C^* \cap D_t^*$. The conclusion of this is that it is impossible for $\{D_t : t \in T\}$ to shatter C if $\{D_t^*, t \in T\}$ does not shatter C^* . Thus, if we can show that $\{D_t^* : t \in T\}$ does not shatter C^* , we have established the desired VC index bound.

Let us first try to shatter $\{p_1^*, p_2^*\}$ with sets of the form D_t^* . To do this, we will need the intervals (l_1, u_1) and (l_2, u_2) to overlap some but also both have non-zero segments which don't overlap. One way to do this is to have $l_1 < l_2 < u_1 < u_2$. The other equivalent possibility happens when the indices 1 and 2 are swapped. Because of this equivalence, we will just use the initial choice of indices. Now let $T^* = \{t_1, \ldots, t_4\}$, where $t_1 < l_1, l_1 < t_2 < l_2, l_2 < d_2$ $t_3 < u_1$, and $u_1 < t_4 < u_2$. Now it is fairly easy to see that D_t^* , as t ranges over T^* , shatters $\{p_1^*, p_2^*\}$. Specifically, D_t^* with $t = t_1$ picks out the null set, t_2 picks out p_1, t_3 picks out $\{p_1, p_2\}$, and t_4 picks out p_2 . We can also see that any other non-equivalent arrangement will not result in shattering. Next we need to see if we can add (l_3, u_3) to this in such a manner that there will exist a $T^* = \{t_1, \ldots, t_8\}$ such that D_t^* , for $t \in T^*$, shatters C^* . To do this, we need (l_3, u_3) to have a portion which does not intersect with either (l_1, u_1) or (l_2, u_2) , another portion which intersects only with (l_1, u_1) , another that intersects only with (l_2, u_2) , and another portion which intersects with both (l_1, u_1) and (l_2, u_2) simultaneously. The impossibility of doing this becomes apparent when visualizing first two intervals above the number line. It is not hard to find an (l_3, u_3) which overlaps each of the non-null subsets, but once we try to have the nterval reach either below l_1 or above u_2 , we can't do this without encompassing at least one of (l_1, u_1) or (l_2, u_2) . Thus it is impossible to find any points $\{p_1^*, p_2^*, p_3^*\}$ and a corresponding T^* which is able to shatter C^* . Hence \mathcal{F}_2 is VC with VC index ≤ 3 .

We now need to prove that \mathcal{F}_2 is pointwise measureable. Let

$$\mathcal{G}_2 = \left\{ 1_{\{l \le t < u\}} (t - l) : t \in [0, 1] \cap \mathbb{Q} \right\},\$$

where \mathbb{Q} is the set of rationals. Fix $t \in [0, 1]$, and let $\{t_n\} \subset [0, 1] \cap \mathbb{Q}$ be a sequence such that $t_n \geq t$ for all $n \geq 1$ and $t_n \to t$, as $n \to \infty$. Then it is easy to verify that $f_{t_n}(l, u) \to f_t(l, u)$ for all $(l, u) \in \mathbb{R}^2$. Since t was arbitrary, we have just shown that every function in \mathcal{F}_2 is the pointwise limit of a sequence of functions in \mathcal{G}_2 . Since \mathcal{G}_2 is a countable set, we have now verified that \mathcal{F}_2 is pointwise measureable.

Last, every element in \mathcal{G} is bounded between zero and one for all $l, u \in [0, 1]$

and therefore \mathcal{G} has a bounded envelope, which is the constant function equal to 1 for all $l, u \in [0, 1]$. The proof is now complete.

Proof of Proposition 3.1.

If F is continuous on t for all x, the statement of the theorem is equivalent to saying: $\mathbb{P}(F(T,X) \leq t \mid L = l, U = u, X = x) = 1\{F(l,x) \leq t \leq F(u,x)\}\frac{t-F(l,x)}{F(u,x)-F(l,x)} + 1\{F(u,x) \leq t\}$. In virtue of Assumption 1, it suffices to show the following

$$\begin{split} \mathbb{P}(F(T,X) \leq t \mid l \leq T \leq u, X = x) \\ = 1\{F(l,x) \leq t \leq F(u,x)\} \frac{t - F(l,x)}{F(u,x) - F(l,x)} + 1\{F(u,x) \leq t\} \end{split}$$

This holds because

$$\begin{split} \mathbb{P}(F(T,X) &\leq t \mid l \leq T \leq u, X = x) = \\ &= \frac{\mathbb{P}(F(T,x) \leq t, l \leq T \leq u \mid X = x)}{\mathbb{P}(l \leq T \leq u \mid X = x)} = \\ &= \frac{\mathbb{P}(F(T,x) \leq t, F(l,x) \leq F(T,x) \leq F(u,x) \mid X = x)}{\mathbb{P}(l \leq T \leq u \mid X = x)} = \\ &= \frac{\mathbb{P}(F(T,x) \leq t, F(l,x) \leq F(T,x) \leq F(u,x) \mid X = x)}{F(u,x) - F(l,x)} = \\ &= 1\{F(l,x) \leq t \leq F(u,x)\}\frac{t - F(l,x)}{F(u,x) - F(l,x)} + 1\{F(u,x) \leq t\} \end{split}$$

7. Proof of Proposition 3.2.

As split 1 and split 2 are independent, the marginal laws equal the conditional ones. Conditional on \hat{F}_1

$$\mathbb{P}(\hat{F}_{1}(T,X) \leq t \mid l \leq T \leq u, X = x) = \frac{\mathbb{P}(\hat{F}_{1}(T,x) \leq t, l \leq T \leq u \mid X = x)}{\mathbb{P}(l \leq T \leq u \mid X = x)}$$

=
$$\frac{\mathbb{P}(\hat{F}_{1}(T,x) \leq t, \hat{F}_{1}(l,x) \leq \hat{F}_{1}(T,x) \leq \hat{F}_{1}(u,x) \mid X = x)}{\mathbb{P}(l \leq T \leq u \mid X = x)} =$$

=
$$\frac{1}{F(u,x) - F(l,x)} 1\{\hat{F}_{1}(l,x) \leq t \leq \hat{F}_{1}(u,x)\} \mathbb{P}(\hat{F}_{1}(l,X) \leq \hat{F}_{1}(T,X) \leq t \mid X = x)$$

+ 1{ $\hat{F}_1(u, x) \leq t$ } Define $F(t, x) = \hat{F}_1(t, x) + r(t, x)$ Formally, the statement

$$\sup_{t \le \tau, x \in \mathbb{R}^p} \left| \hat{S}_1(t, x) - S(t, x) \right| = \mathcal{O}_P(r_N)$$

means that

$$\forall \varepsilon \quad \exists N_{\varepsilon}, \delta_{\varepsilon} \quad \text{ such that } P\left(\sup_{t \leq \tau, x \in \mathbb{R}^{p}} |r(t, x)| \geq \delta_{\varepsilon} r_{N}\right) \leq \varepsilon \quad \forall N > N_{\varepsilon}$$

Focusing on the $\mathbb P$ term

$$\mathbb{P}\left(\hat{F}_{1}(l,X) \leq \hat{F}_{1}(T,X) \leq t \mid X=x\right) =$$

$$= \mathbb{P}\left(\hat{F}_{1}(l,X) + r(T,x) \leq \hat{F}_{1}(T,X) + r(T,x) \leq t + r(T,x) \mid X=x\right) =$$

$$= \mathbb{P}\left(\hat{F}_{1}(l,X) + r(T,x) \leq F(T,x) \leq t + r(T,x) \mid X=x\right) =$$

$$= \mathbb{P}\left(\hat{F}_{1}(l,X) + \delta_{\varepsilon}r_{N} \leq F(T,x) \leq t + \delta_{\varepsilon}r_{N} \mid X=x\right) = t - \hat{F}_{1}(l,x)$$

And therefore we have

$$\mathbb{P}(\hat{F}_1(T,X) \le t \mid l \le T \le u, X = x) = \\ = \frac{1}{F(u,x) - F(l,x)} \mathbb{1}\{\hat{F}_1(l,x) \le t \le \hat{F}_1(u,x)\}(t - \hat{F}_1(l,x)) + \mathbb{1}\{\hat{F}_1(u,x) \le t\}$$

Now using Taylor expansion theorem

$$\frac{1}{\hat{F}_1(u,x) - \hat{F}_1(l,x)} = \frac{1}{F(u,x) - F(l,x)} - \frac{r(l,x) - r(u,x)}{(F(u,x) - F(l,x))^2} + \cdots$$

8. Proof of Lemma 3.4.

Using iterated expectations, LHS is

$$EE\left[1\{\Phi \le \cdot \le \Upsilon\} \mid \Lambda, \Upsilon\right]$$

Because of the assumption, we have that the conditional density function of Φ given Λ, Υ is

$$g(\phi, \lambda, \upsilon) = \frac{1}{\upsilon - \lambda} \mathbb{1}\{\lambda \le \phi \le \upsilon\}$$

We can compute the inner conditional expectation as

$$\begin{split} E\left[1\{\Phi \leq \cdot \leq \Upsilon\} \mid \Lambda, \Upsilon\right] &= E\left[1\{\Phi \leq \cdot \leq \Upsilon\}1\{\Lambda \leq \cdot \leq \Upsilon\} \mid \Lambda, \Upsilon\right] \\ &= \int 1\{\phi \leq \cdot \leq \Upsilon\}1\{\Lambda \leq \cdot \leq \Upsilon\}g(\phi, \Lambda, \Upsilon)d\phi \\ &= 1\{\Lambda \leq \cdot \leq \Upsilon\}\int 1\{\phi \leq \cdot \leq \Upsilon\}\frac{1}{\Upsilon - \Lambda}1\{\Lambda \leq \phi \leq \Upsilon\}d\phi \\ &= 1\{\Lambda \leq \cdot \leq \Upsilon\}\frac{\cdot - \Lambda}{\Upsilon - \Lambda} \end{split}$$

9. Proof of Corollary 3.1.

Just in this proof, we denote by G the theoretical cdf of $\psi(F_i)$. First,

$$|V_{N+1} - \psi(F_{N+1})| = \left| \left| \widehat{F}_1(X_{N+1}, T_{N+1}) - \frac{1}{2} \right| - \left| F(X_{N+1}, T_{N+1}) - \frac{1}{2} \right| \right|$$
$$\leq \left| \widehat{F}_1(X_{N+1}, T_{N+1}) - F(X_{N+1}, T_{N+1}) \right|$$

Now,

$$\left| \widehat{G}(V_{N+1}) - G(\psi(F_{N+1})) \right| = \left| \widehat{G}(V_{N+1}) - G(V_{N+1}) \right| + |G(V_{N+1}) - G(\psi(F_{N+1}))| \le \mathcal{O}_P\left(\max\{\eta_n^{\frac{1}{3}}, n^{-\frac{1}{2}}\} \right) + W |V_{N+1} - \psi(F_{N+1})| = \mathcal{O}_P\left(\max\{r_{N-n}^{2/3}, r_{N-n}, n^{-1/2}\} \right)$$

Then $\left|\widehat{G}(V_{N+1}) - G(\psi(F_{N+1}))\right| = O_P\left(\max\{r_n^{2/3}, n^{-1/2}\}\right)$ as $r_n < r_n^{2/3}$ for $r_n < 1$

Finally, we have

$$P\left(1 - \widehat{G}(V_{N+1}) > O_P\left(n^{-\frac{1}{2}}\right) + \alpha\right) + O_P\left(n^{-\frac{1}{2}}\right)$$

= $P\left(1 - G(\psi(F_{N+1})) > O_P\left(n^{-\frac{1}{2}}\right) + O_P\left(\max\{r_n^{2/3}, n^{-1/2}\}\right) + \alpha\right) + O_P\left(n^{-\frac{1}{2}}\right)$
= $1 - \alpha + O_P\left(\max\{r_n^{2/3}, n^{-1/2}\}\right)$

10. Proof of Corollary 7

Fix $t \in [0, 1]$. On the one hand, we have $1\{\Phi \leq t\} = 1\{\Phi \leq t \leq \Upsilon\} + 1\{\Upsilon \leq t\}$ and therefore $G(t) = P(\Phi \leq t) = E1\{\Phi \leq t\} = E1\{\Phi \leq t \leq \Upsilon\} + E1\{\Upsilon \leq t\}.$

On the other hand, write $g_t(l, u) = 1\{u \leq t\} + 1\{l \leq t < u\}\frac{t-l}{u-l}$ and therefore what we want to show holds iff $E1\{\Upsilon \leq t\} + E\left[1\{\Lambda \leq t < \Upsilon\}\frac{t-\Lambda}{\Upsilon-\Lambda}\right] = E1\{\Phi \leq t \leq \Upsilon\} + E1\{\Upsilon \leq t\}$ iff $E\left[1\{\Lambda \leq t < \Upsilon\}\frac{t-\Lambda}{\Upsilon-\Lambda}\right] = E1\{\Phi \leq t \leq \Upsilon\}$, and the last is true in virtue of Lemma 3.4.

11. Proof of Theorem 11

We denote $P(\cdot) = \mathbb{P}(\cdot \mid (X_i, L_i, U_i) : i \in \mathcal{I}_1)$. Denoting $V_i = \psi(\Phi_i)$,

$$P\left(T_{N+1} \in \widehat{\mathcal{C}}_{(1-\alpha)}^{\text{split},*}(X_{N+1}) \mid \mathbb{I}_n\right) = P\left(\psi(\widehat{F}_1(T_{N+1}, X_{N+1})) \le \widehat{Q}_{\mathcal{I}_2}^* \mid \mathbb{I}_n\right) = P\left(V_{N+1} \le \widehat{Q}_{\mathcal{I}_2}^* \mid \mathbb{I}_n\right)$$
$$P\left(V_{N+1} \le \widehat{Q}_{\mathcal{I}_2}^* \mid \mathbb{I}_n\right) = P\left(\frac{1}{n}\sum_{i=1}^n \mathbb{1}\left\{\widehat{V}_i^* \ge V_{N+1}\right\} > \alpha \mid \mathbb{I}_n\right)$$

Now for all v > 0

$$\hat{V}_i^* \le v \iff \left| \Phi_i^* - \frac{1}{2} \right| < v \iff \Phi_i^* < \frac{1}{2} + v \text{ and } \Phi_i^* > \frac{1}{2} - v$$

Therefore,

$$1\left\{\hat{V}_{i}^{*} \leq v\right\} = 1\left\{\Phi_{i}^{*} < \frac{1}{2} + v\right\} - 1\left\{\Phi_{i}^{*} < \frac{1}{2} - v\right\}$$

So that

$$\frac{1}{n}\sum_{i=1}^{n} 1\left\{\hat{V}_{i}^{*} \leq v\right\} = \mathbb{I}_{n}^{*}\left(\frac{1}{2}+v\right) - \mathbb{I}_{n}^{*}\left(\frac{1}{2}-v\right) =:\mathbb{I}_{n}^{*}\left(v^{+}\right) - \mathbb{I}_{n}^{*}\left(v^{-}\right)$$
$$P\left(v \leq \hat{Q}_{\mathcal{I}_{2}}^{*} \mid \mathbb{I}_{n}\right) = P\left(1 + \mathbb{I}_{n}^{*}\left(v^{-}\right) - \mathbb{I}_{n}^{*}\left(v^{+}\right) > \alpha \mid \mathbb{I}_{n}\right)$$

If we add $\mathbb{I}_n(v^+)$ subtract $1 - \mathbb{I}_n(v^-)$, and multiply by $-\sqrt{n}$ both sides of the inequality we have for v > 0

$$\begin{split} &P\left(v \leq \hat{Q}_{\mathcal{I}_{2}}^{*} \mid \mathbb{I}_{n}\right) \\ &= P\left(\sqrt{n}\left(\mathbb{I}_{n}^{*}\left(v^{+}\right) - \mathbb{I}_{n}\left(v^{+}\right)\right) - \sqrt{n}\left(\mathbb{I}_{n}^{*}\left(v^{-}\right) - \mathbb{I}_{n}\left(v^{-}\right)\right) < \sqrt{n}\left(1 - \alpha + \mathbb{I}_{n}\left(v^{-}\right) - \mathbb{I}_{n}\left(v^{+}\right)\right) - \mathbb{I}_{n}\left(v^{+}\right)\right) + \sqrt{n}\left(\mathbb{I}_{n}^{*}\left(v^{-}\right) - \mathbb{I}_{n}\left(v^{+}\right)\right) \mid \mathbb{I}_{n}\right) \\ &= P\left(\sqrt{n}\left(\mathbb{I}_{n}\left(v^{+}\right) - \mathbb{G}\left(v^{+}\right)\right) < \sqrt{n}\left(1 - \alpha + \mathbb{I}_{n}\left(v^{-}\right) - \mathbb{I}_{n}\left(v^{+}\right)\right) + \sqrt{n}\left(\mathbb{I}_{n}^{*}\left(v^{-}\right) - \mathbb{I}_{n}\left(v^{-}\right)\right) \mid \mathbb{I}_{n}\right) + o_{P}(1) \\ &= P\left(\sqrt{n}\left(\mathbb{I}_{n}\left(v^{+}\right) - G\left(v^{+}\right)\right) - \sqrt{n}\left(1 - \alpha + \mathbb{I}_{n}\left(v^{-}\right) - \mathbb{I}_{n}\left(v^{+}\right)\right) < \sqrt{n}\left(\mathbb{I}_{n}\left(v^{-}\right) - \mathbb{I}_{n}\left(v^{-}\right)\right) \mid \mathbb{I}_{n}\right) + o_{P}(1) \\ &\overset{\text{Cor. 3.3}}{=} P\left(\sqrt{n}\left(\mathbb{I}_{n}\left(v^{+}\right) - G\left(v^{+}\right)\right) - \sqrt{n}\left(1 - \alpha + \mathbb{I}_{n}\left(v^{-}\right) - \mathbb{I}_{n}\left(v^{+}\right)\right) < \sqrt{n}\left(\mathbb{I}_{n}\left(v^{-}\right) - G\left(v^{-}\right)\right)\right) + o_{P}(1) \\ &= P\left(\left(\mathbb{I}_{n}\left(v^{+}\right) - G\left(v^{+}\right)\right) - \left(\mathbb{I}_{n}\left(v^{-}\right) - G\left(v^{-}\right)\right) < 1 - \alpha + \mathbb{I}_{n}\left(v^{-}\right) - \mathbb{I}_{n}\left(v^{+}\right)\right) + o_{P}(1) \\ &= P\left(\left(\mathbb{I}_{n}\left(v^{+}\right) - G\left(v^{-}\right)\right) - \left(\mathbb{I}_{n}\left(v^{-}\right) - \mathbb{I}_{n}\left(v^{+}\right)\right) + o_{P}(1) \\ &= P\left(\mathbb{G}_{n}\left(v^{+}\right) - \mathbb{G}_{n}\left(v^{-}\right) + o_{P}(1) < 1 - \alpha + \mathbb{I}_{n}\left(v^{-}\right) - \mathbb{G}_{n}\left(v^{-}\right) - \mathbb{I}_{n}\left(v^{+}\right) + \mathbb{G}_{n}\left(v^{-}\right)\right) + o_{P}(1) \\ &= P\left(\left(\mathbb{G}_{n}\left(v^{+}\right) - \mathbb{G}_{n}\left(v^{-}\right) < 1 - \alpha + o_{P}(1)\right) + o_{P}(1) \\ &= P\left(\left(\mathbb{I}_{n}\sum_{i=1}^{n} 1\left\{V_{i} \le v\right\} < 1 - \alpha + o_{P}(1)\right) + o_{P}(1) \\ &= P\left(\left(\mathbb{I}_{n}\sum_{i=1}^{n} 1\left\{V_{i} \le v\right\} < 1 - \alpha + o_{P}(1)\right) + o_{P}(1) \\ &= \text{Therefore}, \end{split}$$

$$P\left(V_{N+1} \le \hat{Q}_{\mathcal{I}_2}^* \mid \mathbb{I}_n\right) = P\left(\frac{1}{n}\sum_{i=1}^n 1\left\{V_i \le V_{N+1}\right\} < 1 - \alpha + o_P(1)\right) + o_P(1)$$

The probability in LHS of the previous line involves randomness just due to resampling. In addition, what we are showing in reality is conditional on split 1: both probabilities involved in last statement are conditional on the data indexed by \mathcal{I}_1 Now by taking expectations over the proper training set on both sides of equal sign we arrive to an equality involving $\mathbb P.$ The rest of the proof goes on as in Theorem 2 in [8].