

Vanishing of the quantum reduction of the Deligne exceptional series representations of negative integer level

*Minoru Wakimoto

Abstract

In this paper we show that, for the Deligne exceptional series representations of negative integer level of affine Lie algebras, the quantum Hamiltonian reduction vanishes except for the cases where the nilpotent element is conjugate to $e_{-\theta}$ or $e_{-\theta_s}$.

Contents

1	Introduction	1
2	Vanishing of the numerators of quantized characters	3
3	Example $\sim B_2$ and $K = -1$	5
3.1	Quantum reduction w.r.to $f = e_{-\theta}$	5
3.2	Quantum reduction w.r.to $f = e_{-\theta_s}$	7
4	Example $\sim D_4$	8
4.1	The case $K = -1$	9
4.2	The case $K = -2 \sim$ Denominator identity for $W(\widehat{D}_4, e_{-\theta})$	12

1 Introduction

In this paper we study the vanishing of the space obtained by the quantum Hamiltonian reduction of the Deligne exceptional series representations of negative integer level, as a continuation from [4]. Let \mathfrak{g} be the non-twisted affinization of a finite-dimensional simple Lie algebra $\bar{\mathfrak{g}}$ with a Cartan subalgebra $\bar{\mathfrak{h}}$. Let $\bar{\Delta}$ (resp. $\bar{\Delta}^+$) be the set of all (resp. positive all) roots and \bar{W} be the Weyl group of $(\bar{\mathfrak{g}}, \bar{\mathfrak{h}})$. Let $(\cdot | \cdot)$ be the standard bilinear form defined in [1], and $\bar{\Delta}_\ell$ (resp. $\bar{\Delta}_s$) be the set of long (resp. short) roots. Let θ (resp. θ_s) denote the highest root (resp. highest short root). For a root $\alpha \in \bar{\Delta}$, let $\bar{\mathfrak{g}}_\alpha$ (resp. e_α) denote the root space (resp. root vector) of α . The coroot of $\alpha \in \bar{\Delta}$ is defined by $\alpha^\vee := \frac{2\alpha}{(\alpha|\alpha)}$, and let $\bar{Q}^\vee := \sum_{\alpha \in \bar{\Delta}} \mathbf{Z}\alpha^\vee$ be the coroot lattice. For the basic notations such as δ and Λ_0 and so on, we follow from [1]. The Cartan subalgebra

*12-4 Karato-Rokkoudai, Kita-ku, Kobe 651-1334, Japan, wakimoto.minoru.314@m.kyushu-u.ac.jp, wakimoto@r6.dion.ne.jp

$\mathfrak{h} := \bar{\mathfrak{h}} + \mathbf{C}\Lambda_0 + \mathbf{C}\delta$ of \mathfrak{g} is identified with its dual space \mathfrak{h}^* by the non-degenerate inner product $(\cdot | \cdot)$. Define the coordinates in \mathfrak{h} by

$$(\tau, z, t) := 2\pi i(-\tau\Lambda_0 + z + t\delta)$$

where $\tau \in \mathbf{C}_+ := \{\tau \in \mathbf{C} ; \text{Im}(\tau) > 0\}$ and $z \in \bar{\mathfrak{h}}$ and $t \in \mathbf{C}$. For an element $\lambda \in \mathfrak{h}$, let $\bar{\lambda}$ denote its $\bar{\mathfrak{h}}$ -component. We put $q := e^{2\pi i\tau}$ as usual.

For $\lambda \in \mathfrak{h}^*$ and $\alpha \in \bar{\Delta}_+$, we put

$$\begin{aligned} F_{\lambda+\rho}^{[\alpha]} &:= \frac{1}{2} q^{\frac{|\lambda+\rho|^2}{2(K+h^\vee)}} \sum_{\gamma \in \bar{Q}^\vee} (\alpha|\gamma) t_\gamma(e^{\lambda+\rho}) \\ &= \frac{1}{2} \sum_{\gamma \in \bar{Q}^\vee} (\alpha|\gamma) e^{\bar{\lambda}+\rho+(K+h^\vee)\gamma} q^{\frac{1}{2(K+h^\vee)}|\bar{\lambda}+\rho+(K+h^\vee)\gamma|^2} \end{aligned} \quad (1.1a)$$

$$A_{\lambda+\rho}^{[\alpha]} := \sum_{w \in \bar{W}} \varepsilon(w) w(F_{\lambda+\rho}^{[\alpha]}) \quad (1.1b)$$

where ρ is the Weyl vector of \mathfrak{g} and $h^\vee = (\rho|\delta)$ is the dual Coxeter number, and t_γ is the linear transformation on \mathfrak{h} defined by the formula (6.5.2) in [1].

For Deligne exceptional series representations, in the case when λ satisfies the conditions (i) \sim (iv) in Theorem 4.1 in [3], the numerator of the character of the irreducible \mathfrak{g} -module $L(\lambda)$ is given by the above $A_{\lambda+\rho}^{[\alpha]}$. We consider the quantum Hamiltonian reduction of these modules. Given an sl_2 -triple (x, e, f) with $[x, e] = e$, $[x, f] = -f$, $[e, f] = x$, the numerator of the character of the $W(\mathfrak{g}, x, f)_K$ -module obtained from the quantum reduction of \mathfrak{g} -module $L(\lambda)$ is $A_{\lambda+\rho}^{[\alpha]}(\tau, -\tau x + H, \frac{\tau}{2}|x|^2)$. In [4], the following theorem is proved:

Theorem 1.1. [4] *Assume that (λ, α) is an element in $\mathfrak{h}^* \times \bar{\Delta}_+$ satisfying the condition*

$$\begin{cases} (\lambda + \rho|\delta) &= K + h^\vee \in \mathbf{N} \\ (\lambda + \rho|\delta - \alpha) &= 0 \end{cases} \quad (1.2)$$

and (x, e, f) is an sl_2 -triple. Choose $\beta \in \bar{\Delta}$ such that

$$\begin{cases} (\beta|x) &\in \mathbf{Z} \\ (\beta|\bar{\mathfrak{h}}^f) &= \{0\} \end{cases} \quad (1.3)$$

Then,

1) *the following formula holds for $H \in \bar{\mathfrak{h}}^f$:*

$$A_{\lambda+\rho}^{[\alpha]}(\tau, -\tau x + H, \frac{\tau}{2}|x|^2) = \frac{1}{4} (\beta|x) \sum_{w \in \bar{W}} \varepsilon(w) (\beta^\vee|w\alpha) f_{\lambda,x,w}(\tau, H) \quad (1.4)$$

where

$$f_{\lambda,x,w}(\tau, H) := \sum_{\gamma \in \bar{Q}^\vee} e^{2\pi i(\bar{\lambda}+\rho+(K+h^\vee)\gamma|w^{-1}H)} q^{\frac{1}{2(K+h^\vee)}|\bar{\lambda}+\rho+(K+h^\vee)\gamma-(K+h^\vee)w^{-1}x|^2} \quad (1.5)$$

2) the RHS of (1.4) does not depend on the choice of β satisfying the condition (1.3).

We note that the RHS of (1.5) can be written as follows since $(x|H) = 0$:

$$f_{\lambda,x,w}(\tau, H) = \sum_{\gamma \in \overline{Q}^\vee} e^{2\pi i(\overline{\lambda+\rho}+(K+h^\vee)\gamma-(K+h^\vee)w^{-1}x|w^{-1}H)} q^{\frac{1}{2(K+h^\vee)}|\overline{\lambda+\rho}+(K+h^\vee)\gamma-(K+h^\vee)w^{-1}x|^2} \quad (1.6)$$

In [4], the modular transformation properties of $A_{\lambda+\rho}^{[\alpha]}$ and the functions

$$A_{\lambda+\rho}^{[\alpha](-)}\left(\tau, -\tau x + H, \frac{\tau}{2}|x|^2\right) := \frac{1}{4}(\beta|x) \sum_{w \in \overline{W}} \varepsilon(w) (\beta^\vee|w\alpha) f_{\lambda,x,w}^{(-)}(\tau, H) \quad (1.7a)$$

$$A_{\lambda+\rho}^{[\alpha](*)}\left(\tau, -\tau x + H, \frac{\tau}{2}|x|^2\right) := \frac{1}{4}(\beta|x) \sum_{w \in \overline{W}} \varepsilon(w) (\beta^\vee|w\alpha) f_{\lambda,x,w}^{(*)}(\tau, H) \quad (1.7b)$$

are studied, where

$$\begin{aligned} f_{\lambda,x,w}^{(-)}(\tau, H) &:= \sum_{\gamma \in \overline{Q}^\vee} e^{2\pi i(\overline{\lambda+\rho}+(K+h^\vee)\gamma-(K+h^\vee)w^{-1}x|w^{-1}(H+x))} \\ &\quad \times q^{\frac{1}{2(K+h^\vee)}|\overline{\lambda+\rho}+(K+h^\vee)\gamma-(K+h^\vee)w^{-1}x|^2} \end{aligned} \quad (1.8a)$$

$$\begin{aligned} f_{\lambda,x,w}^{(*)}(\tau, H) &:= q^{-\frac{K+h^\vee}{2}|x|^2} \sum_{\gamma \in \overline{Q}^\vee} e^{2\pi i(\overline{\lambda+\rho}+(K+h^\vee)\gamma-(K+h^\vee)w^{-1}x|w^{-1}(H+x+\tau x))} \\ &\quad \times q^{\frac{1}{2(K+h^\vee)}|\overline{\lambda+\rho}+(K+h^\vee)\gamma-(K+h^\vee)w^{-1}x|^2} \end{aligned} \quad (1.8b)$$

In this paper we will show that the RHS of (1.4) vanishes if the nilpotent element f is of the following form:

$$f = \sum_{j=1}^n e_{-\beta_j} \quad (n \geq 2) \quad (1.9)$$

namely, the surviving cases are only $f = e_{-\theta}$ and $f = e_{-\theta_s}$. In §3.2, we will see that the quantum Hamiltonian reduction with respect to $f = e_{-\theta_s}$ does not necessarily vanish in the case B_2 .

The author is grateful to Professor Victor Kac for recalling my attention to our 2018's work [3] and fruitful collaboration in [4].

2 Vanishing of the numerators of quantized characters

Let us begin this section with noticing a simple fact which can be shown very easily.

Note 2.1. Let $\overline{\mathfrak{g}}$ be a finite-dimensional simple Lie algebra of rank ≥ 2 . Then, for $\alpha, \beta \in \overline{\Delta}$ such that $\alpha \neq \beta$, either $(\alpha|\beta^\vee) \neq 2$ or $(\alpha^\vee|\beta) \neq 2$.

Lemma 2.1. *Let (x, e, f) be an sl_2 -triple such that $f = \sum_{j=1}^n e_{-\beta_j}$ ($n \geq 2$). Take $\beta' \in \{\beta_1, \dots, \beta_n\}$ and put $w_0 := r_{\beta'}$. Then*

- 1) $w_0 x = w_0^{-1} x = x - \beta'^\vee$ and $w_0 H = w_0^{-1} H = H$ for $\forall H \in \bar{\mathfrak{h}}^f$.
- 2) $f_{\lambda, x, w_0 w}(\tau, H) = f_{\lambda, x, w}(\tau, H)$ for $\forall w \in \bar{W}$ and $\forall H \in \bar{\mathfrak{h}}^f$.

Proof. 1) is clear. 2) By (1.6), one has

$$\begin{aligned} & f_{\lambda, x, w_0 w}(\tau, H) \\ = & \sum_{\gamma \in \bar{Q}^\vee} e^{2\pi i(\bar{\lambda} + \rho + (K+h^\vee)\gamma - (K+h^\vee)w^{-1}w_0^{-1}x | w^{-1}w_0^{-1}H)} q^{\frac{1}{2(K+h^\vee)}|\bar{\lambda} + \rho + (K+h^\vee)\gamma - (K+h^\vee)w^{-1}w_0^{-1}x|^2} \\ = & \sum_{\gamma \in \bar{Q}^\vee} e^{2\pi i(\bar{\lambda} + \rho + (K+h^\vee)\gamma - (K+h^\vee)w^{-1}(x - \beta'^\vee) | w^{-1}H)} q^{\frac{1}{2(K+h^\vee)}|\bar{\lambda} + \rho + (K+h^\vee)\gamma - (K+h^\vee)w^{-1}(x - \beta'^\vee)|^2} \end{aligned}$$

Putting $\gamma' := \gamma + w^{-1}\beta'^\vee$, this equation is rewritten as follows:

$$\begin{aligned} & = \sum_{\gamma' \in \bar{Q}^\vee} e^{2\pi i(\bar{\lambda} + \rho + (K+h^\vee)\gamma' - (K+h^\vee)w^{-1}x | w^{-1}H)} q^{\frac{1}{2(K+h^\vee)}|\bar{\lambda} + \rho + (K+h^\vee)\gamma' - (K+h^\vee)w^{-1}x|^2} \\ & = f_{\lambda, x, w}(\tau, H), \end{aligned}$$

proving 2). □

Theorem 2.1. *Let $\bar{\mathfrak{g}}$ be a simple Lie algebra of rank ≥ 2 . Assume that (λ, α) is an element in $\mathfrak{h}^* \times \bar{\Delta}_+$ satisfying the condition (1.2) and $(x, e, f = \sum_{j=1}^n e_{-\beta_j})$ is an sl_2 -triple such that $n \geq 2$. Then*

$$A_{\lambda+\rho}^{[\alpha]} \left(\tau, -\tau x + H, \frac{\tau}{2} |x|^2 \right) = 0$$

Proof. Choose $\beta, \beta' \in \{\beta_1, \dots, \beta_n\}$ such that $\beta \neq \beta'$ and $(\beta^\vee | \beta') \neq 2$, and put $w_0 := r_{\beta'}$. Then, replacing w with $w_0 w$ in (1.4) and using Lemma 2.1 and noticing that $w_0 \beta^\vee = w_0^{-1} \beta^\vee = \beta^\vee - (\beta^\vee | \beta') \beta'^\vee$, one has

$$\begin{aligned} A_{\lambda+\rho}^{[\alpha]} \left(\tau, -\tau x + H, \frac{\tau}{2} |x|^2 \right) &= \frac{1}{4} \sum_{w \in \bar{W}} \varepsilon(w_0 w) (\beta^\vee | w_0 w \alpha) f_{\lambda, w_0 w}(\tau, H) \\ &= -\frac{1}{4} \sum_{w \in \bar{W}} \varepsilon(w) (w_0^{-1} \beta^\vee | w \alpha) f_{\lambda, w}(\tau, H) \\ &= -\frac{1}{4} \sum_{w \in \bar{W}} \varepsilon(w) (\beta^\vee | w \alpha) f_{\lambda, w}(\tau, H) + (\beta^\vee | \beta') \underbrace{\frac{1}{4} \sum_{w \in \bar{W}} \varepsilon(w) (\beta'^\vee | w \alpha) f_{\lambda, w}(\tau, H)}_{A_{\lambda+\rho}^{[\alpha]}(\tau, -\tau x + H, \frac{\tau}{2} |x|^2)} \\ &= [-1 + (\beta^\vee | \beta')] A_{\lambda+\rho}^{[\alpha]} \left(\tau, -\tau x + H, \frac{\tau}{2} |x|^2 \right) \end{aligned}$$

So one has

$$\underbrace{[2 - (\beta^\vee | \beta')]}_{\neq 0} A_{\lambda+\rho}^{[\alpha]} \left(\tau, -\tau x + H, \frac{\tau}{2} |x|^2 \right) = 0,$$

proving Theorem 2.1. \square

The calculation in §3.2 shows that $A_{\lambda+\rho}^{[\alpha]}(\tau, -\tau x + H, \frac{\tau}{2}|x|^2)$ for $\alpha = \theta_s$ does not necessarily vanish. From Theorem 4.1 in [3] and the above Theorem 2.1, we immediately obtain the following:

Corollary 2.1. *Let $\bar{\mathfrak{g}}$ be a simple Lie algebra of rank ≥ 2 . Assume that (λ, α) is an element in $\mathfrak{h}^* \times \bar{\Delta}_+$ satisfying the conditions (i) \sim (iv) in Theorem 4.1 in [3] and $(x, e, f = \sum_{j=1}^n e_{-\beta_j})$ is an sl_2 -triple such that $n \geq 2$. Then the quantum Hamiltonian reduction $H(\lambda)$ of the \mathfrak{g} -module $L(\lambda)$ with respect to this sl_2 -triple vanishes.*

3 Example $\sim B_2$ and $K = -1$

We consider the simple Lie algebra B_2 with the Dynkin diagram $\begin{array}{c} \alpha_1 \quad \alpha_2 \\ \circ \rightleftarrows \circ \end{array}$ and inner product $((\alpha_i | \alpha_j))_{i,j=1,2} = \begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix}$. The Weyl vector of B_2 is $\bar{\rho} = \frac{3}{2}\alpha_1 + 2\alpha_2$ with its square length $|\bar{\rho}|^2 = \frac{5}{2}$ and the dual Coxeter number $h^\vee = 3$. The highest root is $\theta = \alpha_1 + 2\alpha_2$ and the highest short root is $\theta_s = \alpha_1 + \alpha_2$. The coroot lattice $\bar{Q}^\vee = \mathbf{Z}\alpha_1^\vee + \mathbf{Z}\alpha_2^\vee$ is written as

$$\begin{aligned} \bar{Q}^\vee &= \{j\alpha_1 + k(\alpha_1 + 2\alpha_2) \ ; \ j, k \in \mathbf{Z}\} \\ &= \{j(\alpha_1 + \alpha_2) + k\alpha_2 \ ; \ j, k \in \mathbf{Z}, \ j + k \in 2\mathbf{Z}\}. \end{aligned}$$

The fundamental integral forms $\bar{\Lambda}_i$ ($i = 1, 2$) defined by $(\bar{\Lambda}_i | \alpha_j) = \delta_{i,j}$ are related to the simple roots as follows:

$$\begin{cases} \alpha_1 &= 2\bar{\Lambda}_1 - 2\bar{\Lambda}_2 \\ \alpha_2 &= -\bar{\Lambda}_1 + 2\bar{\Lambda}_2 \end{cases} \quad \begin{cases} \bar{\Lambda}_1 &= \alpha_1 + \alpha_2 \\ \bar{\Lambda}_2 &= \frac{1}{2}(\alpha_1 + 2\alpha_2) \end{cases}$$

As usual, for each $\alpha \in \bar{\Delta}$, we choose an element e_α in the root space \mathfrak{g}_α such that $(e_\alpha | e_{-\alpha}) = 1$ namely $[e_\alpha, e_{-\alpha}] = \alpha$.

3.1 Quantum reduction w.r.to $f = e_{-\theta}$

In this section, we consider the quantum Hamiltonian reduction with respect to the sl_2 -triple $(x, e, f) = (\frac{1}{2}\theta, \frac{1}{2}e_\theta, e_{-\theta})$ for which $\bar{\mathfrak{h}}^f = \mathbf{C}\alpha_1$.

First we consider the weight $\lambda = -\Lambda_0$, for which $\lambda + \rho = 2\Lambda_0 + \bar{\rho}$ and $(\lambda + \rho | \delta - \theta) = 0$. Then, by computation letting $\alpha = \beta = \theta$ in (1.4) and (1.7a) and (1.7b), we obtain the following:

$$A_{-\Lambda_0+\rho}^{[\theta]}(\tau, -\tau x + z\alpha_1, \frac{\tau}{4}) = \frac{-i}{2} [\vartheta_{00} + \vartheta_{01}](\tau, 0) \cdot \vartheta_{11}(\tau, 2z) \quad (3.1a)$$

$$A_{-\Lambda_0+\rho}^{[\theta](-)}(\tau, -\tau x + z\alpha_1, \frac{\tau}{4}) = \frac{-i}{2} [\vartheta_{00} + \vartheta_{01}](\tau, 0) \cdot \vartheta_{11}(\tau, 2z) \quad (3.1b)$$

$$A_{-\Lambda_0+\rho}^{[\theta](*)}(\tau, -\tau x + z\alpha_1, \frac{\tau}{4}) = \frac{-i}{2} [\vartheta_{00} - \vartheta_{01}](\tau, 0) \cdot \vartheta_{11}(\tau, 2z) \quad (3.1c)$$

where $\vartheta_{ab}(\tau, z)$ are the Mumford's theta functions ([6]).

Next we consider the weight $\lambda = -\Lambda_0 + \bar{\Lambda}_2$, for which $\lambda + \rho = 2\Lambda_0 + \bar{\Lambda}_1 + 2\bar{\Lambda}_2$ and $(\lambda + \rho | \delta - (\alpha_1 + \alpha_2)) = 0$. Then, by computation letting $\alpha = \alpha_1 + \alpha_2$ and $\beta = \theta$ in (1.4) and (1.7a) and (1.7b), we obtain the following:

$$A_{-\Lambda_0+\bar{\Lambda}_2+\rho}^{[\alpha_1+\alpha_2]}(\tau, -\tau x + z\alpha_1, \frac{\tau}{4}) = -i \vartheta_{10}(\tau, 0) \cdot \vartheta_{11}(\tau, 2z) \quad (3.2)$$

$$\begin{aligned} A_{-\Lambda_0+\bar{\Lambda}_2+\rho}^{[\alpha_1+\alpha_2](-)}(\tau, -\tau x + z\alpha_1, \frac{\tau}{4}) &= A_{-\Lambda_0+\bar{\Lambda}_2+\rho}^{[\alpha_1+\alpha_2](*)}(\tau, -\tau x + z\alpha_1, \frac{\tau}{4}) \\ &= -A_{-\Lambda_0+\bar{\Lambda}_2+\rho}^{[\alpha_1+\alpha_2]}(\tau, -\tau x + z\alpha_1, \frac{\tau}{4}) \end{aligned}$$

The modular transformation of these functions are easily calculated by the transformation properties of ϑ_{ab} as follows:

Note 3.1. Define the functions $\psi_i^{(1)}(\tau, z)$ ($i = 1, 2, 3$) by

$$\begin{cases} \psi_1^{(1)}(\tau, z) & := A_{-\Lambda_0+\rho}^{[\theta]}(\tau, -\frac{\tau}{2}\theta + z\alpha_1, \frac{\tau}{4}) \\ \psi_2^{(1)}(\tau, z) & := A_{-\Lambda_0+\rho}^{[\theta](*)}(\tau, -\frac{\tau}{2}\theta + z\alpha_1, \frac{\tau}{4}) \\ \psi_3^{(1)}(\tau, z) & := A_{-\Lambda_0+\bar{\Lambda}_2+\rho}^{[\alpha_1+\alpha_2]}(\tau, -\frac{\tau}{2}\theta + z\alpha_1, \frac{\tau}{4}) \end{cases}$$

Then

- 1) (i) $\psi_1^{(1)}(\tau, z) + \psi_2^{(1)}(\tau, z) = -i \vartheta_{00}(\tau, 0) \cdot \vartheta_{11}(\tau, 2z)$
(ii) $\psi_1^{(1)}(\tau, z) - \psi_2^{(1)}(\tau, z) = -i \vartheta_{01}(\tau, 0) \cdot \vartheta_{11}(\tau, 2z)$
(iii) $\psi_3^{(1)}(\tau, z) = -i \vartheta_{10}(\tau, 0) \cdot \vartheta_{11}(\tau, 2z)$
- 2) (i) $[\psi_1^{(1)} + \psi_2^{(1)}](-\frac{1}{\tau}, \frac{z}{\tau}) = -\tau e^{\frac{4\pi iz^2}{\tau}} [\psi_1^{(1)} + \psi_2^{(1)}](\tau, z)$
(ii) $[\psi_1^{(1)} - \psi_2^{(1)}](-\frac{1}{\tau}, \frac{z}{\tau}) = -\tau e^{\frac{4\pi iz^2}{\tau}} \psi_3^{(1)}(\tau, z)$
(iii) $\psi_3^{(1)}(-\frac{1}{\tau}, \frac{z}{\tau}) = -\tau e^{\frac{4\pi iz^2}{\tau}} [\psi_1^{(1)} - \psi_2^{(1)}](\tau, z)$
- 3) (i) $[\psi_1^{(1)} + \psi_2^{(1)}](\tau + 1, z) = e^{\frac{\pi i}{4}} [\psi_1^{(1)} - \psi_2^{(1)}](\tau, z)$
(ii) $[\psi_1^{(1)} - \psi_2^{(1)}](\tau + 1, z) = e^{\frac{\pi i}{4}} [\psi_1^{(1)} + \psi_2^{(1)}](\tau, z)$
(iii) $\psi_3^{(1)}(\tau + 1, z) = i \psi_3^{(1)}(\tau, z)$

Remark 3.1. The modular transformation properties of the functions $\psi_i^{(1)}$'s in the above Note 3.1 are quite different from the ones of $A_{\lambda+\rho}^{[\alpha]}$'s in [4]. This is because the condition $(\lambda + \rho|\delta - \alpha) = 0$ is not assumed in the calculation of the modular transformation formulas in [4]. Also these modular transformation properties of the numerators in Note 3.1 are not in consistency with those of the denominators given in [4]. This incoincidence of the modular transformation properties between numerators and denominators seems to suggest that the condition (iv) in Theorem 4.1 in [4] is not satisfied in this case namely in the case B_2 and $K = -1$.

3.2 Quantum reduction w.r.to $f = e_{-\theta_s}$

In this section, we consider the quantum Hamiltonian reduction with respect to the sl_2 -triple $(x, e, f) = (\alpha_1 + \alpha_2, e_{\alpha_1 + \alpha_2}, e_{-(\alpha_1 + \alpha_2)})$ for which $\bar{\mathfrak{h}}^f = \mathbf{C}\alpha_2$.

First we consider the weight $\lambda = -\Lambda_0$, for which $\lambda + \rho = 2\Lambda_0 + \bar{\rho}$ and $(\lambda + \rho|\delta - \theta) = 0$. Then, by computation letting $\alpha = \theta$ and $\beta = \alpha_1 + \alpha_2 = \theta_s$ in (1.4) and (1.7a) and (1.7b), we obtain the following:

$$A_{-\Lambda_0+\rho}^{[\theta]}(\tau, -\tau\theta_s + z\alpha_2, \frac{\tau}{2}) = \sum_{\substack{j, k \in \mathbf{Z} \\ \text{s.t. } j+k \in \mathbf{Z}_{\text{even}}}} [e^{4\pi i(j+\frac{1}{4})z} - e^{-4\pi i(j+\frac{1}{4})z}] q^{(j+\frac{1}{4})^2 + (k+\frac{1}{4})^2} \quad (3.3a)$$

$$A_{-\Lambda_0+\rho}^{[\theta](-)}(\tau, -\tau\theta_s + z\alpha_2, \frac{\tau}{2}) = -A_{\lambda+\rho}^{[\theta]}(\tau, -\tau\theta_s + z\alpha_2, \frac{\tau}{2}) \quad (3.3b)$$

$$A_{-\Lambda_0+\rho}^{[\theta](*)}(\tau, -\tau\theta_s + z\alpha_2, \frac{\tau}{2}) = - \sum_{\substack{j, k \in \mathbf{Z} \\ \text{s.t. } j+k \in \mathbf{Z}_{\text{odd}}}} [e^{4\pi i(j+\frac{1}{4})z} - e^{-4\pi i(j+\frac{1}{4})z}] q^{(j+\frac{1}{4})^2 + (k+\frac{1}{4})^2} \quad (3.3c)$$

Next we consider the weight $\lambda = -\Lambda_0 + \bar{\Lambda}_2$, for which $\lambda + \rho = 2\Lambda_0 + \bar{\Lambda}_2 + 2\bar{\Lambda}_2$ and $(\lambda + \rho|\delta - (\alpha_1 + \alpha_2)) = 0$. Then, by computation letting $\alpha = \beta = \alpha_1 + \alpha_2 = \theta_s$ in (1.4) and (1.7a) and (1.7b), we obtain the following:

$$A_{-\Lambda_0+\bar{\Lambda}_2+\rho}^{[\alpha_1+\alpha_2]}(\tau, -\tau\theta_s + z\alpha_2, \frac{\tau}{2}) = \theta_{0,1}^{(-)}(\tau, 0) \theta_{1,1}^{(-)}(\tau, 2z) \quad (3.4)$$

where $\theta_{j,m}^{(\pm)}(\tau, z) := \sum_{k \in \mathbf{Z}} (\pm 1)^k e^{2\pi i m(k+\frac{j}{2m})z} q^{m(k+\frac{j}{2m})^2}$ is the Jacobi's theta function.

The modular transformation of these functions are easily calculated by the transformation properties of the Jacobi's theta functions as follows:

Note 3.2. Define the functions $\psi_i^{(2)}(\tau, z)$ ($i = 1, 2, 3$) by


$$\begin{cases} \psi_1^{(2)}(\tau, z) & := A_{-\Lambda_0+\rho}^{[\theta]}(\tau, -\tau\theta_s + z\alpha_2, \frac{\tau}{2}) \\ \psi_2^{(2)}(\tau, z) & := A_{-\Lambda_0+\rho}^{[\theta](*)}(\tau, -\tau\theta_s + z\alpha_2, \frac{\tau}{2}) \\ \psi_3^{(2)}(\tau, z) & := A_{-\Lambda_0+\bar{\Lambda}_2+\rho}^{[\alpha_1+\alpha_2]}(\tau, -\tau\theta_s + z\alpha_2, \frac{\tau}{2}) \end{cases}$$

Then

- 1) (i) $\psi_1^{(2)}(\tau, z) + \psi_2^{(2)}(\tau, z) = \theta_{\frac{1}{2},1}^{(-)}(\tau, 0) [\theta_{\frac{1}{2},1}^{(-)} - \theta_{-\frac{1}{2},1}^{(-)}](\tau, 2z)$
- (ii) $\psi_1^{(2)}(\tau, z) - \psi_2^{(2)}(\tau, z) = \theta_{\frac{1}{2},1}^{(+)}(\tau, 0) [\theta_{\frac{1}{2},1}^{(+)} - \theta_{-\frac{1}{2},1}^{(+)}](\tau, 2z)$
- (iii) $\psi_3^{(2)}(\tau, z) = \theta_{0,1}^{(-)}(\tau, 0) \theta_{1,1}^{(-)}(\tau, 2z)$
- 2) (i) $[\psi_1^{(2)} + \psi_2^{(2)}](-\frac{1}{\tau}, \frac{z}{\tau}) = -i\tau e^{\frac{2\pi iz^2}{\tau}} [\psi_1^{(2)} + \psi_2^{(2)}](\tau, z)$
- (ii) $[\psi_1^{(2)} - \psi_2^{(2)}](-\frac{1}{\tau}, \frac{z}{\tau}) = -\tau e^{\frac{2\pi iz^2}{\tau}} \psi_3^{(2)}(\tau, z)$
- (iii) $\psi_3^{(2)}(-\frac{1}{\tau}, \frac{z}{\tau}) = -\tau e^{\frac{2\pi iz^2}{\tau}} [\psi_1^{(2)} - \psi_2^{(2)}](\tau, z)$
- 3) (i) $[\psi_1^{(2)} + \psi_2^{(2)}](\tau + 1, z) = e^{\frac{\pi i}{4}} [\psi_1^{(2)} - \psi_2^{(2)}](\tau, z)$
- (ii) $[\psi_1^{(2)} - \psi_2^{(2)}](\tau + 1, z) = e^{\frac{\pi i}{4}} [\psi_1^{(2)} + \psi_2^{(2)}](\tau, z)$
- (iii) $\psi_3^{(2)}(\tau + 1, z) = i\psi_3^{(2)}(\tau, z)$

Remark 3.2. The modular transformation properties of $\psi_i^{(1)}$'s in Note 3.1 are quite similar with those of $\psi_i^{(2)}$'s in Note 3.2 except for only difference between the factors $e^{\frac{4\pi iz^2}{\tau}}$ and $e^{\frac{2\pi iz^2}{\tau}}$.

4 Example $\sim D_4$



We consider the simple Lie algebra D_4 with the Dynkin diagram

and inner product $((\alpha_i|\alpha_j))_{i,j=1,2,3,4} = \begin{pmatrix} 2 & -1 & 0 & 0 \\ -1 & 2 & -1 & -1 \\ 0 & -1 & 2 & 0 \\ 0 & -1 & 0 & 2 \end{pmatrix}$. The highest root is $\theta = \alpha_1 +$

$2\alpha_2 + \alpha_3 + \alpha_4$. The central charge of the quantum Hamiltonian reduction of a highest weight \widehat{D}_4 -module of level K with respect to the sl_2 -triple (x, e, f) , where $f = e_{-\theta}$ and $x = \frac{1}{2}\theta$, is given by

$$c(K) = -\frac{6(K-1)(K+2)}{K+6}. \quad (4.1)$$

We define the coordinates in the Cartan subalgebra \mathfrak{h} of \widehat{D}_4 by

$$2\pi i \left(-\tau\Lambda_0 + z_1 \frac{\alpha_1}{2} + z_2 \frac{\theta}{2} + z_3 \frac{\alpha_3}{2} + z_4 \frac{\alpha_4}{2} + t\delta \right) = (\tau, z_1, z_2, z_3, z_4, t) \quad (4.2)$$

Then the space $\overline{\mathfrak{h}}^f$ is written as follows:

$$\overline{\mathfrak{h}}^f = \left\{ H = z_1 \frac{\alpha_1}{2} + z_3 \frac{\alpha_3}{2} + z_4 \frac{\alpha_4}{2} \ ; \ z_1, z_2, z_3 \in \mathbf{C} \right\} \quad (4.3)$$

4.1 The case $K = -1$

In this section, we consider the functions $f_{ijk}^{(\pm)}$ and g_{ijk} ($i, j, k \in \{0, 1\}$) defined by

$$\begin{aligned}
f_{000}^{(\pm)}(\tau, z_1, z_3, z_4) &:= \text{ch}_{\Lambda_0}^{(1)} \text{ch}_{\Lambda_0}^{(3)} \text{ch}_{\Lambda_0}^{(4)} \chi_{1,1}^{(5,3)} \pm \text{ch}_{\Lambda_1}^{(1)} \text{ch}_{\Lambda_1}^{(3)} \text{ch}_{\Lambda_1}^{(4)} \chi_{2,1}^{(5,3)} \\
f_{100}^{(\pm)}(\tau, z_1, z_3, z_4) &:= \text{ch}_{\Lambda_0}^{(1)} \text{ch}_{\Lambda_1}^{(3)} \text{ch}_{\Lambda_1}^{(4)} \chi_{1,1}^{(5,3)} \pm \text{ch}_{\Lambda_1}^{(1)} \text{ch}_{\Lambda_0}^{(3)} \text{ch}_{\Lambda_0}^{(4)} \chi_{2,1}^{(5,3)} \\
f_{010}^{(\pm)}(\tau, z_1, z_3, z_4) &:= \text{ch}_{\Lambda_1}^{(1)} \text{ch}_{\Lambda_0}^{(3)} \text{ch}_{\Lambda_1}^{(4)} \chi_{1,1}^{(5,3)} \pm \text{ch}_{\Lambda_0}^{(1)} \text{ch}_{\Lambda_1}^{(3)} \text{ch}_{\Lambda_0}^{(4)} \chi_{2,1}^{(5,3)} \\
f_{001}^{(\pm)}(\tau, z_1, z_3, z_4) &:= \text{ch}_{\Lambda_1}^{(1)} \text{ch}_{\Lambda_1}^{(3)} \text{ch}_{\Lambda_0}^{(4)} \chi_{1,1}^{(5,3)} \pm \text{ch}_{\Lambda_0}^{(1)} \text{ch}_{\Lambda_0}^{(3)} \text{ch}_{\Lambda_1}^{(4)} \chi_{2,1}^{(5,3)} \\
f_{011}^{(\pm)}(\tau, z_1, z_3, z_4) &:= \text{ch}_{\Lambda_1}^{(1)} \text{ch}_{\Lambda_0}^{(3)} \text{ch}_{\Lambda_0}^{(4)} \chi_{2,3}^{(5,3)} \pm \text{ch}_{\Lambda_0}^{(1)} \text{ch}_{\Lambda_1}^{(3)} \text{ch}_{\Lambda_1}^{(4)} \chi_{2,2}^{(5,3)} \\
f_{101}^{(\pm)}(\tau, z_1, z_3, z_4) &:= \text{ch}_{\Lambda_0}^{(1)} \text{ch}_{\Lambda_1}^{(3)} \text{ch}_{\Lambda_0}^{(4)} \chi_{2,3}^{(5,3)} \pm \text{ch}_{\Lambda_1}^{(1)} \text{ch}_{\Lambda_0}^{(3)} \text{ch}_{\Lambda_1}^{(4)} \chi_{2,2}^{(5,3)} \\
f_{110}^{(\pm)}(\tau, z_1, z_3, z_4) &:= \text{ch}_{\Lambda_0}^{(1)} \text{ch}_{\Lambda_0}^{(3)} \text{ch}_{\Lambda_1}^{(4)} \chi_{2,3}^{(5,3)} \pm \text{ch}_{\Lambda_1}^{(1)} \text{ch}_{\Lambda_1}^{(3)} \text{ch}_{\Lambda_0}^{(4)} \chi_{2,2}^{(5,3)} \\
f_{111}^{(\pm)}(\tau, z_1, z_3, z_4) &:= \text{ch}_{\Lambda_1}^{(1)} \text{ch}_{\Lambda_1}^{(3)} \text{ch}_{\Lambda_1}^{(4)} \chi_{2,3}^{(5,3)} \pm \text{ch}_{\Lambda_0}^{(1)} \text{ch}_{\Lambda_0}^{(3)} \text{ch}_{\Lambda_0}^{(4)} \chi_{2,2}^{(5,3)}
\end{aligned}$$

and

$$\begin{aligned}
g_{000}(\tau, z_1, z_3, z_4) &:= \text{ch}_{\Lambda_1}^{(1)} \text{ch}_{\Lambda_1}^{(3)} \text{ch}_{\Lambda_1}^{(4)} \chi_{1,1}^{(5,3)} - \text{ch}_{\Lambda_0}^{(1)} \text{ch}_{\Lambda_0}^{(3)} \text{ch}_{\Lambda_0}^{(4)} \chi_{2,1}^{(5,3)} \\
g_{100}(\tau, z_1, z_3, z_4) &:= \text{ch}_{\Lambda_1}^{(1)} \text{ch}_{\Lambda_0}^{(3)} \text{ch}_{\Lambda_0}^{(4)} \chi_{1,1}^{(5,3)} - \text{ch}_{\Lambda_0}^{(1)} \text{ch}_{\Lambda_1}^{(3)} \text{ch}_{\Lambda_1}^{(4)} \chi_{2,1}^{(5,3)} \\
g_{010}(\tau, z_1, z_3, z_4) &:= \text{ch}_{\Lambda_0}^{(1)} \text{ch}_{\Lambda_1}^{(3)} \text{ch}_{\Lambda_0}^{(4)} \chi_{1,1}^{(5,3)} - \text{ch}_{\Lambda_1}^{(1)} \text{ch}_{\Lambda_0}^{(3)} \text{ch}_{\Lambda_1}^{(4)} \chi_{2,1}^{(5,3)} \\
g_{001}(\tau, z_1, z_3, z_4) &:= \text{ch}_{\Lambda_0}^{(1)} \text{ch}_{\Lambda_0}^{(3)} \text{ch}_{\Lambda_1}^{(4)} \chi_{1,1}^{(5,3)} - \text{ch}_{\Lambda_1}^{(1)} \text{ch}_{\Lambda_1}^{(3)} \text{ch}_{\Lambda_0}^{(4)} \chi_{2,1}^{(5,3)} \\
g_{011}(\tau, z_1, z_3, z_4) &:= \text{ch}_{\Lambda_0}^{(1)} \text{ch}_{\Lambda_1}^{(3)} \text{ch}_{\Lambda_1}^{(4)} \chi_{2,3}^{(5,3)} - \text{ch}_{\Lambda_1}^{(1)} \text{ch}_{\Lambda_0}^{(3)} \text{ch}_{\Lambda_0}^{(4)} \chi_{2,2}^{(5,3)} \\
g_{101}(\tau, z_1, z_3, z_4) &:= \text{ch}_{\Lambda_1}^{(1)} \text{ch}_{\Lambda_0}^{(3)} \text{ch}_{\Lambda_1}^{(4)} \chi_{2,3}^{(5,3)} - \text{ch}_{\Lambda_0}^{(1)} \text{ch}_{\Lambda_1}^{(3)} \text{ch}_{\Lambda_0}^{(4)} \chi_{2,2}^{(5,3)} \\
g_{110}(\tau, z_1, z_3, z_4) &:= \text{ch}_{\Lambda_1}^{(1)} \text{ch}_{\Lambda_1}^{(3)} \text{ch}_{\Lambda_0}^{(4)} \chi_{2,3}^{(5,3)} - \text{ch}_{\Lambda_0}^{(1)} \text{ch}_{\Lambda_0}^{(3)} \text{ch}_{\Lambda_1}^{(4)} \chi_{2,2}^{(5,3)} \\
g_{111}(\tau, z_1, z_3, z_4) &:= \text{ch}_{\Lambda_0}^{(1)} \text{ch}_{\Lambda_0}^{(3)} \text{ch}_{\Lambda_0}^{(4)} \chi_{2,3}^{(5,3)} - \text{ch}_{\Lambda_1}^{(1)} \text{ch}_{\Lambda_1}^{(3)} \text{ch}_{\Lambda_1}^{(4)} \chi_{2,2}^{(5,3)}
\end{aligned}$$

where $\text{ch}_{\lambda}^{(\ell)} := \text{ch}_{L(\lambda; A_1^{(1)})}(\tau, z_{\ell})$ ($\ell = 1, 3, 4$) is the normalized character of the irreducible $A_1^{(1)}$ -module $L(\lambda)$ and $\chi_{r,s}^{(p,q)}$ is the normalized character of the irreducible Virasoro module with the central charge $z^{(p,q)} = 1 - \frac{6(p-q)^2}{pq}$ and the vacuum anomaly $h_{r,s}^{(p,q)} = \frac{(rp - sq)^2 - (p - q)^2}{4pq}$.

The modular transformation properties of these functions are easily computed by using transformation properties of $A_1^{(1)}$ -characters and Virasoro characters given in [7] and [8], and obtained as follows:

Lemma 4.1. We put $a := \frac{1}{\sqrt{5}} \sin \frac{2\pi}{5}$, $b := \frac{1}{\sqrt{5}} \sin \frac{\pi}{5}$ and

$$J := \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & 1 & -1 \\ 1 & -1 & -1 & 1 \end{pmatrix}$$

$$A := aJ = \begin{pmatrix} a & a & a & a \\ a & a & -a & -a \\ a & -a & a & -a \\ a & -a & -a & a \end{pmatrix}, \quad B := bJ = \begin{pmatrix} b & b & b & b \\ b & b & -b & -b \\ b & -b & b & -b \\ b & -b & -b & b \end{pmatrix}$$

Then the S -transformation of $f_{ijk}^{(\pm)}$ and g_{ijk} are as follows:

$$\begin{pmatrix} f_{000}^{(+)}|_S \\ f_{100}^{(+)}|_S \\ f_{010}^{(+)}|_S \\ f_{001}^{(+)}|_S \\ f_{111}^{(+)}|_S \\ f_{011}^{(+)}|_S \\ f_{101}^{(+)}|_S \\ f_{110}^{(+)}|_S \end{pmatrix} = \left(\begin{array}{c|c} A & B \\ \hline -B & A \end{array} \right) \begin{pmatrix} g_{000} \\ g_{100} \\ g_{010} \\ g_{001} \\ g_{111} \\ g_{011} \\ g_{101} \\ g_{110} \end{pmatrix}, \quad \begin{pmatrix} g_{000}|_S \\ g_{100}|_S \\ g_{010}|_S \\ g_{001}|_S \\ g_{111}|_S \\ g_{011}|_S \\ g_{101}|_S \\ g_{110}|_S \end{pmatrix} = \left(\begin{array}{c|c} A & -B \\ \hline B & A \end{array} \right) \begin{pmatrix} f_{000}^{(+)} \\ f_{100}^{(+)} \\ f_{010}^{(+)} \\ f_{001}^{(+)} \\ f_{111}^{(+)} \\ f_{011}^{(+)} \\ f_{101}^{(+)} \\ f_{110}^{(+)} \end{pmatrix}$$

$$\begin{pmatrix} f_{000}^{(-)}|_S \\ f_{100}^{(-)}|_S \\ f_{010}^{(-)}|_S \\ f_{001}^{(-)}|_S \\ f_{111}^{(-)}|_S \\ f_{011}^{(-)}|_S \\ f_{101}^{(-)}|_S \\ f_{110}^{(-)}|_S \end{pmatrix} = \left(\begin{array}{c|c} A & B \\ \hline B & -A \end{array} \right) \begin{pmatrix} f_{000}^{(-)} \\ f_{100}^{(-)} \\ f_{010}^{(-)} \\ f_{001}^{(-)} \\ f_{111}^{(-)} \\ f_{011}^{(-)} \\ f_{101}^{(-)} \\ f_{110}^{(-)} \end{pmatrix}$$

Lemma 4.2. The T -transformation of $f_{ijk}^{(\pm)}$ and g_{ijk} are as follows:

$$f_{ijk}^{(\pm)}|_T = \begin{cases} e^{-\frac{\pi i}{5}} f_{000}^{(\mp)} & \text{if } (i, j, k) = (000) \\ e^{\frac{4\pi i}{5}} f_{ijk}^{(\mp)} & \text{if } (i, j, k) = (100), (010), (001) \\ e^{\frac{\pi i}{5}} f_{ijk}^{(\mp)} & \text{if } (i, j, k) = (011), (101), (110) \\ -e^{\frac{\pi i}{5}} f_{111}^{(\mp)} & \text{if } (i, j, k) = (111) \end{cases}$$

$$g_{ijk}|_T = \begin{cases} -e^{\frac{3\pi i}{10}} g_{000} & \text{if } (i, j, k) = (000) \\ e^{\frac{3\pi i}{10}} g_{ijk} & \text{if } (i, j, k) = (100), (010), (001) \\ e^{\frac{7\pi i}{10}} g_{ijk} & \text{if } (i, j, k) = (011), (101), (110) \\ -e^{\frac{7\pi i}{10}} g_{111} & \text{if } (i, j, k) = (111) \end{cases}$$

The characters of D_4 -Deligne series of level $K = -1$ are written by these functions $f_{ijk}^{(+)}$'s as follows:

Proposition 4.1.

$$\begin{aligned} 1) \quad \text{ch}_{H(-\Lambda_0)}^w &= f_{000}^{(+)} \\ 2) \quad \text{ch}_{H(\Lambda)}^w &= \begin{cases} f_{100}^{(+)} & \text{if } \Lambda = -\Lambda_0 + \bar{\Lambda}_3 + \bar{\Lambda}_4 \\ f_{010}^{(+)} & \text{if } \Lambda = -\Lambda_0 + \bar{\Lambda}_1 + \bar{\Lambda}_4 \\ f_{001}^{(+)} & \text{if } \Lambda = -\Lambda_0 + \bar{\Lambda}_1 + \bar{\Lambda}_3 \end{cases} \\ 3) \quad \text{ch}_{H(\Lambda)}^w &= \begin{cases} f_{011}^{(+)} & \text{if } \Lambda = -\Lambda_0 + \bar{\Lambda}_1 \\ f_{101}^{(+)} & \text{if } \Lambda = -\Lambda_0 + \bar{\Lambda}_3 \\ f_{110}^{(+)} & \text{if } \Lambda = -\Lambda_0 + \bar{\Lambda}_4 \end{cases} \\ 4) \quad \text{ch}_{H(\Lambda)}^w &= f_{111}^{(+)} \quad \text{if } \Lambda = -\Lambda_0 + \bar{\Lambda}_2 \end{aligned}$$

Proof. 1) is due to Kawasetsu's results [5]. For 2) ~ 4), we compute the character of $L(\Lambda)$'s by using Theorem 1.1. Then, letting $\tau \rightarrow \tau + 1$, we get the signed characters of $L(\Lambda)$'s. For these signed characters we compute their polar parts, namely the terms of q with non-negative power, and their modular transformation explicitly. Then, comparing these data with those for $f_{ijk}^{(-)}$'s, we conclude that both coincide, namely we obtain the relation between signed characters and $f_{ijk}^{(-)}$'s. Letting $\tau \rightarrow \tau - 1$ in these formulas, we obtain the relation between characters and $f_{ijk}^{(+)}$'s as desired. \square

The asymptotic behavior of these characters as $\tau \downarrow 0$ is given by the following:

Proposition 4.2.

$$\begin{aligned} 1) \quad (i, j, k) = (000), (100), (010), (001) &\implies f_{ijk}^{(+)}(\tau) \stackrel{\tau \downarrow 0}{\sim} \frac{1}{\sqrt{5}} \sin \frac{\pi}{5} e^{\frac{3}{10} \cdot \frac{\pi i}{\tau}} \\ 2) \quad (i, j, k) = (011), (101), (110), (111) &\implies f_{ijk}^{(+)}(\tau) \stackrel{\tau \downarrow 0}{\sim} \frac{1}{\sqrt{5}} \sin \frac{2\pi}{5} e^{\frac{3}{10} \cdot \frac{\pi i}{\tau}} \end{aligned}$$

Proof. Applying Theorem 4.17 in [7] to $L(\Lambda_i, A_1^{(1)})$, we have

$$\text{ch}_{L(\Lambda_j, A_1^{(1)})}(\tau, 0, 0) \stackrel{\tau \downarrow 0}{\sim} \frac{1}{\sqrt{2}} e^{\frac{\pi i}{12\tau}} \quad (i = 0, 1)$$

And, for the asymptotic behavior of the Virasoro character, we have

$$\begin{cases} \chi_{1,1}^{(5,3)}(\tau) \stackrel{\tau \downarrow 0}{\sim} \sqrt{\frac{2}{5}} \sin \frac{\pi}{5} \cdot e^{\frac{\pi i}{20\tau}} \\ \chi_{2,1}^{(5,3)}(\tau) \stackrel{\tau \downarrow 0}{\sim} \sqrt{\frac{2}{5}} \sin \frac{\pi}{5} \cdot e^{\frac{\pi i}{20\tau}} \end{cases} \quad \begin{cases} \chi_{2,2}^{(5,3)}(\tau) \stackrel{\tau \downarrow 0}{\sim} \sqrt{\frac{2}{5}} \sin \frac{2\pi}{5} \cdot e^{\frac{\pi i}{20\tau}} \\ \chi_{2,3}^{(5,3)}(\tau) \stackrel{\tau \downarrow 0}{\sim} \sqrt{\frac{2}{5}} \sin \frac{2\pi}{5} \cdot e^{\frac{\pi i}{20\tau}} \end{cases}$$

by Theorem 7.1.5 in [8]. Using these and definition of $f_{ijk}^{(+)}$'s, we obtain the asymptotics of $f_{ijk}^{(+)}$'s as desired. \square

4.2 The case $K = -2 \sim$ Denominator identity for $W(\widehat{D}_4, e_{-\theta})$

By (4.1), one has $c(K) = 0$ if $K = -2$, so one may expect there exists the trivial representation of the minimal W-algebra $W(\widehat{D}_4, e_{-\theta})_K$ when $K = -2$, namely one may expect to obtain the denominator identity from the quantum Hamiltonian reduction of the \widehat{D}_4 -module $L(-2\Lambda_0)$. The denominator of the minimal W-algebra $W(\widehat{D}_4, e_{-\theta})$ is given by

$$\begin{aligned} \overset{w}{R}(\tau, H) &= \frac{1}{\eta(\tau)^3} \prod_{i=1,3,4} \vartheta_{11}(\tau, z_i) \cdot \vartheta_{01}\left(\tau, \frac{z_1 + z_3 + z_4}{2}\right) \\ &\times \vartheta_{01}\left(\tau, \frac{-z_1 + z_3 + z_4}{2}\right) \vartheta_{01}\left(\tau, \frac{z_1 - z_3 + z_4}{2}\right) \vartheta_{01}\left(\tau, \frac{z_1 + z_3 - z_4}{2}\right) \end{aligned} \quad (4.4)$$

where $H = \sum_{i=1,3,4} z_i \frac{\alpha_i}{2} \in \bar{\mathfrak{h}}^f$. Then, working out this calculation, we obtain the following:

Proposition 4.3. (Denominator identity of $W(\widehat{D}_4, e_{-\theta})$)

$$\begin{aligned} 1) \quad & \frac{i}{\eta(\tau)^3} \left[\prod_{j=1,3,4} \vartheta_{11}(\tau, z_j) \right] \vartheta_{01}\left(\tau, \frac{z_1 + z_3 + z_4}{2}\right) \vartheta_{01}\left(\tau, \frac{-z_1 + z_3 + z_4}{2}\right) \\ & \times \vartheta_{01}\left(\tau, \frac{z_1 - z_3 + z_4}{2}\right) \vartheta_{01}\left(\tau, \frac{z_1 + z_3 - z_4}{2}\right) \\ & = \sum_{(m_0, m_1, m_3, m_4)} \theta_{m_0,4}(\tau, 0) \prod_{j=1,3,4} [\theta_{m_j,4} - \theta_{-m_j,4}](\tau, z_j) \\ & \quad \cap \\ & \quad \{(4222), (1111), (1133), (3131), (3113)\} \\ & - \sum_{(m_0, m_1, m_3, m_4)} \theta_{m_0,4}(\tau, 0) \prod_{j=1,3,4} [\theta_{m_j,4} - \theta_{-m_j,4}](\tau, z_j) \\ & \quad \cap \\ & \quad \{(0222), (3333), (3311), (1313), (1331)\} \end{aligned} \quad (4.5a)$$

$$\begin{aligned}
2) \quad & \frac{i}{\eta(\tau)^3} \left[\prod_{j=1,3,4} \vartheta_{11}(\tau, z_j) \right] \vartheta_{00}\left(\tau, \frac{z_1 + z_3 + z_4}{2}\right) \vartheta_{00}\left(\tau, \frac{-z_1 + z_3 + z_4}{2}\right) \\
& \quad \times \vartheta_{00}\left(\tau, \frac{z_1 - z_3 + z_4}{2}\right) \vartheta_{00}\left(\tau, \frac{z_1 + z_3 - z_4}{2}\right) \\
= & - \sum_{(m_0, m_1, m_3, m_4)} (-1)^{m_1} \theta_{m_0,4}(\tau, 0) \prod_{j=1,3,4} [\theta_{m_j,4} - \theta_{-m_j,4}](\tau, z_j) \\
& \quad \cap \\
& \quad \{(4222), (1111), (1133), (3131), (3113)\} \\
& + \sum_{(m_0, m_1, m_3, m_4)} (-1)^{m_1} \theta_{m_0,4}(\tau, 0) \prod_{j=1,3,4} [\theta_{m_j,4} - \theta_{-m_j,4}](\tau, z_j) \quad (4.5b) \\
& \quad \cap \\
& \quad \{(0222), (3333), (3311), (1313), (1331)\}
\end{aligned}$$

$$\begin{aligned}
3) \quad & \frac{i}{\eta(\tau)^3} \left[\prod_{j=1,3,4} \vartheta_{11}(\tau, z_j) \right] \vartheta_{11}\left(\tau, \frac{z_1 + z_3 + z_4}{2}\right) \vartheta_{11}\left(\tau, \frac{-z_1 + z_3 + z_4}{2}\right) \\
& \quad \times \vartheta_{11}\left(\tau, \frac{z_1 - z_3 + z_4}{2}\right) \vartheta_{11}\left(\tau, \frac{z_1 + z_3 - z_4}{2}\right) \\
= & - \sum_{(m_0, m_1, m_3, m_4)} \theta_{m_0,4}(\tau, 0) \prod_{j=1,3,4} [\theta_{m_j,4} - \theta_{-m_j,4}](\tau, z_j) \\
& \quad \cap \\
& \quad \{(4222), (1333), (1311), (3313), (3331)\} \\
& + \sum_{(m_0, m_1, m_3, m_4)} \theta_{m_0,4}(\tau, 0) \prod_{j=1,3,4} [\theta_{m_j,4} - \theta_{-m_j,4}](\tau, z_j) \quad (4.5c) \\
& \quad \cap \\
& \quad \{(0222), (3111), (3133), (1131), (1113)\}
\end{aligned}$$

$$\begin{aligned}
4) \quad & \frac{i}{\eta(\tau)^3} \left[\prod_{j=1,3,4} \vartheta_{11}(\tau, z_j) \right] \vartheta_{10}\left(\tau, \frac{z_1 + z_3 + z_4}{2}\right) \vartheta_{10}\left(\tau, \frac{-z_1 + z_3 + z_4}{2}\right) \\
& \quad \times \vartheta_{10}\left(\tau, \frac{z_1 - z_3 + z_4}{2}\right) \vartheta_{10}\left(\tau, \frac{z_1 + z_3 - z_4}{2}\right) \\
= & - \sum_{(m_0, m_1, m_3, m_4)} (-1)^{m_1} \theta_{m_0,4}(\tau, 0) \prod_{j=1,3,4} [\theta_{m_j,4} - \theta_{-m_j,4}](\tau, z_j) \\
& \quad \cap \\
& \quad \{(4222), (1333), (1311), (3313), (3331)\} \\
& + \sum_{(m_0, m_1, m_3, m_4)} (-1)^{m_1} \theta_{m_0,4}(\tau, 0) \prod_{j=1,3,4} [\theta_{m_j,4} - \theta_{-m_j,4}](\tau, z_j) \quad (4.5d) \\
& \quad \cap \\
& \quad \{(0222), (3111), (3133), (1131), (1113)\}
\end{aligned}$$

Proof. We consider $\Lambda = -2\Lambda_0$, which satisfies $(\Lambda + \rho|\delta - \alpha) = 0$ where $\alpha = \sum_{i=1}^4 \alpha_i$. Then, making a very long calculation by letting $\Lambda := -2\Lambda_0$ and $\beta := \theta$ in (1.4) and (1.5), we obtain the formula (4.5a). But we do not know whether $\Lambda = -2\Lambda_0$ satisfies the condition (iv) in

Theorem 4.1 in [3], namely whether the RHS of (1.4) gives the right numerator of the quantum reduction of $L(-2\Lambda_0)$. So, at this stage, the formula (4.5a) is a “candidate” of the denominator identity. In order to confirm that (4.5a) is a right formula, we multiply $1/\eta(\tau)^4$ to both sides of (4.5a) and compute and compare their polar parts and modular transformations. Then we see that they just coincide and the formula (4.5a) is established. 2) follows from 1) by letting $z_i \rightarrow z_i + 1$, and 3) follows from 1) by letting $z_i \rightarrow z_i + \tau$, and 4) follows from 2) by letting $z_i \rightarrow z_i + \tau$. \square

References

- [1] V. G. Kac : Infinite-Dimensional Lie Algebras, 3rd edition, Cambridge University Press, 1990.
- [2] V. G. Kac and M. Wakimoto : Modular and conformal invariance constraints in representation theory of affine algebras, *Advances in Math.* 70 (1988), 156-236.
- [3] V. G. Kac and M. Wakimoto : On characters of irreducible highest weight modules of negative integer level over affine Lie algebras, in *Progress of Mathematics* 326 “Lie Groups, Geometry, and Representation Theory”, pp.235-252, Birkhäuser, 2018.
- [4] V. G. Kac and M. Wakimoto : On modular invariance of quantum affine W-algebras, in preparation.
- [5] Kazuya Kawasetsu : W-algebras with non-admissible levels and the Deligne exceptional series, *International Math. Research Notices*, Vol.2018 No.3, pp.641-676, 2018.
- [6] D. Mumford : *Tata Lectures on Theta I*, *Progress in Math.* 28, Birkhäuser Boston, 1983.
- [7] M. Wakimoto : *Infinite-Dimensional Lie Algebras*, Translation of *Mathematical Monographs* Vol.195, American Mathematical Society 2001.
- [8] M. Wakimoto : *Lectures on Infinite-Dimensional Lie Algebra*, World Scientific, 2001.