

# A REMARK ON THE LINEARIZATION OF NONDEGENERATE NAMBU STRUCTURES OF COORDER 1

FLORIAN ZEISER

ABSTRACT. In this note we show that Nambu structures of coorder 1 can always be linearized if they admit a closed integrable differential form. In particular, we show that a unimodular Poisson structure whose isotropy Lie algebra at a singular point is  $\mathfrak{sl}(2, \mathbb{R})$ , can always be linearized.

Nambu structures were introduced by Nambu [1] and expanded on by Takhtajan [2], in an attempt to generalize Hamiltonian mechanics. A Nambu structure of order  $q$  on a smooth manifold  $M$  is given by a  $q$ -vector field  $\Pi \in \mathfrak{X}^q(M)$  which is preserved by all its Hamiltonian vector fields, i.e.

$$\mathcal{L}_{X_{f_1, \dots, f_{q-1}}} \Pi = 0 \quad \text{where } X_{f_1, \dots, f_{q-1}} := \Pi(df_1, \dots, df_{q-1}, \cdot) \text{ and } f_1, \dots, f_{q-1} \in C^\infty(M).$$

We call  $p := \dim(M) - q$  the coorder of  $\Pi$ . In particular, we recover the definition of a Poisson structure for  $q = 2$ . Hence, they are also known as Nambu-Poisson structures. To describe normal forms of Nambu structures we distinguish the cases  $q = 2$  and  $q \geq 3$ . The case of vector fields, i.e. when  $q = 1$ , is not treated here. For more details on Nambu structures we refer to [3].

### The Poisson case:

Weinstein showed in [4] that, locally around a point, any Poisson structure decomposes as product of a symplectic manifold and a manifold with vanishing Poisson structure. Hence the existence of local normal forms for Poisson structures reduces to the question of linearization, i.e. given a Poisson structure  $\pi$  on  $M$  and  $p \in M$  with  $\pi(p) = 0$ , the isotropy Lie algebra  $\mathfrak{g}_p$  on  $T_p^*M$  is defined by

$$[d_p g, d_p h] := d_p \pi(dg, dh), \quad g, h \in C^\infty(M).$$

Any Lie algebra  $(\mathfrak{g}, [\cdot, \cdot])$  admits an associated Poisson structure  $\pi_{\mathfrak{g}}$  on the dual space  $\mathfrak{g}^*$  defined by

$$\pi_{\mathfrak{g}, \xi}(X, Y) := \xi([X, Y]) \quad \text{where } \xi \in \mathfrak{g}^*, X, Y \in \mathfrak{g}.$$

Applying this construction to the isotropy Lie algebra  $\mathfrak{g}_p$  of  $\pi$  at  $p \in M$ , one obtains the linear Poisson structure  $\pi_{\mathfrak{g}_p}$  on  $\mathfrak{g}_p^* = T_p^*M$ , which plays the role of the first order approximation of  $\pi$  at  $p$ .

**Definition.** A Poisson structure  $\pi$  on  $M$  is called **linearizable** around a zero  $p \in M$  of  $\pi$  if there exists a smooth Poisson diffeomorphism

$$\Phi : (O, \pi) \xrightarrow{\sim} (V, \pi_{\mathfrak{g}_p}), \quad \text{with } \Phi(p) = 0,$$

for some open neighborhoods  $p \in O \subset M$  and  $0 \in V \subset \mathfrak{g}_p^*$ .

The question of linearization has been of interest ever since Weinstein's work (see [5, 6, 7, 8, 3, 9, ?, 10]). Of particular interest for this work is the following:

**Proposition 1** (Weinstein [4]). *There exists a Poisson structure  $\pi$  on  $\mathbb{R}^3$  with  $\pi(0) = 0$  and  $\mathfrak{g}_0 = \mathfrak{sl}_2(\mathbb{R})$  which is not linearizable.*

We refer to [10] for a more detailed outline and references on the linearization problem.

### The Nambu case:

If the order  $q$  of the Nambu structure  $\Pi$  is at least three, then  $\Pi$  induces a singular foliation by leaves of dimension  $q$  and 0 respectively, and  $\Pi$  is decomposable around regular points (Alekseevsky & Guha [11], Gautheron [12], Nakanishi [13]).

Around singular points, i.e.  $p \in M$  with  $\Pi(p) = 0$ , we can ask again the question of linearization. In this case  $\Pi$  gives rise to an  $q$ -ary Lie algebra  $\mathfrak{F}$  (or Filippov algebra [14]) on  $T_p^*M$  by

$$[d_p f_1, \dots, d_p f_q] := d_p \Pi(df_1, \dots, df_q), \quad f_1, \dots, f_q \in C^\infty(M).$$

Contrary to the Lie algebra case, not every  $q$ -ary Lie algebra (for  $q \geq 3$ ) gives rise to a Nambu structure on its dual. However, those induced by Nambu structures do and we obtain a linear Nambu structure  $\Pi_l$  on  $T_p M$  by

$$\Pi_l(\alpha_1, \dots, \alpha_q)(Z) := [\alpha_1, \dots, \alpha_q](Z) \quad \text{where } Z \in T_p M, \alpha_1, \dots, \alpha_q \in T_p^* M.$$

As in the Poisson case we can ask, for a given Nambu structure  $\Pi$  and  $p \in M$  with  $\Pi(p) = 0$ , does there exist a smooth diffeomorphism of Nambu structures

$$\Phi : (O, \Pi) \rightarrow (V, \Pi_l)$$

between open neighborhoods of  $p \in M$  and  $0 \in T_p M$ , respectively?

**Definition.** *If such a  $\Phi$  exists, then we call  $\Pi$  linearizable at  $p \in M$ .*

Linear Nambu structures have been classified by Dufour & Zung [15] and Medeiros [16]. Both make use of a dual notion for Nambu structures due to Dufour & Zung.

**Proposition 2** (Dufour & Zung [15]). *Let  $\Pi \in \mathfrak{X}^q(M)$  and  $\mu \in \Omega^m(M)$  a volume form on  $M$ . Set*

$$\omega := \iota_\Pi \mu.$$

*Then  $\Pi$  is Nambu if and only if  $\omega$  satisfies for all  $\xi \in \mathfrak{X}^{m-q-1}(M)$  the equations*

$$(1) \quad \iota_\xi \omega \wedge \omega = 0 \quad \text{and} \quad \iota_\xi \omega \wedge d\omega = 0.$$

**Definition.** *We call  $\omega \in \Omega^p(M)$  satisfying (1) an integrable differential form of order  $p$ .*

Note that this is coherent with the notion of coorder, i.e. the coorder of a Nambu structure is the order of an associated integrable differential form.

**Remark.** *Integrable differential forms have been studied in the context of first integrals and singular foliations (see [17, 18, 19, 20, 21, 22, 23, 16]).*

By Dufour & Zung [15] and Medeiros [16], every linear Nambu structure  $\Pi_l$  of order  $q \geq 3$  on an  $m$ -dimensional vector space  $V$  belongs to one of the following two types

- **Type 1:**

$$\Pi_l = \sum_{i=1}^{r+1} \pm x_i \partial_{x_1} \wedge \cdots \wedge \widehat{\partial_{x_i}} \wedge \cdots \wedge \partial_{x_{q+1}} + \sum_{i=1}^s \pm x_{q+1+i} \partial_{x_1} \wedge \cdots \wedge \widehat{\partial_{x_{r+i}}} \wedge \cdots \wedge \partial_{x_{q+1}}$$

with  $0 \leq r \leq q+1$  and  $0 \leq s \leq \min(m-q-1, q+1-r)$ ;

- **Type 2:**

$$\Pi_l = \partial_{x_1} \wedge \cdots \wedge \partial_{x_{q-1}} \wedge \left( \sum_{i,j=q}^m b_i^j x_j \partial_{x_i} \right) \quad \text{for } b_i^j \in \mathbb{R}.$$

In this article, we focus on linear Nambu structures of Type 1. We call a linear Nambu structure of Type 1 nondegenerate if  $r = q+1$ . Using the standard volume form  $\mu_{std} = dx_1 \wedge \cdots \wedge dx_m$  on  $\mathbb{R}^m$ , we obtain for the associated integrable differential form the expression

$$(2) \quad \iota_{\Pi_l} \mu_{std} = dx_{q+2} \wedge \cdots \wedge dx_m \wedge df \quad \text{where} \quad f = \frac{1}{2} \left( \sum_{i=1}^l x_i^2 - \sum_{i=l+1}^{q+1} x_i^2 \right)$$

up to permutation. The signature  $(l, q+1-l)$  of  $f$  is, up to permutation, a discrete invariant of  $\Pi$ .

**Theorem 3** (Zung [24]). *Let  $\Pi$  be a Nambu structure on  $M$  of order  $q \geq 3$  and  $p \in M$  such that  $\Pi(p) = 0$ . If the linear part is nondegenerate of Type 1 with signature different from  $(2, \star)$  and  $(\star, 2)$ , then  $\Pi$  is linearizable at  $p$ .*

For Nambu structures whose linear part is nondegenerate of Type 1 of signature  $(2, \star)$  and  $(\star, 2)$  Dufour & Zung [15, Section 5] gave examples of coorder 2 which are not linearizable. For the coorder 1 case there are even earlier counterexamples for linearizability, using Proposition 2.

**Example** ([19]). Consider  $\alpha \in \Omega^1(\mathbb{R}^n)$  given by

$$\alpha := df + \frac{g(f)}{x_1^2 + x_2^2}(x_2 dx_1 - x_1 dx_2)$$

where  $f$  is the function from (2) with  $l = 2$  and  $g \in C^\infty(\mathbb{R})$  with  $g(x) = 0$  for  $x \leq 0$  and  $g(x) > 0$  else. Note that  $\alpha$  satisfies (1) and hence induces a Nambu structure  $\Pi$  of coorder 1 with

$$X = X_{x_3, \dots, x_n} = \pm \left( x_1 \frac{\partial}{\partial x_2} - x_2 \frac{\partial}{\partial x_1} + \frac{l(f)}{x_1^2 + x_2^2} \left( x_1 \frac{\partial}{\partial x_1} + x_2 \frac{\partial}{\partial x_2} \right) \right)$$

The Hamiltonian vector field  $X$  admits flow lines spiraling towards the cone  $C := \{f = 0\}$ , hence inducing holonomy at the leaf  $C \setminus \{0\}$  of the regular foliation on  $\mathbb{R}^n \setminus \{0\}$ . Since the foliation induced by the linear Nambu structure does not have holonomy, there can not exist a linearization. For  $n = 3$ , under a duality as in Proposition 2, this corresponds to Weinstein's example to show Proposition 1.

Here we show that Nambu structures which admit a closed integrable 1-form are always linearizable.

**Main Theorem.** Let  $\Pi$  be a Nambu structure on  $M^m$  of order  $m - 1$  with  $m \geq 3$  and  $p \in M$  such that  $\Pi(p) = 0$ . Assume the linearization  $\Pi_l$  is nondegenerate of type 1 and there exists a volume form  $\mu$  around  $p \in M$  such that

$$(3) \quad d\iota_{\Pi}\mu = 0.$$

Then  $\Pi$  is linearizable at  $p$ .

The Main theorem, along with Theorem 3, leads to the question:

**Question 1.** Let  $\Pi$  be a Nambu structure on  $M$  of order  $q \geq 3$  and  $p \in M$  such that  $\Pi(p) = 0$ . Assume the linear part is nondegenerate of Type 1 with signature  $(2, \star)$  or  $(\star, 2)$  and there exists a volume form  $\mu$  around  $p \in M$  such that

$$d\iota_{\Pi}\mu = 0.$$

Is  $\Pi$  always linearizable at  $p$  in this case?

For the Poisson case, the condition (3) is well-known. Recall from [25] that a Poisson structure  $\pi$  on an orientable manifold  $M^m$  is called unimodular if there exists a volume form  $\mu \in \Omega^m(M)$  with

$$d\iota_{\pi}\mu = 0.$$

Hence for  $m = 3$  the Main Theorem above yields:

**Corollary.** Let  $\pi$  be Poisson structure on  $M^3$  which is unimodular locally around a zero  $p \in M$  with  $\mathfrak{g}_p \simeq \mathfrak{sl}_2(\mathbb{R})$ . Then  $\pi$  is linearizable around  $p$ .

Taking into account the isomorphism  $\mathfrak{sl}_2(\mathbb{R}) \simeq \mathfrak{su}(1, 1)$  and the examples of non-linearizable Poisson structures for semisimple Lie algebras of real rank 1 with non-semisimple compact part by Monnier & Zung [15][Theorem 4.3.9] one is inclined to ask:

**Question 2.** Let  $\pi$  be a Poisson structure on  $M$  and  $p \in M$  such that  $\pi(p) = 0$ . Assume the isotropy Lie algebra  $\mathfrak{g}_p$  is semisimple of real rank 1 with non-semisimple compact part. If  $\pi$  is unimodular around  $p$ , is  $\pi$  linearizable at  $p$ ?

The proof of the Main Theorem is obtained in two steps. We first linearize the singular foliation using an isochore Morse Lemma due to Colin de Verdière & Vey [26]. Then we use a Moser argument to linearize the associated Nambu structure. We want to point out that the linearization of the associated singular foliation was already shown by Moussu [19, Theorem C] under the condition that the foliation is without holonomy.

**Acknowledgements.** I would like to thank Ioan Marcut for making me aware of the reference [26].

**Proof of the Main Theorem.** Let  $\mu \in \Omega^n(\mathbb{R}^n)$  be a volume form such that

$$d\iota_{\Pi}\mu = 0.$$

Since  $\mathbb{R}^n$  has trivial first de-Rham cohomology, there exists a function  $g \in C^\infty(\mathbb{R}^n)$  such that

$$dg = \iota_{\Pi}\mu \quad \text{and} \quad g(0) = 0.$$

Note that  $\Pi$  is uniquely determined by the pair  $(g, \mu)$ . By assumption, the second order Taylor polynomial of  $g$  gives, up to a linear change of coordinates, rise to a non-degenerate quadratic form  $f$  on  $\mathbb{R}^n$  as described in (2) for  $q + 1 = n$ .

**Theorem ([26]).** *There exist open neighborhoods  $O, O' \subset \mathbb{R}^n$  and  $U \subset \mathbb{R}$  of the origins respectively, a smooth function  $h \in C^\infty(U)$  and a diffeomorphism  $\phi : O \rightarrow O'$  such that*

$$(f, h(f)\mu_{std}) = \phi^*(g, \mu).$$

Since  $\Pi$  is uniquely determined by the pair  $(f, h(f)\mu_{std})$  we have

$$\Pi = k(f)\Pi_l$$

where  $k(f) = h(f)^{-1}$ , and due to the condition on the first jet we have  $k(0) = 1$ . We finish the proof using a Moser argument. For  $t \in [0, 1]$  consider the family of  $(n - 1)$ -vector fields given by

$$\Pi_t = t\Pi_l + (1 - t)\Pi_l = (1 + t(k(f) - 1))\Pi_l.$$

Note that this is a family of Nambu structures since  $[\Pi_l, f] = 0$ . Denote by  $E$  be the Euler vector field on  $\mathbb{R}^n$  and  $r \in C^\infty(\mathbb{R} \times [0, 1])$ . We make the Ansatz:

$$\Phi_t^*(\Pi_t) = \Pi_l,$$

where  $\Phi_t$  is the time- $t$  flow with respect to the vector field  $r_t(f)E$ . From the Ansatz we obtain

$$0 = \mathcal{L}_{r_t(f)E}\Pi_t + \partial_t\Pi_t = \left( r_t(f) \left( t(2f\partial_x(k)(f) - (n - 2)(k(f) - 1)) - (n - 2) \right) + (k(f) - 1) \right) \Pi_l.$$

Sufficiently close to the origin in  $\mathbb{R}^n$  the function

$$r_t(f) := \frac{k(f) - 1}{(n - 2)(1 + t(1 - k(f))) - 2tf\partial_x(k)(f)}$$

is well-defined for all  $t \in [0, 1]$ . Hence the flow  $\Phi_t$  of  $r_t(f)E$  is well defined for all  $t \in [0, 1]$  sufficiently close to the origin and satisfies

$$\Phi_t^*\Pi_t = \Pi_l.$$

Therefore  $\Phi_1$  maps  $\Pi$  to  $\Pi_l$ .

## REFERENCES

- [1] Y. Nambu. Generalized Hamiltonian dynamics. *Phys. Rev. D* (3), 7:2405–2412, 1973.
- [2] L. Takhtajan. On foundation of the generalized Nambu mechanics. *Comm. Math. Phys.*, 160(2):295–315, 1994.
- [3] J.-P. Dufour and N. T. Zung. *Poisson structures and their normal forms*, volume 242 of *Progress in Mathematics*. Birkhäuser Verlag, Basel, 2005.
- [4] A. Weinstein. The local structure of Poisson manifolds. *J. Differential Geom.*, 18(3):523–557, 1983.
- [5] J. F. Conn. Normal forms for smooth Poisson structures. *Ann. of Math. (2)*, 121(3):565–593, 1985.
- [6] P. Dazord. Stabilité et linéarisation dans les variétés de Poisson. In *Symplectic geometry and mechanics (Balaruc, 1983)*, Travaux en Cours, pages 59–75. Hermann, Paris, 1985.
- [7] Alan Weinstein. Poisson geometry of the principal series and nonlinearizable structures. *J. Differential Geom.*, 25(1):55–73, 1987.
- [8] P. Monnier and N. T. Zung. Levi decomposition for smooth Poisson structures. *J. Differential Geom.*, 68(2):347–395, 2004.
- [9] M. Crainic and R. L. Fernandes. A geometric approach to Conn’s linearization theorem. *Ann. of Math. (2)*, 173(2):1121–1139, 2011.
- [10] I. Mărcuț and F. Zeiser. The poisson linearization problem for  $\mathfrak{sl}_2(\mathbb{C})$ . part i: Poisson cohomology, 2022.
- [11] D. Alekseevsky and P. Guha. On decomposability of Nambu-Poisson tensor. *Acta Math. Univ. Comenian. (N.S.)*, 65(1):1–9, 1996.
- [12] P. Gautheron. Some remarks concerning Nambu mechanics. *Lett. Math. Phys.*, 37(1):103–116, 1996.

- [13] N. Nakanishi. On Nambu-Poisson manifolds. *Rev. Math. Phys.*, 10(4):499–510, 1998.
- [14] V. T. Filippov.  $n$ -Lie algebras. *Sibirsk. Mat. Zh.*, 26(6):126–140, 191, 1985.
- [15] J.-P. Dufour and N. T. Zung. Linearization of Nambu structures. *Compositio Math.*, 117(1):77–98, 1999.
- [16] A. S. de Medeiros. Singular foliations and differential  $p$ -forms. *Ann. Fac. Sci. Toulouse Math. (6)*, 9(3):451–466, 2000.
- [17] W.-T. Wu and G. Reeb. *Sur les espaces fibrés et les variétés feuilletées*. Institut de mathématique (Strasbourg), 1952.
- [18] I. Kupka. The singularities of integrable structurally stable Pfaffian forms. *Proc. Nat. Acad. Sci. U.S.A.*, 52:1431–1432, 1964.
- [19] R. Moussu. Existence d'intégrales premières pour un germe de forme de Pfaff  $C^\infty$ , non dégénéré. *Bol. Soc. Brasil. Mat.*, 7(2):111–120, 1976.
- [20] R. Moussu. Sur l'existence d'intégrales premières pour un germe de forme de Pfaff. *Ann. Inst. Fourier (Grenoble)*, 26(2):xi, 171–220, 1976.
- [21] A. S. de Medeiros. Structural stability of integrable differential forms. In *Geometry and topology (Proc. III Latin Amer. School of Math., Inst. Mat. Pura Aplicada CNPq, Rio de Janeiro, 1976)*, volume Vol. 597 of *Lecture Notes in Math.*, pages 395–428. Springer, Berlin-New York, 1977.
- [22] C. Camacho. Structural stability theorems for integrable differential forms on 3-manifolds. *Topology*, 17(2):143–155, 1978.
- [23] R. Moussu. Classification  $C^\infty$  des équations de Pfaff complètement intégrables à singularité isolée. *Invent. Math.*, 73(3):419–436, 1983.
- [24] N. T. Zung. New results on the linearization of Nambu structures. *J. Math. Pures Appl. (9)*, 99(2):211–218, 2013.
- [25] A. Weinstein. The modular automorphism group of a Poisson manifold. *J. Geom. Phys.*, 23(3-4):379–394, 1997.
- [26] Y. Colin de Verdière and J. Vey. Le lemme de Morse isochore. *Topology*, 18(4):283–293, 1979.

Florian Zeiser,  
 UNIVERSITY OF ILLINOIS AT URBANA-CHAMPAIGN, 61801 URBANA, UNITED STATES  
*E-mail address:* `fzeiser@illinois.edu`