

# CONTINUITY METHOD FOR THE MABUCHI SOLITON ON THE EXTREMAL FANO MANIFOLDS

TOMOYUKI HISAMOTO AND SATOSHI NAKAMURA

**ABSTRACT.** We run the continuity method for Mabuchi's generalization of Kähler-Einstein metrics, assuming the existence of an extremal Kähler metric. It gives an analytic proof (without minimal model program) of the recent existence result obtained by Apostolov, Lahdili and Nitta. Our key observation is the boundedness of the energy functionals along the continuity method. The same argument can be applied to general  $g$ -solitons and  $g$ -extremal metrics.

## CONTENTS

1. Introduction	1
2. Preliminary	3
3. Continuity method for $g$ -solitons	8
4. Estimate of $M_g$ along the continuity method	10
5. The closeness of $\mathcal{T}$	13
References	17

## 1. INTRODUCTION

The celebrated work of Chen-Donaldson-Sun [14] and Tian [34] states that a Fano manifold admits a Kähler-Einstein metric if and only if it is K-polystable. There are several generalization of Kähler-Einstein metrics, which may exist even if the manifold is obstructed by Futaki's invariant (introduced in [20]). One is the extremal Kähler metric introduced by [12]. It is defined in terms of the scalar curvature so that make sense for general polarized manifolds. In [30] Mabuchi introduced an analogous notion in terms of the Ricci potential function. If the automorphism group is finite, it is classically known that these two notions coincide. In general, however, these are different metrics and one may only ask if the existence conditions are equivalent. The one direction is relatively clear; if the Fano manifold admits a Mabuchi soliton, there exists an extremal metric. (See e.g. [25, 29, 36]). The recent result [2] states that the converse is also true under a sharp assumption (see Theorem 1.4). The proof is intricate as it exploits the deep result

---

2010 *Mathematics Subject Classification.* Primary 53C25; Secondary 53C55, 58E11.

*Key words and phrases.* Calabi's extremal metric, Mabuchi's soliton, complex Monge-Ampere equation.

of the minimal model program in algebraic geometry. The aim of the present paper is to provide a direct analytic proof for the existence of the Mabuchi soliton. The strategy is to bound the associated K-energy functional such that one may run the continuity method for the Mabuchi soliton. The boundedness and the related compactness argument is inevitable since the monotonicity property of the energy along such path might not be expected. Unlike the Kähler-Einstein case, we cannot provide a simple comparison for the two canonical energy functionals  $D_g$  and  $M_g$ , either.

Our argument can be applied to general  $g$ -soliton metric in the sense of [11], [22]. In order to describe the general situation, we fix a compact torus  $T \subset \text{Aut}(X)$  so that obtain the associated weight polytope  $P$  as the image of the moment map  $\mu_\omega: X \rightarrow \mathfrak{t}^*$ . We take a smooth positive function  $g: P \rightarrow \mathbb{R}$  and denote the composition by  $g_\omega := g \circ \mu_\omega$ .

**Definition 1.1** ([22], Definition). A  $T$ -invariant Kähler metric  $\omega$  is called a  $g$ -soliton if it satisfies

$$(1.1) \quad \text{Ric}(\omega) - \omega = \sqrt{-1} \partial \bar{\partial} \log g_\omega.$$

It is called  $g$ -extremal if the scalar curvature satisfies

$$(1.2) \quad S_\omega - n = 1 - g_\omega.$$

When  $g$  is an affine function, each  $g$ -soliton and  $g$ -extremal metric yields the Mabuchi soliton and the extremal metric. In the case  $g = e^l$  for some affine function  $l$ ,  $g$ -soliton yields the Kähler-Ricci soliton.

As it was observed in Mabuchi's original paper [30], the constant

$$(1.3) \quad m_X := \sup_X (1 - g_\omega)$$

is independent of the metrics and  $m_X < 1$  if there exists a  $g$ -soliton.

In order to find the Mabuchi soliton, [28] introduced the following variant of the continuity method. Let us take a reference  $\omega_0$  and represent each metric by a Kähler potential  $\varphi$  such that  $\omega = \omega_0 + \sqrt{-1} \partial \bar{\partial} \varphi$ . The continuity path is then described as the Monge-Ampère equation:

$$(1.4) \quad g_\varphi \omega_\varphi^n = e^{-t\varphi + \rho_0} \omega_0^n \quad \text{for } t \in [0, 1].$$

The solution at  $t = 1$  is nothing but a  $g$ -soliton. Such a continuity path is originated from [3], where the path is formulated for the Kähler-Einstein metrics. See also [35] for the Kähler-Ricci soliton. Our main result is the following.

**Theorem 1.2.** Let  $\mathcal{T}$  be the set of  $t \in [0, 1]$  such that the equation (1.4) has a  $T$ -invariant solution. In general  $\mathcal{T}$  is non-empty and open. If one assume that  $X$  admits a  $g$ -extremal metric and that the Mabuchi constant enjoys  $m_X < 1$ , then  $\mathcal{T} = [0, 1]$ .

Recall that the  $g$ -extremal metric can be characterized as a critical point of the  $g$ -Mabuchi functional  $M_g$  (see (2.24) for the definition). The following is a key to prove the closedness of  $\mathcal{T}$ .

**Theorem 1.3.** Assume  $m_X < 1$  and fix any  $\varepsilon \in (0, 1)$ . Let  $\omega_t$  be a  $T$ -invariant solution of (1.4) at  $t \in (\varepsilon, 1]$ . There exists a constant  $C > 0$  independent of  $\varepsilon$  and  $t$  satisfying

$$M_g(\omega_t) \leq C\varepsilon^{-1}.$$

We emphasize that the critical point of  $M_g$  is a  $g$ -extremal metric which is different from the  $g$ -soliton unless  $g \equiv 1$ . In fact there exists another K-energy functional in the literatures, which we distinguish as  $F_g$ . The energy  $F_g$  is defined as the free energy of the Monge-Ampère measure with  $g$ -density:  $g_\omega \omega^n$ , and the critical point is the  $g$ -soliton (see Remark 2.2 below). The difficulty of general  $g$  case is caused by the difference between  $M_g$  and  $F_g$ .

As a consequence of Theorem 1.2, we have the following.

**Theorem 1.4.** For any Fano manifold  $X$  the followings are equivalent.

- (1)  $X$  admits a  $g$ -soliton.
- (2) It satisfies  $m_X < 1$  and is uniformly  $g$ -relatively D-stable.
- (3) It satisfies  $m_X < 1$  and admits a  $g$ -extremal metric in  $c_1(X)$ .

Equivalence of (1) and (2) is proved by [25] and [22], assuming  $m_X < 1$ . In [36] it was shown that at least if  $g$  is affine  $m_X < 1$  follows from the stability. [2] showed the implication (3)  $\Rightarrow$  (2) in an algebraic way. Note that the assumption  $m_X < 1$  in (3) is necessary, as it was pointed out by the second author in [29, Section 9.5]. In fact there exists a Fano 3-fold which admits an extremal metric in  $c_1(X)$  but  $m_X \geq 1$ .

**Acknowledgment.** The authors would like to thank Vestislav Apostolov, Eiji Inoue, Yasufumi Nitta and Yi Yao for helpful discussion and comments. The second author expresses his gratitude to Shigetoshi Bando for encouragements for the problem discussed in this article. The first author is supported by JSPS Grant-in-Aid for Scientific Research (C) No. 21K03229. The second author is supported by JSPS Grant-in-Aid for Science Research Activity Start-up No. 21K20342 and JSPS Grant-in-Aid for Early-Career Scientists No. 24K16917.

## 2. PRELIMINARY

**2.1.  $g$ -solitons and  $g$ -extremal metrics.** Throughout this paper we consider an  $n$ -dimensional Fano manifold  $X$ . We fix a compact torus  $T \subset \text{Aut}(X)$  and a  $T$ -invariant Kähler metric  $\omega_0 \in 2\pi c_1(X)$ . Let

$$(2.1) \quad \mathcal{H}(X, \omega_0)^T := \left\{ \varphi \in C^\infty(X, \mathbb{R}) : \omega_\varphi := \omega_0 + \sqrt{-1} \partial \bar{\partial} \varphi > 0 \text{ and } \varphi \text{ is } T\text{-invariant} \right\}$$

be the set of  $T$ -invariant Kähler potentials. Here each  $\sigma \in T_{\mathbb{C}}$  sends  $\varphi$  to  $\sigma[\varphi]$  which is defined by  $\sigma^*\omega_\varphi = \omega_0 + \sqrt{-1}\partial\bar{\partial}\sigma[\varphi]$ . The Ricci potential is a smooth function  $\rho_\varphi$  characterized by the property

$$(2.2) \quad \text{Ric}(\omega_\varphi) - \omega_\varphi = \sqrt{-1}\partial\bar{\partial}\rho_\varphi \quad \text{and} \quad \int_X (e^{\rho_\varphi} - 1)\omega_\varphi^n = 0.$$

We say that  $\omega_\varphi$  is Kähler-Einstein if  $\rho_\varphi = 0$ . In terms of the scalar curvature  $S_\varphi := \text{Tr}_{\omega_\varphi} \text{Ric}(\omega_\varphi)$ , it is equivalent to say  $S_\varphi = n$ .

For a convenient let us fix an isomorphism  $T \simeq (\mathbb{S}^1)^r$  with some non-negative integer  $r$ . For each  $\alpha = 1, \dots, r$ , let  $\xi_\alpha$  be the holomorphic vector field on  $X$  generating the action of the  $\alpha$ -th factor of  $T_{\mathbb{C}} \simeq (\mathbb{C}^*)^r$ . Once the metric  $\varphi \in \mathcal{H}(X, \omega_0)^T$  is fixed, each vector  $\xi \in \mathfrak{t}_{\mathbb{C}}$  defines the Hamilton function which is characterized by the properties  $\theta_\xi(\varphi) \in C^\infty(X; \mathbb{R})$  such that

$$(2.3) \quad i_\xi \omega_\varphi = \sqrt{-1}\partial\bar{\partial}\theta_\xi(\varphi) \quad \text{and} \quad \int_X \theta_\xi(\varphi)\omega_\varphi^n = 0.$$

We set  $\theta_\alpha := \theta_{\xi_\alpha}$ . The moment map  $\mu_\omega: X \rightarrow \mathbb{R}^r$  is then described as  $\mu_\varphi(x) = (\theta_1(x), \dots, \theta_r(x))$ . It is well-known that the image  $P := \mu_\varphi(X)$  defines a convex polytope in  $\mathbb{R}^r$  and independent of the choice of  $\varphi \in \mathcal{H}(X, \omega_0)^T$ .

We will also fix a smooth positive function  $g: P_X \rightarrow \mathbb{R}$ . and put  $g_\varphi := g \circ \mu_\varphi$ . In what follows it is always normalized such that

$$(2.4) \quad \int_X (g_\varphi - 1)\omega_\varphi^n = 0$$

holds. Thanks to the compactness of  $X$  there exists a uniform constant  $C > 0$  such that

$$(2.5) \quad C^{-1} \leq g_\varphi \leq C$$

**Definition 2.1** ([22], Definition). A  $T$ -invariant Kähler metric  $\omega$  is called a  $g$ -soliton if it satisfies

$$(2.6) \quad \text{Ric}(\omega) - \omega = \sqrt{-1}\partial\bar{\partial}\log g_\omega,$$

or equivalently,  $e^{\rho_\omega} = g_\omega$ . It is called  $g$ -extremal if the scalar curvature satisfies

$$(2.7) \quad S_\omega - n = 1 - g_\omega.$$

The definition of  $g$ -soliton generalizes the notion of the Kähler-Einstein metric, the Kähler-Ricci soliton, and the Mabuchi soliton, as it was explained in the introduction. The  $g$ -extremal Kähler metric gives a generalization of extremal Kähler metrics. In the original work [11] consideration of general  $g$ -density is motivated by the optimal transport theory.

Let us consider the particular case when  $g$  is an affine function. In this case  $\omega_\varphi$  is  $g$ -soliton if and only if  $e^{\rho_\varphi} - 1$  is contained in the space of (the above normalized) Hamilton functions

of  $T_{\mathbb{C}}$ . The vector field  $\eta$  corresponds to  $\theta_{\eta}(\varphi) = 1 - e^{\rho\varphi}$  is algebraically characterized by the property

$$(2.8) \quad \text{Fut}(\xi) + \langle \xi, \eta \rangle = 0,$$

where  $\text{Fut}(\xi)$  denotes the Futaki invariant. Similarly  $g$ -extremality defines a vector field with the property  $\theta_{\eta}(\varphi) = S(\omega_{\varphi}) - n$ . As it was classically known, these two notions are equivalent and we call  $\eta$  *the extremal vector field*. One can also check that  $\eta$  is contained in the center of the Lie algebra of all holomorphic vector fields. As the first obstruction to the  $g$ -soliton metric, Mabuchi introduced the invariant

$$(2.9) \quad m_X := \sup_{x \in X} \theta_{\eta}(\varphi)(x),$$

which is actually independent of the choice of  $\varphi \in \mathcal{H}(X, \omega_0)^T$ . Existence of the Mabuchi soliton forces  $m_X < 1$ , as the positivity of the affine function  $g(\theta_1, \dots, \theta_r) := 1 - \sum \eta^{\alpha} \theta_{\alpha}$  on  $P$ .

**2.2. Energy functionals.** Let us briefly review the geometry of the space  $\mathcal{H}^T = \mathcal{H}(X, \omega_0)^T$  and the canonical energy functionals there. As it was first observed by Mabuchi, the tangent space of  $\mathcal{H}^T$  at  $\varphi$  is naturally identified with the smooth functions space  $C^{\infty}(X; \mathbb{R})^T$  which equips the canonical  $L^p$  metric

$$(2.10) \quad \|u\|_p := \left[ \frac{1}{V} \int_X |u|^p \omega_{\varphi}^n \right]^{\frac{1}{p}},$$

where  $V = \int_X \omega_{\varphi}^n$  is the volume. Recent development of variational approach to the Monge-Ampère equation reveals that the topology of  $\mathcal{H}^T$  induced by the  $L^1$ -metric is rather important. (See [9], [10] for this point.) We denote by  $d_1$  the distance function of the above  $L^1$ -metric structure. It is a bit confusing but note that the  $d_1$ -convergence is much stronger than the ordinal convergence  $\int_X |\varphi_j - \varphi| \omega_0^n \rightarrow 0$  of  $L^1$  functions. The  $d_1$ -topology is closely related to the Monge-Ampère energy

$$(2.11) \quad E(\varphi) = \frac{1}{(n+1)V} \sum_{i=0}^n \int_X \varphi \omega_0^i \wedge \omega_{\varphi}^{n-i},$$

which is pluripotential generalization of the Dirichlet energy and indeed characterized by the property

$$(2.12) \quad d_{\varphi} E = V^{-1} \omega_{\varphi}^n.$$

In fact the energy  $E(\varphi) \in [-\infty, \infty)$  is defined for an arbitrary  $\omega_0$ -plurisubharmonic (psh in short) function  $\varphi$ . Extending the classical idea of [5], in [7] the authors also extended the definition of the Monge-Ampère product  $V^{-1} \omega_{\varphi}^n$  to general  $\omega_0$ -psh functions, as a measure  $\text{MA}(\varphi)$  which contains no mass on any pluripolar set. The differential formula (2.12) then

still holds in the appropriate sense. See the textbook [21] for the exposition. According to the seminal work of Darvas [15, 16], the space of finite energy  $\omega_0$ -psh functions

$$(2.13) \quad \mathcal{E}^1(X, \omega_0)^T := \left\{ \varphi \in \text{PSH}(X, \omega_0)^T : \int_X \omega_\varphi^n = V, \quad \int_X \varphi \omega_\varphi^n > -\infty \right\}$$

precisely gives the completion of  $\mathcal{H}(X, \omega_0)^T$  endowed with the distance  $d_1$ .

According to the theory of [11], one can also define  $V^{-1}g_\varphi\omega_\varphi^n$  as a non-pluripolar probability measure  $\text{MA}_g(\varphi)$ , for general  $\omega_0$ -psh  $\varphi$  and for a fixed positive function  $g: P \rightarrow \mathbb{R}$  which was discussed in the previous subsection. At the same time they defined the  $g$ -twisted Monge-Ampère energy which enjoys the property

$$(2.14) \quad d_\varphi E_g = V^{-1}g_\varphi\omega_\varphi^n,$$

at least for smooth  $\varphi$ . If one sets  $g \equiv 1$  it recovers the original Monge-Ampère energy. It is then straightforward to extend the definition of the classical energy  $I$  and  $J$  of Aubin to this setting. For a smooth  $\varphi$ , they are written down as

$$(2.15) \quad I_g(\varphi) = \frac{1}{V} \int_X \varphi (g_0 \omega_0^n - g_\varphi \omega_\varphi^n),$$

$$(2.16) \quad J_g(\varphi) = \frac{1}{V} \int_X \varphi g_0 \omega_0^n - E_g(\varphi).$$

A simple integration part yields  $0 \leq I \leq (n+1)(I-J) \leq nI$ . According to (2.5), one has the uniform estimate  $\frac{1}{C}I \leq I_g \leq CI$ ,  $\frac{1}{C}J \leq J_g \leq CJ$  and  $\frac{1}{C}(I-J) \leq (I_g - J_g) \leq C(I-J)$ . The reader can consult with [22] for the detail computations.

Now we introduce the canonical energy functionals  $D$  and  $M$  which are originally introduced to study the Kähler-Einstein and the constant scalar curvature Kähler metric. The principal term of each energy is the  $L$ -functional and the relative entropy (of the two probability measure  $\mu, \mu_0$ ), defined respectively as

$$(2.17) \quad L(\varphi) = -\log \frac{1}{V} \int_V e^{-\varphi + \rho_0} \omega_0^n \quad \text{and} \quad \text{Ent}(\mu|\mu_0) = \int_X \log \left( \frac{\mu}{\mu_0} \right) \mu.$$

The D-energy and the Mabuchi's K-energy are defined as

$$(2.18) \quad D(\varphi) := L(\varphi) - E(\varphi) \quad \text{and}$$

$$(2.19) \quad M(\varphi) := \text{Ent}(\text{MA}(\varphi)|\text{MA}(0)) + \frac{1}{V} \int_X \varphi \omega_\varphi^n - E(\varphi) + \frac{1}{V} \int_X \rho_0 (\omega_0^n - \omega_\varphi^n).$$

It is not so hard to check that the critical points of each energy give the Kähler-Einstein and cscK metrics, respectively. According to [6], these two energies can be related from the thermodynamical point of view. The reader can also consult with [9]. The Legendre duality between the functions and measures introduces the pluricomplex energy of the probability measure  $\mu$  as

$$(2.20) \quad E^*(\mu) := \sup_{u \in \mathcal{E}(X, \omega_0)} \left[ E(u) - \int_X u d\mu \right] \in (-\infty, \infty].$$

Then the Helmholtz free energy of the given probability measure  $\mu$  is defined to be

$$(2.21) \quad F(\mu) := \text{Ent}(\mu|\mu_0) - E^*(\mu).$$

In this context  $D$  and  $M$  appear in the form

$$(2.22) \quad \text{Ent}(\mu|\mu_0) = \sup_{f \in C^0(X; \mathbb{R})} \left[ \int_X f d\mu - \log \int_X e^f d\mu_0 \right],$$

$$(2.23) \quad M(\varphi) = F(\text{MA}(\varphi)),$$

where the background measure is chosen to be  $\mu_0 = \text{MA}(0)$ . The above relation is essential in showing the equivalence of the coercivity of two energies  $D$  and  $M$ .

Let us discuss the  $g$ -twisted version of the canonical energies, which are defined as

$$(2.24) \quad D_g = D + E - E_g \quad \text{and} \quad M_g = M + E - E_g.$$

Critical points of  $D_g$  and  $M_g$  gives  $g$ -solitons and  $g$ -extremal metric, respectively.

**Remark 2.2.** One can also define the  $g$ -twisted free energy  $F_g(\varphi) := F(\text{MA}_g(\varphi))$  which is explicitly written down to the form

$$(2.25) \quad F_g(\varphi) := \text{Ent}(\text{MA}_g(\varphi)|\text{MA}_g(0)) + \frac{1}{V} \int_X \varphi g_\varphi \omega_\varphi^n - E_g(\varphi) + \frac{1}{V} \int_X \rho_0 (g_0 \omega_0^n - g_\varphi \omega_\varphi^n).$$

The functional plays an important role in the study of  $g$ -solitons, however, the critical point of  $F_g$  does not give the extremal metrics but  $g$ -solitons. See [22] for the detail. The fact  $F_g \neq M_g$  for general  $g$  causes difficulty in comparing  $D_g$  with  $M_g$ . This is why we need to develop another approach in showing (3)  $\Rightarrow$  (1) of Theorem 1.4.

We here prepare for the later use an explicit difference between  $M_g$  and  $D_g$ .

**Lemma 2.3.** For any  $\varphi \in \mathcal{H}^T$ ,

$$M_g(\varphi) - D_g(\varphi) = -\frac{1}{V} \int_X \rho_\varphi \omega_\varphi^n + \frac{1}{V} \int_X \rho_0 \omega_0^n.$$

In particular,  $M_g(\varphi) - D_g(\varphi) \geq \frac{1}{V} \int_X \rho_0 \omega_0^n$ .

*Proof.* By the definition of  $D_g$  and  $M_g$ , it suffice to show the equality for  $g \equiv 1$ . The equality for  $g \equiv 1$  was firstly observed by Ding-Tian [18]. The inequality follows from the fact that  $e^x \geq 1 + x$  and the normalization of the Ricci potential.  $\square$

Nextly we introduce the coercivity property of the energy, which corresponds to algebraic (so-called uniform) stability and ensures the existence of a critical point. See [26] for the validity of the following definition which is modified by a (possibly non-maximal) torus.

**Definition 2.4.** We say that a functional  $F: \mathcal{H}^T \rightarrow \mathbb{R}$  is  $T_{\mathbb{C}}$ -coercive if there exist uniform constants  $\delta, C > 0$  such that for any  $\varphi \in \mathcal{H}^T$

$$(2.26) \quad F(\varphi) \geq \delta \inf_{\sigma \in T_{\mathbb{C}}} J(\sigma[\varphi]) - C$$

holds, where  $\sigma[\varphi]$  is defined by  $\sigma^*\omega_{\varphi} = \omega_0 + \sqrt{-1}\partial\bar{\partial}\sigma[\varphi]$ .

Note that we can also use the functional  $I_g - J_g$  to define the coercivity, since  $J$  and  $I_g - J_g$  are equivalent, as it was explained before.

**Theorem 2.5.** A Fano manifold  $X$  admits a  $g$ -soliton in  $\mathcal{H}^T$  if and only if  $D_g$  is  $T_{\mathbb{C}}$ -coercive. It admits a  $g$ -extremal metric in  $\mathcal{H}^T$  if and only if  $M_g$  is  $T_{\mathbb{C}}$ -coercive.

*Proof.* For the  $g$ -soliton, this is due to [22], Theorem 3.5. See also [31]. For the  $g$ -extremal metric, we can adapt the argument of [31] which generalizes He's extension [24] of Chen-Chen's result [13]. We also refer the proof of [2] Theorem 5. Indeed more general weighted constant scalar curvature Kähler metric case is treated in [1, 23].  $\square$

In view of the inequality in Lemma 2.3 and above theorems, we can conclude that if a Fano manifold admits a  $g$ -soliton it also admits a  $g$ -extremal metric. This proves the direction (1)  $\Rightarrow$  (3) in Theorem 1.4.

### 3. CONTINUITY METHOD FOR $g$ -SOLITONS

**3.1.  $g$ -twisted Laplacian.** We start from discussing the linearized equation for the continuity method. Let  $\Delta_{\varphi} := -\bar{\partial}^*\bar{\partial}$  be the geometric Laplacian with respect to the Kähler metric  $\varphi \in \mathcal{H}^T$ , acting on smooth functions space  $C^{\infty}(X; \mathbb{R})$ . The adjoint operator  $\bar{\partial}^*$  is taken with respect to the natural Hermitian inner product

$$(3.1) \quad \langle u, v \rangle_{\varphi} = \frac{1}{V} \int_X u \bar{v} \omega_{\varphi}^n.$$

Define the  $g$ -twisted Laplacian for  $\varphi \in \mathcal{H}^T$  acting on functions as

$$(3.2) \quad \Delta_{g,\varphi} f = -\frac{1}{g_{\varphi}} \bar{\partial}^*(g_{\varphi} \bar{\partial} f).$$

**Lemma 3.1.** We have the following properties for  $\Delta_{g,\varphi}$ .

- (1) The operator  $\Delta_{g,\varphi}$  is elliptic and self-adjoint with respect to the  $g$ -twisted Hermitian inner product

$$(3.3) \quad \langle u, v \rangle_{g,\varphi} := \frac{1}{V} \int_X u \bar{v} g_{\varphi} \omega_{\varphi}^n.$$

- (2) The kernel of  $\Delta_{g,\varphi}$  consists of the constant functions.
- (3) Let  $g_{i\bar{j}}$  be the metric tensor of  $\omega_{\varphi}$ . Then

$$\Delta_{g,\varphi} f = \Delta_{\varphi} f + \langle \bar{\partial} \log g_{\varphi}, \bar{\partial} f \rangle = \Delta_{\varphi} f + g^{i\bar{j}} \partial_{\bar{j}} f \partial_i \log g_{\varphi}.$$



(4) For any  $\varphi \in \mathcal{H}^T$ , let  $\varphi_t$  be a path in  $\mathcal{H}^T$  with  $\varphi_0 = \varphi$  and  $\frac{d}{dt}\big|_{t=0}\varphi_t =: u$ . Then

$$(3.4) \quad \frac{d}{dt}\bigg|_{t=0} (g_{\varphi_t} \omega_{\varphi_t}^n) = (\Delta_{g,\varphi} u) g_{\varphi} \omega_{\varphi}^n.$$

*Proof.* The claims (1), (2) and (3) follow from the definition of  $\Delta_{g,\varphi}$  immediately. We give a proof of (4). Recall that  $\xi_1, \dots, \xi_r$  are the generating vector fields for  $T_{\mathbb{C}} \simeq (\mathbb{C}^*)^r$ . Let  $(\theta_1, \dots, \theta_r)$  be the standard coordinate of the moment polytope  $P$ . In view of (3), it suffice to show

$$(3.5) \quad \frac{d}{dt}\bigg|_{t=0} g_{\varphi_t} = \langle \bar{\partial} g_{\varphi}, \bar{\partial} u \rangle.$$

By simple calculations, we get

$$\frac{d}{dt}\bigg|_{t=0} g_{\varphi_t} = \sum_{\alpha=1}^r \frac{\partial g_{\varphi}}{\partial \theta_{\alpha}} \xi_{\alpha}(u) \quad \text{and} \quad \langle \bar{\partial} g_{\varphi}, \bar{\partial} u \rangle = \sum_{\alpha=1}^r \frac{\partial g_{\varphi}}{\partial \theta_{\alpha}} \overline{\xi_{\alpha}(u)}.$$

Since any metric in  $\mathcal{H}^T$  is invariant under the action of the imaginary part of  $\xi_{\alpha}$ , we have  $\xi_{\alpha}(u) = \overline{\xi_{\alpha}(u)}$ . This completes the proof.  $\square$

**3.2. Continuity method.** In this subsection we prove the set  $\mathcal{T}$  in Theorem 1.2 is non-empty and open. Recall we consider the continuity method (1.4) to construct a  $g$ -soliton. Let  $\mathcal{T}$  be the set of  $t \in [0, 1]$  such that the equation (1.4) has a solution in  $\mathcal{H}^T$ . For this purpose, we consider the following modified equation

$$(3.6) \quad g_{\varphi} \omega_{\varphi}^n = e^{-t\varphi - E_g(\varphi) + \rho_0} \omega_0^n \quad \text{for } t \in [0, 1].$$

Let  $\mathcal{T}^*$  be the set of  $t \in [0, 1]$  such that the equation (3.6) has a solution in  $\mathcal{H}^T$ . For any  $t \in (0, 1]$  there is a one-to-one correspondence between the solution of (1.4) and the solution of (3.6). The linearized operator associated from (1.4), however, is not invertible at  $t = 0$  as discussed below.

**3.2.1. Solution at  $t = 0$ .** According to [11], Theorem 1.2, there exists a  $T$ -invariant continuous solution  $\varphi$  of  $g_{\varphi} \omega_{\varphi}^n = e^{\rho_0} \omega_0$ . The regularity result [33] shows that  $\varphi$  is smooth. One can also apply the argument of [22], Proposition 32 for the smoothness. Up to addition of a constant  $\varphi$  satisfies (3.6) at  $t = 0$ . Therefore  $0 \in \mathcal{T}^*$ .

**3.2.2. The openness of  $\mathcal{T}^*$  at  $t = 0$ .** For any  $\varphi \in \mathcal{H}^T$ , define

$$\Phi^*(\varphi) = \log \left( \frac{g_{\varphi} \omega_{\varphi}^n}{\omega_0^n} \right) + E_g(\varphi) - \rho_0.$$

By lemma 3.1, the linearized operator at  $t = 0$  is  $\delta\Phi^*(u) = \Delta_{g,\varphi} u + V^{-1} \int_X u g_{\varphi} \omega_{\varphi}^n$ , where  $u$  is any variation at a solution  $\varphi$  of (3.6) at  $t = 0$ . Since the kernel of  $\delta\Phi^*$  is trivial, the implicit function theorem shows the openness of  $\mathcal{T}^*$  at  $t = 0$ .

3.2.3. *The openness of  $\mathcal{T} \cap (0, 1)$ .* For  $\varphi \in \mathcal{H}^T$ , define

$$\Phi(\varphi) = \log \left( \frac{g_\varphi \omega_\varphi^n}{\omega_0^n} \right) + t\varphi - \rho_0.$$

By Lemma 3.1, the linearized operator is  $\delta\Phi(u) = \Delta_{g,\varphi}u + tu$ , where  $u$  is any variation at a solution  $\varphi$  of (1.4) at  $t$ . The openness of  $\mathcal{T} \cap (0, 1)$  follows from the next proposition combined with the implicit function theorem.

**Proposition 3.2.** Let  $\varphi_t \in \mathcal{H}^T$  be the solution of (1.4) at  $t \in (0, 1)$ . The first eigenvalue of the operator  $\Delta_{g,\varphi_t} + t$  is negative.

*Proof.* Let  $g_{i\bar{j}}$  be the metric tensor for  $\omega_{\varphi_t}$  and  $\lambda$  be the first non-zero eigenvalue of  $\Delta_{g,\varphi_t}$ , and  $u \in C^\infty(X, \mathbb{R})$  be the eigenvector. Put  $f = \log g_{\varphi_t}$  for simplicity. Applying  $\nabla_{\bar{k}}$  to the equation  $\Delta_{g,\varphi_t}u = \lambda u$  we have

$$(3.7) \quad \lambda \nabla_{\bar{k}}u = g^{i\bar{j}} \nabla_i \nabla_{\bar{j}} \nabla_{\bar{k}}u - R_{\bar{k}}^{\bar{p}} \nabla_{\bar{p}}u + g^{i\bar{j}} \nabla_{\bar{j}}u \nabla_{\bar{k}} \nabla_i f + g^{i\bar{j}} \nabla_{\bar{k}} \nabla_{\bar{j}} f \nabla_i u$$

$$(3.8) \quad < g^{i\bar{j}} \nabla_i \nabla_{\bar{j}} \nabla_{\bar{k}}u - t \nabla_{\bar{k}}u + g^{i\bar{j}} \nabla_{\bar{k}} \nabla_{\bar{j}}u \nabla_i f$$

$$(3.9) \quad = e^{-f} \nabla_i \left( e^f \nabla_{\bar{j}} \nabla_{\bar{k}}u \right) - t \nabla_{\bar{k}}u.$$

In the above inequality we used the equation (1.4) in the form

$$(3.10) \quad \text{Ric}(\omega_{\varphi_t}) - \sqrt{-1} \partial \bar{\partial} \log g_{\varphi_t} = (1-t)\omega_0 + t\omega_{\varphi_t} > t\omega_{\varphi_t}.$$

If one multiplies  $g^{m\bar{l}} \nabla_m u g_{\varphi_t} \omega_{\varphi_t}$  to the both sides then a simple integration by part shows

$$\lambda \int_X |\bar{\nabla}u|^2 g_{\varphi_t} \omega_{\varphi_t}^n < - \int_X |\bar{\nabla} \bar{\nabla}u|^2 g_{\varphi_t} \omega_{\varphi_t}^n - t \int_X |\bar{\nabla}u|^2 g_{\varphi_t} \omega_{\varphi_t}^n.$$

Therefore  $\lambda + t < 0$ . □

Combining the above results together, we conclude that  $\mathcal{T}$  is non-empty and open.

#### 4. ESTIMATE OF $M_g$ ALONG THE CONTINUITY METHOD

In this section we prove Theorem 1.3. Let  $\varphi_t \in \mathcal{H}^T$  be a solution of the equation (1.4) at  $t$ . In order to obtain the upper bound of  $M_g(\varphi_t)$ , it suffices to control  $D_g(\varphi_t) - \int_X \rho_\varphi \omega_\varphi^n$ , in view of Lemma 2.3. We put  $\omega_t := \omega_{\varphi_t}$ ,  $\rho_t := \rho_{\varphi_t}$ ,  $g_t := g_{\varphi_t}$  and  $\Delta_t := \Delta_{\varphi_t}$  for simplicity.

**Lemma 4.1.** Let  $\varphi_t \in \mathcal{H}^T$  be a solution of (1.4) at  $t$ . One has the relation

$$(4.1) \quad \rho_t + (1-t)\varphi_t - \log g_t - L(\varphi_t) = 0.$$

*Proof.* From the equation (1.4) and definition of the Ricci potential, we observe

$$(4.2) \quad 0 = -\sqrt{-1} \partial \bar{\partial} \log g_{\varphi_t} + \text{Ric}(\omega_t) - \text{Ric}(\omega_0) + \sqrt{-1} \partial \bar{\partial} (\rho_0 - t\varphi_t)$$

$$(4.3) \quad = \sqrt{-1} \partial \bar{\partial} \left( \rho_t + (1-t)\varphi_t - \log g_{\varphi_t} \right).$$

Thus  $\rho_t + (1-t)\varphi_t - \log g_{\varphi_t} = c_t$  for some constant  $c_t$  depending on  $t$ . The normalization of  $\rho_t$  and the equation (1.4) forces

$$(4.4) \quad c_t = -\log \frac{1}{V} \int_X e^{-(1-t)\varphi_t} g_{\varphi_t} \omega_t^n = -\log \frac{1}{V} \int_X e^{-\varphi_t + \rho_0} \omega_0^n.$$

The right-hand side is the  $L$ -functional introduced by (2.17).  $\square$

Integrating the both sides of Lemma 4.1, one obtains

$$(4.5) \quad -\frac{1}{V} \int_X \rho_t \omega_t^n \leq \frac{1-t}{V} \int_X \varphi_t \omega_t^n - \log(\inf_X g) - L(\varphi_t).$$

The first term is bounded from above, as follows.

**Lemma 4.2.** For a fixed  $\varepsilon \in (0, 1)$  the solution  $\varphi_t$  at  $t > \varepsilon$  satisfies

$$(4.6) \quad \frac{1}{V} \int_X \varphi_t \omega_t^n \leq \frac{e^{\sup \rho_0 - 1}}{\inf_X g} \varepsilon^{-1}.$$

*Proof.* Using the equation (1.4) of the continuity method, we transrate the volume form into the form

$$\begin{aligned} \frac{1}{V} \int_X \varphi_t \omega_t^n &\leq \frac{1}{V} \int_{\{\varphi_t > 0\}} \varphi_t g_t^{-1} e^{-t\varphi_t + \rho_0} \omega_0^n \\ &\leq \frac{e^{\sup \rho_0}}{V \inf_X g} \int_{\{\varphi_t > 0\}} \varphi_t e^{-t\varphi_t} \omega_0^n. \end{aligned}$$

This completes the proof, since for any  $t > \varepsilon$  the function  $\mathbb{R}_{>0} \ni x \mapsto xe^{-tx}$  is bounded from above by the constant  $(e\varepsilon)^{-1}$ .  $\square$

Although the third term  $L(\varphi_t)$  in the right hand side in (4.5) is not bounded from above, this is cancelled by the same term in  $D_g$ . The remaining part is in fact uniformly bounded from above regardless of  $m_X$  or  $\varepsilon$ .

**Lemma 4.3.** The functional  $I_g - J_g$  is non-decreasing along the continuity path (1.4).

*Proof.* The same result for  $I - J$  is proved in [4] page 28. If one Differentiates (1.4), Lemma 3.1 implies

$$(4.7) \quad \Delta_{g,t} \dot{\varphi}_t + t \dot{\varphi}_t + \varphi_t = 0.$$

So we may compute as

$$(4.8) \quad \frac{d}{dt}(I_g - J_g)(\varphi_t) = -\frac{1}{V} \int_X \varphi_t \Delta_{g,t} \dot{\varphi}_t g_t \omega_t^n$$

$$(4.9) \quad = \frac{1}{V} \int_X (\Delta_{g,t} \dot{\varphi}_t + t \dot{\varphi}_t) \Delta_{g,t} \dot{\varphi}_t g_t \omega_t^n$$

$$(4.10) \quad = \frac{1}{V} \int_X (\Delta_{g,t} \dot{\varphi}_t + t \dot{\varphi}_t)^2 \omega_t^n - \frac{t}{V} \int_X \dot{\varphi}_t (\Delta_{g,t} \dot{\varphi}_t + t \dot{\varphi}_t) g_t \omega_t^n.$$

The second term in the last line is non-negative, by Proposition 3.2.  $\square$

**Proposition 4.4.** Along the solution  $\varphi_t$ , the  $g$ -twisted Monge-Ampère energy is non-negative:  $E_g(\varphi_t) \geq 0$ .

*Proof.* Thanks to Lemma 4.3, as in the argument of Proposition 5.2, one can extend the solution  $\varphi_t$  at  $t$  to a family of solutions  $\varphi_s$  for  $s \in [0, t]$ . It is then sufficient to prove the formula

$$(4.11) \quad E_g(\varphi_t) = \frac{1}{t} \int_0^t \left( I_g(\varphi_s) - J_g(\varphi_s) \right) ds,$$

because  $I_g - J_g$  is non-negative. This is also well-known to the experts but we give a proof of (4.11) for the convenience to the reader. From the general expression

$$(4.12) \quad I_g(\varphi) - J_g(\varphi) = E_g(\varphi) - \frac{1}{V} \int_X \varphi g_\varphi \omega_\varphi^n,$$

it is reduced to the differential formula

$$(4.13) \quad t \frac{d}{dt} E_g(\varphi_t) = -\frac{1}{V} \int_X \varphi_t g_t \omega_t^n.$$

Let us differentiate the Monge-Ampère energy along the continuity path. We denote  $\dot{\rho}_t := \frac{d}{dt} \rho_t$  and  $\dot{\varphi}_t := \frac{d}{dt} \varphi_t$  for simplicity. If one differentiates the identity of Lemma 4.1 first, he observes

$$(4.14) \quad \dot{\rho}_t = \varphi_t - (1-t)\dot{\varphi}_t - \langle \bar{\partial} \log g_t, \bar{\partial} \dot{\varphi}_t \rangle + \frac{1}{V} \int_X \dot{\varphi}_t e^{\rho_t} \omega_t^n.$$

On the other hand, the definition of the Ricci potential yields

$$(4.15) \quad \dot{\rho}_t = \Delta_t \dot{\varphi}_t - \dot{\varphi}_t + \frac{1}{V} \int_X \dot{\varphi}_t e^{\rho_t} \omega_t^n,$$

where  $\Delta_t$  is the geometric Laplacian with respect to  $\omega_t$ . Using the two equations (4.14) and (4.15), one can compute as

$$(4.16) \quad \frac{d}{dt} E_g(\varphi_t) = \frac{1}{V} \int_X \dot{\varphi}_t g_t \omega_t^n$$

$$(4.17) \quad = \frac{1}{V} \int_X \left( -\varphi_t + (1-t)\dot{\varphi}_t + \langle \bar{\partial} \log g_t, \bar{\partial} \dot{\varphi}_t \rangle + \Delta_t \dot{\varphi}_t \right) g_t \omega_t^n$$

$$(4.18) \quad = -\frac{1}{V} \int_X \varphi_t g_t \omega_t^n + (1-t) \frac{d}{dt} E_g(\varphi_t).$$

In the last line we used Lemma 3.1, (3), (4) which tells

$$(4.19) \quad 0 = \frac{d}{dt} \int_X g_t \omega_t^n = \int_X \left( \Delta_t \dot{\varphi}_t + \langle \bar{\partial} \log g_t, \bar{\partial} \dot{\varphi}_t \rangle \right) g_t \omega_t^n.$$

It shows (4.13).  $\square$

Summarizing the above argument, we obtain the following: Fix any  $\varepsilon \in (0, 1)$ . Let  $\varphi_t \in \mathcal{H}^T$  be a solution of the equation (1.4) at  $t > \varepsilon$ . Then one has the estimate

$$M_g(\varphi_t) \leq \frac{e^{\sup \rho_0 - 1}}{\inf g} \varepsilon^{-1} - \log(\inf_X g) + \frac{1}{V} \int_X \rho_0 \omega_0^n.$$

This completes the proof of Theorem 1.3.

## 5. THE CLOSEDNESS OF $\mathcal{T}$

In this last section we complete the proof of Theorem 1.2. It remains to show the closedness of  $\mathcal{T}$ , when there exists an extremal Kähler metric. Thanks to Theorem 2.5, we may assume the coercivity of  $M_g$ . As in the previous section, we will write as  $\omega_t := \omega_{\varphi_t}$ ,  $\rho_t := \rho_{\varphi_t}$  and  $g_t := g_{\varphi_t}$ .

**5.1. Control of  $I_g - J_g$ .** In the situation of Theorem 1.3, the coercivity of  $M_g$  implies the boundedness of  $I_g - J_g$  modulo the action of the torus. The first step is to show that one can in fact control  $I_g - J_g$  along the continuity path. It extends [28], Lemma 3.3 to the general  $g$  case.

The following lemma rephrases the equivalence of the two expressions for the (modified) Futaki invariant, in terms of the energies.

**Lemma 5.1.** Let  $\xi \in \mathfrak{t}_{\mathbb{C}}$  and  $\{\sigma_s\}_{s \in \mathbb{R}}$  the one-parameter subgroup generated by the real part  $\operatorname{Re}(\xi)$ . Fix any  $\varphi \in \mathcal{H}^T$ . Then,

$$(5.1) \quad \frac{d}{ds} M_g(\sigma_s[\varphi]) = \frac{d}{ds} D_g(\sigma_s[\varphi]).$$

If the functional  $M_g$  is bounded from below on  $\mathcal{H}^T$ , then the both sides vanish identically.

*Proof.* Since  $\sigma_s^* \omega_{\varphi} = \omega_{\sigma_s[\varphi]}$  and  $\sigma_s^* \rho_{\varphi} = \rho_{\sigma_s[\varphi]}$ , (5.1) follows from Lemma 2.3. We put  $\psi := \sigma_s[\varphi]$  and starts from the expression

$$(5.2) \quad \frac{d}{ds} D_g(\sigma_s[\varphi]) = \frac{1}{V} \int_X \operatorname{Re}(\theta_{\xi}(\psi))(e^{\rho_{\psi}} - g_{\psi}) \omega_{\psi}^n,$$

where  $\theta_{\xi}(\psi)$  is the Hamilton function. If one differentiates the defining equation of the Ricci potential along  $\xi$ , he observes

$$(5.3) \quad \Delta_{\psi} \theta_{\xi}(\psi) + \theta_{\xi}(\psi) + \xi \rho_{\psi} = \frac{1}{V} \int_X \theta_{\xi}(\psi) e^{\rho_{\psi}} \omega_{\psi}^n$$

(see e.g. [32], Lemma 3.1). Therefore we derive

$$(5.4) \quad \frac{d}{ds} D_g(\sigma_s[\varphi]) = \frac{1}{V} \int_X \operatorname{Re}(\xi)(\rho_{\psi} - \log g_{\psi}) g_{\psi} \omega_{\psi}^n.$$

The right-hand side is the real part of the  $g$ -twisted Futaki invariant (see [22], Definition 4.1). In particular, it does not depend on the choice of  $\psi$  and  $s$ . If  $M_g$  is bounded from below, therefore, it identically vanishes.  $\square$

**Proposition 5.1.** Assume the functional  $M_g$  is bounded from below on  $\mathcal{H}^T$ . Let  $\varphi_t \in \mathcal{H}^T$  be a solution of (1.4) at  $t \in [0, 1)$ . Then we have

$$\inf_{\sigma \in T_{\mathbb{C}}} (I_g - J_g)(\sigma[\varphi_t]) = (I_g - J_g)(\varphi_t).$$

*Proof.* In the same notation as the above lemma, a direct computation using (4.12) shows

$$(5.5) \quad \frac{d}{ds} \Big|_{s=0} (I_g - J_g)(\sigma_s[\varphi_t]) = -\frac{1}{V} \int_X \frac{d}{ds} \Big|_{s=0} \sigma_s[\varphi_t](\Delta_{g,\varphi_t}\varphi_t) g_t \omega_t^n$$

$$(5.6) \quad = \frac{1}{V} \int_X \langle \bar{\partial} \operatorname{Re}(\theta_\xi(\varphi_t)), \bar{\partial} \varphi_t \rangle g_t \omega_t^n.$$

If one takes notice of the definition of the Hamilton function and Lemma 4.1, it is equivalent to

$$(5.7) \quad -\frac{1}{(1-t)V} \int_X \operatorname{Re}(\xi)(\rho_t - \log g_t) g_t \omega_t^n = -\frac{1}{1-t} \frac{d}{ds} \Big|_{s=0} D_g(\sigma_s[\varphi]).$$

According to Lemma 5.1, it implies that  $I_g - J_g$  is critical at  $\sigma = \operatorname{id}$ .

We will show that for any fixed  $\varphi \in \mathcal{H}^T$ , the function  $s \mapsto f(s) := (I_g - J_g)(\sigma_s[\varphi])$  is properly convex (so that the infimum is achieved by  $\sigma = \operatorname{id}$ ). Similar argument for the  $J$ ,  $D$ , and  $M$  has appeared in [25], Theorem 2.5. See also [22], Lemma 29 and 38 for the detail of the fiber integration below. We consider the complex variable  $s \in \mathbb{C}$  and define  $\Omega(z, s) = \sigma_s^*(\omega_\varphi(z))$  as a smooth semipositive form on the product space  $X \times \mathbb{C}$ . If regards the pulled-back form  $p_1^*\omega_0$  by the first projection as a reference form, One has a potential function  $\Psi(z, s)$  such that  $\Omega = p_1^*\omega_0 + \sqrt{-1}\partial\bar{\partial}\Psi$ . In terms of the fiber integration the second variation of the Monge-Ampère energy (Recall that the first variation was given by (2.12)) is expressed as

$$(5.8) \quad \sqrt{-1}\partial\bar{\partial} E_g(\Psi) = \frac{1}{V} \int_X g_\Psi \Omega^{n+1},$$

where the operator  $\partial$  is taken for  $n+1$  variables  $(z, s)$ . It actually vanishes since  $\Omega^{n+1} = \sigma_s^*(\omega_\varphi^{n+1}) = 0$ . On the other hand, the integration of  $\Psi$  against the  $g$ -twisted Monge-Ampère measure on the product space is computed as

$$(5.9) \quad -\sqrt{-1}\partial\bar{\partial} \int_X \Psi g_\Psi \Omega^n = -\int_X \sqrt{-1}\partial\bar{\partial} \Psi \wedge g_\Psi \Omega^n$$

$$(5.10) \quad = -\int_X g_\Psi \Omega^{n+1} + \int_X p_1^*\omega_0 \wedge g_\Psi \Omega^n$$

$$(5.11) \quad = \int_X p_1^*\omega_0 \wedge g_\Psi \Omega^n.$$

The last term is obviously non-negative. It implies that  $(I_g - J_g)(\sigma_s[\varphi])$  is subharmonic in  $s$ . Properness of  $f(s)$  follows from the fact that the slope at infinity of  $f(s)$  is positive unless  $\operatorname{Re}(\xi) = 0$ .  $\square$

**5.2. Closedness of  $\mathcal{T}$ .** Let us now prove the closedness of  $\mathcal{T}$  so that conclude Theorem 1.2.

**Proposition 5.2.** *Suppose that the functional  $M_g$  is coercive on  $\mathcal{H}^T$ . Take  $t_j \in \mathcal{T}$  which converges to  $t_\infty \in \mathbb{R}$ , and  $\varphi_j \in \mathcal{H}^T$  as the solution of (1.4) at  $t = t_j$ . Replaced with a*

subsequence if necessary,  $\varphi_j$  converges to some  $\varphi_\infty \in \mathcal{E}^1(X, \omega_0)^T$  in the  $d_1$ -topology and the limit function  $\varphi_\infty$  is in fact a smooth solution of (1.4) at  $t = t_\infty$ . It implies  $t_\infty \in \mathcal{T}$ .

*Proof.* In the sequel all the constants  $C$  are uniform in  $j$ . We write  $\omega_j := \omega_{\varphi_j}$  and  $g_j := g_{\varphi_j}$ .

As we have already mentioned, the functionals  $I, J$ , and  $I_g - J_g$  are interchangeable with each other. According to Theorem 1.3 and Proposition 5.1, the coercivity of  $M_g$  implies the uniform bound  $J(\varphi_j) \leq C$ .

Uniform bound of the  $I$ -functional, combined with Lemma 4.2 yields  $\int_X \varphi_j \omega_0^n \leq C$ . Then by the standard Green function argument we may obtain the uniform bound of  $\sup_X \varphi_j$ . Let  $G_{\omega_0}$  be the Green function for the background metric  $\omega_0$ , which satisfies  $\inf_X G_{\omega_0} \geq -B$  for some positive constant  $B$ . Since  $\Delta_{\omega_0} \varphi_j \geq -n$ , the representation formula implies

$$(5.12) \quad \sup_X \varphi_j \leq \frac{1}{V} \int_X \varphi_i(y) \omega_0^n(y) + \sup_{x \in X} \int_X (G_{\omega_0}(x, y) + B) (-2\Delta_{\omega_0} \varphi_i(y)) \omega_i^n(y) \leq C.$$

The lower bound  $\sup_X \varphi_i \geq -C$  is rather clear since

$$(5.13) \quad \int_X e^{-t_j \varphi_j + \rho_0} \omega_0^n = \int_X g_j \omega_j^n = \int_X \omega_0^n.$$

Thus we obtain

$$(5.14) \quad \left| \sup_X \varphi_i \right| \leq C.$$

The above Green function argument implies the general inequality  $\sup_X \varphi \leq V^{-1} \int_X \varphi \omega_0^n + C$  so the uniform estimate of  $I(\varphi_i)$  now implies  $|\int_X \varphi_j \omega_j^n| < C$ . If look back to the expression of  $M_g$  (which is bounded from above by Theorem 1.3), we obtain the uniform bound of the relative entropy

$$(5.15) \quad \text{Ent}(\text{MA}(\varphi_j) | \text{MA}(0)) \leq C.$$

This is a key to extract a convergent subsequence. Indeed by the fundamental compactness result [9], Theorem 2.17 and Proposition 2.6, there exists a convergent subsequence and the limit in  $(\mathcal{E}^1)^T = \mathcal{E}^1(X, \omega_0)^T$ . As an abuse of notation, we denote the subsequence by the same symbol  $\varphi_j$ . According to [16], the convergence  $\varphi_j \rightarrow \varphi_\infty$  in  $d_1$ -topology implies  $\varphi_j \rightarrow \varphi_\infty$  in  $L^1(X, \omega_0^n)$ ,  $E(\varphi_j) \rightarrow E(\varphi_\infty)$ , and vice versa. Finiteness of the Monge-Ampère energy forces the singularities of the psh functions very mild. Indeed, we can apply the uniform version of the Skoda integrability theorem ([38], Corollary 3.2) to the sequence so that

$$(5.16) \quad \int_X e^{-pt_j \varphi_j} \omega_0^n, \quad \int_X e^{-pt_\infty \varphi_\infty} \omega_0^n \leq C$$

holds for any  $p > 1$ . The effective version of Demailly-Kollàr's semi-continuity theorem for log canonical thresholds ([19], Main Theorem 0.2, (2)). See also [9], Proposition 1.4)

shows the convergence  $e^{-t_j \varphi_j} \rightarrow e^{-t_\infty \varphi_\infty}$  in  $L^p(\omega_0^n)$  for all  $p > 1$ . It follows the convergence of probability measures

$$(5.17) \quad g_j \omega_j^n = e^{-t_j \varphi_j + \rho_0} \omega_0^n \rightarrow e^{-t_\infty \varphi_\infty + \rho_0} \omega_0^n.$$

We also claim the convergence of probability measures

$$(5.18) \quad g_j \omega_j^n = \text{MA}_g(\varphi_j) \rightarrow \text{MA}_g(\varphi_\infty).$$

Note that the limit measure is a priori singular and  $g_{\varphi_\infty}$  itself does not make sense. See [11] for the detail. It is convenient to use the symmetric  $I$ -functional

$$(5.19) \quad I(\psi_1, \psi_2) := \int_X (\psi_1 - \psi_2)(\omega_{\psi_1}^n - \omega_{\psi_2}^n)$$

which is defined for  $\psi_1, \psi_2 \in \mathcal{E}^1(X, \omega_0)$ . As [8], Lemma 3.13, we observe

$$\begin{aligned} \sup_{u \in \mathcal{E}_B^1} \left| \int_X u g_{\psi_1} \omega_{\psi_1}^n - \int_X u g_{\psi_2} \omega_{\psi_2}^n \right| &\leq C \sup_{u \in \mathcal{E}_B^1} \left| \int_X u \omega_{\psi_1}^n - \int_X u \omega_{\psi_2}^n \right| \\ &\leq CI(\psi_1, \psi_2)^{1/2} \end{aligned}$$

for the weak compact set

$$(5.20) \quad (\mathcal{E}^1)_B^T := \left\{ \varphi \in \mathcal{E}^1(X, \omega_0)^T : \sup_X \varphi \leq 0, E(\varphi) \geq -B \right\}.$$

Following [8], Proposition 5.6, we may exploit the above inequality to control the ( $g$ -twisted) pluricomplex energy of the probability measure

$$(5.21) \quad E_g^*(\mu) := \sup_{u \in (\mathcal{E}^1)^T} \left[ E_g(u) - \int_X u d\mu \right].$$

Indeed, the convergence  $\varphi_i \rightarrow \varphi_\infty$  in  $d_1$ -topology is equivalent to  $I(\varphi_j, \varphi_\infty) \rightarrow 0$ . Using the above inequality, we deduce  $E_g^*(\text{MA}_g(\varphi_j)) \rightarrow E_g^*(\text{MA}_g(\varphi_\infty))$ , which in particular implies the weak convergence (5.18).

From (5.17) and (5.18), we obtain the weak solution which satisfies

$$(5.22) \quad \text{MA}_g(\varphi_\infty) = e^{-t_\infty \varphi_\infty + \rho_0} \omega_0^n.$$

Since the right-hand side has  $L^p$ -density for some  $p > 1$ , one can apply [11], Theorem 1.2 (or more directly Kołodziej's a priori estimate [27] for the Monge-Ampère equation) so that  $\varphi_\infty$  is continuous. By the regularity result of [33],  $\varphi_\infty$  is then smooth. We can also apply the argument of [22], Proposition 32 for the smoothness of  $\varphi_\infty$ . At any rate it implies  $t_\infty \in \mathcal{T}$ .  $\square$



## REFERENCES

- [1] V. Apostolov, S. Jubert and A. Lahdili: *Weighted K-stability and coercivity with applications to extremal Kähler and Sasaki metrics*. *Geom. Topol.* **27** (2023), no.8, 3229–3302. 8
- [2] V. Apostolov, A Lahdili, and Y. Nitta: *Mabuchi Kähler solitons versus extremal Kähler metrics and beyond*. [arXiv:2407.01871](https://arxiv.org/abs/2407.01871). 1, 3, 8
- [3] T. Aubin: *Réduction du cas positif de l'équation de Monge-Ampère sur les variétés kählériennes compactes à la démonstration d'une inégalité*. *J. Funct. Anal.* **57** (1984), no.2, 143–153. 2
- [4] S. Bando and T. Mabuchi: *Uniqueness of Einstein Kähler metrics modulo connected group actions*. In *Algebraic geometry, Sendai, 1985* (T. Oda, ed.), *adv. Stud. Pure Math.* **10**, Kinokuniyua, Tokyo, 1987, 11–40. 11
- [5] E. Bedford and A. Taylor: *The Dirichlet problem for a complex Monge-Ampère equation*. *Invent. Math.* **37** (1976), no. 1, 1–44. 5
- [6] R. J. Berman: *A thermodynamical formalism for Monge-Ampère equations, Moser-Trudinger inequalities and Kähler-Einstein metrics*. *Adv. Math.* **248** (2013), 1254–1297. 6
- [7] S. Boucksom, P. Eyssidieux, V. Guedj, and A. Zeriahi: *Monge-Ampère equations in big cohomology classes*. *Acta Math.* **205** (2010), no. 2, 199–262. 5
- [8] R. J. Berman, S. Boucksom, V. Guedj, and A. Zeriahi: *A variational approach to complex Monge-Ampère equations*. *Publ. Math. Inst. Hautes Études Sci.* **117** (2013), 179–245. 16
- [9] R. J. Berman, S. Boucksom, P. Eyssidieux, V. Guedj, and A. Zeriahi: *Kähler-Einstein metrics and the Kähler-Ricci flow on log Fano varieties*. *J. Reine Angew. Math.* **751** (2019), 27–89. 5, 6, 15
- [10] S. Boucksom T. Hisamoto and M. Jonsson: *Uniform K-stability, Duistermaat-Heckman measures and singularities of pairs*. *Ann. Inst. Fourier (Grenoble)* **67** no. 2 (2017), 743–841. 5
- [11] R. J. Berman and D. Witt-Nyström: *Complex optimal transport and the pluripotential theory of Kähler-Ricci solitons*, [arXiv:1401.8264](https://arxiv.org/abs/1401.8264). 2, 4, 6, 9, 16
- [12] E. Calabi: *Extremal Kähler metrics II*. *Differential geometry and complex analysis*, I. Chavel and H.M. Farkas eds., 95–114, Springer-Verlag, Berlin-Heidelberg-NewYork, (1985). 1
- [13] X. X Chen and J. Cheng: *On the constant scalar curvature Kähler metrics I, II*. *J. Amer. Math. Soc.* **34** (2021), no. 4, 909–936. 937–1009. 8
- [14] X. X. Chen, S. K. Donaldson and S. Sun: *Kähler-Einstein metrics on Fano manifolds I, II, III*. *J. Amer. Math. Soc.* **28** (2015), no. 1, 183–197, 199–234, 235–278. 1
- [15] T. Darvas: *The Mabuchi completion of the space of Kähler potentials*. *Amer. J. Math.* **139** (2017), no. 5, 1275–1313. 6
- [16] T. Darvas: *The Mabuchi geometry of finite energy classes*. *Adv. Math.* **285** (2015), 182–219. 6, 15
- [17] T. Darvas and Y. A. Rubinstein: *Tian's properness conjectures and Finsler geometry of the space of Kähler metrics*. *J. Amer. Math. Soc.* **30** (2017), no. 2, 347–387.
- [18] W. Ding and G. Tian: *The generalized Moser-Trudinger inequality*. In *Nonlinear Analysis and Microlocal Analysis, Proceedings of the International Conference at Nankai Institute of Mathematics* (K.C. Chang et al., eds.), World Scientific, (1992), 57–70. 7
- [19] J. P. Demailly and J. Kollàr: *Semi-continuity of complex singularity exponents and Kähler-Einstein metrics on Fano orbifolds*. *Ann. Sci. École Norm. Sup. (4)* **34** (2001), no. 4, 525–556. 15
- [20] A. Futaki: *An obstruction to the existence of Einstein Kähler metrics*. *Invent. Math.* **73** (1983), no.3, 437–443. 1
- [21] V. Guedj and A. Zeriahi: *Degenerate complex Monge-Ampère equations*. *EMS Tracts in Mathematics* **26** (2017). 6

- [22] J. Han and C. Li: *On the Yau-Tian-Donaldson conjecture for generalized Kähler-Ricci soliton equations*. *Comm. Pure Appl. Math.* **76** (2023), 1793–1867 2, 3, 4, 6, 7, 8, 9, 13, 14, 16
- [23] J. Han and Y. Liu: *On the existence of weighted-cscK metrics*. [arXiv:2406.10939](https://arxiv.org/abs/2406.10939). 8
- [24] W. He: *On Calabi’s extremal metric and properness*. *Trans. Amer. Math. Soc.* **372** (2019), no. 8, 5595–5619. 8
- [25] T. Hisamoto: *Mabuchi’s soliton metric and relative D-stability*. *Amer. J. Math.* **145** (2023), no. 3, 765–806. 1, 3, 14
- [26] T. Hisamoto: *Stability and coercivity for toric polarizations*. [arXiv:1610.07998](https://arxiv.org/abs/1610.07998). 7
- [27] S. Kołodziej: *The complex Monge-Ampère equation*. *Acta Math.* **180** (1998), no. 1, 69–117. 16
- [28] Y. Li and B. Zhou: *Mabuchi metrics and properness of the modified Ding functional*. *Pacific J. Math.* **302** (2019), no. 2, 659–692. 2, 13
- [29] T. Mabuchi: *Test configurations, stabilities and canonical Kähler metrics—complex geometry by the energy method*. *SpringerBriefs in Mathematics*, Springer, Singapore, (2021). 1, 3
- [30] T. Mabuchi: *Kähler-Einstein metrics for manifolds with nonvanishing Futaki character*. *Tohoku Math. J.* (2) **53** (2001), 171–182. 1, 2
- [31] Y. Nakagawa and S. Nakamura: *Modified extremal Kähler metrics and multiplier Hermitian-Einstein metrics*. [arXiv:2405.05604](https://arxiv.org/abs/2405.05604). 8
- [32] Y. Nakagawa and S. Nakamura: *Multiplier Hermitian-Einstein metrics on Fano manifolds of KSM-type*. *Tohoku Math. J.* **76** (1) (2024), no. 1. 127–152. 13
- [33] G. Székelyhidi and V. Tosatti: *Regularity of weak solutions of a complex Monge-Ampère equation*. *Anal. PDE* **4** (2011), no. 3, 369–378. 9, 16
- [34] G. Tian: *K-stability and Kähler-Einstein metrics*. *Comm. Pure Appl. Math.* **68** (2015), 1085–1156. 1
- [35] G. Tian and X. Zhu: *Uniqueness of Kähler-Ricci solitons*. *Acta Math.*, **184** (2000), no. 2, 271–305. 2
- [36] Y. Yao: *Relative Ding Stability and an Obstruction to the Existence of Mabuchi Solitons*. *J. Geom. Anal.*, **32** (2022), no. 105. 1, 3
- [37] S. T. Yau, *On the Ricci curvature of a compact Kähler manifold and the complex Monge-Ampère equation, I*. *Comm. Pure Appl. Math.*, **31** (1978), 339–411.
- [38] A. Zeriahi: *Volume and capacity of sublevel sets of a Lelong class of plurisubharmonic functions*. *Indiana Univ. Math. J.* **50** (2001), no. 1, 671–703. 15

T. HISAMOTO: DEPARTMENT OF MATHEMATICAL SCIENCES, TOKYO METROPOLITAN UNIVERSITY,  
1-1 MINAMIOSAWA, HACHIOJI-SHI, TOKYO 192-0397, JAPAN

*Email address:* hisamoto@tmu.ac.jp

S. NAKAMURA: DEPARTMENT OF MATHEMATICS, TOKYO INSTITUTE OF TECHNOLOGY, 2-12-1  
OOKAYAMA, MEGURO-KU, TOKYO 152-8551, JAPAN

*Email address:* s.nakamura@math.titech.ac.jp