

EXTENDING THE SCIENCE FICTION AND THE LOEHR–WARRINGTON FORMULA

DONGHYUN KIM AND JAESEONG OH

ABSTRACT. We introduce the Macdonald piece polynomial $I_{\mu,\lambda,k}[X; q, t]$, which is a vast generalization of the Macdonald intersection polynomial in the science fiction conjecture by Bergeron and Garsia. We demonstrate a remarkable connection between $I_{\mu,\lambda,k}$, ∇s_λ , and the Loehr–Warrington formula LW_λ , thereby obtaining the Loehr–Warrington conjecture as a corollary. To connect $I_{\mu,\lambda,k}$ and ∇s_λ , we employ the plethystic formula for the Macdonald polynomials of Garsia–Haiman–Tesler, and to connect $I_{\mu,\lambda,k}$ and LW_λ , we use our new findings on the combinatorics of P -tableaux together with the column exchange rule. We also present an extension of the science fiction conjecture and the Macdonald positivity by exploiting $I_{\mu,\lambda,k}$.

1. INTRODUCTION

In an effort to extend the science fiction conjecture by [BG99], we introduce the Macdonald piece polynomial $I_{\mu,\lambda,k}$, which possesses remarkable properties. To provide a comprehensive overview of the primary results, we discuss the historical context and background of Macdonald polynomials, the Loehr–Warrington conjecture, and the science fiction conjecture.

1.1. Macdonald polynomials. The *Macdonald P polynomials*, denoted as $P_\mu[X; q, t]$, are symmetric functions over $\mathbb{Q}(q, t)$ which form a basis for the ring of symmetric functions. Macdonald introduced them in 1988 [Mac88] as a unified generalization of two significant families of symmetric functions, the *Jack polynomials* and the *Hall–Littlewood polynomials*. Since then, Macdonald polynomials have been established as fundamental objects with deep connections across various areas of mathematics. These include their appearance and relationships with affine Hecke algebras, quantum groups, elliptic Hall algebras, Hilbert scheme, affine Springer fiber, the Calogero–Sutherland model, diagonal coinvariants, and combinatorics.

An important question posed by Macdonald was the *Macdonald positivity conjecture*. This conjecture asserts that the *modified Macdonald polynomial*, defined through a plethystic substitution to the *Macdonald J polynomial* in [Mac88],

$$\tilde{H}_\mu[X; q, t] = t^{n(\mu)} J_\mu \left[\frac{X}{1-t^{-1}}; q, t^{-1} \right],$$

is Schur positive. To address this conjecture, Garsia and Haiman introduced the bigraded \mathfrak{S}_n module R_μ in 1993, now called the *Garsia–Haiman module*. This module is associated with a partition $\mu \vdash n$ and is defined as the subspace of the polynomial ring $\mathbb{C}[x_1, \dots, x_n, y_1, \dots, y_n]$ of $2n$ variables, spanned by the partial derivatives of the *generalized Vandermonde determinant*

$$\Delta_\mu := \det \left(x_k^{i-1} y_k^{j-1} \right)_{\substack{1 \leq k \leq n \\ (i,j) \in \mu}}$$

associated with μ [GH93, GH96].

In his groundbreaking work [Hai01], Haiman utilized the geometry of the Hilbert scheme of n points in the plane \mathbb{C}^2 to prove that

$$\text{grFrob}(R_\mu; q, t) = \tilde{H}_\mu[X; q, t], \quad (1.1)$$

thereby settling the Macdonald positivity conjecture. Another consequence of Haiman's result is the *n!* theorem, which states that

$$\dim(R_\mu) = n!. \quad (1.2)$$

Despite Haiman's resolution of Schur positivity, an explicit combinatorial formula for the Schur coefficients remains elusive.

Problem 1.1. Find a combinatorial description for the Schur coefficient $\tilde{K}_{\lambda, \mu}(q, t)$ of the Macdonald polynomials:

$$\tilde{H}_\mu[X; q, t] = \sum_{\lambda} \tilde{K}_{\lambda, \mu}(q, t) s_{\lambda},$$

where s_{λ} is the Schur function.

This open problem is one of the most fundamental and challenging questions in the theory of Macdonald polynomials and has been a driving force behind many major breakthroughs. One such advancement is the Haglund–Haiman–Loehr formula [HHL05a], which provides an explicit monomial expansion for the Macdonald polynomials in terms of words and two statistics, inv_{μ} and maj_{μ} (see Theorem 5.1).

1.2. Extending science fiction conjecture. One possible approach to resolving Problem 1.1 is to decompose \tilde{H}_{μ} into pieces that are Schur positive, and then find a combinatorial description of the Schur coefficients for each piece. The celebrated HHL formula breaks down \tilde{H}_{μ} into LLT polynomials, which are known to be Schur positive [GH07]. However, describing the Schur coefficients of LLT polynomials remains a challenging open problem. Another attempt involved using k -Schur functions [LLM03], whose Schur coefficients were explicitly described in [BMPS19]. However, it is still unclear whether \tilde{H}_{μ} can be expressed positively in terms of k -Schur functions. In this paper, we introduce an alternative way to decompose \tilde{H}_{μ} , generalizing the way proposed in the science fiction conjecture.

Bergeron and Garsia studied the Garsia–Haiman modules using inductive and combinatorial methods [BG99]. They introduced a collection of conjectures referred to as the *science fiction conjecture*, which include the existence of a certain basis for Garsia–Haiman modules and the elegant properties of those bases while intersecting Garsia–Haiman modules. To elaborate on the science fiction conjecture, let μ be a partition with n (removable) corners. For $1 \leq i \leq n$, let $\mu^{(i)}$ denote the partition obtained from μ by removing the i -th corner. The science fiction conjecture then implies the following two assertions, which are analogous to (1.1) and (1.2), respectively:

(1) The bigraded Frobenius characteristic of the intersection of the modules $\bigcap_{i=1}^k R_{\mu^{(i)}}$ is given by

$$\text{grFrob}\left(\bigcap_{i=1}^k R_{\mu^{(i)}}; q, t\right) = I_{\mu^{(1)}, \dots, \mu^{(k)}}[X; q, t] := \sum_{i=1}^k \left(\prod_{j \neq i} \frac{T_{\mu^{(j)}}}{T_{\mu^{(j)}} - T_{\mu^{(i)}}} \right) \tilde{H}_{\mu^{(i)}}[X; q, t], \quad (1.3)$$

(2) the dimension of the intersection $\bigcap_{i=1}^k R_{\mu^{(i)}}$ is given by

$$\dim \left(\bigcap_{i=1}^k R_{\mu^{(i)}} \right) = \frac{n!}{k}. \quad (1.4)$$

Here, $T_\mu := \prod_{(i,j) \in \mu} t^{i-1} q^{j-1}$. We refer to the symmetric function $I_{\mu^{(1)}, \dots, \mu^{(k)}}$ in (1.3) as the *Macdonald intersection polynomial*, and the second part as the *$n!/k$ conjecture*. The Macdonald intersection polynomial naturally refines the usual Macdonald polynomials. As such, the combinatorics of Macdonald intersection polynomials may lead to progress on Problem 1.1. The authors, along with Seung Jin Lee, recently provided an explicit positive monomial expansion for the Macdonald intersection polynomials for two partitions [KLO22], refining the celebrated HHL formula and providing a step toward Butler’s conjecture [But94].

In this paper, we further generalize the science fiction conjecture. Especially, we generalize the Macdonald intersection polynomial in (1.3) and the $n!/k$ conjecture (1.4). To be precise, like in the science fiction conjecture, let μ be a partition with n removable corners $\{c_1, \dots, c_n\}$ and let $1 \leq k \leq n-1$. For a subset $S \in \binom{[n]}{k}$ of $[n]$ of size k , let μ^S be the partition obtained from μ by deleting corners $\{c_i : i \in S\}$. Let $R_{n,k} := ((n-k)^k)$ be the rectangular partition, and let λ be a partition inside $R_{n,k}$. We define the *Macdonald piece polynomial* $I_{\mu,\lambda,k}$ indexed by μ , k , and λ as

$$I_{\mu,\lambda,k}[X; q, t] := (-1)^{|\lambda|} \sum_{S \in \binom{[n]}{k}} \frac{s_\lambda[z_S] \prod_{j \in S^c} z_j}{\prod_{\substack{i \in S \\ j \in S^c}} (z_j - z_i)} \tilde{H}_{\mu^S}[X]. \quad (1.5)$$

Here, we let $z_i = T_{\mu^{(i)}}/T_\mu = q^{-\text{coarm}(c_i)} t^{-\text{coleg}(c_i)}$ and $S^c = \{1, 2, \dots, n\} \setminus S$. Additionally, $f[z_S]$ denotes the polynomial f in the z_i ’s where i runs through S . For $k=1$ and the empty partition $\lambda = \emptyset$, $I_{\mu,\lambda,k}$ recovers the Macdonald intersection polynomial $I_{\mu^{(1)}, \dots, \mu^{(n)}}$.

In Section 2, using the novel symmetric function $I_{\mu,\lambda,k}$, we define another symmetric function $\tilde{H}_{\mu^S}^\lambda$ (See (2.2)). We conjecture that it is the graded Frobenius characteristic of the intersection of certain Garsia–Haiman modules (see Conjecture 2.3), largely generalizing the science fiction conjecture given in (1.3). In the appendix A, as a counterpart for the $n!/k$ conjecture (1.4), numbers corresponding to each partition and their conjectural properties are outlined.

In addition to its crucial role in generalizing the science fiction conjecture, Macdonald piece polynomial $I_{\mu,\lambda,k}$ enjoys a remarkable property. We unveil a surprising connection with ∇_{S_λ} and the Loehr–Warrington formula IW_λ , where the Loehr–Warrington conjecture follows as a corollary. We first give a background of the Loehr–Warrington conjecture.

1.3. The Loehr–Warrington conjecture. The theory of Macdonald polynomials has seen significant advancement due to its deep connections with algebraic geometry, operators on symmetric functions, and Catalan combinatorics. A key result that exemplifies these connections is Haiman’s $(n+1)^{n-1}$ theorem. By leveraging the geometric structure of the Hilbert scheme of n points in the plane \mathbb{C}^2 , Haiman proved that the dimension of the *diagonal coinvariant ring*,

$$\text{DR}_n := \mathbb{C}[x_1, \dots, x_n, y_1, \dots, y_n] / \langle \mathbb{C}[x_1, \dots, x_n, y_1, \dots, y_n]_+^{\otimes n} \rangle,$$

is $(n+1)^{n-1}$. Furthermore, he established that the bigraded Frobenius characteristic of this ring coincides with the symmetric function ∇e_n , where ∇ is the eigenoperator on Macdonald polynomials defined by $\nabla \tilde{H}_\mu = T_\mu \tilde{H}_\mu$, and e_n is the elementary symmetric function.

The celebrated *Shuffle Theorem*, conjectured in [HHL⁺05b] and proved in [CM18], further underscores the interplay between Macdonald polynomials, operators on symmetric functions, and Catalan combinatorics. This theorem provides an explicit combinatorial expansion for ∇e_n in terms of the combinatorics of labeled Dyck paths, showcasing the rich combinatorial structure. In an earlier work, the authors, along with Seung Jin Lee, offered a new proof of the shuffle theorem by connecting the Macdonald intersection polynomial $I_{\mu^{(1)}, \dots, \mu^{(n)}}$ to ∇e_{n-1} [KLO23].

Over the past two decades, the shuffle theorem has been extended the rational shuffle theorem [Mel21] and the Delta theorem [DM22, BHM⁺23]. The Loehr–Warrington conjecture [LW08], the final outstanding conjecture in this series, was very recently proved in [BHM⁺21], providing a combinatorial formula for ∇s_λ in terms of *nested Dyck paths*. We use another formulation expressing the combinatorial formula in the Loehr–Warrington conjecture in terms of certain *P-tableaux* [Ges84]:

$$\text{LW}_\lambda := q^{\text{adj}(\lambda)} \sum_{T \in \mathcal{T}(\lambda)} q^{\text{dinv}(T)} t^{\text{area}(T)} x^T.$$

See Section 4.1 for precise definition for the set $\mathcal{T}(\lambda)$ of certain *P-tableaux*, and the statistics *dinv* and *area*. It turns out the combinatorics of *P-tableau* beautifully match with the combinatorics of Macdonald piece polynomials.

1.4. Main results. Before stating our main results, we introduce a notation that we will use throughout this paper. For a partition λ inside a rectangle $R(n, k)$, we define $\tilde{\lambda}$ to be a conjugate partition of $(n - k - \lambda_k, \dots, n - k - \lambda_1)$ where we regard $\lambda_i = 0$ if $i > \ell(\lambda)$. The notation $\tilde{\lambda}$ is dependent on an ambient rectangle that λ lives in, which will be obvious from the context.

Now we state our main results. Note that the straightforward implication of Theorem 1.2 (b) and (c) is the Loehr–Warrington conjecture (Corollary 1.3).

Theorem 1.2. *Let μ be a partition with n corners, and $1 \leq k < n$. Then for a partition λ inside a rectangle $R(n, k)$ the symmetric function $I_{\mu, \lambda, k}$ satisfies the following:*

(a) For $N > |\mu| - |\tilde{\lambda}| - k$, we have

$$e_N^\perp I_{\mu, \lambda, k} = 0. \tag{1.6}$$

(b) We have

$$\frac{1}{T_{\mu^{[n]}}} e_{|\mu| - |\tilde{\lambda}| - k}^\perp I_{\mu, \lambda, k} = \nabla s_\lambda. \tag{1.7}$$

In particular, $e_{|\mu| - |\tilde{\lambda}| - k}^\perp I_{\mu, \lambda, k}$ does not depend on the partition μ (up to a constant).

(c) We have

$$\frac{1}{T_{\mu^{[n]}}} e_{|\mu| - |\tilde{\lambda}| - k}^\perp I_{\mu, \lambda, k} = (-1)^{\text{adj}(\tilde{\lambda})} \text{LW}_{\tilde{\lambda}}. \tag{1.8}$$

Here, e_n^\perp denotes the adjoint operator to the multiplication by e_n with respect to the Hall inner product. The precise definitions of the Loehr–Warrington formula LW_λ and $\text{adj}(\lambda)$ will be given in Section 4.

Corollary 1.3. (*The Loehr–Warrington conjecture*) We have $\nabla s_\lambda = (-1)^{\text{adj}(\lambda)} \text{LW}_\lambda$.

We conjecture that $(-1)^{\text{adj}(\bar{\lambda})} I_{\mu,\lambda,k}$ is Schur positive. We do not know the monomial positivity either, and Theorem 1.2 (c) can be thought of as supporting evidence. Finding a combinatorial description of the monomial expansion for $I_{\mu,\lambda,k}$ is a challenging open problem, as one would need to discover an ambient formula that includes the Loehr–Warrington formula as a small portion.

We also study the specialization of the Macdonald piece polynomials at $q = t = 1$. In particular, we describe the h -expansion in terms of $W_{\lambda,\mu} := [h_\mu](\nabla s_\lambda|_{q,t=1})$, the coefficient of h_μ in the h -expansion of $\nabla s_\lambda|_{q,t=1}$. In [KLO23], the authors proved that $W_{(1^n),\mu}$ is given by the Kreweras number, but we do not yet have a combinatorial description of $W_{\lambda,\mu}$ in general beyond the case $\lambda = (1^n)$. Now, we state a vast generalization of [KLO23, Theorem 1.3].

Theorem 1.4. *We have*

$$I_{\mu,\lambda,k}[X; 1, 1] = \sum_{\nu \subseteq \lambda} (-1)^{|\lambda| - |\nu|} d_{\lambda,\nu}^{(k)} \sum_{\tau} W_{\bar{\nu},\tau} h_{(\tau+1)^{|\mu|-k-|\bar{\nu}|}}$$

where

$$d_{\lambda,\nu}^{(k)} = \det \left(\begin{pmatrix} \lambda_i + k - i \\ \nu_j + k - j \end{pmatrix} \right)_{1 \leq i, j \leq k}$$

and $\bar{\nu}$ is taken regarding ν is inside a rectangle $R(n, k)$.

Example 1.5. Let $\mu = (4, 3, 2, 1)$, $k = 2$, $n = 4$ and $\lambda = (2, 1)$. Then we have

$$d_{(2,1),(2,1)}^{(2)} = 1, \quad d_{(2,1),(1,1)}^{(2)} = 3, \quad d_{(2,1),(2)}^{(2)} = 1, \quad d_{(2,1),(1)}^{(2)} = 3, \quad \text{and} \quad d_{(2,1),0}^{(2)} = 2.$$

From

$$\begin{aligned} \nabla s_{(2,2)}|_{q,t=1} &= -h_{(1,1,1,1)} + h_{(2,1,1)} + h_{(2,2)} - h_{(3,1)}, & \nabla s_{(2,1)}|_{q,t=1} &= -3h_{(1,1,1)} + 4h_{(2,1)} - h_{(3)}, \\ \nabla s_{(1,1)}|_{q,t=1} &= 2h_{(1,1)} - h_{(2)}, & \nabla s_{(2)}|_{q,t=1} &= -h_{(1,1)} + h_{(2)}, & \text{and} & \quad \nabla s_{(1)}|_{q,t=1} = h_{(1)}, \end{aligned}$$

we obtain

$$\begin{aligned} I_{\mu,\lambda,k}[X; 1, 1] &= (h_{(2,1,1,1,1,1,1,1)} - 3(-h_{(2,2,1,1,1,1,1)} + h_{(3,1,1,1,1,1,1)})) - (2h_{(2,2,1,1,1,1,1)} - h_{(3,1,1,1,1,1,1)}) \\ &\quad + 3(-3h_{(2,2,2,1,1)} + 4h_{(3,2,1,1,1)} - h_{(4,1,1,1,1,1)}) - 2(-h_{(2,2,2,2)} + h_{(3,2,2,1)} + h_{(3,3,1,1)} - h_{(4,2,1,1)}) \\ &= h_{(2,1,1,1,1,1,1,1)} + h_{(2,2,1,1,1,1,1)} - 2h_{(3,1,1,1,1,1,1)} - 9h_{(2,2,2,1,1)} + 12h_{(3,2,1,1,1)} - 3h_{(4,1,1,1,1,1,1)} \\ &\quad + 2h_{(2,2,2,2)} - 2h_{(3,2,2,1)} - 2h_{(3,3,1,1)} + 2h_{(4,2,1,1)}. \end{aligned}$$

1.5. organization. This paper is organized as follows. In Section 2, we introduce the necessary concepts in symmetric function theory and then present the extension of the science fiction conjecture (Conjecture 2.3), thereby motivating the importance of the Macdonald piece polynomial. In Section 3, we establish Theorem 1.2 (a) and (b) utilizing the plethystic formula for Macdonald polynomials by Garsia–Tesler–Haiman. In Section 4, we recall the Loehr–Warrington formula given in [LW08] and present its deformation in terms of the operators. In Section 5, we explore generalized Macdonald polynomials for filled diagrams and the column exchange rule introduced in [KLO22]. In Section 6, we finalize the proof of Theorem 1.2 by showing that the combinatorics of the Macdonald piece polynomial elegantly matches

with the Loehr–Warrington formula. In Section 7, we prove Theorem 1.4 on the h -expansion of Macdonald piece polynomials at $q = t = 1$. In Section 8, we discuss some future questions arising from this work. Finally, Appendix A includes data with interesting numbers, ratios of dimensions.

ACKNOWLEDGEMENT

The authors are grateful to François Bergeron, James Haglund, Seung Jin Lee and Brendon Rhoades for helpful conversations. The authors also appreciate to Raymond Chou for sharing his SAGE code for computing Garsia–Haiman modules. D. Kim was supported by NRF grants partially supported by Science Research Center Program through the National Research Foundation of Korea(NRF) Grant funded by the Korean Government and NRF grants 2022R1I1A1A01070620. J. Oh was supported by KIAS Individual Grant (HP083401) at June E Huh Center for Mathematical Challenges in Korea Institute for Advanced Study.

2. EXTENDING THE SCIENCE FICTION CONJECTURE WITH MACDONALD PIECE POLYNOMIALS

2.1. Preliminaries. A partition $\mu = (\mu_1, \dots, \mu_\ell)$ is a weakly decreasing sequence of positive integers. We say that μ is a partition of n , denoted by $\mu \vdash n$, if $|\mu| := \sum_i \mu_i = n$. We denote the length of μ , i.e., the number of its parts, by $\ell(\mu)$. We use *Young diagrams* in French notation to display a partition as

$$\mu = \{(i, j) \in \mathbb{Z}_{\geq 1} \times \mathbb{Z}_{\geq 1} : j \leq \mu_i\},$$

where each element of this set represents a *cell* in the diagram. The *conjugate partition* of μ , denoted by $\mu' = (\mu'_1, \mu'_2, \dots)$, is obtained by reflecting the Young diagram of μ along the diagonal $y = x$. We denote the empty partition by \emptyset .

For a cell u in a partition μ , we define the *arm (coarm)* of u , denoted by $\text{arm}_\mu(u)$ ($\text{coarm}_\mu(u)$), as the number of cells strictly to the right (left) of u in the same row. Similarly, we define the *leg (coleg)* of u , denoted by $\text{leg}_\mu(u)$ ($\text{coleg}_\mu(u)$), as the number of cells strictly above (below) u in the same column.

A *diagram* is a collection of cells in the first quadrant. A partition can be considered as a left-justified diagram, where the number of cells in each row is weakly decreasing from bottom to top. We denote a cell in a diagram in the i -th row, and j -th column by (i, j) . We usually represent a diagram by

$$D = [D^{(1)}, D^{(2)}, \dots],$$

where $D^{(j)} = \{i : (i, j) \in D\}$ is the row numbers of the j -th column of D . For a diagram D , the *size* $|D|$ is defined as the number of cells in D .

Let \mathbf{Sym} be the graded ring of symmetric functions in infinite variables $X = x_1, x_2, \dots$ over the ground field $\mathbb{C}(q, t)$. We use the notations from [Mac88] for families of symmetric functions indexed by a partition: m_λ for the *monomial symmetric function*, h_λ for the *homogeneous symmetric function*, e_λ for the *elementary symmetric function*, p_λ for the *power sum symmetric function*, s_λ for the *Schur function*, \tilde{H}_λ for the *(modified) Macdonald polynomial*. The (Hall) inner product on symmetric functions, denoted by $\langle -, - \rangle$, is defined as $\langle s_\lambda, s_\mu \rangle = \delta_{\lambda, \mu}$.

2.2. Extending the science fiction conjecture and the Macdonald positivity. We fix a partition μ with n corners $\{c_1, c_2, \dots, c_n\}$ and let $\mu^{(i)}$ be a partition obtained by removing a corner c_i . Recall that the science fiction conjecture suggests that

$$I_{\mu^{(1)}, \dots, \mu^{(n)}}[X; q, t] := \sum_{i=1}^n \left(\prod_{j \neq i} \frac{z_j}{z_j - z_i} \right) \tilde{H}_{\mu^{(i)}} = \text{grFrob} \left(\cap_{i=1}^n R_{\mu^{(i)}} \right), \quad (2.1)$$

where $z_i = T_{\mu^{(i)}}/T_\mu$. As mentioned before, $I_{\mu^{(1)}, \dots, \mu^{(n)}}[X; q, t]$ is a special case of the Macdonald piece polynomial $I_{\mu, \lambda, k}$ obtained by letting $\lambda = \emptyset$ and $k = 1$. Our goal is to further generalize (2.1) and refine the Macdonald positivity as well (Conjecture 2.3).

Now we consider partitions $\mu^S = \mu \setminus \{c_i : i \in S\}$ for $S \in \binom{[n]}{k}$. Note that $I_{\mu, \lambda, k}$ can be regarded as a linear combination of \tilde{H}_{μ^S} . We present the converse by representing each \tilde{H}_{μ^S} as a linear combination of $I_{\mu, \lambda, k}$.

Lemma 2.1. *We have*

$$\tilde{H}_{\mu^S} = \sum_{\lambda \subseteq R(n, k)} \frac{s_{\tilde{\lambda}}[z_{S^c}]}{\prod_{j \in S^c} z_j} I_{\mu, \lambda, k}$$

where $z_i = T_{\mu^{(i)}}/T_\mu = q^{-\text{coarm}(c_i)} t^{-\text{coleg}(c_i)}$.

Proof. Consider a matrix M whose rows are indexed by $\lambda \subseteq R(n, k)$ and columns are indexed by $S \in \binom{[n]}{k}$. Its entries are given by

$$M_{\lambda, S} = (-1)^{|\lambda|} \frac{s_{\tilde{\lambda}}[z_S] \prod_{j \in S^c} z_j}{\prod_{\substack{i \in S \\ j \in S^c}} (z_j - z_i)}.$$

Then let M' be a matrix whose columns are indexed by $\lambda \subseteq R(n, k)$ and rows are indexed by $S \in \binom{[n]}{k}$. Its entries are given by

$$M'_{S, \lambda} = \frac{s_{\tilde{\lambda}}[z_{S^c}]}{\prod_{j \in S^c} z_j}.$$

By Lemma 3.2 we have that MM' equals the identity matrix. \square

From now on we regard z_1, z_2, \dots, z_n as indeterminates and let

$$\tilde{H}_{\mu^S}[X; q, t, z] := \sum_{\lambda \subseteq k \times (n-k)} \frac{s_{\tilde{\lambda}}[z_{S^c}]}{\prod_{j \in S^c} z_j} I_{\mu, \lambda, k}.$$

Define Φ to be a specialization map on functions in z_1, z_2, \dots, z_n given by letting $z_i = T_{\mu^{(i)}}/T_\mu = q^{-\text{coarm}(c_i)} t^{-\text{coleg}(c_i)}$. By Lemma 2.1, we trivially have $\Phi(\tilde{H}_{\mu^S}[X; q, t, z]) = \tilde{H}_{\mu^S}$.

Let $\pi_{i, j}^1$ be the operator acting on functions in z_1, z_2, \dots, z_n defined by

$$\pi_{i, j}^1 f(z_1, z_2, \dots, z_n) = \frac{z_j f - z_i f|_{z_i \leftrightarrow z_j}}{z_j - z_i}.$$

Denoting $S = \{i_1 < i_2 < \dots < i_k\}$ and $S^c = \{j_1 < j_2 < \dots < j_{n-k}\}$, for a partition $\lambda \subseteq R(n, k)$ we define $\pi_{\lambda, S}$ to be a sequence of operators given by

$$\pi_{\lambda, S} = \prod_{(r, s) \in \lambda} \pi_{i_r, j_s}.$$

¹This operator is motivated by Butler's symmetric function [But94] and for $j = i + 1$, $\pi_{i, j}$ coincides with the Demazure operator π_i .

We may consider $\tilde{H}_{\mu^S}[X; q, t, z]$ as a function in z_1, z_2, \dots, z_n regarding $I_{\mu, \lambda, k}$ as constants. In this sense the operators $\pi_{i,j}$ act on $\tilde{H}_{\mu^S}[X; q, t, z]$ by

$$\pi_{\lambda, S}(\tilde{H}_{\mu^S}[X; q, t, z]) = \sum_{v \subseteq R(n, k)} \left(\pi_{\lambda, S} \frac{s_{\tilde{v}}[z_{S^c}]}{\prod_{j \in S^c} z_j} \right) I_{\mu, v, k}.$$

Finally, we define

$$\tilde{H}_{\mu^S}^{\lambda} := \Phi \left(\pi_{\lambda, S}(\tilde{H}_{\mu^S}[X; q, t, z]) \right). \quad (2.2)$$

Now we give a conjectural module theoretic interpretation of $\tilde{H}_{\mu^S}^{\lambda}$. To that end, we define a notation.

Definition 2.2. For $S \in \binom{[n]}{k}$ and a partition $\lambda \in R(n, k)$, denote by

$$S = \{i_1 < i_2 < \dots < i_k\}, \quad S^c = \{j_1 < j_2 < \dots < j_{n-k}\}.$$

We define $S(\lambda)$ to be a set of $S' \in \binom{[n]}{k}$ satisfying the following: Denote by

$$S \setminus S' = \{i_{a_1} < i_{a_2} < \dots < i_{a_r}\}, \quad S' \cap S^c = \{j_{b_1} > j_{b_2} > \dots > j_{b_r}\},$$

then we have $b_s \leq \lambda_{a_s}$ for $1 \leq s \leq r$. For example in the extreme, $S(\emptyset) = \{S\}$ and $S(R(n, k)) = \binom{[n]}{k}$. For example in the middle, letting $k=2$, $n=4$ and $\lambda = (2, 1)$, we have $S(\lambda) = \{\{1, 2\}, \{1, 3\}, \{2, 3\}, \{2, 4\}, \{3, 4\}\}$.

Conjecture 2.3. We have

$$\tilde{H}_{\mu^S}^{\lambda} = \text{grFrob} \left(\bigcap_{S' \in S(\lambda)} R_{\mu^{S'}} \right).$$

Letting λ to be a full rectangle $R(n, k)$ the right-hand side is the Frobenius characteristic of the full intersection. It is easy to check that the left-hand side $\tilde{H}_{\mu^S}^{R(n, k)}$ is independent of the choice of $S \in \binom{[n]}{k}$. Moreover, setting $k=1$ and $\lambda = R(n, 1)$, Conjecture 2.3 simply reduces to (2.1).

For partitions $\lambda^{(1)} \subset \lambda^{(2)}$, we trivially have $S(\lambda^{(1)}) \subset S(\lambda^{(2)})$. Therefore Conjecture 2.3 implies that

$$\tilde{H}_{\mu^S}^{\lambda^{(1)}} - \tilde{H}_{\mu^S}^{\lambda^{(2)}}$$

is Schur positive. Given a sequence $\emptyset = \lambda^{(1)} \subset \lambda^{(2)} \subset \dots \subset \lambda^{(\ell)} = R(n, k)$, we may obtain a sequence of symmetric functions that grows from $\tilde{H}_{\mu^S}^{R(n, k)}$ to $\tilde{H}_{\mu^S}^{\emptyset} = \tilde{H}_{\mu^S}$ in a Schur positive sense. This refines the Macdonald positivity.

3. PROOF OF THEOREM 1.2 (A) AND (B)

3.1. Technical Lemmas. We begin by recalling several technical lemmas. To state the first lemma, we review the following notation

$$M = (1-q)(1-t), \quad B_{\mu} = \sum_{c \in \mu} q^{\text{coarm}_{\mu}(c)} t^{\text{coleg}_{\mu}(c)}, \quad D_{\mu} = MB_{\mu} - 1,$$

$$\tilde{h}_{\mu} = \prod_{c \in \mu} (q^{\text{arm}_{\mu}(c)} - t^{\text{leg}_{\mu}(c)+1}), \quad \tilde{h}'_{\mu} = \prod_{c \in \mu} (t^{\text{leg}_{\mu}(c)} - q^{\text{arm}_{\mu}(c)+1}).$$

We will use brackets to denote plethystic substitutions. We use two different minus signs $-$ and ϵ to denote

$$p_k[-X] = -p_k[X], \quad \text{and} \quad p_k[\epsilon X] = (-1)^k p_k[X].$$

Furthermore, for a symmetric function f , define Π'_f as

$$\Pi'_f[X; q, t] = \nabla^{-1} f[X - \epsilon],$$

and $\text{rev}(f)$ as the q, t -reversal

$$\text{rev}(f) = f|_{q \rightarrow q^{-1}, t \rightarrow t^{-1}}.$$

In [GT96], the $*$ -inner product $\langle f, g \rangle_*$ is defined by

$$\langle p_\lambda, p_\mu \rangle_* = \begin{cases} (-1)^{|\lambda| - \ell(\lambda)} z_\lambda \prod_{i=1}^{\ell(\lambda)} (1 - q^{\lambda_i}) (1 - t^{\lambda_i}) & \text{if } \mu = \lambda, \\ 0 & \text{otherwise.} \end{cases}$$

The $*$ -inner product exhibits numerous useful properties. In particular, it commutes with the q, t -reversal: for symmetric functions f and g of homogeneous degree m , we have

$$\langle \text{rev}(f), \text{rev}(g) \rangle_* = (qt)^m \text{rev}(\langle f, g \rangle_*). \quad (3.1)$$

In addition, the modified Macdonald polynomials form an orthogonal basis with respect to the $*$ -inner product [GT96, Theorem 1.1]:

$$\langle \tilde{H}_\lambda, \tilde{H}_\mu \rangle_* = \begin{cases} \tilde{h}_\mu \tilde{h}'_\mu & \text{if } \mu = \lambda, \\ 0 & \text{otherwise.} \end{cases} \quad (3.2)$$

The following lemma [KLO23, Equation (3.4)] provides a formula for the Macdonald polynomial skewed by an elementary symmetric function. The proof of the lemma uses Garsia, Haiman, and Tesler's plethystic formula [GHT99, Theorem I.2].

Lemma 3.1. [KLO23] *For a partition $\mu \vdash n$, we have*

$$e_{n-m}^\perp \tilde{H}_\mu = (qt)^m T_\mu \sum_{\lambda \vdash m} \text{rev}(\Pi'_{\tilde{H}_\lambda}[D_\mu; q, t]) \frac{T_\lambda \tilde{H}_\lambda}{\tilde{h}_\lambda \tilde{h}'_\lambda}.$$

In the proof relating the Macdonald intersection polynomial $I_{\mu^{(1)}, \dots, \mu^{(k)}}$ and ∇e_{k-1} , a folklore lemma [KLO23, Lemma 3.3] played a key role. It turns out that this well-known lemma can be generalized to Lemma 3.2.

For λ and μ , we denote their coordinate-wise sum as $\lambda + \mu$ and concatenate them to form the vector (λ, μ) . Additionally, let st_k represent the staircase $(k-1, k-2, \dots, 1, 0)$. For $\lambda \subseteq R(n, k)$ and $\mu \subseteq R(n, n-k)$, we examine the expression $(\lambda + \text{st}_k, \mu + \text{st}_{n-k})$. In this case, we treat λ (a vector of length k) and μ (a vector of length $n-k$) by appending zeros as necessary. It can be easily verified that $(\lambda + \text{st}_k, \mu + \text{st}_{n-k})$ forms a rearrangement of the staircase st_n if and only if $\mu = \tilde{\lambda}$. For instance, when $n = 6$ and $k = 3$, we find $R(n, k) = (3, 3, 3)$. Given $\lambda = (3, 1)$, we compute $\tilde{\lambda} = (2, 2, 1)$, which leads to $\lambda + \text{st}_k = (5, 2, 0)$ and $\tilde{\lambda} + \text{st}_{n-k} = (4, 3, 1)$. Therefore, the combined vector $(\lambda + \text{st}_k, \tilde{\lambda} + \text{st}_{n-k}) = (5, 2, 0, 4, 3, 1)$ rearranges to form the staircase st_n .

Lemma 3.2. *Let z_1, \dots, z_n be variables. Let λ be a partition contained within the rectangle $R(n, k)$. For a partition μ of size $|\mu| \leq k(n-k) - |\lambda|$,*

$$\sum_{S \in \binom{[n]}{k}} \frac{s_\lambda[z_S] s_\mu[z_{S^c}]}{\prod_{i \in S, j \in S^c} (z_j - z_i)} = (-1)^{|\lambda|} \delta_{\tilde{\lambda}, \mu}. \quad (3.3)$$

Proof. Recall Jacobi's bi-alternant formula

$$s_\lambda(z_1, \dots, z_n) = \frac{a_{\lambda + \text{st}_n}(z_1, z_2, \dots, z_n)}{a_{\text{st}_n}(z_1, z_2, \dots, z_n)},$$

for the Schur polynomials, where $a_\lambda(z_1, \dots, z_n)$ is the alternating polynomial defined by

$$a_\lambda(z_1, \dots, z_n) = \det \left(z_j^{\lambda_i} \right)_{1 \leq i, j \leq n}.$$

In particular, $a_{\text{st}_n}(z_1, \dots, z_n)$ is the usual Vandermonde determinant. The left-hand side of (3.3) becomes

$$\sum_{S \in \binom{[n]}{k}} \frac{s_\lambda[z_S] s_\mu[z_{S^c}]}{\prod_{i \in S, j \in S^c} (z_j - z_i)} = \sum_{S \in \binom{[n]}{k}} \frac{a_{\lambda + \text{st}_k}(z_S) a_{\mu + \text{st}_{n-k}}(z_{S^c})}{a_{\text{st}_k}(z_S) a_{\text{st}_{n-k}}(z_{S^c}) \prod_{i \in S, j \in S^c} (z_j - z_i)} = \frac{\sum_{S \in \binom{[n]}{k}} \text{sgn}(S) a_{\lambda + \text{st}_k}(z_S) a_{\mu + \text{st}_{n-k}}(z_{S^c})}{a_{\text{st}_n}(z_{[n]})} \quad (3.4)$$

where $\text{sgn}(S) = (-1)^{\sum_{r=k+1}^n r - \sum_{s \in S} s}$.

The Laplace expansion of a determinant by complementary minors, we have

$$a_{(\lambda + \text{st}_k, \mu + \text{st}_{n-k})}(z_{[n]}) = \sum_{S \in \binom{[n]}{k}} \text{sgn}(S) a_{\lambda + \text{st}_k}(z_S) a_{\mu + \text{st}_{n-k}}(z_{S^c}).$$

By substituting this into (3.4), we obtain

$$\sum_{S \in \binom{[n]}{k}} \frac{s_\lambda[z_S] s_\mu[z_{S^c}]}{\prod_{i \in S, j \in S^c} (z_j - z_i)} = \frac{a_{(\lambda + \text{st}_k, \mu + \text{st}_{n-k})}(z_{[n]})}{a_{\text{st}_n}(z_{[n]})}.$$

Since the numerator $a_{(\mu + \text{st}_{n-k}, \lambda + \text{st}_k)}$ is an alternating polynomial of degree $|\lambda| + |\mu| + |\text{st}_k| + |\text{st}_{n-k}| \leq |\text{st}_n|$, it is nonzero if and only if the rearrangement of $(\mu + \text{st}_{n-k}, \lambda + \text{st}_k)$ is exactly the staircase st_n . In that case, we have

$$\frac{a_{(\lambda + \text{st}_k, \mu + \text{st}_{n-k})}(z_{[n]})}{a_{\text{st}_n}(z_{[n]})} = (-1)^{|\lambda|} \frac{a_{\text{st}_n}(z_{[n]})}{a_{\text{st}_n}(z_{[n]})} = (-1)^{|\lambda|}.$$

□

Remark 3.3. Lemma 3.2 can be seen as an application of the Atiyah–Bott localization theorem for equivariant cohomologies [AB84]. More precisely, let M be a smooth manifold that admits an action by a compact connected Lie group G . If the fixed points M^G consist of isolated points p with associated normal bundles ν_p in M with Euler classes $e(\nu_p)$, then the localization formula asserts that

$$\int_M \phi = \sum_{p \in M^G} \int_p \frac{i^* \phi}{e(\nu_p)}, \quad (3.5)$$

where $i : M^G \hookrightarrow M$.

If we let M be the Grassmannian $\text{Gr}(n, k)$ with the natural torus $G = (\mathbb{C}^*)^n$ action, then the fixed points are in one-to-one correspondence with k -subsets of $[n]$. Recall that the cohomology ring of the Grassmannian has an integral basis described by Schur functions. Localizing the Schur function indexed by the full-rectangle $R(n, k)$ in (3.5) proves Lemma 3.2.

3.2. Proof of Theorem 1.2 (a) and (b). Let μ be a partition of N with n corners $\{c_1, \dots, c_n\}$. Fix $0 \leq k \leq n$ and let λ be a partition inside the rectangular partition $R(n, k)$. Let $m = k(n - k) - |\lambda| = |\tilde{\lambda}|$. By the definition of $I_{\mu, \lambda, k}$, for an integer ℓ ,

$$e_{N-k-\ell}^\perp I_{\mu, \lambda, k} = \sum_{S \in \binom{[n]}{k}} \frac{s_\lambda[z_S] \prod_{j \in S^c} z_j}{\prod_{i \in S, j \in S^c} (z_j - z_i)} e_{N-k-\ell}^\perp \tilde{H}_\mu^S. \quad (3.6)$$

By Lemma 3.1, we have

$$e_{N-k-\ell}^\perp I_{\mu, \lambda, k} = (qt)^\ell T_{\mu^{[m]}} \sum_{v \vdash \ell} \sum_{S \in \binom{[m]}{k}} \frac{s_\lambda[z_S]}{\prod_{i \in S, j \in S^c} (z_j - z_i)} \text{rev}(\Pi'_{\tilde{H}_v} [D_{\mu^S}(q, t); q, t]) \frac{T_v \tilde{H}_v}{\tilde{h}_v \tilde{h}'_v}. \quad (3.7)$$

Here, we used $T_{\mu^{[m]}} = T_\mu^S \prod_{j \in S^c} z_j$. Note that

$$\text{rev}(\Pi'_{\tilde{H}_v} [D_{\mu^S}(q, t); q, t]) = T_v \tilde{H}_v \left[D_{\mu^{[m]}}(q^{-1}, t^{-1}) - \epsilon + \frac{M}{qt} z_{S^c}; q^{-1}, t^{-1} \right]$$

is a polynomial in z_{S^c} of degree $|v| = \ell$. If, on the right-hand side of (3.7), the degree of the polynomial in z_{S^c} is less than m , or equivalently if $\ell < m$, then by Lemma 3.2, it vanishes. This proves Theorem 1.2 (a).

Proceed to prove Theorem 1.2 (b). Now let $\ell = m$ in (3.7). Note that the leading term of $\text{rev}(\Pi'_{\tilde{H}_v} [D_{\mu^S}(q, t); q, t])$ is equal to

$$T_v \tilde{H}_v \left[\frac{M}{qt} z_{S^c}; q^{-1}, t^{-1} \right] = (qt)^{-m} \omega \tilde{H}_v [Mz_{S^c}; q, t],$$

by the symmetry relation $\omega \tilde{H}_\mu = T_\mu \text{rev}(\tilde{H}_\mu)$. Again, by Lemma 3.2, we only need to consider this leading term. Let us rewrite the right-hand side of (3.7) as

$$\begin{aligned} & T_{\mu^{[m]}} \nabla \left(\sum_{v \vdash \ell} \sum_{S \in \binom{[m]}{k}} \frac{s_\lambda[z_S]}{\prod_{i \in S, j \in S^c} (z_j - z_i)} \frac{\tilde{H}_v \omega \tilde{H}_v [Mz_{S^c}; q, t]}{\tilde{h}_v \tilde{h}'_v} \right) \\ &= T_{\mu^{[m]}} \nabla \left(\sum_{\rho \vdash \ell} \sum_{v \vdash \ell} \sum_{S \in \binom{[m]}{k}} \frac{s_\lambda[z_S]}{\prod_{i \in S, j \in S^c} (z_j - z_i)} \frac{\tilde{K}_{\rho', v}(q, t) \omega \tilde{H}_v [Mz_{S^c}; q, t]}{\tilde{h}_v \tilde{h}'_v} s_{\rho'} \right) \\ &= T_{\mu^{[m]}} \nabla \left(\sum_{\rho \vdash \ell} \sum_{S \in \binom{[m]}{k}} \frac{s_\lambda[z_S] s_{\rho'}[z_{S^c}]}{\prod_{i \in S, j \in S^c} (z_j - z_i)} s_{\rho'} \right). \end{aligned}$$

Here, in the first equation, $\tilde{K}_{\lambda, \mu}(q, t)$ is the *Macdonald–Kostka polynomial*, the Schur coefficient of Macdonald polynomials

$$\tilde{H}_\mu[X; q, t] = \sum_{\lambda \vdash n} \tilde{K}_{\lambda, \mu}(q, t) s_\lambda.$$

In the last equation, we used [Hag04, (2.67)]:

$$s_\lambda = \sum_{\beta \vdash |\lambda|} \frac{\tilde{K}_{\lambda', \beta}(q, t) \tilde{H}_\beta[MX; q, t]}{\tilde{h}_\beta \tilde{h}'_\beta}.$$

Now applying Lemma 3.2 again we obtain

$$T_{\mu^{[m]}} \nabla \left(\sum_{\rho \vdash \ell} \sum_{S \in \binom{[m]}{k}} \frac{s_\lambda[z_S] s_{\rho'}[z_{S^c}]}{\prod_{i \in S, j \in S^c} (z_j - z_i)} s_{\rho'} \right) = T_{\mu^{[m]}} \nabla s_{\tilde{\lambda}}.$$

4. LOEHR-WARRINGTON FORMULA

In this section, we recall the Loehr–Warrington formula, denoted by LW_λ , introduced in [LW08]. We then present our original contribution: Jacobi-Trudi type formula for LW_λ (Proposition 4.6), expressed using the operators defined in Definition 4.3. By leveraging the relations among these operators (as established in Lemma 4.7), we modify the Jacobi-Trudi type formula, ultimately arriving at the expression presented in (4.5). In the following sections, we complete the proof of Theorem 1.2 (c) by connecting the Macdonald intersection polynomial to (4.5).

4.1. The Loehr–Warrington formula. Throughout this section, we fix a poset \mathbf{P} on $\mathbb{Z}_{\geq 0} \times \mathbb{Z}_{\geq 1}$ defined as follows². For $(a, b), (c, d) \in \mathbb{Z}_{\geq 0} \times \mathbb{Z}_{\geq 1}$ we say $(a, b) \prec_{\mathbf{P}} (c, d)$ in \mathbf{P} if and only if $a + 1 < c$ or $a + 1 = c$ and $b \geq d$. Otherwise, we write $(a, b) \not\prec_{\mathbf{P}} (c, d)$. We also give a total ordering on $\mathbb{Z}_{\geq 0} \times \mathbb{Z}_{\geq 1}$ by a (mixed) lexicographic ordering. Define $(a, b) \prec_{\text{lex}} (c, d)$ if $a < c$, or $a = c$ and $b > d$. Note that the ordering of the second coordinate is reversed. For example, in terms of order of \mathbf{P} , we have $(4, 3) \prec_{\mathbf{P}} (5, 3)$ and $(4, 2) \not\prec_{\mathbf{P}} (5, 3)$. On the other hand, in terms of lexicographic order, we have $(4, 3) \prec_{\text{lex}} (5, 3)$ and $(4, 3) \prec_{\text{lex}} (4, 2)$.

Consider a tuple $L = (L_1, \dots, L_r)$ such that each L_i is a finite sub(multi)set of $\mathbb{Z}_{\geq 0} \times \mathbb{Z}_{\geq 1}$. We define the *diagonal inversion* dinv of L by

$$\begin{aligned} \text{dinv}(L) &= \sum_{i < j} \sum_{\substack{(a,b) \in L_i \\ (a',b') \in L_j}} \chi((a,b) \prec_{\text{lex}} (a',b')) \chi((a,b) \not\prec_{\mathbf{P}} (a',b')) \\ &= \sum_{i < j} \sum_{\substack{(a,b) \in L_i \\ (a',b') \in L_j}} (\chi(a = a') \chi(b > b') + \chi(a + 1 = a') \chi(b < b')). \end{aligned}$$

For a diagram D , let T be a filling of D with elements from $\mathbb{Z}_{\geq 0} \times \mathbb{Z}_{\geq 1}$. Denote $T(i, j)$ to be a filling in the cell $(i, j) \in D$. We use $T(i, j)_1$ (resp. $T(i, j)_2$) to refer to the first (resp. second) entry of $T(i, j)$. We define the area of T as:

$$\text{area}(T) = \sum_{(i,j) \in D} T(i, j)_1.$$

We use the notation $\text{dinv}(T)$ to denote $\text{dinv}(L_1, \dots, L_r)$, where L_i represents the set of entries in the i -th column of T .

We fix a partition $\lambda \subseteq R(n, k) = k \times (n - k)$. Let s be the size of the *Durfee square*, which is the maximal number such that $\lambda_s \geq s$ (if does not exist, $s = 0$). We define a *dinv adjustment* of λ denoted by $\text{adj}(\lambda)$ as

$$\text{adj}(\lambda) = \sum_{i=1}^s (\lambda_i - i).$$

Consider a n -vector

$$(\lambda_1 + k - 1, \lambda_2 + k - 2, \dots, \lambda_k, k, k + 1, \dots, n - 1)$$

where we regard $\lambda_i = 0$ if $i > \ell(\lambda)$, and define $v(\lambda)$ to be a vector obtained by sorting entries in the above vector in a weakly increasing order. Then we define the *bottom* of λ , denoted by $\text{bot}(\lambda)$, as an n -vector

²Our poset \mathbf{P} is both $(3+1)$ -free and $(2+2)$ -free, which is the condition in the famous Stanley–Stembridge conjecture regarding the e -positivity of chromatic symmetric functions.

given by

$$\text{bot}(\lambda) = (s+1, s+2, \dots, s+n) - v(\lambda),$$

where the subtraction is performed element-wise. We also define a *pivot* of λ , denoted by $\text{piv}(\lambda)$, to be a vector (a_1, a_2, \dots, a_s) of length s where a_i is a number satisfying

$$v(\lambda)_{a_i} = v(\lambda)_{a_i+1} = \lambda_{s+1-i} + k - (s+1-i),$$

i.e., the indices where $v(\lambda)$ is non increasing. Lastly we associate a diagram $D(\lambda)$ to the partition λ as

$$D(\lambda) = [[\text{bot}(\lambda)_1, s], [\text{bot}(\lambda)_2, s], \dots, [\text{bot}(\lambda)_n, s]]$$

and define a set $\mathcal{T}(\lambda)$ consisting of filling T of $D(\lambda)$ satisfying the following conditions:

- $T(i+1, j) \succ_{\mathbf{P}} T(i, j)$ and $T(i, j-1) \not\succeq_{\mathbf{P}} T(i, j)$
- for each $j > k-s$ if $j \in \text{piv}(\lambda)$ we have $T(\text{bot}(\lambda)_j, j)_1 > 0$
- for each $j > k-s$ if $j \notin \text{piv}(\lambda)$ we have $T(\text{bot}(\lambda)_j, j)_1 = 0$.

Note that the first condition says that T (after taking appropriate reflection) is a \mathbf{P} -tableau defined in [Gas96]. The *Loehr–Warrington formula* LW_λ is defined by

$$\text{LW}_\lambda := q^{\text{adj}(\lambda)} \sum_{T \in \mathcal{T}(\lambda)} q^{\text{dinv}(T)} t^{\text{area}(T)} x^T, \quad (4.1)$$


where $x^T = \prod x_{T(i,j)_2}$.

Example 4.1. Let $k=2$ and $n=5$, and consider $\lambda = (3, 2)$ within a 2×3 rectangle. To compute $\text{adj}(\lambda)$, we first observe that $\lambda_2 \geq 2$, and $\lambda_3 < 3$. Thus, the q -adjustment $\text{adj}(\lambda)$ is calculated as $(\lambda_1 - 1) + (\lambda_2 - 2) = 2$. The vector $v(\lambda)$ is the weakly increasing rearrangement $(2, 2, 3, 4, 4)$ of $(\lambda_1 + 1, \lambda_2 + 0, k + 1, \dots, n) = (3 + 1, 2 + 0, 2, 3, 4)$. The bottom of λ is given by $\text{bot}(\lambda) = (3, 4, 5, 6, 7) - (2, 2, 3, 4, 4) = (1, 2, 2, 2, 3)$. The pivots are 1 and 4 since $v(\lambda)_1 = v(\lambda)_2$ and $v(\lambda)_4 = v(\lambda)_5$. Finally, $D(\lambda) = [[1, 2], [2, 2], [2, 2], [2, 2], [3, 2]] = [\{1, 2\}, \{2\}, \{2\}, \{2\}, \emptyset]$.

Remark 4.2. In [LW08], the Loehr–Warrington formula LW_λ is described in terms of nested labeled Dyck paths. We demonstrate how $T \in \mathcal{T}(\lambda)$ can be naturally associated with such nested labeled Dyck paths. Specifically, we read the entries of each row of T from right to left and construct a Dyck path that satisfies the following condition: the number of boxes between the i -th upstep and the diagonal $y = x$ equals the first coordinate of the i -th entry (counted from the right) in T , while the second coordinate corresponds to its label. Finally, the starting point for each Dyck path is determined as follows: the Dyck path corresponding to the i -th row of T (counting from the bottom) begins at (a, a) , where $a = \text{piv}(\lambda)_i - i$. Refer to Figure 1 for an example where $\lambda = (3, 2)$.

4.2. Jacobi–Trudi type formula for the Loehr–Warrington formula. We provide a Jacobi–Trudi type formula for LW_λ in terms of the operators $\mathfrak{h}_m, \bar{\mathfrak{h}}_m$, and $\hat{\mathfrak{h}}_m$, defined below.

Definition 4.3. We define \mathbf{C}_m to be a set of all \mathbf{P} -chains $\{(a_1, b_1) \prec_{\mathbf{P}} (a_2, b_2) \prec_{\mathbf{P}} \dots \prec_{\mathbf{P}} (a_m, b_m)\}$ where each $(a_\ell, b_\ell) \in \mathbb{Z}_{\geq 0} \times \mathbb{Z}_{\geq 1}$. Then $\bar{\mathbf{C}}_m$ (resp $\hat{\mathbf{C}}_m$) is a subset of \mathbf{C}_m consisting of elements satisfying $a_1 = 0$ (resp $a_1 > 0$). Obviously, $\mathbf{C}_m = \bar{\mathbf{C}}_m \cup \hat{\mathbf{C}}_m$.

FIGURE 1. From $T \in \mathcal{F}(\lambda)$ to the nested labeled Dyck paths.

Let $\mathbf{y} = \{y_{i,j}\}_{i \in \mathbb{Z}_{\geq 0}, j \in \mathbb{Z}_{\geq 1}}$ be a set of indeterminates. For a multiset A of elements in $\mathbb{Z}_{\geq 0} \times \mathbb{Z}_{\geq 1}$, let y^A be the monomial whose exponent of $y_{i,j}$ is given by the number of elements (i, j) in A . We define the operators \mathfrak{h}_m , $\bar{\mathfrak{h}}_m$, and $\hat{\mathfrak{h}}_m$ acting on a polynomial ring $\mathbb{F}[\mathbf{y}]$ (\mathbb{F} is a ground field containing $\mathbb{C}(q)$) by describing their actions on a monomial as follows:

$$\mathfrak{h}_m \cdot y^A = \sum_{L \in \mathbf{C}_m} q^{\text{dinv}(L,A)} y^{(L,A)}, \quad \bar{\mathfrak{h}}_m \cdot y^A = \sum_{L \in \bar{\mathbf{C}}_m} q^{\text{dinv}(L,A)} y^{(L,A)}, \quad \hat{\mathfrak{h}}_m \cdot y^A = \sum_{L \in \hat{\mathbf{C}}_m} q^{\text{dinv}(L,A)} y^{(L,A)},$$

where $y^{(L,A)} = y^L y^A$. We then extend this linearly to define operators \mathfrak{h}_n , $\bar{\mathfrak{h}}_n$, and $\hat{\mathfrak{h}}_n$. If $m < 0$ then they are all zero operators.

For example, we have

$$\hat{\mathfrak{h}}_2 \bar{\mathfrak{h}}_3 \mathfrak{h}_4 \cdot 1 = \sum_{L \in \hat{\mathbf{C}}_2 \times \bar{\mathbf{C}}_3 \times \mathbf{C}_4} q^{\text{dinv}(L)} y^L$$

where $y^L = y^{L_1} y^{L_2} y^{L_3}$ for $L = (L_1, L_2, L_3)$.

We say two operators are the same if they act on $\mathbb{F}[\mathbf{y}]$ in the same way. For example, $\mathfrak{h}_m = \hat{\mathfrak{h}}_m + \bar{\mathfrak{h}}_m$, which is trivial from $\mathbf{C}_m = \bar{\mathbf{C}}_m \cup \hat{\mathbf{C}}_m$.

The operators \mathfrak{h}_n , $\bar{\mathfrak{h}}_n$, and $\hat{\mathfrak{h}}_n$ do not generally commute (for example, see Lemma 4.7). Nevertheless, operators of the same type do commute (Corollary 4.5). The following lemma, which is a step for proving Corollary 4.5, is inspired by the symmetry of chromatic quasisymmetric functions. Though the proof parallels [SW16, Lemma 4.4, Theorem 4.5] using an involution on proper colorings (or P -tabloids), we provide a proof for the sake of completeness.

Lemma 4.4. *For nonnegative integers n, m , there exists a bijection*

$$\text{SW} : \mathbf{C}_n \times \mathbf{C}_m \rightarrow \mathbf{C}_m \times \mathbf{C}_n$$

such that for $L \in \mathbf{C}_n \times \mathbf{C}_m$, $y^L = y^{\text{SW}(L)}$ and $\text{dinv}(L) = \text{dinv}(\text{SW}(L))$.

Proof. Let $L = (L_1, L_2) \in \mathbf{C}_n \times \mathbf{C}_m$. Consider a directed graph $G(L)$ on the elements of $L_1 \cup L_2$ (with possible repeated entries), where edges are defined as $(a, b) \rightarrow (c, d)$ for pairs satisfying $(a, b) <_{\text{lex}} (c, d) <_{\text{lex}} (a+1, b)$. Or equivalently, when $(a, b) <_{\text{lex}} (c, d)$ and $(a, b) \not\prec_P (c, d)$. It is straightforward to see that $G(L)$ is bipartite since elements in L_1 are not connected to each other, and the same applies to elements in L_2 .

We claim that every connected component of $G(L)$ is a directed path of the form $(a_1, b_1) <_{\text{lex}} (a_2, b_2) <_{\text{lex}} \cdots <_{\text{lex}} (a_j, b_j)$, where the membership of (a_i, b_i) alternates between L_1 and L_2 . To prove the claim, we first show that each vertex has an out-degree at most one. Suppose not. Without loss of generality, assume $(a, b) \in L_1$ has out-degree more than one. Then there exist $(c, d), (c', d') \in L_2$ such that $(a, b) \rightarrow (c, d)$ and $(a, b) \rightarrow (c', d')$. Therefore, we have $(a, b) <_{\text{lex}} (c, d) <_{\text{lex}} (a+1, b)$ and $(a, b) <_{\text{lex}} (c', d') <_{\text{lex}} (a+1, b)$. This contradicts the fact that $(c, d) \in L_2$ and $(c', d') \in L_2$ are comparable in \mathbf{P} . Similarly, each vertex has an in-degree at most one. Finally, there cannot be a directed cycle because every directed edge is increasing in terms of the $<_{\text{lex}}$ ordering. Combining all these proves the claim.

For a connected component (path) P of $G(L)$, let

$$\text{SW}(P) := \begin{cases} (P_1, P_2) & \text{if } P \text{ has an odd number of vertices,} \\ (P_2, P_1) & \text{if } P \text{ has an even number of vertices,} \end{cases}$$

where $P_1 = P \cap L_1$ and $P_2 = P \cap L_2$. Then define $\text{SW}(L) := (\cup_P \text{SW}(P)_1, \cup_P \text{SW}(P)_2)$, where the union runs over all connected components of $G(L)$. The map SW does not change the elements in $L_1 \cup L_2$, thus $y^L = y^{\text{SW}(L)}$. It is clear to see that $\text{SW}(L) \in \mathbf{C}_m \times \mathbf{C}_n$ and that SW is involutive. Finally, note that dinv counts the number of directed edges from an element in L_1 to L_2 . For each connected component P in $G(L)$, this number is preserved by SW . Thus total dinv statistic of L is preserved under the involution SW . \square

Corollary 4.5. *We have $\mathfrak{h}_n \mathfrak{h}_m = \mathfrak{h}_m \mathfrak{h}_n$, $\bar{\mathfrak{h}}_n \bar{\mathfrak{h}}_m = \bar{\mathfrak{h}}_m \bar{\mathfrak{h}}_n$, and $\hat{\mathfrak{h}}_n \hat{\mathfrak{h}}_m = \hat{\mathfrak{h}}_m \hat{\mathfrak{h}}_n$.*

Proof. The map SW in Lemma 4.4 directly explains that $\mathfrak{h}_n \mathfrak{h}_m = \mathfrak{h}_m \mathfrak{h}_n$ and it is easy to check that SW induces bijections $\bar{\mathbf{C}}_n \times \bar{\mathbf{C}}_m \rightarrow \bar{\mathbf{C}}_m \times \bar{\mathbf{C}}_n$ and $\hat{\mathbf{C}}_n \times \hat{\mathbf{C}}_m \rightarrow \hat{\mathbf{C}}_m \times \hat{\mathbf{C}}_n$. We conclude $\bar{\mathfrak{h}}_n \bar{\mathfrak{h}}_m = \bar{\mathfrak{h}}_m \bar{\mathfrak{h}}_n$ and $\hat{\mathfrak{h}}_n \hat{\mathfrak{h}}_m = \hat{\mathfrak{h}}_m \hat{\mathfrak{h}}_n$. \square

Consider an n by n square matrix $W = (W_{i,j})$ whose entry $W_{i,j}$ is an operator acting on $\mathbb{F}[y]$. We let $\det(W)$ to be the operator defined by

$$\det(W) = \sum_{\sigma \in \mathfrak{S}_n} (-1)^{\text{sgn}(\sigma)} W_{\sigma(1),1} W_{\sigma(2),2} \cdots W_{\sigma(n),n}.$$

For example, we have $\det \begin{pmatrix} \mathfrak{h}_2 & \mathfrak{h}_3 \\ \mathfrak{h}_4 & \mathfrak{h}_5 \end{pmatrix} = \mathfrak{h}_2 \mathfrak{h}_5 - \mathfrak{h}_4 \mathfrak{h}_3$. Upon this notation, we provide a Jacobi–Trudi type formula for the Loehr–Warrington formula LW_λ .

Proposition 4.6 (Jacobi–Trudi type formula). *For integers $k < n$ and a partition $\lambda \subseteq R(n, k)$ we associate an n by n square matrix $W(\lambda)$*

$$W(\lambda)_{i,j} = \begin{cases} \mathfrak{h}_{v(\lambda)_{j-i+1}} & \text{if } j \leq k-s \\ \bar{\mathfrak{h}}_{v(\lambda)_{j-i+1}} & \text{if } j > k-s \text{ and } j \text{ is an entry of } \text{piv}(\lambda) \\ \hat{\mathfrak{h}}_{v(\lambda)_{j-i+1}} & \text{if } j > k-s \text{ and } j \text{ is not an entry of } \text{piv}(\lambda), \end{cases}$$

where s is the size of the Durfee square of λ . Then we have

$$q^{-\text{adj}(\lambda)} \text{LW}_\lambda = \det(W(\lambda)) \cdot 1|_{y_{i,j} = t^i x_j}$$

Proof. For a permutation $\sigma \in \mathfrak{S}_n$, let $D(\sigma)$ be the diagram

$$D(\sigma) = [[\text{bot}(\lambda)_1, s+1-\sigma(1)], [\text{bot}(\lambda)_2, s+2-\sigma(2)], \dots, [\text{bot}(\lambda)_n, s+n-\sigma(n)]].$$

Note that $D(\text{id}) = D(\lambda)$. Let C_σ be the set of fillings A of $D(\sigma)$ with elements in $\mathbb{Z}_{\geq 0} \times \mathbb{Z}_{\geq 1}$ such that

- $A(i+1, j) >_{\mathbf{P}} A(i, j)$,
- for each $j > k-s$, if $j \in \text{piv}(\lambda)$, we have $A(\text{bot}(\lambda)_j, j)_1 > 0$,
- for each $j > k-s$, if $j \notin \text{piv}(\lambda)$, we have $A(\text{bot}(\lambda)_j, j)_1 = 0$.

If $s+i-\sigma(i) < \text{bot}(\lambda)_i$ for some i , we consider that there is no filling of $D(\sigma)$.

By definition, it is straightforward to see that

$$\det(W(\lambda)) \cdot 1 = \sum_{\sigma \in \mathfrak{S}_n} (-1)^{\text{sgn}(\sigma)} \prod_{i=1}^n W(\lambda)_{\sigma(i), i} \cdot 1 = \sum_{\sigma \in \mathfrak{S}_n} (-1)^{\text{sgn}(\sigma)} \sum_{A \in C_\sigma} q^{\text{dinv}(A)} y^A. \quad (4.2)$$

We construct a sign-reversing involution $\widetilde{\text{SW}}$ on the set

$$B_\lambda := \{(A, \sigma) : \sigma \in \mathfrak{S}_n, A \in C_\sigma\} \setminus \{(T, \text{id}) : T \in \mathcal{T}(\lambda)\}$$

such that

$$\text{sgn}(\sigma) = -\text{sgn}(\sigma'), \quad y^A = y^{A'}, \quad \text{and} \quad \text{dinv}(A) = \text{dinv}(A'),$$

where we denoted $(A', \sigma') = \widetilde{\text{SW}}(A, \sigma)$. Via such involution, all terms in (4.2) except for the ones from $\{(T, \text{id}) : T \in \mathcal{T}(\lambda)\}$ cancels. This implies that we have

$$\det(W(\lambda)) \cdot 1 = \sum_{T \in \mathcal{T}(\lambda)} q^{\text{dinv}(T)} y^T,$$

which completes the proof.

We proceed to construct $\widetilde{\text{SW}}$. For a given $(A, \sigma) \in B_\lambda$, a cell (i, j) in $D(\sigma)$ is called *bad* in A if either of the following conditions holds:

- $(i, j-1)$ is empty with $i \geq \text{bot}(\lambda)_{j-1}$,
- $A(i, j-1) >_{\mathbf{P}} A(i, j)$.

For $(A, \sigma) \in B_\lambda$, there must be at least one bad cell. Let $r = r(A)$ be the smallest i such that there is a bad cell (i, j) . Once r is determined, let c be the largest j such that the cell (r, j) is bad. Define the sets

$$S_{c-1}(A) = \{A(i, c-1) : r \leq i \leq s+(c-1)-\sigma(c-1)\}, \quad \text{and} \quad S_c(A) = \{A(i, c) : r < i \leq s+c-\sigma(c)\}.$$

We then define $\widetilde{\text{SW}}(A, \sigma) := (A', \sigma(c-1, c))$, where A' is obtained from A by applying the map SW (as defined in Lemma 4.4) to $(S_{c-1}(A), S_c(A))$ while leaving other parts unchanged, and $(c-1, c)$ is the transposition swapping $c-1$ and c . We show that A' is in $C_{\sigma'}$ where $\sigma' = \sigma(c-1, c)$. It suffices to show $A'(r-1, c-1) <_{\mathbf{P}} A'(r, c-1)$ and $A'(r, c) <_{\mathbf{P}} A'(r+1, c)$ so that each column is still \mathbf{P} -chain. Note that $A'(r, c) = A(r, c)$ and

$A'(r+1, c)$ equals $A(r+1, c)$ or $A(r, c-1)$, implying $A'(r, c) <_{\mathbf{P}} A'(r+1, c)$. If $A(r-1, c-1) \not\prec_{\mathbf{P}} A(r+1, c)$, together with $A(r-1, c) \not\prec_{\mathbf{P}} A(r-1, c-1)$ from the minimality of r , we have

$$\begin{aligned} A(r-1, c) &<_{\mathbf{P}} A(r, c) <_{\mathbf{P}} A(r+1, c) \\ A(r-1, c) &\not\prec_{\mathbf{P}} A(r-1, c-1) \not\prec_{\mathbf{P}} A(r+1, c) \end{aligned}$$

which is impossible. Since $A'(r-1, c-1) = A(r-1, c-1)$ and $A'(r, c-1)$ equals $A(r, c-1)$ or $A(r+1, c)$, we conclude $A'(r-1, c-1) <_{\mathbf{P}} A'(r, c-1)$.

It is straightforward to see that (r, c) remains a bad cell in A' , whose minimality is trivial. This guarantees that the map $\widetilde{S\bar{W}}$ is indeed an involution. Moreover, it is evident that the map $\widetilde{S\bar{W}}$ is sign-reversing and preserves y^A and $\text{div}(A)$. □

4.3. Reformulation of Loehr–Warrington formula. We reformulate the Loehr–Warrington formula LW_{λ} from Proposition 4.6. Our main tool is Lemma 4.7, whose proof will be provided in Section 4.4. For a matrix A , let $T_{ij}(A)$ be the matrix obtained by moving the j -th column to the position of the i -th column and shifting the ℓ -th column to the $(\ell+1)$ -th position for $i < \ell < j$. For example, we have

$$A = \begin{pmatrix} | & | & | & | & | \\ c_1 & c_2 & c_3 & c_4 & c_5 \\ | & | & | & | & | \end{pmatrix} \longrightarrow T_{25}A = \begin{pmatrix} | & | & | & | & | \\ c_1 & c_5 & c_2 & c_3 & c_4 \\ | & | & | & | & | \end{pmatrix}.$$

Lemma 4.7. *For an integer vector $v = (v_1, v_2, \dots, v_n)$ of length $n \geq 2$, consider an $n \times n$ matrix of operators $V = (V_{i,j})$ given by*

$$V_{i,j} = \begin{cases} \hat{h}_{v_i+j-1} & \text{if } j < n \\ \bar{h}_{v_i+n-1} & \text{if } j = n \end{cases}$$

Then we have $(-q)^{n-1} \det V = \det T_{1,n}(V)$.

Now we modify the matrix $W(\lambda)$ given in Proposition 4.6. Let s be the maximal number such that $\lambda_s \geq s$. Set $W^{(0)} = W(\lambda)$ and then construct $W^{(1)}, \dots, W^{(s)}$ recursively as follows:

- (Step 1) Given $W^{(i)}$, let $V^{(i+1)} = T_{ab}(W^{(i)})$ where $a = k - s + i$ and $b = \text{piv}(\lambda)_i$.
- (Step 2) Add the $(\text{piv}(\lambda)_i + 1)$ -th column to the $(k - s + i)$ -th column in $V^{(i+1)}$ to obtain $W^{(i+1)}$.

By Lemma 4.7, after performing Step 1, the determinant multiplies by $(-q)^{\text{piv}(\lambda)_i - (k-s+i)} = (-q)^{\lambda_{s-i+1} - (s-i+1)}$. The property of a determinant together with Corollary 4.5 implies that Step 2 does not change the determinant. We conclude that

$$(-q)^{\text{adj}(\lambda)} \det(W(\lambda)) = \det(W^{(s)}). \quad (4.3)$$

It is easy to check that $W^{(s)}$ is given by

$$W_{i,j}^{(s)} = \begin{cases} h_{\lambda_{k+1-j+j-i}} & \text{if } j \leq k \\ \hat{h}_{j-i} & \text{if } j > k. \end{cases} \quad (4.4)$$

Finally, by Proposition 4.6, we conclude

$$LW_\lambda = (-1)^{\text{adj}(\lambda)} \det(W^{(s)}) \cdot 1 \Big|_{y_{i,j}=t^i x_j}. \quad (4.5)$$

Example 4.8. Let $\lambda = (3, 2)$ as in Example 4.1. Recall that $\nu(\lambda) = (2, 2, 3, 4, 4)$. Then $W^{(0)} = W(\lambda)$, $W^{(1)}$ and $W^{(2)}$ can be computed as below:

$$\begin{aligned} W^{(0)} &= \begin{pmatrix} \bar{h}_2 & \hat{h}_2 & \hat{h}_3 & \bar{h}_4 & \hat{h}_4 \\ \vdots & \vdots & \vdots & \vdots & \vdots \end{pmatrix} \xrightarrow{\text{Step1}} \begin{pmatrix} \bar{h}_2 & \hat{h}_2 & \hat{h}_3 & \bar{h}_4 & \hat{h}_4 \\ \vdots & \vdots & \vdots & \vdots & \vdots \end{pmatrix} \xrightarrow{\text{Step2}} \begin{pmatrix} \mathfrak{h}_2 & \hat{h}_2 & \hat{h}_3 & \bar{h}_4 & \hat{h}_4 \\ \vdots & \vdots & \vdots & \vdots & \vdots \end{pmatrix} = W^{(1)}, \\ W^{(1)} &= \begin{pmatrix} \mathfrak{h}_2 & \hat{h}_2 & \hat{h}_3 & \bar{h}_4 & \hat{h}_4 \\ \vdots & \vdots & \vdots & \vdots & \vdots \end{pmatrix} \xrightarrow{\text{Step1}} \begin{pmatrix} \mathfrak{h}_2 & \bar{h}_4 & \hat{h}_2 & \hat{h}_3 & \hat{h}_4 \\ \vdots & \vdots & \vdots & \vdots & \vdots \end{pmatrix} \xrightarrow{\text{Step2}} \begin{pmatrix} \mathfrak{h}_2 & \mathfrak{h}_4 & \hat{h}_2 & \hat{h}_3 & \hat{h}_4 \\ \vdots & \vdots & \vdots & \vdots & \vdots \end{pmatrix} = W^{(2)}. \end{aligned}$$

4.4. Proof of Lemma 4.7. In this section we prove Lemma 4.7. In particular, we prove Lemma 4.9 which naturally implies Lemma 4.7.

Lemma 4.9. For an integer vector $\nu = (\nu_1, \nu_2, \dots, \nu_n)$ of length $n \geq 2$, consider n by n matrices V and V' given as

$$V_{i,j} = \begin{cases} \hat{h}_{\nu_i} & \text{if } j = 1 \\ \bar{h}_{\nu_i+n-1} & \text{if } j = 2 \end{cases}, \quad V'_{i,j} = \begin{cases} \bar{h}_{\nu_i+n-1} & \text{if } j = 1 \\ \hat{h}_{\nu_i} & \text{if } j = 2. \end{cases}$$

and $V_{i,j} = V'_{i,j} = \mathfrak{h}_{\nu_i+j-2}$ for $j > 2$. Then we have $q \det(V) + \det(V') = 0$.

Proof. We proceed with induction on n . For the base case $n = 2$, we prove the claim by imitating the argument in Lemma 4.4. Let $\nu = (\nu_1, \nu_2) = (a, b)$. Then, we need to prove

$$q \det(V) + \det(V') = q(\hat{h}_a \bar{h}_{b+1} - \hat{h}_b \bar{h}_{a+1}) + (\bar{h}_{a+1} \hat{h}_b - \bar{h}_{b+1} \hat{h}_a) = 0, \quad (4.6)$$

which is equivalent to

$$q \hat{h}_a \bar{h}_{b+1} + \bar{h}_{a+1} \hat{h}_b = q \hat{h}_b \bar{h}_{a+1} + \bar{h}_{b+1} \hat{h}_a. \quad (4.7)$$

To that end, we define a bijection

$$\widetilde{SW}: \hat{\mathbf{C}}_a \times \bar{\mathbf{C}}_{b+1} \cup \bar{\mathbf{C}}_{a+1} \times \hat{\mathbf{C}}_b \rightarrow \hat{\mathbf{C}}_b \times \bar{\mathbf{C}}_{a+1} \cup \bar{\mathbf{C}}_{b+1} \times \hat{\mathbf{C}}_a$$

as follows. Let $L = (L_1, L_2) \in \hat{\mathbf{C}}_a \times \bar{\mathbf{C}}_{b+1} \cup \bar{\mathbf{C}}_{a+1} \times \hat{\mathbf{C}}_b$. Consider a directed graph on the elements of L , where edges are defined as $(a, b) \rightarrow (c, d)$ for those satisfying $(a, b) <_{\text{lex}} (c, d)$ and $\text{dinv}(\{(a, b)\}, \{(c, d)\}) = 1$. As we have seen in Lemma 4.4, every connected component of this directed graph forms a directed path. Note that in L there is only one element of the form $(0, *)$. Let $P = (P_1, P_2)$ be the directed path containing this element, where $P_1 = L_1 \cap P$ and $P_2 = L_2 \cap P$. Then, let $(L'_1, L'_2) = SW(L_1 \setminus P_1, L_2 \setminus P_2)$. Then, $\widetilde{SW}(L)$ is defined by

$$\widetilde{SW}(L) = \begin{cases} (P_1 \cup L'_1, P_2 \cup L'_2) & \text{if } P \text{ is an odd path,} \\ (P_2 \cup L'_1, P_1 \cup L'_2) & \text{if } P \text{ is an even path.} \end{cases}$$

Since the involution \widetilde{SW} preserves the dinv statistic, and swapping the even path changes an additional ± 1 to the dinv , it is straightforward to see that

$$\text{dinv}(L) + \chi(L \in \hat{\mathbf{C}}_a \times \bar{\mathbf{C}}_{b+1}) = \text{dinv}(\widetilde{SW}(L)) + \chi(\widetilde{SW}(L) \in \hat{\mathbf{C}}_b \times \bar{\mathbf{C}}_{a+1}).$$

This proves (4.7), thus (4.6) follows.

Now let $n \geq 3$ and assume the claim is true for any number smaller than n . We define n by n matrices W, W', Z and Z' as

$$\begin{aligned} W_{i,1} &= \bar{h}_{v_i+1}, & W_{i,2} &= \hat{h}_{v_i+n-2}, & W'_{i,1} &= \hat{h}_{v_i+n-2}, & W'_{i,2} &= \bar{h}_{v_i+1}, \\ Z_{i,1} &= \bar{h}_{v_i+n-2}, & Z_{i,2} &= \hat{h}_{v_i+1}, & Z'_{i,1} &= \hat{h}_{v_i+1}, & Z'_{i,2} &= \bar{h}_{v_i+n-2}, \end{aligned}$$

and $W_{i,j} = W'_{i,j} = Z_{i,j} = Z'_{i,j} = h_{v_i+j-2}$ for $j > 2$. We claim the following

$$q \det(V) + \det(V') = -\det(W) - q \det(W') \quad (4.8)$$

$$\det(W) = -\det(Z) \quad (4.9)$$

$$\det(W') = -\det(Z') \quad (4.10)$$

$$\det(Z) = -q \det(Z') \quad (4.11)$$

which finishes the proof.

To prove (4.8), pick any $1 \leq a < b \leq n$. Then it is enough to show

$$q \det \begin{pmatrix} V_{a,1} & V_{a,2} \\ V_{b,1} & V_{b,2} \end{pmatrix} + \det \begin{pmatrix} V'_{a,1} & V'_{a,2} \\ V'_{b,1} & V'_{b,2} \end{pmatrix} = -\det \begin{pmatrix} W_{a,1} & W_{a,2} \\ W_{b,1} & W_{b,2} \end{pmatrix} - q \det \begin{pmatrix} W'_{a,1} & W'_{a,2} \\ W'_{b,1} & W'_{b,2} \end{pmatrix}$$

which is equivalent to (after a rearrangement of terms)

$$\left(q \det \begin{pmatrix} \hat{h}_{v_a} & \bar{h}_{v_a+1} \\ \hat{h}_{v_b+n-2} & \bar{h}_{v_b+n-1} \end{pmatrix} + \det \begin{pmatrix} \bar{h}_{v_a+1} & \hat{h}_{v_a} \\ \bar{h}_{v_b+n-1} & \hat{h}_{v_b+n-2} \end{pmatrix} \right) - \left(q \det \begin{pmatrix} \hat{h}_{v_b} & \bar{h}_{v_b+1} \\ \hat{h}_{v_a+n-2} & \bar{h}_{v_a+n-1} \end{pmatrix} + \det \begin{pmatrix} \bar{h}_{v_b+1} & \hat{h}_{v_b} \\ \bar{h}_{v_a+n-1} & \hat{h}_{v_a+n-2} \end{pmatrix} \right) = 0$$

Each summand on the left-hand side is zero due the induction hypothesis for $n = 2$.

Now we show (4.9). Starting from W , we apply following column operations:

- subtract the n -th column from the second column, then multiply (-1) to the second column
- swap the first and the second columns
- subtract the third column from the second column, then multiply (-1) to the second column.

It is easy to check that the resulting matrix is Z and $\det(W) = -\det(Z)$ follows. Similar argument also shows (4.10).

Lastly, we show (4.11). Let $I = \{1, 2, \dots, n\} \setminus \{3, 4\}$ and pick any $A \subset \{1, 2, \dots, n\}$ such that $|A| = n - 2$. Then denoting $Z_{A,I}$ to be a submatrix given by restricting row indices to A and column indices to I , induction hypothesis implies $\det(Z_{A,I}) + q \det(Z'_{A,I}) = 0$. This completes the proof. \square

Corollary 4.10. *With the same notations as in Lemma 4.9, define n by n matrices \hat{V} and \hat{V}' by*

- $\hat{V}_{i,j} = V_{i,j}$ and $\hat{V}'_{i,j} = V'_{i,j}$ for $j = 1, 2$
- $\hat{V}_{i,j} = \hat{V}'_{i,j} = \hat{h}_{v_i+j-2}$ for $j > 2$.

Then we have $q \det(\hat{V}) + \det(\hat{V}') = 0$.

Proof. Let $\Phi : \mathbb{F}[y] \rightarrow \mathbb{F}[y]$ be a linear operator defined by $\Phi(y^A) = y^A$ if $\sum_j A_{0,j} = 1$, otherwise $\Phi(y^A) = 0$. Then Φ sends $(q \det(V) + \det(V')) \cdot 1$ to $(q \det(\hat{V}) + \det(\hat{V}')) \cdot 1$. Therefore we deduce $(q \det(\hat{V}) + \det(\hat{V}')) \cdot 1 = 0$ and this is enough to imply $q \det(\hat{V}) + \det(\hat{V}') = 0$ as an operator. \square

Proof of Lemma 4.7. We define n by n matrices $V^{(1)}, V^{(2)}, \dots, V^{(n)}$ given as

$$V_{i,j}^{(\ell)} = \begin{cases} \hat{h}_{v_i+j-1} & \text{if } j < \ell \\ \bar{h}_{v_i+n-1} & \text{if } j = \ell \\ \hat{h}_{v_i+j-2} & \text{if } j > \ell. \end{cases}$$

We claim that $\det(V^{(\ell)}) = -q \det(V^{(\ell+1)})$. Let $I = \{\ell, \ell+1, \dots, n\}$ and pick any $A \subset \{1, 2, \dots, n\}$ such that $|A| = n - \ell + 1$. Then we have $q \det(V_{A,I}^{(\ell+1)}) + \det(V_{A,I}^{(\ell)}) = 0$ by the previous corollary. Therefore the claim is proved and we obtain $\det(V^{(1)}) = (-q)^{n-1} \det(V^{(n)})$. \square

5. GENERALIZED MACDONALD POLYNOMIALS FOR FILLED DIAGRAMS

5.1. Macdonald polynomials for filled diagrams and HHL formula. Given a diagram D , we regard a word $\sigma \in \mathbb{Z}_{\geq 1}^{|D|}$ as an assignment of integer entries to the cells of D , proceeding from left to right within each row, and from the top row to the bottom row. In this way, we consider $\sigma(u)$ for $u \in D$ as an integer assigned for u in the assignment of σ in D . The *content* $\text{cont}_D(\sigma, i)$ of σ in the i -th row of D is defined by the (infinite) vector whose a -th component is given by the number of a 's in the assignment of σ in the i -th row of D .

The celebrated result of Haglund–Haiman–Loehr [HHL05a] gives Macdonald polynomials using two statistics inv_μ and maj_μ . We will use slightly different convention for inv , which we now explain. For a diagram D , and cells $u = (i, j)$ and $v = (i', j')$ of D , we say that a pair (u, v) is an *attacking pair* if either

$$(1) \ i = i' \text{ and } j < j' \quad \text{or} \quad (2) \ i = i' + 1 \text{ and } j > j'.$$

For a word $\sigma \in \mathbb{Z}_{\geq 1}^{|D|}$, we consider a pair (u, v) of cells in D to be an *inversion pair* of σ if (u, v) is an attacking pair and $\sigma(u) > \sigma(v)$. We denote the set of inversion pairs of σ by $\text{Inv}_D(\sigma)$ and define $\text{inv}_D(\sigma)$ to be

$$\text{inv}_D(\sigma) := q^{|\text{Inv}_D(\sigma)|}.$$

A cell u is *descent* of σ if $\sigma(u) > \sigma(v)$, where the cell v is the cell right below u , i.e. $u = (i, j)$ and $v = (i-1, j)$. Define $\text{Des}_D(\sigma)$ to be the set of descents of σ .

A *filling* of D is a function $f : D \rightarrow \mathbb{F}$, where we take \mathbb{F} as a field containing $\mathbb{C}(q, t)$. Then a *filled diagram* (D, f) is a pair of a diagram and a filling on it. We define $\text{maj}_{(D,f)}(\sigma)$ as the product of $f(u)$ over all cells u which are descents of σ , i.e.,

$$\text{maj}_{(D,f)}(\sigma) := \prod_{u \in \text{Des}_D(\sigma)} f(u).$$

Note that bottom cells of D cannot be a descent so the value $f(u)$ for bottom cell u is redundant in a definition of $\text{maj}_{(D,f)}$. Therefore we usually represent a filled diagram (D, f) omitting values on bottom

cells. Finally, $\text{stat}_{(D,f)}(w)$ is defined by the product of inv_D and $\text{maj}_{(D,f)}$,

$$\text{stat}_{(D,f)}(\sigma) := \text{inv}_D(\sigma) \text{maj}_{(D,f)}(\sigma).$$

In [KLO22], a generalization of modified Macdonald polynomial of a filled diagram (D, f) is defined by

$$\tilde{H}_{(D,f)} := \sum_{\sigma \in \mathbb{Z}_{\geq 1}^{|D|}} \text{stat}_{(D,f)}(\sigma) x^\sigma.$$

We also call $\tilde{H}_{(D,f)}$ as the *Macdonald polynomial* for a filled diagram (D, f) . Indeed, this generalizes the concept of the celebrated HHL formula [HHL05a, HHL08], which gives the combinatorial formula for the modified Macdonald polynomial. To state, we recall some notations first. Previously we defined $\text{arm}_\mu(u)$ and $\text{leg}_\mu(u)$ for a partition μ and these were generalized to a more general shape [HHL08].

Let β be a diagram $\beta = [[\beta_1], [\beta_2], \dots, [\beta_k]]$ for some positive integers β_i 's. For a cell $u \in \beta$, the arm_β and leg_β are defined as follows:

$$\text{leg}_\beta(u) = \beta_i - j = (\text{number of cells strictly north of } u),$$

$$\text{arm}_\beta(u) = |\{i' \in \{1, 2, \dots, i-1\} : j-1 \leq \beta_{i'} < \beta_i\}| + |\{i' \in \{i, i+1, \dots, k\} : j \leq \beta_{i'} \leq \beta_i\}|.$$

In other words, $\text{arm}_\beta(u)$ represents the number of cells strictly to the right of u in columns of height $\beta_{i'} \leq \beta_i$ or strictly to the left of the cell just below u in columns of height $\beta_{i'} < \beta_i$. Define the *standard filling* f_β^{st} on β by

$$f_\beta^{\text{st}}(u) = q^{-\text{arm}_\beta(u)} t^{\text{leg}_\beta(u)}.$$

Now we can give the celebrated HHL formula using this standard filling.

Theorem 5.1. [HHL05a, HHL08] *Let μ be a partition and β be the diagram obtained by rearranging columns of μ . Then we have*

$$\tilde{H}_\mu = \sum_{\sigma \in \mathbb{Z}_{\geq 1}^{|\beta|}} \text{stat}_{(\beta, f_\beta^{\text{st}})}(\sigma) x^\sigma. \quad (5.1)$$

Haglund, Haiman, and Loehr [HHL05a] proved (5.1) for $\beta = \mu$, and later generalized it to any rearrangements β in [HHL08].

Theorem 5.1 shows an example of two different filled diagrams giving the same Macdonald polynomials. Another (rather simple) operation on filled diagrams, called the *cycling rule*, also gives the same Macdonald polynomials. Given a diagram $D = [D^{(1)}, \dots, D^{(\ell)}]$, consider the diagram

$$D' = [D^{(2)}, \dots, D^{(\ell)}, D^{(1)} + 1],$$

where $I+1 = \{a+1 : a \in I\}$ for an interval I . In other words, D' is obtained by moving the leftmost column of D to the end on the right and shifting it up by one cell. The cells in D and D' correspond naturally in a bijective manner. For a filling f of D , we can define a filling f' of D' inherited from f through this natural bijection. We define $\text{Cyc}(D, f) := (D', f')$ and it is direct to see

$$\tilde{H}_{(D,f)}[X] = \tilde{H}_{(D',f')}[X]. \quad (5.2)$$

5.2. Column exchange rule. In [KLO22], the authors together with Seung Jin Lee introduced a *column exchange rule* (Proposition 5.3), which we exploit in the next section.

Definition 5.2. Let (D, f) be a filled diagram satisfying the following for some positive integer j :

- denoting $D = [D^{(1)}, D^{(2)}, \dots]$ there exist integers $n \geq m$ such that $D^{(j)} = [1, n]$ and $D^{(j+1)} = [1, m]$,
- if $n > m$ we have $\frac{f(2,j)}{f(2,j+1)} = \dots = \frac{f(m,j)}{f(m,j+1)} = q^{-1}f(m+1, j)$, and
- if $n = m$ we have $\frac{f(2,j)}{f(2,j+1)} = \dots = \frac{f(m,j)}{f(m,j+1)}$.

Then we define $S_j(D, f)$ to be a filled diagram (D', f') given by: $D' = [D^{(1)}, \dots, D^{(j+1)}, D^{(j)}, \dots]$ and

- $f'(i, c) = f(i, c)$ if $c \neq j, j+1$,
- $f'(i, j) = f(i, j+1)$, and
- $f'(i, j+1) = f(i, j)$ if $i \neq m+1$ and $f'(i, j+1) = q^{-1}f(i, j)$ if $i = m+1$.

As it is obvious that the map S_j is injective, we can naturally define S_j^{-1} . We say that two filled diagrams are column-equivalent $(D, f) \equiv (D', f')$, if (D', f') is obtained from (D, f) by applying a sequence of maps of the form S_j or S_j^{-1} . An example of the application of the map S_j will be given in the next section (see bottom of Figure 3).

Proposition 5.3. *If $(D, f) \equiv (D', f')$ then there exists a bijection $\phi: \mathbb{Z}^{|D|} \rightarrow \mathbb{Z}^{|D'|}$ satisfying*

- (ϕ 1) $\text{stat}_{(D,f)}(\sigma) = \text{stat}_{(D',f')}(\phi(\sigma))$
- (ϕ 2) $\text{cont}_D(\sigma, i) = \text{cont}_{D'}(\phi(\sigma), i)$ for all i .
- (ϕ 3) $\text{iDes}(\text{std}(\sigma)) = \text{iDes}(\text{std}(\phi(\sigma)))$ where std denotes the standardization of a word.

Indeed, if $(D, f) \equiv (D', f')$, property (ϕ 1) together with the condition $x^\sigma = x^{\phi(\sigma)}$, a weaker condition than (ϕ 2), are sufficient to imply $\tilde{H}_{(D,f)} = \tilde{H}_{(D',f')}$. The map ϕ also satisfies (ϕ 2) and (ϕ 3). Later, we use these additional properties to derive a more refined identity, see (6.6) for example.

6. PROOF OF THEOREM 1.2 (C)

In this section, we show Theorem 1.2 (c). In particular we will connect $\left(e_{|\mu|-|\lambda|-k}^\perp I_{\mu, \tilde{\lambda}, k}\right)$ with the Loehr–Warrington formula through (4.5). By the shape independence in Theorem 1.2 (b), it is sufficient to provide a combinatorial formula for $\left(e_{|\mu|-|\lambda|-k}^\perp I_{\mu, \tilde{\lambda}, k}\right)$ for a specific partition μ of n corners. We fix μ to be the augmented staircase $\delta = \delta_{n,N} := \underbrace{(n, \dots, n)}_N, n-1, \dots, 1$ for a large enough N ($N > |\lambda|$ suffices). We outline the proof strategy:

- In Section 6.1, we define auxiliary filled diagrams (δ_S, f_S) , (δ_S, f_S^z) and $(\text{Rec}^{\leq m}, g_S^z)$
- In Section 6.2, we go over a technical process to obtain (6.8) from $\left(e_{|\delta|-|\lambda|-k}^\perp I_{\delta, \tilde{\lambda}, k}\right)$, imitating arguments in [KLO23, Section 6]
- In Section 6.3, we illustrate a clear connection between (6.8) and (reformulated) Jacobi-Trudi type formula (4.5) for the Loehr–Warrington formula LW_λ .

6.1. Deformation of filled diagrams. Let the corners of δ be indexed by c_1, c_2, \dots, c_n from top to bottom. Additionally, for a k -subset $S = \{i_1 < i_2 < \dots < i_k\}$ of $[n]$, let δ^S denote the partition obtained from δ by deleting the corners c_i 's for $i \in S$. We rearrange columns of δ^S by moving the i_1, i_2, \dots, i_k -th columns all the way to the left in this order and denote by β . Now we define $(\delta_S, f_S) := \text{Cyc}^k(\beta, f_\beta^{\text{st}})$ (the diagram δ_S should not be confused with δ^S). According to Theorem 5.1 and (5.2), we may use (δ_S, f_S) for the computation of \tilde{H}_{μ^S} i.e. $\tilde{H}_{\mu^S} = \tilde{H}_{(\delta_S, f_S)}$. For example, Figure 2 illustrates the described process for $\delta = \delta_{4,3} = (4, 4, 4, 3, 2, 1)$ and $S = \{2, 3\}$. The first figure shows $(\delta^S, f_{\delta^S}^{\text{st}})$ and the second shows $(\beta, f_\beta^{\text{st}})$ for the described rearrangement β . Applying the operator Cyc twice, we obtain (δ_S, f_S) as shown in the third figure.

Now, we introduce indeterminates z_1, z_2, \dots , and define (δ_S, f_S^z) which is a z -deformation (δ_S, f_S) . Denote $S = \{i_1 < i_2 < \dots < i_k\}$ and $S^c = \{j_1 < j_2 < \dots < j_{n-k}\}$. Then the filling f_S^z is given by

$$f_S^z(a, b) = q^{\frac{z_{N+n+1-a}}{z_{j_b}}} \quad \text{for } b \leq n-k, \quad f_S^z(a, n-k+b) = \begin{cases} q^{\frac{z_{N+n+1-a}}{z_{i_b}}} & \text{if } N+n+1-a \in S^c \\ q^{\frac{z_{N+n+1-a}}{z_{i_b}}} & \text{otherwise} \end{cases} \quad \text{for } b \leq k.$$

The fourth figure in Figure 2 shows (δ_S, f_S^z) corresponding to (δ_S, f_S) in the third. A straight calculation shows that under the specialization given by

$$z_j = \begin{cases} q^{1-j} t^{j+1-N-n} & \text{if } j \leq n \\ q^{-n} t^{j+1-N-n} & \text{if } j > n \end{cases} \quad (6.1)$$

(δ_S, f_S^z) recovers as (δ_S, f_S) .




FIGURE 2. The process to obtain (δ_S, f_S^z) .

Lastly for a positive integer $m \leq N$, we define $(\text{Rec}^{\leq m}, g_S^z)$ to be a sub filled diagram of (δ_S, f_S^z) obtained by restricting to the first m rows. Note that the diagram $\text{Rec}^{\leq m} = \underbrace{[[1, m], \dots, [1, m]]}_{n-k} \underbrace{[[2, m], \dots, [2, m]]}_k$ is independent of the choice of S . In Figure 3, top figure shows $(\text{Rec}^{\leq m}, g_S^z)$ obtained from the fourth figure in Figure 2.

For $S \in \binom{[n]}{k}$, the filled diagrams (δ_S, f_S^z) are not in general column-equivalent to each other. However, after truncating bottom rows, they are column-equivalent as shown in the following lemma.




FIGURE 3. An example of $\text{Rec}^{\leq m}$ (top figure) and an illustration of the proof of Lemma 6.1 (bottom figures).

Lemma 6.1. For $m \geq 2$, let (D'_S, f'_S) be a sub filled diagram of (δ_S, f_S) obtained by restricting to the m -th row and above. Then for any $I, J \in \binom{[n]}{k}$ we have

$$(D'_I, f'_I) \equiv (D'_J, f'_J).$$

Proof. It suffices to prove the case when $m = 2$. Consider a filled diagram (D, f) given by $D = (\underbrace{n, \dots, n}_{N-1}, n-1, \dots, 1)$ and $f(a, b) = \frac{q^{z_{N+n-a}}}{z_j}$. Then $(D'_S, f'_S) \equiv (D, f)$ for any $S \in \binom{[n]}{k}$, as illustrated in the bottom of Figure 3. The left side shows (D, f) , and after applying the operators S_3 and S_2 , we obtain the filled diagram on the right. This diagram is derived from (δ_S, f_S^z) in Figure 2 by deleting the first row. \square

6.2. Reducing to the $\text{Rec}^{\leq N}$. In this section we compute $(e_{|\delta|-k-|\lambda|}^\perp I_{\delta, \lambda, k})$ starting from filled diagrams (δ_S, f_S) for the computation of \tilde{H}_{δ_S} . Our goal is to obtain (6.8). This section consists of a technical process where the idea of most of the arguments was given in [KLO23, Section 6].

Definition 6.2. Given a diagram D and a positive integer $\ell \leq |D|$, we define $\text{Sub}(D, \ell)$ to be the set of assignments of ℓ cells of D by positive integers. We think of $R \in \text{Sub}(D, \ell)$ as a function $R : D' \rightarrow \mathbb{Z}_{\geq 1}$ for some subdiagram $D' \subseteq D$ with $|D'| = \ell$. We let $\text{rsum}(R)$ (resp $\text{csum}(R)$) to be a vector whose i -th entry is a number of cells in the i -th row (resp column) of D' . We also define an induced full assignment $\bar{R} : D \rightarrow \mathbb{Z}_{\geq 1}$ given by assigning $1, 2, \dots, |D| - \ell$ for cells in $D \setminus D'$ in a reverse reading order, and by letting $\bar{R}(c) = R(c) + |D| - \ell$ for cells $c \in D'$. See Figure 4 for an example. We let $x^R = \prod_{c \in D'} x_{R(c)}$ and lastly for a filled diagram (D, f) , we abuse the notation for $\text{stat}_{(D, f)}(R)$ instead of $\text{stat}_{(D, f)}(\bar{R})$.

			1
4			1
3			

6	5	4	7
10	3	2	7
9	1		

FIGURE 4. The left shows $R \in \text{Sub}(D, 4)$ for $D = \text{Rec}^{\leq 3}$ and the right show the induced full assignment \bar{R} . We have $\text{rsum}(R) = (1, 2, 1)$, $\text{csum} = (2, 0, 0, 2)$ and $x^R = x_1^2 x_3 x_4$.

Lemma 6.3. *We have*

$$e_{|\delta|-k-|\lambda|}^\perp I_{\delta, \bar{\lambda}, k} = (-1)^{|\bar{\lambda}|} \sum_{S \in \binom{[m]}{k}} \frac{s_{\bar{\lambda}}[w_S] \prod_{j \in S^c} w_j}{\prod_{\substack{i \in S \\ j \in S^c}} (w_j - w_i)} \sum_{R \in \text{Sub}(\delta_S, |\lambda|)} \text{stat}_{(\delta_S, f_S)}(R) x^R \quad (6.2)$$

where $w_i = T_{\delta^{(i)}} / T_\delta$.

Proof. For any nonnegative integer m , it is elementary to see

$$e_{|\delta_S|-k-m}^\perp \left(\sum_{\sigma \in \mathbb{Z}_{\geq 1}^{|\delta_S|}} \text{stat}_{(\delta_S, f_S)}(\sigma) x^\sigma \right) = \sum_{R \in \text{Sub}(\delta_S, m)} \text{stat}_{(\delta_S, f_S)}(R) x^R,$$

and the proof follows. \square

For indeterminates z_1, z_2, \dots we consider the z -deformation of the right hand side of (6.2):

$$(-1)^{|\bar{\lambda}|} \sum_{S \in \binom{[m]}{k}} \frac{s_{\bar{\lambda}}[z_S] \prod_{j \in S^c} z_j}{\prod_{\substack{i \in S \\ j \in S^c}} (z_j - z_i)} \sum_{R \in \text{Sub}(\delta_S, |\lambda|)} \text{stat}_{(\delta_S, f_S^z)}(R) x^R. \quad (6.3)$$

Since $w_i = T_{\delta^{(i)}} / T_\delta = q^{1-i} t^{i+1-N-n}$, we have $w_i = z_i$ under the specialization (6.1). Therefore we have

$$(6.3) \xrightarrow{\text{Specialize by (6.1)}} \left(e_{|\delta|-k-|\bar{\lambda}|}^\perp I_{\delta, \bar{\lambda}, k} \right) \quad (6.4)$$

Now our goal is to reduce (6.3) to (6.8).

The following lemma follows from a simple computation so we leave it as an exercise to readers.

Lemma 6.4. *Let w'_0 and w_0 be the longest permutations of length $|\delta_S|$ and $|\text{Rec}^{\leq N}|$. The following are true:*

- (1) *the quantity $(\prod_{j \in S^c} z_j) \text{stat}_{(\delta_S, f_S^z)}(w'_0)$ is independent of the choice of $S \in \binom{[m]}{k}$*
- (2) *the quantity $(\prod_{j \in S^c} z_j) \text{stat}_{(\text{Rec}^{\leq N}, g_S^z)}(w_0)$ is independent of the choice of $S \in \binom{[m]}{k}$.*
- (3) *the quantity $(\prod_{j \in S^c} z_j) \text{stat}_{(\delta_S, f_S^z)}(w'_0)$ specializes to $T_{\delta^{[m]}}$ under (6.1).*

For the rest of the paper, we denote the quantity in Lemma 6.4 (1) and (2) by C_1 and C_2 respectively.

Definition 6.5. Let $D = \text{Rec}^{\leq m}$ for some $m \leq N$ and consider $R \in \text{Sub}(D, \ell)$ denoted by $R : D' \rightarrow \mathbb{Z}_{>0}$. We say that a cell $c = (i, j) \in D'$ is bad if $c' = (i+1, j) \in \text{Des}_D(\bar{R})$ where \bar{R} is the associated full assignments. Note that

every cell $c \in D'$ located in a top row is bad. We define $\gamma(R)$ to be a vector whose i -th entry is the number of bad cells in the i -th row. For example, we have $\gamma(R) = (1, 0, 1)$ for R in Figure 4

Now we prove Lemma 6.6 and 6.7, key ingredients to obtain (6.8).

Lemma 6.6. *For vectors α and γ in $\mathbb{Z}_{\geq 0}^m$, we have*

- (1) $\left(\sum_{\substack{R \in \text{Sub}(\text{Rec}^{\leq m}, \ell) \\ \text{rsum}(R) = \alpha, \gamma(R) = \gamma}} \text{stat}_{(\text{Rec}^{\leq m}, g_S^z)}(R) x^R \right)$ is symmetric in variables z_S and symmetric in variables z_{S^c} ,
(2) if $|\alpha| - |\gamma| < |\lambda|$ we have

$$\sum_{S \in \binom{[m]}{k}} \frac{s_{\tilde{\lambda}}[z_S] \prod_{j \in S^c} z_j}{\prod_{\substack{i \in S \\ j \in S^c}} (z_j - z_i)} \sum_{\substack{R \in \text{Sub}(\text{Rec}^{\leq m}, \ell) \\ \text{rsum}(R) = \alpha, \gamma(R) = \gamma}} \text{stat}_{(\text{Rec}^{\leq m}, g_S^z)}(R) x^R = 0. \quad (6.5)$$

Proof. We denote $A_{\alpha, S}$ and $A_{\alpha, S}^\gamma$ to be

$$A_{\alpha, S} = \sum_{\substack{R \in \text{Sub}(\text{Rec}^{\leq m}, \ell) \\ \text{rsum}(R) = \alpha}} \text{stat}_{(\text{Rec}^{\leq m}, g_S^z)}(R) x^R, \quad A_{\alpha, S}^\gamma = \sum_{\substack{R \in \text{Sub}(\text{Rec}^{\leq m}, \ell) \\ \text{rsum}(R) = \alpha, \gamma(R) = \gamma}} \text{stat}_{(\text{Rec}^{\leq m}, g_S^z)}(R) x^R.$$

For $S = \{i_1 < \dots < i_k\}$ and $S^c = \{j_1 < \dots < j_{n-k}\}$, we have $A_{\alpha, S}^\gamma = A_{\alpha, [n-k+1, n]}^\gamma(z_{j_1}, \dots, z_{j_{n-k}}, z_{i_1}, \dots, z_{i_k})$ by the property of g_S^z . Therefore it is enough to show (1) when $S = [n-k+1, n]$. Set $S = [n-k+1, n]$ and we first show that $A_{\alpha, S}$ is symmetric in variables z_S and symmetric in variables z_{S^c} . Let (D', g') be the filled diagram obtained by applying the map S_1 (see Definition 5.2) to $(\text{Rec}^{\leq m}, g_S^z)$. Then $D' = \text{Rec}^{\leq m}$ and the filling g' can be obtained from g_S^z by swapping z_1 and z_2 . By Proposition 5.3, we have

$$\sum_{\substack{R \in \text{Sub}(\text{Rec}^{\leq m}, \ell) \\ \text{rsum}(R) = \alpha}} \text{stat}_{(\text{Rec}^{\leq m}, g_S^z)}(R) x^R = \sum_{\substack{R \in \text{Sub}(D', \ell) \\ \text{rsum}(R) = \alpha}} \text{stat}_{(D', g')}(R) x^R. \quad (6.6)$$

The right hand side is obtained from the left hand side by swapping z_1 and z_2 , this shows $A_{\alpha, S}$ is symmetric in z_1 and z_2 . Applying the similar argument, we conclude that $A_{\alpha, S}$ is symmetric in variables z_S and symmetric in variables z_{S^c} .

Now pick any $R \in \text{Sub}(\text{Rec}^{\leq m}, \ell)$ with $\text{rsum}(R) = \alpha$ and $\gamma(R) = \gamma$. On the $(i+1)$ -th row there are $(\alpha_i - \gamma_i)$ many cells that do not belong to $\text{Des}_{\text{Rec}^{\leq m}}(\tilde{R})$. Since the filling $g_S^z(i+1, j)$ on the $(i+1)$ -th row is of the form $\frac{q z_{N+n-i}}{z_a}$ for some $1 \leq a \leq n$, we can write

$$\text{stat}_{(\text{Rec}^{\leq m}, g_S^z)}(R) = \text{stat}_{(\text{Rec}^{\leq m}, g_S^z)}(w_0) \left(\frac{q^c \prod_{i=1}^n z_i^{d_i}}{\prod_{i=1}^{N-1} z_{N+n-i}^{\alpha_i - \gamma_i}} \right)$$

for some c and d_i 's, where w_0 is the longest permutation of length $|\text{Rec}^{\leq m}|$. We conclude that there exists a polynomial $P_{\alpha, S}^\gamma(z_1, z_2, \dots, z_n)$ of degree $(|\alpha| - |\gamma|)$ such that

$$A_{\alpha, S}^\gamma = \text{stat}_{(\text{Rec}^{\leq m}, g_S^z)}(w_0) \left(\frac{P_{\alpha, S}^\gamma}{\prod_{i=1}^{N-1} z_{N+n-i}^{\alpha_i - \gamma_i}} \right).$$

From

$$\frac{A_{\alpha, S}}{\text{stat}_{(\text{Rec}^{\leq m}, g_S^z)}(w_0)} = \frac{\sum_{\gamma} A_{\alpha, S}^\gamma}{\text{stat}_{(\text{Rec}^{\leq m}, g_S^z)}(w_0)} = \sum_{\gamma} \frac{P_{\alpha, S}^\gamma}{\prod_{i=1}^{N-1} z_{N+n-i}^{\alpha_i - \gamma_i}},$$

the left hand side is symmetric in variables z_S and symmetric in variables z_{S^c} , which implies that each $P_{\alpha,S}^\gamma$ is symmetric in variables z_S and symmetric in variables z_{S^c} . This establishes (1).

To prove (2), from the symmetry established in (1), we may simply write $P_{\alpha,S}^\gamma = P_\alpha^\gamma(z_{S^c}, z_S)$ for $P_\alpha^\gamma = P_{\alpha, [n-k+1, n]}^\gamma$. Since $C = \prod_{j \in S^c} z_j \left(\text{stat}_{(\text{Rec}^{\leq m}, g_S^z)}(w_0) \right)$ is independent of the choice of $S \in \binom{[n]}{k}$ by Lemma 6.4 (2), the left hand side of (6.5) equals

$$\frac{C}{\prod_{i=1}^{N-1} z_{N+n-i}^{\alpha_i - \gamma_i}} \left(\sum_{S \in \binom{[n]}{k}} \frac{s_{\tilde{\lambda}}[z_S] P_\alpha^\gamma(z_S^c, z_S)}{\prod_{i \in S} (z_j - z_i)} \right).$$

Note that P_α^γ is of degree $|\alpha| - |\gamma|$ (in variables z_1, z_2, \dots, z_n). Therefore if $|\alpha| - |\gamma| < |\lambda|$, the above vanishes by Lemma 3.2. \square

Lemma 6.7. *Let α be a vector such that $\alpha_m = 0$ for some $m \leq N$ and $\sum_{i=1}^{m-1} \alpha_i < |\lambda|$. Then we have*

$$\sum_{S \in \binom{[n]}{k}} \frac{s_{\tilde{\lambda}}[z_S] \prod_{j \in S^c} z_j}{\prod_{i \in S} (z_j - z_i)} \sum_{\substack{R \in \text{Sub}(\delta_S, \ell) \\ \text{rsum}(R) = \alpha}} \text{stat}_{(\delta_S, f_S^z)}(R) x^R = 0. \quad (6.7)$$

Proof. Recall that $(\text{Rec}^{\leq m}, g_S^z)$ is a sub filled diagram of (δ_S, f_S^z) given by restricting to the first m rows. We denote (D_S, f_S') to be a remaining sub filled diagram of (δ_S, f_S^z) (obtained by restricting to the $(m+1)$ -th row and above). Note that $R \in \text{Sub}(\delta_S, \ell)$ naturally induces $R_1 \in \text{Sub}(\text{Rec}^{\leq m}, \ell')$ and $R_2 \in \text{Sub}(D_S, \ell'')$ by restriction. Since $\alpha_m = 0$, there exists a constant C (independent of S) such that

$$\text{stat}_{(\delta_S, f_S^z)}(R) = C \text{stat}_{(\text{Rec}^{\leq m}, g_S^z)}(R_1) \text{stat}_{(D_S, f_S')} (R_2).$$

Therefore we have

$$\sum_{\substack{R \in \text{Sub}(\delta_S, \ell) \\ \text{rsum}(R) = \alpha}} \text{stat}_{(\delta_S, f_S^z)}(R) x^R = C \left(\sum_{\substack{R_1 \in \text{Sub}(\text{Rec}^{\leq m}, \ell') \\ \text{rsum}(R_1) = \alpha^{(1)}}} \text{stat}_{(\text{Rec}^{\leq m}, g_S^z)}(R_1) x^{R_1} \right) \left(\sum_{\substack{R_2 \in \text{Sub}(D_S, \ell'') \\ \text{rsum}(R_2) = \alpha^{(2)}}} \text{stat}_{(D_S, f_S')} (R_2) x^{R_2} \right)$$

where $\alpha^{(1)} = (\alpha_1, \alpha_2, \dots, \alpha_m)$ and $\alpha^{(2)} = (\alpha_{m+1}, \dots)$. Note that $\left(\sum_{\substack{R_2 \in \text{Sub}(D_S, \ell'') \\ \text{rsum}(R_2) = \alpha^{(2)}}} \text{stat}_{(D_S, f_S')} (R_2) x^{R_2} \right)$ is independent of the choice of S by Lemma 6.1 and Proposition 5.3. Now Lemma 6.6 (2) completes the proof as $|\alpha^{(1)}| < |\lambda|$. \square

By the previous lemma, in (6.3), it is enough to consider $R \in \text{Sub}(\delta_S, |\lambda|)$ such that $\text{rsum}(R) = \alpha$ satisfies $\alpha_i = 0$ for $i \geq N$. Otherwise we have $\sum_{i=1}^{N-1} \alpha_i < |\lambda|$ and there exists $\alpha_m = 0$ for $m \leq N$ as $N > |\lambda|$. We can naturally regard such R as an element of $\text{Sub}(\text{Rec}^{\leq N}, |\lambda|)$ by restriction. The transition ratio of stat is given by

$$\frac{C_1}{C_2} = \frac{\text{stat}_{(\delta_S, f_S^z)}(w_0')}{\text{stat}_{(\text{Rec}^{\leq N}, g_S^z)}(w_0)} = \frac{\text{stat}_{(\delta_S, f_S^z)}(R)}{\text{stat}_{(\text{Rec}^{\leq N}, g_S^z)}(R)}.$$

Therefore (6.3) becomes

$$(-1)^{|\tilde{\lambda}|} \frac{C_1}{C_2} \sum_{S \in \binom{[n]}{k}} \frac{s_{\tilde{\lambda}}[z_S] \prod_{j \in S^c} z_j}{\prod_{i \in S} (z_j - z_i)} \sum_{R \in \text{Sub}(\text{Rec}^{\leq N}, |\lambda|)} \text{stat}_{(\text{Rec}^{\leq N}, g_S^z)}(R) x^R$$

and by Lemma 6.6 (2), this equals to

$$(-1)^{|\tilde{\lambda}|} \frac{C_1}{C_2} \sum_{S \in \binom{[n]}{k}} \frac{s_{\tilde{\lambda}}[z_S] \prod_{j \in S^c} z_j}{\prod_{\substack{i \in S \\ j \in S^c}} (z_j - z_i)} \sum_{\substack{R \in \text{Sub}(\text{Rec}^{\leq N}, |\lambda|) \\ \gamma(R)=0}} \text{stat}_{(\text{Rec}^{\leq N}, g_S^z)}(R) x^R. \quad (6.8)$$

6.3. Toward the Jacobi-Trudi type formula. In this section, we complete the proof of Theorem 1.2 (c), starting from (6.8). Let B_S to be

$$B_S = \frac{1}{\text{stat}_{(\text{Rec}^{\leq N}, g_S^z)}(w_0)} \sum_{\substack{R \in \text{Sub}(\text{Rec}^{\leq N}, |\lambda|) \\ \gamma(R)=0}} \text{stat}_{(\text{Rec}^{\leq N}, g_S^z)}(R) x^R.$$

Then (6.8) equals to

$$(-1)^{|\tilde{\lambda}|} C_1 \times \sum_{S \in \binom{[n]}{k}} \frac{s_{\tilde{\lambda}}[z_S] B_S}{\prod_{j \in S^c} (z_j - z_i)}. \quad (6.9)$$

Note that B_S is symmetric in variables z_S and symmetric in variable z_{S^c} by Lemma 6.4 (2) and Lemma 6.6 (1). Therefore letting $B = B_{|n-k+1, n|}$, we can write $B_S = B(z_{S^c}, z_S)$. The following lemma says that understanding a monomial coefficient of B is crucial.

Lemma 6.8. *Let λ be a partition inside $(n-k) \times k$ rectangle and $P(z_1, z_2, \dots, z_n)$ be a polynomial function of degree $|\lambda|$ that is symmetric in variables z_1, z_2, \dots, z_{n-k} and symmetric in variables $z_{n-k+1}, z_{n-k+2}, \dots, z_n$. Consider an n by n matrix V given by*

$$V_{i,j} = \begin{cases} (\lambda_{n-k+1-j}) + j - i & \text{if } j \leq n-k \\ j - i & \text{if } j > n-k. \end{cases}$$

Then we have

$$\sum_{S \in \binom{[n]}{k}} \frac{s_{\tilde{\lambda}}[z_S] P(z_{S^c}, z_S)}{\prod_{j \in S^c} (z_j - z_i)} = (-1)^{|\tilde{\lambda}|} \sum_{\sigma \in \mathfrak{S}_n} (-1)^{\text{sgn}(\sigma)} \left[\prod_{i=1}^n z_i^{V_{\sigma(i), i}} \right] P(z_1, z_2, \dots, z_n)$$

where notation $[z_1^{c_1} z_2^{c_2} \dots z_n^{c_n}] P(z_1, z_2, \dots, z_n)$ on the right-hand side represents the operation of taking the coefficient.

Proof. We denote $\langle -, - \rangle^{(1)}$ (resp $\langle -, - \rangle^{(2)}$) to be the usual Hall inner product on symmetric functions with variables z_1, z_2, \dots, z_{n-k} (resp $z_{n-k+1}, z_{n-k+2}, \dots, z_n$). First we expand P in terms of the Schur functions in variables $z_{n-k+1}, z_{n-k+2}, \dots, z_n$:

$$P = \sum_{\mu} Q_{\mu}(z_1, z_2, \dots, z_{n-k}) s_{\mu}(z_{n-k+1}, z_{n-k+2}, \dots, z_n).$$

Letting $s_{\tilde{\lambda}} s_{\mu} = \sum_{\nu} c_{\nu} s_{\nu}$, by Lemma 3.2 we have

$$\sum_{S \in \binom{[n]}{k}} \frac{s_{\tilde{\lambda}}(z_S) Q_{\mu}(z_{S^c}) s_{\mu}(z_S)}{\prod_{j \in S^c} (z_j - z_i)} = (-1)^{|\tilde{\lambda}| + |\mu|} \langle Q_{\mu}, \sum_{\nu} c_{\nu} s_{\nu} \rangle^{(1)} = (-1)^{|\tilde{\lambda}| + |\mu|} \langle Q_{\mu}, s_{\tilde{\mu}/(\tilde{\lambda})'} \rangle^{(1)}$$

where the last equality used $\sum_{\nu} c_{\nu} s_{\nu} = s_{\square}(\omega(s_{\tilde{\lambda}} s_{\mu}))^{\perp} = s_{\tilde{\mu}/(\tilde{\lambda})}'$ for a rectangle $\square = R(n, k)$. We may only consider the case when μ is inside $R(n, n-k)$. Since $Q_{\mu} = \langle P, s_{\mu} \rangle^{(2)}$, expanding s_{μ} in terms of h using Jacobi-Trudi formula gives

$$Q_{\mu} = \sum_{\tau \in \mathfrak{S}_k} \left[\prod_{i=1}^k z_{n-k+i}^{V_{\tau(i), i}} \right] P$$

where $V' = V_{A, [n-k+1, n]}$ is a submatrix for some $A \in \binom{[n]}{k}$. Again using a Jaboci-Trudi formula for $s_{\bar{\mu}/(\bar{\lambda})'}$ gives

$$\langle Q_{\mu}, s_{\bar{\mu}/(\bar{\lambda})'} \rangle^{(1)} = \sum_{\tau \in \mathfrak{S}_{n-k}} \left[\prod_{i=1}^{n-k} z_i^{V''_{\tau(i), i}} \right] Q_{\mu}$$

where $V'' = V_{A^c, [1, n-k]}$. Now summing over all possible μ inside a rectangle $R(n, n-k)$ is the same as summing over all $A \in \binom{[n]}{k}$ and the proof is complete. \square

Now we describe a monomial coefficient of $B = B_{[n-k+1, n]}$. For $R \in \text{Sub}(\text{Rec}^{\leq N}, |\lambda|)$ denoted by $R : D \rightarrow \mathbb{Z}_{\geq 1}$, we associate a tuple $\eta(R) = (L_1, L_2, \dots, L_n)$ where L_i is a (finite) subset of $\mathbb{Z}_{\geq 0} \times \mathbb{Z}_{\geq 1}$ given by: $(c, d) \in L_i$ if and only if $(c+1, i) \in D$ and $R(c+1, i) = d$. For example, $\eta(R) = (\{(0, 3), (1, 4)\}, \{\}, \{\}, \{(1, 1), (2, 1)\})$ for R in Figure 4. Note that $|\eta(R)_i| = \text{rsum}(R)_i$

Lemma 6.9. *Given $R \in \text{Sub}(\text{Rec}^{\leq N}, |\lambda|)$ and $\gamma(R) = 0$, let $\alpha = \text{rsum}(R)$ and $\beta = \text{csum}(R)$. Then*

- (1) denoting $\eta(R) = (L_1, L_2, \dots, L_n)$, we have $L_i \in \mathbf{C}_{\beta_i}$ if $i \leq n-k$ and $L_i \in \hat{\mathbf{C}}_{\beta_i}$ if $i > n-k$
- (2) for $S = \{n-k+1, n-k+2, \dots, n\}$ we have

$$\text{stat}_{(\text{Rec}^{\leq N}, g_S^z)}(R) = \frac{\text{stat}_{(\text{Rec}^{\leq N}, g_S^z)}(w_0)}{q^{n|\lambda|} \prod_{i=1}^{N-1} z_{N+n-i}^{\alpha_i}} q^{\text{dinv}(\eta(R))} \prod_{i=1}^n z_i^{\beta_i}.$$

Proof. We denote $R : D \rightarrow \mathbb{Z}_{\geq 1}$ for some $D \subseteq \text{Rec}^{\leq N}$ with $|D| = \lambda$.

(1) Let $L_i = \{(a_1, b_1) \prec_{\text{lex}} \dots \prec_{\text{lex}} (a_{\beta_i}, b_{\beta_i})\}$. If $a_{\ell} + 1 < a_{\ell+1}$ then $(a_{\ell}, b_{\ell}) \prec_p (a_{\ell+1}, b_{\ell+1})$. If $a_{\ell} + 1 = a_{\ell+1}$ and $b_{\ell} < b_{\ell+1}$ then the cell $(a_{\ell} - 1, i)$ is bad. Therefore we have $b_{\ell} \geq b_{\ell+1}$, thus $(a_{\ell}, b_{\ell}) \prec_p (a_{\ell+1}, b_{\ell+1})$. We conclude $L_i \in \mathcal{P}_{\beta_i}$. If $i > n-k$, i -th column of $\text{Rec}^{\leq N}$ does not have a cell in the first row, we have $L_i \in \hat{\mathbf{C}}_{\beta_i}$.

(2) For every $c = (i, j) \in D$, $(i+1, j) \notin \text{Des}_{\text{Rec}^{\leq N}}(\bar{R})$ and we have $g_S^z(i+1, j) = \frac{z_{N+n-i}}{z_j}$. There are n -many cells u such that (u, c) is an attacking pair, so total $n|\lambda|$ many attacking pairs will be lost. Extra contribution among attacking pairs within R will be counted by $\text{dinv}(\eta(R))$. We conclude

$$\text{maj}_{(\text{Rec}^{\leq N}, g_S^z)}(\bar{R}) = \frac{\text{maj}_{(\text{Rec}^{\leq N}, g_S^z)}(w_0)}{\frac{\prod_{i=1}^{N-1} z_{N+n-i}^{\alpha_i}}{\prod_{i=1}^n z_i^{\beta_i}}} \quad \text{inv}_{(\text{Rec}^{\leq N})}(\bar{R}) = \frac{\text{inv}_{(\text{Rec}^{\leq N})}(w_0)}{q^{n|\lambda|}} q^{\text{dinv}(\eta(R))}.$$

\square

Recall the operators \mathfrak{h}_m , $\bar{\mathfrak{h}}_m$ and $\hat{\mathfrak{h}}_m$ acting on $\mathbb{F}[\mathbf{y}]$, polynomial ring of \mathbf{y} variables (Definition 4.3). We let Φ be an evaluation map given by: $y_{i,j} \rightarrow z_{N+n-i-1}^{-1} x_j$ if $0 \leq i \leq N-2$ and $y_{i,j} \rightarrow 0$ if $i > N-2$. Then by Lemma 6.9 we have

$$\left[\prod_{i=1}^n z_i^{\beta_i} \right] B = \frac{1}{q^{n|\lambda|}} \Phi \left(\prod_{i=1}^{n-k} \mathfrak{h}_{\beta_i} \prod_{i=n-k+1}^n \hat{\mathfrak{h}}_{\beta_i} \cdot 1 \right).$$

Note that the specialization $y_{i,j} \rightarrow 0$ if $i > N-2$ is because $\text{Rec}^{\leq N}$ has N rows and R with $\gamma(R) = 0$ is not allowed to assign a top row cell. By Lemma 6.8, (6.9) equals

$$\frac{C_1}{q^{n|\lambda|}} \Phi(\det(W) \cdot 1). \quad (6.10)$$

where $W = W^{(s)}$ (k is replaced by $n-k$) defined in (4.4).

Proof of Theorem 1.2 (c). In the expansion $\det(W) \cdot 1$, a variable $y_{i,j}$ for $i > N - 2$ does not appear as N is sufficiently large. We may only consider the specialization $y_{i,j} \rightarrow z_{N+n-i-1}^{-1} x_j$ and again specializing with (6.1), gives $y_{i,j} \rightarrow q^n t^i x_j$. Note that $\det(W) \cdot 1$ is of (homogenous) degree $|\lambda|$ in \mathbf{y} 's, comparing with (4.5) and using Lemma 6.4 (3) we have

$$(6.10) \xrightarrow{\text{Specialize by (6.1)}} \frac{T_{\delta^{|\lambda|}}(q^n)^{|\lambda|}}{q^{n|\lambda|}} \left(\det(W) \cdot 1 \Big|_{y_{i,j}=t^i x_j} \right) = T_{\delta^{|\lambda|}}(-1)^{\text{adj}(\lambda)} \text{LW}_\lambda.$$

Together with (6.4), the proof is complete. \square

7. MACDONALD PIECE POLYNOMIAL $I_{\mu,\lambda,k}$ AT $q = t = 1$

For a partition λ such that $\ell(\lambda) \leq k$, let $s_\lambda^b(x_1, x_2, \dots, x_k) := s_\lambda(x_1 - 1, x_2 - 1, \dots, x_k - 1)$. Then we have

$$s_\lambda(x_1, x_2, \dots, x_k) = \sum_{\nu \subseteq \lambda} d_{\lambda,\nu}^{(k)} s_\nu(x_1 - 1, x_2 - 1, \dots, x_k - 1) \quad (7.1)$$

where $d_{\lambda,\mu}^{(k)} = \det \left(\binom{\lambda_i + k - i}{\mu_j + k - j} \right)_{1 \leq i, j \leq k}$ [Sta]. Note that $d_{\lambda,\mu}^{(k)}$ is a number of certain non-intersecting lattice paths by Lindström–Gessel–Viennot lemma, therefore a nonnegative integer. Lastly, we denote $[h_\mu](\nabla s_\lambda|_{q,t=1})$ by $W_{\lambda,\mu}$.

Proof of Theorem 1.4. Consider

$$\bar{I}_{\mu,\lambda,k}[X; q, t] := (-1)^{|\lambda|} \sum_{S \in \binom{[n]}{k}} \frac{s_\lambda^b(z_S) \prod_{j \in S^c} z_j}{\prod_{j \in S} (z_j - z_i)} \tilde{H}_{\mu^S}[X]$$

where $z_i = q^{-\text{coarm}(c_i)} t^{-\text{coleg}(c_i)}$. We will prove the following:

$$\bar{I}_{\mu,\lambda,k}[X; 1, 1] = \sum_{\tau} W_{\tilde{\lambda}, \tau} h_{(\tau+1|\mu-k-\tilde{\lambda})}. \quad (7.2)$$

Expanding $s_\lambda^b(z_S)$ in terms of Schur functions as follows

$$s_\lambda^b(z_S) = \sum_{\nu \subseteq \lambda} d'_{\lambda,\nu} s_\nu(z_S),$$

we have

$$\bar{I}_{\mu,\lambda,k}[X; q, t] = \sum_{\nu \subseteq \lambda} (-1)^{|\lambda|-|\nu|} d'_{\lambda,\nu} I_{\mu,\nu,k}[X; q, t].$$

Note that $d'_{\lambda,\lambda} = 1$. Therefore, by Theorem 1.2 (a) and (b) we have

$$e_N^\perp(\bar{I}_{\mu,\lambda,k}[X; q, t]) = \begin{cases} 0 & \text{if } |\mu| - k - |\tilde{\lambda}| < N \\ \nabla s_{\tilde{\lambda}} & \text{if } |\mu| - k - |\tilde{\lambda}| = N. \end{cases} \quad (7.3)$$

Together with the following elementary fact

$$e_N^\perp h_\nu = \begin{cases} 0 & \text{if } \ell(\nu) < N \\ h_{\nu-(1^N)} & \text{if } \ell(\nu) = N, \end{cases}$$

we conclude

$$[h_\nu] (\bar{I}_{\mu,\lambda,k}[X; 1, 1]) = \begin{cases} 0 & \text{if } \ell(\nu) > |\mu| - k - |\tilde{\lambda}| \\ W_{\tilde{\lambda}, \nu'} & \text{if } \ell(\nu) = |\mu| - k - |\tilde{\lambda}|, \end{cases}$$

where $\nu' = \nu - (1^{|\mu| - k - |\tilde{\lambda}|})$.

Now it suffices to show $[h_\nu] (\bar{I}_{\mu,\lambda,k}[X; 1, 1]) = 0$ for $\ell(\nu) < |\mu| - k - |\tilde{\lambda}|$. Recall that for a partition $\mu \vdash n$, we have,

$$\tilde{H}_\mu[X; q, 1] = \sum_{\nu \vdash n} (q-1)^{n-\ell(\nu)} A_{\mu,\nu}(q) h_\nu[X],$$

for some $A_{\mu,\nu}(q) \in \mathbb{Z}[q]$ [GHQR19, Proposition 1.1]. From the definition of $\bar{I}_{\mu,\lambda,k}[X; q, t]$, we have,

$$[h_\tau] (\bar{I}_{\mu,\lambda,k}[X; q, 1]) = (-1)^{|\lambda|} \sum_{S \in \binom{[n]}{k}} \frac{s_\lambda^b(z_S) \prod_{j \in S^c} z_j}{\prod_{\substack{i \in S \\ j \in S^c}} (z_j - z_i)} (q-1)^{|\mu| - k - \ell(\nu)} A_{\mu^S, \tau}(q),$$

where we abuse our notation to write $z_i = q^{-\text{coleg}(c_i)}$. Note that

$$\frac{s_\lambda^b(z_S) \prod_{j \in S^c} z_j}{\prod_{\substack{i \in S \\ j \in S^c}} (z_j - z_i)}$$

has a pole of degree $k(n-k) - |\lambda| = |\tilde{\lambda}|$ at $q = 1$. Since $|\mu| - k - \ell(\tau) > |\tilde{\lambda}|$, each term

$$\frac{s_\lambda^b(z_S) \prod_{j \in S^c} z_j}{\prod_{\substack{i \in S \\ j \in S^c}} (z_j - z_i)} (q-1)^{|\mu| - \ell(\nu)} A_{\mu^S, \tau}(q)$$

vanishes at $q = 1$.

By (7.1) we have

$$I_{\mu,\lambda,k}[X; 1, 1] = \sum_{\nu \subseteq \lambda} (-1)^{|\lambda| - |\nu|} d_{\lambda,\nu}^{(k)} (\bar{I}_{\mu,\nu,k}[X; 1, 1]).$$

Now (7.2) completes the proof. \square

8. FUTURE DIRECTION

We list some future questions from this work.

- We conjectured that $I_{\mu,\lambda,k}$ is signed Schur positive and $\tilde{H}_{\mu^S}^\lambda$ is Schur positive. Therefore, finding positive monomial expansion formulas (possibly of the HHL type) for these symmetric functions would be the next challenge. As we have seen that $\tilde{H}_{\mu^S}^\lambda$ refines the Macdonald positivity, finding a positive monomial expansion would be a significant step toward addressing Problem 1.1. Additionally, we may further attempt to extend Conjecture 2.3, by expressing not only the intersection but also the intersection together with the difference of Garsia–Haiman modules.
- The arguments presented in Section 4 remain valid when we change a poset \mathbf{P} to any unit interval order. It would be interesting to find a generalization of $I_{\mu,\lambda,k}$ that matches with the combinatorics of the generalized LW_λ .

- The Macdonald polynomials have fascinating connections to the geometry of Hilbert schemes [Hai01] and affine Springer fibers [Mel20]. However, understanding our main results from a geometric perspective remains elusive. We hope that our findings provide new insights and contribute to the theory of Macdonald polynomials.

APPENDIX A. DATA FOR THE DIMENSION OF $\tilde{H}_{\mu^s}^\lambda$

Recall the symmetric function $\tilde{H}_{\mu^s}^\lambda$ defined in (2.2). Note that we can express $\tilde{H}_{\mu^s}^\lambda$ in terms of $I_{\mu,v,k}$'s, and each $I_{\mu,v,k}$ has a fixed h -expansion formula (up to a multiplication by $h_{(1^m)}$) at $q = t = 1$ by Theorem 1.4. Therefore, the 'relative dimension'

$$\text{RD}(\lambda) := \frac{\langle \tilde{H}_{\mu^s}^\lambda[X; 1, 1], e_{(1^{|\mu-k})} \rangle}{(|\mu| - k)!}$$

is a well-defined number that only depends on λ . Note that $\text{RD}((n)) = \frac{1}{n+1}$ as proved in [BG99, KLO23], which is also compatible with (1.4). We provide experimental data for $\text{RD}(\lambda)$ for $|\lambda| \leq 6$. According to Conjecture 2.3, these numbers are the relative dimension of a certain intersection of Garsia–Haiman modules.

λ	(1)	(2)	(1, 1)	(3)	(2, 1)	(1, 1, 1)	(4)	(3, 1)	(2, 2)	(2, 1, 1)	(1, 1, 1, 1)
$\text{RD}(\lambda)$	$\frac{1}{2}$	$\frac{1}{3}$	$\frac{5}{12}$	$\frac{1}{4}$	$\frac{1}{4}$	$\frac{3}{8}$	$\frac{1}{5}$	$\frac{7}{40}$	$\frac{17}{72}$	$\frac{77}{360}$	$\frac{251}{720}$
λ	(5)	(4, 1)	(3, 2)	(3, 1, 1)	(2, 2, 1)	(2, 1, 1, 1)	(1, 1, 1, 1, 1)				
$\text{RD}(\lambda)$	$\frac{1}{6}$	$\frac{2}{15}$	$\frac{19}{120}$	$\frac{7}{48}$	$\frac{47}{240}$	$\frac{139}{720}$	$\frac{95}{288}$				

λ	(6)	(5, 1)	(4, 2)	(4, 1, 1)	(3, 3)	(3, 2, 1)	(3, 1, 1, 1)
$\text{RD}(\lambda)$	$\frac{1}{7}$	$\frac{3}{28}$	$\frac{7}{60}$	$\frac{23}{210}$	$\frac{37}{240}$	$\frac{1}{8}$	$\frac{437}{3360}$
λ	(2, 2, 2)	(2, 2, 1, 1)	(2, 1, 1, 1, 1)	(1, 1, 1, 1, 1, 1)			
$\text{RD}(\lambda)$	$\frac{413}{2160}$	$\frac{749}{4320}$	$\frac{5419}{30240}$	$\frac{19087}{60480}$			

For the final remarks, we display a series of observations, which we do not have a proof, according to the data above.

- (1) (Pieri rule) For a partition λ , let λ^+ be the set of partitions obtained from λ by adding a cell. Then we have

$$\left(|\lambda^+| - \frac{1}{2}\right) \text{RD}(\lambda) = \sum_{\mu \in \lambda^+} \text{RD}(\mu).$$

- (2) (one column shapes) We have

$$\text{RD}((1^{n-1})) = \frac{N_n}{H_n},$$

where the numerator is the *Nörlund number* given in [Slo, A002657]. and the denominator is the *Hirzebruch number* given in [Slo, A091137].

- (3) (staircase shapes) Let $\text{st}_n = (n-1, n-2, \dots, 1)$ be the staircase partition. Then we have

$$\text{RD}(\text{st}_n) = \frac{1}{2^n}.$$

(4) (sum of hooks) Let Hook_n be the set of hook shape partition of size n . Then we have

$$\sum_{\lambda \in \text{Hook}_n} \text{RD}(\lambda) = 1 - \frac{1}{2^n}.$$

REFERENCES

- [AB84] M. F. Atiyah and R. Bott. The moment map and equivariant cohomology. *Topology*, 23(1):1–28, 1984.
- [BG99] Francois Bergeron and Adriano Garsia. Science fiction and Macdonald’s polynomials. *CRM Proceedings & Lecture Notes, American Mathematical Society*, 22:1–52, 1999.
- [BHM⁺21] Jonah Blasiak, Mark Haiman, Jennifer Morse, Anna Pun, and George Seelinger. Dens, nests and the loehr-warrington conjecture. *arXiv preprint arXiv:2112.07070*, 2021.
- [BHM⁺23] Jonah Blasiak, Mark Haiman, Jennifer Morse, Anna Pun, and George H. Seelinger. A proof of the extended delta conjecture. *Forum Math. Pi*, 11:Paper No. e6, 28, 2023.
- [BMPS19] Jonah Blasiak, Jennifer Morse, Anna Pun, and Daniel Summers. Catalan functions and k -Schur positivity. *J. Amer. Math. Soc.*, 32(4):921–963, 2019.
- [But94] Lynne M. Butler. Subgroup lattices and symmetric functions. *Mem. Amer. Math. Soc.*, 112(539):vi+160, 1994.
- [CM18] Erik Carlsson and Anton Mellit. A proof of the shuffle conjecture. *J. Amer. Math. Soc.*, 31(3):661–697, 2018.
- [DM22] Michele D’Adderio and Anton Mellit. A proof of the compositional delta conjecture. *Adv. Math.*, 402:Paper No. 108342, 17, 2022.
- [Gas96] Vesselin Gasharov. Incomparability graphs of $(3 + 1)$ -free posets are s -positive. In *Proceedings of the 6th Conference on Formal Power Series and Algebraic Combinatorics (New Brunswick, NJ, 1994)*, volume 157, pages 193–197, 1996.
- [Ges84] Ira M. Gessel. Multipartite P -partitions and inner products of skew Schur functions. In *Combinatorics and algebra (Boulder, Colo., 1983)*, volume 34 of *Contemp. Math.*, pages 289–317. Amer. Math. Soc., Providence, RI, 1984.
- [GH93] Adriano M. Garsia and Mark Haiman. A graded representation model for Macdonald’s polynomials. *Proc. Nat. Acad. Sci. U.S.A.*, 90(8):3607–3610, 1993.
- [GH96] A. M. Garsia and M. Haiman. Some natural bigraded S_n -modules and q, t -Kostka coefficients. volume 3, pages Research Paper 24, approx. 60. 1996. The Foata Festschrift.
- [GH07] Ian Grojnowski and Mark Haiman. Affine Hecke algebras and positivity of LLT and Macdonald polynomials. *preprint*, 2007.
- [GHQR19] Adriano M Garsia, James Haglund, Dun Qiu, and Marino Romero. e -positivity results and conjectures. *arXiv preprint arXiv:1904.07912*, 2019.
- [GHT99] A. M. Garsia, M. Haiman, and G. Tesler. Explicit plethystic formulas for Macdonald q, t -Kostka coefficients. volume 42, pages Art. B42m, 45. 1999. The Andrews Festschrift (Maratea, 1998).
- [GT96] A. M. Garsia and G. Tesler. Plethystic formulas for Macdonald q, t -Kostka coefficients. *Adv. Math.*, 123(2):144–222, 1996.
- [Hag04] J. Haglund. A proof of the q, t -Schröder conjecture. *Int. Math. Res. Not.*, (11):525–560, 2004.
- [Hai01] Mark Haiman. Hilbert schemes, polygraphs and the Macdonald positivity conjecture. *J. Amer. Math. Soc.*, 14(4):941–1006, 2001.
- [HHL05a] J. Haglund, M. Haiman, and N. Loehr. A combinatorial formula for Macdonald polynomials. *J. Amer. Math. Soc.*, 18(3):735–761, 2005.
- [HHL⁺05b] J. Haglund, M. Haiman, N. Loehr, J. B. Remmel, and A. Ulyanov. A combinatorial formula for the character of the diagonal coinvariants. *Duke Math. J.*, 126(2):195–232, 2005.
- [HHL08] J. Haglund, M. Haiman, and N. Loehr. A combinatorial formula for nonsymmetric Macdonald polynomials. *Amer. J. Math.*, 130(2):359–383, 2008.
- [KLO22] Donghyun Kim, Seung Jin Lee, and Jaeseong Oh. Toward Butler’s conjecture. *arXiv preprint arXiv:2212.09419*, 2022.
- [KLO23] Donghyun Kim, Seung Jin Lee, and Jaeseong Oh. Shuffle formula in science fiction for macdonald polynomials. *arXiv preprint arXiv:2306.14371*, 2023.
- [LLM03] L. Lapointe, A. Lascoux, and J. Morse. Tableau atoms and a new Macdonald positivity conjecture. *Duke Math. J.*, 116(1):103–146, 2003.

- [LW08] Nicholas A. Loehr and Gregory S. Warrington. Nested quantum Dyck paths and $\nabla(s_\lambda)$. *Int. Math. Res. Not. IMRN*, (5):Art. ID rnm 157, 29, 2008.
- [Mac88] Ian Macdonald. A new class of symmetric functions. *Séminaire Lotharingien de Combinatoire*, 20:B20a–41, 1988.
- [Mel20] Anton Mellit. Poincaré polynomials of character varieties, Macdonald polynomials and affine Springer fibers. *Ann. of Math. (2)*, 192(1):165–228, 2020.
- [Mel21] Anton Mellit. Toric braids and (m, n) -parking functions. *Duke Math. J.*, 170(18):4123–4169, 2021.
- [Slo] Sloane Neil. The On-Line Encyclopedia of Integer Sequences. <http://oeis.org>.
- [Sta] Richard Stanley. Plugging $1-x$ into schur polynomials. MathOverflow. URL:<https://mathoverflow.net/q/228999> (version: 2016-01-22).
- [SW16] John Shareshian and Michelle L. Wachs. Chromatic quasisymmetric functions. *Adv. Math.*, 295:497–551, 2016.

DEPARTMENT OF MATHEMATICAL SCIENCES, SEOUL NATIONAL UNIVERSITY, SEOUL 151-247, KOREA

Email address: hyun920310@snu.ac.kr

JUNE E HUH CENTER FOR MATHEMATICAL CHALLENGES, KOREA INSTITUTE FOR ADVANCED STUDY, SEOUL 02455, KOREA

Email address: jsOh@kias.re.kr