

THE QUASI-POLYNOMIALITY OF $\text{mod } q$ PERMUTATION REPRESENTATIONS FOR A LINEAR FINITE GROUP ACTION ON A LATTICE

RYO UCHIUMI AND MASAHIKO YOSHINAGA

ABSTRACT. For given linear action of a finite group on a lattice and a positive integer q , we prove that the $\text{mod } q$ permutation representation is a quasi-polynomial in q . Additionally, we establish several results that can be considered as $\text{mod } q$ -analogues of results by Stapledon for equivariant Ehrhart quasi-polynomials. We also prove a reciprocity-type result for multiplicities of irreducible decompositions.

CONTENTS

1. Introduction	1
1.1. Quasi-polynomials	1
1.2. Equivariant Ehrhart theory	2
1.3. Towards an equivariant version of characteristic quasi-polynomials	3
2. Quasi-polynomiality	3
2.1. Group action and representation	3
2.2. Multiplicities of irreducible decompositions	4
2.3. Permutation representations	6
2.4. Number of orbits	7
2.5. Reciprocity for the multiplicities	7
2.6. Examples	8
References	12

1. INTRODUCTION

1.1. **Quasi-polynomials.** Let R be a commutative ring. A function $f : \mathbb{Z}_{(>0)} \rightarrow R$ is called a *quasi-polynomial* if there exist a positive integer $\tilde{n} \in \mathbb{Z}_{>0}$ and polynomials $g_1(t), \dots, g_{\tilde{n}}(t) \in R[t]$ such that

$$f(q) = g_r(q), \quad \text{if } q \equiv r \pmod{\tilde{n}} \quad (1 \leq r \leq \tilde{n}).$$

The positive integer \tilde{n} is called a *period* and each polynomial g_r is called the *constituent* of f . The quasi-polynomial f has degree d if all the constituents have degree d . Moreover, the quasi-polynomial f has the *gcd-property* if the polynomial g_r depends on r only through $\gcd\{\tilde{n}, r\}$. In other words, $g_{r_1} = g_{r_2}$ if $\gcd\{\tilde{n}, r_1\} = \gcd\{\tilde{n}, r_2\}$.

Quasi-polynomials play important roles in many areas of mathematics. They appear frequently as counting functions (in this case, $R = \mathbb{Z}$). In particular, the following two notions have been actively studied.

Date: August 30, 2024.

2010 Mathematics Subject Classification. 05E18, 20C10.

Key words and phrases. quasi-polynomials; finite groups; lattices.

R. U. was supported by JST SPRING, Grant Number JPMJSP2138.

M. Y. was partially supported by JSPS KAKENHI, Grant Number JP23H00081.

Example 1.1 (The Ehrhart quasi-polynomial). Let \mathcal{P} be a rational polytope in \mathbb{R}^n . For $q \in \mathbb{Z}_{>0}$, define

$$L_{\mathcal{P}}(q) := \#(q\mathcal{P} \cap \mathbb{Z}^n).$$

Then $L_{\mathcal{P}}(q)$ is a quasi-polynomial ([1, Theorem 3.23]), known as the Ehrhart quasi-polynomial.

Example 1.2 (The characteristic quasi-polynomial). Let $L \simeq \mathbb{Z}^\ell$ be a lattice and $L^\vee := \text{Hom}_{\mathbb{Z}}(L, \mathbb{Z})$ be the dual lattice. Given $\alpha_1, \dots, \alpha_n \in L^\vee$, we can associate a hyperplane arrangement $\mathcal{A} := \{H_1, \dots, H_n\}$ in $\mathbb{R}^\ell \simeq L \otimes \mathbb{R}$, where

$$H_i := \{x \in L \otimes \mathbb{R} \mid \alpha_i(x) = 0\}.$$

For a positive integer $q \in \mathbb{Z}_{>0}$, define the mod q complement of the arrangement by

$$\begin{aligned} M(\mathcal{A}; q) &:= (L/qL)^\ell \setminus \bigcup_{i=1}^n \bar{H}_i \\ &= \{\bar{x} \in L/qL \mid \alpha_i(x) \not\equiv 0 \pmod{q} \text{ for all } i \in \{1, \dots, n\}\}. \end{aligned}$$

It is known ([4, Theorem 2.4] and [5, Theorem 3.1] for non-central case) that

$$\chi_{\text{quasi}}(\mathcal{A}; q) := \#M(\mathcal{A}; q)$$

is a quasi-polynomial.

Roughly speaking, the notion of characteristic quasi-polynomial is a mod q -version of the Ehrhart quasi polynomial. However, the characteristic quasi-polynomials possess some additional properties. First, the constituents $g_r(t)$ ($r \in \{1, \dots, \tilde{n}\}$) of the characteristic quasi-polynomial $\chi_{\text{quasi}}(\mathcal{A}; q)$ satisfy the gcd-property. Secondly, the first constituent $g_1(t)$ (and equivalently, $g_r(t)$ for r coprime to \tilde{n}) is known to be equal to the characteristic polynomial $\chi(\mathcal{A}, t)$ of the arrangement \mathcal{A} (see [7]). Furthermore, $g_{\tilde{n}}(t)$ is the characteristic polynomial of the associated toric arrangement [6, 11]. The characteristic quasi-polynomial is an important concept, not only in the context of enumerative problems but also in its connections with arithmetic matroids and toric arrangements [2, 6].

1.2. Equivariant Ehrhart theory. In [10], Stapledon proposed an equivariant version of Ehrhart theory. Let $L \simeq \mathbb{Z}^n$ be a lattice and let Γ be a finite group acting linearly on L via $\rho : \Gamma \rightarrow \text{GL}(L)$. Let \mathcal{P} be a Γ -invariant lattice polytope. For a positive integer $q \in \mathbb{Z}_{>0}$, the group Γ acts on the lattice points $q\mathcal{P} \cap L$. Let $\chi_{q\mathcal{P}}$ denote the character of this permutation representation. Stapledon proved representation-theoretic analogues of several classical results in Ehrhart theory. For example, the map

$$F : \mathbb{Z}_{>0} \longrightarrow R(\Gamma), \quad q \longmapsto F(q) = \chi_{q\mathcal{P}}$$

is a quasi-polynomial of degree $\dim \mathcal{P}$ with the leading coefficient $\frac{\text{vol} \mathcal{P}}{\#\Gamma} \chi_{\mathbb{R}}$, where $\chi_{\mathbb{R}}$ is the regular (standard) character of Γ ([10, Theorem 5.7, Corollary 5.9]). It was also proved that the number of Γ -orbits in $q\mathcal{P} \cap L$ is a quasi-polynomial in q .

Stapledon also proved the following reciprocity. Let $F^*(q) = \chi_{q\mathcal{P}^*}$ be the permutation representation of the lattice points in the interior of $q\mathcal{P}$. Then, from the Ehrhart reciprocity, the relation

$$(1.1) \quad F^*(q) = (-1)^{\dim \mathcal{P}} \det(\rho) F(-q)$$

holds. Note that when Γ is the trivial group, these results recover the classical results in Ehrhart theory.

1.3. Towards an equivariant version of characteristic quasi-polynomials. It is natural to consider $\text{mod } q$ -version of the equivariant Ehrhart quasi-polynomial, namely, the equivariant characteristic quasi-polynomials for an arrangement invariant under a group action. For simplicity, in this paper, we do not consider hyperplanes, and instead focus solely on the $\text{mod } q$ permutation representation. Let Γ be a finite group and let $L \simeq \mathbb{Z}^\ell$ be a lattice. Suppose that Γ acts linearly on L . Then the action of Γ on $L_q := L/qL$ is naturally induced for each $q \in \mathbb{Z}_{>0}$. One of our problems is how the permutation character χ_{L_q} of L_q depends on q . In the case where Γ is the Weyl group and L is a lattice associated with a root system, there are several known results (e.g. [3, 8]), especially for $q \equiv 1 \pmod{h}$, where h is the Coxeter number. In this paper, we consider a general finite group action on a lattice. The main result of this paper is the following.

Theorem 1.3 (Theorem 2.5 below). *Consider the function $F : \mathbb{Z}_{>0} \rightarrow R(\Gamma)$ defined by $q \mapsto \chi_{L_q}$. Then F is a quasi-polynomial of degree ℓ ($= \text{rank } L$) with the gcd-property.*

We will also prove that the number of Γ -orbits is a quasi-polynomial with the gcd-property (Corollary 2.8).

Let χ_1, \dots, χ_k be irreducible characters of Γ . Then χ_{L_q} can be expressed as

$$(1.2) \quad \chi_{L_q} = m(\chi_1; q) \cdot \chi_1 + \dots + m(\chi_k; q) \cdot \chi_k,$$

with $m(\chi_k; q) \in \mathbb{Z}$. Theorem 1.3 is equivalent to that each $m(\chi_i; q)$ is a \mathbb{Z} -valued quasi-polynomial in q (Corollary 2.2).

There are several relations among these quasi-polynomials. In particular, there is a reciprocity-type relation between $m(\chi_i; q)$ and $m(\chi_i \otimes \delta_\rho; q)$, where $\delta_\rho = \det \rho$. More precisely, we have (Theorem 2.10)

$$m(\chi_i \otimes \delta_\rho; q) = (-1)^\ell m(\chi_i; -q).$$

This implies the following relation (Corollary 2.11):

$$(1.3) \quad F(-q) = (-1)^\ell \delta_\rho F(q).$$

Although the formula (1.3) appears similar to (1.1), they are different in nature. It is important to note that (1.1) represents a reciprocity between $F(q)$ and $F^*(q)$, whereas (1.3) is a self-duality of $F(q)$.

We will also provide several explicit examples.

2. QUASI-POLYNOMIALITY

2.1. Group action and representation. We recall several notions and basic facts about representations of finite groups [9].

Let Γ be a finite group. Let V be a finite-dimensional vector space over \mathbb{C} , and let $\text{GL}(V)$ denote the group of linear isomorphisms of V onto itself. A (linear) representation of Γ on V is a homomorphism $\rho : \Gamma \rightarrow \text{GL}(V)$. In this paper, we assume that ρ is injective. The space V is called the representation space of ρ .

The character $\chi_\rho : \Gamma \rightarrow \mathbb{C}$ of the representation ρ is the function defined by $\gamma \mapsto \text{tr } \rho(\gamma)$, where tr denotes the trace function. The character χ_ρ is constant on each conjugacy class. A function $\phi : \Gamma \rightarrow \mathbb{C}$ is called a class function if ϕ is constant on each conjugacy class. For functions $\phi, \psi : \Gamma \rightarrow \mathbb{C}$, define the inner product (ϕ, ψ) by

$$(\phi, \psi) = \frac{1}{\#\Gamma} \sum_{\gamma \in \Gamma} \phi(\gamma) \overline{\psi(\gamma)},$$

where \bar{z} denotes the complex conjugate of $z \in \mathbb{C}$. Let χ_1, \dots, χ_k be the set of all irreducible characters of Γ . Then χ_1, \dots, χ_k form an orthonormal basis of the space of class functions. In particular, $(\chi_i, \chi_j) = \delta_{ij}$. Thus, if a class function χ is expressed as a linear combination of irreducible characters $\chi = m_1 \chi_1 + \dots + m_k \chi_k$, then we have $m_i = (\chi, \chi_i)$.

Let Γ' be a subgroup of Γ . The restriction of a class function $\chi : \Gamma \rightarrow \mathbb{C}$ to Γ' is clearly a class function on Γ' , which is denoted by $\text{Res}_{\Gamma'}^{\Gamma} \chi : \Gamma' \rightarrow \mathbb{C}$. Conversely, for a class function $\varphi : \Gamma' \rightarrow \mathbb{C}$, define the *induced function* $\text{Ind}_{\Gamma'}^{\Gamma} \varphi : \Gamma \rightarrow \mathbb{C}$ by

$$(2.1) \quad (\text{Ind}_{\Gamma'}^{\Gamma} \varphi)(\gamma) = \frac{1}{\#\Gamma'} \sum_{\substack{\eta \in \Gamma \\ \eta^{-1}\gamma\eta \in \Gamma'}} \varphi(\eta^{-1}\gamma\eta).$$

These two operators are related by the following Frobenius reciprocity:

$$(2.2) \quad (\chi, \text{Ind}_{\Gamma'}^{\Gamma} \varphi) = (\text{Res}_{\Gamma'}^{\Gamma} \chi, \varphi).$$

Recall that the *representation ring* $R(\Gamma)$ of Γ is $\bigoplus_V \mathbb{Z}[V]/\sim$, where V runs all finite-dimensional representations of Γ , and \sim is an equivalence relation generated by $[V] \sim [V']$ for isomorphic representations $V \simeq V'$ and $[V_1 \oplus V_2] \sim [V_1] + [V_2]$. The multiplication is defined by $[V_1] \cdot [V_2] = [V_1 \otimes V_2]$. The character gives a natural isomorphism of abelian groups

$$R(\Gamma) \simeq \mathbb{Z}\chi_1 \oplus \cdots \oplus \mathbb{Z}\chi_k.$$

The trivial representation $\rho_{\mathbf{1}}$ is the unit element in $R(\Gamma)$. The character of $\rho_{\mathbf{1}}$ is denoted by $\mathbf{1}$.

Suppose that Γ acts on a finite set X . Let $\mathbb{C}X$ denote the vector space based on X , that is, $\mathbb{C}X = \bigoplus_{x \in X} \mathbb{C}x$. This gives rise to a natural representation $\rho_X : \Gamma \rightarrow \text{GL}(\mathbb{C}X)$, which is called the *permutation representation* of X . In the case of $X = \Gamma$ with action defined by the left multiplication, it is called the *regular (standard) representation*, denoted by $\rho_{\mathbf{R}}$. Note that the character $\chi_{\mathbf{R}}$ of the regular representation satisfies the following.

$$\chi_{\mathbf{R}} = \sum_{i=1}^k \chi_i(1)\chi_i, \quad \chi_{\mathbf{R}}(\gamma) = \begin{cases} \#\Gamma & \text{if } \gamma = 1; \\ 0 & \text{otherwise.} \end{cases}$$

For $x \in X$, the Γ -orbit $\Gamma(x)$ and the *isotropy subgroup* Γ_x are defined as follows:

$$\begin{aligned} \Gamma(x) &= \{ \gamma x \in X \mid \gamma \in \Gamma \}, \\ \Gamma_x &= \{ \gamma \in \Gamma \mid \gamma x = x \}. \end{aligned}$$

2.2. Multiplicities of irreducible decompositions. Let L be a lattice, and $\{\beta_1, \dots, \beta_{\ell}\}$ be a \mathbb{Z} -basis of L , that is, $L = \mathbb{Z}\beta_1 \oplus \cdots \oplus \mathbb{Z}\beta_{\ell} \simeq \mathbb{Z}^{\ell}$. We identify an element $x = x_1\beta_1 + \cdots + x_{\ell}\beta_{\ell}$ of L with the row vector $x = (x_1, \dots, x_{\ell})$ of \mathbb{Z}^{ℓ} .

Let Γ be a finite group. Let $\rho : \Gamma \rightarrow \text{GL}(L)$ be a group homomorphism. Let us denote the representation matrix of $\rho(\gamma)$ by R_{γ} , and we consider the right multiplication, namely,

$$\rho(\gamma) : L \rightarrow L, \quad x \mapsto xR_{\gamma}.$$

For $q \in \mathbb{Z}_{>0}$, define $\mathbb{Z}_q := \mathbb{Z}/q\mathbb{Z}$. We will consider the following q -reduction of $x = (x_1, \dots, x_{\ell}) \in \mathbb{Z}^{\ell}$:

$$[x]_q := ([x_1]_q, \dots, [x_{\ell}]_q) \in \mathbb{Z}_q^{\ell},$$

where $[x_i]_q = x_i + q\mathbb{Z} \in \mathbb{Z}_q$. We similarly consider the q -reduction of an integer matrix $A = (a_{ij})_{ij}$:

$$[A]_q := ([a_{ij}]_q)_{ij}.$$

Let $\varphi : \mathbb{Z}^{\ell} \rightarrow \mathbb{Z}^{\ell}$ be a \mathbb{Z} -homomorphism represented by a $\ell \times \ell$ integer matrix A . We can define the induced morphism $\varphi_q : \mathbb{Z}_q^{\ell} \rightarrow \mathbb{Z}_q^{\ell}$ by

$$x \mapsto x[A]_q.$$

Let $L_q := L/qL \simeq (\mathbb{Z}/q\mathbb{Z})^{\ell}$. The action of Γ on L_q is induced by $\rho(\gamma)_q : L_q \rightarrow L_q$. Let χ_{L_q} denote the character of the permutation representation of L_q , and consider its irreducible decomposition:

$$\chi_{L_q} = m(\chi_1; q) \cdot \chi_1 + \cdots + m(\chi_k; q) \cdot \chi_k,$$

where $m(\chi_i; q)$ denotes the multiplicity of χ_i in χ_{L_q} . Since $\chi_{L_q}(\gamma)$ is equal to the number of elements in L_q fixed by $\gamma \in \Gamma$, we have

$$(2.3) \quad m(\chi_i; q) = (\chi_i, \chi_{L_q}) = \frac{1}{\#\Gamma} \sum_{\gamma \in \Gamma} \chi_i(\gamma) \overline{\chi_{L_q}(\gamma)} = \frac{1}{\#\Gamma} \sum_{\gamma \in \Gamma} \chi_i(\gamma) \cdot \#L_q^\gamma,$$

where $L_q^\gamma := \{x \in L_q \mid \gamma x = x\}$. Thus, by studying the properties of $\#L_q^\gamma$, we can determine how $m(\chi_i; q)$ depends on q . Note that for the trivial character $\mathbf{1}$, the multiplicity $m(\mathbf{1}; q)$ represents the number of Γ -orbits of L_q , according to Burnside's lemma.

The fixed point set L_q^γ is expressed as

$$\begin{aligned} L_q^\gamma &= \{x \in L_q \mid \gamma x = x\} \\ &= \{x \in L_q \mid x[R_\gamma]_q = x\} \\ &= \{x \in L_q \mid x[R_\gamma - I_\ell]_q = 0\}, \end{aligned}$$

where I_ℓ is the identity matrix of size ℓ . Therefore, L_q^γ is equal to the kernel of the induced morphism $(\rho(\gamma) - \text{id})_q$. The cardinality of the kernel of this induced morphism is known to be quasi-monomial, as shown in [4]:

Lemma 2.1 ([4, Lemma 2.1]). *Let $\varphi : \mathbb{Z}^\ell \rightarrow \mathbb{Z}^\ell$ be a \mathbb{Z} -homomorphism. Then The cardinality of the kernel of this induced morphism $\varphi_q : \mathbb{Z}_q^\ell \rightarrow \mathbb{Z}_q^\ell$ is a quasi-monomial in q . Furthermore, suppose φ is represented by a matrix A . Then the quasi-monomial $\#\ker \varphi_q$ can be expressed as*

$$(2.4) \quad \#\ker \varphi_q = \left(\prod_{j=1}^r \gcd\{e_j, q\} \right) q^{\ell-r},$$

where $r := \text{rank } A$ and $e_1, \dots, e_r \in \mathbb{Z}_{>0}$, with $e_1 \mid e_2 \mid \dots \mid e_r$, are the elementary divisors of A . Hence, the quasi-monomial $\#\ker \varphi_q$ has the gcd-property and the minimum period e_r . If $r = 0$, we consider e_0 to be 1.

Proof. Here, we only review quasi-monomiality. For further details, see [4, Lemma 2.1].

Since $\#\ker \varphi_q = q^\ell / \#\text{im } \varphi_q$, we will study $\#\text{im } \varphi_q$. Consider the Smith normal form

$$SAT = \begin{pmatrix} e_1 & & & \\ & \ddots & & \\ & & e_r & \\ & & & 0 \end{pmatrix}, \quad r = \text{rank } A, \quad e_1, \dots, e_r, \in \mathbb{Z}_{>0}, \quad e_1 \mid e_2 \mid \dots \mid e_r,$$

where S and T are $\ell \times \ell$ unimodular matrices. Since unimodularity is preserved under q -reductions, we may assume that A is a diagonal matrix $\text{diag}(e_1, \dots, e_r, 0, \dots, 0)$ from the outset. Then, for $x = (x_1, \dots, x_\ell) \in \mathbb{Z}_q^\ell$, we have

$$\varphi_q(x) = ([e_1]_q x_1, \dots, [e_r]_q x_r, 0, \dots, 0)$$

and hence $\text{im } \varphi_q = [e_1]_q \mathbb{Z}_q \times \dots \times [e_r]_q \mathbb{Z}_q$. Therefore,

$$\#\text{im } \varphi_q = \frac{q^r}{\prod_{j=1}^r \gcd\{e_j, q\}},$$

and we obtain (2.4). \square

Corollary 2.2. *The multiplicity $m(\chi_i; q)$ of χ_i in χ_{L_q} is a quasi-polynomial in q . More explicitly,*

$$(2.5) \quad m(\chi_i; q) = \frac{1}{\#\Gamma} \sum_{\gamma \in \Gamma} \chi_i(\gamma) \cdot \left(\prod_{j=1}^{r(\gamma)} \gcd\{e_{\gamma,j}, q\} \right) q^{\ell-r(\gamma)},$$

where $r(\gamma) := \text{rank}(R_\gamma - I_\ell)$ and $e_{\gamma,1}, \dots, e_{\gamma,r(\gamma)} \in \mathbb{Z}_{>0}$ with $e_{\gamma,1} \mid e_{\gamma,2} \mid \dots \mid e_{\gamma,r(\gamma)}$, are the elementary divisors of $R_\gamma - I_\ell$.

Proof. The equation (2.5) is given by (2.3) and (2.4). \square

Next, we present some properties of $m(\chi_i; q)$.

Proposition 2.3. *The quasi-polynomial $m(\chi_i; q)$ has the gcd-property with a period*

$$\tilde{n} := \text{lcm} \{ e_{\gamma, r(\gamma)} \mid \gamma \in \Gamma \}.$$

Furthermore, the minimum period of the quasi-polynomial $m(\mathbf{1}; q)$ is equal to \tilde{n} .

Note that if $\chi_i \neq \mathbf{1}$, we do not know if \tilde{n} is the minimum period.

Proof. Let $\gamma \in \Gamma$ be an element that is not the identity, and let $e_{\gamma, 1}, \dots, e_{\gamma, r(\gamma)}$ be the elementary divisors of $R_\gamma - I_\ell$. Since $e_{\gamma, j}$ divides \tilde{n} for $j \in \{1, \dots, r(\gamma)\}$, we have

$$\prod_{j=1}^{r(\gamma)} \gcd\{e_{\gamma, j}, q\} = \prod_{j=1}^{r(\gamma)} \gcd\{e_{\gamma, j}, \tilde{n}, q\} = \prod_{j=1}^{r(\gamma)} \gcd\{e_{\gamma, j}, \gcd\{\tilde{n}, q\}\}.$$

Hence $m(\chi_i; q)$ depends on q only through $\gcd\{\tilde{n}, q\}$, which means that \tilde{n} is a period of $m(\chi_i; q)$.

Let $g_1(t), \dots, g_{\tilde{n}}(t) \in \mathbb{Z}[t]$ denote the constituents of the quasi-polynomial $m(\mathbf{1}; q)$. Since \tilde{n} is divisible by $e_{\gamma, r(\gamma)}$ for all $\gamma \in \Gamma$, we have

$$(2.6) \quad g_{\tilde{n}}(t) = \frac{1}{\#\Gamma} \sum_{\gamma \in \Gamma} \left(\prod_{j=1}^{r(\gamma)} e_{\gamma, j} \right) t^{\ell - r(\gamma)}$$

from equation (2.5). Suppose that $r < \tilde{n}$. Then there exists $\gamma \in \Gamma$ such that $\gcd\{e_{\gamma, r(\gamma)}, r\} \neq e_{\gamma, r(\gamma)}$. Since $\gcd\{e_{\gamma, j}, r\} \leq e_{\gamma, j}$ for any $\gamma \in \Gamma$ and $j \in \{1, \dots, r(\gamma)\}$, we conclude that

$$g_r(t) = \frac{1}{\#\Gamma} \sum_{\gamma \in \Gamma} \left(\prod_{j=1}^{r(\gamma)} \gcd\{e_{\gamma, j}, q\} \right) t^{\ell - r(\gamma)} \neq g_{\tilde{n}}(t)$$

by the equations (2.5) and (2.6), and hence r is not a period. Therefore, \tilde{n} is the minimum period of $m(\mathbf{1}; q)$. \square

Proposition 2.4. *The leading term of the quasi-polynomial $m(\chi_i; q)$ is $\frac{\chi_i(1)}{\#\Gamma} q^\ell$.*

Proof. Since ρ is injective, $r(\gamma) = 0$ holds if and only if γ is the identity. Therefore, by Corollary 2.2, the leading term of $m(\chi_i; q)$ is q^ℓ with the coefficient $\frac{\chi_i(1)}{\#\Gamma}$. \square

2.3. Permutation representations. Since each multiplicity $m(\chi_i; q)$ is a quasi-polynomial, the quasi-polynomiality of the function $F : q \mapsto \chi_{L_q}$ follows immediately. The following theorem is the main result of this paper.

Theorem 2.5 (Restatement of Theorem 1.3). *Consider the function $F : \mathbb{Z}_{>0} \rightarrow R(\Gamma)$ defined by $q \mapsto \chi_{L_q}$. Then F is a quasi-polynomial of degree ℓ . Furthermore, F has the gcd-property, the minimum period \tilde{n} , and the leading coefficient of the quasi-polynomial χ_{L_q} is $\frac{\chi_R}{\#\Gamma}$.*

Proof. By equation (2.5), we have

$$F(q) = \chi_{L_q} = \sum_{i=1}^k m(\chi_i; q) \cdot \chi_i = \frac{1}{\#\Gamma} \sum_{i=1}^k \sum_{\gamma \in \Gamma} \chi_i(\gamma) \cdot \left(\prod_{j=1}^{r(\gamma)} \gcd\{e_{\gamma, j}, q\} \right) \cdot \chi_i \cdot q^{\ell - r(\gamma)} \in R(\Gamma)[q],$$

hence F is a quasi-polynomial with the gcd-property. Since $m(\mathbf{1}; q)$ has the minimum period \tilde{n} , F also has the minimum period \tilde{n} .

By Proposition 2.4, the leading term of each multiplicity $m(\chi_i; q)$ is $\frac{\chi_i(1)}{\#\Gamma} q^\ell$. Thus, we have

$$\sum_{i=1}^k \frac{\chi_i(1)}{\#\Gamma} \cdot \chi_i \cdot q^\ell = \frac{\chi_R}{\#\Gamma} q^\ell$$

as the leading term of F . \square

2.4. Number of orbits. In this section, we prove the quasi-polynomiality of the number of Γ -orbits. First, we describe the permutation character $\chi_{\Gamma(x)}$ on the Γ -orbit $\Gamma(x)$ of $x \in L_q$.

Lemma 2.6. *Let $\Gamma(x)$ denote the Γ -orbit of $x \in L_q$. Then we have*

$$\chi_{\Gamma(x)}(\gamma) = \#\Gamma(x)^\gamma = \left(\text{Ind}_{\Gamma_x}^{\Gamma} \mathbf{1}\right)(\gamma).$$

Proof. An element ηx of $\Gamma(x)$ is fixed by γ if and only if $\eta^{-1}\gamma\eta$ fixes x . Thus, the cardinality of $\Gamma(x)^\gamma$ is

$$\#\Gamma(x)^\gamma = \frac{\#\{\eta \in \Gamma \mid \eta^{-1}\gamma\eta \in \Gamma_x\}}{\#\Gamma_x}.$$

On the other hand, it follows directly that the above expression is equal to $\left(\text{Ind}_{\Gamma_x}^{\Gamma} \mathbf{1}\right)(\gamma)$:

$$\left(\text{Ind}_{\Gamma_x}^{\Gamma} \mathbf{1}\right)(\gamma) = \frac{1}{\#\Gamma_x} \sum_{\substack{\eta \in \Gamma \\ \eta^{-1}\gamma\eta \in \Gamma_x}} \mathbf{1}(\eta^{-1}\gamma\eta) = \frac{\#\{\eta \in \Gamma \mid \eta^{-1}\gamma\eta \in \Gamma_x\}}{\#\Gamma_x}.$$

□

For a 1-dimensional character λ of Γ and $q \in \mathbb{Z}_{>0}$, let $f_{L/\Gamma}(\lambda; q)$ denote the number of Γ -orbit of L_q whose isotropy subgroup is contained in the subgroup $\lambda^{-1}(1)$ of Γ . Using the Frobenius reciprocity (2.2), we obtain the following lemma.

Lemma 2.7. *For a 1-dimensional character λ of Γ and $q \in \mathbb{Z}_{>0}$, we have*

$$f_{L/\Gamma}(\lambda; q) = (\lambda, \chi_{L_q}) = m(\lambda; q).$$

Proof. Note that the second equality is the definition of $m(\lambda; q)$.

Note that the permutation character χ_{L_q} can be decomposed into a sum of all permutation characters of Γ -orbit of L_q :

$$\chi_{L_q} = \sum_{\Gamma(x) : \Gamma\text{-orbit}} \chi_{\Gamma(x)}.$$

By Lemma 2.6 and Frobenius reciprocity (2.2), we have

$$(\lambda, \chi_{L_q}) = \sum_{\Gamma(x) : \Gamma\text{-orbit}} (\lambda, \chi_{\Gamma(x)}) = \sum_{\Gamma(x) : \Gamma\text{-orbit}} (\lambda, \text{Ind}_{\Gamma_x}^{\Gamma} \mathbf{1}) = \sum_{\Gamma(x) : \Gamma\text{-orbit}} \left(\text{Res}_{\Gamma_x}^{\Gamma} \lambda, \mathbf{1}\right).$$

Since $\text{Res}_{\Gamma_x}^{\Gamma} \lambda$ is an irreducible character of Γ_x , the orthogonality of irreducible characters implies that

$$\left(\text{Res}_{\Gamma_x}^{\Gamma} \lambda, \mathbf{1}\right) = \begin{cases} 1 & \Gamma_x \subseteq \lambda^{-1}(1); \\ 0 & \text{otherwise.} \end{cases}$$

Therefore, we have $(\lambda, \chi_{L_q}) = f_{L/\Gamma}(\lambda; q)$. □

Corollary 2.8. *The function $f_{L/\Gamma}(\lambda; -) : \mathbb{Z}_{>0} \rightarrow \mathbb{Z}$ is a quasi-polynomial of degree ℓ and it has the gcd-property.*

Proof. This follows from Corollary 2.2 and Lemma 2.7. □

2.5. Reciprocity for the multiplicities. Let $\rho : \Gamma \rightarrow \text{GL}(L)$ be a representation and R_γ the representation matrix of $\rho(\gamma)$. Define the *reciprocity character* $\delta_\rho : \Gamma \rightarrow \mathbb{C}$ by

$$\delta_\rho(\gamma) := (-1)^{r(\gamma)},$$

where $r(\gamma) = \text{rank}(R_\gamma - I_\ell)$. The following lemma shows that $\delta_\rho(\gamma) = \det R_\gamma$ and that δ_ρ is an irreducible character of Γ .

Lemma 2.9 ([10, Lemma 5.5]). *Let $R \in \text{GL}_n(\mathbb{R})$ be a real matrix of finite order. Let $r := \text{rank}(R - I_n)$. Then $\det R = (-1)^r$.*

Proof. Since R is finite order, it is diagonalizable (in \mathbb{C}), and we can write $R = PDP^{-1}$, where $P, D \in \text{GL}_n(\mathbb{C})$ with D diagonal. Clearly, $\text{rank}(R - I_n) = \text{rank}(D - I_n)$. Thus, r is the number of diagonal entries of D that are not equal to 1. Since R is a real matrix, the set of eigenvalues is closed under complex conjugation. The finiteness of the order implies that all the eigenvalues have absolute value 1. Therefore, the diagonal entries of D are as follows (with multiplicities):

$$1^{p_1}, (-1)^{p_2}, \alpha_1^{q_1}, \bar{\alpha}_1^{q_1}, \alpha_2^{q_2}, \bar{\alpha}_2^{q_2}, \dots, \alpha_m^{q_m}, \bar{\alpha}_m^{q_m},$$

with $p_i, q_j \in \mathbb{Z}$ and $|\alpha_j| = 1$. Hence, we have

$$r = p_2 + 2(q_1 + q_2 + \dots + q_m),$$

and $\det D = (-1)^{p_2}$. Thus, $\det R = (-1)^r$. \square

The quasi-polynomials $m(\chi_i \otimes \delta_\rho; q)$ and $m(\chi_i; q)$ are connected by the following formula.

Theorem 2.10 (Reciprocity theorem). *The following formula holds for an irreducible character χ_i of Γ :*

$$(2.7) \quad m(\chi_i \otimes \delta_\rho; q) = (-1)^\ell m(\chi_i; -q).$$

Proof. Note that since χ_i is an irreducible character, $\chi_i \otimes \delta_\rho$ is also irreducible. Using (2.5), we compute $m(\chi_i \otimes \delta_\rho; q)$ as follows (omitting $\prod_{j=1}^{r(\gamma)}$):

$$(\chi_i \otimes \delta_\rho)(\gamma) \cdot q^{\ell-r(\gamma)} = \chi_i(\gamma)(-1)^{r(\gamma)} q^{\ell-r(\gamma)} = \chi_i(\gamma)(-1)^\ell (-q)^{\ell-r(\gamma)}$$

for each $\gamma \in \Gamma$. This implies (2.7). \square

Note that the map $F(q) = \chi_{L_q}$ can be extended to $F : \mathbb{Z} \rightarrow R(\Gamma)$ as a quasi-polynomial.

Corollary 2.11. *The quasi-polynomial $F : \mathbb{Z} \rightarrow R(\Gamma)$ satisfies $F(q) = (-1)^\ell \delta_\rho F(-q)$.*

Proof. By Theorem 2.10, it follows that

$$\begin{aligned} F(q) &= \chi_{L_q} = \sum_{i=1}^k m(\chi_i; q) \cdot \chi_i \\ &= \sum_{i=1}^k (-1)^\ell m(\chi_i \otimes \delta_\rho; -q) \cdot \chi_i \\ &= (-1)^\ell \sum_{i=1}^k m(\chi_i; -q) (\chi_i \otimes \delta_\rho) \\ &= (-1)^\ell \delta_\rho F(-q). \end{aligned}$$

\square

2.6. Examples. We present some simple examples involving cyclic groups and symmetric groups.

Example 2.12. Let $\Gamma := \mathbb{Z}/6\mathbb{Z}$ be a cyclic group of order 6 generated by σ . Let $\chi : \Gamma \rightarrow \mathbb{C}$ be the function that sends σ to $\zeta_6 := e^{\frac{2\pi\sqrt{-1}}{6}}$. Then the irreducible characters of Γ are $\{\chi, \dots, \chi^5, \chi^6 = \mathbf{1}\}$, where $\mathbf{1}$ is the character of the trivial representation of Γ . Consider the action of Γ on $L := \mathbb{Z}^2$ given by

$$\sigma \mapsto R_\sigma := \begin{pmatrix} 0 & 1 \\ -1 & 1 \end{pmatrix}.$$

To compute χ_{L_q} , we need to compute the rank and the elementary divisors of $R_{\sigma^i} - I_\ell$ for each $i \in \{1, \dots, 5\}$. They are as follows:

$$r(\sigma^i) = 2 \text{ for all } i \in \{1, \dots, 5\}, \quad (e_{\sigma^1,1}, e_{\sigma^1,2}) = (e_{\sigma^5,1}, e_{\sigma^5,2}) = (1, 1),$$

$$(e_{\sigma^2,1}, e_{\sigma^2,2}) = (e_{\sigma^4,1}, e_{\sigma^4,2}) = (1, 3), \quad (e_{\sigma^3,1}, e_{\sigma^3,2}) = (2, 2).$$

Hence, we obtain the multiplicity $m(\chi^j; q)$ as follows:

$$m(\chi^1; q) = m(\chi^5; q) = \begin{cases} \frac{1}{6}(q^2 - 1) & \gcd\{6, q\} = 1; \\ \frac{1}{6}(q^2 - 4) & \gcd\{6, q\} = 2; \\ \frac{1}{6}(q^2 - 3) & \gcd\{6, q\} = 3; \\ \frac{1}{6}(q^2 - 6) & \gcd\{6, q\} = 6, \end{cases}$$

$$m(\chi^2; q) = m(\chi^4; q) = \begin{cases} \frac{1}{6}(q^2 - 1) & \gcd\{6, q\} = 1; \\ \frac{1}{6}(q^2 + 2) & \gcd\{6, q\} = 2; \\ \frac{1}{6}(q^2 - 3) & \gcd\{6, q\} = 3; \\ \frac{1}{6}q^2 & \gcd\{6, q\} = 6, \end{cases}$$

$$m(\chi^3; q) = \begin{cases} \frac{1}{6}(q^2 - 1) & \gcd\{6, q\} = 1; \\ \frac{1}{6}(q^2 - 4) & \gcd\{6, q\} = 2; \\ \frac{1}{6}(q^2 + 3) & \gcd\{6, q\} = 3; \\ \frac{1}{6}q^2 & \gcd\{6, q\} = 6, \end{cases}$$

$$m(\mathbf{1}; q) = \begin{cases} \frac{1}{6}(q^2 + 5) & \gcd\{6, q\} = 1; \\ \frac{1}{6}(q^2 + 8) & \gcd\{6, q\} = 2; \\ \frac{1}{6}(q^2 + 9) & \gcd\{6, q\} = 3; \\ \frac{1}{6}(q^2 + 12) & \gcd\{6, q\} = 6. \end{cases}$$

In this case, since $\delta_\rho = \mathbf{1}$, it follows that $m(\chi^j; q) = m(\chi^j; -q)$ for $j \in \{1, \dots, 6\}$.

We also obtain χ_{L_q} as

$$\chi_{L_q} = \begin{cases} \frac{1}{6}(\chi_R q^2 + 6(\mathbf{1}) - \chi_R) & \gcd\{6, q\} = 1; \\ \frac{1}{6}(\chi_R q^2 + 12(\mathbf{1}) + 6(\chi^2 + \chi^4) - 4\chi_R) & \gcd\{6, q\} = 2; \\ \frac{1}{6}(\chi_R q^2 + 12(\mathbf{1}) + 6\chi^3 - 3\chi_R) & \gcd\{6, q\} = 3; \\ \frac{1}{6}(\chi_R q^2 + 18(\mathbf{1}) + 6(\chi^2 + \chi^3 + \chi^4) - 6\chi_R) & \gcd\{6, q\} = 6, \end{cases}$$

where $\chi_R = \chi + \dots + \chi^6$ is the regular character of Γ .

Example 2.13. As in the previous example, we consider the cyclic group $\Gamma = \mathbb{Z}/6\mathbb{Z}$. The action of Γ on $L := \mathbb{Z}^3$ is given by

$$\sigma \longmapsto R_\sigma := \begin{pmatrix} -1 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix}.$$

By computing in the same way, we obtain the following:

$$\begin{aligned} r(\sigma^1) = r(\sigma^5) = 3, \quad r(\sigma^2) = r(\sigma^4) = 2, \quad r(\sigma^3) = 1, \\ (e_{\sigma^1,1}, e_{\sigma^1,2}, e_{\sigma^1,3}) = (e_{\sigma^5,1}, e_{\sigma^5,2}, e_{\sigma^5,3}) = (1, 1, 6), \\ (e_{\sigma^2,1}, e_{\sigma^2,2}) = (e_{\sigma^4,1}, e_{\sigma^4,2}) = (1, 3), \quad e_{\sigma^3,1} = 2, \end{aligned}$$

and

$$\begin{aligned} m(\chi^1; q) = m(\chi^5; q) &= \begin{cases} \frac{1}{6}(q^3 - q^2 - q + 1) & \gcd\{6, q\} = 1; \\ \frac{1}{6}(q^3 - 2q^2 - q + 2) & \gcd\{6, q\} = 2; \\ \frac{1}{6}(q^3 - q^2 - 3q + 3) & \gcd\{6, q\} = 3; \\ \frac{1}{6}(q^3 - 2q^2 - 3q + 6) & \gcd\{6, q\} = 6, \end{cases} \\ m(\chi^2; q) = m(\chi^4; q) &= \begin{cases} \frac{1}{6}(q^3 + q^2 - q - 1) & \gcd\{6, q\} = 1; \\ \frac{1}{6}(q^3 + 2q^2 - q - 2) & \gcd\{6, q\} = 2; \\ \frac{1}{6}(q^3 + q^2 - 3q - 3) & \gcd\{6, q\} = 3; \\ \frac{1}{6}(q^3 + 2q^2 - 3q - 6) & \gcd\{6, q\} = 6, \end{cases} \\ m(\chi^3; q) &= \begin{cases} \frac{1}{6}(q^3 - q^2 + 2q - 2) & \gcd\{6, q\} = 1; \\ \frac{1}{6}(q^3 - 2q^2 + 2q - 4) & \gcd\{6, q\} = 2; \\ \frac{1}{6}(q^3 - q^2 + 6q - 6) & \gcd\{6, q\} = 3; \\ \frac{1}{6}(q^3 - 2q^2 + 6q - 12) & \gcd\{6, q\} = 6, \end{cases} \\ m(\mathbf{1}; q) &= \begin{cases} \frac{1}{6}(q^3 + q^2 + 2q + 2) & \gcd\{6, q\} = 1; \\ \frac{1}{6}(q^3 + 2q^2 + 2q + 4) & \gcd\{6, q\} = 2; \\ \frac{1}{6}(q^3 + q^2 + 6q + 6) & \gcd\{6, q\} = 3; \\ \frac{1}{6}(q^3 + 2q^2 + 6q + 12) & \gcd\{6, q\} = 6. \end{cases} \end{aligned}$$

In this case, $\delta_p = \chi^3$. Then, we have $m(\chi^1; q) = -m(\chi^4; -q)$ and $m(\chi^3; q) = -m(\mathbf{1}; -q)$.

We also obtain χ_{L_q} as follows:

$$\chi_{L_q} = \begin{cases} \frac{1}{6}(\chi_{\mathbb{R}}q^3 + (\mathbf{1} - \chi_{15} + \chi_{24} - \chi^3)q^2 + (2(\mathbf{1}) - \chi_{15} - \chi_{24} + 2\chi^3)q \\ \quad - (2(\mathbf{1}) + \chi_{15} - \chi_{24} - 2\chi^3)) & \gcd\{6, q\} = 1; \\ \frac{1}{6}(\chi_{\mathbb{R}}q^3 + 2(\mathbf{1} - \chi_{15} + \chi_{24} - \chi^3)q^2 + (2(\mathbf{1}) - \chi_{15} - \chi_{24} + 2\chi^3)q \\ \quad - 2(2(\mathbf{1}) + \chi_{15} - \chi_{24} - 2\chi^3)) & \gcd\{6, q\} = 2; \\ \frac{1}{6}(\chi_{\mathbb{R}}q^3 + (\mathbf{1} - \chi_{15} + \chi_{24} - \chi^3)q^2 + 3(2(\mathbf{1}) - \chi_{15} - \chi_{24} + 2\chi^3)q \\ \quad - 3(2(\mathbf{1}) + \chi_{15} - \chi_{24} - 2\chi^3)) & \gcd\{6, q\} = 3; \\ \frac{1}{6}(\chi_{\mathbb{R}}q^3 + 2(\mathbf{1} - \chi_{15} + \chi_{24} - \chi^3)q^2 + 3(2(\mathbf{1}) - \chi_{15} - \chi_{24} + 2\chi^3)q \\ \quad - 6(2(\mathbf{1}) + \chi_{15} - \chi_{24} - 2\chi^3)) & \gcd\{6, q\} = 6, \end{cases}$$

where $\chi_{15} := \chi^1 + \chi^5$ and $\chi_{24} := \chi^2 + \chi^4$.

Example 2.14. Let $\Gamma := \mathfrak{S}_3$ be the symmetric group of degree 3, which is also the Weyl group of type A_2 . The group Γ has three irreducible characters: the trivial character $\mathbf{1}$, the determinant character δ and the 2-dimensional character χ . Consider the lattice $L := \mathbb{Z}(e_1 - e_2) \oplus \mathbb{Z}(e_2 - e_3)$. The group Γ acts on L as a permutation of $\{e_1, e_2, e_3\}$.

Note that we only need to calculate the rank and the elementary divisors for the representative of each conjugacy class. Choose the representatives $\tau := (1\ 2)$ and $\sigma := (1\ 2\ 3)$. The representation matrices are given by

$$R_{\tau} = \begin{pmatrix} -1 & 1 \\ 0 & 1 \end{pmatrix}, \quad R_{\sigma} = \begin{pmatrix} 0 & -1 \\ 1 & -1 \end{pmatrix}$$

Thus, we have

$$r(\tau) = 1, \quad r(\sigma) = 2, \quad e_{\tau,1} = 1, \quad (e_{\sigma,1}, e_{\sigma,2}) = (1, 3).$$

Therefore, we obtain

$$m(\mathbf{1}; q) = \begin{cases} \frac{1}{6}(q^2 + 3q + 2) & \gcd\{3, q\} = 1; \\ \frac{1}{6}(q^2 + 3q + 6) & \gcd\{3, q\} = 3, \end{cases}$$

$$m(\delta; q) = \begin{cases} \frac{1}{6}(q^2 - 3q + 2) & \gcd\{3, q\} = 1; \\ \frac{1}{6}(q^2 - 3q + 6) & \gcd\{3, q\} = 3, \end{cases}$$

$$m(\chi; q) = \begin{cases} \frac{1}{6}(2q^2 - 2) & \gcd\{3, q\} = 1; \\ \frac{1}{6}(2q^2 - 6) & \gcd\{3, q\} = 3. \end{cases}$$

In this case, $\delta_{\rho} = \delta$. Hence, we have $m(\mathbf{1}; q) = m(\delta; -q)$ and $m(\chi; q) = m(\chi; -q)$.

We also obtain χ_{L_q} as

$$\chi_{L_q} = \begin{cases} \frac{1}{6}(\chi_{\mathbb{R}}q^2 + 3(\mathbf{1} - \delta)q + 2(\mathbf{1} - \delta + \chi)) & \gcd\{3, q\} = 1; \\ \frac{1}{6}(\chi_{\mathbb{R}}q^2 + 3(\mathbf{1} - \delta)q + 6(\mathbf{1} - \delta + \chi)) & \gcd\{3, q\} = 3, \end{cases}$$

where $\chi_R = \mathbf{1} + \delta + 2\chi$. As Haiman mentions in [3, §7.4], the multiplicity $m(\mathbf{1}; q)$ is equal to the Ehrhart quasi-polynomial $L_{\overline{A_0}}(q) = \#(q\overline{A_0} \cap L)$ of the fundamental alcove $\overline{A_0}$ of type A_2 .

REFERENCES

- [1] M. Beck and S. Robins, “Computing the Continuous Discretely”, Undergraduate Texts in Mathematics, Springer, 2007.
- [2] M. D’Adderio, L. Moci. Arithmetic matroids, the Tutte polynomial and toric arrangements. *Adv. Math.* **232**, (2013) 335–67.
- [3] M. D. Haiman, Conjectures on the quotient ring by diagonal invariants, *J. Algebraic Combin.* **3** (1994), 17–76.
- [4] H. Kamiya, A. Takemura, H. Terao, Periodicity of hyperplane arrangements with integral coefficients modulo positive integers, *J. Algebraic Combin.* **27** (2008), no. 3, 317–330.
- [5] H. Kamiya, A. Takemura, H. Terao, Periodicity of Non-Central Integral Arrangements Modulo Positive Integers, *Ann. Comb.* **15** (2011), no. 3, 449–464.
- [6] Y. Liu, T. N. Tran, M. Yoshinaga, G -Tutte polynomials and abelian Lie group arrangements, *Int. Math. Res. Not. IMRN* (2021), no. 1, 152–190.
- [7] P. Orlik and H. Terao, Arrangements of hyperplanes, Grundlehren der Mathematischen Wissenschaften, 300. Springer-Verlag, Berlin, 1992. xviii+325 pp.
- [8] B. Rhoades, Parking structures: Fuss analogs, *J. Algebraic Combin.* **40** (2014), 417–473.
- [9] J.-P. Serre, Linear representations of finite groups. Springer, 1977.
- [10] A. Stapledon, Equivariant Ehrhart theory, *Adv. Math.* **226** (2011), no. 4, 3622–3654.
- [11] T. N. Tran, M. Yoshinaga, Combinatorics of certain abelian Lie group arrangements and chromatic quasi-polynomials. *J. Combin. Theory Ser. A* **165** (2019), 258–272.