De Broglie-Bohm Quantum Mechanics

Antony Valentini^{[1](#page-0-0)}

Theoretical Physics, Blackett Laboratory, Imperial College London, Prince Consort Road, London SW7 2AZ, United Kingdom.

We provide an overview of the de Broglie-Bohm pilot-wave formulation of quantum mechanics, emphasising its applications to field theory, high-energy physics, gravitation, and cosmology.

1 Introduction

- 2 Low-energy many-body systems
- 3 Quantum 'measurements'

4 Quantum relaxation

- 5 Beyond quantum mechanics
	- 5.1 Nonlocality and nonequilibrium signalling
	- 5.2 Breaking uncertainty
	- 5.3 Subquantum measurement
	- 5.4 Subquantum information and subquantum computation

6 Quantum field theory and high-energy physics

- 6.1 Scalar field
	- 6.1.1 Emergent quantum theory and Lorentz invariance
	- 6.1.2 Remarks on Lorentz and Galilean invariance
	- 6.1.3 Quantum field relaxation

6.2 Fermions

6.2.1 Grassmann field theory

- 6.2.2 Dirac sea theory
- 6.3 Electromagnetic field. Scalar QED
- 6.4 Non-Abelian gauge theories
	- 6.4.1 QCD
	- 6.4.2 Electroweak Theory. Spontaneous Symmetry Breaking
- 7 The early universe
	- 7.1 Quantum fields in curved spacetime
	- 7.2 Suppression of quantum relaxation on expanding space
- 8 Quantum gravity and quantum cosmology
	- 8.1 Pilot-wave theory and the Wheeler-DeWitt equation
	- 8.2 Dynamical consistency. Preferred foliation
	- 8.3 Pilot-wave quantum cosmology
	- 8.4 Problem of probability. Beyond the Born rule
- 9 Conclusion
- Bibliography

To appear in: *Encyclopedia of Mathematical Physics, 2nd edition*, Amsterdam: Elsevier.

¹email: a.valentini@imperial.ac.uk

1 Introduction

The de Broglie-Bohm pilot-wave formulation of quantum mechanics was first proposed by de Broglie in 1927, then revived and extended by Bohm in 1952 (de Broglie 1928; Bohm 1952a,b; Holland 1993; Bacciagaluppi and Valentini 2009; Valentini 2025). It is a deterministic dynamics of trajectories for individual quantum systems, where each trajectory is guided by a physical 'pilot wave' in configuration space. The theory is nonlocal, as required by Bell's theorem, and it resolves the quantum measurement problem (Bell 1987). For ensembles of systems, it reproduces the statistical predictions of quantum mechanics if the initial conditions obey the Born rule (Bohm 1952a,b). For more general initial conditions, the Born rule is violated and we obtain new physics outside the domain of quantum mechanics (Valentini 1991a,b, 1992). The validity of the Born rule today can be explained by a past process of dynamical relaxation to 'quantum equilibrium'. A more general nonequilibrium physics may, however, have existed in the early universe, and could exist today in some exotic circumstances. According to pilot-wave theory, quantum physics is only a special case of a wider physics which should be accessible at least in principle.

Employing units $\hbar = 1$, consider a general system with configuration-space wave function $\psi(q, t)$. The Schrödinger equation

$$
i\frac{\partial \psi}{\partial t} = \hat{H}\psi\tag{1}
$$

implies the continuity equation

$$
\frac{\partial |\psi|^2}{\partial t} + \partial_q \cdot j = 0 , \qquad (2)
$$

where $\partial_q \cdot j = 2 \operatorname{Re} \left(i \psi^* \hat{H} \psi \right)$ and ∂_q is a gradient in configuration space. For a given Hamiltonian \hat{H} , we can write $j = j[\psi] = j(q, t)$ in terms of ψ (where j is determined up to a divergence-free term) (Struyve and Valentini 2009). We can then postulate the de Broglie equation of motion

$$
\frac{dq}{dt} = v(q, t) = \frac{j(q, t)}{|\psi(q, t)|^2}
$$
\n(3)

for individual trajectories $q(t)$. In this theory, ψ is a physical field on configuration space guiding the motion of an individual system. For a given wave function $\psi(q, t)$, the actual trajectory $q(t)$ is determined by the initial condition $q(0)$ via the first-order equation of motion [\(3\)](#page-1-0).

Now consider an ensemble of systems with the same wave function ψ . The systems follow the velocity field v, hence a distribution $\rho(q, t)$ of configurations satisfies the continuity equation

$$
\frac{\partial \rho}{\partial t} + \partial_q \cdot (\rho v) = 0 \tag{4}
$$

This takes the same form as [\(2\)](#page-1-1) or

$$
\frac{\partial |\psi|^2}{\partial t} + \partial_q \cdot (|\psi|^2 v) = 0 \tag{5}
$$

(with $j = |\psi|^2 v$). It follows that if $\rho = |\psi|^2$ initially then $\rho = |\psi|^2$ at all times. This is the state of 'quantum equilibrium', with probability density given by the Born rule. In this state, the theory yields the same statistical predictions as quantum mechanics (Bohm 1952a,b).

In principle, however, there can exist initial 'quantum nonequilibrium' ensembles with $\rho \neq |\psi|^2$ (Valentini 1991a,b, 1992). Such ensembles violate the statistical predictions of quantum mechanics. However, in appropriate conditions, such ensembles relax to the equilibrium state $\rho = |\psi|^2$, a process that may have taken place in the early universe, explaining why we observe the Born rule today (Valentini and Westman 2005). Even so, there could exist residual nonequilibrium effects, for example in relic particles from the early universe (Valentini 2007; Underwood and Valentini 2015).

The Schrödinger equation (1) is a field equation on configuration space. It can be derived from a Lagrangian $L = \int \mathcal{L} dq$ where

$$
\mathcal{L} = \frac{i}{2} \left(\psi^* \dot{\psi} - \psi \dot{\psi}^* \right) - \frac{1}{2} \psi^* \hat{H} \psi - \frac{1}{2} \psi \hat{H} \psi^* \tag{6}
$$

is invariant under global phase transformations

$$
\psi \to \psi e^{i\theta} \tag{7}
$$

(where θ is an arbitrary constant). Noether's theorem then implies a local conservation law, which coincides with [\(2\)](#page-1-1) (Struyve and Valentini 2009). Thus the de Broglie velocity [\(3\)](#page-1-0) originates from the global symmetry [\(7\)](#page-2-0) of \mathcal{L} .

We have assumed for simplicity that ψ has only one component. For a system with spin, ψ can have more than one component. The theory is easily generalised to such cases.

2 Low-energy many-body systems

We now consider a system of low-energy spinless particles. The wave function $\psi = \psi(\mathbf{x}_1, \mathbf{x}_2, ..., \mathbf{x}_N, t)$ obeys the Schrödinger equation

$$
i\frac{\partial\psi}{\partial t} = -\sum_{n=1}^{N} \frac{1}{2m_n} \nabla_n^2 \psi + V\psi \;, \tag{8}
$$

which implies a continuity equation

$$
\frac{\partial |\psi|^2}{\partial t} + \sum_{n=1}^{N} \mathbf{\nabla}_n \cdot \left(|\psi|^2 \frac{\mathbf{\nabla}_n S}{m_n} \right) = 0 , \qquad (9)
$$

where $\psi = |\psi| e^{iS}$. We can then identify the de Broglie velocity

$$
\frac{d\mathbf{x}_n}{dt} = \mathbf{v}_n = \frac{1}{m_n} \nabla_n S \tag{10}
$$

of the nth particle. Equations [\(8\)](#page-2-1) and [\(10\)](#page-3-0) define a deterministic dynamics of a many-body system. If the ith and jth particles are entangled, the time evolution of x_i generally depends instantaneously on x_j . The dynamics is nonlocal.

Taking the time derivative of [\(10\)](#page-3-0) and using [\(8\)](#page-2-1) we can write

$$
m_n \frac{d^2 \mathbf{x}_n}{dt^2} = -\nabla_n (V + Q) , \qquad (11)
$$

with a 'quantum potential'

$$
Q = -\sum_{n=1}^{N} \frac{1}{2m_n} \frac{\nabla_n^2 |\psi|}{|\psi|} . \tag{12}
$$

Bohm regarded [\(11\)](#page-3-1) as the equation of motion, with in-principle arbitrary initial conditions for position and momentum. In contrast, in de Broglie's original dynamics, [\(10\)](#page-3-0) is the equation of motion, with in-principle arbitrary initial conditions for position only. Bohm's dynamics has been shown to be unstable and not physically tenable (Colin and Valentini 2014). Even so the form [\(11\)](#page-3-1) is sometimes useful when discussing the classical limit, which in simple cases can be characterised by $Q \approx 0$ (Holland 1993).

We can now consider an ensemble of similar many-body systems with the same wave function ψ . An arbitrary distribution $\rho(\mathbf{x}_1, \mathbf{x}_2, ..., \mathbf{x}_N, t)$ satisfies the continuity equation

$$
\frac{\partial \rho}{\partial t} + \sum_{n=1}^{N} \mathbf{\nabla}_n \cdot (\rho \mathbf{v}_n) = 0 \tag{13}
$$

(cf. [\(4\)](#page-1-3) with $q = (\mathbf{x}_1, \mathbf{x}_2, ..., \mathbf{x}_N)$ and $\partial_q = (\mathbf{\nabla}_1, \mathbf{\nabla}_2, ..., \mathbf{\nabla}_N)$). This takes the same form as [\(9\)](#page-2-2). Thus, as in the general case, if $\rho = |\psi|^2$ initially then $\rho = |\psi|^2$ at later times. This is the state of quantum equilibrium with the Born rule. Again, in principle there can exist initial quantum nonequilibrium ensembles with $\rho \neq |\psi|^2$.

The theory is easily generalised to a system of N (distinguishable) spin- $1/2$ particles with a multi-component wave function $\psi_{s_1s_2...s_N}(\mathbf{x}_1, \mathbf{x}_2, ..., \mathbf{x}_N, t)$ (with spin indices $s_1, s_2, ..., s_N = +, -$). We have the Schrödinger equation (or Pauli equation)

$$
i\frac{\partial\psi}{\partial t} = \sum_{n=1}^{N} \left(-\frac{1}{2m_n} (\boldsymbol{\sigma}_n \cdot \boldsymbol{\nabla}_n)^2 + V \right) \psi
$$
 (14)

(suppressing spin indices), where the vector Pauli spin matrix $\sigma_n = (\sigma_x, \sigma_y, \sigma_z)_n$ acts on the *n*th spin index of ψ . The wave function generates particle trajectories $\mathbf{x}_n(t)$. The de Broglie velocities \mathbf{v}_n follow as usual from the quantum continuity equation. From [\(14\)](#page-3-2) we find a quantum density $\psi^{\dagger} \psi$ and velocities

$$
\mathbf{v}_n = \frac{1}{m_n} \frac{1}{\psi^{\dagger} \psi} \operatorname{Im} \psi^{\dagger} \mathbf{\nabla}_n \psi \tag{15}
$$

(where $\psi^{\dagger} \psi$ is shorthand for $\sum_{s_1s_2...s_N} \psi^*_{s_1s_2...s_N} \psi_{s_1s_2...s_N}$). For a general ensemble with distribution $\rho(\mathbf{x}_1, \mathbf{x}_2, ..., \mathbf{x}_N, t)$ we have

$$
\frac{\partial \rho}{\partial t} + \sum_{n=1}^{N} \mathbf{\nabla}_n \cdot (\rho \mathbf{v}_n) = 0.
$$
 (16)

In this model the particles are not spinning but simply follow trajectories in space (Bell 1966; 1987, chaps. 1, 15 and 17). For example, for a single particle in an external magnetic field **B**, the Hamiltonian includes a term $\mu \sigma \cdot \mathbf{B}$, where μ is the magnetic moment, and the gauge-invariant de Broglie velocity includes a term $-(e/m)\mathbf{A}^2$ $-(e/m)\mathbf{A}^2$ If the particle enters a Stern-Gerlach apparatus, it will be deflected upwards or downwards, depending on its initial position, corresponding to 'spin up' and 'spin down'.

For N identical particles we have the symmetry condition

$$
\psi_{s_1...s_i...s_j...s_N}(\mathbf{x}_1,...,\mathbf{x}_i,...,\mathbf{x}_j,...,\mathbf{x}_N,t) = \pm \psi_{s_1...s_j...s_N}(\mathbf{x}_1,...,\mathbf{x}_j,...,\mathbf{x}_i,...,\mathbf{x}_N,t),
$$
\n(17)

for distinct i, $j = 1, ..., N$, with + for bosons and − for fermions. This ensures that the set of 3-space velocities is invariant under exchange $i \leftrightarrow j$ of labels (Bacciagaluppi 2003; Sebens 2016).

3 Quantum 'measurements'

We consider a so-called quantum 'measurement' for a general observable $\hat{\omega}$. The system (with configuration q) is coupled to an apparatus pointer y via the interaction Hamiltonian

$$
\hat{H}_{\rm I} = a\hat{\omega}\hat{p}_y \tag{18}
$$

where a is the coupling and $\hat{p}_y = -i\partial_y$. During the interaction a is so large we can neglect the rest of the Hamiltonian. The initial system wave function $\psi_0(q) = \sum_n c_n \phi_n(q)$ is a superposition of eigenstates $(\hat{\omega}\phi_n = \omega_n \phi_n)$, and the initial pointer packet $g_0(y)$ is narrowly-peaked around $y = 0$ with width Δy . The initial joint wave function $\Psi_0(q, y) = \psi_0(q)g_0(y)$ evolves to

$$
\Psi(q, y, t) = \sum_{n} c_n \phi_n(q) g_0(y - a\omega_n t) . \qquad (19)
$$

The nth term shifts by $a\omega_n t$ in y-space. At sufficiently large times the 'branches' have negligible overlap.

²For pilot-wave theory in external electromagnetic fields see Holland (1993).

In standard quantum mechanics it is usual to postulate a random 'collapse' to the nth branch with probability $p_n = |c_n|^2$. No such postulate is needed in pilot-wave theory. The configuration $Q(t) = (q(t), y(t))$ can occupy (the support of) only one of the non-overlapping terms in [\(19\)](#page-4-1). Inspection of the de Broglie equations of motion for Q shows that, after the branches have separated, the velocity \dot{Q} is determined by the occupied nth branch only. Furthermore, because each term in [\(19\)](#page-4-1) separates in q and y, the system velocity \dot{q} is determined by $\phi_n(q)$ only. At the end of the experiment the system has an effective 'collapsed' wave function

$$
\psi_{\text{coll}} = \phi_n(q) \tag{20}
$$

The probability for this to occur is just the probability p_n for $Q(t)$ to occupy the nth branch. For an equilibrium ensemble, with distribution $P(Q, t) = |\Psi(Q, t)|^2$, this is found by integrating $|\Psi(Q, t)|^2$ over the support of the *n*th term, yielding $p_n = |c_n|^2$. For a nonequilibrium ensemble, with $P(Q, t) \neq |\Psi(Q, t)|^2$, in general $p_n \neq |c_n|^2$. Thus in equilibrium we obtain the usual Born rule for 'measurement' outcomes, while in nonequilibrium the Born rule is violated.

After the described experiment, there are still empty branches of Ψ in other regions of configuration space. But these have no effect on the system for as long as they remain separated from the occupied branch. Subsequent interaction between y and the environment ensures that a later reoverlap (with respect to all relevant degrees of freedom) is unlikely (Bohm 1952b, p. 182). For most practical purposes, the 'collapse' is irreversible.[3](#page-5-0)

A so-called 'measurement of $\hat{\omega}$ ' reduces to an approximate position measurement for y (to indicate the occupied branch). It is important to recognise that the 'outcome' ω_n is not usually equal to the value of any previouslyexisting property of the system, and therefore the quantum 'measurement' is not usually a correct measurement. For example, for a particle in one dimension with initial wave function $\psi_0(x) \propto e^{ipx} + e^{-ipx}$, a so-called 'measurement of momentum' $(\hat{\omega} = \hat{p})$ yields outcomes $\pm p$, while the actual initial momentum $p_0 = \partial S_0/\partial x = 0$. Similarly, for the same initial state, a 'measurement of kinetic energy' $(\hat{\omega} = \hat{p}^2/2m)$ yields a value $p^2/2m$ while the actual value is again 0 (in this case both before and after the interaction). It might be said that these measurements are 'unfaithful', which is to say they are not correct measurements. Notable exceptions are quantum measurements of position $(\hat{\omega} = \hat{x})$, which do correctly indicate the (previously-existing) particle position.

In general quantum 'measurements' are not correct measurements. They are simply experiments that bring about a final state where the system has an effective wave function ϕ_n . Even so, in quantum equilibrium, there is no observable difference from standard quantum mechanics: the distribution of pointer positions is the same. It is only the interpretation that differs.

³ In principle the empty branches can play a role if a reoverlap somehow occurs. The scenario of 'Wigner's friend' and more recent extensions thereof can be understood only if we take into account the empty branches (Bacciagaluppi and Valentini 2009, p. 147; Lazarovici and Hubert 2019).

4 Quantum relaxation

We can explain the emergence of the Born rule by a dynamical process of quantum relaxation. Consider an ensemble of systems with wave function $\psi(q, t)$, whose configurations q have a distribution $\rho(q,t)$. Because ρ and $|\psi|^2$ obey the same continuity equation, the ratio $f = \rho / |\psi|^2$ obeys an analogue of the classical Liouville theorem,

$$
\frac{df}{dt} = 0\,,\tag{21}
$$

where $d/dt = \partial/\partial t + v \cdot \partial_q$ is the time derivative along trajectories with de Broglie velocity v. Despite this, quantum relaxation can occur for coarse-grained densities $\bar{\rho}$ and $|\psi|^2$ (averaged over small coarse-graining cells),

$$
\bar{\rho}(q,t) \to \overline{|\psi(q,t)|^2} \tag{22}
$$

This can be quantified by a decrease of the coarse-grained H-function

$$
\bar{H}(t) = \int dq \ \bar{\rho} \ln(\bar{\rho}/\overline{|\psi|^2}) \tag{23}
$$

(minus the relative entropy of $\bar{\rho}$ with respect to $\overline{|\psi|^2}$), where $\bar{H} \ge 0$ and $\bar{H} = 0$ if and only if $\bar{\rho} = |\psi|^2$. This satisfies a coarse-graining H-theorem (Valentini 1991a)

$$
\bar{H}(t) \le \bar{H}(0) \tag{24}
$$

assuming there is no initial fine-grained structure,

$$
\bar{\rho}(q,0) = \rho(q,0) , \quad \overline{|\psi(q,0)|^2} = |\psi(q,0)|^2 . \tag{25}
$$

In fact, $\bar{H}(t)$ strictly decreases $(\bar{H}(t) < \bar{H}(0))$ when ρ develops fine-grained structure ($\rho \neq \bar{\rho}$), which generally occurs when v varies over the coarse-graining cells (Valentini 1992).

For an ensemble undergoing efficient quantum relaxation, $\bar{H}(t) \rightarrow 0$ and $\bar{\rho} \rightarrow |\psi|^2$ (at least to a good approximation). To show that this actually occurs requires numerical simulations (Valentini and Westman 2005). The relaxation timescale depends on the system as well as on the initial conditions.

An example is shown in Fig. 1, for a two-dimensional oscillator whose wave function is a superposition of 25 energy states (Abraham, Colin, and Valentini 2014). Plotting H against t (Fig. 2) we find an approximately exponential decay,

$$
\bar{H}(t) \approx \bar{H}_0 e^{-t/\tau} \tag{26}
$$

with a best-fit timescale $\tau \simeq 6$ (in our units), an order of magnitude larger than the quantum timescale $\Delta t = \hbar / \Delta E \simeq 0.5$.

The timescale τ generally depends on the number M of energy modes in ψ as well as on the coarse-graining length ε , tending to be smaller for larger M and for larger ε . For particles in a two-dimensional box, there is strong evidence for

relaxation of initial nonequilibrium ($\rho \neq |\psi|^2$)

Figure 1: Quantum relaxation for a two-dimensional oscillator (Abraham, Colin, and Valentini 2014). The initial density $\rho(x, y, 0)$ is a Gaussian. The wave function $\psi(x, y, t)$ has time period 2π (units $\hbar = m = \omega = 1$).

an approximate scaling $\tau \propto 1/M$ (at fixed ε) and weak evidence for $\tau \sim \varepsilon^{-1/4}$ (at fixed M) (Towler, Russell, and Valentini 2012). Efficient relaxation has been found for M as low as $M = 4$ (Abraham, Colin, and Valentini 2014).

In Fig. 2 the best-fit 'residue' $c \approx 0.02$ is comparable to the numerical error in H, so the results are consistent with $H(t) \to 0$ as $t \to \infty$. A small but nonzero residue $\bar{H}(t) \to \bar{H}_{\text{res}} > 0$ has been demonstrated for a case with $M = 4$. Residues can occur when the trajectories show significant confinement (not fully exploring the support of $|\psi|^2$). Whether or not there is significant confinement depends on the initial phases in ψ , and is less likely to occur when M is larger (Abraham, Colin, and Valentini 2014). Laboratory systems have a long and violent astrophysical history. In the past M will have been very large, and so we can expect equilibrium to have been reached to very high accuracy.

Simulations of quantum relaxation have also been carried out for Dirac fermions (Colin 2012), for initial extreme nonequilibrium (Underwood 2018), for systems with small perturbations (Kandhadai and Valentini 2019), and for coupled systems (Lustosa, Colin, and Perez Bergliaffa 2021; Lustosa, Pinto-Neto, and Valentini 2023). In general, for relaxing systems the trajectories tend to be quite erratic, and neighbouring trajectories quickly diverge. Chaos plays a role in quantum relaxation (Efthymiopoulos, Contopoulos, and Tzemos 2017; Drezet 2021).

Some authors derive the Born rule seen in the laboratory today from the Born rule applied to the beginning of the universe (Dürr, Goldstein, and Zanghì

Figure 2: Approximate exponential decay of $\bar{H}(t)$ for a two-dimensional oscillator (Abraham, Colin, and Valentini 2014). The error is estimated by running simulations with different grids (solid curves). The dashed curve shows a best fit to an exponential function $a \exp[-b(t/2\pi)] + c$.

1992; Dürr and Teufel 2009). However, this approach is conceptually circular: the Born rule is assumed at $t = 0$ in order to derive the Born rule at later times (Valentini 1996, 2020). It aids clarity to bear in mind that, in a deterministic theory of individual systems, initial conditions are contingencies not fixed by any law or principle. In pilot-wave theory, whether the universe began in equilibrium or nonequilibrium is ultimately an empirical question (Section 7.2).

5 Beyond quantum mechanics

According to pilot-wave theory the Born rule can be broken, resulting in new physics outside the domain of standard quantum theory.

5.1 Nonlocality and nonequilibrium signalling

Pilot-wave dynamics is nonlocal. Consider for example a pair of entangled spin-1/2 particles A and B. If particle A enters a Stern-Gerlach apparatus with orientation m_A , while particle B enters a second Stern-Gerlach apparatus with orientation \mathbf{m}_B , then the (upwards or downwards) motion of particle A depends nonlocally on the 'measurement setting' m_B of the second apparatus (Bell 1966, 1987). Such nonlocal effects are generic for entangled systems, and are required by Bell's theorem. Changing the local Hamiltonian H_B at one wing of an entangled state instantaneously affects the particle motion $x_A(t)$ at the other wing, no matter how widely separated the particles may be.

Now consider an ensemble of entangled pairs in the singlet state. For an equilibrium ensemble with $\rho(\mathbf{x}_A, \mathbf{x}_B, t) = \sum_{s_1 s_2} |\psi_{s_1 s_2}(\mathbf{x}_A, \mathbf{x}_B, t)|^2$, and for given measurement settings \mathbf{m}_A , \mathbf{m}_B , we find an even ratio of 'spin' outcomes +, − at A (as in quantum mechanics). Under a change $\mathbf{m}_B \to \mathbf{m}'_B$ of setting at B, some of the outcomes at A will change. A fraction $\nu_A(+,-)$ of the ensemble makes the transition + \rightarrow - at A, while a fraction $\nu_A(-, +)$ makes the reverse transition $-\rightarrow +$ at A ^{[4](#page-9-0)}. In equilibrium we have 'detailed balancing':

$$
\nu_A(+,-) = \nu_A(-,+) \ . \tag{27}
$$

Despite the change in some individual outcomes at A, the marginal distribution at A remains the same. This is emergent statistical locality: in equilibrium the nonlocal effects average to zero. In contrast, for a nonequilibrium ensemble with $\rho(\mathbf{x}_A, \mathbf{x}_B, t) \neq \sum_{s_1s_2} |\psi_{s_1s_2}(\mathbf{x}_A, \mathbf{x}_B, t)|^2$, in general

$$
\nu_A(+,-) \neq \nu_A(-,+)\,,\tag{28}
$$

in which case the marginal distribution at A changes and there is a net statistical nonlocal signal from B to A. In nonequilibrium, the nonlocal effects need not average to zero and statistical locality can be violated (Valentini 1991b, 2002a). Similar arguments apply to any nonlocal deterministic hidden-variables theory (Valentini 2002b).

5.2 Breaking uncertainty

The uncertainty principle is also a peculiarity of equilibrium and is generally violated for nonequilibrium ensembles. For example, for a single particle in one dimension, the outcome p of a quantum 'measurement' of momentum generally depends on the initial particle position $x(0)$ within the support of the initial wave function $\psi(x, 0)$. The distribution of outcomes p then depends on the distribution $\rho(x,0)$ of initial positions. For $\rho(x,0) = |\psi(x,0)|^2$ the statistical spread Δp matches the prediction of quantum mechanics and is therefore consistent with the usual uncertainty lower bound $\Delta x \Delta p \geq \hbar/2$. For $\rho(x,0) \neq |\psi(x,0)|^2$ the spread Δp will be anomalous and the uncertainty principle can be violated (Valentini 1991b, 2025).

5.3 Subquantum measurement

Nonequilibrium systems can be employed to perform 'subquantum measurements' whereby we measure the position x of a particle without disturbing its wave function $\psi_0(x)$ (Valentini 2002c). For simplicity we work in one dimension. We assume the particle has an initial equilibrium probability density $\rho_0(x) = |\psi_0(x)|^2$. A standard quantum measurement of \hat{x} will reduce the (effective) wave function to a narrow packet centred on the measured value. This

⁴The actual values of the fractions depend on \mathbf{m}_A , \mathbf{m}_B , \mathbf{m}'_B and have been calculated in a simplified model of spin (Valentini 2002a).

can be avoided if the initial pointer coordinate y has a narrow nonequilibrium probability density $\pi_0(y) \neq [g_0(y)]^2$. If $\pi_0(y)$ is arbitrarily narrow, the initial position x_0 can be measured without disturbing $\psi_0(x)$, to arbitrary accuracy.

This is done by switching on the interaction Hamiltonian [\(18\)](#page-4-2), with $\hat{\omega} = \hat{x}$, for a short time only. During that time we have a Schrödinger equation $\frac{\partial \Psi}{\partial t} =$ $-ax\partial\Psi/\partial y$, so the initial wave function $\Psi_0(x, y) = \psi_0(x)g_0(y)$ evolves into

$$
\Psi(x, y, t) = \psi_0(x)g_0(y - axt) . \tag{29}
$$

The continuity equation $\partial |\Psi|^2 / \partial t + ax \partial |\Psi|^2 / \partial y = 0$ implies the simple de Broglie velocities $\dot{x} = 0$, $\dot{y} = ax$ and trajectories

$$
x(t) = x_0, \ y(t) = y_0 + ax_0t \ . \tag{30}
$$

Now, for $at \to 0$ (with a fixed), $\Psi(x, y, t) \to \psi_0(x)g_0(y)$ and the particle wave function $\psi_0(x)$ is undisturbed. Even so, for arbitrarily small at, the pointer $y(t)$ contains precise information about x_0 , which will be visible to the experimenter if $\pi_0(y)$ is sufficiently narrow. From the continuity equation $\partial P/\partial t + ax \partial P/\partial y =$ 0, the initial distribution $P_0(x, y) = |\psi_0(x)|^2 \pi_0(y)$ evolves into

$$
P(x, y, t) = |\psi_0(x)|^2 \pi_0(y - axt) . \qquad (31)
$$

Let $\pi_0(y)$ be highly localised around $y = 0$, with $\pi_0(y) = 0$ for $|y| > w/2$. From a standard (faithful) measurement of y we may deduce that x lies in the interval $(y/at - w/2at, y/at + w/2at)$ (where $P(x, y, t) \neq 0$ only if $|y - axt| < w/2$). Taking simultaneous limits $at \to 0$ and $w \to 0$, with $w/at \to 0$, the midpoint $y/at \to x_0$ (since $y = y_0 + ax_0t$ and $|y_0| < w/2$), while the error $w/2at \to 0$. We then have an exact measurement of x_0 , with no disturbance of $\psi_0(x)$.

An arbitrarily fine time sequence of such measurements can also track the trajectory $x(t)$ without disturbing the evolving wave function $\psi(x, t)$.

5.4 Subquantum information and subquantum computation

In quantum mechanics non-orthogonal quantum states $|\psi_1\rangle$, $|\psi_2\rangle$ ($\langle \psi_1|\psi_2\rangle \neq$ 0) cannot be reliably distinguished for a single system (Nielsen and Chuang 2000). This theorem breaks down in nonequilibrium pilot-wave theory (Valentini 2002c). The associated sets of trajectories $q_1(t)$, $q_2(t)$ generally differ, hence subquantum monitoring of trajectories can render non-orthogonal states distinguishable.

The E91 quantum-cryptographic protocol (Ekert 1991) is rendered insecure if subquantum measurements allow an eavesdropper to predict quantum 'measurement' outcomes at each wing of a (bipartite) entangled state that is being employed to generate a shared secret key. The B92 protocol (Bennett 1992) is rendered insecure if subquantum measurements allow an eavesdropper to distinguish non-orthogonal states transmitted between two parties. Thus the discovery of quantum nonequilibrium systems would undermine the security of the quantum internet.

The ability to distinguish non-orthogonal states reliably also allows new and potentially more powerful forms of computation. These could be realised in nonlinear quantum mechanics (Abrams and Lloyd 1998) or in nonequilibrium pilot-wave theory (Valentini 2002c). However, to evaluate the power of 'subquantum computing' we need to quantify how the nonequilibrium resources scale (polynomially or exponentially) with the size of the computational task. Such studies have yet to be carried out.

6 Quantum field theory and high-energy physics

We now consider pilot-wave field theory and high-energy physics on flat spacetime. This is formulated with a preferred rest frame and a preferred time parameter t . Nonlocality acts instantaneously along hypersurfaces of constant t . Lorentz invariance emerges as an effective symmetry for equilibrium ensembles.

6.1 Scalar field

Employing units $c = 1$ we first consider a real massive scalar field (Bohm, Hiley, and Kaloyerou 1987; Valentini 1992; Holland 1993). This has a classical Lagrangian $L = \int \mathcal{L} d^3x$ with

$$
\mathcal{L} = \frac{1}{2} (\dot{\phi}^2 - (\nabla \phi)^2 - m^2 \phi^2) . \tag{32}
$$

The system has a configuration $q = \phi(\mathbf{x})$. In the functional Schrödinger picture, the wave functional $\Psi[\phi, t] = \langle \phi | \Psi(t) \rangle$ (where $|\phi\rangle$ is a field eigenstate) satisfies the Schrödinger equation^{[5](#page-11-0)}

$$
i\frac{\partial\Psi}{\partial t} = \int d^3x \; \frac{1}{2} \left(-\frac{\delta^2}{\delta\phi^2} + (\nabla\phi)^2 + m^2\phi^2 \right) \Psi \; . \tag{33}
$$

This implies a continuity equation

$$
\frac{\partial |\Psi|^2}{\partial t} + \int d^3x \; \frac{\delta}{\delta \phi} \left(|\Psi|^2 \frac{\delta S}{\delta \phi} \right) = 0 \tag{34}
$$

(with $\Psi = |\Psi| e^{iS}$) and a de Broglie velocity $v = \delta S/\delta \phi$. We then have the equation of motion

$$
\frac{\partial \phi}{\partial t} = \frac{\delta S}{\delta \phi} \tag{35}
$$

for the trajectory $q(t) = \phi(\mathbf{x}, t)$. If field elements $\phi(\mathbf{x})$, $\phi(\mathbf{x}')$ at different points $\mathbf{x} \neq \mathbf{x}'$ are entangled, the time evolution of $\phi(\mathbf{x})$ generally depends instantaneously on $\phi(\mathbf{x}')$. The nonlocal dynamics singles out an absolute simultaneity labelled by the global time parameter t .

⁵The functional derivative $\delta\Psi/\delta\phi(\mathbf{x})$ is defined by $\delta\Psi = \int d^3\mathbf{x} \left[\delta\Psi/\delta\phi(\mathbf{x}) \right] \delta\phi(\mathbf{x})$ for arbitrary infinitesimal $\delta\phi(\mathbf{x})$.

For a general ensemble of fields with the same wave functional Ψ , the probability distribution $P[\phi, t]$ evolves by the continuity equation

$$
\frac{\partial P}{\partial t} + \int d^3x \frac{\delta}{\delta \phi} \left(P \frac{\delta S}{\delta \phi} \right) = 0 \tag{36}
$$

6.1.1 Emergent quantum theory and Lorentz invariance

For an equilibrium ensemble, with $P[\phi, t] = |\Psi(\phi, t)|^2$, we recover the usual statistical predictions of quantum field theory. Reference to [\(35\)](#page-11-1) may be dropped. We may write [\(33\)](#page-11-2) for a quantum state $|\Psi(t)\rangle$ acted on by linear operators $\hat{\phi}(\mathbf{x}), \hat{\pi}(\mathbf{x}),$ where $\hat{\phi}(\mathbf{x}) \rightarrow \phi(\mathbf{x}), \hat{\pi}(\mathbf{x}) \rightarrow -i\delta/\delta\phi(\mathbf{x})$ in the basis $|\phi(\mathbf{x})\rangle$. Transforming to the Heisenberg picture, operators $\hat{\omega}(\mathbf{x}, t)$ have a time evolution $\partial_t \hat{\omega} = -i[\hat{\omega}, \hat{H}]$ with Hamiltonian

$$
\hat{H} = \int d^3x \; \frac{1}{2} \left(\hat{\pi}^2 + (\nabla \hat{\phi})^2 + m^2 \hat{\phi}^2 \right) \; . \tag{37}
$$

We find $\partial_t \hat{\phi} = \hat{\pi}$ and the Lorentz-covariant Klein-Gordon equation

$$
\partial_t^2 \hat{\phi} - \nabla^2 \hat{\phi} + m^2 \hat{\phi} = 0.
$$
 (38)

Lorentz invariance emerges as an equilibrium symmetry, along with statistical locality and uncertainty.

For a quantum nonequilibrium ensemble, with $P[\phi, t] \neq |\Psi(\phi, t)|^2$, the usual statistical results of quantum field theory break down, with potentially observable implications for the early universe (Section 7). Furthermore, nonlocal signalling is possible and Lorentz invariance is broken.

At the fundamental level we have an Aristotelian spacetime $E \times E^3$ with a preferred state of rest, while in quantum equilibrium we obtain an effective Minkowski spacetime M^4 . The emergent symmetry group depends, however, on the structure of the field Hamiltonian. For example if, in [\(37\)](#page-12-0), we replace $(\nabla \hat{\phi})^2$ by $-(\nabla \hat{\phi})^2$, we obtain [\(38\)](#page-12-1) with a plus sign in the second term. The symmetries of the operator wave equation then correspond to an emergent Euclidean spacetime E^4 . As far as we know, for the Hamiltonians found in nature the emergent spacetime is Lorentzian. We might ask if there are deeper reasons for this. In some models of particle physics and quantum gravity, Lorentz invariance breaks down at high energies (Kostelecký and Mewes 2002; Hořava 2009). In a pilot-wave formulation of such models, Lorentz invariance emerges in equilibrium at low energies only.

6.1.2 Remarks on Lorentz and Galilean invariance

Alternatively, some workers attempt to construct a fundamentally Lorentzinvariant pilot-wave theory. This program has not succeeded. The dynamics and the quantum equilibrium distribution can be defined consistently only with a preferred rest frame (Hardy 1992; Berndl et al. 1996). The definition of the preferred frame might be formally Lorentz covariant (D¨urr et al. 1999; Tumulka 2007; Dürr et al. 2014). For example the preferred frame might coincide with the mean rest frame (Vigier 1985; Dürr et al. 2014). While the equations may appear formally Lorentz covariant, physically there is still a preferred frame.

Some workers regard the low-energy particle theory as Galilean covariant, and appeal to this purported symmetry to constrain the form of the guidance equation [\(10\)](#page-3-0) (Dürr, Goldstein, and Zanghì 1992). However, because the dynamics has a law of motion for velocity and not for acceleration, the natural kinematics is Aristotelian with a preferred state of rest (Valentini 1997). Galilean covariance is then a fictitious symmetry of the low-energy theory – just as covariance under transformation to a uniformly-accelerated frame is (well known to be) a fictitious symmetry of Newtonian mechanics. Furthermore, for systems of particles and more generally, the de Broglie velocity [\(3\)](#page-1-0) originates from the global symmetry [\(7\)](#page-2-0) on configuration space and not from any symmetry in space or spacetime. A natural Aristotelian kinematics is also consistent with a preferred simultaneity defined by nonequilibrium nonlocal signalling (Valentini 2008).

6.1.3 Quantum field relaxation

The theory of quantum relaxation is readily extended to fields.

It is convenient to work in Fourier space with $\Psi[\phi_{\mathbf{k}}, t]$ a function of the Fourier components $\phi_{\mathbf{k}}$. Considering a box of volume V with periodic boundary conditions, we have a discrete system. We can write $\phi_{\mathbf{k}} = \frac{\sqrt{V}}{(2\pi)^{3/2}} (q_{\mathbf{k}1} + iq_{\mathbf{k}2}),$ with two real degrees of freedom $q_{\mathbf{k}r}$ ($r = 1, 2$) for each field mode **k**. For an independent (unentangled) field mode, and taking $m = 0$, the effective wave function $\psi_{\mathbf{k}}(q_{\mathbf{k}1}, q_{\mathbf{k}2}, t)$ satisfies a two-dimensional Schrödinger equation

$$
i\frac{\partial\psi_{\mathbf{k}}}{\partial t} = -\frac{1}{2}\left(\frac{\partial^2}{\partial q_{\mathbf{k}1}^2} + \frac{\partial^2}{\partial q_{\mathbf{k}2}^2}\right)\psi_{\mathbf{k}} + \frac{1}{2}k^2\left(q_{\mathbf{k}1}^2 + q_{\mathbf{k}2}^2\right)\psi_{\mathbf{k}}\,,\tag{39}
$$

while the de Broglie equations of motion for q_{k1} , q_{k2} read

$$
\frac{dq_{\mathbf{k}1}}{dt} = \frac{\partial s_{\mathbf{k}}}{\partial q_{\mathbf{k}1}}, \quad \frac{dq_{\mathbf{k}2}}{dt} = \frac{\partial s_{\mathbf{k}}}{\partial q_{\mathbf{k}2}} \tag{40}
$$

(with $\psi_{\mathbf{k}} = |\psi_{\mathbf{k}}| e^{is_{\mathbf{k}}}\rangle$. A general (marginal) distribution $\rho_{\mathbf{k}}(q_{\mathbf{k}1}, q_{\mathbf{k}2}, t)$ obeys

$$
\frac{\partial \rho_{\mathbf{k}}}{\partial t} + \frac{\partial}{\partial q_{\mathbf{k}1}} \left(\rho_{\mathbf{k}} \frac{\partial s_{\mathbf{k}}}{\partial q_{\mathbf{k}1}} \right) + \frac{\partial}{\partial q_{\mathbf{k}2}} \left(\rho_{\mathbf{k}} \frac{\partial s_{\mathbf{k}}}{\partial q_{\mathbf{k}2}} \right) = 0.
$$
 (41)

This system is mathematically equivalent to a two-dimensional oscillator of mass $m = 1$ and angular frequency $\omega = k$. Relaxation results for the oscillator (Section 4) then apply to a single field mode (Valentini 2007). On expanding space field relaxation is suppressed at long wavelengths, with potentially important implications for the early universe (Section 7).

6.2 Fermions

Bosons are usually described in terms of fields as above. For fermions, there are both field-theoretic and particle-trajectory approaches.^{[6](#page-14-0)}

6.2.1 Grassmann field theory

A pilot-wave theory of a massive spin-1/2 Grassmann field (Valentini 1992, 1996) is conveniently written in terms of a two-component van der Waerden field, which is equivalent to the more usual four-component Dirac field (Brown 1958; Sakurai 1967).

The two-component complex field ϕ_{α} ($\alpha = 1, 2$) has a classical Lagrangian $L = \int \mathcal{L} d^3x$ with

$$
\mathcal{L} = \dot{\phi}_{\alpha}^* \dot{\phi}_{\alpha} - (\boldsymbol{\sigma} \cdot \boldsymbol{\nabla} \phi)_{\alpha}^* (\boldsymbol{\sigma} \cdot \boldsymbol{\nabla} \phi)_{\alpha} - m^2 \phi_{\alpha}^* \phi_{\alpha} , \qquad (42)
$$

where σ^i are Pauli spin matrices and α is summed over. This yields the classical wave equation

$$
\partial_t^2 \phi - (\boldsymbol{\sigma} \cdot \boldsymbol{\nabla})^2 \phi + m^2 \phi = 0 \tag{43}
$$

(suppressing indices). A four-component field ψ obeying the Dirac equation can be constructed from linear combinations of ϕ with its first space and time derivatives.

In pilot-wave theory $\phi_{\alpha}, \phi_{\alpha}^*$ are taken to be complex Grassmann fields. These anticommute,

$$
\{\phi_{\alpha}(\mathbf{x}), \phi_{\beta}(\mathbf{y})\} = \{\phi_{\alpha}^*(\mathbf{x}), \phi_{\beta}^*(\mathbf{y})\} = \{\phi_{\alpha}^*(\mathbf{x}), \phi_{\beta}(\mathbf{y})\} = 0 \tag{44}
$$

(where $\{a, b\} \equiv ab + ba$), with derivatives defined from the left $(\overrightarrow{\delta}/\delta\phi)$ and from the right $(\overleftarrow{\delta}/\delta\phi)$ (Berezin 1966). We have the Schrödinger equation for $\Psi = \Psi[\phi_\alpha, \phi^*_\alpha, t]$:

$$
i\frac{\partial\Psi}{\partial t} = \int d^3x \, \left(-\frac{\overrightarrow{\delta}}{\delta\phi^*_{\alpha}} \left(\Psi \frac{\overleftarrow{\delta}}{\delta\phi_{\alpha}} \right) + (\boldsymbol{\sigma} \cdot \boldsymbol{\nabla}\phi)^*_{\alpha} (\boldsymbol{\sigma} \cdot \boldsymbol{\nabla}\phi)_{\alpha} \Psi + m^2 \phi^*_{\alpha} \phi_{\alpha} \Psi \right) \,. \tag{45}
$$

This implies a continuity equation

$$
\frac{\partial |\Psi|^2}{\partial t} + \int d^3x \, \left(\frac{\overrightarrow{\delta}}{\delta \phi^*_{\alpha}} \left(|\Psi|^2 \frac{S \overleftarrow{\delta}}{\delta \phi_{\alpha}} \right) + \left(|\Psi|^2 \frac{\overrightarrow{\delta} S}{\delta \phi^*_{\alpha}} \right) \frac{\overleftarrow{\delta}}{\delta \phi_{\alpha}} \right) = 0 \ ,
$$

←−δ

from which we can identify the de Broglie velocities

$$
\frac{\partial \phi_{\alpha}^{*}}{\partial t} = \frac{S^{'\overline{\delta}}}{\delta \phi_{\alpha}} , \qquad \frac{\partial \phi_{\alpha}}{\partial t} = \frac{\overrightarrow{\delta} S}{\delta \phi_{\alpha}^{*}} . \tag{46}
$$

Grassmann variables can be represented by anticommuting matrices. We may then regard the matrix entries as the evolving objects.

 6 Though widely seen as problematic, some authors consider particle-trajectory models for bosons as well.

6.2.2 Dirac sea theory

An alternative theory of (high-energy) fermions has been developed in terms of particle trajectories (Bohm, Hiley, and Kaloyerou 1987; Bohm and Hiley 1993; Colin 2003; Colin and Struyve 2007). In this model there are trajectories for particles filling the (negative-energy) Dirac sea as well as for particles with positive energies. Anti-particles are 'holes' in the Dirac sea. We confine ourselves to the free theory. Inclusion of external fields and interactions is straightforward.

The trajectories are guided by a multi-component wave function $\psi_{\alpha_1\alpha_2...\alpha_n}$... $(\mathbf{x}_1, \mathbf{x}_2, ..., \mathbf{x}_n, ..., t)$ (with $\alpha_n = 1, ..., 4$) obeying the many-body Dirac equation

$$
i\frac{\partial\psi}{\partial t} = \sum_{n} \left(-i\alpha_n \cdot \nabla_n + \beta_n m \right) \psi \tag{47}
$$

(suppressing spinor indices), where the 4×4 matrices α_n , β_n act on the *n*th spinor index of ψ . This implies a continuity equation

$$
\frac{\partial(\psi^{\dagger}\psi)}{\partial t} + \sum_{n} \nabla_n \cdot (\psi^{\dagger} \alpha_n \psi) = 0 \tag{48}
$$

(here $\psi^{\dagger}\psi$ is shorthand for $\sum_{\alpha_1\alpha_2...\alpha_n...\psi_{\alpha_1\alpha_2...\alpha_n...\psi_{\alpha_1\alpha_2...\alpha_n...}}$). We then have a quantum equilibrium density $\rho = \psi^{\dagger} \psi$ and the de Broglie guidance equation

$$
\frac{d\mathbf{x}_n}{dt} = \frac{\psi^\dagger \mathbf{\alpha}_n \psi}{\psi^\dagger \psi} \; . \tag{49}
$$

More general distributions $\rho \neq \psi^{\dagger} \psi$ are in principle possible. A general density ρ evolves by

$$
\frac{\partial \rho}{\partial t} + \sum_{n} \mathbf{\nabla}_n \cdot (\rho \mathbf{v}_n) = 0 \tag{50}
$$

with $\mathbf{v}_n = \psi^\dagger \alpha_n \psi / \psi^\dagger \psi$.

The above theory of fermions, with deterministic trajectories, was proposed by Bohm, Hiley, and Kaloyerou (1987). The theory was derived by Colin (2003) as the continuum limit of a discrete model of fermion numbers evolving stochastically on a lattice (Bell 1986, 1987). Dürr et al. $(2004, 2005)$ studied the same limit of Bell's model but obtained a different result: a theory of particle trajectories with stochastic jumps at points where particles are created or annihilated. The difference stems from Dürr et al.'s reading of Bell's 'fermion number', which was mistakenly taken to be the number of particles plus the number of anti-particles (instead of Colin's conventional reading as the number of particles minus the number of anti-particles).

6.3 Electromagnetic field. Scalar QED

The pilot-wave theory of the electromagnetic field is often written in the Coulomb gauge (with a divergence-free vector potential, $\nabla \cdot \mathbf{A} = 0$, and a Coulomb potential $\phi = A^0$) (Bohm 1952b; Kaloyerou 1994). However, on Aristotelian

spacetime with a preferred state of rest, it is more natural to employ a purely spatial gauge field A with no time component (Valentini 1992, 1996). This is equivalent to working in the temporal gauge $(A^0 = 0)^7$ $(A^0 = 0)^7$. Both gauges are non-Lorentz-covariant and hence free of the unphysical ghost states (some with negative norm) which appear in manifestly Lorentz-covariant quantum gauge theories (Leibbrandt 1987). In a theory with a preferred frame, ghosts are automatically avoided.

We can represent the electromagnetic field by a 3-vector field $\mathbf{A}(\mathbf{x},t)$ with classical Lagrangian $L = \int \mathcal{L} d^3x$ where

$$
\mathcal{L} = \frac{1}{2}(\dot{\mathbf{A}}^2 - \mathbf{B}^2)
$$
 (51)

 $(E = -\partial_t \mathbf{A}$ and $\mathbf{B} = \nabla \times \mathbf{A}$. The field is subject to time-independent gauge transformations

$$
\mathbf{A}(\mathbf{x},t) \to \mathbf{A}(\mathbf{x},t) - \nabla \Lambda(\mathbf{x}) . \tag{52}
$$

The wave functional $\Psi = \Psi[\mathbf{A}, t]$ is a function, not on the space of gaugedependent fields A, but on the space of equivalence classes of fields connected by gauge transformations. It then satisfies the constraint

$$
\nabla \cdot \frac{\delta \Psi}{\delta \mathbf{A}} = 0 \;, \tag{53}
$$

which follows by setting $\delta \Psi = 0$ under an arbitrary infinitesimal gauge change $\delta \mathbf{A} = -\boldsymbol{\nabla} \Lambda.$

The Schrödinger equation is

$$
i\frac{\partial \Psi}{\partial t} = \int d^3x \, \frac{1}{2} \left(-\frac{\delta^2}{\delta \mathbf{A}^2} + \mathbf{B}^2 \right) \Psi \tag{54}
$$

(which preserves [\(53\)](#page-16-1) in time). This implies a continuity equation

$$
\frac{\partial |\Psi|^2}{\partial t} + \int d^3x \; \frac{\delta}{\delta \mathbf{A}} \cdot \left(|\Psi|^2 \frac{\delta S}{\delta \mathbf{A}} \right) = 0 \tag{55}
$$

and the guidance equation

$$
\frac{\partial \mathbf{A}}{\partial t} = \frac{\delta S}{\delta \mathbf{A}} \tag{56}
$$

In quantum equilibrium, again, reference to [\(56\)](#page-16-2) may be dropped, and the field statistics obey the Born rule. We then recover standard QED in the temporal gauge. In the Heisenberg picture, [\(53\)](#page-16-1) becomes

$$
\nabla \cdot \partial_t \hat{\mathbf{A}} | \Psi \rangle = 0 \tag{57}
$$

We have equal-time commutation relations^{[8](#page-16-3)}

$$
\left[\hat{A}_i(\mathbf{x},t),\partial_t\hat{A}_j(\mathbf{x}',t)\right] = i\delta_{ij}\delta^3(\mathbf{x}-\mathbf{x}')\tag{58}
$$

⁷We can always set $A^0 = 0$ by a gauge transformation $A^0 \to A^0 + \partial \Lambda / \partial t$, $\mathbf{A} \to \mathbf{A} - \nabla \Lambda$ with $\Lambda(\mathbf{x},t) = -\int_0^t dt' \, A^0(\mathbf{x},t').$

 8 On Aristotelian spacetime the spatial indices i, j can be trivially raised or lowered with no sign change.

 $(i, j = 1, 2, 3)$ and the operator field equation

$$
\partial_t^2 \hat{\mathbf{A}} + \nabla \times (\nabla \times \hat{\mathbf{A}}) = 0.
$$
 (59)

Equations (57) – (59) define (free) QED in the temporal gauge, which is equivalent to the more usual forms of QED (in the Coulomb gauge or a Lorentzcovariant gauge) (Leibbrandt 1987; Valentini 1996).

Interactions can be introduced in the usual way via local gauge invariance. For simplicity we consider interaction with a charged scalar field (scalar QED) (Valentini 1992). The gauge transformations

$$
\phi \to \phi e^{ie\Lambda}, \quad \phi^* \to \phi^* e^{-ie\Lambda}, \quad \mathbf{A} \to \mathbf{A} - \nabla \Lambda \tag{60}
$$

(with $\Lambda = \Lambda(\mathbf{x})$ and e the charge) imply that $\Psi = \Psi[\phi, \phi^*, \mathbf{A}, t]$ is subject to the constraint

$$
\nabla \cdot \frac{\delta \Psi}{\delta \mathbf{A}} = ie \left(\phi^* \frac{\delta \Psi}{\delta \phi^*} - \phi \frac{\delta \Psi}{\delta \phi} \right) \tag{61}
$$

(again obtained by setting $\delta \Psi = 0$ under an infinitesimal gauge change). The Schrödinger equation is

$$
i\frac{\partial\Psi}{\partial t} = \int d^3 \mathbf{x} \left(-\frac{1}{2} \frac{\delta^2}{\delta \mathbf{A}^2} + \frac{1}{2} \mathbf{B}^2 - \frac{\delta^2}{\delta \phi^* \delta \phi} + m^2 \phi^* \phi + (\mathbf{D}\phi)^* \cdot (\mathbf{D}\phi) \right) \Psi , \tag{62}
$$

where $\mathbf{D}\phi = \nabla \phi + ie\mathbf{A}\phi$ is the gauge-covariant derivative. From the associated continuity equation we find the guidance equations

$$
\frac{\partial \phi}{\partial t} = \frac{\delta S}{\delta \phi^*}, \quad \frac{\partial \phi^*}{\partial t} = \frac{\delta S}{\delta \phi}, \quad \frac{\partial \mathbf{A}}{\partial t} = \frac{\delta S}{\delta \mathbf{A}}.
$$

In quantum equilibrium we recover standard (scalar) QED in the temporal gauge.

To describe interactions with electrons and positrons we can introduce a gauge coupling of \bf{A} to a spin-1/2 complex Grassmann field or (in a particle description of fermions) to the many-body Dirac equation. Again, in quantum equilibrium, we necessarily recover standard QED in the temporal gauge.

6.4 Non-Abelian gauge theories

In the standard $SU(3)\times SU(2)\times U(1)$ model of particle physics, interactions are usually mediated by 4-vector gauge fields on Minkowski spacetime, as defined in a manifestly Lorentz-covariant quantum field theory. In pilot-wave theory it is more natural to employ 3-vector gauge fields on Aristotelian spacetime (Valentini 1992, 1996). In quantum equilibrium we recover the usual theories of electroweak and strong interactions written in the ghost-free temporal gauge (with all time components set to zero).

6.4.1 QCD

In quantum chromodynamics we have 3-vector gluon fields $\mathbf{A}^{a}(\mathbf{x},t)$ $(a = 1,...,8)$ with a classical Lagrangian $L = \int \mathcal{L} d^3x$ where^{[9](#page-18-0)}

$$
\mathcal{L} = \frac{1}{2} (\dot{A}_i^a)^2 - \frac{1}{4} (F_{ij}^a)^2 \,, \tag{63}
$$

with

$$
F_{ij}^a = -\partial_i A_j^a + \partial_j A_i^a + gf^{abc} A_i^b A_j^c , \qquad (64)
$$

where the fields are subject to time-independent (infinitesimal) SU(3) gauge transformations

$$
\mathbf{A}^a \to \mathbf{A}^a - f^{abc} \Lambda^b \mathbf{A}^c - \frac{1}{g} \nabla \Lambda^a \ . \tag{65}
$$

Here Λ^a are arbitrary (infinitesimal) functions on space, f^{abc} are the structure constants, and g is the gauge coupling. For free gluon fields the wave functional $\Psi = \Psi[\mathbf{A}^a, t]$ is then constrained by

$$
\nabla \cdot \frac{\delta \Psi}{\delta \mathbf{A}^a} = gf^{abc} \mathbf{A}^b \cdot \frac{\delta \Psi}{\delta \mathbf{A}^c}
$$
 (66)

(setting $\delta \Psi = 0$ under transformations [\(65\)](#page-18-1)).

The Schrödinger equation is

$$
i\frac{\partial\Psi}{\partial t} = \int d^3x \, \frac{1}{2} \left(-\frac{\delta^2}{(\delta A_i^a)^2} + \frac{1}{2} (F_{ij}^a)^2 \right) \Psi \,. \tag{67}
$$

The associated continuity equation for $|\Psi|^2$ implies the de Broglie velocities

$$
\frac{\partial \mathbf{A}^a}{\partial t} = \frac{\delta S}{\delta \mathbf{A}^a} \,. \tag{68}
$$

To describe interactions with quarks we can introduce an SU(3) gauge coupling to colour triplets of spin-1/2 Grassmann fields or (in a particle description of fermions) to colour triplets of the many-body Dirac equation. Again, in quantum equilibrium, we recover standard QCD in the temporal gauge.

6.4.2 Electroweak Theory. Spontaneous Symmetry Breaking

A pilot-wave formulation of $SU(2) \times U(1)$ electroweak theory is also readily obtained from the standard theory in the temporal gauge. We now have four massless 3-vector gauge fields W_i (i = 1, 2, 3) and **B** (not to be confused with the magnetic field). These are related to the W-boson and Z^0 -boson fields W, W^{\dagger}, Z , and to the electromagnetic field A , by the usual relations

$$
\mathbf{W} = \frac{1}{\sqrt{2}} (\mathbf{W}_1 - i\mathbf{W}_2) , \qquad \mathbf{W}^{\dagger} = \frac{1}{\sqrt{2}} (\mathbf{W}_1 + i\mathbf{W}_2) , \qquad (69)
$$

⁹Repeated indices are summed.

$$
\mathbf{Z} = \mathbf{W}_3 \cos \theta_W - \mathbf{B} \sin \theta_W , \qquad \mathbf{A} = \mathbf{W}_3 \sin \theta_W + \mathbf{B} \cos \theta_W , \qquad (70)
$$

where θ_W is the weak mixing angle. The fields W, W^{\dagger} , Z become effectively massive after spontaneous symmetry breaking. In quantum equilibrium we necessarily recover standard electroweak theory in the temporal gauge, with time components $W^0 = (W^{\dagger})^0 = Z^0 = A^0 = 0.$

The pilot-wave account of spontaneous symmetry breaking is, again, as in the standard theory written in the temporal gauge. For example, the Abelian Higgs model is just scalar QED with the Hamiltonian in [\(62\)](#page-17-1) supplemented by a self-interaction term $\lambda |\phi|^4$. As usually understood, in appropriate conditions the system has a continuous set of stable ground states with non-zero (constant) expectation values ϕ_0 for the field ϕ , where an arbitrary choice of ground state effectively breaks the gauge symmetry. Expanding ϕ around ϕ_0 , the term $(ie \mathbf{A}\phi)^* \cdot (ie \mathbf{A}\phi)$ in the original Hamiltonian yields an effective mass for the field **A**, arising from the coupling between **A** and the constant field ϕ_0 . The effective particle content – massive neutral vector bosons and massive neutral scalar bosons – is usually shown in the unitary gauge. But the Higgs model in the temporal gauge is equivalent to the model in the unitary gauge (Kim et al. 1990). In quantum equilibrium the pilot-wave formulation, with 3-vector gauge fields, is then necessarily equivalent to the standard theory.

7 The early universe

Quantum field relaxation presumably took place in the early universe. To describe this we need to generalise pilot-wave field theory to a background curved spacetime. On expanding space we find that quantum relaxation can be suppressed at long wavelengths, so that early violations of the Born rule may be imprinted on the cosmic microwave background (CMB) at large scales.

7.1 Quantum fields in curved spacetime

Pilot-wave field theory is readily generalised to a classical curved spacetime background (Valentini 2004).

We assume spacetime to be globally hyperbolic, so it can be foliated (nonuniquely) by spacelike hypersurfaces $\Sigma(t)$ with a global time t. The spacetime line element can then be written as

$$
d\tau^2 = (N^2 - N_i N^i)dt^2 - 2N_i dx^i dt - g_{ij} dx^i dx^j , \qquad (71)
$$

with a lapse function N and shift vector N^i , where g_{ij} is the 3-metric on $\Sigma(t)$. We may set $N^i = 0$ (provided lines $x^i = \text{const.}$ do not meet singularities).

For a massless, minimally-coupled real scalar field ϕ , we have a classical Lagrangian $L = \int \mathcal{L}d^3x$ with

$$
\mathcal{L} = \frac{1}{2} N \sqrt{g} \left(\frac{1}{N^2} \dot{\phi}^2 - g^{ij} \partial_i \phi \partial_j \phi \right) , \qquad (72)
$$

where $g = \det g_{ij}$. The wave functional $\Psi[\phi, t]$ satisfies the Schrödinger equation

$$
i\frac{\partial\Psi}{\partial t} = \int d^3x \; \frac{1}{2}N\sqrt{g} \left(-\frac{1}{g}\frac{\delta^2}{\delta\phi^2} + g^{ij}\partial_i\phi\partial_j\phi \right) \Psi \; . \tag{73}
$$

The associated continuity equation for $|\Psi|^2$ implies the de Broglie velocity

$$
\frac{\partial \phi}{\partial t} = \frac{N}{\sqrt{g}} \frac{\delta S}{\delta \phi} \tag{74}
$$

An arbitrary distribution $P[\phi, t]$ satisfies the same continuity equation,

$$
\frac{\partial P}{\partial t} + \int d^3x \, \frac{\delta}{\delta \phi} \left(P \frac{N}{\sqrt{g}} \frac{\delta S}{\delta \phi} \right) = 0 \; . \tag{75}
$$

As usual, $P = |\Psi|^2$ at some initial time implies $P = |\Psi|^2$ at all times.

If Ψ is entangled across the hypersurface $\Sigma(t)$, the time evolution of ϕ at a spatial point x^i can depend instantaneously (with respect to t) on ϕ at remote points $(x')^i \neq x^i$. For nonequilibrium ensembles, there will be statistical nonlocal signals across $\Sigma(t)$. The theory is physically consistent if we assume there is a preferred foliation (Valentini 2008).

7.2 Suppression of quantum relaxation on expanding space

Quantum relaxation in the early universe has been studied for a massless field ϕ on flat expanding space, with spacetime metric

$$
d\tau^2 = dt^2 - a^2 \delta_{ij} dx^i dx^j \tag{76}
$$

and with a scale factor $a(t) \propto t^{1/2}$ (a radiation-dominated expansion) (Colin and Valentini 2013, 2015). In Fourier space, for an unentangled field mode (q_{k1}, q_{k2}) , we again obtain Schrödinger and de Broglie equations of the form (39) and (40) for a two-dimensional oscillator but with mass $m = a³$ and angular frequency $\omega = k/a$ (Valentini 2007). This system is equivalent to a standard oscillator, of constant mass and angular frequency, with t replaced by a 'retarded time' $t_{\text{ret}} = t_{\text{ret}}(t, k)$ (Colin and Valentini 2013). Quantum relaxation depends on the relation between the physical wavelength $\lambda_{\text{phys}} = a(2\pi/k)$ and the Hubble radius $H^{-1} = a/a$. For $\lambda_{\text{phys}} \ll H^{-1}$ we find $t_{\text{ret}}(t, k) \to t$ and the usual rapid relaxation (Fig. 1). For $\lambda_{\text{phys}} \gtrsim H^{-1}$ we find $t_{\text{ret}}(t, k) < t$ and relaxation is suppressed.

The suppression of quantum relaxation can be quantified by the mean square

$$
\langle |\phi_{\mathbf{k}}|^2 \rangle = \langle |\phi_{\mathbf{k}}|^2 \rangle_{\text{QT}} \xi(k) , \qquad (77)
$$

where $\langle \ldots \rangle$ and $\langle \ldots \rangle_{\mathrm{QT}}$ are respectively nonequilibrium and equilibrium expectation values and $\xi(k) < 1$. Numerical simulations show that $\xi(k)$ tends to decrease as k decreases (ignoring small oscillations), with a good fit to the curve

$$
\xi(k) = \tan^{-1}(c_1 \frac{k}{\pi} + c_2) - \frac{\pi}{2} + c_3 , \qquad (78)
$$

where the parameters c_1 , c_2 , c_3 depend on the initial state and on the time interval (Colin and Valentini 2015, 2016).

In inflationary cosmology, during a very early approximately exponential expansion, perturbations $\phi_{\mathbf{k}}$ of the inflaton field generate primordial curvature perturbations $\mathcal{R}_{\mathbf{k}} \propto \phi_{\mathbf{k}}$, which later generate small temperature anisotropies in the CMB (Liddle and Lyth 2000). By measuring statistical properties of the CMB we can constrain the primordial power spectrum

$$
\mathcal{P}_{\mathcal{R}}(k) = \frac{4\pi k^3}{V} \left\langle |\mathcal{R}_{\mathbf{k}}|^2 \right\rangle \propto \frac{4\pi k^3}{V} \left\langle |\phi_{\mathbf{k}}|^2 \right\rangle \ . \tag{79}
$$

Thus from measurements of the CMB we can set observational bounds on corrections to the Born rule in the very early universe (Valentini 2010).

Assuming that quantum relaxation occurred during a pre-inflationary era, and making the simplifying assumption that the spectrum is unaffected by the transition from pre-inflation to inflation, the usual quantum spectrum $\mathcal{P}^{\mathcal{Q}T}_{\mathcal{R}}(k)$ is replaced by

$$
\mathcal{P}_{\mathcal{R}}(k) = \mathcal{P}_{\mathcal{R}}^{\text{QT}}(k)\xi(k) , \qquad (80)
$$

with a power deficit at small k of the form [\(78\)](#page-20-0). CMB data show hints of a power deficit at small k (Aghanim et al. 2016). However, current data neither support nor rule out the prediction [\(78\)](#page-20-0) (Vitenti, Peter, and Valentini 2019). Further predictions, such as nonequilibrium violations of statistical isotropy, are needed to obtain sufficient evidence for or against the model.

According to inflationary cosmology, the matter in our universe was created at early times by inflaton decay. If the early inflaton field violates the Born rule, so will its decay products (Underwood and Valentini 2015). Such early nonequilibrium particles may have survived to the present day, possibly as a component of dark matter. If those particles decay or annihilate, the resulting spectral lines could be anomalous (Underwood and Valentini 2020). The decay or annihilation products (probably photons) may show violations of the Born rule in simple quantum experiments.

8 Quantum gravity and quantum cosmology

The pilot-wave theory of gravitation, with de Broglie-Bohm trajectories associated with the time-independent Wheeler-DeWitt equation, was first discussed for quantum cosmology by Vink (1992) and then developed for quantum gravity by Horiguchi (1994) .^{[10](#page-21-0)} It has since been extensively applied to cosmology (Pinto-Neto 2005; Pinto-Neto and Fabris 2013; Pinto-Neto 2021).

8.1 Pilot-wave theory and the Wheeler-DeWitt equation

Beginning with the classical Einstein-Hilbert action, canonical quantum gravity applies the foliation [\(71\)](#page-19-0) of spacetime and quantises the spatial 3-geometry with

 10 An alternative approach attempts to associate trajectories with a time-dependent gravitational Schrödinger equation (Valentini 1992, 1996; Roser and Valentini 2014).

metric g_{ij} . In the presence of a scalar matter field ϕ with potential $\mathcal{V}(\phi)$, this yields the Wheeler-DeWitt equation (units $\hbar = c = 16\pi G = 1$) (Kiefer 2012)

$$
\left(-G_{ijkl}\frac{\delta^2}{\delta g_{ij}\delta g_{kl}} - \sqrt{g}R + \hat{\mathcal{H}}_{\phi}\right)\Psi = 0
$$
\n(81)

for the wave functional $\Psi = \Psi[g_{ij}, \phi]$, where

$$
G_{ijkl} = \frac{1}{2}g^{-1/2}(g_{ik}g_{jl} + g_{il}g_{jk} - g_{ij}g_{kl})
$$
\n(82)

and

$$
\hat{\mathcal{H}}_{\phi} = \frac{1}{2} \sqrt{g} \left(-\frac{1}{g} \frac{\delta^2}{\delta \phi^2} + g^{ij} \partial_i \phi \partial_j \phi \right) + \sqrt{g} \mathcal{V} . \tag{83}
$$

The ordering in the kinetic term ' $G\delta^2/\delta g^2$ ' is ambiguous. We also have a constraint

$$
-2D_j \frac{\delta \Psi}{\delta g_{ij}} + \partial^i \phi \frac{\delta \Psi}{\delta \phi} = 0
$$
\n(84)

(where D_i is a spatial covariant derivative), which follows by setting $\delta \Psi = 0$ under arbitrary infinitesimal spatial diffeomorphisms. This ensures that Ψ is a function of the coordinate-independent 3-geometry and not of the coordinatedependent 3-metric.

Writing [\(81\)](#page-22-0) with the operator ordering ' $(\delta/\delta g)G(\delta/\delta g)$ ', inserting $\Psi =$ $|\Psi| e^{iS}$ and taking the imaginary part, we obtain

$$
\frac{\delta}{\delta g_{ij}} \left(|\Psi|^2 2NG_{ijkl} \frac{\delta S}{\delta g_{kl}} \right) + \frac{\delta}{\delta \phi} \left(|\Psi|^2 \frac{N}{\sqrt{g}} \frac{\delta S}{\delta \phi} \right) = 0 \tag{85}
$$

This is not a continuity equation but an infinity of equations (one per space point x^i). Even so we may take it to define a natural velocity field in configuration space, and so identify the de Broglie guidance equations

$$
\frac{\partial g_{ij}}{\partial t} = 2NG_{ijkl}\frac{\delta S}{\delta g_{kl}}\ ,\quad \frac{\partial \phi}{\partial t} = \frac{N}{\sqrt{g}}\frac{\delta S}{\delta \phi}\ .\tag{86}
$$

Alternatively, these can be derived by setting the classical canonical momenta equal to the phase gradient.^{[11](#page-22-1)} Equations [\(86\)](#page-22-2) and [\(81\)](#page-22-0), with the constraint [\(84\)](#page-22-3), are taken to define the dynamics of an individual system.

8.2 Dynamical consistency. Preferred foliation

The consistency of the above dynamics has been questioned. For given initial conditions on a spacelike slice, the 4-geometry traced out by the evolving 3-metric should not depend on the arbitrary lapse and shift functions N, N^i . Otherwise the initial-value problem would not be well posed. Shtanov (1996)

¹¹For $N_i \neq 0$ the right-hand sides of [\(86\)](#page-22-2) have additional terms $D_iN_j + D_jN_i$ and $N^i\partial_i\phi$ respectively.

argued that the resulting 4-geometry depends on N and that the theory breaks foliation invariance. We might then make a case for including a specific choice for N as part of the theory. In contrast, work by Pinto-Neto and Santini (2002) appears to demonstrate that the 4-geometry is in fact independent of N, N^i . By writing the dynamics as a classical Hamiltonian system (with an additional 'quantum potential density'), and applying the analogue of a well-known theorem from the Hamiltonian dynamics of classical gravitation, Pinto-Neto and Santini argue that the evolving 3-metric yields the same 4-geometry for all choices of N, N^i – a 4-geometry that is, however, non-Lorentzian. The argument seems to imply that the resulting spacetime has an effective preferred foliation (for a given wave functional Ψ , and for a given initial 3-geometry). Local Lorentz invariance is broken, as we would expect in a nonlocal theory.

These results seem consistent with the first-order (Aristotelian) structure of pilot-wave dynamics, which naturally defines a preferred state of rest (Valentini 1997). Furthermore, nonequilibrium entangled systems generate statistical nonlocal signals (Section 5.1), which may be employed to define an absolute simultaneity (Valentini 2008). Thus, general features of pilot-wave theory already point to an underlying preferred foliation of spacetime.

8.3 Pilot-wave quantum cosmology

As a simple example consider a flat expanding universe, with line element [\(76\)](#page-20-1), containing a homogenous matter field ϕ with potential $\mathcal{V}(\phi)$. With appropriate factor ordering, the Wheeler-DeWitt equation for $\Psi(a, \phi)$ reads (Brizuela, Kiefer, and Krämer 2016)

$$
\frac{1}{2m_{\rm P}^2} \frac{1}{a} \frac{\partial}{\partial a} \left(a \frac{\partial \Psi}{\partial a} \right) - \frac{1}{2a^2} \frac{\partial^2 \Psi}{\partial \phi^2} + a^4 \mathcal{V} \Psi = 0 \;, \tag{87}
$$

where $m_P^2 = 3/4\pi G$ is the (rescaled) Planck mass squared. Inserting $\Psi = |\Psi| e^{iS}$ and taking the imaginary part, we find a continuity equation

$$
-\frac{1}{m_{\rm P}^2} \frac{\partial}{\partial a} \left(a^2 \left| \Psi \right|^2 \frac{1}{a} \frac{\partial S}{\partial a} \right) + \frac{\partial}{\partial \phi} \left(a^2 \left| \Psi \right|^2 \frac{1}{a^3} \frac{\partial S}{\partial \phi} \right) = 0 \tag{88}
$$

for a static density $a^2 |\Psi|^2$ with a natural velocity field

$$
\dot{a} = -\frac{1}{m_{\rm P}^2} \frac{1}{a} \frac{\partial S}{\partial a} \ , \quad \dot{\phi} = \frac{1}{a^3} \frac{\partial S}{\partial \phi} \ . \tag{89}
$$

We can identify [\(89\)](#page-23-0) as the de Broglie guidance equations. Again, the same equations follow if we set the classical canonical momenta equal to the phase gradient.

De Broglie-Bohm quantum cosmology has been extensively applied to bouncing cosmological models (alternatives to inflation with a contracting era followed by an expanding phase) and to the theory of cosmological perturbations. Other applications include demonstrations of singularity avoidance, explanations for cosmic acceleration, and a clear treatment of the quantum-to-classical transition in the early universe. For reviews see Pinto-Neto (2005), Pinto-Neto and Fabris (2013), and Pinto-Neto (2021).

8.4 Problem of probability. Beyond the Born rule

Most work in pilot-wave quantum cosmology is focussed on properties of trajectories, without developing a theory of ensembles or probabilities. This is because Ψ is non-normalisable and cannot define a Born-rule density. This is a general feature of solutions to the Wheeler-DeWitt equation, whose math-ematical structure is analogous to a Klein-Gordon equation.^{[12](#page-24-0)} The integral $\int dq |\Psi|^2$ over configuration space is comparable to the integral $\int d^3x \int dt |\phi|^2$ for a Klein-Gordon field ϕ over spacetime and necessarily diverges (even after factoring out coordinate dependence) (Isham 1993; Kuchař 2011). Even so, we can make sense of probability in pilot-wave quantum gravity (Valentini 2021, 2023).

First, we can define a theoretical ensemble with a normalised probability density P , even though P can never be equal to the (non-normalised) Born-rule density $|\Psi|^2$. For example, in our model of quantum cosmology, we can write a general continuity equation

$$
\frac{\partial P}{\partial t} + \frac{\partial}{\partial a} (P\dot{a}) + \frac{\partial}{\partial \phi} (P\dot{\phi}) = 0 , \qquad (90)
$$

with de Broglie velocities [\(89\)](#page-23-0). This can be applied to a general theoretical ensemble with wave function Ψ . Because P is by construction normalised, it can never relax to $|\Psi|^2$ (or to $a^2 |\Psi|^2$ in our model). Thus the deep quantumgravity regime is always in a state of quantum nonequilibrium.

Second, the Born rule can emerge in a semiclassical regime, with an effective time-dependent Schrödinger equation $i\partial \psi/\partial t = \hat{H}\psi$ for a normalisable wave functional $\psi[\phi, t]$, where ϕ represents quantum fields propagating on a classical spacetime background and \hat{H} is the effective Hamiltonian. This approximation arises with a Wheeler-DeWitt wave functional $\Psi[g_{ij}, \phi] \approx \Psi_{WKB}[g_{ij}]\psi[\phi, g_{ij}],$ where $\Psi_{WKB}[g_{ij}]$ is a WKB state and the effective $\psi[\phi, t]$ is just $\psi[\phi, g_{ij}(t)]$ evaluated along a classical background trajectory $g_{ij}(t)$. In this regime we find the usual (coarse-grained) relaxation $\rho \rightarrow |\psi|^2$ to the Born rule.

The effective Hamiltonian \hat{H} has tiny quantum-gravitational corrections, some of which render the Born rule unstable (Valentini 2021, 2023). The cor-rections are derived from [\(81\)](#page-22-0) by a semiclassical expansion of Ψ (Kiefer and Singh 1991; Brizuela, Kiefer, and Krämer 2016). There are both Hermitian and non-Hermitian terms.^{[13](#page-24-1)} Applying the same expansion to the de Broglie guid-ance equation [\(86\)](#page-22-2) for ϕ , the field velocity takes the same form as before but

¹²Owing to the indefinite 'DeWitt metric' G_{ijkl} .

¹³There is no consensus as to whether the non-Hermitian terms are an artifact or a real physical effect. Some authors advocate eliminating them by appropriate redefinitions of the wave function (Kiefer and Wichmann 2018).

with S equal to the phase of ψ . Writing $\hat{H} = \hat{H}_1 + i\hat{H}_2$ (with \hat{H}_1 , \hat{H}_2 Hermi-tian), we then have the usual continuity equation [\(4\)](#page-1-3) for ρ (cf. [\(75\)](#page-20-2)) while the continuity equation [\(5\)](#page-2-3) for $|\psi|^2$ now has a source term $s = 2 \operatorname{Re} \left(\psi^* \hat{H}_2 \psi \right)$ on the right-hand side. The continuity equations for ρ and $|\psi|^2$ no longer match. Initial equilibrium $\rho = |\psi|^2$ can therefore evolve into final nonequilibrium $\rho \neq |\psi|^2$ on an estimated timescale

$$
\tau_{\text{noneq}} \approx \frac{1}{2 \left| \left\langle \hat{H}_2 \right\rangle \right|} \,. \tag{91}
$$

For a scalar field in the vicinity of an evaporating Schwarzchild black hole of mass $M(t)$, the Hamiltonian $\hat{H}_{\mathbf{k}}$ of a field mode has a non-Hermitian correction $i\hat{H}_2$ with (Kiefer, Müller, and Singh 1994)

$$
\hat{H}_2 \simeq -\frac{1}{12} \kappa \left(\frac{m_\text{P}}{M}\right)^4 \hat{H}_\mathbf{k} , \qquad (92)
$$

where κ is a numerical factor and here $m_{\rm P} = \sqrt{\hbar c/G} \simeq 10^{-5}$ g is the standard Planck mass. We then find (Valentini 2021, 2023)

$$
\tau_{\text{noneq}} \sim \frac{48\pi}{\kappa} t_{\text{P}} \left(\frac{M}{m_{\text{P}}}\right)^5 \tag{93}
$$

(where t_P is the Planck time). The effect is significant when M approaches m_P . The final burst of Hawking radiation could violate the Born rule.[14](#page-25-0) In principle such violations might be observed in radiation from exploding primordial black holes.

9 Conclusion

The de Broglie-Bohm pilot-wave formulation of quantum mechanics avoids the notorious quantum measurement problem while highlighting the nonlocality of quantum physics. Quantum 'measurements' are understood to be physical processes like any other, obeying the same underlying physical laws. There is no need for a fundamental division between 'system' and 'apparatus', and physical processes may be described in objective terms even in the absence of 'observers'. Pilot-wave theory may be applied across the full range of known physics, including high-energy physics, quantum field theory, gravitation, and cosmology. It has proved to be especially useful in cosmology, an area where textbook quantum mechanics is difficult to apply since there is no outside observer.

The internal logic of pilot-wave theory changes our perspective on the Born rule, which can no longer be regarded as a law of physics. Instead it corresponds to a state of statistical equilibrium, broadly analogous to thermal equilibrium, which may be understood as having arisen by dynamical relaxation. Quantum

 $14A$ breakdown of the Born rule in Hawking radiation has also been suggested as a possible resolution of the information-loss puzzle (Valentini 2004, 2007; Kandhadai and Valentini 2020).

physics is seen to be merely a special case of a much wider physics that includes nonlocal signalling, violations of the uncertainty principle, and the breaking of other standard quantum constraints. Thus pilot-wave theory provides a natural extension of quantum physics to a new nonequilibrium regime.

Evidence for quantum nonequilibrium physics could be found in the early universe, though this is still an active area of research with as yet no clear conclusions. Relic nonequilibrium systems from early times might still exist today. Exotic gravitational effects could potentially create nonequilibrium where there was none before. Should nonequilibrium systems be discovered and put to practical use, there would be radical technological implications, including for cryptography and computing.

Pilot-wave theory is a distinctive approach to the theory of motion and to physics generally. It provides a novel perspective on the structure of spacetime, as well as on the foundations of quantum gravity and the nature of black holes. While much remains to be understood, this formulation of quantum mechanics continues to provide insights into the nature of quantum physics and to suggest new lines of research that would not be available to other formulations.

Bibliography

Abraham, E., Colin, S., and Valentini, A. (2014). Long-time relaxation in pilot-wave theory. *Journal of Physics A* 47, 395306.

Abrams, D. S. and Lloyd, S. (1998). Nonlinear quantum mechanics implies polynomial-time solution for NP-complete and #P problems. *Physical Review Letters* 81, 3992–3995.

Aghanim, N. *et al*. (Planck Collaboration) (2016). *Planck* 2015 results. XI. CMB power spectra, likelihoods, and robustness of parameters. *Astronomy and Astrophysics* 594, A11.

Bacciagaluppi, G. (2003). Derivation of the symmetry postulates for identical particles from pilot-wave theories. [arXiv:quant-ph/0302099.](http://arxiv.org/abs/quant-ph/0302099)

Bacciagaluppi, G. and Valentini, A. (2009). *Quantum theory at the crossroads: Reconsidering the 1927 Solvay conference*. Cambridge: Cambridge University Press.

Bell, J. S. (1966). On the problem of hidden variables in quantum mechanics. *Reviews of Modern Physics* 38, 447–452.

Bell, J. S. (1986). Quantum field theory without observers. *Physics Reports* 137, 49–54.

Bell, J. S. (1987). *Speakable and unspeakable in quantum mechanics*. Cambridge: Cambridge University Press.

Bennett, C. H. (1992). Quantum cryptography using any two nonorthogonal states. *Physical Review Letters* 68, 3121–3124.

Berndl, K., Dürr, D., Goldstein, S., and Zanghì, N. (1996). Nonlocality, Lorentz invariance, and Bohmian quantum theory. *Physical Review A* 53, 2062– 2073.

Bohm, D. (1952a). A suggested interpretation of the quantum theory in terms of 'hidden' variables. I. *Physical Review* 85, 166–179.

Bohm, D. (1952b). A suggested interpretation of the quantum theory in terms of 'hidden' variables. II. *Physical Review* 85, 180–193.

Bohm, D., Hiley, B. J., and Kaloyerou, P. N. (1987). An ontological basis for the quantum theory. *Physics Reports* 144, 321–375.

Bohm, D. and Hiley, B. J. (1993). *The undivided universe: an ontological interpretation of quantum theory*. London: Routledge.

Brizuela, D., Kiefer, C., and Krämer, M. (2016). Quantum-gravitational effects on gauge-invariant scalar and tensor perturbations during inflation: the de Sitter case. *Physical Review D* 93, 104035.

Brown, L. M. (1958). Two-component fermion theory. *Physical Review* 111, 957–964.

Colin, S. (2003). A deterministic Bell model. *Physics Letters A* 317, 349– 358.

Colin, S. (2012). Relaxation to quantum equilibrium for Dirac fermions in the de Broglie-Bohm pilot-wave theory. *Proceedings of the Royal Society A* 468, 1116–1135.

Colin, S. and Struyve, W. (2007). A Dirac sea pilot-wave model for quantum field theory. *Journal of Physics A* 40, 7309–7341.

Colin, S. and Valentini, A. (2013). Mechanism for the suppression of quantum noise at large scales on expanding space. *Physical Review D* 88, 103515.

Colin, S. and Valentini, A. (2014). Instability of quantum equilibrium in Bohm's dynamics. *Proceedings of the Royal Society A* 470, 20140288.

Colin, S. and Valentini, A. (2015). Primordial quantum nonequilibrium and large-scale cosmic anomalies. *Physical Review D* 92, 043520.

Colin, S. and Valentini, A. (2016). Robust predictions for the large-scale cosmological power deficit from primordial quantum nonequilibrium. *International Journal of Modern Physics D* 25, 1650068.

de Broglie, L. (1928). La nouvelle dynamique des quanta. In *Electrons et ´* photons: Rapports et discussions du cinquième conseil de physique, pp 105–132. Paris: Gauthier-Villars. [English translation: Bacciagaluppi, G. and Valentini, A. (2009).]

Drezet, A. (2021). Justifying Born's rule $P_{\alpha} = |\Psi_{\alpha}|^2$ using deterministic chaos, decoherence, and the de Broglie-Bohm quantum theory. *Entropy* 23, 1371.

Dürr, D., Goldstein, S., and Zanghì, N. (1992). Quantum equilibrium and the origin of absolute uncertainty. *Journal of Statistical Physics* 67, 843–907.

Dürr, D., Goldstein, S., Münch-Berndl, K., and Zanghi, N. (1999). Hypersurface Bohm-Dirac models. *Physical Review A* 60, 2729–2736.

Dürr, D., Goldstein, S., Tumulka, R., and Zanghì, N. (2004). Bohmian mechanics and quantum field theory. *Physical Review Letters* 93, 090402.

Dürr, D., Goldstein, S., Tumulka, R., and Zanghì, N. (2005). Bell-type quantum field theories. *Journal of Physics A* 38, R1–R43.

Dürr, D. and Teufel, S. (2009). *Bohmian mechanics: the physics and mathematics of quantum theory*. Heidelberg: Springer-Verlag.

Dürr, D., Goldstein, S., Norsen, T., Struyve, W., and Zanghì, N. (2014). Can Bohmian mechanics be made relativistic? *Proceedings of the Royal Society A* 470, 20130699.

Efthymiopoulos, C., Contopoulos, G., and Tzemos, A. C. (2017). Chaos in de Broglie-Bohm quantum mechanics and the dynamics of quantum relaxation. *Annales de la Fondation Louis de Broglie* 42, 133–159.

Ekert, A. K. (1991). Quantum cryptography based on Bell's theorem. *Physical Review Letters* 67, 661–663.

Hardy, L. (1992). Quantum mechanics, local realistic theories, and Lorentzinvariant realistic theories. *Physical Review Letters* 68, 2981–2984.

Holland, P. R. (1993). *The quantum theory of motion: an account of the de Broglie-Bohm causal interpretation of quantum mechanics*. Cambridge: Cambridge University Press.

Hoˇrava, P. (2009). Quantum gravity at a Lifshitz point. *Physical Review D* 79, 084008.

Horiguchi, T. (1994). Quantum potential interpretation of the Wheeler– DeWitt equation. *Modern Physics Letters A* 9, 1429–1443.

Isham, C. J. (1993). Canonical quantum gravity and the problem of time. In Ibort, L. A. & Rodriguez, M. A. (eds.) *Integrable systems, quantum groups, and quantum field theories*, pp 157–288. London: Kluwer.

Kaloyerou, P. N. (1994). The causal interpretation of the electromagnetic field. *Physics Reports* 244, 287–358.

Kandhadai, A. and Valentini, A. (2019). Perturbations and quantum relaxation. Foundations of Physics 49, 1–23.

Kandhadai, A. and Valentini, A. (2020). Mechanism for nonlocal information flow from black holes. *International Journal of Modern Physics A* 35, 2050031.

Kiefer, C. (2012). *Quantum gravity*. Oxford: Oxford University Press.

Kiefer, C., Müller, R., and Singh, T. P. (1994). Quantum gravity and nonunitarity in black hole evaporation. *Modern Physics Letters A* 9, 2661–2670.

Kiefer, C. and Singh, T. P. (1991). Quantum gravitational corrections to the functional Schrödinger equation. *Physical Review D* 44, 1067–1076.

Kiefer, C. and Wichmann, D. (2018). Semiclassical approximation of the Wheeler-DeWitt equation: arbitrary orders and the question of unitarity. *General Relativity and Gravitation* 50, 66.

Kim, S. K., Namgung, W., Soh, K. S., and Yee, J. H. (1990). Equivalence between the Weyl, Coulomb, and unitary gauges in the functional Schrödinger picture. *Physical Review D* 41, 3792–3795.

Kosteleck´y, V. A. and Mewes, M. (2002). Signals for Lorentz violation in electrodynamics. *Physical Review D* 66, 056005.

Kuchař, K. V. (2011). Time and interpretations of quantum gravity. *International Journal of Modern Physics D* 20, 3–86.

Lazarovici, D. and Hubert, M. (2019). How quantum mechanics can consistently describe the use of itself. *Scientific Reports* 9, 470.

Leibbrandt, G. (1987). Introduction to noncovariant gauges. *Reviews of Modern Physics* 59, 1067–1119.

Liddle, A. R. and Lyth, D. H. (2000). *Cosmological inflation and large-scale structure*. Cambridge: Cambridge University Press.

Lustosa, F. B., Colin, S., and Perez Bergliaffa, S. E. (2021). Quantum relaxation in a system of harmonic oscillators with time-dependent coupling. *Proceedings of the Royal Society A* 477, 20200606.

Lustosa, F. B., Pinto-Neto, N., and Valentini, A. (2023). Evolution of quantum non-equilibrium for coupled harmonic oscillators. *Proceedings of the Royal Society A* 479, 20220411.

Nielsen, M. A. and Chuang, I. L. (2000). *Quantum computation and quantum information*. Cambridge: Cambridge University Press.

Pinto-Neto, N. (2005). The Bohm interpretation of quantum cosmology. *Foundations of Physics* 35, 577–603.

Pinto-Neto, N. (2021). The de Broglie-Bohm quantum theory and its application to quantum cosmology. *Universe* 7, 134.

Pinto-Neto, N. and Fabris, J. C. (2013). Quantum cosmology from the de Broglie–Bohm perspective. *Classical and Quantum Gravity* 30, 143001.

Pinto-Neto, N. and Sergio Santini, E. (2002). The consistency of causal quantum geometrodynamics and quantum field theory. *General Relativity and Gravitation* 34, 505–532.

Roser, P. and Valentini, A. (2014). Classical and quantum cosmology with York time. *Classical and Quantum Gravity* 31, 245001.

Sakurai, J. J. (1967). *Advanced quantum mechanics*. Reading, Massachusetts: Addison-Wesley.

Sebens, C. T. (2016). Constructing and constraining wave functions for identical quantum particles. *Studies in History and Philosophy of Modern Physics* 56, 48–59.

Shtanov, Yu. V. (1996). Pilot wave quantum cosmology. *Physical Review D* 54, 2564–2570.

Struyve, W. and Valentini, A. (2009). De Broglie-Bohm guidance equations for arbitrary Hamiltonians. *Journal of Physics A* 42, 035301.

Towler, M. D., Russell, N. J., and Valentini, A. (2012). Time scales for dynamical relaxation to the Born rule. *Proceedings of the Royal Society A* 468, 990–1013.

Tumulka, R. (2007). The 'unromantic pictures' of quantum theory. *Journal of Physics A* 40, 3245–3273.

Underwood, N. G. (2018). Extreme quantum nonequilibrium, nodes, vorticity, drift and relaxation retarding states. *Journal of Physics A* 51, 055301.

Underwood, N. G. and Valentini, A. (2015). Quantum field theory of relic nonequilibrium systems. *Physical Review D* 92, 063531.

Underwood, N. G. and Valentini, A. (2020). Anomalous spectral lines and relic quantum nonequilibrium. *Physical Review D* 101, 043004.

Valentini, A. (1991a). Signal-locality, uncertainty, and the subquantum Htheorem. I. *Physics Letters A* 156, 5–11.

Valentini, A. (1991b). Signal-locality, uncertainty, and the subquantum Htheorem, II. *Physics Letters A* 158, 1–8.

Valentini, A. (1992). On the pilot-wave theory of classical, quantum and subquantum physics. PhD thesis, International School for Advanced Studies, Trieste, Italy. [\[http://hdl.handle.net/20.500.11767/4334\]](http://hdl.handle.net/20.500.11767/4334)

Valentini, A. (1996). Pilot-wave theory of fields, gravitation and cosmology. In Cushing, J. T., Fine, A. & Goldstein, S. (eds.) *Bohmian mechanics and quantum theory: an appraisal*, pp 45–66. Dordrecht: Kluwer.

Valentini, A. (1997). On Galilean and Lorentz invariance in pilot-wave dynamics. *Physics Letters A* 228, 215–222.

Valentini, A. (2002a). Signal-locality and subquantum information in deterministic hidden-variables theories. In Placek, T. & Butterfield, J. (eds.) *Non-locality and modality*, pp 81–103. Dordrecht: Kluwer.

Valentini, A. (2002b). Signal-locality in hidden-variables theories. *Physics Letters A* 297, 273–278.

Valentini, A. (2002c). Subquantum information and computation. *Pramana— Journal of Physics* 59, 269–277.

Valentini, A. (2004). Black holes, information loss, and hidden variables. [arXiv:hep-th/0407032.](http://arxiv.org/abs/hep-th/0407032)

Valentini, A. (2007). Astrophysical and cosmological tests of quantum theory. *Journal of Physics A* 40, 3285–3303.

Valentini, A. (2008). Hidden variables and the large-scale structure of spacetime. In Craig, W. L. & Smith, Q. (eds.) *Einstein, relativity and absolute simultaneity*, pp 125–155. London: Routledge.

Valentini, A. (2010). Inflationary cosmology as a probe of primordial quantum mechanics. *Physical Review D* 82, 063513.

Valentini, A. (2020). Foundations of statistical mechanics and the status of the Born rule in de Broglie-Bohm pilot-wave theory. In Allori, V. (ed.) *Statistical mechanics and scientific explanation: determinism, indeterminism and laws of nature*, pp 423–477. Singapore: World Scientific.

Valentini, A. (2021). Quantum gravity and quantum probability. [arXiv:2104.07966.](http://arxiv.org/abs/2104.07966)

Valentini, A. (2023). Beyond the Born rule in quantum gravity. *Foundations of Physics* 53, 6.

Valentini, A. (2025). *Introduction to quantum foundations and pilot-wave theory*. Oxford: Oxford University Press.

Valentini, A. and Westman, H. (2005). Dynamical origin of quantum probabilities. *Proceedings of the Royal Society A* 461, 253–272.

Vigier, J. P. (1985). Nonlocal quantum potential interpretation of relativistic actions at a distance in many-body problems. In Tarozzi, G. & van der Merwe, A. (eds.) *Open questions in quantum physics: invited papers on the foundations of microphysics*, pp 297–322. Dordrecht: Reidel.

Vink, J. C. (1992). Quantum potential interpretation of the wave function of the universe. *Nuclear Physics B* 369, 707–728.

Vitenti, S., Peter, P. and Valentini, A. (2019). Modeling the large-scale power deficit with smooth and discontinuous primordial spectra. *Physical Review D* 100, 043506.