ON PROPAGATION OF INFORMATION IN QUANTUM MECHANICS AND MAXIMAL VELOCITY BOUNDS

ISRAEL MICHAEL SIGAL AND XIAOXU WU

ABSTRACT. We revisit key notions related to evolution of quantum information in quantum mechanics and prove uniform bounds on the maximal speed of propagation of quantum information for states and observables with exponential error bounds. Our results imply, in particular, a quantum mechanical version of the Lieb-Robinson bound, which is known to yield various constraints on propagation of quantum information. We propose a novel approach to proving maximal speed bounds.

1. INTRODUCTION

1.1. **Problem and results.** Study of evolution of information in condensed matter physics is an active, robust area of research with many profound results. At the same time, perhaps, due to difficulty of experimental implementation, with exception of a few works on quantum open systems ([8,9]), this fundamental issue was not tackled in the original setting of quantum mechanics, i.e. at zero particle density. In this paper, we address this subject in a systematic way.

Investigation of propagation of quantum information has begun in the context of condensed matter physics with the discovery ([7, 14, 28–32, 45, 47]) that the Lieb-Robinson bound obtained for lattice spin systems in statistical mechanics can be used to derive general constraints on propagation of quantum information. Time bounds on quantum messaging, creation and propagation of corrections and entanglement, state transport and control, quantum simulation algorithms believe propagation raised in these papers were improved and extended significantly in [6, 11, 12, 14–20, 22, 24, 25, 36–42, 44, 46, 48–52, 54, 56, 59, 61–66], see the survey papers [4, 25, 51] and brief reviews in [18, 19, 38].

A different approach was introduced in [58] and extended in [1-3, 23, 33, 35, 55, 57, 60]. Dealing originally with scattering theory in quantum mechanics, it was extended to many-body systems proving light-cone bounds on the propagation in bose gases ([18, 19, 42, 59]), the problem which was open since the groundbreaking work of Lieb and Robinson ([43]).

Date: September 4, 2024.

²⁰²⁰ Mathematics Subject Classification. 35Q40 (primary); 81P45 (secondary).

Key words and phrases. Maximal propagation speed; Lieb-Robinson bounds; quantum mechanics; quantum information; quantum light cones, quantum observables quantum states.

In this paper, we consider the evolution of states and observables in the quantummechanical context and prove the uniform maximal velocity bound with exponential tails yielding existence of the effective light cone (LC) (modulo exponentially small leakage) in quantum mechanical systems. In particular, we prove the Lieb-Robinson bound implying the simultaneous measurability of evolving observables as long as their light cones do not intersect.

1.2. Setup. Consider a quantum system with a state space \mathcal{H} and a Hamiltonian H, a self-adjoint operator on \mathcal{H} . We suppose that $\mathcal{H} = L^2(\Lambda)$, where Λ is either \mathbb{R}^n or \mathbb{Z}^n or a bounded subset (box) in \mathbb{R}^n or \mathbb{Z}^n . To fix ideas, in what follows, we take $\Lambda = \mathbb{R}^n$.

A specific operator H we have in mind is the Schrödinger-type operator

$$H = \omega(p) + V(x), \tag{1.1}$$

where $\omega(k)$ is a real smooth positive function, $p := -i\nabla$ is the momentum operator and the potential V(x) is a real function s.t. H is self-adjoint on the domain of $\omega(p)$, i.e. V(x) is $\omega(p)$ -bounded with the relative bound < 1.

One can recognize information by the properties that it is transmittable, deletable¹, localizable and measurable (or at least detectable).

For quantum mechaniccal systems, the second property requires extending the state space \mathcal{H} to the space, S_1^+ , of positive, trace-class operators, ρ , acting on \mathcal{H} . The original state space \mathcal{H} is identified with the subspace of rank one projections. Quantum information related to a given system is encoded in density operators describing it.

The evolution of density operators, is given by the von Neumann equation (vNE) (here and in the rest of this section we set $\hbar = 1$)

$$\frac{\partial \rho_t}{\partial t} = -i[H, \rho_t], \quad \text{with} \quad \rho_{t=0} = \rho_0.$$
(1.2)

This equation preserves the rank of projections and, when reduced to the rank-one projections, is equivalent to the Schrödinger equation.

The initial value problem (1.2) has a unique solution which generates the automorphism on S_1^+ :

$$\alpha_t'(\rho) := e^{-iHt} \rho e^{iHt}.$$
(1.3)

This formula allows one reduce many results on propagation of states and observables to estimates on the Schrödinger evolution e^{-iHt} .

However, the framework of the vNE is much broader and entails a different take on the evolution problem (i.e. the semiclassics for the vNE leads to Liouville's equation, rather than Newton's one), and it is foundational for the theory of open quantum dynamics.

¹One should be able to erase parts of information one processes (i.e. irrelevant or inaccessible parts)

1.3. **MVB.** We begin with a key result concerning the propagator e^{-itH} , which implies a variety of bounds on propagation of quantum information.

In what follows, we always assume (without specifying this) that the quantum Hamiltonian H we deal with is self-adjoint. For self-adjoint operators on $L^2(\mathbb{R}^n)$, we use the notation $\sup A = \sup_{u \in D(A), ||u|| = 1} \langle u, Au \rangle$. For families of bounded operators,

we use the notion of analyticity in the sense of Kato (resolvent sense, see [53], Section XII).

Let T_{ξ} be the unitary operator of multiplication by the function $e^{-i\xi \cdot x}$. For a selfadjoint operator H on $L^2(\mathbb{R}^n)$, we introduce the operator family $H_{\xi} := T_{\xi}HT_{\xi}^{-1}$. Now, we assume

(A) The family $H_{\xi} := T_{\xi}HT_{\xi}^{-1}$, $\xi \in \mathbb{R}^n$, has an analytic continuation in ξ from \mathbb{R}^n to \mathcal{S}_a^n in the sense of Kato and for this continuation $\operatorname{Im} H_{\zeta} = \frac{1}{2i}(H_{\zeta} - H_{\zeta}^*), \, \zeta \in \mathcal{S}_a^n$, are bounded operators.

We fix $\mu \in (0, a)$ and define the number

$$c := \sup_{\xi \in \mathbb{R}^n, b \in \mathcal{S}^{n-1}} \sup(\mathrm{Im} H_{\xi + i\mu b})/\mu.$$
(1.4)

In what follows, X and Y denote open subsets of \mathbb{R}^n , $X^c := \mathbb{R}^n - X$, d_{XY} , the distance between X and Y and χ_X , the characteristic function of X, as well as the operator of multiplication by this function. Moreover, depending on the context, $\|\cdot\|$ stands either for the norm in $L^2(\mathbb{R}^n)$, or the operator norm on $L^2(\mathbb{R}^n)$. We have the following result:

Theorem 1.1 (Light cone (maximal propagation velocity) bound). Let Condition (A) hold and let $\mu \in (0, a)$. Then, for any $\mu' \in (0, \mu)$ and for any two disjoint sets X and Y in \mathbb{R}^n , we have,

$$\|\chi_X e^{-iHt}\chi_Y\| \leqslant C e^{-\mu'(d_{XY} - c'|t|)},\tag{1.5}$$

where $c' = \frac{\mu c}{\mu'}$, with c given in Eq. (1.4), and C > 0 is a constant depending on $\frac{\mu}{\mu'} - 1, \mu$ and n.

This theorem is proven in Section 2. We call inequality (1.5) the *uniform* maximal velocity bound (uMVB).

Condition (A) implies that for any $a' \in [0, a)$,

$$\sup_{\zeta \in \mathcal{S}^n_{a'}} \sup \operatorname{Im} H_{\zeta} < \infty. \tag{1.6}$$

Indeed, since $\operatorname{Im} H_{\zeta+\xi} = T_{\zeta} \operatorname{Im} H_{\xi} T_{\zeta}^{-1} \quad \forall \xi \in \mathbb{R}^n \text{ and } \zeta \in \mathcal{S}_a^n$, we have that $\sup_{\zeta \in \mathcal{S}_{a'}^n} \operatorname{Sup} \operatorname{Im} H_{\zeta}$ is independent of $\operatorname{Re} \zeta$ and therefore

$$\sup_{\zeta \in \mathcal{S}_{a'}^n} \sup \operatorname{Im} H_{\zeta} = \sup_{\operatorname{Im} \zeta \in [0, a']^n} \sup \operatorname{Im} H_{\zeta} < \infty.$$
(1.7)

For the Schrödinger-type operator (1.1) on $L^2(\mathbb{R}^n)$, Assumption (A) follows from the following condition

 $\omega(k)$ has an analytic continuation $(\omega(\zeta))$ from \mathbb{R}^n to the polystrip \mathcal{S}_a^n , for some a > 0, and $\operatorname{Im}\omega(\zeta)$ is a bounded function on \mathcal{S}_a^n . (1.8)

Note that the Schrödinger operator $H = -\Delta + V$ does not satisfy the second part of Condition (1.8) since $\text{Im}(p + \zeta)^2 = 2p \cdot \text{Im}\zeta$ is not bounded. On the other hand, the semi-relativistic Hamiltonian

$$H = \sqrt{-\Delta + m^2} + V$$

obeys Condition (1.8) with a = m.

Earlier, state-dependent, power-decay MVB were proven, in connection with quantum scattering theory, in [2, 33, 34, 58, 60].

1.4. Localization. A key notion in analysis of evolution of quantum information is that of localization. It is reasonable to consider states localized in bounded sets, say, states created in an apparatus in a lab. With this motivation, we say that

• a state ρ is *localized* in X, if in ρ , the probability for the system to be in X is equal to 1:

$$\operatorname{Tr}(\chi_X \rho) = 1 \quad \text{or} \quad \operatorname{Tr}(\chi_{X^c} \rho) = 0. \tag{1.9}$$

Remark 1.2. By linearity, this notion could be readily extended to the one of *locally perturbed states*.

1.5. Light cone for evolution of states and observables.

Corollary 1.3. Suppose Condition (A) holds. Then, for any density operator ρ_0 localized in X, the probability that its evolution $\rho_t = \alpha'_t(\rho_0)$ is localized in a disjoint set Y is bounded as

$$\operatorname{Tr}(\chi_Y \alpha'_t(\rho_0)) \leqslant C e^{-\mu'(d_{XY} - c't)},\tag{1.10}$$

where μ, c, μ', c' are as in Theorem 1.1 and $C = C(\frac{\mu}{\mu'} - 1, \mu, n) > 0.$

This corollary says that the probability that ρ_t spills outside the light cone

$$\Lambda_{X,c} := \{ (x,t) : d_X(x) < ct \}$$
(1.11)

of X is exponentially small.

The second key ingredient in the general theory is the notion of observables. These are operators on \mathcal{H} representing actual physical quantities and their probability distributions. An average of a physical quantity (say, momentum) represented by an observable A (say, $p = -i\nabla$) in a state ρ is given by $\text{Tr}(A\rho)$. There is a duality between states and observables given by the coupling

$$\rho(A) = \operatorname{Tr}(A\rho), \tag{1.12}$$

which can be considered as either a linear, positive functional of A or a convex one of ρ . In what follows, we identify density operators ρ with linear positive functionals $\rho : A \to \rho(A) := \text{Tr}(A\rho)$.

The evolution of observables is determined by the Heisenberg equation

$$\partial_t A_t = i [H, A_t]. \tag{1.13}$$

Given initial conditions $A_t|_{t=0} = A$, (1.13) generates the (Heisenberg) automorphism

$$A_t \equiv \alpha_t(A) = e^{itH} A e^{-itH}.$$
(1.14)

The Heisenberg equation (or representation) is equivalent to the vNE and, since observables form C^* algebra, often, is more convenient to work with. The duality between states and observables extends to respective evolutions

$$\operatorname{Tr}(\alpha_t(A)\rho) = \operatorname{Tr}(A\alpha_t'(\rho)). \tag{1.15}$$

It is natural to have observables which act locally, i.e. in some set, but leave states outside this set unchanged. Thus, we introduce

• an observable A acts on X iff it is of the form

$$A = \chi_X A \chi_X + \chi_{X^c}, \tag{1.16}$$

where, recall, χ_X and χ_{X^c} stand for multiplication operators by the corresponding cut-off functions (so that $\chi_X + \chi_{X^c} = 1$).

As suggested by the term, (1.16) implies that $\chi_{X^c}A\psi = \chi_{X^c}\psi$ and $A\psi = A\chi_X\psi + \chi_{X^c}\psi$. Note that if A and B act on X and Y, respectively, then

 $[A, B] = 0, \quad \text{whenever } X \cap Y = \emptyset. \tag{1.17}$

We call the smallest set on which an observable A acts the *action domain* of this observable and denote it by actA. Many notions related to and statements about evolution of quantum information can be formulated in parallel for states and observables.

The next useful result on localization of the Heisenberg evolution parallels Corollary 1.3. It shows that the evolution $A_t = \alpha_t(A)$ acts, up to exponentially small tails, within the light cone of its initial action domain of A.

We define an approximation of the evolution $A_t = \alpha_t(A)$ in the set U as

$$A_{t,U} := \chi_U A_t \chi_U + \chi_{U^c}. \tag{1.18}$$

Clearly, $A_{t,U}$ act on the set U. For a subset $X \subset \mathbb{R}^n$, let X_η be the η -neighborhood of X :

$$X_{\eta} = \{ x \in \mathbb{R}^{n} : d_{X}(x) < \eta \}.$$
(1.19)

Theorem 1.4 (Light cone approximation of Heisenberg evolution). Suppose the Condition (A) holds. Then there exist c > 0 and C = C(c, n) > 0 s.t. for any

 $\eta \ge 1$ and every open set X and every operator A acting on X, the evolution $A_t \equiv \alpha_t(A)$ satisfies

$$|A_t - A_{t,X_\eta}|| \le C e^{-\mu(\eta - ct)} ||A||.$$
(1.20)

This theorem is proven in Section 4. It says that, up to exponentially small tails, the evolution of an operator acting on X acts inside the *c*-light cone of X.

One can also define the localization of the evolution A_t to a set U as $A_{t,U} = e^{itH_U}Ae^{-itH_U}$ as an evolution of A with a Hamiltonian H_U supported in U, but the proof of this version is more involved.

Our next result has no parallel for quantum states and is an analogue of one of the key results of quantum information theory.

Theorem 1.5 (Quantum-mechanical Lieb-Robinson bound). Suppose Condition (A) holds and let $X, Y \subset \mathbb{R}^n$ with $d_{XY} > 0$. Then, there exist c > 0 and C = C(n, c) > 0 s.t. for every pair operators A and B acting on X and Y, respectively, we have the following estimate

$$\|[\alpha_t(A), B]\| \le C e^{-\mu(d_{XY} - ct)} \|A\| \|B\|.$$
(1.21)

A proof of Theorem 1.5 is given in Section 3. We call (1.21) the quantummechanical Lieb-Robinson bound (LRBqm). Recall that the physical importance of commuting quantum observables is that they can be measured simultaneously.

Theorem 1.5 yields that, for any state ρ and for all $|t| < d_{XY}/c$,

$$|\rho([\alpha_t(A), B])| \leq C ||A|| ||B|| e^{-\mu(d_{XY} - ct)}.$$
(1.22)



FIGURE 1. Light cone diagram of A and B

Estimate (1.22) shows that, with the probability approaching 1 exponentially, as $t \to \infty$, an evolving observable $A_t = \alpha_t(A)$ commutes with any other observable acting outside the light cone

$$\{ (x,t) \mid \operatorname{dist}(x,\operatorname{act} A) \leqslant ct \},\$$

of $\operatorname{act} A$, where $\operatorname{act} A$ is the smallest set on which A acts (see Fig. 1).

Recall that the expectation $-\rho([A(t), B]^2) \equiv -\operatorname{tr}([A(t), B]^2\rho)$ is called the outof-time-order correlations (OTOC). The inequality $-\rho([A(t), B]^2) \leq ||[A(t), B]||^2$ and Theorem 1.5 imply

Corollary 1.6 (OTOC estimate). Suppose the Condition (A) holds and let μ and c be as in Theorem 1.5. Then $OTOC - \rho([A(t), B]^2)$ satisfies the estimate

$$-\rho([A(t), B]^2) \leqslant C e^{-2\mu(d_{XY} - ct)} (\|A\| \|B\|)^2.$$
(1.23)

Estimate (1.23) extends an estimate of OTOC in the finite dimensions,

$$\frac{1}{D}\operatorname{tr}([A(t),B]^2),\tag{1.24}$$

where $D = \dim \mathcal{H}$, see e.g. [38], to the infinite-dimensional case.

In Appendix A, we extend our approach to differentiable deformations H_{ξ} obtaining power bounds on the error terms. We expect that it could be extended to open quantum systems, where estimates of evolving states and observables cannot be reduced to estimate of the Schrödinger evolution and one has to estimate instead the von Neumann-Lindblad semigroup, e^{Lt} , see [8], and to many-body (condensed matter) systems as suggested by our results on N-particle Schrödinger dynamcis presented in Section B.

This paper is organized as follows. Theorems 1.1, 1.4 and 1.5 are proven in Sections 2, 4 and 3, respectively. In Appendices A and B, we present extensions of our technologies to differentiable families H_{ξ} and to N-particle systems.

<u>Notation.</u> We use the abbreviation $\|\cdot\|$ for both $\|\cdot\|_{L^2(\mathbb{R}^n)}$ and $\|\cdot\|_{L^2(\mathbb{R}^n)\to L^2(\mathbb{R}^n)}$. Throughout this paper, C will denote a constant and may vary from one line to another. We write \lesssim or \gtrsim whenever $A \leq CB$ or $CA \geq B$ for some constant C > 0. We write $A \leq_a B$ or $A \gtrsim_a B$ if $A \leq C_a B$ or $C_a A \geq B$ for some constant $C_a > 0$ which depends on parameter a.

As usual,
$$\partial_x^{\alpha} = \prod_{j=1}^n \partial_{x_j}^{\alpha_j}$$
, for $\alpha = (\alpha_1, \cdots, \alpha_n)$, $x = (x_1, \cdots, x_n)$ and $|\alpha| = \sum_{j=1}^n \alpha_j$.

In what follows, C stands for a generic constant which changes from equation to equation and are independent of variable parameters, such as distances between sets and their sizes, etc.

2. Proof of Theorem 1.1

Proof of Theorem 1.1. Let $n \ge 2$. Given $b \in S^{n-1}$, we define the subspace $R_b^{\perp} = b^{\perp} = \{k \in \mathbb{R}^n : k \cdot b = 0\}$ and the map $\varphi_b : \mathbb{C} \times R_b^{\perp} \to \mathbb{C}^n$ as

$$\varphi_b(z,k^{\perp}) := zb + k^{\perp}. \tag{2.1}$$

Thus, points k of \mathbb{R}^n are parameterized by $(\lambda, k^{\perp}) \in \mathbb{R} \times R_b^{\perp}$. We have

Lemma 2.1. Let $n \ge 2$. Under Assumption (A), for any $b \in S^{n-1}$, the function φ_b has the range in \mathcal{S}^n_a and is analytic in z. Moreover, $\omega \circ \varphi_b$ has an analytic continuation in z from $\mathbb{R} \times R^{\perp}_b$ to $\mathcal{S}_a \times R^{\perp}_b$.

Proof. φ is clearly analytic in z. Let $k^z \equiv \varphi_b(z, k^{\perp}) := zb + k^{\perp}$, or in coordinates $k_j^z = zb_j + k_j^{\perp}, j = 1, \cdots, n$. We have $|\text{Im}k_j^z| = |\text{Im}z||b_j| \leq |\text{Im}z|$. Hence, $z \in S_a$ implies $k^z \in S_a^n$ and therefore $\omega(k^{\lambda}) \equiv \omega(\varphi_b(\lambda, k^{\perp}))$ has an analytic continuation from $\mathbb{R} \times R_b^{\perp}$ to $S_a \times R_b^{\perp}$.

 $\xi^{\lambda} = \lambda b + \xi^{\perp}$ gives a parameterization of $\xi \in \mathbb{R}^n$ by $(\lambda, \xi^{\perp}) \in \mathbb{R} \times \mathbb{R}_b^{\perp}$. For notational convenience, we extend the definition of

$$\xi^z = \varphi_b(z, \xi^\perp) = zb + \xi^\perp \tag{2.2}$$

to n = 1, by setting in this case b = 1 and $\xi^{\perp} = 0$, i.e. $\xi^z = \varphi_b(z, \xi^{\perp}) = z$. In what follows, b is a vector in S^{n-1} for n > 1 and b = 1 for n = 1.

Recall the *n*-parameter unitary group

$$T_{\xi} = e^{-ix \cdot \xi}, \xi \in \mathbb{R}^n, \tag{2.3}$$

and introduce the transformation

$$U_{t,\xi} := T_{\xi} e^{-itH} T_{\xi}^{-1}.$$
 (2.4)

By the unitarity of T_{ξ} , we have

$$U_{t,\xi} = e^{-iH_{\xi}t}, \qquad \text{where } H_{\xi} := T_{\xi}HT_{\xi}^{-1}.$$
 (2.5)

Proposition 2.2. Under Assumption (A), the operators $U_{t,\xi}$ and $U_{t,\xi^{\lambda}}$ have analytic continuations in ξ and λ from \mathbb{R}^n to \mathcal{S}^n_a and from \mathbb{R} to \mathcal{S}_a , as bounded operators.

Proof. By Assumption (A), H_{ξ} has an analytic continuation in ξ from \mathbb{R}^n to \mathcal{S}^n_a , in the sense of Kato, and the operator $\operatorname{Im} H_{\zeta} = \frac{1}{2i}(H_{\zeta} - H_{\zeta}^*)$ is bounded for every $\zeta \in \mathcal{S}^n_a$. Hence, the relations $H_{\zeta} = H_r + iH_i$, where $H_r = \operatorname{Re} H_{\zeta}$ and $H_i = \operatorname{Im} H_{\zeta}$, and

$$(H_{\zeta} - i\lambda)^{-1} = (H_r + iH_i - i\lambda)^{-1} = R_i^{1/2} \left[R_i^{1/2} H_r R_i^{1/2} - i \right]^{-1} R_i^{1/2}, \qquad (2.6)$$

where $R_i = (\lambda - H_i)^{-1}$, imply that $\sigma(-iH_{\zeta}) \subset \{\zeta \in \mathbb{C}^n : |\operatorname{Re} \zeta| \leq C\}, \forall \zeta \in \mathcal{S}_a^n$, for $C = ||\operatorname{Im} H_{\zeta}|| > 0$, and, for any $\lambda \in \mathbb{C}$ with $|\operatorname{Re} \lambda| > C$, we have the estimate

$$\|(H_{\zeta} - i\lambda)^{-1}\| \le (|\operatorname{Re} \lambda| - \|\operatorname{Im} H_{\xi^z}\|)^{-1}.$$
 (2.7)

Hence, by the Hille-Yosida theorem, H_{ζ} generates the bounded evolution $U_{t,\zeta} = e^{-iH_{\zeta}t}$, $t \in \mathbb{R}$, which is analytic as an operator-function of $\zeta \in \mathcal{S}_a^n$, by a standard theory.

Let
$$U_t \equiv e^{-itH}$$
 and $U_{t,\xi} := T_{\xi} U_t T_{\xi}^{-1}$. We have

Lemma 2.3 (Key lemma). Under Condition (A), we have

$$\chi_X U_t \chi_Y = \chi_X T_{\zeta}^{-1} U_{t,\zeta} T_{\zeta} \chi_Y, \qquad \forall \zeta \in \mathcal{S}_a^n.$$
(2.8)

Proof. Using the invertibility of T_{ξ} and the definition $U_{t,\xi} = T_{\xi}U_tT_{\xi}^{-1}$, we rewrite $\chi_X U_t \chi_Y$ as

$$\chi_X U_t \chi_Y = \chi_X T_{\xi}^{-1} T_{\xi} U_t T_{\xi}^{-1} T_{\xi} \chi_Y = \chi_X T_{\xi}^{-1} U_{t,\xi} T_{\xi} \chi_Y, \qquad \forall \, \xi \in \mathbb{R}^n.$$
(2.9)

The right-hand side of Eq. (2.9) has an analytic continuation in ξ from \mathbb{R}^n to \mathcal{S}^n_a and is independent of $\xi \in \mathbb{R}^n$. Hence, its analytic continuation is independent of $\zeta \in \mathcal{S}^n_a$ and (2.8) follows.

Relation (2.8) implies, for $\zeta \in \mathcal{S}_a^n$,

$$\|\chi_X U_t \chi_Y\| \le \|\chi_X T_{\zeta}^{-1}\| \|U_{t,\zeta}\| \|\chi_Y T_{\zeta}\|.$$
(2.10)

Let $\zeta = \xi + i\mu b, \xi \in \mathbb{R}^n, \mu \in (0, a)$ and $b \in S^{n-1}$. Then estimate (2.10) implies

$$\|\chi_X U_t \chi_Y\| \le e^{-\mu(r_X - r_Y)} \|U_{t,\zeta}\|,$$
(2.11)

where

$$r_Y := \sup_{y \in Y} b \cdot y$$
 and $r_X := \inf_{x \in X} b \cdot x.$ (2.12)

We can cover X and Y by small balls. Hence, we begin with $Y = B_r(y_0)$ and $X = B_r(x_0)$ with $r = \frac{\epsilon}{2} d_{XY}$ for some $y_0 \in Y$, $x_0 \in X$ and $\epsilon \in (0, 1)$.

Translate both balls by y_0 to place y_0 at the origin and x_0 at $x_0 - y_0$. Let S_{y_0} denote the corresponding shift operator. Then we have

$$S_{y_0}\chi_X U_t \chi_Y S_{y_0}^{-1} = S_{y_0}\chi_X S_{y_0}^{-1} S_{y_0} U_t S_{y_0}^{-1} S_{y_0} \chi_Y S_{y_0}^{-1} = \chi_{X^{y_0}} U_t^{y_0} \chi_{Y^{y_0}}, \qquad (2.13)$$

where $X^{y_0} = X - y_0 = B_r(x_0 - y_0)$, $Y^{y_0} = Y - y_0 = B_r(0)$ and $U_t^{y_0} := S_{y_0}U_t S_{y_0}^{-1}$. Thus, it suffices to estimate the right-hand side of Eq. (2.13).

Now, we skip the superindex y_0 , so we are back to Eq. (2.8), but with $X = B_r(x_0 - y_0)$ and $Y = B_r(0)$.

Let $\zeta = \xi + i\mu b$, where $b = \frac{x_0 - y_0}{|x_0 - y_0|}$. (For each pair of points $x_0 \in X$ and $y_0 \in Y$, we choose a different analytic deformation b.) Since $|x_0 - y_0| \ge d_{XY}$, by the definition of r_X and r_Y , we have

$$r_X - r_Y \ge (1 - \epsilon) |x_0 - y_0| \ge (1 - \epsilon) d_{XY}.$$
 (2.14)

Eqs. (2.11) and (2.14) yield

$$\|\chi_X U_t \chi_Y\| \le \|U_{t,\zeta}\| e^{-\mu(1-\epsilon)d_{x_0 y_0}}.$$
(2.15)

Note that the complex deformation $U_{t,\zeta}$ of the evolution operator changes from one pair of balls to another. The next proposition provides a uniform estimates of various such deformations.

Proposition 2.4. Let Assumption (A) be satisfied, and let μ and c be as in Eq. (1.4). Then we have the estimate

$$\|U_{t,\xi+i\mu b}\| \leqslant e^{\mu t c}, \qquad \forall \, \xi \in \mathbb{R}^n.$$
(2.16)

I. M. SIGAL AND X. WU

Proof of Proposition 2.4. Take $g \in L^2(\mathbb{R}^n)$. We denote $g_{t,\zeta} := U_{t,\zeta}g, \zeta \in \mathcal{S}_a^n$, and compute

$$\partial_t \|g_{t,\zeta}\|^2 = -i\langle g_{t,\zeta}, (H_\zeta - H_\zeta^*)g_{t,\zeta}\rangle = 2\langle g_{t,\zeta}, (\operatorname{Im} H_\zeta) g_{t,\zeta}\rangle.$$
(2.17)

For $\zeta = \xi + i\mu b$, by Eq. (1.4), we have, $\text{Im}H_{\zeta} \leq \mu c$. This together with Eq. (2.17) implies

$$|\partial_t \| g_{t,\zeta} \|^2 | \leq 2\mu c \| g_{t,\zeta} \|^2.$$
(2.18)

Since $g_{t,\zeta}|_{t=0} = g$, this gives $||g_{t,\zeta}|| \leq ||g|| e^{\mu ct}$ yielding (2.16).

Eqs. (2.15) and (2.16) yield

$$\|\chi_X U_t \chi_Y\| \le e^{\mu ct} e^{-\mu (1-\epsilon) d_{x_0 y_0}}.$$
(2.19)

Now, we return to arbitrary disjoint sets X and Y. We cover X and Y by the balls, $B_j^X = B_r(x_j), j = 1, \dots, N_1$ and $B_k^Y = B_k(y_k), k = 1, \dots, N_2$, in \mathbb{R}^n of the radius $r = \frac{\epsilon}{2}d_{XY}$, centered at x_j and y_k , respectively. N_1 and N_2 could be either finite or infinite. With this cover, we associate partitions of unity

$$\chi_X = \sum_{j=1}^{N_1} \chi_j^2 \quad \text{and } \chi_Y = \sum_{k=1}^{N_2} \tilde{\chi}_k,$$
(2.20)

where χ_j and $\tilde{\chi}_k$ satisfy $\operatorname{supp}(\chi_j) \subset B_r(x_j)$ and $\operatorname{supp}(\tilde{\chi}_k) \subset B_r(y_k), j = 1, \cdots, N_1$ and $k = 1, \cdots, N_2$. For each $g \in L^2(\mathbb{R}^n)$, we estimate, using Eqs. (2.20),

$$\|\chi_X e^{-itH} \chi_Y g\|^2 = \sum_{k=1}^{N_1} \|\chi_k e^{-iHt} \chi_Y g\|^2 \leqslant \sum_{k=1}^{N_1} \left(\sum_{j=1}^{N_2} \|\chi_k e^{-iHt} \tilde{\chi}_j g\| \right)^2.$$
(2.21)

By (2.19), we have

$$\|\chi_X e^{-itH} \chi_Y g\|^2 \leq C e^{2\mu ct} M(g),$$
 (2.22)

where M(g) is given by, with $\mu' = \mu(1 - \epsilon)$,

$$M(g) := \sum_{k=1}^{N_1} \left(\sum_{j=1}^{N_2} e^{-\mu' d_{x_k y_j}} \| \tilde{\chi}_j g \| \right)^2$$
$$= \sum_{k=1}^{N_1} \sum_{j=1}^{N_2} \sum_{l=1}^{N_2} e^{-\mu' (d_{x_k y_j} + d_{x_k y_l})} \| \tilde{\chi}_j g \| \| \tilde{\chi}_l g \|.$$
(2.23)

To estimate M(g), we use arithmetic mean inequality to obtain

$$M(g) \leqslant \sum_{k=1}^{N_1} \sum_{j=1}^{N_2} \sum_{l=1}^{N_2} e^{-\mu'(d_{x_k y_j} + d_{x_k y_l})} \left(\frac{\|\tilde{\chi}_j g\|^2 + \|\tilde{\chi}_l g\|^2}{2}\right).$$
(2.24)

By the symmetry with respect to j and l in the right-hand side of (2.24), (2.24) implies

$$M(g) \leq \sum_{k=1}^{N_1} \sum_{j=1}^{N_2} \sum_{l=1}^{N_2} e^{-\mu'(d_{x_k y_j} + d_{x_k y_l})} \|\tilde{\chi}_j g\|^2 = \sum_{j=1}^{N_2} \|\tilde{\chi}_j g\|^2 C_{XY},$$
(2.25)

where

$$C_{XY} := \sum_{k=1}^{N_1} \sum_{l=1}^{N_2} e^{-\mu'(d_{x_k y_j} + d_{x_k y_l})}.$$
(2.26)

First, we sum over l. To this end, for each k, we decompose \mathbb{R}^d into the spherical shells

$$\Lambda_m(x_k) = \{x : r_m \leqslant d_{xx_k} < r_{m+1}\},\tag{2.27}$$

where $m = 0, \dots, r_m = r_0 + m\epsilon d_{XY}$, with $r_0 = (1 - \epsilon)d_{XY}$. We sum first over the balls covering a given shell and then over the shells. Each shell $\Lambda_m(x_k)$ is covered by at most \tilde{N}_m balls B_l^Y , with $\tilde{N}_m = C_n r_m^{n-1}$ for some constant $C_n > 0$. This gives

$$C_{X,Y} \leqslant \sum_{k=1}^{N_1} \sum_{m=0}^{\infty} C_n r_m^{n-1} e^{-\mu'(d_{x_k y_j} + r_m)}.$$
(2.28)

To evaluate the sum over the shells, we use that

$$\sum_{m=0}^{\infty} r_m^{n-1} e^{-\mu' r_m} \leqslant C \sum_{m=0}^{\infty} e^{-(1-\epsilon/2)\mu' r_m} \\ \leqslant C e^{-(1-\epsilon/2)\mu' r_0} = C e^{-(1-\epsilon/2)(1-\epsilon)\mu' d_{XY}}.$$
(2.29)

We conclude that there is a $C_{n,\epsilon,\mu'} > 0$ depending on n, ϵ and $\mu', \text{ s.t.}$

$$C_{X,Y} \leqslant C_{n,\epsilon,\mu'} \sum_{k=1}^{N_1} d_{XY}^{n-1} e^{-\mu'(d_{x_k y_j} + d_{XY})}.$$
(2.30)

Next, to estimate the sum over x_k , we introduce the spherical shells centered at y_j

$$\Lambda_m(y_j) = \{x : r_m \le d_{xy_j} < r_{m+1}\},\tag{2.31}$$

where $r_m = r_0 + m\epsilon d_{XY}$, $m = 0, \cdots$, with $r_0 = (1 - \epsilon)d_{XY}$. Similarly, following (2.30), we obtain

$$C_{X,Y} \leqslant C_{n,\epsilon,\mu'}^2 d_{XY}^{2(n-1)} e^{-2\mu' d_{XY}}.$$
(2.32)

This, together with Eqs. (2.25) and (2.26), implies

$$M(g) \leqslant C_{n,\epsilon,\mu'}^2 d_{XY}^{2(n-1)} e^{-2\mu' d_{XY}} \|g\|^2.$$
(2.33)

Therefore, using (2.22), and (2.33), we conclude that with $\mu'' = (1 - 2\epsilon)\mu$,

$$\|\chi_X e^{-itH} \chi_Y g\| \le C_{n,\epsilon,\mu'} C'_{n,\epsilon,\mu} e^{\mu ct} e^{-\mu'' d_{XY}} \|g\|,$$
(2.34)

where $C'_{n,\epsilon,\mu} := \sup_{u \ge 0} u^{(n-1)} e^{-\epsilon \mu u}$. Estimate (2.34) yields

$$\|\chi_X U_t \chi_Y\| \le C e^{-\mu''(d_{XY} - c''t)}, \qquad \mu'' = (1 - 2\epsilon)\mu, \ c' = \frac{c}{1 - 2\epsilon}$$
 (2.35)

with the constant C depending on $\epsilon = 1 - \frac{\mu''}{\mu}$, n and μ . This implies (1.5).

3. Localization of observables and proof of Theorem 1.5

We introduce a mathematically convenient notion of localized observables. We say that an observable A is *localized* in X if

$$A\chi_{X^c} = \chi_{X^c}A = 0, \text{ or } A = \chi_X A\chi_X.$$
(3.1)

Since $\chi_{X^c} = \mathbb{1} - \chi_X$, Eq. (1.16) implies, for any operator A acting on X,

$$A = \hat{A}_X + 1, \qquad \text{where } \hat{A}_X = \chi_X A \chi_X - \chi_X. \tag{3.2}$$

By the definition, the observable \tilde{A}_X is localized in X.

Proof of Theorem 1.5. Since A and B act on X and Y, respectively, by (3.2), they are of the form $A = \tilde{A}_X + \mathbb{1}$ and $B = \tilde{B}_Y + \mathbb{1}$, where \tilde{A}_X and \tilde{B}_Y are localized in X and Y, respectively. Let $A_t = \alpha_t(A)$ and $\tilde{A}_{t,X} = \alpha_t(\tilde{A}_X)$. Then $A_t = \tilde{A}_{t,X} + \mathbb{1}$ and

$$[A_t, B] = [\tilde{A}_{t,X}, \tilde{B}_Y]. \tag{3.3}$$

Using that $\tilde{A}_X = \chi_X \tilde{A}_X \chi_X$ and $\tilde{B}_Y = \chi_Y \tilde{B}_Y \chi_Y$ and using Theorem 1.1, we obtain

$$\begin{aligned} \|[\tilde{A}_{t,X}, \tilde{B}_Y]\| \leqslant \|\tilde{A}_X \chi_X e^{-itH} \chi_Y \tilde{B}_Y\| + \|\tilde{B}_Y \chi_Y e^{itH} \chi_X \tilde{A}_X\| \\ \leqslant C e^{-\mu(d_{XY} - ct)} \|\tilde{A}_X\| \|\tilde{B}_X\|. \end{aligned}$$
(3.4)

Since $X^c \neq \emptyset$ and since ||Au|| = ||u|| for u supported in X^c , we have $||A|| \ge 1$. Hence,

$$\|\tilde{A}_X\| \le \|\chi_X A \chi_X\| + \|\chi_X\| \le \|A\| + 1 \le 2\|A\|$$
(3.5)

and similarly for \tilde{B} . These inequalities together with (3.4) yield

$$\|[\tilde{A}_{t,X}, \tilde{B}_Y]\| \le C e^{-\mu (d_{XY} - ct)} \|A\| \|B\|.$$
(3.6)

Relations (3.3) and (3.6) yield (1.21).

4. MVB FOR EVOLUTION OF OBSERVABLES: PROOF OF THEOREM 1.4

Proof of Theorem 1.4. Since A acts on X, by (3.2), it can be be written as $A = \tilde{A}_X + \mathbb{1}$, where the operator \tilde{A}_X is localized in X, i.e. satisfies $\tilde{A}_X = \chi_X \tilde{A}_X \chi_X$. Hence the operator family A_{t,X_n} defined in (1.18) can be written as

$$A_{t,X_{\eta}} := \chi_{X_{\eta}} \alpha_t(A_X) \chi_{X_{\eta}} + \mathbb{1}.$$

$$(4.1)$$

By the definition, $A_{t,X_{\eta}}$ acts on X_{η} . To prove (1.20), we use that $A_t = \alpha_t(A_X + \mathbb{1}) = \alpha_t(\tilde{A}_X) + \mathbb{1}$, to write

$$A_t - A_{t,X_\eta} = \alpha_t(\tilde{A}_X) - \chi_{X_\eta} \alpha_t(\tilde{A}_X) \chi_{X_\eta}$$

= $\chi_{X_\eta} \alpha_t(\tilde{A}_X) \chi_{X_\eta^c} + \chi_{X_\eta^c} \alpha_t(\tilde{A}_X).$ (4.2)

Using this relation and Theorem 1.1, we arrive at (1.20).

Remark 4.1. Theorem 1.4 yields a natural (but slightly longer) proof of Theorem 1.5: $A_{t,X_{\eta}}$ commutes with B as long as $\eta < d_{XY}$ and therefore $[A_t, B_t] = [R_t^A, B]$, where $R_t^A = A_t - A_{t,X_{\eta}}$, which leads to an estimate of $[A_t, B_t]$ through an estimate of the remainder R_t^A .

Acknowledgment. The first author is grateful to Jeremy Faupin, Marius Lemm and Jingxuan Zhang, and both authors, to Avy Soffer, for enjoyable and fruitful collaborations. The research of I.M.S. is supported in part by NSERC Grant NA7901. X. W. is supported by 2022 *Australian Laureate Fellowships* grant FL220100072. Her research was also supported in part by NSERC Grant NA7901. Parts of this work were done while the second author was at the Fields Institute for Research in Mathematical Sciences, Toronto.

As we were completing this paper, we came across a very recent preprint [10] whose approach is similar to ours.

Declarations.

- Conflict of interest: The Authors have no conflicts of interest to declare that are relevant to the content of this article.
- Data availability: Data sharing is not applicable to this article as no datasets were generated or analysed during the current study.

APPENDIX A. DIFFERENTIABILITY AND POWER ESTIMATES

In this appendix, we consider a self-adjoint operator H on $L^2(\Lambda)$ under a weaker assumption than the analyticity assumption (A) of Introduction.

With the definition (2.3) and $H_{\xi} = T_{\xi}HT_{\xi}^{-1}, \ \xi \in \mathbb{R}^n$, we assume

(Diff) The family $H_{\xi}, \xi \in \mathbb{R}^n$, is *m* times differentiable, with all derivatives yielding bounded operators.

Given $b \in S^{n-1}$ and $z = \lambda + i\mu$ and recalling the notation $\xi^z = zb + \xi^{\perp}$ (see (2.2)), we define the number \tilde{c} :

$$\tilde{c} := \sum_{k=1}^{m} \frac{1}{k!} \operatorname{Re}(i\mu)^{k-1} \sup_{b \in S^{n-1}} \sup(b \cdot \nabla_{\xi})^k H_{\xi^{\lambda}}.$$
(A.1)

Theorem A.1. Suppose that Assumption (Diff) hold for some $m \ge n$ and let X and Y be two bounded, disjoint sets. Then, for every $\tilde{c}' > \tilde{c}$, there exists a constant C > 0, depending on $\tilde{c}' - \tilde{c}$ and n, such that

$$\|\chi_X e^{-iHt}\chi_Y\| \leqslant CtM(d_{XY} - t\tilde{c}')^{-m-1+n},$$
(A.2)

for all $1 \leq t \leq d_{XY}/\tilde{c}'$, where constant M is given by

$$M := 1 + \sup_{b \in S^{n-1}} \| (b \cdot \nabla_{\xi})^{m+1} H_{\xi} \|.$$
(A.3)

Remark A.2. For Hamiltonians of form (1.1), condition (Diff) follows from the condition

$$\partial^{\alpha}\omega(k) \lesssim 1 \text{ for } 1 \leq |\alpha| \leq m+1 \text{ for some } m \geq 1.$$
 (A.4)

The proof of Theorem A.1 is based on the following proposition.

Proposition A.3. Let $X = B_r(x_0)$ and $Y = B_r(y_0)$ with $r = \epsilon/2, \epsilon \in (0, 1)$, and let $\xi^z = zb + \xi^{\perp}$ (see (2.16)) with $z = \lambda + i\mu \in \mathbb{C}^+$ and $b = \frac{x_0 - y_0}{|x_0 - y_0|} \in S^{n-1}$. For H_{ξ} , m + 1 times boundedly differentiable, instead of (2.8), we have for all $\mu \in (0, 1)$ and $d_{x_0y_0} - \epsilon - t\tilde{c} \ge 0$,

$$\chi_X U_t \chi_Y = \chi_X T_{\xi^z}^{-1} \tilde{U}_{t,\xi^z} T_{\xi^z} \chi_Y + Rem, \tag{A.5}$$

where \tilde{U}_{t,ξ^z} is the almost analytic extension of U_{t,ξ^z} defined as

$$\tilde{U}_{t,\xi^z} = e^{-i\tilde{H}_{\xi^z}t},\tag{A.6}$$

where

$$\tilde{H}_{\xi^z} = \sum_{k=0}^m \frac{1}{k!} (b \cdot \nabla_{\xi})^k H_{\xi^\lambda}(i\mu)^k, \tag{A.7}$$

and Rem is a bounded operator satisfying

$$\|Rem\| \lesssim_{\mu} \frac{tM}{(d_{x_0y_0} - \epsilon - t\tilde{c})^{m+1}},\tag{A.8}$$

with \tilde{c} and M defined in Eqs. (A.1) and (A.3), respectively.

Remark A.4. (i) For \tilde{H}_{ξ^z} given by (A.7), we have

$$\tilde{c} = \sup_{b \in S^{n-1}} \sup \frac{\mathrm{Im}\tilde{H}_{\xi^z}}{\mu}.$$
(A.9)

(ii) For $\mu \in (0, 1)$ sufficiently small, $\xi^z = zb + \xi^{\perp}$ with $z \in \mathbb{C}^+$ and $b \in S^{n-1}$, we have

$$\tilde{c} = \sup_{b \in S^{n-1}} \sup \left(b \cdot \nabla_{\xi} H_{\xi^{\lambda}} \right) + O(\mu).$$
(A.10)

(iii) We can also consider the speed $\tilde{c}(b) = \sup(\mathrm{Im}\tilde{H}_{\xi^z})/\mu$ in a direction $b \in S^{n-1}$.

We derive this proposition from the following two lemmas.

Lemma A.5. Let f(z) be a differentiable function in the strip S_a , for some a, which is independent of Re z. Then

$$f(z) = f(x) - \int_0^y (\bar{\partial}_z f)(x+is)ds, \qquad (A.11)$$

where z = x + y.

Proof. Let z = x + iy. By the fundamental theorem of Calculus, we have

$$f(z) = f(x) - i \int_0^y (\partial_y f)(x + is) ds.$$
 (A.12)

Furthermore, since f(z) is independent of x, we have $\partial_x f(x+iy) = 0$. These two relations and the definition $\overline{\partial}_z = \partial_x + i\partial_y$ imply (A.11).

Lemma A.6. For any $f \in C^{m+1}(\mathbb{R})$, define an almost analytic extension of f as

$$\tilde{f}(z) = \sum_{k=0}^{m} f^{(k)}(x) \frac{(iy)^k}{k!},$$
(A.13)

where $f^{(k)} = \frac{d^k}{dx^k}f$ and z = x + iy. Then \tilde{f} satisfies the estimate

$$|\bar{\partial}_z \tilde{f}(z)| \leq \frac{1}{m!} |f^{(m+1)}(x)| |y|^m.$$
 (A.14)

Proof. Eq. (A.13) follows from the straightforward computation. With $\bar{\partial}_z = \partial_x + i\partial_y$, $\bar{\partial}_z \tilde{f}(z)$ reads

$$\bar{\partial}_{z}\tilde{f}(z) = \sum_{k=0}^{m} f^{(k+1)}(x)\frac{(iy)^{k}}{k!} + i\sum_{k=1}^{m} f^{(k)}(x)\frac{i(iy)^{k-1}}{(k-1)!}$$
$$= \frac{1}{m!}f^{(m+1)}(x)(iy)^{m},$$
(A.15)

which gives (A.14).

Proof of Proposition A.3. Recall that $X = B_r(x_0 - y_0)$ and $Y = B_r(0)$, where $r = \epsilon/2$. Now, let $g(\lambda) = \chi_X T_{\xi^{\lambda}}^{-1} f(\lambda) T_{\xi^{\lambda}} \chi_Y$, where $f(\lambda) := U_{t,\xi^{\lambda}}$. We define

$$\tilde{g}(z) = \chi_X T_{\xi^z}^{-1} \tilde{f}(z) T_{\xi^z} \chi_Y,$$
(A.16)

where $\tilde{f}(z)$ is the almost analytic extension of $f(\lambda)$ in λ constructed in Eq. (A.6).

To compute $\bar{\partial}_z g(z)$, we note that, by Lemma A.6,

$$\bar{\partial}_z \tilde{H}_{\xi^z} = \frac{1}{m!} \partial_\lambda^{m+1} H_{\xi^\lambda} (i\mu)^m. \tag{A.17}$$

This and the Duhamel principle yield

$$\bar{\partial}_z f(z) = \frac{-i}{m!} \int_0^t e^{-i\tilde{H}_{\xi^z}(t-s)} \bar{\partial}_z \tilde{H}_{\xi^z} e^{-i\tilde{H}_{\xi^z}s} ds$$
$$= -i \int_0^t e^{-i\tilde{H}_{\xi^z}(t-s)} \partial_\lambda^{m+1} H_{\xi^\lambda} e^{-i\tilde{H}_{\xi^z}s} ds (i\mu)^m, \qquad (A.18)$$

Since $\chi_X T_{\xi^z}^{-1}$ and $T_{\xi^z} \chi_Y$ are analytic in z, we have, by the Leibnitz rule and Eq. (A.18), that

$$\bar{\partial}_{z}\tilde{g}(z) = \chi_{X}T_{\xi^{z}}^{-1}\bar{\partial}_{z}f(z)T_{\xi^{z}}\chi_{Y}$$
$$= \frac{-i}{m!}\chi_{X}T_{\xi^{z}}^{-1}\int_{0}^{t}R_{t,s}(z)(i\mu)^{m}ds,$$
(A.19)

where, with $z = \lambda + i\mu$,

$$R_{t,u}(z) := \chi_X T_{\xi^z}^{-1} e^{-i\tilde{H}_{\xi^z}(t-u)} \hat{c}_{\lambda}^{m+1} H_{\xi^{\lambda}} e^{-i\tilde{H}_{\xi^z} u} T_{\xi^z} \chi_Y.$$
(A.20)

Next, we claim that $\tilde{g}(z)$ is independent of Re z. Indeed, we have

$$\tilde{g}(z) = \chi_X T_{\xi^z}^{-1} T_\eta \tilde{f}(z) T_\eta^{-1} T_\eta T_{\xi^z} \chi_Y,$$
(A.21)

where $\eta = \alpha b \in \mathbb{R}^n$ with $\alpha \in \mathbb{R}$. Using that T_η commutes with ∂_{ξ} , we find

$$T_{\eta}\partial_{\lambda}^{k}H_{\xi^{\lambda}}T_{\eta}^{-1} = \partial_{\lambda}^{k}[T_{\eta}H_{\xi^{\lambda}}T_{\eta}^{-1}] = \partial_{\lambda}^{k}H_{\xi^{\lambda+a}}, \qquad k = 0, \cdots, m.$$
(A.22)

This, together with Eqs. (A.7), (A.6) and $\tilde{f}(z) = \tilde{U}_{t,\xi^z}$, yields that $T_{\eta}\tilde{f}(z)T_{\eta}^{-1} = \tilde{f}(z+\alpha)$. The last relation, together with (A.21) and the group property $T_{\eta}T_{\xi^z} = T_{\xi^{z+\alpha}}$, implies

$$\tilde{g}(z) = \tilde{g}(z+\alpha), \quad \forall \alpha \in \mathbb{R},$$
(A.23)

which shows that $\tilde{g}(z)$ is independent of Re z. Hence, Lemma A.5 applies to $\tilde{g}(z)$ and yields

$$g(\lambda) = \tilde{g}(\lambda) = \tilde{g}(z) + Rem, \tag{A.24}$$

where, by (A.19),

$$Rem := i \int_0^\mu (\bar{\partial}_z \tilde{g})(x+is) ds$$
$$= \frac{1}{m!} \int_0^\mu \int_0^t R_{t,u}(\lambda+is)(is)^m du ds.$$
(A.25)

Our next goal is to estimate this reminder.

To estimate $R_{t,u}(\lambda + is)$, we proceed as in Proposition 2.4 to obtain

$$\|e^{-i\tilde{H}_{\xi^z}t}\| \leqslant e^{\mu t\tilde{c}},\tag{A.26}$$

where $\mu = \text{Im}z$ and the constant \tilde{c} is defined in Eq. (A.1). Next, from Eq. (A.20) we find,

$$\|R_{t,u}(\lambda+is)\| \leq \|\chi_X T_{\xi^{\lambda+is}}^{-1}\| \|e^{-i\tilde{H}_{\xi^{\lambda+is}}(t-u)}\| \|\partial_{\lambda}^{m+1} H_{\xi^{\lambda}}\| \|e^{-i\tilde{H}_{\xi^{z}}u}\| \|T_{\xi^{\lambda+is}}\chi_Y\|.$$
(A.27)

Proceeding as (2.10)-(2.15), using (A.26) and (A.3) and assuming $d_{XY} - \epsilon - t\tilde{c} \ge 0$, we obtain

$$\|R_{t,u}(\lambda+is)\| \leqslant e^{-s(d_{x_0y_0}-\epsilon-t\tilde{c})}M.$$
(A.28)

This together with Eqs. (A.20) and (A.25) and estimates (A.28) and, for all $|x_0 - y_0| - \epsilon - t\tilde{c} > 0$ and $\mu \in (0, 1)$,

$$\frac{1}{m!} \int_{0}^{\mu} e^{-s(d_{x_{0}y_{0}}-\epsilon-t\tilde{c})} s^{m} ds \lesssim_{\mu} \frac{1}{(d_{x_{0}y_{0}}-\epsilon-t\tilde{c})^{m+1}},$$
(A.29)

yields

$$\|Rem\| \lesssim_{\mu} \frac{tM}{(d_{x_0y_0} - \epsilon - t\tilde{c})^{m+1}},\tag{A.30}$$

where M is given by (A.3), and subsequently

$$g(\lambda) = \tilde{g}(z) + Rem, \tag{A.31}$$

with Rem satisfying (A.30), yielding (A.5)-(A.8). \Box

Proof of Theorem A.1. Let X, Y and b be the same as in Proposition A.3. We estimate the first term on the right-hand side of (A.5) as in the proof of Theorem 1.1. Similarly to (2.15), we find

$$\|\chi_X T_{\xi^z}^{-1} \tilde{U}_{t,\xi^z} T_{\xi^z} \chi_Y \| \le \|\tilde{U}_{t,\xi^z}\| e^{-\mu' d_{x_0 y_0}},\tag{A.32}$$

with $\xi^z = zb + \xi^{\perp}$ (see (2.2)), $b = \frac{x_0 - y_0}{|x_0 - y_0|}$, \tilde{U}_{t,ξ^z} defined in Eq. (A.6) and $\mu' = (1-\epsilon)\mu$. Next, applying (A.26) and setting $\tilde{c}' = \tilde{c}/(1-\epsilon)$, we conclude that

$$\|\chi_X T_{\xi^z}^{-1} \tilde{U}_{t,\xi^z} T_{\xi^z} \chi_Y \| \leqslant e^{-\mu' (d_{x_0 y_0} - \tilde{c}' t)}.$$
(A.33)

Eq. (A.5), together with estimate (A.33), yields

$$\|\chi_X U_t \chi_Y\| \lesssim_{\mu} e^{-(1-\epsilon)\mu(d_{x_0 y_0} - \tilde{c}t)} + \frac{\langle t \rangle M}{(d_{x_0 y_0} - \epsilon - t\tilde{c})^{m+1}}.$$
(A.34)

We take $\mu = \frac{1}{2}$. Then, for $1 < t < \frac{d_{x_0 y_0}}{\tilde{c}'}$, $\tilde{c}' = \tilde{c} + 2\epsilon > \tilde{c}$, we have

$$\|\chi_X e^{-itH}\chi_Y\| \lesssim \frac{tM}{(d_{x_0y_0} - t\tilde{c}')^{m+1}}$$
 (A.35)

Now, we return to general compact sets X and Y. Covering X and Y with balls $B_r(x_j)$, $j = 1, \dots, N_1$, and $B_r(y_k)$, $k = 1, \dots, N_2$ and proceeding as in

(2.20)-(2.26) and using (A.35), we obtain

$$\|\chi_X e^{-itH} \chi_Y g\|^2 \leq C \sum_{j=1}^{N_2} t^2 M^2 \|\tilde{\chi}_j g\|^2 \tilde{C}_{XY}, \qquad \forall \, g \in L^2(\mathbb{R}^n),$$
(A.36)

where

$$\tilde{C}_{XY} := \sum_{k=1}^{N_1} \sum_{l=1}^{N_2} (d_{x_k y_j} - t\tilde{c}')^{-m-1} (d_{x_k y_l} - t\tilde{c}')^{-m-1}.$$
(A.37)

Using the shell argument, as in (2.27)-(2.32), we arrive at

$$\|\chi_X e^{-itH} \chi_Y g\|^2 \lesssim_n t^2 M^2 (d_{XY} - t\tilde{c}')^{-2m-2+2n} \|g\|^2,$$
(A.38)

which yields (A.2).

APPENDIX B. *N*-PARTICLE DYNAMICS

For the N-particle problem, consider the quantum Hamiltonian for N identical bosons

$$H_N := \sum_{j=1}^N (\omega_1(p_j) + v(x_j)) + \frac{1}{2} \sum_{i \neq j} w(x_i - x_j)$$
(B.1)

on $L^2_{sym}(\mathbb{R}^{dN})$, where $L^2_{sym}(\mathbb{R}^{dN})$ is either bosonic space of symmetric functions of ferminoic one associated with certain representations of the symmetric group S_N of permutations of N indices (see e.g. [26]).

The operator H_N is of the form (1.1), with

$$\omega(k) = \sum_{j=1}^{N} \omega_1(k_j) \text{ and } V(x) = \sum_{j=1}^{N} v(x_j) + \frac{1}{2} \sum_{i \neq j} w(x_i - x_j),$$
(B.2)

and $x = (x_1, \dots, x_N) \in \mathbb{R}^{dN}$, $k = (k_1, \dots, k_N) \in \mathbb{R}^{dN}$. Now, we define $\chi_X(x)$ as the characteristic function of the set $X^N := \underbrace{X \times \dots \times X}_{X}$:

N-fold product

$$\tilde{\chi}_X(x) \equiv \chi_{X^N}(x) := \prod_{j=1}^N \chi_X(x_j).$$
(B.3)

Theorem B.1. Let H_N be as in Eq. (B.1) and let ω_1 and V (defined in Eq. (B.2)) satisfy Condition (1.8) (with n = d). Let X and Y be two disjoint sets. Then, for any $\mu' \in (0, \mu)$, we have

$$\|\tilde{\chi}_X e^{-itH_N} \tilde{\chi}_Y\| \leqslant C e^{-\mu' N(d_{XY} - c_1't)},\tag{B.4}$$

where X is the multiplication operator by the function $\chi_X(x)$ in (B.3), C > 0 is a constant depending on $\epsilon = 1 - \frac{\mu'}{\mu}$, d, N and μ , with $c'_1 = \frac{c_1}{1-\epsilon}$ and

$$c_1 := \sup_{\xi \in \mathbb{R}^d, \ b \in S^{d-1}} \frac{\operatorname{Im}\omega_1(\xi + i\mu b)}{\mu} < \infty.$$
(B.5)

Eq. (B.4) is one of the simplest many-body estimates. More refined and more difficult estimate to prove would be one with $\chi_X(x)$ and $\chi_Y(y)$ replaced in (B.4) by $\chi_X^{(k)}(x) \equiv \chi_{X^k}(x) = \prod_{j=1}^k \chi_X(x_j)$ and $\chi_Y^{(l)}(y) \equiv \chi_{Y^l}(y) = \prod_{j=1}^l \chi_Y(y_j)$, with $1 \leq k \leq l < N$.

Proof of Theorem B.1. We follow the proof of Theorem 1.1, with the following modifications:

$$\xi = (\xi_1, \cdots, \xi_N) \in \mathbb{R}^{dN}, \, \xi_j \in \mathbb{R}^d, \, b = (b_1, \cdots, b_N), b_j \in S^{n-1}, \, \forall j, \qquad (B.6)$$

$$\zeta = (\zeta_1, \cdots, \zeta_N) \in \mathbb{R}^{dN}, \, \zeta_j = \xi_j + i\mu b_j, \, \mu \in (0, a), \, b \cdot x = \sum_{j=1}^N b_j \cdot x_j. \quad (B.7)$$

As in the proof of Theorem 1.1, we reduce proving Eq. (B.4) for disjoint sets X and Y to Eq. (B.4) for the polyballs $X = \prod_{j=1}^{N} X_j, X_j := B_r(x_{0j} - y_{0j})$ and $Y = \prod_{j=1}^{N} Y_0, Y_0 := B_r(0)$, with $r = \frac{\epsilon}{2} d_{XY}$, with $x_{0j} \in X$ and $y_{0j} \in Y, \forall j$, and $\epsilon \in (0, 1)$.

Similarly to (2.11) and (2.16), we have

$$\|\tilde{\chi}_X e^{-iH_N t} \tilde{\chi}_Y\| \leqslant e^{-\mu(r_X - r_Y)} e^{\mu tc},\tag{B.8}$$

with

$$c = \frac{1}{\mu} \sup_{\substack{b \in \bigotimes_{1} S^{d-1}}} \sup_{\xi \in \mathbb{R}^{dN}} \operatorname{Im} \sum_{j=1}^{N} \omega(\xi_j + i\mu b_j),$$
(B.9)

$$r_Y = \sup_{y \in Y} b \cdot y, \quad r_X = \inf_{x \in X} b \cdot x.$$
(B.10)

Let $b = (b_1, \dots, b_N)$, with $b_j = \frac{x_{0j} - y_{0j}}{|x_{0j} - y_{0j}|}, j = 1, \dots, N$. We claim that

$$r_X - r_Y \ge (1 - \epsilon) \sum_{j=1}^N |x_{0j} - y_{0j}| \ge (1 - \epsilon) N d_{XY}.$$
 (B.11)

Indeed, by the definitions of X and Y and Eq. (B.10), we have

$$r_Y \leqslant \sum_{j=1}^N |y_j| \leqslant \frac{\epsilon N}{2} d_{XY} \tag{B.12}$$

and

$$r_X = \sum_{j=1}^N b \cdot (x_{0j} - y_{0j}) + \inf_{x \in X} b \cdot (x - (x_{0j} - y_{0j}))$$

$$\geq \sum_{j=1}^N (|x_{0j} - y_{0j}| - \frac{\epsilon}{2} d_{XY}).$$
(B.13)

Hence,

$$r_X - r_Y \ge \sum_{j=1}^N (|x_{0j} - y_{0j}| - \epsilon d_{XY}) \ge (1 - \epsilon) \sum_{j=1}^N |x_{0j} - y_{0j}|,$$
(B.14)

as claimed.

Furthermore, we compute

$$c = \frac{1}{\mu} \sum_{j=1}^{N} \sup_{\xi_j \in \mathbb{R}^d, \, b_j \in S^{d-1}} \operatorname{Im}\omega_1(\xi_j + i\mu b_j) = Nc_1,$$
(B.15)

where, recall,

$$c_1 := \sup_{\lambda \in \mathbb{R}^n, b \in S^{n-1}} \frac{\omega_1(\lambda + i\mu b)}{\mu}.$$
(B.16)

Combining (B.8), (B.11) and (B.15), we arrive at

$$\|\tilde{\chi}_X e^{-itH_N} \tilde{\chi}_Y\| \leqslant e^{-\mu(1-\epsilon)N(d_{XY}-c_1't)},\tag{B.17}$$

where $c'_1 = \frac{c_1}{1-\epsilon}$, which implies, with $\mu'' = (1-\epsilon)\mu$,

$$\|\tilde{\chi}_X e^{-itH_N} \tilde{\chi}_Y\| \leqslant e^{-\mu'' N(d_{XY} - c_1't)},\tag{B.18}$$

with $X = \prod_{j=1}^{N} X_j$, $X_j := B_r(x_{0j})$ and $Y = \prod_{j=1}^{N} Y_0$, $Y_0 := B_r(y_{0j})$, where $r = \frac{\epsilon}{2} d_{XY}$, for some $x_0 \neq y_0$, with $x_{0j} \in X$ and $y_{0j} \in Y$, $\forall j$, and $\epsilon \in (0, 1)$. Proceeding as in (2.20)-(2.35) in each variable $x_j \in \mathbb{R}^d$, $j = 1, \dots, N$, we arrive at (1.5) by taking $\mu' = \mu(1 - 2\epsilon)$ for any $\epsilon \in (0, 1)$, and the constant C in (B.4) depends on $\epsilon = 1 - \frac{\mu'}{\mu}, d, N$ and μ .

References

- [1] J. Arbunich, J. Faupin, F. Pusateri, and I. M. Sigal, *Maximal speed of quantum propagation* for the Hartree equation, Comm. Partial Differential Equations 48 (2023), no. 4.
- [2] J. Arbunich, F. Pusateri, I. M. Sigal, and A. Soffer, Maximal speed of quantum propagation, Lett. Math. Phys. 111 (2021), 1–16.
- [3] J.-F. Bony, J. Faupin, and I. M. Sigal, Maximal velocity of photons in non-relativistic QED, Adv. Math 231 (2012), no. 5, 3054–3078.
- [4] S. Bose, Quantum communication through spin chain dynamics: an introductory overview, Contemporary Physics 48 (2007), no. 1, 13–30.
- [5] O. Bratteli and D. W. Robinson, Operator algebras and quantum statistical mechanics. 2, Second, Texts and Monographs in Physics, Springer-Verlag, Berlin, 1997.
- S. Bravyi and M. B. Hastings, Topological quantum order: Stability under local perturbations, J. Math. Phys. 51 (2010), no. 9.
- [7] S. Bravyi, M. B. Hastings, and F. Verstraete, Lieb-Robinson bounds and the generation of correlations and topological quantum order, Phys. Rev. Lett. 97 (2006), no. 5, 050401.
- [8] S. Breteaux, J. Faupin, M. Lemm, and I. M. Sigal, Maximal speed of propagation in open quantum systems, The Physics and Mathematics of Elliott Lieb, 2022, pp. 109–130.
- [9] S. Breteaux, J. Faupin, M. Lemm, D. Ouyang, I. M. Sigal, and J. Zhang, *Light cones for open quantum systems in the continuum*, Reviews in Mathematical Physics (2024).
- [10] C. Cedzich, A. Joye, A.H. Werner, and R.F. Werner, Exponential tail estimates for quantum lattice dynamics, Preprint, arXiv 2408.02108v1 (2024).
- [11] C.-F. Chen and A. Lucas, Finite speed of quantum scrambling with long range interactions, Phys. Rev. Lett. 123 (2019), no. 25, 250605.
- [12] N. Defenu, T. Donner, T. Macrì, G. Pagano, S. Ruffo, and A. Trombettoni, Long-range interacting quantum systems, Rev. Mod. Phys. 95 (2023), no. 3, 035002.
- [13] J. Dereziński and C. Gérard, Asymptotic completeness in quantum in field theory: massive Pauli-Fierz hamiltonians, Rev. Math. Phys. 11 (1999), no. 04, 383–450.
- [14] J. Eisert and T. J. Osborne, General entanglement scaling laws from time evolution, Phys. Rev. Lett. 97 (2006), no. 15, 150404.
- [15] Z. Eldredge, Z.-X. Gong, J. T Young, A. H. Moosavian, M. Foss-Feig, and A. V. Gorshkov, Fast quantum state transfer and entanglement renormalization using long-range interactions, Phys. Rev. Lett. 119 (2017), no. 17, 170503.
- [16] D. V. Else, F. Machado, Ch. Nayak, and N. Y. Yao, Improved Lieb-Robinson bound for many-body Hamiltonians with power-law interactions, Phys. Rev. A 101 (2020), no. 2, 022333.
- [17] J. M. Epstein and K. B. Whaley, Quantum speed limits for quantum-information-processing task, Phys. Rev. A 95 (2017), no. 4, 042314.
- [18] J. Faupin, M. Lemm, and I. M. Sigal, Maximal speed for macroscopic particle transport in the Bose-Hubbard model, Phys. Rev. Lett. 128 (2022), no. 15, 150602.
- [19] _____, On Lieb-Robinson bounds for the Bose-Hubbard model, Commun. Math. Phys. 394 (2022), no. 3, 1011–1037.
- [20] J. Faupin and I. M. Sigal, On Rayleigh scattering in non-relativistic quantum electrodynamics, Commun. Math. Phys. 328 (2014), 1199–1254.
- [21] O. Fawzi and R. Renner, Quantum conditional mutual information and approximate Markov chains, Commun. Math. Phys. 340 (2015), no. 2, 575–611.
- [22] M. Foss-Feig, Z. Gong, C. W. Clark, and A. V. Gorshkov, Nearly-linear light cones in long-range interacting quantum systems, Phys. Rev. Lett. 114 (2015), no. 15, 157201.
- [23] J. Fröhlich, M. Griesemer, and B. Schlein, Asymptotic completeness for Rayleigh scattering, Ann. Henri Poincaré 3 (2002), 107–170.

- [24] M. Gebert, B. Nachtergaele, J. Reschke, and R. Sims, Lieb-Robinson bounds and strongly continuous dynamics for a class of many-body fermion systems in R^d, Ann. Henri Poincaré 21 (2020), no. 11, 3609–3637.
- [25] C. Gogolin and J. Eisert, Equilibration, thermalisation, and the emergence of statistical mechanics in closed quantum systems, Rep. Prog. Phys. 79 (2016), no. 5, 056001.
- [26] S. J. Gustafson and I. M. Sigal, Mathematical Concepts of Quantum Mechanics, 3-rd ed., Universitext, Springer, 2020.
- [27] W. A. Harrison, Electronic structure and the properties of solids: the physics of the chemical bond, Courier Corporation, 2012.
- [28] M. B. Hastings, Lieb-Schultz-Mattis in higher dimensions, Phys. Rev. B 69 (2004), no. 10, 104431.
- [29] _____, Locality in quantum and Markov dynamics on lattices and networks, Phys. Rev. Lett. **93** (2004), no. 14, 140402.
- [30] _____, An area law for one-dimensional quantum systems, J. Stat. Mech. Theory Exp. 8 (2007), P08024, 14.
- [31] _____, Quantum belief propagation, an algorithm for thermal quantum systems, Phys. Rev. B **76** (2007), no. 20, 201102(R).
- [32] M. B. Hastings and T. Koma, Spectral gap and exponential decay of correlations, Commun. Math. Phys. 265 (2006), 781–804.
- [33] I. Herbst and E. Skibsted, Free channel Fourier transform in the long-range n-body problem., J. d'Analyse Math 65 (1995), no. 1, 297–332.
- [34] W. Hunziker and I. M. Sigal, n-body quantum problem, J. Math. Phys. 41 (2000), no. 6, 3448–3510.
- [35] W. Hunziker, I. M. Sigal, and A. Soffer, *Minimal escape velocities*, Comm. Partial Differential Equations 24 (1999), no. 11-12, 2279–2295.
- [36] M. Kliesch, C. Gogolin, and J. Eisert, Lieb-Robinson bounds and the simulation of timeevolution of local observables in lattice systems, In Many-Electron Approaches in Physics, Chemistry and Mathematics, 301-318. Springer, 2014.
- [37] T. Kuwahara and Lemm. M., Enhanced Lieb-Robinson bounds for a class of Bose-Hubbard type Hamiltonians, Preprint, arXiv 2405.04672 (2024).
- [38] T. Kuwahara and K. Saito, Absence of fast scrambling in thermodynamically stable longrange interacting systems, Phys. Rev. Lett. 126 (2021), no. 3, 030604.
- [39] T. Kuwahara, T. V. Vu, and K. Saito, Optimal light cone and digital quantum simulation of interacting bosons, Preprint, arXiv 2206.14736 (2022).
- [40] T. Kuwahara and K. Saito, Strictly linear light cones in long-range interacting systems of arbitrary dimensions, Phys. Rev. X 10 (2020Jul), no. 3, 031010.
- [41] _____, Lieb-Robinson bound and almost-linear light cone in interacting boson systems, Phys. Rev. Lett. **127** (2021), no. 7, 070403.
- [42] M. Lemm, C. Rubiliani, I. M. Sigal, and J. Zhang, Information propagation in long-range quantum many-body systems, Phys. Rev. A 108 (2023), no. 6, L060401.
- [43] E. H. Lieb and D. W. Robinson, The finite group velocity of quantum spin systems, Commun. Math. Phys. 28 (1972), no. 3, 251–257.
- [44] T. Matsuta, T. Koma, and T. Nakamura, Improving the Lieb-Robinson bound for long-range interactions, Ann. Henri Poincaré 18 (2017), 519–528.
- [45] B. Nachtergaele, Y. Ogata, and R. Sims, Propagation of correlations in quantum lattice systems, J. Stat. Phys. 124 (2006), 1–13.
- [46] B. Nachtergaele, B. Schlein, R. Sims, S. Starr, and V. Zagrebnov, On the existence of the dynamics for anharmonic quantum oscillator systems, Rev. Math. Phys. 22 (2010), no. 02, 207–231.

- [47] B. Nachtergaele and R. Sims, Lieb-Robinson bounds and the exponential clustering theorem, Commun. Math. Phys. 265 (2006), 119–130.
- [48] B. Nachtergaele, R. Sims, and A. Young, Quasi-locality bounds for quantum lattice systems.
 i. Lieb-Robinson bounds, quasi-local maps, and spectral flow automorphisms, J. Math. Phys.
 60 (2019), no. 6, 061101.
- [49] B. Nachtergaele, A. Vershynina, and V. Zagrebnov, Lieb-Robinson bounds and existence of the thermodynamic limit for a class of irreversible quantum dynamics, Contemp. Math 552 (2011), 161–175.
- [50] B. Nachtergaele, H. Raz, B. Schlein, and R. Sims, Lieb-Robinson bounds for harmonic and anharmonic lattice systems, Comm. Math. Phys. 286 (2009), no. 3, 1073–1098. MR2472026
- [51] B. Nachtergaele and R. Sims, Much ado about something: Why Lieb-Robinson bounds are useful, Preprint, arXiv 1102.0835 (2010).
- [52] D. Poulin, Lieb-Robinson bound and locality for general Markovian quantum dynamics, Phys. Rev. Lett. 104 (2010), no. 19, 190401.
- [53] M. Reed and B. Simon, Methods of modern mathematical physics. IV. Analysis of operators, Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York, 1978.
- [54] D. A. Roberts and B. Swingle, Lieb-Robinson bound and the butterfly effect in quantum field theories, Phys. Rev. Lett. 117 (2016), no. 9, 091602.
- [55] W. Schlag, On pointwise decay of waves, J. Math. Phys. 62 (2021), no. 6, 061509.
- [56] N. Schuch, S. K. Harrison, T. J. Osborne, and J. Eisert, Information propagation for interacting-particle systems, Phys. Rev. A 84 (2011), no. 3, 032309.
- [57] I. M. Sigal, On long range scattering, Duke Math. J. 60 (1990), 473–496.
- [58] I. M. Sigal and A. Soffer, Long-range many-body scattering, Invent. Math 99 (1990), no. 1, 115–143.
- [59] I. M. Sigal and J. Zhang, On propagation of information in quantum many-body systems, Preprint, arXiv 2212.14472 (2023).
- [60] E. Skibsted, Propagation estimates for N-body Schrödinger operators, Commun. Math. Phys. 142 (1991), no. 1, 67–98.
- [61] M. C. Tran, Yuan Su, D. Carney, and J. M. Taylor, Faster digital quantum simulation by symmetry protection, PRX QUANTUM 2 (2021), no. 1, 010323.
- [62] M. C. Tran, C.-F. Chen, A. Ehrenberg, A. Y. Guo, A. Deshpande, Y. Hong, Z.-X. Gong, A. V. Gorshkov, and A. Lucas, *Hierarchy of linear light cones with long-range interactions*, Phys. Rev. X **10** (2020), no. 3, 031009.
- [63] M. C. Tran, A. Y. Guo, C. L. Baldwin, A. Ehrenberg, A. V. Gorshkov, and A. Lucas, *Lieb-Robinson light cone for power-law interactions*, Phys. Rev. Lett. **127** (2021), no. 16, 160401.
- [64] M. C. Tran, A. Y. Guo, A. Deshpande, A. Lucas, and A. V. Gorshkov, Optimal state transfer and entanglement generation in power-law interacting systems, Phys. Rev. X 11 (2021), no. 3, 031016.
- [65] Z. Wang and K. R. Hazzard, Tightening the Lieb-Robinson Bound in Locally Interacting Systems, PRX Quantum 1 (2020), no. 1, 010303.
- [66] C. Yin and A. Lucas, Finite speed of quantum information in models of interacting bosons at finite density, Phys. Rev. X 12 (2022), no. 2, 021039.

Department of Mathematics, University of Toronto, Toronto, ON M5S 2E4, Canada

Email address: im.sigal@utoronto.ca

Mathematical Sciences Institute, Australian National University, Acton ACT 2601, Australia

Email address: Xiaoxu.Wu@anu.edu.au