

# Bi-slant Riemannian maps to Kenmotsu manifolds and some optimal inequalities

Adeeba Zaidi<sup>1</sup> and Gauree Shanker<sup>1\*</sup>

<sup>1</sup>Department of Mathematics and Statistics, Central University of Punjab, Bathinda, 151401, Punjab, India.

\*Corresponding author(s). E-mail(s): [gauree.shanker@cup.edu.in](mailto:gauree.shanker@cup.edu.in);  
Contributing authors: [adeebazaidi.az25@gmail.com](mailto:adeebazaidi.az25@gmail.com);

## Abstract

In this paper, we introduce bi-slant Riemannian maps from Riemannian manifolds to Kenmotsu manifolds which are the natural generalizations of invariant, anti-invariant, semi-invariant, slant, semi-slant and hemi-slant Riemannian maps, with nontrivial examples. We study these maps and give some curvature relations for  $(\text{range } F_*)^\perp$ . We construct Chen-Ricci inequalities, DDVV inequalities and further some optimal inequalities involving Casorati curvatures from bi-slant Riemannian manifolds to Kenmotsu space forms.

**Keywords:** Contact Manifolds, Kenmotsu manifolds, Riemannian maps, bi-slant Riemannian maps, Chen-Ricci inequality, DDVV conjecture, Casorati curvature.

**MSC Classification:** 53B20 , 53C25 , 53D10 , 54C05

## 1 Introduction

In the field of submanifold theory, it is quite interesting to know the relationship between intrinsic and extrinsic properties of a submanifold. In Riemannian geometry, the intrinsic properties of Riemannian manifolds are essentially the Riemannian invariants, which in turn influence the general form behavior of the manifold. In 1993, Chen [1] established the relationship between intrinsic and extrinsic invariants. This relationship takes the form of various inequalities and connects the main intrinsic and extrinsic invariants. Chen [2] also established a relationship between the squared mean curvature and Ricci curvature of a submanifold in the form of an inequality, known as

the Chen-Ricci inequality. This was the beginning of the Chen invariants and Chen-Ricci relations, which have since become a topic of interest among geometers (see[3–5] etc). Later, In 2005, Chen proved optimal relationship between Riemannian submersions and minimal immersions in [6] and [7]. After that many optimal inequalities relations on Riemannian submersions are investigated by various geometers (see[8–10]etc).

On the other hand, instead of concentrating on the sectional curvature with the extrinsic squared mean curvature, Casorati introduced Casorati curvature as the normalized square of the length of the second fundamental form. The notion of Casorati curvature extends the concept of the principal direction of a hypersurface of a Riemannian manifold. Given the critical role and significance of Casorati curvature in the visual perception of shapes and the appearance of manifolds, many authors have taken an interest in obtaining optimal inequalities for the Casorati curvatures of submanifolds in various ambient spaces (see [11–13] etc). In 2020, Lee et al [14] obtained optimal inequalities for Riemannian maps and Riemannian submersions for complex space forms. Recently, Gündüzalp et al studied Casorati inequalities for pointwise slant Riemannian maps [15], pointwise hemi-slant Riemannian maps [16], pointwise semi-slant Riemannian maps [17] from complex space forms.

Motivated by these work, we study bi-slant Riemannian maps from Riemannian manifolds to Kenmotsu manifolds and established some inequalities. This article is organized as follows. Section 2 covers the basic terminologies, definitions, and properties of Riemannian maps and Kenmotsu manifolds that are essential throughout this article. Section 3 introduces the definition of bi-slant Riemannian maps from Riemannian manifolds to Kenmotsu manifolds, considering both cases when the Reeb vector field is in  $range F_*$  and  $(range F_*)^\perp$ . Examples of such maps are constructed, followed by a study of some properties and an inequality related to these maps. In Section 4, we derive Ricci-Chen inequalities for bi-slant Riemannian maps from Riemannian manifolds to Kenmotsu space forms for both cases of Reeb vector fields and study DDVV inequality for such maps. In Section 5, we establish optimal inequalities for the same maps from Riemannian manifolds to Kenmotsu space forms involving Casorati curvatures, and we also discuss the inequalities for subcases for these maps.”.

## 2 Preliminaries

In this section, basic definitions and terminologies are recalled that are needed throughout this article.

A  $(2m + 1)$ - dimensional differentiable manifold  $M$  is said to have an almost-contact metric structure  $(\psi, \xi, \eta, g)$  if it admits a tensor field  $\psi$  of type  $(1, 1)$ , a vector field (characteristic vector field or Reeb vector field)  $\xi$ , a 1-form  $\eta$  and a Riemannian metric  $g$ , satisfying [18]

$$\psi^2 = -I + \eta \otimes \xi, \quad \psi\xi = 0, \quad \eta \circ \psi = 0, \quad \eta(\xi) = 1, \quad (1)$$

$$g(\psi X, \psi Y) = g(X, Y) - \eta(X)\eta(Y), \quad (2)$$

$$g(\psi X, Y) = -g(X, \psi Y), \quad \eta(X) = g(X, \xi) \quad (3)$$

for any vector fields  $X, Y \in \Gamma(TM)$ . Moreover,  $M$  with  $(\psi, \xi, \eta, g)$  is called almost contact metric manifold. An almost contact metric manifold  $M$  is called a Kenmotsu manifold, if [19]

$$(\nabla_X \psi)Y = g(\psi X, Y)\xi - \eta(Y)\psi X, \quad (4)$$

$$\nabla_X \xi = X - \eta(X)\xi, \quad (5)$$

where  $\nabla$  is the Levi-Civita connection of the Riemannian metric  $g$  on  $M$ . Also, for a Kenmotsu manifold, we have

$$R(\psi X, \psi Y)Z - R(X, Y)Z = g(Y, Z)X - g(X, Z)Y + g(Y, \psi Z)\psi X - g(X, \psi Z)\psi Y. \quad (6)$$

A Kenmotsu manifold with constant  $\psi$ -holomorphic sectional curvature  $c$  is called a Kenmotsu space form and denoted by  $M(c)$ . The curvature tensor  $R$  of a Kenmotsu space form  $M(c)$  is given by, [19]

$$\begin{aligned} R(X, Y)Z = & \frac{c-3}{4} [g(Y, Z)X - g(X, Z)Y] + \frac{c+1}{4} [\eta(X)\eta(Z)Y - \eta(Y)\eta(Z)X \\ & + \eta(Y)g(X, Z)\xi - \eta(X)g(Y, Z)\xi - g(\psi X, Z)\psi Y + g(\psi Y, Z)\psi X \\ & + 2g(\psi Y, X)\psi Z] \end{aligned} \quad (7)$$

for all  $X, Y \in \Gamma(TM)$ .

Further, consider  $F : (\overline{M}^{\overline{m}}, g_1) \rightarrow (M^m, g_2)$  be a smooth map between smooth finite dimensional Riemannian manifolds  $\overline{M}^{\overline{m}}$  and  $M^m$  such that  $0 < \text{rank} F < \min\{\overline{m}, m\}$  and if  $F_* : T_p \overline{M} \rightarrow T_{F(p)} M$  denotes the differential map at  $p \in \overline{M}$ , and  $F(p) \in M$ , then  $T_p \overline{M}$  and  $T_{F(p)} M$  split orthogonally with respect to  $g_1(p)$  and  $g_2(F(p))$ , respectively, as [20]

$$T_p \overline{M} = \ker F_{*p} \oplus (\ker F_{*p})^\perp = \mathcal{V}_p \oplus \mathcal{H}_p,$$

where  $\mathcal{V}_p = \ker F_{*p}$  and  $\mathcal{H}_p = (\ker F_{*p})^\perp$  are vertical and horizontal parts of  $T_p \overline{M}$  respectively. Since  $0 < \text{rank} F < \min\{\overline{m}, m\}$ , we have  $(\text{range} F_{*p})^\perp \neq 0$ ; therefore,  $T_{F(p)} M$  can be written as  $T_{F(p)} M = \text{range} F_{*p} \oplus (\text{range} F_{*p})^\perp$ , then the map  $F : (\overline{M}^{\overline{m}}, g_1) \rightarrow (M^m, g_2)$  is called a Riemannian map at  $p \in \overline{M}$ , if  $g_2(F_* X, F_* Y) = g_1(X, Y)$  for all vector fields  $X, Y \in \Gamma(\ker F_{*p})^\perp$ . The second fundamental form of the map  $F$  is given by [21]

$$(\nabla F_*)(X, Y) = \overset{M}{\nabla_X^F} F_* Y - F_*(\nabla_X^{\overline{M}} Y), \quad (8)$$

where  $\overset{M}{\nabla_X^F} F_* Y \circ F = \nabla_{F_* X}^M F_* Y$ , while  $\nabla^M$  is the Levi-Civita connection on  $M$  and  $\overset{M}{\nabla^F}$  is the pullback connection of  $\nabla^M$  along  $F$ , provided that  $(\nabla F_*)(X, Y)$  has no components in  $\text{range} F_*$ , if  $X, Y \in \Gamma((\ker F_*)^\perp)$ .

Let  $\nabla^{F^\perp}$  be a linear connection on  $(F_*(T\overline{M}))^\perp$  and  $\nabla_X^{F^\perp}V$  be the orthogonal projection of  $\nabla_X^M V$  onto  $(range F_*)^\perp$ ,  $X \in \Gamma((ker F_*)^\perp)$ ,  $V \in \Gamma((range F_*)^\perp)$ , then [21]

$$\nabla_X^M V = -S_V F_* X + \nabla_X^{F^\perp} V, \quad (9)$$

where  $-S_V F_* X$  is the tangential component of  $\nabla_X^M V$ . Noted that  $\nabla_X^M V$  is obtained from the pullback connection of  $\nabla^M$ . It can be seen that  $(S_V F_* X)_p$  depends only on  $V_p$  and  $F_{*p}(X_p)$ , and  $S_V F_* X$  is bi-linear in  $V$  and  $F_* X$ . By direct computations, we have

$$g_2(S_V F_* X, F_* Y) = g_2(V, (\nabla F_*)(X, Y)) \quad (10)$$

for  $X, Y \in \Gamma((ker F_*)^\perp)$  and  $V \in \Gamma((range F_*)^\perp)$ . The curvature tensor  $R^{F^\perp}$  of the sub-bundle  $(range F_*)^\perp$  is defined by

$$R^{F^\perp}(F_*(X), F_*(Y))V = \nabla_X^{F^\perp} \nabla_Y^{F^\perp} V - \nabla_Y^{F^\perp} \nabla_X^{F^\perp} V - \nabla_{[X, Y]}^{F^\perp} V.$$

Further, let  $R^{\overline{M}}$ ,  $R^M$  and  $R^{F^\perp}$  be the curvature tensors of  $\nabla^{\overline{M}}$ ,  $\nabla^M$  and  $(range F_*)^\perp$ , respectively. Then we have the Gauss equation and the Ricci equation for the Riemannian map  $F$  given by [21]

$$g_2(R^M(F_* X, F_* Y)F_* Z, F_* H) = g_1(R^{\overline{M}}(X, Y)Z, H) + g_2((\nabla F_*)(X, Z), (\nabla F_*)(Y, H)) - g_2((\nabla F_*)(Y, Z), (\nabla F_*)(X, H)), \quad (11)$$

$$g_2(R^M(F_* X, F_* Y)V_1, V_2) = g_2(R^{F^\perp}(F_* X, F_* Y)V_1, V_2) + g_2([S_{V_2}, S_{V_1}]F_* X, F_* Y), \quad (12)$$

where  $X, Y, Z, H \in \Gamma((ker F_*)^\perp)$ ,  $V_1, V_2 \in \Gamma((range F_*)^\perp)$  and  $[S_{V_2}, S_{V_1}] = S_{V_2} S_{V_1} - S_{V_1} S_{V_2}$ . Let  $\{e_1, \dots, e_r\}$  and  $\{v_{r+1}, \dots, v_s\}$  be the orthonormal basis of  $(ker F_*)^\perp_p$  and  $(range F_*)^\perp_{F_* p}$ ,  $p \in \overline{M}$ , respectively. Then the Ricci curvature  $\mathcal{R}ic^{(ker F_*)^\perp}$ , scalar curvature  $\tau^{(ker F_*)^\perp}$  and normalized scalar curvature  $\varrho^{(ker F_*)^\perp}$  on  $((ker F_*)^\perp)_p$  are given by

$$\begin{aligned} \mathcal{R}ic^{(ker F_*)^\perp}(X) &= \sum_{i=1}^r g_1(R^{\overline{M}}(e_i, X)X, e_i), \quad \forall X \in (ker F_*)^\perp, \\ \tau^{(ker F_*)^\perp} &= \sum_{1 \leq i < j \leq r} g_1(R^{\overline{M}}(e_i, e_j)e_j, e_i), \\ \varrho^{(ker F_*)^\perp} &= \frac{2\tau^{(ker F_*)^\perp}}{r(r-1)}, \end{aligned}$$

whereas, a new scalar curvature  $\tau^\perp$  of sub-bundle  $(range F_*)^\perp$  called normal scalar curvature, is defines as

$$\tau^{(range F_*)^\perp} = \sqrt{\sum_{1 \leq i < j \leq r} \sum_{1 \leq \alpha < \beta \leq m} g_2(R^\perp(e_i, e_j)v_\alpha, v_\beta)^2}$$

and normalized normal scalar curvature  $\varrho^{(range F_*)^\perp}$  is given by

$$\varrho^{(range F_*)^\perp} = \frac{2\tau^{(range F_*)^\perp}}{r(r-1)}. \quad (13)$$

Also, for the Riemannian map  $F$ , we can write [22]

$$\zeta_{ij}^\alpha = g_2((\nabla F_*)(e_i, e_j), v_\alpha), \quad i, j = 1, \dots, r, \quad \alpha = r+1, \dots, m \quad (14)$$

$$\|\zeta\|^2 = \sum_{i,j=1}^r g_2((\nabla F_*)(e_i, e_j), (\nabla F_*)(e_i, e_j)), \quad (15)$$

$$trace \zeta = \sum_{i=1}^r (\nabla F_*)(e_i, e_i), \quad (16)$$

$$\|trace \zeta\|^2 = g_2(trace \zeta, trace \zeta). \quad (17)$$

### 3 Bi-slant Riemannian Maps to Kenmotsu manifolds

In this section, we define bi-slant Riemannian maps from Riemannian manifolds to Kenmotsu manifolds. We study the geometry of these maps and construct some non trivial examples.

**Definition 1.** Let  $F : (\overline{M}, g_1) \rightarrow (M, \psi, \xi, \eta, g_2)$  be a Riemannian map from a Riemannian manifold  $\overline{M}$  to Kenmotsu manifold  $M$  of dimensions  $\overline{m}$  and  $m$ , respectively. The map  $F$  is said to be bi-slant Riemannian map if there exists two slant distribution  $\mathcal{D}_{\theta_1}$  and  $\mathcal{D}_{\theta_2}$  in  $range F_*$ , such that

1. if  $\xi \in \Gamma(range F_*)$ ,

$$range F_* = \mathcal{D}_{\theta_1} \oplus \mathcal{D}_{\theta_2} \oplus \{\xi\}.$$

In this case, for  $X \in \Gamma((ker F_*)^\perp)$ ,  $F_*(X) = \sigma F_*(X) + \pi F_*(X) + \eta(U)\xi$ ;

2. if  $\xi \in \Gamma((range F_*)^\perp)$ ,

$$range F_* = \mathcal{D}_{\theta_1} \oplus \mathcal{D}_{\theta_2}.$$

In this case, for  $X \in \Gamma((ker F_*)^\perp)$ ,  $F_*(X) = \sigma F_*(X) + \pi F_*(X)$ ,

where  $\psi \mathcal{D}_{\theta_i} \perp \mathcal{D}_{\theta_j}$  for  $i \neq j = 1, 2$ , and  $\sigma, \pi$  are orthogonal projections of  $F_*X$  on  $\mathcal{D}_{\theta_1}$  and  $\mathcal{D}_{\theta_2}$ , respectively. Whereas,  $\theta_1, \theta_2$  are the slant angles for distributions  $\mathcal{D}_{\theta_1}$  and  $\mathcal{D}_{\theta_2}$ , respectively.

If the dimensions of  $\mathcal{D}_{\theta_1}$  and  $\mathcal{D}_{\theta_2}$  are  $2r_1$  and  $2r_2$ , then we have the following particular cases of Riemannian maps to Kenmotsu manifolds:

Cases	Dimensions of the distributions	$\theta_1$	$\theta_2$	Riemannian maps
1	$2r_1 = 0$ and $2r_2 \neq 0$	-	0	Invariant
2	$2r_1 = 0$ and $2r_2 \neq 0$	-	$\pi/2$	Anti-invariant
3	$2r_1, 2r_2 \neq 0$	0	$\pi/2$	Semi-invariant
4	$2r_1 = 0$ and $2r_2 \neq 0$	0	$\in (0, \pi/2)$	Proper slant
5	$2r_1 = 0, 2r_2 \neq 0$	0	$\in (0, \pi/2)$	Semi-slant
6	$2r_1 = 0, 2r_2 \neq 0$	$\pi/2$	$\in (0, \pi/2)$	Hemi-slant

However, a bi-slant Riemannian map is said to be proper, if  $0 < \theta_1, \theta_2 < \pi/2$ . Again, If  $X \in \Gamma(\ker F_*)$ , then

$$\psi F_* X = P F_* X + Q F_* X, \quad (18)$$

where,  $P F_* X \in \Gamma(\text{range } F_*)$  and  $Q F_* X \in \Gamma((\text{range } F_*)^\perp)$ . If  $V \in \Gamma((\text{range } F_*)^\perp)$ , then

$$\psi V = \phi V + \omega V, \quad (19)$$

where  $\phi U \in \Gamma(\text{range } F_*)$  and  $\omega U \in \Gamma((\text{range } F_*)^\perp)$ . Also,  $(\text{range } F_*)^\perp = Q \mathcal{D}_{\theta_1} \oplus Q \mathcal{D}_{\theta_2} \oplus \mu$ , where  $\mu$  is the complementary distribution to  $Q \mathcal{D}_{\theta_1} \oplus Q \mathcal{D}_{\theta_2}$  in  $(\text{range } F_*)^\perp$ .

**Example 1.** Let  $M = \{(u_1, u_2, u_3, v_1, v_2, v_3, w) \in \mathbb{R}^7 \mid w \neq 0\}$  be a Riemannian manifold with contact metric structure  $(\psi, \xi, \eta, g_2)$  define by  $\eta = dz$ ,  $\xi = \frac{\partial}{\partial w}$ ,  $\psi(\frac{\partial}{\partial u_1}) = \frac{\partial}{\partial u_2}$ ,  $\psi(\frac{\partial}{\partial u_2}) = -\frac{\partial}{\partial u_1}$ ,  $\psi(\frac{\partial}{\partial u_3}) = \frac{\partial}{\partial v_3}$ ,  $\psi(\frac{\partial}{\partial v_3}) = -\frac{\partial}{\partial u_3}$ ,  $\psi(\frac{\partial}{\partial v_1}) = \frac{\partial}{\partial v_2}$ ,  $\psi(\frac{\partial}{\partial v_2}) = -\frac{\partial}{\partial v_1}$ ,  $\psi(\frac{\partial}{\partial w}) = 0$  and  $g_2 = du_1^2 + \frac{4}{9}(du_2^2 + du_3^2) + \frac{3}{16}(dv_2 + 3dv_2^2) + dv_3^2 + dw^2$ . Then, from (1)–(5) we can see that  $M$  is a Kenmotsu manifold. Further, let  $F : (\overline{M} = \mathbb{R}^7, g_1) \rightarrow (M, \psi, \xi, \eta, g_2)$  be a Riemannian map defines by

$$F_*(x_1, x_2, x_3, x_4, x_5, x_6, x_7) = (x_1, x_3 - \frac{1}{\sqrt{2}}x_4, \frac{1}{\sqrt{2}}x_3 - \frac{1}{2}x_4, -\frac{1}{\sqrt{3}}x_5 + x_6, \frac{1}{\sqrt{6}}x_5 - \frac{1}{\sqrt{2}}x_6, 0, x_7),$$

where  $\overline{M}$  is a Riemannian manifold with Cartesian co-ordinates  $(x_1, x_2, x_3, x_4, x_5, x_6, x_7)$ , and  $g_1$  is a usual basis on  $\mathbb{R}^7$ . Then, by simple calculation we get  $(\ker F_*)^\perp = \text{span}\{X_1 = \frac{\partial}{\partial x_1}, X_2 = \frac{1}{\sqrt{2}}\frac{\partial}{\partial x_3} - \frac{1}{2}\frac{\partial}{\partial x_4}, X_3 = \frac{1}{3}\frac{\partial}{\partial x_5} - \frac{1}{\sqrt{3}}\frac{\partial}{\partial x_6}, X_4 = \frac{\partial}{\partial x_7}\}$  and  $\text{range } F_* = \text{span}\{F_*X_1 = \frac{\partial}{\partial u_1}, F_*X_2 = \frac{3}{2\sqrt{2}}\frac{\partial}{\partial u_2} + \frac{3}{4}\frac{\partial}{\partial u_3}, F_*X_3 = -\frac{4}{3\sqrt{2}}\frac{\partial}{\partial v_1} + \frac{2}{3}\frac{\partial}{\partial v_2}, F_*X_4 = \frac{\partial}{\partial w} = \xi\}$ , where  $\mathcal{D}_{\theta_1} = \text{span}\{F_*X_1, F_*X_2\}$  and  $\mathcal{D}_{\theta_2} = \text{span}\{F_*X_3\}$ . Here, we can see  $g_1(X_i, X_j) = g_2(F_*X_i, F_*X_j)$ ,  $i, j = 1, \dots, 4$  and  $\theta_1 = \cos^{-1}(\frac{2}{3})$  and  $\theta_2 = \frac{\pi}{2}$ . Hence,  $F$  is a bi-slant Riemannian map with slant angles  $\cos^{-1}(\frac{2}{3})$  and  $\frac{\pi}{2}$ .

**Example 2.** Let  $M = \{(u_1, u_2, u_3, u_4, v_1, v_2, v_3, v_4, w) \in \mathbb{R}^9 \mid w \neq 0\}$  be a Riemannian manifold with contact metric structure  $(\psi, \xi, \eta, g_2)$  define by  $\eta = dz$ ,  $\xi = \frac{\partial}{\partial w}$ ,  $\psi(\frac{\partial}{\partial u_1}) = \frac{\partial}{\partial u_3}$ ,  $\psi(\frac{\partial}{\partial u_3}) = -\frac{\partial}{\partial u_1}$ ,  $\psi(\frac{\partial}{\partial u_2}) = \frac{\partial}{\partial u_4}$ ,  $\psi(\frac{\partial}{\partial u_4}) = -\frac{\partial}{\partial u_2}$ ,  $\psi(\frac{\partial}{\partial v_1}) = \frac{\partial}{\partial v_3}$ ,  $\psi(\frac{\partial}{\partial v_3}) = -\frac{\partial}{\partial v_1}$ ,  $\psi(\frac{\partial}{\partial v_2}) = \frac{\partial}{\partial v_4}$ ,  $\psi(\frac{\partial}{\partial v_4}) = -\frac{\partial}{\partial v_2}$ ,  $\psi(\frac{\partial}{\partial w}) = 0$  and  $g_2 = \sum_{i=1}^2 (du_i^2 + dv_i^2) + \sum_{j=3}^4 \frac{1}{2}(du_j^2 + \frac{1}{(\beta^2 + \gamma^2)} dv_j^2) + dw^2$ . Then, from (1)–(5) we can see that  $M$  is a Kenmotsu manifold. Further, let  $F : (\overline{M} = \mathbb{R}^7, g_1) \rightarrow (M, \psi, \xi, \eta, g_2)$  be a Riemannian map defines by

$$F_*(x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8, x_9) = (x_1, 0, \sin\alpha(x_3 - x_4), \cos\alpha(x_3 - x_4), 0, x_6, \beta(x_7 + x_8), \gamma(x_7 + x_8), x_9),$$

where  $\alpha \in (0, \pi/2)$ ,  $0 \neq \beta, \gamma \in \mathbb{R}$  and  $\overline{M}$  is a Riemannian manifold with Cartesian co-ordinates  $(x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8, x_9)$  and  $g_2$  is usual inner product on  $\mathbb{R}^9$ . Then

by simple calculation, we get  $(\ker F_*)^\perp = \text{span}\{X_1 = \frac{\partial}{\partial x_1}, X_2 = \frac{\partial}{\partial x_3} - \frac{\partial}{\partial x_4}, X_3 = \frac{\partial}{\partial x_6}, X_4 = \frac{\partial}{\partial x_7} + \frac{\partial}{\partial x_8}, X_5 = \frac{\partial}{\partial x_9}\}$  and  $\text{range} F_* = \text{span}\{F_*X_1 = \frac{\partial}{\partial u_1}, F_*X_2 = 2\sin\alpha\frac{\partial}{\partial u_3} + 2\cos\alpha\frac{\partial}{\partial u_4}, F_*X_3 = \frac{\partial}{\partial v_2}, F_*X_4 = 2\beta\frac{\partial}{\partial v_3} + 2\gamma\frac{\partial}{\partial v_4}, F_*X_5 = \frac{\partial}{\partial w} = \xi\}$ , where  $\mathcal{D}_{\theta_1} = \text{span}\{F_*X_1, F_*X_2\}$  and  $\mathcal{D}_{\theta_2} = \text{span}\{F_*X_3, F_*X_4\}$ . Here, we can see  $g_1(X_i, X_j) = g_2(F_*X_i, F_*X_j), i, j = 1, \dots, 5$  and  $\theta_1 = \cos^{-1}\sin\alpha$  and  $\theta_2 = \cos^{-1}\left(\frac{\gamma}{\sqrt{\beta^2+\gamma^2}}\right)$ . Hence,  $F$  is a bi-slant Riemannian map with slant angles  $\cos^{-1}\sin\alpha$  and  $\cos^{-1}\left(\frac{\gamma}{\sqrt{\beta^2+\gamma^2}}\right)$ .

**Lemma 1.** Let  $F : (\overline{M}, g_1) \rightarrow (M, \psi, \xi, \eta, g_2)$  be a bi-slant Riemannian map from a Riemannian manifold  $\overline{M}$  to Kenmotsu manifold  $M$  with slant angles  $\theta_1$  and  $\theta_2$  for the distribution  $\mathcal{D}_{\theta_1}$  and  $\mathcal{D}_{\theta_2}$ , respectively, then

$$P^2 F_*(X) = -\cos^2\theta_i(\psi^2 F_*X), \quad i = 1, 2, \quad (20)$$

where  $\theta_i = \begin{cases} \theta_1, & \text{if } F_*X \in \mathcal{D}_{\theta_1}, \\ \theta_2, & \text{if } F_*X \in \mathcal{D}_{\theta_2}. \end{cases}$

Using the above lemma we have the following result.

**Lemma 2.** Let  $F : (\overline{M}, g_1) \rightarrow (M, \psi, \xi, \eta, g_2)$  be a bi-slant Riemannian map from a Riemannian manifold  $\overline{M}$  to Kenmotsu manifold  $M$  with slant angles  $\theta_1$  and  $\theta_2$  for the distribution  $\mathcal{D}_{\theta_1}$  and  $\mathcal{D}_{\theta_2}$ , respectively, then

1.  $\phi Q F_*X = \sin^2\theta_i \psi^2 F_*X$ ,
2.  $Q P F_*X + \omega Q F_*X = 0$ ,
3.  $g_2(P F_*X, P F_*Y) = \cos^2\theta_i g_2(\psi F_*X, \psi F_*Y)$ ,
4.  $g_2(Q F_*X, Q F_*Y) = \sin^2\theta_i g_2(\psi F_*X, \psi F_*Y)$ ,
5.  $g_2(\omega Q F_*X, \omega Q F_*Y) = \sin^2\theta_i \cos^2\theta_i g_2(\psi F_*X, \psi F_*Y)$ ,

where  $X, Y \in \Gamma((\ker F_*)^\perp)$  and  $\theta_i = \begin{cases} \theta_1, & \text{if } F_*X \in \mathcal{D}_{\theta_1}, \\ \theta_2, & \text{if } F_*X \in \mathcal{D}_{\theta_2} \end{cases}$ .

**Theorem 3.** Let  $F : (\overline{M}, g_1) \rightarrow (M, \psi, \xi, \eta, g_2)$  be a bi-slant Riemannian map from a Riemannian manifold  $\overline{M}$  to Kenmotsu manifold  $M$  with  $\xi \in \Gamma(\text{range} F_*)$ , then  $\text{range} F_*$  defines a totally geodesic foliation if and only if

$$Q(S_{QF_*Y}F_*X - F_*(\nabla_X^* F_* P F_*Y)) = \omega((\nabla F_*)(X, *F_* P F_*Y) - \nabla_X^{F_*\perp} Q F_*Y),$$

where  $X, Y \in \Gamma((\ker F_*)^\perp)$ .

*Proof.* Let  $X, Y \in \Gamma((\ker F_*)^\perp)$ , since  $F$  is a Riemannian map and  $M$  is a Kenmotsu manifold, from (1), (2), (4), (5) and (18), we get

$$\nabla_{F_*X} F_*Y = -\psi \nabla_{F_*X} P F_*Y - \psi \nabla_{F_*X} Q F_*Y - g_2(X, Y)\xi + F_*X\eta(F_*Y). \quad (21)$$

Let  $*F_*$  be the adjoint of the map  $F$ , from (8), (9) and (21), we have

$$\nabla_{F_*X} F_*Y = -\psi((\nabla F_*)(X, *F_* P F_*Y) - F_*(\nabla_X^* F_* P F_*Y)) - \psi(-S_{QF_*Y}F_*X$$

$$+ \nabla_X^{F^\perp} QF_*Y) - g_2(X, Y)\xi + F_*X\eta(F_*Y). \quad (22)$$

If  $\xi \in \Gamma(\text{range}F_*)$  and  $(\text{range}F_*)$  defines a totally geodesic foliation, then from (22) we have the result.  $\square$

**Theorem 4.** Let  $F : (\overline{M}, g_1) \rightarrow (M, \psi, \xi, \eta, g_2)$  be a bi-slant Riemannian map from a Riemannian manifold  $\overline{M}$  to Kenmotsu manifold  $M$  with  $\xi \in \Gamma(\text{range}F_*)$ , then  $(\text{range}F_*)^\perp$  defines a totally geodesic foliation if and only if

$$g_2(S_{\omega V_1}F_*X, \phi V_2 - F_*(\nabla^*F_*\phi V_1)) = g_2([V_1, F_*X], V_2) + g_2((\nabla F_*)(X, *F_*\phi V_1) + \nabla_X^{F^\perp}\omega V_1, \omega V_2)$$

where  $X \in \Gamma((\ker F_*)^\perp)$  and  $V_1, V_2 \in \Gamma((\text{range}F_*)^\perp)$ .

*Proof.* Let  $V_1, V_2 \in \Gamma(\text{range}F_*)^\perp$  and  $X \in \Gamma((\ker F_*)^\perp)$ , then we have

$$g_2(\nabla_{V_1}V_2, F_*X) = g_2([V_1, F_*X] + \nabla_X^F V_1, V_2). \quad (23)$$

Since  $M$  is a Sasakian manifold, from (2) and (23), we have

$$g_2(\nabla_{V_1}V_2, F_*X) = g_2([V_1, F_*X], V_2) + g_2(\nabla_{F_*X}\psi V_1, \psi V_2), \quad (24)$$

if  $*F_*$  is a adjoint map of  $F$ , using (8), (9) and (19) in (24), we get

$$g_2(\nabla_{V_1}V_2, F_*X) = g_2([V_1, F_*X], V_2) + g_2(F_*(\nabla^*F_*\phi V_1) - S_{\omega V_1}F_*X, \phi V_2) + g_2((\nabla F_*)(X, *F_*\phi V_1) + \nabla_X^{F^\perp}\omega V_1, \omega V_2). \quad (25)$$

Hence, from (25), we have the result.  $\square$

**Theorem 5.** Let  $F : (\overline{M}, g_1) \rightarrow (M, \psi, \xi, \eta, g_2)$  be a bi-slant Riemannian map from a Riemannian manifold  $\overline{M}$  to Kenmotsu manifold  $M$  with  $\xi \in \Gamma(\text{range}F_*)$ , then  $F$  is totally geodesic if and only if

(1)

$$(\cos^2\theta_2 - \cos^2\theta_1)g_2(\nabla_X^F\sigma F_*Y, F_*Z) = \cos^2\theta_2g_2(\nabla_X^F F_*Y, V) + g_2(\nabla_X^{F^\perp}QPF_*Y, V) + g_2(S_{QF_*Y}F_*X, \phi V) + g_2(\nabla_X^{F^\perp}QF_*Y, \omega V)$$

(2)  $\ker F_*$  is totally geodesics.

(3)  $(\ker F_*)^\perp$  is totally geodesics.

where  $X, Y \in \Gamma((\ker F_*)^\perp)$ .

*Proof.* Since  $F$  is a Riemannian map and  $M$  is a Kenmotsu manifold, from (4), (5) and (8), we can write

$$g_2((\nabla F_*)(X, Y), V) = g_2(\nabla_X^F F_*Y, V) = -g_2(\psi \nabla_X^F \psi F_*Y, F_*Z),$$



now, using (4), (18) and (19) in above equation, we get

$$g_2((\nabla F_*)(X, Y), V) = -g_2(\nabla_X^F P^2 F_* Y, V) + g_2(\nabla_X^{F^\perp} Q P F_* Y, V) + g_2(S_{Q F_* Y} F_* X, \phi V) + g_2(\nabla_X^{F^\perp} Q F_* Y, \omega V), \quad (26)$$

substituting  $F_*(Y) = \sigma F_*(Y) + \pi F_*(Y) + \eta(U)\xi$  in (26), further using lemma 3.2 and simplifying, we get

$$g_2((\nabla F_*)(X, Y), V) = \cos^2 \theta_1 g_2(\nabla_X^F \sigma F_* Y, V) + \cos^2 \theta_2 g_2(\nabla_X^F \pi F_* Y, V) + g_2(\nabla_X^{F^\perp} Q P F_* Y, V) + g_2(S_{Q F_* Y} F_* X, \phi V) + g_2(\nabla_X^{F^\perp} Q F_* Y, \omega V)$$

again, substituting  $\pi F_* Y = F_* Y - \sigma F_* Y - \eta(F_* Y)\xi$  and using (5) in above equation, we have

$$g_2((\nabla F_*)(X, Y), V) = (\cos^2 \theta_1 - \cos^2 \theta_2) g_2(\nabla_X^F \sigma F_* Y, V) + \cos^2 \theta_2 g_2(\nabla_X^F F_* Y, V) + g_2(\nabla_X^{F^\perp} Q P F_* Y, V) + g_2(S_{Q F_* Y} F_* X, \phi V) + g_2(\nabla_X^{F^\perp} Q F_* Y, \omega V), \quad (27)$$

that implies (1), similarly we can obtain (2) and (3).  $\square$

Further, Let  $X, Y, Z, H \in \Gamma((\ker F_*)^\perp)$  and  $(\text{range } F_*)^\perp$  is totally geodesic, then we can write

$$g_2(R^M(\psi F_* X, \psi F_* Y) Q F_* Z, Q F_* H) = g_2(R^M(P F_* X, P F_* Y) Q F_* Z, Q F_* H) + g_2(R^M(Q F_* X, Q F_* Y) Q F_* Z, Q F_* H).$$

Since  $M$  is a Kenmotsu manifold, from (6) get

$$g_2(R^M(\psi F_* X, \psi F_* Y) Q F_* Z, Q F_* H) = g_2(R^M(F_* X, F_* Y) Q F_* Z, Q F_* H) - g_2(\psi F_* Y, Q F_* Z)(\psi F_* X, Q F_* H) + g_2(\psi F_* X, Q F_* Z)(\psi F_* Y, Q F_* H),$$

using (12) in above equation, we get

$$g_2(R^M(\psi F_* X, \psi F_* Y) Q F_* Z, Q F_* H) = g_2(R^{F^\perp}(F_* X, F_* Y) Q F_* Z, Q F_* H) + g_2([S_{Q F_* H}, S_{Q F_* Z}] F_* X, F_* Y), - g_2(\psi F_* Y, Q F_* Z)(\psi F_* X, Q F_* H) + g_2(\psi F_* X, Q F_* Z)(\psi F_* Y, Q F_* H)$$

and also from (12), we have

$$g_2(R^M(P F_* X, P F_* Y) Q F_* Z, Q F_* H) = g_2(R^{F^\perp}(P F_* X, P F_* Y) Q F_* Z, Q F_* H) + g_2([S_{Q F_* H}, S_{Q F_* Z}] P F_* X, P F_* Y),$$

hence, we conclude that

$$\begin{aligned}
g_2(R^M(QF_*X, QF_*Y)QF_*Z, QF_*H) &= g_2(R^{F^\perp}(F_*X, F_*Y)QF_*Z, QF_*H) \\
&\quad - g_2(R^{F^\perp}(PF_*X, PF_*Y)QF_*Z, QF_*H) \\
&\quad + g_2([S_{QF_*H}, S_{QF_*Z}]F_*X, F_*Y) \\
&\quad - g_2([S_{QF_*H}, S_{QF_*Z}]PF_*X, PF_*Y) \\
&\quad - g_2(QF_*Y, QF_*Z)g_2(QF_*X, QF_*H) \\
&\quad + g_2(QF_*X, QF_*Z)g_2(QF_*Y, QF_*H) \quad (28)
\end{aligned}$$

**Theorem 6.** Let  $F : (\overline{M}, g_1) \rightarrow (M, \psi, \xi, \eta, g_2)$  be a bi-slant Riemannian map from a Riemannian manifold  $\overline{M}$  to Kenmotsu manifold  $M$  with  $\xi \in \Gamma(\text{range}F_*)$ ,  $\mu = \{0\}$  and  $(\text{range}F_*)^\perp$  is totally geodesic, then we have

$$\begin{aligned}
\mathcal{R}ic^{(\text{range}F_*)^\perp}(V_1, V_2) &\geq g_2(R^{F^\perp}(F_*e_j, F_*X)V_2, v_i) + g_2([S_{v_i}, S_{V_2}]F_*e_j, F_*X) \\
&\quad + (1 - m + r)g_2(V_1, V_2), \quad (29)
\end{aligned}$$

where  $X, Y \in \Gamma((\ker F_*)^\perp)$  and  $V_1 = QF_*X, V_2 = QF_*Y \in \Gamma((\text{range}F_*)^\perp)$ . Also,  $\{e_1, \dots, e_r\}$  and  $\{v_{r+1}, \dots, v_m\}$  be the orthonormal basis of  $(\ker F_*)^\perp$  and  $(\text{range}F_*)^\perp_{F_*p}$ ,  $p \in \overline{M}$ , respectively such that  $\{F_*e_1, \dots, F_*e_r\}$  is the orthonormal basis of  $(\text{range}F_*)_{F_*p}$ . The equality case arises when  $F$  is totally geodesic, in this case

$$\begin{aligned}
\mathcal{R}ic^{(\text{range}F_*)^\perp}(V_1, V_2) &= g_2(R^{F^\perp}(F_*e_j, F_*X)V_2, v_i) - g_2(R^{F^\perp}(PF_*e_j, PF_*X)V_2, v_i) \\
&\quad + (1 - m + r)g_2(V_1, V_2) \quad (30)
\end{aligned}$$

*Proof.* Since  $\mu = \{0\}$ , then  $(\text{range}F_*)^\perp = Q\mathcal{D}_{\theta_1} \oplus Q\mathcal{D}_{\theta_2}$ . Also,  $(\text{range}F_*)^\perp$  is totally geodesic, then from (28), we have (29). If  $F$  is totally geodesic, then  $S_V F_*X = 0$ , from (28) we have (30).  $\square$

**COROLLARY 3.1.** Let  $F : (\overline{M}, g_1) \rightarrow (M, \psi, \xi, \eta, g_2)$  be a bi-slant totally geodesic Riemannian map from a Riemannian manifold  $\overline{M}$  to Kenmotsu manifold  $M$  with  $\xi \in \Gamma(\text{range}F_*)$ ,  $\mu = \{0\}$ , then we have (30).

*Proof.* Using theorem 3.7 and (28), we have the required result.  $\square$

## 4 Chen-Ricci and DDVV inequalities for bi-slant Riemannian maps to Kenmotsu space form

In this section, we established Chen-Ricci inequalities for the bi-slant Riemannian maps from Riemannian manifolds to Kenmotsu space forms.

Let  $F : \overline{M}^{\overline{m}} \rightarrow M(c)^m$  be a Riemannian map from a Riemannian manifold to Kenmotsu space Form  $M(c)$  with  $\xi \in \Gamma(\text{range}F_*)$ , where  $\overline{m}$  and  $m$  are the dimensions of  $\overline{M}$  and  $M$  respectively. Further, let  $\{e_1, \dots, e_r\}$  and  $\{v_{r+1}, \dots, v_m\}$  be the orthonormal basis of  $(\ker F_*)^\perp$  and  $(\text{range}F_*)^\perp_{F_*p}$ ,  $p \in \overline{M}$ , respectively, then  $\{F_*e_1, \dots, F_*e_r\}$  is the

orthonormal basis of  $(range F_*)_{F_*p}$  such that  $r = 2r_1 + 2r_2 + 1$  (if  $\xi \in \Gamma(range F_*)$ ) or  $r = 2r_1 + 2r_2$  (if  $\xi \in \Gamma((range F_*)^\perp)$ ), where dimensions of  $\mathcal{D}_{\theta_1}$  and  $\mathcal{D}_{\theta_2}$  are  $2r_1$  and  $2r_2$ , respectively. Then we can consider canonical bi-slant orthonormal frame as

$$\begin{aligned} F_*e_1, F_*e_2 &= sec\theta_1 PF_*e_1, \dots, F_*e_{2r_1-1}, F_*e_{2r_1} = sec\theta_1 PF_*e_{2r_1-1}, F_*e_{2r_1+1}, F_*e_{2r_1+2} \\ &= sec\theta_2 PF_*e_{2r_1+1}, \dots, F_*e_{2r_1+2r_2-1}, F_*e_{2r_1+2r_2} = sec\theta_2 PF_*e_{2r_1+2r_2-1}, F_*e_{2r_1+2r_2+1} = \xi, \end{aligned}$$

hence, we have

$$g_2^2(\psi F_*e_i, F_*e_{i+1}) = \begin{cases} \cos^2\theta_1, & \text{for } i = 1, \dots, 2r_1 - 1, \\ \cos^2\theta_2, & \text{for } i = 2r_1 + 1, \dots, 2r_1 + 2r_2 - 1, \end{cases}$$

then,

$$\sum_{i,j=1}^r g_2^2(\psi F_*e_i, F_*e_{i+1}) = 2(r_1 \cos^2\theta_1 + r_2 \cos^2\theta_2).$$

Similarly, if  $\xi \in \Gamma((range F_*)^\perp)$ , we have same results for  $g_2^2(\psi F_*e_i, F_*e_{i+1})$  and  $\sum_{i,j=1}^r g_2^2(\psi F_*e_i, F_*e_{i+1})$ .

**Remark:** Throughout this article, if  $\xi \in \Gamma(range F_*)$ , we consider  $r = 2r_1 + 2r_2 + 1$  and if  $\xi \in \Gamma((range F_*)^\perp)$  then  $r = 2r_1 + 2r_2$ .

**Theorem 7.** Let  $F : \overline{M} \rightarrow M(c)$  be a bi-slant Riemannian map from a Riemannian manifold  $\overline{M}$  to Kenmotsu space Form  $M(c)$  with  $\xi \in \Gamma(range F_*)$  and  $rank F = r < m$  then we have

$$4Ric^{(ker F_*)^\perp}(X) \leq (c-3)(r-1) - 2(c+1) + ||trace\zeta||^2 + 3(c+1) \sum_{i=1}^r g_2^2(\psi F_*X, F_*e_i), \quad (31)$$

where  $X$  is a unit horizontal vector field on  $\overline{M}$ , provided,

$$g_2^2(\psi F_*e_1, e_i) = \begin{cases} \cos^2\theta_1, & \text{if } F_*e_1 \in \mathcal{D}_{\theta_1} \\ \cos^2\theta_2, & \text{if } F_*e_1 \in \mathcal{D}_{\theta_2} \end{cases}$$

Moreover;

1. The equality case of (31) arises for  $X \in \Gamma((ker F_*)^\perp)$  if and only if

$$\begin{cases} (\nabla F_*)(X, Y) = 0 \quad \forall Y \in \Gamma((ker F_*)^\perp) \text{ orthogonal to } X, \\ (\nabla F_*)(X, X) = \frac{1}{2}trace\zeta, \end{cases} \quad (32)$$

2. The equality case of (31) true for all  $X \in \Gamma((ker F_*)^\perp)$  if and only if  $\zeta = 0$ . or  $r = 3$  and  $\zeta_{11}^\alpha = \zeta_{22}^\alpha, \alpha = r + 1, \dots, m$ .

*Proof.* Let  $M(c)$  be a Kenmotsu space form with constant sectional curvature  $c$ , then from (7) and (11), we have

$$\begin{aligned}
g_1(R^{\overline{M}}(X, Y)Z, H) &= \frac{c-3}{4} \left\{ g_1(Y, Z)g_1(X, H) - g_1(X, Z)g_1(Y, H) \right\} \\
&\quad + \frac{c+1}{4} \left\{ \eta(F_*X)\eta(F_*Z)g_1(Y, H) - \eta(F_*Y)\eta(F_*Z)g_1(X, H) \right. \\
&\quad + \eta(F_*Y)\eta(F_*H)g_1(X, Z) - \eta(F_*X)\eta(F_*H)g_1(Y, Z) \\
&\quad - g_2(\psi F_*X, F_*Z)g_2(\psi F_*Y, F_*H) + g_2(\psi F_*Y, F_*Z)g_2(\psi F_*X, F_*H) \\
&\quad \left. + 2g_2(\psi F_*Y, F_*X)g_2(\psi F_*Z, F_*H) \right\} - g_2((\nabla F_*)(X, Z), (\nabla F_*)(Y, H)) \\
&\quad + g_2((\nabla F_*)(Y, Z), (\nabla F_*)(X, H)),
\end{aligned} \tag{33}$$

where  $X, Y, Z, H \in \Gamma((ker F_*)^\perp)$ . let  $\{e_1, \dots, e_{r_1}, e_{r_1+1}, \dots, e_{2r_1+2r_2+1=r}\}$  with  $e_1 = X$  and  $\{v_{r+1}, \dots, v_s\}$  be the orthonormal basis of  $(ker F_*)^\perp$  and  $(range F_*)^\perp_{F_*p}$ ,  $p \in \overline{M}$ , respectively, where  $dim(\mathcal{D}_{\theta_1}) = 2r_1$  and  $dim(\mathcal{D}_{\theta_2}) = 2r_2$ , then  $\{F_*e_1, \dots, F_*e_{2r_1}, F_*e_{2r_1+1}, \dots, F_*e_{2r_1+2r_2+1=r} = \xi\}$  is the orthonormal basis of  $(range F_*)^\perp_{F_*p}$ . From (14)–(17) and (33), we get

$$\frac{c-3}{4}r(r-1) + \frac{c+1}{4} \left\{ -2r+2+3 \sum_{i,j=1}^r g_2^2(\psi F_*e_i, F_*e_j) \right\} = \|\zeta\|^2 - \|\text{trace} \zeta\|^2 - 2\tau^{(ker F_*)^\perp}(p) \tag{34}$$

Also, we can see that [22]

$$\begin{aligned}
\|\zeta\|^2 &= \frac{1}{2} \|\text{trace} \zeta\|^2 + \frac{1}{2} \sum_{\alpha=r+1}^m (\zeta_{11}^\alpha - \zeta_{22}^\alpha - \dots - \zeta_{rr}^\alpha)^2 + 2 \sum_{\alpha=r+1}^m \sum_{i=2}^r (\zeta_{1i}^\alpha)^2 \\
&\quad - 2 \sum_{\alpha=r+1}^m \sum_{2 \leq i < j \leq r} \left\{ \zeta_{ii}^\alpha \zeta_{jj}^\alpha - (\zeta_{ij}^\alpha)^2 \right\}, \quad 6(r_1 \cos^2 \theta_1 + r_2 \cos^2 \theta_2)
\end{aligned} \tag{35}$$

and

$$\begin{aligned}
\tau^{(ker F_*)^\perp}(p) &= \frac{c-3}{8}(r-1)(r-2) + \frac{c+1}{4} \left\{ -r+2+3 \sum_{2 \leq i < j \leq r} g_2^2(\psi F_*e_i, F_*e_j) \right\} \\
&\quad - \sum_{\alpha=r+1}^m \sum_{2 \leq i < j \leq r} \left\{ (\zeta_{ij}^\alpha)^2 - \zeta_{ii}^\alpha \zeta_{jj}^\alpha \right\} + Ric^{(ker F_*)^\perp}(X),
\end{aligned} \tag{36}$$

substituting the value of (35) and (36) in (34), we get

$$\begin{aligned}
\mathcal{R}ic^{(ker F_*)^\perp}(X) &= \frac{c-3}{4}(r-1) - \frac{c+1}{2} + \frac{1}{4} \|\text{trace} \zeta\|^2 + \frac{3(c+1)}{4} \sum_{i=1}^r g_2^2(\psi F_* X, F_* e_i) \\
&\quad - \frac{1}{4} \sum_{\alpha=r+1}^m \left( \zeta_{11}^\alpha - \zeta_{22}^\alpha - \dots - \zeta_{rr}^\alpha \right)^2 - \sum_{\alpha=r+1}^m \sum_{i=2}^r (\zeta_{1i}^\alpha)^2 \\
&\leq \frac{c-3}{4}(r-1) - \frac{c+1}{2} + \frac{1}{4} \|\text{trace} \zeta\|^2 + \frac{3(c+1)}{4} \sum_{i=1}^r g_2^2(\psi F_* X, F_* e_i),
\end{aligned} \tag{37}$$

hence, from we have (31). The equality case of (31) holds if and only if

$$\begin{aligned}
\zeta_{11}^\alpha &= \zeta_{22}^\alpha + \dots + \zeta_{r-1r-1}^\alpha \\
\text{and } \zeta_{1i}^\alpha &= 0, \quad i = 1, \dots, r-1; \alpha = r+1, \dots, m;
\end{aligned}$$

hence we have (32). Further, we consider that the equality case of (31) holds for all  $X \in \Gamma((ker F_*)^\perp)$ , then for all  $\alpha$ , we have

$$\zeta_{ij}^\alpha = 0, \quad i \neq j,$$

$$\begin{aligned}
2\zeta_{ii}^\alpha &= \zeta_{11}^\alpha + \zeta_{22}^\alpha + \dots + \zeta_{r-1r-1}^\alpha, \quad i = 1, \dots, r-1; \\
\text{or } (r-3) \left( \zeta_{11}^\alpha + \zeta_{22}^\alpha + \dots + \zeta_{r-1r-1}^\alpha \right) &= 0,
\end{aligned}$$

which implies that either  $r = 3$  or  $(\zeta_{11}^\alpha + \zeta_{22}^\alpha + \dots + \zeta_{r-1r-1}^\alpha = 0$ . If  $r = 3$ , then  $\zeta_{11}^\alpha = \zeta_{22}^\alpha + \zeta_{33}^\alpha$ . Also, assuming  $F_* e_3 = \xi$ , we get  $\zeta_{33}^\alpha = 0$ . Hence, the equality case of (31) holds for all  $X \in \Gamma((ker F_*)^\perp)$ , either  $r = 3$  and  $\zeta_{11}^\alpha = \zeta_{22}^\alpha$  or  $\zeta = 0$ . Since the converse case is trivial so we have the result.  $\square$

Similarly, when  $\xi \in \Gamma((range F_*)^\perp)$  we have the following result.

**Theorem 8.** *Let  $F : \overline{M} \rightarrow M(c)$  be a bi-slant Riemannian map from a Riemannian manifold  $\overline{M}$  to Kenmotsu space Form  $M(c)$  with  $\xi \in \Gamma((range F_*)^\perp)$  and  $\text{rank} F = r < m$  then we have*

$$4\mathcal{R}ic^{(ker F_*)^\perp}(X) \leq (c-3)(r-1) + \|\text{trace} \zeta\|^2 + 3(c+1) \sum_{i=1}^r g_2^2(\psi F_* X, F_* e_i), \tag{38}$$

where  $X$  is a unit horizontal vector field on  $\overline{M}$ , provided,

$$g_2^2(\psi F_* e_1, e_i) = \begin{cases} \cos^2 \theta_1, & \text{if } F_* e_1 \in \mathcal{D}_{\theta_1}, \\ \cos^2 \theta_2, & \text{if } F_* e_1 \in \mathcal{D}_{\theta_2}. \end{cases}$$

Moreover,

1. The equality case of (38) arises for  $X \in \Gamma((kerF_*)^\perp)$  if and only if

$$\begin{cases} (\nabla F_*)(X, Y) = 0 \quad \forall Y \in \Gamma((kerF_*)^\perp) \text{ orthogonal to } X, \\ (\nabla F_*)(X, X) = \frac{1}{2} \text{trace} \zeta. \end{cases} \quad (39)$$

2. The equality case of (38) true for all  $X \in \Gamma((kerF_*)^\perp)$  if and only if  $\zeta = 0$  or  $\dim((kerF_*)^\perp) = 2$  and  $\zeta_{11}^\alpha = \zeta_{22}^\alpha, \alpha = r+1, \dots, m$ .

## DDVV inequalities from Riemannian manifolds to Kenmotsu space forms

**Theorem 9.** Let  $F : (\overline{M}, g_1) \rightarrow M(c)$  be a bi-slant Riemannian map from a Riemannian manifold  $\overline{M}$  to Kenmotsu space form  $M(c)$  with  $\xi \in \Gamma(\text{range}F_*)$  and  $\text{rank}F = r < m$ , then normalized normal scalar curvature  $\varrho^{(\text{range}F_*)^\perp}$  and normalized scalar curvature  $\varrho^{(kerF_*)^\perp}$  satisfy

$$\varrho^{(\text{range}F_*)^\perp} + \varrho^{(kerF_*)^\perp} \leq \frac{1}{r^2} \|\text{trace} \zeta\|^2 + \frac{c-3}{4} - \frac{c+1}{2r} + \frac{3(c+1)}{2r(r-1)} (r_1 \cos^2 \theta_1 + r_2 \cos^2 \theta_2), \quad (40)$$

where  $\theta_1, \theta_2$  are the slant angles and  $2r_1, 2r_2$  are the dimensions of distributions  $\mathcal{D}_{\theta_1}$  and  $\mathcal{D}_{\theta_2}$ , respectively.

*Proof.* Let  $\{e_1, \dots, e_{2r_1}, e_{2r_1+1}, \dots, e_{2r_1+2r_2+1=r}\}$  and  $\{v_{r+1}, \dots, v_s\}$  be the orthonormal basis of  $(kerF_*)_p^\perp$  and  $(\text{range}F_*)_{F_*p}^\perp$ ,  $p \in \overline{M}$ , respectively, where  $\dim(\mathcal{D}_{\theta_1}) = 2r_1$  and  $\dim(\mathcal{D}_{\theta_2}) = 2r_2$ , then  $\{F_*e_1, \dots, F_*e_{2r_1}, F_*e_{2r_1+1}, \dots, F_*e_{2r_1+2r_2+1=r} = \xi\}$  is the orthonormal basis of  $(\text{range}F_*)_{F_*p}$ . Then from (7), (11) and (12), we have

$$\sum_{\alpha, \beta=r+1}^m \sum_{i, j=1}^r g_2(R^{F^\perp}(F_*e_i, F_*e_j)V_\alpha, V_\beta) = - \sum_{\alpha, \beta=r+1}^m \sum_{i, j=1}^r g_2([S_{v_\beta}, S_{v_\alpha}]F_*e_i, F_*e_j)$$

using the above result in (13), we get

$$\varrho^{(\text{range}F_*)^\perp} = \frac{2}{r(r-1)} \sqrt{\sum_{1 \leq i < j \leq r} \sum_{1 \leq \alpha < \beta \leq m} g_2^2([S_{v_\beta}, S_{v_\alpha}]F_*e_i, F_*e_j)}, \quad (41)$$

using (10) in (41) and solving further, we get

$$\varrho^{(\text{range}F_*)^\perp} = \frac{2}{r(r-1)} \sqrt{\sum_{1 \leq i < j \leq r} \sum_{1 \leq \alpha < \beta \leq m} \left( \sum_{k=1}^r (\zeta_{jk}^\alpha \zeta_{ik}^\beta - \zeta_{ik}^\alpha \zeta_{jk}^\beta) \right)^2}, \quad (42)$$

Further, from [23] (page no. 2), we have

$$\sum_{\alpha=r+1}^m \sum_{1 \leq i < j \leq r} (h_{ii}^\alpha - h_{jj}^\alpha)^2 + 2r \sum_{\alpha=r+1}^m \sum_{1 \leq i < j \leq r} (h_{ij}^\alpha)^2$$

$$\geq 2r \left( \sum_{1 \leq \alpha < \beta \leq m} \sum_{1 \leq i < j \leq r} \left( \sum_{k=1}^r (h_{jk}^\alpha h_{ik}^\beta - h_{ik}^\alpha h_{jk}^\beta) \right)^2 \right)^{\frac{1}{2}},$$

where  $(h_{ij}^\alpha)(i, j = 1, \dots, n$  and  $\alpha = r+1, \dots, m)$  be the entries of the second fundamental form of manifold  $B$ , an  $n$ -dimensional manifold isometrically immersed into the space form  $\overline{B}^{n+m}(c)$  of constant sectional curvature  $c$ , under the orthonormal frames of both the tangent bundle and the normal bundle. It can be verify that above inequality is also true for  $\zeta_{ij}^\alpha$ . Hence using the above inequality for  $\zeta_{ij}^\alpha$  in (42), we get

$$r^2(r-1)\varrho^{(rangeF_*)^\perp} \leq \sum_{\alpha=r+1}^m \sum_{1 \leq i < j \leq r} (\zeta_{ii}^\alpha - \zeta_{jj}^\alpha)^2 + 2r \sum_{\alpha=r+1}^m \sum_{1 \leq i < j \leq r} (\zeta_{ij}^\alpha)^2. \quad (43)$$

Further, we can compute

$$(r-1)\|\text{trace}\zeta\|^2 = \sum_{\alpha=r+1}^m \sum_{i,j=1}^r \zeta_{ii}^\alpha \zeta_{jj}^\alpha = \sum_{\alpha=r+1}^m \sum_{1 \leq i < j \leq r} \left( (\zeta_{ii}^\alpha - \zeta_{jj}^\alpha)^2 + 2r\zeta_{ii}^\alpha \zeta_{jj}^\alpha \right), \quad (44)$$

and

$$\begin{aligned} \tau^{(kerF_*)^\perp} &= \frac{(r-1)}{4} \left( \frac{c-3}{2} r - c - 1 \right) + \frac{3(c+1)}{4} (r_1 \cos^2 \theta_1 + r_2 \cos^2 \theta_2) \\ &+ \sum_{\alpha=r+1}^m \sum_{1 \leq i < j \leq r} \left( \zeta_{ii}^\alpha \zeta_{jj}^\alpha - (\zeta_{ij}^\alpha)^2 \right). \end{aligned} \quad (45)$$

From (43), (44) and (45), we get the required result.  $\square$

Similarly, when  $\xi \in \Gamma((rangeF_*)^\perp)$  we have the following result.

**Theorem 10.** *Let  $F : (\overline{M}, g_1) \rightarrow M(c)$  be a bi-slant Riemannian map from a Riemannian manifold  $\overline{M}$  to Kenmotsu space form  $M(c)$  with  $\xi \in \Gamma((rangeF_*)^\perp)$  and  $\text{rank}F = r < m$ , then normalized normal scalar curvature  $\varrho^{(rangeF_*)^\perp}$  and normalized scalar curvature  $\varrho^{(kerF_*)^\perp}$  satisfy*

$$\varrho^{(rangeF_*)^\perp} + \varrho^{(kerF_*)^\perp} \leq \frac{1}{r^2} \|\text{trace}\zeta\|^2 + \frac{c-3}{4} + \frac{3(c+1)}{2r(r-1)} (r_1 \cos^2 \theta_1 + r_2 \cos^2 \theta_2),$$

where  $\theta_1, \theta_2$  are the slant angles and  $2r_1, 2r_2$  are the dimensions of distributions  $\mathcal{D}_{\theta_1}$  and  $\mathcal{D}_{\theta_2}$ , respectively.

**COROLLARY 4.1.** *Let  $F : (\overline{M}, g_1) \rightarrow M(c)$  be a bi-slant Riemannian map from a Riemannian manifold  $\overline{M}$  to Kenmotsu space form  $M(c)$  with  $\xi \in \Gamma(rangeF_*)$  and  $\text{rank}F = r < m$ , then for normalized normal scalar curvature  $\varrho^{(rangeF_*)^\perp}$  and normalized scalar curvature  $\varrho^{(kerF_*)^\perp}$ , we have the following relations where  $\theta_1, \theta_2$  are the slant angles and  $2r_1, 2r_2$  are the dimensions of distributions  $\mathcal{D}_{\theta_1}$  and  $\mathcal{D}_{\theta_2}$ , respectively.*

Riemannian Maps	$r_1$ & $r_2$	$\theta_1$	$\theta_2$	$\varrho^{(range F_*)^\perp} + \varrho^{(ker F_*)^\perp}$
Invariant	$r_1 = 0$	-	0	$\leq \frac{1}{r^2} \ \text{trace} \zeta\ ^2 + \frac{c-3}{4} + \frac{c+1}{r}$ ,
Anti-invariant	$r_1 = 0$	-	$\frac{\pi}{2}$	$\leq \frac{1}{r^2} \ \text{trace} \zeta\ ^2 + \frac{c-3}{4}$ ,
Semi-invariant	$r_1, r_2 \neq 0$	0	$\frac{\pi}{2}$	$\leq \frac{1}{r^2} \ \text{trace} \zeta\ ^2 + \frac{c-3}{4} + \frac{c+1}{2r} \left\{ \frac{3r_1}{r-1} - \frac{1}{2} \right\}$ ,
Proper slant	$r_1 = 0$	-	$\in (0, \frac{\pi}{2})$	$\leq \frac{1}{r^2} \ \text{trace} \zeta\ ^2 + \frac{c-3}{4} + \frac{c+1}{4r} (3\cos^2\theta_2 - 1)$ ,
Semi-slant	$r_1, r_2 \neq 0$	0	$\in (0, \frac{\pi}{2})$	$\leq \frac{1}{r^2} \ \text{trace} \zeta^{\mathcal{H}}\ ^2 + \frac{c-3}{4} + \frac{c+1}{2r} \left\{ \frac{3(r_1+r_2\cos^2\theta_2)}{r-1} - \frac{1}{2} \right\}$ ,
Hemi-slant	$r_1, r_2 \neq 0$	$\frac{\pi}{2}$	$\in (0, \frac{\pi}{2})$	$\leq \frac{1}{r^2} \ \text{trace} \zeta\ ^2 + \frac{c-3}{4} + \frac{c+1}{2r} \left\{ \frac{3r_2\cos^2\theta_2}{r-1} - \frac{1}{2} \right\}$ ,

COROLLARY 4.2. Let  $F : \overline{M} \rightarrow M(c)$  be a Riemannian map from a Riemannian manifold  $\overline{M}$  to Kenmotsu space form  $M(c)$  with  $\xi \in \Gamma((range F_*)^\perp)$  and  $rank F = r < m$ , then for normalized normal scalar curvature  $\varrho^{(range F_*)^\perp}$  and normalized scalar curvature  $\varrho^{(ker F_*)^\perp}$ , we have the following relations

Riemannian Maps	$r_1$ & $r_2$	$\theta_1$	$\theta_2$	$\varrho^{(range F_*)^\perp} + \varrho^{(ker F_*)^\perp}$
Invariant	$r_1 = 0$	-	0	$\leq \frac{1}{r^2} \ \text{trace} \zeta\ ^2 + \frac{c-3}{4} + \frac{3(c+1)}{4(r-1)}$ ,
Anti-invariant	$r_1 = 0$	-	$\frac{\pi}{2}$	$\leq \frac{1}{r^2} \ \text{trace} \zeta\ ^2 + \frac{c-3}{4}$ ,
Semi-invariant	$r_1, r_2 \neq 0$	0	$\frac{\pi}{2}$	$\leq \frac{1}{r^2} \ \text{trace} \zeta\ ^2 + \frac{c-3}{4} + \frac{3(c+1)r_1}{2r(r-1)}$ ,
Proper slant	$r_1 = 0$	-	$\in (0, \frac{\pi}{2})$	$\leq \frac{1}{r^2} \ \text{trace} \zeta\ ^2 + \frac{c-3}{4} + \frac{3(c+1)}{4(r-1)} \cos^2\theta_2$ ,
Semi-slant	$r_1, r_2 \neq 0$	0	$\in (0, \frac{\pi}{2})$	$\leq \frac{1}{r^2} \ \text{trace} \zeta\ ^2 + \frac{c-3}{4} + \frac{3(c+1)}{2r(r-1)} (r_1 + r_2 \cos^2\theta_2)$ ,
Hemi-slant	$r_1, r_2 \neq 0$	$\frac{\pi}{2}$	$\in (0, \frac{\pi}{2})$	$\leq \frac{1}{r^2} \ \text{trace} \zeta\ ^2 + \frac{c-3}{4} + \frac{3(c+1)}{2r(r-1)} r_2 \cos^2\theta_2$ ,

where  $\theta_1, \theta_2$  are the slant angles and  $2r_1, 2r_2$  are the dimensions of distributions  $\mathcal{D}_{\theta_1}$  and  $\mathcal{D}_{\theta_2}$ , respectively.

## 5 Optimal inequalities for bi-slant Riemannian maps to Kenmotsu space form involving Cosorati curvature

In this section, we obtain inequalities for bi-slant Riemannian maps from Riemannian manifolds to Kenmotsu space forms involving Cosorati curvatures in both cases when  $\xi \in \Gamma(range F_*)$  or  $\xi \in \Gamma((range F_*)^\perp)$ .

Let  $\mathcal{C}^{(ker F_*)^\perp}$  be the second fundamental form of the horizontal space  $(ker F_*)^\perp$  over manifold  $(M, \psi, \eta, \xi, g_2)$ , called the Casorati curvature of  $(ker F_*)^\perp$ . Hence, we have

$$\mathcal{C}^{(ker F_*)^\perp} = \frac{1}{r} \sum_{\alpha=r+1}^m \sum_{i,j=1}^r \left( c_{ij}^\alpha \right)^2. \quad (46)$$



Now, consider that  $\mathcal{L}^{(kerF_*)^\perp}$  be the  $s$  dimensional subspace of horizontal space  $(kerF_*)^\perp$ ,  $s \geq 2$ , with an orthogonal basis  $\{e_1, \dots, e_s\}$ , then the Cosorati curvature  $\mathcal{C}^{(kerF_*)^\perp}(\mathcal{L}^{(kerF_*)^\perp})$  of  $\mathcal{L}^{(kerF_*)^\perp}$  is defined as

$$\mathcal{C}^{(kerF_*)^\perp}(\mathcal{L}^{(kerF_*)^\perp}) = \frac{1}{s} \sum_{\alpha=r+1}^m \sum_{i,j=1}^s (\zeta_{ij}^\alpha)^2, \quad (47)$$

and the normalized  $\delta_C^{(kerF_*)^\perp}$ -Cosorati curvatures  $\delta_C^{(kerF_*)^\perp}(r-1)$  and  $\hat{\delta}_C^{(kerF_*)^\perp}(r-1)$  of  $(kerF_*)^\perp$  are defined as

$$[\delta_C^{(kerF_*)^\perp}(r-1)]_p = \frac{1}{2} \mathcal{C}_p^{(kerF_*)^\perp} + \frac{r+1}{2r} \text{inf} \{ \mathcal{C}^{(kerF_*)^\perp}(\mathcal{L}^{(kerF_*)^\perp}) | \mathcal{L}^{(kerF_*)^\perp} \text{ a hyperplane of } (kerF_*)^\perp \} \quad (48)$$

$$[\hat{\delta}_C^{(kerF_*)^\perp}(r-1)]_p = 2\mathcal{C}_p^{(kerF_*)^\perp} - \frac{2r-1}{2r} \text{inf} \{ \mathcal{C}^{(kerF_*)^\perp}(\mathcal{L}^{(kerF_*)^\perp}) | \mathcal{L}^{(kerF_*)^\perp} \text{ a hyperplane of } (kerF_*)^\perp \} \quad (49)$$

**Lemma 11.** [24] Let  $\Xi = \{(x_1, \dots, x_n) \in \mathbb{R}^n : x_1 + \dots + x_n = k\}$  be a hyperplane of  $\mathbb{R}^n$  and  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  be a quadratic form given by

$$f(x_1, \dots, x_n) = b \sum_{i=1}^{n-1} (x_i)^2 + d(y_i)^2 - 2 \sum_{1 \leq i < j \leq n} x_i x_j, \quad b, d > 0.$$

Then the constrained extremum problem  $\min_{(x_1, \dots, x_n) \in \Xi} f$  has the following solution

$$x_1 = \dots = x_{n-1} = \frac{k}{b+1}, \quad x_n = \frac{k}{d+1} = \frac{k(n-1)}{(b+1)d} = (b-n+2) \frac{k}{b+1}$$

provided that  $d = \frac{n-1}{b-n+2}$ .

**Theorem 12.** Let  $F : \bar{M} \rightarrow M(c)$  be a bi-slant Riemannian map from a Riemannian manifold  $\bar{M}$  to Kenmotsu space form  $M(c)$  with  $\xi \in \Gamma(\text{range}F_*)$  and  $3 \leq r = \text{rank}F < \min\{\bar{m}, m\}$ . Then, the normalized  $\mathcal{C}^{(kerF_*)^\perp}$ -Casorati curvatures  $\delta_C^{(kerF_*)^\perp}(r-1)$  and  $\hat{\delta}_C^{(kerF_*)^\perp}(r-1)$  of  $(kerF_*)^\perp$  satisfy

$$\varrho^{(kerF_*)^\perp} \leq \delta_C^{(kerF_*)^\perp}(r-1) + \frac{c-3}{4} - \frac{(c+1)}{2r} + \frac{3(c+1)}{2r(r-1)}(r_1 \cos^2 \theta_1 + r_2 \cos^2 \theta_2), \quad (50)$$

$$\varrho^{(kerF_*)^\perp} \leq \hat{\delta}_C^{(kerF_*)^\perp}(r-1) + \frac{c-3}{4} - \frac{(c+1)}{2r} + \frac{3(c+1)}{2r(r-1)}(r_1 \cos^2 \theta_1 + r_2 \cos^2 \theta_2), \quad (51)$$

where,  $\theta_1, \theta_2$  are the slant angles and  $2r_1, 2r_2$  are the dimensions of distributions  $\mathcal{D}_{\theta_1}$  and  $\mathcal{D}_{\theta_2}$ , respectively. Further, the case of equality holds in any of the above two inequalities at a point  $p \in \overline{M}$  if and only if with respect to suitable orthonormal bases  $\{e_1, \dots, e_r\}$  on  $(\ker F_*)^\perp$  and  $\{v_{r+1}, \dots, v_m\}$  on  $\Gamma((\text{range } F_*)_{F_*p})^\perp, \zeta = 0$ .

*Proof.* Since  $\overline{M}$  is a Riemannian manifold and  $M(c)$  is the Kenmotsu space form such that  $3 \leq r = \text{rank } F < \min\{\overline{m}, m\}$ , then from (7), (11) and (14)-(17), we have

$$\begin{aligned} 2\tau^{(\ker F_*)^\perp}(p) &= \frac{c-3}{4}r(r-1) + \frac{c+1}{4}\left\{2-2r+6(r_1\cos^2\theta_1+r_2\cos^2\theta_2)\right. \\ &\quad \left.- \sum_{\alpha=r+1}^m \sum_{i,j=1}^r (\zeta_{ij}^\alpha)^2 + \|\text{trace } \zeta\|^2\right\}, \end{aligned} \quad (52)$$

using (46) in (52), we get

$$\frac{c-3}{4}r(r-1) + \frac{c+1}{4}\left\{2-2r+6(r_1\cos^2\theta_1+r_2\cos^2\theta_2)\right\} = r\mathcal{C}^{(\ker F_*)^\perp} - \|\text{trace } \zeta\|^2 + 2\tau^{(\ker F_*)^\perp}(p). \quad (53)$$

Now, we consider following quadratic polynomial in term of the component of the second fundamental form of  $F$

$$\begin{aligned} \mathcal{P}^{(\ker F_*)^\perp} &= \frac{1}{2}r(r-1)\mathcal{C}^{(\ker F_*)^\perp} + \frac{1}{2}(r-1)(r+1)\mathcal{C}^{(\ker F_*)^\perp}(\mathcal{L}^{(\ker F_*)^\perp}) - 2\tau^{(\ker F_*)^\perp}(p) \\ &\quad + \frac{c-3}{4}r(r-1) + \frac{c+1}{4}\left\{2-2r+6(r_1\cos^2\theta_1+r_2\cos^2\theta_2)\right\}, \end{aligned} \quad (54)$$

where  $\mathcal{L}^{(\ker F_*)^\perp}$  is hyperplane of  $(\ker F_*)^\perp$ , without loss of generality, we assume that the hyperplane  $\mathcal{L}^{(\ker F_*)^\perp}$  is spanned by  $e_1, \dots, e_{r-1}$ , from (53) and (54), we have

$$\begin{aligned} \mathcal{P}^{(\ker F_*)^\perp} &= \sum_{\alpha=r+1}^m \left\{ 2(r+1) \sum_{1 \leq i < j \leq r-1} (\zeta_{ij}^\alpha)^2 - 2 \sum_{1 \leq i < j \leq r-1} \zeta_{ii}^\alpha \zeta_{jj}^\alpha + (r+1) \sum_{i=1}^{r-1} (\zeta_{ir}^\alpha)^2 \right. \\ &\quad \left. + r \sum_{i=1}^{r-1} (\zeta_{ii}^\alpha)^2 + \frac{r-1}{2} (\zeta_{rr}^\alpha)^2 \right\}, \\ &\geq \sum_{\alpha=r+1}^m \left\{ r \sum_{i=1}^{r-1} (\zeta_{ii}^\alpha)^2 + \frac{r-1}{2} (\zeta_{rr}^\alpha)^2 - 2 \sum_{1 \leq i < j \leq r-1} \zeta_{ii}^\alpha \zeta_{jj}^\alpha \right\}. \end{aligned}$$

Now, for  $\alpha = r+1, \dots, m$ , considering the quadratic form  $f_\alpha : \mathbb{R}^n \rightarrow \mathbb{R}$ , defined by

$$f_\alpha(\zeta_{11}^\alpha, \dots, \zeta_{rr}^\alpha) = r \sum_{i=1}^{r-1} (\zeta_{ii}^\alpha)^2 + \frac{r-1}{2} (\zeta_{rr}^\alpha)^2 - 2 \sum_{1 \leq i < j \leq r} \zeta_{ii}^\alpha \zeta_{jj}^\alpha,$$

then the constrained extremum problem  $\min f_\alpha$ , subjected to

$$\zeta^\alpha : \zeta_{11}^\alpha + \dots + \zeta_{rr}^\alpha = k^\alpha,$$

where  $k^\alpha$  is a real constant. Then from Lemma 5.1, we get  $b = r$ ,  $d = \frac{r-1}{2}$  and the critical points  $(\zeta_{11}^\alpha, \dots, \zeta_{rr}^\alpha)$  given by

$$\zeta_{11}^\alpha = \zeta_{22}^\alpha = \dots = \zeta_{r-1,r-1}^\alpha = \frac{k^\alpha}{r+1}, \quad \zeta_{rr}^\alpha = \frac{2k^\alpha}{r+1}$$

is a global minimum point, also  $f_\alpha(\zeta_{11}^\alpha, \dots, \zeta_{rr}^\alpha) = 0$ , which implies that  $\mathcal{P}^{(ker F_*)^\perp} \geq 0$ . Hence, we have

$$\begin{aligned} \varrho^{(ker F_*)^\perp} &\leq \frac{1}{2} \mathcal{C}^{(ker F_*)^\perp} + \frac{r+1}{2r} \mathcal{C}^{(ker F_*)^\perp} (\mathcal{L}^{(ker F_*)^\perp}) + \frac{c-3}{4} - \frac{c+1}{2r} \\ &\quad + \frac{3(c+1)}{2r(r-1)} (r_1 \cos^2 \theta_1 + r_2 \cos^2 \theta_2), \end{aligned} \quad (55)$$

for all hyperplane  $\mathcal{L}^{(ker F_*)^\perp}$  of  $(ker F_*)^\perp$ . Similarly considering another quadratic polynomial

$$\begin{aligned} \mathcal{Q}^{(ker F_*)^\perp} &= 2r(r-1) \mathcal{C}^{(ker F_*)^\perp} - \frac{1}{2} (r-1)(2r-1) \mathcal{C}^{(ker F_*)^\perp} (\mathcal{L}^{(ker F_*)^\perp}) - 2\tau^{(ker F_*)^\perp}(p) \\ &\quad + \frac{c-3}{4} r(r-1) + \frac{c+1}{4} \left\{ 2 - 2r + 6(r_1 \cos^2 \theta_1 + r_2 \cos^2 \theta_2) \right\}, \end{aligned}$$

proceeding the similar way as above, we get  $\mathcal{Q}^{(ker F_*)^\perp} \geq 0$ , therefore we have

$$\varrho^{(ker F_*)^\perp} \leq 2\mathcal{C}_p^{(ker F_*)^\perp} - \frac{2r-1}{2r} \mathcal{C}^{(ker F_*)^\perp} (\mathcal{L}^{(ker F_*)^\perp}) + \frac{c-3}{4} - \frac{c+1}{2r} \quad (56)$$

$$+ \frac{3(c+1)}{2r(r-1)} (r_1 \cos^2 \theta_1 + r_2 \cos^2 \theta_2), \quad (57)$$

for all hyperplanes  $\mathcal{L}^{(ker F_*)^\perp}$  of  $(ker F_*)^\perp$ . Now taking the infimum in (55) and the supremum in (56) over all hyperplanes  $\mathcal{L}^{(ker F_*)^\perp}$ , we get (50) and (51). Also analyzing the equality case for  $\mathcal{P}^{(ker F_*)^\perp} \geq 0$ . and  $\mathcal{Q}^{(ker F_*)^\perp} \geq 0$ , we get the equality conditions

$$\begin{cases} \zeta_{11}^\alpha = \zeta_{22}^\alpha = \dots = \zeta_{r-1,r-1}^\alpha = \frac{1}{2} \zeta_{rr}^\alpha, \\ \zeta_{ij}^\alpha = 0, \quad i, j = 1, \dots, r; i \neq j. \end{cases}$$

Since,  $\zeta_{rr}^\alpha = 0$  as  $F_* e_r = \xi$ , we conclude the result.  $\square$

**COROLLARY 5.1.** *Let  $F : \overline{M} \rightarrow M(c)$  be a Riemannian map from a Riemannian manifold  $\overline{M}$  to Kenmotsu space form  $M(c)$  with  $\xi \in \Gamma(\text{range } F_*)$  and  $3 \leq r = \text{rank } F <$*

$\min\{\overline{m}, m\}$ . Then, for normalized  $\mathcal{C}^{(kerF_*)^\perp}$ -Casorati curvatures  $\delta_C^{(kerF_*)^\perp}(r-1)$  and  $\hat{\delta}_C^{(kerF_*)^\perp}(r-1)$  of  $(kerF_*)^\perp$  and different Riemannian maps, we have Table 1

Riemannian Maps	$r_1 \& r_2$	$\theta_1$	$\theta_2$	normalized $\mathcal{C}^{(kerF_*)^\perp}$ -Casorati curvatures relation
Invariant	$r_1 = 0$	-	0	$\varrho^{(kerF_*)^\perp} \leq \delta_C^{(kerF_*)^\perp}(r-1) + \frac{c-3}{4} + \frac{c+1}{r}$ , $\varrho^{(kerF_*)^\perp} \leq \hat{\delta}_C^{(kerF_*)^\perp}(r-1) + \frac{c-3}{4} + \frac{c+1}{r}$ ,
Anti-invariant	$r_1 = 0$	-	$\frac{\pi}{2}$	$\varrho^{(kerF_*)^\perp} \leq \delta_C^{(kerF_*)^\perp}(r-1) + \frac{c-3}{4}$ , $\varrho^{(kerF_*)^\perp} \leq \hat{\delta}_C^{(kerF_*)^\perp}(r-1) + \frac{c-3}{4}$ ,
Semi-invariant	$r_1, r_2 \neq 0$	0	$\frac{\pi}{2}$	$\varrho^{(kerF_*)^\perp} \leq \delta_C^{(kerF_*)^\perp}(r-1) + \frac{c-3}{4} + \frac{c+1}{2r} \left\{ \frac{3r_1}{r-1} - \frac{1}{2} \right\}$ , $\varrho^{(kerF_*)^\perp} \leq \hat{\delta}_C^{(kerF_*)^\perp}(r-1) + \frac{c-3}{4} + \frac{c+1}{2r} \left\{ \frac{3r_1}{r-1} - \frac{1}{2} \right\}$ ,
Proper slant	$r_1 = 0$	-	$\in (0, \frac{\pi}{2})$	$\varrho^{(kerF_*)^\perp} \leq \delta_C^{(kerF_*)^\perp}(r-1) + \frac{c-3}{4} + \frac{c+1}{4r} (3\cos^2\theta_2 - 1)$ , $\varrho^{(kerF_*)^\perp} \leq \hat{\delta}_C^{(kerF_*)^\perp}(r-1) + \frac{c-3}{4} + \frac{c+1}{4r} (3\cos^2\theta_2 - 1)$ ,
Semi-slant	$r_1, r_2 \neq 0$	0	$\in (0, \frac{\pi}{2})$	$\varrho^{(kerF_*)^\perp} \leq \delta_C^{(kerF_*)^\perp}(r-1) + \frac{c-3}{4} + \frac{c+1}{2r} \left\{ \frac{3(r_1+r_2\cos^2\theta_2)}{r-1} - \frac{1}{2} \right\}$ , $\varrho^{(kerF_*)^\perp} \leq \hat{\delta}_C^{(kerF_*)^\perp}(r-1) + \frac{c-3}{4} + \frac{c+1}{2r} \left\{ \frac{3(r_1+r_2\cos^2\theta_2)}{r-1} - \frac{1}{2} \right\}$ ,
Hemi-slant	$r_1, r_2 \neq 0$	$\frac{\pi}{2}$	$\in (0, \frac{\pi}{2})$	$\varrho^{(kerF_*)^\perp} \leq \delta_C^{(kerF_*)^\perp}(r-1) + \frac{c-3}{4} + \frac{c+1}{2r} \left\{ \frac{3r_2\cos^2\theta_2}{r-1} - \frac{1}{2} \right\}$ , $\varrho^{(kerF_*)^\perp} \leq \hat{\delta}_C^{(kerF_*)^\perp}(r-1) + \frac{c-3}{4} + \frac{c+1}{2r} \left\{ \frac{3r_2\cos^2\theta_2}{r-1} - \frac{1}{2} \right\}$ ,

**Table 1** Normalized  $\mathcal{C}^{(kerF_*)^\perp}$ -Casorati curvatures relation, when  $\xi \in \Gamma(rangeF_*)$ .

where  $\theta_1, \theta_2$  are the slant angles and  $2r_1, 2r_2$  are the dimensions of distributions  $\mathcal{D}_{\theta_1}$  and  $\mathcal{D}_{\theta_2}$ , respectively. Further, the case of equality holds in any of the above two inequalities at a point  $p \in \overline{M}$  if and only if with respect to suitable orthonormal bases  $\{e_1, \dots, e_r\}$  on  $(kerF_*)^\perp_p$  and  $\{v_{r+1}, \dots, v_m\}$  on  $\Gamma((rangeF_*)_{F_*p})^\perp$ ,  $\zeta^\alpha = 0$ .

Further, if  $\xi \in \Gamma((rangeF_*)^\perp)$ , then we have the following result, which can be prove same as above theorem.

**Theorem 13.** Let  $F : \overline{M} \rightarrow M(c)$  be a bi-slant Riemannian map from a Riemannian manifold  $\overline{M}$  to Kenmotsu space form  $M(c)$  with  $\xi \in \Gamma((rangeF_*)^\perp)$  and  $3 \leq r = \text{rank}F < \min\{\overline{m}, m\}$ . Then, the normalized  $\mathcal{C}^{(kerF_*)^\perp}$ -Casorati curvatures

$\delta_C^{(kerF_*)^\perp}(r-1)$  and  $\hat{\delta}_C^{(kerF_*)^\perp}(r-1)$  of  $(kerF_*)^\perp$  satisfy

$$\varrho^{(kerF_*)^\perp} \leq \delta_C^{(kerF_*)^\perp}(r-1) + \frac{c-3}{4} + \frac{3(c+1)}{2r(r-1)}(r_1 \cos^2 \theta_1 + r_2 \cos^2 \theta_2), \quad (58)$$

$$\varrho^{(kerF_*)^\perp} \leq \hat{\delta}_C^{(kerF_*)^\perp}(r-1) + \frac{c-3}{4} + \frac{3(c+1)}{2r(r-1)}(r_1 \cos^2 \theta_1 + r_2 \cos^2 \theta_2), \quad (59)$$

where  $\theta_1, \theta_2$  are the slant angles and  $2r_1, 2r_2$  are the dimensions of distributions  $\mathcal{D}_{\theta_1}$  and  $\mathcal{D}_{\theta_2}$ , respectively. Further, the case of equality holds in any of the above two inequalities at a point  $p \in \overline{M}$  if and only if with respect to suitable orthonormal bases  $\{e_1, \dots, e_r\}$  on  $(kerF_*)^\perp_p$  and  $\{v_{r+1}, \dots, v_m\}$  on  $\Gamma((rangeF_*)_{F_*p})^\perp$ , the components of  $\zeta$  satisfy

$$\begin{cases} \zeta_{11}^\alpha = \zeta_{22}^\alpha = \dots = \zeta_{r-1r-1}^\alpha = \frac{1}{2}\zeta_{rr}^\alpha, \\ \zeta_{ij}^\alpha = 0, \quad i \neq j = 1, \dots, r. \end{cases} \quad (60)$$

**COROLLARY 5.2.** Let  $F : \overline{M} \rightarrow M(c)$  be a Riemannian map from a Riemannian manifold  $\overline{M}$  to Kenmotsu space form  $M(c)$  with  $\xi \in \Gamma((rangeF_*)^\perp)$  and  $3 \leq r = \text{rank}F < \min\{\overline{m}, m\}$ . Then, for normalized  $\mathcal{C}^{(kerF_*)^\perp}$ -Casorati curvatures  $\delta_C^{(kerF_*)^\perp}(r-1)$  and  $\hat{\delta}_C^{(kerF_*)^\perp}(r-1)$  of  $(kerF_*)^\perp$  and different Riemannian maps we have Table 2

where  $\theta_1, \theta_2$  are the slant angles and  $2r_1, 2r_2$  are the dimensions of distributions  $\mathcal{D}_{\theta_1}$  and  $\mathcal{D}_{\theta_2}$ , respectively. Further, the case of equality holds in any of the above two inequalities at a point  $p \in \overline{M}$  if and only if with respect to suitable orthonormal bases  $\{e_1, \dots, e_r\}$  on  $(kerF_*)^\perp_p$  and  $\{v_{r+1}, \dots, v_m\}$  on  $\Gamma((rangeF_*)_{F_*p})^\perp$ , the components of  $\zeta$  satisfy

$$\begin{cases} \zeta_{11}^\alpha = \zeta_{22}^\alpha = \dots = \zeta_{r-1r-1}^\alpha = \frac{1}{2}\zeta_{rr}^\alpha, \\ \zeta_{ij}^\alpha = 0, \quad i \neq j = 1, \dots, r. \end{cases} \quad (61)$$

Riemannian Maps	$r_1$ & $r_2$	$\theta_1$	$\theta_2$	normalized $\mathcal{C}^{(kerF_*)^\perp}$ -Casorati curvatures relation
Invariant	$r_1 = 0$	-	0	$\varrho^{(kerF_*)^\perp} \leq \delta_C^{(kerF_*)^\perp} (r-1) + \frac{c-3}{4} + \frac{3(c+1)}{4(r-1)},$ $\varrho^{(kerF_*)^\perp} \leq \hat{\delta}_C^{(kerF_*)^\perp} (r-1) + \frac{c-3}{4} + \frac{3(c+1)}{4(r-1)},$
Anti-invariant	$r_1 = 0$	-	$\frac{\pi}{2}$	$\varrho^{(kerF_*)^\perp} \leq \delta_C^{(kerF_*)^\perp} (r-1) + \frac{c-3}{4},$ $\varrho^{(kerF_*)^\perp} \leq \hat{\delta}_C^{(kerF_*)^\perp} (r-1) + \frac{c-3}{4},$
Semi-invariant	$r_1, r_2 \neq 0$	0	$\frac{\pi}{2}$	$\varrho^{(kerF_*)^\perp} \leq \delta_C^{(kerF_*)^\perp} (r-1) + \frac{c-3}{4} + \frac{3(c+1)r_1}{2r(r-1)},$ $\varrho^{(kerF_*)^\perp} \leq \hat{\delta}_C^{(kerF_*)^\perp} (r-1) + \frac{c-3}{4} + \frac{3(c+1)r_1}{2r(r-1)},$
Proper slant	$r_1 = 0$	-	$\in (0, \frac{\pi}{2})$	$\varrho^{(kerF_*)^\perp} \leq \delta_C^{(kerF_*)^\perp} (r-1) + \frac{c-3}{4} + \frac{3(c+1)}{4(r-1)} \cos^2 \theta_2,$ $\varrho^{(kerF_*)^\perp} \leq \hat{\delta}_C^{(kerF_*)^\perp} (r-1) + \frac{c-3}{4} + \frac{3(c+1)}{4(r-1)} \cos^2 \theta_2,$
Semi-slant	$r_1, r_2 \neq 0$	0	$\in (0, \frac{\pi}{2})$	$\varrho^{(kerF_*)^\perp} \leq \delta_C^{(kerF_*)^\perp} (r-1) + \frac{c-3}{4} + \frac{3(c+1)}{2r(r-1)} (r_1 + r_2 \cos^2 \theta_2),$ $\varrho^{(kerF_*)^\perp} \leq \hat{\delta}_C^{(kerF_*)^\perp} (r-1) + \frac{c-3}{4} + \frac{3(c+1)}{2r(r-1)} (r_1 + r_2 \cos^2 \theta_2),$
Hemi-slant	$r_1, r_2 \neq 0$	$\frac{\pi}{2}$	$\in (0, \frac{\pi}{2})$	$\varrho^{(kerF_*)^\perp} \leq \delta_C^{(kerF_*)^\perp} (r-1) + \frac{c-3}{4} + \frac{3(c+1)}{2r(r-1)} r_2 \cos^2 \theta_2,$ $\varrho^{(kerF_*)^\perp} \leq \hat{\delta}_C^{(kerF_*)^\perp} (r-1) + \frac{c-3}{4} + \frac{3(c+1)}{2r(r-1)} r_2 \cos^2 \theta_2,$

**Table 2** Normalized  $\mathcal{C}^{(kerF_*)^\perp}$ -Casorati curvatures relation, when  $\xi \in \Gamma((rangeF_*)^\perp)$ .

**Acknowledgements.** The first author is thankful to UGC for providing financial assistance in terms of MANF scholarship vide letter with UGC-Ref. No. 4844/(CSIR-UGC NET JUNE 2019). The second author is thankful to DST Gov. of India for providing financial support in terms of DST-FST label-I grant vide sanction number SR/FST/MS-I/2021/104(C).

## Declarations

**Author contributions.** All authors contributed equally.

**Data Availability.** There is no associate data.

**Conflict of interest.** There is no conflict of interest.

## References

- [1] Chen, B.-Y.: Some pinching and classification theorems for minimal submanifolds. *Arch. Math. (Basel)* **60**(6), 568-578 (1993).
- [2] Chen, B.-Y.: Relations between Ricci curvature and shape operator for submanifolds with arbitrary codimensions. *Glasg. Math. J.* **41**(1), 33-41 (1999).
- [3] Kaur, H., Shanker, G., Kaur, R., Mustafa, A.: Generalized Chen Inequality for CR-warped products of locally conformal Kähler manifolds, *Honam Mathematical J.* **46**(1), 47-59 (2024).
- [4] Kılıç, E., Tripathi, M.M., Gülbahar, M.: Chen-Ricci inequalities for submanifolds of Riemannian and Kaehlerian product manifolds. In *Annales Polonici Mathematici*. Instytut Matematyczny Polskiej Akademii Nauk **116**, 37-56 (2016).
- [5] Tripathi, M.M.: Chen-Ricci inequality for submanifolds of contact metric manifolds. *Journal of Advanced Mathematical Studies*, **1**(1-2), 111-135 (2008).
- [6] Chen, B.-Y.: Riemannian submersions, minimal immersions and cohomology class. *Proc. Japan Acad. Ser. A Math. Sci.* **81**(10), 162-167 (2005).
- [7] Chen, B.-Y.: Examples and classification of Riemannian submersions satisfying a basic equality. *Bull. Austral. Math. Soc.* **72**(3), 391-402 (2005).
- [8] Akyol, M.A.: Chen inequalities for slant Riemannian submersions from cosymplectic space forms. *Filomat*, **37**(11), 3615-3629 (2023).
- [9] Lone, M.A., Wani, T.A.: On optimal inequalities for anti-invariant Riemannian submersions from conformal Sasakian space forms. *Advances in Applied Clifford Algebras*, **34**(1), 8 (2024).
- [10] Önen Poyraz, N., Gündüzalp, Y., Akyol, M.A.: Some inequalities for bi-slant Riemannian submersions in complex space forms. *International Journal of Geometric Methods in Modern Physics* **21**(8), 2450150-1660 (2024).
- [11] Lone, M.A., Shahid, M.H., Vîlcu, G.E.: On Casorati curvatures of submanifolds in pointwise Kenmotsu space forms. *Math. Phys. Anal. Geom.* **22**(2) (2019).
- [12] Siddiqui, A.N.: Optimal Casorati inequalities on bi-slant submanifolds of generalized Sasakian space forms. *Tamkang Journal of Mathematics* **49**(3), 235-255 (2018).
- [13] Lee, C.W., Lee, J.W., Vîlcu, G.E.: A new proof for some optimal inequalities involving generalized normalized  $\delta$ -Casorati curvatures. *Journal of Inequalities and Applications* **2015**(310), 1-9 (2015).

- [14] Lee, C. W., Lee, J. W., Şahin, B., Vilcu, G. E.: Optimal inequalities for Riemannian maps and Riemannian submersions involving Casorati curvatures. *Annali di Matematica Pura ed Applicata (1923-)* **200**, 1277-1295 (2021).
- [15] Gündüzalp, Y., Akyol, M. A.: Pointwise slant Riemannian maps from Kähler manifolds. *Journal of Geometry and Physics* **179**, 104589 (2022).
- [16] Gündüzalp, Y., Akyol, M.A., Şahin, B.: Pointwise hemi-slant Riemannian maps into almost Hermitian manifolds and Casorati inequalities. *International Journal of Maps in Mathematics*, **7**(1), 76-96 (2024).
- [17] Gündüzalp, Y., Akyol, M. A.: Pointwise semi-slant Riemannian (PSSR) maps from almost Hermitian manifolds. *Filomat* **37**(13), 4271-4286 (2023).
- [18] Blair, D. E.: *Lecture notes in Mathematics*. Springer-Verlag Berlin Heidelberg, New York (1976).
- [19] Kenmotsu, K.: A class of almost contact Riemannian manifolds. *Tohoku Mathematical Journal, Second Series* **24**(1), 93-103 (1972).
- [20] Fisher A. E.: Riemannian maps between Riemannian manifolds. *Contemp. Math.* **132**, 331-336 (1992).
- [21] Şahin B.: *Riemannian Submersions, Riemannian Maps in Hermitian Geometry and their Applications*. Cambridge, Elsevier Academic Press (2017).
- [22] Lee, J.W., Lee, C.W., Şahin, B., Vilcu, G.E.: Chen-Ricci inequalities for Riemannian maps and their applications. (2022)
- [23] Lu, Z.: Normal scalar curvature conjecture and its applications. *Journal of Functional Analysis* **261**(5): 1284-1308 (2011).
- [24] Tripathi, M.M.: Inequalities for algebraic Casorati curvatures and their applications. arXiv preprint arXiv:1607.05828 (2016).
- [25] Zaidi, A., Shanker, G., Yadav, J.: Conformal pointwise slant Riemannian maps from or to Kähler manifolds. *Journal of Geometry and Physics* **198**, 105113 (2024).
- [26] Zaidi, A., Shanker, G., Yadav, A.: Conformal anti-invariant Riemannian map from or to Sasakian manifold. *Lobachevskii Journal of Mathematics* **44**(4), 1509-1518 (2023).
- [27] Zaidi A., Shanker G.: Clairaut anti-invariant Riemannian maps to trans-Sasakian manifolds. *Filomat* **38**(11), 3777-3789 (2024).