

# A formula for the $\alpha$ -Futaki character

Kartick Ghosh\*

September 4, 2024

## Abstract

Álvarez-Cónsul–García-Fernández–García-Prada introduced the Kähler-Yang-Mills equations. They also introduced the  $\alpha$ -Futaki character, an analog of the Futaki invariant, as an obstruction to the existence of the Kähler-Yang-Mills equations. The equations depend on a coupling constant  $\alpha$ . Solutions of these equations with coupling constant  $\alpha > 0$  are of utmost importance. In this paper, we provide a formula for the  $\alpha$ -Futaki character on certain ample line bundles over toric manifolds. We then show that there are no solutions with  $\alpha > 0$  on certain ample line bundles over certain toric manifolds and compute the value of  $\alpha$  if a solution exists. We also relate our result to the existence result of Keller-Friedman in dimension-two.

## 1 Introduction

The Kähler-Yang-Mills equations, introduced in ([2], [14]), are a system of equations coupling the scalar curvature of Kähler metric and a Hermitian-Yang-Mills connections of vector bundles. Suppose  $E$  is a holomorphic vector bundle over a Kähler manifold  $X$ . Then for a Kähler metric  $\omega$  on  $X$  and a Hermitian metric  $H$  on  $E$ , the Kähler-Yang-Mills equations are

$$\begin{aligned} \sqrt{-1} \bigwedge_{\omega} F_H &= \lambda Id, \\ \alpha_0 S(\omega) - \alpha_1 \bigwedge_{\omega}^2 \text{tr}(F_H \wedge F_H) &= c. \end{aligned} \tag{1}$$

Here  $\sqrt{-1} \bigwedge_{\omega} F_H \omega^n = n \sqrt{-1} F_H \wedge \omega^{n-1}$  and  $\lambda, c$  are topological constants. The equations depend on the coupling constants  $\alpha_0, \alpha_1$ . In this setting,  $\frac{\alpha_1}{\alpha_0} = \alpha$ .

If  $X$  is a compact Riemann surface, the term  $\text{tr}(F_H \wedge F_H)$  vanishes and hence the system decouples. So for compact Riemann surfaces, the existence problem is equivalent to the uniformisation theorem and the Narasimhan-Seshadri theorem([20], [8]).

The existence problem in higher dimensions is very difficult because in this case, the system is truly a system of coupled non-linear partial differential equations. Despite this difficulty, there are some existence results: for small  $\alpha$  by perturbing a constant scalar curvature Kähler metric and a Hermitian-Yang-Mills connection([2]), applying the dimensional reduction of

---

\*kartick@math.tifr.res.in, School of Mathematics, TIFR Mumbai.

the Kähler-Yang-Mills equations from  $\Sigma \times \mathbb{P}^1$  to  $\Sigma$ , where  $\Sigma$  is a compact Riemann surface ([3],[4],[15],[5],[24]), simultaneous deformation of complex structure of  $X$  and  $E$  ([16]), on line bundles over ruled manifolds using the Calabi ansatz ([18]).

Álvarez-Cónsul–García-Fernández–García-Prada also introduced the  $\alpha$ -Futaki character  $\mathcal{F}$ , an analog of the classical Futaki invariant, which is an obstruction to the existence of the Kähler-Yang-Mills equations (1). This character does not depend on the choice of a metric  $H$  on  $E$  and a metric  $\omega$  on  $X$ . The computation of this character is known only in some specific cases ([4],[5]). Also, there is no formula for this character analogous to Mabuchi's formula for the Futaki invariant on toric manifolds ([19]) and residue formula for the Futaki invariant ([22], [13]).

Álvarez-Cónsul–García-Fernández–García-Prada remarked in [2] that even for line bundles, the existence problem is difficult (page 2807 in [2]). In this paper, we give a formula of the  $\alpha$ -Futaki invariant for certain ample line bundles on toric manifolds.

Suppose  $(X, \omega)$  is a toric manifold and  $L$  is an ample line bundle over it. All the Kähler forms will be torus invariant. Take a metric  $h$  on  $L$  such that its curvature  $\beta = \sqrt{-1}F_h$  is Kähler. We will be concerned with only those ample bundles  $L$  such that there exists a metric  $\omega' \in [\omega]$  which solves the equation

$$n\beta \wedge \omega'^{n-1} = \lambda \omega'^n.$$

This equation is known as the  $J$ -equation which was introduced by Donaldson ([9]), Chen ([6]) and studied on toric manifolds by Yao ([23]) and Collins-Székelyhidi ([7]).

If we calculate the invariant  $\mathcal{F}$  with this choice of metrics  $h$  and  $\omega'$ , then we get our main theorem.

**Theorem 1.1.** *Let  $P$  and  $Q$  be the moment polytopes of  $\omega'$  and  $\beta$  respectively. Then there is a diffeomorphism  $U : P \rightarrow Q$  such that  $\text{tr}(DU) = \lambda$  and the following holds*

$$(2\pi)^{-n} \langle \mathcal{F}, \xi_i \rangle = \frac{\alpha_0}{2} \int_{\partial P} (x_i + c_i) d\sigma + \alpha_1 \int_P (x_i + c_i) \left( \sum_{m \in M_2} m \right) d\mu.$$

Here  $M_2$  is the set of all  $(2 \times 2)$  principal minor of  $DU$ ,  $\xi_i$  is a holomorphic vector field on the associated principal  $\mathbb{C}^*$ -bundle of  $L$ ,  $d\mu$  is the Lebesgue measure on  $P$ ,  $d\sigma$  is a positive measure on  $\partial P$  normalized so that on a face  $l_i = 0$  we have  $d\sigma \wedge dl_i = \pm d\mu$ ,  $c_i$  is a constant such that  $\int_P (x_i + c_i) d\mu = 0$ .

We explain the notations in section (2) and section (3). As an immediate corollary of this theorem, we get the following.

**Corollary 1.2.** *If Solution of Kähler-Yang-Mills equations exist with  $\alpha_0 \alpha_1 \neq 0$  then  $\langle \mathcal{F}, \xi_i \rangle = 0 \forall i$ . Therefore, we get*

$$\frac{\int_{\partial P} (x_i + c_i) d\sigma}{\int_P (x_i + c_i) \sum_{m \in M_2} m d\mu} = \frac{\int_{\partial P} (x_j + c_j) d\sigma}{\int_P (x_j + c_j) \sum_{m \in M_2} m d\mu},$$

for all  $1 \leq i, j \leq n$ .

On  $\mathbb{P}^2 \# \overline{\mathbb{P}^2}$ , we compute the  $\alpha$ -Futaki character for certain ample line bundles. We prove the following theorem.

**Theorem 1.3.** *Suppose  $L$  is an ample line bundle on  $\mathbb{P}^2 \# \overline{\mathbb{P}^2}$  with  $c_1(L) = a[H_2] - [E_2]$  and  $[\omega] = b[H_2] - [E_2]$  satisfying  $2\frac{ab-1}{b^2-1} > 1$ . Then there is no solution of the Kähler-Yang-Mills equations with  $\frac{\alpha_1}{\alpha_0} > 0$ . Furthermore, If solution exists then  $\frac{\alpha_1}{\alpha_0} = -\frac{(b^2-1)}{(b-a)^2}$ .*

We explain the notations in 4. For some of these line bundles, there do exist a solution of the Kähler-Yang-Mills equations with the prescribed value of  $\frac{\alpha_1}{\alpha_0}$ . The existence result is due to Keller-Friedman([18]).

**Theorem 1.4** (Keller-Friedman). *There exists metrics  $h$  on the line bundle corresponding to the integral cohomology class  $(8k+3)[H_2] - [E_2]$  and  $\omega \in 3[H_2] - [E_2]$  such that they satisfy the Kähler-Yang-Mills equations with  $\frac{\alpha_1}{\alpha_0} = -\frac{1}{8k^2}$  (Here  $k \in \mathbb{N}$ ).*

We also compute the  $\alpha$ -Futaki character on some ample line bundles on  $\mathbb{P}^3 \# \overline{\mathbb{P}^3}$ .

**Theorem 1.5.** *Suppose  $L$  is an ample line bundle on  $\mathbb{P}^3 \# \overline{\mathbb{P}^3}$  with  $c_1(L) = a[H_3] - [E_3]$  and  $[\omega] = b[H_3] - [E_3]$  satisfying  $3\frac{ab^2-1}{b^3-1} > 2$ . Then there is no solution of Kähler-Yang-Mills equations with  $\frac{\alpha_1}{\alpha_0} > 0$ . Furthermore, If solution exists then  $\frac{\alpha_1}{\alpha_0} = -\frac{(3b+1)(b-1)^3(b^2+b+1)}{3b(b-a)^2(b^3+b^2-b+1)}$ .*

The author introduced coupled Kähler-Einstein and Hermitian-Yang-Mills equations([17]). Suppose  $(X, \omega)$  is a compact Kähler  $n$ -manifold with  $-\sqrt{-1}\omega$  as the curvature form of the Chern connection of  $(K_X^{-1}, h)$ . One can view the metric  $h$  as a volume form  $\Omega_h$ . Let  $\pi : E \rightarrow X$  be a holomorphic vector bundle and  $H$  be a Hermitian metric on the bundle. The coupled Kähler-Einstein and Hermitian-Yang-Mills equations are

$$\begin{aligned} \sqrt{-1} \bigwedge_{\omega} F_H &= \lambda Id \\ \frac{\alpha_0}{2} \left( \frac{\Omega_h}{\int_X \Omega_h} - \frac{\omega^n}{\int_X \omega^n} \right) - \alpha_1 \frac{2}{(n-2)!} \text{tr}(F_H \wedge F_H) \wedge \omega^{n-2} &= \tilde{C} \text{vol}_{\omega}, \end{aligned} \quad (2)$$

where  $\alpha_0, \alpha_1$  are coupling constants and  $\tilde{C}, \lambda$  are topological constants. One can show the following.

**Lemma 1.6** (Lemma 3.1.1 in [17]). *Suppose  $(X, \omega)$  is a Fano manifold and  $f$  is the Ricci potential that is,  $\text{Ric}(\omega) - \omega = \sqrt{-1} \partial \bar{\partial} f$  normalized so that  $\int_X e^f \omega^n = \int_X \omega^n = V$ . Then*

$$\int_X \theta_v S(\omega) \omega^n = V \int_X \theta_v \left( \frac{\Omega_h}{\int_X \Omega_h} - \frac{\omega^n}{\int_X \omega^n} \right)$$

where  $\theta_v$  is the Hamiltonian function corresponding to a holomorphic vector field  $v$  (that is,  $i_v \omega = \partial \bar{\theta}_v$ ) and  $h$  is a metric on  $K_X^{-1}$  whose curvature is  $\omega$ . The function  $\theta_v$  is normalized by requiring  $\int_X \theta_v \omega^n = 0$ .

With the help of this lemma, one can identify the  $\alpha$ -Futaki character as an obstruction for the existence of coupled Kähler-Einstein and Hermitian-Yang-Mills equations albeit when  $\omega \in c_1(X)$  and  $X$  is Fano. The author proved an existence result for line bundles over

$\mathbb{P}(\mathcal{O} \oplus \mathcal{O}(-1)) \rightarrow \mathbb{P}^1$ . In this paper, we prove that these line bundles are not ample. Our paper is organized as follows: In section two, we discuss about some basic facts of toric Kähler geometry, Futaki invariants, and  $J$ -equation on toric manifolds. In section three, we recall the  $\alpha$ -Futaki character and prove theorem 1.1 and corollary 1.2. In section four, we prove theorem 1.3, 1.4, 1.5.

**Acknowledgements:** The author would like to thank Prof. Vamsi Pritham Pingali for various useful conversations about the project. The author would also like to thank him for reading a draft of the paper and giving valuable feedbacks. The author would like to thank Prof. Ved V. Datar for a fruitful discussion at IIT Guwahati during the 38th RMS annual conference. The author would like to thank the organizers of the 38th RMS annual conference for the invitation. The author was supported by a post-doctoral fellowship from the Tata Institute of Fundamental Research, Mumbai.

## 2 Preliminaries

In this section, we briefly discuss some basic notions of toric Kähler geometry, the Futaki invariant and the  $J$ -equation on toric manifolds.

### 2.1 Kähler geometry of toric manifolds

**Definition 1.** A Kähler manifold  $(X, \omega)$  is called a toric manifold if it admits an effective Hamiltonian action of the torus  $(S^1)^n$  which extends to a holomorphic action of the complexified torus  $(\mathbb{C}^*)^n$  with a free open dense orbit  $\approx (\mathbb{C}^*)^n$ .

The Hamiltonian action induces a moment map  $\mu : X \rightarrow \mathbb{R}^n$ , where the dual of the Lie algebra of the torus is identified with  $\mathbb{R}^n$ . By the Atiyah-Guillemin-Sternberg theorem, the closure of the image is a convex polytope. The size of the polytope is determined by the Kähler class while its position is determined by a choice of moment map or normalization of the Kähler potential. There is a one-to-one correspondence between toric manifolds and a class of convex polytopes known as Delzant polytope.

**Definition 2.** A convex polytope is said to be a Delzant polytope if every vertex of the polytope has a neighbourhood which is  $SL(n, \mathbb{Z})$  equivalent to  $\{y \in \mathbb{R}^n | y_i \geq 0; i = 1, \dots, n\}$ .

A Delzant polytope is said to be integral if all the vertex points are in  $\mathbb{Z}^n$ . A Delzant polytope  $P$  is defined by

$$P = \{y \in \mathbb{R}^n | l_i(y) = \langle y, v_i \rangle + \lambda_i > 0; i = 1, \dots, N\}, \quad (3)$$

where  $v_i \in \mathbb{Z}^n$  are primitive integral vectors and  $\lambda_i \in \mathbb{R}$ . All the irreducible toric divisors  $D_i$  are given by the inverse images of the facets  $l_i = 0$  under the moment map. The cohomology class of the Kähler form  $\omega$  is given by

$$[\omega] = \sum_{i=1}^N \lambda_i [D_i],$$

where  $[D_i]$  are Poincarè dual of  $D_i \in H_{2n-2}(X, \mathbb{Z})$ . If the Delzant polytope is integral then the Kähler class is integral and hence first Chern class of some ample line bundle.

Torus invariant Kähler metrics take a very special form on toric manifolds. A torus  $((S^1)^n)$  invariant metric  $\omega$  is of the form  $\omega|_{(\mathbb{C}^*)^n} = \sqrt{-1} \partial \bar{\partial} F$ ; where  $F : \mathbb{R}^n \rightarrow \mathbb{R}$  is a strictly convex function. The function  $F$  is called a Kähler potential of  $\omega$ . Consider the coordinates  $x_i = \log|z_i|^2$  and  $\theta_i$ , where  $(z_1, \dots, z_n)$  are holomorphic coordinates of  $(\mathbb{C}^*)^n$ . In this coordinates, the torus invariant metric takes the form

$$\omega|_{(\mathbb{C}^*)^n} = \sqrt{-1} \frac{\partial^2 F}{\partial x_i \partial x_j} dx^i \wedge d\theta^j.$$

Observe that the positivity of the form  $\omega$  is equivalent to the strict convexity of the Kähler potential. The moment map up to translation is given by  $\nabla F$ . Let us consider the coordinate  $y = \nabla F$ . Since  $F$  is strictly convex, we can take its Legendre transform. The Legendre transform of  $F$  satisfies

$$G(y) = \langle x, y \rangle + F(x).$$

The function  $G$  is called the symplectic potential of  $\omega$  and it is a strictly convex function on the moment polytope.

**Theorem 2.1.** *Let  $(X, \omega)$  be a toric Kähler manifold. Suppose the corresponding moment polytope is defined by (3). Then every torus invariant metric in the class  $[\omega]$  has a symplectic potential of the form*

$$G = \sum_{i=1}^N l_i(y) \log l_i(y) + v,$$

where  $v$  is smooth upto boundary.

## 2.2 The Futaki invariant

Futaki introduced an invariant as an obstruction to the existence of Kähler-Einstein metric ([12]). Later Calabi and Futaki generalized the invariants to any compact Kähler manifolds. This generalized invariant is an obstruction to the existence of Kähler metrics with constant scalar curvature. It is also a character of the Lie algebra of holomorphic vector fields.

**Definition 3.** *The Futaki invariant is given by*

$$\mathcal{Fut}(v) = \int_X \phi_v s(\omega) \omega^n,$$

where  $s(\omega)$  is the scalar curvature of  $\omega$ ,  $\phi_v$  is a holomorphy potential of  $v$  (i.e.,  $\bar{\partial} \phi_v = i_v \omega$ ). Moreover, if  $[\omega] = c_1(X)$ , then

$$\mathcal{Fut}_{c_1}(v) = \int_X v(h_\omega) \omega^n,$$

where  $h_\omega$  satisfies  $\text{Ric}(\omega) - \omega = \sqrt{-1} \partial \bar{\partial} h_\omega$ .

The map  $\mathcal{Fut}$  does not depend on the choice of  $\omega$ .

## 2.3 The $J$ -equation on toric manifolds

The  $J$ -equation was introduced by Donaldson([9]) and Chen([6]). Given a Kähler form  $\beta$  on a compact Kähler manifold  $X$ , it asks for a Kähler metric  $\omega$  such that

$$n\beta \wedge \omega^{n-1} = \lambda\omega^n, \quad (4)$$

where  $n = \dim_{\mathbb{C}} X$  and  $\lambda$  is a topological constant.

On Toric manifolds, Collins-Székelyhidi([7]) proved that  $J$ -equation is solvable iff certain inequalities are met. Yao also studied the  $J$ -equation on toric manifolds([23]).

Suppose  $\beta, \omega$  are two torus invariant Kähler metrics on toric manifolds  $X$ . They induce two moment maps  $\mu_{\omega} : X \rightarrow \bar{\mathcal{P}}_{\omega}$  and  $\mu_{\beta} : X \rightarrow \bar{\mathcal{P}}_{\beta}$ . Then there exists a transition map

$$U : \bar{\mathcal{P}}_{\omega} \rightarrow \bar{\mathcal{P}}_{\beta}. \quad (5)$$

Suppose  $u$  is a symplectic potential for  $\omega$  and  $f$  is a Kähler potential for  $\beta$ , then on the interior of  $P$ ,  $U$  is given by

$$U = \nabla f \circ \nabla u.$$

From the work of Yao(page 1586 in [23]), we have

$$tr_{\omega}\beta = \frac{n\beta \wedge \omega^{n-1}}{\omega^n} = tr(DU|_{\mu_{\omega}}); \quad \frac{\beta^n}{\omega^n} = det DU|_{\mu_{\omega}}; \quad n(n-1) \frac{\beta^2 \wedge \omega^{n-2}}{\omega^n} = \sum_{m \in M_2} m, \quad (6)$$

where  $M_2$  is the set of all  $(2 \times 2)$  principal minors of  $DU|_{\mu_{\omega}}$ .

## 3 The $\alpha$ -Futaki character with toric symmetry

In this section, we recall the  $\alpha$ -Futaki character introduced by Alvarez-Consul-Garcia-Fernandez-Garcia-Prada(see page 2775 in [2]).

Suppose  $\pi : \mathcal{E} \rightarrow X$  is the associated principal  $GL(r, \mathbb{C})$ -bundle of a holomorphic vector bundle  $E$  over  $X$ , where  $r = rank(E)$ . Suppose  $I$  is an integrable almost complex structure on  $\mathcal{E}$  that preserves the vertical bundle  $V\mathcal{E}(= ker(d\pi))$  and  $\check{I}$  is the unique integrable almost complex structure on the manifold  $X$  determined by  $I$ . Any vector field  $\xi \in LieAut(\mathcal{E}, I)$  covers a real holomorphic vector field  $\check{\xi}$  on  $(X, \check{I})$  which can be written as

$$\check{\xi} = \eta_{\phi_1} + \check{I}\eta_{\phi_2} + \beta,$$

for any Kähler form  $\omega$  in some Kähler class in  $(X, \check{I})$ , where  $\eta_{\phi_i}$  is the Hamiltonian vector field of  $\phi_i \in C_0^{\infty}(X, \omega)$  on  $(X, \omega)$  for  $i = 1, 2$ , and  $\beta$  is the dual of a 1-form which is harmonic with respect to the Kähler metric  $\omega(., \check{I}.)$ .

$$\begin{aligned} \mathcal{F}_I : LieAut(\mathcal{E}, I) &\rightarrow \mathbb{C} \\ \langle \mathcal{F}_I, \xi \rangle &= -4\sqrt{-1}\alpha_1 \int_X tr \left( \theta_H \xi \wedge (\sqrt{-1} \wedge_{\omega} F_H - \lambda Id) \right) \frac{\omega^n}{n!} \\ &+ \int_X \phi \left( \alpha_0 S_{\omega, \check{I}} \omega^n + n(n-1)\alpha_1 tr(\sqrt{-1} F_H \wedge \sqrt{-1} F_H \wedge \omega^{n-2}) \right) \\ &- \int_X \phi(4\alpha_1 \lambda) tr(\sqrt{-1} F_H \wedge \omega^{n-1}), \end{aligned} \quad (7)$$

where  $\phi = \phi_1 + \sqrt{-1}\phi_2$ . The functional  $\mathcal{F}_I$  does not depend on the choice of  $\omega$  and  $H$  and  $\mathcal{F}_I$  vanishes whenever the coupled equations have a solution. We omit the  $I$  in  $\mathcal{F}_I$  from now onwards. .

**Fact 1.** *If  $E$  is a line bundle then  $\mathcal{E}$  is a principal  $\mathbb{C}^*$ -bundle. It is a fact that any holomorphic vector field on  $X$  always lifts to a  $\mathbb{C}^*$ -invariant holomorphic vector field on a principal  $\mathbb{C}^*$ -bundle(see page 2806 in [2]). We will use this fact in the next subsection.*

### 3.1 Formula for certain ample line bundles on toric manifolds

Suppose  $L$  is an ample line bundle and  $\omega$  is a Kähler form on a toric manifold  $X$ . Also assume that there is a metric  $h$  on  $L$  such that for its curvature  $\sqrt{-1}F_h = \beta > 0$ , there is a metric  $\omega' \in [\omega]$  that solves the  $J$ -equation. We get the formula if we calculate the  $\alpha$ -Futaki character with  $h$  and  $\omega'$ . We denote by  $P, Q$ , the moment polytope of  $(X, \omega')$  and  $(X, \beta)$  respectively.

The manifold  $X$  has a free open dense subset  $(\mathbb{C}^*)^n$ . We denote the holomorphic coordinate of  $(\mathbb{C}^*)^n$  by  $(z_1, \dots, z_n)$ . The  $(\mathbb{C}^*)^n$ -invariant holomorphic vector fields  $v_i = z_i \frac{\partial}{\partial z_i}$  naturally extend to the toric manifold  $X$ . Suppose  $\xi_i$  is a lift of the vector field  $v_i$  on the associated principal  $\mathbb{C}^*$  bundle  $\mathcal{L}$ (using the fact 1). Then we have the following formula.

**Theorem 3.1.** *There is a diffeomorphism  $U : P \rightarrow Q$  such that  $\text{tr}(DU) = \lambda$  and the following holds*

$$(2\pi)^{-n} \langle \mathcal{F}, \xi_i \rangle = \frac{\alpha_0}{2} \int_{\partial P} (x^i + c_i) d\sigma + \alpha_1 \int_P (x^i + c_i) \left( \sum_{m \in M_2} m \right) d\mu.$$

Here  $M_2$  is the set of all  $(2 \times 2)$  principal minor of  $DU$ ,  $d\mu$  is the Lebesgue measure on  $P$ ,  $d\sigma$  is a positive measure on  $\partial P$  normalized so that on a face  $l_i = 0$  we have  $d\sigma \wedge dl_i = \pm d\mu$ , and  $\int_P (x_i + c_i) d\mu = 0$ .

*Proof.* Since for  $\sqrt{-1}F_h$ ,  $\omega'$  solves the  $J$ -equation, we have  $\sqrt{-1} \wedge_{\omega'} F_h = \lambda$ . Putting this on 7, we get

$$\langle \mathcal{F}, \xi_i \rangle = \int_X \phi_i (\alpha_0 S(\omega') \omega'^m + \alpha_1 n(n-1) (\sqrt{-1}F_h \wedge \sqrt{-1}F_h \wedge \omega'^{m-2})), \quad (8)$$

where  $i_{v_i} \omega' = \bar{\partial} \phi_i$  and  $\int_X \phi_i \omega'^m = 0$ . Now we write down how some terms correspond to similar terms in the polytope  $P$ . The top form  $\omega'^m$  corresponds to the measure  $(2\pi)^n d\mu$  on  $P$ , where  $d\mu$  is the Lebesgue measure on  $P$ . The function  $\phi_i \in C_0^\infty(X, \omega')$  corresponds to the affine function  $x^i + c_i$  with normalisation condition  $\int_P (x^i + c_i) d\mu = 0$ . Suppose  $u$  is the symplectic potential of  $\omega'$ , then from Abreu's equation([1]), we have

$$S(\omega') = -\frac{1}{2} \sum_{k,l} \frac{\partial^2 u^{kl}}{\partial x_k \partial x_l}.$$

Hence the first term of 8 becomes

$$\begin{aligned} & \int_X \phi_i S(\omega') \omega'^m \\ &= \frac{(2\pi)^n}{2} \int_{\partial P} (x^i + c_i) d\sigma. \end{aligned} \quad (9)$$

We have used Donaldson's formula([10], Lemma 4.37 in [21]): If  $u$  is a symplectic potential and  $h : P \rightarrow \mathbb{R}$  is a continuous convex function smooth on the interior, then

$$\int_P u^{kl} h_{kl} d\mu = \int_{\partial P} h d\sigma - \int_P h S(\omega') d\mu,$$

where  $d\mu$  is the Lebesgue measure on  $P$  and  $d\sigma$  is a positive measure on the boundary  $\partial P$  normalized so that on a face defined by  $l_i(y) = 0$  we have  $d\sigma \wedge dl_i = \pm d\mu$ .

Now putting 6 and 9 in 8, we get

$$(2\pi)^{-n} \langle \mathcal{F}, \xi_i \rangle = \frac{\alpha_0}{2} \int_{\partial P} (x^i + c_i) d\sigma + \alpha_1 \int_P (x^i + c_i) \left( \sum_{m \in M_2} m \right) d\mu$$

□

As an immediate consequence of the theorem, we get

**Corollary 3.2.** *If Solution of Kähler-Yang-Mills equations exist with  $\alpha_0 \alpha_1 \neq 0$  then  $\langle \mathcal{F}, \xi_i \rangle = 0 \ \forall i$ . Therefore, we get*

$$\frac{\int_{\partial P} (x_i + c_i) d\sigma}{\int_P (x_i + c_i) \sum_{m \in M_2} m d\mu} = \frac{\int_{\partial P} (x_j + c_j) d\sigma}{\int_P (x_j + c_j) \sum_{m \in M_2} m d\mu},$$

for all  $1 \leq i, j \leq n$ .

## 4 Solution of the $J$ -equation on $\mathbb{P}^n \# \overline{\mathbb{P}^n}$

Fang and Lai([11]) studied the  $J$ -equation on  $\mathbb{P}^n \# \overline{\mathbb{P}^n}$  imposing Calabi symmetry on both the metrics. Yao ([23]) cast Fang and Lai's solution in the toric setting as an example of his approach. The Calabi symmetry requires  $U(n)$  symmetry instead of  $U(1)^n$  symmetry as in the toric case. So to make the metrics to have  $U(n)$  symmetry, the symplectic potentials need to have more symmetries.

We take moment polytopes of  $\mathbb{P}^n \# \overline{\mathbb{P}^n}$  in the following form

$$P = \{x \in \mathbb{R}^n | x^i > 0, 1 < \sum_{i=1}^n x^i < b\},$$

where  $b$  determines the Kähler class. In this case, the Kähler class is  $[\omega] = b[H_n] - [E_n]$ , where  $H_n$  is the pull-back of the hyperplane divisor on  $\mathbb{P}^n$  and  $E_n$  is the exceptional divisor. Suppose  $[\beta] = a[H_n] - [E_n]$  and the corresponding polytope is  $Q$ .

One takes that the symplectic potentials of  $\omega$  and  $\beta$  have specific forms

$$u = \sum_{i=1}^n x^i \log x^i - X \log X + h(X), \quad X = \sum_{i=1}^n x^i,$$

where  $h$  is a strictly convex function defined on  $[1, b]$  satisfying

$$h(X) - (X - 1) \log(X - 1) - (b - X) \log(b - X) \in C^\infty[1, b]$$



and

$$v = \sum_{i=1}^n x^i \log x^i - X \log X + \theta(X),$$

where  $\theta$  is defined on  $[1, a]$  satisfying similar condition as  $h$ .

We denote the Legendre transform of  $h$  and  $\theta$  by  $p$  and  $\eta$  respectively. We define the map  $f(X) = \eta'(h'(X))$ . Then  $f : [1, b] \rightarrow [1, a]$  is a diffeomorphism. The transition map  $U : P \rightarrow Q$  in this case is given by

$$U^i(x) = f(X) \frac{x^i}{X} \quad \text{for } 1 \leq i \leq n.$$

Whenever  $n \frac{ab^{n-1}-1}{b^{n-1}} > n-1$ , the  $J$ -equation has a solution given by

$$\tilde{f}(r) = Ar + \frac{B}{r^{n-1}}, \quad (10)$$

where  $A = \frac{ab^{n-1}-1}{b^{n-1}}$  and  $B = \frac{b^n-ab^{n-1}}{b^{n-1}}$ .

Hence we have the following.

**Theorem 4.1** ([11], Theorem 2.3). *On  $\mathbb{P}^n \# \overline{\mathbb{P}^n}$ , we have two Kähler classes  $b[H_n] - [E_n]$ ,  $a[H_n] - [E_n]$ . Suppose  $\beta \in a[H_n] - [E_n]$  is a Kähler form. If  $n \frac{ab^{n-1}-1}{b^{n-1}} > (n-1)$ , then the equation 4 has a solution  $\omega \in a[H_n] - [E_n]$  given by 10 and the corresponding map  $U : P \rightarrow Q$  is given by*

$$U^i(x^1, \dots, x^n) = Ax^i + \frac{Bx^i}{X^n},$$

where  $X = \sum_{i=1}^n x^i$ ,  $A = \frac{ab^{n-1}-1}{b^{n-1}}$ ,  $B = \frac{b^n-ab^{n-1}}{b^{n-1}}$ ,  $P$  moment polytope of  $\omega$ ,  $Q$  moment polytope of  $\beta$ .

**Remark 1.** *Observe that one can work with Kähler classes of the form  $a[H_n] - b[E_n]$  with  $a > b > 0$  in the theorem 4.1 instead of Kähler classes of the form  $p[H_n] - [E_n]$  with  $p > 1$ . This will change the boundary values of the corresponding ODE (see Proposition 2.1 in [11]).*

## 4.1 The $\alpha$ -Futaki character on $\mathbb{P}^2 \# \overline{\mathbb{P}^2}$

We will work with Kähler classes of the following form  $p[H_2] - [E_2]$ , where  $H_2$  is the pull back of the hyperplane divisor from  $\mathbb{P}^2$ ,  $E_2$  is the exceptional divisor, and  $p > 1$ .

**Theorem 4.2.** *Suppose  $L$  is an ample line bundle on  $\mathbb{P}^2 \# \overline{\mathbb{P}^2}$  with  $c_1(L) = a[H_2] - [E_2]$  and  $[\omega] = b[H_2] - [E_2]$  satisfying  $2 \frac{ab-1}{b^2-1} > 1$ . Then there is no solution to the Kähler-Yang-Mills equations with  $\frac{\alpha_1}{\alpha_0} > 0$ . Furthermore, If a solution exists then  $\frac{\alpha_1}{\alpha_0} = -\frac{2b^2(b^2-1)}{(b^2-ab)^2}$ .*

*Proof.* From 4.1, we know that there is a solution of the  $J$ -equation and a corresponding map between moment polytopes. The map  $U : P_2 \rightarrow Q_2$  is given by

$$U(x^1, x^2) = \left( Ax^1 + \frac{Bx^1}{(x^1+x^2)^2}, Ax^2 + \frac{Bx^2}{(x^1+x^2)^2} \right),$$

where  $P_2 = \{(x^1, x^2) \in \mathbb{R}^2 | x^1 \geq 0, x^2 \geq 0, 1 \leq x^1 + x^2 \leq b\}$ ,  $Q_2 = \{(x^1, x^2) \in \mathbb{R}^2 | x^1 \geq 0, x^2 \geq 0, 1 \leq x^1 + x^2 \leq a\}$ ,  $A = \frac{ab-1}{b^2-1}$ , and  $B = \frac{b^2-ab}{b^2-1}$ .

A simple calculation gives us

$$DU = \begin{bmatrix} A + \frac{B(x^2-x^1)}{(x^1+x^2)^3} & -\frac{2Bx^1}{(x^1+x^2)^3} \\ -\frac{2Bx^2}{(x^1+x^2)^3} & A + \frac{B(x^1-x^2)}{(x^1+x^2)^3} \end{bmatrix}$$

and

$$\text{Det}(DU) = A^2 - \frac{B^2}{(x^1 + x^2)^4}.$$

A routine calculation gives

$$\begin{aligned} \int_{P_2} d\mu &= \frac{b^2 - 1}{2} \\ \int_{\partial P_2} d\sigma &= 3b - 1 \\ \int_{P_2} x^1 d\mu &= \frac{b^3 - 1}{6} = \int_{P_2} x^2 d\mu \\ \int_{\partial P_2} x^1 d\sigma &= b^2 = \int_{\partial P_2} x^2 d\sigma \\ c_1 &= -\frac{\int_{P_2} x^1 d\mu}{\int_{P_2} d\mu} = -\frac{b^2 + b + 1}{3(b + 1)} = c_2 \\ \int_{P_2} (x^1 + c_1) \det(DU) dx^1 dx^2 &= B^2 \frac{(b-1)^3}{6b^2} = \frac{(b^2 - ab)^2 (b-1)}{6b^2 (b+1)^2} \end{aligned} \tag{11}$$

Now if there is a solution to the Kähler-Yang-Mills equations then the  $\alpha$ -Futaki character vanishes. Hence using theorem 3.1, we get  $\frac{\alpha_1}{\alpha_0} = -\frac{(b^2-1)}{(b-a)^2}$ .

Hence there is no solution to the Kähler-Yang-Mills equations for  $\frac{\alpha_1}{\alpha_0} > 0$  and furthermore if there is a solution then  $\frac{\alpha_1}{\alpha_0} = -\frac{(b^2-1)}{(b-a)^2}$ .  $\square$

**Remark 2.** *If we use Kähler classes of the form  $a[H] - b[E]$  with  $a > b > 0$  as stated in 1, we will get similar results, that is,  $\frac{\alpha_1}{\alpha_0} < 0$ .*

Keller and Friedman constructed solution of Kähler-Yang-Mills equations on some line bundles over  $\mathbb{P}^2 \# \overline{\mathbb{P}^2}$ .

**Theorem 4.3** (Keller-Friedman). *There exists metrics  $h$  on the line bundle corresponding to the integral cohomology class  $(8k+3)[H_2] - [E_2]$  and  $\omega \in 3[H_2] - [E_2]$  such that they satisfy the Kähler-Yang-Mills equations with  $\frac{\alpha_1}{\alpha_0} = -\frac{1}{8k^2}$  (Here  $k \in \mathbb{N}$ ).*

*Proof.* In [18], Keller and Friedman proved that:

Suppose  $M = \mathbb{P}(\mathcal{O} \oplus \mathcal{L}) \rightarrow \Sigma$  is a ruled surface with  $\mathcal{L}$  of degree  $k \in \mathbb{Z}_+$ . Fix  $k' \in \mathbb{Z}_+$ ,  $k_1 \in \mathbb{Z}$ ,  $k_2 \in \mathbb{Z}^*$ . Consider the integral classes  $L = 2[E_0] + k'[C]$  and  $E = 2(k_1 - k_2)[E_0] + (k_1 k' + k_2(2k + k'))[C]$ . Then there exists a solution of Kähler-Yang-Mills equations with

$\frac{\alpha_1}{\alpha_0} = -\frac{(2-s_\Sigma)k+2k'}{8k_2^2(k+k')}$ , where  $E_0$  is the zero section,  $C$  denotes fiber,  $s_\Sigma = \frac{2(1-h)}{k}$ ,  $h = \text{genus}(\Sigma)$ .

Since  $\mathbb{P}^2 \# \overline{\mathbb{P}^2} = \mathbb{P}(\mathcal{O} \oplus \mathcal{O}(1)) \rightarrow \mathbb{P}^1$ , we take  $\Sigma = \mathbb{P}^1$  and  $k = 1$ . The relations between  $[E_2], [H_2]$ , and  $C, E_0$  are  $[C] = [H_2] - [E_2], [E_0] = [H_2]$ . Hence for  $L$  to be of the form  $a[H_2] - [E_2]$ , we must have  $k' = 1$ . Now  $E$  becomes  $2(k_1 - k_2)[E_0] + (k_1 + 3k_2)[C] = (3k_1 + k_2)[H_2] - (k_1 + 3k_2)[E_2]$ . Now putting  $k_2 = -k$  and  $k_1 = 1 + 3k$  with  $k \in \mathbb{N}$ , we get  $E = (8k + 3)[H_2] - [E_2]$ . Finally, we compute  $\frac{\alpha_1}{\alpha_0} = -\frac{1}{8k^2}$ . Observe that our theorem 4.2 also asserts that  $\frac{\alpha_1}{\alpha_0}$  has to be  $-\frac{3^2-1}{(3-8k-3)^2} = -\frac{1}{8k^2}$ .  $\square$

**Remark 3.** *The above analysis also indicates that why Keller-Friedman could not find solution of Kähler-Yang-Mills equations with  $\frac{\alpha_1}{\alpha_0} > 0$ .*

## 4.2 The $\alpha$ -Futaki character on $\mathbb{P}^3 \# \overline{\mathbb{P}^3}$

We will work with Kähler classes of the form  $p[H_3] - [E_3]$ , where  $H_3$  is the pullback of the hyperplane divisor from  $\mathbb{P}^3$  and  $E_3$  is the exceptional divisor,  $p > 1$ .

**Theorem 4.4.** *Suppose  $L$  is an ample line bundle on  $\mathbb{P}^3 \# \overline{\mathbb{P}^3}$  with  $c_1(L) = a[H_3] - [E_3]$  and  $[\omega] = b[H_3] - [E_3]$  satisfying  $3\frac{ab^2-1}{b^3-1} > 2$ . Then there is no solution to Kähler-Yang-Mills equations with  $\frac{\alpha_1}{\alpha_0} > 0$ . Furthermore, If a solution exists then  $\frac{\alpha_1}{\alpha_0} = -\frac{(3b+1)(b-1)^3(b^2+b+1)}{3b(b-a)^2(b^3+b^2-b+1)}$ .*

*Proof.* From 4.1, we know that there is a solution of the  $J$ -equation and a corresponding map between moment polytopes. So we have a map  $U : P_3 \rightarrow Q_3$  given by

$$U(x^1, x^2, x^3) = \left( Ax^1 + \frac{Bx^1}{(x^1 + x^2 + x^3)^3}, Ax^2 + \frac{Bx^2}{(x^1 + x^2 + x^3)^3}, Ax^3 + \frac{Bx^3}{(x^1 + x^2 + x^3)^3} \right),$$

where  $P_3 = \{(x^1, x^2, x^3) \in \mathbb{R}^3 | x^i \geq 0 \ i = 1, 2, 3; 1 \leq x^1 + x^2 + x^3 \leq b\}$ ,  $Q_3 = \{(x^1, x^2, x^3) \in \mathbb{R}^3 | x^i \geq 0 \ i = 1, 2, 3; 1 \leq x^1 + x^2 + x^3 \leq a\}$ ,  $A = \frac{ab^2-1}{b^3-1}$ , and  $B = \frac{b^3-ab^2}{b^3-1}$ .

A simple calculation yields

$$DU = \begin{bmatrix} A - \frac{3Bx^1}{X^4} + \frac{B}{X^3} & -\frac{3Bx^1}{X^4} & -\frac{3Bx^1}{X^4} \\ -\frac{3Bx^2}{X^4} & A - \frac{3Bx^2}{X^4} + \frac{B}{X^3} & -\frac{3Bx^2}{X^4} \\ -\frac{3Bx^3}{X^4} & -\frac{3Bx^3}{X^4} & A - \frac{3Bx^3}{X^4} + \frac{B}{X^3} \end{bmatrix},$$

where  $X = x^1 + x^2 + x^3$ . Suppose  $M_1, M_2, M_3$  are all the  $2 \times 2$  principal minor of  $DU$ . Then

$$M_1 + M_2 + M_3 = 3\left(A^2 - \frac{B^2}{X^6}\right).$$

A routine calculation yields

$$\begin{aligned}
\int_{P_3} d\mu &= \frac{b^3 - 1}{6} \\
\int_{\partial P_3} d\sigma &= 2b^2 - 1 \\
\int_{P_3} x^i d\mu &= \frac{b^4 - 1}{24}, \quad i = 1, 2, 3 \\
\int_{\partial P_3} x^i d\sigma &= \frac{3b^3 - 1}{6}, \quad i = 1, 2, 3 \\
\int_{P_3} (x^i + c_i) 3(A^2 - \frac{B^2}{X^6}) d\mu &= \frac{B^2(b^4 - 2b^3 + 2b - 1)}{8b^3}, \quad i = 1, 2, 3
\end{aligned} \tag{12}$$

where  $c_i = -\frac{(b^2+1)(b+1)}{4(b^2+b+1)}$ .

Now if there is a solution to the Kähler-Yang-Mills equations then the  $\alpha$ -Futaki character vanishes. Hence using theorem 3.1, we get  $\frac{\alpha_1}{\alpha_0} = -\frac{(3b+1)(b-1)^3(b^2+b+1)}{3b(b-a)^2(b^3+b^2-b+1)}$ . Hence there is no solution of the Kähler-Yang-Mills equations for  $\frac{\alpha_1}{\alpha_0} > 0$  and furthermore, if there is a solution then  $\frac{\alpha_1}{\alpha_0} = -\frac{(3b+1)(b-1)^3(b^2+b+1)}{3b(b-a)^2(b^3+b^2-b+1)}$ .  $\square$

Finally, we prove that the line bundles that we constructed in [17] over  $\mathbb{P}(\mathcal{O} \oplus \mathcal{O}(-1)) \rightarrow \mathbb{P}^1$  are not ample.

**Proposition 4.5.** *Suppose  $X = \mathbb{P}(\mathcal{O} \oplus \mathcal{O}(-1)) \rightarrow \mathbb{P}^1$  and  $L$  is the line bundle corresponding to the second integral cohomology class  $\delta = \frac{a\omega + b\gamma}{2\pi}$ , where  $\omega$  is a Kähler metric,  $\gamma$  is a tracefree (with respect to  $\omega$ )  $(1, 1)$  form,  $a = \frac{m_1 + m_2 \log 3}{2 + 3 \log 3}$ , and  $b = \frac{2m_2 - 3m_1}{2 + 3 \log 3}$  for  $(m_1, m_2) \neq (0, 0)$ . These line bundles are not ample.*

*Proof.* First we recall the expression for  $\omega$  and  $\gamma$  (for details see [17]). The expressions are as follows

$$\begin{aligned}
\omega &= (1 + \tau)\omega_{FS} + \phi(\tau) \frac{\sqrt{-1}dw \wedge d\bar{w}}{|w|^2}, \\
\gamma &= \omega_{FS} - \frac{\phi(\tau)}{1 + \tau} \frac{\sqrt{-1}dw \wedge d\bar{w}}{|w|^2},
\end{aligned}$$

where  $\phi(\tau) : [0, 2] \rightarrow \mathbb{R}$  is a function,  $\tau : X \rightarrow [0, 2]$  is a map,  $w$  fiber coordinate, and  $\omega_{FS}$  is the Fubini-Study metric on  $X$ .

Let's assume that the line bundles are ample. Then from Nakai-Moishezon criterion, we get

$$\int_{\mathbb{P}^1} \delta > 0, \quad \int_C \delta > 0, \quad \int_X \delta^2 > 0,$$

where  $C$  denotes fiber.

A routine calculations yields

$$a + b > 0, \quad 2a - b \log 3 > 0, \quad b^2 \log 3 > 4a^2.$$

Writing these inequalities in terms of  $m_1, m_2$ , we get

$$m_1 < (1 + \log \sqrt{3})m_2, \quad m_1 > 0, \quad (9 \log 3 - 4)m_1^2 > 20m_1m_2 \log 3.$$

Now

$$\begin{aligned} m_1 &< (1 + \log \sqrt{3})m_2 < \frac{9 \log 3 - 4}{20 \log 3}m_1 \\ 20 \log 3 &< 9 \log 3 - 4 \\ 4 + 11 \log 3 &< 0. \end{aligned} \tag{13}$$

A contradiction. □

## References

- [1] Miguel Abreu. Kähler geometry of toric varieties and extremal metrics. *Internat. J. Math.*, 9(6):641–651, 1998.
- [2] Luis Álvarez Cónsul, Mario García-Fernández, and Oscar García-Prada. Coupled equations for Kähler metrics and Yang-Mills connections. *Geom. Topol.*, 17(5):2731–2812, 2013.
- [3] Luis Álvarez-Cónsul, Mario Garcia-Fernandez, and Oscar García-Prada. Gravitating vortices, cosmic strings, and the Kähler-Yang-Mills equations. *Comm. Math. Phys.*, 351(1):361–385, 2017.
- [4] Luis Álvarez-Cónsul, Mario Garcia-Fernandez, Oscar García-Prada, and Vamsi Pritham Pingali. Gravitating vortices and the Einstein-Bogomol’nyi equations. *Math. Ann.*, 379(3-4):1651–1684, 2021.
- [5] Luis Álvarez-Cónsul, Mario Garcia-Fernandez, Oscar García-Prada, Vamsi Pritham Pingali, and Chengjian Yao. Obstructions to the existence of solutions of the self-dual Einstein-Maxwell-Higgs equations on a compact surface. *Bull. Sci. Math.*, 183:Paper No. 103233, 14, 2023.
- [6] Xiuxiong Chen. On the lower bound of the Mabuchi energy and its application. *Internat. Math. Res. Notices*, (12):607–623, 2000.
- [7] Tristan C. Collins and Gábor Székelyhidi. Convergence of the  $J$ -flow on toric manifolds. *J. Differential Geom.*, 107(1):47–81, 2017.
- [8] S. K. Donaldson. A new proof of a theorem of Narasimhan and Seshadri. *J. Differential Geom.*, 18(2):269–277, 1983.
- [9] S. K. Donaldson. Moment maps and diffeomorphisms. volume 3, pages 1–15. 1999. Sir Michael Atiyah: a great mathematician of the twentieth century.
- [10] S. K. Donaldson. Scalar curvature and stability of toric varieties. *J. Differential Geom.*, 62(2):289–349, 2002.

- [11] Hao Fang and Mijia Lai. Convergence of general inverse  $\sigma_k$ -flow on Kähler manifolds with Calabi ansatz. *Trans. Amer. Math. Soc.*, 365(12):6543–6567, 2013.
- [12] A. Futaki. An obstruction to the existence of Einstein Kähler metrics. *Invent. Math.*, 73(3):437–443, 1983.
- [13] Akito Futaki. *Kähler-Einstein metrics and integral invariants*, volume 1314 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1988.
- [14] Mario Garcia-Fernandez. Coupled equations for kähler metrics and yang-mills connections (thesis), 2011.
- [15] Mario Garcia-Fernandez, Vamsi Pritham Pingali, and Chengjian Yao. Gravitating vortices with positive curvature. *Adv. Math.*, 388:Paper No. 107851, 42, 2021.
- [16] Mario Garcia-Fernandez and Carl Tipler. Deformation of complex structures and the coupled Kähler-Yang-Mills equations. *J. Lond. Math. Soc. (2)*, 89(3):779–796, 2014.
- [17] Kartick Ghosh. Coupled Kähler-Einstein and Hermitian-Yang-Mills equations. *Bull. Sci. Math.*, 183:Paper No. 103232, 33, 2023.
- [18] Julien Keller and Christina W. Tønnesen Friedman. Nontrivial examples of coupled equations for Kähler metrics and Yang-Mills connections. *Cent. Eur. J. Math.*, 10(5):1673–1687, 2012.
- [19] Toshiki Mabuchi. Einstein-Kähler forms, Futaki invariants and convex geometry on toric Fano varieties. *Osaka J. Math.*, 24(4):705–737, 1987.
- [20] M. S. Narasimhan and C. S. Seshadri. Stable and unitary vector bundles on a compact Riemann surface. *Ann. of Math. (2)*, 82:540–567, 1965.
- [21] Gábor Székelyhidi. *An introduction to extremal Kähler metrics*, volume 152 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2014.
- [22] Gang Tian. Kähler-Einstein metrics on algebraic manifolds. In *Transcendental methods in algebraic geometry (Cetraro, 1994)*, volume 1646 of *Lecture Notes in Math.*, pages 143–185. Springer, Berlin, 1996.
- [23] Yi Yao. The J-flow on toric manifolds. *Acta Math. Sin. (Engl. Ser.)*, 31(10):1582–1592, 2015.
- [24] L. Álvarez Cónsul, M. Garcia-Fernandez, O. García-Prada, V. P. Pingali, and C. J. Yao. Gravitating vortices and symplectic reduction by stages, 2024.