

Stability and instability of constant stationary solutions to some integro-differential equations

Yuming Chen¹, Vitali Vougalter^{2 *}

¹ Department of Mathematics, Wilfrid Laurier University
Waterloo, Ontario, N2L 3C5, Canada
e-mail: ychen@wlu.ca

^{2 *} Department of Mathematics, University of Toronto
Toronto, Ontario, M5S 2E4, Canada
e-mail: vitali@math.toronto.edu

Abstract. We consider some reaction-diffusion equations describing systems with the non-local consumption of resources and the intraspecific competition. Sharp conditions on the coefficients are obtained to ensure the stability and instability of nontrivial constant stationary solutions.

Keywords: reaction-diffusion equation, stationary solution, stability, Fourier transform

AMS subject classification: 35K55, 35K57, 45K05

1 Introduction

In the first part of this article we consider the following integro-differential equation,

$$\frac{\partial u}{\partial t} = \Delta u + ku^2 \left(1 - au - b \int_{\mathbb{R}^n} \phi(x-y)u(y,t)dy \right), \quad (1.1)$$

where $(x, t) \in \mathbb{R}^n \times \mathbb{R}$, $n \in \mathbb{N}$, $u \in \mathcal{C} := C(\mathbb{R}^n, \mathbb{R})$. Here the integral kernel $\phi(x) \geq 0$ for $x \in \mathbb{R}^n$, which is dependent only upon the radial variable $|x|$ or it is a product of even functions (see e.g. expressions (1.9), (1.10), (1.11)) and is normalized as $\int_{\mathbb{R}^n} \phi(x)dx = 1$. The parameters $k, b > 0$ and $a \geq 0$. Obviously, the only nontrivial solution constant in space and time the equation (1.1) possesses is

$$u(x, t) = \frac{1}{a+b}. \quad (1.2)$$

Linearization of problem (1.1) around (1.2) with the ansatz $u(x, t) = \frac{1}{a+b} + \psi(x)e^{-\lambda t}$ yields the spectral problem on $L^2(\mathbb{R}^n)$

$$\mathcal{L}\psi = \lambda\psi, \quad (1.3)$$

where

$$\mathcal{L}\psi = -\Delta\psi + k_1\psi + k_2 \int_{\mathbb{R}^n} \phi(x-y)\psi(y)dy$$

with $k_1 = \frac{ka}{(a+b)^2}$ and $k_2 = \frac{kb}{(a+b)^2}$. The spectrum of the operator \mathcal{L} is denoted as $\sigma(\mathcal{L})$. Problem (1.3) is said to be *spectrally stable* if $\sigma(\mathcal{L}) \subseteq [0, \infty)$ and *spectrally unstable* if there exists $\lambda \in \sigma(\mathcal{L})$ such that $\lambda < 0$.

We denote the standard Fourier transform of a function $\psi(x) \in L^2(\mathbb{R}^n)$ as

$$\widehat{\psi}(p) = \frac{1}{(2\pi)^{\frac{n}{2}}} \int_{\mathbb{R}^n} \psi(x)e^{-ip \cdot x} dx, \quad p \in \mathbb{R}^n,$$

which enables us to write the quadratic form of the linearized operator as

$$(\mathcal{L}\psi, \psi) = \int_{\mathbb{R}^n} \Phi(p)|\widehat{\psi}(p)|^2 dp,$$

where

$$\Phi(p) = |p|^2 + k_1 + k_2(2\pi)^{\frac{n}{2}}\widehat{\phi}(p).$$

Clearly problem (1.3) is spectrally stable if for the Fourier image of the integral kernel involved in equation (1.1) we have $\widehat{\phi}(p) \geq 0$ for $p \in \mathbb{R}^n$ a.e.. Spectral instability arises when there is a region $\Omega \subset \mathbb{R}^n$ with nonempty interior such that $\Phi(p) < 0$ for $p \in \Omega$ a.e., which occurs due to the negativity of $\widehat{\phi}(p)$. Then we can choose a trial function $\psi(x) \in L^2(\mathbb{R}^n)$ such that its Fourier image $\widehat{\psi}(p)$ is compactly supported inside Ω . The negativity of the resulting quadratic form $(\mathcal{L}\psi, \psi)$ along with the min-max principle (see e.g. [8, p.306]) implies that the operator \mathcal{L} has negative spectrum.

We consider the following types of integral kernels whose Fourier transforms can be easily computed. The first one is

$$\phi_1(x) = \frac{\alpha}{2}e^{-\alpha|x|}, \quad x \in \mathbb{R}, \alpha > 0 \quad (1.4)$$

and hence $\widehat{\phi}_1(p) = \frac{\alpha^2}{\sqrt{2\pi}(p^2 + \alpha^2)}$, $p \in \mathbb{R}$; the second one is the two dimensional generalization of ϕ_1 ,

$$\phi_2(x, y) = \phi_1(x)\phi_1(y) = \frac{\alpha^2}{4}e^{-\alpha|x|}e^{-\alpha|y|}, \quad x, y \in \mathbb{R}, \alpha > 0 \quad (1.5)$$

with $\widehat{\phi}_2(p) = \frac{\alpha^4}{2\pi(p_1^2 + \alpha^2)(p_2^2 + \alpha^2)}$, $p = (p_1, p_2) \in \mathbb{R}^2$; the third one is the Gaussian kernel

$$\phi_3(x) = \left(\frac{\alpha}{\pi}\right)^{\frac{n}{2}} e^{-\alpha|x|^2}, \quad x \in \mathbb{R}^n, n \in \mathbb{N}, \alpha > 0. \quad (1.6)$$

One can get $\widehat{\phi}_3(p) = \frac{1}{(2\pi)^{\frac{n}{2}}} e^{-\frac{|p|^2}{4\alpha}}$, $p \in \mathbb{R}^n$. The fourth one is the three dimensional generalization of ϕ_1 ,

$$\phi_4(x) = \frac{\alpha^3}{8\pi}e^{-\alpha|x|}, \quad x \in \mathbb{R}^3, \alpha > 0. \quad (1.7)$$

Thus using the formula for the heat kernel of the square root of the Laplacian (see e.g. [7, p.169]) we easily obtain $\widehat{\phi}_4(p) = \frac{\alpha^4}{(2\pi)^{\frac{3}{2}}(\alpha^2+|p|^2)^2}$, $p \in \mathbb{R}^3$. Hence these choices of kernels (1.4)-(1.7) having positive Fourier images in the whole space yield spectral stability for problem (1.3). On the other hand if we choose the integral kernel

$$\phi_5(x) = \frac{1}{2N}\chi_{[-N, N]}(x), \quad x \in \mathbb{R}, \quad N > 0, \quad (1.8)$$

where χ_A stands for the characteristic function of a set A , we arrive at

$$\widehat{\phi}_5(p) = \frac{1}{\sqrt{2\pi}} \frac{\sin pN}{pN}, \quad p \in \mathbb{R},$$

which is sign indefinite. Now we form products out of the kernels above, namely,

$$\phi_5(x)\phi_1(y) = \frac{1}{2N}\chi_{[-N, N]}(x)\frac{\alpha}{2}e^{-\alpha|y|}, \quad x, y \in \mathbb{R}, \quad (1.9)$$

$$\phi_5(x)\phi_3(y) = \frac{1}{2N}\chi_{[-N, N]}(x)\sqrt{\frac{\alpha}{\pi}}e^{-\alpha y^2}, \quad x, y \in \mathbb{R}, \quad (1.10)$$

$$\phi_5(x)\phi_4(y) = \frac{1}{2N}\chi_{[-N, N]}(x)\frac{\alpha^3}{8\pi}e^{-\alpha|y|}, \quad x \in \mathbb{R}, y \in \mathbb{R}^3. \quad (1.11)$$

Since the linearized operator involved in (1.3) is similar to the one studied in [1] (see also [3, 5]), we arrive at the following statement analogous to Theorem 2.8 of [1] (see also Chapter 3 of [5]).

Theorem 1. *Let $k_1 = 0$ and the kernel ϕ be one of (1.8), (1.9), (1.10) and (1.11). Then problem (1.3) is spectrally stable if $\frac{1}{k_2} \geq -N^2 \frac{\sin z_1}{z_1^3}$ and is spectrally unstable if $\frac{1}{k_2} < -N^2 \frac{\sin z_1}{z_1^3}$, where z_1 is the unique solution of the equation $\tan z = \frac{1}{3}z$ on the interval $(\pi, \frac{3\pi}{2})$.*

Emergence and propagation of patterns in nonlocal reaction-diffusion equations arising in the theory of speciation was covered in [10]. Doubly nonlocal reaction-diffusion equations and the emergence of species were discussed in [2]. Asymptotic analysis of SIR epidemic model with nonlocal diffusion and general nonlinear incidence was performed in [4]. The article [6] deals with the diffusion-driven blow-up for a nonlocal Fisher-KPP type model. A discretization method for nonlocal diffusion type equations was developed in [9].

In the second part of this article we study the following integro-differential equation,

$$\frac{\partial u}{\partial t} = d\Delta u + u^2 \left(a - \int_{\mathbb{R}^n} \phi(x-y)u(y,t)dy \right) - bu, \quad (1.12)$$

where $(x, t) \in \mathbb{R}^n \times \mathbb{R}$, $n \in \mathbb{N}$, $u \in \mathcal{C}$, the integral kernel ϕ satisfies the same requirements as those for (1.1), the diffusion coefficient $d > 0$ and the parameters $a, b > 0$. We assume that $\frac{a^2}{4} \geq b$ such that equation (1.12) admits two nontrivial real-valued solutions constant in space and time given by

$$c_1 = \frac{a}{2} + \sqrt{\frac{a^2}{4} - b}, \quad c_2 = \frac{a}{2} - \sqrt{\frac{a^2}{4} - b} \quad (1.13)$$

and, in particular $c_{1,2} = \frac{a}{2}$ when $\frac{a^2}{4} = b$. Linearization of equation (1.12) around a solution in (1.13) with the ansatz $u(x, t) = c_{1,2} + \psi(x)e^{-\lambda t}$ yields the spectral problem on $L^2(\mathbb{R}^n)$

$$\mathcal{L}_k \psi = \lambda \psi, \quad k = 1, 2, \quad (1.14)$$

where

$$\mathcal{L}_k \psi = -d\Delta \psi + c_k^2 \int_{\mathbb{R}^n} \phi(x-y)\psi(y)dy - b\psi, \quad k = 1, 2,$$

such that problem (1.14) is spectrally stable if $\sigma(\mathcal{L}_k) \subseteq [0, \infty)$ and is spectrally unstable if there exists $\lambda \in \sigma(\mathcal{L}_k)$ such that $\lambda < 0$. Using the Fourier transform we write the quadratic form

$$(\mathcal{L}_k \psi, \psi) = \int_{\mathbb{R}^n} \Phi_k(p) |\widehat{\psi}(p)|^2 dp,$$

where

$$\Phi_k(p) = d|p|^2 + (2\pi)^{\frac{n}{2}} c_k^2 \widehat{\phi}(p) - b, \quad k = 1, 2.$$

By considering the integral kernels mentioned above we show that problem (1.14) is distinct from (1.3). In fact, (1.14) can be spectrally unstable even if the Fourier image of the integral kernel involved in (1.12) is positive in the whole space. More precisely, we have the following results.

Theorem 2. *Let the kernel ϕ be given by (1.4).*

- (i) *When $\frac{a^2}{4} > b$, problem (1.14) with $k = 1$ is spectrally stable if $d \geq \frac{(c_1 - \sqrt{c_1^2 - b})^2}{\alpha^2}$ and is spectrally unstable if $0 < d < \frac{(c_1 - \sqrt{c_1^2 - b})^2}{\alpha^2}$, and problem (1.14) with $k = 2$ is spectrally unstable.*
- (ii) *When $\frac{a^2}{4} = b$ problem (1.14) with $k = 1, 2$ is spectrally stable if $d \geq \frac{b}{\alpha^2}$ and is spectrally unstable if $0 < d < \frac{b}{\alpha^2}$.*

Theorem 3. *Let the kernel ϕ be given by (1.5).*

- (i) *When $\frac{a^2}{4} > b$, problem (1.14) with $k = 1$ is spectrally stable if $d \geq \frac{x^*}{\alpha^2}$ and is spectrally unstable if $0 < d < \frac{x^*}{\alpha^2}$, where x^* is defined in Lemma 7, and Problem (1.14) with $k = 2$ is spectrally unstable.*
- (ii) *When $\frac{a^2}{4} = b$, problem (1.14) with $k = 1, 2$ is spectrally stable if $d \geq \frac{a^2}{4\alpha^2}$ and is spectrally unstable if $0 < d < \frac{a^2}{4\alpha^2}$.*

Theorem 4. *Let the kernel ϕ be given by (1.6).*

- (i) *When $\frac{a^2}{4} > b$, problem (1.14) with $k = 1$ is spectrally stable if $d \geq s_0(a, b) \frac{c_1^2}{4\alpha}$ and is spectrally unstable if $0 < d < s_0(a, b) \frac{c_1^2}{4\alpha}$, where $s_0(a, b)$ is defined in Lemma 8, and problem (1.14) with $k = 2$ is spectrally unstable.*

- (ii) When $\frac{a^2}{4} = b$ problem (1.14) with $k = 1, 2$ is spectrally stable if $d \geq \frac{b}{4\alpha}$ and is spectrally unstable if $0 < d < \frac{b}{4\alpha}$.

Theorem 5. Let the kernel ϕ be given by (1.7).

- (i) When $\frac{a^2}{4} > b$, problem (1.14) with $k = 1$ is spectrally stable if $d \geq \frac{2x^*}{\alpha^2}$ and is spectrally unstable if $0 < d < \frac{2x^*}{\alpha^2}$, where x^* is defined in Lemma 7, and problem (1.14) with $k = 2$ is spectrally unstable.
- (ii) When $\frac{a^2}{4} = b$, problem (1.14) with $k = 1, 2$ is spectrally stable if $d \geq \frac{a^2}{2\alpha^2}$ and is spectrally unstable if $0 < d < \frac{a^2}{2\alpha^2}$.

These results are proved in the next section.

2 Proofs of Theorems 2–5

For technical purpose, we first establish the following a little bit trivial result, which proves the statements of Theorems 2–5 in the case where $\frac{a^2}{4} > b$ and $k = 2$.

Lemma 6. When $\frac{a^2}{4} > b$, we have $c_1^2 - b > 0$ and $c_2^2 - b < 0$ and hence problem (1.14) with the kernels given by (1.4)–(1.7) and $k = 2$ is spectrally unstable. Moreover, when $\frac{a^2}{4} = b$ both $c_1^2 - b$ and $c_2^2 - b$ are equal to zero.

Proof. By means of (1.13),

$$c_1^2 - b = \left(\frac{a}{2} - \sqrt{b} + \sqrt{\frac{a^2}{4} - b} \right) (c_1 + \sqrt{b}),$$

which is positive if $\frac{a^2}{4} > b$ and vanishes if $\frac{a^2}{4} = b$. Similarly

$$c_2^2 - b = \left(\frac{a}{2} - \sqrt{b} - \sqrt{\frac{a^2}{4} - b} \right) \left(\frac{a}{2} + \sqrt{b} - \sqrt{\frac{a^2}{4} - b} \right),$$

which clearly vanishes when $\frac{a^2}{4} = b$. For $\frac{a^2}{4} > b$, we expand the second term in the product above as

$$\sqrt{\frac{a}{2} + \sqrt{b}} \left(\sqrt{\frac{a}{2} + \sqrt{b}} - \sqrt{\frac{a}{2} - \sqrt{b}} \right) > 0$$

and the first one as

$$\sqrt{\frac{a}{2} - \sqrt{b}} \left(\sqrt{\frac{a}{2} - \sqrt{b}} - \sqrt{\frac{a}{2} + \sqrt{b}} \right) < 0.$$

Since $\int_{\mathbb{R}^n} \phi(x) dx = 1$, we have $\widehat{\phi}(0) = \frac{1}{(2\pi)^{\frac{n}{2}}}$ and it follows that $\Phi_k(0) = c_k^2 - b < 0$ when $k = 2$ and $\frac{a^2}{4} > b$. Hence by continuity for the kernels given by formulas (1.4)–(1.7), the function $\Phi_2(p)$ is negative in a neighborhood Ω of the origin with nonempty interior. Let us choose

the trial function $\widehat{\psi}(p) \in L^2(\mathbb{R}^n)$ to be compactly supported inside Ω . Then the quadratic form $(\mathcal{L}_2\psi, \psi) < 0$ which by means of the min-max principle implies the existence of negative spectrum and spectral instability of the problem (1.14) with $k = 2$ and the kernels given by (1.4)–(1.7). \blacksquare

We first prove Theorem 2.

Proof of Theorem 2. When the kernel is given by (1.4), we have

$$\Phi_k(p) = dp^2 + c_k^2 \frac{\alpha^2}{p^2 + \alpha^2} - b, \quad p \in \mathbb{R}, k = 1, 2.$$

Let us first consider the case of two distinct constant nontrivial stationary solutions of equation (1.12), i.e., $\frac{a^2}{4} > b$. When $k = 1$, clearly, it is sufficient to investigate the existence of intervals of negativity for the function

$$N_1(p) = dp^4 + (d\alpha^2 - b)p^2 + \alpha^2(c_1^2 - b), \quad p \in \mathbb{R}.$$

We distinguish the two cases of large and small diffusion coefficient. If $d \geq \frac{b}{\alpha^2}$, by Lemma 6, $N_1(p)$ is strictly positive on \mathbb{R} . This implies spectral stability for problem (1.14) with $k = 1$. Now suppose $0 < d < \frac{b}{\alpha^2}$. Then $N_1(p)$ is minimal at $p^2 = -\frac{d\alpha^2 - b}{2d}$ and a straightforward computation yields the minimal value

$$\frac{\alpha^2}{4d} \left[-\alpha^2 d^2 + 4d \left(c_1^2 - \frac{b}{2} \right) - \frac{b^2}{\alpha^2} \right].$$

The quadratic polynomial in d inside the brackets above has two roots

$$d_1 = \frac{(c_1 + \sqrt{c_1^2 - b})^2}{\alpha^2}, \quad d_2 = \frac{(c_1 - \sqrt{c_1^2 - b})^2}{\alpha^2}.$$

It can be easily verified that $0 < d_2 < \frac{b}{\alpha^2} < d_1$. Therefore, when $d \geq d_2$ we have $\Phi_1(p)$ nonnegative on \mathbb{R} and therefore, spectral stability for problem (1.14) with $k = 1$. When $0 < d < d_2$ there are intervals of negativity of $\Phi_1(p)$ on the real line, which by means of the min-max principle yields the existence of negative spectrum and therefore, spectral instability for problem (1.14) with $k = 1$.

Finally, we study the situation when both nontrivial constant solutions of (1.12) coincide, i.e., $\frac{a^2}{4} = b$. Then $c_1 = c_2 = \frac{a}{2}$. In this case

$$\Phi_k(p) = \frac{dp^2 + d\alpha^2 - b}{1 + \frac{\alpha^2}{p^2}}, \quad p \in \mathbb{R}, k = 1, 2.$$

Hence for $d \geq \frac{b}{\alpha^2}$, $\Phi_k(p)$ is nonnegative on \mathbb{R} which implies spectral stability for our problem and for $0 < d < \frac{b}{\alpha^2}$ it is negative in the neighborhood of the origin, which means spectral instability for (1.14). \blacksquare

Before moving on, we need the following technical lemma.

Lemma 7. Assume $\frac{a^2}{4} > b$. Consider the function $f(x) = 27c_1^2x^2 - (b + 2x)^3$, $x \in (0, b)$. f has the minimum only at $x_1 = \frac{9c_1^2 - 4b - 3c_1\sqrt{9c_1^2 - 8b}}{8}$ and is monotonically increasing on (x_1, b) . Moreover, in the interval $(0, b)$ the equation $f(x) = 0$ admits the unique solution $x^* \in (x_1, b)$.

Proof. A simple calculation yields

$$f'(x) = -6[4x^2 - (9c_1^2 - 4b)x + b^2],$$

which has two real zeros via Lemma 6

$$x_1 = \frac{9c_1^2 - 4b - 3c_1\sqrt{9c_1^2 - 8b}}{8}, \quad x_2 = \frac{9c_1^2 - 4b + 3c_1\sqrt{9c_1^2 - 8b}}{8}.$$

Since both the product and the sum of x_1 and x_2 are positive, we have $x_{1,2} > 0$. Apparently, it can be verified that $0 < x_1 < b < x_2$. Near the endpoints of the $(0, b)$ interval $f(0^+) = -b^3 < 0$ and $f(b^-) = 27b^2(c_1^2 - b) > 0$ via Lemma 6. Thus $f(x)$ considered on $(0, b)$ attains its minimal value, which is negative at x_1 , is strictly increasing for $x > x_1$, and has the unique zero at some point $x^* \in (x_1, b)$. ■

Next, we prove Theorem 3.

Proof of Theorem 3. When the kernel function is given by (1.5), we have

$$\Phi_k(p) = d|p|^2 + c_k^2 \frac{\alpha^4}{(p_1^2 + \alpha^2)(p_2^2 + \alpha^2)} - b, \quad p = (p_1, p_2) \in \mathbb{R}^2, k = 1, 2. \quad (2.1)$$

We start with the case of $\frac{a^2}{4} > b$ and therefore, problem (1.12) has two distinct constant stationary solutions c_1 and c_2 . When $k = 1$, in polar coordinates $p = (|p|, \theta)$, $|p| \in [0, \infty)$, $\theta \in [0, 2\pi)$, the function $\Phi_1(p)$ obviously equals

$$d|p|^2 + c_1^2 \frac{\alpha^4}{\frac{1}{4}|p|^4 \sin^2 2\theta + \alpha^2|p|^2 + \alpha^4} - b,$$

which is minimal for $\theta = \frac{\pi}{4}, \frac{3\pi}{4}, \frac{5\pi}{4}, \frac{7\pi}{4}$, namely when $p_1^2 = p_2^2$. In this case we introduce the new variable $q = p_1^2 \in [0, \infty)$ and arrive at the expression

$$\frac{(2dq - b)(q + \alpha^2)^2 + c_1^2\alpha^4}{(q + \alpha^2)^2}.$$

Since its denominator is strictly positive, it makes sense to consider

$$N_2(q) = (2dq - b)(q + \alpha^2)^2 + c_1^2\alpha^4,$$

which is positive at the origin by means of Lemma 6. We easily get

$$\frac{dN_2(q)}{dq} = 2(q + \alpha^2)(3dq + d\alpha^2 - b).$$

Thus when the diffusion coefficient is large enough, namely $d \geq \frac{b}{\alpha^2}$, the function $N_2(q)$ is monotonically increasing on the interval $(0, \infty)$. This implies its strict positivity and the spectral stability of (1.14) with $k = 1$. For small values of the diffusion coefficient, $0 < d < \frac{b}{\alpha^2}$ the function $N_2(q)$ attains its minimal value at

$$q_0 = \frac{b - d\alpha^2}{3d}.$$

A straightforward computation yields

$$N_2(q_0) = \frac{27c_1^2\alpha^4d^2 - (b + 2d\alpha^2)^3}{27d^2}.$$

Let us introduce the new variable $x = d\alpha^2$, $0 < x < b$. This will turn the numerator of the expression in the right side of $N_2(q_0)$ into the function $f(x)$ studied in Lemma 7. Therefore, problem (1.14) with $\frac{a^2}{4} > b$ and $k = 1$ is spectrally stable for $d \geq \frac{x^*}{\alpha^2}$ and is spectrally unstable when $0 < d < \frac{x^*}{\alpha^2}$.

Now consider the degenerate case that $\frac{a^2}{4} = b$. In this case,

$$\Phi_k(p) = d|p|^2 + \frac{a^2}{4} \frac{\alpha^4}{(p_1^2 + \alpha^2)(p_2^2 + \alpha^2)} - \frac{a^2}{4}, \quad p = (p_1, p_2) \in \mathbb{R}^2, k = 1, 2. \quad (2.2)$$

By changing the variables to the polar coordinates we easily obtain

$$\Phi_k(p) = d|p|^2 + \frac{a^2\alpha^4}{|p|^4 \sin^2 2\theta + 4\alpha^2|p|^2 + 4\alpha^4} - \frac{a^2}{4},$$

which is minimal when $p_1^2 = p_2^2$. Using the variable $q = p_1^2 \in [0, \infty)$ we arrive at

$$\frac{(8dq - a^2)(q + \alpha^2)^2 + a^2\alpha^4}{4(q + \alpha^2)^2}.$$

Due to the positivity of the denominator in the expression above, one needs to study

$$N_3(q) = (8dq - a^2)(q + \alpha^2)^2 + a^2\alpha^4, \quad q \in [0, \infty).$$

Obviously, $N_3(0) = 0$. A straightforward computation yields

$$\frac{dN_3(q)}{dq} = 2(q + \alpha^2)(12dq + 4d\alpha^2 - a^2).$$

Therefore, when the diffusion coefficient $d \geq \frac{a^2}{4\alpha^2}$, we have $N_3(q) \geq N_3(0) = 0$, $q \in [0, \infty)$ and hence problem (1.14) with $\frac{a^2}{4} = b$ and $k = 1, 2$ is spectrally stable. On the other hand, for $0 < d < \frac{a^2}{4\alpha^2}$, the function $N_3(q)$ considered on the nonnegative semi-axis has the minimal value at

$$q_{0,0} = \frac{a^2 - 4d\alpha^2}{12d} > 0.$$

Thus $N_3(q)$ is monotonically decreasing on the interval $(0, q_{0,0})$ from its zero value at the origin and therefore has the intervals of negativity, which yields the spectral instability for problem (1.14) with $\frac{a^2}{4} = b$ for the small values of the diffusion coefficient $0 < d < \frac{a^2}{4\alpha^2}$. ■

In order to study spectral stability for stationary solutions of our integro-differential equation in the case of the Gaussian kernel, we will need the following elementary result.

Lemma 8. *Assume $\frac{a^2}{4} > b$. Then the function $g(s) = -s \ln s + s - \frac{b}{c_1^2}$, $s \in (0, 1)$ is strictly increasing and it has a unique zero $s_0 = s_0(a, b)$.*

Proof. Near the endpoints of the interval $(0, 1)$, $g(0^+) = -\frac{b}{c_1^2} < 0$ and $g(1^-) = 1 - \frac{b}{c_1^2} > 0$ via Lemma 6. On this interval the function is continuously differentiable such that $g'(s) = -\ln s > 0$, $s \in (0, 1)$. Therefore, it is strictly increasing and has a unique zero at some point $s_0 = s_0(a, b) \in (0, 1)$. ■

Now we come to prove Theorem 4.

Proof of Theorem 4. When the kernel is given by (1.6), we obtain

$$\Phi_k(p) = \Phi_k(|p|) = d|p|^2 + c_k^2 e^{-\frac{|p|^2}{4\alpha}} - b, \quad p \in \mathbb{R}^n, k = 1, 2.$$

First we turn our attention to the situation where $\frac{a^2}{4} > b$. In the case of $k = 1$, $\Phi_k(0) = c_1^2 - b > 0$ via Lemma 6 and we compute

$$\frac{\partial \Phi_1(|p|)}{\partial |p|} = 2|p| \left(d - \frac{c_1^2}{4\alpha} e^{-\frac{|p|^2}{4\alpha}} \right), \quad |p| \in [0, \infty). \quad (2.3)$$

Thus when $d \geq \frac{c_1^2}{4\alpha}$, the derivative above is nonnegative, $\Phi_1(p) > 0$, $p \in \mathbb{R}^n$ and hence problem (1.14) with $k = 1$ is spectrally stable. On the other hand, when $0 < d < \frac{c_1^2}{4\alpha}$, formula (2.3) yields the single critical point outside the origin for $\Phi_1(|p|)$,

$$|p_0| = 2\sqrt{\alpha} \sqrt{\ln \frac{c_1^2}{4\alpha d}}. \quad (2.4)$$

We compute the second derivative $\frac{\partial^2 \Phi_1}{\partial |p|^2}(|p_0|) = \frac{c_1^2 |p_0|^2}{4\alpha^2} e^{-\frac{|p_0|^2}{4\alpha}} > 0$ such that $|p_0|$ given by (2.4) is the minimal point of the function $\Phi_1(|p|)$ on the semi-axis $(0, \infty)$. A straightforward computation yields $\Phi_1(|p_0|) = c_1^2 g(s)$ with $g(s)$ defined in Lemma 8 and $s = \frac{4\alpha d}{c_1^2}$ such that $s \in (0, 1)$. Thus by means of Lemma 8 when $d \geq s_0(a, b) \frac{c_1^2}{4\alpha}$ and therefore, when $s \geq s_0(a, b)$, we have $\Phi_1(|p_0|) > 0$ and problem (1.14) with $\frac{a^2}{4} > b$ and $k = 1$ is spectrally stable. But when $0 < d < s_0(a, b) \frac{c_1^2}{4\alpha}$ such that $0 < s < s_0(a, b)$ problem (1.14) with $k = 1$ is spectrally unstable due to the existence of the regions of negativity of $\Phi_1(p)$, which completes the proof of Theorem 4 in the case of $\frac{a^2}{4} > b$.

Now, we study the degenerate case of $\frac{a^2}{4} = b$. Then

$$\Phi_k(p) = \Phi_k(|p|) = d|p|^2 + c_1^2 e^{-\frac{|p|^2}{4\alpha}} - b, \quad p \in \mathbb{R}^n, k = 1, 2,$$

and $c_1 = c_2 = \frac{a}{2} = \sqrt{b}$. Note that $\Phi_1(0) = c_1^2 - b = 0$ via Lemma 6 and the derivative with respect to the radial variable is given by (2.3). Hence when $d \geq \frac{b}{4\alpha}$ the derivative (2.3) is positive outside the origin such that the function $\Phi_1(p) > 0$ for $|p| \in (0, \infty)$, which implies the spectral stability. On the other hand, when $0 < d < \frac{b}{4\alpha}$ the derivative given by (2.3) is negative for small values of $|p| \in (0, \infty)$. Therefore, problem (1.14) with $\frac{a^2}{4} = b$ and $k = 1, 2$ is spectrally unstable in this case due to the existence of the regions of negativity of $\Phi_1(p)$. ■

Finally, we prove Theorem 5 in an analogous way as for Theorem 3.

Proof of Theorem 5. In the case of the kernel function given by (1.7),

$$\Phi_k(p) = d|p|^2 + c_k^2 \frac{\alpha^4}{(\alpha^2 + |p|^2)^2} - b, \quad p \in \mathbb{R}^3, \quad k = 1, 2.$$

By comparing the expression above with formula (2.1) in the case of $p_1^2 = p_2^2$ we draw the conclusion that when $\frac{a^2}{4} > b$ and $d \geq \frac{2x^*}{\alpha^2}$ with x^* defined in Lemma 7, the function $\Phi_1(p)$ is nonnegative in the space of three dimensions and problem (1.14) with $k = 1$ is spectrally stable. Moreover, $0 < d < \frac{2x^*}{\alpha^2}$ yields regions of negativity for $\Phi_1(p)$ and therefore, spectral instability. In the degenerate case of $\frac{a^2}{4} = b$ the expression for $\Phi_k(p)$, $k = 1, 2$ is given by

$$d|p|^2 + \frac{a^2}{4} \frac{\alpha^4}{(\alpha^2 + |p|^2)^2} - \frac{a^2}{4}, \quad p \in \mathbb{R}^3.$$

Let us compare it with formula (2.2) when $p_1^2 = p_2^2$. It follows that when $d \geq \frac{a^2}{2\alpha^2}$ there is spectral stability and when $0 < d < \frac{a^2}{2\alpha^2}$ the problem (1.14) with $k = 1, 2$ and $\frac{a^2}{4} = b$ is spectrally unstable. ■

Acknowledgement

Y.C. was partially supported by the NSERC of Canada (Nos. RGPIN-2019-05892, RGPIN-2024-05593).

References

- [1] N. Apreutesei, N. Bessonov, V. Volpert, V. Vougalter, *Spatial structures and generalized travelling waves for an integro-differential equation*, Discrete Contin. Dyn. Syst. Ser. B, **13** (2010), no. 3, 537–557.
- [2] M. Banerjee, V. Vougalter, V. Volpert, *Doubly nonlocal reaction-diffusion equations and the emergence of species*, Appl. Math. Model., **42** (2017), 591–599.
- [3] H. Berestycki, G. Nadin, B. Perthame, L. Ryzhik, *The non-local Fisher-KPP equation: travelling waves and steady states*, Nonlinearity, **22** (2009), no. 12, 2813–2844.

- [4] S. Djilali, Y. Chen, S. Bentout, *Asymptotic analysis of SIR epidemic model with nonlocal diffusion and generalized nonlinear incidence functional*, Math. Methods Appl. Sci., **46** (2023), no. 5, 6279–6301.
- [5] S. Genieys, V. Volpert, P. Auger, *Pattern and waves for a model in population dynamics with nonlocal consumption of resources*, Math. Model. Nat. Phenom., **1** (2006), no. 1, 65–82.
- [6] N.I. Kavallaris, E. Latos, T. Suzuki, *Diffusion-driven blow-up for a nonlocal Fisher-KPP type model*, SIAM J. Math. Anal., **55** (2023), no. 3, 2411–2433.
- [7] E.H. Lieb, M. Loss, *Analysis*, Grad. Stud. Math., **14**, American Mathematical Society, Providence, RI, (1997), 278 pp.
- [8] E.H. Lieb, B. Simon, A.S. Wightman, *Studies in mathematical physics. Essays in honor of Valentine Bargmann*. Princeton Series in Physics. Princeton University Press (1976), 460 pp.
- [9] D. Mezzanotte, D. Occorsio, M.G. Russo, E. Venturino, *A discretization method for nonlocal diffusion type equations*, Ann. Univ. Ferrara, Sez. VII, Sci. Mat., **68** (2022), no. 2, 505–520.
- [10] V. Volpert, V. Vougalter, *Emergence and propagation of patterns in nonlocal reaction-diffusion equations arising in the theory of speciation*, Lecture Notes in Math., **2071** Springer, Heidelberg, (2013), 331–353.