

# ON THE SIZE OF SETS AVOIDING A GENERAL STRUCTURE

RUNZE WANG

ABSTRACT. Given a finite abelian group  $G$  and a subset  $S \subseteq G$ , we let  $N_{G, S}$  be the smallest integer  $N$  such that for any subset  $A \subseteq G$  with  $N$  elements, we have  $g + S \subseteq A$  for some  $g \in G$ . Using the probabilistic method, we prove that

$$\frac{|H_G(S)| - 1}{|H_G(S)|} |G| + \left\lceil \left( \frac{|G|}{|H_G(S)|} \right)^{1 - |H_G(S)|/|S|} \right\rceil \leq N_{G, S} \leq \left\lfloor \frac{|S| - 1}{|S|} |G| \right\rfloor + 1,$$

where  $H_G(S)$  is the stabilizer of  $S$ .

Problems about avoiding structures, especially avoiding arithmetic progressions, are well-known and have been extensively studied. For example, the famous Roth's theorem, which is about avoiding three-term arithmetic progressions, was proved in [3] and has been refined in [1, 2, 4, 5]. In this succinct paper, we take the avoided structure to be a general set.

For a finite abelian group  $(G, +)$ , an element  $g \in G$ , and a subset  $S \subseteq G$ , we define  $g + S$  to be  $\{g + s : s \in S\}$ , and define the *stabilizer* of  $S$  to be

$$H_G(S) = \{g' \in G : g' + S = S\}.$$

It is easy to check that  $H_G(S)$  is a subgroup of  $G$ , and  $S$  is the union of some cosets of  $H_G(S)$ .

Given a finite abelian group  $G$  and a subset  $S \subseteq G$ , we let  $N_{G, S}$  denote the smallest integer  $N \geq |S|$  such that for any subset  $A \subseteq G$  with  $N$  elements, we have  $g + S \subseteq A$  for some  $g \in G$ . Thus, for any  $M \leq N_{G, S} - 1$ , there exists a subset  $B \subseteq G$  with  $M$  elements, such that  $g + S \not\subseteq B$  for any  $g \in G$ . Roughly speaking, this means the additive structure of  $S$  is avoided in  $B$ .

Firstly we prove the following bounds on  $N_{G, S}$ , and the lower bound will be improved later.

**Theorem 1.** *We have*

$$\frac{|H_G(S)| - 1}{|H_G(S)|} |G| + 1 \leq N_{G, S} \leq \left\lfloor \frac{|S| - 1}{|S|} |G| \right\rfloor + 1.$$

*Proof.* For the lower bound, we can construct a subset  $B \subseteq G$  with  $\frac{|H_G(S)| - 1}{|H_G(S)|} |G|$  elements by excluding one element from each coset of  $H_G(S)$ , then we will have  $g + S \not\subseteq B$  for any  $g \in G$ .

For the upper bound, let us assume for some subset  $A \subseteq G$  with  $\left\lfloor \frac{|S| - 1}{|S|} |G| \right\rfloor + 1$  elements, we have  $g + S \not\subseteq A$  for any  $g \in G$ , which means  $(g + S) \cap (G \setminus A) \neq \emptyset$  for any  $g \in G$ . For each  $\alpha \in G \setminus A$ , we have

$$|\{g \in G : \alpha \in g + S\}| = |\{g \in G : g \in \alpha - S\}| = |\alpha - S| = |S|,$$

which implies

$$\begin{aligned}
|\{g \in G : (g + S) \cap (G \setminus A) \neq \emptyset\}| &\leq \sum_{\alpha \in G \setminus A} |\{g \in G : \alpha \in g + S\}| \\
&= |G \setminus A| |S| \\
&= \left( |G| - \left( \left\lfloor \frac{|S|-1}{|S|} |G| \right\rfloor + 1 \right) \right) |S| \\
&< |G|,
\end{aligned}$$

contradicting the assumption that  $(g + S) \cap (G \setminus A) \neq \emptyset$  for any  $g \in G$ . So for any subset  $A \subseteq G$  with  $\left\lfloor \frac{|S|-1}{|S|} |G| \right\rfloor + 1$  elements, we can find  $g + S$  in  $A$  for some  $g \in G$ , and thus  $N_{G, S} \leq \left\lfloor \frac{|S|-1}{|S|} |G| \right\rfloor + 1$ .  $\square$

We have a direct corollary.

**Corollary 2.** *If  $S$  is a coset of some subgroup of  $G$ , then*

$$N_{G, S} = \frac{|S|-1}{|S|} |G| + 1.$$

*Proof.* If  $S$  is a coset of a subgroup, then  $|H_G(S)| = |S|$ , and the equalities in Theorem 1 will be attained.  $\square$

Let  $T_G(S)$  be a transversal of  $G/H_G(S)$ , which means  $T_G(S)$  contains exactly one element from each coset of  $H_G(S)$ , so  $|T_G(S)| = \frac{|G|}{|H_G(S)|}$ . For a subset  $A \subseteq G$ , it is easy to see that the following two statements are equivalent.

- There exists  $g \in G$ , such that  $g + S \subseteq A$ .
- There exists  $g' \in T_G(S)$ , such that  $g' + S \subseteq A$ .

Using the probabilistic method, we prove another lower bound on  $N_{G, S}$ . We will use this result as a lemma to prove a better lower bound in Theorem 4, which is our final goal.

**Lemma 3.** *We have*

$$N_{G, S} \geq |T_G(S)|^{-1/|S|} |G| = |H_G(S)|^{1/|S|} |G|^{1-1/|S|}$$

*Proof.* Suppose  $N \geq |S|$  is an integer such that for any subset  $A \subseteq G$  with  $N$  elements, there exists  $g \in G$  such that  $g + S \subseteq A$ . We randomly choose a set  $X$  from all  $N$ -element subsets of

$G$ , then

$$\begin{aligned}
\mathbb{P}(\exists g \in G \text{ s.t. } g + S \subseteq X) &= \mathbb{P}(\exists g' \in T_G(S) \text{ s.t. } g' + S \subseteq X) \\
&\leq \sum_{g' \in T_G(S)} \mathbb{P}(g' + S \subseteq X) \\
&= |T_G(S)| \frac{\binom{|G|-|S|}{N-|S|}}{\binom{|G|}{N}} \\
&\leq |T_G(S)| \left( \frac{N}{|G|} \right)^{|S|}.
\end{aligned}$$

If  $N < |T_G(S)|^{-1/|S|}|G|$ , then  $\mathbb{P}(\exists g \in G \text{ s.t. } g + S \subseteq X) < 1$ , which means there is some  $N$ -element set  $A \subseteq G$  such that  $g + S \not\subseteq A$  for any  $g \in G$ , contradiction. So  $N \geq |T_G(S)|^{-1/|S|}|G|$ , and thus  $N_{G, s} \geq |T_G(S)|^{-1/|S|}|G| = |H_G(S)|^{1/|S|}|G|^{1-1/|S|}$ .  $\square$

Combining the ideas in Theorem 1 and Lemma 3, we prove the following result. In the proof, we take  $G' := G/H_G(S)$  and  $S' := S/H_G(S)$ , then  $H_{G'}(S')$  will be trivial, and by Lemma 3, we have  $N_{G', s'} \geq |G'|^{1-1/|S'|}$ . And because  $N_{G', s'}$  is an integer, we know  $N_{G', s'} \geq \lceil |G'|^{1-1/|S'|} \rceil$ .

**Theorem 4.** *We have*

$$N_{G, s} \geq \frac{|H_G(S)| - 1}{|H_G(S)|} |G| + \left\lceil \left( \frac{|G|}{|H_G(S)|} \right)^{1-|H_G(S)|/|S|} \right\rceil. \quad (0.1)$$

*Proof.* We shall construct a subset  $B \subseteq G$  with  $\frac{|H_G(S)| - 1}{|H_G(S)|} |G| + \left\lceil \left( \frac{|G|}{|H_G(S)|} \right)^{1-|H_G(S)|/|S|} \right\rceil - 1$  elements, and show that  $g + S \not\subseteq B$  for any  $g \in G$ .

By Lemma 3, we know that there is a subset  $B' \subseteq G'$  with  $\lceil |G'|^{1-1/|S'|} \rceil - 1$  elements, such that  $g' + S' \not\subseteq B'$  for any  $g' \in G'$ . We let

$$B_1 = \{b \in G : b + H_G(S) \in B'\},$$

so

$$|B_1| = (\lceil |G'|^{1-1/|S'|} \rceil - 1) |H_G(S)|.$$

Then, there are  $|G'| - |B'|$  cosets of  $H_G(S)$  which are not in  $B'$ , we denote these cosets by  $H_1, H_2, \dots, H_{|G'|-|B'|}$ . In each  $H_i$ , we randomly pick an element  $h_i$ , and let  $K_i$  be  $H_i \setminus \{h_i\}$ . We let  $B_2$  be the union of  $K_i$ , so

$$B_2 = \bigcup_{i=1}^{|G'|-|B'|} K_i,$$

and

$$|B_2| = (|G'| - |B'|)(|H_G(S)| - 1) = (|G'| - (\lceil |G'|^{1-1/|S'|} \rceil - 1))( |H_G(S)| - 1).$$

Now, we take  $B$  to be  $B_1 \cup B_2$ , then

$$\begin{aligned} |B| &= |G'|(|H_G(S)| - 1) + \lceil |G'|^{1-1/|S'|} \rceil - 1 \\ &= \frac{|H_G(S)| - 1}{|H_G(S)|} |G| + \left\lceil \left( \frac{|G|}{|H_G(S)|} \right)^{1-|H_G(S)|/|S|} \right\rceil - 1. \end{aligned}$$

And we need to show that  $g + S \not\subseteq B$  for any  $g \in G$ .

- If for some  $g \in G$ , we have  $g + S \subseteq B_1$ , then  $g' := g + H_G(S) \in G'$  and  $g' + S' \subseteq B'$ , contradicting the definition of  $B'$ .
- If for some  $g \in G$ , we have  $g + S \subseteq B$  and  $(g + S) \cap B_2 \neq \emptyset$ , then again we have a contradiction, because  $g + S$  is a union of  $H_G(S)$  cosets, but  $B_2$  is a union of  $H_G(S)$  cosets with punched holes.

So  $g + S \not\subseteq B$  for any  $g \in G$ . □

We need to check the lower bound obtained in Theorem 4 is better than the one in Lemma 3. Although this is intuitive, we have a formal verification given by the following proposition, where  $g$ ,  $h$ , and  $s$  play the roles of  $|G|$ ,  $|H_G(S)|$ , and  $|S|$  respectively.

**Proposition 5.** *Let  $g$ ,  $h$ ,  $s \geq 1$  be three real numbers with  $g \geq h$ , then*

$$\frac{h-1}{h}g + \left(\frac{g}{h}\right)^{1-h/s} \geq h^{1/s}g^{1-1/s}.$$

*Proof.* We can fix  $g$  and  $h$ , and take  $s$  as a variable. Note that actually we should have  $h \leq s \leq g$ , but for calculation convenience, let us take  $1 \leq s < \infty$ . Let  $f(s) = \frac{h-1}{h}g + \left(\frac{g}{h}\right)^{1-h/s} - h^{1/s}g^{1-1/s}$ . It turns out  $f'(s) \leq 0$ , so  $f(s)$  is decreasing on  $[1, \infty)$ . And if  $s$  is taken to be  $\infty$ , then  $f(\infty) = 0$ . So we always have  $f(s) \geq 0$ , and thus  $\frac{h-1}{h}g + \left(\frac{g}{h}\right)^{1-h/s} \geq h^{1/s}g^{1-1/s}$ . □

Note that if  $S$  is a coset of some subgroup of  $G$ , then the lower bound in Theorem 4 is the same as the one in Theorem 1; if  $|H_G(S)| = 1$  or  $H_G(S) = S = G$ , then the lower bound in Theorem 4 is the same as the one in Lemma 3.

We finish this short paper with an example.

Let us take  $G$  to be  $C_{2024}$ , the cyclic group of order  $2024 = 2^3 \cdot 11 \cdot 23$ , and take  $S$  to be the union of  $n$  cosets of the subgroup of order eight. So  $|S| = 8n$ , and if we restrict  $n$  to be in  $[1, 10]$ ,

then  $|H_G(S)|$  must be eight. Then by Theorem 1, Corollary 2, and Theorem 4, we have

$$N_{G, S} \left\{ \begin{array}{ll} = 1772 & \text{if } n = 1, \\ \in [1787, 1898] & \text{if } n = 2, \\ \in [1812, 1940] & \text{if } n = 3, \\ \in [1835, 1961] & \text{if } n = 4, \\ \in [1855, 1974] & \text{if } n = 5, \\ \in [1872, 1982] & \text{if } n = 6, \\ \in [1886, 1988] & \text{if } n = 7, \\ \in [1898, 1993] & \text{if } n = 8, \\ \in [1908, 1996] & \text{if } n = 9, \\ \in [1917, 1999] & \text{if } n = 10. \end{array} \right.$$

## REFERENCES

- [1] J. Bourgain. Roth's theorem on progressions revisited. *J. Anal. Math.*, 104:155–192, 2008.
- [2] D. R. Heath-Brown. Integer sets containing no arithmetic progressions. *J. London Math. Soc. (2)*, 35(3):385–394, 1987.
- [3] K. F. Roth. On certain sets of integers. *J. London Math. Soc.*, 28:104–109, 1953.
- [4] T. Sanders. On Roth's theorem on progressions. *Ann. of Math. (2)*, 174(1):619–636, 2011.
- [5] E. Szemerédi. Integer sets containing no arithmetic progressions. *Acta Math. Hungar.*, 56(1-2):155–158, 1990.

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF MEMPHIS, MEMPHIS, TN 38152, USA  
*Email address:* runze.w@hotmail.com