

MONOTONICITY FORMULAS FOR CAPILLARY SURFACES

GUOFANG WANG, CHAO XIA, AND XUWEN ZHANG

ABSTRACT. In this paper, we establish monotonicity formulas for capillary surfaces in the half-space \mathbb{R}_+^3 and in the unit ball \mathbb{B}^3 and extend the result of Volkman [27] for surfaces with free boundary. As applications, we obtain Li-Yau-type inequalities for the Willmore energy of capillary surfaces, and extend Fraser-Schoen's optimal area estimate for minimal free boundary surfaces in \mathbb{B}^3 [10] to the capillary setting, which is different to another optimal area estimate proved by Brendle in [5].

Keywords: monotonicity formula, Li-Yau inequality, Willmore energy, optimal area estimate, minimal surface, capillary surface

MSC 2020: 53C42, 53A10, 49Q15

CONTENTS

1. Introduction	1
1.1. Main Result	2
1.2. Organization of the Paper	5
2. Monotonicity Formula in the Half-Space	5
2.1. Set-ups	5
2.2. A Monotonicity Formula	6
3. Willmore Energy and Li-Yau-Type Inequality for Capillary Immersions	12
3.1. Capillary Immersions	12
3.2. Willmore Energy in the Half-Space	14
3.3. Willmore Energy in the Unit Ball	17
4. Monotonicity Identities in the Unit Ball	18
4.1. Set-ups	18
4.2. Simon-Type Monotonicity Formulas	20
4.3. Applications of Eqn. 4.15	28
References	29

1. INTRODUCTION

For a 2-dimensional immersed, open surface $\Sigma \subset \mathbb{R}^{n+1}$, it is proved by Simon [25] that for $0 < \sigma < \rho < \infty$, $a \in \mathbb{R}^{n+1}$,

$$g_a(\rho) - g_a(\sigma) = \frac{1}{\pi} \int_{\Sigma \cap B_\rho(a) \setminus B_\sigma(a)} \left| \frac{1}{4} \vec{\mathbf{H}} + \frac{(x-a)^\perp}{|x-a|^2} \right|^2 d\mathcal{H}^2, \quad (1.1)$$

where

$$g_a(r) := \frac{\mathcal{H}^2(\Sigma \cap B_r(a))}{\pi r^2} + \frac{1}{16\pi} \int_{\Sigma \cap B_r(a)} |\vec{\mathbf{H}}|^2 d\mathcal{H}^2 + \frac{1}{2\pi r^2} \int_{\Sigma \cap B_r(a)} \vec{\mathbf{H}} \cdot (x - a) d\mathcal{H}^2.$$

This is known as Simon's monotonicity identity and is later generalized to hold for integral 2-varifolds by Kuwert-Schätzle [17]. As an interesting application, the monotonicity formula yields an alternative proof (by taking $\rho \rightarrow \infty$ and $\sigma \rightarrow 0^+$) of the Li-Yau inequality ([20]):

$$\pi \Theta_{max} \leq \frac{1}{16} \int_{\mathbb{R}^{n+1}} |\vec{\mathbf{H}}|^2 d\mu,$$

where Θ_{max} denotes the maximal density of an integral 2-varifold μ , whose generalized mean curvature in \mathbb{R}^{n+1} is square integrable. Note that for an immersion of a 2-dimensional compact orientable closed smooth surface $F : \Sigma \rightarrow \mathbb{R}^{n+1}$, let μ_g be the induced area measure of Σ with respect to the pull-back metric $g = F^* g_{\text{euc}}$, then its image as integral 2-varifold is given by

$$\mu := F(\mu_g) = \left(x \mapsto \mathcal{H}^0(F^{-1}(x)) \right) \mathcal{H}^2 \llcorner F(\Sigma).$$

Hence from the Li-Yau inequality, we easily see that the Willmore energy

$$\mathcal{W}(\mu) := \frac{1}{4} \int_{\mathbb{R}^{n+1}} |\vec{\mathbf{H}}|^2 d\mu \geq 4\pi,$$

and $F : \Sigma \rightarrow \mathbb{R}^{n+1}$ is an embedding if

$$\mathcal{W}(\mu) < 8\pi.$$

We refer the interested readers to the monographs [18, 22] for an overview of the study of Willmore energy.

Recently, Volkman [27] generalized this theory to free boundary surfaces in the unit ball $\mathbb{B}^{n+1} = \{x \in \mathbb{R}^{n+1} : |x| < 1\}$ (indeed, he proved it in the context of integral free boundary 2-varifolds). He obtained a monotonicity identity similar to (1.1) (see Section 4) and consequently a Li-Yau-type inequality. As an application, he established the Willmore energy for integral free boundary 2-varifolds in \mathbb{B}^{n+1} , and proved that its lower bound is given exactly by 2π .

1.1. Main Result. In this paper, we deal with the capillary counterpart of the above results. We first prove a Simon-type monotonicity formula for capillary hypersurfaces in the half-space

$$\mathbb{R}_+^3 := \{x \in \mathbb{R}^3 : x_3 < 1\}$$

that are formulated in the weak sense, see Section 2.1 for the precise definition.

Theorem 1.1 (Simon-type monotonicity formula). *Given $\theta \in (0, \pi)$, let V be a rectifiable 2-varifold supported on \mathbb{R}_+^3 with its weight measure denoted by μ and let W a rectifiable 2-varifold supported on $\partial\mathbb{R}_+^3$ with its weight measure denoted by η , satisfying a contact*

angle condition as in Definition 2.1 (and adopt the notations in Section 2.1 below). For any $a \in \mathbb{R}^3$, consider the functions $g_a(r), \hat{g}_a(r)$ defined by

$$\begin{aligned} g_a(r) &:= \frac{\mu(B_r(a))}{\pi r^2} - \cos \theta \frac{\eta(B_r(a))}{\pi r^2} + \frac{1}{16\pi} \int_{B_r(a)} |\vec{\mathbf{H}}|^2 d\mu + \frac{1}{2\pi r^2} \int_{B_r(a)} \vec{\mathbf{H}} \cdot (x - a) d\mu, \\ \hat{g}_a(r) &:= \frac{\mu(\hat{B}_r(a))}{\pi r^2} - \cos \theta \frac{\eta(\hat{B}_r(a))}{\pi r^2} + \frac{1}{16\pi} \int_{\hat{B}_r(a)} |\vec{\mathbf{H}}|^2 d\mu + \frac{1}{2\pi r^2} \int_{\hat{B}_r(a)} \vec{\mathbf{H}} \cdot (x - \tilde{a}) d\mu. \end{aligned}$$

Then for any $0 < \sigma < \rho < \infty$, we have

$$\begin{aligned} &(g_a(\rho) + \hat{g}_a(\rho)) - (g_a(\sigma) + \hat{g}_a(\sigma)) \\ &= \frac{1}{\pi} \int_{B_\rho(a) \setminus B_\sigma(a)} \left| \frac{1}{4} \vec{\mathbf{H}} + \frac{(x - a)^\perp}{|x - a|^2} \right|^2 d\mu + \frac{1}{\pi} \int_{\hat{B}_\rho(a) \setminus \hat{B}_\sigma(a)} \left| \frac{1}{4} \vec{\mathbf{H}} + \frac{(x - \tilde{a})^\perp}{|x - \tilde{a}|^2} \right|^2 d\mu \\ &\quad - 2 \cos \theta \int_{B_\rho(a) \setminus B_\sigma(a)} \frac{a_3^2}{|x - a|^4} d\eta - 2 \cos \theta \int_{\hat{B}_\rho(a) \setminus \hat{B}_\sigma(a)} \frac{\tilde{a}_3^2}{|x - \tilde{a}|^4} d\eta, \end{aligned} \quad (1.2)$$

where $(x - a)^\perp := \nu_\Sigma \cdot (x - a) \nu_\Sigma$.

As an application, a Li-Yau-type inequality together with the characterization of the equality case is obtained for capillary immersions in the half-space. For the precise definition of capillary immersions, see Section 3.1. We will adopt the notations that are used in Section 3.1.

The Willmore energy of capillary immersion F is defined in the usual sense, that is,

Definition 1.2. Given $\theta \in (0, \pi)$ and a θ -capillary immersion $F : \Sigma \rightarrow \overline{\mathbb{R}_+^3}$. The Willmore energy of the capillary immersion F is defined as

$$\mathcal{W}(F) := \frac{1}{4} \int |\vec{\mathbf{H}}|^2 d\mu.$$

$\mathcal{W}(F)$ is a conformal invariant with respect to conformal diffeomorphisms. Detailed discussions regarding the definition of the Willmore functional for capillary surfaces are provided in Section 3.2.

Theorem 1.3. Given $\theta \in (0, \pi)$, let $F : \Sigma \rightarrow \overline{\mathbb{R}_+^3}$ be a compact θ -capillary immersion, satisfying (3.1). Then we have

(1) the Li-Yau-type inequality

$$\mathcal{W}(F) \geq 4(1 - \cos \theta) \pi \Theta^2(\mu, x_0) \quad (1.3)$$

holds for every $x_0 \in F(\partial\Sigma)$, where $\Theta^2(\mu, x)$ denotes the density of μ at x_0 .

(2) the sharp estimate on Willmore energy:

$$\mathcal{W}(F) \geq 2(1 - \cos \theta) \pi, \quad (1.4)$$

and if

$$\mathcal{W}(F) < 4(1 - \cos \theta) \pi,$$

then $F : \partial\Sigma \subset \partial\mathbb{R}_+^3$ must be an embedding.

(3) For $\theta \in [\frac{\pi}{2}, \pi)$, if

$$W(F) < 4\pi,$$

then $F : \Sigma \rightarrow \overline{\mathbb{R}^3_+}$ is an embedding.

Moreover, equality in (1.4) holds if and only if $F(\Sigma)$ is a θ -spherical cap.

Inequality (1.4) was proved using a flow method by the first and the second author in [28, Corollary 1.3] for convex capillary surfaces in $\overline{\mathbb{R}^3_+}$, while the contact angle θ is restricted within the range $(0, \frac{\pi}{2}]$. Therefore Theorem 1.3 complements the full range of contact angle and moreover removes the assumption of the convexity.

Due to the conformal invariance of the Willmore functional, Theorem 1.3 implies the same inequality as (1.4) for capillary surfaces in the unit ball, which interestingly implies a new optimal area estimate for minimal capillary surfaces in \mathbb{B}^3 .

Theorem 1.4. *Given $\theta \in (0, \pi)$, let $F : \Sigma \rightarrow \overline{\mathbb{B}^3}$ be a θ -capillary minimal immersion, satisfying (3.1). Then there holds*

$$2|\Sigma| - \cos \theta |T| \geq 2(1 - \cos \theta)\pi. \quad (1.5)$$

Equality holds if and only if $F(\Sigma)$ is a totally geodesic disk in $\overline{\mathbb{B}^3}$. Here $|T|$ is the area of the “wetting” part defined in Section 3.1.

As a special case of the isoperimetric inequality for minimal submanifolds proved by Brendle [4], any capillary minimal surface Σ in the unit ball satisfies the following optimal estimate

$$|\Sigma| \geq \sin^2 \theta \pi = \sin^2 \theta |\mathbb{B}^2|$$

with equality holding if and only if Σ is a flat disk. In fact, the higher dimensional counterparts also hold, see [5, Theorem 5.5]. Restricting $\theta = \frac{\pi}{2}$, i.e., in the free boundary case, this area lower bound was obtained in [10] and [3]. (1.5) now provides a new optimal area estimate for minimal capillary surfaces in the unit ball. It seems that both area estimates do not imply each other. In view of [5], it would be interesting to seek for a higher dimensional generalization of (1.5).

In the last part of the paper, Section 4, we establish a monotonicity formula (4.7) for capillary surfaces in \mathbb{B}^3 . This monotonicity formula is a generalization of Volkman’s theorem for surfaces with free boundary in the unit ball [27, Theorem 3.1]. Due to the extra 2-varifold W supported on the unit sphere (as in Theorem 1.1), our monotonicity formula (4.7) becomes more involved than his. We refer the readers to Theorem 4.3 below. Theorem 1.4 can also be proved directly by this monotonicity formula.

We end the introduction with a short summary of the fruitful results of the study of capillary surfaces. The existence, regularity, and geometric properties of capillary surfaces have attracted more and more attention from differential geometers and geometric analysts, see the nice book of Finn [9] for a through introduction. Here we mention some recent progress on this topic. In \mathbb{R}^3 , optimal boundary regularity for solutions to the capillarity problems (volume-constrained local minimizers of the free energy functional) was obtained by Taylor [26], while the partial regularity was obtained recently by De Philippis-Maggi [7, 8] in \mathbb{R}^n ($n \geq 4$), even in the anisotropic setting. Quite recently, using the Min-Max method, the existence of capillary minimal or CMC hypersurfaces

in compact 3-manifolds with boundary has been shown independently by De Masi-De Philippis [6] and Li-Zhou-Zhu [19]. In terms of the classical differential geometry, Hong-Saturnino [12] carried out curvature and index estimates for compact and non-compact capillary surfaces. In [13], we proved a Heintze-Karcher-type inequality and give the characterization of smooth CMC capillary hypersurfaces in the half-space or in a wedge, namely, the Alexandrov-type theorem. The anisotropic counterpart was tackled in [14, 15] as well. See also [29] for a non-smooth generalization.

1.2. Organization of the Paper. In Section 2 we first introduce the capillary surface in the half-space in the setting of varifolds and then provide the proof of Theorem 1.1, In Section 3.2 we use the monotonicity formula to estimate the Willmore energy for capillary surfaces in the half-space and prove the Li-Yau inequality (1.3) and Theorem 1.3. The Willmore energy for capillary surfaces in the unit ball is discussed in Section 3.3, together with one proof of Theorem 1.4. Finally, in Section 4, we prove the Simon-type monotonicity formulas in the unit ball, and then provide an alternative proof of Theorem 1.4.

2. MONOTONICITY FORMULA IN THE HALF-SPACE

2.1. Set-ups. We will be working on the Euclidean space \mathbb{R}^3 , with the Euclidean metric denoted by g_{euc} and the corresponding Levi-Civita connection denoted by ∇ . $\mathbb{R}_+^3 = \{x : x_3 > 0\}$ is the open upper half-space and $E_3 := (0, 0, 1)$.

Let $\xi : \mathbb{R}^3 \rightarrow \partial\mathbb{R}_+^3$ denote the unique point projection onto the hyperplane $\partial\mathbb{R}_+^3$. It is easy to see that for any $x \in \mathbb{R}^3$, $\xi(x) = x - x_3 E_3$, and hence ξ is a smooth map. Define

$$\tilde{x} := 2\xi(x) - x = x - 2x_3 E_3$$

to be the reflection of x across $\partial\mathbb{R}_+^3$. Given a point $a \in \mathbb{R}^3$, define

$$r = |x - a|, \quad \tilde{r} = |\tilde{x} - a|,$$

it is clear that $\tilde{r} = |\tilde{x} - a| = |x - \tilde{a}|$.

We refer to [24] for the background materials in Geometric Measure Theory and consider the following weak formulation of capillary surfaces.

Definition 2.1. Given $\theta \in (0, \pi)$, let V be a rectifiable 2-varifold supported on $\overline{\mathbb{R}_+^3}$ with its weight measure denoted by μ and let W a rectifiable 2-varifold supported on $\partial\mathbb{R}_+^3$ with its weight measure denoted by η . (V, W) is said to satisfy a *contact angle condition* θ if there exists a μ -measurable vector field $\vec{\mathbf{H}} \in \mathcal{L}^1(\mathbb{R}^3, \mu)$ with $\vec{\mathbf{H}}(x) \in T_x \partial\mathbb{R}_+^3$ for μ -a.e. $x \in \partial\mathbb{R}_+^3$, such that for every $X \in C_c^1(\mathbb{R}^3; \mathbb{R}^3)$ with X tangent to $\partial\mathbb{R}_+^3$,

$$\int_{\mathbb{R}^3} \text{div}_\Sigma X d\mu - \cos \theta \int_{\partial\mathbb{R}_+^3} \text{div}_{\partial\mathbb{R}_+^3} X d\eta = - \int_{\mathbb{R}^3} \vec{\mathbf{H}} \cdot X d\mu, \quad (2.1)$$

where $\Sigma := \text{spt } \mu$ is countably 2-rectifiable, for simplicity we also set $T := \text{spt } \eta$. In particular, if X vanishes along $\partial\mathbb{R}_+^3$, there holds that

$$\int_{\mathbb{R}^3} \text{div}_\Sigma X d\mu = - \int_{\mathbb{R}^3} \vec{\mathbf{H}} \cdot X d\mu.$$

This definition was introduced in [6], which follows from the one initiated in [16], with the boundary part weakened to be just a 2-varifold.

Notice that the first variation formula (2.1) is valid if X is merely a Lipschitz vector field.

In this section, we consider those V, W that are integral 2-varifolds with $\mu(\partial\mathbb{R}_+^3) = 0$ and $\vec{\mathbf{H}} \in \mathcal{L}^2(\mathbb{R}^3, \mu)$. It follows that $\vec{\mathbf{H}} \perp \Sigma$ for μ -a.e. $x \in \Sigma$ thanks to the well known Brakke's perpendicular theorem ([2, Sect. 5.8]), so that for any vector $v \in \mathbb{R}^3$, one has

$$2 \left| \frac{1}{4} \vec{\mathbf{H}} + v^\perp \right|^2 = \frac{1}{8} |\vec{\mathbf{H}}|^2 + 2|v^\perp|^2 + \vec{\mathbf{H}} \cdot v, \quad (2.2)$$

where v^\perp denotes the normal part of v with respect to the approximate tangent space $T_x \Sigma$.

2.2. A Monotonicity Formula. A general monotonicity formula for the pairs of varifolds satisfying a contact condition has been established in [6] (see also [16]) with none sharp constants, which generalizes a previous result in [11] for rectifiable free boundary varifolds. We are interested in the Simon-type monotonicity formula (1.1) for the capillary case with optimal constants.

Proof of Theorem 1.1. We follow closely the reflection idea in [11]. See also [16]. Define $l(s) := \left(\left(\frac{1}{s\sigma} \right)^2 - \frac{1}{\rho^2} \right)_+$, where $s_\sigma := \max\{s, \sigma\}$. $l(s)$ is then a Lipschitz cut-off function and it is easy to see that

$$l(s) = \begin{cases} \left(\frac{1}{\sigma^2} - \frac{1}{\rho^2} \right), & 0 < s \leq \sigma, \\ \left(\frac{1}{s^2} - \frac{1}{\rho^2} \right), & \sigma < s \leq \rho, \\ 0, & \rho < s. \end{cases}$$

We wish to find a suitable vector field to test (2.1). The construction is inspired by [11] and is nowadays quiet standard. Precisely, we define

$$\begin{aligned} X_1(x) &= \left(\left(\frac{1}{|x-a|_\sigma} \right)^2 - \frac{1}{\rho^2} \right)_+ (x-a), \\ X_2(x) &= \left(\left(\frac{1}{|x-\tilde{a}|_\sigma} \right)^2 - \frac{1}{\rho^2} \right)_+ (x-\tilde{a}), \\ X(x) &:= X_1(x) + X_2(x). \end{aligned}$$

For $x \in \partial\mathbb{R}_+^3$, since $x = \tilde{x}$ we simply have

$$|x-a|_\sigma = |\tilde{x}-a|_\sigma = |x-\tilde{a}|_\sigma,$$

and hence

$$X(x) = \left(\left(\frac{1}{|x-a|_\sigma} \right)^2 - \frac{1}{\rho^2} \right)_+ (2x - (a + \tilde{a})) \in \partial\mathbb{R}_+^3.$$

Let $\hat{B}_r(a) = \{x : |x - \tilde{a}| < r\}$ and consider the partitions of \mathbb{R}^3 :

$$\begin{aligned} \mathcal{F}_1 &:= \{B_\sigma(a), B_\rho(a) \setminus B_\sigma(a), \mathbb{R}^3 \setminus B_\rho(a)\}, \\ \mathcal{F}_2 &:= \{\hat{B}_\sigma(a), \hat{B}_\rho(a) \setminus \hat{B}_\sigma(a), \mathbb{R}^3 \setminus \hat{B}_\rho(a)\}. \end{aligned}$$

We wish to test (2.1) by $X(x)$. To this end, we compute

$$\int_A \operatorname{div}_\Sigma X_i d\mu - \cos \theta \int_A \operatorname{div}_{\partial\mathbb{R}_+^3} X_i d\eta, \text{ and } \int_A \vec{\mathbf{H}} \cdot X_i d\mu$$

for all sets $A \in \mathcal{F}_i$, $i = 1, 2$, separately.

For X_2 : on $0 \leq |x - \tilde{a}| \leq \sigma$, direct computation shows that

$$\nabla X_2(x) = \left(\frac{1}{\sigma^2} - \frac{1}{\rho^2} \right) \operatorname{Id},$$

and hence

$$\begin{aligned} \int_{\hat{B}_\sigma(a)} \operatorname{div}_\Sigma X_2 d\mu - \cos \theta \int_{\hat{B}_\sigma(a)} \operatorname{div}_{\partial\mathbb{R}_+^3} X_2 d\eta &= \left(\frac{2}{\sigma^2} - \frac{2}{\rho^2} \right) (\mu(\hat{B}_\sigma(a)) - \cos \theta \eta(\hat{B}_\sigma(a))), \\ \int_{\hat{B}_\sigma(a)} \vec{\mathbf{H}} \cdot X_2 d\mu &= \left(\frac{1}{\sigma^2} - \frac{1}{\rho^2} \right) \int_{\hat{B}_\sigma(a)} \vec{\mathbf{H}} \cdot (x - \tilde{a}) d\mu. \end{aligned}$$

On $\sigma < |x - \tilde{a}| \leq \rho$, we have $X_2(x) = \left(\frac{1}{|x - \tilde{a}|^2} - \frac{1}{\rho^2} \right) (x - \tilde{a})$. By noticing that

$$\nabla \left(\frac{1}{|x - \tilde{a}|^2} \right) = -2 \frac{1}{|x - \tilde{a}|^4} (x - \tilde{a}),$$

we have

$$\begin{aligned} \nabla X_2(x) &= \left(\frac{1}{|x - \tilde{a}|^2} - \frac{1}{\rho^2} \right) \operatorname{Id} - 2 \frac{1}{|x - \tilde{a}|^4} (x - \tilde{a}) \otimes (x - \tilde{a}), \\ \operatorname{div} X_2(x) &= 3 \left(\frac{1}{|x - \tilde{a}|^2} - \frac{1}{\rho^2} \right) - \frac{2}{|x - \tilde{a}|^2} = \frac{1}{|x - \tilde{a}|^2} - \frac{3}{\rho^2}, \\ \operatorname{div}_\Sigma X_2(x) &= \operatorname{div} X_2(x) - \nabla X_2[\nu_\Sigma] \cdot \nu_\Sigma = -\frac{2}{\rho^2} + 2 \left| \frac{(x - \tilde{a})^\perp}{|x - \tilde{a}|^2} \right|^2, \\ \operatorname{div}_{\partial\mathbb{R}_+^3} X_2(x) &= \operatorname{div} X_2(x) - \nabla X_2[E_3] \cdot E_3 = -\frac{2}{\rho^2} + \frac{2\tilde{a}_3^2}{|x - \tilde{a}|^4}. \end{aligned}$$

It follows that

$$\begin{aligned} &\int_{\hat{B}_\rho(a) \setminus \hat{B}_\sigma(a)} \operatorname{div}_\Sigma X_2 d\mu - \cos \theta \int_{\hat{B}_\rho(a) \setminus \hat{B}_\sigma(a)} \operatorname{div}_{\partial\mathbb{R}_+^3} X_2 d\eta \\ &= -\frac{2}{\rho^2} (\mu(\hat{B}_\rho(a) \setminus \hat{B}_\sigma(a)) - \cos \theta \eta(\hat{B}_\rho(a) \setminus \hat{B}_\sigma(a))) + 2 \int_{\hat{B}_\rho(a) \setminus \hat{B}_\sigma(a)} \left| \frac{(x - \tilde{a})^\perp}{|x - \tilde{a}|^2} \right|^2 d\mu \\ &\quad - 2 \cos \theta \int_{\hat{B}_\rho(a) \setminus \hat{B}_\sigma(a)} \frac{\tilde{a}_3^2}{|x - \tilde{a}|^4} d\eta, \\ &\int_{\hat{B}_\rho(a) \setminus \hat{B}_\sigma(a)} \vec{\mathbf{H}} \cdot X_2 d\mu = -\frac{1}{\rho^2} \int_{\hat{B}_\rho(a) \setminus \hat{B}_\sigma(a)} \vec{\mathbf{H}} \cdot (x - \tilde{a}) d\mu + \int_{\hat{B}_\rho(a) \setminus \hat{B}_\sigma(a)} \vec{\mathbf{H}} \cdot \frac{x - \tilde{a}}{|x - \tilde{a}|^2} d\mu. \end{aligned}$$

Combining these computations, we obtain

$$\begin{aligned} & \int_{\mathbb{R}^3} \operatorname{div}_{\Sigma} X_2 d\mu - \cos \theta \int_{\mathbb{R}^3} \operatorname{div}_{\partial\mathbb{R}_+^3} X_2 d\eta \\ &= \frac{2}{\sigma^2} (\mu(\hat{B}_{\sigma}(a)) - \cos \theta \eta(\hat{B}_{\sigma}(a))) - \frac{2}{\rho^2} (\mu(\hat{B}_{\rho}(a)) - \cos \theta \eta(\hat{B}_{\rho}(a))) \\ & \quad + 2 \int_{\hat{B}_{\rho}(a) \setminus \hat{B}_{\sigma}(a)} \left| \frac{(x - \tilde{a})^{\perp}}{|x - \tilde{a}|^2} \right|^2 d\mu - 2 \cos \theta \int_{\hat{B}_{\rho}(a) \setminus \hat{B}_{\sigma}(a)} \frac{a_3^2}{|x - \tilde{a}|^4} d\eta, \end{aligned}$$

and

$$\begin{aligned} \int_{\mathbb{R}^3} \vec{\mathbf{H}} \cdot X_2 d\mu &= \frac{1}{\sigma^2} \int_{\hat{B}_{\sigma}(a)} \vec{\mathbf{H}} \cdot (x - \tilde{a}) d\mu - \frac{1}{\rho^2} \int_{\hat{B}_{\rho}(a)} \vec{\mathbf{H}} \cdot (x - \tilde{a}) d\mu \\ & \quad + \int_{\hat{B}_{\rho}(a) \setminus \hat{B}_{\sigma}(a)} \vec{\mathbf{H}} \cdot \frac{x - \tilde{a}}{|x - \tilde{a}|^2} d\mu. \end{aligned}$$

Similar computations hold for X_1 and we obtain

$$\begin{aligned} & \int_{\mathbb{R}^3} \operatorname{div}_{\Sigma} X_1 d\mu - \cos \theta \int_{\mathbb{R}^3} \operatorname{div}_{\partial\mathbb{R}_+^3} X_1 d\eta \\ &= \frac{2}{\sigma^2} (\mu(B_{\sigma}(a)) - \cos \theta \eta(B_{\sigma}(a))) - \frac{2}{\rho^2} (\mu(B_{\rho}(a)) - \cos \theta \eta(B_{\rho}(a))) \\ & \quad + 2 \int_{B_{\rho}(a) \setminus B_{\sigma}(a)} \left| \frac{(x - a)^{\perp}}{|x - a|^2} \right|^2 d\mu - 2 \cos \theta \int_{B_{\rho}(a) \setminus B_{\sigma}(a)} \frac{a_3^2}{|x - a|^4} d\eta, \end{aligned}$$

and

$$\begin{aligned} \int_{\mathbb{R}^3} \vec{\mathbf{H}} \cdot X_1 d\mu &= \frac{1}{\sigma^2} \int_{B_{\sigma}(a)} \vec{\mathbf{H}} \cdot (x - a) d\mu - \frac{1}{\rho^2} \int_{B_{\rho}(a)} \vec{\mathbf{H}} \cdot (x - a) d\mu \\ & \quad + \int_{B_{\rho}(a) \setminus B_{\sigma}(a)} \vec{\mathbf{H}} \cdot \frac{x - a}{|x - a|^2} d\mu. \end{aligned}$$

On the other hand, thanks to (2.2), we have

$$\begin{aligned} \int_{B_{\rho}(a) \setminus B_{\sigma}(a)} \vec{\mathbf{H}} \cdot \frac{x - a}{|x - a|^2} d\mu &= 2 \int_{B_{\rho}(a) \setminus B_{\sigma}(a)} \left| \frac{1}{4} \vec{\mathbf{H}} + \frac{(x - a)^{\perp}}{|x - a|^2} \right|^2 d\mu \\ & \quad - \frac{1}{8} \int_{B_{\rho}(a) \setminus B_{\sigma}(a)} |\vec{\mathbf{H}}|^2 d\mu - 2 \int_{B_{\rho}(a) \setminus B_{\sigma}(a)} \left| \frac{(x - a)^{\perp}}{|x - a|^2} \right|^2 d\mu, \\ \int_{\hat{B}_{\rho}(a) \setminus \hat{B}_{\sigma}(a)} \vec{\mathbf{H}} \cdot \frac{x - \tilde{a}}{|x - \tilde{a}|^2} d\mu &= 2 \int_{\hat{B}_{\rho}(a) \setminus \hat{B}_{\sigma}(a)} \left| \frac{1}{4} \vec{\mathbf{H}} + \frac{(x - \tilde{a})^{\perp}}{|x - \tilde{a}|^2} \right|^2 d\mu \\ & \quad - \frac{1}{8} \int_{\hat{B}_{\rho}(a) \setminus \hat{B}_{\sigma}(a)} |\vec{\mathbf{H}}|^2 d\mu - 2 \int_{\hat{B}_{\rho}(a) \setminus \hat{B}_{\sigma}(a)} \left| \frac{(x - \tilde{a})^{\perp}}{|x - \tilde{a}|^2} \right|^2 d\mu. \end{aligned}$$

Putting these facts into the first variation formula (2.1) and recalling the definition of $g_a(r), \hat{g}_a(r)$, we deduce (1.2). \square

Remark 2.2. We remark that in Theorem 1.1:

- (1) $B_r(a) \cap \partial\mathbb{R}_+^3 = \hat{B}_r(a) \cap \partial\mathbb{R}_+^3$, and hence $\eta(B_r(a)) = \eta(\hat{B}_r(a))$.

(2) $a_3^2 = \tilde{a}_3^2$ and hence the corresponding terms involving them are the same.

Proposition 2.3. *Under the assumptions of Theorem 1.1, the following statements hold:*

(1) *Given $\theta \in [\frac{\pi}{2}, \pi)$. For every $a \in \mathbb{R}^3$, the tilde-density*

$$\tilde{\Theta}^2(\mu - \cos \theta \eta, a) := \lim_{r \searrow 0} \left(\frac{(\mu - \cos \theta \eta)(B_r(a))}{\pi r^2} + \frac{(\mu - \cos \theta \eta)(\hat{B}_r(a))}{\pi r^2} \right)$$

exists. Moreover, the function $x \mapsto \tilde{\Theta}(\mu - \cos \theta \eta, x)$ is upper semi-continuous in \mathbb{R}^3 ;

(2) *Given $\theta \in (0, \pi)$. For every $a \in \partial \mathbb{R}^3$, the tilde-density $\tilde{\Theta}^2(\mu - \cos \theta \eta, a)$ exists, and the function $x \mapsto \tilde{\Theta}(\mu - \cos \theta \eta, x)$ is upper semi-continuous in $\partial \mathbb{R}_+^3$.*

Proof. Define

$$R(r) := \frac{1}{2\pi r^2} \int_{B_r(a)} \vec{\mathbf{H}} \cdot (x - a) d\mu + \frac{1}{2\pi r^2} \int_{\hat{B}_r(a)} \vec{\mathbf{H}} \cdot (x - \tilde{a}) d\mu,$$

and

$$\begin{aligned} G_\theta(r) &:= \frac{(\mu - \cos \theta \eta)(B_r(a))}{\pi r^2} + \frac{(\mu - \cos \theta \eta)(\hat{B}_r(a))}{\pi r^2} \\ &\quad + \frac{1}{16\pi} \int_{B_r(a)} |\vec{\mathbf{H}}|^2 d\mu + \frac{1}{16\pi} \int_{\hat{B}_r(a)} |\vec{\mathbf{H}}|^2 d\mu + R(r), \end{aligned}$$

then from (1.2) and thanks to $\theta \in [\pi/2, \pi)$, we know that $G_\theta(r)$ is monotonically nondecreasing, so that $\lim_{r \rightarrow 0^+} G_\theta(r)$ exists.

For $R(r)$, we estimate with Hölder inequality:

$$\begin{aligned} |R(r)| &\leq \left(\frac{\mu(B_r(a))}{\pi r^2} \right)^{\frac{1}{2}} \left(\frac{1}{4\pi} \int_{B_r(a)} |\vec{\mathbf{H}}|^2 d\mu \right)^{\frac{1}{2}} \\ &\quad + \left(\frac{\mu(\hat{B}_r(a))}{\pi r^2} \right)^{\frac{1}{2}} \left(\frac{1}{4\pi} \int_{\hat{B}_r(a)} |\vec{\mathbf{H}}|^2 d\mu \right)^{\frac{1}{2}}. \end{aligned} \tag{2.3}$$

Moreover, for $\frac{1}{4} < \epsilon < \frac{1}{2}$, Young's inequality gives

$$\begin{aligned} |R(r)| &\leq \epsilon \frac{\mu(B_r(a))}{\pi r^2} + \frac{1}{16\pi\epsilon} \int_{B_r(a)} |\vec{\mathbf{H}}|^2 d\mu \\ &\quad + \epsilon \frac{\mu(\hat{B}_r(a))}{\pi r^2} + \frac{1}{16\pi\epsilon} \int_{\hat{B}_r(a)} |\vec{\mathbf{H}}|^2 d\mu. \end{aligned}$$

By virtue of the monotonicity of $G_\theta(r)$, we obtain

$$\begin{aligned}
& \frac{(\mu - \cos \theta \eta)(B_\sigma(a))}{\pi \sigma^2} + \frac{(\mu - \cos \theta \eta)(\hat{B}_\sigma(a))}{\pi \sigma^2} \\
& \leq \frac{(\mu - \cos \theta \eta)(B_\rho(a))}{\pi \rho^2} + \frac{(\mu - \cos \theta \eta)(\hat{B}_\rho(a))}{\pi \rho^2} \\
& \quad + \frac{1}{16\pi} \int_{B_\rho(a)} |\vec{\mathbf{H}}|^2 d\mu + \frac{1}{16\pi} \int_{\hat{B}_\rho(a)} |\vec{\mathbf{H}}|^2 d\mu + |R(\rho)| + |R(\sigma)| \\
& \leq (1 + \epsilon) \left(\frac{(\mu - \cos \theta \eta)(B_\rho(a))}{\pi \rho^2} + \frac{(\mu - \cos \theta \eta)(\hat{B}_\rho(a))}{\pi \rho^2} \right) \\
& \quad + \frac{1 + 2\epsilon^{-1}}{16\pi} \left(\int_{B_\rho(a)} |\vec{\mathbf{H}}|^2 d\mu + \int_{\hat{B}_\rho(a)} |\vec{\mathbf{H}}|^2 d\mu \right) \\
& \quad + \epsilon \left(\frac{(\mu - \cos \theta \eta)(B_\sigma(a))}{\pi \sigma^2} + \frac{(\mu - \cos \theta \eta)(\hat{B}_\sigma(a))}{\pi \sigma^2} \right).
\end{aligned}$$

In particular, since $\frac{1}{4} < \epsilon < \frac{1}{2}$, this yields for $0 < r < R < \infty$

$$\begin{aligned}
& \frac{(\mu - \cos \theta \eta)(B_r(a))}{\pi r^2} + \frac{(\mu - \cos \theta \eta)(\hat{B}_r(a))}{\pi r^2} \\
& \leq 3 \left(\frac{(\mu - \cos \theta \eta)(B_R(a))}{\pi R^2} + \frac{(\mu - \cos \theta \eta)(\hat{B}_R(a))}{\pi R^2} \right) + \frac{9}{8\pi} \int_{\mathbb{R}^3} |\vec{\mathbf{H}}|^2 d\mu < \infty,
\end{aligned} \tag{2.4}$$

which, together with (2.3), gives

$$\lim_{r \rightarrow 0^+} R(r) = 0,$$

implying that the tilde-density $\tilde{\Theta}(\mu - \cos \theta \eta, a)$ exists, and also

$$\begin{aligned}
\tilde{\Theta}(\mu - \cos \theta \eta, a) & \leq \left(\frac{(\mu - \cos \theta \eta)(B_\rho(a))}{\pi \rho^2} + \frac{(\mu - \cos \theta \eta)(\hat{B}_\rho(a))}{\pi \rho^2} \right) \\
& \quad + \frac{1}{16\pi} \left(\int_{B_\rho(a)} |\vec{\mathbf{H}}|^2 d\mu + \int_{\hat{B}_\rho(a)} |\vec{\mathbf{H}}|^2 d\mu \right) + R(\rho).
\end{aligned}$$

Thus for a fixed $R > 0$ and a sequence of points in \mathbb{R}^3 such that $x_j \rightarrow a$, we have: for $0 < \rho < R$, it holds that

$$\begin{aligned}
& \frac{(\mu - \cos \theta \eta)(\overline{B_\rho(a)})}{\pi \rho^2} + \frac{(\mu - \cos \theta \eta)(\widehat{B}_\rho(a))}{\pi \rho^2} \\
& \geq \limsup_{j \rightarrow \infty} \left(\frac{(\mu - \cos \theta \eta)(\overline{B_\rho(x_j)})}{\pi \rho^2} + \frac{(\mu - \cos \theta \eta)(\widehat{B}_\rho(x_j))}{\pi \rho^2} \right) \\
& \geq \limsup_{j \rightarrow \infty} \left(\tilde{\Theta}(\mu - \cos \theta \eta, x_j) - \frac{1}{16\pi} \left(\int_{B_\rho(x_j)} |\vec{\mathbf{H}}|^2 d\mu + \int_{\widehat{B}_\rho(x_j)} |\vec{\mathbf{H}}|^2 d\mu \right) - R(\rho) \right) \\
& \stackrel{(2.4)}{\geq} \limsup_{j \rightarrow \infty} \tilde{\Theta}(\mu - \cos \theta \eta, x_j) \\
& \quad - C \left(\frac{(\mu - \cos \theta \eta)(B_R(a)) + (\mu - \cos \theta \eta)(\widehat{B}_R(a))}{\pi R^2} + \mathcal{W}(\mu) \right)^{\frac{1}{2}} \|\vec{\mathbf{H}}\|_{L^2(B_{2\rho}(a))}.
\end{aligned}$$

Letting $\rho \rightarrow 0^+$, this gives

$$\tilde{\Theta}^2(\mu - \cos \theta \eta, a) \geq \limsup_{j \rightarrow \infty} \tilde{\Theta}(\mu - \cos \theta \eta, x_j),$$

which completes the proof of (1).

To prove (2), notice that for any $a \in \partial \mathbb{R}_+^3$, $a_3 = \tilde{a}_3 = 0$, and hence as $\theta \in (0, \pi)$, the function $G_\theta(r)$ is also monotonically nondecreasing for any such a . Following the proof of (1), we conclude (2). \square

Remark 2.4. Suppose that μ, η are compactly supported, we may thus use (2.3) to find that: for $\theta \in [\frac{\pi}{2}, \pi)$, $a \in \mathbb{R}^3$,

$$\lim_{r \rightarrow \infty} |R(r)| = 0,$$

and obtain

$$\lim_{r \rightarrow \infty} (g_a(r) + \hat{g}_a(r)) = \frac{1}{8\pi} \int_{\mathbb{R}^3} |\vec{\mathbf{H}}|^2 d\mu.$$

Thus by letting $\sigma \rightarrow 0^+$ and $\rho \rightarrow \infty$ in (1.2), we obtain

$$\begin{aligned}
& \frac{1}{\pi} \int_{\mathbb{R}^3} \left| \frac{1}{4} \vec{\mathbf{H}} + \frac{(x-a)^\perp}{|x-a|^2} \right|^2 d\mu + \frac{1}{\pi} \int_{\mathbb{R}^3} \left| \frac{1}{4} \vec{\mathbf{H}} + \frac{(x-\tilde{a})^\perp}{|x-\tilde{a}|^2} \right|^2 d\mu \\
& \quad - 2 \cos \theta \int_{\mathbb{R}^3} \frac{a_3^2}{|x-a|^4} d\eta - 2 \cos \theta \int_{\mathbb{R}^3} \frac{\tilde{a}_3^2}{|x-\tilde{a}|^4} d\eta \\
& = \frac{1}{8\pi} \int_{\mathbb{R}^3} |\vec{\mathbf{H}}|^2 d\mu - \tilde{\Theta}^2(\mu - \cos \theta \eta, a).
\end{aligned} \tag{2.5}$$

This is also true for any $a \in \mathbb{R}^3$ and any $\theta \in (0, \frac{\pi}{2})$, provided that $\tilde{\Theta}^2(\mu - \cos \theta \eta, a)$ exists.

Similarly, for $\theta \in (0, \pi)$ and $a \in \partial \mathbb{R}_+^3$, we have

$$\frac{2}{\pi} \int_{\mathbb{R}^3} \left| \frac{1}{4} \vec{\mathbf{H}} + \frac{(x-a)^\perp}{|x-a|^2} \right|^2 d\mu = \frac{1}{8\pi} \int_{\mathbb{R}^3} |\vec{\mathbf{H}}|^2 d\mu - \tilde{\Theta}^2(\mu - \cos \theta \eta, a). \tag{2.6}$$

3. WILLMORE ENERGY AND LI-YAU-TYPE INEQUALITY FOR CAPILLARY IMMERSIONS

3.1. Capillary Immersions. Given a compact orientable smooth surface Σ , with non-empty boundary $\partial\Sigma$. Let $F : \Sigma \rightarrow \overline{\mathbb{R}_+^3}$ be an orientation preserving proper immersion, that is, F is smooth on the interior of Σ and C^2 up to the boundary, such that $F(\text{int}\Sigma) \subset \mathbb{R}_+^3$ and $F(\partial\Sigma) \subset \partial\mathbb{R}_+^3$. In this way, F induces an immersion of $\partial\Sigma$ into $\partial\mathbb{R}_+^3$.

Fix a global unit normal field ν to Σ along F , which determines an orientation on Σ and an induced orientation on $\partial\Sigma$ given by a tangential vector field τ along $\partial\Sigma$. Denote by μ the unit conormal to $\partial\Sigma$ in Σ so that $\{\tau, \nu, \mu\}$ is compatible with $\{E_1, E_2, E_3\}$ of \mathbb{R}^3 . Let $\bar{\nu}$ be the unit normal to $\partial\Sigma$ in $\partial\mathbb{R}_+^3$ so that $\{\bar{\nu}, -E_3\}$ compatible with $\{\nu, \mu\}$. Given $\theta \in (0, \pi)$, we say that the proper immersion $F : \Sigma \rightarrow \overline{\mathbb{R}_+^3}$ is a *capillary immersion* (or *θ -capillary immersion*) with contact angle θ if the angle determined by μ and $\bar{\nu}$ is constant and equals to θ along $\partial\Sigma$, that is, for any $x \in \partial\Sigma$,

$$\mu(x) = \sin \theta(-E_3) + \cos \theta \bar{\nu}(x).$$

Abuse of terminology, for the immersion $F : \partial\Sigma \rightarrow \partial\mathbb{R}_+^3$, we use T to denote the ‘‘enclosed domain’’ by $F(\partial\Sigma)$, which means the topological boundary ∂T is given by $F(\partial\Sigma)$ and it has induced orientation by τ . We use $|T|$ to denote the oriented area of T , which is defined to be

$$|T| = \sum_i \text{sgn}(T_i) |T_i|,$$

where each T_i is a bounded domain such that ∂T_i is a simply closed curve and $\text{sgn}(T_i) = +1$ if $\bar{\nu}$ points outward T_i , while $\text{sgn}(T_i) = -1$ if $\bar{\nu}$ points inward T_i . $|T|$ is the signed area of the so-called ‘‘wetting part’’.

Now given a θ -capillary immersion $F : \Sigma \rightarrow \overline{\mathbb{R}_+^3}$ for $\theta \in (0, \pi) \setminus \{\frac{\pi}{2}\}$. Let $g = F^*g_{\text{euc}}$ be the pull-back metric and μ_g be the induced area measure on Σ . The induced varifold of Σ is given by $V_g = \mu_g \otimes \delta_{T_p\Sigma}$, where $T_p\Sigma$ is the tangent space and its image as varifold is given by $V = F_{\#}(V_g)$, which is an integral 2-varifold in \mathbb{R}^3 with weight measure $\mu := (x \mapsto \mathcal{H}^0(F^{-1}(x))) \mathcal{H}^2 \llcorner F(\Sigma)$, see [24, Section 15].

Define the following measure:

$$\eta := \text{wind}(x) \mathcal{H}^2,$$

where $\text{wind}(x)$ is the winding number of $F(\partial\Sigma)$ about $x \in \partial\mathbb{R}^3$. Throughout the paper, in terms of capillary immersion, we always assume that

$$\text{wind}(x) \geq 0 \text{ for any } x \in \partial\mathbb{R}_+^3. \quad (3.1)$$

With this assumption, η is a positive Radon measure, and we may relate to it naturally the 2-varifold:

$$W = \eta \otimes \delta_{T_x \partial\mathbb{R}_+^3}.$$

Let $T = \text{spt}\eta \subset \partial\mathbb{R}_+^3$. By definition of winding number, the mass $\mathbb{M}(W)$ of W is exactly the oriented area of T (In fact, $\text{sgn}(T_i)$ will be always +1 thanks to (3.1)).

Proposition 3.1. *The pair (V, W) above satisfies the contact angle condition θ with square integrable generalized mean curvature as in Definition 2.1.*

Proof. In the case that Σ is an open surface and F is a properly immersion, the fact that V has square integrable generalized mean curvature in \mathbf{R}^3 has already been observed in [18, Section 2.1]. Here we prove the capillary version.

Let us consider any $X \in C_c^1(\mathbf{R}^3; \mathbf{R}^3)$ which is tangent to $\partial\mathbf{R}_+^3$, and let $\{f_t : \mathbf{R}^3 \rightarrow \mathbf{R}^3\}_{|t|<\epsilon}$ be the induced variation of X , that is,

$$f_0 = \text{id}, f_t(\partial\mathbf{R}_+^3) \subset \partial\mathbf{R}_+^3, \text{ and } \frac{d}{dt} \Big|_{t=0} f_t = X,$$

which induces a family of immersions: $\{\psi_t\}_{|t|<\epsilon}$, where $\psi_t = f_t \circ F : \Sigma \rightarrow \overline{\mathbf{R}_+^3}$. $\Psi : (-\epsilon, \epsilon) \times \Sigma \rightarrow \overline{\mathbf{R}_+^3}$, given by $\Psi(t, p) = \psi_t(p)$, is called an admissible variation. Let $\xi(p) = \frac{\partial\Psi}{\partial t}(0, p)$ for $p \in \Sigma$, then $\xi(p) = X(F(p))$.

For this family of immersions, let μ_{g_t} be the induced area measure of Σ with respect to the pull-back metric $g_t = \psi_t^* g_{\text{euc}}$, the area functional is then given by

$$\mathcal{A} : (-\epsilon, \epsilon) \rightarrow \mathbb{R}, \quad \mathcal{A}(t) = \int_{\Sigma} d\mu_{g_t}.$$

The wetted area functional $\mathcal{W}(t) : (-\epsilon, \epsilon) \rightarrow \mathbb{R}$ is defined by

$$\mathcal{W}(t) = \int_{[0,t] \times \partial\Sigma} \Psi^* \Omega,$$

where Ω is the volume form of $\partial\mathbf{R}_+^3$. We define the free energy functional by

$$\mathcal{F} : (-\epsilon, \epsilon) \rightarrow \mathbb{R}, \quad \mathcal{F}(t) = \mathcal{A}(t) - \cos\theta \mathcal{W}(t),$$

from the well-known first variation formula (see e.g., [1, (2.1)]) and the capillary condition, we find (recall that $\xi(p) = X(F(p))$)

$$\begin{aligned} \mathcal{F}'(0) &= - \int_{\Sigma} \mathbf{H}_{\Sigma}(p) \langle \xi(p), \nu(p) \rangle d\mu_g \\ &= - \int_{F(\Sigma)} \sum_{p \in F^{-1}(x)} \mathbf{H}_{\Sigma}(p) \langle \xi(p), \nu(p) \rangle d\mathcal{H}^2(x) \\ &=: - \int \langle X, \vec{\mathbf{H}} \rangle d\mu. \end{aligned}$$

On the other hand, it is not difficult to see that

$$A(t) = \mathbb{M}((f_t)_{\#} V),$$

and because $\mathbb{M}(W)$ is the oriented area,

$$\mathcal{W}'(0) = \frac{d}{dt} \Big|_{t=0} \mathbb{M}((f_t)_{\#} W).$$

By virtue of the well-known first variation formula for varifolds [24], we get

$$(\delta V - \cos\theta \delta W)[X] = \mathcal{F}'(0) = - \int \langle X, \vec{\mathbf{H}} \rangle d\mu,$$

as desired. \square

Remark 3.2. We point out that condition (3.1) is irredundant, because without this constraint, one could simply construct an example as Fig. 1, so that we could find some point $x \in F(\partial\Sigma)$ at which we no longer expect the upper-semi continuity of the tilde density (Proposition 2.3(1)) to be valid.

Precisely, given $\theta \in (\frac{\pi}{2}, \pi)$, let $F : \Sigma \rightarrow \overline{\mathbb{R}^3_+}$ be a θ -capillary immersion with $F(\partial\Sigma)$ as in Fig. 1. At the point x , because of the orientation of $F(\partial\Sigma)$, we have

$$\lim_{r \rightarrow 0^+} \frac{\eta(B_r(x))}{\pi r^2} = -\frac{1}{2},$$

and hence

$$\tilde{\Theta}^2(\mu - \cos \theta \eta, x) = 1 + \cos \theta < 1,$$

implying that the function $\cdot \mapsto \tilde{\Theta}(\mu - \cos \theta \eta, \cdot)$ is not upper semi-continuous at x .




FIGURE 1. Example.

3.2. Willmore Energy in the Half-Space. The Willmore energy $\mathcal{W}(F)$ of a smooth immersed compact connected orientable surface $F : \Sigma \rightarrow \mathbb{R}^3$ with boundary $\partial\Sigma$, which may have several connected components, is usually proposed by

$$\mathcal{W}_0(F) := \frac{1}{4} \int_{\Sigma} |\mathbf{H}_{\Sigma}|^2 d\mu_g + \int_{\partial\Sigma} \kappa_g, \quad (3.2)$$

where κ_g denotes the geodesic curvature of $\partial\Sigma$ as a submanifold of Σ . The reason for proposing (3.2) is to keep the peculiarity of the Willmore functional—the invariance under the conformal transformation of the ambient space. Thanks to the well-known Gauß-Bonnet theorem, the Willmore energy may be rewritten as

$$\mathcal{W}_0(F) = \frac{1}{4} \int_{\Sigma} |\mathring{A}|^2 + 2\pi\chi(\Sigma), \quad (3.3)$$

where \mathring{A} is the traceless second fundamental form and $\chi(\Sigma)$ is the Euler characteristic, it is clear that the first term in (3.3) is conformal invariant, while the second term is topological invariant. See for example [23].

The point in the above argument is that, in order to keep the conformal invariance, one may indeed add or subtract a topological quantity to the Willmore energy functional. This insight motivates us to drop the term $\int_{\Sigma} \kappa_g$ when considering the Willmore energy of the surface Σ that is capillary immersed into $\overline{\mathbb{R}_+^3}$, since this term is just a multiple of some topological quantity. Indeed, for any θ -capillary immersion $F : \Sigma \rightarrow \overline{\mathbb{R}_+^3}$, it holds locally

$$\kappa_g = \nabla_{\tau} \mu_{\Sigma} \cdot \tau = \nabla_{\tau} (\cos \theta \bar{\nu} - \sin \theta E_3) \cdot \tau = \cos \theta \tilde{\kappa}_g,$$

where $\tilde{\kappa}_g$ is the geodesic curvature of $F(\partial\Sigma)$ as an immersion in $\partial\mathbb{R}_+^3$, using the Gauß-Bonnet theorem, one sees

$$\int_{\partial\Sigma} \kappa_g = \cos \theta \int_{\partial\Sigma} \tilde{\kappa}_g = 2\pi \cos \theta \operatorname{ind}(F(\partial\Sigma)),$$

where $\operatorname{ind}(F(\partial\Sigma))$ is rotation index of immersed plane curve $F(\partial\Sigma)$, which is again a topological invariant.

Therefore, in this paper, we regard the Willmore functional for capillary hypersurfaces in \mathbb{R}_+^{n+1} as

$$\begin{aligned} \mathcal{W}(F) &= \mathcal{W}_0(F) - 2\pi \cos \theta \operatorname{ind}(F(\partial\Sigma)) \\ &= \frac{1}{4} \int_{\Sigma} |\mathbf{H}_{\Sigma}|^2 d\mu_g. \end{aligned} \quad (3.4)$$

$\mathcal{W}(F)$ is again a conformal invariant with respect to conformal diffeomorphisms of \mathbb{R}^3 .

It matches the Willmore functional for capillary hypersurfaces in the unit ball as well, which will be discussed in the next subsection.

We may rewrite the Willmore energy for the θ -capillary immersion as:

$$\mathcal{W}(F) = \frac{1}{4} \int_{\Sigma} |\mathbf{H}_{\Sigma}|^2 d\mu_g = \frac{1}{4} \int |\tilde{\mathbf{H}}|^2 d\mu.$$

The first part of Theorem 1.3 amounts to be a direct application of the monotonicity formula we obtained above.

Proof of Theorem 1.3, (1)-(3). Let us consider the point $x_0 \in \partial\Sigma \subset \partial\mathbb{R}_+^3$. In this case, $\tilde{x}_0 = x_0$ and hence $B_r(x_0) = \hat{B}_r(x_0)$. It follows that

$$\lim_{r \searrow 0^+} \frac{\eta(B_r(x_0)) + \eta(\hat{B}_r(x_0))}{\pi r^2} = N(x_0),$$

for some positive integer $N(x_0) = \mathcal{H}^0(F^{-1}(x_0))$, since $F : \partial\Sigma \rightarrow \partial\mathbb{R}_+^3$ is an immersion. Thus we find

$$\tilde{\Theta}^2(\mu - \cos \theta \eta, x_0) = 2\Theta^2(\mu, x_0) - \cos \theta \cdot N(x_0). \quad (3.5)$$

Moreover, since μ is supported on the upper half-space, $\Theta^2(\mu, x_0) = \frac{N(x_0)}{2}$. Hence

$$\tilde{\Theta}^2(\mu - \cos \theta \eta, x_0) = (1 - \cos \theta)N(x_0). \quad (3.6)$$

By virtue of the above observation, we infer easily from the monotonicity identity (2.6) that

$$\mathcal{W}(F) = 4 \int_{\mathbb{R}^3} \left| \frac{1}{4} \vec{\mathbf{H}} + \frac{(x - x_0)^\perp}{|x - x_0|^2} \right|^2 d\mu + 4\pi\Theta^2(\mu, x_0) - 2 \cos \theta \pi N(x_0). \quad (3.7)$$

Thus, we obtain the Li-Yau-type inequality:

$$\mathcal{W}(F) \geq 4(1 - \cos \theta)\pi\Theta^2(\mu, x_0) = 2(1 - \cos \theta)\pi N(x_0),$$

which implies easily that

$$\mathcal{W}(F) \geq 2(1 - \cos \theta)\pi,$$

the sharp estimate on the lower bound of Willmore energy.

If $\mathcal{W}(F) < 4(1 - \cos \theta)\pi$, for any $x_0 \in \partial\Sigma$, the Li-Yau-type inequality (1.3) implies that $N(x_0) = 1$, and hence $F : \partial\Sigma \rightarrow \partial\mathbb{R}_+^3$ must be an embedding.

Let us now consider the point $y_0 \in \Sigma \setminus \partial\Sigma$. Observe that

$$\tilde{\Theta}^2(\mu - \cos \theta \eta, y_0) = \Theta^2(\mu, y_0).$$

Letting $a = y_0$ in (2.5) and recalling the definition of $\mathcal{W}(F)$, since $\theta \in [\frac{\pi}{2}, \pi)$, we obtain readily

$$2\pi\Theta^2(\mu, y_0) \leq \mathcal{W}(F),$$

and hence $\Theta^2(\mu, y_0) = 1$ if $\mathcal{W}(F) < 4\pi$, in which case $F : \Sigma \rightarrow \overline{\mathbb{R}_+^3}$ must be an embedding. This completes the proof. \square

Remark 3.3. When $F : \Sigma \rightarrow \overline{\mathbb{R}_+^3}$ is an embedding, one can infer easily from (3.5) that

$$\tilde{\Theta}^2(\mu - \cos \theta \eta, x_0) = 1 - \cos \theta,$$

for $x_0 \in \partial\Sigma$. Therefore, it may be convenient to define

$$\frac{\tilde{\Theta}^2(\mu - \cos \theta \eta, x_0)}{1 - \cos \theta}$$

as the *capillary density* of Σ at a boundary point $x_0 \in \partial\Sigma$.

A further investigation gives the following characterization of the situation when $\mathcal{W}(F)$ attains its minima, which also reveals the sharpness of (1.4).

Theorem 3.4. *Given $\theta \in (0, \pi)$, let $F : \Sigma \rightarrow \overline{\mathbb{R}_+^3}$ be a capillary immersion. If $\mathcal{W}(F) = 2\pi(1 - \cos \theta)$, then $F(\Sigma)$ must be a θ -cap in $\overline{\mathbb{R}_+^3}$, i.e., a part of the sphere intersecting $\partial\mathbb{R}_+^{n+1}$ at the angle θ .*

Proof. We begin by noticing that $F : \partial\Sigma \rightarrow \partial\mathbb{R}_+^3$ is an embedding thanks to Theorem 1.3. In particular we learn from (3.7) that

$$\frac{1}{4} \vec{\mathbf{H}}(x) + \frac{(x - x_0)^\perp}{|x - x_0|^2} = 0 \quad (3.8)$$

for μ -a.e. $x \in F(\Sigma)$ and any $x_0 \in F(\partial\Sigma)$. It is not difficult to observe that there must be some $y \in F(\Sigma \setminus \partial\Sigma)$ such that $\vec{\mathbf{H}}(y) \neq 0$, otherwise for μ -a.e. $x \in F(\Sigma)$, one has (up to a translation, we assume that $x_0 = 0$)

$$x \cdot \nu(x) = 0,$$

implying that the enclosing region of $F(\Sigma)$ with $\partial\mathbb{R}_+^3$ is a cone (see e.g. [21, Proposition 28.8]), which is not possible. Consequently, by letting $x = y$ in (3.8), we obtain for any $x_0 \in F(\partial\Sigma)$ that

$$\left| x_0 - \left(y - \frac{2\nu(y)}{|\vec{\mathbf{H}}(y)|} \right) \right|^2 = \frac{4}{|\vec{\mathbf{H}}(y)|^2},$$

which shows that $F(\partial\Sigma)$ is indeed a 1-dimensional sphere on $\partial\mathbb{R}_+^3$, say \mathfrak{s} .

To proceed, we consider the unique spherical cap in $\overline{\mathbb{R}_-^3}$, say $C_{\pi-\theta}$, which intersects $\partial\mathbb{R}_+^3$ along \mathfrak{s} with the constant angle $(\pi - \theta)$. Let $V_{C_{\pi-\theta}}$ denote the naturally induced varifold of $C_{\pi-\theta}$, with weight measure denoted by $\mu_{C_{\pi-\theta}}$, which is exactly given by $\mu_{C_{\pi-\theta}} = \mathcal{H}^2 \llcorner C_{\pi-\theta}$. Write $\tilde{\mu} = \mu + \mu_{C_{\pi-\theta}}$, $\tilde{\Sigma} = \text{spt}(\tilde{\mu}) = F(\Sigma) \cup C_{\pi-\theta}$.

Since F is a θ -capillary immersion with $F : \partial\Sigma \rightarrow \partial\mathbb{R}_+^3$ an embedding, and because of the construction of $C_{\pi-\theta}$, we see that the integral varifold $\tilde{\mu}$ has the following variational structure: for any $X \in C_c^1(\mathbb{R}^3; \mathbb{R}^3)$,

$$\int \text{div}_{\tilde{\Sigma}} X d\tilde{\mu} = - \int_{\tilde{\Sigma}} X \cdot \vec{\mathbf{H}} d\tilde{\mu},$$

where $\vec{\mathbf{H}}$ is the generalized mean curvature vector of $\tilde{\Sigma}$, which is square integrable on $\tilde{\Sigma}$.

Moreover, a direct computation shows that for the spherical cap $C_{\pi-\theta}$,

$$\mathcal{W}(C_{\pi-\theta}) = \frac{1}{4} \int_{C_{\pi-\theta}} |\vec{\mathbf{H}}|^2 d\mathcal{H}^2 = 2(1 - \cos(\pi - \theta))\pi = 2(1 + \cos \theta)\pi,$$

which, together with $\mathcal{W}(F) = 2(1 - \cos \theta)\pi$, yields

$$\mathcal{W}(\tilde{\mu}) = \frac{1}{4} \int |\vec{\mathbf{H}}|^2 d\tilde{\mu} = 4\pi.$$

Here $\mathcal{W}(\tilde{\mu})$ is the Willmore energy of $\tilde{\mu}$ in \mathbb{R}^3 , see [17, Appendix A]. It is worth noting that a simple modification of [27, Proposition 4.3] in conjunction with [18, Proposition 2.1.1] shows that for any compactly supported integral 2-varifold with square integrable generalized mean curvature, if its Willmore energy is 4π , then its support is exactly a closed round 2-sphere. Therefore we conclude that $\tilde{\Sigma}$ must be a closed round 2-sphere, and hence $F(\Sigma)$ is a θ -capillary spherical cap as desired. \square

The last part of the assertion in Theorem 1.3 is simply given by Theorem 3.4.

3.3. Willmore Energy in the Unit Ball. Due to the conformal invariance of the Willmore functional, the results above can be transferred directly to hold for capillary surfaces in the unit ball.

It is well-known that \mathcal{W}_0 , and hence \mathcal{W} defined in (3.4), are invariant under conformal transformations. Therefore the Willmore functional for capillary surfaces in the unit ball is the same as \mathcal{W} , i.e.,

$$\mathcal{W}(F) := \frac{1}{4} \int_{\Sigma} |\vec{\mathbf{H}}|^2 d\mu_g + \int_{\partial\Sigma} \kappa_g - 2\pi \cos \theta \text{ind}(F(\partial\Sigma)).$$

Here $\text{ind}(F(\partial\Sigma))$ is the rotation index of the immersed plane curve $\phi \circ F(\partial\Sigma)$ where $\phi : \mathbb{S}^3 \setminus \{p\} \rightarrow \mathbb{R}^3$ is the stereographic projection for some $p \notin \partial\Sigma$.

Now in this case, the geodesic curvature is given by

$$\kappa_g = \nabla_\tau \mu_\Sigma \cdot \tau = \nabla_\tau (\cos \theta \bar{\nu} + \sin \theta \bar{N}(x)) \cdot \tau = \cos \theta \tilde{\kappa}_g + \sin \theta,$$

where $\bar{N}(x) = x$ is the outer unit normal of \mathbb{B}^3 and $\tilde{\kappa}_g$ is the geodesic curvature of $F : \partial\Sigma \rightarrow \mathbb{S}^2$. By the Gauß-Bonnet theorem, we have

$$2\pi \operatorname{ind}(F(\partial\Sigma)) = \int_{\partial\Sigma} \tilde{\kappa}_g + |T|,$$

where $|T|$ is the oriented area of T . Altogether implying that

$$\mathcal{W}(F) = \frac{1}{4} \int_\Sigma |\bar{\mathbf{H}}|^2 d\mu_g + \sin \theta |\partial\Sigma| - \cos \theta |T|.$$

Corollary 3.5. *Given $\theta \in (0, \pi)$, let $F : \Sigma \rightarrow \overline{\mathbb{B}^3}$ be a θ -capillary immersion. Then, there holds*

$$\mathcal{W}(F) \geq 2(1 - \cos \theta)\pi.$$

Equality holds if and only if $F(\Sigma)$ is a spherical cap or a totally geodesic disk in $\overline{\mathbb{B}^3}$.

Proof. As explained above, the inequality holds thanks to Theorem 1.3 and it suffices to consider the characterization of equality case. Observe that in the half-space case, equality is achieved if and only if the capillary surfaces are spherical. Since umbilicity preserves under conformal transformations, we readily see that in the half-ball case, $F(\Sigma)$ should be either a spherical cap or a totally geodesic disk. \square

If $\theta = \frac{\pi}{2}$, Corollary 3.5 was proved in [27]. Now we show how it simply implies Theorem 1.4.

Proof of Theorem 1.4. From Corollary 3.5, we find

$$\sin \theta |\partial\Sigma| - \cos \theta |T| \geq 2(1 - \cos \theta)\pi.$$

From $\operatorname{div}_\Sigma x = 2$ in Σ we have

$$\begin{aligned} 2|\Sigma| &= \int_{\partial\Sigma} \langle x, \mu_\Sigma \rangle = \int_{\partial\Sigma} \langle \cos \theta \bar{\nu} + \sin \theta \bar{N}(x), x \rangle \\ &= \sin \theta |\partial\Sigma|. \end{aligned}$$

(1.5) follows clearly. \square

4. MONOTONICITY IDENTITIES IN THE UNIT BALL

4.1. Set-ups. Let $\mathbb{B}^3 \subset \mathbb{R}^3$ be the Euclidean unit ball centered at the origin, $\mathbb{S}^2 := \partial\mathbb{B}^3$ denotes the corresponding unit sphere.

In this section we adopt the following notations: let $\xi : \mathbb{R}^3 \setminus \{0\} \rightarrow \mathbb{R}^3$ denote the spherical inversion with respect to \mathbb{S}^2 . Fix $x_0 \in \mathbb{R}^3$ and $r > 0$, $B_r(x_0)$ denotes the open ball of radius r centered at x_0 , and

$$\hat{B}_r(x_0) = B_{\frac{r}{|x_0|}}(\xi(x_0))$$

denotes the ball of radius $\frac{r}{|x_0|}$ centered at $\xi(x_0)$. A direct computation then shows that: for $x \in \mathbb{S}^2$,

$$|x_0||x - \xi(x_0)| = |x - x_0|,$$

in other words, $B_r(x_0) \cap \mathbb{S}^2 = \hat{B}_r(x_0) \cap \mathbb{S}^2$.

Similar with Definition 2.1, we consider the following weak formulation of capillary surfaces in the unit ball.

Definition 4.1. Given $\theta \in (0, \pi)$, let V be a rectifiable 2-varifold supported on $\overline{\mathbb{B}^3}$ with its weight measure denoted by μ , let W be a rectifiable 2-varifold supported on \mathbb{S}^2 with its weight measure denoted by η .

(V, W) is said to *satisfy the contact angle condition* θ if there exists a μ -measurable vector field $\vec{\mathbf{H}} \in \mathcal{L}^1(\mathbb{R}^3, \mu)$ with $\vec{\mathbf{H}}(x) \in T_x \mathbb{S}^2$ for μ -a.e. $x \in \mathbb{S}^2$, such that for every $X \in C_c^1(\mathbb{R}^3; \mathbb{R}^3)$ with X tangent to \mathbb{S}^2 , it holds that

$$\int_{\mathbb{R}^3} \operatorname{div}_{\Sigma} X d\mu - \cos \theta \int_{\mathbb{S}^2} \operatorname{div}_{\mathbb{S}^2} X d\eta = - \int_{\mathbb{B}^3} \vec{\mathbf{H}} \cdot X d\mu, \quad (4.1)$$

where $\Sigma := \operatorname{spt}(\mu)$ is countably 2-rectifiable.

An important proposition for the pairs of varifolds satisfying contact angle condition is that they have bounded first variation and satisfy the following first variation formula (see [6, Proposition 3.1]).

Proposition 4.2. *Given $\theta \in [\frac{\pi}{2}, \pi)$, let (V, W) be as in Definition 2.1. Then $V - \cos \theta W$ has bounded first variation. More precisely, there exists a positive Radon measure σ_V on \mathbb{S}^2 such that*

$$\begin{aligned} \int_{\mathbb{R}^3} \operatorname{div}_{\Sigma} X d\mu - \cos \theta \int_{\mathbb{S}^2} \operatorname{div}_{\mathbb{S}^2} X d\eta &= - \int_{\mathbb{B}^3} \vec{\mathbf{H}} \cdot X d\mu \\ &+ 2 \int_{\mathbb{S}^2} X(x) \cdot x d(\mu - \cos \theta \eta) + \int_{\mathbb{S}^2} X(x) \cdot x d\sigma_V \end{aligned} \quad (4.2)$$

for every $X \in C_c^1(\mathbb{R}^3; \mathbb{R}^3)$.

If Σ is in fact a smooth capillary surface embedded in \mathbb{B}^3 , then we may use a classical computation to see that the above formula is true with $\sigma_V = \sin \theta \mathcal{H}^2 \llcorner \partial \Sigma$. This motivates us to define $\gamma := \frac{1}{\sin \theta} \sigma_V$ as the generalized boundary measure of V and $\operatorname{spt} \gamma$ as the generalized boundary of V (which are different from the ones obtained from Lebesgue's decomposition theorem).

In all follows, we consider only those V that are integral 2-varifolds with $\mu(\mathbb{S}^2) = 0$ and $\vec{\mathbf{H}} \in \mathcal{L}^2(\mathbb{R}^3, \mu)$, it follows that (2.2) holds. Moreover, we may rewrite (4.2) as

$$\begin{aligned} \int_{\mathbb{R}^3} \operatorname{div}_{\Sigma} X d\mu - \cos \theta \int_{\mathbb{S}^2} \operatorname{div}_{\mathbb{S}^2} X d\eta &= - \int_{\mathbb{B}^3} \vec{\mathbf{H}} \cdot X d\mu \\ &- 2 \cos \theta \int_{\mathbb{S}^2} X(x) \cdot x d\eta + \sin \theta \int_{\mathbb{S}^2} X(x) \cdot x d\gamma \end{aligned} \quad (4.3)$$

for every $X \in C_c^1(\mathbb{R}^3; \mathbb{R}^3)$. An immediate consequence is that we may use a compactly supported vector field which coincides with the position vector field in a neighborhood of \mathbb{B}^3 to test (4.3) and obtain

$$2\mu(\mathbb{R}^3) = - \int_{\mathbb{B}^3} \vec{\mathbf{H}} \cdot x d\mu + \sin \theta \gamma(\mathbb{S}^2). \quad (4.4)$$

4.2. Simon-Type Monotonicity Formulas. Let us first recall that in [27], a Simon-type monotonicity formula is proved for integral 2-varifold μ with $\mu(\mathbb{S}^2) = 0$, that has free boundary in $\overline{\mathbb{B}^3}$. Precisely, it is proved that for $x_0 \neq 0$,

$$\begin{aligned} & \frac{1}{\pi} \int_{B_\rho(x_0) \setminus B_\sigma(x_0)} \left| \frac{1}{4} \vec{\mathbf{H}} + \frac{(x - x_0)^\perp}{|x - x_0|^2} \right|^2 d\mu + \frac{1}{\pi} \int_{\hat{B}_\rho(x_0) \setminus \hat{B}_\sigma(x_0)} \left| \frac{1}{4} \vec{\mathbf{H}} + \frac{(x - \xi(x_0))^\perp}{|x - \xi(x_0)|^2} \right|^2 d\mu \\ &= (g_{x_0}(\rho) + \hat{g}_{x_0}(\rho)) - (g_{x_0}(\sigma) + \hat{g}_{x_0}(\sigma)), \end{aligned} \quad (4.5)$$

where

$$\begin{aligned} g_{x_0}(r) &:= \frac{\mu(B_r(x_0))}{\pi r^2} + \frac{1}{16\pi} \int_{B_r(x_0)} |\vec{\mathbf{H}}|^2 d\mu + \frac{1}{2\pi r^2} \int_{B_r(x_0)} \vec{\mathbf{H}} \cdot (x - x_0) d\mu, \\ \hat{g}_{x_0}(r) &:= g_{\xi(x_0)}\left(\frac{r}{|x_0|}\right) - \frac{|x_0|^2}{\pi r^2} \int_{\hat{B}_r(x_0)} \left(|x - \xi(x_0)|^2 + (x - \xi(x_0))^T \cdot x \right) d\mu \\ &\quad - \frac{|x_0|^2}{2\pi r^2} \int_{\hat{B}_r(x_0)} \vec{\mathbf{H}} \cdot (|x - \xi(x_0)|^2 x) d\mu + \frac{1}{2\pi} \int_{\hat{B}_r(x_0)} \vec{\mathbf{H}} \cdot x d\mu + \frac{\mu(\hat{B}_r(x_0))}{\pi}, \end{aligned}$$

and $(\cdot)^T$ denotes the orthogonal projection of a vector onto the approximate tangent space $T_x \Sigma$. For $x_0 = 0$, he obtained

$$\frac{1}{\pi} \int_{B_\rho(0) \setminus B_\sigma(0)} \left| \frac{1}{4} \vec{\mathbf{H}} + \frac{1}{\pi} \frac{x^\perp}{|x|^2} \right|^2 d\mu = (g_0(\rho) + \hat{g}_0(\rho)) - (g_0(\sigma) + \hat{g}_0(\sigma)), \quad (4.6)$$

where

$$\begin{aligned} g_0(r) &:= \frac{\mu(B_r(0))}{\pi r^2} + \frac{1}{16\pi} \int_{B_r(0)} |\vec{\mathbf{H}}|^2 d\mu + \frac{1}{2\pi r^2} \int_{B_r(0)} \vec{\mathbf{H}} \cdot x d\mu, \\ \hat{g}_0(r) &:= -\frac{\min(r^{-2}, 1)}{2\pi} \left(2\mu(\mathbb{R}^3) + \int_{\mathbb{R}^3} \vec{\mathbf{H}} \cdot x d\mu \right). \end{aligned}$$

Volkman's proof is based on using a properly chosen vector field to test the first variation of a free boundary varifold. This particular choice can be dated back to [3]. We follow to use this vector field to test (4.1) and obtain the following result.

Theorem 4.3. *Given $\theta \in [\frac{\pi}{2}, \pi)$, let (V, W) be as in Definition 4.1. Suppose in addition that V is an integral 2-varifold with $\mu(\mathbb{S}^2) = 0$, then it holds that: for $x_0 \neq 0$, and for every $0 < \sigma < \rho < \infty$,*

$$\begin{aligned} & \frac{1}{\pi} \int_{B_\rho(x_0) \setminus B_\sigma(x_0)} \left| \frac{1}{4} \vec{\mathbf{H}} + \frac{(x - x_0)^\perp}{|x - x_0|^2} \right|^2 d\mu + \frac{1}{\pi} \int_{\hat{B}_\rho(x_0) \setminus \hat{B}_\sigma(x_0)} \left| \frac{1}{4} \vec{\mathbf{H}} + \frac{(x - \xi(x_0))^\perp}{|x - \xi(x_0)|^2} \right|^2 d\mu \\ & - \frac{\cos \theta}{\pi} \int_{B_\rho(x_0) \setminus B_\sigma(x_0)} \left(\frac{x - x_0}{|x - x_0|^2} \cdot x \right)^2 d\eta - \frac{\cos \theta}{\pi} \int_{\hat{B}_\rho(x_0) \setminus \hat{B}_\sigma(x_0)} \left(\frac{x - \xi(x_0)}{|x - \xi(x_0)|^2} \cdot x \right)^2 d\eta \\ &= (g_{x_0, \theta}(\rho) + \hat{g}_{x_0, \theta}(\rho)) - (g_{x_0, \theta}(\sigma) + \hat{g}_{x_0, \theta}(\sigma)), \end{aligned} \quad (4.7)$$

where

$$\begin{aligned} g_{x_0, \theta}(r) &:= g_{x_0}(r) - \frac{\cos \theta \eta(B_r(x_0))}{\pi r^2}, \\ \hat{g}_{x, \theta}(r) &:= \hat{g}_{x_0}(r) - \frac{\cos \theta |x_0|^2 \eta(\hat{B}_r(x_0))}{\pi r^2} \\ &\quad + \frac{\cos \theta |x_0|^2}{\pi r^2} \int_{\hat{B}_r(x_0)} |x - \xi(x_0)|^2 d\eta - \frac{\cos \theta \eta(\hat{B}_r(x_0))}{\pi}. \end{aligned}$$

And for $x_0 = 0$,

$$\frac{1}{\pi} \int_{B_\rho(0) \setminus B_\sigma(0)} \left| \frac{1}{4} \vec{\mathbf{H}} + \frac{1}{\pi} \frac{x^\perp}{|x|^2} \right|^2 d\mu = (g_0(\rho) + \hat{g}_{0, \theta}(\rho)) - (g_0(\sigma) + \hat{g}_{0, \theta}(\sigma)), \quad (4.8)$$

where

$$\hat{g}_{0, \theta}(r) := - \frac{\min(r^{-2}, 1)}{2\pi} \sin \theta \gamma(\mathbb{S}^2).$$

Proof. For every $0 < \sigma < \rho < \infty$, we continue to use the Lipschitz cut-off function l defined in the proof of Theorem 1.1. For $x_0 \in \mathbb{R}^3$, let

$$X_1(x) := l(|x - x_0|)(x - x_0).$$

Case 1. $x_0 \neq 0$.

Define

$$\begin{aligned} X_2(x) &:= \left(\left(\frac{1}{|x - \xi(x_0)| \frac{\sigma}{|x_0|}} \right)^2 - \frac{|x_0|^2}{\rho^2} \right)_+ (x - \xi(x_0)) \\ &\quad + \left(\frac{\min(|x_0| |x - \xi(x_0)|, \rho)^2}{\rho^2} - \frac{\min(|x_0| |x - \xi(x_0)|, \sigma)^2}{\sigma^2} \right) x, \end{aligned}$$

and

$$X := X_1 + X_2.$$

As verified in [27, (7)], X is an admissible vector field to test (4.1) for a.e. $0 < \sigma < \rho < \infty$. Moreover, the terms $\int_{\mathbb{R}^3} \operatorname{div}_\Sigma X d\mu$ and $-\int_{\mathbb{R}^3} \vec{\mathbf{H}} \cdot X d\mu$ are explicitly computed. That is, if $\theta = \frac{\pi}{2}$, testing (4.1) with such X , one gets exactly (4.5). Therefore for the case $\theta \in (\frac{\pi}{2}, \pi)$, it suffices to compute

$$-\cos \theta \int_{\mathbb{S}^2} \operatorname{div}_{\mathbb{S}^2} X d\eta.$$

To this end, we consider the following decomposition of \mathbb{R}^3 :

$$\begin{aligned} \mathcal{F}_1 &:= \{B_\sigma(x_0), B_\rho(x_0) \setminus B_\sigma(x_0), \mathbb{R}^3 \setminus B_\rho(x_0)\}, \\ \mathcal{F}_2 &:= \{\hat{B}_\sigma(x_0), \hat{B}_\rho(x_0) \setminus \hat{B}_\sigma(x_0), \mathbb{R}^3 \setminus \hat{B}_\rho(x_0)\}, \end{aligned}$$

and we shall compute

$$-\cos \theta \int_A \operatorname{div}_{\mathbb{S}^2} X_i d\eta$$

for all sets $A \in \mathcal{F}_i$, $i=1,2$, separately.

For X_1 , a direct computation shows that

$$\nabla X_1 = \begin{cases} \left(\frac{1}{\sigma^2} - \frac{1}{\rho^2} \right) \text{Id}, & 0 \leq |x - x_0| \leq \sigma, \\ \left(\frac{1}{|x-x_0|^2} - \frac{1}{\rho^2} \right) \text{Id} - 2 \frac{x-x_0}{|x-x_0|^2} \otimes \frac{x-x_0}{|x-x_0|^2}, & \sigma < |x - x_0| \leq \rho \\ 0, & \rho < |x - x_0|, \end{cases}$$

so that on S^2 ,

$$\begin{aligned} \text{div}_{\mathbb{S}^2} X_1(x) &= \text{div} X_1(x) - \nabla X_1[x] \cdot x \\ &= \begin{cases} \frac{2}{\sigma^2} - \frac{2}{\rho^2}, & 0 \leq |x - x_0| \leq \sigma, \\ -\frac{2}{\rho^2} + 2 \left(\frac{x-x_0}{|x-x_0|^2} \cdot x \right)^2, & \sigma < |x - x_0| \leq \rho, \\ 0 & \rho < |x - x_0|, \end{cases} \end{aligned}$$

and it is easy to deduce that

$$\begin{aligned} & -\cos \theta \int_{\mathbb{S}^2} \text{div}_{\mathbb{S}^2} X_1 d\eta \\ &= -\cos \theta \left(\frac{2}{\sigma^2} \eta(B_\sigma(x_0)) - \frac{2}{\rho^2} \eta(B_\rho(x_0)) + 2 \int_{B_\rho(x_0) \setminus B_\sigma(x_0)} \left(\frac{x-x_0}{|x-x_0|^2} \cdot x \right)^2 d\eta \right). \end{aligned} \quad (4.9)$$

For X_2 , we may compute directly to find:

As $0 \leq |x - \xi(x_0)| \leq \frac{\sigma}{|x_0|}$,

$$X_2 = |x_0|^2 \left(\frac{1}{\sigma^2} - \frac{1}{\rho^2} \right) (x - \xi(x_0)) - |x_0|^2 |x - \xi(x_0)|^2 \left(\frac{1}{\sigma^2} - \frac{1}{\rho^2} \right) x;$$

as $\frac{\sigma}{|x_0|} < |x - \xi(x_0)| \leq \frac{\rho}{|x_0|}$,

$$X_2 = \left(\frac{1}{|x - \xi(x_0)|^2} - \frac{|x_0|^2}{\rho^2} \right) (x - \xi(x_0)) - x + \frac{|x_0|^2 |x - \xi(x_0)|^2}{\rho^2} x;$$

and $X_2 \equiv 0$ as $\frac{\rho}{|x_0|} < |x - \xi(x_0)|$.

Thus ∇X_2 vanishes on $\mathbb{R}^3 \setminus \hat{B}_\rho(x_0)$; as $0 < |x - \xi(x_0)| \leq \frac{\sigma}{|x_0|}$,

$$\nabla X_2 = \left(\frac{1}{\sigma^2} - \frac{1}{\rho^2} \right) |x_0|^2 \left\{ (1 - |x - \xi(x_0)|^2) \text{Id} - 2x \otimes (x - \xi(x_0)) \right\},$$

and as $\frac{\sigma}{|x_0|} \leq |x - \xi(x_0)| \leq \frac{\rho}{|x_0|}$,

$$\begin{aligned} \nabla X_2 &= \left(\frac{1}{|x - \xi(x_0)|^2} - \frac{|x_0|^2}{\rho^2} \right) \text{Id} - 2 \frac{x - \xi(x_0)}{|x - \xi(x_0)|^2} \otimes \frac{x - \xi(x_0)}{|x - \xi(x_0)|^2} \\ &\quad - \left(1 - \frac{|x_0|^2 |x - \xi(x_0)|^2}{\rho^2} \right) \text{Id} + 2 \frac{|x_0|^2}{\rho^2} x \otimes (x - \xi(x_0)). \end{aligned}$$

Therefore it is not difficult to compute that: as $0 \leq |x - \xi(x_0)| \leq \frac{\sigma}{|x_0|}$,

$$\text{div}_{\mathbb{S}^2} X_2(x) = 2|x_0|^2 \left(1 - |x - \xi(x_0)|^2 \right) \left(\frac{1}{\sigma^2} - \frac{1}{\rho^2} \right),$$

and as $\frac{\sigma}{|x_0|} < |x - \xi(x_0)| < \frac{\rho}{|x_0|}$,

$$\operatorname{div}_{\mathbb{S}^2} X_2(x) = -\frac{2|x_0|^2(1 - |x - \xi(x_0)|^2)}{\rho^2} - 2 + 2 \left(\frac{x - \xi(x_0)}{|x - \xi(x_0)|^2} \cdot x \right)^2.$$

It follows that

$$\begin{aligned} & -\cos \theta \int_{\mathbb{S}^2} \operatorname{div}_{\mathbb{S}^2} X_2 d\eta \\ &= -\cos \theta \left[\left(\frac{2|x_0|^2}{\sigma^2} \eta(\hat{B}_\sigma(x_0)) - \frac{2|x_0|^2}{\sigma^2} \int_{\hat{B}_\sigma(x_0)} |x - \xi(x_0)|^2 d\eta + 2\eta(\hat{B}_\sigma(x_0)) \right) \right. \\ & \quad \left. - \left(\frac{2|x_0|^2}{\rho^2} \eta(\hat{B}_\rho(x_0)) - \frac{2|x_0|^2}{\rho^2} \int_{\hat{B}_\rho(x_0)} |x - \xi(x_0)|^2 d\eta + 2\eta(\hat{B}_\rho(x_0)) \right) \right. \\ & \quad \left. + 2 \int_{\hat{B}_\rho(x_0) \setminus \hat{B}_\sigma(x_0)} \left(\frac{x - \xi(x_0)}{|x - \xi(x_0)|^2} \cdot x \right)^2 d\eta \right]. \end{aligned} \quad (4.10)$$

Recall that if $\theta = \frac{\pi}{2}$, testing (4.1) with such X , one gets exactly (4.5). Now for $\theta \in (\frac{\pi}{2}, \pi)$, taking (4.9) and (4.10) into consideration, we thus obtain (4.7) for a.e. σ and ρ , and an approximation argument shows that this indeed holds for every σ and ρ .

Case 2. $x_0 = 0$.

We continue to use X_1 defined in **Case 1** (with $x_0 = 0$) and define

$$\begin{aligned} X_2(x) &:= \left(\frac{\min(1, \rho)^2}{\rho^2} - \frac{\min(1, \sigma)^2}{\sigma^2} \right) x, \\ X &:= X_1 + X_2. \end{aligned}$$

A direct computation shows that X is an admissible vector field to test (4.1). Indeed, we have $X = X_1 + X_2 \equiv 0$ on \mathbb{S}^2 for every $0 < \sigma < \rho < \infty$. Thus

$$-\cos \theta \int_{\mathbb{S}^2} \operatorname{div}_{\mathbb{S}^2} X d\eta = 0.$$

Recall that as $\theta = \frac{\pi}{2}$, testing (4.1) with such X one gets exactly (4.6), therefore for $\theta \in (\frac{\pi}{2}, \pi)$, we shall get the same identity. Moreover, thanks to (4.4) and invoking the definition of \hat{g}_0 , we may rearrange this and deduce (4.8) as desired. \square

Proposition 4.4. *Given $\theta \in [\pi/2, \pi)$. For every $x_0 \in \mathbb{R}^3$, the tilde-density*

$$\begin{aligned} & \tilde{\Theta}^2(\mu - \cos \theta \eta, x_0) \\ &:= \begin{cases} \lim_{r \searrow 0} \left(\frac{(\mu - \cos \theta \eta)(B_r(x_0))}{\pi r^2} + \frac{(\mu - \cos \theta \eta)(\hat{B}_r(x_0))}{\pi |x_0|^{-2} r^2} \right), & x_0 \neq 0, \\ \lim_{r \searrow 0} \frac{\mu(B_r(0))}{\pi r^2}, & x_0 = 0, \end{cases} \end{aligned}$$

exists. Moreover, the function $x \mapsto \tilde{\Theta}(\mu - \cos \theta \eta, x)$ is upper semi-continuous in \mathbb{R}^3 .

Proof. We prove for the first assertion:

Case 1. $x_0 \neq 0$.

Define

$$\begin{aligned}
R_{x_0}(r) &:= \frac{1}{2\pi r^2} \int_{B_r(x_0)} \vec{\mathbf{H}} \cdot (x - x_0) d\mu + \frac{|x_0|^2}{2\pi r^2} \int_{\hat{B}_r(x_0)} \vec{\mathbf{H}} \cdot (x - \xi(x_0)) d\mu \\
&\quad - \frac{|x_0|^2}{\pi r^2} \int_{\hat{B}_r(x_0)} \left(|x - \xi(x_0)|^2 + (x - \xi(x_0))^T \cdot x \right) d\mu \\
&\quad - \frac{|x_0|^2}{2\pi r^2} \int_{\hat{B}_r(x_0)} \vec{\mathbf{H}} \cdot (|x - \xi(x_0)|^2 x) d\mu, \\
R_{x_0, \theta}(r) &:= R_{x_0}(r) + \frac{\cos \theta |x_0|^2}{\pi r^2} \int_{\hat{B}_r(x_0)} |x - \xi(x_0)|^2 d\eta - \frac{\cos \theta \eta(\hat{B}_r(x_0))}{\pi} \\
&\quad + \frac{1}{2\pi} \int_{\hat{B}_r(x_0)} \vec{\mathbf{H}} \cdot x d\mu + \frac{\mu(\hat{B}_r(x_0))}{\pi},
\end{aligned}$$

then

$$\begin{aligned}
G_{x_0, \theta}(r) &:= \frac{(\mu - \cos \theta \eta)(B_r(x_0))}{\pi r^2} + \frac{|x_0|^2 (\mu - \cos \theta \eta)(\hat{B}_r(x_0))}{\pi r^2} \\
&\quad + \frac{1}{16\pi} \int_{B_r(x_0)} |\vec{\mathbf{H}}|^2 d\mu + \frac{1}{16\pi} \int_{\hat{B}_r(x_0)} |\vec{\mathbf{H}}|^2 d\mu + R_{x_0, \theta}(r)
\end{aligned}$$

is monotonically non-decreasing thanks to (4.7) and the fact that $\theta \in [\frac{\pi}{2}, \pi)$. Therefore

$$\lim_{r \rightarrow 0^+} G_{x_0, \theta}(r)$$

exists.

Now we estimate with Hölder inequality:

$$\begin{aligned}
|R_{x_0, \theta}(r)| &\leq \left(\frac{\mu(B_r(x_0))}{\pi r^2} \right)^{\frac{1}{2}} \left(\frac{1}{4\pi} \int_{B_r(x_0)} |\vec{\mathbf{H}}|^2 d\mu \right)^{\frac{1}{2}} \\
&\quad + \left(\frac{\mu(\hat{B}_r(x_0))}{\pi |x_0|^{-2} r^2} \right)^{\frac{1}{2}} \left(\frac{1}{4\pi} \int_{\hat{B}_r(x_0)} |\vec{\mathbf{H}}|^2 d\mu \right)^{\frac{1}{2}} + \frac{\mu(\hat{B}_r(x_0))}{\pi} \\
&\quad + \left(\frac{\mu(\hat{B}_r(x_0))}{\pi |x_0|^{-2} r^2} \right)^{\frac{1}{2}} \left(\frac{\mu(\hat{B}_r(x_0))}{\pi} \right)^{\frac{1}{2}} + \left(\frac{\mu(\hat{B}_r(x_0))}{\pi} \right)^{\frac{1}{2}} \left(\frac{1}{4\pi} \int_{\hat{B}_r(x_0)} |\vec{\mathbf{H}}|^2 d\mu \right)^{\frac{1}{2}} \\
&\quad - \frac{2 \cos \theta \eta(\hat{B}_r(x_0))}{\pi} + \frac{1}{2\pi} \int_{\hat{B}_r(x_0)} |\vec{\mathbf{H}}| d\mu + \frac{\mu(\hat{B}_r(x_0))}{\pi},
\end{aligned} \tag{4.11}$$

where we have used the fact that $\text{spt}\mu \subset \overline{\mathbb{B}^3}$ so that $|x| \leq 1$. Moreover, for $\frac{1}{8} < \epsilon < \frac{1}{4}$, Young's inequality gives

$$\begin{aligned}
|R_{x_0, \theta}(r)| &\leq \epsilon \frac{\mu(B_r(x_0))}{\pi r^2} + \frac{1}{16\pi\epsilon} \int_{B_r(x_0)} |\vec{\mathbf{H}}|^2 d\mu \\
&\quad + \epsilon \frac{\mu(\hat{B}_r(x_0))}{\pi|x_0|^{-2}r^2} + \frac{1}{16\pi\epsilon} \int_{\hat{B}_r(x_0)} |\vec{\mathbf{H}}|^2 d\mu \\
&\quad + \epsilon \frac{\mu(\hat{B}_r(x_0))}{\pi|x_0|^{-2}r^2} + \frac{1}{4\epsilon} \frac{\mu(\hat{B}_r(x_0))}{\pi} + \frac{\mu(\hat{B}_r(x_0))}{4\pi} + \frac{1}{4\pi} \int_{\hat{B}_r(x_0)} |\vec{\mathbf{H}}|^2 d\mu \\
&\quad + 2 \frac{(\mu - \cos \theta \eta)(\hat{B}_r(x_0))}{\pi} + \frac{1}{2\pi} \int_{\hat{B}_r(x_0)} |\vec{\mathbf{H}}| d\mu \\
&\leq 2\epsilon \left(\frac{\mu(B_r(x_0))}{\pi r^2} + \frac{\mu(\hat{B}_r(x_0))}{\pi|x_0|^{-2}r^2} \right) + \frac{\epsilon^{-1}}{16\pi} \left(\int_{B_r(x_0) \cup \hat{B}_r(x_0)} |\vec{\mathbf{H}}|^2 d\mu \right) \\
&\quad + \frac{11 + \epsilon^{-1}}{4\pi} (\mu - \cos \theta \eta)(\hat{B}_r(x_0)) + \frac{3}{8\pi} \int_{\hat{B}_r(x_0)} |\vec{\mathbf{H}}|^2 d\mu,
\end{aligned}$$

where we have used again $\theta \in [\frac{\pi}{2}, \pi)$ and the fact that

$$\frac{1}{2\pi} \int_{\hat{B}_r(x_0)} |\vec{\mathbf{H}}| d\mu \leq \frac{\mu(\hat{B}_r(x_0))}{2\pi} + \frac{1}{8\pi} \int_{\hat{B}_r(x_0)} |\vec{\mathbf{H}}|^2 d\mu.$$

By virtue of the monotonicity of $G_{x_0, \theta}(r)$, we obtain for $0 < \sigma < \rho < \infty$

$$\begin{aligned}
&\frac{(\mu - \cos \theta \eta)(B_\sigma(x_0))}{\pi\sigma^2} + \frac{(\mu - \cos \theta \eta)(\hat{B}_\sigma(x_0))}{\pi|x_0|^{-2}\sigma^2} \\
&\leq \frac{(\mu - \cos \theta \eta)(B_\sigma(x_0))}{\pi\sigma^2} + \frac{(\mu - \cos \theta \eta)(\hat{B}_\sigma(x_0))}{\pi|x_0|^{-2}\sigma^2} \\
&\quad + \frac{1}{16\pi} \int_{B_\rho(x_0)} |\vec{\mathbf{H}}|^2 d\mu + \frac{1}{16\pi} \int_{\hat{B}_\rho(x_0)} |\vec{\mathbf{H}}|^2 d\mu + |R_{x_0, \theta}(\rho)| + |R_{x_0, \theta}(\sigma)| \\
&\leq (1 + 2\epsilon) \left(\frac{(\mu - \cos \theta \eta)(B_\rho(x_0))}{\pi\rho^2} + \frac{(\mu - \cos \theta \eta)(\hat{B}_\rho(x_0))}{\pi|x_0|^{-2}\rho^2} \right) \\
&\quad + \frac{1 + 2\epsilon^{-1}}{16\pi} \int_{B_\rho(x_0) \cup \hat{B}_\rho(x_0)} |\vec{\mathbf{H}}|^2 d\mu + \frac{11 + \epsilon^{-1}}{2\pi} (\mu - \cos \theta \eta)(\mathbb{R}^3) \\
&\quad + \frac{3}{4\pi} \int_{\mathbb{R}^3} |\vec{\mathbf{H}}|^2 d\mu + 2\epsilon \left(\frac{(\mu - \cos \theta \eta)(B_\sigma(x_0))}{\pi\sigma^2} + \frac{(\mu - \cos \theta \eta)(\hat{B}_\sigma(x_0))}{\pi|x_0|^{-2}\sigma^2} \right).
\end{aligned}$$

In particular, since $\frac{1}{8} \leq \epsilon \leq \frac{1}{4}$, this yields for a fixed $R > 0$ that

$$\begin{aligned} & \frac{(\mu - \cos \theta \eta)(B_r(x_0))}{\pi r^2} + \frac{(\mu - \cos \theta \eta)(\hat{B}_r(x_0))}{\pi |x_0|^{-2} r^2} \\ & \leq 3 \left(\frac{(\mu - \cos \theta \eta)(B_R(x_0))}{\pi R^2} + \frac{(\mu - \cos \theta \eta)(\hat{B}_R(x_0))}{\pi |x_0|^{-2} R^2} \right) \\ & \quad + \frac{23}{4\pi} \int_{\mathbb{R}^3} |\vec{\mathbf{H}}|^2 d\mu + \frac{19}{\pi} (\mu - \cos \theta \eta)(\mathbb{R}^3) < \infty \end{aligned} \quad (4.12)$$

for every $0 < r < R$.

This, in conjunction with (4.11), yields

$$\lim_{r \rightarrow 0^+} |R_{x_0, \theta}(r)| = 0,$$

and also implies that the tilde-density $\tilde{\Theta}(\mu - \cos \theta \eta, x_0)$ exists. Consequently

$$\begin{aligned} \tilde{\Theta}(\mu - \cos \theta \eta, x_0) & \leq \left(\frac{(\mu - \cos \theta \eta)(B_r(x_0))}{\pi r^2} + \frac{(\mu - \cos \theta \eta)(\hat{B}_r(x_0))}{\pi |x_0|^{-2} r^2} \right) \\ & \quad + \frac{1}{16\pi} \int_{B_r(x_0)} |\vec{\mathbf{H}}|^2 d\mu + \frac{1}{16\pi} \int_{\hat{B}_r(x_0)} |\vec{\mathbf{H}}|^2 d\mu + R_{\theta, x_0}(\rho). \end{aligned}$$

Case 2. $x_0 = 0$.

Define

$$R_{0, \theta}(r) := \frac{1}{2\pi r^2} \int_{B_r(0)} \vec{\mathbf{H}} \cdot x d\mu,$$

then

$$G_{0, \theta}(r) := \frac{\mu(B_r(0))}{\pi r^2} + \frac{1}{16\pi} \int_{B_r(0)} |\vec{\mathbf{H}}|^2 d\mu - \frac{\min(r^{-2}, 1)}{2\pi} \sin \theta \gamma(\mathbb{S}^2) + R_{0, \theta}(r)$$

is monotonically non-decreasing thanks to (4.8), thus

$$\lim_{r \rightarrow 0^+} G_{0, \theta}(r)$$

exists.

Using Hölder inequality and Young's inequality, we obtain for $\frac{1}{4} \leq \epsilon \leq \frac{1}{2}$,

$$\begin{aligned} |R_{0, \theta}(r)| & \leq \left(\frac{\mu(B_r(0))}{\pi r^2} \right)^{\frac{1}{2}} \left(\frac{1}{4\pi} \int_{B_r(0)} |\vec{\mathbf{H}}|^2 d\mu \right)^{\frac{1}{2}} \\ & \leq \epsilon \frac{\mu(B_r(0))}{\pi r^2} + \frac{1}{16\pi \epsilon} \int_{B_r(0)} |\vec{\mathbf{H}}|^2 d\mu. \end{aligned} \quad (4.13)$$

By virtue of the monotonicity of $G_{0, \theta}(r)$, we obtain for $0 < \sigma < \rho \leq 1$

$$\begin{aligned} \frac{\mu(B_\sigma(0))}{\pi \sigma^2} & \leq \frac{\mu(B_\rho(0))}{\pi \rho^2} + \frac{1}{16\pi} \int_{B_\rho(0)} |\vec{\mathbf{H}}|^2 d\mu + |R_{0, \theta}(\rho)| + |R_{0, \theta}(\sigma)| \\ & \leq (1 + \epsilon) \frac{\mu(B_\rho(0))}{\pi \rho^2} + \frac{1 + 2\epsilon^{-1}}{16\pi} \int_{B_\rho(0)} |\vec{\mathbf{H}}|^2 d\mu + \epsilon \frac{\mu(B_\sigma(0))}{\pi \sigma^2}, \end{aligned}$$

where we have used that $\hat{g}_{0,\theta}(r) \equiv -\frac{\sin \theta}{2\pi} \gamma(\mathbb{S}^2)$ for every $0 < r \leq 1$ in the first inequality. In particular, since $\frac{1}{4} \leq \epsilon \leq \frac{1}{2}$ this yields for any $0 < r < 1$:

$$\frac{\mu(B_r(0))}{\pi r^2} \leq 3 \frac{\mu(B_1(0))}{\pi} + \frac{9}{8\pi} \int_{\mathbb{R}^3} |\vec{\mathbf{H}}|^2 d\mu < \infty, \quad (4.14)$$

which, together with (4.13), yields

$$\lim_{r \rightarrow 0^+} |R_{0,\theta}(r)| = 0,$$

and also implies that the tilde-density $\tilde{\Theta}(\mu - \cos \theta \eta, 0)$ exists. Consequently for $0 < r < 1$,

$$\tilde{\Theta}(\mu, 0) \leq \frac{\mu(B_r(0))}{\pi r^2} + \frac{1}{16\pi} \int_{B_r(0)} |\vec{\mathbf{H}}|^2 d\mu + R_{0,\theta}(r).$$

Now we verify the upper semi-continuity by definition, for a sequence of points $x_j \rightarrow x_0$ and for $0 < \rho < \frac{1}{2}$, we have

$$\begin{aligned} & \frac{(\mu - \cos \theta)(\overline{B_\rho(x_0)})}{\pi \rho^2} + \frac{(\mu - \cos \theta)(\overline{\hat{B}_\rho(x_0)})}{\pi |x_0|^{-2} \rho^2} \\ & \geq \limsup_{j \rightarrow \infty} \left(\frac{(\mu - \cos \theta)(\overline{B_\rho(x_j)})}{\pi \rho^2} + \frac{(\mu - \cos \theta)(\overline{\hat{B}_\rho(x_j)})}{\pi |x_j|^{-2} \rho^2} \right) \\ & \geq \limsup_{j \rightarrow \infty} \left(\tilde{\Theta}(\mu - \cos \theta \eta, x_j) - \frac{1}{16\pi} \left(\int_{B_\rho(x_j)} |\vec{\mathbf{H}}|^2 d\mu + \int_{\hat{B}_\rho(x_j)} |\vec{\mathbf{H}}|^2 d\mu \right) - R_{x_j,\theta}(\rho) \right) \\ & \geq \limsup_{j \rightarrow \infty} \tilde{\Theta}(\mu - \cos \theta \eta, x_j) - C \left(\frac{(\mu - \cos \theta \eta)(B_{\frac{1}{2}}(x_0) \cup \hat{B}_{\frac{1}{2}}(x_0))}{\pi (\frac{1}{2})^2} \right. \\ & \quad \left. + \int_{\mathbb{R}^3} |\vec{\mathbf{H}}|^2 d\mu + (\mu - \cos \theta \eta)(\mathbb{R}^3) \right) \cdot \left(\|\vec{\mathbf{H}}\|_{L^2(B_{2\rho}(x_0))} + \left(\frac{\mu(\hat{B}_\rho(x_0))}{\pi} \right)^{\frac{1}{2}} \right) \\ & \quad - 2 \frac{(\mu - \cos \theta \eta)(\hat{B}_\rho(x_0))}{\pi}, \end{aligned}$$

where we have used (4.12), (4.14) for the last inequality, here we interpret $\hat{B}_r(0) = \emptyset$ and $\frac{(\mu - \cos \theta)(\hat{B}_\rho(0))}{\pi |0|^{-2} \rho^2} = 0$. Letting $\rho \rightarrow 0^+$, this gives

$$\tilde{\Theta}(\mu - \cos \theta \eta, x_0) \geq \limsup_{j \rightarrow \infty} \tilde{\Theta}(\mu - \cos \theta \eta, x_j),$$

which completes the proof of the second assertion. \square

Remark 4.5. Since $\text{spt} \mu$ and $\text{spt} \eta$ are compact, we see from the definition that

$$\begin{aligned} \lim_{r \rightarrow \infty} R_{x_0,\theta}(r) &= \frac{1}{2\pi} \int_{\mathbb{R}^3} \vec{\mathbf{H}} \cdot x d\mu + \frac{(\mu - \cos \theta \eta)(\mathbb{R}^3)}{\pi}, \quad x_0 \neq 0, \\ \lim_{r \rightarrow \infty} R_{0,\theta}(r) &= 0, \end{aligned}$$

and obtain from the Simon-type monotonicity formulas (letting $\sigma \rightarrow 0^+$, $\rho \rightarrow \infty$)

$$\begin{aligned}
& \frac{1}{\pi} \int_{\mathbb{R}^3} \left| \frac{1}{4} \vec{\mathbf{H}} + \frac{(x-x_0)^\perp}{|x-x_0|^2} \right|^2 + \left| \frac{1}{4} \vec{\mathbf{H}} + \frac{(x-\xi(x_0))^\perp}{|x-\xi(x_0)|^2} \right|^2 d\mu \\
& - \frac{\cos \theta}{\pi} \int_{\mathbb{R}^3} \left(\frac{x-x_0}{|x-x_0|^2} \cdot x \right)^2 + \left(\frac{x-\xi(x_0)}{|x-\xi(x_0)|^2} \cdot x \right)^2 d\eta \\
& = \lim_{\rho \rightarrow \infty} G_{x_0, \theta}(\rho) - \lim_{\sigma \rightarrow 0^+} G_{x_0, \theta}(\sigma) \\
& = \frac{1}{8\pi} \int_{\mathbb{R}^3} |\vec{\mathbf{H}}|^2 d\mu + \frac{1}{2\pi} \int_{\mathbb{R}^3} \vec{H} \cdot x d\mu + \frac{(\mu - \cos \theta \eta)(\mathbb{R}^3)}{\pi} - \tilde{\Theta}(\mu - \cos \theta \eta, x_0) \\
& = \frac{1}{8\pi} \int_{\mathbb{R}^3} |\vec{\mathbf{H}}|^2 d\mu + \frac{\sin \theta \gamma(\mathbb{S}^2) - 2 \cos \theta \eta(\mathbb{R}^3)}{2\pi} - \tilde{\Theta}(\mu - \cos \theta \eta, x_0)
\end{aligned} \tag{4.15}$$

for $x_0 \neq 0$, and for $x_0 = 0$:

$$\frac{1}{\pi} \int_{\mathbb{R}^3} \left| \frac{1}{4} \vec{\mathbf{H}} + \frac{x^\perp}{|x|^2} \right|^2 d\mu = \frac{1}{16\pi} \int_{\mathbb{R}^3} |\vec{\mathbf{H}}|^2 d\mu + \frac{\sin \theta \gamma(\mathbb{S}^2)}{2\pi} - \tilde{\Theta}(\mu, 0).$$

4.3. Applications of Eqn. 4.15. As a by-product of the monotonicity formula, we may establish the lower bound for the Willmore functional, which is conformal invariant and was proved in the previous Section. Moreover, we may obtain the optimal area estimate for minimal capillary surfaces in the unit ball in a rather direct way, which needs not go through the Willmore functional.

Another proof of Theorem 1.4. In the case that $x_0 \in \mathbb{S}^2$, we clearly have $\xi(x_0) = x_0$. Besides, for any $x \in \mathbb{S}^2$, there holds $|x - x_0|^4 = 4(1 - (x \cdot x_0))^2$, and hence a direct computation shows that

$$\left(\frac{x-x_0}{|x-x_0|^2} \cdot x \right)^2 + \left(\frac{x-\xi(x_0)}{|x-\xi(x_0)|^2} \cdot x \right)^2 = \frac{1}{2} \text{ on } \mathbb{S}^2.$$

Taking this into account, the Simon-type monotonicity formula (4.15) then reads

$$\begin{aligned}
& \frac{1}{\pi} \int_{\mathbb{R}^3} \left| \frac{1}{4} \vec{\mathbf{H}} + \frac{(x-x_0)^\perp}{|x-x_0|^2} \right|^2 + \left| \frac{1}{4} \vec{\mathbf{H}} + \frac{(x-\xi(x_0))^\perp}{|x-\xi(x_0)|^2} \right|^2 d\mu \\
& = \frac{1}{8\pi} \int_{\mathbb{R}^3} |\vec{\mathbf{H}}|^2 d\mu + \frac{\sin \theta \gamma(\mathbb{S}^2) - \cos \theta \eta(\mathbb{R}^3)}{2\pi} - \tilde{\Theta}(\mu - \cos \theta \eta, x_0).
\end{aligned}$$

Now since $F: \Sigma \rightarrow \mathbb{R}^3$ be a capillary minimal immersion, and also recall (4.4), we obtain

$$\frac{1}{\pi} \int_{\mathbb{R}^3} \left| \frac{(x-x_0)^\perp}{|x-x_0|^2} \right|^2 + \left| \frac{(x-\xi(x_0))^\perp}{|x-\xi(x_0)|^2} \right|^2 d\mu = \frac{2|\Sigma| - \cos \theta |T|}{2\pi} - \tilde{\Theta}(\mu - \cos \theta \eta, x_0).$$

As (3.6) we have $\tilde{\Theta}(\mu - \cos \theta \eta, x_0) = (1 - \cos \theta)N(x_0)$, where $N(x_0) = \mathcal{H}^0(F^{-1}(x_0))$. Theorem 1.4 then follows. \square

REFERENCES

- [1] Abdelhamid Ainouz and Rabah Souam, *Stable capillary hypersurfaces in a half-space or a slab*, Indiana Univ. Math. J. **65** (2016), no. 3, 813–831. MR3528820
- [2] Kenneth A. Brakke, *The motion of a surface by its mean curvature.*, Math. Notes (Princeton), Princeton University Press, Princeton, NJ, 1978. MR485012
- [3] Simon Brendle, *A sharp bound for the area of minimal surfaces in the unit ball*, Geom. Funct. Anal. **22** (2012), no. 3, 621–626. MR2972603
- [4] ———, *The isoperimetric inequality for a minimal submanifold in Euclidean space*, J. Amer. Math. Soc. **34** (2021), no. 2, 595–603. MR4280868
- [5] ———, *Minimal hypersurfaces and geometric inequalities*, Ann. Fac. Sci. Toulouse Math. (6) **32** (2023), no. 1, 179–201. MR4574744
- [6] Luigi De Masi and Guido De Philippis, *Min-max construction of minimal surfaces with a fixed angle at the boundary*, 2021. arXiv:2111.09913.
- [7] G. De Philippis and F. Maggi, *Regularity of free boundaries in anisotropic capillarity problems and the validity of Young's law*, Arch. Ration. Mech. Anal. **216** (2015), no. 2, 473–568. MR3317808
- [8] Guido De Philippis and Francesco Maggi, *Dimensional estimates for singular sets in geometric variational problems with free boundaries*, J. Reine Angew. Math. **725** (2017), 217–234. MR3630122
- [9] Robert Finn, *Equilibrium capillary surfaces*, Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 284, Springer-Verlag, New York, 1986. MR816345
- [10] Ailana Fraser and Richard Schoen, *The first Steklov eigenvalue, conformal geometry, and minimal surfaces*, Adv. Math. **226** (2011), no. 5, 4011–4030. MR2770439
- [11] Michael Grüter and Jürgen Jost, *Allard type regularity results for varifolds with free boundaries*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **13** (1986), no. 1, 129–169. MR863638
- [12] Han Hong and Artur B. Saturnino, *Capillary surfaces: Stability, index and curvature estimates*, J. Reine Angew. Math. **803** (2023), 233–265. MR4649183
- [13] Xiaohan Jia, Guofang Wang, Chao Xia, and Xuwen Zhang, *Heintze-Karcher inequality and capillary hypersurfaces in a wedge*, 2022. to appear in **Ann. Sc. Norm. Super. Pisa Cl. Sci.** doi: [10.2422/2036-2145.202212_001](https://doi.org/10.2422/2036-2145.202212_001), arXiv:2209.13839.
- [14] ———, *Heintze-Karcher inequality for anisotropic free boundary hypersurfaces in convex domains*, 2023. to appear in **J. Math. Study**, Special Issue for the 100th Anniversary of Mathematics of XMU, arXiv:2311.01162.
- [15] Xiaohan Jia, Guofang Wang, Chao Xia, and Xuwen Zhang, *Alexandrov's theorem for anisotropic capillary hypersurfaces in the half-space*, Arch. Ration. Mech. Anal. **247** (2023), no. 2, Paper No. 25, 19. MR4562813
- [16] Takashi Kagaya and Yoshihiro Tonegawa, *A fixed contact angle condition for varifolds*, Hiroshima Math. J. **47** (2017), no. 2, 139–153. MR3679887
- [17] Ernst Kuwert and Reiner Schätzle, *Removability of point singularities of Willmore surfaces*, Ann. of Math. (2) **160** (2004), no. 1, 315–357. MR2119722
- [18] ———, *The Willmore functional*, CRM Series, vol. 13, Ed. Norm., Pisa, 2012. MR2882586
- [19] Chao Li, Xin Zhou, and Jonathan J. Zhu, *Min-max theory for capillary surfaces*, 2021. arXiv:2111.09924.
- [20] Peter Li and Shing Tung Yau, *A new conformal invariant and its applications to the Willmore conjecture and the first eigenvalue of compact surfaces*, Invent. Math. **69** (1982), no. 2, 269–291. MR674407
- [21] Francesco Maggi, *Sets of finite perimeter and geometric variational problems*, Cambridge Studies in Advanced Mathematics, vol. 135, Cambridge University Press, Cambridge, 2012. An introduction to geometric measure theory. MR2976521
- [22] Fernando C. Marques and André Neves, *The Willmore conjecture*, Jahresber. Dtsch. Math.-Ver. **116** (2014), no. 4, 201–222. MR3280571
- [23] Reiner Schätzle, *The Willmore boundary problem*, Calc. Var. Partial Differential Equations **37** (2010), no. 3–4, 275–302. MR2592972

- [24] Leon Simon, *Lectures on geometric measure theory*, Proceedings of the Centre for Mathematical Analysis, Australian National University, vol. 3, Australian National University, Centre for Mathematical Analysis, Canberra, 1983. MR756417
- [25] ———, *Existence of surfaces minimizing the Willmore functional*, Comm. Anal. Geom. **1** (1993), no. 2, 281–326. MR1243525
- [26] Jean E. Taylor, *Boundary regularity for solutions to various capillarity and free boundary problems*, Comm. Partial Differential Equations **2** (1977), no. 4, 323–357. MR487721
- [27] Alexander Volkman, *A monotonicity formula for free boundary surfaces with respect to the unit ball*, Comm. Anal. Geom. **24** (2016), no. 1, 195–221. MR3514558
- [28] Guofang Wang, Liangjun Weng, and Chao Xia, *Alexandrov–Fenchel inequalities for convex hypersurfaces in the half-space with capillary boundary*, Math. Ann. (2023). <https://doi.org/10.1007/s00208-023-02571-4>.
- [29] Chao Xia and Xuwen Zhang, *"Alexandrov-type theorem for singular capillary CMC hypersurfaces in the half-space"*, 2023. [arXiv:2304.01735](https://arxiv.org/abs/2304.01735).

(G.W) MATHEMATISCHES INSTITUT, UNIVERSITÄT FREIBURG, ERNST-ZERMELO-STR.1, 79104,
FREIBURG, GERMANY
Email address: guofang.wang@math.uni-freiburg.de

(C.X) SCHOOL OF MATHEMATICAL SCIENCES, XIAMEN UNIVERSITY, 361005, XIAMEN, P.R. CHINA
Email address: chaoxia@xmu.edu.cn

(X.Z) MATHEMATISCHES INSTITUT, UNIVERSITÄT FREIBURG, ERNST-ZERMELO-STR.1, 79104,
FREIBURG, GERMANY
Email address: xuwen.zhang@math.uni-freiburg.de