

A Deceptively Simple Quadratic Recurrence

STEVEN FINCH

October 8, 2024

ABSTRACT. Standard techniques for treating linear recurrences no longer apply for quadratic recurrences. It is not hard to determine asymptotics for a specific parametrized model over a wide domain of values (all $p \neq 1/2$ here). The gap between theory and experimentation seems insurmountable, however, at a single outlier ($p = 1/2$).

Fix $0 < p < 1$ and define a quadratic recurrence [1]

$$a_0 = 0, \quad a_k = (1 - p) + p a_{k-1}^2 \quad \text{for } k \geq 1$$

which arises in the study of random Galton-Watson binary tree heights. Clearly

$$\lim_{k \rightarrow \infty} a_k = r = \begin{cases} 1 & \text{if } 0 < p \leq \frac{1}{2}, \\ \frac{1-p}{p} & \text{if } \frac{1}{2} < p < 1 \end{cases}$$

and $0 < r \leq 1$. We wish initially to prove that the convergence rate of $\{a_k\}$ is exponential if and only if $p \neq 1/2$. More precisely,

$$0 < \lim_{k \rightarrow \infty} \frac{r - a_k}{(2rp)^k} = r \prod_{j=0}^{\infty} \frac{r + a_j}{2r} < 1.$$

The case $p = 1/2$ is more difficult. We examine Schoenfield's [2, 3] analysis in deriving the asymptotic expansion

$$a_k \sim 1 - \frac{2}{k} + \frac{2 \ln(k) + C}{k^2} - \frac{2 \ln(k)^2 + (2C - 2) \ln(k) + (\frac{1}{2}C^2 - C + 1)}{k^3} + \dots$$

and in calculating the constant $C = 2(1.76799378\dots) = 3.53598757\dots$. This interesting case occurs in optimal stopping theory as well [4].

⁰Copyright © 2024 by Steven R. Finch. All rights reserved.

1. SUBCRITICAL

Assume that $0 < p < 1/2$. First, note that $0 \leq a_k < 1$ for all k by induction ($a_k \geq 1 - p > 0$ is obvious; supposing $0 \leq a_{k-1} < 1$, we obtain $a_k < (1 - p) + p = 1$). Now, writing $b_k = 1 - a_k$, we have $b_0 = 1$, $0 < b_k \leq 1$ and

$$\begin{aligned} b_k &= p(1 - a_{k-1}^2) = p(1 - a_{k-1})(1 + a_{k-1}) \\ &= p b_{k-1}(2 - b_{k-1}) \\ &< 2p b_{k-1} < (2p)^2 b_{k-2} < (2p)^3 b_{k-3} \end{aligned}$$

thus $b_k < (2p)^k$ for all k . Observe that

$$\begin{aligned} b_k &= 2p b_{k-1} \left(1 - \frac{b_{k-1}}{2}\right) \\ &= (2p)^2 b_{k-2} \left(1 - \frac{b_{k-2}}{2}\right) \left(1 - \frac{b_{k-1}}{2}\right) \\ &= (2p)^3 b_{k-3} \left(1 - \frac{b_{k-3}}{2}\right) \left(1 - \frac{b_{k-2}}{2}\right) \left(1 - \frac{b_{k-1}}{2}\right) \\ &= (2p)^k \prod_{j=0}^{k-1} \left(1 - \frac{b_j}{2}\right) \end{aligned}$$

hence

$$\begin{aligned} C &= \lim_{k \rightarrow \infty} \frac{1 - a_k}{(2p)^k} = \lim_{k \rightarrow \infty} \frac{b_k}{(2p)^k} = \prod_{j=0}^{\infty} \left(1 - \frac{b_j}{2}\right) \\ &= \prod_{j=0}^{\infty} \left(1 - \frac{1 - a_j}{2}\right) = \prod_{j=0}^{\infty} \frac{1 + a_j}{2} \end{aligned}$$

exists and is nonzero since

$$\sum_{j=0}^{\infty} \frac{b_j}{2} < \frac{1}{2} \sum_{j=0}^{\infty} (2p)^j$$

converges. This completes the proof. The expression for C as an infinite product turns out to be useful for high precision estimates of C , given p (see Table 1).

2. SUPERCRITICAL

The following lemmata is needed for $i \geq 1$:

$$r^i p = r^{i-1}(1 - p)$$

and is true because $r p = ((1 - p)/p)p = 1 - p$.

Assume that $1/2 < p < 1$. First, note that $0 \leq a_k < r$ for all k by induction ($a_k \geq 1 - p > 0$ is obvious; supposing $0 \leq a_{k-1} < r$, we obtain

$$a_k < (1 - p) + p r^2 = p r + (1 - p)r = r$$

by lemmata, $i = 1$ & 2). Now, writing $b_k = r - a_k$, we have $b_0 = r$, $0 < b_k \leq r$ and

$$\begin{aligned} b_k &= r - (1 - p) - p a_{k-1}^2 && \text{(by lemmata, } i = 1) \\ &= r - r p - p a_{k-1}^2 && \\ &= r(1 - p) - p a_{k-1}^2 && \\ &= r^2 p - p a_{k-1}^2 && \text{(by lemmata, } i = 2) \\ &= p (r^2 - a_{k-1}^2) && \\ &= p (r - a_{k-1})(r + a_{k-1}) && \\ &= p b_{k-1} [2r - (r - a_{k-1})] && \\ &= p b_{k-1} (2r - b_{k-1}) && \\ &< 2 r p b_{k-1} < (2 r p)^2 b_{k-2} < (2 r p)^3 b_{k-3} \end{aligned}$$

thus $b_k < (2 r p)^k b_0 = r(2 r p)^k$ for all k . Observe that

$$\begin{aligned} b_k &= 2 r p b_{k-1} \left(1 - \frac{b_{k-1}}{2r}\right) \\ &= (2 r p)^2 b_{k-2} \left(1 - \frac{b_{k-2}}{2r}\right) \left(1 - \frac{b_{k-1}}{2r}\right) \\ &= (2 r p)^3 b_{k-3} \left(1 - \frac{b_{k-3}}{2r}\right) \left(1 - \frac{b_{k-2}}{2r}\right) \left(1 - \frac{b_{k-1}}{2r}\right) \\ &= r(2 r p)^k \prod_{j=0}^{k-1} \left(1 - \frac{b_j}{2r}\right) \end{aligned}$$

because $b_0 = r$; hence

$$\begin{aligned} C &= \lim_{k \rightarrow \infty} \frac{r - a_k}{(2 r p)^k} = \lim_{k \rightarrow \infty} \frac{b_k}{(2 r p)^k} = r \prod_{j=0}^{\infty} \left(1 - \frac{b_j}{2r}\right) \\ &= r \prod_{j=0}^{\infty} \left(1 - \frac{r - a_j}{2r}\right) = r \prod_{j=0}^{\infty} \frac{r + a_j}{2r} \end{aligned}$$

exists and is nonzero since

$$\sum_{j=0}^{\infty} \frac{b_j}{2r} < \frac{1}{2} \sum_{j=0}^{\infty} (2 r p)^j$$

converges. This completes the proof. The expression for C as an infinite product turns out to be useful for high precision estimates of C , given p (see Table 1).

Table 1. *Numerical estimates of C : no closed-form expressions are known*

p	C	p	C
1/5	0.423894537869731...	3/5	0.158431105979816...
1/4	0.392906852755779...	2/3	0.161059687971223...
1/3	0.322119375942447...	3/4	0.130968950918593...
2/5	0.237646658969724...	4/5	0.105973634467432...

3. CRITICAL

Assume that $p = 1/2$. While studying

$$a_0 = 0, \quad a_k = \frac{1}{2} (1 + a_{k-1}^2) \quad \text{for } k \geq 1$$

in the limit as $k \rightarrow \infty$, we shall defer to the standard texts [5, 6, 7] just once: our starting point will be

$$1 - a_k \sim \frac{2}{k + \ln(k) + \frac{1}{2}C}$$

or equivalently

$$a_k \sim 1 - \frac{2}{k} + \frac{2 \ln(k) + C}{k^2}$$

for some constant C . On the basis of numerical experimentation, Schoenfield [2, 3] hypothesized that the next terms of the asymptotic series must be of the form

$$\frac{c_{3,2} \ln(k)^2 + c_{3,1} \ln(k) + c_{3,0}}{k^3} + \frac{c_{4,3} \ln(k)^3 + c_{4,2} \ln(k)^2 + c_{4,1} \ln(k) + c_{4,0}}{k^4} + \dots$$

He actually went as far as $c_{20,0}/k^{20}$, but we shall stop at $c_{4,0}/k^4$ for brevity's sake. The challenge is to express each coefficient $c_{i,j}$ as a polynomial in C .

Letting $x = c_{3,2}$, $y = c_{3,1}$, $z = c_{3,0}$ for convenience, we replace a_{k+1} by

$$1 - \frac{2}{k+1} + \frac{2 \ln(k+1) + C}{(k+1)^2} + \frac{x \ln(k+1)^2 + y \ln(k+1) + z}{(k+1)^3}$$

and expand in powers of k and $\ln(k)$:

$$-\frac{2}{k+1} \sim -\frac{2}{k} + \frac{2}{k^2} - \frac{2}{k^3} + \frac{2}{k^4} - \frac{2}{k^5} + \dots,$$

$$\frac{2 \ln(k+1)}{(k+1)^2} \sim \left(\frac{2}{k^2} - \frac{4}{k^3} + \frac{6}{k^4} - \frac{8}{k^5} + \dots \right) \ln(k) + \left(\frac{2}{k^3} - \frac{5}{k^4} + \frac{26}{3k^5} - \dots \right),$$

$$\begin{aligned}\frac{C}{(k+1)^2} &\sim \frac{C}{k^2} - \frac{2C}{k^3} + \frac{3C}{k^4} - \frac{4C}{k^5} + \dots, \\ \frac{x \ln(k+1)^2}{(k+1)^3} &\sim x \left(\frac{1}{k^3} - \frac{3}{k^4} + \frac{6}{k^5} - \dots \right) \ln(k)^2 + x \left(\frac{2}{k^4} - \frac{7}{k^5} + \dots \right) \ln(k) + x \left(\frac{1}{k^5} - \dots \right), \\ \frac{y \ln(k+1)}{(k+1)^3} &\sim y \left(\frac{1}{k^3} - \frac{3}{k^4} + \frac{6}{k^5} - \dots \right) \ln(k) + y \left(\frac{1}{k^4} - \frac{7}{2k^5} + \dots \right), \\ \frac{z}{(k+1)^3} &\sim z \left(\frac{1}{k^3} - \frac{3}{k^4} + \frac{6}{k^5} - \dots \right).\end{aligned}$$

Upon rearrangement, a_{k+1} becomes

$$\begin{aligned}1 - 3x \frac{\ln(k)^2}{k^4} + (6 + 2x - 3y) \frac{\ln(k)}{k^4} + (-3 + 3C + y - 3z) \frac{1}{k^4} + x \frac{\ln(k)^2}{k^3} \\ + (-4 + y) \frac{\ln(k)}{k^3} + (-2C + z) \frac{1}{k^3} + 2 \frac{\ln(k)}{k^2} + (2 + C) \frac{1}{k^2} - 2 \frac{1}{k}.\end{aligned}$$

Performing an analogous substitution in a_k , the expression $(1 + a_k^2)/2$ becomes

$$\begin{aligned}1 + (2 - 2x) \frac{\ln(k)^2}{k^4} + (2C - 2y) \frac{\ln(k)}{k^4} + \left(\frac{C^2}{2} - 2z \right) \frac{1}{k^4} + x \frac{\ln(k)^2}{k^3} \\ + (-4 + y) \frac{\ln(k)}{k^3} + (-2C + z) \frac{1}{k^3} + 2 \frac{\ln(k)}{k^2} + (2 + C) \frac{1}{k^2} - 2 \frac{1}{k}.\end{aligned}$$

Matching coefficients, we obtain

$$\begin{aligned}-3x = 2 - 2x \quad \text{hence} \quad c_{3,2} = x = -2; \\ 6 + 2(-2) - 3y = 2C - 2y \quad \text{hence} \quad c_{3,1} = y = -2C + 2; \\ -3 + 3C + (-2C + 2) - 3z = \frac{C^2}{2} - 2z \quad \text{hence} \quad c_{3,0} = z = -\frac{C^2}{2} + C - 1\end{aligned}$$

as was to be shown.

Let $t = c_{4,3}$, $u = c_{4,2}$, $v = c_{4,1}$, $w = c_{4,0}$. We return to replacing a_{k+1} but with higher precision:

$$\begin{aligned}\frac{t \ln(k+1)^3}{(k+1)^4} &\sim t \left(\frac{1}{k^4} - \frac{4}{k^5} + \dots \right) \ln(k)^3 + t \left(\frac{3}{k^5} - \dots \right) \ln(k)^2, \\ \frac{u \ln(k+1)^2}{(k+1)^4} &\sim u \left(\frac{1}{k^4} - \frac{4}{k^5} + \dots \right) \ln(k)^2 + u \left(\frac{2}{k^5} - \dots \right) \ln(k), \\ \frac{v \ln(k+1)}{(k+1)^4} &\sim v \left(\frac{1}{k^4} - \frac{4}{k^5} + \dots \right) \ln(k) + v \left(\frac{1}{k^5} - \dots \right),\end{aligned}$$

$$\frac{w}{(k+1)^4} \sim w \left(\frac{1}{k^4} - \frac{4}{k^5} + \dots \right).$$

The new terms in the rearranged a_{k+1} become

$$\begin{aligned} & -4t \frac{\ln(k)^3}{k^5} + (6x + 3t - 4u) \frac{\ln(k)^2}{k^5} + (-8 - 7x + 6y + 2u - 4v) \frac{\ln(k)}{k^5} \\ & + \left(\frac{20}{3} - 4C + x - \frac{7}{2}y + 6z + v - 4w \right) \frac{1}{k^5}. \end{aligned}$$

Analogous substitutions in $(1 + a_k^2)/2$ and matching coefficients give rise to equations

$$-4t = 2x - 2t,$$

$$6x + 3t - 4u = Cx + 2y - 2u,$$

$$-8 - 7x + 6y + 2u - 4v = Cy + 2z - 2v,$$

$$\frac{20}{3} - 4C + x - \frac{7}{2}y + 6z + v - 4w = Cz - 2w$$

i.e.,

$$-4t = 2(-2) - 2t \quad \text{hence} \quad c_{4,3} = t = 2;$$

$$6(-2) + 3(2) - 4u = C(-2) + 2(-2C + 2) - 2u \quad \text{hence} \quad c_{4,2} = u = 3C - 5;$$

$$-8 - 7(-2) + 6(-2C + 2) + 2(3C - 5) - 4v = C(-2C + 2) + 2 \left(-\frac{C^2}{2} + C - 1 \right) - 2v$$

hence

$$c_{4,1} = v = \frac{3}{2}C^2 - 5C + 5;$$

$$\frac{20}{3} - 4C + (-2) - \frac{7}{2}(-2C + 2) + 6 \left(-\frac{C^2}{2} + C - 1 \right) + \left(\frac{3}{2}C^2 - 5C + 5 \right) - 4w = C \left(-\frac{C^2}{2} + C - 1 \right) - 2w$$

hence

$$c_{4,0} = w = \frac{1}{4}C^3 - \frac{5}{4}C^2 + \frac{5}{2}C - \frac{5}{3}.$$

To find $c_{5,0}$ would require all expansions for $\ln(k)^j/k^i$ to order $i = 6 > j$.

Schoenfield [2, 3] used an elaborate method and his $c_{20,0}/k^{20}$ series for numerically calculating the constant C to over 1000 digits. A less accurate method involves computing $a_{100000000}$ exactly via recursion, setting this equal to our $c_{4,0}/k^4$ series and then solving:

$$C = 3.535987572272308\dots$$

In an unrelated thread, Schoenfield employed a similarly intricate procedure to evaluate a quadratic threshold constant $\lambda = 0.399524667096799\dots$ due to Somos [8, 9].

4. CLOSING WORDS

The recursion for $p = 1/2$ appears elsewhere in disguised form. Letting

$$a_k = 1 - 2\alpha_k$$

we have

$$\begin{aligned} 1 - 2\alpha_k &= \frac{1}{2} [1 + (1 - 2\alpha_{k-1})^2] \\ &= \frac{1}{2} (2 - 4\alpha_{k-1} + 4\alpha_{k-1}^2) = 1 - 2\alpha_{k-1} + 2\alpha_{k-1}^2 \end{aligned}$$

therefore

$$\alpha_0 = \frac{1}{2}, \quad \alpha_k = \alpha_{k-1}(1 - \alpha_{k-1}) \quad \text{for } k \geq 1.$$

Clearly $\alpha_k = (1 - a_k)/2$. As a consequence of the preceding,

$$\alpha_k \sim \frac{1}{k} - \frac{\ln(k) + c}{k^2} + \frac{\ln(k)^2 + (2c - 1)\ln(k) + (c^2 - c + \frac{1}{2})}{k^3} - + \dots$$

where $c = C/2 = 1.767993786136154\dots$. The series $\sum_{k=0}^{\infty} \alpha_k$ diverges akin to the harmonic series [10] and the constant

$$s_1 = \alpha_0 + \sum_{k=1}^{\infty} \left(\alpha_k - \frac{1}{k} \right) = -1.60196478\dots$$

provably exists. In contrast,

$$s_2 = \sum_{k=0}^{\infty} \alpha_k^2 = \frac{1}{2}$$

because, by induction,

$$\sum_{k=0}^n \alpha_k^2 = \sum_{k=0}^{n-1} \alpha_k^2 + \alpha_n^2 = \left(\frac{1}{2} - \alpha_n \right) + \alpha_n^2 = \frac{1}{2} - \alpha_n(1 - \alpha_n) = \frac{1}{2} - \alpha_{n+1}$$

and $\alpha_{n+1} \rightarrow 0$. Closed-form expressions for

$$s_m = \sum_{k=0}^{\infty} \alpha_k^m = \begin{cases} 0.159488853036112\dots & \text{if } m = 3 \\ 0.068977706072225\dots & \text{if } m = 4 \\ 0.032622409767106\dots & \text{if } m = 5 \\ 0.015934111084642\dots & \text{if } m = 6 \\ 0.007884618832013\dots & \text{if } m = 7 \\ 0.003923447888623\dots & \text{if } m = 8 \end{cases}$$

remain unknown. The constants s_m appear in an alternative formula for c . A technique called “bootstrapping” is instructive here [11, 12]:

$$\begin{aligned}
\frac{1}{\alpha_k} &= \frac{1}{\alpha_{k-1}} \frac{1}{1 - \alpha_{k-1}} \\
&= \frac{1}{\alpha_{k-1}} (1 + \alpha_{k-1} + \alpha_{k-1}^2 + \alpha_{k-1}^3 + \alpha_{k-1}^4 + \cdots) \\
&= \frac{1}{\alpha_{k-1}} + 1 + \alpha_{k-1} + \alpha_{k-1}^2 + \alpha_{k-1}^3 + \cdots \\
&= \frac{1}{\alpha_{k-2}} + 2 + (\alpha_{k-1} + \alpha_{k-2}) + (\alpha_{k-1}^2 + \alpha_{k-2}^2) + (\alpha_{k-1}^3 + \alpha_{k-2}^3) + \cdots \\
&= \frac{1}{\alpha_{k-3}} + 3 + (\alpha_{k-1} + \alpha_{k-2} + \alpha_{k-3}) + (\alpha_{k-1}^2 + \alpha_{k-2}^2 + \alpha_{k-3}^2) + (\alpha_{k-1}^3 + \alpha_{k-2}^3 + \alpha_{k-3}^3) + \cdots \\
&= \frac{1}{\alpha_0} + k + \sum_{j=0}^{k-1} \alpha_j + \sum_{j=0}^{k-1} \alpha_j^2 + \sum_{j=0}^{k-1} \alpha_j^3 + \cdots \\
&\sim 2 + k + (\ln(k) + \gamma + s_1) + s_2 + s_3 + \cdots
\end{aligned}$$

where γ is the Euler-Mascheroni constant. The promised formula

$$c = 2 + \gamma + \sum_{m=1}^{\infty} s_m$$

unfortunately is not computationally helpful, owing to our limited understanding of the sequence $s_1, s_2, s_3, s_4, \dots$.

5. ACKNOWLEDGEMENTS

Robert Israel and Anthony Quas gave the simple proof that C exists for $0 < p < 1/2$, reproduced here from [1]. The analogous formula corresponding to $1/2 < p < 1$ is new, as far as I know. A significant portion of the logistic recurrence asymptotics appear in [13] – I had overlooked this fact – and their claim that $\exp(c-1) \approx 2.15768$ is noteworthy. Philippe Flajolet is still deeply missed by all who knew him.

REFERENCES

- [1] S. R. Finch, Subcritical Galton-Watson trees, *Mathematical Constants II*, Cambridge Univ. Press, 2019, pp. 511–519; MR3887550.
- [2] J. E. Schoenfeld, Magma program for determining terms of OEIS A245771, <http://oeis.org/A245771/a245771.txt>
- [3] N. J. A. Sloane, On-Line Encyclopedia of Integer Sequences, A245771.

- [4] S. R. Finch, Optimal stopping constants, *Mathematical Constants*, Cambridge Univ. Press, 2003, pp. 361–369; MR2003519.
- [5] L. Moser, On a problem of Cayley, *Scripta Math.* 22 (1956) 289–292.
- [6] J. P. Gilbert and F. Mosteller, Recognizing the maximum of a sequence, *J. Amer. Statist. Assoc.* 61 (1966) 35–73; MR0198637.
- [7] T. S. Ferguson, *Optimal Stopping and Applications*, unpublished manuscript (2000), <http://www.math.ucla.edu/~tom/Stopping/Contents.html>.
- [8] J. E. Schoenfeld, Magma program for determining terms of OEIS A268107, <http://oeis.org/A268107/a268107.txt>.
- [9] N. J. A. Sloane, On-Line Encyclopedia of Integer Sequences, A268107.
- [10] P. Flajolet and A. Odlyzko, The average height of binary trees and other simple trees, *J. Comput. System Sci.* 25 (1982) 171–213; MR0680517.
- [11] P. Flajolet, Analyses of tree height, Séminaire de Combinatoire énumérative et analytique, Oct. 2010, Paris Institut Henri Poincaré, slide 24; <http://algo.inria.fr/flajolet/Publications/Slides/IHP-OCT10-slides.pdf>.
- [12] P. Flajolet and R. Sedgewick, *Analytic Combinatorics*, Cambridge Univ. Press, 2009, pp. 536–537; MR2483235.
- [13] E. Ionascu and P. Stanica, Effective asymptotics for some nonlinear recurrences and almost doubly-exponential sequences, *Acta Math. Univ. Comenian.* 73 (2004) 75–87; MR2076045.
- [14] S. R. Finch, Generalized logistic maps and convergence, arXiv:2409.15175.
- [15] S. R. Finch, Iterated radical expansions and convergence, arXiv:2410.02114.

Steven Finch
MIT Sloan School of Management
Cambridge, MA, USA
steven_finch_math@outlook.com