

Extended symmetry of higher Painlevé equations of even periodicity and their rational solutions

H. Aratyn¹, J.F. Gomes², G.V. Lobo², and A.H. Zimerman²

¹ Department of Physics, University of Illinois at Chicago, 845 W. Taylor St.
Chicago, Illinois 60607-7059, USA

² Instituto de Física Teórica-UNESP, Rua Dr Bento Teobaldo Ferraz 271,
Bloco II, 01140-070 São Paulo, Brazil

September 10, 2024

Abstract

The structure of extended affine Weyl symmetry group of higher Painlevé equations of N periodicity depends on whether N is even or odd. We find that for even N , the symmetry group $\widehat{A}_{N-1}^{(1)}$ contains the conventional Bäcklund transformations $s_j, j = 1, \dots, N$, the group of automorphisms consisting of cycling permutations but also reflections on a periodic circle of N points, which is a novel feature uncovered in this paper.

The presence of reflection automorphisms is connected to existence of degenerated solutions and for $N = 4$ we explicitly show how the reflection automorphisms around even points cause degeneracy of a class of rational solutions obtained on the orbit of translation operators of $\widehat{A}_3^{(1)}$. We obtain the closed expressions for solutions and their degenerated counterparts in terms of determinants of Kummer polynomials.

1 Introduction

This paper analyses the higher even N Painlevé equations:

$$g_{i,x} = \sum_{k=1}^{\frac{N}{2}-1} g_i g_{i+2k} \left(- \sum_{r=1}^k g_{i+2r-1} + \sum_{s=k}^{\frac{N}{2}-1} g_{i+2s+1} \right) - \frac{\alpha_i + \alpha_{i+2} + \dots + \alpha_{i+N-2}}{2x} g_i + \frac{\alpha_i}{2x}, \quad i = 1, \dots, N, \quad (1.1)$$

with g_i and α_i satisfying conditions

$$\sum_{r=1}^{N/2} g_{2r} = 1, \quad \sum_{r=1}^{N/2} g_{2r+1} = 1, \quad \sum_i^N \alpha_i = 2, \quad (1.2)$$

that are derived from the formalism of dressing chain equations (2.2) of even N periodicity, meaning that $g_{i+N} = g_i, \alpha_{i+N} = \alpha_i$, see also the formulation of higher Painlevé equations in [21] in terms of variable z such that $x = -z^2/2$.

The higher Painlevé equations (1.1) are manifestly invariant under the Weyl group of Bäcklund symmetry transformations $s_i, i = 1, 2, \dots, N$ (3.1) as well as the group of

automorphisms $\{1, \pi, \dots, \pi^{N-1}, \widehat{P}_n, n = 1, \dots, N\}$ where π is a cyclic permutation defined in (3.1) and reflection automorphisms \widehat{P}_n are generated by reflections P_n on a periodic circle of N points (3.2) and (3.3) and induce actions of \widehat{P}_n on $g_i(x), \alpha_i$ (3.4) and on the Bäcklund symmetry transformations (3.10).

The reflections around even points $n = 2i$ of the periodic circle with the period N are shown to produce degenerated solutions (solutions that share the same parameters α_i as original solutions), when acting on solutions generated from the particular seed solution by translation operators.

The paper is organized as follows. After we derive higher $\widehat{A}_{N-1}^{(1)}$ symmetric Painlevé equations from the dressing chain equations of even periodicity in section 2, we analyze their symmetries in section 3. In this section, we introduce reflection transformations $P_n, n = 1, \dots, N$ that close under multiplication with cyclic permutations $\{1, \pi, \pi^2, \dots, \pi^{N-1}\}$ of the extended Weyl group $\widehat{A}_{N-1}^{(1)}$ and form with them a group of automorphisms of higher Painlevé equations of even N periodicity. We refer to these transformations as reflection automorphisms and observe how they naturally split into even and odd reflections depending whether the reflection is around even or odd point of $1, \dots, N$ periodic circle. All reflection automorphisms square to one and transform the underlying variable x to $-x$. We describe representation of reflections automorphisms on the $A_{N-1}^{(1)}$ symmetries $s_i, i = 1, \dots, N$ and the abelian subgroup of $A_{N-1}^{(1)}$ consisting of translation operators $T_i, i = 1, \dots, N$. These results path the way to an explanation how the even reflections introduce degeneracy among a class of rational solutions of Painlevé equations, which is explicitly done for the $N = 4$ model.

Working with the $N = 4$ case we present in section 4 an explicit construction of a class of rational solutions of Painlevé equations that reside on the orbit of translation operators of the extended affine Weyl group $A_3^{(1)}$. These solutions are given by two components $F_m^k(x, \mathbf{a}), G_m^k(x, \mathbf{a})$ and are labeled by two independent positive integer parameters that enter the formalism as powers of the translation operators. We find an explicit determinant expressions for $F_m^k(x, \mathbf{a}), G_m^k(x, \mathbf{a})$ in terms of Kummer polynomials in section 5 for all orbit solutions generated from a particular seed solution. The technique we use is based on the fact that the actions of translation operators give rise to the recurrence relations. We are able to solve these recurrence relations in terms of determinants of Kummer polynomials constructed explicitly in this section. It is crucial to emphasize the role played by one of the odd reflection automorphism that gives rise to the “duality” relation that ensures the intrinsic consistency of rational solutions by permitting two dual formulations based on two related Kummer polynomial definitions.

In section 6, we discuss the degeneracy that exists for a class of rational solutions of $N = 4$ linking it to actions of the two even reflection automorphisms that are shown to map the rational solutions to different solutions that share the values of the parameters of the $N = 4$ Painlevé equations.

The two remaining odd reflection automorphisms are discussed in section 7. The \widehat{P}_1 automorphism is shown to connect two dual solutions. The invariance under \widehat{P}_1 ensures validity the recursive relations and accordingly proves the intrinsic consistency of construction of the determinant solutions based on these recursive relations.

In section 8, we show examples of how our formalism extends to $N = 6, 8$ by utilizing presence of reflection automorphisms as a tool to finding a class of relevant solutions.

In appendix A we illustrate some of our findings by explicitly constructing braids and matrix representations of reflection automorphisms for $N = 4$.

2 Dressing chain derivation of Painlevé equations for even periodicity

We start with a formalism of the dressing chain equations of even periodicity. The conventional definition of a dressing chain of N periodicity is

$$(j_n + j_{n+1})_z = -j_n^2 + j_{n+1}^2 + \alpha_n, \quad n = 1, \dots, N, \quad j_{N+i} = j_i, \quad (2.1)$$

however for even N this expression requires, for consistency, an imposition of a quadratic constraint that can be introduced as modification of the dressing chain formulation. Such modification was proposed on basis of Dirac reduction method in [6], where the authors put forward a system of dressing chain equations of even periodicity defined as :

$$(j_i + j_{i+1})_z = -j_i^2 + j_{i+1}^2 + \alpha_i + (-1)^{i+1} \frac{(j_i + j_{i+1})}{\Phi} \Psi, \quad i = 1, 2, \dots, N, \quad j_{N+i} = j_i, \quad (2.2)$$

where

$$\Psi = \sum_{k=1}^N (-1)^{k+1} \left(j_k^2 - \frac{1}{2} \alpha_k \right), \quad \Phi = \sum_{k=1}^N j_k = z. \quad (2.3)$$

Remarkably, the dressing chain equations (2.2) can be rewritten entirely in terms of $f_i = j_i + j_{i+1}$ without any references to j_i . It needs to be emphasized that such elimination of j_i variables while expressing the dressing chain equations in terms of f_i requires inserting the definition of Ψ from (2.3) into equations (2.2). This is because the relation $f_i = j_i + j_{i+1}$ is not invertible for even N . In contrast when N is taken to be odd such relation is invertible as illustrated by example of $N = 3$ with the inverse relation given by:

$$j_1 = \frac{1}{2}(f_1 - f_2 + f_3), \quad j_2 = \frac{1}{2}(f_2 - f_3 + f_1), \quad j_3 = \frac{1}{2}(f_3 - f_1 + f_2),$$

and after inserting these relations back into equations (2.1) one obtains the symmetric Painlevé IV equations $f_{i,z} = f_i(f_{i+1} - f_{i-1}) + \alpha_i, i = 1, 2, 3$.

In the case of equations (2.2) introducing new variables

$$g_i(x) = \frac{f_i}{z} = \frac{j_i + j_{i+1}}{z}, \quad x = -z^2/2, \quad (2.4)$$

we are able to derive the N even Painlevé equations (1.1), where g_i and α_i satisfy conditions (1.2). The equations (1.1) agree with higher Noumi-Yamada equations [21] with $A_l^{(1)}$ symmetry for $l = 2n + 1$, which were originally written in terms of variable z . One of the advantages of using x is that it makes it possible to uncover a new symmetry structure of N automorphisms. These automorphisms preserve the $A_{N-1}^{(1)}$ Weyl symmetry group of Bäcklund transformations of equations (1.1) and will explain presence of denegeneracy among rational solutions. In case of equations with $N = 4$ we obtain a regular Painlevé V equation directly from (1.1) as Painlevé V equation is conventionally expressed using the variable x (see below the equation (4.2) for the derivation).

3 Symmetries of higher Painlevé equations for even N

Equations (1.1) are invariant under the extended affine $A_{N-1}^{(1)}$ Weyl group of Bäcklund transformations, s_i and an automorphism π :

$$\begin{aligned} s_i : g_{i\pm 1} &\rightarrow g_{i\pm 1} \mp \frac{\gamma_i}{g_i}, \quad g_i \rightarrow g_i, \quad g_j \rightarrow g_j, \quad j \neq i, i \pm 1, \\ \gamma_i &\rightarrow -\gamma_i, \quad \gamma_{i\pm 1} \rightarrow \gamma_{i\pm 1} + \gamma_i, \quad \gamma_j \rightarrow \gamma_j, \quad j \neq i, i \pm 1, \quad i = 1, \dots, N, \\ \pi : \pi(g_i) &= g_{i+1}, \quad \pi(\gamma_i) = \gamma_{i+1}, \end{aligned} \quad (3.1)$$

where for convenience we introduced the parameters

$$\gamma_i = \frac{\alpha_i}{2x}$$

with the normalization $\sum_i^N \gamma_i = 1/x$ that corresponds to normalization $\sum_i^N \alpha_i = 2$ for the parameters α_i .

In addition to the extended affine Weyl group of Bäcklund transformations the equations (1.1) are invariant under automorphisms that are generated by reflections around the fixed point n on a periodic circle $1, 2, 3, \dots, N+1, N+2, \dots$ with a periodic condition $k = k + N$ for any point k . These automorphisms also map $x \rightarrow -x$. We denote these automorphisms as P_n^N but will drop the upper script N , when it will be obvious from the context and will use only the P_n symbol. We distinguish between two types of automorphisms depending on whether n is even or odd. Accordingly, the two classes of reflections are :

1. Reflections around even points $n = 2i$ denoted by $P_{2i}, i = 0, 1, \dots, \frac{N}{2}$ that act as follows:

$$P_{2i}(k) = 2i - k, \quad k = 0, 1, 2, 3, 4, 5, \dots, \quad k + N = k \quad (3.2)$$

on a periodic line of points with a period of N . Each reflection is equivalent to a single transposition that interchanges k with $2i - k$. There are always two fixed points, namely, i and $i + \frac{N}{2}$ that according to

$$P_{2i}(i) = 2i - i = i, \quad P_{2i}(i + \frac{N}{2}) = 2i - i - \frac{N}{2} = i - \frac{N}{2} \sim i + \frac{N}{2},$$

that are being transformed into themselves. The symbol \sim is to indicate that the periodic condition $k = k + N$ was used. The remaining $N - 2$ points fall into $N/2 - 1$ (even/even and odd/odd) pairs that are mapped into each other under even reflections.

2. Reflections around odd points $n = 2i + 1$ denoted by $P_{2i+1}, i = 0, 1, \dots, \frac{N}{2} - 1$ that act through

$$P_{2i+1}(k) = 2i + 1 - k, \quad k = 0, 1, 2, 3, 4, 5, \dots, \quad k + N = k \quad (3.3)$$

on a periodic line of points $k = 1, 2, 3, 4, \dots$ with a period of N . Each of these reflections is equivalent to transpositions of $N/2$ pairs of even/odd points that are being interchanged into each other.

All these reflections obviously square to one:

$$P_n^2 = 1.$$

All the above reflections naturally induce transformations of α_k, g_k through:

$$\begin{aligned} \widehat{P}_n(g_k(x)) &= g_{P_n(k)}(-x) = g_{n-k}(-x), \\ \widehat{P}_n(\alpha_k) &= \alpha_{P_n(k)} = \alpha_{n-k}, \quad n = 2i, 2i + 1 \end{aligned} \quad (3.4)$$

that keep equations (1.1) invariant when also $x \rightarrow -x$. The straightforward although formal way to prove the invariance of the Painlevé equations (1.1) is to introduce transformations :

$$\widehat{P}_n(j_k) = \sqrt{-1}j_{n-k+1}, \quad \widehat{P}_n(\alpha_k) = \alpha_{n-k}, \quad \widehat{P}_n(z) = \sqrt{-1}z. \quad (3.5)$$

of the dressing equations (2.2) that generate transformations (3.4) together with $x \rightarrow -x$. It follows from these definitions that

$$\widehat{P}_n(\Psi) = (-1)^n \Psi, \quad \widehat{P}_n(\Phi) = \Phi,$$

when the use is made of periodic conditions. As a result the dressing chain equations (2.2) remain invariant under the transformations (3.5). This in turn proves invariance of equations (1.1) under automorphisms (3.4) that are generated by reflections on a periodic circle. Note that the imaginary factor present in definition (3.5) goes away when the transformation is applied on quantities x and $g_k(x)$.

We find from the definitions of reflections the following relations for obtaining the automorphisms π and its powers out of products of reflections \widehat{P}_n :

$$\pi = \widehat{P}_{n+1}\widehat{P}_n, \quad \pi^{-1} = \widehat{P}_n\widehat{P}_{n+1}, \quad \pi^2 = \widehat{P}_{n+2}\widehat{P}_n, \quad \pi^{-2} = \widehat{P}_n\widehat{P}_{n+2}, \quad n = 2i, 2i + 1. \quad (3.6)$$

More generally we find the following identities

$$\widehat{P}_{n_1}\widehat{P}_{n_2} = \pi^{n_1-n_2}, \quad \widehat{P}_{n_1}\pi^{n_2} = \widehat{P}_{n_1-n_2}, \quad (3.7)$$

which show that reflections and π automorphisms close under multiplications and form the group with identity $1 = \pi^0 = \widehat{P}_n^2$. This structure is consistent with the reflection automorphisms transforming $x \rightarrow -x$, despite the fact that π is not acting on x . We will refer to reflections and π and its powers as group of automorphisms with cyclic permutations $\{1, \pi, \pi^2, \dots, \pi^{N-1}\}$ forming a subgroup of these automorphisms.

Even/odd reflections commute among themselves

$$[P_{2i}, P_{2j}] = 0, \quad [P_{2i+1}, P_{2j+1}] = 0, \quad i \neq j$$

while we have the following conjugation transformations:

$$\begin{aligned} P_{2i+1}P_{2i}P_{2i+1} &= P_{2i+2}, & P_{2i+1}P_{2i}P_{2i+3} &= P_{2i} \\ P_{2i}P_{2i+1}P_{2i} &= P_{2i+1}, & P_{2i}P_{2i+1}P_{2i+2} &= P_{2i-1} \end{aligned} \quad (3.8)$$

for the mixed even/odd reflections.

Additional important identities are :

$$\widehat{P}_n\pi\widehat{P}_n = \pi^{-1}, \quad \widehat{P}_n\pi^{-1}\widehat{P}_n = \pi, \quad n = 2i, 2i + 1. \quad (3.9)$$

It is important that reflections preserve the Bäcklund symmetries through the following conjugation formula:

$$\widehat{P}_n s_m \widehat{P}_n = s_{P_n(m)}, \quad n = 2i, 2i + 1, \quad n, m = 1, \dots, N \quad (3.10)$$

where $P_n(m)$ is the point obtained from m by acting with the reflection P_n .

The above situation differs from the one which is encountered in the case of odd N . First, in such case the corresponding Painlevé equations remain expressed in terms of the variable z that causes a presence of imaginary factors when dealing with the transformation defined in (3.5). Furthermore the “even” and “odd” reflections are no longer clearly distinguishable. For example, for $N = 3$ trying to apply the above definitions of reflections will result in a similar action for the “even” reflection:

$$P_2(1) = 1, \quad P_2(2) = 3, \quad P_2(3) = 2,$$

as for the “odd” reflections:

$$P_1(1) = 3, \quad P_2(2) = 2, \quad P_2(3) = 1, \quad P_3(1) = 1, \quad P_3(2) = 1, \quad P_3(3) = 3.$$

We notice in all three cases the existence of one fixed point with the remaining two points being transposed into each other. Although such reflections are not inducing directly any symmetry of the symmetric Painlevé IV equations (without introducing a transformation of the variable z involving imaginary factor), a similar but albeit more general automorphisms have been recently discussed in [5].

4 A special case of $N = 4$ and Painlevé V equations

For $N = 4$ the equations (1.1) simplify to :

$$\begin{aligned} g_{i,x} &= g_i g_{i+2} (g_{i-1} - g_{i+1}) - \frac{\alpha_i + \alpha_{i+2}}{2x} g_i + \frac{\alpha_i}{2x} \\ &= g_i g_{i+2} (g_{i-1} - g_{i+1}) + \frac{\alpha_i}{2x} g_{i+2} - \frac{\alpha_{i+2}}{2x} g_i, \quad i = 1, 2, 3, 4 \end{aligned} \quad (4.1)$$

We impose everywhere in this section the conventional periodicity conditions : $g_i = g_{i+4}$, $\alpha_i = \alpha_{i+4}$.

We write $g_1 = F, g_2 = G, g_3 = 1 - F, g_4 = 1 - G$ and $\alpha_4 = 2 - \sum_{i=1}^3 \alpha_i$, with F, G satisfying equations:

$$\begin{aligned} F_x &= F(F-1)(2G-1) - \frac{\alpha_1 + \alpha_3}{2x} F + \frac{\alpha_1}{2x}, \\ G_x &= -G(G-1)(2F-1) + \frac{\alpha_1 + \alpha_3 - 2}{2x} G + \frac{\alpha_2}{2x}, \end{aligned} \quad (4.2)$$

derived from equations (4.1). In this setting the $A_3^{(1)}$ extended affine Weyl group of Bäcklund transformations, $\pi, s_i, i = 1, \dots, 4$ emerges as group of symmetry operations on equations (4.2) and can be obtained from relations (3.1) for $N = 4$.

Below we will determine explicitly solutions of equations (4.2) for the special values of the parameters $\alpha_1, \alpha_2, \alpha_3$:

$$\alpha_1 = a + 2k, \quad \alpha_2 = -2k, \quad \alpha_3 = -2m, \quad m, k \in \mathbb{Z}_+, \quad (4.3)$$

where \mathbb{Z}_+ contains positive integers and zero. In the notation of four-component parameters $\alpha_i, i = 1, 2, 3, 4$ we have

$$\alpha_i = 2 \left(\frac{a}{2} + k, -k, -m, 1 - \frac{a}{2} + m \right),$$

with a being an arbitrary variable. For the values of parameters given in (4.3) the solutions of equations (4.2) can be obtained by action of translation operators on a class of seed solutions as will be presented in the next few subsections.

Let us finally comment on connection of equations (4.2) to the Painlevé V equation that can be recovered by eliminating one of the two variables. Eliminating G from equations (4.2) we find that $y = F(F-1)^{-1}$ satisfies the Painlevé V equation:

$$y_{xx} = -\frac{y_x}{x} + \left(\frac{1}{2y} + \frac{1}{y-1} \right) y_x^2 + \frac{(y-1)^2}{x^2} \left(\alpha y + \frac{\beta}{y} \right) + \frac{\gamma}{x} y + \delta \frac{y(y+1)}{y-1}.$$

with

$$\begin{aligned} \alpha &= \frac{1}{8} \alpha_3^2, \quad \beta = -\frac{1}{8} \alpha_1^2, \quad \gamma = \frac{\alpha_4 - \alpha_2}{2}, \quad \delta = -\frac{1}{2} \\ \alpha &= \frac{1}{2} (m)^2, \quad \beta = -\frac{1}{2} \left(\frac{a}{2} + k \right)^2, \quad \gamma = 1 - \frac{a}{2} + m + k, \quad m, k \in \mathbb{Z}_+. \end{aligned}$$

Eliminating instead F from equations (4.2) we find that $y = G(G-1)^{-1}$ satisfies the Painlevé V equation with coefficients:

$$\alpha = \frac{1}{8} \alpha_4^2, \quad \beta = -\frac{1}{8} \alpha_2^2, \quad \gamma = \frac{\alpha_1 - \alpha_3}{2}, \quad \delta = -\frac{1}{2}$$

4.1 Seed solutions and translation operators

There are essentially two fundamental seed solutions to equations (4.2):

$$F = \frac{1}{2}, \quad G = \frac{1}{2}, \quad (\mathbf{a}, 1 - \mathbf{a}, \mathbf{a}, 1 - \mathbf{a}), \quad (4.4)$$

$$F = 1, \quad G = 0, \quad (\mathbf{a}, 0, 0, 2 - \mathbf{a}) \quad (4.5)$$

with an arbitrary constant \mathbf{a} . The first of these seed solutions given in (4.4) is invariant under π^2 . This class of seed solutions gives rise to Umemura polynomials and is not associated with degeneracy. It is illustrative to recall [18], that in the setting of the even dressing chain (2.2) the seed solution (4.4) is represented by solution $j_i(z) = (z/4)(1, 1, 1, 1)$ with only positive components, while the seed solution (4.5) is represented by solution $j_i(z) = (z/4)(1, 1, -1, 1)$ with one negative and three positive components. In this presentation, we will only study solutions (4.5) to obtain closed expressions for the special function solutions generated from it by Bäcklund transformations. Acting with π will generate from the solution (4.5) equivalent solutions:

$$F = 1, \quad G = 1, \quad (2 - \mathbf{a}, \mathbf{a}, 0, 0), \quad (4.6)$$

$$F = 0, \quad G = 1, \quad (0, 2 - \mathbf{a}, \mathbf{a}, 0), \quad (4.7)$$

$$F = 0, \quad G = 0, \quad (0, 0, 2 - \mathbf{a}, \mathbf{a}), \quad (4.8)$$

which therefore do not require a separate treatment. We will refer to the seed solution (4.5) using the following (Physics inspired) notation:

$$|F = 1, G = 0\rangle_{(\mathbf{a}, 0, 0, 2 - \mathbf{a})}. \quad (4.9)$$

Within the $A_3^{(1)}$ extended affine Weyl group one defines an abelian subgroup of translation operators defined as $T_i = r_{i+3}r_{i+2}r_{i+1}r_i, i = 1, 2, 3, 4$, where $r_i = r_{4+i} = s_i$ for $i = 1, 2, 3$ and $r_4 = \pi$. The translation operators commute among themselves, $T_i T_j = T_j T_i$, and generate the following translations when acting on the α_i parameters:

$$T_i(\alpha_i) = \alpha_i + 2, \quad T_i(\alpha_{i-1}) = \alpha_{i-1} - 2, \quad T_i(\alpha_j) = \alpha_j, \quad j = i + 1, j = i + 2.$$

The inverse translation operators are defined as $T_i^{-1} = r_i^{-1}r_{i+1}^{-1}r_{i+2}^{-1}r_{i+3}^{-1}$, with $r_i^{-1} = r_{4+i}^{-1} = s_i$ for $i = 1, 2, 3$ and $r_4^{-1} = \pi^{-1}$. They satisfy $T_i^{-1}T_i = 1$ and accordingly they generate the following α_i shifts

$$T_i^{-1}(\alpha_i) = \alpha_i - 2, \quad T_i^{-1}(\alpha_{i-1}) = \alpha_{i-1} + 2, \quad T_i^{-1}(\alpha_j) = \alpha_j, \quad j = i + 1, j = i + 2.$$

We notice that the Bäcklund transformations s_2, s_3 generate infinities when applied on the solution (4.5) because of $G = 0$ and $1 - F = 0$. To avoid these singularities only the following solutions are permitted to be generated out of the the seed solution (4.5) by use of translations [18]:

$$T_1^n T_2^{-k} T_4^m |F = 1, G = 0\rangle_{\alpha_{\mathbf{a}}}, \quad n \in \mathbb{Z}, \quad k, m \in \mathbb{Z}_+, \quad \alpha_{\mathbf{a}} = (\mathbf{a}, 0, 0, 2 - \mathbf{a}).$$

In this way we obtain, by action of translation operators, new solutions $T_1^n T_2^{-k} T_4^m (F = 1)$ and $T_1^n T_2^{-k} T_4^m (G = 0)$ of equations (4.2) with a new parameter:

$$T_1^n T_2^{-k} T_4^m (\alpha_{\mathbf{a}}) = (\mathbf{a} + 2n + 2k, -2k, -2m, 2 - \mathbf{a} - 2n + 2m). \quad (4.10)$$

One notices that the action of T_1^n merely produces a shift of a parameter \mathbf{a} and as shown in [18] leaves the configuration $F = 1, G = 0$ unchanged:

$$T_1^n |F = 1, G = 0\rangle_{\alpha_{\mathbf{a}}} = |F = 1, G = 0\rangle_{\alpha_{\mathbf{a}+2n}}. \quad (4.11)$$

We can therefore, replace the action of T_1 by appropriately redefining \mathbf{a} and accordingly restrict our discussion to the solutions of F, G coupled equations of the form :

$$\begin{aligned}\mathbb{T}(k, m; \mathbf{a}) &= T_2^{-k} T_4^m |F = 1, G = 0\rangle_{\alpha_{\mathbf{a}}}, \quad k, m \in \mathbb{Z}_+, \\ \alpha_{k, m, \mathbf{a}} &= T_2^{-k} T_4^m (\alpha_{\mathbf{a}}) = (\mathbf{a} + 2k, -2m, -2m, 2 - \mathbf{a} + 2m),\end{aligned}\tag{4.12}$$

where we listed both the solution and its corresponding parameter $\alpha_{k, m, \mathbf{a}}$ generated by combined actions of translation operators T_2^{-k} and T_4^m , with k, m being positive integers, acting on the seed solution (4.5). In section 5 we will find closed expressions for solutions $\mathbb{T}(k, m; \mathbf{a})$ introduced in relation (4.12) in terms of Kummer polynomials.

4.2 Action of automorphisms for the $N = 4$ model

Taking a look at the $N = 4$ equations (4.1) we can easily see that they are manifestly invariant under actions generated by four (two even and two odd) automorphisms $P_i, i = 1, 2, 3, 4$:

$$P_4 \rightarrow: x \rightarrow -x, g_1 \leftrightarrow g_3, \alpha_1 \leftrightarrow \alpha_3 \tag{4.13}$$

$$P_2 \rightarrow: x \rightarrow -x, g_2 \leftrightarrow g_4, \alpha_2 \leftrightarrow \alpha_4 \tag{4.14}$$

$$P_1 \rightarrow: x \rightarrow -x, g_1 \leftrightarrow g_4, g_2 \leftrightarrow g_3, \alpha_1 \leftrightarrow \alpha_4, \alpha_2 \leftrightarrow \alpha_3 \tag{4.15}$$

$$P_3 \rightarrow: x \rightarrow -x, g_1 \leftrightarrow g_2, g_3 \leftrightarrow g_4, \alpha_1 \leftrightarrow \alpha_2, \alpha_3 \leftrightarrow \alpha_4, \tag{4.16}$$

The above automorphisms are associated with reflections on a circle $k = 1, 2, 3, 4$ with periodic conditions $k = 5 = 1, k = 6 = 2, \dots$ obtained when we add periodic condition $k+4 = k$ to the line of points $k = 1, 2, 3, 4, 5, \dots$ introduced around equations (3.2) and (3.3) for a general even N . As in the case of general even N there is an even number of even and two odd automorphisms for $N = 4$. Namely, there are even reflections P_2, P_4 introduced in relations (3.2) that act as simple transpositions of two points. The automorphisms P_1, P_3 are odd reflections introduced in (3.3). They all act on α_k, g_k via relations (3.4).

Example 4.1. It follows that for $i = 1$ the reflection $P_{2i+1} = P_3$ acts as

$$P_3(1) = 2, \quad P_3(2) = 1, \quad P_3(3) = 0 = 4, \quad P_3(4) = -1 = 3$$

and induces automorphism (4.16).

Similarly for $i = 2$ the reflection $P_{2i+1} = P_5 \sim P_1$ acts as

$$P_5(1) = 4, \quad P_5(2) = 3, \quad P_5(3) = 2, \quad P_5(4) = 1.$$

and induces the automorphism (4.15). For even reflections it appears that they are effectively transpositions of two lines. For example for P_{2i} with $i = 1$ we have using the periodicity $i + N \sim i$:

$$P_2(1) = 1, \quad P_2(2) = 0 \sim 4, \quad P_2(3) = -1 \sim 3, \quad P_2(4) = -2 \sim 2.$$

Thus P_2 is transposing points 2 and 4 while 1 and 3 are fixed points and consequently induces \widehat{P}_2 from relation (4.14). Similarly for P_{2i} with $i = 2$ we have

$$P_4(1) = 3, \quad P_4(2) = 2, \quad P_4(3) = 1, \quad P_4(4) = 0 \sim 4.$$

Correspondingly \widehat{P}_4 acts as shown in relation (4.13).

Below are few obvious identities following from the basic definitions:

$$P_2 P_3 = P_4 P_1, \quad P_4 P_3 = P_2 P_5$$

or more generally

$$P_{2i}P_{2i+1} = P_{2i+2}P_{2i+3}$$

etc. Also it follows that

$$\widehat{P}_{2i+1}\widehat{P}_2\widehat{P}_{2i+1} = \widehat{P}_4, \quad \widehat{P}_{2i}\widehat{P}_1\widehat{P}_{2i} = \widehat{P}_3,$$

etc. We refer to Appendix A for a convenient matrix representation of such identities.

Furthermore, the product of all four reflections $\widehat{P}_4\widehat{P}_2\widehat{P}_1\widehat{P}_3$ is an identity due to

$$P_4P_2P_1P_3(k) = -(2 - (1 - (3 - k))) = k - 4 = k.$$

4.3 Conjugations of Bäcklund and translations operators with reflections

Conjugation of Bäcklund transformations by reflections is described by the general formula (3.10) adopted to $N = 4$ case:

$$\widehat{P}_n s_m \widehat{P}_n = s_{P_n(m)}, \quad n = 2i, 2i + 1, \quad n, m = 1, 2, 3, 4 \quad (4.17)$$

where $P_n(m)$ is the point obtained from m by acting with the reflection P_n . Repeated conjugation yields:

$$\widehat{P}_{n+1}\widehat{P}_n s_m \widehat{P}_n \widehat{P}_{n+1} = s_{P_{n+1}P_n(m)} = s_{m+1}$$

which in view of relations (3.6) reproduces a basic relation $\pi s_m = s_{m+1}\pi$.

We are able to extend the above results to include conjugations by reflections of the translation operators. They are described by the following two relations :

$$\widehat{P}_n T_i^{-1} \widehat{P}_n = T_{P_{n+1}(i)}, \quad \widehat{P}_n T_i \widehat{P}_n = T_{P_{n+1}(i)}^{-1}. \quad (4.18)$$

Several relevant for $N = 4$ examples that follow from the above identity are listed below:

$$\widehat{P}_2 T_2^{-1} \widehat{P}_2 = T_1, \quad \widehat{P}_2 T_4 \widehat{P}_2 = T_3^{-1}, \quad (4.19)$$

$$\widehat{P}_0 T_2^{-1} \widehat{P}_0 = T_3, \quad \widehat{P}_0 T_4 \widehat{P}_0 = T_1^{-1}. \quad (4.20)$$

The transformations of translation operators by odd reflections are as follows :

$$\widehat{P}_1 T_2^{-1} \widehat{P}_1 = T_4, \quad \widehat{P}_1 T_4 \widehat{P}_1 = T_2^{-1}, \quad (4.21)$$

for \widehat{P}_1 reflection and

$$\widehat{P}_3 T_2^{-1} \widehat{P}_3 = T_2, \quad \widehat{P}_3 T_4 \widehat{P}_3 = T_4^{-1}, \quad (4.22)$$

for \widehat{P}_3 reflection. The first of these results given in relation (4.21) is the duality map: $k \leftrightarrow m$ that also maps $x \rightarrow -x$ and $\mathbf{a} \rightarrow 2 - \mathbf{a}$, since $\alpha_1 \leftrightarrow \alpha_4$.

5 Closed expressions for solutions on the orbit of translation operators.

In this section we find a closed expression for the general solution $\mathbb{T}(k, m; \mathbf{a})$ introduced in relation (4.12). We start by first considering the two simpler cases of T_4 orbit and the T_2 orbits.

5.1 The T_4 and the T_2 orbits and Kummer polynomials

The T_4 orbit and the T_2 orbits included in the definition of (4.12) are described as :

1. the $k = 0$ case that maintains $G = 0$, while it transforms F by T_4 ,
2. the $m = 0$ case that maintains $F = 1$ and transforms G by T_2 .

The T_4 orbit:

$$\mathbb{T}(0, m; \mathbf{a}) = T_4^m |F = 1, G = 0\rangle_{\alpha_{\mathbf{a}}} = (F_m(x, \mathbf{a}), G_m(x, \mathbf{a}) = 0),$$

is governed by the recurrence relation

$$F_m = 1 + \frac{xm}{xF_{m-1} + \mathbf{a}/2 - m}, \quad (5.1)$$

obtained from the T_4 transformation rule

$$\begin{aligned} T_4(F) &= 1 - G + (\gamma_1 + \gamma_4)/(F - \gamma_4/(1 - G)), \\ T_4(G) &= F - \gamma_4/(1 - G) + (\gamma_1 + \gamma_2 + \gamma_4)/(G - (\gamma_1 + \gamma_4)/(F - \gamma_4/(1 - G))). \end{aligned} \quad (5.2)$$

The above recurrence relation for F_m has a solution :

$$F_m = \frac{N_m(x, \mathbf{a} + 2)}{N_m(x, \mathbf{a})}, \quad (5.3)$$

in terms of Kummer polynomials $N_m(x, \mathbf{a})$:

$$N_n(x, \mathbf{a} + 2) = 2^n x^n (-x)^{-\mathbf{a}/2} U(-\frac{\mathbf{a}}{2}, -\frac{\mathbf{a}}{2} + n + 1, x), \quad (5.4)$$

where

$$U(b, b + n + 1, x) = x^{-b} \sum_{s=0}^{s=n} \binom{k}{s} (b)_s x^{-s}, \quad (5.5)$$

is a Kummer polynomial $U(a, b, x)$ in x of degree n when $a - b + 1 = -n$, $n \in \mathbb{Z}_+$. Generally $U(a, b, x)$ solves Kummer's equation:

$$x \frac{d^2 w}{dx^2} + (b - x) \frac{dw}{dx} - aw = 0.$$

Starting with $N_0(x, \mathbf{a}) = 1$ and applying the first of the basic two recurrence relations:

$$N_{m+1}(x, \mathbf{a}) = 2xN_m(x, \mathbf{a}) + (\mathbf{a} - 2)N_m(x, \mathbf{a} - 2), \quad (5.6)$$

$$2mN_{m-1}(x, \mathbf{a}) = N_m(x, \mathbf{a} + 2) - N_m(x, \mathbf{a}) = \frac{dN_m(x, \mathbf{a})}{dx} \quad m = 0, 1, 2, \dots, \quad (5.7)$$

that can be derived from relation (5.4), one first obtains $N_1(x, \mathbf{a}) = 2x + \mathbf{a} - 2$ and eventually arrives at a general expression:

$$N_m(x, \mathbf{a}) = \sum_{p=0}^{p=m} \binom{m}{p} (2x)^p (\mathbf{a} - 2)(\mathbf{a} - 4) \cdots (\mathbf{a} - 2(m - p)). \quad (5.8)$$

For the T_2^{-1} orbit:

$$\mathbb{T}(k, 0; \mathbf{a}) = T_2^{-k} |F = 1, G = 0\rangle_{\alpha_{\mathbf{a}}}, \quad (5.9)$$

we obtain from expression for the action by T_2^{-1} :

$$T_2^{-1}(F, G) = \left(G - \frac{\gamma_1}{F} + \frac{1/x - \gamma_2}{1 - F + \frac{\gamma_1 + \gamma_4}{1 - G + \frac{\gamma_1}{F}}}, 1 - F + \frac{\gamma_1 + \gamma_4}{1 - G + \frac{\gamma_1}{F}} \right), \quad (5.10)$$

a chain of transformations that results in a recurrence relation:

$$G^k = \frac{2k}{2x(1 - G^{k-1}) + (a + 2k - 2)}. \quad (5.11)$$

The solution is given this time by

$$G_k = 1 - \frac{R_k(x, a - 2)}{R_k(x, a)} = \frac{2kR_{k-1}(x, a)}{R_k(x, a)}, \quad (5.12)$$

in terms of Kummer polynomials:

$$\begin{aligned} R_n(x, a) &= \sum_{p=0}^{p=n} \binom{n}{p} (2x)^p a(a+2)(a+4) \cdots (a+2(n-p-1)) \\ &= 2^n x^n x^{a/2} U\left(\frac{a}{2}, \frac{a}{2} + n + 1, x\right), \end{aligned} \quad (5.13)$$

The above expression can alternatively be obtained from the initial condition $R_0(x, a) = 1$ by applying the recursion relation (5.14) from the two basic recursive relations satisfied by polynomials $R_k(x, a)$:

$$R_{k+1}(x, a) = 2xR_k(x, a) + aR_k(x, a + 2), \quad (5.14)$$

$$2kR_{k-1}(x, a) = R_k(x, a) - R_k(x, a - 2) = \frac{dR_k(x, a)}{dx}. \quad (5.15)$$

Polynomials denoted by $N_k(x, a)$ for the T_4 -orbit and $R_k(x, a)$ for the T_2 -orbit, are both of order k , and are related to each other through relation:

$$R_k(-x, a) = (-1)^k N_k(x, 2 - a). \quad (5.16)$$

that is consistent with P_1 reflection that induces $x \rightarrow -x$, $F \rightarrow 1 - G$, $G \rightarrow 1 - F$ and $\alpha_1 \leftrightarrow \alpha_4$, $\alpha_2 \leftrightarrow \alpha_3$. For α_i given in relation (4.3) this is equivalent to

$$\widehat{P}_1 : k \leftrightarrow m, a \leftrightarrow 2 - a, x \leftrightarrow -x. \quad (5.17)$$

5.2 A General Case of $T_4^m T_2^{-k}$ orbit and the generalized higher-type Kummer polynomials

We now consider the full orbit given as in (4.12) by :

$$T_2^{-k} T_4^m |F = 1, G = 0\rangle_{\alpha_a} = (F_m^k(x, a), G_m^k(x, a)), \quad k, m \in \mathbb{Z}_+.$$

Here $F_m^k(x, a)$, $G_m^k(x, a)$ are solutions to equations (4.2)

$$\begin{aligned} \frac{d}{dx} F_m^k &= F_m^k (F_m^k - 1) (2G_m^k - 1) - \frac{a + 2k - 2m}{2x} F_m^k + \frac{a + 2k}{2x}, \\ \frac{d}{dx} G_m^k &= -G_m^k (G_m^k - 1) (2F_m^k - 1) + \frac{a + 2k - 2m - 2}{2x} G_m^k + \frac{-2k}{2x}, \end{aligned} \quad (5.18)$$

with parameters given in (4.3).

See also the reference [10] for use of the translation operators to obtain solution of Painlevé V equation in terms of Laguerre polynomials and [19] for a more recent discussion.

Here we are able to find the Kummer polynomial representation for both F_m^k and G_m^k as given in the main result of this section by the following expression:

$$\begin{aligned} F_m^k(x, a) &= \frac{R_{mk}^{(m-1)}(x, a - 2) R_{m(k+1)}^{(m-1)}(x, a)}{R_{mk}^{(m-1)}(x, a) R_{m(k+1)}^{(m-1)}(x, a - 2)}, \\ G_m^k(x, a) &= 2k \frac{R_{m(k+1)}^{(m-1)}(x, a - 2) R_{(m+1)(k-1)}^{(m)}(x, a)}{R_{mk}^{(m-1)}(x, a - 2) R_{(m+1)k}^{(m)}(x, a)}, \quad m, k = 0, 1, 2, \dots \end{aligned} \quad (5.19)$$

The symbols $R_n^{(0)}(x, \mathbf{a}) = R_n(x, \mathbf{a})$ are regular Kummer polynomials of the n -th order defined in relation (5.13). The symbols $R_n^{(m)}(x, \mathbf{a})$ are generalized Kummer polynomials of the n -th order of the m -th type. They are obtained from Kummer polynomials by a set of recurrence relations :

$$R_{(m-1)(k+1)}^{(m-2)}(x, \mathbf{a} - 2)R_{(m+1)k}^{(m)}(x, \mathbf{a}) = \frac{1}{2m} [R_{mk}^{(m-1)}(x, \mathbf{a} - 2)R_{m(k+1)}^{(m-1)}(x, \mathbf{a}) - R_{mk}^{(m-1)}(x, \mathbf{a})R_{m(k+1)}^{(m-1)}(x, \mathbf{a} - 2)], \quad (5.20)$$

that are valid for $m = 1, 2, 3, \dots$ and $k = 0, 1, \dots$ with initial variables being $R_0^{(-1)}(x, \mathbf{a} - 2) = 1$ and $R_k^{(0)}(x, \mathbf{a}) = R_k(x, \mathbf{a})$. The master recurrence relation (5.20) gives rise to the determinant expressions for the mk order and the $m - 1$ -type Kummer-like polynomials $R_{mk}^{(m-1)}(x, \mathbf{a})$. Their determinant expressions are obtained by employing the Desnanot-Jacobi identity:

$$|M| \times |M_{1,n}^{1,n}| = |M_n^n| \times |M_1^1| - |M_1^n| \times |M_n^1| \quad (5.21)$$

which is valid for any a $(k + 1) \times (k + 1)$ matrix M . The notation in identity (5.21) is such that M_i^j is a matrix M with j -th row and i -th column removed. When the Desnanot-Jacobi identity is applied on the above recurrence relation (5.20) (up to a constant C_{m-1} determined below) one finds mk order and the $m - 1$ -type Kummer-like polynomial $R_{mk}^{(m-1)}(x, \mathbf{a})$ being the determinant of the $m \times m$ matrix M with matrix elements to be given by:

$$R_{mk}^{(m-1)}(x, \mathbf{a}) = C_{m-1}|M|, \quad M_{ij} = R_{k+i-1}(x, \mathbf{a} - 2(m - j)), \quad i, j = 1, \dots, m, \quad (5.22)$$

where $R_n(x, b)$ are the original Kummer polynomials (5.13) and the constants satisfy the relation:

$$C_m = \frac{1}{2m} \frac{C_{m-1}^2}{C_{m-2}}, \quad m = 2, 3, 4, \dots \quad (5.23)$$

It follows for the first few cases that

$$C_0 = 1, \quad C_1 = \frac{1}{2}, \quad C_2 = \frac{1}{2 \cdot 2} \left(\frac{1}{2}\right)^2 = \frac{1}{16}.$$

Generally, we find from relation (5.23):

$$C_m = \frac{1}{2^{m(m+1)/2}} \prod_{j=0}^{m-2} \frac{1}{(m-j)^{j+1}}, \quad m = 2, 3, 4, \dots \quad (5.24)$$

The value of C_m is such that the identity $R_0^{(m)}(x, \mathbf{a}) = 1, m = 0, 1, 2, \dots$ holds as follows from the above determinant expressions.

From the determinant expression (5.22) we find as a special example $R_{2k}^{(1)}(x, \mathbf{a})$ to be given by:

$$R_{2k}^{(1)}(x, \mathbf{a}) = \frac{1}{2} \begin{vmatrix} R_k(x, \mathbf{a} - 2) & R_k(x, \mathbf{a}) \\ R_{k+1}(x, \mathbf{a} - 2) & R_{k+1}(x, \mathbf{a}) \end{vmatrix} = \frac{1}{2} (R_2(x, \mathbf{a})R_1(x, \mathbf{a} - 2) - R_1(x, \mathbf{a})R_2(x, \mathbf{a} - 2)). \quad (5.25)$$

One can check explicitly that

$$R_2^{(1)}(x, \mathbf{a}) = N_2(x, \mathbf{a} + 2), \quad (5.26)$$

which is a special case of “duality” relation (5.32) to be introduced below and shown to play an important role in a general scheme illustrating the power of symmetry generated by \widehat{P}_1 automorphism.

Other basic examples of $R_{mk}^{(m-1)}$ are given by the following determinant expressions for $m = 3$ and $m = 4$:

$$R_{3k}^{(2)}(x, a) = \frac{1}{16} \begin{vmatrix} R_k(x, a-4) & R_k(x, a-2) & R_k(x, a) \\ R_{k+1}(x, a-4) & R_{k+1}(x, a-2) & R_{k+1}(x, a) \\ R_{k+2}(x, a-4) & R_{k+2}(x, a-2) & R_{k+2}(x, a) \end{vmatrix} \quad k = 1, 2, 3, \dots, \quad (5.27)$$

$$R_{4k}^{(3)}(x, a) = \frac{1}{3} \left(\frac{1}{16} \right)^2 \begin{vmatrix} R_k(x, a-2 \cdot 3) & R_k(x, a-2 \cdot 2) & R_k(x, a-2) & R_k(x, a) \\ R_{k+1}(x, a-2 \cdot 3) & R_{k+1}(x, a-2 \cdot 2) & R_{k+1}(x, a-2) & R_{k+1}(x, a) \\ R_{k+2}(x, a-2 \cdot 3) & R_{k+2}(x, a-2 \cdot 2) & R_{k+2}(x, a-2) & R_{k+2}(x, a) \\ R_{k+3}(x, a-2 \cdot 3) & R_{k+3}(x, a-2 \cdot 2) & R_{k+3}(x, a-2) & R_{k+3}(x, a) \end{vmatrix}. \quad (5.28)$$

It follows from the above determinant expressions that Kummer polynomials of the p -th kind and of the n -th order are described by a general formula:

$$R_n^{(p)}(x, a) = \sum_{k=0}^n \binom{n}{k} (2x)^k \left(a - a_1^{(n,k,p)} \right) \dots \left(a - a_{n-k}^{(n,k,p)} \right), \quad p \geq 0, n > 0, \quad R_0^{(p)}(x, a) = 1, \quad (5.29)$$

and thus fully defined by roots $a_i^{(n,k,p)}$ of the monomials for $i = 1, \dots, n-k$.

Using equation : $dR_n(x, a)/dx = R_n(x, a) - R_n(x, a-2)$ we find that

$$\begin{aligned} R_n(x, a-6) &= R_n(x, a) - 3R_n'(x, a) + 3R_n''(x, a) - R_n'''(x, a), \\ R_n(x, a-4) &= R_n(x, a) - 2R_n'(x, a) + R_n''(x, a), \quad R_n(x, a-2) = R_n(x, a) - R_n'(x, a), \end{aligned}$$

and accordingly the determinant in equation (5.28) can be rewritten as Wronskian of Kummer polynomials:

$$R_{4k}^{(3)}(x, a) = \frac{1}{3} \left(\frac{1}{16} \right)^2 \begin{vmatrix} R_k'''(x, a) & R_k''(x, a) & R_k'(x, a) & R_k(x, a) \\ R_{k+1}'''(x, a) & R_{k+1}''(x, a) & R_{k+1}'(x, a) & R_{k+1}(x, a) \\ R_{k+2}'''(x, a) & R_{k+2}''(x, a) & R_{k+2}'(x, a) & R_{k+2}(x, a) \\ R_{k+3}'''(x, a) & R_{k+3}''(x, a) & R_{k+3}'(x, a) & R_{k+3}(x, a) \end{vmatrix}, \quad (5.30)$$

with all polynomials now taken at the same point a .

There also exists a dual master recurrence relation:

$$\begin{aligned} &\frac{1}{2k} \left(R_{mk}^{(m-1)}(x, a-2) R_{(m+1)k}^{(m)}(x, a) - R_{mk}^{(m-1)}(x, a) R_{k(m+1)}^{(m)}(x, a-2) \right) \\ &= R_{m(k+1)}^{(m-1)}(x, a-2) R_{(k-1)(m+1)}^{(m)}(x, a). \end{aligned} \quad (5.31)$$

Applying the duality relation

$$R_{km}^{(m-1)}(x, a-2) = N_{km}^{(k-1)}(x, a), \quad (5.32)$$

that exchanges $m \leftrightarrow k$ and $a \rightarrow a-2$ and introduces the higher type order generalization of Kummer polynomials $N_k(x, a)$ defined in (5.4) we obtain the dual version of the recurrence relations (5.20) :

$$\begin{aligned} &\frac{1}{2k} \left(N_{mk}^{(k-1)}(x, a) N_{(m+1)k}^{(k-1)}(x, a+2) - N_{mk}^{(k-1)}(x, a+2) N_{k(m+1)}^{(k-1)}(x, a) \right) \\ &= N_{m(k+1)}^{(k)}(x, a) N_{(k-1)(m+1)}^{(k-2)}(x, a+2), \end{aligned} \quad (5.33)$$

that via Desnanot-Jacobi identity (5.21) leads to expression for the generalized higher type Kummer polynomials $N_k(x, a)$:

$$N_{km}^{(k-1)}(x, a) = C_{k-1} |M|, \quad M_{ij} = N_{m+i-1}(x, a+2(j-1)), \quad (5.34)$$

where the normalization constant C_{k-1} is defined in (5.23). For example

$$N_{3m}^{(2)}(x, (x, \mathbf{a})) = \frac{1}{16} \begin{vmatrix} N_m(x, \mathbf{a}) & N_m(x, \mathbf{a} + 2) & N_m(x, \mathbf{a} + 4) \\ N_{m+1}(x, \mathbf{a}) & N_{m+1}(x, \mathbf{a} + 2) & N_{m+1}(x, \mathbf{a} + 4) \\ N_{m+2}(x, \mathbf{a}) & N_{m+2}(x, \mathbf{a} + 2) & N_{m+2}(x, \mathbf{a} + 4) \end{vmatrix} \quad m = 1, 2, 3, \dots \quad (5.35)$$

Thus we have determinant expressions (5.22) and (5.34) that are dual to each other according to the duality relation (5.32) that can also be viewed as generated by \widehat{P}_1 transformation $m \leftrightarrow k, x \rightarrow -x, \mathbf{a} \rightarrow 2 - \mathbf{a}$. In view of the duality relations (5.32) we have the following relation

$$R_{km}^{(m-1)}(-x, -\mathbf{a}) = (-1)^{km} R_{km}^{(k-1)}(x, \mathbf{a}). \quad (5.36)$$

To understand better relation (5.36) it is illustrative to introduce notation $R_{m,k}(x, \mathbf{a}) = R_{km}^{(m-1)}(x, \mathbf{a})$. In this new notation the relation (5.36) describes a condition for commutation of indices $m, k \rightarrow k, m : R_{k,m}(x, \mathbf{a}) = (-1)^{km} R_{m,k}(-x, -\mathbf{a})$.

Also, thanks to duality relation (5.32), we have an alternative expression for our solutions as :

$$\begin{aligned} F_m^k(x, \mathbf{a}) &= \frac{N_{mk}^{(k-1)}(x, \mathbf{a}) N_{m(k+1)}^{(k)}(x, \mathbf{a} + 2)}{N_{mk}^{(k-1)}(x, \mathbf{a} + 2) N_{m(k+1)}^{(k)}(x, \mathbf{a})}, \\ G_m^k(x, \mathbf{a}) &= 2k \frac{N_{m(k+1)}^{(k)}(x, \mathbf{a}) N_{(k-1)(m+1)}^{(k-2)}(x, \mathbf{a} + 2)}{N_{mk}^{(k-1)}(x, \mathbf{a}) N_{(m+1)k}^{(k-1)}(x, \mathbf{a} + 2)}. \end{aligned} \quad (5.37)$$

6 Even $\widehat{P}_2, \widehat{P}_4$ reflections and degeneracy

General remark that applies to automorphisms $\widehat{P}_4, \widehat{P}_2, \widehat{P}_3$ is that implementing them requires enforcing the constant $a/2$ to be an integer, chosen here to be $l \in \mathbb{Z}$.

6.1 Action of the \widehat{P}_2 automorphism

For \widehat{P}_2 the transformation $\alpha_2 \leftrightarrow \alpha_4$ for $\alpha_2 = -2k$ and $\alpha_4 = 2 - \mathbf{a} + 2m = 2(1 - l + m)$ is equivalent to that of

$$l \rightarrow 1 + m + k, \quad k \rightarrow l - 1 - m,$$

for $l > m$.

Accordingly \widehat{P}_2 transformations are such that

$$\begin{aligned} \widehat{P}_2(F_m^k(x, l)) &= F_m^{l-1-m}(-x, 1 + m + k) \\ \widehat{P}_2(G_m^k(x, l)) &= G_m^{l-1-m}(-x, 1 + m + k), \quad \mathbf{a} = 2l \end{aligned}$$

and are expected to be solutions to equations (4.2) with the same parameters as solutions $F_m^k(x, \mathbf{a}), 1 - G_m^k(x, \mathbf{a})$. Are these functions equal to the original solutions $F_m^k(x, \mathbf{a}), 1 - G_m^k(x, \mathbf{a})$ or are they degenerated solutions is the main question. We will fully answer it in the subsection 6.3 but first we will discuss how it works on an explicit example:

Example 6.1. Here we calculate a solution $T_2^{-k} T_4^m |F = 1, G = 0\rangle_{\alpha_{\mathbf{a}=4}}$ using the expression for solutions found in equation (5.19) for $k = m = 1$ with $l = 2$ ($\mathbf{a} = 2l = 4$):

$$\begin{aligned} F_1^1(x, \mathbf{a} = 4) &= \frac{R_1(x, \mathbf{a} - 2) R_2(x, \mathbf{a})}{R_1(x, \mathbf{a}) R_2(x, \mathbf{a} - 2)} \Big|_{\mathbf{a}=4} = \frac{(2x + 2)((2x)^2 + 16x + 24)}{(2x + 4)((2x)^2 + 8x + 8)} \\ G_1^1(x, \mathbf{a} = 4) &= 2 \frac{R_2(x, \mathbf{a} - 2) R_0^{(1)}(x, \mathbf{a})}{R_1(x, \mathbf{a} - 2) R_2^{(1)}(x, \mathbf{a})} \Big|_{\mathbf{a}=4} = 2 \frac{R_2(x, \mathbf{a} - 2)}{R_1(x, \mathbf{a} - 2) N_2(x, \mathbf{a} + 2)} \Big|_{\mathbf{a}=4} \\ &= 2 \frac{((2x)^2 + 8x + 8)}{(2x + 2)((2x)^2 + 16x + 8)}, \end{aligned} \quad (6.1)$$

where we used the identities (5.26) and $R_0^{(m)}(x, \mathbf{a}) = 1$. The above two solutions in relations (6.1) satisfy the equations (5.18) with $k = m = 1$, ($\alpha_1 = \mathbf{a} + 2 = 6, \alpha_2 = -2, \alpha_3 = -2$). We write these parameters as:

$$\alpha_i = 2(k + l, -k, -m, 1 - l + m) = 2(3, -1, -1, 0), \quad (6.2)$$

Under the \widehat{P}_2 transformation the parameter m is invariant and we have the following expressions for the \mathbf{a} , k transformations :

$$\begin{aligned} \bar{\mathbf{a}} &= \mathbf{a} + 2(1 + k + m - l) = 4 + 2 = 6 \\ \bar{k} &= k - (1 + k + m - l) = 1 - 1 = 0 \end{aligned}$$

Thus we are looking for the transformed solutions given by:

$$\begin{aligned} F_m^{\bar{k}}(-x, \bar{\mathbf{a}}/2) &= F_1^0(-x, 6), \\ G_m^{\bar{k}}(-x, \bar{\mathbf{a}}/2) &= G_1^0(-x, 6), \end{aligned}$$

which we calculate using equation (5.19) to obtain:

$$\begin{aligned} F_1^0(-x, 6) &= \frac{R_0(-x, 4)R_1(-x, 6)}{R_0(-x, 6)R_1(-x, 4)} = \frac{6 - 2x}{4 - 2x} = \frac{x - 3}{x - 2}, \\ G_1^0(-x, 6) &= 2 \cdot 0 \cdot \frac{R_1(-x, 4)R_{-2}^{(1)}(-x, 6)}{R_0(-x, 4)R_0^{(1)}(-x, 6)} = 0. \end{aligned} \quad (6.3)$$

One checks explicitly that $F = (x - 3)/(x - 2)$ and $G = 1$ solve the equation (4.2) with $\alpha_1 = 6, \alpha_2 = -2, \alpha_3 = -2$ that agrees with the parameters (6.2) of the original solution. We will see in subsection 6.3 that this degeneracy is a general feature of the even \widehat{P}_{2i} automorphisms.

Generally we can rewrite

$$\begin{aligned} F_m^{l-1-m}(-x, 2(1 + m + k)) &= \frac{R_{m(l-1-m)}^{(m-1)}(-x, 2(k + m))R_{m(l-m)}^{(m-1)}(-x, 2(1 + m + k))}{R_{m(l-1-m)}^{(m-1)}(-x, 2(1 + m + k))R_{m(l-m)}^{(m-1)}(-x, 2(k + m))} \\ &= \frac{N_{m(l-1-m)}^{(m-1)}(x, 2(1 - k - m))N_{m(l-m)}^{(m-1)}(x, -2(m + k))}{N_{m(l-1-m)}^{(m-1)}(x, -2(m + k))N_{m(l-m)}^{(m-1)}(x, 2(1 - k - m))} \end{aligned} \quad (6.4)$$

where we used the identity

$$R_{mk}^{(m-1)}(x, \mathbf{a}) = (-1)^{mk} N_{mk}^{(m-1)}(-x, 2 - \mathbf{a}), \quad R_{mk}^{(m-1)}(-x, \mathbf{a}) = (-1)^{mk} N_{mk}^{(m-1)}(x, 2 - \mathbf{a}), \quad (6.5)$$

Expression (6.4) describes a closed expression for the degenerated partner of :

$$F_m^k(x, 2l) = \frac{R_{mk}^{(m-1)}(x, 2(l - 1))R_{m(k+1)}^{(m-1)}(x, 2l)}{R_{mk}^{(m-1)}(x, 2l)R_{m(k+1)}^{(m-1)}(x, 2(l - 1))},$$

which is a solution given in (5.19) for $m, k, a = 2l$.

For the the remaining part of solution we find that the degenerated solution is:

$$\begin{aligned} &G_m^{l-1-m}(-x, 2(1 + m + k)) \\ &= 2(l - 1 - m) \frac{R_{m(l-m)}^{(m-1)}(-x, 2(m + k))R_{(l-m-2)(m+1)}^{(m)}(-x, 2(1 + m + k))}{R_{m(l-1-m)}^{(m-1)}(-x, 2(m + k))R_{(m+1)(l-1-m)}^{(m)}(-x, 2(1 + m + k))} \\ &= -2(l - 1 - m) \frac{N_{m(l-m)}^{(m-1)}(x, 2(1 - m - k))N_{(l-m-2)(m+1)}^{(m)}(x, -2(m + k))}{N_{m(l-1-m)}^{(m-1)}(x, 2(1 - m - k))N_{(m+1)(l-1-m)}^{(m)}(x, -2(m + k))}, \end{aligned} \quad (6.6)$$

where we again used equation (6.5).

Remember that $\widehat{P}_2(G) = 1 - G_m^{l-1-m}(-x, 2(1+m+k))$. Thus

$$\widehat{P}_2(G) = 1 + 2(l-1-m) \frac{N_{m(l-m)}^{(m-1)}(x, 2(1-m-k))N_{(l-m-2)(m+1)}^{(m)}(x, -2(m+k))}{N_{m(l-1-m)}^{(m-1)}(x, 2(1-m-k))N_{(m+1)(l-1-m)}^{(m)}(x, -2(m+k))}. \quad (6.7)$$

Together with (6.4), formula (6.7) solves (4.2) with the identical parameters $\alpha_i = 2(l+k, -k, -m, 1-l+m)$ as the original solution (5.19) with $\mathbf{a} = 2l$.

Example 6.2. We consider the case of

$$k = 1, m = 1, l = 4, a = 2 \cdot l = 8 \rightarrow \alpha_i = 2(l+k, -k, -m, 1-l+m) = 2(5, -1, -1, -2), \quad (6.8)$$

We find $\bar{k} = l - 1 - m = 2$, $\bar{l} = (1 + m + k) = 3$. Applying the formulas (6.4) and (6.7) we obtain the \widehat{P}_2 transformed degenerated solutions :

$$\begin{aligned} F_1^2(-x, 2 \cdot 3) &= \frac{N_2(x, -2 \cdot 1)N_3(x, -2 \cdot 2)}{N_2(x, -2 \cdot 2)N_3(x, -2 \cdot 1)} \\ &= \frac{(x^2 - 4x + 6)(x^3 - 9x^2 + 36x - 60)}{(x^2 - 6x + 12)(x^3 - 6x^2 + 18x - 24)} \\ 1 - G_1^2(-x, 2 \cdot 3) &= 1 - (-2 \cdot 2) \frac{N_3(x, -2)N_2^{(1)}(x, -4)}{N_2(x, -2)N_4^{(1)}(x, -4)} \\ &= \frac{(x^2 - 6x + 12)(x^4 - 8x^3 + 24x^2 - 24x + 12)}{(x^2 - 4x + 6)(72 + 54x^2 + x^4 - 12x^3 - 96x)}. \end{aligned}$$

Inserting $x = -z^2/2$ and multiplying by z we obtain from the above two results the expressions (3.15) for $q(z), p(z)$ of Example 3.1 of [20] that also solve the relevant Painlevé with the parameters $\alpha_i = 2(5, -1, -1, -2)$.

6.2 Action of the \widehat{P}_4 automorphism

The \widehat{P}_4 generated transformation $\alpha_1 \leftrightarrow \alpha_3$ for $\alpha_1 = 2(l+k)$ and $\alpha_3 = -2m$ is equivalent to that of

$$l \rightarrow -k - m, \quad k \rightarrow k, \quad m \rightarrow -l - k.$$

Thus it must hold $-l - k \geq 0$ or $l \leq -k$.

Accordingly, we obtain:

$$\begin{aligned} \widehat{P}_4(F_m^k(x, 2l)) &= F_{-l-k}^k(-x, 2(-k-m)), \\ \widehat{P}_4(G_m^k(x, 2l)) &= G_{-l-k}^k(-x, 2(-k-m)) \end{aligned} \quad (6.9)$$

The first of equations (6.9) yields after substituting the relation (5.19) for $G_m^k(x, \mathbf{a})$:

$$\begin{aligned} G_{-l-k}^k(-x, -k-m) &= 2k \frac{R_{(-l-k)(k+1)}^{(-l-k-1)}(-x, 2(-k-m-1))R_{(-l-k+1)(k-1)}^{(-l-k)}(-x, 2(-k-m))}{R_{(-l-k)k}^{(-l-k-1)}(-x, 2(-k-m-1))R_{(-l-k+1)k}^{(-l-k)}(-x, 2(-k-m))} \\ &= -2k \frac{N_{(-l-k)(k+1)}^{(-l-k-1)}(x, 2(2+k+m))N_{(-l-k+1)(k-1)}^{(-l-k)}(x, 2(1+k+m))}{N_{(-l-k)k}^{(-l-k-1)}(x, 2(2+k+m))N_{(-l-k+1)k}^{(-l-k)}(x, 2(1+k+m))} \end{aligned}$$

From $\widehat{P}_4(F) = F_{-l-k}^k(-x, -2(k+m))$ we get

$$\begin{aligned} F_{-l-k}^k(-x, -2(k+m)) &= \frac{R_{(-l-k)k}^{(-l-k-1)}(-x, -2(1+k+m))R_{(-l-k)(k+1)}^{(-l-k-1)}(-x, -2(k+m))}{R_{(-l-k)k}^{(-l-k-1)}(-x, -2(k+m))R_{(l-k)(k+1)}^{(-l-k-1)}(-x, -2(1+k+m))} \\ &= \frac{N_{(-l-k)k}^{(-l-k-1)}(x, 2(2+k+m))N_{(-l-k)(k+1)}^{(-l-k-1)}(x, 2(1+k+m))}{N_{(-l-k)k}^{(-l-k-1)}(x, 2(1+k+m))N_{(l-k)(k+1)}^{(-l-k-1)}(x, 2(2+k+m))} \end{aligned}$$

We will apply these results in subsection 6.3 and Example 6.4, where we will study degeneracy introduced by even reflection automorphisms and answer the question whether the results for $\widehat{P}_4(F_m^k(x, 2l))$ and $\widehat{P}_4(G_m^k(x, 2l))$ are equal to $1 - F_m^k(x, 2l)$ and $G_m^k(x, 2l)$ that share the same parameters.

6.3 Degeneracy induced by even reflections \widehat{P}_2 and \widehat{P}_4

We now present a general discussion on $\widehat{P}_4, \widehat{P}_2$ reflections that acting on solutions $T_2^{-k}T_4^m|g_1 = 1, g_2 = 0\rangle_a$ produce degenerated solutions (solutions that share the same parameters α_i as original solutions).

The result will follow if we are able to show the following equalities:

$$\widehat{P}_2 T_2^{-k_3} T_4^{m_3} |g_1 = 1, g_2 = 0\rangle_{a_3} = \pi s_1 T_2^{-k_1} T_4^{m_1} |g_1 = 1, g_2 = 0\rangle_{a_1}, \quad (6.10)$$

$$\widehat{P}_4 T_2^{-k_4} T_4^{m_4} |g_1 = 1, g_2 = 0\rangle_{a_4} = \pi^{-1} s_4 T_2^{-k_2} T_4^{m_2} |g_1 = 1, g_2 = 0\rangle_{a_2}, \quad (6.11)$$

for some values of k_i, m_i and a_i , since we have previously established in [20] that both solutions on the right hand sides are different from $T_2^{-k}T_4^m|g_1 = 1, g_2 = 0\rangle_a$, while they share the identical parameters.

First, we will show that the two relations given in equations (6.10) and (6.11) are equivalent. Assume that equation (6.10) holds and multiply the equation (6.10) by \widehat{P}_2 . Due to relation $\pi = \widehat{P}_2 \widehat{P}_1$ we obtain that equation (6.10) is equivalent to:

$$\begin{aligned} T_2^{-k_3} T_4^{m_3} |g_1 = 1, g_2 = 0\rangle_{a_3} &= \widehat{P}_1 s_1 T_2^{-k_1} T_4^{m_1} |g_1 = 1, g_2 = 0\rangle_{a_1} \\ &= s_4 \widehat{P}_1 T_2^{-k_1} T_4^{m_1} |g_1 = 1, g_2 = 0\rangle_{a_1} \end{aligned}$$

where in the last equation we used that $\widehat{P}_1 s_1 \widehat{P}_1 = s_4$. Multiplying both sides by s_4 and then using that $\widehat{P}_1 \widehat{P}_2 = \pi^{-1}$ we get :

$$\begin{aligned} \pi^{-1} s_4 \widehat{P}_2 T_2^{-k_3} T_4^{m_3} |g_1 = 1, g_2 = 0\rangle_{a_3} &= \widehat{P}_1 \widehat{P}_2 \widehat{P}_1 s_1 T_2^{-k_1} T_4^{m_1} |g_1 = 1, g_2 = 0\rangle_{a_1} \\ &= \widehat{P}_4 T_2^{-k_1} T_4^{m_1} |g_1 = 1, g_2 = 0\rangle_{a_1}, \end{aligned}$$

where we used the identity $\widehat{P}_1 \widehat{P}_2 \widehat{P}_1 = \widehat{P}_4$ that follows from (3.8). Thus equation (6.11) follows from equation (6.10) and we will only need to show that one of these two equations holds.

We now embark on a proof of equation (6.10). First, we recall that

$$\begin{aligned} \widehat{P}_2 |g_1 = 1, g_3 = 0, g_2 = 0, g_4 = 1\rangle &= |g_1 = 1, g_3 = 0, g_4 = 0, g_2 = 1\rangle, \\ \widehat{P}_2(a, 0, 0, 2 - a) &= (a, 2 - a, 0, 0), \end{aligned}$$

then using (4.19) we get

$$\begin{aligned} \widehat{P}_2 T_2^{-k_3} T_4^{m_3} |g_1 = 1, g_3 = 0, g_2 = 0, g_4 = 1\rangle_{a_3} \\ = T_1^{k_3} T_3^{-m_3} |g_1 = 1, g_3 = 0, g_4 = 0, g_2 = 1\rangle_{(a_3, 2 - a_3, 0, 0)}. \end{aligned} \quad (6.12)$$

Next we consider parameters of solution (6.12) with k_3, m_3 and a_3 :

$$T_1^{k_3} T_3^{-m_3}(a_3, 2 - a_3, 0, 0) = (a_3 + 2k_3, 2 - a_3 + 2m_3, -2m_3, -2k_3),$$

that will agree with the parameters $(a + 2k, -2k, -2m, 2 - a + 2m)$ of the solution $T_2^{-k}T_4^m|g_1 = 1, g_3 = 0, g_2 = 0, g_4 = 1\rangle$ if $a = 2l$, $a_3 = 2l_3$ and if it holds that

$$k_3 = l - m - 1, \quad l_3 = 1 + m + k, \quad m_3 = m. \quad (6.13)$$

We will see that for these values we have a degeneracy.

Let us compare the two sides of expressions (6.10). Since we have an identity $\pi = \widehat{P}_2 \widehat{P}_1$ we can rewrite the equation (6.10) as

$$\widehat{P}_1 s_1 T_2^{-k_1} T_4^{m_1} |g_1 = 1, g_2 = 0\rangle_{a_1} = T_2^{-k_3} T_4^{m_3} |g_1 = 1, g_2 = 0\rangle_{a_3},$$

or

$$s_1 T_2^{-k_1} T_4^{m_1} |g_1 = 1, g_2 = 0\rangle_{a_1} = \widehat{P}_1 T_2^{-k_3} T_4^{m_3} |g_1 = 1, g_2 = 0\rangle_{a_3} = T_4^{k_3} T_2^{-m_3} \widehat{P}_1 |g_1 = 1, g_2 = 0\rangle_{a_3}. \quad (6.14)$$

The left hand side of equation (6.14) can be rewritten as

$$\begin{aligned} T_1^{-k_1} T_4^{m_1} s_1 |g_1 = 1, g_2 = 0\rangle_{a_1} &= T_1^{-k_1} T_4^{m_1} s_1 T_1^{a_1/2} |g_1 = 1, g_2 = 0\rangle_{a=0} \\ &= T_1^{-k_1} T_4^{m_1} T_2^{a_1/2} s_1 |g_1 = 1, g_2 = 0\rangle_{a=0} = T_1^{-k_1} T_4^{m_1} T_2^{a_1/2} |g_1 = 1, g_2 = 0\rangle_{(0,0,0,2)} \\ &= T_1^{-k_1-1} T_4^{m_1} T_2^{a_1/2} T_1 |g_1 = 1, g_2 = 0\rangle_{(0,0,0,2)}, \end{aligned} \quad (6.15)$$

where we used the relations $s_i T_i s_i = T_{i+1}$, $s_i T_j = T_j s_i$, $j \neq i, i+1$ that hold between the s_i transformations and the translation operators [18]. Also we used that s_1 is an identity when acting on $|g_1 = 1, g_2 = 0\rangle_{(0,0,0,2)}$.

On the right hand side of equation (6.14) we have

$$T_4^{k_3} T_2^{-m_3} \widehat{P}_1 |g_1 = 1, g_2 = 0\rangle_{a_3} = T_4^{k_3} T_2^{-m_3} T_1^{-a_3/2} |g_1 = 0, g_2 = 1\rangle_{(2,0,0,0)}, \quad (6.16)$$

where we used that $\widehat{P}_1 T_1 \widehat{P}_1 = T_1^{-1}$ and

$$\begin{aligned} \widehat{P}_1 (g_1 = 1, g_2 = 0, g_3 = 0, g_4 = 1) &= (g_4 = 1, g_3 = 0, g_2 = 0, g_1 = 1) = (1, 0, 0, 1), \\ \widehat{P}_1 (a, 0, 0, 2 - a) &= (2 - a, 0, 0, a). \end{aligned}$$

Comparing the powers of T_1, T_4, T_2 in expressions (6.15) and (6.16) we obtain three conditions:

$$l_3 = 1 + k_1, \quad m_1 = k_3, \quad l_1 = m_3, \quad (6.17)$$

with $a_i = 2l_i$, $i = 1, 3$. For these conditions being satisfied the relation (6.10) holds both for the parameters and the solutions provided that

$$T_1 |g_1 = 1, g_2 = 0\rangle_{(0,0,0,2)} = \widehat{P}_1 |g_1 = 1, g_2 = 0\rangle_{(0,0,0,2)} = |g_1 = 1, g_2 = 0\rangle_{(2,0,0,0)},$$

but since T_1 only changes the parameters but not the underlying F, G variables, this relation is an identity and therefore also the relation (6.10) holds. Comparing relations (6.13) and (6.17) we find a direct relation:

$$l - m - 1 = m_1, \quad k_1 = m + k, \quad m = l_1,$$

between the parameters k_1, m_1, l_1 and k, m, l . These are the exact values [20] for which it holds that solutions $\pi s_1 T_2^{-k_1} T_4^{m_1} |F = 1, G = 0\rangle_{\alpha_b}$ and $T_2^{-k} T_4^m |F = 1, G = 0\rangle_{\alpha_a}$ are different while the four parameters of the two solutions are equal [20]:

$$\pi s_1 (\alpha_{k_1, m_1; b}) = \alpha_{k, m; a}.$$

Example 6.3.

$$\widehat{P}_2 : (F, G) \rightarrow (\text{degenerate}(F), \text{degenerate}(G))$$

Let $m = 2$ and $k = 1$ and $l = 3$, the corresponding solution is obtained from relation (5.19) to be

$$\begin{aligned} T_2^{-1} T_4^2 |F = 1, G = 0\rangle_{\alpha_{a=6}} &= \left(F = \frac{(x^2 + 4x + 6)(x^3 + 9x^2 + 36x + 60)}{(x^2 + 6x + 12)(18x + 6x^2 + x^3 + 24)} \right. \\ &\left. , G = 2 \frac{(x^2 + 6x + 6)(18x + 6x^2 + x^3 + 24)}{(x^2 + 4x + 6)(12x^3 + 54x^2 + 96x + 72 + x^4)} \right), \end{aligned}$$

with $\alpha_i = 2(l+k, -k, -m, 1-l+m) = 2(4, -1, -2, 0) = 2\pi s_1(l_1+k_1, -k_1, -m_1, 1-l_1+m_1)$. The unique solution for l_1, k_1, m_1 is: $l_1 = -2, k_1 = 3, m_1 = 0$. We use the right hand side of relation (6.10) to write the degenerated solution as :

$$\begin{aligned} \pi s_1 T_2^{-3} |F = 1, G = 0\rangle_{\alpha_a = -4} &= \left(\frac{-8x + 12 + x^2}{-6x + 6 + x^2}, 1 \right), \\ \pi s_1 2(l_1 + k_1, -k_1, -m_1, 1 - l_1 + m_1) &= 2\pi s_1(1, -3, 0, 3) = 2(4, -1, -2, 0). \end{aligned} \quad (6.18)$$

We will now reproduce the degenerated solution (6.18) using the \widehat{P}_2 transposition and the left hand side of relation (6.10). Starting again as above with $m = 2, k = 1$ and $a = 6$ or $l = 3$ so that $\alpha_i = 2(l+k, -k, -m, 1-l+m) = 2(4, -1, -2, 0)$ with $l-1-m = 0$ and $F_m^{l-1-m}(-x, 2(1+m+k)) = F_2^0(-x, 2 \cdot 4)$:

$$\begin{aligned} F_2^0(-x, 2 \cdot 4) &= \frac{N_0^{(1)}(x, 2(-2))N_{2(3-2)}^{(1)}(x, -2(3))}{N_0^{(1)}(x, -2(3))N_{2(3-2)}^{(1)}(x, 2(-2))} = \frac{N_2^{(1)}(x, -6)}{N_2^{(1)}(x, -4)} \\ &= \frac{-32x + 48 + 4x^2}{-24x + 24 + 4x^2} = \frac{-8x + 12 + x^2}{-6x + 6 + x^2}. \end{aligned}$$

Since $l-1-m = 0$ then $G_m^{l-1-m}(-x, 2(1+m+k)) = 0$, but then the \widehat{P}_2 transformed solution is $\widehat{P}_2(G) = 1 - G_m^{l-1-m}(-x, 2(1+m+k))$ which is equal to 1.

The above functions $F_2^0(-x, 2 \cdot 4) = (-8x + 12 + x^2)/(-6x + 6 + x^2)$ and $\widehat{P}_2(G) = 1$ solve the Hamiltonian equations (4.2) with $\alpha_1 = 8, \alpha_2 = -2, \alpha_3 = -4$ coefficients so that again we obtain acting with \widehat{P}_2 automorphism a solution for the same parameters $\alpha_i = 2(4, -1, -2, 0)$ as original F_m^k, G_m^k solution.

Example 6.4. Here we illustrate how the action of \widehat{P}_4 generates degenerated solutions according to a simple prescription:

$$\widehat{P}_4 : (F, G) \rightarrow (\text{degenerate}(F), \text{degenerate}(G)).$$

Recall that as shown in subsection (6.2) we have

$$\widehat{P}_4 : \alpha_1 \leftrightarrow \alpha_3 \sim l \rightarrow -m - k \in \mathbb{Z}_+, \quad m \rightarrow -l - k \in \mathbb{Z}_+,$$

and

$$\begin{aligned} \widehat{P}_4(F_m^k(x, l)) &= 1 - F_{-l-k}^k(-x, -m - k), \\ \widehat{P}_4(G_m^k(x, l)) &= G_{-l-k}^k(-x, -m - k), \quad a = 2l. \end{aligned}$$

Accordingly, for $m = 1, k = 1, l = -1$ or $a = -2$ we obtain

$$\begin{aligned} F_1^1(x, -2) &= \frac{(x-2)^2 x}{(x-1)(x^2 - 4x + 2)} \xrightarrow{\widehat{P}_4} 1 - F_0^1(-x, -2) = 0, \\ G_1^1(x, -4) &= \frac{(x^2 - 4x + 2)}{(x-2)(x^2 - 2x + 2)} \xrightarrow{\widehat{P}_4} G_0^1(-x, -2) = -\frac{1}{x+2}. \end{aligned}$$

Both above results are solutions of the F, G equations (4.2) with the same parameters.

Although equations (6.10) and (6.11) follow from each other they hold for different values of the parameter l such that $a = 2l$. To show this we discuss the equality between parameters shared between the left hand side of equation (6.10) and the parameter of the solution $T_2^k T_4^m |g_1 = 1, g_2 = 0\rangle_a$. The common parameters are :

$$T_2^{-k} T_4^m(a, 0, 0, 2-a) = \widehat{P}_2 T_2^{-k_2} T_4^{m_2}(a_2, 0, 0, 2-a_2),$$

or

$$\begin{aligned}(l+k, -k, -m, 1-l+m) &= \widehat{P}_2(l_2+k_2, -k_2, -m_2, 1-l_2+m_2) \\ &= (l_2+k_2, 1-l_2+m_2, -m_2, -k_2),\end{aligned}$$

the solution to these conditions is

$$m_2 = m, l = k_1 + m + 1, l_2 = k + m_2 + 1 = k + m + 1,$$

which implies that both l and l_2 are positive.

$$l > m, l_2 > m.$$

Repeating the same analysis for the degeneracy connected with \widehat{P}_4 we start with equality

$$T_2^{-k}T_4^m(\mathbf{a}, 0, 0, 2-\mathbf{a}) = \widehat{P}_4T_2^{-k_0}T_4^{m_0}(\mathbf{a}_0, 0, 0, 2-\mathbf{a}_0),$$

that can be rewritten as

$$\begin{aligned}(l+k, -k, -m, 1-l+m) &= \widehat{P}_4(l_0+k_0, -k_0, -m_0, 1-l_0+m_0) \\ &= (-m_0, -k_0, l_0+k_0, 1-l_0+m_0).\end{aligned}$$

This time it is the k parameter which remains unchanged $k = k_0$ while the expressions for l, l_0 become $l = -k - m_0, l_0 = -k - m$. Thus the \widehat{P}_4 degeneracy occurs when both l, l_0 are negative and satisfy $l < -k_0, l < k$.

Thus to cover all cases of both positive and negative values of the parameter l one needs to consider degeneracies induced by both \widehat{P}_2 and \widehat{P}_4 .

7 The odd \widehat{P}_1 and \widehat{P}_3 automorphisms

7.1 The action of \widehat{P}_1 automorphism

Consider the two \widehat{P}_1 transformations

$$G_m^k(x, \mathbf{a}) \xrightarrow{\widehat{P}_1} G_k^m(-x, -\mathbf{a} + 2) = 1 - F_m^k(x, \mathbf{a}), \quad (7.1)$$

and

$$F_m^k(x, \mathbf{a}) \xrightarrow{\widehat{P}_1} F_k^m(-x, -\mathbf{a} + 2) = 1 - G_m^k(x, \mathbf{a}), \quad (7.2)$$

where we took into consideration that \widehat{P}_1 transforms $x \rightarrow -x, a \rightarrow 2 - a$ and $m \leftrightarrow k$ (which is equivalent to $\alpha_1 \leftrightarrow \alpha_4, \alpha_2 \leftrightarrow \alpha_3$). Simultaneously, we have $\widehat{P}_1(F) = 1 - G$ and $\widehat{P}_1(G) = 1 - F$ that we inserted on the right hand sides.

Using identities (5.36) from subsection 5.2 we find for $G_k^m(-x, -\mathbf{a} + 2)$ from (5.19) using relation (5.36):

$$\begin{aligned}G_k^m(-x, -\mathbf{a} + 2) &= -2m \frac{R_{k(m+1)}^{(m)}(x, \mathbf{a})R_{(k+1)(m-1)}^{(m-2)}(x, \mathbf{a} - 2)}{R_{mk}^{(m-1)}(x, \mathbf{a})R_{(k+1)m}^{(m-1)}(x, \mathbf{a} - 2)} \\ &= 1 - F_m^k(x, \mathbf{a}) = 1 - \frac{R_{mk}^{(m-1)}(x, \mathbf{a} - 2)R_{m(k+1)}^{(m-1)}(x, \mathbf{a})}{R_{mk}^{(m-1)}(x, \mathbf{a})R_{m(k+1)}^{(m-1)}(x, \mathbf{a} - 2)},\end{aligned} \quad (7.3)$$

or after multiplying both sides by $R_{km}^{(m-1)}(x, \mathbf{a})/R_{m(k+1)}^{(m-1)}(x, \mathbf{a})$:

$$\begin{aligned}&\frac{R_{(m-1)(k+1)}^{(m-2)}(x, \mathbf{a} - 2)R_{(m+1)k}^{(m)}(x, \mathbf{a})}{R_{m(k+1)}^{(m-1)}(x, \mathbf{a})R_{m(k+1)}^{(m-1)}(x, \mathbf{a} - 2)} \\ &= \frac{1}{2m} \left(\frac{R_{mk}^{(m-1)}(x, \mathbf{a} - 2)}{R_{m(k+1)}^{(m-1)}(x, \mathbf{a} - 2)} - \frac{R_{km}^{(m-1)}(x, \mathbf{a})}{R_{m(k+1)}^{(m-1)}(x, \mathbf{a})} \right),\end{aligned} \quad (7.4)$$

in which we recognize the master recurrence relation (5.20) !

Looking back on the remaining relation in equation (7.2) it can be shown that it is equivalent to recurrence relation (5.33) dual to the master recurrence relation (5.20).

This completes the \widehat{P}_1 symmetry discussion showing how it nicely fits the structure of solutions we have generated and fully characterized by recursion relations. Thus the \widehat{P}_1 automorphism is verified as long as the fundamental recursion relations hold and its invariance can be considered a consistency check on the formalism.

7.2 The \widehat{P}_3 automorphism

The automorphism \widehat{P}_3 transforms $x \rightarrow -x$ and $\alpha_1 \leftrightarrow \alpha_2$ and $\alpha_3 \leftrightarrow \alpha_4$. For $\alpha_i = 2(l+k, -k, -m, 1-l+m)$ it follows that $\widehat{P}_3(\alpha_i) = 2(-k, l+k, 1-l+m, -m) = 2(\bar{l} + \bar{k}, -\bar{k}, -\bar{m}, 1-\bar{l} + \bar{m})$ which implies

$$k \rightarrow \bar{k} = -l - k, \quad m \rightarrow \bar{m} = l - 1 - m.$$

Note that $\alpha_1 + \alpha_2 = 2l$ and $\alpha_3 + \alpha_4 = 2(1-l)$ are invariant under \widehat{P}_3 and so this time l is invariant! Note that for \bar{k} to be positive. we must have $l < 0$ with $|l| \geq k$. For \bar{m} to be positive we must have $l > 0$ and $l \geq m + 1$. These are contradictory requirements and it looks that the \widehat{P}_3 transformed solution is not obtained by a simply transformation or relabeling of parameters. However the transformation

$$\widehat{P}_3 : x \rightarrow -x, F \rightarrow G, G \rightarrow F, \alpha_1 \leftrightarrow \alpha_2, \alpha_3 \leftrightarrow \alpha_4,$$

is realized as

$$\begin{aligned} \widehat{P}_3 T_2^{-k} T_4^m |F=1, G=0\rangle_{(a,0,0,2-a)} &= T_2^k T_4^{-m} |F=0, G=1\rangle_{(0,a,2-a,0)} \\ &= \widehat{P}_3 (F_m^k(x, 2l), G_m^k(x, 2l)) = (G_m^k(-x, 2l), F_m^k(-x, 2l)). \end{aligned}$$

Substituting $k \rightarrow \bar{k} = -l - k$, $m \rightarrow \bar{m} = l - 1 - m$ with $\bar{l} = l$ into equation

$$\begin{aligned} F_x &= F(F-1)(2G-1) - \frac{l+k-m}{x} F + \frac{l+k}{x}, \\ G_x &= -G(G-1)(2F-1) + \frac{l+k-m-1}{x} G - \frac{k}{x}, \end{aligned} \tag{7.5}$$

followed by $G \leftrightarrow F$ and $x \rightarrow -x$ transformation will leave equations (7.5) invariant verifying the invariance under \widehat{P}_3 automorphism.

8 Extending the method to $N > 4$

We consider few cases of solutions for higher $N = 6$ and $N = 8$ generated from specific seed solutions and illustrate the power of the method based on automorphisms to determine the explicit form of solutions. For convenience to describe the seed solution in such cases we will be using the notation based on variables $j_i(z)$ of the even dressing chain (2.2).

8.1 $N = 6$ and $N = 8$ and the seed solutions $j_i(z) = z/2, i = 1, 2, 3$ and $j_i(z) = (-1)^{i+1} z/2, i = 4, \dots, N$

In this subsection we will consider solutions constructed from seed solutions for even N that of the type $j(z) = (z/2)(1, 1, 1, -1, 1, -1)$ and $j(z) = (z/2)(1, 1, 1, -1, 1, -1, 1, -1)$. Here we analyses such models for $N = 6$ and $N = 8$ but the construction extends easily to all higher N .

We first consider a basic seed solution $j_i = \frac{z}{2}(1, 1, 1, -1, 1, -1)$ with the parameters $\alpha_i = (2 - \mathbf{a}_2, \mathbf{a}_2, 0, 0, 0, 0)$ of $N = 6$ dressing equations (2.2). This seed solution solves the higher Painlevé equations (1.1) with $g_1 = 1, g_3 = g_5 = 0, g_2 = 1, g_4 = g_6 = 0$ and will be denoted by $|g_1 = 1, g_2 = 1\rangle_{(2-\mathbf{a}, \mathbf{a}, 0, 0, 0, 0)}$. For the above solution it holds that $j_3 + j_4 = 0, j_4 + j_5 = 0, j_5 + j_6 = 0, j_6 + j_1 = 0$. As discussed in [18], to avoid division by zero we need to exclude any action by $s_i, s_{i-1}\pi^{-1}, s_{i+1}\pi$ with $i = 3, 4, 5, 6$. For similar reasons we also need to exclude the translation operators $T_i, T_{i+1}^{-1}, i = 3, 4, 5, 6$. In conclusions we are led to the only finite solutions of the form:

$$T_1^{n_1} T_2^{n_2} T_3^{-n_3} |g_1 = 1, g_2 = 1\rangle_{(2-\mathbf{a}, \mathbf{a}, 0, 0, 0, 0)}, \quad n_1, n_3 \in \mathbb{Z}_+, \quad n_2 \in \mathbb{Z},$$

with the parameters

$$(2 - \mathbf{a} + 2n_1 - 2n_2, \mathbf{a} + 2n_2 + 2n_3, -2n_3, 0, 0, 0). \quad (8.1)$$

One sees that the effect of T_2 is only to redefine \mathbf{a} by a shift with $2n_2$ as T_1 did in the case of the $N = 4$ model. According to the transformation rule (4.18) we find that the above three translation operators are being connected via the automorphism \widehat{P}_3 :

$$\widehat{P}_3 T_3^{-1} \widehat{P}_3 = T_1, \quad \widehat{P}_3 T_2 \widehat{P}_3 = T_2^{-1} \quad (8.2)$$

We can now investigate the recurrence relations emerging on the orbit of T_3^{-1} to express it in terms of Kummer polynomials and build the complete solutions utilizing fully the \widehat{P}_3 automorphism to ensure invariance under: $x \rightarrow -x, a \rightarrow 2 - a, n_2 \rightarrow n_4, n_4 \rightarrow n_2$.

Extending the solution to $N = 8$ using analogous configuration of $j_i(z) = z/2, i = 1, 2, 3$ and $j_i(z) = (-1)^{i+1} z/2, i = 4, \dots, 8$ we can now repeat the above analysis to arrive at the same form of solutions

$$T_1^{n_1} T_2^{n_2} T_3^{-n_3} |g_1 = 1, g_2 = 1\rangle_{(2-\mathbf{a}, \mathbf{a}, 0, 0, 0, 0, 0, 0)}, \quad n_1, n_3 \in \mathbb{Z}_+, \quad n_2 \in \mathbb{Z},$$

with the parameters

$$(2 - \mathbf{a} + 2n_1 - 2n_2, \mathbf{a} + 2n_2 + 2n_3, -2n_3, 0, 0, 0, 0, 0). \quad (8.3)$$

The translation operators will continue to be connected via the automorphism \widehat{P}_3 as in relation (8.2).

8.2 $N = 6$ and the seed solutions $j(z) = (z/2)(1, 1, -1, 1, 1, -1)$

As shown in [18] the configuration :

$$j_i = \frac{z}{2}(1, 1, -1, 1, 1, -1), \quad \alpha_i = (2 - \mathbf{a}_4, 0, 0, \mathbf{a}_4, 0, 0), \quad (8.4)$$

is another example of the seed solutions of $N = 6$ dressing chain equations (2.2) with two negative components of $j_i(z)$ that this time have two positive components between them.

As shown in [18] the general rational solutions are then obtained by acting with

$$T_1^{n_1} T_4^{n_4} T_2^{-n_2} T_5^{-n_5}, \quad n_1, n_2, n_4, n_5 \in \mathbb{Z}_+,$$

on the seed solution (8.4) with the parameters:

$$(2 - \mathbf{a}_4 + 2n_1 + 2n_2, -2n_2, -2n_4, \mathbf{a}_4 + 2n_4 + 2n_5, -2n_5, -2n_1). \quad (8.5)$$

According to the transformation rule (4.18) we find that the automorphisms \widehat{P}_5 and \widehat{P}_2 will connect the above translation operators as follows:

$$\begin{aligned} \widehat{P}_5 T_1 \widehat{P}_5 &= T_5^{-1}, & \widehat{P}_5 T_4 \widehat{P}_5 &= T_2^{-1} \\ \widehat{P}_2 T_1 \widehat{P}_2 &= T_2^{-1}, & \widehat{P}_2 T_4 \widehat{P}_2 &= T_5^{-1} \end{aligned} \quad (8.6)$$

This should allow us to construct general solutions out of Kummer-like polynomials.


9 Summary and conclusions

The even periodicity higher Painlevé equations possess extended symmetry structure when compared with the odd periodicity counterparts. The additional reflection automorphisms that we have introduced in this work split on reflection around even and odd points and subsequently play different roles in determining rational solutions. The reflections around even points generate new solutions while maintaining the parameters of Painlevé equations unchanged thus they are directly engaged in generating degeneracy. The reflections around odd points act as duality transformations that maintain the value of solutions although formally changing its appearance by mapping into the dual formulation. The duality property plays crucial role in determining the structure of solutions and provide a useful framework for merging solutions of simple orbits of one single translation operator into orbits created by actions of two independent translation operators. Developing these formalism embed higher number of translation operators will be crucial for complete characterization of more complex solutions including Umemura solutions, We palne to complete proofs for N byond N =4. It would be important to establish theory of discrete Painleve structures that are naturally e emerging from use of translation operators Proofs given for $N = 4$ and arguments presented by looking for particular solutions will be established for higher N ≥ 4 .


A Matrix and braids representation of reflection automorphisms for $N = 4$


It is illuminating to describe the reflection automorphisms in the language of braids and matrices. We give our discussion for $N = 4$ but it can easily be generalized to higher N .

We start with braids and separate automorphisms into two classes. The first class consists of those automorphisms that can be graphically represented by braids assigned to a single segment with upper/lower boundaries consisting of four points (1,2,3,4). This layout allows for crossing of two lines (13 for P_4 and 24 for P_2) while the two remaining lines do not transpose as shown on the next two figures.



The next class of automorphisms P_3 and P_1 act graphically over two segments by transposing lines with each transposition being allowed only one single crossing of lines in every segment.





Automorphisms $P_{2i+1}, i = 0, 1$ transpose the line 1 into the even lines 4 and 2. If these lines cross in the first segment, then the crossings of the remaining lines 23 and 41 occurs in the second segment.

Next, let us represent the reflection automorphisms as matrices acting on columns consisting of points 1, 2, 3, 4. All matrix elements will be either 0 or 1. The even reflections will have two diagonal elements equal to 1 corresponding to the two fixed points:

$$P_2 : \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix} \begin{pmatrix} 1 \\ 2 \\ 3 \\ 4 \end{pmatrix} = \begin{pmatrix} 1 \\ 4 \\ 3 \\ 2 \end{pmatrix}$$

$$P_4 : \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 \\ 2 \\ 3 \\ 4 \end{pmatrix} = \begin{pmatrix} 3 \\ 2 \\ 1 \\ 4 \end{pmatrix}$$

Thus $Tr(P_2) = Tr(P_4) = 2$. The odd reflections are traceless: $Tr(P_1) = Tr(P_3) = 0$:

$$P_1 : \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} 1 \\ 2 \\ 3 \\ 4 \end{pmatrix} = \begin{pmatrix} 4 \\ 3 \\ 2 \\ 1 \end{pmatrix}$$

$$P_3 : \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} 1 \\ 2 \\ 3 \\ 4 \end{pmatrix} = \begin{pmatrix} 2 \\ 1 \\ 4 \\ 3 \end{pmatrix}$$

The determinants are $|P_i| = -(-1)^i$ with even matrices having one eigenvalue -1 and the remaining three eigenvalues $+1$, the odd matrices have two eigenvalues equal to -1 and two equal to $+1$. The matrices are all symmetric $P_i^T = P_i$ and orthogonal. All rows are orthogonal and $P_i^2 = 1$.

The even matrices commute and the odd matrices commute among themselves. Also

$$P_2P_4 = P_1P_3 = P_4P_2 = P_3P_1 = P_1P_2P_3P_4 = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix} = \pi^2 = \pi^{-2}$$

and

$$P_1P_2 = P_2P_3 = P_3P_4 = P_4P_1 = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \end{pmatrix} = \pi^{-1}$$

And

$$P_4P_3 = P_3P_2 = P_2P_1 = P_1P_4 = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix} = \pi$$

Eigenvalues of π and π^{-1} are $-1, +1, -I, +I$ so the determinant is -1 . Eigenvalues of π^2 are $-1, 1, -1, 1$ so that the determinant is $+1$. All π^i are traceless.

It is easy to verify explicitly that reflections together with automorphisms π^n close under multiplications and therefore form a group $\{P_{n_1}, \pi^{n_2}\}$ with the identity matrix 1. For example, we find easily in the above matrix representation that

$$P_n\pi = P_{n-1}$$

and that the product $P_2P_4P_1P_3$ is equal to 1 as found by general arguments in the text.

References

- [1] Gromak V. I.; Laine I.; Shimomura S. “Painlevé Differential Equations in the Complex Plane” (2002) de Gruyter Studies in Mathematics, Volume 28
- [2] Veselov A. P.; Shabat A. B. *Funct. Anal. Appl.* (1993) **27** 81–96
- [3] Adler V. E. *Functional Analysis and Its Applications* (1993) **27:2**, 141–143
- [4] Noumi M.; Yamada Y. *Communications in Mathematical Physics* (1998) **199** 281-295
- [5] Alves V. C. C. ; Aratyn H.; Gomes J. F.; Zimerman A. H. *J. Phys. A: Math. Theor.* (2021)**54** 195701
- [6] Aratyn H.; Gomes J. F.; Zimerman A. H. *Open Communications in Nonlinear Mathematical Physics* (2022) **2**, <https://doi.org/10.46298/ocnmp.10161>
- [7] Noumi M.; Yamada Y. *Physics Letters* (1998) **A247** 65-69
- [8] Watanabe H. *Hokkaido Math. J.* (1995) **24** no.2, 231-267
- [9] Kitaev A. V.; Law C. K.; McLeod J. B. *Differential Integral Equations* (1994) **7** 967-1000
- [10] Masuda T.; Ohta Y.; Kajiwara K. *Nagoya Mathematical Journal* (2002) **168** 1-25
- [11] Kajiwara K.; Yamamoto K.; Ohta Y. *Phys. Lett.* **A232** (1997) 189-199
- [12] Matsuda K. Rational Solutions of the Noumi and Yamada System of type $A_5^{(1)}$, arXiv:0708.2960 (2007)
- [13] Masuda T. *Tohoku Math. J.* (2004) **56** 467-490
- [14] Olver F. W. J.; Lozier D. W.; Boisvert R. F.; Clark C. W. *NIST Handbook of Mathematical Functions* (2010) Cambridge University Press
- [15] Forrester P. J.; Witte N. S. *Communications in Mathematical Physics* (2001) **219** 357-398
- [16] Aratyn H.; Gomes J. F.; Ferreira L. A.; Zimerman A. H. *Phys. Lett. B* (1993) **316** 85
- [17] Masuda T, Ohta Y and Kajiwara K, *Nagoya Mathematical Journal* **168** 1-25 (2002)
- [18] Aratyn H, Gomes J F, Lobo G V and Zimerman A H 2023 *Symmetry*, **15** 249, 2023
- [19] Clarkson P A and Dunning C, *Stud Appl Math.* **152**, 453-507, 2024

- [20] H. Aratyn, J. F. Gomes, G. V. Lobo, A. H. Zimerman, Open Communications in Nonlinear Mathematical Physics, April 15, 2024, Proceedings: OCNMP Conference, Bad Ems (Germany), 23-29 June 2024 <https://doi.org/10.46298/ocnmp.13201>
- [21] Noumi M and Yamada Y, Higher order Painlevé equations of type $A_l^{(1)}$, Funkcialaj Ekvacioj **41**, 483-503, 1998, DOI:10.48550/arXiv.math/9808003