

# Decoherence due to Casimir effect?

Anirudh Gundhi\*

*Department of Physics, University of Trieste, Strada Costiera 11, 34151 Trieste, Italy and  
Istituto Nazionale di Fisica Nucleare, Trieste Section, Via Valerio 2, 34127 Trieste, Italy*

(Dated: September 16, 2024)

Open system dynamics of an electron is studied in the presence of radiation field, confined between two parallel conducting plates. It has been suggested in previous works that the quantized zero-point modes of this field lead to finite decoherence effects, possibly due to the Casimir force. However, in this work it is shown that the off-diagonal elements of the reduced density matrix of the electron are suppressed due to the sudden switching on of the interaction with the environment, and would not be observationally relevant in typical scenarios. The work clarifies important theoretical aspects of the setup and argues that any irreversible loss of coherence, in general, should not be ascribed to vacuum fluctuations. It can only originate from emission of bremsstrahlung by the electron, due to the acceleration caused by the effective Coulomb potential of all the infinite image charges.

## I. INTRODUCTION

Previous works [1–7] have considered the intriguing possibility of observing decoherence effects solely due to the presence of vacuum fluctuations<sup>1</sup>. Any such loss of coherence would also compete with the models of wavefunction collapse [11–14] in predicting the quantum-to-classical transition of perfectly isolated systems. However, it was shown in [3, 6, 7] that there can be no observable decoherence due to vacuum fluctuations alone. The overall effect of the vacuum is merely to dress the bare electron with a cloud of virtual photons [3, 6]. The QED vacuum state depends upon the position of the charged particle but has no *memory* of its trajectory. It cannot acquire which-path information about the particle and results in no irreversible loss of coherence. Therefore, the presence of zero-point modes cannot be deduced by measuring the loss of coherence for the free electron.

One can infer the presence of vacuum fluctuations, however, by measuring the stress two parallel and perfectly conducting plates exert on each other [15–19]. Simplest way to understand the Casimir effect is to notice that the boundary conditions imposed by the plates do not allow for all the zero-point modes to be present, since the allowed mode-wavelengths are constrained by the separation between the plates. Thus, there is a favorable energy gradient in bringing the plates from infinity to a finite separation, resulting in the attractive Casimir force. A review of the subject can be found in [20, 21].

The question then arises whether the Casimir force can also *monitor* the motion of a charged particle and thus lead to decoherence. In other words, it remains to be understood whether the role of the environment played by vacuum fluctuations changes from being trivial, in empty space, to something significant and observable, when the radiation field is confined between conducting

plates. Such effects have been studied in [22–24] (single conducting plate) and in [25] (where the case of parallel conducting plates was also considered explicitly). It was concluded that the zero-point fluctuations of the radiation field result in some decoherence at the level of the electron, and that this effect might be related to the Casimir force.

Instead, in this work it will be shown that the suppression of the off-diagonal elements of the reduced density matrix is unphysical and due to the sudden switching on of the interaction between the electron and the radiation field. There is no physical mechanism through which the electron’s which-path information can be lost irreversibly to the environment, especially if the plates are treated as ideal conductors and effects such as ohmic resistance to the image currents, as considered in [26], are ignored<sup>2</sup>. Nevertheless, the initial *jolt*, resulting from the sudden switching on of the interaction, leads to peculiar features in the reduced density matrix which are typically not found in other open systems. These features are explained by realizing that part of the initial jolt in the radiation field would affect the dynamics even at later times, since it is reflected back and forth by the conducting plates on either side of the electron. Although it would have little observational relevance in typical scenarios, the dynamics might still serve as a special reference for this widely discussed technical aspect of open quantum systems, dealing with the transient effects due to the sudden switching on of the system-environment (S-E) interaction [30–37].

In summary, it is here that from a physical point of view, in which such an initial jolt would be absent, decoherence can only result from the emission of bremsstrahlung, induced by the force exerted on the electron by the image charges, but not from the vacuum fluctuations of the radiation field.

---

\* [anirudh.gundhi@units.it](mailto:anirudh.gundhi@units.it)

<sup>1</sup> An introduction to open quantum systems can be found in [8, 9]. See also [10] for a discussion on decoherence in electrodynamics.

<sup>2</sup> Non-ideal conductors [27] or dynamical boundary conditions [28, 29] might also lead to irreversible decoherence.

## II. IMAGE CHARGES




FIG. 1. Infinite number of positive (blue) and negative (red) image charges induced by the plates, located at  $x = -L/2$  and  $x = L/2$ , to cancel the Coulomb field of the electron (center) along their surface.

The electric field along the surface of an idealized perfect conductor is zero. It is well-known from standard electrostatics that if an electron is placed between two parallel conducting plates, as shown in Fig. 1, this requirement can be met with the help of an infinite number of image charges, whose position is determined by the position of the electron between the plates. If the electron is placed at a coordinate  $x$ , the position-state of the image charges is given by

$$\begin{aligned} |+\rangle_x &= |\cdots, -(2n-1)L - x, \cdots, (2n-1)L - x, \cdots\rangle, \\ |-\rangle_x &= |\cdots, -2nL + x, \cdots, 2nL + x, \cdots\rangle, \end{aligned} \quad (1)$$

where  $n \geq 1$  is an integer.

From an open quantum system point of view, these infinite image charges belong to the environment. One might therefore ask whether or not there would be any direct positional decoherence due to the image charges themselves, since the position state of the images is perfectly correlated to the position state of the electron. This is the subject of this section, while decoherence due to vacuum fluctuations is studied in the following ones.

To find an answer for the loss in coherence, one might look at the off-diagonal elements of the reduced density matrix  $\hat{\rho}_r$ . The standard expression for  $\hat{\rho}_r$  is given by

$$\hat{\rho}_r(t) = \sum_E \langle E | \Psi_t \rangle \langle \Psi_t | E \rangle, \quad (2)$$

where  $|\Psi_t\rangle$  represents the full S-E state at time  $t$ , and  $|E\rangle$  the environmental basis states. In general, due to the S-E interaction, the state of the environment  $|\mathcal{E}\rangle$  becomes correlated to the position of the system such that  $|\Psi_t\rangle = \int dx \psi_t(x) |\mathcal{E}_x\rangle |x\rangle$ , where  $\psi_t(x)$  is the probability amplitude for the system to be at a position  $x$  at time  $t$ . In such a scenario, the off-diagonal elements of  $\hat{\rho}_r$  are given by

$$\langle x | \hat{\rho}_r(t) | x' \rangle = \langle \mathcal{E}_{x'} | \mathcal{E}_x \rangle \psi_t(x) \psi_t^*(x'). \quad (3)$$

If the environmental states for two different states of the system are completely orthogonal, the system is generally understood to have decohered completely. If one follows this line of reasoning strictly, the decoherence kernel

$\mathcal{D}(x, x')$  due to image charges will be given by

$$\mathcal{D}(x, x') := \langle \mathcal{E}_{x'} | \mathcal{E}_x \rangle = {}_{x'} \langle +, - | +, - \rangle_x. \quad (4)$$

It is clear that  $\mathcal{D}(x, x') = 0$  whenever  $x' \neq x$  and  $\mathcal{D}(x, x') = 1$  for  $x' = x$ . Thus, one might conclude that the image charges measure the electron's position perfectly and instantaneously.

However, this suppression of the off-diagonal elements represents *false decoherence*. In order to observe loss of fringe contrast in a double-slit experiment, the environment must be able to acquire which-path information rather than which-position information about the system. The interference fringes appear because the electron reaches a given point on the detector screen while being in a superposition of different paths. As it approaches the screen and is detected at a given point, the state of the image charges would only be correlated to the position at which the electron is detected on the screen, and would not have any *memory* of which slit the electron passed through<sup>3</sup>. In other words,  $\mathcal{D} \rightarrow 1$  for the two paths that pass through different slits but end at a given point on the screen where the electron is detected. Therefore, there would be no loss in fringe contrast due to the images. The situation here is analogous to empty space where it is understood why there are no decoherence effects due to the particle's Coulomb field itself [38, 39]. For a loss in fringe contrast, the environment should continuously monitor the system, like a thermal bath, where the photons scattering off the electron at different times can effectively measure its full trajectory between two fixed points. Instead, if  $\mathcal{D}$  depends only on the position of the system, which-path information cannot be acquired.

While the image charges cannot directly decohere the electron, there is an indirect way by which they can. It is due to the acceleration caused by the image charges, resulting in bremsstrahlung, which carries away which-path information about the electron. The effective potential at the electron's location can be computed by summing over the Coulomb potential due to all the infinite images. In doing so, one might first notice that the distance between a given negative image charge and the electron is  $|\pm 2nL + x - x| = 2nL$ . Since it is independent of the electron's position, the Coulomb potential due to the negative images cannot influence the electron's dynamics, as it simply adds a constant to the Hamiltonian. It is therefore sufficient to compute the Coulomb potential

<sup>3</sup> It might be argued that by *looking* at the images one can deduce which slit the electron passed through, and therefore the interference pattern should be lost. However, in this line of reasoning, an extra environment is introduced which can look at the images themselves and then remember the outcome. This should not be done, since the only environment we have is that of the image charges.

due to the positive image charges only. It is given by

$$V_{\text{im}} = -\alpha\hbar c \sum_{n=1}^{\infty} \left[ \frac{1}{(2n-1)L-2x} + \frac{1}{(2n-1)L+2x} \right], \quad (5)$$

where  $\alpha = e^2/(4\pi\epsilon_0\hbar c)$  is the fine structure constant. This sum is divergent, but not the dependence on  $x$ . To see this, the expression above is re-written as

$$\begin{aligned} V_{\text{im}} &= -\alpha\hbar c \sum_{n=0}^{\infty} \frac{2(2n+1)L}{(2n+1)^2L^2 - 4x^2} \\ &= -\alpha\hbar c \sum_{n=0}^{\infty} \left[ \frac{2}{(2n+1)L} \left( 1 - \frac{4x^2}{(2n+1)^2L^2} \right)^{-1} \right]. \end{aligned} \quad (6)$$

From the setup shown in Fig. 1, it can be seen that  $-L/2 < x < L/2$  and therefore the term  $(1 - 4x^2/((2n+1)L)^2)^{-1}$  can be written as a geometric series. The zero<sup>th</sup> order term in  $x$  of this expansion would again just give an irrelevant (divergent) constant contribution to the Hamiltonian and can therefore be ignored. The remaining  $x$ -dependent part of the potential is given by

$$\begin{aligned} V_{\text{im}}(x) &= \frac{-2\alpha\hbar c}{L} \sum_{n=0}^{\infty} \frac{1}{2n+1} \times \frac{4x^2}{(2n+1)^2L^2 - 4x^2} = \\ &= \frac{\alpha\hbar c}{2L} [H(-1/2 - x/L) + H(-1/2 + x/L) + \ln(16)], \end{aligned} \quad (7)$$

where,  $H(x)$  is the standard Harmonic number function generalized to the domain of real numbers. Note that the expression for the effective potential  $V_{\text{im}}$  has been previously derived in [40], using a more formal technique, but not by explicitly summing over the Coulomb potential due to all the image charges.




FIG. 2. Exact Coulomb potential due to the infinite image charges, Eq. (7), at a location  $x$  between the plates.

It can be seen from Fig. 2 that the electron would be in an unstable equilibrium at  $x = 0$  and inevitably attracted to either one of the two plates. This is inevitable, because in quantum mechanics one cannot have both the

position and velocity to be zero simultaneously<sup>4</sup>. The effect of the images on the electron dynamics if far from being negligible. It is clear that the electron would experience strong acceleration, especially near  $x = \pm L/2$ , and thus decohere due to which-path information being carried away by bremsstrahlung. To conclude this section, a simple scenario is analyzed in which the effects of the image charges can be computed easily, and contrasted with the predictions in empty space.

To study decoherence effects over large time scales, one might introduce an external harmonic potential potential,

$$V_{\text{ext}}(x) = \frac{1}{2}m\Omega^2x^2, \quad (8)$$

which restricts the electron dynamics in the neighborhood of the origin. This is because the electron would otherwise run-away towards the plates. In such a scenario, it is useful to expand  $V_{\text{im}}(x)$  in powers of  $x/L$  and to keep the leading order term. The potential due to the images (7) can be written as

$$V_{\text{im}}(x) = \frac{\alpha\hbar c}{L} \sum_{n=1}^{\infty} \frac{\psi(2n, 1/2)}{(2n)!} \left( \frac{x}{L} \right)^{2n}, \quad (9)$$

where  $\psi(2n, 1/2) := d^{2n}\psi(x)/dx^{2n}$ , evaluated at  $x = 1/2$ ,  $\psi(x)$  being the digamma function. Therefore, for a large enough  $\Omega$ , the overall effect of the image charges is to lower the frequency of the harmonic oscillator (since  $\psi(2, 1/2) = -16.8288$ )

$$\begin{aligned} V_{\text{eff}} &= V_{\text{im}} + V_{\text{ext}} = \frac{1}{2}m\Omega_{\text{eff}}^2x^2, \\ \Omega_{\text{eff}}^2 &= \Omega^2 + \frac{\alpha\hbar c\psi(2, 1/2)}{mL^3}, \end{aligned} \quad (10)$$

and thus red shift the spectrum. To leading order in  $1/L$ , the emission of bremsstrahlung between the plates is the same as in empty space. This is simply because in the limit  $L \rightarrow \infty$  we should recover the dynamics as in empty space. Therefore, to leading order, decoherence in the presence of parallel plates can be described by the master equation for bremsstrahlung [42]

$$\begin{aligned} \partial_t \hat{\rho}_r(t) &= -\frac{i}{\hbar} \left[ \frac{\hat{p}^2}{2m} + \frac{1}{2}m\Omega_{\text{eff}}^2\hat{x}^2, \hat{\rho}_r(t) \right] \\ &\quad - \frac{i\alpha\Omega_{\text{eff}}^2}{3mc^2} [\hat{x}, \{\hat{p}, \hat{\rho}_r(t)\}] - \frac{\alpha\Omega_{\text{eff}}^3}{3c^2} [x [\hat{x}, \hat{\rho}_r(t)]] . \end{aligned} \quad (11)$$

We see that since  $\Omega_{\text{eff}} < \Omega$ , the overall effect of the image charges is to increase the coherence length, simply because the plates effectively lower the frequency of the

<sup>4</sup> In this context, one may find the discussion on the quantum mechanical treatment of the inverted oscillator in [41] relevant.

external harmonic potential  $V_{\text{ext}}$ . Nevertheless, it is clear that the image potential exerts an additional force on the electron, which is absent in empty space, leading to decoherence rates which are different compared to those in empty space. The rate can be lower or higher, depending upon the setup. For other experimental scenarios, one might use the exact potential derived in Eq. (7) and compute the decoherence rate using the full noise kernel, presented in the next section. Further details concerning the effect of image charges on the electron dynamics will not be discussed. The main objective of this section is to highlight how the effective potential inevitably accelerates the electron, leading to irreversible decoherence.

### III. ZERO-POINT MODES

The final aspect of the analysis is to understand whether vacuum fluctuations, confined between conducting plates, can lead to observable decoherence. To answer this question, one must distinguish between true and unphysical decoherence, and study whether the environment *remembers* the state of the system, even after its interaction with the system is switched-off after some finite time. If it does not, the general implication would be that the environment is only correlated to the state of the system at a given time, and not to its history/trajectory. The situation would then be identical to that of Sec. II, where it was explained why environments of this type do not lead to a loss in fringe contrast.

#### A. Master equation

Up to second order in the charge  $e$ , using the influence-functional formalism [43, 44], the time evolution of the reduced density matrix of the electron was recently derived in [7], starting from the non-relativistic QED Lagrangian. It reads

$$\begin{aligned} \partial_t \hat{\rho}_r = & -\frac{i}{\hbar} \left[ \hat{H}_s, \hat{\rho}_r(t) \right] \\ & -\frac{1}{\hbar} \int_0^{t-t_i} d\tau \mathcal{N}(t; t-\tau) [\hat{x}, [\hat{x}_{H_s}(-\tau), \hat{\rho}_r(t)]] \\ & +\frac{i}{2\hbar} \int_0^{t-t_i} d\tau D(t; t-\tau) [\hat{x}, \{\hat{x}_{H_s}(-\tau), \hat{\rho}_r(t)\}]. \end{aligned} \quad (12)$$

In the equation above,  $H_s$  is the system part of the full Hamiltonian  $H$

$$H := H_s + H_{\text{EM}} + H_{\text{int}}, \quad (13)$$

where  $H_{\text{EM}}$  governs the free evolution of the radiation field and  $H_{\text{int}}$  governs the system-environment (S-E) interaction

$$H_{\text{EM}} = \int d^3r (\mathbf{\Pi}^2 + c^2 \mathbf{B}^2), \quad H_{\text{int}} = e\mathbf{r}\mathbf{\Pi}(\mathbf{r}, t). \quad (14)$$

The system Hamiltonian is given by

$$\hat{H}_s = \frac{\hat{p}^2}{2m} + V_{\text{eff}}(\hat{x}) + V_{\text{EM}}(\hat{x}), \quad (15)$$

where  $V_{\text{eff}}$  is the effective potential due to the images and some external potential that might be applied, and  $V_{\text{EM}} = e^2 \omega_{\text{max}}^3 \hat{x}^2 / (3\pi^2 \epsilon_0 c^3)$  is a divergent contribution, that scales with the UV cutoff  $\omega_{\text{max}}$ , originating from the Legendre transformation relating the QED Lagrangian and the Hamiltonian [1, 7]. The role played by  $V_{\text{EM}}$  is to cancel the divergence coming from the dissipation kernel  $D$  in the master equation (12), as shown in [7].

For the motion of the electron considered along the  $x$ -axis, the noise kernel  $\mathcal{N}$  and the dissipation kernel  $D$  in Eq. (12) are defined to be

$$\begin{aligned} \mathcal{N}(t_1; t_2) & := \frac{e^2}{2\hbar} \left\langle \{ \hat{\Pi}^x(t_1), \hat{\Pi}^x(t_2) \} \right\rangle_0, \\ D(t_1; t_2) & := \frac{ie^2}{\hbar} \left[ \hat{\Pi}^x(t_1), \hat{\Pi}^x(t_2) \right]_0 \theta(t_1 - t_2). \end{aligned} \quad (16)$$

The kernels involve the freely evolved conjugate momentum operator of the environment  $\hat{\Pi}^x(t)$ , while the expectation value in the kernels is taken with respect to the initial state of the environment. In this case we take it to be the ground state  $|0\rangle$  of the free electromagnetic (EM) field, assuming that there are no photons at the initial time  $t_i = 0$  and that the initial S-E state is completely uncorrelated, i.e.,  $\hat{\rho}(t_i) = \hat{\rho}_s(t_i) \otimes |0\rangle\langle 0|$ . The kernel  $\mathcal{N}$  describes decoherence while  $D$  describes energy lost by the system to the environment. Since it is important to distinguish between the decoherence effects due to bremsstrahlung and those due to zero-point modes of the EM field, we imagine a situation in which the electron is kept steady in a superposition of  $x$  and  $x'$ . In such a scenario,  $\hat{x}_{H_s}(-\tau)$ , which is the Heisenberg time-evolved position operator with respect to the system Hamiltonian  $\hat{H}_s$ , is given by  $\hat{x} - \hat{p}\tau/m$ . Moreover, in the absence of acceleration, the dissipation kernel would not give any contribution to the master equation as the electron can only lose energy to the environment via radiation emission upon acceleration (to second order in  $e$ ).

Thus, the master equation describing decoherence reduces to

$$\partial_t \hat{\rho}_r|_{\text{dec}} = -\frac{1}{\hbar} \int_0^{t-t_i} d\tau \mathcal{N}(t; t-\tau) [\hat{x}, [\hat{x}, \hat{\rho}_r(t)]] . \quad (17)$$

The decoherence kernel in Eq. (4) for the problem at hand can also be computed more directly, without referring to the influence functional formalism, as it will be shown later. However, computing decoherence effects via Eq. (17) has the advantage of discussing any possible connection between the Casimir force and decoherence in a straightforward way. Therefore, in what follows, the decoherence kernel  $\mathcal{D}$  is derived using both the approaches.

## B. Decoherence

The freely evolved momentum operator, same as the transverse electric field operator of the free EM field, in the presence of conducting plates, is given by [25, 45]

$$\begin{aligned} \hat{\Pi}(\mathbf{x}_{\parallel}, x, t) = & \\ & -i\sqrt{\frac{2\hbar}{\epsilon_0 L}} \sum_{n=0}^{\infty} f(n) \int \frac{d^2 k_{\parallel}}{2\pi} \sqrt{\frac{\omega_n}{2}} e^{i(\mathbf{k}_{\parallel} \cdot \mathbf{x}_{\parallel} - \omega_n t)} \times \\ & \left[ \hat{a}_1(\mathbf{k}_{\parallel}, n) (\hat{\mathbf{k}}_{\parallel} \times \hat{\mathbf{x}}) \sin\left(\frac{n\pi(x+L/2)}{L}\right) + \right. \\ & \left. \hat{a}_2(\mathbf{k}_{\parallel}, n) \left\{ \frac{i\hat{\mathbf{k}}_{\parallel} n\pi c}{\omega_n L} \sin\left(\frac{n\pi(x+L/2)}{L}\right) \right. \right. \\ & \left. \left. - \frac{\hat{\mathbf{x}} k_{\parallel} c}{\omega_n} \cos\left(\frac{n\pi(x+L/2)}{L}\right) \right\} \right] + \text{HC}. \quad (18) \end{aligned}$$

Here,  $f(n)$  is a weighing function such that  $f(0) = 1/\sqrt{2}$  and  $f(n) = 1 \forall n > 0$  [15],  $\hat{\mathbf{k}}_{\parallel}$  is a unit vector along the plates,  $\omega_n^2/c^2 := k_{\parallel}^2 + n^2\pi^2/L^2$ , and  $\hat{a}_1, \hat{a}_2$  are the annihilation operators corresponding to the two modes of the EM field. It can also be seen that  $\hat{\Pi}_{\parallel} = 0$  along the plates at  $x = \pm L/2$ , thus satisfying the appropriate boundary conditions.

We are mainly interested in studying decoherence for the electron in a superposition along the  $\hat{\mathbf{x}}$  axis. Thus, only the  $\hat{\mathbf{x}}$  component of  $\hat{\Pi}$  enters the noise kernel in Eq. (17). For electron superpositions prepared near the origin, on length scales  $|x - x'| \ll L$  (superpositions over a distance  $L$  will also be discussed later), the two-point correlation  $\langle 0 | \hat{\Pi}^x(1) \hat{\Pi}^x(2) | 0 \rangle$  is given by

$$\begin{aligned} \langle 0 | \hat{\Pi}^x(1) \hat{\Pi}^x(2) | 0 \rangle = & \\ \frac{\hbar c^2}{\epsilon_0 L} \sum_{n=-\infty}^{\infty} \int \frac{d^2 k_{\parallel}}{(2\pi)^2} \frac{k_y^2 + k_z^2}{2\omega_n} \cos^2(n\pi/2) e^{i(\mathbf{k}_{\parallel} \cdot \Delta \mathbf{x}_{\parallel} - \omega_n \tau)} & \quad (19) \end{aligned}$$

where  $\Delta \mathbf{x}_{\parallel} := \mathbf{x}_{\parallel}^{(1)} - \mathbf{x}_{\parallel}^{(2)}$  and  $\tau := t^{(1)} - t^{(2)}$ . The surface integral above can be turned into a volume integral by introducing  $\delta_n(k_x) := \delta(k_x - n\pi/L)$  such that

$$\langle 0 | \hat{\Pi}^x(1) \hat{\Pi}^x(2) | 0 \rangle = \frac{\hbar c^2}{\epsilon_0 L} \sum_{n \text{ even}} \int \frac{d^3 k}{(2\pi)^2} \frac{k^2 - k_x^2}{2\omega} e^{i(\mathbf{k}_{\parallel} \cdot \Delta \mathbf{x}_{\parallel} - \omega \tau)} \delta_n(k_x). \quad (20)$$

Further, using the properties of the Dirac comb, one can write

$$\sum_{n \text{ even}} \delta_n = \sum_{n=-\infty}^{\infty} \delta_{2n} = \frac{L}{2\pi} \sum_{m=-\infty}^{\infty} e^{imk_x L}. \quad (21)$$

Using this identity, the two-point correlation becomes

$$\langle 0 | \hat{\Pi}^x(1) \hat{\Pi}^x(2) | 0 \rangle = \frac{\hbar c^2}{2\pi \epsilon_0} \sum_{m=-\infty}^{\infty} \hat{\square}_m \int \frac{d^3 k}{(2\pi)^2 2\omega} e^{i(\mathbf{k} \cdot \Delta \mathbf{x}_m - \omega \tau)} e^{-k/k_{\max}}, \quad (22)$$

where  $\Delta \mathbf{x}_m := \{x_m, \Delta \mathbf{x}_{\parallel}\}$ ,  $x_m := mL$ ,  $\hat{\square}_m := -\partial_{\tau}^2/c^2 + \partial_{x_m}^2$ , and the UV cutoff has been introduced in the calculations by inserting  $\exp(-k/k_{\max})$  inside the integral. Further, the dependence on  $\Delta \mathbf{x}_{\parallel}$  can be ignored not only because we are mainly interested in the dynamics along the  $\mathbf{x}$ -axis, where  $\Delta \mathbf{x}_{\parallel} = 0$ , but also because  $\Delta \mathbf{x}_{\parallel} \ll c\tau$  for a non-relativistic particle. The integral can now be easily evaluated and gives

$$\langle 0 | \hat{\Pi}^x(1) \hat{\Pi}^x(2) | 0 \rangle = \frac{\hbar c}{\pi^2 \epsilon_0} \sum_m \frac{1}{(m^2 L^2 - c^2(\tau - i\epsilon)^2)^2}, \quad (23)$$

where  $\epsilon := 1/(k_{\max} c)$ . The noise kernel is thus given by

$$\mathcal{N}(\tau) = \frac{e^2 c}{2\pi^2 \epsilon_0} \sum_{m=-\infty}^{\infty} \frac{1}{(m^2 L^2 - c^2(\tau - i\epsilon)^2)^2} + \text{HC}. \quad (24)$$

Some important observations can already be made at this stage. The  $m = 0$  term in the summation is independent of  $L$  and is the only piece that survives in the limit  $L \rightarrow \infty$ . It must therefore correspond to the noise kernel  $\mathcal{N}_0$  of empty space without any conducting plates. Indeed, it can be seen that the  $m = 0$  term matches the noise kernel derived in [7] (c.f. Eqs. (47) - (49) therein). When the acceleration of the charged particle can be neglected, the suppression of the off-diagonal elements of the density matrix due to  $\mathcal{N}_0$  is false [3, 7]. It is thus not relevant to observable decoherence effects. Thus, the observationally relevant part  $\mathcal{N}_{\text{ob}}(\tau)$  is given by

$$\mathcal{N}_{\text{ob}}(\tau) = \frac{e^2 c}{\epsilon_0 \pi^2} \sum_{m=1}^{\infty} \frac{1}{(m^2 L^2 - c^2(\tau - i\epsilon)^2)^2} + \text{HC}. \quad (25)$$

Another interesting aspect emerges if we consider the large  $L$  limit or, more correctly, the limit  $c\tau \ll L$  which is given by

$$\mathcal{N}_{\text{Cas}} = \frac{16e^2}{\hbar \epsilon_0} \times \frac{\pi^2 \hbar c}{720 L^4}. \quad (26)$$

The expression is written in a suggestive way since  $\pi^2 \hbar c / (720 L^4)$  is nothing but the Casimir energy density (c.f. pg. 13 in [21]). One might therefore attribute the leading order contribution to decoherence to the Casimir effect. This, however, should not be done. If expression (26) is used in Eq. (17), the off-diagonal elements of the density matrix would be suppressed as

$$\langle x' | \hat{\rho}_r(t) | x \rangle \stackrel{\text{Cas.}}{=} \langle x' | \hat{\rho}_r(0) | x \rangle \exp\left\{ \frac{-4\pi^3 \alpha c^2 t^2 (x - x')^2}{90 L^4} \right\}. \quad (27)$$

We see that within the validity of the  $1/L$  expansion of  $\mathcal{N}_{\text{ob}}$ , i.e.  $ct \ll L$ , there will be practically no loss of coherence over any length scale  $|x - x'| \leq L$ . One might, however, extrapolate the consequences to the time

domain  $ct \gg L$  and conclude that eventually coherence will be lost over time due to the plates. This should not be done, since the behavior of the full noise kernel is very different from the leading order term.

To see this, the full expression for  $\mathcal{N}_{\text{ob}}$  in Eq. (25) needs to be integrated. Doing that, the evolution of the off-diagonal elements, and thus the decoherence kernel (Eq. (4)), is obtained to be

$$\begin{aligned} \rho_r(x', x, t) &= \exp\left(-\frac{(x' - x)^2}{\hbar} \mathcal{N}_2(t)\right) \rho_r(x', x, 0), \\ \mathcal{N}_2(t) &:= \int_0^t dt' \int_0^{t'} d\tau \mathcal{N}_{\text{ob}}(\tau) \\ &= \sum_{m=1}^{\infty} \frac{e^2}{2\pi^2 \epsilon_0 m^3 L^3} \times t \ln\left(\left|\frac{1 + ct/mL}{1 - ct/mL}\right|\right), \end{aligned} \quad (28)$$

where the limit  $\epsilon \rightarrow 0$  has been taken at the end. Before going into the properties of the decoherence kernel  $\mathcal{D}$ , it is instructive to derive it using a more direct approach. As mentioned in Sec. II, for a particle held in a superposition of  $x$  and  $x'$ ,  $\mathcal{D}(x, x', t)$  is given by the overlap of the environmental states corresponding to the two particle locations. Given the interaction Hamiltonian in Eq. (14), to leading order in  $e$ , the state of the environment (in the interaction picture) is given by

$$|\mathcal{E}(x)\rangle_t = \exp\left(\frac{-ie x}{\hbar} \int_0^t dt' \hat{\Pi}^x(x, t')\right) |0\rangle. \quad (29)$$

Since the operator  $\hat{\Pi}^x$  given in Eq. (18) is simply a linear sum of creation and annihilation operators,  $|\mathcal{E}(x)\rangle_t$  is a coherent state  $|\alpha(x, t)\rangle$ . To work out the overlap between the coherent states corresponding to different particle locations, the use of dipole approximation is well justified if the superpositions are prepared around  $x = 0$ . Therefore

$$\begin{aligned} \langle \alpha(x', t) | \alpha(x, t) \rangle &= \prod_{\mathbf{k}_{\parallel}} \prod_n \langle \alpha_{\mathbf{k}_{\parallel} n}(x', t) | \alpha_{\mathbf{k}_{\parallel} n}(x, t) \rangle = \\ &\exp\left\{ \frac{-e^2(x' - x)^2}{4\pi^2 \hbar \epsilon_0 L} \sum_{n=-\infty}^{\infty} \int d\mathbf{k}_{\parallel} \times \right. \\ &\quad \left. \times \frac{(k_{\parallel} c)^2 \cos^2(n\pi/2) \sin^2(\omega_n t/2)}{\omega_n^3} \right\}, \end{aligned} \quad (30)$$

where, the relation for the coherent states,

$$\langle \beta | \alpha \rangle = \exp(-(|\beta|^2 + |\alpha|^2 - 2\beta^* \alpha)/2), \quad (31)$$

has been used. Using the Dirac comb (21), the integral above can be evaluated exactly and gives

$$\begin{aligned} \mathcal{D}(x, x', t) &= \\ &\exp\left\{ \frac{-e^2(x' - x)^2}{2\pi^2 \hbar \epsilon_0 L^3} \sum_{m=1}^{\infty} \frac{t}{m^3} \ln\left(\left|\frac{1 + ct/mL}{1 - ct/mL}\right|\right) \right\}, \end{aligned} \quad (32)$$

where again, the contribution from the  $m = 0$  term has been discarded for the reasons described before. We see that the expression for the decoherence kernel is indeed the same when derived using the influence functional formalism (Eq. (28)), or when it is obtained by computing the overlap between the environmental states correlated to different electron positions.




FIG. 3. Time evolution of the decoherence kernel showing suppression of the off-diagonal elements at times  $t = mL/c$ , and an otherwise constant curve for  $ct \gg L$ .

#### IV. INITIAL JOLT

In this section it will be argued that all the features in  $\mathcal{D}$ , as depicted in Fig. 3, are due to the sudden switching on of the S-E interaction. These effects should not be considered physical, since in a realistic scenario there is no such sharp instant of time at which the electron starts interacting with the EM field vacuum.

A sudden switching on of the S-E interaction results in a disturbance in the state of the environment, which is sometimes referred to as a *jolt* (c.f. [34], the references therein, and [37]). In general, after the initial jolt has passed away, an equilibrium between the system and the environment is established. However, in the setup that is being considered in this work, part of the initial jolt would always influence the dynamics and the S-E equilibrium is never established. This can be seen from the periodic fall in  $\mathcal{D}$  at times  $t_m = mL/c$ .

A possible explanation of these features is that the jolt in the EM field at time  $t = 0$  propagates, gets reflected by the plates, and arrives at the location of the electron again. Since the superpositions are assumed to be near the center, this jolt would arrive at the electron location again at times  $t_m$ , resulting in a temporary fall in coherence<sup>5</sup>.

The same features can also be equivalently described in terms of the image charges. It can be seen from Eq. (1)

<sup>5</sup> Sudden switching on of the S-E interaction might also lead to emission of real photons. However, possible irreversible decoherence due to scattering of such photons is a fourth order effect in  $e$  and would thus be absent in  $\mathcal{D}(x, x', t)$  shown here.

that for an electron located at the center  $x = 0$ , the nearest images are at a distance  $L$ , the next nearest at  $2L$ , and so on. As the initial jolt is sourced by the electron at the center, the images must source a counter-jolt in order to maintain the boundary conditions. That is, by the time the jolt from the electron arrives at one of the plates, a counter jolt must arrive at the same plate from the nearest image to nullify the electric field along the plate. Subsequently, the jolt from the nearest images reaches the electron at  $t = L/c$ , from the next nearest images at  $t = 2L/c$  and so on<sup>6</sup>, explaining the fall in  $\mathcal{D}$  at times  $t_m$ .

The reasoning above indicates that the sharp falls in the off-diagonal elements would not be present in the absence of a jolt. Therefore, in such a more realistic scenario, the decoherence kernel  $\mathcal{D}$  would then only have a steady value, given by the constant upper bound in Fig. 3. This asymptotic value can be calculated analytically. If  $ct \gg L$ , the main contribution to  $\mathcal{D}$  comes from the modes for which  $mL \ll ct$ , since the contribution of the higher modes is greatly suppressed due to the  $1/mL$  dependence. In this limit  $\ln\{(1+mL/ct)/(1-mL/ct)\} \approx mL/ct$  and therefore

$$\mathcal{D}(x, x', t) \approx \exp\left\{\frac{-2\alpha(x' - x)^2}{\pi L^2} \sum_{m=1}^{\infty} \frac{1}{m^2}\right\}. \quad (33)$$

The asymptotic expression for the coherence length is rather simple and involves both the fine structure constant  $\alpha$  and the separation length  $L$ . Further, since  $\sum_m 1/m^2 = \pi^2/6$ , the off-diagonal elements would be unaffected over length scales  $L/\sqrt{\alpha}$ <sup>7</sup>, which basically means no decoherence at all.

Therefore, it is important to realize that the environment under consideration does not have a typical behavior. For instance, in the environment of thermal photons, the off-diagonal elements decay exponentially in time (ignoring dissipation, c.f. Fig. 3.8 in [8]). Thus, the extrapolation of Eq. (27) in thinking that the Casimir force leads to a continuous irreversible loss of coherence, like ordinary environments, does not hold.

It was shown in [7] that if one considers a time dependent coupling  $q(t)$  with the environment, i.e.,  $q(t) = -ef(t)$  such that  $f(t) = 1$  for most of the dynamics between the initial time  $t = 0$  and the final time  $t = T$ , while  $f(0) = f(T) = 0$ , the noise kernel in Eq. (17), and defined in Eq. (16), transforms as  $\mathcal{N} \rightarrow \tilde{\mathcal{N}}$ , with

$$\tilde{\mathcal{N}} = f(t_1)f(t_2)\mathcal{N}(t_1; t_2) = f(t_1)f(t_2)\mathcal{N}(t_1 - t_2). \quad (34)$$

<sup>6</sup> The method of images can be applied not only to electrostatics, but also in dealing with the radiation field. For instance, the well-known Casimir force was derived by explicitly referring to images in [46].

<sup>7</sup> Although on such length scales the validity of the dipole approximation can be questioned. However, for any superpositions on scales much less than  $L$ , it is clear that no coherence is lost over time. The case of wider superpositions will also be discussed explicitly later.

It was further shown at a very general level that if  $\mathcal{N}_2$  approaches a constant value on some time scale, which in this case is  $L/c$ , and if the S-E interaction is switched on adiabatically over this time scale, then  $\mathcal{N}_2$  transforms as  $\mathcal{N}_2 \rightarrow \tilde{\mathcal{N}}_2$  (c.f. Eq. (27) in [7]) with

$$\tilde{\mathcal{N}}_2(T) = \frac{\mathcal{N}_2}{2} (f^2(0) + f^2(T)) = 0, \quad (35)$$

implying no irreversible decoherence. These considerations imply that in a typical scenario in which the S-E interaction is not switched on suddenly, there would be no irreversible loss of coherence at the level of the electron and thus no loss in the fringe contrast in a double slit type of experiment. The reasoning and the arguments presented in this section are confirmed in Sec. VI, where the decoherence kernel is computed after adiabatically switching on the S-E interaction.

## V. SUPERPOSITIONS OVER LARGER LENGTH SCALES

For particle superpositions near the center, the analysis shows that there is no loss in coherence over time and that the features in  $\mathcal{D}(x, x', t)$  are solely due to the sudden switching on of the interactions. One may still question the validity of these conclusions for superpositions prepared over large length scales. It should first be clear that when considering superpositions along the  $x$ -axis, the results will not be affected by the  $y$  and  $z$  dependence of the noise kernel. This is because the dependence on the difference between the  $y$  and  $z$  coordinates, of the non-relativistic particle superpositions, can be ignored since  $\Delta \mathbf{x}_{\parallel} \ll c\tau$ . This can be seen easily from Eq. (22)<sup>8</sup>.

With this in mind, the decoherence kernel for the electron in a superposition over large length scales,  $x = -L/2$  and  $x' = L/2$ , is now computed. In this case, the two-point correlation is given by

$$\langle 0 | \hat{\Pi}^x(-L/2, t_1) \hat{\Pi}^x(L/2, t_2) | 0 \rangle = \frac{\hbar c^2}{\epsilon_0 L} \sum_{n=-\infty}^{\infty} \int \frac{d^2 k_{\parallel}}{(2\pi)^2} \frac{k_y^2 + k_z^2}{2\omega_n} (-1)^n e^{i(\mathbf{k}_{\parallel} \cdot \Delta \mathbf{x}_{\parallel} - \omega_n \tau)}. \quad (36)$$

As before, the surface integral can be converted into a volume integral by inserting a Dirac comb, and then summing over  $\sum_n (-1)^n \delta_n$ . This sum can be written as

$$\sum_{n=-\infty}^{\infty} (-1)^n \delta_n = 2 \sum_{n=-\infty}^{\infty} \delta_{2n} - \sum_{n=-\infty}^{\infty} \delta_n. \quad (37)$$

<sup>8</sup> See also the discussion in Sec. 2.1.5 (and the footnote below) of [50], where it is mentioned that corrections to the dipole approximation are only relevant when relativistic effects become important.

The identity involving the Dirac comb can now be applied to both the terms on the RHS of the equation above. With little algebra, it is easy to see that it gives

$$\begin{aligned} \mathcal{N}(\tau) &= \frac{e^2 c}{\epsilon_0 \pi^2} \sum_{m=-\infty}^{\infty} \frac{1}{(m^2 L^2 - c^2(\tau - i\epsilon)^2)^2} + \text{HC} \\ &\quad \frac{-e^2 c}{\epsilon_0 \pi^2} \sum_{m=-\infty}^{\infty} \frac{1}{(m^2 (2L)^2 - c^2(\tau - i\epsilon)^2)^2} + \text{HC}. \end{aligned} \quad (38)$$

The decoherence kernel is then immediately obtained to be

$$\begin{aligned} \mathcal{D}(-L/2, L/2, t) &= \\ \exp \left\{ \frac{-e^2 (x' - x)^2}{\pi^2 \hbar \epsilon_0} \sum_{m=1}^{\infty} \frac{t}{m^3} \times \right. \\ &\quad \left. \left[ \frac{1}{L^3} \ln \left( \left| \frac{mL + ct}{mL - ct} \right| \right) - \frac{1}{(2L)^3} \ln \left( \left| \frac{2mL + ct}{2mL - ct} \right| \right) \right] \right\}. \end{aligned} \quad (39)$$

Consistent with the description above, it can be seen that there will be falls in coherence not only at time intervals  $t = 2mL/c$ , since it takes the time  $2L/c$  for the jolt sourced near one of the plates to return to the same plate, but also at time intervals  $t = mL/c$ , since the jolt sourced from  $x = -L/2$  can temporarily become correlated to the electron at  $x' = L/2$ . And again, even for the largest possible superposition, one sees that all the features in the off-diagonal elements can be ascribed to the sudden switching on of the interaction. The same and consistent conclusions are reached if  $\mathcal{D}(-L/2, 0, t)$  or  $\mathcal{D}(0, L/2, t)$  is computed.

## VI. ADIABATIC SWITCHING ON

To confirm the physical interpretation, one can compute the off-diagonal elements of the density matrix, but this time after adiabatically switching on the S-E interaction. This can be modeled by evolving the state of the environment as

$$|\mathcal{E}(x)\rangle_t^{\text{ad}} = \exp \left\{ \frac{-ie x}{\hbar} \int_0^t dt' \hat{\Pi}^x(x, t') (1 - e^{-t'/T}) \right\} |0\rangle. \quad (40)$$

Here,  $T$  represents the time scale over which the S-E interaction is switched on and  $t$  is some late time at which the interaction is fully switched on. After computing the integral, one should first take the limit  $t \gg T$  and then the limit  $T \rightarrow \infty$ . If the limits are taken in the opposite order, the S-E interaction would never be switched on. After the interaction is fully switched on at some time  $t$ ,

the decoherence kernel is given by

$$\begin{aligned} \mathcal{D}^{\text{ad}}(x, x', t) &= \langle \mathcal{E}(x') | \mathcal{E}(x) \rangle_t^{\text{ad}} = \\ \exp \left\{ \frac{-e^2 (x' - x)^2}{16\pi^2 \hbar \epsilon_0 L} \sum_{n=-\infty}^{\infty} \int d\mathbf{k}_{\parallel} \frac{(k_{\parallel} c)^2 \cos^2(n\pi/2)}{\omega_n^3} \right\}. \end{aligned} \quad (41)$$

Here, the calculations are performed for the case where the superpositions are prepared near the center. The main effect of a sudden jolt is not related to the length scale over which the superpositions are prepared and the conclusions of adiabatically switching on the interaction would also apply to  $\mathcal{D}(L/2, -L/2, t)$ .

It is already interesting to notice that  $\mathcal{D}$  in Eq. (41) does not depend on time  $t$ , indicating that the oscillations in Fig. 3 do not appear in the absence of a sudden jolt. As before, the integral can be computed by inserting a Dirac comb and using the identity (21). It gives

$$\begin{aligned} \mathcal{D}^{\text{ad}}(x, x', t) &= \\ \exp \left\{ \frac{-e^2 (x' - x)^2}{8\pi^2 \hbar c \epsilon_0} \times \right. \\ &\quad \left. \times \sum_{m=1}^{\infty} \int_0^{\infty} dk k \int_0^{\pi} d\theta \sin \theta (1 - \cos^2 \theta) e^{imLk \cos \theta} \right\} \\ &= \exp \left\{ \frac{-2\alpha (x' - x)^2}{\pi L^2} \sum_{m=1}^{\infty} \frac{1}{m^2} \right\}. \end{aligned} \quad (42)$$

It can be clearly seen that the oscillations have disappeared from the decoherence kernel altogether (compared to Eq. (32)) and that its value is the same as that of the asymptotic curve in Fig. 3. The off-diagonal elements are not suppressed over any length scale  $|x - x'| \leq L$ . What is even more important to notice is that after the S-E interaction is fully switched on, the overlap between the environmental states is stationary in time. Thus, the vacuum of the interacting radiation field effectively acts like an environment which is only correlated to the position of the electron. The situation then becomes identical to the one described in Sec. II. This implies that there would be no loss in the fringe contrast due to vacuum fluctuations, as they are not able to resolve the two paths which end at the same point on the detector screen.

## VII. CONCLUSIONS

In this work, decoherence due to the vacuum fluctuations of the EM field, confined between parallel conducting plates, is analyzed. It is shown that the loss in coherence that one may find is a consequence of the sudden switching on of the S-E interaction and not of the nature of the vacuum fluctuations themselves, and thus not related to the Casimir force. To study the time evolution of the reduced density matrix, it is standard practice, for technical convenience, to start from an initially uncorrelated S-E state. However, this initial condition, in

general, is only justified when the interaction between the system and the environment is also switched off initially, since S-E interaction leads to S-E correlations.

Therefore, if one takes the vacuum state of the free EM field as the initial state of the environment, uncorrelated to the initial state of the electron, one should also switch off their interaction initially. It is well-known from standard quantum mechanics that to go from the vacuum state of the free EM field (i.e. the bare vacuum), to the vacuum state of the interacting radiation field (which is the main subject of the analysis), the interaction must be switched on adiabatically. In other words, the gap between the starting point (dictated by technical convenience), and the actual physical situation of interest, must be bridged by adiabatically switching on the interaction. When this is done, it is shown in this work that the zero-point modes of the vacuum do not lead to any

decoherence effects at the level of the electron. Decoherence due to zero point modes might still be relevant in a situation where the charged particle suddenly enters and leaves a region confined between conducting plates. However, in typical scenarios, the only irreversible loss of coherence in the presence of parallel plates would be due to the effective Coulomb potential of all the infinite image charges, which accelerates the electron resulting in bremsstrahlung.

## VIII. ACKNOWLEDGEMENTS

I thank Angelo Bassi and fellow group members for several discussions. This work was financially supported by the University of Trieste, INFN and EIC Pathfinder project QuCoM (GA No. 101046973).

- 
- [1] P. M. V. B. Barone and A. O. Caldeira, *Phys. Rev. A* **43**, 57 (1991).
  - [2] E. Santos, *Physics Letters A* **188**, 198 (1994).
  - [3] L. Diósi, *Physics Letters A* **197**, 183 (1995).
  - [4] L. H. Ford, *Phys. Rev. A* **56**, 1812 (1997).
  - [5] H.-P. Breuer and F. Petruccione, in *Relativistic Quantum Measurement and Decoherence*, edited by H.-P. Breuer and F. Petruccione (Springer Berlin Heidelberg, Berlin, Heidelberg, 2000) pp. 31–65.
  - [6] W. G. Unruh, in *Relativistic Quantum Measurement and Decoherence*, edited by H.-P. Breuer and F. Petruccione (Springer Berlin Heidelberg, Berlin, Heidelberg, 2000) pp. 125–140.
  - [7] A. Gundhi and A. Bassi, *Phys. Rev. A* **107**, 062801 (2023).
  - [8] E. Joos, H. Zeh, D. Giulini, C. Kiefer, J. Kupsch, and I. Stamatescu, *Decoherence and the Appearance of a Classical World in Quantum Theory*, Physics and astronomy online library (Springer, 2003).
  - [9] M. A. Schlosshauer, *Decoherence and the Quantum-To-Classical Transition* (Springer-Verlag Berlin Heidelberg, 2007).
  - [10] C. Kiefer, *Phys. Rev. D* **46**, 1658 (1992).
  - [11] A. Bassi and G. Ghirardi, *Phys. Rep.* **379**, 257 (2003).
  - [12] A. Bassi, K. Lochan, S. Satin, T. P. Singh, and H. Ulbricht, *Rev. Mod. Phys.* **85**, 471 (2013).
  - [13] M. Arndt and K. Hornberger, *Nature Phys.* **10**, 271 (2014).
  - [14] M. Carlesso, S. Donadi, L. Ferialdi, M. Paternostro, H. Ulbricht, and A. Bassi, *Nature Phys.* **18**, 243 (2022).
  - [15] H. B. G. Casimir, *Indag. Math.* **10**, 261 (1948).
  - [16] E. S. Sabisky and C. H. Anderson, *Phys. Rev. A* **7**, 790 (1973).
  - [17] S. K. Lamoreaux, *Phys. Rev. Lett.* **78**, 5 (1997).
  - [18] S. K. Lamoreaux, *Phys. Rev. Lett.* **81**, 5475 (1998).
  - [19] S. K. Lamoreaux, *Phys. Rev. A* **59**, R3149 (1999).
  - [20] L. Spruch, An overview of long-range casimir interactions, in *Long-Range Casimir Forces: Theory and Recent Experiments on Atomic Systems*, edited by F. S. Levin and D. A. Micha (Springer US, Boston, MA, 1993) pp. 1–71.
  - [21] K. A. Milton, *The Casimir effect: Physical manifestations of zero-point energy* (2001).
  - [22] L. H. Ford, *Phys. Rev. D* **47**, 5571 (1993).
  - [23] H. Yu and L. H. Ford, *Phys. Rev. D* **70**, 065009 (2004).
  - [24] F. D. Mazzitelli, J. P. Paz, and A. Villanueva, *Phys. Rev. A* **68**, 062106 (2003).
  - [25] J.-T. Hsiang and D.-S. Lee, *Phys. Rev. D* **73**, 065022 (2006).
  - [26] J. R. Anglin and W. H. Zurek, in *31st Rencontres de Moriond: Dark Matter and Cosmology, Quantum Measurements and Experimental Gravitation* (1996) pp. 263–270, [arXiv:quant-ph/9611049](https://arxiv.org/abs/quant-ph/9611049).
  - [27] Y. Levinson, *Journal of Physics A: Mathematical and General* **37**, 3003 (2004).
  - [28] D. A. R. Dalvit and P. A. Maia Neto, *Phys. Rev. Lett.* **84**, 798 (2000).
  - [29] C.-H. Wu and D.-S. Lee, *Phys. Rev. D* **71**, 125005 (2005).
  - [30] V. Hakim and V. Ambegaokar, *Phys. Rev. A* **32**, 423 (1985).
  - [31] R. T. Skodje and J. R. Cary, *Computer Physics Reports* **8**, 221 (1988).
  - [32] H. Grabert, P. Schramm, and G. L. Ingold, *Phys. Rept.* **168**, 115 (1988).
  - [33] W. G. Unruh and W. H. Zurek, *Phys. Rev. D* **40**, 1071 (1989).
  - [34] B. L. Hu, J. P. Paz, and Y. Zhang, *Phys. Rev. D* **45**, 2843 (1992).
  - [35] M. Rosenau da Costa, A. O. Caldeira, S. M. Dutra, and H. Westfahl, *Phys. Rev. A* **61**, 022107 (2000).
  - [36] S. R. Das, D. A. Galante, and R. C. Myers, *JHEP* **02**, 167 (2015).
  - [37] C. Agón and A. Lawrence, *JHEP* **04**, 008 (2018).
  - [38] D. Giulini, C. Kiefer, and H. D. Zeh, *Phys. Lett. A* **199**, 291 (1995).
  - [39] C. Kiefer, Decoherence in quantum field theory and quantum gravity, in *Decoherence and the Appearance of a Classical World in Quantum Theory* (Springer Berlin Heidelberg, Berlin, Heidelberg, 2003) pp. 181–225.
  - [40] G. Barton, *Proc. Roy. Soc. Lond. A* **320**, 251 (1970).
  - [41] G. Barton, *Annals of Physics* **166**, 322 (1986).

- [42] A. Gundhi, O. Angeli, and A. Bassi, (2024), [arXiv:2403.14752](https://arxiv.org/abs/2403.14752) [quant-ph].
- [43] R. Feynman and F. Vernon, *Annals of Physics* **24**, 118 (1963).
- [44] E. A. Calzetta and B.-L. B. Hu, *Nonequilibrium Quantum Field Theory*, Cambridge Monographs on Mathematical Physics (Cambridge University Press, 2008).
- [45] G. Barton, *Journal of Physics A: Mathematical and General* **24**, 991 (1991).
- [46] L. S. Brown and G. J. Maclay, *Phys. Rev.* **184**, 1272 (1969).
- [47] H.-P. Breuer, E.-M. Laine, J. Piilo, and B. Vacchini, *Rev. Mod. Phys.* **88**, 021002 (2016).
- [48] N. Megier, A. Smirne, and B. Vacchini, *Phys. Rev. Lett.* **127**, 030401 (2021).
- [49] A. Smirne, N. Megier, and B. Vacchini, *Phys. Rev. A* **106**, 012205 (2022).
- [50] Dalibard, J., Dupont-Roc, J., and Cohen-Tannoudji, C., *J. Phys. France* **43**, 1617 (1982).