Extending a result of Carlitz and McConnel to polynomials which are not permutations

Bence Csajbók[∗]

Abstract

Let D denote the set of directions determined by the graph of a polynomial f of $\mathbb{F}_q[x]$, where q is a power of the prime p. If D is contained in a multiplicative subgroup M of \mathbb{F}_q^{\times} $_{q}^{\times}$, then by a result of Carlitz and McConnel it follows that $f(x) = ax^{p^k} + b$ for some $k \in \mathbb{N}$. Of course, if $D \subseteq M$, then $0 \notin D$ and hence f is a permutation. If we assume the weaker condition $D \subseteq M \cup \{0\}$, then f is not necessarily a permutation, but Sziklai conjectured that $f(x) = ax^{p^k} + b$ follows also in this case. When q is odd, and the index of M is even, then a result of Ball, Blokhuis, Brouwer, Storme and Sz˝onyi combined with a result of McGuire and Göloğlu proves the conjecture. Assume deg $f \geq 1$. We prove that if the size of $D^{-1}D = \{d^{-1}d': d \in D \setminus \{0\}, d' \in D\}$ is less than $q - \deg f + 2$, then f is a permutation of \mathbb{F}_q . We use this result to verify the conjecture of Sziklai.

1 Introduction

Let \mathbb{F}_q denote the finite field of $q = p^n$ elements, where p is a prime. If f is an $\mathbb{F}_q \to \mathbb{F}_q$ function then the affine q-set $\{(x, f(x)) : x \in \mathbb{F}_q\} \subseteq \text{AG}(2, q)$ is called the graph of f, the subset $D_f := \frac{\left(\frac{f(x)-f(y)}{x-y}\right) \cdot x, y \in \mathbb{F}_q, x \neq 0}{\left(\frac{f(x)-f(y)}{x-y}\right)}$ y} of \mathbb{F}_q is called the set of directions, or slopes, determined by (the graph of) f. Each $\mathbb{F}_q \to \mathbb{F}_q$ function can be uniquely represented by a polynomial of $\mathbb{F}_q[x]$ and of degree at most $q-1$, so we will consider polynomials instead of function. For a subset A of \mathbb{F}_q we will denote $A \setminus \{0\}$ by A^* .

^{*}The author is supported by the János Bolyai Research Scholarship of the Hungarian Academy of Sciences.

When $d > 1$ is a divisor of $q - 1$ then put

$$
M_d := \{ x^d : x \in \mathbb{F}_q^* \}.
$$

The following result was proved first by Carlitz in the case of $d = 2$ [\[5\]](#page-4-0) and then generalised by McConnel for other divisors d of $q-1$ [\[6\]](#page-4-1).

Result 1.1. If $D_f \subseteq M_d$, then f is of the form $f(x) = a + bx^{p^k}$.

There are several generalisations and different proofs of this result, due to Bruen and Levinger [\[3\]](#page-4-2), Grundhöfer [\[8\]](#page-5-0), Lenstra [\[4\]](#page-4-3), see [\[9,](#page-5-1) Section 9] for a survey on these results and their relation with the Paley graph.

Since $0 \notin D_f$ yields $f(x) \neq f(y)$ for each $x \neq y$, it is obvious that only permutations can satisfy the condition $D \subseteq M_d$. This is not the case anymore if we allow $D \subseteq M_d \cup \{0\}$. In [\[10,](#page-5-2) pg. 114] Sziklai conjectured that $f(x) = ax^{p^k} + b$ holds also when one replaces M_d with $M_d \cup \{0\}$ in the statement of Result [1.1.](#page-1-0) To present what is known regarding this conjecture, we need to recall some definitions. An $\mathbb{F}_q \to \mathbb{F}_q$ function f is called additive if $f(a + b) = f(a) + f(b)$ for each $a, b \in \mathbb{F}_q$. Such functions correspond to polynomials $a_0x + a_1x^p + \ldots + a_{n-1}x^{p^{n-1}} \in \mathbb{F}_q[x]$. If f is additive and $\alpha \in \mathbb{F}_q$, then we we will call $f + \alpha$ an affine polynomial. An important result on directions is the following, due to Ball, Blokhuis, Brouwer, Storme, Szőnyi $|2|$ and Ball $|1|$.

Result 1.2. If $|D_f| \leq (q+1)/2$, then f is an affine polynomial.

If f is additive, then for the affine polynomial $g = f + \alpha$ it holds that $D_g = \{f(x)/x : x \in \mathbb{F}_q^*\}$ $\binom{4}{q} = D_f$. The following result is due to McGuire and Göloğlu [\[7\]](#page-5-3).

Result 1.3. If q is odd, f is additive and $D_f \subseteq M_2 \cup \{0\}$, then $f(x) = ax^{p^k}$.

If $D_q \subseteq M_2 \cup \{0\}$, then $|D_q| \leq (q+1)/2$ and hence by Result [1.2](#page-1-1) $q = f + \alpha$ for some additive f with $D_f = D_q$, thus Result [1.3](#page-1-2) proves Sziklai's conjecture in case of odd q and even d . In [\[7\]](#page-5-3) the authors used Kloosterman sums and it seems to be that their technique cannot be used to prove the conjecture when d is odd.

Our key result is Theorem [2.1.](#page-2-0) It's proof was inspired by the recent manuscript [\[11\]](#page-5-4) of C. H. Yip who used Result [1.2](#page-1-1) in a clever way to strengthen Result [1.1.](#page-1-0) In Corollary [2.5](#page-4-6) we extend Yip's result to polynomials which are

not necessarily permutations. Our notation is standard, if $A, B \subseteq \mathbb{F}_q$ then $A^{-1} = \{a^{-1} : a \in A^*\}\$ and $AB = \{ab : a \in A, b \in B\}$. If $c \in \mathbb{F}_q$, then $A - c = \{a - c : a \in A\}$. In Theorem [2.2](#page-3-0) and Corollary [2.3](#page-3-1) we present conditions on the size of $D_f^{-1}D_f$ which ensure that f is a permutation. In Corollary [2.4](#page-4-7) we prove Sziklai's conjecture.

2 New Results

Our first result shows how a simple combinatorial property of the graph of f implies an algebraic property of D_f .

Theorem 2.1. Assume that the line of equation $y = mx + b$ meets the graph of $f \in \mathbb{F}_q[x]$ in k points for some $1 < k < q$. Then

$$
|(D_f - m)^{-1}(D_f - m)| \ge q - k + 2.
$$

Proof. Put $q(x) := f(x) - mx - b$. Then q has exactly k distinct roots. Let a be one of them and define $h(x) := g(x + a)$. Clearly h has exactly k distinct roots and 0 is one of them. Also,

$$
D_h = \left\{ \frac{h(x) - h(y)}{x - y} : x, y \in \mathbb{F}_q, x \neq y \right\} =
$$

$$
\left\{ \frac{f(x + a) - f(y + a) - m(x - y)}{(x + a) - (y + a)} : x, y \in \mathbb{F}_q, x \neq y \right\} = D_f - m.
$$

Denote by $r_1, r_2, \ldots, r_{k-1}$ the distinct non-zero roots of h.

We claim that

$$
H := \left\{ \frac{x}{x - y} : x, y \in \mathbb{F}_q, h(x) \neq 0, h(y) = 0 \right\} \subseteq D_h^{-1} D_h,
$$
 (1)

and

$$
|H| \ge q - k + 1. \tag{2}
$$

To prove [\(1\)](#page-2-1) take an element $x/(x-y)$ from H (i.e., $h(x) \neq 0$, $h(y) = 0$) and put $c = x/h(x)$. Since $h(0) = 0$, we have

$$
c = \frac{x}{h(x)} = \frac{x - 0}{h(x) - h(0)} \in D_h^{-1}.
$$

Also, since $h(y) = 0$, we have

$$
\frac{x}{x-y} = c \frac{h(x) - h(y)}{x-y} \in D_h^{-1}D_h.
$$

To prove [\(2\)](#page-2-2) first note that $0, 1 \in D_h^{-1}D_h$, $0 \notin H$ and $1 \in H$. Assume that $\alpha \in \mathbb{F}_q \setminus \{0, 1\}$ is not contained in H. Then the solutions in x of the equations

$$
x/(x - r_i) = \alpha, \quad i \in \{1, 2, \dots, k - 1\},\tag{3}
$$

are in $\{r_1, r_2, \ldots, r_{k-1}\}$. If x is a solution of $x/(x - r_i) = \alpha$, then $x =$ $r_i\alpha/(\alpha - 1)$, so if $\alpha \notin H$, then

$$
\alpha/(\alpha-1)\{r_1,r_2,\ldots,r_{k-1}\}\subseteq \{r_1,r_2,\ldots,r_{k-1}\},\,
$$

and hence

$$
\alpha/(\alpha-1)\{r_1,r_2,\ldots,r_{k-1}\}=\{r_1,r_2,\ldots,r_{k-1}\}.
$$

Put $\beta = \alpha/(\alpha - 1)$ and $R = \{r_1, r_2, \ldots, r_{k-1}\}.$ Then $\beta R = R$. Clearly there are at most $k-1$ such βs : $r_1/r_1, r_2/r_1, \ldots, r_{k-1}/r_1$, but $\beta \neq 1$ and hence there are at most $k - 2$ values of $\alpha \in \mathbb{F}_q \setminus \{0, 1\}$ for which [\(3\)](#page-3-2) does not have a solution with $h(x) \neq 0$. It follows that $|H| \geq (q-2) - (k-2) + 1$ (because of $1 \in H$). Since $0 \in D_h^{-1}D_h \setminus H$, we have $|D_h^{-1}D_h| \geq |H| + 1$, which proves the assertion. □

Theorem 2.2. Let $f \in \mathbb{F}_q[x]$ be a polynomial of degree k for some $0 < k < q$. If $|D_f^{-1}D_f| < q-k+2$, then f is a permutation of \mathbb{F}_q .

Proof. If $k = 1$, then f is a permutation. Assume $k > 1$. If f was not a permutation, then there would be a line with slope 0 meeting U_f in at least two and at most k points (since $f(x) = c$ has at most deg f roots for each $c \in \mathbb{F}_q$). By Theorem [2.1](#page-2-0) it follows that $|D_f^{-1}D_f| \ge q - k + 2$, contradicting the assumption. \Box

Corollary 2.3. If $|D_f^{-1}D_f| \leq (q+1)/2$ holds for some $\mathbb{F}_q \to \mathbb{F}_q$ non-constant function f, then f permutes \mathbb{F}_q .

Proof. f is non-constant, hence there is a non-zero element in D_f , so D_f^{-1} $f^ \neq$ \emptyset and $|D_f| \leq |D_f^{-1}D_f|$. By Result [1.2](#page-1-1) f can be represented by an affine polynomial of degree at most $q/p \leq q/2$. Then

$$
|D_f^{-1}D_f| \le \frac{q+1}{2} < q - q/2 + 2 \le q - \deg f + 2,
$$

and the result follows from Theorem [2.2.](#page-3-0)

 \Box

The next result proves Conjecture 18.11 form [\[10,](#page-5-2) pg. 114].

Corollary 2.4. If $D_f \subseteq M_d \cup \{0\}$, then $f(x) = ax^{p^k} + b$.

Proof. $D_f^{-1}D_f \subseteq M_d^{-1}(M_d \cup \{0\}) = M_d \cup \{0\}$ and hence $|D_f^{-1}D_f| \leq |M_d| + 1 \leq$ $(q+1)/2$. By Corollary [2.3](#page-3-1) f is a constant function (that is, $a=0$), or it is a permutation and hence $0 \notin D_f$. The assertion follows from Result [1.1.](#page-1-0) \Box

The next result weakens the condition on f from [\[11,](#page-5-4) Theorem 1.2], since we do not require f to be a permutation.

Corollary 2.5. If for some $\mathbb{F}_q \to \mathbb{F}_q$ function f it holds that $|D_f^{-1}D_fD_f^{-1}$ $\vert f^{\text{--}1}\vert \leq$ $(q+1)/2$, then $f(x) = ax^{p^k} + b$.

Proof. If f is the constant function then the result trivially holds. If it is not, then D_f^{-1} $f_f^{-1} \neq \emptyset$ and hence $|D_f^{-1}D_f| \leq |D_f^{-1}D_fD_f^{-1}$ $|f^{-1}| \leq (q+1)/2$, thus f is a permutation by Corollary [2.3.](#page-3-1) Then the result follows from [\[11,](#page-5-4) Theorem 1.2 and from the fact that $1 \in D_f^{-1}D_f$ and hence $D_f^{-1} \subseteq D_f^{-1}D_fD_f^{-1}$ f^{-1} , thus $D_f^{-1} \cup D_f^{-1} D_f D_f^{-1} = D_f^{-1} D_f D_f^{-1}$ $\frac{-1}{f}$.

References

- [1] S. Ball: The number of directions determined by a function over a finite field, J. Combin. Theory Ser. A, 104(2) (2003), 341–350.
- [2] A. Blokhuis, S. Ball, A. E. Brouwer, L. Storme, and T. SZŐNYI: On the number of slopes of the graph of a function defined on a finite field, J. Combin. Theory Ser. A, 86(1) (1999), 187–196.
- [3] A. Bruen and B. Levinger: A theorem on permutations of a finite field, Canadian J. Math. 25 (1973), 1060–1065.
- [4] H. W. LENSTRA, JR.: Automorphisms of finite fields. J. Number Theory, 34(1) (1990), 33–40.
- [5] L. Carlitz: A theorem on permutations in a finite field. Proc. Amer. Math. Soc. 11 (1960), 456–459.
- [6] R. McConnel: Pseudo-ordered polynomials over a finite field, Acta Arith. 8 (1963), 127–151.
- [7] G. McGuire and F. Göloğlu: On theorems of Carlitz and Payne on permutation polynomials over finite fields with an application to $x^{-1} + L(x)$, Finite Fields Appl. 27 (2014), 130-142.
- [8] T. GRUNDHÖFER: Über Abbildungen mit eingeschränktem Differenzenprodukt auf einem endlichen Körper. Arch. Math. (Basel), $37(1)$ (1981), 59–62.
- [9] G. A. Jones: Paley and the Paley graphs. In Isomorphisms, symmetry and computations in algebraic graph theory, volume 305 of Springer Proc. Math. Stat., pages 155–183. Springer, Cham, 2020.
- [10] P. Sziklai: Polynomials in finite geometry, 2008, [https://www.academia.edu/69422637/Polynomials](https://www.academia.edu/69422637/Polynomials_in_finite_geometry) in finite geometry
- [11] C. H. Yip: A strengthening of McConnel's theorem on permutations over finite fields, 2024,<https://arxiv.org/abs/2407.21362>

Bence Csajbók ELTE Eötvös Loránd University, Budapest, Hungary Department of Computer Science 1117 Budapest, Pázmány P. stny. 1/C, Hungary bence.csajbok@ttk.elte.hu