

THE GEOMETRIC CAUCHY PROBLEM FOR CONSTANT-RANK SUBMANIFOLDS

MATTEO RAFFAELLI

ABSTRACT. Given a smooth s -dimensional submanifold S of \mathbb{R}^{m+c} and a smooth distribution $\mathcal{D} \supset TS$ of rank m along S , we study the following geometric Cauchy problem: to find an m -dimensional rank- s submanifold M of \mathbb{R}^{m+c} (that is, an m -submanifold with constant index of relative nullity $m - s$) such that $M \supset S$ and $TM|_S = \mathcal{D}$. In particular, under some reasonable assumption and using a constructive approach, we show that a solution exists and is unique in a neighborhood of S .

1. INTRODUCTION AND MAIN RESULT

Given a manifold $Q^{q \geq 3}$ and some class \mathcal{A} of $(m < q)$ -dimensional submanifolds, the *geometric Cauchy problem for the class \mathcal{A}* is to find all members of \mathcal{A} passing from a given $(s < m)$ -dimensional submanifold S of Q with prescribed tangent bundle along S . This problem, which represents a far-reaching generalization of the classical Björling problem for minimal surfaces in \mathbb{R}^3 [18, section 22.6], has recently been examined, always with $s = 1$, for several combinations of Q and \mathcal{A} ; see, e.g., [24, 5, 3, 16, 8, 23, 6, 2, 1].

In this paper, we are interested in the geometric Cauchy problem for *constant-rank submanifolds*, that is, smooth embedded submanifolds of \mathbb{R}^{m+c} whose Gauss map has constant rank, or, equivalently, whose second fundamental form has constant index of relative nullity [7]. These make one of the simplest and most natural classes of submanifolds and, for this reason, also one of the most studied; see, e.g., [27, 25, 10, 17, 9, 29, 13, 11] and [14, Chapter 7].

Problem 1.1 (Geometric Cauchy problem for constant-rank submanifolds). Let S be a connected s -dimensional smooth embedded submanifold of \mathbb{R}^{m+c} , and let $\mathcal{D} \supset TS$ be a smooth distribution of rank m along S , that is, a smooth rank- m subbundle of the ambient tangent bundle over S . Find all rank- s submanifolds of \mathbb{R}^{m+c} containing S and whose tangent bundle along S is precisely \mathcal{D} .

Some special cases of Problem 1.1 have already been considered: the case $m = 2$ is classical [15, pp. 198–200]; the case of rank-one hypersurfaces ($s = c = 1$) was first addressed by Markina and the author in [22]; and the rank-one case in arbitrary codimension was solved by the author in [26]. The interest in Problem 1.1 is at least partially motivated by the desire to find new methods, alternative to the Gauss parametrization [12, 28], to describe constant-rank submanifolds. The purpose of

Date: September 5, 2024.

2020 Mathematics Subject Classification. Primary 53A07; Secondary 53B20, 53B25.

Key words and phrases. Constant nullity, distributions along submanifolds, index of relative nullity, ruled submanifold, vector cross product.

this paper is to present a solution of Problem 1.1 without any assumption on s , m , or c .

To state our main result, let \mathcal{D}^\perp be the distribution of rank c whose fiber at $p \in S$ is the orthogonal complement of \mathcal{D}_p in \mathbb{R}^{m+c} , let π^\top denote orthogonal projection onto \mathcal{D} , and let ϕ_p be the map

$$\begin{aligned} T_p S \times \mathcal{D}_p^\perp &\rightarrow \mathcal{D}_p \\ (v, n) &\mapsto \pi^\top \nabla_v N, \end{aligned}$$

where ∇ is the Euclidean connection, and where N is any extension of n to a local section of \mathcal{D}^\perp . It is easy to see that ϕ_p is well defined.

Theorem 1.2. *Suppose that there exists a section N^* of \mathcal{D}^\perp such that the shape operator $A^* := A_{N^*}$ of S in direction N^* is nonsingular, i.e.,*

$$\text{rank } A^*|_p = s \quad \forall p \in S.$$

The geometric Cauchy problem for rank- s submanifolds of \mathbb{R}^{m+c} has a solution if and only if $\phi_p(T_p S, N_p^) = \text{Im } \phi_p$ for all $p \in S$. Moreover:*

- (i) *The solution is locally unique, i.e., any two solutions coincide on an open set containing S .*
- (ii) *In this open set, the unique solution is given by*

$$\{p + x \mid p \in S, x \in \mathcal{D}_p \cap (\text{Im } \phi_p)^\perp\},$$

where $^\perp$ denotes orthogonal complement in \mathbb{R}^{m+c} .

- (iii) *Given a local parametrization $\xi: A \rightarrow \mathbb{R}^{m+c}$ of S , the solution can be locally parametrized as follows. Let (E_1, \dots, E_m) be a smooth orthonormal frame for $\mathcal{D}|_{\xi(A)}$ whose first s elements span TS . For any $j = 1, \dots, m-s$, let G_i be the group of permutations of the set $\{1, \dots, \widehat{i}, \dots, s, s+j\}$ obtained by omitting the element i , and, denoting by ϕ_i^k the coordinate function $\langle \phi(E_i, N^*), E_k \rangle = \langle \nabla_{E_i} N^*, E_k \rangle$, let*

$$(1) \quad X_j = \sum_{i \in \{1, \dots, s, s+j\}} (-1)^h \sum_{\lambda \in G_i} \text{sgn}(\lambda) \phi_1^{\lambda(1)} \dots \phi_s^{\lambda(s+j)} E_i,$$

where $h = i$ for $i = 1, \dots, s$ and $h = s+1$ otherwise. Then

$$\sigma(a, b^1, \dots, b^{m-s}) = \xi(a) + b^1 X_1(a) + \dots + b^{m-s} X_{m-s}(a)$$

is the desired parametrization.

Remark 1.3. The section N^* does not need to be global. The theorem remains valid if one merely assumes the existence of a local section $N^*(p)$ in a neighborhood U_p of each point $p \in S$, along with the compatibility condition

$$N^*(p) = \pm N^*(q) \quad \forall p, q \text{ such that } U_p \cap U_q \neq \emptyset.$$

Remark 1.4. If the distribution \mathcal{D} is obtained by restricting to S the tangent bundle of an arbitrary submanifold $M \supset S$, then the map $\phi(\cdot, N^*)$ coincides with the restriction to TS of the shape operator of M with respect to N^* . In this case the theorem provides conditions for the existence of a constant-rank, first-order approximation of M along S .

Remark 1.5. When $s = 1$, formula (1) reduces to $X_j = \phi^1 E_{1+j} - \phi^{1+j} E_1$, and we thus retrieve the third item in [26, Theorem 1.3]. The formula is new from $s = 2$, when it reduces to $X_j = -(\phi_1^2 \phi_2^{2+j} - \phi_1^{2+j} \phi_2^2) E_1 + (\phi_1^1 \phi_2^{2+j} - \phi_1^{2+j} \phi_2^1) E_2 - (\phi_1^1 \phi_2^2 - \phi_1^2 \phi_2^1) E_{j+2}$.

Clearly, when $c = 1$, the condition on ϕ_p is automatically satisfied, and Problem 1.1 becomes generically well posed.

Corollary 1.6. *Suppose that $c = 1$. If there exists a section N^* of \mathcal{D}^\perp such that the shape operator $A^* = A_{N^*}$ of S in direction N^* is nonsingular, then the geometric Cauchy problem for rank- s hypersurfaces of \mathbb{R}^{m+1} has a solution.*

The proof of Theorem 1.2 will be given in section 3. It is based on the well-known fact that any m -dimensional rank- s submanifold of \mathbb{R}^{m+c} admits a foliation by totally geodesic leaves, of dimension $m - s$, along which the tangent space is constant; the converse holds under the additional assumption that the foliation is unique. We will exploit this fact in the next section, where we will derive a sufficient criterion for an $(m - s)$ -ruled submanifold to be rank- s . Such criterion extends a classical result about developable surfaces in \mathbb{R}^3 [20, Theorem 54.1]; see also [30].

2. CONSTANT-RANK SUBMANIFOLDS

The purpose of this section is to generalize [26, Lemma 3.7] to $(m - s)$ -ruled submanifolds.

Definition 2.1. Let M be an m -dimensional embedded submanifold of \mathbb{R}^{m+c} . We say that M is $(m - s)$ -ruled if it admits a foliation by $(m - s)$ -dimensional totally geodesic submanifolds of \mathbb{R}^{m+c} , called *rulings*. In particular, we say that M is *properly $(m - s)$ -ruled* when the foliation is unique.

Given a s -dimensional embedded submanifold S of \mathbb{R}^{m+c} , the following lemma is the key to extending S to an $(m - s)$ -ruled submanifold. Recall that two embedded submanifolds S, S' of a smooth manifold are said to *intersect transversely* if for each $p \in S \cap S'$, the tangent spaces $T_p S$ and $T_p S'$ together span the ambient tangent space.

Lemma 2.2. *Let \mathcal{F} be a distribution of rank $m - s$ along S such that*

$$\dim(T_p S \oplus \mathcal{F}_p) = m \quad \forall p \in S.$$

Then the subset

$$\{p + x \mid p \in S, x \in \mathcal{F}_p\}$$

defines an $(m - s)$ -ruled submanifold $M_{\mathcal{F}}$ in a neighborhood of S . Conversely, any $(m - s)$ -ruled submanifold containing S and whose rulings intersect S transversely admits a description of this form.

Proof. We first prove a local version of the lemma. Let $\xi: A \rightarrow \mathbb{R}^{m+c}$ be a local parametrization of S , let (X_1, \dots, X_{m-s}) be a local frame for \mathcal{F} over A , and, for $B \subset \mathbb{R}^{m-s}$ open, define $\sigma: A \times B \rightarrow \mathbb{R}^{m+c}$ by

$$\sigma(a, b^1, \dots, b^{m-s}) = \xi(a) + b^1 X_1(a) + \dots + b^{m-s} X_{m-s}(a).$$

Since the partial derivatives of σ are linearly independent at $(a, 0)$, they remain linearly independent in some neighborhood U_a of $(a, 0)$ in $A \times B$. Moreover, after shrinking U_a and B if necessary, we may assume that $\sigma(U_a)$ is embedded and there exists a neighborhood V_a of $\xi(A)$ in \mathbb{R}^{m+c} such that

$$\sigma(U_a) = V_a \cap \sigma(A \times B);$$

to see this, note that otherwise, by making B arbitrarily close to 0, the assumption that S is embedded would be violated. Let $M := \cup_{a \in A} U_a$, and let $q \in M$. Since

M is an immersed submanifold, the rank theorem [21, Theorem 4.12] implies the existence of a slice chart for M in \mathbb{R}^{m+c} around q . Hence M satisfies the local slice criterion for embedded submanifolds [21, Theorem 5.8], which is the desired local conclusion.

To prove the full statement, let $\{\xi_i: A_i \rightarrow \mathbb{R}^{m+c}\}_{i=1}^r$ be a set of local parametrizations of S covering S , and let $\{\sigma_i: A_i \times B_i \rightarrow \mathbb{R}^{m+c}\}_{i=1}^r$ be the corresponding maps σ , as defined earlier. Then we know that for each i there exists a neighborhood U_i of $(A_i, 0)$ in $A_i \times B_i$ such that $\sigma_i(U_i)$ is embedded. Shrinking U_1, \dots, U_r along the second factor if necessary, we may assume that there are neighborhoods V_1, \dots, V_r of $\xi_1(A_1), \dots, \xi_r(A_r)$ in \mathbb{R}^{m+c} such that

$$\sigma_i(U_i) = V_i \cap (\cup_{i=1}^r \sigma_i(U_i)) \quad \forall i = 1, \dots, r.$$

As before, this implies the existence of a slice chart for $M_i = \sigma_i(U_i)$ in \mathbb{R}^{m+c} around every $q \in M_i$. Hence $M := \cup_i M_i$ is the desired $(m-s)$ -ruled submanifold. \square

Next we define constant-rank submanifolds and give two alternative characterizations of them.

Definition 2.3. Let M be an m -dimensional embedded submanifold of \mathbb{R}^{m+c} . The *index of relative nullity* of M at a point $p \in M$ is the dimension of the kernel Δ of the second fundamental form α at p :

$$\Delta = \{x \in T_p M \mid \alpha(x, \cdot) = 0\}.$$

We say that M is a *constant-rank* submanifold if the index of relative nullity is constant. In particular, we say that M is *rank- s* when the index is equal to $m-s$.

Theorem 2.4 ([19, Lemma 3.1]). *The following statements are equivalent:*

- (1) M is rank- s .
- (2) The Gauss map $M \rightarrow Gr(m, m+c)$ has rank s everywhere; here $Gr(m, m+c)$ denotes the Grassmannian of m -dimensional subspaces of \mathbb{R}^{m+c} .
- (3) M is properly $(m-s)$ -ruled with constant tangent space along the $(m-s)$ -rulings.

Thus, under very mild conditions, rank- s submanifolds containing S are locally in one-to-one correspondence with distributions \mathcal{F} of rank $m-s$ along S satisfying

$$T_p M_{\mathcal{F}} = T_{p+x} M_{\mathcal{F}} \quad \forall x \in \mathcal{F}_p \text{ such that } p+x \in M_{\mathcal{F}}.$$

The next lemma turns this condition into a relation involving the ambient covariant derivative along S and represents the sought generalization of [26, Lemma 3.7].

Lemma 2.5. *The tangent space of $M_{\mathcal{F}}$ is constant along $x \in \mathcal{F}_p$ if and only if*

$$\nabla_v X \in T_p M_{\mathcal{F}} \quad \forall v \in T_p S,$$

where X is any extension of x to a local section of \mathcal{F} .

Proof. Let $\xi: A \rightarrow \mathbb{R}^{m+c}$ be a local parametrization of S around p , and let $(X_j)_{j=1}^{m-s}$ be a local frame for \mathcal{F} over A . Then the map $\sigma: A \times \mathbb{R}^{m-s} \rightarrow \mathbb{R}^{m+c}$ defined by

$$\sigma(a, b^1, \dots, b^{m-s}) = \xi(a) + b^1 X_1(a) + \dots + b^{m-s} X_{m-s}(a)$$

is a local parametrization of $M_{\mathcal{F}}$ in a neighborhood of p . Since the tangent space $T_{(a, 0, \dots, 0, b^j, 0, \dots, 0)} \sigma$ of σ at $(a, 0, \dots, 0, b^j, 0, \dots, 0)$ is spanned by

$$\partial_1 \xi(a) + b^j \partial_1 X_j(a), \dots, \partial_s \xi(a) + b^j \partial_s X_j(a), X_1(a), \dots, X_{m-s}(a),$$

it follows that

$$T_{(a,0,\dots,0,b^j,0,\dots,0)}\sigma = T_{(a,0)}\sigma = \text{span}(\partial_1\xi(a), \dots, \partial_s\xi(a), X_1(a), \dots, X_{m-s}(a))$$

if and only if $\partial_i X_j(a) \in T_{(a,0)}\sigma$ for every $i = 1, \dots, s$; by linearity, this is equivalent to all directional derivatives of X_j being in $T_{(a,0)}\sigma$ at a . Now, to complete the proof, it is enough to note that any of the elements of the frame $(X_j)_{j=1}^{m-s}$ may be chosen to coincide with x at p . \square

Remark 2.6. It is easy to see, using orthogonality of $T_p M_{\mathcal{F}}$ and $(T_p M_{\mathcal{F}})^\perp$, that the condition $\nabla_v X \in T_p M_{\mathcal{F}}$ in Lemma 2.5 only depends on X through $X_p = x \in T_p M_{\mathcal{F}}$. Hence the lemma continues to hold if “local section of \mathcal{F} ” is replaced by “local section of $TM_{\mathcal{F}}$ ”.

Remark 2.7. By the properties of the covariant derivative, Lemma 2.5 implies that the tangent space of $M_{\mathcal{F}}$ is constant along every $x \in \mathcal{F}$ if and only if

$$\nabla_{v_i} X_j \in T_p M_{\mathcal{F}} \quad \forall i, j,$$

where (v_1, \dots, v_s) is a basis of $T_p S$ and (X_1, \dots, X_{m-s}) a local frame for \mathcal{F} around p .

3. PROOF OF THE MAIN RESULT

In this section we prove Theorem 1.2. The proof of the first two items is based on the following lemma.

Lemma 3.1. *Suppose that there exists a section N^* of \mathcal{D}^\perp such that the shape operator A^* of S in direction N^* is nonsingular at p . Then*

$$T_p S \cap \phi_p(T_p S, N_p^*)^\perp = \{0\}.$$

Proof. Let π_S denote orthogonal projection onto TS , so that

$$A_N = \pi_S \circ \phi(\cdot, N).$$

Suppose, towards a contradiction, that there exists a *nonzero* vector $v \in T_p S$ such that

$$\langle v, \phi(w, N^*) \rangle = 0 \quad \forall w \in T_p S.$$

Since $\phi(\cdot, N)$ is self-adjoint with respect to the metric, this implies

$$A^*(v) = 0,$$

contradicting the assumption that A^* has full rank at p . \square

Proof of Theorem 1.2(i)–(ii). According to Lemmas 2.2 and 2.5, we need to find an $(m-s)$ -dimensional subspace \mathcal{F}_p of \mathcal{D}_p satisfying $T_p S \cap \mathcal{F}_p = \{0\}$ and

$$\nabla_v x \in \mathcal{D}_p \quad \forall (v, x) \in T_p S \times \mathcal{F}_p,$$

where x is being extended arbitrarily to a local section X of \mathcal{D} . Note that the latter condition is equivalent to

$$(2) \quad \langle \nabla_v x, n \rangle = 0 \quad \forall (v, x, n) \in T_p S \times \mathcal{F}_p \times \mathcal{D}_p^\perp;$$

using orthogonality of x and n , we can rewrite (2) as

$$\langle x, \phi_p(v, n) \rangle = 0 \quad \forall (v, x, n) \in T_p S \times \mathcal{F}_p \times \mathcal{D}_p^\perp$$

or, more succinctly, as

$$\mathcal{F}_p \subset (\text{Im } \phi_p)^\perp.$$

Suppose that there exists a section N^* of \mathcal{D}^\perp such that the shape operator A^* of S in direction N^* is nonsingular at p , and set

$$\mathcal{F}_p^* := \mathcal{D}_p \cap \phi_p(T_p S, N_p^*)^\perp.$$

Then, since $A_{N^*} = \pi_S \circ \phi(\cdot, N^*)$, we deduce that \mathcal{F}_p^* is a subspace of dimension $m - s$; besides, it has trivial intersection with $T_p S$ by Lemma 3.1. Observing that

$$(\text{Im } \phi_p)^\perp \subset \phi_p(T_p S, N_p^*)^\perp,$$

we conclude that \mathcal{F}^* can only define a solution of our problem when $\text{Im } \phi_p = \phi_p(T_p S, N_p^*)$.

Next we check that \mathcal{F}^* does indeed define a solution. To this end, let $M^* := M_{\mathcal{F}^*}$ be an embedded submanifold containing S , whose existence is guaranteed by Lemma 2.2. First note that, by construction, the shape operator A^* of S is nothing but the restriction to TS of a shape operator of M^* . Hence, as the tangent space is constant along \mathcal{F}^* , the index of relative nullity of M^* equals $m - s$ along S . By the same reason, extending N^* via parallel translation in the ambient space to a unit normal vector field along M^* , we have that

$$A_{p+x}^*(v) = A_p^*(v) \quad \forall v \in \mathcal{D}_p \equiv T_{p+x} M^*,$$

which implies that the index of relative nullity of M^* is constant, as claimed. \square

It remains to prove Theorem 1.2(iii). To set the stage, we first recall the notion of $((m - 1)$ -fold) vector cross product.

Let V be an m -dimensional oriented inner product space, and let (e_1, \dots, e_m) be an orthonormal basis of V . A *vector cross product on V* is a multilinear map $\cdot \times \cdots \times \cdot : V^{m-1} \rightarrow V$ such that

$$\begin{aligned} \langle e_1 \times \cdots \times e_{m-1}, e_1 \rangle &= \cdots = \langle e_1 \times \cdots \times e_{m-1}, e_{m-1} \rangle = 0, \\ \langle e_1 \times \cdots \times e_{m-1}, e_1 \times \cdots \times e_{m-1} \rangle &= \det(\langle e_i, e_j \rangle)_{i,j=1}^{m-1}, \\ (e_1, \dots, e_{m-1}, e_1 \times \cdots \times e_{m-1}) &\text{ is positively oriented.} \end{aligned}$$

It is well known that there is only one vector cross product on V . It is given by

$$v_1 \times \cdots \times v_{m-1} = \star(v_1 \wedge \cdots \wedge v_{m-1}),$$

where \star denotes the Hodge star operator and \wedge the wedge product on V ; see [4]. More generally, if v_1, \dots, v_{m-1} are continuous sections of a smooth oriented vector bundle over a smooth manifold, then both \star and \wedge are globally defined, and the previous equation defines their cross product.

Our proof of Theorem 1.2(iii) is based on the following lemma.

Lemma 3.2. *Let W be an s -dimensional subspace of V , and let ρ be a linear map $W \rightarrow V$; without loss of generality, we may assume that the first s elements of (e_1, \dots, e_m) span W and the first r elements span $\text{coim } \rho = (\ker \rho)^\perp$. Suppose that the orthogonal complement $\varphi(W)^\perp$ of $\varphi(W)$ in V has trivial intersection with $\text{coim } \rho$. Then*

$$x_j = \varphi(e_1) \times \cdots \times \varphi(e_r) \times e_{r+1} \times \cdots \times \widehat{e_{r+j}} \times \cdots \times e_m, \quad j = 1, \dots, m - r$$

defines an orthogonal basis of $\varphi(W)^\perp$; here the hat indicates that e_{r+j} is omitted.

Proof. By construction, the vector x_j belongs to the intersection between $\varphi(W)^\perp$ and the subspace spanned by e_1, \dots, e_r, e_{r+j} . Since $\varphi(W)^\perp$ has trivial intersection with $\text{span}(e_1, \dots, e_r)$, it follows that x_1, \dots, x_{m-r} all lie in orthogonal subspaces, showing that they indeed span $\varphi(W)^\perp$. \square

Proof of Theorem 1.2(iii). Let (E_1, \dots, E_m) be a smooth orthonormal frame for $\mathcal{D}|_{\xi(A)}$ whose first s elements span TS . Equipping $\mathcal{D}|_{\xi(A)}$ with the orientation naturally induced by that frame, the vector cross product becomes well defined on $\mathcal{D}|_{\xi(A)}$. Then, for each $j = 1, \dots, m - s$, let

$$(3) \quad X_j = \phi(E_1, N^*) \times \cdots \times \phi(E_s, N^*) \times E_{s+1} \times \cdots \times \widehat{E_{s+j}} \times \cdots \times E_m,$$

where the hat indicates that E_{s+j} is omitted. Since $\text{coim } \phi = TS$ by assumption, we deduce from Lemmas 3.1 and 3.2 that (X_1, \dots, X_{m-s}) is an orthogonal frame for $\phi(TS, N^*)^\perp$.

It remains to compute (3) in coordinates. Let $\phi_i^k = \langle \phi(E_i, N^*), E_k \rangle$. Then

$$\phi(E_i, N^*) = \phi_i^1 E_1 + \cdots + \phi_i^m E_m.$$

Substitution into (3) gives

$$\begin{aligned} X_j = & (\phi_1^1 E_1 + \cdots + \phi_1^s E_s + \phi_1^{s+j} E_{s+j}) \times \cdots \times (\phi_s^1 E_1 + \cdots + \phi_s^s E_s + \phi_s^{s+j} E_{s+j}) \\ & \times E_{s+1} \times \cdots \times \widehat{E_{s+j}} \times \cdots \times E_m, \end{aligned}$$

which implies (1). \square

REFERENCES

1. Juan A. Aledo, José A. Gálvez, and Pablo Mira, *A D'Alembert formula for flat surfaces in the 3-sphere*, J. Geom. Anal. **19** (2009), no. 2, 211–232. MR 2481959
2. David Brander and Martin Svensson, *The geometric Cauchy problem for surfaces with Lorentzian harmonic Gauss maps*, J. Differential Geom. **93** (2013), no. 1, 37–66. MR 3019511
3. David Brander and Peng Wang, *On the Björling problem for Willmore surfaces*, J. Differential Geom. **108** (2018), no. 3, 411–457. MR 3770847
4. Robert B. Brown and Alfred Gray, *Vector cross products*, Comment. Math. Helv. **42** (1967), 222–236. MR 222105
5. Antonio Bueno, *The Björling problem for prescribed mean curvature surfaces in \mathbb{R}^3* , Ann. Global Anal. Geom. **56** (2019), no. 1, 87–96. MR 3962027
6. R. M. B. Chaves, M. P. Dussan, and M. Magid, *Björling problem for timelike surfaces in the Lorentz-Minkowski space*, J. Math. Anal. Appl. **377** (2011), no. 2, 481–494. MR 2769151
7. Shiing-shen Chern and Nicolaas H. Kuiper, *Some theorems on the isometric imbedding of compact Riemann manifolds in euclidean space*, Ann. of Math. (2) **56** (1952), 422–430. MR 50962
8. Adriana A. Cintra, Francesco Mercuri, and Irene I. Onnis, *The Björling problem for minimal surfaces in a Lorentzian three-dimensional Lie group*, Ann. Mat. Pura Appl. (4) **195** (2016), no. 1, 95–110. MR 3453594
9. M. Dajczer, L. Florit, and R. Tojeiro, *Euclidean hypersurfaces with genuine deformations in codimension two*, Manuscripta Math. **140** (2013), no. 3-4, 621–643. MR 3019143
10. M. Dajczer and Th. Vlachos, *The infinitesimally bendable Euclidean hypersurfaces*, Ann. Mat. Pura Appl. (4) **196** (2017), no. 6, 1961–1979. MR 3714748
11. Marcos Dajczer and Luis A. Florit, *Genuine rigidity of Euclidean submanifolds in codimension two*, Geom. Dedicata **106** (2004), 195–210. MR 2079843
12. Marcos Dajczer and Detlef Gromoll, *Gauss parametrizations and rigidity aspects of submanifolds*, J. Differential Geom. **22** (1985), no. 1, 1–12. MR 826420
13. Marcos Dajczer and Pedro Morais, *Isometric rigidity in codimension 2*, Michigan Math. J. **58** (2009), no. 3, 759–770. MR 2595563
14. Marcos Dajczer and Ruy Tojeiro, *Submanifold theory: Beyond an introduction*, Universitext, Springer, New York, 2019. MR 3969932

15. Manfredo P. do Carmo, *Differential geometry of curves & surfaces*, second ed., Dover Publications, Mineola, NY, 2016. MR 3837152
16. M. P. Dussan, A. P. Franco Filho, and M. Magid, *The Björling problem for timelike minimal surfaces in \mathbb{R}_1^4* , Ann. Mat. Pura Appl. (4) **196** (2017), no. 4, 1231–1249. MR 3673665
17. Luis A. Florit and Guilherme M. de Freitas, *Classification of codimension two deformations of rank two Riemannian manifolds*, Comm. Anal. Geom. **25** (2017), no. 4, 751–797. MR 3731641
18. Alfred Gray, Elsa Abbena, and Simon Salamon, *Modern differential geometry of curves and surfaces with Mathematica®*, third ed., Studies in Advanced Mathematics, Chapman & Hall/CRC, Boca Raton, FL, 2006. MR 2253203
19. Philip Hartman, *On isometric immersions in Euclidean space of manifolds with non-negative sectional curvatures*, Trans. Amer. Math. Soc. **115** (1965), 94–109. MR 202094
20. Erwin Kreyszig, *Introduction to differential geometry and Riemannian geometry*, Mathematical Expositions, no. 16, University of Toronto Press, Toronto, 1968. MR 226507
21. John M. Lee, *Introduction to smooth manifolds*, second ed., Graduate Texts in Mathematics, no. 218, Springer, New York, 2013. MR 2954043
22. Irina Markina and Matteo Raffaelli, *Flat approximations of hypersurfaces along curves*, Manuscripta Math. **160** (2019), no. 3–4, 315–325. MR 4016641
23. Antonio Martínez and Francisco Milán, *The geometric Cauchy problem for the hyperbolic Hessian one equation*, Nonlinear Anal. **125** (2015), 323–333. MR 3373588
24. Francisco Milán, *The indefinite affine Cauchy problem*, Nonlinear Anal. **196** (2020), Paper No. 111819, 11 pp. MR 4074625
25. Christos-Raent Onti, *On complete conformally flat submanifolds with nullity in Euclidean space*, Results Math. **75** (2020), no. 3, Paper No. 106, 7 pp. MR 4119521
26. Matteo Raffaelli, *The geometric Cauchy problem for rank-one submanifolds*, Comm. Anal. Geom., to appear.
27. ———, *Classification of rank-one submanifolds*, Results Math. **78** (2023), no. 6, Paper No. 208, 13 pp. MR 4630556
28. Umberto Sbrana, *Sulla deformazione infinitesima delle ipersuperficie*, Ann. Mat. Pura Appl. **15** (1908), 329–348.
29. Francisco Vittono, *On the nullity distribution of a submanifold of a space form*, Math. Z. **272** (2012), no. 1–2, 1–16. MR 2968211
30. Kentaro Yano, *On the torse-forming directions in Riemannian spaces*, Proc. Imp. Acad. Tokyo **20** (1944), 340–345. MR 14777

SCHOOL OF MATHEMATICS, GEORGIA INSTITUTE OF TECHNOLOGY, ATLANTA, GEORGIA 30332

Email address: raffaelli@math.gatech.edu