# Parameter estimation for fractional stochastic heat equations : Berry-Esséen bounds in CLTs

Soukaina Douissi \*and Fatimah Alshahrani †

Tuesday 10<sup>th</sup> September, 2024

Abstract: The aim of this work is to estimate the drift coefficient of a fractional heat equation driven by an additive space-time noise using the Maximum likelihood estimator (MLE). In the first part of the paper, the first N Fourier modes of the solution are observed continuously over a finite time interval [0,T]. The explicit upper bounds for the Wasserstein distance for the central limit theorem of the MLE is provided when  $N \to \infty$  and/or  $T \to \infty$ . While in the second part of the paper, the N Fourier modes are observed at uniform time grid :  $t_i = i\frac{T}{M}$ , i = 0, ..., M, where M is the number of time grid points. The consistency and asymptotic normality are studied when  $T, M, N \to +\infty$  in addition to the rate of convergence in law in the CLT.

Mathematics Subject Classification 2020: 62F12; 60F05; 60G15; 60H15; 60H07.

**Keywords**: Fractional stochastic partial differential equations; Parameter estimation; Rate of normal convergence of the MLE, Malliavin calculus, Wasserstein distance.

#### 1 Introduction

The aim of this study is to solve the drift parameter estimation problem  $\theta > 0$  of the following fractional stochastic heat equation :

$$\begin{cases}
dY_{\theta}(t,x) = -\theta \left(-\Delta\right)^{\alpha} Y_{\theta}(t,x) dt + \sum_{n \in \mathbb{N}} \delta_{n}^{-\gamma} e_{n}(x) dw_{n}(t), & 0 \leqslant t \leqslant T, \quad x \in G. \\
Y_{\theta}(0,x) = Y_{0} \quad x \in G,
\end{cases} \tag{1}$$

where T>0,  $\theta>0$ ,  $\alpha>0$ ,  $\gamma\geqslant 0$ , G is a bounded and smooth domain in  $\mathbb{R}^d$ ,  $d\geqslant 1$ .  $\Delta$  is the Laplace operator on G with zero boundary conditions and  $H^r(G)$  for  $r\in\mathbb{R}$ , denotes the corresponding Sobolev spaces. The initial condition of (1) is such that  $Y_0\in H^r(G)$  for some  $r\in\mathbb{R}$ . The process  $\{Y_\theta(t,x),x\in G,t\in[0,T]\}$  is defined on a filtered probability space  $(\Omega,\mathcal{F},\{\mathcal{F}_t\}_{t\geqslant 0},\mathbf{P})$  on which we consider a family  $\{w_j,j\geqslant 1\}$  of independent standard Brownian motions.

The set  $\{e_n, n \in \mathbb{N}\}$  are the eigenfunctions of  $\Delta$  that form a complete orthonormal system in  $L^2(G)$ , the corresponding eigenvalues  $h_n, n \in \mathbb{N}$  can be sorted such that  $0 < -h_1 \leqslant -h_2 \leqslant \dots$ 

Let us introduce the notation  $\delta_n := (-h_n)^{1/2}$ ,  $n \in \mathbb{N}$ , so, there exists a positive constant  $\bar{\sigma}$  such that, see [1]

$$\lim_{n \to +\infty} |\delta_n|^2 n^{-2/d} = \bar{\sigma}. \tag{2}$$

<sup>\*</sup>Ecole Nationale des Sciences Appliquées, BP 575, Avenue Abdelkrim Khattabi, 40000, Guéliz-Marrakech Email:s.douissi@uca.ma

<sup>&</sup>lt;sup>†</sup>Department of Mathematical Sciences, College of Science, Princess Nourah bint Abdulrahman University, P.O. Box 84428, Riyadh 11671, Saudi Arabia. Email:fmalshahrani@pnu.edu.sa

Assuming that  $2(\gamma - r) > d$ , then (see for instance [2, 3, 4]) equation (1) has a unique solution  $Y_{\theta}$  weak in the PDE sense and strong in the probability sense.

In the following, it is assumed that  $r \ge 0$  and  $2\gamma > d$ . Let  $y_n, n \ge 1$  be the Fourier coefficient of the solution  $Y_{\theta}$  of (1) with respect to  $e_n, n \ge 1$ , i.e.  $y_n(t) = (Y_{\theta}(t), e_n)_0, n \in \mathbb{N}$ . Let  $H^N$  be the finite dimensional subspace of  $L^2(G)$  generated by  $\{e_n, n = 1, ..., N\}$  and denote  $P_N$  the projection operator of  $L^2(G)$  into  $H^N$  and put  $Y_{\theta}^N = P_N Y_{\theta}$ . Each Fourier mode  $y_n, n \ge 1$  follows the dynamics of an Ornstein-Uhlenbeck process given by

$$dy_n(t) = -\theta \delta_n^{2\alpha} y_n(t) dt + \delta_n^{-\gamma} dw_n(t), \quad y_n(0) = (Y_0, e_n), \quad t \geqslant 0.$$
(3)

In the following, we will denote by  $\mathbf{P}_{T,N}^{\theta}$  the probability measure on  $C([0,T];H^N)$  generated by  $Y_{\theta}^N$ . We will also fix in the rest of the paper, a parameter value  $\theta_0 > 0$ . Then since the measures  $\{\mathbf{P}_{T,N}^{\theta}, \theta > 0\}$  are equivalent, hence applying Girsanov's theorem we obtain the following Likelihood Ratio or Radon-Nikodym derivative:

$$\frac{d\mathbf{P}_{T,N}^{\theta}}{d\mathbf{P}_{T,N}^{\theta_0}}(Y_{\theta}^{N}) = \exp\left(-(\theta - \theta_0) \sum_{n=1}^{N} \delta_n^{2\alpha + \gamma} \int_0^T y_n(t) dy_n(t) - \frac{(\theta^2 - \theta_0^2)}{2} \sum_{n=1}^{N} \delta_n^{4\alpha + 2\gamma} \int_0^T y_n^2(t) dt\right).$$

Maximizing the log-likelihood ratio with respect to  $\theta$  gives the Maximum Likelihood Estimator (MLE) for  $\theta$ 

$$\hat{\theta}_{T,N} := -\frac{\sum_{n=1}^{N} \delta_n^{2\alpha + 2\gamma} \int_0^T y_n(t) dy_n(t)}{\sum_{n=1}^{N} \delta_n^{4\alpha + 2\gamma} \int_0^T y_n^2(t) dt}, \quad N \in \mathbb{N}, \quad T > 0.$$
(4)

Moreover, using (3), we get : 
$$\theta - \hat{\theta}_{T,N} = \frac{\sum\limits_{n=1}^{N} \delta_n^{2\alpha+\gamma} \int_0^T y_n(t) dw_n(t)}{\sum\limits_{n=1}^{N} \delta_n^{4\alpha+2\gamma} \int_0^T y_n^2(t) dt}, \quad N \in \mathbb{N}, \quad T > 0.$$

The consistency and asymptotic normality of the MLE  $\hat{\theta}_{T,N}$  have been studied in several papers when  $N \to \infty$  and/or  $T \to \infty$ , see for instance [5, 6] and [7]. However, to utilize the asymptotic distribution of an estimator it is essential that the rate of convergence is known. For the best of our knowledge, no result of the Berry-Esséen type is known for the distribution of the MLE  $\hat{\theta}_{T,N}$  of the drift parameter  $\theta$  of the SPDE (1).

The aim of this work is to study the rate of convergence for the central limit theorem of the MLE of  $\theta$  in its continuous and discrete versions. The article is presented as follows: Section 2 consist of some of the essential tools needed from the analysis on Wiener space and Malliavin calculus.

In Section 3, we provide explicit bounds of the convergence of the MLE to a Gaussian random variable for the Wasserstein distance, considering three different scenarios: when  $N \to +\infty$  and T is fixed, when  $T \to +\infty$  and N is fixed and when both  $N, T \to +\infty$ , see Theorem 2 and Corollary 5. The key of the proof is the fact that the numerator of  $\theta_0 - \hat{\theta}_{T,N}$  is a second chaos random sequence that depends on both N and T. A rigorous study of the third and fourth cumulant of the numerator along with the application of the Optimal fourth moment theorem [8] lead to a CLT with the rates of its convergence in law. It is also worth noting that the proof of the consistency part relies mainly on the properties of multiple Wiener integrals in conjunction with Borelli-Catelli Lemma.

In Section 4, we study the asymptotic properties of an approximate version of the MLE  $\hat{\theta}_{T,N}$ , which we denote by  $\tilde{\theta}_{T,N,M}$ , defined by  $\tilde{\theta}_{T,N,M} := -\frac{\sum_{n=1}^{N} \delta_n^{2\alpha+2\gamma} \sum_{i=1}^{M} y_n(t_{i-1})[y_n(t_i)-y_n(t_{i-1})]}{\Delta_M \sum_{n=1}^{N} \delta_n^{4\alpha+2\gamma} \sum_{i=1}^{M} y_n^2(t_{i-1})}$ , where we simply discretized the

numerator and denominator of  $\hat{\theta}_{N,T}$ , considering that the first N Fourier modes are now observed at discrete time instant  $t_i = i \frac{T}{M}$ , i = 0,...,M where M denotes the number of time grid points. We proved that

when  $\frac{T}{M}N^{2\alpha/d} \to 0$  as  $T, M, N \to +\infty$ , the estimator  $\tilde{\theta}_{T,N,M}$  is consistent and when  $\frac{T^{3/2}N^{\frac{3\alpha}{d}+\frac{1}{2}}}{M} \to 0$  as  $T, M, N \to +\infty$ , then  $\tilde{\theta}_{T,N,M}$  is asymptotically Gaussian. We also derive the rate of the convergence in law of  $\tilde{\theta}_{T,N,M}$  for the Wasserstein distance, see Theorem 13 for more details.

#### 2 Elements of analysis on Wiener space.

In this section, we give a concise overview of some elements from Malliavin calculus and elements of Gaussian analysis. For additional details about this topic, the interested reader is referred to the following references [9], and [10].

Consider W a Brownian motion defined on a probability space  $(\Omega, \mathcal{F}, \mathbf{P})$ . The Wiener integral of a deterministic function  $g \in L^2([0,T])$ ,  $\int_0^T g(s)dW(s)$  can also be denoted by W(g). The Hilbert space  $\mathcal{H} := L^2([0,T])$  is endowed with the inner product

$$\mathbf{E}[W(f)W(g)] = \int_0^T f(s)g(s)ds = \langle f, g \rangle_{L^2([0,T])}$$

In the following, we will consider that  $\mathcal{F}$  is generated by W and  $L^2(\Omega) := L^2(\Omega, \mathcal{F}, \mathbf{P})$ . For all  $p \ge 1$ ,  $\mathcal{H}_p^W$  denotes the p-th Wiener chaos of W is the closed linear subspace of  $L^2(\Omega)$  generated by  $\{H_p(W(g)), g \in L^2([0,T]), \|g\|_{L^2([0,T])} = 1\}$ , where  $H_p$  is the p-th Hermite polynomial defined by:

$$H_p(x) = (-1)^p e^{\frac{x^2}{2}} \frac{d^p}{dx^p} e^{-\frac{x^2}{2}}, \quad p \geqslant 1,$$

and  $H_0(x) = 1$ .

In the rest of this section, the notations  $\mathcal{H}^{\otimes p}$  and  $\mathcal{H}^{\odot p}$  for any integer  $p \geqslant 1$ , mean the p-th tensor product and the p-th symmetric tensor product of  $\mathcal{H}$  respectively.

The linear mapping defined by  $I_p^W(g^{\otimes p}) := p! H_p(W(g))$  is an isometry between  $\mathcal{H}^{\odot p} = L_s^2([0,T]^p)$  equipped with the norm  $\sqrt{p!} \|.\|_{\mathcal{H}^{\odot p}}$  and the Wiener chaos of order p under  $L^2(\Omega)$ 's norm. For  $h \in \mathcal{H}^{\otimes p}$ ,  $I_p^W(h)$  is a multiple Wiener-Itô integral of order p with respect to W and we can write

$$I_p^W(h) = \int_{[0,T]^p} h(x_1,...,x_p)dW(x_1)...dW(x_p).$$

Multiple Wiener integrals have many properties, we recall the main ones that we will need in our analysis. We start with the **isometry property**, which states that if  $h_1 \in \mathcal{H}^{\odot p}$  and  $h_2 \in \mathcal{H}^{\odot q}$ , for  $p, q \geqslant 1$ . Then the following holds:

$$\mathbf{E}\left[I_p^W(h_1)I_q^W(h_2)\right] = \begin{cases} p! \times \langle h_1, h_2 \rangle_{L^2(\mathbb{R}^p)} & \text{if } p = q \\ 0 & \text{if } p \neq q. \end{cases}$$
 (5)

The second property we will be using is the so-called **product formula**: Let  $p, q \ge 1$ . If  $h_1 \in \mathcal{H}^{\odot p}$  and  $h_2 \in \mathcal{H}^{\odot q}$  then

$$I_{p}(h_{1})I_{q}(h_{2}) = \sum_{l=0}^{p \wedge q} l! \binom{p}{l} \binom{q}{l} I_{p+q-2l}(h_{1} \widetilde{\otimes}_{l} h_{2}).$$
 (6)

where  $h_1 \otimes_l h_2$  is the contraction of  $h_1$  and  $h_2$  of order l which is an element of  $\mathcal{H}^{\otimes (p+q-2l)}$  defined by

$$(h_1 \otimes_l h_2)(r_1, \dots, r_{p-l}, s_1, \dots, s_{q-l})$$

$$:= \int_{[0,T]^{p+q-2l}} h_1(r_1, \dots, r_{p-l}, v_1, \dots, v_l) h_2(s_1, \dots, s_{q-l}, v_1, \dots, v_l) dv_1 \cdots dv_l.$$

The notation  $h_1 \approx_l h_2$  in (6) means the symmetrization of  $h_1 \otimes_l h_2$  which is defined for a function h by

$$\tilde{h}(y_1, ..., y_q) = \frac{1}{q!} \sum_{\sigma} h(y_{\sigma(1)}, ..., y_{\sigma(q)})$$

where all permutations  $\sigma$  of  $\{1,...,q\}$  are included in the sum. The special case p=q=1 in (6) is useful in practice can be written as follows:

$$I_1(h_1)I_1(h_2) = 2^{-1}I_2(h_1 \otimes h_2 + h_2 \otimes h_1) + \langle h_1, h_2 \rangle_{\mathcal{H}}.$$
 (7)

Another important property in Wiener chaos is **the hypercontractivity**: If  $q \ge 1$ , then for any  $p \ge 2$ , there exists C(p,q) depending only on p and q such that, for every  $X \in \bigoplus_{l=1}^{q} \mathcal{H}_{l}$ ,

$$\left(\mathbf{E}\left[|X|^{p}\right]\right)^{1/p} \leqslant C(p,q) \left(\mathbf{E}\left[|X|^{2}\right]\right)^{1/2}.$$
(8)

The constants C(p,q) above are known with some precision when  $F \in \mathcal{H}_q$ ;  $C(p,q) = (p-1)^{q/2}$ , see [9]. We also recall the **Optimal fourth moment theorem** which will be used in order to derive rates of convergences in CLTs.

Let N denote the standard normal random variable. Consider a random sequence  $X_n \in \mathcal{H}_q$ ,  $q \ge 1$  such that  $Var[X_n] = 1$  and  $(X_n)_n$  converges in distribution to N. The Fourth moment theorem proved in [?] claims that this convergence is equivalent to  $\lim_n \mathbf{E}[X_n^4] = 3$ . The following optimal estimate for  $d_{TV}(X_n, \mathcal{N})$ , known as the Optimal fourth moment theorem, was proved in [8]: with the sequence  $X_n$  as above, assuming convergence, there exist two constants c, C > 0 independent of n, such that

$$c \max \left\{ \mathbf{E} \left[ X_n^4 \right] - 3, \left| \mathbf{E} \left[ X_n^3 \right] \right| \right\} \leqslant d_{TV} \left( X_n, \mathcal{N} \right) \leqslant C \max \left\{ \mathbf{E} \left[ X_n^4 \right] - 3, \left| \mathbf{E} \left[ X_n^3 \right] \right| \right\}. \tag{9}$$

Notice that optimal bound (9) holds with  $d_{TV}$  replaced by  $d_W$ , see ([11], Remark 2.2). We recall that, for two random variables F and G, the former metrics are respectively given by

$$d_{TV}\left(F,G\right) := \sup_{A \in \mathcal{B}(\mathbb{R})} \left| \mathbf{P}\left[F \in A\right] - \mathbf{P}\left[G \in A\right]\right|, \quad d_{W}\left(F,G\right) := \sup_{h \in Lip(1)} \left| \mathbf{E}[h(F)] - \mathbf{E}[h(G)]\right|, \tag{10}$$

where Lip(1) is the set of all Lipschitz functions with Lipschitz constant  $\leq 1$ . We also recall, the definition of the cumulant of order  $p \geq 2$ ,  $k_p(F)$  of a random variable  $F = I_2(h)$ ,

$$k_p(F) = 2^{p-1}(p-1)!\langle \underbrace{h \otimes_1 \dots \otimes_1 h}_{p-1 \text{ copies of } h}, h \rangle_{\mathcal{H}^{\otimes 2}}.$$

In particular for p = 3, 4, we obtain estimates of the third and fourth cumulant respectively as follows:

$$k_{3}(F) = \mathbf{E}[F^{3}] = 8\langle h, h \otimes_{1} h \rangle_{\mathcal{H}^{\otimes 2}} \text{ and } k_{4}(F) = \mathbf{E}[F^{4}] - 3\mathbf{E}[F^{2}]^{2}$$

$$= 16 \left( \|h \otimes_{1} h\|_{\mathcal{H}^{\otimes 2}}^{2} + 2\|h\widetilde{\otimes_{1}}h\|_{\mathcal{H}^{\otimes 2}}^{2} \right)$$

$$\leq 48\|h \otimes_{1} h\|_{\mathcal{H}^{\otimes 2}}^{2}.$$
(11)

# 3 Berry-Esséen bounds for the MLE when N and T are large: Continuous time observations

For simplicity, we set  $Y_0 = 0$  hence  $y_n(0) = 0$  for all  $n \ge 1$ . Because equation (3) is linear, it is immediate to see that its solution can be expressed explicitly as follows:

$$y_n(t) = \delta_n^{-\gamma} \int_0^t e^{-\theta_0 \delta_n^{2\alpha} (t-u)} dw_n(u), \quad n = 1, ..., N.$$

Define:

$$E_{T,N} := \sum_{n=1}^{N} \delta_n^{2\alpha + \gamma} \int_0^T y_n(t) dw_n(t), \quad \psi_{T,N}^{\theta} := \frac{T}{2\theta_0} \sum_{n=1}^{N} \delta_n^{2\alpha} \sim \frac{\bar{\sigma} dT N^{\frac{2\alpha}{d} + 1}}{(4\alpha + 2d)\theta_0} \quad \text{as} \quad N, T \to +\infty.$$
 (13)

Indeed, by (2), we have  $\delta_n \sim \bar{\sigma}^{1/2} n^{1/d}$  as  $n \to +\infty$ , thus  $\sum_{n=1}^N \delta_n^{2\alpha} \sim \bar{\sigma}^{\alpha} \frac{N^{\frac{2\alpha}{d}+1}}{\frac{2\alpha}{d}+1}$ , as  $N \to +\infty$ . By (13), we can write

$$\theta_0 - \hat{\theta}_{T,N} = \frac{\sum_{n=1}^{N} \delta_n^{2\alpha + \gamma} \int_0^T y_n(t) dw_n(t)}{\sum_{n=1}^{N} \delta_n^{4\alpha + 2\gamma} \int_0^T y_n^2(t) dt} = \frac{E_{T,N}}{\langle E_{T,N} \rangle}.$$
 (14)

The numerator term of (14) is a second chaos random variable, indeed for every  $n \ge 1$ ,

$$\int_0^T y_n(t) dw_n(t) = \delta_n^{-\gamma} \int_0^T \int_0^t e^{-\theta_0 \delta_n^{2\alpha} (t-u)} dw_n(u) dw_n(t) = I_2^{w_n}(g^n),$$

where  $g^n(s,t) := \delta_n^{2\alpha} e^{-\theta_0 \delta_n^{2\alpha} |t-s|} \mathbf{1}_{[0,T]^2}(t,s)$ , n = 1,...,N. Therefore, using the linear property of multiple integrals, we can express  $E_{T,N}$  as follows

$$E_{T,N} = I_2\left(\frac{g_{T,N}}{2}\right) = \frac{1}{2} \sum_{n=1}^{N} I_2^{w_n}(g^n)$$
(15)

where  $g_{T,N} = (g^1, ..., g^N)$ . For the denominator term, using the product formula (7), we get for every n = 1, ..., N

$$y_n^2(t) = \delta_n^{-2\gamma} e^{-2\theta_0 \delta_n^{2\alpha} t} I_2^{w_n} \left( e^{\theta_0 \delta_n^{2\alpha} (u+v)} \mathbf{1}(u,v)_{[0,t]^2} \right) + \frac{\delta_n^{-2\gamma}}{2\theta_0 \delta_n^{2\alpha}} \left( 1 - e^{-2\theta_0 \delta_n^{2\alpha} t} \right).$$

Integrating over [0, T],

$$\begin{split} \int_{0}^{T} y_{n}^{2}(t)dt &= \delta_{n}^{-2\gamma} I_{2}^{w_{n}} \left( e^{\theta_{0}\delta_{n}^{2\alpha}(u+v)} \int_{u\vee v}^{T} e^{-2\theta_{0}\delta_{n}^{2\alpha}s} ds \right) + \frac{\delta_{n}^{-2\gamma}}{2\theta_{0}\delta_{n}^{2\alpha}} \left( T + \frac{e^{-2\theta_{0}\delta_{n}^{2\alpha}T} - 1}{2\theta_{0}\delta_{n}^{2\alpha}} \right) \\ &= \delta_{n}^{-2\gamma} I_{2}^{w_{n}} \left( \frac{e^{-\theta_{0}\delta_{n}^{2\alpha}|u-v|}}{2\theta_{0}\delta_{n}^{2\alpha}} \mathbf{1}(u,v)_{[0,T]^{2}} \right) - \delta_{n}^{-2\gamma} I_{2}^{w_{n}} \left( \frac{e^{-2\theta_{0}\delta_{n}^{2\alpha}T} e^{\theta_{0}\delta_{n}^{2\alpha}(u+v)}}{2\theta_{0}\delta_{n}^{2\alpha}} \mathbf{1}(u,v)_{[0,T]^{2}} \right) \\ &+ \frac{\delta_{n}^{-2\gamma}}{2\theta_{0}\delta_{n}^{2\alpha}} \left( T + \frac{e^{-2\theta_{0}\delta_{n}^{2\alpha}T} - 1}{2\theta_{0}\delta_{n}^{2\alpha}} \right) \end{split}$$

Thus

$$\langle E_{T,N} \rangle = \sum_{n=1}^{N} \delta_{n}^{4\alpha+2\gamma} \int_{0}^{T} y_{n}^{2}(t) dt = \sum_{n=1}^{N} \delta_{n}^{2\alpha} I_{2}^{w_{n}} \left( \frac{e^{-\theta_{0}|u-v|}}{2\theta_{0}} \mathbf{1}(u,v)_{[0,T]^{2}} \right) - \sum_{n=1}^{N} \delta_{n}^{2\alpha} I_{2}^{w_{n}} \left( \frac{e^{-2\theta_{0}\delta_{n}^{2\alpha}T} e^{\theta_{0}\delta_{n}^{2\alpha}(u+v)}}{2\theta_{0}} \mathbf{1}(u,v)_{[0,T]^{2}} \right) + \sum_{n=1}^{N} \frac{\delta_{n}^{2\alpha}}{2\theta_{0}} \left( T + \frac{e^{-2\theta_{0}\delta_{n}^{2\alpha}T} - 1}{2\theta_{0}\delta_{n}^{2\alpha}} \right)$$

$$= I_{2}(\frac{g_{T,N}}{2\theta_{0}}) - I_{2}(l_{T,N}) + \lambda_{T,N} = \frac{E_{T,N}}{\theta_{0}} - I_{2}(l_{T,N}) + \lambda_{T,N}$$

$$(16)$$

with  $l_{T,N} := (l^1, ..., l^N)$  where  $l^n(u, v) := \frac{\delta_n^{2\alpha}}{2\theta_0} e^{-2\theta_0 \delta_n^{2\alpha} T} e^{\theta_0 \delta_n^{2\alpha} (u+v)} \mathbf{1}(u, v)_{[0,T]^2}$  and

$$\begin{split} \lambda_{T,N} &= \frac{1}{2\theta_0} \sum_{n=1}^N \delta_n^{2\alpha} \left( T + \frac{e^{-2\theta_0 \delta_n^{2\alpha} T} - 1}{2\theta_0 \delta_n^{2\alpha}} \right) = \frac{T}{2\theta_0} \sum_{n=1}^N \delta_n^{2\alpha} + \frac{1}{4\theta_0} \sum_{n=1}^N (e^{-2\theta_0 \delta_n^{2\alpha} T} - 1) \\ &= \psi_{T,N}^{\theta_0} + \frac{1}{4\theta_0} \sum_{n=1}^N (e^{-2\theta_0 \delta_n^{2\alpha} T} - 1). \end{split}$$

By (14), (13) and (16), we can write, for every  $N \ge 1$ , T > 0,

$$\theta_0 - \hat{\theta}_{T,N} := \frac{\frac{1}{2} I_2(g_{T,N})}{\frac{1}{2\theta_0} I_2(g_{T,N}) - I_2(l_{T,N}) + \lambda_{T,N}}.$$
(17)

Note that the strong consistency of  $\hat{\theta}_{T,N}$  when  $N \to \infty$  and T fixed or  $T \to \infty$  and N fixed, and when both  $N, T \to \infty$  has been already proved in [5, 6] and [7], respectively; here we propose a different approach based on the Borel-Cantelli's Lemma and the properties of multiple Wiener integrals recalled in the preliminaries.

**Theorem 1** Suppose that  $\alpha > 0$ ,  $\theta_0 > 0$ ,  $\gamma \geqslant 0$ ,  $d \geqslant 1$  in equation (1). Let  $\hat{\theta}_{T,N}$  be the MLE given in (4) and  $\psi_{T,N}^{\theta}$  given in (13). Then, for every T > 0 fixed (resp.  $N \geqslant 1$  fixed), we have

$$\hat{\theta}_{T,N} \longrightarrow \theta_0$$
, almost surely as  $N \to +\infty$  (resp.  $T \to +\infty$ ).

**Proof.** Assume that T > 0 is fixed, we can write from (17),

$$\theta_0 - \hat{\theta}_{T,N} := \frac{\frac{1}{2} I_2(g_{T,N}) / \psi_{T,N}^{\theta_0}}{\left(\frac{1}{2\theta_0} I_2(g_{T,N}) - I_2(l_{T,N}) + \lambda_{T,N}\right) / \psi_{T,N}^{\theta_0}}.$$
(18)

We have

$$\mathbf{E}\left[\left(\frac{1}{2}I_{2}(g_{T,N})\right)^{2}\right] = \frac{1}{2}\|g_{T,N}\|^{2} = \frac{1}{2}\sum_{n=1}^{N}\delta_{n}^{4\alpha}\int_{0}^{T}\int_{0}^{T}e^{-2\theta_{0}\delta_{n}^{2\alpha}|u-v|}dudv = \sum_{n=1}^{N}\delta_{n}^{4\alpha}\int_{0}^{T}\int_{0}^{u}e^{-2\theta_{0}\delta_{n}^{2\alpha}(u-v)}dudv$$

$$= \sum_{n=1}^{N}\delta_{n}^{4\alpha}\int_{0}^{T}\int_{0}^{u}e^{-2\theta_{0}\delta_{n}^{2\alpha}w}dwdu = \sum_{n=1}^{N}\delta_{n}^{4\alpha}\int_{0}^{T}\left(\frac{1-e^{-2\theta_{0}\delta_{n}^{2\alpha}u}}{2\theta_{0}\delta_{n}^{2\alpha}}du\right)$$

$$= \psi_{T,N}^{\theta_{0}} + \frac{1}{2\theta_{0}}\sum_{n=1}^{N}(e^{-2\theta_{0}\delta_{n}^{2\alpha}T} - 1) = \lambda_{T,N}. \tag{19}$$

Thus, there exists a constant  $C(\alpha, d, \theta_0, T)$  such that

$$\mathbf{E}\left[\left(\frac{\frac{1}{2}I_{2}(g_{T,N})}{\psi_{T,N}^{\theta_{0}}}\right)^{2}\right] = \frac{1}{\psi_{T,N}^{\theta_{0}}} + \frac{1}{(\psi_{T,N}^{\theta_{0}})^{2}} \frac{1}{2\theta_{0}} \sum_{n=1}^{N} (e^{-2\theta_{0}\delta_{n}^{2\alpha}T} - 1) \leqslant \frac{1}{\psi_{T,N}^{\theta_{0}}} + \frac{N}{2\theta_{0}(\psi_{T,N}^{\theta_{0}})^{2}} \leqslant \frac{C(\alpha,\theta_{0},d,T)}{N^{\frac{2\alpha}{d}+1}}.$$

It follows that for every  $\varepsilon > 0$ , we have by Markov's inequality

$$\sum_{N=1}^{+\infty} \mathbf{P}\left(\left|\frac{\frac{1}{2}I_2(g_{T,N})}{\psi_{T,N}^{\theta_0}}\right| > \varepsilon\right) \leqslant \frac{1}{\varepsilon^2} \sum_{N=1}^{+\infty} \mathbf{E}\left[\left(\frac{\frac{1}{2}I_2(g_{T,N})}{\psi_{T,N}^{\theta_0}}\right)^2\right] \leqslant \frac{C(\alpha,\theta_0,d,T)}{\varepsilon^2} \sum_{N=1}^{+\infty} \frac{1}{N^{\frac{2\alpha}{d}+1}} < +\infty.$$

Hence, from Borel-Cantelli's Lemma, for every T > 0, we have  $\frac{\frac{1}{2}I_2(g_{T,N})}{\psi_{N,T}^{\theta_0}} \longrightarrow 0$  a.s as  $N \to +\infty$ . On the other hand,

$$||l_{T,N}||^2 = \frac{1}{4\theta_0^2} \sum_{n=1}^N \delta_n^{4\alpha} \int_0^T \int_0^T e^{2\theta_0 \delta_n^{2\alpha} (u+v)} du dv = \frac{1}{16\theta_0^4} \sum_{n=1}^N (1 - e^{-2\theta_0 \delta_n^{2\alpha} T})^2 \leqslant \frac{N}{16\theta_0^4}.$$

We obtain therefore

$$\mathbf{E}\left[\left(\frac{I_{2}(l_{T,N})}{\psi_{T,N}^{\theta_{0}}}\right)^{2}\right] = \frac{2\|l_{T,N}\|^{2}}{(\psi_{T,N}^{\theta_{0}})^{2}} \leqslant C(\alpha, d, \theta_{0}, T) \frac{1}{N^{\frac{4\alpha}{d} + 1}}$$

Hence, for every  $\varepsilon > 0$ , we have

$$\sum_{N=1}^{+\infty} \mathbf{P} \left( \left| \frac{I_2(l_{T,N})}{\psi_{T,N}^{\theta_0}} \right| > \varepsilon \right) \leqslant \frac{1}{\varepsilon^2} \sum_{N=1}^{+\infty} \mathbf{E} \left[ \left( \frac{I_2(l_{T,N})}{\psi_{T,N}^{\theta_0}} \right)^2 \right] \leqslant \frac{C(\alpha,\theta_0,d,T)}{\varepsilon^2} \sum_{N=1}^{+\infty} \frac{1}{N^{\frac{4\alpha}{d}+1}} < +\infty.$$

Hence, from Borel-Cantelli's Lemma, for every T>0, we have  $\frac{I_2(l_{T,N})}{\psi_{T,N}^{\theta_0}} \longrightarrow 0$  a.s as  $N\to +\infty$ . For the last deterministic term of the denominator, we have

$$\frac{\lambda_{T,N}}{\psi_{T,N}^{\theta_0}} = 1 + \frac{1}{4\theta_0^2 \psi_{T,N}^{\theta_0}} \sum_{n=1}^N \left( e^{-2\theta_0 \delta_n^{2\alpha} T} - 1 \right) \longrightarrow 1 \text{ as } N \to +\infty.$$

Therefore the desired result follows.

Assume now that  $N \ge 1$  is fixed, we can write

$$\theta_0 - \hat{\theta}_{T,N} = \frac{\sum_{n=1}^{N} \delta_n^{2\alpha + \gamma} \int_0^T y_n(t) dw_n(t)}{T} \times \frac{T}{\sum_{n=1}^{N} \delta_n^{4\alpha + 2\gamma} \int_0^T y_n^2(t) dt}$$

We have for any fixed  $N \ge 1$ ,

• 
$$\frac{1}{T} \sum_{n=1}^{N} \delta_n^{2\alpha+\gamma} \int_0^T y_n(t) dw_n(t) \to 0$$
 a.s. as  $T \to +\infty$ .

• 
$$\frac{1}{T} \sum_{n=1}^{N} \delta_n^{4\alpha+2\gamma} \int_0^T y_n^2(t) dt \to \frac{1}{2\theta_0} \sum_{n=1}^{N} \delta_n^{2(\alpha+\gamma)}$$
 a.s. as  $T \to +\infty$ .

In fact, using Ito's formula gives for all  $n \ge 1$ ,

$$y_n^2(T) = -2\theta_0 \delta_n^{2\alpha} \int_0^T y_n^2(t)dt + 2\delta_n^{-\gamma} \int_0^T y_n(t)dw_n(t) + T$$

Thus,

$$\frac{\delta_n^{-\gamma}}{T} \int_0^T y_n(t) dw_n(t) = \frac{y_n^2(T)}{2T} + \frac{\theta_0 \delta_n^{2\alpha}}{T} \int_0^T y_n^2(t) dt - \frac{1}{2}.$$

We set  $z_n(t) := \int_{-\infty}^t e^{-\theta_0 \delta_n^{2\alpha}(t-u)} dw_n(t)$  which is Gaussian, stationary and ergodic, thus by the ergodic theorem

$$\frac{1}{T} \int_0^T z_n^2(t) dt \to \mathbf{E}[z_n^2(0)] = \frac{1}{2\theta_0 \delta_n^{2\alpha}}$$

a.s. as  $T \to +\infty$ . Since  $z_n(t) = y_n(t) + e^{-\theta_0 \delta_n^{2\alpha} t} \varepsilon_n(0)$  where  $\varepsilon_n(0) := \int_{-\infty}^0 e^{\theta_0 \delta_n^{2\alpha} u} dw_n(u)$ . It follows that for all  $n \ge 1$ ,

$$\frac{1}{T} \int_0^T y_n^2(t) dt \to \frac{1}{2\theta_0 \delta_n^{2\alpha}}$$

a.s. as  $T \to +\infty$ . Moreover, it was proved in [12] in Lemma 6.7 that for any  $n \geqslant 1$ , we have for any  $\alpha > 0$ ,  $\frac{z_n(T)}{T^\alpha} \to 0$  a.s. as  $T \to +\infty$ . It follows that  $\frac{y_n^2(T)}{T} \to 0$  a.s. as  $T \to +\infty$  for any  $n \geqslant 1$ . Therefore, we get for any  $n \geqslant 1$ 

- $\frac{1}{T} \int_0^T y_n(t) dw_n(t) \to 0$  a.s. as  $T \to +\infty$ .
- $\frac{1}{T} \int_0^T y_n^2(t) dt \to \frac{1}{2\theta_0 \delta_2^{2\alpha}}$  a.s. as  $T \to +\infty$ .

Then the desired result follows.

**Theorem 2** Suppose that  $\alpha > 0$ ,  $\theta_0 > 0$ ,  $\gamma \ge 0$ ,  $d \ge 1$  in equation (1). Let  $\hat{\theta}_{T,N}$  be the MLE given in (4) and  $\psi_{T,N}^{\theta_0}$  given in (13). Then there exists a positive constant  $C(\alpha,d,\theta_0)$  depending on  $\alpha$  the power of the Laplacian, d is the space dimension and  $\theta_0$  such that for any  $N \ge 1$  and any T > 0,

$$d_{W}\left(\sqrt{\psi_{T,N}^{\theta_{0}}}(\theta_{0}-\hat{\theta}_{T,N}),Z\right) \leqslant C(\alpha,d,\theta_{0}) \times \begin{cases} \frac{1}{\sqrt{T}} \frac{1}{N^{\frac{2\alpha}{d}}} & \text{if } 2\alpha < d\\ \frac{1}{\sqrt{T}} \frac{1}{N} & \text{if } 2\alpha = d\\ \frac{1}{\sqrt{T}} \frac{1}{N^{\frac{\alpha}{d}+\frac{1}{2}}} & \text{if } 2\alpha > d, \end{cases}$$

$$(20)$$

where  $Z \sim \mathcal{N}(0,1)$ . Consequently as  $T \to +\infty$ ,  $N \to +\infty$ 

$$\sqrt{T}N^{\frac{\alpha}{d}+\frac{1}{2}}\left(\theta_0-\hat{\theta}_{T,N}\right) \xrightarrow{law} \mathcal{N}\left(0,\frac{(4\alpha/d+2)\theta_0}{\bar{\sigma}^{\alpha}}\right).$$

**Proof.** We recall that from (14) and (17), we have  $\theta_0 - \hat{\theta}_{T,N} = \frac{E_{T,N}}{\langle E_{T,N} \rangle} = \frac{\frac{1}{2}I_2(g_{T,N})}{\frac{1}{2\theta_0}I_2(g_{T,N}) - I_2(l_{T,N}) + \lambda_{T,N}}$ . On the other hand using the calculus (19), the following estimate holds

$$\mathbf{E}\left[\left(\frac{E_{T,N}}{\sqrt{\psi_{T,N}^{\theta_0}}}\right)^2\right] = \frac{\lambda_{N,T}}{\psi_{T,N}^{\theta_0}} = 1 + \frac{1}{4\theta_0^2 \psi_{T,N}^{\theta_0}} \sum_{n=1}^N (e^{-2\theta_0 \delta_n^{2\alpha} T} - 1).$$

Therefore,

$$\left| \mathbf{E} \left[ \left( \frac{E_{T,N}}{\sqrt{\psi_{T,N}^{\theta_0}}} \right)^2 \right] - 1 \right| = \left| \frac{\lambda_{N,T}}{\psi_{T,N}^{\theta_0}} - 1 \right| \leqslant \frac{C(\alpha, d, \theta_0)}{TN^{\frac{2\alpha}{d}}}, \tag{21}$$

where  $C(\alpha, d, \theta)$  is a constant depending on  $\alpha$ , d and  $\theta$ . Moreover, since  $\frac{E_{T,N}}{\sqrt{\psi_{T,N}^{\theta_0}}} \in \mathcal{H}_2$  and  $\mathbf{E}\left[\left(\frac{E_{T,N}}{\sqrt{\psi_{T,N}^{\theta_0}}}\right)^2\right] \leqslant 1$ . The above the thin have the depending on  $\alpha$ , d and d. Moreover, since  $\frac{E_{T,N}}{\sqrt{\psi_{T,N}^{\theta_0}}} \in \mathcal{H}_2$  and  $\mathbf{E}\left[\left(\frac{E_{T,N}}{\sqrt{\psi_{T,N}^{\theta_0}}}\right)^2\right] \leqslant 1$ .

1, then by the hypercontractivity property, we have  $\|\frac{E_{T,N}}{\sqrt{\psi_{T,N}^{\theta_0}}}\|_{L^4} \leqslant 3\|\frac{E_{T,N}}{\sqrt{\psi_{T,N}^{\theta_0}}}\|_{L^2} \leqslant 3.$ 

**Lemma 3** Consider  $\langle E_{T,N} \rangle$  defined in (16) and  $\psi_{T,N}^{\theta_0}$  defined in (13). Then, there exists a constant depending on  $\alpha$ , d and  $\theta_0$ ,  $C(\alpha, d, \theta_0)$  such that for any  $N \geqslant 1$  and T > 0, we have

$$\|\frac{\langle E_{T,N} \rangle}{\psi_{T,N}^{\theta_0}} - 1\|_{L^2} \leqslant C(\alpha, d, \theta_0) \times b_{N,T}, \text{ where } b_{N,T} := \begin{cases} \frac{1}{\sqrt{T}} \frac{1}{N^{\frac{2\alpha}{d}}} & \text{if } 2\alpha < d \\ \frac{1}{\sqrt{T}} \frac{1}{N} & \text{if } 2\alpha = d \\ \frac{1}{\sqrt{T}} \frac{1}{N^{\frac{\alpha}{d} + \frac{1}{2}}} & \text{if } 2\alpha > d. \end{cases}$$
(22)

**Proof.** Indeed, using (16), (21) and Minkowski's inequality, we can write

$$\|\frac{< E_{T,N}>}{\psi_{T,N}^{\theta_0}} - 1\|_{L^2} \leqslant \|\frac{E_{T,N}}{\theta_0\psi_{T,N}^{\theta_0}}\|_{L^2} + \|\frac{I_2(h_{N,T})}{\psi_{T,N}^{\theta_0}}\|_{L^2} + |\frac{\lambda_{N,T}}{\psi_{T,N}^{\theta_0}} - 1| \leqslant \frac{1}{\theta_0} \frac{1}{\sqrt{\psi_{T,N}^{\theta_0}}} + \frac{C(\alpha,d,\theta_0)}{TN^{\frac{2\alpha}{d}+\frac{1}{2}}} + \frac{C(\alpha,d,\theta_0)}{TN^{\frac{2\alpha}{d}}}.$$

where we used for the estimate of  $\|\frac{I_2(l_{T,N})}{\psi_{T,N}^{\theta_0}}\|_{L^2}$ , the fact that

$$||l_{T,N}||^2 = \frac{1}{4\theta_0^2} \sum_{n=1}^N \delta_n^{4\alpha} \int_0^T \int_0^T e^{2\theta_0 \delta_n^{2\alpha} (u+v)} du dv = \frac{1}{16\theta_0^4} \sum_{n=1}^N (1 - e^{-2\theta_0 \delta_n^{2\alpha} T})^2 \leqslant \frac{N}{16\theta_0^4}.$$

We obtain therefore

$$\mathbf{E}\left[\left(\frac{I_{2}(l_{T,N})}{\psi_{T,N}^{\theta_{0}}}\right)^{2}\right] = \frac{2\|l_{T,N}\|^{2}}{(\psi_{T,N}^{\theta_{0}})^{2}} \leqslant C(\alpha,d,\theta_{0}) \frac{1}{T^{2}} \frac{1}{N^{\frac{4\alpha}{d}+1}}.$$

**Lemma 4** Consider  $E_{T,N}$ ,  $\psi_{T,N}^{\theta_0}$  defined respectively in (13), then the following estimates hold

$$\kappa_3\left(\frac{E_{T,N}}{\sqrt{\psi_{T,N}^{\theta_0}}}\right) \leqslant \frac{3}{\theta_0\sqrt{\psi_{T,N}^{\theta_0}}}, \quad and \quad \left|\kappa_4\left(\frac{E_{T,N}}{\sqrt{\psi_{T,N}^{\theta_0}}}\right)\right| \leqslant \frac{18}{\theta_0^2\psi_{T,N}^{\theta_0}}$$

**Proof.** we have by (11)

$$\kappa_3 \left( \frac{E_{T,N}}{\sqrt{\psi_{T,N}^{\theta_0}}} \right) = \kappa_3 \left( \frac{I_2(g_{T,N})}{2\sqrt{\psi_{T,N}^{\theta_0}}} \right) = \mathbf{E} \left[ \left( \frac{I_2(g_{T,N})}{2\sqrt{\psi_{T,N}^{\theta_0}}} \right)^3 \right] = \sum_{n=1}^N \mathbf{E} \left[ \left( \frac{I_2(g^n)}{2\sqrt{\psi_{T,N}^{\theta_0}}} \right)^3 \right] \\
= \frac{1}{(\psi_{T,N}^{\theta_0})^{3/2}} \sum_{n=1}^N \langle g^n, g^n \otimes_1 g^n \rangle_{\mathcal{H}^{\otimes 2}}.$$

Moreover, we have for every n = 1, ..., N

$$\langle g^{n}, g^{n} \otimes_{1} g^{n} \rangle_{\mathcal{H}^{\otimes 2}} = \int_{0}^{T} \int_{0}^{T} \int_{0}^{T} g^{n}(x, y) g^{n}(x, t) g^{n}(y, t) dt dx dy$$

$$= \delta_{n}^{6\alpha} \int_{[0, T]^{3}} e^{-\theta_{0} \delta_{n}^{2\alpha} |x - y|} e^{-\theta_{0} \delta_{n}^{2\alpha} |x - t|} e^{-\theta_{0} \delta_{n}^{2\alpha} |y - t|} dt dx dy$$

$$= 3! \delta_{n}^{6\alpha} \int_{0}^{T} \int_{0}^{y} \int_{0}^{t} e^{-\theta_{0} \delta_{n}^{2\alpha} (y - x)} e^{-\theta_{0} \delta_{n}^{2\alpha} (t - x)} e^{-\theta_{0} \delta_{n}^{2\alpha} (t - y)} dt dx dy$$

$$= 3! \delta_{n}^{6\alpha} \int_{0}^{T} \int_{0}^{y} \int_{0}^{t} e^{-\theta_{0} \delta_{n}^{2\alpha} (2t - 2x)} dt dx dy$$

$$= \frac{3 \delta_{n}^{4\alpha}}{\theta_{0}} \int_{0}^{T} \left( \frac{1 - e^{-2\theta_{0} \delta_{n}^{2\alpha} t}}{2\theta_{0} \delta_{n}^{2\alpha}} - t e^{-2\theta_{0} \delta_{n}^{2\alpha} t} \right) dt.$$

Thus,

$$\kappa_3 \left( \frac{E_{T,N}}{\sqrt{\psi_{T,N}^{\theta_0}}} \right) \leqslant \frac{3T}{2\theta_0 (\psi_{T,N}^{\theta_0})^{3/2}} \sum_{n=1}^N \delta_n^{2\alpha} = \frac{3}{\theta_0 \sqrt{\psi_{T,N}^{\theta_0}}}.$$

We also have by (12)

$$\begin{split} |\kappa_4\left(\frac{E_{T,N}}{\sqrt{\psi_{T,N}^{\theta_0}}}\right)| &= |\kappa_4\left(\frac{I_2(g_{T,N})}{2\sqrt{\psi_{T,N}^{\theta_0}}}\right)| \\ &\leqslant \frac{3}{(\psi_{T,N}^{\theta_0})^2} \sum_{n=1}^N \|g^n \otimes_1 g^n\|_{\mathcal{H}^{\otimes 2}}^2 \\ &= \frac{3}{(\psi_{T,N}^{\theta_0})^2} \sum_{n=1}^N \int_{[0,T]^4} g^n(x_1,x_2) g^n(x_2,x_3) g^n(x_3,x_4) g^n(x_4,x_1) dx_1 dx_2 dx_3 dx_4 \\ &= \frac{3}{(\psi_{T,N}^{\theta_0})^2} \sum_{n=1}^N 4! \delta_n^{8\alpha} \int_0^T \int_0^{x_4} \int_0^{x_3} \int_0^{x_2} e^{-\theta_0 \delta_n^{2\alpha} (2x_4 - 2x_1)} dx_1 dx_2 dx_3 dx_4 \\ &\leqslant \frac{18}{\theta_0^2 \psi_{T,N}^{\theta_0}}. \end{split}$$

■ Consequently, there exists a positive constant  $C(\theta_0)$  depending only on  $\theta_0$  such that for every  $N \ge 1$ , T > 0, we have

$$d_W\left(\frac{E_{T,N}}{\sqrt{\psi_{N,T}^{\theta_0}}}, Z\right) \leqslant \frac{C(\theta_0)}{\sqrt{\psi_{N,T}^{\theta_0}}},$$

where  $Z \sim \mathcal{N}(0,1)$ . On the other hand,

$$\sqrt{\psi_{T,N}^{\theta_0}}(\theta_0 - \hat{\theta}_{T,N}) = \frac{E_{T,N}/\sqrt{\psi_{T,N}^{\theta_0}}}{\langle E_{T,N} \rangle / \psi_{T,N}^{\theta_0}}.$$

Thus,

$$\begin{split} d_{W}\left(\sqrt{\psi_{T,N}^{\theta_{0}}}(\theta_{0}-\hat{\theta}_{T,N}),Z\right) &= d_{W}\left(\frac{E_{T,N}/\sqrt{\psi_{T,N}^{\theta_{0}}}}{< E_{T,N} > /\psi_{T,N}^{\theta_{0}}},Z\right) \\ &\leqslant d_{W}\left(\frac{E_{T,N}}{\sqrt{\psi_{T,N}^{\theta_{0}}}},Z\right) + d_{W}\left(\frac{E_{T,N}/\sqrt{\psi_{T,N}^{\theta_{0}}}}{< E_{T,N} > /\psi_{T,N}^{\theta_{0}}},\frac{E_{T,N}}{\sqrt{\psi_{T,N}^{\theta_{0}}}}\right) \\ &\leqslant d_{W}\left(\frac{E_{T,N}}{\sqrt{\psi_{T,N}^{\theta_{0}}}},Z\right) + \mathbf{E}\left[\left|(\frac{E_{T,N}/\sqrt{\psi_{T,N}^{\theta_{0}}}}{< E_{T,N} > /\psi_{T,N}^{\theta_{0}}} - \frac{E_{T,N}}{\sqrt{\psi_{T,N}^{\theta_{0}}}}\right|\right] \\ &\leqslant d_{W}\left(\frac{E_{T,N}}{\sqrt{\psi_{T,N}^{\theta_{0}}}},Z\right) + \left\|\frac{E_{T,N}/\sqrt{\psi_{T,N}^{\theta_{0}}}}{< E_{T,N} > /\psi_{T,N}^{\theta_{0}}}\right\|_{L^{2}} \times \left\|1 - \frac{< E_{T,N} >}{\psi_{T,N}^{\theta_{0}}}\right\|_{L^{2}} \\ &\leqslant d_{W}\left(\frac{E_{T,N}}{\sqrt{\psi_{T,N}^{\theta_{0}}}},Z\right) + \left\|\frac{E_{T,N}}{\sqrt{\psi_{T,N}^{\theta_{0}}}}\right\|_{L^{4}} \times \left\|\frac{\psi_{T,N}^{\theta_{0}}}{< E_{T,N} >} \right\|_{L^{4}} \times \left\|1 - \frac{< E_{T,N} >}{\psi_{T,N}^{\theta_{0}}}\right\|_{L^{2}}. \end{split}$$

The desired result follows from the previous calculus along with Lemma 14 which implies the existence of a constant  $C(\theta_0)$  such that  $\|\frac{\psi_{T,N}^{\theta_0}}{\langle E_{T,N} \rangle}\|_{L^4} < C(\theta_0)$ . The following corollary is a direct consequence of Theorem 2.

Corollary 5 Suppose that  $\alpha > 0$ ,  $\theta_0 > 0$ ,  $\gamma \geqslant 0$ ,  $d \geqslant 1$  in equation (1). Let  $\hat{\theta}_{T,N}$  be the MLE given in (4). Then, we have

• If  $N \to +\infty$  and T is fixed, then there exists a constant  $C(\alpha, d, \theta_0, T)$  such that

$$d_{W}\left(N^{\frac{\alpha}{d}+\frac{1}{2}}(\theta_{0}-\hat{\theta}_{T,N}), \mathcal{N}\left(\frac{(4\alpha/d+2)\theta_{0}}{\bar{\sigma}^{\alpha}T}\right)\right) \leqslant C(\alpha,d,\theta_{0},T) \times \begin{cases} \frac{1}{N^{\frac{2\alpha}{d}}} & \text{if } 2\alpha < d\\ \frac{1}{N} & \text{if } 2\alpha = d\\ \frac{1}{N^{\frac{\alpha}{d}+\frac{1}{2}}} & \text{if } 2\alpha > d. \end{cases}$$

$$(23)$$

• If  $T \to +\infty$  and N is fixed, then there exists a constant  $C(\alpha, d, \theta_0, N)$  such that

$$d_{W}\left(\sqrt{T}(\theta_{0} - \hat{\theta}_{T,N}), \mathcal{N}\left(0, \frac{2\theta_{0}}{\sum\limits_{n=1}^{N} \delta_{n}^{2\alpha}}\right)\right) \leqslant \frac{C(\alpha, d, \theta_{0}, N)}{\sqrt{T}}.$$
(24)

• If  $T \to +\infty$  and  $N \to +\infty$ , then there exists a constant  $C(\alpha, d, \theta_0)$  such that

$$d_{W}\left(\sqrt{T}N^{\frac{\alpha}{d}+\frac{1}{2}}(\theta_{0}-\hat{\theta}_{T,N}), \mathcal{N}\left(\frac{(4\alpha/d+2)\theta_{0}}{\bar{\sigma}^{\alpha}}\right)\right) \leqslant C(\alpha,d,\theta_{0}) \times \begin{cases} \frac{1}{\sqrt{T}}\frac{1}{N^{\frac{2\alpha}{d}}} & \text{if } 2\alpha < d\\ \frac{1}{\sqrt{T}}\frac{1}{N} & \text{if } 2\alpha = d\\ \frac{1}{\sqrt{T}}\frac{1}{N^{\frac{\alpha}{d}+\frac{1}{2}}} & \text{if } 2\alpha > d. \end{cases}$$

$$(25)$$

**Example 6** If the power of the Laplacian in equation (1) equals 1:  $\alpha = 1$ , then equation (1) becomes the classical stochastic heat equation with an additive space-time white noise as follows:

$$\begin{cases} dY_{\theta}(t,x) - \theta \Delta Y_{\theta}(t,x) dt = \sum_{n \in \mathbb{N}} \delta_n^{-\gamma} e_n(x) dw_n(t), & x \in G \\ Y_{\theta}(0,x) = 0, & x \in G. \end{cases}$$

Then, in this case, if  $G \subset \mathbb{R}$ , namely d=1, there exists a constant  $C(\theta_0)$  depending only on  $\theta_0$  such that the following bound of convergence in  $d_W$  holds for the MLE  $\hat{\theta}_{T,N}$ :

$$d_W\left(\sqrt{T}N^{3/2}(\theta_0 - \hat{\theta}_{T,N}), \mathcal{N}\left(0, \frac{6\theta_0}{\bar{\sigma}}\right)\right) \leqslant \frac{C(\theta_0)}{\sqrt{T}N^{3/2}}.$$

#### 4 Approximate maximum likelihood estimator

In this section, we study the approximative MLE of the parameter  $\theta$ , defined by :

$$\widetilde{\theta}_{T,N,M} := -\frac{\sum_{n=1}^{N} \delta_n^{2\alpha+2\gamma} \sum_{i=1}^{M} y_n(t_{i-1}) \left[ y_n(t_i) - y_n(t_{i-1}) \right]}{\Delta_M \sum_{n=1}^{N} \delta_n^{4\alpha+2\gamma} \sum_{i=1}^{M} y_n^2(t_{i-1})}$$
(26)

$$= -\frac{\sum_{n=1}^{N} \delta_n^{2\alpha} \sum_{i=1}^{M} v_n(t_{i-1}) \left[ v_n(t_i) - v_n(t_{i-1}) \right]}{\Delta_M \sum_{n=1}^{N} \delta_n^{4\alpha} \sum_{i=1}^{M} v_n^2(t_{i-1})},$$
(27)

which is a discretized version of the MLE given by (4), where

$$y_n(t) = \delta_n^{-\gamma} \int_0^t e^{-\theta_0 \delta_n^{2\alpha}(t-s)} dw_n(s), \qquad v_n(t) = \int_0^t e^{-\theta_0 \delta_n^{2\alpha}(t-s)} dw_n(s), \quad n = 1, ..., N.$$

In what follows, we assume that the Fourier modes  $y_n(t), n \ge 1$ , are observed at a uniform time grid:

$$0 = t_0 < t_1 < \dots < t_M = T$$
, with  $\Delta_M := t_i - t_{i-1} = \frac{T}{M}, i = 1, \dots, M$ .

We are interested in studying the consistency and the rate of convergence of the distribution of  $\widetilde{\theta}_{T,N,M}$  as  $N, M, T \to \infty$ . For this aim, let us introduce the following sequences

$$S_{N,M} := \Delta_{M} \sum_{n=1}^{N} \delta_{n}^{4\alpha} \sum_{i=1}^{M} v_{n}^{2}(t_{i-1}), \text{ and } \Lambda_{N,M} := \sum_{n=1}^{N} \delta_{n}^{2\alpha} \sum_{i=1}^{M} e^{-\theta_{0} \delta_{n}^{2\alpha} t_{i}} v_{n}(t_{i-1}) \left[ \eta_{n}(t_{i}) - \eta_{n}(t_{i-1}) \right]$$

$$= \sum_{n=1}^{N} \delta_{n}^{2\alpha} \sum_{i=1}^{M} e^{-\theta_{0} \delta_{n}^{2\alpha} (t_{i} + t_{i-1})} \eta_{n}(t_{i-1}) \left[ \eta_{n}(t_{i}) - \eta_{n}(t_{i-1}) \right]$$

$$=: \sum_{n=1}^{N} B_{n,M}, \tag{28}$$

where  $\eta_n(t) = \int_0^t e^{\theta_0 \delta_n^{2\alpha} s} dw_n(s)$ . Thus,

$$-\widetilde{\theta}_{T,N,M} = \frac{\sum_{n=1}^{N} \delta_n^{2\alpha} \left( e^{-\delta_n^{2\alpha} \theta_0 \Delta_M} - 1 \right) \sum_{i=1}^{M} v_n^2 \left( t_{i-1} \right)}{\Delta_M \sum_{n=1}^{N} \delta_n^{4\alpha} \sum_{i=1}^{M} v_n^2 \left( t_{i-1} \right)} + \frac{\Lambda_{N,M}}{S_{N,M}}.$$
 (29)

Hence

$$\theta_{0} - \widetilde{\theta}_{T,N,M} = \frac{\sum_{n=1}^{N} \delta_{n}^{2\alpha} \left( e^{-\delta_{n}^{2\alpha} \theta_{0} \Delta_{M}} - 1 + \delta_{n}^{2\alpha} \theta_{0} \Delta_{M} \right) \sum_{i=1}^{M} v_{n}^{2} \left( t_{i-1} \right)}{\Delta_{M} \sum_{n=1}^{N} \delta_{n}^{4\alpha} \sum_{i=1}^{M} v_{n}^{2} \left( t_{i-1} \right)} + \frac{\Lambda_{N,M}}{S_{N,M}}$$
(30)

$$=: \frac{R_{N,M}}{S_{N,M}} + \frac{\Lambda_{N,M}}{S_{N,M}}.$$
 (31)

## 4.1 Consistency of the estimator $\widetilde{\theta}_{T,N,M}$

Proposition 7 Assume that

$$\Delta_M N^{\frac{2\alpha}{d}} \to 0$$
, as  $N, M, T \to +\infty$ . (32)

Then the estimator  $\widetilde{\theta}_{T,N,M}$  is weakly consistent, namely the following convergence holds in probability

$$\widetilde{\theta}_{T,N,M} \to \theta_0$$
, as  $N, M, T \to +\infty$ .

**Proof.** Following the decomposition (31), we have for any  $\varepsilon > 0$ 

$$\mathbf{P}\left(|\widetilde{\theta}_{T,N,M} - \theta_0| > \varepsilon\right) \leqslant \mathbf{P}\left(\left|\frac{R_{N,M}}{S_{N,M}}\right| > \varepsilon/2\right) + \mathbf{P}\left(\left|\frac{\Lambda_{N,M}}{S_{N,M}}\right| > \varepsilon/2\right)$$
(33)

Moreover, for any arbitrary fixed  $\delta \in (0, \varepsilon/2)$ , we have

$$\mathbf{P}\left(|\widetilde{\theta}_{T,N,M} - \theta_0| > \varepsilon\right) \leqslant \mathbf{P}\left(\left|\frac{R_{N,M}}{\psi_{T,N}^{\theta_0}}\right| > \delta\right) + \mathbf{P}\left(\left|\frac{\Lambda_{N,M}}{\psi_{T,N}^{\theta_0}}\right| > \delta\right) + 2\mathbf{P}\left(\left|\frac{S_{N,M}}{\psi_{T,N}^{\theta_0}} - 1\right| > \frac{\varepsilon - 2\delta}{\varepsilon}\right)$$
(34)

We recall that

$$\psi_{T,N}^{\theta_0} = \frac{T}{2\theta_0} \sum_{n=1}^{N} \delta_n^{2\alpha} \sim \frac{\bar{\sigma} dT N^{\frac{2\alpha}{d}+1}}{(4\alpha + 2d)\theta_0} \text{ as } N, T \to +\infty.$$

For each term included in the inequality (34), the second moment need to be estimated by Markov's inequality.

**Lemma 8** Consider the sequence  $R_{N,M}$  defined in (31) and assume that the assumption (32) holds, then there exists a constant  $C(\theta_0, \alpha, d, \bar{\sigma})$  such that for any N, T, M large, we have

$$\mathbf{E}\left[\left(\frac{R_{N,M}}{\psi_{T,N}^{\theta_0}}\right)^2\right] \leqslant C(\theta_0, \alpha, d, \bar{\sigma}) \Delta_M^2 N^{4\alpha/d}.$$

**Proof.** Recall that the sequence  $R_{N,M}$  defined in (31) can be written as follows

$$R_{N,M} := \sum_{n=1}^{N} s_n \sum_{i=1}^{M} v_n^2(t_{i-1}),$$

where

$$s_n := \delta_n^{2\alpha} \left( e^{-\delta_n^{2\alpha} \theta_0 \Delta_M} - 1 + \delta_n^{2\alpha} \theta_0 \Delta_M \right), \quad n = 1, ..., N.$$

Therefore,

$$\begin{split} \mathbf{E}\left[R_{N,M}^{2}\right] &= \sum_{n,l=1}^{N} s_{n} s_{l} \sum_{i,j=1}^{M} \mathbf{E}\left[v_{n}^{2}(t_{i-1})v_{l}^{2}(t_{j-1})\right] \\ &= \sum_{\substack{n,l=1\\n \neq l}}^{N} s_{n} s_{l} \sum_{i,j=1}^{M} \mathbf{E}\left[v_{n}^{2}(t_{i-1})\right] \mathbf{E}\left[v_{l}^{2}(t_{j-1})\right] + \sum_{n=1}^{N} s_{n}^{2} \mathbf{E}\left[\left(\sum_{i=1}^{M} v_{n}^{2}(t_{i-1})\right)^{2}\right]. \end{split}$$

Besides, using Isserlis' theorem, we obtain for any n = 1, ..., N,

$$\mathbf{E}\left[\left(\sum_{i=1}^{M} v_n^2(t_{i-1})\right)^2\right] = \left(\sum_{i=1}^{M} \mathbf{E}[v_n^2(t_{i-1})]\right)^2 + 2\sum_{i=1}^{M} \left(\mathbf{E}[v_n^2(t_{i-1})]\right)^2 + 2\sum_{\substack{i,j=1\\i\neq j}}^{M} \left(\mathbf{E}[v_n(t_{i-1})v_n(t_{j-1})]\right)^2.$$

Therefore,

$$\mathbf{E}\left[R_{N,M}^{2}\right] = \left[\left(\sum_{n=1}^{N} s_{n} \sum_{i=1}^{M} \mathbf{E}[v_{n}^{2}(t_{i-1})]\right)^{2} + 2\sum_{n=1}^{N} s_{n}^{2} \sum_{i,j=1}^{M} \left(\mathbf{E}[v_{n}(t_{i-1})v_{n}(t_{j-1})]\right)^{2}\right]$$

$$:= A_{N,M,T} + B_{N,M,T}.$$
(35)

Since  $|1 - e^{-x} - x|/x^2 \le 1$  for all x > 0, by applying Mean Value Theorem twice, we have for any n = 1, ..., N

$$s_n := \delta_n^{2\alpha} \left[ e^{-\delta_n^{2\alpha} \theta_0 \Delta_M} - 1 + \delta_n^{2\alpha} \theta_0 \Delta_M \right] \leqslant \theta_0^2 \delta_n^{6\alpha} \Delta_M^2.$$

Besides, we also have the following estimate, for any n = 1, ..., N, we have

$$\mathbf{E}[v_n^2(t)] = \int_0^t e^{-2\theta_0 \delta_n^{2\alpha}(t-s)} ds = \frac{1}{2\theta_0 \delta_n^{2\alpha}} (1 - e^{-2\theta_0 \delta_n^{2\alpha}t}) \leqslant \frac{1}{2\theta_0 \delta_n^{2\alpha}}.$$

Therefore, we can write

$$\frac{4\theta_0^2 A_{N,M,T}}{T^2 \left(\sum\limits_{n=1}^N \delta_n^{2\alpha}\right)^2} \leqslant \frac{4\theta_0^2}{T^2 \left(\sum\limits_{n=1}^N \delta_n^{2\alpha}\right)^2} \left(\sum\limits_{n=1}^N \theta_0^2 \delta_n^{6\alpha} \Delta_M^2 \sum\limits_{i=1}^M \frac{1}{2\theta_0 \delta_n^{2\alpha}}\right)^2 \leqslant \frac{M^2 \Delta_M^4 \theta_0^4}{T^2 \left(\sum\limits_{n=1}^N \delta_n^{2\alpha}\right)^2} \left(\sum\limits_{n=1}^N \delta_n^{4\alpha}\right)^2.$$

Using the fact that  $\delta_n \sim \bar{w}^{1/2} n^{1/d}$  as  $n \to +\infty$ , the following estimate holds for  $A_{N,M,T}$  for N,M,T large

$$\frac{4\theta_0^2 A_{N,M,T}}{T^2 \left(\sum\limits_{n=1}^N \delta_n^{2\alpha}\right)^2} \leqslant \theta_0^4 \left(\frac{2\alpha+d}{\alpha+d}\right)^2 \bar{\sigma}^{2\alpha} \Delta_M^2 N^{4\alpha/d}.$$

For the sequence  $B_{N,M,T}$ , we have using Cauchy-Schwarz inequality and the estimates above,

$$\frac{4\theta_0^2 B_{N,M,T}}{T^2 \left(\sum_{n=1}^N \delta_n^{2\alpha}\right)^2} \leqslant \frac{2}{T^2 \left(\sum_{n=1}^N \delta_n^{2\alpha}\right)^2} \sum_{n=1}^N s_n^2 \sum_{i,j=1}^M \frac{1}{\delta_n^{4\alpha}} \leqslant \frac{2M^2 \Delta_M^4 \theta_0^4}{T^2 \left(\sum_{n=1}^N \delta_n^{2\alpha}\right)^2} \sum_{n=1}^N \delta_n^{8\alpha}$$

as previsouly, using the fact that  $\delta_n \sim \bar{\sigma}^{1/2} n^{1/d}$  as  $n \to +\infty$ , the following estimate holds for  $B_{N,M,T}$  for N, M, T large

$$\frac{4\theta_0^2 B_{N,M,T}}{T^2 \left(\sum_{n=1}^N \delta_n^{2\alpha}\right)^2} \leqslant \left(\frac{2\alpha + d}{8\alpha + d}\right)^2 \bar{\sigma}^{2\alpha} 2\theta_0^4 \Delta_M^2 N^{\frac{4\alpha}{d} - 1}.$$

The desired result follows.

**Lemma 9** Consider the sequence  $S_{N,M}$  defined in (31) and assume that assumption (32) holds true. Then, there exists a constant  $C(\theta_0, \alpha, \bar{\sigma}, d)$  such that for N, M, T large we have

$$\mathbf{E}\left[\left|\frac{S_{N,M}}{\psi_{T,N}^{\theta_0}} - 1\right|^2\right] \leqslant C(\theta_0, \alpha, \bar{\sigma}, d) \max\left(\frac{1}{TN^{\frac{2\alpha}{d}+1}}, \frac{1}{T^2N^{\frac{4\alpha}{d}}}\right).$$

**Proof.** Recall that  $S_{N,M} = \Delta_M \sum_{n=1}^N \delta_n^{4\alpha} \sum_{i=1}^M v_n^2(t_{i-1})$ . Then, by the product formula (6), we obtain

$$S_{N,M} = \Delta_M \left[ \sum_{n=1}^N \delta_n^{4\alpha} \sum_{i=1}^M I_2^{w_n} \left( e^{-2\theta_0 \delta_n^{2\alpha} t_{i-1}} e^{\theta_0 \delta_n^{2\alpha} (u+v)} \mathbf{1}(u,v)_{[0,t_{i-1}]^2} \right) + \sum_{n=1}^N \delta_n^{4\alpha} \sum_{i=1}^M \frac{1}{2\theta_0 \delta_n^{2\alpha}} (1 - e^{-2\theta_0 \delta_n^{2\alpha} t_{i-1}}) \right].$$

Therefore, we can write

$$\frac{S_{N,M}}{\psi_{T,N}^{\theta_0}} - 1 := C_{N,M,T} + (D_{N,M,T} - 1). \tag{36}$$

Thus, using the inequality  $(a+b)^2 \leq 2a^2 + 2b^2$ , we obtain

$$\mathbf{E}\left[\left|\frac{S_{N,M}}{\psi_{T,N}^{\theta_0}} - 1\right|^2\right] \leqslant 2\mathbf{E}[C_{N,M,T}^2] + 2|D_{N,M,T} - 1|^2.$$

For the seek of simplicity, let us denote  $\varphi_n^{\theta_0}(t) := e^{-2\theta_0 \delta_n^{2\alpha} t} e^{\theta_0 \delta_n^{2\alpha} (u+v)} \mathbf{1}(u,v)_{[0,t]^2}$ , we have

$$\begin{split} &\mathbf{E}\left[C_{N,M,T}^{2}\right] \\ &= \frac{4\theta_{0}^{2}\Delta_{M}^{2}}{T^{2}(\sum\limits_{n=1}^{N}\delta_{n}^{2\alpha})^{2}}\left[\sum\limits_{n=1}^{N}\delta_{n}^{8\alpha}\sum\limits_{i=1}^{M}\mathbf{E}\left[I_{2}^{w_{n}}(\varphi_{n}^{\theta_{0}}(t_{i-1}))^{2}\right] + \sum\limits_{n=1}^{N}\delta_{n}^{8\alpha}\sum\limits_{i,j=1}^{M}\mathbf{E}\left[I_{2}^{w_{n}}(\varphi_{n}^{\theta_{0}}(t_{i-1}))I_{2}^{w_{n}}(\varphi_{n}^{\theta_{0}}(t_{j-1}))\right]\right] \\ &:= C_{N,M,T,1} + C_{N,M,T,2}. \end{split}$$

For the sequence  $C_{N,M,T,1}$ , using the isometry property (5) the following estimate holds

$$C_{N,M,T,1} = \frac{2\Delta_M^2}{T^2(\sum_{n=1}^N \delta_n^{2\alpha})^2} \sum_{n=1}^N \delta_n^{8\alpha} \sum_{i=1}^M \frac{1}{\delta_n^{4\alpha}} (1 - e^{-2\theta_0 \delta_n^{2\alpha t_{i-1}}})^2 \leqslant \frac{2M\Delta_M^2}{T^2(\sum_{n=1}^N \delta_n^{2\alpha})^2} \sum_{n=1}^N \delta_n^{4\alpha} (1 - e^{-2\theta_0 \delta_n^{2\alpha t_{i-1}}})^2 \leqslant \frac{2M\Delta_M^2}{T^2(\sum_{n=1}^N \delta_n^{2\alpha})^2} \sum_{n=1}^N \delta_n^{4\alpha} (1 - e^{-2\theta_0 \delta_n^{2\alpha t_{i-1}}})^2 \leqslant \frac{2M\Delta_M^2}{T^2(\sum_{n=1}^N \delta_n^{2\alpha})^2} \sum_{n=1}^N \delta_n^{4\alpha} (1 - e^{-2\theta_0 \delta_n^{2\alpha t_{i-1}}})^2 \leqslant \frac{2M\Delta_M^2}{T^2(\sum_{n=1}^N \delta_n^{2\alpha})^2} \sum_{n=1}^N \delta_n^{4\alpha} (1 - e^{-2\theta_0 \delta_n^{2\alpha t_{i-1}}})^2 \leqslant \frac{2M\Delta_M^2}{T^2(\sum_{n=1}^N \delta_n^{2\alpha})^2} \sum_{n=1}^N \delta_n^{4\alpha} (1 - e^{-2\theta_0 \delta_n^{2\alpha t_{i-1}}})^2 \leqslant \frac{2M\Delta_M^2}{T^2(\sum_{n=1}^N \delta_n^{2\alpha})^2} \sum_{n=1}^N \delta_n^{4\alpha} (1 - e^{-2\theta_0 \delta_n^{2\alpha t_{i-1}}})^2 \leqslant \frac{2M\Delta_M^2}{T^2(\sum_{n=1}^N \delta_n^{2\alpha})^2} \sum_{n=1}^N \delta_n^{4\alpha} (1 - e^{-2\theta_0 \delta_n^{2\alpha t_{i-1}}})^2 \leqslant \frac{2M\Delta_M^2}{T^2(\sum_{n=1}^N \delta_n^{2\alpha})^2} \sum_{n=1}^N \delta_n^{4\alpha} (1 - e^{-2\theta_0 \delta_n^{2\alpha t_{i-1}}})^2 \leqslant \frac{2M\Delta_M^2}{T^2(\sum_{n=1}^N \delta_n^{2\alpha})^2} \sum_{n=1}^N \delta_n^{4\alpha} (1 - e^{-2\theta_0 \delta_n^{2\alpha t_{i-1}}})^2 \leqslant \frac{2M\Delta_M^2}{T^2(\sum_{n=1}^N \delta_n^{2\alpha})^2} \sum_{n=1}^N \delta_n^{2\alpha} (1 - e^{-2\theta_0 \delta_n^{2\alpha t_{i-1}}})^2 \leqslant \frac{2M\Delta_M^2}{T^2(\sum_{n=1}^N \delta_n^{2\alpha})^2} \sum_{n=1}^N \delta_n^{2\alpha} (1 - e^{-2\theta_0 \delta_n^{2\alpha t_{i-1}}})^2 \leqslant \frac{2M\Delta_M^2}{T^2(\sum_{n=1}^N \delta_n^{2\alpha t_{i-1}})^2} \leqslant \frac{2M\Delta_M^2}{T^2$$

For the term  $C_{N,M,T,2}$ , we have

$$\begin{split} C_{N,M,T,2} &= \frac{4\theta_0^2 \Delta_M^2}{T^2 (\sum\limits_{n=1}^N \delta_n^{2\alpha})^2} \sum_{n=1}^N \delta_n^{8\alpha} \sum_{i,j=1}^M <\varphi_n^{\theta_0}(t_{i-1}), \varphi_n^{\theta_0}(t_{j-1}) >_{L^2([0,T]^2)} \\ &= \frac{8\theta_0^2 \Delta_M^2}{T^2 (\sum\limits_{n=1}^N \delta_n^{2\alpha})^2} \sum_{n=1}^N \delta_n^{8\alpha} \sum_{i=1}^{M-1} \sum_{j=i+1}^M e^{-2\theta_0 \delta_n^{2\alpha}(t_{j-1} - t_{i-1})} \left( \int_0^{t_{i-1}} e^{-2\theta_0 \delta_n^{2\alpha} u} du \right)^2 \\ &= \frac{2\Delta_M^2}{T^2 (\sum\limits_{n=1}^N \delta_n^{2\alpha})^2} \sum_{n=1}^N \delta_n^{4\alpha} \sum_{i=1}^{M-1} \sum_{j=i+1}^M e^{-2\theta_0 \delta_n^{2\alpha} \Delta_M(j-i)} (1 - e^{-2\theta_0 \delta_n^{2\theta_0} t_{i-1}})^2 \\ &\leqslant \frac{2\Delta_M^2}{T^2 (\sum\limits_{n=1}^N \delta_n^{2\alpha})^2} \sum_{n=1}^N \delta_n^{4\alpha} \sum_{i=1}^{M-1} \sum_{r=1}^{M-i} e^{-2\theta_0 \delta_n^{2\alpha} \Delta_M r} \\ &\leqslant \frac{2M\Delta_M^2}{T^2 (\sum\limits_{n=1}^N \delta_n^{2\alpha})^2} \sum_{n=1}^N \delta_n^{4\alpha} \left( \frac{1 - e^{-2\theta_0 \delta_n^{2\alpha} T}}{1 - e^{-2\theta_0 \delta_n^{2\alpha} \Delta_M}} \right) \\ &\leqslant \frac{M\Delta_M}{\theta_0 T^2 (\sum\limits_{n=1}^N \delta_n^{2\alpha})^2} \sum_{n=1}^N \delta_n^{2\alpha} \frac{\Delta_M \delta_n^{2\alpha} 2\theta_0}{(1 - e^{-2\theta_0 \delta_n^{2\alpha} \Delta_M})} \sim c(\theta_0, \alpha, d, \bar{w}) \frac{1}{TN^{\frac{2\alpha}{d}+1}}, \end{split}$$

where we used the fact that under the assumption  $\Delta_M N^{\frac{2\alpha}{d}} \to 0$ , as  $N, M \to +\infty$ , we have

$$\delta_n^{2\alpha} \frac{\Delta_M \delta_n^{2\alpha} 2\theta_0}{(1 - e^{-2\theta_0 \delta_n^{2\alpha} \Delta_M})} \sim \delta_n^{2\alpha}, \text{ as } n, M \to +\infty.$$

Moreover, under the assumption  $\Delta_M N^{\frac{2\alpha}{d}} \to 0$ , as  $N, M, T \to +\infty$ , we get  $\frac{1}{MN} = o\left(\frac{1}{TN^{\frac{2\alpha}{d}+1}}\right)$ .

For the left hand term  $D_{N,M,T}$  in (36), using similar arguments as previously, we have

$$\begin{split} D_{N,M,T} - 1 &= \frac{\Delta_M}{T \sum_{n=1}^N \delta_n^{2\alpha}} \sum_{n=1}^N \delta_n^{2\alpha} \sum_{i=1}^M (1 - e^{-2\theta_0 \delta_n^{2\alpha} t_{i-1}}) - 1 \\ &= \frac{1}{T \sum_{n=1}^N \delta_n^{2\alpha}} \left[ \Delta_M \sum_{n=1}^N \delta_n^{2\alpha} \sum_{n=1}^M ((1 - e^{-2\theta_0 \delta_n^{2\alpha} t_{i-1}}) - 1) \right] \\ &= -\frac{\Delta_M}{T \sum_{n=1}^N \delta_n^{2\alpha}} \sum_{n=1}^N \delta_n^{2\alpha} \left( \frac{1 - e^{-2\theta_0 \delta_n^{2\alpha} T}}{1 - e^{-2\theta_0 \delta_n^{2\alpha} \Delta_M}} \right). \end{split}$$

Hence

$$|D_{N,M,T} - 1|^2 \leqslant c(\theta_0, \alpha, d, \bar{\sigma}) \frac{1}{T^2 N^{4\alpha/d}}.$$

**Lemma 10** For any N, M, T, the following estimate holds for the sequence  $\Lambda_{N,M}$ ,

$$\left| \mathbf{E} \left[ \left( \frac{\Lambda_{N,M}}{\sqrt{T \sum_{n=1}^{N} \delta_n^{2\alpha}}} \right)^2 \right] - \frac{1}{2\theta_0} \right| \leq \Delta_M \frac{\sum_{n=1}^{N} \delta_n^{4\alpha}}{\sum_{n=1}^{N} \delta_n^{2\alpha}} + \frac{N}{(2\theta_0)^2 T \sum_{n=1}^{N} \delta_n^{2\alpha}}.$$
 (37)

Moreover, if N, M, T satisfy  $\Delta_M N^{\frac{2\alpha}{d}} \to 0$ , as  $N, M, T \to +\infty$ , then

$$\mathbf{E}\left[\left(\frac{\Lambda_{N,M}}{\sqrt{T\sum_{n=1}^{N}\delta_{n}^{2\alpha}}}\right)^{2}\right] \longrightarrow \frac{1}{2\theta_{0}},\tag{38}$$

as  $N, M, T \to +\infty$ .

**Proof.** Since  $\eta_n$ , n = 1, ..., N are Gaussian independent processes with independent increments, we can write

$$\mathbf{E}\left[\left(\frac{\Lambda_{N,M}}{\sqrt{T}}\right)^{2}\right] = \frac{1}{T}\sum_{n=1}^{N}\mathbf{E}\left(B_{n,M}^{2}\right).$$

Next, since  $\eta_n(t_{i-1})$  and  $\eta_n(t_i) - \eta_n(t_{i-1})$  are independent, we have

$$\begin{split} &\frac{1}{T}\mathbf{E}\left(B_{n,M}^{2}\right) \\ &= \frac{\delta_{n}^{4\alpha}}{T}\sum_{i,j=1}^{M}e^{-\theta_{0}\delta_{n}^{2\alpha}(t_{i}+t_{i-1}+t_{j}+t_{j-1})}\mathbf{E}\left[\eta_{n}(t_{i-1})\left(\eta_{n}(t_{i})-\eta_{n}(t_{i-1})\right)\eta_{n}(t_{j-1})\left(\eta_{n}(t_{j})-\eta_{n}(t_{j-1})\right)\right] \\ &= \frac{\delta_{n}^{4\alpha}}{T}\sum_{i=1}^{M}e^{-2\theta_{0}\delta_{n}^{2\alpha}(t_{i}+t_{i-1})}\mathbf{E}\left[\eta_{n}(t_{i-1})^{2}\left(\eta_{n}(t_{i})-\eta_{n}(t_{i-1})\right)^{2}\right] \\ &= \frac{\delta_{n}^{4\alpha}}{T}\sum_{i=1}^{M}e^{-2\theta_{0}\delta_{n}^{2\alpha}(t_{i}+t_{i-1})}\mathbf{E}\left[\eta_{n}(t_{i-1})^{2}\right]\mathbf{E}\left[\left(\eta_{n}(t_{i})-\eta_{n}(t_{i-1})\right)^{2}\right] \\ &= \frac{\delta_{n}^{4\alpha}}{T}\sum_{i=1}^{M}e^{-2\theta_{0}\delta_{n}^{2\alpha}(t_{i}+t_{i-1})}\left(\frac{e^{2\theta_{0}\delta_{n}^{2\alpha}t_{i-1}}-1}{2\theta_{0}\delta_{n}^{2\alpha}}\right)\left(\frac{e^{2\theta_{0}\delta_{n}^{2\alpha}t_{i}}-e^{2\theta_{0}\delta_{n}^{2\alpha}t_{i-1}}}{2\theta_{0}\delta_{n}^{2\alpha}}\right) \\ &= \frac{\left(1-e^{-2\theta_{0}\delta_{n}^{2\alpha}\Delta_{M}}\right)}{(2\theta_{0})^{2}\Delta_{M}}\frac{1}{M}\sum_{i=1}^{M}\left(1-e^{-2\theta_{0}\delta_{n}^{2\alpha}\Delta_{M}}\right)\left(\frac{1-e^{-2\theta_{0}\delta_{n}^{2\alpha}T}}{M(1-e^{-2\theta_{0}\delta_{n}^{2\alpha}\Delta_{M}})}\right) \\ &= \frac{\left(1-e^{-2\theta_{0}\delta_{n}^{2\alpha}\Delta_{M}}\right)}{(2\theta_{0})^{2}\Delta_{M}}-\frac{\left(1-e^{-2\theta_{0}\delta_{n}^{2\alpha}T}\right)}{(2\theta_{0})^{2}T}. \end{split}$$

This implies,

$$\begin{split} \sum_{n=1}^{N} \left[ \frac{1}{T} \mathbf{E} \left( B_{n,M}^2 \right) - \frac{\delta_n^{2\alpha}}{2\theta_0} \right] &= \sum_{n=1}^{N} \left[ \frac{\left( 1 - e^{-2\theta_0 \delta_n^{2\alpha} \Delta_M} - 2\theta_0 \delta_n^{2\alpha} \Delta_M \right)}{(2\theta_0)^2 \Delta_M} - \frac{\left( 1 - e^{-2\theta_0 \delta_n^{2\alpha} T} \right)}{(2\theta_0)^2 T} \right] \\ &\leqslant \sum_{n=1}^{N} \left[ \frac{\left| 1 - e^{-2\theta_0 \delta_n^{2\alpha} \Delta_M} - 2\theta_0 \delta_n^{2\alpha} \Delta_M \right|}{(2\theta_0)^2 \Delta_M} + \frac{1}{(2\theta_0)^2 T} \right] \\ &\leqslant \sum_{n=1}^{N} \left[ \delta_n^{4\alpha} \Delta_M + \frac{1}{(2\theta_0)^2 T} \right], \end{split}$$

where the last inequality follows from the fact that  $|1 - e^{-x} - x|/x^2 \le 1$  for all x > 0, by applying Mean Value Theorem twice.

Therefore,

$$\left| \frac{1}{T \sum_{n=1}^{N} \delta_n^{2\alpha}} \mathbf{E} \left( B_{n,M}^2 \right) - \frac{1}{2\theta_0} \right| \leq \Delta_M \frac{\sum_{n=1}^{N} \delta_n^{4\alpha}}{\sum_{n=1}^{N} \delta_n^{2\alpha}} + \frac{N}{(2\theta_0)^2 T \sum_{n=1}^{N} \delta_n^{2\alpha}}, \tag{39}$$

which proves (37). The limit (38), is a direct consequence of (37) and the fact that when  $N, M, T \to +\infty$ , the following hold true

$$\Delta_M \frac{\sum_{n=1}^N \delta_n^{4\alpha}}{\sum_{n=1}^N \delta_n^{2\alpha}} \sim \frac{T}{M} \bar{\sigma}^{\alpha} N^{\frac{2\alpha}{d}} \left( \frac{2\alpha + d}{4\alpha + d} \right).$$

$$\frac{N}{(2\theta_0)^2 T \sum_{n=1}^N \delta_n^{2\alpha}} \sim \frac{(2\alpha + d)}{T N^{\frac{2\alpha}{d}} d(2\theta_0)^2 \bar{\sigma}^{\alpha}}.$$

The desired result follows.

# 4.2 Asymptotic normality of the estimator $\widetilde{\theta}_{T,N,M}$

Lemma 11 We have

$$\kappa_3 \left( \frac{\Lambda_{N,M}}{\sqrt{T \sum_{n=1}^N \delta_n^{2\alpha}}} \right) = 0 \text{ and } \kappa_4 \left( \frac{\Lambda_{N,M}}{\sqrt{T \sum_{n=1}^N \delta_n^{2\alpha}}} \right) \leqslant \frac{1}{M \left( 2\theta_0 \right)^2} \frac{\sum_{n=1}^N \delta_n^{4\alpha}}{\left( \sum_{n=1}^N \delta_n^{2\alpha} \right)^2}$$
(40)

**Proof.** Using  $\mathbf{E}[\Lambda_{N,M}] = 0$  and the fact that  $\eta_n$ , n = 1, ..., N are Gaussian independent processes with independent increments, we can write

$$\kappa_3 \left( \frac{1}{\sqrt{T \sum_{n=1}^N \delta_n^{2\alpha}}} \Lambda_{N,M} \right) = \mathbf{E} \left[ \left( \frac{1}{\sqrt{T \sum_{n=1}^N \delta_n^{2\alpha}}} \Lambda_{N,M} \right)^3 \right] = \frac{1}{\left( T \sum_{n=1}^N \delta_n^{2\alpha} \right)^{3/2}} \sum_{n=1}^N \mathbf{E} \left( B_{n,M}^3 \right).$$

Moreover, since  $\eta_n(t_{i-1})$  and  $\eta_n(t_i) - \eta_n(t_{i-1})$  are independent, then  $\mathbf{E}\left(B_{n,M}^3\right) = 0$ , which implies (40). Since  $B_{n,M}$ ,  $n = 1, \ldots, N$  are independent and  $\mathbf{E}\left(B_{n,M}\right) = 0$ ,

$$\mathbf{E}\left(\Lambda_{N,M}^{4}\right) = 3\sum_{n\neq j=1}^{N} \mathbf{E}\left(B_{n,M}^{2}\right) \mathbf{E}\left(B_{j,M}^{2}\right) + \sum_{n=1}^{N} \mathbf{E}\left(B_{n,M}^{4}\right)$$

and  $\mathbf{E}\left[\Lambda_{N,M}^2\right] = \sum_{n=1}^N \mathbf{E}\left(B_{n,M}^2\right)$  . Hence

$$\kappa_{4} \left( \Lambda_{N,M} \right) = \mathbf{E} \left[ \Lambda_{N,M}^{4} \right] - 3 \left[ \mathbf{E} \left( \Lambda_{N,M}^{2} \right) \right]^{2}$$

$$= \sum_{k=1}^{N} \mathbf{E} \left( B_{n,M}^{4} \right) - 3 \sum_{n=1}^{N} \left[ \mathbf{E} \left( B_{n,M}^{2} \right) \right]^{2}$$

$$= \sum_{n=1}^{N} \left( \mathbf{E} \left( B_{n,M}^{4} \right) - 3 \left[ \mathbf{E} \left( B_{n,M}^{2} \right) \right]^{2} \right).$$

On the other hand, similar arguments as in [13] imply

$$\left| \mathbf{E} \left( B_{n,M}^4 \right) - 3 \left[ \mathbf{E} \left( B_{n,M}^2 \right) \right]^2 \right| \leq \left( \delta_n^{2\alpha} \right)^4 M \frac{\left( 1 - e^{-2\theta_0 \delta_n^{2\alpha} \Delta_M} \right)^2}{\left( 2\theta_0 \delta_n^{2\alpha} \right)^4} \leq \left( \delta_n^{2\alpha} \right)^2 M \frac{\Delta_M^2}{\left( 2\theta_0 \right)^2},$$

where the last inequality follows from the fact that  $|1 - e^{-x}|/x \le 1$  for all x > 0, by Mean Value Theorem. Therefore,

$$\kappa_4 \left( \frac{\Lambda_{N,M}}{\sqrt{T \sum_{n=1}^N \delta_n^{2\alpha}}} \right) \leq \frac{1}{M (2\theta_0)^2} \frac{\sum_{n=1}^N \delta_n^{4\alpha}}{\left(\sum_{n=1}^N \delta_n^{2\alpha}\right)^2},$$

which implies (40)

**Theorem 12** There exists a constant  $C(\alpha, \theta_0, d)$  such that for N, M, T large, we have

$$d_W\left(\frac{\Lambda_{N,M}}{\sqrt{\psi_{T,N}^{\theta_0}}}, Z\right) \leqslant C(\alpha, \theta_0, d) \times \frac{1}{MN}. \tag{41}$$

where  $\psi_{T,N}^{\theta_0} = \frac{T}{2\theta_0} \sum_{n=1}^{N} \delta_n^{2\alpha}$  and  $Z \sim \mathcal{N}(0,1)$ . Consequently, as  $N, M, T \to +\infty$ .

$$\frac{\Lambda_{N,M}}{\sqrt{T}N^{\frac{\alpha}{d}+\frac{1}{2}}} \xrightarrow{law} \mathcal{N}\left(0, \frac{\bar{\sigma}^{\alpha}d}{\theta_0(4\alpha+2d)}\right).$$

**Proof.** Theorem (12) is a direct consequence of (9), Lemma (11) and the fact that for N, M large, we have

$$\frac{1}{M} \frac{\sum_{n=1}^{N} \delta_n^{4\alpha}}{\left(\sum_{n=1}^{N} \delta_n^{2\alpha}\right)^2} \sim \frac{1}{MN} \frac{1}{d} \frac{(2\alpha + d)^2}{4\alpha + d}.$$

 $\blacksquare$ By the decomposition (31), we can write

$$\sqrt{\psi_{T,N}^{\theta_0}} \left( \theta_0 - \tilde{\theta}_{T,N,M} \right) = \sqrt{\psi_{T,N}^{\theta_0}} \frac{R_{N,M}}{S_{N,M}} + \frac{\Lambda_{N,M} / \sqrt{\psi_{T,N}^{\theta_0}}}{S_{N,M} / \psi_{T,N}^{\theta_0}}.$$

Therefore, we can write using the fact that  $d_W(X+Y,Z) \leq E[|X|] + d_W(Y,Z)$  for all X,Y,Z random variables such that  $X \in L^1(\Omega)$ .

$$d_{W}\left(\sqrt{\psi_{T,N}^{\theta_{0}}}\left(\theta_{0}-\tilde{\theta}_{T,N,M}\right),Z\right)\leqslant\sqrt{\psi_{T,N}^{\theta_{0}}}\mathbf{E}\left[\left|\frac{R_{N,M}}{S_{N,M}}\right|\right]+d_{W}\left(\frac{\Lambda_{N,M}/\sqrt{\psi_{T,N}^{\theta_{0}}}}{S_{N,M}/\psi_{T,N}^{\theta_{0}}},Z\right).$$
(42)

On the other hand, using triangular inequality for the distance  $d_W$  we get

$$\begin{split} d_{W}\left(\frac{\Lambda_{N,M}/\sqrt{\psi_{T,N}^{\theta_{0}}}}{S_{N,M}/\psi_{T,N}^{\theta_{0}}},Z\right) & \leqslant d_{W}\left(\frac{\Lambda_{N,M}}{\sqrt{\psi_{T,N}^{\theta_{0}}}},Z\right) + d_{W}\left(\frac{\Lambda_{N,M}/\sqrt{\psi_{T,N}^{\theta_{0}}}}{S_{N,M}/\psi_{T,N}^{\theta_{0}}},\frac{\Lambda_{N,M}}{\sqrt{\psi_{T,N}^{\theta_{0}}}}\right) \\ & \leqslant d_{W}\left(\frac{\Lambda_{N,M}}{\sqrt{\psi_{T,N}^{\theta_{0}}}},Z\right) + \mathbf{E}\left[\left|\frac{\Lambda_{N,M}/\sqrt{\psi_{T,N}^{\theta_{0}}}}{S_{N,M}/\psi_{T,N}^{\theta_{0}}} - \frac{\Lambda_{N,M}}{\sqrt{\psi_{T,N}^{\theta_{0}}}}\right|\right] \\ & \leqslant d_{W}\left(\frac{\Lambda_{N,M}}{\sqrt{\psi_{T,N}^{\theta_{0}}}},Z\right) + \left\|\frac{\Lambda_{N,M}/\sqrt{\psi_{T,N}^{\theta_{0}}}}{S_{N,M}/\psi_{T,N}^{\theta_{0}}}\right\|_{L^{2}} \times \|1 - \frac{S_{N,M}}{\psi_{T,N}^{\theta_{0}}}\|_{L^{2}} \\ & \leqslant d_{W}\left(\frac{\Lambda_{N,M}}{\sqrt{\psi_{T,N}^{\theta_{0}}}},Z\right) + \|\frac{\Lambda_{N,M}}{\sqrt{\psi_{T,N}^{\theta_{0}}}}\|_{L^{4}} \times \|\frac{\psi_{T,N}^{\theta_{0}}}{S_{N,T}}\|_{L^{4}} \times \|1 - \frac{S_{N,M}}{\psi_{T,N}^{\theta_{0}}}\|_{L^{2}}. \end{split}$$

Notice that the sequence  $\Lambda_{N,M}$  is a second chaos random variable, indeed by (28), we can write

$$\Lambda_{N,M} = \sum_{n=1}^{N} \delta_n^{2\alpha} \sum_{i=1}^{M} I_1^{w_n} \left( e^{-\theta_0 \delta_n^{2\alpha} (t_{i-1} - \cdot)} \mathbf{1}(\cdot)_{[0,t_{i-1}]} \right) I_1^{w_n} \left( e^{-\theta_0 \delta_n^{2\alpha} (t_{i-1} - \cdot)} \mathbf{1}(\cdot)_{[t_{i-1},t_i]} \right) \\
= \sum_{n=1}^{N} I_2^{w_n} \left[ \delta_n^{2\alpha} \sum_{i=1}^{M} \left( e^{-\theta_0 \delta_n^{2\alpha} (t_{i-1} - \cdot)} \mathbf{1}(\cdot)_{[0,t_{i-1}]} \otimes e^{-\theta_0 \delta_n^{2\alpha} (t_{i-1} - \cdot)} \mathbf{1}(\cdot)_{[t_{i-1},t_i]} \right) \right].$$

Then, by the hypercontractivity property (8), we get  $\|\frac{\Lambda_{N,M}}{\sqrt{\psi_{T,N}^{\theta_0}}}\|_{L^4} \leqslant 3\|\frac{\Lambda_{N,M}}{\sqrt{\psi_{T,N}^{\theta_0}}}\|_{L^2} < +\infty$ .

On the other hand, by (47) in Lemma 14, we have  $\|\frac{\psi_{T,N}^{\theta_0}}{S_{N,T}}\|_{L^4} < +\infty$ . Finally, by Lemma 9, the following estimates hold true

$$\|\frac{S_{N,M}}{\psi_{T,N}^{\theta_0}} - 1\|_{L^2} \leqslant C(\alpha, \theta_0, \bar{\sigma}, d) \times b_{N,T}$$
(43)

with  $b_{N,T}$  is the same bound in (22). A comparison, between the bound (41) obtained in Lemma 12 and (22), we can easily checn that under the assumption  $\Delta_M N^{\frac{2\alpha}{d}} \to 0$  as  $N, M, T \to +\infty$ , we get

$$\frac{1}{MN} = \begin{cases}
o\left(\frac{1}{\sqrt{T}} \frac{1}{N^{\frac{2\alpha}{d}}}\right) & \text{if } 2\alpha < d \\
o\left(\frac{1}{\sqrt{T}} \frac{1}{N}\right) & \text{if } 2\alpha = d \\
o\left(\frac{1}{\sqrt{T}} \frac{1}{N^{\frac{\alpha}{d} + \frac{1}{2}}}\right) & \text{if } 2\alpha > d.
\end{cases}$$
(44)

Now, by (42) it remains to control the convergence to zero of the term  $\sqrt{\psi_{T,N}^{\theta_0}} \mathbf{E}\left[\left|\frac{R_{N,M}}{S_{N,M}}\right|\right]$  when  $T, M, N \to +\infty$ . Indeed, by Cauchy-Schwartz inequality, we have

$$\sqrt{\psi_{T,N}^{\theta_0}} \mathbf{E} \left[ \left| \frac{R_{N,M}}{S_{N,M}} \right| \right] \leqslant \mathbf{E} \left[ \left( \frac{R_{N,M}}{\sqrt{\psi_{T,N}^{\theta_0}}} \right)^2 \right]^{1/2} \times \mathbf{E} \left[ \left( \frac{\psi_{T,N}^{\theta_0}}{S_{N,M}} \right)^2 \right]^{1/2}.$$

By by (47) in Lemma 14, we have  $\|\frac{\psi_{T,N}^{\theta_0}}{S_{N,T}}\|_{L^2} < +\infty$ . On the other hand, from (35) of Lemma 8, we have  $\mathbf{E}[R_{N,M}^2] = A_{N,M,T} + B_{N,M,T}$ . Moreover,

$$\frac{A_{N,M,T}}{\psi_{T,N}^{\theta_0}} \leqslant \frac{\theta_0^4 M^2 \Delta_M^4}{T\left(\sum_{n=1}^N \delta_n^{2\alpha}\right)} \left(\sum_{n=1}^N \delta_n^{4\alpha}\right)^2 \leqslant c(\theta_0, \alpha, \bar{\sigma}, d) \frac{T^3}{M^2} N^{\frac{6\alpha}{d} + 1},$$

and

$$\frac{B_{N,M,T}}{\psi_{T,N}^{\theta_0}} \leqslant \frac{2\theta_0^4 M^2 \Delta_M^4}{T\left(\sum_{n=1}^N \delta_n^{2\alpha}\right)} \sum_{n=1}^N \delta_n^{8\alpha} \leqslant c(\theta_0, \alpha, \bar{\sigma}, d) \frac{T^3}{M^2} N^{\frac{6\alpha}{d}}.$$

We conclude that if

$$\frac{T^{3/2}N^{\frac{3\alpha}{d}+\frac{1}{2}}}{M} \to 0 \text{ as } N, M, T \to +\infty.$$

Then, there exists a constant  $c(\theta_0, \alpha, \bar{\sigma}, d) > 0$  such that

$$\sqrt{\psi_{T,N}^{\theta_0}} \mathbf{E} \left[ \left| \frac{R_{N,M}}{S_{N,M}} \right| \right] \leqslant c(\theta_0, \alpha, \bar{\sigma}, d) \frac{T^{3/2} N^{\frac{3\alpha}{d} + \frac{1}{2}}}{M} \to 0$$

$$\tag{45}$$

as  $M, N, T \to +\infty$ .

**Theorem 13** Consider the estimator  $\tilde{\theta}_{T,N,M}$  defined in (26). Then, there exists a constant  $C(\alpha, \theta_0, \bar{\sigma}, d)$  such that

$$d_W\left(\sqrt{\psi_{T,N}^{\theta_0}}\left(\theta_0 - \tilde{\theta}_{T,N,M}\right), Z\right) \leqslant C(\alpha, \theta_0, \bar{\sigma}, d) \max\left(\frac{T^{3/2} N^{\frac{3\alpha}{d} + \frac{1}{2}}}{M}, b_{N,T}\right)$$

where  $b_{N,T}$  is defined in (22) and  $Z \sim \mathcal{N}(0,1)$ . In particular, if

$$\frac{T^{3/2}N^{\frac{3\alpha}{d}+\frac{1}{2}}}{M} \rightarrow 0 \quad as \quad N,M,T \rightarrow +\infty.$$

Then, as  $T, M, N \to +\infty$ ,

$$\sqrt{T}N^{\frac{\alpha}{d}+\frac{1}{2}}\left(\theta_0-\tilde{\theta}_{T,N,M}\right) \xrightarrow{law} \mathcal{N}\left(0,\frac{(4\alpha/d+2)\theta_0}{\bar{\sigma}^{\alpha}}\right).$$

#### 5 Appendix

**Lemma 14** Let  $N \ge 1$  and consider  $\psi_{T,N}^{\theta_0}$  defined in (13), then for every  $p, T_0 > 0$ , we have

$$\sup_{T\geqslant T_0} \mathbf{E} \left[ \left( \frac{1}{\psi_{T,N}^{\theta_0}} \sum_{i=1}^N \delta_i^{4\alpha} \int_0^T v_i^2(s) ds \right)^{-p} \right] < \infty, \tag{46}$$

Moreover, for every p > 0, there exists  $M_0 \ge 1$  such that

$$\sup_{M\geqslant M_0} \mathbf{E} \left[ \left( \frac{1}{T \sum_{i=1}^N \delta_j^{2\alpha}} \Delta_M \sum_{i=1}^N \delta_i^{4\alpha} \sum_{k=0}^{M-1} v_i^2(t_k) \right)^{-p} \right] < +\infty, \tag{47}$$

**Proof.** The proof of assertion (46) relies on similar arguments as in the proof of Lemma 1 in [14]. For the discrete case (47) Let T > 0. Recall that for k = 1, ..., N, the processes  $v_k(t) := \int_0^t e^{-\theta_0 \delta_k^{2\alpha}(t-u)} dw_k(u)$ ,  $t \in [0, T]$ , where  $w_1, ..., w_n$  are N independent Brownian motions. Let  $w = (w_1, ..., w_N)$  and define  $\mathcal{F}_t^w = \sigma(\bigcup_{i=1}^N \mathcal{F}_t^{w_i})$  where  $\mathcal{F}_t^{w_i} = \sigma\{w_i(u), u \leq t\}$  denotes the sigma-field generated by  $w_i$  for i = 1, ..., N. Fix  $N \geqslant 1$  and p > 0 and let  $M_0 \geqslant 1$  an integer such that for every  $M \geqslant M_0$  and M > 2p, we have

$$\int_{0}^{\frac{T}{M}} e^{-2\theta_0 \delta_i^{2\alpha} u} du \geqslant \frac{1}{4\theta_0 \delta_i^{2\alpha}}, \quad i = 1, ..., N.$$

$$\tag{48}$$

Then, consider the following processes defined for every i = 1, ..., N as

$$Y_{i,t} := \sum_{k=0}^{M-1} v_i(t_k) \mathbf{1}_{]t_{k-1},t_k]}(t),$$

where for k=0,...,M  $t_k=k\Delta_M$ ,  $\Delta_M=\frac{T}{M}\to 0$  and  $M\Delta_M\to +\infty$  as  $M\to +\infty$ . We can easily check that

$$\frac{1}{T} \int_0^T Y_{i,t}^2 dt = \frac{1}{M} \sum_{k=0}^{M-1} v_i^2(t_k).$$

We can follow the same steps of the continuous case with  $Y_{i,t}$  instead of  $v_i(t)$ . The only remaining part to be proved is a lower bound for all  $M \ge M_0$  of the term for any k = 1, ..., N

$$A_{M,N,k} := \mathbf{E} \left[ \frac{\sum_{i=1}^{N} \delta_i^{4\alpha}}{T \sum_{j=1}^{N} \delta_j^{4\alpha}} \int_{(k-1)T/M}^{kT/M} Y_{i,t}^2 dt \mid \mathcal{F}_{(k-1)T/M}^w \right]. \tag{49}$$

Or for k = 1, ..., N, we have for any  $M \ge M_0$ ,

$$A_{M,N,k} = \frac{1}{M} \frac{\sum_{i=1}^{N} \delta_{i}^{4\alpha}}{\sum_{j=1}^{N} \delta_{j}^{2\alpha}} \int_{(k-1)T/M}^{kT/M} \mathbf{E}[v_{i}^{2}(t_{k}) \mid \mathcal{F}_{(k-1)T/M}^{w_{i}}] \geqslant \frac{1}{M} \frac{\sum_{i=1}^{N} \delta_{i}^{4\alpha}}{\sum_{j=1}^{N} \delta_{j}^{2\alpha}} \left[ \int_{(k-1)T/M}^{kT/M} e^{-2\theta_{0}\delta_{i}^{2\alpha}(\frac{kT}{M} - u)} du \right]$$

$$\geqslant \frac{1}{M} \frac{\sum_{i=1}^{N} \delta_{i}^{4\alpha}}{\sum_{j=1}^{N} \delta_{j}^{2\alpha}} \int_{0}^{T/M} e^{-2\theta_{0}\delta_{i}^{2\alpha}u} du \geqslant \frac{1}{M} \frac{\sum_{i=1}^{N} \delta_{i}^{4\alpha}}{\sum_{j=1}^{N} \delta_{j}^{2\alpha}} \frac{1}{4\theta_{0}\delta_{i}^{2\alpha}} = \frac{1}{4\theta_{0}M}.$$

which yields the proof of (47).

## 6 Acknowlegment

This research project was funded by Princess Nourah bint Abdulrahman University Researchers Supporting Project number (PNURSP2024R358), Princess Nourah bint Abdulrahman University, Riyadh, Saudi Arabia.

#### References

- [1] Shubin, M.A. Pseudodifferential Operators and Spectral Theory, 2nd edn. Springer, Berlin. 2001.
- [2] Chow, P. Stochastic Partial Differential Equations. Applied Mathematics and Nonlinear Science Series. Chapman & Hall, Boca Raton, 2007.
- [3] Lototsky, S.V., Rozovsky, B.L. Stochastic Partial Differential Equations. Springer, New York, 2017.
- [4] Lototsky, S.V., Rozovsky, B.L. Stochastic Evolution Systems. Linear Theory and Applications to Non-linear Filtering. *Probability Theory and Stochastic Modelling*, **89**, 2nd edn. Springer, New York, 2018.
- [5] Cialenco, I. Statistical inference for SPDEs: an overview. Stat Inference Stoch Process. 2018 21 309-329. https://doi.org/10.1007/s11203-018-9177-9.
- [6] Cialenco, I., Xu, L. Hypothesis testing for stochastic PDEs driven by additive noise. *Stoch. Process.* Appl., **125**(3), (2015) 819–866.

- [7] Cialenco, I., Delgado-Vences, F., Kim, H. J. Drift estimation for discretely sampled SPDEs. Stoch PDE: Anal. Comp., 8, (2001) 895–920. https://doi.org/10.1007/s40072-019-00164-4.
- [8] Nourdin, I. and Peccati, G. The optimal fourth moment theorem. *Proc. Amer. Math. Soc.*, **143** (2015) 3123-3133.
- [9] Nourdin, I. and Peccati, G. Normal approximations with Malliavin calculus: from Stein's method to universality. In *Cambridge Tracts in Mathematics* 192. Cambridge University Press, Cambridge. 2012.
- [10] Nualart, D. The Malliavin calculus and related topics. Springer-Verlag, Berlin. 2006.
- [11] Douissi, S., Es-Sebaiy, K., Kerchev, G., Nourdin I. Berry-esseen bounds of second moment estimators for gaussian processes observed at high frequency, *Electron. J. Statist.16*(1): (2022) 636–670.
- [12] Hu, Y., Nualart, D., Zhou, H. Parameter estimation for fractional Ornstein-Uhlenbeck processes of general Hurst parameter. *Statistical Inference for Stochastic Processes*, **22**(1), (2019) 111–142.
- [13] Es-Sebaiy, K., Al-Foraih, M., Alazemi, F. Wasserstein bounds in CLT of approximative MCE and MLE of the drift parameter for Ornstein-Uhlenbeck processes observed at high frequency. *J. Inequal. Appl.* (2023), 62. https://doi.org/10.1186/s13660-023-02976-4.
- [14] Es-Sebaiy, K., Al-Foraih, M., Alazemi, F. Wasserstein Bounds in the CLT of the MLE for the drift Coefficient of a Stochastic Partial Differential Equation. Fractal Fract., 5, (2021) 187.