

Highlights

Bilinearization-reduction approach to the classical and nonlocal semi-discrete modified Korteweg-de Vries equations with nonzero backgrounds

Xiao Deng, Hongyang Chen, Song-Lin Zhao, Guanlong Ren

- The quasi double Casoratian solutions to the classical and nonlocal semi-discrete modified Korteweg-de Vries equations with nonzero backgrounds are found.
- Solitonic, periodic and rational solutions are obtained by using the quasi double Casoratians.

Bilinearization-reduction approach to the classical and nonlocal semi-discrete modified Korteweg-de Vries equations with nonzero backgrounds

Xiao Deng^a, Hongyang Chen^a, Song-Lin Zhao^b and Guanlong Ren^{c,*}

^aResearch Center for Data Hub and Security, Zhejiang Lab, Hangzhou, 311100, Zhejiang Province, China

^bSchool of Mathematical Sciences, Zhejiang University of Technology, Hangzhou, 310023, Zhejiang Province, China

^cResearch Center for Space Computing System, Zhejiang Lab, Hangzhou, 311100, Zhejiang Province, China

ARTICLE INFO

Keywords:

semi-discrete modified Korteweg-de Vries equation
Ablowitz-Ladik equation
nonzero backgrounds
Casoratian

ABSTRACT

Quasi double Casoratian solutions are derived for a bilinear system reformulated from the coupled semi-discrete modified Korteweg-de Vries equations with nonzero backgrounds. These solutions, when applied with the classical and nonlocal reduction techniques, also satisfy the corresponding classical and nonlocal semi-discrete modified Korteweg-de Vries equations with nonzero backgrounds. They can be expressed explicitly, allowing for an easy investigation of the dynamics of systems. As illustrative examples, the dynamics of solitonic, periodic and rational solutions with a plane wave background are examined for the focusing semi-discrete Korteweg-de Vries equation and the defocusing reverse-space-time complex semi-discrete Korteweg-de Vries equation.

1. Introduction

This paper aims to investigate solutions with nonzero backgrounds for the (classical real) semi-discrete modified Korteweg-de Vries (sd-mKdV) equation (Hirota, 1972)

$$\partial_t Q_n = (1 - \delta Q_n Q_n)(Q_{n+1} - Q_{n-1}) \quad (1)$$

and its classical and nonlocal analogous (see eqs.(5)). Here, $Q_n = Q(n, t)$ is a function of discretized-space and time $(n, t) \in \mathbb{Z} \times \mathbb{R}$, ∂_t denotes the derivative to time t and $\delta = \pm 1$. This equation is termed focusing sd-mKdV when $\delta = -1$ and defocusing sd-mKdV when $\delta = 1$.

The sd-mKdV equation, which can be derived from the Ablowitz-Ladik (AL) spectral problem (Ablowitz and Ladik, 1976) and treated as a modified Volterra equation (Wadati, 1976), is an important integrable spatial discretization of the mKdV equation (Wadati, 1973). The mKdV equation, which is also referred as the 3rd (reduced) equation in Ablowitz-Kaup-Newell-Segur (AKNS) hierarchy (Ablowitz et al., 1973), plays a vital role in mathematically modeling various nonlinear physical phenomena, such as hydromagnetic waves in plasma (Kakutani and Ono, 1969) and optical pulses in nonlinear media (Leblond and Sanchez, 2003; Leblond and Mihalache, 2013) (see Zhang et al. (2014) for more examples and references therein). Furthermore, the sd-mKdV equation itself arises in nonlinear models describing wave propagation in ladder type network (Hirota, 1972), motions of discrete curves (Doliwa and Santini, 1995) and motions of discrete surfaces (Hisakado and Wadati, 1995). As an integrable system, the sd-mKdV equation is also characterized with rich mathematical structures like Lax pair, infinite many conservation laws and infinite many symmetries (Ablowitz and Ladik, 1976; Fu et al., 2015).

When a nonlocal version of the nonlinear Schrödinger (NLS) equation, which is able to describe wave phenomena in nonlinear parity-time (PT) symmetric media and admits the PT symmetry (Sarma et al., 2014), is proposed by Ablowitz and Musslimani (2013), nonlocal integrable systems attract much attention and become a hot research topic. Like many other nonlocal integrable systems, nonlocal (real and complex) sd-mKdV equations can be obtained from a coupled system with nonlocal reductions and inherit many characteristics found in classical integrable systems (Ablowitz and Musslimani, 2017).

*Corresponding author

✉ rengl@zhejianglab.org (G. Ren)

ORCID(s): 0009-0001-2573-4276 (G. Ren)

Many methods (Common, 1993; Mukaihira and Nakamura, 2000; Zhang et al., 2012; Zhu et al., 2013; Wang and Ma, 2010; Ankiewicz et al., 2010; Sun and Zhao, 2019; Feng et al., 2020) are applied to solve the sd-mKdV equation and its analogous exactly. Among these, the bilinearization-reduction approach proposed by Chen et al. (2018) is a powerful technique in solving the classical and nonlocal integrable systems. This method first constructs double Wronskian/Casoratian solutions for a coupled integrable system using associated bilinear forms, and then enforces these solutions to satisfy the classical and nonlocal reduced systems (Zhang, 2023) by directly imposing reduction constraints. This approach has been successfully applied to various integrable systems with zero background, such as AKNS hierarchy (Chen et al., 2018), negative order AKNS hierarchy (Wang et al., 2020), derivative NLS equations (Shi et al., 2019) and sdNLS equations (Deng et al., 2018) (for more examples see Zhang (2023) and reference therein). Recently, this method has also been implemented for NLS equations with nonzero backgrounds (Zhang et al., 2023) and sdNLS equations with nonzero backgrounds (Deng et al., 2024). This progress motivates us to apply the bilinearization-reduction approach to integrable sd-mKdV equations with nonzero backgrounds. Here, solutions with nonzero backgrounds are characterized by the property that they do not tend to zero as $|n| \rightarrow \infty$.

This paper is organized as follows. In Sec.2, the unreduced system of the coupled sd-mKdV equations, along with its integrable reductions are presented. Sec.3 presents the Casoratian solutions of the unreduced system, which is derived from its associated bilinearized form with nonzero backgrounds. Sec.4 details the classical and nonlocal reduction techniques applied to the Casoratian solutions. The dynamics of solitonic, periodic and rational solutions are illustrated for the focusing sd-mKdV equation and the nonlocal complex defocusing sd-mKdV equation with space-time reversed in Sec.5. The last section is devoted to conclusions.

2. The unreduced system and its reductions

The unreduced system of the coupled sd-mKdV equations reads

$$\partial_t Q_n = (1 - Q_n R_n)(Q_{n+1} - Q_{n-1}), \quad (2a)$$

$$\partial_t R_n = (1 - Q_n R_n)(R_{n+1} - R_{n-1}), \quad (2b)$$

which can be derived from a Lax pair (Ablowitz and Ladik, 1976), i.e., the compatibility condition of the Ablowitz-Ladik (AL) spectral problem

$$\Theta_{n+1} = \mathcal{M}_n \Theta_n, \quad \mathcal{M}_n = \begin{bmatrix} z & Q_n \\ R_n & z^{-1} \end{bmatrix}, \quad \Theta_n = \begin{bmatrix} \theta_{1,n} \\ \theta_{2,n} \end{bmatrix}, \quad (3)$$

and the following time evolution equation

$$\partial_t \Theta_n = \mathcal{N}_n \Theta_n, \quad \mathcal{N}_n = \frac{1}{2} \begin{bmatrix} z^2 - z^{-2} - 2Q_n R_{n-1} & 2Q_n z + 2Q_{n-1} z^{-1} \\ 2R_{n-1} z + 2R_n z^{-1} & z^{-2} - z^2 - 2R_n Q_{n-1} \end{bmatrix}. \quad (4)$$

It admits the following reductions

$$\partial_t Q_n = (1 - \delta Q_n Q_n)(Q_{n+1} - Q_{n-1}), \quad R_n = \delta Q_n, \quad (5a)$$

$$\partial_t Q_n = (1 - \delta Q_n Q_n^*)(Q_{n+1} - Q_{n-1}), \quad R_n = \delta Q_n^*, \quad (5b)$$

$$\partial_t Q_n = (1 - \delta Q_n Q_{-n}(-t))(Q_{n+1} - Q_{n-1}), \quad R_n = \delta Q_{-n}(-t), \quad (5c)$$

$$\partial_t Q_n = (1 - \delta Q_n Q_{-n}^*(-t))(Q_{n+1} - Q_{n-1}), \quad R_n = \delta Q_{-n}^*(-t). \quad (5d)$$

Here, complex conjugate is denoted by $*$ and the function Q with reverse-space-time is denoted by $Q_{-n}(-t) = Q(-n, -t)$. Eqs. (5a) and (5b), which are obtained with classical reductions, are classical real and complex sd-mKdV equations, respectively. Eqs. (5c) and (5d), which are obtained with nonlocal reduction techniques, are nonlocal real and complex sd-mKdV equations with reverse-space-time, respectively.

3. Bilinearization and Casoratian solutions

In this section, the construction of the Casoratian solutions to the unreduced system (2) is shown.

Using the following variable transformation

$$Q_n = G_n / F_n, \quad R_n = H_n / F_n, \quad (6)$$

the unreduced system (2) is transformed to a bilinear system

$$(1 - q_n r_n) F_{n+1} F_{n-1} + G_n H_n - F_n^2 = 0, \quad (7a)$$

$$D_t G_n \cdot F_n = (1 - q_n r_n)(G_{n+1} F_{n-1} - G_{n-1} F_{n+1}), \quad (7b)$$

$$D_t H_n \cdot F_n = (1 - q_n r_n)(F_{n+1} H_{n-1} - F_{n-1} H_{n+1}), \quad (7c)$$

where (q_n, r_n) is a simple solution pair of the unreduced system (2) and served as nonzero backgrounds. The so-called Hirota bilinear operator (Hirota, 2004) is denoted by D_t and defined as

$$D_t^m G \cdot F = (\partial_t - \partial_{t'})^m G(t) F(t')|_{t'=t}. \quad (8)$$

Note that when $q_n = r_n = 0$, this bilinearized system degenerates to that with zero background, which has been shown in Feng et al. (2020).

Following Deng et al. (2024), the quasi double Casoratians are constructed as follows:

$$F(A, \Phi_n, \Psi_n) = \alpha_n^{(m+1)/2} |\Phi_{n+1}, A^2 \Phi_{n+1}, \dots, A^{2m} \Phi_{n+1}; \Psi_n, A^2 \Psi_n, \dots, A^{2m} \Psi_n|, \quad (9a)$$

$$G(A, \Phi_n, \Psi_n) = \alpha_n^{(m+1)/2} |\Phi_n, A \Phi_{n+1}, A^3 \Phi_{n+1}, \dots, A^{2m+1} \Phi_{n+1}; A \Psi_n, A^3 \Psi_n, \dots, A^{2m-1} \Psi_n|, \quad (9b)$$

$$H(A, \Phi_n, \Psi_n) = \alpha_n^{(m+1)/2} |A \Phi_{n+1}, A^3 \Phi_{n+1}, \dots, A^{2m-1} \Phi_{n+1}; \Psi_{n+1}, A \Psi_n, A^3 \Psi_n, \dots, A^{2m+1} \Psi_n|. \quad (9c)$$

Here, $\alpha_n = 1 - q_n r_n$, the component entries Φ_n and Ψ_n are $(2m+2)$ -th order column vectors and given by matrix equations

$$\begin{bmatrix} \Phi_{n+1} \\ \Psi_{n+1} \end{bmatrix} = M_n \begin{bmatrix} \Phi_n \\ \Psi_n \end{bmatrix}, \quad M_n = \alpha_n^{-1/2} \begin{bmatrix} A & q_n I_{2m+2} \\ r_n I_{2m+2} & A^{-1} \end{bmatrix}, \quad (10a)$$

$$2\partial_t \begin{bmatrix} \Phi_n \\ \Psi_n \end{bmatrix} = N_n \begin{bmatrix} \Phi_n \\ \Psi_n \end{bmatrix}, \quad N_n = \begin{bmatrix} A^2 - A^{-2} + (r_n q_{n-1} - q_n r_{n-1}) I_{2m+2} & 2A q_n + 2A^{-1} q_{n-1} \\ 2A r_{n-1} + 2A^{-1} r_n & A^{-2} - A^2 - (r_n q_{n-1} - q_n r_{n-1}) I_{2m+2} \end{bmatrix}, \quad (10b)$$

in which $A \in \mathbb{C}_{(2m+2) \times (2m+2)}$, $|A| \neq 0$ and I_μ is a μ -th order identity matrix.

Utilizing these Casoratians, the solutions to the bilinearized form (7) can be derived, which yields the following theorem.

Theorem 1. *The bilinear system (7) has double Casoratian solutions*

$$F_n = F(A, \Phi_n, \Psi_n), \quad G_n = G(A, \Phi_n, \Psi_n), \quad H_n = H(A, \Phi_n, \Psi_n), \quad (11)$$

where their vector entries Φ_n and Ψ_n satisfy matrix equations (10) with the given matrix A . These solutions also give the Casoratian solutions Q_n and R_n to the unreduced system (2) through variable transformation (6). Matrix A together with its associated vector entries and its similar matrices together with their associated vector entries lead to the same solutions Q_n and R_n .

A brief proof can be find in Appendix A.

4. Reductions

In this section, the classical and nonlocal reductions to the Casoratian solutions given by theorem 1 are presented. The constraints between the matrices $A = e^B$ and T induced by associated reductions are discussed, followed by a detail construction of the component vectors Φ_n and Ψ_n .

4.1. Reductions of the double Casoratians

In what follows, the detail reduction procedures to equation (5a) are presented.

Equation (5a) is reduced from the unreduced system (2) by requiring $R_n = \delta Q_n$, hence the nonzero backgrounds are assumed to satisfy $r_n = \delta q_n$. By imposing constraint

$$\Psi_n = T \Phi_n, \quad (12)$$

and assuming

$$A^{-1}T = TA, \quad TT = \delta I_{2m+2}, \quad (13)$$

one can verify that the matrix system reduces to the following system

$$\Phi_{n+1} = \alpha_n^{-1/2} A \Phi_n + \alpha_n^{-1/2} q_n T \Phi_n, \quad (14a)$$

$$2\partial_t \Phi_n = (A^2 - A^{-2})\Phi_n + 2(Aq_n + A^{-1}q_{n-1})T\Phi_n \quad (14b)$$

with $\alpha_n = 1 - \delta q_n^2$. If introducing B such that $A = e^B$, the assumption (13) becomes

$$-BT = TB, \quad TT = \delta I_{2m+2}. \quad (15)$$

Using matrix equation (10a) and constraint (12), the double Casoratian F_n can be written as

$$F_n = |A\Phi_n, \dots, A^{2m+1}\Phi_n; \Psi_n, \dots, A^{2m}\Psi_n| = |A\Phi_n, \dots, A^{2m+1}\Phi_n; T\Phi_n, \dots, A^{2m}T\Phi_n|. \quad (16)$$

With constraints (12) and (13), one can see that

$$F_n = |A^{2m+1}T|^{-1} |A^{2m}T\Phi_n, \dots, T\Phi_n; \delta A^{2m+1}\Phi_n, \dots, \delta A\Phi_n| = (-\delta)^{m+1} |A^{2m+1}T|^{-1} F_n. \quad (17)$$

Similarly, we can also derive that $H_n = -(-\delta)^m |A^{2m+1}T|^{-1} G_n$. They give the relation between Q_n and R_n by

$$R_n = \frac{H_n}{F_n} = \frac{-(-\delta)^m |A^{2m+1}T|^{-1} G_n}{(-\delta)^{m+1} |A^{2m+1}T|^{-1} F_n} = \delta G_n / F_n = \delta Q_n, \quad (18)$$

which implies that Q_n is a solution of the sd-mKdV equation (5a).

This reduction procedure can also be applied to equations (5b), (5c) and (5d). We skip the details and list main results in Table 1. Note that in this table, vector Φ_n for (5b), (5c) and (5d) is determined by

$$\Phi_{n+1} = \alpha_n^{-1/2} A \Phi_n + \alpha_n^{-1/2} q_n T \Phi_n^*, \quad (19a)$$

$$2\partial_t \Phi_n = (A^2 - A^{-2} - \delta q_n q_{n-1}^* + \delta q_{n-1} q_n^*) \Phi_n + 2(Aq_n + A^{-1}q_{n-1})T\Phi_n^* \quad (19b)$$

with $\alpha_n = 1 - \delta |q_n|^2$,

$$\Phi_{n+1} = \alpha_n^{-1/2} A \Phi_n + \alpha_n^{-1/2} q_n T \Phi_{1-n}(-t), \quad (20a)$$

$$2\partial_t \Phi_n = (A^2 - A^{-2} - \delta q_n q_{1-n}(-t) + \delta q_{n-1} q_{-n}(-t)) \Phi_n + 2(Aq_n + A^{-1}q_{n-1})T\Phi_{1-n}(-t) \quad (20b)$$

with $\alpha_n = 1 - \delta q_n q_{-n}(-t)$ and

$$\Phi_{n+1} = \alpha_n^{-1/2} A \Phi_n + \alpha_n^{-1/2} q_n T \Phi_{1-n}^*(-t), \quad (21a)$$

$$2\partial_t \Phi_n = (A^2 - A^{-2} - \delta q_n q_{1-n}^*(-t) + \delta q_{n-1} q_{-n}^*(-t)) \Phi_n + 2(Aq_n + A^{-1}q_{n-1})T\Phi_{1-n}^*(-t) \quad (21b)$$

with $\alpha_n = 1 - \delta q_n q_{-n}^*(-t)$, respectively.

The above results are summarized as the following theorem.

Theorem 2. *Solutions of the four sd-mKdV equations in (5) are given by variable transformation*

$$Q_n = \frac{G_n}{F_n}, \quad (22)$$

in which G_n and F_n are double Casoratians defined in formulae (11) with their vector entries Φ_n and Ψ_n given by Table 1.

Remark 1. By defining $\hat{T} = \gamma A^\mu T$, the constraint (12) becomes $\gamma A^\mu \Psi_n = \hat{T} \Phi_n$, the assumption (13) yields $A^{-1} \hat{T} = \hat{T} A$ and $\hat{T} \hat{T} = \gamma^2 \delta I_{2m+2}$ and the Casoratians change accordingly, where μ is an arbitrary integer and γ is an arbitrary nonzero complex number. Similar discussions can be applied to other equations in (5), which allow us to obtain Casoratians presented in Feng et al. (2020) when requiring $q_n = r_n = 0$ and choosing γ and μ properly.

Table 1
Reduction constraints for equations (5)

equation	(q_n, r_n)	constraints	F_n, G_n, H_n	Φ_n
(5a)	$r_n = \delta q_n$	$\Psi_n = T\Phi_n$ $A^{-1}T = TA, TT = \delta I_{2m+2}$ $-BT = TB, TT = \delta I_{2m+2}$	$F_n = (-\delta)^{m+1} A^{2m+1}T ^{-1}F_n$ $H_n = -(-\delta)^m A^{2m+1}T ^{-1}G_n$	(14)
(5b)	$r_n = \delta q_n^*$	$\Psi_n = T\Phi_n^*$ $A^{-1}T = TA^*, TT^* = \delta I_{2m+2}$ $-BT = TB^*, TT^* = \delta I_{2m+2}$	$F_n = (-\delta)^{m+1} A^{2m+1}T ^{-1}F_n^*$ $H_n = -(-\delta)^m A^{2m+1}T ^{-1}G_n^*$	(19)
(5c)	$r_n = \delta q_{-n}(-t)$	$\Psi_n = T\Phi_{1-n}(-t)$ $AT = TA, TT = -\delta I_{2m+2}$ $BT = TB, TT = -\delta I_{2m+2}$	$F_n = \delta^{m+1} T ^{-1}F_{-n}(-t)$ $H_n = \delta^m T ^{-1}G_{-n}(-t)$	(20)
(5d)	$r_n = \delta q_{-n}^*(-t)$	$\Psi_n = T\Phi_{1-n}^*(-t)$ $AT = TA^*, TT^* = -\delta I_{2m+2}$ $BT = TB^*, TT^* = -\delta I_{2m+2}$	$F_n = \delta^{m+1} T ^{-1}F_{-n}^*(-t)$ $H_n = \delta^m T ^{-1}G_{-n}^*(-t)$	(21)

Table 2
Matrices B and T in constraints for equations (5)

equation	δ	T	B
(5a)	δ	$T_1 = T_4 = \mathbf{0}_{m+1}, T_2 = \delta T_3 = \mathbf{I}_{m+1}$	$K_1 = \mathbf{K}_{m+1}, K_4 = -\mathbf{K}_{m+1}$
(5b)	δ	$T_1 = T_4 = \mathbf{0}_{m+1}, T_2 = \delta T_3 = \mathbf{I}_{m+1}$	$K_1 = \mathbf{K}_{m+1}, K_4 = -\mathbf{K}_{m+1}^*$
(5c)	1	$T_1 = -T_4 = i\mathbf{I}_{m+1}, T_2 = T_3 = \mathbf{0}_{m+1}$	$K_1 = \mathbf{K}_{m+1}, K_4 = \mathbf{H}_{m+1}$
	-1	$T_1 = -T_4 = \mathbf{I}_{m+1}, T_2 = T_3 = \mathbf{0}_{m+1}$	
(5d)	δ	$T_1 = T_4 = \mathbf{0}_{m+1}, T_2 = -\delta T_3 = \mathbf{I}_{m+1}$	$K_1 = \mathbf{K}_{m+1}, K_4 = \mathbf{K}_{m+1}^*$

4.2. Matrices B and T

This subsection discusses the explicit forms of B and T in theorem 2. Assuming matrices B and T admit the following specific form

$$B = \begin{bmatrix} K_1 & \mathbf{0} \\ \mathbf{0} & K_4 \end{bmatrix}, \quad T = \begin{bmatrix} T_1 & T_2 \\ T_3 & T_4 \end{bmatrix}, \quad (23)$$

where T_μ and K_μ are $(m+1) \times (m+1)$ matrices, some solutions to the constraints given by Table 1 are presented in Table 2.

Additionally, the constraint for equation (5d) with $\delta = 1$ admits a real solution of form (23) with

$$K_1 = \mathbf{K}_{m+1}, \quad K_4 = \mathbf{H}_{m+1}, \quad \mathbf{K}_{m+1}, \mathbf{H}_{m+1} \in \mathbb{R}_{(m+1) \times (m+1)}, \quad (24a)$$

$$T_1 = \pm T_4 = \mathbf{I}_{m+1}, \quad T_2 = T_3 = \mathbf{0}_{m+1}. \quad (24b)$$

Using theorem 1 and $A = e^B$, it is easy to see that matrix B and its similar matrices can lead to the same solution Q_n of the reduced equations (5), which allow us to further restrict the discussions of matrix B to its canonical form, namely matrix \mathbf{K}_{m+1} is assumed to be

$$\mathbf{K}_{m+1} = \text{Diag}(J_{\mu_1}(k_1), J_{\mu_2}(k_2), \dots, J_{\mu_s}(k_s)), \quad (25)$$

where $J_\mu(k)$ is Jordan block defined by

$$J_h(k) = \begin{bmatrix} k & 0 & 0 & \dots & 0 & 0 \\ 1 & k & 0 & \dots & 0 & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 0 & \dots & 1 & k \end{bmatrix}_{\mu \times \mu}, \quad (26)$$

with $\sum_{j=1}^s \mu_j = m + 1$.

There are two extreme cases of the canonical forms gained much attention. The first case assumes that all $\mu_j = 1$, which results to following diagonal matrix

$$\mathbf{K}_{m+1} = \text{Diag}(k_1, k_2, \dots, k_{m+1}), \quad (27)$$

while the second case assumes that $\mu_1 = m + 1$, resulting to the following Jordan matrix

$$\mathbf{K}_{m+1} = J_{m+1}(k_1). \quad (28)$$

4.3. Vectors Φ_n and Ψ_n

To construct component vectors Φ_n and Ψ_n , the following system

$$\begin{bmatrix} \phi_{n+1} \\ \psi_{n+1} \end{bmatrix} = \alpha_n^{-1/2} \begin{bmatrix} e^k & q_n \\ r_n & e^{-k} \end{bmatrix} \begin{bmatrix} \phi_n \\ \psi_n \end{bmatrix}, \quad (29a)$$

$$2 \begin{bmatrix} \phi_n \\ \psi_n \end{bmatrix}_t = \begin{bmatrix} e^{2k} - e^{-2k} + (r_n q_{n-1} - q_n r_{n-1}) & 2e^k q_n + 2e^{-k} q_{n-1} \\ 2e^k r_{n-1} + 2e^{-k} r_n & e^{-2k} - e^{2k} - (r_n q_{n-1} - q_n r_{n-1}) \end{bmatrix} \begin{bmatrix} \phi_n \\ \psi_n \end{bmatrix}, \quad (29b)$$

which can be seen as a scalar version of the matrix system (10), is considered.

Assume that the nonzero backgrounds are given by

$$q_n = a_0, \quad r_n = b_0, \quad (30)$$

then the system (29) admits the following solution pair

$$\phi_n(k, c, d) = ce^{\lambda n + \eta t} + de^{-(\lambda n + \eta t)}, \quad (31a)$$

$$\psi_n(k, c, d) = -c\xi(k)e^{\lambda n + \eta t} + d\xi(-k)e^{-(\lambda n + \eta t)}, \quad (31b)$$

with

$$e^\lambda = e^{\lambda(k)} = \frac{e^k + e^{-k} + \sqrt{(e^k - e^{-k})^2 + 4a_0 b_0}}{2\sqrt{1 - a_0 b_0}}, \quad (31c)$$

$$\eta = \eta(k) = \frac{1}{2}(e^k + e^{-k})\sqrt{(e^k - e^{-k})^2 + 4a_0 b_0}, \quad (31d)$$

$$\xi = \xi(k) = \frac{e^k - e^{-k} - \sqrt{(e^k - e^{-k})^2 + 4a_0 b_0}}{2a_0}. \quad (31e)$$

Here c and d are constants (or functions of k). To simplify discussions, assumptions $b_0 = \delta a_0$ and $a_0 \in \mathbb{R}$ are taken.

In what follows, the construction of vectors Φ_n and Ψ_n using the above solution pair $\phi_n(k, c, d)$ and $\psi_n(k, c, d)$ is shown.

4.3.1. The unreduced case

Defining

$$\Phi_n = (\phi_n(k_1, c_1, d_1), \phi_n(k_2, c_2, d_2), \dots, \phi_n(k_{2m+2}, c_{2m+2}, d_{2m+2}))^T, \quad (32a)$$

$$\Psi_n = (\psi_n(k_1, c_1, d_1), \psi_n(k_2, c_2, d_2), \dots, \psi_n(k_{2m+2}, c_{2m+2}, d_{2m+2}))^T, \quad (32b)$$

the resulting double Casoratians provide solutions of the unreduced system (2) via the transform (6) with regards to matrix $B = \text{Diag}(k_1, k_2, \dots, k_{2m+2})$.

While defining

$$\Phi_n = (\phi_n(k_1, c_1, d_1), \frac{\partial_{k_1}}{1!} \phi_n(k_1, c_1, d_1), \dots, \frac{\partial_{k_1}^{2m+1}}{(2m+1)!} \phi_n(k_1, c_1, d_1))^T, \quad (33a)$$

$$\Psi_n = (\psi_n(k_1, c_1, d_1), \frac{\partial_{k_1}}{1!} \psi_n(k_1, c_1, d_1), \dots, \frac{\partial_{k_1}^{2m+1}}{(2m+1)!} \psi_n(k_1, c_1, d_1))^T, \quad (33b)$$

the resulting double Casoratians give solutions corresponding to matrix $B = J_{2m+2}(k_1)$.

Table 3

Vectors Φ_n and Ψ_n when matrix T is block skew diagonal

equation	Φ_n	Ψ_n
(5a)	$(\Phi_n^+, \Psi_n^+)^T$	$(\Psi_n^+, \delta\Phi_n^+)^T$
(5b)	$(\Phi_n^+, \Psi_n^{+*})^T$	$(\Psi_n^+, \delta\Phi_n^{+*})^T$
(5d)	$(\Phi_n^+, \Psi_{1-n}^{+*}(-t))^T$	$(\Psi_n^+, -\delta\Phi_{1-n}^{+*}(-t))^T$

4.3.2. The reduced cases

To discuss the reduced equations (5), we define

$$\Phi_n = (\Phi_n^+, \Phi_n^-)^T, \quad \Phi_n^\pm = (\Phi_{1,n}^\pm, \dots, \Phi_{m+1,n}^\pm)^T, \quad (34a)$$

$$\Psi_n = (\Psi_n^+, \Psi_n^-)^T, \quad \Psi_n^\pm = (\Psi_{1,n}^\pm, \dots, \Psi_{m+1,n}^\pm)^T, \quad (34b)$$

with $\Phi_{j,n}^\pm$ and $\Psi_{j,n}^\pm$ being scalar functions.

For equations (5a), (5b) and (5d), the matrix T in constraints is block skew diagonal, which allows us to express Φ_n and Ψ_n with Φ_n^+ and Ψ_n^+ . The detail expressions are listed in Table 3. In this scenario, vectors Φ_n^+ and Ψ_n^+ are given by

$$\Phi_{j,n}^+ = \phi_n(k_j^+, c_j^+, d_j^+), \quad \Psi_{j,n}^+ = \psi_n(k_j^+, c_j^+, d_j^+), \quad j = 1, 2, \dots, m+1, \quad (35)$$

when $\mathbf{K}_{m+1} = \text{Diag}(k_1^+, k_2^+, \dots, k_{m+1}^+)$ is a diagonal matrix, or are given by

$$\Phi_{j,n}^+ = \frac{\partial_{k_1^+}^{j-1} \phi(k_1^+, c_1^+, d_1^+)}{(j-1)!}, \quad \Psi_{j,n}^+ = \frac{\partial_{k_1^+}^{j-1} \psi(k_1^+, c_1^+, d_1^+)}{(j-1)!}, \quad j = 1, 2, \dots, m+1, \quad (36)$$

when $\mathbf{K}_{m+1} = J_{m+1}(k_1^+)$ is a Jordan matrix.

For equation (5c) and the special reduction (24), the matrix T in constraints is block diagonal, which enable us to express Ψ_n in terms of Φ_n with appropriate restrictions between the parameters c and d . The detail expressions and restrictions are listed in Table 4. In this scenario, vector Φ_n is given by

$$\Phi_{j,n}^+ = \phi_n(k_j^+, c_j^+, d_j^+), \quad \Phi_{j,n}^- = \psi_n(h_j^-, c_j^-, d_j^-), \quad j = 1, 2, \dots, m+1, \quad (37)$$

when both $\mathbf{K}_{m+1} = \text{Diag}(k_1^+, k_2^+, \dots, k_{m+1}^+)$ and $\mathbf{H}_{m+1} = \text{Diag}(h_1^-, h_2^-, \dots, h_{m+1}^-)$ are diagonal, or is given by

$$\Phi_{j,n}^+ = \frac{\partial_{k_1^+}^{j-1} \phi(k_1^+, c_1^+, d_1^+)}{(j-1)!}, \quad \Phi_{j,n}^- = \frac{\partial_{h_1^-}^{j-1} \psi(h_1^-, c_1^-, d_1^-)}{(j-1)!}, \quad j = 1, 2, \dots, m+1, \quad (38)$$

when both $\mathbf{K}_{m+1} = J_{m+1}(k_1^+)$ and $\mathbf{H}_{m+1} = J_{m+1}(h_1^-)$ are Jordan matrices.

5. Dynamics

This section devotes to demonstrating dynamics such as solitons, periodic waves and rational solutions.

5.1. The focusing sd-mKdV equation

5.1.1. solitonic and periodic solutions

When $m = 0$ and $\mathbf{K}_1 = k_1$, the first order solution $Q_n = G_n/F_n$ for the focusing sd-mKdV equation (5a) ($\delta = -1$) can be presented by Casoratians

$$F_n = -\sqrt{1+a_0^2} \left[A_1 e^{2\lambda_1 n + 2\eta_1 t} + B_1 e^{-2\lambda_1 n - 2\eta_1 t} + C_1 \right], \quad (39a)$$

Table 4
Vectors Φ_n and Ψ_n when matrix T is block diagonal

equation	δ	Φ_n	Ψ_n	restriction between c and d
(5c)	1	$(\Phi_n^+, \Phi_n^-)^T$	$(i\Phi_{1-n}^+(-t), -i\Phi_{1-n}^-(-t))^T$	$d_j^+ = i\xi(k_j^+)e^{\lambda(k_j^+)}c_j^+$ $d_j^- = -i\xi(h_j^-)e^{\lambda(h_j^-)}c_j^-$
	-1	$(\Phi_n^+, \Phi_n^-)^T$	$(\Phi_{1-n}^+(-t), -\Phi_{1-n}^-(-t))^T$	$d_j^+ = -\xi(k_j^+)e^{\lambda(k_j^+)}c_j^+$ $d_j^- = \xi(h_j^-)e^{\lambda(h_j^-)}c_j^-$
(24) for (5d)	1	$(\Phi_n^+, \Phi_n^-)^T$	$(\Phi_{1-n}^{+*}(-t), \pm\Phi_{1-n}^{-*}(-t))^T$	$d_j^+ = -\xi(k_j^+)e^{\lambda(k_j^+)}c_j^+$ $d_j^- = \mp\xi(h_j^-)e^{\lambda(h_j^-)}c_j^-$

$$G_n = \sqrt{1 + a_0^2} \left[A_2 e^{2\lambda_1 n + 2\eta_1 t} + B_2 e^{-2\lambda_1 n - 2\eta_1 t} + C_2 \right], \quad (39b)$$

with

$$\lambda_1 = \lambda(k_1), \quad \eta_1 = \eta(k_1), \quad (39c)$$

$$A_1 = c_1^2 e^{\lambda_1} (1 + \xi(k_1)\xi(k_1)), \quad A_2 = c_1^2 e^{\lambda_1} \xi(k_1)(e^{k_1} - e^{-k_1}), \quad (39d)$$

$$B_1 = d_1^2 e^{-\lambda_1} (1 + \xi(-k_1)\xi(-k_1)), \quad B_2 = d_1^2 e^{-\lambda_1} \xi(-k_1)(e^{-k_1} - e^{k_1}), \quad (39e)$$

$$C_1 = 2c_1 d_1 (e^{\lambda_1} + e^{-\lambda_1}), \quad C_2 = c_1 d_1 (e^{-\lambda_1 - k_1} \xi(-k_1) - e^{\lambda_1 - k_1} \xi(k_1) - e^{\lambda_1 + k_1} \xi(-k_1) + e^{-\lambda_1 + k_1} \xi(k_1)). \quad (39f)$$

Here, $\lambda(k)$, $\eta(k)$ and $\xi(k)$ are defined in (31c)-(31e) with $b_0 = -a_0$. It is easy to see that the solution dynamics are controlled by $X_1 = \lambda_1 n + \eta_1 t$. We are interested in two special cases.

The first case is that

$$k_1, c_1, d_1 \in \mathbb{R}, \quad (e^{k_1} - e^{-k_1})^2 > 4a_0^2, \quad (40)$$

which yields real λ_1 , η_1 and Q_n . For a fixed t , we have

$$\lim_{\lambda_1 n \rightarrow \pm\infty} Q_n = a_0, \quad (41)$$

which tells it lives on a nonzero background given by a_0 . Further analyzing the possible maximum (minimum), one can obtain soliton (anti-soliton) characterized by its maximum (minimum) traveling along $X_1 = \frac{1}{2} \ln \left| \frac{d_1}{c_1 \xi(k_1) e^{\lambda_1}} \right|$ with constant traveling speed η_1 / λ_1 . A soliton is shown Fig. 1(a) and 1(b)¹, while an anti-soliton is shown in Fig. 1(c).

The second case is when

$$k_1 \in \mathbb{R}, \quad (e^{k_1} - e^{-k_1})^2 < 4a_0^2, \quad (42)$$

or

$$k_1 \in i\mathbb{R}, \quad (43)$$

which yields that both λ_1 and η_1 are purely imaginary. In this scenario, the solution Q_n is doubly periodic with period $T_1 = |2\pi/\lambda_1|$ in n -direction² and $T_2 = |2\pi/\eta_1|$ in t -direction. An example is shown in Fig. 2.

When $m = 1$ and $\mathbf{K}_2 = \text{diag}(k_1, k_2)$ is diagonal, the second order solution $Q_n = G_n/F_n$ can be investigated by using the following Casoratians

$$F_n = \alpha_n |\Phi_{n+1}, A^2 \Phi_{n+1}, T \Phi_n, A^2 T \Phi_n|, \quad G_n = \alpha_n |\Phi_n, A \Phi_{n+1}, A^3 \Phi_{n+1}, AT \Phi_n|, \quad (44)$$

with

$$\Phi_n = (\phi_n(k_1^+, c_1^+, d_1^+), \phi_n(k_2^+, c_2^+, d_2^+), \psi_n(k_1^+, c_1^+, d_1^+), \psi_n(k_2^+, c_2^+, d_2^+))^T. \quad (45)$$




Fig. 1: Shape and motion of first order solutions (Q_n is presented) for the focusing sd-mKdV equation. (a) and (b) A soliton with $a_0 = 0.2$, $k_1 = 0.2$, $c_1 = -d_1 = 1$. (c) An anti-soliton for $a_0 = 0.2$, $k_1 = 0.3$, $c_1 = d_1 = 1$.




Fig. 2: Shape and motion of first order doubly periodic solutions ($|Q_n|^2$ is presented) for the focusing sd-mKdV equation with $a_0 = 0.2$, $k_1 = 0.1$, $c_1 = d_1 = 1$.

It allows us to get insight into interactions, for example, between two solitons, between soliton and waves and between two waves. Examples are demonstrated in Fig.3 without giving detail expressions.

When $m = 1$ and $\mathbf{K}_2 = J_2(k_1)$ is Jordan matrix, solution $Q_n = G_n/F_n$ can be investigated by using the Casoratians (44) with components

$$\Phi_n = (\phi_n(k_1^+, c_1^+, d_1^+), \partial_{k_1} \phi_n(k_1^+, c_1^+, d_1^+), \psi_n(k_1^+, c_1^+, d_1^+), \partial_{k_1} \psi_n(k_1^+, c_1^+, d_1^+))^T. \quad (46)$$

It allows us to investigate behaviors similar to interaction between two solitons or interaction between soliton and doubly periodic wave. Examples are shown in Fig.4 without giving detail expressions.

5.1.2. Rational solutions

To obtain rational solutions, a new parameter κ is introduced, which satisfies

$$\kappa = \sqrt{(e^k - e^{-k})^2 - 4a_0^2}, \quad e^k = \frac{1}{2} \sqrt{\kappa^2 + 4 + 4a_0^2} + \frac{1}{2} \sqrt{\kappa^2 + 4a_0^2}. \quad (47)$$

The solution pair $\phi_n(k, c, d)$ and $\psi_n(k, c, d)$ given by formula (31) can be rewritten as

$$\hat{\phi}_n(\kappa, c, d) = ce^{\hat{\lambda}n + \hat{\eta}t} + de^{-\hat{\lambda}n - \hat{\eta}t}, \quad (48a)$$

¹We should in principle demonstrate all figures as Fig. 1(a), but in what follows we show them as Fig. 1(b) to visualize the solution dynamics more clearly.

²In principle T is a period only when it is a positive integer due the discretized spatial coordinate n . But we also call it period for convenient.




Fig. 3: Shape and motion of second order solutions (diagonal matrix, $|Q_n|^2$ is presented) for the focusing sd-mKdV equation. (a) Interaction between two solitons with $a_0 = 0.2$, $k_1 = 0.2$, $k_2 = 0.6$, $c_1 = -d_1 = c_2 = d_2 = 1$. (b) Interaction between soliton and doubly periodic wave with $a_0 = 0.2$, $k_1 = 0.1$, $k_2 = 0.5$, $c_1 = -d_1 = c_2 = d_2 = 1$. (c) Interaction between two doubly periodic waves with $a_0 = 0.7$, $k_1 = 0.4$, $k_2 = 0.5$, $c_1 = -d_1 = c_2 = d_2 = 1$.




Fig. 4: Shape and motion of second order solutions (Jordan matrix, $|Q_n|^2$ is presented) for the focusing sd-mKdV equation. (a) soliton-soliton-like Interaction with $a_0 = 0.2$, $k_1 = 0.5$, $c_1 = d_1 = 1$. (b) soliton-wave-like interaction with $a_0 = 0.4$, $k_1 = 0.2$, $c_1 = d_1 = 1$.

$$\hat{\psi}_n(\kappa, c, d) = -c\hat{\xi}(\kappa)e^{\hat{\lambda}n+\hat{\eta}t} - d\hat{\xi}(-\kappa)e^{-\hat{\lambda}n-\hat{\eta}t}, \quad (48b)$$

with

$$e^{\hat{\lambda}} = \frac{\kappa + \sqrt{\kappa^2 + 4 + 4a_0^2}}{\sqrt{4 + 4a_0^2}}, \quad \hat{\eta} = \frac{1}{2}\kappa\sqrt{\kappa^2 + 4 + 4a_0^2}, \quad \hat{\xi}(\kappa) = \frac{\sqrt{\kappa^2 + 4a_0^2} - \kappa}{2a_0}. \quad (48c)$$

Assume that $d(\kappa) = -c(-\kappa)$. Taking a formal Taylor expansion of c and d results to

$$c = \sum_{j=0}^{\infty} c^{(j)}\kappa^j, \quad d = \sum_{j=0}^{\infty} -c^{(j)}(-\kappa)^j. \quad (49)$$

It is easy to verify that functions $\hat{\phi}_n(\kappa, c, d)$ and $\hat{\psi}_n(\kappa, c, d)$ can be formally expanded into Taylor series with only odd order terms of the parameter κ , i.e.,

$$\hat{\phi}_n(\kappa, c, d) = \sum_{j=0}^{\infty} \phi_n^{(2j+1)} \kappa^{2j+1}, \quad \hat{\psi}_n(\kappa, c, d) = \sum_{j=0}^{\infty} \psi_n^{(2j+1)} \kappa^{2j+1}. \quad (50)$$

Redefining the component vectors with

$$\Phi_n = (\phi_n^{(1)}, \phi_n^{(3)}, \dots, \phi_n^{(2m+1)}, \psi_n^{(1)}, \psi_n^{(3)}, \dots, \psi_n^{(2m+1)})^T, \quad (51a)$$

$$\Psi_n = (\psi_n^{(1)}, \psi_n^{(3)}, \dots, \psi_n^{(2m+1)}, -\phi_n^{(1)}, -\phi_n^{(3)}, \dots, -\phi_n^{(2m+1)})^T, \quad (51b)$$

and reconstructing matrix $A = \text{Diag}(e^{K_{m+1}}, e^{-K_{m+1}})$ by using

$$e^{K_{m+1}} = \begin{bmatrix} \zeta_0 & 0 & 0 & \cdots & 0 \\ \zeta_2 & \zeta_0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \zeta_{2m} & \zeta_{2m-2} & \zeta_{2m-4} & \cdots & \zeta_0 \end{bmatrix}, \quad \text{with } \zeta_{2j} = \frac{1}{(2j)!} \partial_{\kappa}^{2j} e^k |_{\kappa=0}, \quad (52)$$

the Casoratians G_n and F_n defined by (11) yeild rational solution by variable transformation $Q_n = G_n/F_n$.

When $m = 0$, the first order rational solution is explicitly presented by

$$Q_n = -a_0 \left(1 + \frac{U_n}{W_n} \right) \quad (53)$$

with

$$W_n = \left[(n + 2(1 + a_0^2)t + 2\frac{c_1}{c_0} \sqrt{1 + a_0^2})^2 + (n + 2(1 + a_0^2)t + 2\frac{c_1}{c_0} \sqrt{1 + a_0^2} - \frac{\sqrt{1 + a_0^2}}{a_0})^2 + \frac{\sqrt{1 + a_0^2}}{a_0} - 1 \right], \quad (54)$$

$$U_n = \frac{1 + a_0^2 + \sqrt{1 + a_0^2}}{a_0}, \quad (55)$$

an example of which is shown in Fig. 5(a). A second order rational solution is demonstrated in Fig.5(b) without giving detail expressions.




Fig. 5: Shape and motion of rational solutions ($|Q_n|^2$ is presented) for the focusing sd-mKdV equation. (a) A first order rational solution with $a_0 = 0.5$ and $c = 1$. (b) A second order rational solution with $a_0 = 0.5$ and $c = 1$.

5.2. The defocusing reverse-space-time complex sd-mKdV equation

5.2.1. solitonic and periodic solutions

When $m = 0$ and $\mathbf{K}_1 = k_1$, the solution $Q_n = G_n/F_n$ of the defocusing reverse-space-time complex sd-mKdV equation (5d) ($\delta = 1$) is determined by Casoratians

$$F_n = -\sqrt{1 - a_0^2} \left[A_1 e^{2(a_1 n + a_2 t)} + B_1 e^{-2(a_1 n + a_2 t)} + C_1 e^{2i(b_1 n + b_2 t)} + D_1 e^{-2i(b_1 n + b_2 t)} \right], \quad (56a)$$

$$G_n = \sqrt{1 - a_0^2} \left[A_2 e^{2(a_1 n + a_2 t)} + B_2 e^{-2(a_1 n + a_2 t)} + C_2 e^{2i(b_1 n + b_2 t)} + D_2 e^{-2i(b_1 n + b_2 t)} \right], \quad (56b)$$

with

$$A_1 = c_1 d_1^* (e^{\lambda_1 - \lambda_1^*} - \xi(k_1) \xi^*(-k_1)), \quad A_2 = c_1 d_1^* \xi^*(-k_1) (e^{k_1^*} - e^{k_1} e^{\lambda_1 - \lambda_1^*}), \quad (56c)$$

$$B_1 = c_1^* d_1 (e^{\lambda_1^* - \lambda_1} - \xi^*(k_1) \xi(-k_1)), \quad B_2 = -c_1^* d_1 \xi^*(k_1) (e^{k_1^*} - e^{k_1} e^{\lambda_1^* - \lambda_1}), \quad (56d)$$

$$C_1 = c_1 c_1^* (e^{\lambda_1 + \lambda_1^*} + \xi(k_1) \xi^*(k_1)), \quad C_2 = -c_1 c_1^* \xi^*(k_1) (e^{k_1^*} - e^{k_1} e^{\lambda_1 + \lambda_1^*}), \quad (56e)$$

$$D_1 = d_1 d_1^* (e^{-\lambda_1 - \lambda_1^*} + \xi(-k_1) \xi^*(-k_1)), \quad D_2 = d_1 d_1^* \xi^*(-k_1) (e^{k_1^*} - e^{k_1} e^{-\lambda_1 - \lambda_1^*}), \quad (56f)$$

$$\lambda_1 = \lambda(k_1) = a_1 + b_1 i, \quad \eta_1 = \eta(k_1) = a_2 + b_2 i. \quad (56g)$$

Here, λ , η and ξ are defined in (31c)-(31e) with $b_0 = a_0$ and $a_0 \in \mathbb{R}$. Similar to the classical situations, this formula allows us to easily analyze the obtained solution Q_n and there are also two interesting cases.

The first case is when

$$k_1 \in i\mathbb{R}, \quad 4 > 4a_0^2 > -(e^{k_1} - e^{-k_1})^2, \quad (57)$$

which yields $b_1 = b_2 = 0$ and $|\xi(k_1)| = 1$. In this case, the solution is reduced to

$$Q_n = -\frac{A_2 e^{X_1} + B_2 e^{-X_1} + C_2 + D_2}{A_1 e^{X_1} + B_1 e^{-X_1} + C_1 + D_1}, \quad \text{with } X_1 = 2(a_1 n + a_2 t), \quad (58)$$

which is a soliton traveling parallel to $X_1 = 0$ with a constant velocity $-a_2/a_1$. For a fixed t , it is easy to see that this solution satisfies

$$\lim_{n \rightarrow \pm\infty} |Q_n|^2 = a_0^2, \quad (59)$$

which means its squared envelope lives on plane characterized by a_0^2 . An example is shown in Fig. 6(a).

The second case is when

$$k_1 \in i\mathbb{R}, \quad 4a_0^2 < -(e^{k_1} - e^{-k_1})^2, \quad a_0^2 < 1, \quad (60)$$

which yields $a_1 = a_2 = 0$. In this scenario, the solution reduces to

$$Q_n = -\frac{A_2 + B_2 + C_2 e^{X_1} + D_2 e^{-X_1}}{A_1 + B_1 + C_1 e^{X_1} + D_1 e^{-X_1}}, \quad \text{with } X_1 = 2i(b_1 n + b_2 t), \quad (61)$$

which is doubly periodic with period $T_1 = \pi/b_1$ in the n -direction and $T_2 = \pi/b_2$ in the t -direction. An example is shown in Fig.6(b).

When $m = 1$ and $\mathbf{K}_2 = \text{diag}(k_1, k_2)$ is diagonal, solution $Q_n = G_n/F_n$ can be investigated by using the following Casoratians

$$F_n = \alpha_n |\Phi_{n+1}, A^2 \Phi_{n+1}, T \Phi_{1-n}^*(-t), A^2 T \Phi_{1-n}^*(-t)|, \quad G_n = \alpha_n |\Phi_n, A \Phi_{n+1}, A^3 \Phi_{n+1}, AT \Phi_{1-n}^*(-t)|, \quad (62a)$$

with

$$\Phi_n = (\phi_n(k_1^+, c_1^+, d_1^+), \phi_n(k_2^+, c_2^+, d_2^+), \psi_{1-n,-t}^*(k_1^+, c_1^+, d_1^+), \psi_{1-n,-t}^*(k_2^+, c_2^+, d_2^+))^T, \quad (62b)$$

which allows us to study interactions between solitons, between periodic wave and soliton and between periodic waves. Examples are demonstrated in Fig.7 without giving detail expressions.




Fig. 6: Shape and motion of first order solutions ($|Q_n|^2$ is presented) for the defocusing reverse-space-time sd-mKdV equation. (a) Soliton with $a_0 = 0.5$, $k_1 = 0.2j$, $c_1 = -d_1 = 1$. (b) Doubly periodic wave with $a_0 = 0.1$, $k_1 = 0.3j$, $c_1 = -d_1 = 1$.




Fig. 7: Shape and motion of second order solutions (diagonal matrix, $|Q_n|^2$ is presented) for the defocusing reverse-space-time sd-mKdV equation. (a) Interaction between two solitons with $a_0 = 0.9$, $k_1 = 0.3j, k_2 = 0.7j$ $c_1 = d_1 = c_2 = d_2 = 1$. (b) Interaction between soliton and periodic wave with $a_0 = 0.6$, $k_1 = 0.3j, k_2 = 0.7j$ $c_1 = d_1 = c_2 = d_2 = 1$. (c) Interaction between two periodic waves with $a_0 = 0.1$, $k_1 = 0.2j, k_2 = 0.6j$ $c_1 = d_1 = c_2 = d_2 = 1$.

When $m = 1$ and $\mathbf{K}_2 = J_2(k_1)$ is a Jordan matrix, solution $Q_n = G_n/F_n$ can be investigated by using the formula (62a) with component vector

$$\Phi_n = (\phi_n(k_1^+, c_1^+, d_1^+), \partial_{k_1} \phi_n(k_1^+, c_1^+, d_1^+), \psi_{1-n,-i}^*(k_1^+, c_1^+, d_1^+), (\partial_{k_1} \psi_{1-n,-i}(k_1^+, c_1^+, d_1^+))^*)^T, \quad (63)$$

which allows us to study, for example, soliton-like interaction. An example is shown in Fig. 8 without giving detail expressions.

5.2.2. Rational solutions

Similar to the classical scenario discussed in Sec.5.1, rational solutions for equation (5d) ($\delta = 1$) can be investigated by a new parameter κ satisfying

$$\kappa = \sqrt{(e^k - e^{-k})^2 + 4a_0^2}, \quad e^k = \frac{1}{2}\sqrt{\kappa^2 + 4 - 4a_0^2} + \frac{1}{2}\sqrt{\kappa^2 - 4a_0^2}. \quad (64)$$




Fig. 8: Shape and motion of a second order solution (Jordan matrix, $|Q_n|^2$ is presented) for the defocusing reverse-space-time sd-mKdV equation with $a_0 = 0.9$, $k_1 = 0.8j$ $c_1 = d_1 = 1$.

The solution pair $\phi_n(k, c, d)$ and $\psi_n(k, c, d)$ given by formula (31) can be rewritten as

$$\hat{\phi}_n(\kappa, c, d) = ce^{\hat{\lambda}n + \hat{\eta}t} + de^{-\hat{\lambda}n - \hat{\eta}t}, \quad (65a)$$

$$\hat{\psi}_n(\kappa, c, d) = -c\hat{\xi}(\kappa)e^{\hat{\lambda}n + \hat{\eta}t} - d\hat{\xi}(-\kappa)e^{-\hat{\lambda}n - \hat{\eta}t}, \quad (65b)$$

with

$$e^{\hat{\lambda}} = \frac{\kappa + \sqrt{\kappa^2 + 4 - 4a_0^2}}{\sqrt{4 - 4a_0^2}}, \quad \hat{\eta} = \frac{1}{2}\kappa\sqrt{\kappa^2 + 4 - 4a_0^2}, \quad \hat{\xi}(\kappa) = \frac{\sqrt{\kappa^2 - 4a_0^2} - \kappa}{2a_0}. \quad (65c)$$

Assume that $d(\kappa) = -c(-\kappa)$, which allows us to take a formal Taylor expansion of c and d and results to eq. (49). Functions $\hat{\phi}_n(\kappa, c, d)$ and $\hat{\psi}_n(\kappa, c, d)$ can be formally expanded into Taylor series with only odd order terms of the parameter κ , namely,

$$\hat{\phi}_n(\kappa, c, d) = \sum_{j=0}^{\infty} \phi_n^{(2j+1)} \kappa^{2j+1}, \quad \hat{\psi}_n(\kappa, c, d) = \sum_{j=0}^{\infty} \psi_n^{(2j+1)} \kappa^{2j+1}. \quad (66)$$

Redefining the Casoratian vectors with

$$\Phi_n = (\phi_n^{(1)}, \phi_n^{(3)}, \dots, \phi_n^{(2m+1)}, \psi_{1-n, -t}^{(1)*}, \psi_{1-n, -t}^{(3)*}, \dots, \psi_{1-n, -t}^{(2m+1)*})^T, \quad (67a)$$

$$\Psi_n = (\psi_n^{(1)}, \psi_n^{(3)}, \dots, \psi_n^{(2m+1)}, -\phi_{1-n, -t}^{(1)*}, -\phi_{1-n, -t}^{(3)*}, \dots, -\phi_{1-n, -t}^{(2m+1)*})^T, \quad (67b)$$

and reconstructing matrix $A = \text{Diag}(e^{K_{m+1}}, e^{K_{m+1}^*})$ by using

$$e^{K_{m+1}} = \begin{bmatrix} \zeta_0 & 0 & 0 & \cdots & 0 \\ \zeta_2 & \zeta_0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \zeta_{2m} & \zeta_{2m-2} & \zeta_{2m-4} & \cdots & \zeta_0 \end{bmatrix}, \quad \text{with } \zeta_{2j} = \frac{1}{(2j)!} \partial_{\kappa}^{2j} e^k |_{\kappa=0}, \quad (68)$$

the Casoratians G_n and F_n defined by (10) yeild rational solutions by variable transformation $Q_n = G_n/F_n$. Two examples of rational solutions are shown in Fig.9 without giving detail expressions.

6. Conclusions

In this paper, the bilinearization-reduction approach (Chen et al., 2018) is implemented to obtain the exact solutions with nonzero backgrounds for the sd-mKdV equation, the complex sd-mKdV equation, the reverse-space-time sd-mKdV equation and the reverse-space-time complex sd-mKdV equation. The first two equations are obtained with local real and complex reductions, while the later two equations are obtained with nonlocal real and complex reductions.




Fig. 9: Shape and motion of rational solutions ($|Q_n|^2$ is presented) for the defocusing reverse-space-time sd-mKdV equation. (a) First order rational solution with $m = 0$, $a_0 = 0.6$ and $c = 1$. (b) Second order rational solution with $m = 0$, $a_0 = 0.5$ and $c = 1 + 2\kappa + 2i\kappa^2$.

The bilinearization-reduction approach starts from transforming the unreduced system (2) of the coupled sd-mKdV equations into a bilinear system (7) with nonzero backgrounds, utilizing the variable transformation (6). This bilinearization admits quasi double Casoratian solutions (11) with a pair of solvable component vectors Φ_n and Ψ_n given by a matrix system (10). It turns out that these Casoratian solutions are also solutions to the unreduced system (2) through the variable transformation (6). When neutralizing nonzero backgrounds, the bilinearization (7) degenerates to a bilinear formula with zero background, which has been discussed in Feng et al. (2020).

For a plane background $(q_n, r_n) = (a_0, \delta a_0)$ with $a_0 \in \mathbb{R}$, the vectors Φ_n and Ψ_n for the unreduced system (2) and its reductions (5) are explicitly constructed by utilizing scalar functions ϕ_n and ψ_n defined in (31), allowing us to obtain Casoratian solutions. These solutions are characterized by the canonical form of matrix A , since its similar matrices can lead to the same solutions (see Theorem 1). To showcase rich dynamics in these solutions, solitons, periodic waves and rational solutions for the focusing sd-mKdV equation and for the defocusing reverse-space-time complex sd-mKdV equation are demonstrated.

Acknowledgments

This article is supported by the NSFC grant (Nos. 12201580, 12071432), Zhejiang Provincial Natural Science Foundation (No. LZ24A010007) and the China Postdoctoral Science Foundation (2022M722921).

The authors would like to thank Kui Chen of Zhejiang Lab for his helpful comments.

Declaration of Interests. The authors report no conflict of interest.

A. Proof of theorem 1

To prove theorem 1, the identity (Freeman and Nimmo, 1983)

$$|M, a, b||M, c, d| - |M, a, c||M, b, d| + |M, a, d||M, b, c| = 0 \quad (69)$$

is utilized. Here M is an $\alpha \times (\alpha - 2)$ matrix, letters a, b, c and d denote α -th order column vectors.

Introducing the following shorthand notation

$$|(\beta, \beta + 2\mu), (\zeta); (\gamma, \gamma + 2\nu), (\xi)| = |A^\beta \Phi_n, A^{\beta+2} \Phi_n, \dots, A^{\beta+2\mu} \Phi_n, A^\zeta \Phi_n; A^\gamma \Phi_n, A^{\gamma+2} \Phi_n, \dots, A^{\gamma+2\nu} \Phi_n, A^\xi \Psi_n|, \quad (70)$$

with parameters β, γ, ζ and ξ denoting integers and parameters μ and ν denoting positive integers, the Casoratians in Theorem 1 can be rewritten as

$$G_n = |A|^{-1}(g_n + (-1)^m q_n f_n), \quad H_n = |A|^{-1}(h_n + (-1)^m r_n f_n), \quad (71a)$$

$$F_n = |(1, 2m + 1); (0, 2m)|, \quad f_n = |(1, 2m + 1); (2, 2m + 2)|, \quad (71b)$$

$$g_n = |(1, 2m + 3); (2, 2m)|, \quad h_n = |(3, 2m + 1); (0, 2m + 2)|, \quad (71c)$$

in which the matrix equation (10a) has been utilized.

A direct calculation yields

$$\begin{aligned} \alpha_n |A| F_{n+1} &= |(3, 2m + 3); (0, 2m)| - (-1)^m q_n h_n - (-1)^m r_n g_n - q_n r_n f_n, \\ |A| F_{n-1} &= f_n, \\ \alpha_n G_{n+1} &= |(1, 2m + 3); (0, 2m - 2)| + (-1)^m q_n |(1, 2m + 1); (0, 2m - 2), (2m + 2)| \\ &\quad - (-1)^m q_n |(1, 2m - 1), (2m + 3); (0, 2m)| + (-1)^m \alpha_n q_n F_n \\ &\quad + q_n q_n |(1, 2m - 1); (0, 2m + 2)| + (-1)^m \alpha_n q_{n+1} F_n, \\ G_{n-1} &= |(-1, 2m + 1); (2, 2m)| + (-1)^m q_{n-1} F_n, \\ 2|A| \partial_t G_n &= 2q_{n-1} r_n g_n + 2(-1)^m q_{n,t} f_n \\ &\quad + |(1, 2m + 1), (2m + 5); (2, 2m)| + |(1, 2m + 3); (0), (4, 2m)| \\ &\quad - |(-1), (3, 2m + 3); (2, 2m)| - |(1, 2m + 3); (2, 2m - 2), (2m + 2)| \\ &\quad + (-1)^m q_n |(1, 2m + 1); (2, 2m), (2m + 4)| + (-1)^m q_n |(1, 2m + 1); (0), (4, 2m + 2)| \\ &\quad - (-1)^m q_n |(-1), (3, 2m + 1); (2, 2m + 2)| - (-1)^m q_n |(1, 2m - 1), (2m + 3); (2, 2m + 2)| \\ &\quad - 2(-1)^m q_{n-1} |(3, 2m + 3); (0, 2m)| + 2(-1)^m q_{n-1} f_n + 2q_n q_{n-1} h_n, \\ 2\partial_t F_n &= |(1, 2m - 1), (2m + 3); (0, 2m)| + |(1, 2m + 1); (-2), (2, 2m)| \\ &\quad - |(-1), (3, 2m + 1); (0, 2m)| - |(1, 2m + 1); (0, 2m - 2), (2m + 2)| \\ &\quad - 2(-1)^m q_n |(1, 2m - 1); (0, 2m + 2)| - 2(-1)^m r_n |(-1, 2m + 1); (2, 2m)|. \end{aligned}$$

Substituting the above calculation into Eq. (7a) results to

$$\begin{aligned} &|A|^2 (\alpha_n F_{n+1} F_{n-1} + G_n H_n - F_n F_n) \\ &= |(3, 2m + 3); (0, 2m)| f_n + g_n h_n - |(3, 2m + 3); (2, 2m + 2)| F_n, \end{aligned}$$

which vanishes by means of eq.(69).

Similarly, Substituting the above calculation into eq. (7b) becomes

$$2|A| (\alpha_n G_{n+1} F_{n-1} - \alpha_n G_{n-1} F_{n+1} - D_t G_n \cdot F_n) = S_1 + S_2 + (-1)^m q_n (S_3 + S_4),$$

where

$$\begin{aligned} S_1 &= -2|(-1, 2m + 1); (2, 2m)| |(3, 2m + 3); (0, 2m)| - |(1, 2m + 3); (0), (4, 2m)| F_n + |(-1), (3, 2m + 3); (2, 2m)| F_n \\ &\quad + |(1, 2m + 1); (-2), (2, 2m)| g_n - |(-1), (3, 2m + 1); (0, 2m)| g_n, \\ S_2 &= 2|(1, 2m + 3); (0, 2m - 2)| f_n - |(1, 2m + 1), (2m + 5); (2, 2m)| F_n + |(1, 2m + 3); (2, 2m - 2), (2m + 2)| F_n \\ &\quad + |(1, 2m - 1), (2m + 3); (0, 2m)| g_n - |(1, 2m + 1); (0, 2m - 2), (2m + 2)| g_n, \\ S_3 &= 2|(-1, 2m + 1); (2, 2m)| h_n - |(1, 2m + 1); (0), (4, 2m + 2)| F_n + |(-1), (3, 2m + 1); (2, 2m + 2)| F_n \\ &\quad + |(1, 2m + 1); (-2), (2, 2m)| f_n - |(-1), (3, 2m + 1); (0, 2m)| f_n, \\ S_4 &= -2|(1, 2m - 1); (0, 2m + 2)| g_n - |(1, 2m + 1); (2, 2m), (2m + 4)| F_n + |(1, 2m - 1), (2m + 3); (2, 2m + 2)| F_n \\ &\quad + |(1, 2m + 1); (0, 2m - 2), (2m + 2)| f_n - |(1, 2m - 1), (2m + 3); (0, 2m)| f_n. \end{aligned}$$

Each S_j vanishes by using the identity (69) twice, thus eq. (7b) holds. Eq. (7c) can be proved in a similar way.

Suppose matrices A and J are similar through $A = P^{-1} J P$, we can introduce $\Phi'_n = P \Phi_n$ and $\Psi'_n = P \Psi_n$, which again satisfy matrix equations (10) with matrix J replacing matrix A . The double Casoratians satisfy $F(J, \Phi'_n, \Psi'_n) = |P| F(A, \Phi_n, \Psi_n)$, $G(J, \Phi'_n, \Psi'_n) = |P| G(A, \Phi_n, \Psi_n)$ and $H(J, \Phi'_n, \Psi'_n) = |P| H(A, \Phi_n, \Psi_n)$. This imply that matrix A , together with its associated vectors Φ_n and Ψ_n , and its similar matrices, together with their associated vectors, lead to same Q_n and R_n .

References

- Ablowitz, M., Musslimani, Z., 2013. Integrable nonlocal nonlinear Schrödinger equation. *Physical Review Letters* 110, 064105 (5pp).
- Ablowitz, M.J., Kaup, D.J., Newell, A.C., Segur, H., 1973. Nonlinear-evolution equations of physical significance. *Physical Review Letters* 31(2), 125.
- Ablowitz, M.J., Ladik, F., 1976. Nonlinear differential-difference equations and fourier analysis. *Journal of Mathematical Physics* 17(6), 1011–1018.
- Ablowitz, M.J., Musslimani, Z.H., 2017. Integrable nonlocal nonlinear equations. *Studies in Applied Mathematics* 139(1), 7–59.
- Ankiewicz, A., Akhmediev, N., Soto-Crespo, J.M., 2010. Discrete rogue waves of the ablowitz-ladik and hirota equations. *Physical Review E—Statistical, Nonlinear, and Soft Matter Physics* 82(2), 026602.
- Chen, K., Deng, X., Lou, S., Zhang, D.J., 2018. Solutions of nonlocal equations reduced from the AKNS hierarchy. *Studies in Applied Mathematics* 141(1), 113–141.
- Common, A.K., 1993. The half-infinite discretized hirota equation and the trigonometric moment problem. *Inverse Problems* 9(6), 641–648.
- Deng, X., Chen, K., Chen, H.Y., Zhang, D.J., 2024. The integrable semi-discrete nonlinear Schrödinger equations with nonzero backgrounds: Bilinearization-reduction approach. *arXiv Preprint*, 1.
- Deng, X., Lou, S., Zhang, D.J., 2018. Bilinearisation-reduction approach to the nonlocal discrete nonlinear Schrödinger equations. *Applied Mathematics and Computation* 332, 477–483.
- Doliwa, A., Santini, P.M., 1995. Integrable dynamics of a discrete curve and the ablowitz-ladik hierarchy. *Journal of Mathematical Physics* 36(3), 1259–1273.
- Feng, W., Zhao, S.L., Sun, Y.Y., 2020. Double Casoratian solutions to the nonlocal semi-discrete modified Korteweg-de Vries equation. *International Journal of Modern Physics B* 34(05), 2050021.
- Freeman, N.C., Nimmo, J.J.C., 1983. Soliton solutions of the Korteweg de-Vries and the Kadomtsev-Petviashvili equations: the Wronskian technique. *Physics Letters A* 95(1), 1–3.
- Fu, W., Qiao, Z., Sun, J., Zhang, D.J., 2015. Continuous correspondence of conservation laws of the semi-discrete akns system. *Journal of Nonlinear Mathematical Physics* 22(3), 321–341.
- Hirota, R., 1972. Exact solution of the modified Korteweg-de Vries equation for multiple collisions of solitons. *Journal of the Physical Society of Japan* 33(5), 1456–1458.
- Hirota, R., 2004. *The direct method in soliton theory*. Cambridge University Press.
- Hisakado, M., Wadati, M., 1995. Integrable dynamics of discrete surfaces. *Journal of the Physical Society of Japan* 64(11), 4129–4134.
- Kakutani, T., Ono, H., 1969. Weak non-linear hydromagnetic waves in a cold collision-free plasma. *Journal of the Physical Society of Japan* 26(5), 1305–1318.
- Leblond, H., Mihalache, D., 2013. Models of few optical cycle solitons beyond the slowly varying envelope approximation. *Physics Reports* 523(2), 61–126.
- Leblond, H., Sanchez, F., 2003. Models for optical solitons in the two-cycle regime. *Physical Review A* 67(1), 013804.
- Mukaihira, A., Nakamura, Y., 2000. Integrable discretization of the modified kdv equation and applications. *Inverse Problems* 16(2), 413–424.
- Sarma, A.K., Miri, M.A., Musslimani, Z.H., Christodoulides, D.N., 2014. Continuous and discrete Schrödinger systems with PT-symmetric nonlinearities. *Physical Review E* 89, 052918.
- Shi, Y., Shen, S., Zhao, S., 2019. Solutions and connections of nonlocal derivative nonlinear Schrödinger equations. *Nonlinear Dynamics* 95, 1257–1267.
- Sun, Y.Y., Zhao, S.L., 2019. Rational solutions for three semi-discrete modified korteweg–de vries-type equations. *Modern Physics Letters B* 33(32), 1950399.
- Wadati, M., 1973. The modified Korteweg-de Vries equation. *Journal of the Physical Society of Japan* 34(5), 1289–1296.
- Wadati, M., 1976. Transformation theories for nonlinear discrete systems. *Supplement of Progress of Theoretical Physics* 59, 36–63.
- Wang, J., Wu, H., Zhang, D.J., 2020. Solutions of the nonlocal (2+1)-D breaking solitons hierarchy and the negative order AKNS hierarchy. *Communications in Theoretical Physics* 72(4), 045002.
- Wang, Z., Ma, W.X., 2010. Discrete jacobi sub-equation method for nonlinear differential-difference equations. *Mathematical methods in the applied sciences* 33(12), 1463–1472.
- Zhang, D.J., 2023. Bilinearization-reduction approach to integrable systems. *Acta Physica Sinica* 72(10), 100203.
- Zhang, D.J., Liu, S.M., Deng, X., 2023. The solutions of classical and nonlocal nonlinear Schrödinger equations with nonzero backgrounds: Bilinearisation and reduction approach. *Open Communications in Nonlinear Mathematical Physics* 3, 10036.
- Zhang, D.J., Zhao, S.L., Sun, Y.Y., Zhou, J., 2014. Solutions to the modified Korteweg-de Vries equation. *Reviews in Mathematical Physics* 26(07), 1430006.
- Zhang, Y., Jin, L.G., Dong, K.H., 2012. Integrable discretizations and soliton solution of kdv and mkdv equations. *Applied Mathematics and Computation* 218(18), 9555–9562.
- Zhu, J.Y., Geng, X.G., Kuang, Y.H., 2013. Complex solution to the discrete mkdv equation and the riemann-hilbert approach. *arXiv preprint*, arXiv:1310.1666.