# Ocneanu Algebra of Seams: Critical Unitary $E_6$ RSOS Lattice Model

Paul A. Pearce<sup>a,b</sup>, Jørgen Rasmussen<sup>b</sup>

<sup>a</sup>School of Mathematics and Statistics, University of Melbourne Parkville, Victoria 3010, Australia

<sup>b</sup>School of Mathematics and Physics, University of Queensland St Lucia, Brisbane, Queensland 4072, Australia

papearce@unimelb.edu.au j.rasmussen@uq.edu.au

#### Abstract

We consider the A series and exceptional  $E_6$  Restricted Solid-On-Solid lattice models as prototypical examples of the critical Yang-Baxter integrable two-dimensional A-D-E lattice models. We focus on type I theories which are characterized by the existence of an extended chiral symmetry in the continuum scaling limit. Starting with the commuting family of column transfer matrices on the torus, we build matrix representations of the Ocneanu graph fusion algebra as integrable seams for arbitrary finite-size lattices with the structure constants specified by Petkova and Zuber. This commutative seam algebra contains the Verlinde, fused adjacency and graph fusion algebras as subalgebras. Our matrix representation of the Ocneanu algebra encapsulates the quantum symmetry of the commuting family of transfer matrices. In the continuum scaling limit, the integrable seams realize the topological defects of the associated conformal field theory and the known toric matrices encode the twisted conformal partition functions.

# Contents

1	1 Introduction		2							
2	2 Critical RSOS models on the torus		4							
	2.1 Critical $A$ - $D$ - $E$ lattice models		4							
	2.2 Transfer matrices, braid limit and Hamilton	nians	4							
3	Ocneanu fusion algebra									
	3.1 Fusion matrices and structure constants .		6							
	3.2 Twisted conformal partition functions		7							
	3.3 Extended toric matrices and quantum symmetry	netry	7							
	3.4 Ocneanu and double graph fusion algebras		8							
4	A-D-E integrable seams									
	4.1 Construction of <i>s</i> -type seams		9							
	4.2 Construction of seams of type $(a, b)$		10							
	4.3 Ocneanu algebra of seams and quantum syn	nmetry	1							
5	5 Conclusion	1	3							

# 1 Introduction

In this short article, we are concerned with two-dimensional critical Restricted Solid-On-Solid (RSOS) lattice models built on the Dynkin diagrams of simply-laced A-D-E Lie algebras as in Figure 1. These models are Yang-Baxter integrable [1]. The first such models were built on the A-type Dynkin diagrams and solved off-criticality in 1984 by Andrews, Baxter and Forrester [2]. The full family of critical A-D-E lattice models was subsequently introduced and studied by Pasquier [3–5]. In the continuum scaling limit, the thermodynamic behaviour of these models is described by Conformal Field Theory (CFT) [6]. The A series is associated with the conformal unitary minimal models  $\mathcal{M}(g-1,g)$  of Belavin, Polyakov and Zamolodchikov [7]. The unitary A-D-E CFTs are in fact (A, G) coset models [8,6]

$$(A_{g-2},G) = \begin{cases} (A_{g-2},A_{g-1}), & g = 4,5,6,\dots \\ (A_{g-2},D_{(g+2)/2}), & g = 6,8,10,\dots \\ (A_{10},E_6), & g = 12 \\ (A_{16},E_7), & g = 18 \\ (A_{28},E_8), & g = 30 \end{cases}$$
(1.1)

where g is the Coxeter number of G and c is the central charge. These theories are rational [9], admitting a finite number of irreducible representations of the Virasoro algebra which close under fusion.

Much effort within CFT has been focussed on the study of boundary conditions. On the cylinder, the boundary conditions of unitary minimal CFTs and their conformal cylinder partition functions are understood [10,11] in terms of nonnegative integer matrix representations (nimreps) of the Verlinde [12] and graph fusion [4] algebras. The irreducible representations are conjugate to conformal boundary conditions [13] labelled by the coset graph nodes  $(r, a) \in (A, G)$ . Much of this structure is common [14,15] between the finite lattice models and the CFT. On the torus with periodic boundary conditions in both directions, the unitary modular invariant partition functions [16] with c < 1 are exhausted by the A-D-E classification of Cappelli, Itzykson and Zuber [17]. Incorporating the left and right chiral copies of the Virasoro algebra, the conformal partition functions are given as sesquilinear forms in Virasoro characters. Twisted boundary conditions are imposed by inserting topological defect seams into the bulk. The resulting twisted partition functions on the torus are understood in terms of the Ocneanu [18] (double) fusion algebra which describes the quantum symmetry and how the left and right chiral halves of the theory are glued together. Indeed, working within CFT, Petkova and Zuber [19] have obtained explicit expressions for the structure constants of this fusion algebra, the toric matrices and the associated twisted conformal partition functions. The algebraic structure behind these results has been elucidated by Coquereaux and collaborators [20–23] who, following Ocneanu [18], identified the Ocenanu algebra with the quotient algebra  $G \otimes_T G := (G \otimes G)/T$  where G denotes the graph algebra of type A, D, E and where T is the ambichiral subalgebra of G specified in [19]. In particular, the authors of [20, 21] give a detailed account of the prototypical type I example of  $E_6$ .

In parallel to the developments coming from CFT, twisted boundary conditions on the torus have also been studied from a purely lattice perspective [24-26] by inserting integrable seam segments into the commuting *row* transfer matrices. These segments give rise to commuting integrable *vertical* seams on the square lattice which, in the continuum scaling limit, become the topological defects of the associated CFT. In this picture, the fusion of the integrable seams is simply implemented by matrix products of the integrable seams. This picture was implicitly confirmed in [26] by numerical studies of the conformal spectra of the commuting row transfer matrices.

In this article, we develop further the lattice approach to fusion of integrable seams on the lattice. As in [27, 28], we turn our attention to the "cross-channel", but here we use the commuting family of column transfer matrices to build the various types of integrable seams on the torus. Adopting the

Lie Algebra	Graph $G$	g	$\operatorname{Exp}(G)$	$\mathrm{Type}/H$	Г
$A_L$	1 2 3 ··· L * •	L+1	$1, 2, \ldots, L$	Ι	$\mathbb{Z}_2$
$D_L (L \text{ even})$	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	2L - 2	$1, 3, \ldots, 2L - 3, L - 1$	Ι	$\mathbb{Z}_2$
$D_L \left( L \text{ odd} \right)$	$1  2  3  \dots L-2  L-1$	2L - 2	$1, 3, \ldots, 2L - 3, L - 1$	$II/A_{2L-3}$	$\mathbb{Z}_2$
$E_6$	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	12	1, 4, 5, 7, 8, 11	Ι	$\mathbb{Z}_2$
$E_7$	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	18	1, 5, 7, 9, 11, 13, 17	$\mathrm{II}/D_{10}$	1
$E_8$	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	30	1, 7, 11, 13, 17, 19, 23, 39	Ι	1

Figure 1: Dynkin diagrams of the classical simply-laced A-D-E Lie algebras. The nodes associated with the identity and the fundamental are shown by \*,  $\Box$  respectively. The fundamental is the unique neighbour of the identity. Also shown are the Coxeter numbers g, exponents Exp(G), the type I or II and the so-called parent graphs  $H \neq G$ . The diagram automorphism group  $\Gamma$  is generated by a single  $\mathbb{Z}_2$  automorphism  $\sigma$ . The  $D_4$  graph is an exception having the noncommutative automorphism group  $\mathbb{S}_3$ . The eigenvalues of G are  $2 \cos \frac{\ell \pi}{g}$  with  $\ell \in \text{Exp}(G)$ . By abuse of notation, we use G to denote the graph, the set of its vertices with cardinality |G| and its adjacency matrix so that 2I - G is the Cartan matrix. The meaning of G should be clear from context.

notation of [19, 26], the composite integrable seams are labelled by  $x = (r, a, b, \kappa)$ . The associated *column* transfer matrices are constructed by matrix products of elementary seam transfer matrices

$$\mathbf{T}_{x}(u) = \mathbf{T}_{r}(u,\xi) \,\widehat{\mathbf{N}}_{a} \,\overline{\widehat{\mathbf{N}}}_{b} \,\boldsymbol{\sigma}^{\kappa}, \qquad x = (r,a,b,\kappa), \qquad r \in A_{g-2}, \quad a,b \in G, \quad \kappa = 0,1 \tag{1.2}$$

Here  $\mathbf{T}_r(u,\xi)$  is the column transfer matrix for an *r*-type seam,  $\xi$  is a column inhomogeneity,  $\widehat{\mathbf{N}}_a$ ,  $\overline{\widehat{\mathbf{N}}}_b$  are graph fusion seams labelled by  $a, b \in G$  (as defined in Section 4) and  $\boldsymbol{\sigma}^{\kappa} = \mathbf{I}$  or  $\boldsymbol{\sigma}$  where the  $\sigma$ -parity  $\kappa = 0, 1$  labels respectively the identity  $\mathbf{I}$  and the  $\mathbb{Z}_2$  graph automorphism seam  $\boldsymbol{\sigma}$ . The seam labels  $\kappa$  do not need to be explicitly added in x except for the D series of graphs.

As pointed out in [26], due to the coset factorization structure [29], the *r*-type seams (with  $r \in A_{g-2}$ ) play a somewhat trivial role. We will therefore usually set r = 1 (no *r*-type seam) and focus on the so-called Wess-Zumino-Witten (WZW) factor associated with the graph *G*. For x = (a, b), we assert that the integrable seams  $\widehat{\mathbf{P}}_{a,b} = \widehat{\mathbf{N}}_a \overline{\widehat{\mathbf{N}}}_b$  yield matrix representations of the Ocneanu fusion algebra for arbitrary system sizes. In particular, for the critical *A*-*D*-*E* lattice models, the Ocneanu algebra encodes the quantum symmetries possessed by the commuting families of column transfer matrices on a finite lattice. Following [20], we focus here on the *A* series and the prototypical type I example of  $E_6$ . Our full findings for the *D* and other *E* critical RSOS lattice models will appear elsewhere.

# 2 Critical RSOS models on the torus

#### 2.1 Critical A-D-E lattice models

Defining quantum dimensions by  $S_a = [a]_x = \frac{x^a - x^{-a}}{x - x^{-1}}$  with  $x = e^{i\lambda}$ , the nondegenerate largest eigenvalue of the adjacency matrix G is  $[2]_x = 2 \cos \lambda$  and the associated Perron-Frobenius eigenvector  $\psi$  is

$$G\psi = [2]_{x}\psi, \qquad \psi = (\psi_{a})_{1 \leqslant a \leqslant |G|} = \begin{cases} \left([1]_{x}, [2]_{x}, \dots, [L]_{x}\right), & G = A_{L} \\ \left([1]_{x}, [2]_{x}, \dots, [\ell]_{x}, \frac{[\ell]_{x}}{[2]_{x}}, \frac{[\ell]_{x}}{[2]_{x}}\right), & G = D_{\ell+2} \\ \left([1]_{x}, [2]_{x}, [3]_{x}, [2]_{x}, [1]_{x}, \frac{[3]_{x}}{[2]_{x}}\right), & G = E_{6} \\ \left([1]_{x}, [2]_{x}, [3]_{x}, [4]_{x}, \frac{[6]_{x}}{[2]_{x}}, \frac{[4]_{x}}{[3]_{x}}, \frac{[4]_{x}}{[2]_{x}}\right), & G = E_{7} \\ \left([1]_{x}, [2]_{x}, [3]_{x}, [4]_{x}, [5]_{x}, \frac{[7]_{x}}{[2]_{x}}, \frac{[5]_{x}}{[3]_{x}}, \frac{[5]_{x}}{[2]_{x}}\right), & G = E_{8} \end{cases}$$

The allowed face weights of the critical A-D-E lattice models are then given by

$$W\begin{pmatrix} d & c \\ a & b \end{pmatrix} u = \begin{bmatrix} u \\ a \end{bmatrix}_{b}^{c} = \frac{\sin(\lambda - u)}{\sin\lambda} \delta_{ac} + \frac{\sin u}{\sin\lambda} \sqrt{\frac{\psi_a \psi_c}{\psi_b \psi_d}} \delta_{bd}, \quad \lambda = \frac{\pi}{g}, \quad 0 \le u \le \lambda$$
(2.2)

where u is the spectral parameter,  $\lambda$  is the crossing parameter and the face orientation is marked by an arc in the bottom-left corner. It is understood that the face weights vanish if  $G_{ab}G_{bc}G_{cd}G_{da} = 0$ . The face weights are invariant under reflections in the two diagonals and satisfy the crossing symmetry

$$W\begin{pmatrix} d & c \\ a & b \end{pmatrix} = \begin{bmatrix} d & c \\ u \end{bmatrix}_{b} = \sqrt{\frac{\psi_{a}\psi_{c}}{\psi_{b}\psi_{d}}} \begin{bmatrix} d & c \\ \lambda - u \end{bmatrix}_{b} = \sqrt{\frac{\psi_{a}\psi_{c}}{\psi_{b}\psi_{d}}} W\begin{pmatrix} c & b \\ d & a \end{bmatrix} \lambda - u$$
(2.3)

#### 2.2 Transfer matrices, braid limit and Hamiltonians

The row and column families of commuting transfer matrices  $T_h(u), T(u)$  of the critical A-D-E models on an  $N \times M$  periodic lattice are defined by

$$\boldsymbol{T}_{h}(\boldsymbol{u}) = \cdots \underbrace{\boldsymbol{u}}_{N} \underbrace{\boldsymbol{u}}_{$$

$$\boldsymbol{T}(u) = \boldsymbol{T}_{v}(u) = \cdots \underbrace{u \quad u \quad u \quad \cdots \quad u}_{M} \cdots \qquad (2.5)$$

where, for convenience, the column transfer matrix is rotated clockwise by 90 degrees. The equality  $T_v(u) = T_h(\lambda - u)$  holds when N = M. Other column transfer matrices (seams) are shown in Figure 2. The braid limits of the allowed A D E free weights are

The braid limits of the allowed A-D-E face weights are

$$B\begin{pmatrix} d & c \\ a & b \end{pmatrix} = \frac{d}{a} \sum_{b}^{c} = -\lim_{u \to -i\infty} \frac{x^{\frac{1}{2}}}{\rho(u)} W\begin{pmatrix} d & c \\ a & b \end{vmatrix} u = i \left( -x^{-\frac{1}{2}} \delta_{ac} + x^{\frac{1}{2}} \sqrt{\frac{\psi_a \psi_c}{\psi_b \psi_d}} \delta_{bd} \right)$$
(2.6a)

$$\overline{B}\begin{pmatrix} d & c \\ a & b \end{pmatrix} = \frac{d}{a} \int_{b}^{c} = -\lim_{u \to i\infty} \frac{x^{-\frac{1}{2}}}{\rho(u)} W\begin{pmatrix} d & c \\ a & b \end{pmatrix} u = i \left( x^{\frac{1}{2}} \delta_{ac} - x^{-\frac{1}{2}} \sqrt{\frac{\psi_a \psi_c}{\psi_b \psi_d}} \, \delta_{bd} \right)$$
(2.6b)

where  $x = e^{i\lambda}$  and  $\rho(u) = \sin(\lambda - u)\sin(\lambda + u)$ . These braid limits relate to the left and right chiral halves of the theory and are related to each other by complex conjugation. Taking the action of the face transfer operators  $X_j(u)$  to be diagonal and using the Temperley-Lieb generators  $e_j$ , these expressions become

$$b_j = -\lim_{u \to -i\infty} \frac{x^{\frac{1}{2}}}{\rho(u)} X_j(u) = i\left(-x^{-\frac{1}{2}}I + x^{\frac{1}{2}}e_j\right)$$
(2.7a)

$$\overline{b}_j = -\lim_{u \to i\infty} \frac{x^{-\frac{1}{2}}}{\rho(u)} X_j(u) = i \left( x^{\frac{1}{2}} I - x^{-\frac{1}{2}} e_j \right) = b_j^{-1}$$
(2.7b)

The Hamiltonian limit of the horizontal transfer matrix (2.4) gives rise to periodic anyonic chains

$$\mathcal{H} = -\sin\lambda \frac{d}{du}\log \mathbf{T}_h(u)\Big|_{u=0} = -\sum_{j=1}^N e_j$$
(2.8)

The simplest case is the Hamiltonian limit of the tricritical Ising  $A_4$  RSOS model [30] corresponding to the Fibonacci golden chain [31, 32].



Figure 2: An  $N \times M$  periodic lattice with (N, M) = (10, 8) showing (i) the row transfer matrix  $T_h^x(u)$  with seam segment  $x = (a, b, \kappa)$ , (ii) the column/seam transfer matrices T(u),  $\widehat{\mathbf{N}}_a$  and  $\overline{\widehat{\mathbf{N}}}_b$  as explained in Section 4 and (iii) the  $\mathbb{Z}_2$  diagram automorphism seam  $\boldsymbol{\sigma}^{\kappa}$ . The composite seam is the matrix product  $\mathbf{T}_x = \widehat{\mathbf{N}}_a \overline{\widehat{\mathbf{N}}}_b \boldsymbol{\sigma}^{\kappa}$ . The seam  $\mathbf{T}_x$  has the same twisted partition function as the seam  $\boldsymbol{\sigma}^{\kappa} \widehat{\mathbf{N}}_a \overline{\widehat{\mathbf{N}}}_b$ . We assume r = 1 so there is no r-type seam shown.

# 3 Ocneanu fusion algebra

#### 3.1 Fusion matrices and structure constants

The fused adjacency matrices (intertwiners)  $n_i$  and graph fusion matrices  $\hat{N}_a$  are defined by the finitely truncated recursions

$$n_1 = I, \quad n_2 = G, \quad n_i n_2 = n_{i-1} + n_{i+1}, \quad n_g = 0, \qquad i \in A_{g-1}$$

$$(3.1)$$

$$\widehat{N}_1 = I, \quad \widehat{N}_2 = G, \quad G \,\widehat{N}_a = \sum_{b \in G} G_{ab} \,\widehat{N}_b, \qquad a \in G$$
(3.2)

For  $G = A_{g-1}$ , the fused adjacency matrices  $n_i$  are in fact the Verlinde [12] fusion matrices  $N_i$ . The fused adjacency and graph fusion matrices satisfy the commutative associative fusion algebras

$$n_i n_j = \sum_{k=1}^{g-1} N_{ij}{}^k n_k, \qquad \widehat{N}_a \, \widehat{N}_b = \sum_{c \in G} \widehat{N}_{ab}{}^c \, \widehat{N}_c, \qquad n_i \, \widehat{N}_a = \sum_{b \in G} n_{ia}{}^b \, \widehat{N}_b \tag{3.3}$$

where the structure constants are

$$N_{ij}{}^{k} = (N_i)_{j}{}^{k} \in \mathbb{N}, \qquad \widehat{N}_{ab}{}^{c} = (\widehat{N}_a)_{b}{}^{c} \in \mathbb{Z}, \qquad n_{ia}{}^{b} = (n_i)_{a}{}^{b} \in \mathbb{N}$$
(3.4)

For type I theories,  $\widehat{N}_{ab}{}^c \in \mathbb{N}$ . So the  $n_i$  and, for type I graphs, the  $\widehat{N}_a$  form nimreps (nonnegative integer matrix representations) of these fusion algebras.

For structure constants larger than 1, it is necessary to introduce so-called bond variables (graphs with multiple edges). Denoting the complex unitary matrices of eigenvectors of  $N_2$ ,  $\hat{N}_2$  by  $S_i^{\ell}$  (modular matrix) and  $\psi_a^{\ell}$  with  $S_1^{\ell}, \psi_1^{\ell} > 0$ , the corresponding Verlinde and Verlinde-type formulas are

$$N_{ij}{}^{k} = \sum_{\ell=1}^{g-1} \frac{S_{i}{}^{\ell} S_{j}{}^{\ell} \overline{S_{k}}^{\ell}}{S_{1}{}^{\ell}}, \quad \widehat{N}_{ab}{}^{c} = \sum_{\ell \in \operatorname{Exp}(G)} \frac{\psi_{a}{}^{\ell} \psi_{b}{}^{\ell} \overline{\psi_{c}{}^{\ell}}}{\psi_{1}{}^{\ell}}, \quad n_{ia}{}^{b} = \sum_{\ell \in \operatorname{Exp}(G)} \frac{S_{i}{}^{\ell} \psi_{a}{}^{\ell} \overline{\psi_{b}{}^{\ell}}}{S_{1}{}^{\ell}}$$
(3.5)

Here bars denote complex conjugation and the sum in the first expression is over  $\ell \in \text{Exp}(A_{g-1})$ . Explicit fusion matrices for  $A_4$  and  $E_6$  are

$$N_{1} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad N_{2} = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix}, \quad N_{3} = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}, \quad N_{4} = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}$$
(3.6)

$$\widehat{N}_{1} = \begin{pmatrix}
1 & 0 & 0 & 0 & 0 & 0 \\
0 & 1 & 0 & 0 & 0 & 0 \\
0 & 0 & 1 & 0 & 0 & 0 \\
0 & 0 & 0 & 1 & 0 & 0 \\
0 & 0 & 0 & 0 & 1 & 0 \\
0 & 0 & 0 & 0 & 0 & 1
\end{pmatrix}, \quad \widehat{N}_{2} = \begin{pmatrix}
0 & 1 & 0 & 0 & 0 & 0 \\
1 & 0 & 1 & 0 & 1 & 0 & 1 \\
0 & 0 & 1 & 0 & 1 & 0 & 1 \\
0 & 0 & 1 & 0 & 1 & 0 & 0 \\
0 & 0 & 1 & 0 & 1 & 0 & 1 \\
0 & 0 & 1 & 0 & 1 & 0 & 0 \\
0 & 0 & 1 & 0 & 1 & 0 & 1 \\
1 & 0 & 1 & 0 & 1 & 0 & 1 \\
1 & 0 & 1 & 0 & 0 & 0 \\
0 & 1 & 0 & 1 & 0 & 0 & 0 \\
0 & 1 & 0 & 1 & 0 & 0 & 0 \\
0 & 0 & 1 & 0 & 0 & 0 & 0 \\
0 & 0 & 1 & 0 & 0 & 0 & 0 \\
0 & 0 & 1 & 0 & 0 & 0 & 0 \\
0 & 0 & 1 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 1 \\
0 & 0 & 1 & 0 & 0 & 0 & 0 \\
0 & 0 & 1 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 1
\end{pmatrix}, \quad \widehat{N}_{6} = \begin{pmatrix}
0 & 0 & 0 & 0 & 0 & 1 & 0 \\
0 & 0 & 1 & 0 & 0 & 0 & 0 \\
0 & 0 & 1 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 1
\end{pmatrix}, \quad (3.7)$$

where  $N_4$  and  $\hat{N}_5$  encode the  $\mathbb{Z}_2$  graph automorphism of  $A_4$  and  $E_6$ . For type I theories, there exists an extended conformal chiral algebra with extended Virasoro characters encoded by the fundamental intertwiner  $C = (n_{i1}{}^a)$ . Accordingly, for the type I graphs, we have

$$n_{i} = \sum_{a \in G} n_{i1}{}^{a} \widehat{N}_{a}, \qquad \widehat{\chi}_{r,a}(q) = \sum_{s=1}^{g-1} n_{s1}{}^{a} \chi_{r,s}(q)$$
(3.8)

The first of these decompositions follows by specializing the compatibility condition in (3.3). For  $E_6$ :

#### **3.2** Twisted conformal partition functions

The twisted partition functions and structure constants of the Ocneanu algebra are given by Petkova and Zuber [19]. Here we focus on  $(A_{g-2}, G)$  cosets with G of type I. For these, the twisted partition functions are given by the sesquilinear forms

$$Z^{(r,a,b,\kappa)} = \sum_{(r',s'),(r'',s'')} N_{r,r'} {}^{r''} [P_{ab}^{(\kappa)}]_{s's''} \chi_{r',s'}(q) \chi_{r'',s''}(\overline{q}) = \sum_{(r',a'),(r'',a'')} N_{r,r'} {}^{r''} [\widehat{P}_{ab}^{(\kappa)}]_{a'a''} \widehat{\chi}_{r',a'}(q) \widehat{\chi}_{r'',a''}(\overline{q})$$
(3.10)

where the second form is new. Here  $r, r', r'' \in A_{g-2}$ ;  $s, s', s'' \in A_{g-1}$ ;  $a, a', a'', b \in G$  and the *toric* and *extended toric* matrices are

$$[P_{ab}^{(\kappa)}]_{s\bar{s}} = \sum_{c \in T_{\kappa}} n_{sa}{}^{c} n_{\bar{s}b}{}^{c}, \qquad [\widehat{P}_{ab}^{(\kappa)}]_{a'a''} = \sum_{c \in T_{\kappa}} \widehat{N}_{ac}{}^{a'} \widehat{N}_{bc}{}^{a''}, \qquad \kappa = 0, 1$$
(3.11)

The label  $\kappa = 0, 1$  is only needed for the *D* series and indicates the absence or presence of the  $\mathbb{Z}_2$  automorphism. The *ambichiral vertices* (common to both left and right chiral halves) are

$$T_{0} = \begin{cases} \{1, 2, \dots, L\}, & G = A_{L} \\ \{1, 3, 5, \dots, 2\ell - 1, 2\ell\}, & G = D_{2\ell} \\ \{1, 5, 6\}, & G = E_{6} \\ \{1, 7\}, & G = E_{8} \end{cases} \qquad T_{1} = \begin{cases} \{2, 4, \dots, 2\ell - 2\}, & G = D_{2\ell} \\ T_{0}, & \text{otherwise} \end{cases}$$
(3.12)

#### 3.3 Extended toric matrices and quantum symmetry

In the following, we fix r = 1. Taking  $x = (a, b, \kappa)$ , the extended toric matrices can be written as

$$\widehat{P}_{ab}^{(\kappa)} = (\widehat{N}_a^{\kappa})^T \widehat{N}_b^{\kappa}, \qquad x = (a, b, \kappa)$$
(3.13)

where the superscript T denotes transpose and, in accord with the tensor structure  $G \otimes_{T_{\kappa}} G$ , the rows of the rectangular matrices  $\widehat{N}_{a}^{\kappa}$  are restricted to those labelled by the vertices of  $T_{\kappa}$ . Alternatively, working with the full  $|G| \times |G|$  matrices, the extended toric matrices are given by the bilinear map

$$\widehat{P}_{ab}^{(\kappa)} = (\widehat{N}_a, \widehat{N}_b)_{\kappa} = \widehat{N}_a \widehat{P}^{(\kappa)} \widehat{N}_b, \qquad (L, R)_{\kappa} = L \widehat{P}^{(\kappa)} R = L \otimes_{T_\kappa} R, \qquad \widehat{P}^{(\kappa)} = \widehat{P}_{11}^{(\kappa)} = (I, I)_{\kappa} \quad (3.14)$$

The projection  $\hat{P}^{(\kappa)} = \hat{P}_{11}^{(\kappa)}$  joins the left and right chiral components through the ambichiral nodes  $T_0$  or, for the *D* series, its complement  $T_1$ .

For  $E_6$ , the 6 × 6 array of extended toric matrices  $\widehat{P}_{ab}$  is shown in Figure 3 with no  $\kappa$  dependence since  $T_0 = T_1$ . In accord with the quantum symmetry exhibited in Figure 3, the 36 toric matrices lie in a convex cone K consisting of nonnegative-integer linear (conical) combinations of the Hilbert basis  $\{\hat{P}_{ab}: a = 1, 2, \dots, 6; b = 1, 2\}$ . The 12 basis elements are in bijection with the vertices of the Ocneanu graph in Figure 4. They are conjugate to the 12  $E_6$  irreducible representations of Vir  $\otimes \overline{\text{Vir}}$ . The remaining 24 toric matrices are expressed as the conical linear combinations

$$\hat{P}_{55} = \hat{P}_{11}, \qquad \hat{P}_{45} = \hat{P}_{21}, \qquad \hat{P}_{26} = \hat{P}_{35} = \hat{P}_{46} = \hat{P}_{31}$$
(3.15a)

$$_{5} = P_{41}, \qquad P_{15} = P_{51}, \qquad P_{16} = P_{56} = P_{65} = P_{61}$$
(3.15b)

$$\hat{P}_{25} = \hat{P}_{41}, \qquad \hat{P}_{15} = \hat{P}_{51}, \qquad \hat{P}_{16} = \hat{P}_{56} = \hat{P}_{65} = \hat{P}_{61} \tag{3.15b}$$

$$\hat{P}_{54} = \hat{P}_{12}, \qquad \hat{P}_{44} = \hat{P}_{22}, \qquad \hat{P}_{23} = \hat{P}_{34} = \hat{P}_{43} = \hat{P}_{32} \tag{3.15c}$$

$$\hat{P}_{24} = \hat{P}_{42}, \qquad \hat{P}_{14} = \hat{P}_{52}, \qquad \hat{P}_{13} = \hat{P}_{53} = \hat{P}_{64} = \hat{P}_{62} \tag{3.15d}$$

$$P_{42}, \qquad P_{14} = P_{52}, \qquad P_{13} = P_{53} = P_{64} = P_{62}$$
 (3.15d)

$$\hat{P}_{33} = \hat{P}_{22} + \hat{P}_{42}, \qquad \hat{P}_{36} = \hat{P}_{21} + \hat{P}_{41}, \qquad \hat{P}_{63} = \hat{P}_{12} + \hat{P}_{52}, \qquad \hat{P}_{66} = \hat{P}_{11} + \hat{P}_{51} \tag{3.15e}$$

The Hilbert basis of extended toric matrices is unique up to the equivalences in (3.15a) to (3.15d).

$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0$	$\begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0$	$\begin{pmatrix} 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 &$	$\begin{pmatrix} 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 &$	$\begin{pmatrix} 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0$	$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0$
$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0$	$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 &$	$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 &$	$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0$	$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 &$	$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 &$
$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 &$	$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 &$	$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 2 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 &$	$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 &$	$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 &$	$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 2 \\ 1 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0$
$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 &$	$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0$	$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 &$	$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 &$	$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0$	$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 &$
$\begin{pmatrix} 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0$	$\begin{pmatrix} 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 &$	$\begin{pmatrix} 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 &$	$\begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0$	$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0$	$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0$
$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0$	$\begin{pmatrix} 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 &$	$\begin{pmatrix} 0 & 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 &$	$\begin{pmatrix} 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 &$	$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0$	$\begin{pmatrix} 1 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0$

Figure 3: The  $6 \times 6$  array of the extended toric matrices  $\hat{P}_{ab}$  of  $E_6$ . Only 12 are linearly independent reflecting the known quantum symmetry. There is no dependence on  $\kappa$  since, in this case,  $T_0 = T_1 = T$ .

#### $\mathbf{3.4}$ Ocneanu and double graph fusion algebras

Algebraically, the Ocneanu fusion algebra is the double graph fusion algebra

$$\widetilde{N}_x \star \widetilde{N}_y = \sum_z \widetilde{N}_{xy}{}^z \widetilde{N}_z, \qquad x = (a, b), \quad y = (a', b''), \quad z = (a'', b'')$$
(3.16)

where  $\star$  denotes the fusion product and  $\widetilde{N}_{xy}^{z}$  are the structure constants. For the A series, with  $i,j,i',j',i'',j'' \in A_{g-1}$  and  $\widehat{P}^{(0)} = I$ , the Ocneanu fusion matrices are the double Verlinde fusion matrices

$$\widetilde{N}_{xy}{}^{z} = N_{ii'}{}^{i''}N_{jj'}{}^{j''}, \qquad \widetilde{N}_{x} = \widetilde{N}_{ij} = N_iN_j, \quad x = (i,j), \quad y = (i',j'), \quad z = (i'',j'')$$
(3.17)

More generally, for type I A-D-E theories with  $a, b, a', b', a'', b'' \in G$ , the Ocneanu algebra (3.16) is given by the double graph fusion matrices

$$\widetilde{N}_{x} = \widetilde{N}_{ab} = \widehat{N}_{a}\widehat{P}^{(\kappa)}\widehat{N}_{b} = \widehat{P}^{(\kappa)}_{ab}, \quad \widetilde{N}_{xy}{}^{z} = \widehat{N}^{\kappa}_{aa''}{}^{a'}\widehat{N}^{\kappa}_{bb'}{}^{b''}, \quad x = (a,b), \ y = (a',b'), \ z = (a'',b'')$$
(3.18)

We define the fusion product  $\star$  of two extended toric matrices by

$$\widehat{P}_{aa'}^{(\kappa)} \star \widehat{P}_{bb'}^{(\kappa')} = (\widehat{N}_a, \widehat{N}_{a'})_{\kappa} \star (\widehat{N}_b, \widehat{N}_{b'})_{\kappa'} := (\widehat{N}_a \widehat{N}_b, \widehat{N}_{a'} \widehat{N}_{b'})_{\kappa+\kappa'} = \sum_{c,c'} \widehat{N}_{ab}{}^c \widehat{N}_{a'b'}{}^{c'} \widehat{P}_{cc'}^{(\kappa+\kappa')}$$
(3.19)

where the  $\sigma$ -parity sum  $\kappa + \kappa'$  is taken mod 2 and we use the independent left and right graph fusion algebras (3.2) and bilinearity. The result can be simplified using the quantum symmetries (3.15). In this notation, the left and right fundamentals are  $\hat{P}_{21}^{(\kappa)} = (\hat{N}_2, \hat{N}_1)_{\kappa} = (\hat{N}_2, I)_{\kappa}$  and  $\hat{P}_{12}^{(\kappa)} = (\hat{N}_1, \hat{N}_2)_{\kappa} = (I, \hat{N}_2)_{\kappa}$ . For example, dropping the  $\kappa$  dependence for  $E_6$ , we find

$$\widehat{P}_{32} \star \widehat{P}_{21} = (\widehat{N}_3, \widehat{N}_2) \star (\widehat{N}_2, \widehat{N}_1) = (\widehat{N}_3 \widehat{N}_2, \widehat{N}_2 \widehat{N}_1) = (\widehat{N}_2 + \widehat{N}_4 + \widehat{N}_6, \widehat{N}_2) = \widehat{P}_{22} + \widehat{P}_{42} + \widehat{P}_{62} \quad (3.20)$$

$$\widehat{P}_{12} \star \widehat{P}_{33} = (\widehat{N}_1, \widehat{N}_2) \star (\widehat{N}_3, \widehat{N}_3) = (\widehat{N}_1 \widehat{N}_3, \widehat{N}_2 \widehat{N}_3) = (\widehat{N}_3, \widehat{N}_2 + \widehat{N}_4 + \widehat{N}_6)$$

$$= (\hat{N}_3, \hat{N}_2) + (\hat{N}_3, \hat{N}_4) + (\hat{N}_3, \hat{N}_6) = \hat{P}_{32} + \hat{P}_{34} + \hat{P}_{36} = \hat{P}_{21} + 2\hat{P}_{32} + \hat{P}_{41}$$
(3.21)

Due to the chiral action of the  $\star$  fusion product multiplying left with left chiral components and right with right chiral components, this realization of the Ocneanu algebra does not yield simple matrix product representations of the algebra.

# 4 A-D-E integrable seams

#### 4.1 Construction of *s*-type seams

The fundamental braid seams  $\mathbf{B}, \overline{\mathbf{B}}$  are built analogously to the transfer matrix T(u) in (2.5) but with the face weights replaced by the complex weights (2.6a) and (2.6b) respectively. These are just the braid limits of the commuting family T(u), suitably normalized, so  $\mathbf{B}, \overline{\mathbf{B}}$  commute with T(u) and with each other. Since  $\overline{\mathbf{B}} = \mathbf{B}^T$ ,  $\mathbf{B}$  is Hermitian. It also has the same real eigenvalues as G. The fused *s*-type braid seams  $\mathbf{n}_s, \bar{\mathbf{n}}_s$  are defined recursively by the analog of (3.1)

$$\mathbf{n}_0 = 0, \qquad \mathbf{n}_1 = I, \qquad \mathbf{n}_2 = \mathbf{B}, \qquad \mathbf{n}_s \mathbf{n}_2 = \mathbf{n}_{s-1} + \mathbf{n}_{s+1}$$
 (4.1a)

$$\mathbf{\bar{n}}_0 = 0, \quad \mathbf{\bar{n}}_1 = I, \quad \mathbf{\bar{n}}_2 = \mathbf{B}, \quad \mathbf{\bar{n}}_s \mathbf{\bar{n}}_2 = \mathbf{\bar{n}}_{s-1} + \mathbf{\bar{n}}_{s+1}$$
(4.1b)

Alternatively, these braid seams are constructed using Wenzl-Jones projectors to build the fused transfer matrices  $\mathbf{T}^{1,s}(u)$  and then taking the braid limits. The fused braid matrices  $\mathbf{n}_s, \bar{\mathbf{n}}_s$  share the same real eigenvalues which are a subset of the eigenvalues of the Verlinde matrices  $N_s$ .

If  $G = A_{g-1}$  then, for arbitrary system size M,  $\mathbf{N}_s = \mathbf{n}_s$  and  $\overline{\mathbf{N}}_s = \overline{\mathbf{n}}_s$  give matrix representations of the Verlinde algebra

$$\mathbf{N}_{i} \mathbf{N}_{j} = \sum_{k=1}^{g-1} N_{ij}{}^{k} \mathbf{N}_{k}, \qquad \overline{\mathbf{N}}_{i} \overline{\mathbf{N}}_{j} = \sum_{k=1}^{g-1} N_{ij}{}^{k} \overline{\mathbf{N}}_{k}$$
(4.2)

In these cases, with x = (i, i'), y = (j, j'), z = (k, k'), the double Verlinde algebra gives a matrix representation of the Ocneanu algebra (3.16) with  $\widetilde{\mathbf{N}}_x = \widetilde{\mathbf{N}}_{ii'} = \mathbf{N}_i \overline{\mathbf{N}}_{i'}$ 

$$\widetilde{\mathbf{N}}_{x}\widetilde{\mathbf{N}}_{y} = \widetilde{\mathbf{N}}_{ii'}\widetilde{\mathbf{N}}_{jj'} = \sum_{k,k'} N_{ij}{}^{k} N_{i'j'}{}^{k'}\widetilde{\mathbf{N}}_{kk'} = \sum_{z} \widetilde{N}_{xy}{}^{z} \widetilde{\mathbf{N}}_{z}$$
(4.3)

### **4.2** Construction of seams of type (a, b)

We now extend the decomposition (3.8) to fused seams

$$\mathbf{n}_s = \sum_{a \in G} n_{s1}{}^a \, \widehat{\mathbf{N}}_a, \qquad \bar{\mathbf{n}}_s = \sum_{b \in G} n_{s1}{}^b \, \overline{\widehat{\mathbf{N}}}_b \tag{4.4}$$

The face weights of the fused seams  $\widehat{\mathbf{N}}_a, \overline{\widehat{\mathbf{N}}}_b$  can be calculated systematically as explained in [26]. However, the form of these face weights is not unique. Indeed, they depend on a choice of gauge. Here we will work in a different gauge to [26]. For arbitrary system size M, the fused seam matrices constructed here provide matrix representations of the same fused adjacency and graph fusion algebras as in (3.3) with the same structure constants

$$\mathbf{n}_{i} \, \mathbf{n}_{j} = \sum_{k=1}^{g-1} N_{ij}{}^{k} \, \mathbf{n}_{k}, \qquad \widehat{\mathbf{N}}_{a} \, \widehat{\mathbf{N}}_{b} = \sum_{c \in G} \widehat{N}_{ab}{}^{c} \, \widehat{\mathbf{N}}_{c}, \qquad \mathbf{n}_{i} \, \widehat{\mathbf{N}}_{a} = \sum_{b \in G} n_{ia}{}^{b} \, \widehat{\mathbf{N}}_{b} \tag{4.5a}$$

$$\bar{\mathbf{n}}_i \, \bar{\mathbf{n}}_j = \sum_{k=1}^{g-1} N_{ij}{}^k \, \bar{\mathbf{n}}_k, \qquad \overline{\widehat{\mathbf{N}}}_a \, \overline{\widehat{\mathbf{N}}}_b = \sum_{c \in G} \widehat{N}_{ab}{}^c \, \overline{\widehat{\mathbf{N}}}_c, \qquad \bar{\mathbf{n}}_i \, \overline{\widehat{\mathbf{N}}}_a = \sum_{b \in G} n_{ia}{}^b \, \overline{\widehat{\mathbf{N}}}_b \tag{4.5b}$$

For our purposes here, we will restrict ourselves to  $E_6$ . In this case, we assert that the graph fusion seam matrices are given by

$$\hat{\mathbf{N}}_a = \mathbf{n}_a, \ a = 1, 2, 3; \quad \hat{\mathbf{N}}_4 = \mathbf{n}_6 - \mathbf{n}_2, \ \hat{\mathbf{N}}_5 = \mathbf{n}_5 - \mathbf{n}_3, \ \hat{\mathbf{N}}_6 = \mathbf{n}_2 + \mathbf{n}_4 - \mathbf{n}_6$$
(4.6a)

$$\overline{\widehat{\mathbf{N}}}_b = \overline{\mathbf{n}}_b, \quad b = 1, 2, 3; \qquad \overline{\widehat{\mathbf{N}}}_4 = \overline{\mathbf{n}}_6 - \overline{\mathbf{n}}_2, \quad \overline{\widehat{\mathbf{N}}}_5 = \overline{\mathbf{n}}_5 - \overline{\mathbf{n}}_3, \quad \overline{\widehat{\mathbf{N}}}_6 = \overline{\mathbf{n}}_2 + \overline{\mathbf{n}}_4 - \overline{\mathbf{n}}_6 \tag{4.6b}$$

With these, it is readily verified using Mathematica [33] that, for even system sizes up to M = 12, all of the relations (4.5) are satisfied and that all of these matrices commute.

For  $N_a$  with a = 4, 5, 6, we find that the allowed face weights are given by

The face weights not explicitly listed vanish. Here  $\hat{\mathbf{N}}_5 = \boldsymbol{\sigma}$  is the  $\mathbb{Z}_2 E_6$  graph automorphism seam. For  $\overline{\hat{\mathbf{N}}}_a$  with a = 4, 5, 6, the face weights are given by the complex conjugates. Reflection in the horizontal implements complex conjugation. We note that  $\widehat{\mathbf{N}}_a$  is real for the ambichiral vertices a = 1, 5, 6.

#### 4.3 Ocneanu algebra of seams and quantum symmetry

For even system sizes up to M = 12, it is now readily confirmed that a *matrix representation* of the  $E_6$  Ocneanu algebra, defined in (3.18) with the fusion product  $\star$  as in (3.19), is given by the double graph fusion algebra of seams

$$\widehat{\mathbf{P}}_{ab} = \widetilde{\mathbf{N}}_x = \widetilde{\mathbf{N}}_{ab} = \widehat{\mathbf{N}}_a \,\widehat{\mathbf{N}}_b, \qquad x = (a, b), \quad y = (a', b'), \quad z = (a'', b'') \tag{4.10a}$$

$$\widetilde{\mathbf{N}}_{x}\widetilde{\mathbf{N}}_{y} = \sum_{z} \widetilde{N}_{xy}{}^{z} \widetilde{\mathbf{N}}_{z}, \qquad \widetilde{\mathbf{N}}_{a,b} \widetilde{\mathbf{N}}_{a'b'} = \sum_{a'',b''} \widehat{N}_{aa'}{}^{a''} \widehat{N}_{bb'}{}^{b''} \widetilde{\mathbf{N}}_{a''b''}, \qquad \widetilde{N}_{xy}{}^{z} = \widehat{N}_{aa'}{}^{a''} \widehat{N}_{bb'}{}^{b''}$$
(4.10b)

In this realization involving fusion of seams, the fusion product  $\star$  is replaced with the usual matrix product. In a sense, the decomposition of fusion products of seams on the lattice is the analog of the local operator product expansion in CFT.

In accord with the quantum symmetry (3.15) exhibited by the 36 "extended toric seams" (see Figure 3), the Hilbert basis of compound seams  $\{\widehat{\mathbf{P}}_{a,b}: a = 1, 2, \ldots, 6; b = 1, 2\}$  is in bijection with the 12 nodes of the Ocneanu graph in Figure 4. The remaining 24 "extended toric seams" are given by the conical linear combinations

$$\widehat{\mathbf{P}}_{5,5} = \widehat{\mathbf{P}}_{1,1}, \qquad \widehat{\mathbf{P}}_{4,5} = \widehat{\mathbf{P}}_{2,1}, \qquad \widehat{\mathbf{P}}_{2,6} = \widehat{\mathbf{P}}_{3,5} = \widehat{\mathbf{P}}_{4,6} = \widehat{\mathbf{P}}_{3,1}$$
(4.11a)

$$\widehat{\mathbf{P}}_{2,5} = \widehat{\mathbf{P}}_{4,1}, \qquad \widehat{\mathbf{P}}_{1,5} = \widehat{\mathbf{P}}_{5,1}, \qquad \widehat{\mathbf{P}}_{1,6} = \widehat{\mathbf{P}}_{5,6} = \widehat{\mathbf{P}}_{6,5} = \widehat{\mathbf{P}}_{6,1}$$
(4.11b)

$$\widehat{\mathbf{P}}_{5,4} = \widehat{\mathbf{P}}_{1,2}, \qquad \widehat{\mathbf{P}}_{4,4} = \widehat{\mathbf{P}}_{2,2}, \qquad \widehat{\mathbf{P}}_{2,3} = \widehat{\mathbf{P}}_{3,4} = \widehat{\mathbf{P}}_{4,3} = \widehat{\mathbf{P}}_{3,2}$$
(4.11c)

$$\hat{\mathbf{P}}_{2,4} = \hat{\mathbf{P}}_{4,2}, \qquad \hat{\mathbf{P}}_{1,4} = \hat{\mathbf{P}}_{5,2}, \qquad \hat{\mathbf{P}}_{1,3} = \hat{\mathbf{P}}_{5,3} = \hat{\mathbf{P}}_{6,4} = \hat{\mathbf{P}}_{6,2}$$
(4.11d)

$$\mathbf{\hat{P}}_{3,3} = \mathbf{\hat{P}}_{2,2} + \mathbf{\hat{P}}_{4,2}, \qquad \mathbf{\hat{P}}_{3,6} = \mathbf{\hat{P}}_{2,1} + \mathbf{\hat{P}}_{4,1}, \qquad \mathbf{\hat{P}}_{6,3} = \mathbf{\hat{P}}_{1,2} + \mathbf{\hat{P}}_{5,2}, \qquad \mathbf{\hat{P}}_{6,6} = \mathbf{\hat{P}}_{1,1} + \mathbf{\hat{P}}_{5,1}$$
(4.11e)

where again this was checked in Mathematica out to even system sizes M = 12. In general, the representations may not be faithful for small values of M. For  $E_6$ , the representations are not faithful for M < 6. The quantum symmetry expresses the rules for fusing left and right chiral seams. The matrix product of any two toric seams  $\hat{\mathbf{P}}_{a,b}$  is easily found using the factorization (4.10a), the independent left and right chiral graph fusion algebras (4.5) and the quantum symmetries (4.11). The  $E_6$  Cayley table of the Ocneanu algebra of seams is shown in Figure 5.



Figure 4: Ocneanu fusion graph  $Oc(E_6) = E_6 \oplus E_6$  of  $E_6$  with 12 vertices showing the two independent copies of  $E_6$  graph colored in black and yellow. The 12 nodes are in bijection with the linearly independent integrable seams  $\hat{\mathbf{P}}_{ab}$ . The red and blue lines (solid or dashed) show the action of the fundamental seams  $\hat{\mathbf{P}}_{21}$  and  $\hat{\mathbf{P}}_{12}$  respectively. This action interlocks four copies of  $E_6$ . The ambichiral seams are  $\hat{\mathbf{P}}_{a1}$  with a = 1, 5, 6.

There is an injective ring homomorphism

$$\phi: \left\langle \widehat{\mathbf{P}}_{a1}, \widehat{\mathbf{P}}_{a2} \,|\, a = 1, \dots, 6 \right\rangle \to \mathbb{Z}[x, y] / \left\langle p_1(x), p_2(x, y) \right\rangle \tag{4.12}$$

where

$$p_1(x) = \prod_{\ell \in \text{Exp}(E_6)} \left( x - 2\cos\frac{\ell\pi}{g} \right) = x^6 - 5x^4 + 5x^2 - 1, \quad p_2(x,y) = y^2 + (x^5 - 5x^3 + 4x)y - 1 \quad (4.13)$$

 $p_2(y, y) = p_1(y)$  and

$$\widehat{\mathbf{P}}_{11} \mapsto 1, \quad \widehat{\mathbf{P}}_{21} \mapsto x, \quad \widehat{\mathbf{P}}_{31} \mapsto x^2 - 1, \quad \widehat{\mathbf{P}}_{41} \mapsto x^5 - 4x^3 + 2x$$

$$(4.14a)$$

$$\widehat{\mathbf{P}}_{51} \mapsto x^4 - 4x^2 + 2, \quad \widehat{\mathbf{P}}_{61} \mapsto -x^5 + 5x^3 - 4x, \qquad \widehat{\mathbf{P}}_{a2} \mapsto \phi(\widehat{\mathbf{P}}_{a1}) y, \quad a = 1, \dots, 6 \qquad (4.14b)$$

The  $E_6$  integrable seams all mutually commute and are simultaneously diagonalizable. The eigenvalues therefore yield a 1-dimensional representation of the  $E_6$  Ocneanu algebra in terms of  $s\ell(2)$  quantum dimensions on Vir  $\otimes \overline{\text{Vir}}$ . For the spin- $\frac{1}{2}$  fundamentals, let us set  $x = q_1 + q_1^{-1}$ ,  $y = q_2 + q_2^{-1}$  with  $q_1 = q^s$ ,  $q_2 = \bar{q}^{\bar{s}} = q^{-\bar{s}}$  and  $q = e^{\pi i/12}$  to ensure the fusion rules truncate correctly. We then need to solve

$$p_1(x) = 0, \qquad p_2(x, y) = 0$$
(4.15)

These equations admit  $6 \times 2 = 12$  solutions for  $x = q^s + q^{-s}$  and  $y = q^{\bar{s}} + q^{-\bar{s}}$  where, explicitly,

$$(s,\bar{s}) \in \{(1,1), (4,4), (5,5), (7,7), (8,8), (11,11), (1,7), (7,1), (4,8), (8,4), (5,11), (11,5)\}$$
(4.16)

As required, this set is symmetric under interchange of s and  $\bar{s}$ . In the  $(s, \bar{s})$  sector, the corresponding

	21	31	41	51	61	12	22	32	42	52	62
21	11+31	$21\!+\!41\!+\!61$	$31\!+\!51$	41	31	22	12+32	22+42+62	32 + 52	42	32
31	21 + 41 + 61	11+2(31)+51	21 + 41 + 61	31	21 + 41	32	22 + 42 + 62	12+2(32)+52	22 + 42 + 62	32	22+42
41	31 + 51	21 + 41 + 61	11+31	21	31	42	32 + 52	22 + 42 + 62	12+32	22	32
51	41	31	21	11	61	52	42	32	22	12	62
61	31	21+41	31	61	11 + 51	62	32	22+42	32	62	12+52
12	22	32	42	52	62	$11\!+\!62$	21 + 32	22 + 31 + 42	$32\!\!+\!\!41$	51 + 62	12 + 52 + 61
22	12+32	22+42+62	32+52	42	32	21+32	$11 + 22 + 31 \\ + 42 + 62$	$\begin{array}{r} 12 + 21 + 2(32) \\ + 41 + 52 + 61 \end{array}$	22+31+42 +51+62	32+41	22+31+42
32	22+42+62	12+2(32)+52	22+42+62	32	22+42	22+31+42	12+21+2(32) +41+52+61	11+2(22)+2(31) +2(42)+51+2(62)	$12+21+2(32) \\ +41+52+61$	22+31+42	21+2(32)+41
42	32+52	22+42+62	12+32	22	32	32+41	22+31+42 +51+62	$\begin{array}{r} 12 + 21 + 2(32) \\ + 41 + 52 + 61 \end{array}$	$11 + 22 + 31 \\ + 42 + 62$	21+32	22+31+42
52	42	32	22	12	62	$51\!+\!62$	32+41	22+31+42	21+32	11+62	12 + 52 + 61
62	32	22+42	32	62	12+52	12+52+61	22+31+42	21+2(32)+41	22+31+42	12 + 52 + 61	11+51+2(62)

Figure 5: Cayley table of the  $E_6$  Ocneanu algebra of seams. In this case, the product of seams is commutative. We use the compact notation  $ab = \widehat{\mathbf{P}}_{ab}$  and omit the identity ab = 11 since trivially (11)(ab) = (ab)(11) = (ab).

quantum dimensions are

$$\begin{array}{ll} (1,1):1 & (1,2):q^{\bar{s}}+q^{-\bar{s}} \\ (2,1):q^{s}+q^{-s} & (2,2):q^{s+\bar{s}}+q^{s-\bar{s}}+q^{-s+\bar{s}}+q^{-s-\bar{s}} \\ (3,1):q^{2s}+1+q^{-2s} & (3,2):q^{2s+\bar{s}}+q^{2s-\bar{s}}+q^{\bar{s}}+q^{-\bar{s}}+q^{-2s+\bar{s}}+q^{-2s-\bar{s}} \\ (4,1):q^{5s}+q^{3s}+q^{-3s}+q^{-5s} & (4,2):q^{5s+\bar{s}}+q^{5s-\bar{s}}+q^{3s+\bar{s}}+q^{3s-\bar{s}}+q^{-3s+\bar{s}}+q^{-3s-\bar{s}}+q^{-5s+\bar{s}}+q^{-5s-\bar{s}} \\ (5,1):q^{4s}+q^{-4s} & (5,2):q^{4s+\bar{s}}+q^{4s-\bar{s}}+q^{-4s+\bar{s}}+q^{-4s-\bar{s}} \\ (6,1):-q^{5s}+q^s+q^{-s}-q^{-5s} & (6,2):-q^{5s+\bar{s}}-q^{5s-\bar{s}}+q^{s+\bar{s}}+q^{s-\bar{s}}+q^{-s+\bar{s}}+q^{-s-\bar{s}}-q^{-5s+\bar{s}}-q^{-5s-\bar{s}} \\ \end{array}$$

Taking the union over the 12 sectors (4.16), we thus find that the distinct eigenvalues of the integrable seams are given by the roots of the minimal polynomials

The toric seam matrices satisfy the polynomial equations obtained by setting the associated minimal polynomial to zero. We observe that some integrable seams have a zero eigenvalue and so are not invertible. Strictly speaking, in this sense, these seams are not symmetries — they are "non-invertible symmetries" [34].

As we will discuss elsewhere, similar characterisations of the Ocneanu algebra of seams apply for the D and other E type theories.

# 5 Conclusion

In this short article, we have developed further the lattice approach to fusion of critical unitary RSOS lattice models. Using Yang-Baxter integrability, it is now possible to construct column transfer matrices

(seams) for each distinct topological defect that appears in the associated CFT in the continuum scaling limit. Like CFT topological defects, the lattice seams commute and propagate freely along the lattice. Remarkably, the lattice seams satisfy Verlinde, graph fusion and Ocneanu algebras. In addition, the CFT quantum symmetry is already exhibited by the finite size integrable seams on the lattice. Importantly, fusion is implemented by simple matrix multiplication of square matrices associated with the integrable seams. Solving the boundary inversion relations for  $T_h(u)$  as in [26], it is seen that the boundary free energies of the s, a and b-type seams vanish whereas this is not the case for the general r-type seams. It would be of interest to find the boundary entropies of the seams as well as to find projectors onto the  $(s, \bar{s})$  and (a, b) sectors of T(u).

Here we only considered type I theories and focussed on the A series and the exceptional  $E_6$  theory of the critical unitary RSOS models on the torus, omitting details and proofs. We will publish details for all the critical A-D-E RSOS models elsewhere. Our new methods provide powerful tools to investigate the Ocneanu algebra of seams for other critical lattice models including models defined on the cylinder. Indeed, it would be of interest to apply the new methods to critical nonunitary A-D-E RSOS models [35], fused A-D-E RSOS models [36] such as the superconformal series [37, 38], as well as the critical dilute A-D-E RSOS lattice models [39–41] including the critical (A, G) series in branch 2 and tricritical (G, A) series in branch 1 [42].

#### Acknowledgments

The Australian Research Council (ARC) is acknowledged for support under Discovery Project DP200102316. This short article is intended for publication in MATRIX Annals. Work on this project was carried out as part of the MATRIX workshop Mathematics and Physics of Integrability (MPI2024): https://www.matrix-inst.org.au/events/mathematics-and-physics-of-integrability Wa thank Leap Perpart 7 when for comments on the menuscript.

We thank Jean-Bernard Zuber for comments on the manuscript.

# References

- [1] R.J. Baxter, Exactly Solved Models in Statistical Mechanics, Academic Press (1982).
- [2] G.E. Andrews, R.J. Baxter, P.J. Forrester, Eight-vertex SOS model and generalized Rogers-Ramanujan-type identities, J. Stat. Phys. 35 (1984) 193–266.
- [3] V. Pasquier, Two-dimensional critical systems labelled by Dynkin diagrams, Nucl. Phys. B285 (1987) 162–172.
- [4] V. Pasquier, Operator content of the ADE lattice models, J. Phys. A: Math. Gen. 20 (1987) 5707– 5717.
- [5] V. Pasquier, Lattice derivation of modular invariant partition functions on the torus, J. Phys. A: Math. Gen. 20 (1987) L1229–L1237.
- [6] P. Di Francesco, P. Mathieu, D. Sénéchal, Conformal Field Theory, Springer (1997).
- [7] A.A. Belavin, A.M. Polyakov, A.B. Zamolodchikov, Infinite conformal symmetry in twodimensional quantum field theory, Nucl. Phys. B241 (1984) 333–380.
- [8] P. Goddard, A. Kent, D. Olive, Virasoro algebras and coset space models, Phys. Lett. B152 (1985) 88–92.

- [9] G. Moore, N. Seiberg, Classical and quantum conformal field theory, Commun. Math. Phys. 123 (1989) 177–254.
- [10] R.E. Behrend, P.A. Pearce, V.B. Petkova, J.-B. Zuber, On the classification of bulk and boundary conformal field theories, Phys. Lett. B444 (1998) 163–166.
- [11] R.E. Behrend, P.A. Pearce, V.B. Petkova, J.-B. Zuber, Boundary conditions in rational conformal field theories, Nucl. Phys. B579 (2000) 707–773.
- [12] E. Verlinde, Fusion rules and modular transformations in 2D conformal field theory, Nucl. Phys. B300 (1988) 360–376.
- [13] R.E. Behrend, P.A. Pearce, Integrable and conformal boundary conditions for sl(2) A-D-E lattice models and unitary minimal conformal field theories, J. Stat. Phys. 102 (2001) 577–640.
- [14] V. Pasquier and H. Saleur, Common Structures Between Finite Systems and Conformal Field Theories Through Quantum Groups Nucl. Phys. B330 (1990) 523–556.
- [15] R.E. Behrend, P.A. Pearce, J.-B. Zuber, Integrable boundaries, conformal boundary conditions and A-D-E fusion rules, J. Phys. A: Math. Gen. 31 (1998) L763–770.
- [16] J.L. Cardy, Operator content of two-dimensional conformally invariant theories, Nucl. Phys. B270 (1986) 186–204.
- [17] A. Cappelli, C. Itzykson, J. B. Zuber. The A-D-E classification of minimal and A<sub>1</sub><sup>(1)</sup> conformal invariant theories, Commun. Math. Phys. **113** (1987) 1–26.
- [18] A. Ocneanu, Paths on Coxeter Diagrams: From Platonic Solids and Singularities to Minimal Models and Subfactors, in Lectures on Operator Theory, Fields Institute, Waterloo, Ontario, April 26–30, 1995, (Notes taken by S. Goto) Fields Institute Monographs, AMS 1999, Rajarama Bhat et al, eds; Quantum symmetries for SU(3) CFT Models, Lectures at Bariloche Summer School, Argentina, Jan 2000, AMS Contemporary Mathematics, R. Coquereaux, A. Garcia and R. Trinchero, eds; The classification of subgroups of quantum SU(N), Bariloche, 2000: https://www.cpt.univ-mrs.fr/~coque/Bariloche2000/Bariloche2000/Bariloche2000.html
- [19] V.B. Petkova, J.-B. Zuber, Generalised twisted partition functions, Physics Letters B504 (2001) 157–164; Conformal field theories, graphs and quantum algebras, MathPhys Odyssey 2001: Integrable Models and Beyond, 415–435.
- [20] R. Coquereaux, *Notes on the quantum tetrahedron*, Moscow Mathematical Journal 2 (2002) 41–80, arXiv:math-ph/0011006.
- [21] R. Coquereaux, G. Schieber, Twisted partition functions for ADE boundary conformal field theories and Ocneanu algebras of quantum symmetries, J. Geom. Phys. 42 (2002) 216–258.
- [22] R. Coquereaux, M. Huerta, Torus structure on graphs and twisted partition functions for minimal and affine models, J. Geom. Phys. 48 (2003) 580–634.
- [23] R. Coquereaux, R. Trinchero, On quntum symmetries of ADE graphs, Adv. Theor. Math. Phys. 8 (2004) 189–216.
- [24] C.H. Otto Chui, C. Mercat, W.P. Orrick, P.A. Pearce, Integrable lattice realizations of conformal twisted boundary conditions, Phys. Lett. B517 (2001) 429–435.

- [25] C.H. Otto Chui, C. Mercat, P.A. Pearce, *Integrable boundaries and universal TBA functional equations*, MathPhys Odyssey 2001: Integrable Models and Beyond, 391–413.
- [26] C.H. Otto Chui, C. Mercat, P.A. Pearce, Integrable and conformal twisted boundary conditions for sl(2) A-D-E lattice models, J. Phys. A: Math. Gen. 36 (2003) 2623–2662.
- [27] J. Belletête, A.M. Gainutdinov, J.L Jacobsen, H. Saleur, T.S. Tavares, *Topological defects in periodic RSOS models and anyonic chains*, arXiv:2003.11293 [math-ph] (2023); *Topological defects in lattice models and affine Temperley-Lieb algebra*, Commun. Math. Phys. 400 (2023) 1203–1254.
- [28] T.S. Tavares, M. Sinha, L. Grans-Samuelsson, A. Roy, H. Saleur, Integrable RG flows on topological defect lines in 2D conformal field theories, arXiv:2408.08241 [hep-th] (2024).
- [29] P. Goddard, A. Kent, and D. Olive, *Virasoro algebras and coset space models*, Phys. Lett. **B152** (1985) 88–92.
- [30] A. Klümper, P.A. Pearce, Analytic calculation of scaling dimensions: Tricritical hard squares and critical hard hexagons, J. Stst. Phys. 64 (1991) 13–76.
- [31] A. Feiguin, S. Trebst, A.W.W. Ludwig, M. Troyer, A. Kitaev, Z. Wang, and M.H. Freedman, *Interacting anyons in topological quantum liquids: The golden chain*, Phys. Rev. Lett. 98 (2007) 160409.
- [32] J. Heymann, T. Quella, *Revisiting the symmetry-resolved entanglement for non-invertible symmetries in* 1+1d conformal field theories, arXiv:2409.02315 [hep-th] (2024).
- [33] Wolfram Research, Mathematica Edition: Version 13, Wolfram Research Inc., Champaign, Illinois (2024).
- [34] S. Schäfer-Nameki, *ICTP lectures on (non-)invertible generalized symmetries*, Physics Reports **1063** (2024) 1–55.
- [35] P.J. Forrester, R.J. Baxter, *Further exact solutions of the eight-vertex SOS model and generalizations of the Rogers-Ramanujan identities*, J. Stat. Phys. **38** (1985) 435–472.
- [36] E. Date, M. Jimbo, A. Kuniba, T. Miwa, M. Okado, *Exactly solvable SOS models: I. Local height probabilities and theta function identities*, Nucl. Phys. B290 [FS20] (1987) 231–273; *Exactly solvable SOS models: II. Proof of star-triangle relation and combinatorial identities*, Adv. Stud. Pure Math. 16 (1988) 17–122.
- [37] P. Goddard, A. Kent, D. Olive, Unitary representations of the Virasoro and super-Virasoro algebras, Commun. Math. Phys. 103 (1986) 105–119.
- [38] C. Richard, P.A. Pearce, Integrable lattice realizations of N = 1 superconformal boundary conditions, Nucl. Phys. B631 (2002) 447–470.
- [39] S.O. Warnaar, B. Nienhuis, K.A. Seaton, New construction of solvable lattice models including an Ising model in a magnetic field, Phys. Rev. Lett. 69 (1992) 710–712.
- [40] P. Roche, On the construction of integrable dilute ADE lattice models, Phys. Lett. B285 (1992) 49–53.
- [41] S.O. Warnaar, P.A. Pearce, K.A. Seaton, B. Nienhuis, Order parameters of the dilute A models, J. Stat. Phys. 74 (1994) 469–531.
- [42] D.L. O'Brien, P.A. Pearce, Lattice realizations of unitary minimal modular invariant partition functions, J. Phys. A Math. Gen. 28 (1995) 4891–4905.