THE NUMBER OF SOLUTIONS OF A RANDOM SYSTEM OF POLYNOMIALS OVER A FINITE FIELD

RITIK JAIN

ABSTRACT. We study the probability distribution of the number of common zeros of a system of m random n-variate polynomials over a finite commutative ring R. We compute the expected number of common zeros of a system of polynomials over R. Then, in the case that R is a field, under a necessary-and-sufficient condition on the sample space, we show that the number of common zeros is binomially distributed.

1. INTRODUCTION

The problem of estimating the number of rational solutions of systems of polynomials over a finite field has a rich history, dating back to the Chevalley-Warning theorem [Che35], which characterizes the number of rational points on certain algebraic varieties. Among other interesting consequences, the Chevalley-Warning theorem implies that finite fields are quasi-algebraically closed (Artin's conjecture), and every conic over a finite field has a rational point [Ser73, Corollary 2.].

How can we study the number of rational points on an arbitrary variety over a finite field? While it might be possible to examine this global behavior using an algebrogeometric tool such as a moduli space, probability provides an elegant and flexible alternative framework. By studying the number of rational points on a suitably chosen, *random* variety, we can gain insights into the number of rational points on an arbitrary variety. Moreover, using limiting arguments, one can obtain deterministic results from a probabilistic setup.

In previous work, the author and collaborators computed the distribution of the number of zeros of a random multivariable polynomial over a finite field \mathbb{F}_q , sampled uniformly from the space $\mathbb{F}_q[\overline{X}_n]_{\overline{d}}$ of polynomials with degree at most d with respect to each variable, relying on elementary algebraic and combinatorial methods [JMW23].

In this paper, we consider several generalizations of the previous result. For one, we will consider *systems* of random multivariable polynomials that may have coefficients in finite commutative rings (which are not necessarily fields). We will also consider the "broadest possible" class of sample spaces over which our results hold.

Notation 1.1. Once and for all, \mathbb{N} denotes the set of positive integers, $\mathbb{Z}_{\geq 0}$ denotes $\mathbb{N} \cup \{0\}$, and $\mathcal{P}(S)$ denotes the power set of a set *S*. Moreover, we take \mathbb{F}_q to be the finite field of

order $q = p^r$ for some prime p and $r \in \mathbb{N}$, R to be a finite commutative ring with unity, and $R[\overline{X}_n]$ to be the space of polynomials in n variables x_1, \dots, x_n over R.

Now, we define the spaces which will be the setting of our discussion.

- **Definition 1.2.** (1) A free *R*-module $M \subseteq R[\overline{X}_n]$ is said to *extend R* if *R* is a proper subset of *M*.
 - (2) A vector space $V \subseteq \mathbb{F}_q[\overline{X}_n]$ is said to *contain functions* if every function $\mathbb{F}_q^n \to \mathbb{F}_q$ is represented by a polynomial in *V*.

The most natural examples of *R*-modules which extend *R* are (i) $R[\overline{X}_n]$ itself, (ii) the subspace of $R[\overline{X}_n]$ with total degree at most $d \ge 1$, and (iii) the subspace of $R[\overline{X}_n]$ with degree at most $d \ge 1$ with respect to each variable. On the other hand, some natural classes of vector spaces which contain functions are (i) $\mathbb{F}_q[\overline{X}_n]$, (ii) the subspace of $\mathbb{F}_q[\overline{X}_n]$ with total degree at most $d \ge n(q-1)$, and (iii) the subspace of $\mathbb{F}_q[\overline{X}_n]$ with degree at most $d \ge n(q-1)$, and (iii) the subspace of $\mathbb{F}_q[\overline{X}_n]$ with degree at most $d \ge n(q-1)$, and (iii) the subspace of $\mathbb{F}_q[\overline{X}_n]$ with degree at most $d \ge n(q-1)$, and (iii) the subspace of $\mathbb{F}_q[\overline{X}_n]$ with degree at most $d \ge n(q-1)$.

Definition 1.3. A point $\mathbf{p} \in \mathbb{R}^n$ is said to be a *zero* of a polynomial $f \in \mathbb{R}[\overline{X}_n]$ if $f(\mathbf{p}) = 0$. If $f_1(\mathbf{p}) = \cdots = f_m(\mathbf{p}) = 0$ for some $f_1, \cdots, f_m \in \mathbb{R}[\overline{X}_n]$, then \mathbf{p} is said to be a *solution* to the system of polynomials f_1, \cdots, f_m .

The following two theorems are our main contributions.

Theorem 1.4. Suppose $K \subseteq R[\overline{X}_n]$ extends R. Then, the average number of common zeros of a system of (not-necessarily distinct) random polynomials $f_1, \dots, f_m \in K$ is given by

 $|R|^{n-m}$.

Theorem 1.5. Let $V \subseteq \mathbb{F}_q[\overline{X}_n]$ be a vector space which contains functions. Then, the number of common zeros of a system of (not-necessarily distinct) random polynomials $f_1, \dots, f_m \in V$ is binomial with parameters q^n and $1/q^m$.

Corollary 1.6. Suppose $K \subseteq R[\overline{X}_n]$ extends R. Then, the average number of common zeros of a system of (not-necessarily distinct) random polynomials $f_1, \dots, f_n \in K$ is exactly 1.

Corollary 1.7. As $q \to \infty$, the number of common zeros of a system of n (not-necessarily distinct) random polynomials in $\mathbb{F}_q[\overline{X}_n]$ follows a Poisson distribution with parameter 1.

In Section 2, the probabilistic background for our discussion is rigorously established. We note that the meanings of Theorems 1.4 and 1.5 change slightly when K and V are finite versus when they have infinite cardinality. To be precise, in the latter case, the two theorems respectively concern *density* rather than probability. This change is necessitated by the fact that there is no uniform distribution on a countably infinite set. In Section 3, we will prove Theorem 1.4 using an algebraic and combinatorial argument. Finally, in Section 4, Theorem 1.5 is proven using elementary linear algebra and probability, and in Section 5, we summarize the main results and give some suggestions for future work.

Acknowledgement. The author would like to thank Han-Bom Moon for many helpful conversations, and his continued support in the writing of this paper.

2. PRELIMINARIES

In this section, we will outline the probabilistic framework for the remainder of our discussion.

For a free, finitely generated *R*-module $M \subseteq R[\overline{X}_n]$ with rank $k \in \mathbb{N}$, define

(1)
$$X_R(M, n, m) : M^m \to \mathbb{Z}_{\geq 0}$$

as the number of distinct zeros of a uniformly chosen set of polynomials $f_1, \dots, f_m \in M$. In this paper, we will compute the expectation of $X_R(M, n, m)$ when M extends R, and its distribution in the case that R is a field and M contains functions.

Let us first clarify the definition of $X_R(M, n, m)$. First, $\mathbb{Z}_{\geq 0}$ and M^m are equipped with the σ -algebras $\mathcal{P}(\mathbb{Z}_{\geq 0})$ and $\mathcal{P}(M^m)$ respectively, giving them the structure of measurable spaces. The probability measure μ on M^m is taken to be the *normalized counting measure*, defined as

$$\mu(A) = \frac{|A|}{|M^m|} = \frac{|A|}{|R|^{mk}}$$

for all subsets $A \subseteq M^m$.

This is a natural choice of measure for two reasons. For one, it is equivalent to sampling polynomials in M with iid uniform coefficients in R with respect to some choice of basis, which is a standard setup. Moreover, since a random polynomial in M is uniquely determined by its sequence of k coefficients, selecting m polynomials in M uniformly is equivalent to uniformly selecting a sequence in R^{mk} . It can be shown that R^{mk} is a compact and Hausdorff additive group, and the unique Haar probability measure on R^{mk} is given by the normalized counting measure.¹ Haar measures enjoy many regularity and invariance properties, thus μ is a desirable probability measure on M^m .

In order to consider the number of zeros of "uniformly random" polynomials when $M \subseteq R[\overline{X}_n]$ is infinitely generated, we must use a weaker measure known as a *density function*, a probability measure which is only finitely additive. This is because one cannot construct a uniform probability measure on a countably infinite set, and in particular, one cannot construct such a measure on any infinite subset of $R[\overline{X}_n]$.

A prototypical example of a density function is the *natural density function* $d_{nat} : \mathcal{P}(\mathbb{N}) \to [0, 1]$ on the set of positive integers, defined as

$$d_{\mathrm{nat}}(A) = \lim_{n \to \infty} \frac{|A \cap \{1, \cdots, n\}|}{n}$$

¹Here, R^{mk} is given the product topology with respect to R given the discrete topology.

for all subsets $A \subseteq \mathbb{N}$ for which the limit exists. In some sense, the natural density of a subset $A \subseteq \mathbb{N}$ can be understood as the probability of a "uniformly selected" $n \in \mathbb{N}$ having the defining property of A. For example, the natural density of even integers in \mathbb{N} is 1/2, and the natural density of squarefree integers in \mathbb{N} is $6/\pi^2$.

Letting $R[\overline{X}_n]_d$ denote the set of polynomials in $R[\overline{X}_n]$ with total degree at most d for some $d \in \mathbb{N}$, we can define an obvious density function $d : \mathcal{P}(R[\overline{X}_n]) \to [0, 1]$ as

(2)
$$\hat{d}(A) = \lim_{d \to \infty} \frac{|A \cap R[X_n]_d|}{|R[\overline{X}_n]_d|}$$

for all subsets $A \subseteq R[\overline{X}_n]$ for which the limit exists. We can generalize this setup to infinitely generated *R*-modules as follows.

Definition 2.1. For an infinitely generated *R*-module $M \subseteq R[\overline{X}_n]$, let $M_t \subseteq M$ be the *R*-module of polynomials in *M* with total degree at most *t* for all $t \in \mathbb{Z}_{\geq 0}$. Then, the *m*-density function $d_m : \mathcal{P}(M^m) \to [0, 1]$ is defined as

$$d_m(A) = \lim_{t \to \infty} \frac{|A \cap M_t^m|}{|M_t^m|} = \lim_{t \to \infty} \frac{|A \cap M_t^m|}{|R|^{mt}}$$

for all subsets $A \subseteq M^m$ for which the limit exists. The image of a subset $A \subseteq M^m$ under d_m is called its *density*.

Note that M_t is free and finitely generated, and if M extends R, so will the M_t for all $t \in \mathbb{N}$, and if M contains functions, then for some $\ell \in \mathbb{N}$, the M_t will contain functions for all $t \ge \ell$.

3. THE AVERAGE NUMBER OF ZEROS OVER A FINITE RING

In this section, we will compute the average number of zeros of a random system of polynomials in an *R*-module $M \subseteq R[X_n]$ that extends *R*.

Lemma 3.1. Suppose $K \subseteq R[\overline{X}_n]$ is a finitely generated *R*-module that extends *R*. Then, for each point $\mathbf{p} \in R$, the space

 $N_{\mathbf{p}} := \{ f \in K \mid f(\mathbf{p}) = 0 \}$

of polynomials in K which vanish at \mathbf{p} has cardinality $|R|^{k-1}$.

Proof. First, since *R* is contained in *K*, *K* contains the multiplicative identity 1_R of *R*. In particular, the linearly independent set $\{1_R\}$ can be extended to a basis

$$\mathcal{B}_K = \{1_R, r_1, \cdots, r_{k-1}\} \subset K$$

of *K*. Now, choose $c_1, \dots, c_{k-1} \in R$ such that, for all $1 \le i \le k-1$, $r_i - c_i$ vanishes at **0**. We claim that

$$\mathcal{B}'_{K} = \{1_{R}, r_{1} - c_{1}, \cdots, r_{k-1} - c_{k-1}\} \subset K$$

$$a_0 \cdot 1_R + \sum_{i=1}^{k-1} a_i (r_i - c_i) = 0$$

for some $a_0, \dots, a_{k-1} \in R$. Then, rearranging,

(3)
$$\left(a_0 - \sum_{i=1}^{k-1} a_i c_i\right) \cdot \mathbf{1}_R + \sum_{i=1}^{k-1} a_i r_i = 0$$

Since $1_R, r_1, \dots, r_{k-1}$ are linearly independent, (3) implies

$$a_0 - \sum_{i=1}^{k-1} a_i c_i = 0, \qquad a_1 = \dots = a_{k-1} = 0$$

But $\sum_{i=1}^{k-1} a_i c_i = 0$, so we also have $a_0 = 0$, therefore \mathcal{B}'_K is a linearly independent set with cardinality rank K = k. In particular, \mathcal{B}'_K is a basis of K, proving the claim.

Then, note that the *R*-module N_0 of polynomials which vanish at $\mathbf{0} := (0, \dots, 0)$ is generated by $\mathcal{B}'_K \setminus \{1_R\}$, so rank $N_0 = k - 1$. Since N_0 is an *R*-module, this implies

$$|N_0| = |R|^{k-1}$$

Then, for all $\mathbf{p} \in \mathbb{R}^n$, define

$$\phi_{\mathbf{p}}: N_{\mathbf{0}} \to N_{\mathbf{p}}, \qquad \phi_{\mathbf{p}}(f) := f - f(\mathbf{p}).$$

Since $\phi_{\mathbf{p}}$ has an inverse map $\phi_{\mathbf{p}}^{-1}(g) := g - g(\mathbf{0})$, it is bijective. Therefore, $|N_{\mathbf{p}}| = |R|^{k-1}$ for all $\mathbf{p} \in \mathbb{R}^{n}$.

Lemma 3.2. Let $K \subseteq R[\overline{X}_n]$ be a finitely generated *R*-module with rank *k* that extends *R*. For a random polynomial $f \in K$ and any point $\mathbf{p} \in R^n$, we have

$$P(f(\mathbf{p})=0) = \frac{1}{|R|}$$

Proof. Defining $N_{\mathbf{p}} := \{f \in K \mid f(\mathbf{p}) = 0\}$ as above, $|N_{\mathbf{p}}| = |R|^{k-1}$ by Lemma 3.1, we have

$$P(f(\mathbf{p}) = 0) = \frac{|N_{\mathbf{p}}|}{|K|} = \frac{|R|^{k-1}}{|R|^k} = \frac{1}{|R|}$$

as desired.

Lemma 3.3. Let $M \subseteq R[\overline{X}_n]$ be an infinitely generated *R*-module that extends *R*, and define $N_{\mathbf{p}} := \{f \in M \mid f(\mathbf{p}) = 0\}$. Then, for all $\mathbf{p} \in R$, the density of $N_{\mathbf{p}}$ in *M* is 1/|R|.

Proof. We have

$$d_1(N_{\mathbf{p}}) = \lim_{t \to \infty} \frac{|N_{\mathbf{p}} \cap M_t|}{|M_t|} = \lim_{t \to \infty} \frac{\{f \in M_t \mid f(\mathbf{p}) = 0\}|}{|R|^t} = \lim_{t \to \infty} \frac{R^{t-1}}{|R|^t} = \frac{1}{|R|}$$

by Lemma 3.2.

Theorem 3.4. Let $K \subseteq R[\overline{X}_n]$ be an *R*-module that extends *R*. Then, the average number of common zeros of a system of (not-necessarily distinct) random polynomials $f_1, \dots, f_m \in K$ is given by

$$|R|^{n-m}$$

Proof. For all $\mathbf{p} \in \mathbb{R}^n$, define the random variable

$$V_{\mathbf{p}} := \begin{cases} 1 & f_1(\mathbf{p}) = \dots = f_m(\mathbf{p}) = 0, \\ 0, & \text{otherwise.} \end{cases}$$

Since the polynomials f_1, \dots, f_m are selected independently, the events $f_i(\mathbf{p}) = 0$ and $f_j(\mathbf{p}) = 0$ are independent for all $i \neq j$, and

$$P(V_{\mathbf{p}} = 1) = \prod_{i=1}^{m} P(f_i(\mathbf{p}) = 0).$$

If *K* is finitely generated, then by Lemma 3.2,

$$P(f_i(\mathbf{p})=0) = \frac{1}{|R|},$$

and if K is infinitely generated, then by Lemma 3.3,

$$P(f_i(\mathbf{p}) = 0) = d_1(N_{\mathbf{p}}) = \frac{1}{|R|}$$

for all $1 \le i \le m$. Therefore, $P(V_p = 1) = 1/|R|^m$ in both cases.

Now, the number of common zeros of the f_i is given by $\sum_{\mathbf{p}\in R^n} V_{\mathbf{p}}$, thus by the linearity of expectation,

(4)
$$\mathbb{E}\left(\sum_{\mathbf{p}\in R^{n}}V_{\mathbf{p}}\right) = \sum_{\mathbf{p}\in R^{n}}\mathbb{E}\left(V_{\mathbf{p}}\right).$$

Since $\mathbb{E}(V_{\mathbf{p}}) = P(V_{\mathbf{p}} = 1) = 1/|R|^m$, (4) simplifies to

$$\sum_{\mathbf{p}\in R^n} \frac{1}{|R|^m} = |R|^{n-m},$$

and the result is obtained.

For all $0 \le r \le |R|^n$, let $A_r \subseteq K^m$ denote the set of *m*-tuples of polynomials with exactly *r* common zeros. If *K* is finitely generated, Theorem 3.4 states

$$\frac{1}{|R|^n + 1} \sum_{r=0}^{|R|^n} \mu(A_r) = |R|^{n-m},$$

6

$$\frac{1}{|R|^n + 1} \sum_{r=0}^{|R|^n} d_m(A_r) = |R|^{n-m}.$$

In the next section, we will obtain a stronger result by restricting to the case of *R* being a field. Namely, we will compute not only the expectation, but the distribution of the number of common zeros.

4. THE DISTRIBUTION OF THE NUMBER OF ZEROS OVER A FINITE FIELD

In this section, we will compute the distribution of the number of common zeros of a system of random polynomials f_1, \dots, f_m sampled from a vector space $V \subseteq \mathbb{F}_q[\overline{X}_n]$ which contains functions.

We denote the set of *n*-tuples in \mathbb{F}_q by

$$\mathbb{F}_q^n = \{\mathbf{p}_1, \cdots, \mathbf{p}_{q^n}\}.$$

Recall that a vector space $V \subseteq \mathbb{F}_q[\overline{X}_n]$ is said to contain functions if every function $\mathbb{F}_q^n \to \mathbb{F}_q$ is represented by a polynomial in *V*. After proving Lemma 4.1, we will give a necessary-and-sufficient criterion for this condition to hold.

Lemma 4.1. Every function $\mathbb{F}_q^n \to \mathbb{F}_q$ is uniquely represented by a polynomial in the space $\mathbb{F}_q[\overline{X}_n]_{\overline{q-1}} \subset \mathbb{F}_q[\overline{X}_n]$ of polynomials with degree at most q-1 with respect to each variable x_i .

Proof. Let *S* denote the set of all functions $\mathbb{F}_q^n \to \mathbb{F}_q$. We will show that the function $\varphi : \mathbb{F}_q[\overline{X}_n]_{\overline{q-1}} \to S$ is injective, where $\varphi(f) : \mathbb{F}_q^n \to \mathbb{F}_q$ is the function induced by a polynomial *f*. Suppose that $f, g \in \mathbb{F}_q[\overline{X}_n]_{\overline{q-1}}$ induce the same function. Then, f - g induces the zero function, that is, f - g is a member of the ideal

(5)
$$I := (x_1^q - x_1, \cdots, x_n^q - x_n).$$

But if $f - g \in I$, then the degree of f - g is at least q with respect to some variable x_i , or f - g = 0. Since every polynomial in the ring $\mathbb{F}_q[\overline{X}_n]_{\overline{q-1}}$ has degree at most q - 1 with respect to each variable, we have f - g = 0, or f = g, implying that φ is injective.

Now, $\mathbb{F}_q[\overline{X}_n]_{\overline{q-1}}$ has dimension $(q-1+1)^n = q^n$, therefore $|\mathbb{F}_q[\overline{X}_n]_{\overline{q-1}}| = q^{q^n}$. On the other hand, $|S| = |\mathbb{F}_q|^{|\mathbb{F}_q^n|} = q^{q^n}$, hence $|\mathbb{F}_q[\overline{X}_n]_{\overline{q-1}}| = |S|$. Since φ is an injective function between finite sets of equal cardinality, it must be a bijection.

In some sense, $\mathbb{F}_q[\overline{X}_n]_{\overline{q-1}}$ is the "smallest" vector space which contains functions.

Proposition 4.2. A vector space $V \subseteq \mathbb{F}_q[\overline{X}_n]$ contains functions if and only if for every $f \in \mathbb{F}_q[\overline{X}_n]_{\overline{q-1}}$, there exists a polynomial $g \in V$ such that $f - g \in I$, where I is defined as in (5).

Proof. First, if *V* contains functions, then for any $f \in \mathbb{F}_q[\overline{X}_n]_{\overline{q-1}}$, there exists a polynomial $g \in V$ which induces the same function $\mathbb{F}_q^n \to \mathbb{F}_q$ as f. Therefore, $f - g \in I$ as desired. On the other hand, if for every $f \in \mathbb{F}_q[\overline{X}_n]_{\overline{q-1}}$, there exists a polynomial $g \in V$ such that $f - g \in I$, then every function $f : \mathbb{F}_q^n \to \mathbb{F}_q$ is represented by $g \in V$.

Lemma 4.3. Let $r \leq q^n$ be a positive integer, and $V \subseteq \mathbb{F}_q[\overline{X}_n]$ be a vector space. The linear map

$$\phi: V \to \mathbb{F}_q^r \qquad \phi(f) = (f(\mathbf{p}_1), \cdots, f(\mathbf{p}_r))$$

is surjective exactly when V contains functions, where $\mathbf{p}_1, \cdots, \mathbf{p}_r$ are distinct points in \mathbb{F}_q^n .

Proof. First, assume *V* contains functions, and $r = q^n$. Then, by definition, *V* contains a polynomial representing every function $\mathbb{F}_q^n \to \mathbb{F}_q$. In particular, for any $(a_1, \dots, a_{q^n}) \in \mathbb{F}_q^{q^n}$, there exists a polynomial *f* such that

$$f(\mathbf{p}_i) = a_i$$

for all $1 \le i \le q^n$. It follows immediately that ϕ is surjective. On the other hand, assume that *V* does not contain functions. Then, there exists a function $\mathbb{F}_q^n \to \mathbb{F}_q$ which does not have a polynomial representative in *V*, hence ϕ cannot be surjective.

Now, suppose *V* contains functions, and $r < q^n$. For any point $(a_1, \dots, a_r) \in \mathbb{F}_q^r$, as shown above, there exists a polynomial $f \in V$ such that

$$(6) f(\mathbf{p}_i) = a_i$$

for all $1 \leq i \leq r$, and $f(\mathbf{p}) = \omega$ for all $\mathbf{p} \in \mathbb{F}_q^n \setminus {\mathbf{p}_1, \cdots, \mathbf{p}_r}$ and an arbitrary $\omega \in \mathbb{F}_q$. In particular, (6) implies that ϕ is surjective. On the other hand, assuming *V* does not contain functions, the map ϕ is not surjective when $r = q^n$, therefore ϕ is not surjective for all $r \leq q^n$. The result is obtained.

Lemma 4.4. Let $f \in V$ be a random polynomial sampled from a finite-dimensional vector space $V \subset \mathbb{F}_q[\overline{X}_n]$ which contains functions. Then, for any $\mathbf{p} \in \mathbb{F}_q^n$,

$$P\left(f(\mathbf{p})=0\right) = \frac{1}{q}.$$

Proof. For all $\mathbf{p} \in \mathbb{F}_q^n$, define the evaluation map

$$\operatorname{ev}_{\mathbf{p}}: V \to \mathbb{F}_q, \qquad \operatorname{ev}_{\mathbf{p}}(f) = f(\mathbf{p}).$$

Since the map ev_p is clearly surjective, the space ker ev_p of polynomials which vanish at p has dimension dim V - 1 by the rank-nullity theorem. Therefore,

$$P(f(\mathbf{p}) = 0) = \frac{|\ker ev_{\mathbf{p}}|}{|V|} = \frac{q^{\dim V - 1}}{q^{\dim V}} = \frac{1}{q}.$$

Lemma 4.5. Let $V \subset \mathbb{F}_q[\overline{X}_n]$ be a finite-dimensional vector space. Then, for any positive integer $r \leq q^n$, there are $q^{\dim V-r}$ polynomials in V which vanish at any r points $\mathbf{p}_1 \cdots \mathbf{p}_r \in \mathbb{F}_q^n$ if and only if V contains functions.

Proof. Note that the space of polynomials vanishing on $\mathbf{p}_1 \cdots \mathbf{p}_r \in \mathbb{F}_q^n$ is precisely the kernel of the map

$$V \to \mathbb{F}_q^r \qquad \phi(f) = (f(\mathbf{p}_1), \cdots, f(\mathbf{p}_r)).$$

By the rank-nullity theorem,

 $\dim \ker \phi = \dim V - \dim \operatorname{im} \phi.$

Now, by Lemma 4.3, we have

(7) $\dim \ker \phi = \dim V - r$

exactly when V contains functions. Finally, ker ϕ is a vector space over \mathbb{F}_q , so (7) implies

$$|\ker\phi| = q^{\dim V - i}$$

as desired.

The following corollary states that, if *f* is a random polynomial from a finite-dimensional vector space *V* which contains functions, then the events of *f* vanishing at any distinct points $\mathbf{p}_1, \dots, \mathbf{p}_r$ are independent.

Corollary 4.6. Let $V \subset \mathbb{F}_q[\overline{X}_n]$ be a finite-dimensional vector space. For a random polynomial $f \in V$, define E_j as the event of f vanishing on a fixed $\mathbf{p}_j \in \mathbb{F}_q^n$. Then, for any $I \subseteq \{1, \dots, q^n\}$, the events $\{E_i\}_{i \in I}$ are independent exactly when V contains functions.

Proof. By Lemma 4.5, for any $I := \{a_1, \dots, a_r\} \subseteq \{1, \dots, q^n\}$, we have

$$P(E_{a_1}, \cdots, E_{a_r}) = \frac{q^{\dim V - r}}{q^{\dim V}} = q^{-r},$$

and $P(E_{a_i}) = q^{-1}$ for any $a_i \in I$. Combining the two,

$$P(E_{a_1},\cdots,E_{a_r})=\prod_{i=1}^r P(E_{a_i}),$$

implying that the events $\{E_i\}_{i \in I}$ are independent as desired.

We are ready to prove the main result.

Theorem 4.7. Let $V \subseteq \mathbb{F}_q[\overline{X}_n]$ be a vector space which contains functions. Then, the number of common zeros of a system of (not-necessarily distinct) random polynomials $f_1, \dots, f_m \in V$ is binomial with parameters q^n and $1/q^m$.

Proof. We first assume that *V* is finite-dimensional. Let $\mathbb{F}_q^n = {\mathbf{p}_1, \cdots, \mathbf{p}_{q^n}}$, and for all $1 \le j \le q^n$, define the indicator random variable

$$X_j = \begin{cases} 1 & f_1(p_j) = \dots = f_m(p_j) = 0, \\ 0, & \text{otherwise.} \end{cases}$$

Note that the X_j are Bernoulli trials, where $X_j = 1$ if the f_1, \dots, f_m simultaneously vanish at \mathbf{p}_j , and $X_j = 0$ if they do not. We claim that the random variables X_1, \dots, X_{q^n} are iid with parameter $1/q^m$.

First, we will show that, for any $1 \le j \le q^n$, X_j has parameter $1/q^m$. By Lemma 4.4, each polynomial f_i vanishes at \mathbf{p}_j with probability 1/q. Since the f_i are chosen independently, we have

$$P(f_1(\mathbf{p}_j) = 0, \cdots, f_m(\mathbf{p}_j) = 0) = \prod_{i=1}^m P(f_i(\mathbf{p}_j)) = \frac{1}{q^m}$$

as desired. On the other hand, the X_i are also independent, since, by the same reasoning as before, the events of two distinct random polynomials f_{a_1}, f_{a_2} vanishing at any point $\mathbf{p} \in \mathbb{F}_q^n$ are independent of one another, and by Corollary 4.6, the events of a fixed random polynomial f_i vanishing at any set of points $\mathbf{p}_1, \dots, \mathbf{p}_r \in \mathbb{F}_q^n$ are independent. The claim is proven, and because

$$\sum_{j=1}^{q^n} X_i$$

is exactly the number of common zeros of f_1, \dots, f_m , the result is also shown.

Now, suppose that *V* is infinite-dimensional. For all $r \in \mathbb{Z}_{\geq 0}$, define the set

 $P_{m,r} = \{(f_1, \cdots, f_m) \in V^m \mid f_1, \cdots, f_m \text{ have } r \text{ common distinct zeros}\}.$

Note that

$$|P_{m,r} \cap V_t^m| = |\{(f_1, \cdots, f_m) \in V_t^m \mid f_1, \cdots, f_m \text{ have } r \text{ common zeros}\}|_{t=1}^{m}$$

therefore

$$d_m(P_{m,r}) = \lim_{t \to \infty} \frac{|P_{m,r} \cap V_t^m|}{|V_t^m|} = \lim_{t \to \infty} P(X_{\mathbb{F}_q}(V_t, n, m) = r).$$

By assumption, there exists some $\ell \in \mathbb{N}$ such that for all $n \ge \ell$, V_n contains functions, thus

$$\lim_{t \to \infty} P(X_{\mathbb{F}_q}(V_t, n, m) = r) = \binom{q^n}{r} (1/q^m)^r (1 - 1/q^m)^{q^n - r}$$

by the finite-dimensional case.

10

$$X_{\mathbb{F}_q}(V, n, m) \sim \operatorname{Bin}\left(q^n, \frac{1}{q^m}\right).$$

Proof. This is immediate from Theorem 4.7.

Corollary 4.9. As $q \to \infty$, the number of common zeros of a system of n (not-necessarily distinct) random polynomials in $\mathbb{F}_q[\overline{X}_n]$ follows a Poisson distribution with parameter 1.

Proof. By Corollary 4.8, as $q \to \infty$, the asymptotic density of the systems of polynomials in $\mathbb{F}_q[\overline{X}_n]^m$ with k solutions is given by

$$\lim_{q \to \infty} P\left(X_{\mathbb{F}_q}(\mathbb{F}_q[\overline{X_n}], n, m) = k\right) = \lim_{q \to \infty} \frac{e^{-q^{n-m}}q^{k(n-m)}}{k!} = \frac{e^{-1}}{k!}$$

as desired.

Remark 4.10. One may wonder if there exists a finite ring R which is not a field such that the distribution of $X_R(M, n, m)$ is binomial with parameters $|R|^n$ and $1/|R|^m$ for some $M \subseteq R[\overline{X}_n]$. The answer to this question is negative for the following reason. Unless R is a finite field, there will always be a function $R^n \to R$ which cannot be represented by a polynomial in $R[\overline{X}_n]$. For example, define the function $\eta : R \to R$ as

$$\eta(r) = \begin{cases} 0_R & r = 0_R, \\ 1_R, & r \neq 0_R. \end{cases}$$

If η were represented as a polynomial $\eta(x) = a_1 x + \cdots + a_n x^n$, we would have

$$\eta(r) = r(a_1 + \dots + a_n r^{n-1}) = 1_R$$

for any nonzero r. That is, every nonzero element in R would be invertible, making R a finite field by Wedderburn's Little Theorem [Fri99]. Therefore, there are no R-modules $M \subseteq R[\overline{X}_n]$ which contain functions unless R is a field.

5. CONCLUSION

We have shown that the average number of zeros of a system of m polynomials from a submodule $M \subset R[\overline{X}_n]$ properly containing R is equal to $|R|^{n-m}$. In the case that there are as many variables as equations, that is, when n = m, the average is 1, regardless of the order of the ring R. In the case that R is a field of order q and M contains functions, we have computed the distribution of the number of solutions. In particular, we have shown that it is a binomial distribution with parameters q^n , $1/q^m$. As a corollary, if n = m, the distribution converges to a Poisson distribution with parameter 1 as $q \to \infty$.

In the future, it will be interesting to explore applications of the above results in cryptography, where random polynomials over finite fields and rings are of great interest. As

an example, the security of many cryptography algorithms depend on the NP-completeness of the problem of finding solutions to a random multivariable system of polynomials over a finite field [LPTV24]. While the above results do not provide a deterministic method of finding the solutions to a system of polynomials, they provide a good heuristic for the number of solutions we can expect. For example, according to Theorem 4.7, a random system of three polynomials $f_1, f_2, f_3 \in \mathbb{F}_{19}[\overline{X}_2]$ has at most one solution with probability ≈ 0.9987 . Therefore, if we obtain a single solution to the system, it is more or less guaranteed that it is the only one.

Multivariable polynomials over finite fields and rings are also of pure mathematical interest, being linked to the factorization of random polynomials over \mathbb{Q} , polynomials over the *p*-adic integers \mathbb{Z}_p via the inverse limit $\lim_{k \to \infty} \mathbb{Z}/p^k \mathbb{Z} = \mathbb{Z}_p$, and the geometry of complex varieties. See [BV19, Shm21, Del74]. In the future, it will be interesting to explore these connections further.

It might be possible to obtain more geometrically natural specializations of Theorems 3.4 and 4.7. For example, what is the distribution of the number of \mathbb{F}_q -rational points on a random *complete intersection* $V \subset \mathbb{A}^n(\mathbb{F}_q)$? How many rational points can we expect on a projective variety $V \subset \mathbb{P}^n(\mathbb{F}_q)$ defined by m random homogenous polynomials $f_1, \dots, f_m \in \mathbb{F}_q[\overline{X_n}]$?

In [JMW23], the latter expectation was found to be

$$\frac{(q^{n+1}-1)\left(q^{\binom{n+d}{d}-1}-1\right)}{(q-1)\left(q^{\binom{n+d}{d}}-1\right)} \approx q^{n-1}$$

in the case of m = 1 polynomials, essentially relying on the argument given in the proof of Lemma 4.4. Can we more generally compute the expectation and probability distribution in the case of projective varieties for all $m \in \mathbb{N}$?

REFERENCES

- [BV19] Emmanuel Breuillard and Péter P. Varjú. Irreducibility of random polynomials of large degree. *Acta Math.* (2) 223 (2019), 195–249. 12
- [Che35] C. Chevalley. Démonstration d'une hypothèse de M. Artin. Abh. Math. Sem. Univ. Hamburg 11 (1935), 73–75. doi:10.1007/BF02940714. 1
- [Del74] Pierre Deligne. La conjecture de Weil. Publications Mathématiques de l'IHÉS (1974), 273–307. 12
- [Fri99] S. Frisch. Polynomial functions on finite commutative rings. In: D. E. Dobbs, M. Fontana, S.-E. Kabbaj (eds.), Advances in Commutative Ring Theory (Fes III Conf. 1997), Lecture Notes in Pure and Appl. Mathematics, vol. 205, Dekker, 1999, pp. 323–336. 11
- [JMW23] R. Jain, H.-B. Moon, and P. Wu. Distribution of the number of zeros of polynomials over a finite field. To appear in *Involve*. 1, 12
- [LPTV24] R. La Scala, F. Pintore, S. K. Tiwari, and A. Visconti. A multistep strategy for polynomial system solving over finite fields and a new algebraic attack on the stream cipher Trivium. *arXiv preprint* arXiv:2304.07820, 2024. https://arxiv.org/abs/2304.07820. 12

THE NUMBER OF SOLUTIONS OF A RANDOM SYSTEM OF POLYNOMIALS OVER A FINITE FIELD 13

- [Ser73] J.P. Serre. A Course in Arithmetic. Graduate Texts in Mathematics. Springer-Verlag, New York, 1973. 6 pp. 1
- [Shm21] Roy Shmueli. The Expected Number of Roots over The Field of p-adic Numbers. *International Mathematics Research Notices* (3) 2023 (2021), 2543–2571. 12