

Tempered space-time fractional negative binomial process

Shilpa^a, Ashok Kumar Pathak^{a,*}, Aditya Maheshwari^b

^aDepartment of Mathematics and Statistics, Central University of Punjab, Bathinda, India

^bOperations Management and Quantitative Techniques Area, Indian Institute of Management Indore, Indore, India

Abstract

In this paper, we define a tempered space-time fractional negative binomial process (TSTFNBP) by subordinating the fractional Poisson process with an independent tempered Mittag-Leffler Lévy subordinator. We study its distributional properties and its connection to partial differential equations. We derive the asymptotic behavior of its fractional order moments and long-range dependence property. It is shown that the TSTFNBP exhibits overdispersion. We also obtain some results related to the first-passage time.

Keywords: Fractional Poisson process, Tempered Mittag-Leffler subordinator, Fractional moments, LRD, PDEs.
2020 MSC: Primary 60G22, 60G51, Secondary 60G55, 60E05

1. Introduction

The applications of subordinated count processes in financial mathematics and actuarial sciences (see Rusakov and Yakubovich (2021), Gillespie (1999), Mendoza-Arriaga and Linetsky (2016)) have created an impetus for sustained enquiry in their theoretical research. In the last two decades, researchers have studied the fractional Poisson processes (FPPs) from a stochastic subordination point of view. In this direction, we find several varieties of the subordinated FPP being studied (see Meerschaert et al. (2011), Orsingher and Polito (2012), Maheshwari and Vellaisamy (2019), Gupta and Kumar (2023), Meoli (2023), Soni and Pathak (2024)). It has led to the development of rich literature in the domain. Several other practical applications of these processes can be found in various disciplines such as economics, finance, actuarial science, physics, infectious diseases modeling, and reliability (see Doukhan et al. (2002), Biard and Sausseureau (2014), Guler Dincer et al. (2022), Di Crescenzo and Meoli (2023), Soni et al. (2024)).

Following the subordination approach, we narrow our attention to the negative binomial (NB) process due to its advantages for modelling overdispersed data. The NB process is a time-changed Poisson process delayed by an independent gamma subordinator. Recently, several fractional variants of the NB process have been explored; for example, Vellaisamy and Maheshwari (2018) introduced the fractional NB process (FNBP) by subordinating the FPP with an independent gamma subordinator and studied its governing partial differential equations (PDEs), Beghin and Vellaisamy (2018) considered a space fractional NB process (SFNBP) and used it in biological modeling. Moreover, several tempered stable extensions of the NB process have been proposed and studied in the literature. The tempered stable subordinator is obtained by exponential tempering of the stable process, which exhibits heavy tail behavior of a stable process at short times and lighter tails at large times, additionally making all its moments finite. Maheshwari (2023) defined a tempered space fractional negative binomial process (TSFNBP) and explored its distributional and long-range dependence (LRD) properties. Recently, Soni et al. (2024) discussed applications of the tempered space fractional Poisson process to the reliability and bivariate shock models. This paper will consider a tempered space-time fractional negative binomial process using the stochastic subordination approach. In the following paragraph, we provide a brief description of the construction of the process and its important properties.

*Corresponding author

Email addresses: shilpa17garg@gmail.com (Shilpa), ashokiitb09@gmail.com (Ashok Kumar Pathak), adityam@iimidr.ac.in (Aditya Maheshwari)

The tempered Mittag-Leffler Lévy subordinator (see Kumar et al. (2019)) is defined as a tempered stable subordinated model delayed by an independent gamma subordinator. Its distribution is semi-heavy tailed, and it is an important feature for studying extreme phenomena. As a result, it can be used in place of the gamma subordinator to construct several time-varying stochastic processes. In this paper, we introduce the FPP time-changed by the tempered Mittag-Leffler Lévy process. We discuss several important characteristics of the process and derive various asymptotic results for the distributional properties and dependence structure. In particular, we derive the probability mass function (pmf) and discuss its connections with PDEs. We derive the asymptotic behavior of its fractional order moments and examine its dependence properties. Some results related to the first-passage time are also explored.

The structure of the article is as follows. In Section 2, we present some preliminary notations, definitions, and results. In Section 3, we define the TSTFNBP and discuss its main distributional characteristics, and we derive the asymptotic behavior of its fractional order moments. In Section 4, we study its dependence properties and some results related to the first-passage time.

2. Preliminaries

This section introduces some notations, definitions, elementary distributions, and results that will be used in the following sections. Let \mathbb{N} , \mathbb{R} , and \mathbb{C} denote the set of all natural, real, and complex numbers, respectively. Let $\mathbb{Z}_+ = \mathbb{N} \cup \{0\}$ denotes the set of all non-negative integers.

2.1. Definitions, some elementary distributions and results

(i) Let $f : [a, b] \subset \mathbb{R} \rightarrow \mathbb{R}$ be such that $f(t)$ is $(n + 1)$ times continuous differentiable for $n < \tau < n + 1$. Then, the Riemann-Liouville fractional derivative of order $\tau > 0$ is defined as (see Podlubny (1999))

$${}_a D_t^\tau f(t) = \left(\frac{d}{dt}\right)^{n+1} \int_a^t (t-u)^{n-\tau} f(u) du.$$

(ii) The generalized Wright function is defined by (Kilbas et al. (2002))

$${}_p \psi_q \left[z \left| \begin{matrix} (\alpha_i, \beta_i)_{1,p} \\ (a_j, b_j)_{1,q} \end{matrix} \right. \right] = \sum_{k=0}^{\infty} \frac{z^k}{k!} \frac{\prod_{i=1}^p \Gamma(\alpha_i + \beta_i k)}{\prod_{j=1}^q \Gamma(a_j + b_j k)}, \quad z, \alpha_i, a_i \in \mathbb{C} \text{ and } \beta_i, b_i \in \mathbb{R}, \quad (2.1)$$

(iii) For $0 < \beta < 1$, let $\{N_\beta(t, \lambda)\}_{t \geq 0}$ be a FPP having parameter $\lambda > 0$. Its one-dimensional distributions are (see Laskin (2003), Meerschaert et al. (2011))

$$p_\beta(n/t, \lambda) = P[N_\beta(t, \lambda) = n] = \frac{(\lambda t^\beta)^n}{n!} \sum_{k=0}^{\infty} \frac{(n+k)!}{k!} \frac{(-\lambda t^\beta)^k}{\Gamma(\beta(k+n)+1)}, \quad n \in \mathbb{Z}_+.$$

(iv) Let $\Upsilon(t) \sim G(\lambda_1, \beta_1 t)$. Its probability density function (pdf) is given by

$$f_G(x, t) = \frac{\lambda_1^{\beta_1 t}}{\Gamma(\beta_1 t)} x^{\beta_1 t - 1} e^{-\lambda_1 x}, \quad x > 0.$$

(v) For $\alpha \in (0, 1)$ and $\mu > 0$, let $S_{\alpha, \mu}(t)$ be a tempered α -stable subordinator (TSS). Then its pdf $g_{\alpha, \mu}(x, t)$ is given by (see Rosiński (2007))

$$g_{\alpha, \mu}(x, t) = e^{-\mu x + \mu^\alpha t} g_\alpha(x, t),$$

where $g_\alpha(x, t)$ is the pdf of α -stable subordinator (see Kumar and Vellaisamy (2015)).

(vi) The tempered Mittag-Leffler Lévy process (TMLLP) $M_{\alpha, \beta_1, \lambda_1, \mu}(t)$ is obtained by subordinating TSS with an independent gamma subordinator as $M_{\alpha, \beta_1, \lambda_1, \mu}(t) := S_{\alpha, \mu}(G_{\lambda_1, \beta_1}(t))$, $\alpha \in (0, 1)$, $\lambda_1, \mu, \beta_1 > 0$, $t \geq 0$. Its pdf $f_{M_{\alpha, \beta_1, \lambda_1, \mu}(t)}$ is given by (see Kumar et al. (2019))

$$f_{M_{\alpha, \beta_1, \lambda_1, \mu}(t, x)} = \lambda_1^{\beta_1 t} e^{-\mu x} \sum_{k=0}^{\infty} \frac{(-1)^k (\lambda_1 - \mu^\alpha)^k \Gamma(\beta_1 t + k) x^{\alpha(\beta_1 t + k) - 1}}{\Gamma(k+1) \Gamma(\beta_1 t) \Gamma(\alpha(\beta_1 t + k))}, \quad \lambda_1 > \mu^\alpha, x > 0 \quad (2.2)$$

The Lévy measure density $\pi(x)$ and the Laplace transform (LT) of the TMLLP are respectively

$$\pi(x) = \frac{\alpha\beta_1}{x} e^{-\mu x} E_{\alpha,1}[(\mu^\alpha - \lambda_1)x^\alpha], \quad \lambda_1 > \mu^\alpha, x > 0, \quad (2.3)$$

where two parameters Mittag-Leffler function $E_{\alpha,\beta}(z)$ is defined as (see Podlubny (1999))

$$E_{\alpha,\beta}(z) = \sum_{k=0}^{\infty} \frac{z^k}{\Gamma(\alpha k + \beta)}, \quad \alpha, \beta > 0 \quad (2.4)$$

and

$$\mathbb{E}\left[e^{-uM_{\alpha,\beta_1,\lambda_1,\mu}(t)}\right] = \left(\frac{\lambda_1}{\lambda_1 - \mu^\alpha + (\mu + u)^\alpha}\right)^{\beta_1 t}. \quad (2.5)$$

3. Tempered space-time fractional negative binomial process

In this section, we define the TSTFNBP process and derive its distributional properties

Definition 3.1. Let $\{N_\beta(t, \lambda)\}_{t \geq 0}$ be an FPP with parameter $\lambda > 0$. The tempered space-time fractional negative binomial process (TSTFNBP), denoted by $\{Q_{\alpha,\beta_1,\mu}^{\lambda_1,\beta}(t, \lambda)\}_{t \geq 0}$, which is obtained by the subordination of the FPP with an independent TMLLP (2.1 (vi)), that is,

$$Q_{\alpha,\beta_1,\mu}^{\lambda_1,\beta}(t, \lambda) := N_\beta(M_{\alpha,\beta_1,\lambda_1,\mu}(t), \lambda), \quad t \geq 0.$$

Let $\lambda_1 > \mu^\alpha$ and $y > 0$, the pmf of $\{Q_{\alpha,\beta_1,\mu}^{\lambda_1,\beta}(t, \lambda)\}_{t \geq 0}$, denoted by $p_{\alpha,\beta_1,\mu}^{\lambda_1,\beta}(n, t) = \mathbb{P}(Q_{\alpha,\beta_1,\mu}^{\lambda_1,\beta}(t, \lambda) = n)$ is derived as

$$\begin{aligned} p_{\alpha,\beta_1,\mu}^{\lambda_1,\beta}(n, t) &= \int_0^\infty p_\beta(n/y, \lambda) f_{M_{\alpha,\beta_1,\lambda_1,\mu}}(t, y) dy \\ &= \int_0^\infty \left(\frac{(\lambda y^\beta)^n}{n!} \sum_{k=0}^{\infty} \frac{(n+k)!}{k!} \frac{(-\lambda y^\beta)^k}{\Gamma(\beta(n+k)+1)} \right) f_{M_{\alpha,\beta_1,\lambda_1,\mu}}(t, y) dy \\ &= \frac{\lambda^n}{n!} \sum_{k=0}^{\infty} \frac{(n+k)!}{k!} \frac{(-\lambda)^k}{\Gamma(\beta(n+k)+1)} \int_0^\infty y^{\beta(n+k)} f_{M_{\alpha,\beta_1,\lambda_1,\mu}}(t, y) dy \\ &= \frac{\lambda^n}{n!} \sum_{k=0}^{\infty} \frac{(n+k)!}{k!} \frac{(-\lambda)^k}{\Gamma(\beta(n+k)+1)} \mathbb{E}\left[(M_{\alpha,\beta_1,\lambda_1,\mu}(t))^{\beta(n+k)}\right], \end{aligned}$$

as $\mathbb{E}\left[(M_{\alpha,\beta_1,\lambda_1,\mu}(t))^\rho\right] < \infty$ for all $\rho > 0$ (see Kumar et al. (2019)).

Remark 3.1. (i) When $\alpha = 1, \mu = 0$, the pmf of TSTFNBP reduces to

$$p_{1,\beta_1,0}^{\lambda_1,\beta}(n, t) = \frac{\lambda^n}{\lambda_1^{\beta n} n! \Gamma(\beta_1 t)} {}_2\psi_1 \left[\begin{matrix} -\lambda \\ \lambda_1^\beta \end{matrix} \middle| \begin{matrix} (n+1, 1) \\ (1+\beta n, \beta) \end{matrix} \right],$$

which is the pmf of the FNBP discussed in Vellaisamy and Maheshwari (2018). In addition, for $\beta = 1$, it leads to the pmf of the $\text{NB}(\beta_1 t, \frac{\lambda}{\lambda_1 + \lambda})$ of the form

$$p_{1,\beta_1,0}^{\lambda_1,1}(n, t) = \binom{n + \beta_1 t - 1}{n} \left(\frac{\lambda_1}{\lambda_1 + \lambda}\right)^{\beta_1 t} \left(\frac{\lambda}{\lambda_1 + \lambda}\right)^n,$$

as discussed in Vellaisamy and Maheshwari (2018).

(ii) When $\mu = 0$, the TSTFNBP corresponds to the generalized fractional negative binomial process defined in Soni and Pathak (2024).

Next, we obtained the governing fractional PDE for the pmf of TSTFNBP $\{Q_{\alpha, \beta_1, \mu}^{\lambda_1, \beta}(t, \lambda)\}_{t \geq 0}$. First, we proceed with the following lemma.

Lemma 3.1. (Vellaisamy and Maheshwari (2018)) For any $\tau \geq 1$, the governing fractional PDE of order τ for the gamma subordinator $\{\Upsilon(t)\}_{t \geq 0}$ is given by

$$\frac{\partial^\tau}{\partial t^\tau} f_G(x, t) = \beta_1 \frac{\partial^{\tau-1}}{\partial t^{\tau-1}} [\log \lambda_1 + \log x - \psi(\beta_1 t)] f_G(x, t), \quad x > 0 \text{ and } f_G(x, 0) = 0,$$

where $\psi(x)$ is the digamma function and $\frac{\partial^\tau}{\partial t^\tau}(\cdot)$ is the Riemann-Liouville fractional differential operator.

The next theorem gives the PDE with respect to time variable satisfying the pdf of the TMLLP.

Theorem 3.1. Let $g_{\alpha, \mu}(x, t)$ be the pdf of the TSS. Then the pdf of TMLLP satisfies the following fractional PDE

$$\frac{\partial^\tau}{\partial t^\tau} f_{M_{\alpha, \beta_1, \lambda_1, \mu}}(x, t) = \beta_1 \frac{\partial^{\tau-1}}{\partial t^{\tau-1}} \left[(\log \lambda_1 - \psi(\beta_1 t)) f_{M_{\alpha, \beta_1, \lambda_1, \mu}}(x, t) + \int_0^\infty g_{\alpha, \mu}(x, y) (\log y) f_G(y, t) dy \right], \quad x > 0, t > 0,$$

with $f_{M_{\alpha, \beta_1, \lambda_1, \mu}}(x, 0) = 0$.

Proof. Consider

$$f_{M_{\alpha, \beta_1, \lambda_1, \mu}}(x, t) = \int_0^\infty g_{\alpha, \mu}(x, y) f_G(y, t) dy.$$

Using the Riemann-Liouville fractional derivative, we obtain

$$\begin{aligned} \frac{\partial^\tau}{\partial t^\tau} f_{M_{\alpha, \beta_1, \lambda_1, \mu}}(x, t) &= \frac{\partial^\tau}{\partial t^\tau} \int_0^\infty g_{\alpha, \mu}(x, y) f_G(y, t) dy \\ &= \int_0^\infty g_{\alpha, \mu}(x, y) \frac{\partial^\tau}{\partial t^\tau} f_G(y, t) dy \\ &= \int_0^\infty g_{\alpha, \mu}(x, y) \left[\beta_1 \frac{\partial^{\tau-1}}{\partial t^{\tau-1}} [\log \lambda_1 + \log y - \psi(\beta_1 t)] f_G(y, t) \right] dy \quad (\text{using Lemma 3.1}) \\ &= \beta_1 \frac{\partial^{\tau-1}}{\partial t^{\tau-1}} \int_0^\infty g_{\alpha, \mu}(x, y) (\log \lambda_1 - \psi(\beta_1 t)) f_G(y, t) dy + \beta_1 \int_0^\infty g_{\alpha, \mu}(x, y) (\log y) \frac{\partial^{\tau-1}}{\partial t^{\tau-1}} f_G(y, t) dy. \quad \square \end{aligned}$$

With the help of Theorem 3.1, we can now obtain the following result.

Theorem 3.2. For $\tau \geq 1$, the pdf $p_{\alpha, \beta_1, \mu}^{\lambda_1, \beta}(n, t)$ of TSTFNBP satisfies

$$\frac{1}{\beta_1} \frac{\partial^\tau}{\partial t^\tau} p_{\alpha, \beta_1, \mu}^{\lambda_1, \beta}(n, t) = \frac{\partial^{\tau-1}}{\partial t^{\tau-1}} \left[(\log \lambda_1 - \psi(\beta_1 t)) p_{\alpha, \beta_1, \mu}^{\lambda_1, \beta}(n, t) + \int_0^\infty \int_0^\infty p_\beta(n/y_1, \lambda) g_{\alpha, \mu}(y_1, y) (\log y) f_G(y, t) dy dy_1 \right],$$

with $p_{\alpha, \beta_1, \mu}^{\lambda_1, \beta}(0, 0) = 1$.

Remark 3.2. When $\mu = 0$, the governing PDE of the TSTFNBP reduces to

$$\frac{1}{\beta_1} \frac{\partial^\tau}{\partial t^\tau} p_{\alpha, \beta_1, 0}^{\lambda_1, \beta}(n, t) = \frac{\partial^{\tau-1}}{\partial t^{\tau-1}} \left[(\log \lambda_1 - \psi(\beta_1 t)) p_{\alpha, \beta_1, 0}^{\lambda_1, \beta}(n, t) + \int_0^\infty \int_0^\infty p_\beta(n/y_1, \lambda) g_\alpha(y_1, y) (\log y) f_G(y, t) dy dy_1 \right], \text{ with } p_{\alpha, \beta_1, 0}^{\lambda_1, \beta}(0, 0) = 1,$$

as reported by Soni and Pathak (2024). Moreover, when $\beta = 1$, it corresponds to the PDE of the SFNB as studied in Beghin and Vellaisamy (2018).

Next, we discuss the asymptotic behavior for the moments for the TMLLP.

Theorem 3.3. Let $q > 0$, the asymptotic behavior of q^{th} order moments of TMLLP is given by

$$\mathbb{E}(M_{\alpha, \beta_1, \lambda_1, \mu}(t))^q \sim \left(\frac{\alpha \beta_1 \mu^{\alpha-1}}{\lambda_1} \right)^q t^q \text{ as } t \rightarrow \infty.$$

Proof. The proof of the theorem can be executed in two parts.

Case 1 – when q is integer: We have the following representation from Kumar et al. (2019)

$$\mathbb{E}(M_{\alpha,\beta_1,\lambda_1,\mu}(t))^q \sim \left(\frac{\alpha\beta_1\mu^{\alpha-1}}{\lambda_1}\right)^q t^q \text{ as } t \rightarrow \infty.$$

Case 2 – when q is non-integer: Assume $0 < q < 1$. With the help of (Kumar et al., 2019, eq. (6)), we have

$$\begin{aligned} \mathbb{E}\left[(M_{\alpha,\beta_1,\lambda_1,\mu}(t))^q\right] &= \frac{-1}{\Gamma(1-q)} \int_0^\infty \frac{d}{du} \left[\frac{\lambda_1}{\lambda_1 - \mu^\alpha + (\mu + u)^\alpha} \right]^{\beta_1 t} u^{-q} du \\ &= \frac{\alpha\beta_1 t \lambda_1^{\beta_1 t}}{\Gamma(1-q)} \int_0^\infty \frac{(u + \mu)^{\alpha-1} u^{-q}}{[\lambda_1 - \mu^\alpha + (\mu + u)^\alpha]^{\beta_1 t + 1}} du \\ &= \frac{\alpha\beta_1 t \lambda_1^{\beta_1 t}}{\Gamma(1-q)} \int_0^\infty \frac{(u + \mu)^{\alpha-1} u^{-q}}{[\lambda_1 - \mu^\alpha + (\mu + u)^\alpha]} e^{-\beta_1 t \ln[\lambda_1 - \mu^\alpha + (\mu + u)^\alpha]} du \\ &= \frac{\alpha\beta_1 t \lambda_1^{\beta_1 t}}{\Gamma(1-q)} \int_\mu^\infty \frac{z^{\alpha-1} (z - \mu)^{-q}}{[\lambda_1 - \mu^\alpha + z^\alpha]} e^{-\beta_1 t \ln[\lambda_1 - \mu^\alpha + z^\alpha]} dz \quad (\text{by letting } u + \mu = z). \end{aligned}$$

By taking $f(z) = \beta_1 \ln(\lambda_1 - \mu^\alpha + z^\alpha)$ and $g(z) = \frac{z^{\alpha-1} (z - \mu)^{-q}}{[\lambda_1 - \mu^\alpha + z^\alpha]}$, we have the Taylor series around μ of the form

$$\begin{aligned} f(z) &= \beta_1 \ln \lambda_1 + \frac{\beta_1 \alpha \mu^{\alpha-1}}{\lambda_1} (z - \mu) + \frac{\beta_1 \alpha \mu^{\alpha-2}}{\lambda_1^2} [(\alpha - 1) \lambda_1 - \alpha \mu^\alpha] (z - \mu)^2 + \dots \\ &= f(\mu) + \sum_{k=0}^{\infty} a_k (z - \mu)^{k+\delta}, \end{aligned}$$

where $\delta = 1$, $f(\mu) = \beta_1 \ln \lambda_1$, $a_0 = \frac{\beta_1 \alpha \mu^{\alpha-1}}{\lambda_1}$, and $a_1 = \frac{\beta_1 \alpha \mu^{\alpha-2}}{\lambda_1^2} [(\alpha - 1) \lambda_1 - \alpha \mu^\alpha]$. Additionally

$$\begin{aligned} g(z) &= (z - \mu)^{-q} \left[\frac{\mu^{\alpha-1}}{\lambda_1} + \frac{(\alpha - 1) \lambda_1 \mu^{\alpha-2} - \alpha \mu^{2\alpha-2}}{\lambda_1^2} (z - \mu) + \frac{\lambda_1^2 (\alpha - 1) (\alpha - 2) \mu^{\alpha-3} - 3\alpha (\alpha - 1) \mu^{2\alpha-3} \lambda_1 + 2\alpha^2 \mu^{2\alpha-3}}{\lambda_1^3} (z - \mu)^2 + \dots \right] \\ &= \sum_{k=0}^{\infty} b_k (z - \mu)^{k+\gamma-1}, \end{aligned}$$

where $\gamma = 1 - q$, $b_0 = \frac{\mu^{\alpha-1}}{\lambda_1}$, and $b_1 = \frac{(\alpha-1)\lambda_1\mu^{\alpha-2} - \alpha\mu^{2\alpha-2}}{\lambda_1^2}$.

Using Laplace-Erdelyi theorem (see (Kumar et al., 2019, Appendix A)), we have that

$$\int_\mu^\infty \frac{z^{\alpha-1} (z - \mu)^{-q}}{\lambda_1 - \mu^\alpha + z^\alpha} e^{-\beta_1 t \ln(\lambda_1 - \mu^\alpha + z^\alpha)} dz \sim e^{-\beta_1 t \ln \lambda_1} \sum_{k=0}^{\infty} \Gamma\left(\frac{k + \gamma}{\delta}\right) \frac{c_k}{t^{\frac{k+\gamma}{\delta}}} = \lambda_1^{-\beta_1 t} \sum_{k=0}^{\infty} \Gamma(k + 1 - q) \frac{c_k}{t^{k+1-q}}.$$

Hence, we obtain

$$\mathbb{E}\left[(M_{\alpha,\beta_1,\lambda_1,\mu}(t))^q\right] \sim \frac{\alpha\beta_1 t}{\Gamma(1-q)} \sum_{k=0}^{\infty} \Gamma(k + 1 - q) \frac{c_k}{t^{k+1-q}}, \quad (3.1)$$

where c_k in terms of coefficients a_k and b_k is given by

$$c_k = \frac{1}{a_0^{\frac{k+\gamma}{\delta}}} \sum_{j=0}^k b_{k-j} \sum_{i=0}^j \binom{-\frac{k+\gamma}{\delta}}{i} \frac{1}{a_0^i} \hat{B}_{(j,i)}(a_1, a_2, \dots, a_{j-i+1})$$

and $\hat{B}_{(j,i)}$ are the partial ordinary Bell polynomials (see Andrews (1998) and Soni et al. (2023)). The dominating term in (3.1), for large t , leads to

$$\mathbb{E} \left[(M_{\alpha, \beta_1, \lambda_1, \mu}(t))^q \right] \sim c_0 \alpha \beta_1 t^q,$$

where $c_0 = \left(\frac{\mu^{\alpha-1}}{\lambda_1} \right)^q \frac{1}{(\alpha \beta_1)^{1-q}}$. Hence

$$\mathbb{E}(M_{\alpha, \beta_1, \lambda_1, \mu}(t))^q \sim \left(\frac{\alpha \beta_1 \mu^{\alpha-1}}{\lambda_1} \right)^q t^q \text{ as } t \rightarrow \infty \text{ for } q \in (0, 1).$$

On similar lines, we obtain for general $q \in (n-1, n)$

$$\begin{aligned} \mathbb{E} \left[(M_{\alpha, \beta_1, \lambda_1, \mu}(t))^q \right] &= \frac{(-1)^n}{\Gamma(n-q)} \int_0^\infty \frac{d^n}{du^n} \left[\frac{\lambda_1}{\lambda_1 - \mu^\alpha + (\mu + u)^\alpha} \right]^{\beta_1 t} u^{n-q-1} du \\ &= \frac{(\alpha \beta_1 t)^n}{\Gamma(n-q)} \int_0^\infty \frac{\lambda_1^{\beta_1 t} (u + \mu)^{n(\alpha-1)}}{[\lambda_1 - \mu^\alpha + (\mu + u)^\alpha]^{\beta_1 t + n}} u^{n-q-1} du \\ &\sim \frac{(\alpha \beta_1 t)^n \lambda_1^{\beta_1 t}}{\Gamma(n-q)} \int_0^\infty \frac{(u + \mu)^{n(\alpha-1)} u^{n-q-1}}{[\lambda_1 - \mu^\alpha + (\mu + u)^\alpha]^n} e^{-\beta_1 t \ln(\lambda_1 - \mu^\alpha + (\mu + u)^\alpha)} du \\ &= \frac{(\alpha \beta_1 t)^n \lambda_1^{\beta_1 t}}{\Gamma(n-q)} \int_\mu^\infty \frac{z^{n(\alpha-1)} (z - \mu)^{n-q-1}}{[\lambda_1 - \mu^\alpha + z^\alpha]^n} e^{-\beta_1 t \ln(\lambda_1 - \mu^\alpha + z^\alpha)} dz \quad (\text{by letting } u + \mu = z). \end{aligned}$$

Let $f(z) = \beta_1 \ln(\lambda_1 - \mu^\alpha + z^\alpha)$ and $g(z) = (z - \mu)^{n-q-1} \left[\frac{z^{\alpha-1}}{\lambda_1 - \mu^\alpha + z^\alpha} \right]^n$. Then, we have

$$\begin{aligned} g(z) &= (z - \mu)^{n-q-1} \left[\left(\frac{\mu^{\alpha-1}}{\lambda_1} \right)^n + n \left(\frac{\mu^{\alpha-1}}{\lambda_1} \right)^{n-1} \frac{(\alpha-1)\lambda_1 \mu^{\alpha-2} - \alpha \mu^{2\alpha-2}}{\lambda_1^2} (z - \mu) + \dots \right] \\ &= \sum_{k=0}^{\infty} b_k (z - \mu)^{k+\gamma-1}, \end{aligned}$$

where $\gamma = n - q$, $b_0 = \left(\frac{\mu^{\alpha-1}}{\lambda_1} \right)^n$, and $b_1 = n \left(\frac{\mu^{\alpha-1}}{\lambda_1} \right)^{n-1} \frac{(\alpha-1)\lambda_1 \mu^{\alpha-2} - \alpha \mu^{2\alpha-2}}{\lambda_1^2}$. With the help of Laplace-Erdelyi theorem (see (Kumar et al., 2019, Appendix A)), we get

$$\int_\mu^\infty \frac{z^{n(\alpha-1)} (z - \mu)^{n-q-1}}{[\lambda_1 - \mu^\alpha + z^\alpha]^n} e^{-\beta_1 t \ln(\lambda_1 - \mu^\alpha + z^\alpha)} dz \sim e^{-t \beta_1 \ln \lambda_1} \sum_{k=0}^{\infty} \Gamma(k + n - q) \frac{d_k}{t^{k+n-q}}.$$

Hence, we have

$$\mathbb{E} \left[(M_{\alpha, \beta_1, \lambda_1, \mu}(t))^q \right] \sim \frac{(\alpha \beta_1 t)^n}{\Gamma(n-q)} \sum_{k=0}^{\infty} \Gamma(k + n - q) \frac{d_k}{t^{k+n-q}},$$

where d_k in terms of coefficient of a_k and b_k is given by

$$d_k = \frac{1}{\alpha_0^{\frac{k+\gamma}{\delta}}} \sum_{j=0}^k b_{k-j} \sum_{i=0}^j \binom{-\frac{k+\gamma}{\delta}}{i} \frac{1}{\alpha_0^i} \hat{B}_{(j,i)}(a_1, a_2, \dots, a_{j-i+1}).$$

The dominating term, for large t , in the above series corresponds to

$$\mathbb{E} \left[(M_{\alpha, \beta_1, \lambda_1, \mu}(t))^q \right] \sim d_0 \frac{(\alpha \beta_1 t)^n}{t^{n-q}},$$

where $d_0 = \left(\frac{\lambda_1}{\mu^{\alpha-1}}\right)^{-q} \frac{1}{(\alpha\beta_1)^{n-q}}$. Therefore

$$\mathbb{E}(M_{\alpha,\beta_1,\lambda_1,\mu}(t))^q \sim \left(\frac{\alpha\beta_1\mu^{\alpha-1}}{\lambda_1}\right)^q t^q \text{ as } t \rightarrow \infty \text{ for } q \in (n-1, n).$$

□

Next, we present the mean, variance, and autocovariance functions for the TSTFNBP.

3.1. Mean, variance, autocovariance and index of dispersion

Theorem 3.4. Let $0 < s \leq t < \infty$, the mean, variance, and autocovariance of the process $\{\mathcal{Q}_{\alpha,\beta_1,\mu}^{\lambda_1,\beta}(t, \lambda)\}_{t \geq 0}$ are given by

$$\begin{aligned} \text{(i)} \quad & \mathbb{E}\left[\mathcal{Q}_{\alpha,\beta_1,\mu}^{\lambda_1,\beta}(t, \lambda)\right] = q_1 \mathbb{E}\left[(M_{\alpha,\beta_1,\lambda_1,\mu}(t))^\beta\right] \sim q_1 \left(\frac{\alpha\beta_1\mu^{\alpha-1}t}{\lambda_1}\right)^\beta \\ \text{(ii)} \quad & \text{Var}\left[\mathcal{Q}_{\alpha,\beta_1,\mu}^{\lambda_1,\beta}(t, \lambda)\right] = q_1 \mathbb{E}\left[(M_{\alpha,\beta_1,\lambda_1,\mu}(t))^\beta\right] - q_1^2 \left[\mathbb{E}\left[(M_{\alpha,\beta_1,\lambda_1,\mu}(t))^\beta\right]\right]^2 + 2c_2 \mathbb{E}\left[(M_{\alpha,\beta_1,\lambda_1,\mu}(t))^{2\beta}\right] \\ \text{(iii)} \quad & \text{Cov}\left[\mathcal{Q}_{\alpha,\beta_1,\mu}^{\lambda_1,\beta}(s, \lambda), \mathcal{Q}_{\alpha,\beta_1,\mu}^{\lambda_1,\beta}(t, \lambda)\right] = q_1 \mathbb{E}\left[(M_{\alpha,\beta_1,\lambda_1,\mu}(s))^\beta\right] + c_1 \mathbb{E}\left[(M_{\alpha,\beta_1,\lambda_1,\mu}(s))^{2\beta}\right] - q_1^2 \mathbb{E}\left[(M_{\alpha,\beta_1,\lambda_1,\mu}(s))^\beta\right] \mathbb{E}\left[(M_{\alpha,\beta_1,\lambda_1,\mu}(t))^\beta\right] \\ & \quad + q_1^2 \beta \mathbb{E}\left[(M_{\alpha,\beta_1,\lambda_1,\mu}(t))^{2\beta} B\left(\beta, 1 + \beta; \frac{M_{\alpha,\beta_1,\lambda_1,\mu}(s)}{M_{\alpha,\beta_1,\lambda_1,\mu}(t)}\right)\right], \end{aligned}$$

where $q_1 = \frac{\lambda}{\Gamma(1+\beta)}$, $c_1 = \beta q_1^2 B(\beta, 1 + \beta)$, $c_2 = \frac{\lambda^2}{\Gamma(2\beta+1)}$, and $B(m, n; x) = \int_0^x t^{m-1} (1-t)^{n-1} dt$ for $0 < x < 1$ is an incomplete beta function.

Proof. We know that (see Laskin (2003))

$$\mathbb{E}\left[N_\beta(t, \lambda)\right] = q_1 t^\beta, \quad \text{Var}\left[N_\beta(t, \lambda)\right] = q_1 t^\beta \left[1 + q_1 t^\beta \left(\frac{\beta B(\beta, 1/2)}{2^{2\beta-1}} - 1\right)\right]. \quad (3.2)$$

Also, from Beghin and Orsingher (2009), we have that

$$\text{Cov}\left[N_\beta(s, \lambda), N_\beta(t, \lambda)\right] = q_1 s^\beta + c_1 s^{2\beta} + q_1^2 \left[\beta t^{2\beta} B(\beta, 1 + \beta; s/t) - (st)^\beta\right], \quad 0 < s \leq t,$$

$$\mathbb{E}\left[N_\beta(s, \lambda)N_\beta(t, \lambda)\right] = q_1 s^\beta + c_1 s^{2\beta} + q_1^2 \left[\beta t^{2\beta} B(\beta, 1 + \beta; s/t)\right],$$

and

$$\text{Var}\left[N_\beta(t, \lambda)\right] = q_1 t^\beta + \frac{(\lambda t^\beta)^2}{\beta} \left(\frac{1}{\Gamma(2\beta)} - \frac{1}{\beta \Gamma^2(\beta)}\right).$$

Using the conditioning argument and with the help of the above quantities for the FPP, we get

$$\begin{aligned} \mathbb{E}\left[\mathcal{Q}_{\alpha,\beta_1,\mu}^{\lambda_1,\beta}(t, \lambda)\right] &= \mathbb{E}\left[N_\beta(M_{\alpha,\beta_1,\lambda_1,\mu}(t))\right] = \mathbb{E}\left[N_\beta(M_{\alpha,\beta_1,\lambda_1,\mu}(t)) | M_{\alpha,\beta_1,\lambda_1,\mu}(t)\right] = q_1 \mathbb{E}\left[(M_{\alpha,\beta_1,\lambda_1,\mu}(t))^\beta\right] \\ \text{Var}\left[\mathcal{Q}_{\alpha,\beta_1,\mu}^{\lambda_1,\beta}(t, \lambda)\right] &= \text{Var}\left[\mathbb{E}\left[N_\beta(M_{\alpha,\beta_1,\lambda_1,\mu}(t)) | M_{\alpha,\beta_1,\lambda_1,\mu}(t)\right]\right] + \mathbb{E}\left[\text{Var}\left[N_\beta(M_{\alpha,\beta_1,\lambda_1,\mu}(t)) | M_{\alpha,\beta_1,\lambda_1,\mu}(t)\right]\right] \\ &= q_1 \mathbb{E}\left[(M_{\alpha,\beta_1,\lambda_1,\mu}(t))^\beta\right] - q_1^2 \left[\mathbb{E}\left[(M_{\alpha,\beta_1,\lambda_1,\mu}(t))^\beta\right]\right]^2 + 2c_2 \mathbb{E}\left[(M_{\alpha,\beta_1,\lambda_1,\mu}(t))^{2\beta}\right] \end{aligned}$$

and

$$\begin{aligned} \mathbb{E}\left[\mathcal{Q}_{\alpha,\beta_1,\mu}^{\lambda_1,\beta}(s, \lambda)\mathcal{Q}_{\alpha,\beta_1,\mu}^{\lambda_1,\beta}(t, \lambda)\right] &= \mathbb{E}\left[\mathbb{E}\left[N_\beta(M_{\alpha,\beta_1,\lambda_1,\mu}(s))N_\beta(M_{\alpha,\beta_1,\lambda_1,\mu}(t)) | M_{\alpha,\beta_1,\lambda_1,\mu}(s), M_{\alpha,\beta_1,\lambda_1,\mu}(t)\right]\right] \\ &= \mathbb{E}\left[q_1 (M_{\alpha,\beta_1,\lambda_1,\mu}(s))^\beta + c_1 (M_{\alpha,\beta_1,\lambda_1,\mu}(s))^{2\beta} + q_1^2 \beta (M_{\alpha,\beta_1,\lambda_1,\mu}(t))^{2\beta} B\left(\beta, 1 + \beta; \frac{M_{\alpha,\beta_1,\lambda_1,\mu}(s)}{M_{\alpha,\beta_1,\lambda_1,\mu}(t)}\right)\right]. \end{aligned}$$

Therefore, with the help of Part (i) and Part (ii), we get the desired expression for the Cov $\left[\mathcal{Q}_{\alpha,\beta_1,\mu}^{\lambda_1,\beta}(s, \lambda), \mathcal{Q}_{\alpha,\beta_1,\mu}^{\lambda_1,\beta}(t, \lambda)\right]$. □

Remark 3.3. Let $\{X(t)\}_{t \geq 0}$ be a stochastic process. We say it is overdispersed if $\text{Var}[X(t)] - \mathbb{E}[X(t)] > 0$ for all $t \geq 0$ (see (Cox and Lewis, 1966, p. 72)). Now, for the process $\{Q_{\alpha, \beta_1, \mu}^{\lambda_1, \beta}(t, \lambda)\}_{t \geq 0}$

$$\begin{aligned} \text{Var} \left[Q_{\alpha, \beta_1, \mu}^{\lambda_1, \beta}(t, \lambda) \right] - \mathbb{E} \left[Q_{\alpha, \beta_1, \mu}^{\lambda_1, \beta}(t, \lambda) \right] &= 2c_2 \mathbb{E} \left[(M_{\alpha, \beta_1, \lambda_1, \mu}(t))^{2\beta} \right] - q_1^2 \left[\mathbb{E} \left[(M_{\alpha, \beta_1, \lambda_1, \mu}(t))^\beta \right] \right]^2 \\ &= \frac{\lambda^2}{\beta} \left[\frac{\mathbb{E} \left[(M_{\alpha, \beta_1, \lambda_1, \mu}(t))^{2\beta} \right]}{\Gamma(2\beta)} - \frac{\left[\mathbb{E} \left[(M_{\alpha, \beta_1, \lambda_1, \mu}(t))^\beta \right] \right]^2}{\beta \Gamma^2(\beta)} \right] \\ &\geq \left(\mathbb{E} \left[(M_{\alpha, \beta_1, \lambda_1, \mu}(t))^\beta \right] \right)^2 \left[\frac{\lambda^2}{\beta} \left(\frac{1}{\Gamma(2\beta)} - \frac{1}{\beta \Gamma^2(\beta)} \right) \right]. \end{aligned}$$

One can observe that $\frac{\lambda^2}{\beta} \left(\frac{1}{\Gamma(2\beta)} - \frac{1}{\beta \Gamma^2(\beta)} \right) > 0$ for $\lambda > 0$ and $\beta \in (0, 1)$ (see Beghin and Macci (2014)) and $\left[\mathbb{E} \left[(M_{\alpha, \beta_1, \lambda_1, \mu}(t))^\beta \right] \right]^2 \leq \mathbb{E} \left[(M_{\alpha, \beta_1, \lambda_1, \mu}(t))^{2\beta} \right]$ is true because of Cauchy-Schwarz inequality and shows the overdispersion of the $\{Q_{\alpha, \beta_1, \mu}^{\lambda_1, \beta}(t, \lambda)\}_{t \geq 0}$.

3.2. Laplace transform

Let $h(x, t)$ be the pdf of the $\mathcal{E}_\beta(M_{\alpha, \beta_1, \lambda_1, \mu}(t))$ and $k_\beta(x, t)$ be the pdf of the inverse stable subordinator $\mathcal{E}_\beta(t)$ with LT $\mathbb{E}[e^{-u\mathcal{E}_\beta(t)}] = E_{\beta, 1}^1(-u^\beta)$ (see Meerschaert and Straka (2013)). Then, the LT of $\mathcal{E}_\beta(M_{\alpha, \beta_1, \lambda_1, \mu}(t))$ can be derived as

$$\mathbb{E} \left[e^{-u\mathcal{E}_\beta(M_{\alpha, \beta_1, \lambda_1, \mu}(t))} \right] = \int_0^\infty e^{-ux} h(x, t) dx = \int_0^\infty \int_0^\infty e^{-ux} k_\beta(x, y) f_{M_{\alpha, \beta_1, \lambda_1, \mu}}(t, y) dy dx \quad (3.3)$$

$$\begin{aligned} &= \int_0^\infty E_{\beta, 1}^1(-uy^\beta) f_{M_{\alpha, \beta_1, \lambda_1, \mu}}(t, y) dy \\ &= \sum_{l=0}^\infty \frac{(-u)^l}{\Gamma(1 + \beta l)} \int_0^\infty y^{\beta l} f_{M_{\alpha, \beta_1, \lambda_1, \mu}}(t, y) dy \\ &= \sum_{l=0}^\infty \frac{(-u)^l}{\Gamma(1 + \beta l)} \mathbb{E} \left[(M_{\alpha, \beta_1, \lambda_1, \mu}(t))^{\beta l} \right]. \end{aligned} \quad (3.4)$$

Using the conditioning arguments, we obtain the LT for TSTFNBP as

$$\begin{aligned} \mathbb{E} \left[e^{-uQ_{\alpha, \beta_1, \mu}^{\lambda_1, \beta}(t, \lambda)} \right] &= \mathbb{E} \left[\mathbb{E} \left[e^{-uN(\mathcal{E}_\beta(M_{\alpha, \beta_1, \lambda_1, \mu}(t)))} | \mathcal{E}_\beta(M_{\alpha, \beta_1, \lambda_1, \mu}(t)) \right] \right] \\ &= \mathbb{E} \left[\mathbb{E} \left[\exp(-\lambda \mathcal{E}_\beta(M_{\alpha, \beta_1, \lambda_1, \mu}(t)) (1 - e^{-u})) / \mathcal{E}_\beta(M_{\alpha, \beta_1, \lambda_1, \mu}(t)) \right] \right] \\ &= \sum_{l=0}^\infty \frac{(-\lambda(1 - e^{-u}))^l}{\Gamma(1 + \beta l)} \mathbb{E} \left[(M_{\alpha, \beta_1, \lambda_1, \mu}(t))^{\beta l} \right]. \end{aligned}$$

The probability generating function of the process $\{Q_{\alpha, \beta_1, \mu}^{\lambda_1, \beta}(t, \lambda)\}_{t \geq 0}$ can be evaluated using LT and is given by

$$\mathbb{E} \left[u^{Q_{\alpha, \beta_1, \mu}^{\lambda_1, \beta}(t, \lambda)} \right] = \sum_{l=0}^\infty \frac{(-\lambda(1 - u))^l}{\Gamma(1 + \beta l)} \mathbb{E} \left[(M_{\alpha, \beta_1, \lambda_1, \mu}(t))^{\beta l} \right].$$

Remark 3.4. When $\alpha = 1, \mu = 0$, the LT of TSTFNBP reduces to

$$\mathbb{E} \left[e^{-uQ_{1, \beta_1, 0}^{\lambda_1, \beta}(t, \lambda)} \right] = \frac{1}{\Gamma(\beta_1 t)} {}_2\psi_1 \left[\frac{-\lambda(1 - e^{-u})}{\lambda_1^\beta} \middle| (1, 1), (\beta_1 t, \beta) \right],$$

which is the LT of the FNBP discussed in Vellaisamy and Maheshwari (2018).

Next, we present the Lévy measure density for the particular case when $\beta = 1$.

3.3. Lévy measure

It is to note that when $\beta = 1$, the TSTFNBP is identical to the TSFNBP defined in Maheshwari (2023). We derive here the Lévy measure density for TSFNBP. Using Lévy density of TMLLP (2.3) and the formula (Ken-Iti, 1999, page 197), the Lévy measure \mathcal{D} for the process $\{Q_{\alpha,\beta_1,\mu}^{\lambda_1,1}(t, \lambda)\}_{t \geq 0}$ can be evaluated as

$$\begin{aligned}
\mathcal{D}(k) &= \int_0^\infty \sum_{i=1}^\infty p_\beta(i/t, \lambda) \delta_{\{i\}}(k) \pi(t) dt \\
&= \int_0^\infty \sum_{i=1}^\infty \frac{(\lambda t)^i}{i!} e^{-\lambda t} \delta_{\{i\}}(k) \frac{\alpha \beta_1}{t} e^{-\mu t} E_{\alpha,1}[(\mu^\alpha - \lambda_1)t^\alpha] dt \\
&= \alpha \beta_1 \sum_{i=1}^\infty \frac{(\lambda)^i}{i!} \delta_{\{i\}}(k) \int_0^\infty e^{-(\mu+\lambda)t} t^{i-1} \sum_{j=0}^\infty \frac{[(\mu^\alpha - \lambda_1)t^\alpha]^j}{\Gamma(\alpha j + 1)} dt \\
&= \alpha \beta_1 \sum_{i=1}^\infty \frac{(\lambda)^i}{i!} \delta_{\{i\}}(k) \sum_{j=0}^\infty \frac{(\mu^\alpha - \lambda_1)^j}{\Gamma(\alpha j + 1)} \int_0^\infty e^{-(\mu+\lambda)t} t^{i-1+\alpha j} dt \\
&= \alpha \beta_1 \sum_{i=1}^\infty \frac{(\lambda)^i}{i!} \delta_{\{i\}}(k) \sum_{j=0}^\infty \frac{(\mu^\alpha - \lambda_1)^j}{\Gamma(\alpha j + 1)} \frac{\Gamma(\alpha j + i)}{(\lambda + \mu)^{\alpha j + i}}. \tag{3.5}
\end{aligned}$$

4. Second-order asymptotic properties and first-passage time

In this section, we will discuss the long-range dependence and first-passage time for the TSTFNBP. We first reproduce the definition of the LRD property (see D'Ovidio and Nane (2014), Maheshwari and Vellaisamy (2016), Kumar et al. (2020)).

Definition 4.1. Let $0 < s < t$, let the correlation function $\text{Corr}[X(s), X(t)]$ for a stochastic process $\{X(t)\}_{t \geq 0}$ satisfies the following relation

$$\lim_{t \rightarrow \infty} \frac{\text{Corr}[X(s), X(t)]}{t^{-d}} = k(s),$$

for some $k(s) > 0$ and $d > 0$. The process $\{X(t)\}_{t \geq 0}$ exhibits the long-range dependence (LRD) property when $d \in (0, 1)$.

4.1. Dependence structure of the TSTFNBP

Lemma 4.1. Let $\beta \in (0, 1)$ and $0 < s < t$, s is fixed. Then the following asymptotic expansion holds for a large t .

- (i) $\mathbb{E}[(M_{\alpha,\beta_1,\lambda_1,\mu}(s))^\beta (M_{\alpha,\beta_1,\lambda_1,\mu}(t))^\beta] \sim \mathbb{E}[(M_{\alpha,\beta_1,\lambda_1,\mu}(s))^\beta] \mathbb{E}[(M_{\alpha,\beta_1,\lambda_1,\mu}(t-s))^\beta]$
- (ii) $\beta \mathbb{E}[(M_{\alpha,\beta_1,\lambda_1,\mu}(t))^{2\beta}] \sim \mathbb{E}[(M_{\alpha,\beta_1,\lambda_1,\mu}(s))^{2\beta}] \mathbb{E}[(M_{\alpha,\beta_1,\lambda_1,\mu}(t-s))^{2\beta}]$.

Proof. Proof of the following lemma is similar to that of Lemma 2 in Maheshwari and Vellaisamy (2016). \square

We will next proof the LRD property for our process.

Theorem 4.1. The process TSTFNBP exhibits the LRD property.

Proof. Using Theorem 3.3 and with the help of Lemma 4.1(ii), the asymptotic behavior of the covariance is

$$\begin{aligned}
\text{Cov} [Q_{\alpha,\beta_1,\mu}^{\lambda_1,\beta}(s, \lambda), Q_{\alpha,\beta_1,\mu}^{\lambda_1,\beta}(t, \lambda)] &\sim q_1 \mathbb{E}[(M_{\alpha,\beta_1,\lambda_1,\mu}(s))^\beta] + c_1 \mathbb{E}[(M_{\alpha,\beta_1,\lambda_1,\mu}(s))^{2\beta}] \\
&\quad - q_1^2 \mathbb{E}[(M_{\alpha,\beta_1,\lambda_1,\mu}(s))^\beta] \left[\mathbb{E}[(M_{\alpha,\beta_1,\lambda_1,\mu}(t))^\beta] - \mathbb{E}[(M_{\alpha,\beta_1,\lambda_1,\mu}(t-s))^\beta] \right] \\
&\sim q_1 \mathbb{E}[(M_{\alpha,\beta_1,\lambda_1,\mu}(s))^\beta] + c_1 \mathbb{E}[(M_{\alpha,\beta_1,\lambda_1,\mu}(s))^{2\beta}] \\
&\quad - q_1^2 \mathbb{E}[(M_{\alpha,\beta_1,\lambda_1,\mu}(s))^\beta] \left[\left(\frac{\alpha \beta_1 \mu^{\alpha-1} t}{\lambda_1} \right)^\beta - \left(\frac{\alpha \beta_1 \mu^{\alpha-1} (t-s)}{\lambda_1} \right)^\beta \right] \\
&\sim q_1 \mathbb{E}[(M_{\alpha,\beta_1,\lambda_1,\mu}(s))^\beta] + c_1 \mathbb{E}[(M_{\alpha,\beta_1,\lambda_1,\mu}(s))^{2\beta}] \quad (\text{since } t^\beta - (t-s)^\beta \sim \beta s t^{\beta-1} \text{ for large value of } t).
\end{aligned}$$

Also, the asymptotic behavior of the variance is as follows

$$\begin{aligned}
\text{Var} \left[\mathcal{Q}_{\alpha, \beta_1, \mu}^{\lambda_1, \beta}(t, \lambda) \right] &\sim q_1 \left(\frac{\alpha \beta_1 \mu^{\alpha-1} t}{\lambda_1} \right)^\beta - q_1^2 \left(\frac{\alpha \beta_1 \mu^{\alpha-1} t}{\lambda_1} \right)^{2\beta} + 2c_2 \left(\frac{\alpha \beta_1 \mu^{\alpha-1} t}{\lambda_1} \right)^{2\beta} \\
&= t^{2\beta} \left[q_1 \left(\frac{\alpha \beta_1 \mu^{\alpha-1}}{\lambda_1 t} \right)^\beta - q_1^2 \left(\frac{\alpha \beta_1 \mu^{\alpha-1}}{\lambda_1} \right)^{2\beta} + 2c_2 \left(\frac{\alpha \beta_1 \mu^{\alpha-1}}{\lambda_1} \right)^{2\beta} \right] \\
&\sim t^{2\beta} \left(\frac{\alpha \beta_1 \mu^{\alpha-1}}{\lambda_1} \right)^{2\beta} (2c_2 - q_1^2) \\
&\sim t^{2\beta} d_1,
\end{aligned}$$

where $d_1 = \left(\frac{\alpha \beta_1 \mu^{\alpha-1}}{\lambda_1} \right)^{2\beta} (2c_2 - q_1^2)$. Therefore, the correlation function can be computed as

$$\begin{aligned}
\text{Corr} \left[\mathcal{Q}_{\alpha, \beta_1, \mu}^{\lambda_1, \beta}(s, \lambda), \mathcal{Q}_{\alpha, \beta_1, \mu}^{\lambda_1, \beta}(t, \lambda) \right] &= \frac{\text{Cov} \left[\mathcal{Q}_{\alpha, \beta_1, \mu}^{\lambda_1, \beta}(s, \lambda), \mathcal{Q}_{\alpha, \beta_1, \mu}^{\lambda_1, \beta}(t, \lambda) \right]}{\sqrt{\text{Var} \left[\mathcal{Q}_{\alpha, \beta_1, \mu}^{\lambda_1, \beta}(s, \lambda) \right]} \sqrt{\text{Var} \left[\mathcal{Q}_{\alpha, \beta_1, \mu}^{\lambda_1, \beta}(t, \lambda) \right]}} \sim \frac{q_1 \mathbb{E} \left[(M_{\alpha, \beta_1, \lambda_1, \mu}(s))^\beta \right] + c_1 \mathbb{E} \left[(M_{\alpha, \beta_1, \lambda_1, \mu}(s))^{2\beta} \right]}{\sqrt{t^{2\beta} d_1} \sqrt{\text{Var} \left[\mathcal{Q}_{\alpha, \beta_1, \mu}^{\lambda_1, \beta}(t, \lambda) \right]}} \\
&= t^{-\beta} \left[\frac{q_1 \mathbb{E} \left[(M_{\alpha, \beta_1, \lambda_1, \mu}(s))^\beta \right] + c_1 \mathbb{E} \left[(M_{\alpha, \beta_1, \lambda_1, \mu}(s))^{2\beta} \right]}{\sqrt{d_1 \text{Var} \left[\mathcal{Q}_{\alpha, \beta_1, \mu}^{\lambda_1, \beta}(t, \lambda) \right]}} \right].
\end{aligned}$$

Hence, for $0 < \beta < 1$ and the decaying power $t^{-\beta}$, the process shows the LRD property. \square

4.2. First-passage time distribution

Finally, we look at the first-passage time distribution of the TSTFNBP. For a stochastic process, it is the time during which a process reaches a certain threshold for the first time.

Let \mathcal{T}_k be the time of first upcrossing of level k and is defined as $\mathcal{T}_k := \inf\{t \geq 0 : \mathcal{Q}_{\alpha, \beta_1, \mu}^{\lambda_1, \beta}(t, \lambda) \geq k\}$. Then the survival function $\Pr\{\mathcal{T}_k > t\}$ can be derived as

$$\begin{aligned}
\Pr\{\mathcal{T}_k > t\} &= \Pr\{\mathcal{Q}_{\alpha, \beta_1, \mu}^{\lambda_1, \beta}(t, \lambda) < k\} = \sum_{n=0}^{k-1} \Pr\{\mathcal{Q}_{\alpha, \beta_1, \mu}^{\lambda_1, \beta}(t, \lambda) = n\} \\
&= \sum_{n=0}^{k-1} \int_0^\infty p_\beta(n/y, \lambda) f_{M_{\alpha, \beta_1, \lambda_1, \mu}(t, y)} dy \\
&= \sum_{n=0}^{k-1} \int_0^\infty \frac{(\lambda y^\beta)^n}{n!} \sum_{l=0}^\infty \frac{(n+l)!}{l!} \frac{(-\lambda y^\beta)^l}{\Gamma(\beta(l+n)+1)} f_{M_{\alpha, \beta_1, \lambda_1, \mu}(t, y)} dy \\
&= \sum_{n=0}^{k-1} \frac{\lambda^n}{n!} \sum_{l=0}^\infty \frac{(n+l)!}{l!} \frac{(-\lambda)^l}{\Gamma(\beta(l+n)+1)} \int_0^\infty y^{\beta(n+l)} f_{M_{\alpha, \beta_1, \lambda_1, \mu}(t, y)} dy \\
&= \sum_{n=0}^{k-1} \frac{\lambda^n}{n!} \sum_{l=0}^\infty \frac{(n+l)!}{l!} \frac{(-\lambda)^l}{\Gamma(\beta(l+n)+1)} \mathbb{E}[(M_{\alpha, \beta_1, \lambda_1, \mu}(t))^{\beta(n+l)}].
\end{aligned}$$

Furthermore, the distribution of \mathcal{T}_k can be written as

$$\begin{aligned}
\Pr\{\mathcal{T}_k < t\} &= \Pr\{\mathcal{Q}_{\alpha, \beta_1, \mu}^{\lambda_1, \beta}(t, \lambda) \geq k\} = \sum_{n=k}^\infty \Pr\{\mathcal{Q}_{\alpha, \beta_1, \mu}^{\lambda_1, \beta}(t, \lambda) = n\} \\
&= \sum_{n=k}^\infty \frac{\lambda^n}{n!} \sum_{l=0}^\infty \frac{(n+l)!}{l!} \frac{(-\lambda)^l}{\Gamma(\beta(l+n)+1)} \mathbb{E}[(M_{\alpha, \beta_1, \lambda_1, \mu}(t))^{\beta(n+l)}].
\end{aligned}$$

Therefore, the density function $\mathcal{P}(n, t) = Pr\{\mathcal{T}_k \in dt\}/dt$ is

$$\begin{aligned}\mathcal{P}(n, t) &= \frac{d}{dt} \sum_{n=k}^{\infty} \frac{\lambda^n}{n!} \sum_{l=0}^{\infty} \frac{(n+l)!}{l!} \frac{(-\lambda)^l}{\Gamma(\beta(l+n)+1)} \mathbb{E}[(M_{\alpha, \beta_1, \lambda_1, \mu}(t))^{\beta(n+l)}] \\ &= \frac{d}{dt} \left(1 - \sum_{n=0}^{k-1} \frac{\lambda^n}{n!} \sum_{l=0}^{\infty} \frac{(n+l)!}{l!} \frac{(-\lambda)^l}{\Gamma(\beta(l+n)+1)} \mathbb{E}[(M_{\alpha, \beta_1, \lambda_1, \mu}(t))^{\beta(n+l)}] \right) \\ &= -\frac{d}{dt} \sum_{n=0}^{k-1} \frac{\lambda^n}{n!} \sum_{l=0}^{\infty} \frac{(n+l)!}{l!} \frac{(-\lambda)^l}{\Gamma(\beta(l+n)+1)} \mathbb{E}[(M_{\alpha, \beta_1, \lambda_1, \mu}(t))^{\beta(n+l)}].\end{aligned}$$

References

- Andrews, G. E. (1998). *The theory of partitions*. Number 2. Cambridge university press.
- Beghin, L. and Macci, C. (2014). Fractional discrete processes: compound and mixed poisson representations. *Journal of Applied Probability*, 51(1):19–36.
- Beghin, L. and Orsingher, E. (2009). Fractional poisson processes and related planar random motions.
- Beghin, L. and Vellaisamy, P. (2018). Space-fractional versions of the negative binomial and polya-type processes. *Methodology and Computing in Applied Probability*, 20(2):463–485.
- Biard, R. and Saussereau, B. (2014). Fractional poisson process: long-range dependence and applications in ruin theory. *Journal of Applied Probability*, 51(3):727–740.
- Cox, D. R. and Lewis, P. A. (1966). The statistical analysis of series of events.
- Di Crescenzo, A. and Meoli, A. (2023). Competing risks and shock models governed by a generalized bivariate poisson process. *Journal of Applied Probability*, 60(2):709–722.
- Doukhan, P., Oppenheim, G., and Taqqu, M. (2002). *Theory and applications of long-range dependence*. Springer Science & Business Media.
- D’Ovidio, M. and Nane, E. (2014). Time dependent random fields on spherical non-homogeneous surfaces. *Stochastic Processes and their Applications*, 124(6):2098–2131.
- Gillespie, T. R. (1999). The stochastically subordinated poisson normal process for modelling financial assets. *Bulletin of the Australian Mathematical Society*, 59(3):527–528.
- Guler Dincer, N., Demir, S., and Yalçın, M. O. (2022). Forecasting covid19 reliability of the countries by using non-homogeneous poisson process models. *New Generation Computing*, pages 1–22.
- Gupta, N. and Kumar, A. (2023). Fractional poisson processes of order k and beyond. *Journal of Theoretical Probability*, 36(4):2165–2191.
- Ken-Iti, S. (1999). *Lévy processes and infinitely divisible distributions*. Cambridge university press.
- Kilbas, A. A., Saigo, M., and Trujillo, J. J. (2002). On the generalized wright function. *Fractional Calculus and Applied Analysis*, 5(4):437–460.
- Kumar, A., Leonenko, N., and Pichler, A. (2020). Fractional risk process in insurance. *Mathematics and Financial Economics*, 14:43–65.
- Kumar, A., Upadhye, N., Wylomanska, A., and Gajda, J. (2019). Tempered mittag-leffler levy processes. *Communications in Statistics-Theory and Methods*, 48(2):396–411.
- Kumar, A. and Vellaisamy, P. (2015). Inverse tempered stable subordinators. *Statistics & Probability Letters*, 103:134–141.
- Laskin, N. (2003). Fractional poisson process. *Communications in Nonlinear Science and Numerical Simulation*, 8(3-4):201–213.
- Maheshwari, A. (2023). Tempered space fractional negative binomial process. *Statistics & Probability Letters*, 196:109799.
- Maheshwari, A. and Vellaisamy, P. (2016). On the long-range dependence of fractional poisson and negative binomial processes. *Journal of Applied Probability*, 53(4):989–1000.
- Maheshwari, A. and Vellaisamy, P. (2019). Fractional poisson process time-changed by lévy subordinator and its inverse. *Journal of Theoretical Probability*, 32(3):1278–1305.
- Meerschaert, M., Nane, E., and Vellaisamy, P. (2011). The fractional poisson process and the inverse stable subordinator. *Electronic Journal of Probability*, 16:1600–1620.
- Meerschaert, M. M. and Straka, P. (2013). Inverse stable subordinators. *Mathematical modelling of natural phenomena*, 8(2):1–16.
- Mendoza-Arriaga, R. and Linetsky, V. (2016). Multivariate subordination of markov processes with financial applications. *Mathematical Finance*, 26(4):699–747.
- Meoli, A. (2023). Some poisson-based processes at geometric times. *Journal of Statistical Physics*, 190(6):107.
- Orsingher, E. and Polito, F. (2012). The space-fractional poisson process. *Statistics & Probability Letters*, 82(4):852–858.
- Podlubny, I. (1999). An introduction to fractional derivatives, fractional differential equations, to methods of their solution and some of their applications. *Math. Sci. Eng.*, 198:340.
- Rosiński, J. (2007). Tempering stable processes. *Stochastic processes and their applications*, 117(6):677–707.
- Rusakov, O. and Yakubovich, Y. (2021). Poisson processes directed by subordinators, stuttering poisson and pseudo-poisson processes, with applications to actuarial mathematics. *Journal of Physics: Conference Series*, 2131(2):022107.
- Soni, R. and Pathak, A. K. (2024). Generalized fractional negative binomial process. *Statistics & Probability Letters*, 207:110021.
- Soni, R., Pathak, A. K., Di Crescenzo, A., and Meoli, A. (2024). Bivariate tempered space-fractional poisson process and shock models. *To appear in Journal of Applied Probability*.
- Soni, R., Vellaisamy, P., and Pathak, A. (2023). A probabilistic generalization of the bell polynomials. *The Journal of Analysis*, pages 1–22.
- Vellaisamy, P. and Maheshwari, A. (2018). Fractional negative binomial and polya processes. *Probability and Mathematical Statistics*, 38(1):77–101.