

Special Moufang sets of finite dimension

Matthias Grüninger

September 12, 2024

Abstract

We prove that a special Moufang sets with abelian root subgroups derive from a quadratic Jordan division algebra if a certain finiteness condition is satisfied.

1 Introduction

The concept of a Moufang set was by introduced by J. Tits in [29]. This concept is essentially equivalent to that of a group with split BN-pair of rank one. We first recall the definition of a Moufang set.

Definition 1.1 A *Moufang set* consists of a set X with $|X| \geq 3$ and a family $(U_x)_{x \in X}$ of subgroups of $\text{Sym}X$ such that

- (MS1) For all $x \in X$, the group U_x fixes x and acts regularly on $X \setminus \{x\}$.
- (MS2) For all $x, y \in X$ and $g \in U_y$ we have $U_x^g = U_{xg}$.

The group $G^\dagger := \langle U_x \mid x \in X \rangle$ is called the *little projective group* of the Moufang set. It acts 2-transitively on X , and for all $x \in X$ the group U_x is a normal subgroup of the stabiliser of x in G^\dagger acting regularly on $X \setminus \{x\}$.

Definition 1.2 A Moufang set $(X, (U_x)_{x \in X})$ is called *proper* if its little projective group is not sharply 2-transitive on X .

The prototype of a Moufang set is the projective line $\text{PG}_1(\mathbb{k})$ over a field \mathbb{k} . In this case we have $G^\dagger = \text{PSL}_2(\mathbb{k})$. As we will explain, one can generalise this by taking a quadratic Jordan division algebra instead of a field.

As shown in [7], every Moufang set can be constructed in the following manner: Let $(U, +)$ be a not necessarily abelian group with $|U| > 1$, $X := U \dot{\cup} \{\infty\}$ and $\tau \in \text{Sym}X$ an element interchanging 0 and ∞ . For $a \in U$ we set

$$\alpha_a : X \rightarrow X : x \mapsto \begin{cases} x + a, & \text{if } x \in U, \\ \infty, & \text{if } x = \infty. \end{cases}$$

Moreover, we set $U_\infty := \{\alpha_a \mid a \in U\}$, $U_0 := U_\infty^\tau$, $U_a := U_0^{\alpha_a}$ for all $a \in U$ and $\mathbb{M}(U, \tau) := (X, (U_x)_{x \in X})$.

This is not always a Moufang set. Condition (MS1) is straightforward, but (MS2) is not always satisfied. There is a nice criterion assuring that $\mathbb{M}(U, \tau)$ is a Moufang set. For $a \in U^\# := U \setminus \{0\}$ set $\mu_a := \alpha_a \alpha_{-a\tau^{-1}} \alpha_{-(-a\tau^{-1})\tau}$ and $h_a := \tau \mu_a$. We call μ_a the μ -map and h_a the *Hua map* associated to a . Note that μ_a interchanges 0 and ∞ and thus h_a fixes both elements.

Theorem 1.3 ([7, Theorem 3.2]) $\mathbb{M}(U, \tau)$ is a Moufang set if and only if $h_a \in \text{Aut}(U)$ for all $a \in U^\#$, i.e. $(b+c)h_a = bh_a + ch_a$ for all $a, b, c \in U$ with $a \neq 0$. If this holds, we have $\mathbb{M}(U, \tau) = \mathbb{M}(U, \mu_a)$ for all $a \in U^\#$.

Note that if $\mathbb{M}(U, \tau)$ is a Moufang set, then $\mathbb{M}(U, \tau) = \mathbb{M}(U, \mu_a)$ for every $a \in U^\#$.

Definition 1.4 Suppose that $\mathbb{M}(U, \tau)$ is a Moufang set. Then the group $H := \langle \mu_a \mu_b \mid a, b \in U^\# \rangle$ is called the *Hua subgroup* of $\mathbb{M}(U, \tau)$.

One can show that $H := G_{0, \infty}^\dagger$, so $\mathbb{M}(U, \tau)$ is proper if and only if $H \neq 1$. By Theorem 1.3 one can see that every quadratic Jordan division algebra gives rise to a Moufang set. Recall the definition of a quadratic Jordan division algebra.

Definition 1.5 Let \mathbb{k} be a field, J a \mathbb{k} -vector space, $Q : J \rightarrow \text{End}_{\mathbb{k}}(J) : a \mapsto Q_a$ a map and $1 \in J^\#$. Then $(J, Q, 1)$ is called a *quadratic Jordan division algebra* if

- (a) The map Q is quadratic, i.e. $Q_{r \cdot a} = r^2 \cdot Q_a$ for all $a \in J$ and all $r \in \mathbb{k}$, and the map $(a, b) \mapsto Q_{a,b} := Q_{a+b} - Q_a - Q_b$ is \mathbb{k} -bilinear.
- (b) Q_a is invertible for all $a \in J^\#$. We set $a^{-1} := aQ_a^{-1}$ for $a \in J^\#$.
- (c) For all $a, b \in J$ we have
 - (QJ1) $Q_1 = \text{id}$.
 - (QJ2) $Q_a V_{a,b} = V_{b,a} Q_a$, where $V_{a,b} \in \text{End}_{\mathbb{k}}(J)$ is defined by $cV_{a,b} := bQ_{a,c}$ for all $c \in J$.
 - (QJ3) $Q_a Q_b = Q_b Q_a Q_b$.

Remark 1.6 The concept of a quadratic Jordan algebra was introduced in [23]. Normally, one requires that (QJ1-3) hold strictly, i.e. continue to hold in all scalar extensions of J . However, by the main theorem of [11], this is not necessary for quadratic Jordan division algebras.

The connection between Moufang sets and quadratic Jordan division algebras is established in the following theorem.

Theorem 1.7 ([7, Theorem 4.1 and 4.2]) Let $(J, Q, 1)$ be a quadratic Jordan division algebra over a field k . For $a \in J^\#$ we set $a\tau := -a^{-1}$. Then $\mathbb{M}(J) := \mathbb{M}(J, \tau)$ is a Moufang set. Moreover, we have $\tau = \mu_1$ and $h_a = Q_a$ for all $a \in J^\#$.

All examples of proper Moufang sets with abelian root groups that are known are isomorphic to $\mathbb{M}(J)$ for some quadratic Jordan division algebra J . Therefore it is a major conjecture that every proper Moufang set with abelian root groups comes from a quadratic Jordan algebra.

If J is a quadratic Jordan division algebra, $\mathbb{M}(J, \tau)$ satisfies the identity $(-a)\tau = -a\tau$ for all $a \in J^\#$. Such Moufang sets are called *special*.

Definition 1.8 A Moufang set $\mathbb{M}(U, \tau)$ is called *special* if $(-a)\tau = -a\tau$ for all $a \in U^\#$.

By [26] a Moufang set with abelian root group is special or improper. Moreover, the projective lines over \mathbb{F}_2 and \mathbb{F}_3 are up to isomorphism the only improper special Moufang sets. Therefore, for Moufang sets with abelian root group, the properties special and proper are almost the same.

Given a proper Moufang set $\mathbb{M}(U, \tau)$ with U abelian, the problem is to recover the Jordan structure. As a first step we can find a field over which the potential Jordan algebra is defined. By [4, Proposition 4.6(5)] either U is torsion-free and uniquely divisible or U is an elementary abelian p -group for some prime p . We say $\text{char}U = 0$ in the first case and $\text{char}U = p$ in the second case. Thus U is a vector space over \mathbb{k} , where $\mathbb{k} = \mathbb{Q}$ if $\text{char}U = 0$ and $\mathbb{k} = \mathbb{F}_p$ if $\text{char}U = p$. This field \mathbb{k} is called the *prime field* of $\mathbb{M}(U, \tau)$.

By Theorem 1.3 we may suppose that $\tau = \mu_e$ for some $e \in U^\#$.

Notation 1.9 Let $\mathbb{M}(U, \tau)$ be a proper Moufang set with U abelian and $\tau = \mu_e$ for some fixed $e \in U^\#$. In the following, h_0 will denote the zero map of U , and we set $h_{a,b} := h_{a+b} - h_a - h_b$ for all $a, b \in U$.

Considering Theorem 1.7, the map $\mathcal{H} : U \rightarrow \text{End}_{\mathbb{k}}(U) : a \mapsto h_a$ is a natural candidate for the quadratic map Q . The map \mathcal{H} depends on the choice of the neutral element $e \in U^\#$. If we choose another element $c \in U^\#$, then for $a \in U$ the Hua map h_a^c is given by $h_a^c = h_c^{-1}h_a$. The triple (U, \mathcal{H}^c, c) with $\mathcal{H}^c : U \rightarrow \text{End}(U) : a \mapsto h_a^c$ is called the *c-isotope* of (U, \mathcal{H}, e) . The triple (U, \mathcal{H}, e) is a quadratic Jordan division algebra if and only if every c -isotope is a quadratic Jordan division algebra.

By [5, Remark after 7.6.1] all conditions for (U, \mathcal{H}, e) to be a quadratic Jordan division algebra but the biadditivity of \mathcal{H} and Axiom (QJ2) are satisfied. In [4] it was proved that these conditions can be replaced by weaker conditions. In [13] the author of this paper proved that these conditions can be replaced by even weaker conditions. For the condition we are going to use in this paper we will need the following definition.

Definition 1.10 Let $\mathbb{M}(U, \tau)$ be a special Moufang set with U abelian and let H be its Hua subgroup. We set $\mathcal{C} := \{T \in \text{End}_H(U) \mid h_{aT} = T^2h_a \text{ for all } a \in U\}$, the *centroid* of $\mathbb{M}(U, \tau)$. Moreover, we set $\mathcal{C}^* := \mathcal{C} \cap \text{GL}(U)$.

By [4][Proposition 4.6(6)] \mathcal{C} contains the prime field of U . Thus if $\text{char}U = 0$, then $|\mathcal{C}|$ is infinite.

We will make use of the following criterion:

Theorem 1.11 *Let $\mathbb{M}(U, \tau)$ be a special Moufang set with U abelian. Suppose that there is a subfield \mathbb{k} of the centroid and an element $1 \neq \lambda \in \mathbb{k}^*$ such that $h_{e, \lambda \cdot a} = \lambda \cdot h_{e, a}$ for all $a \in U^\#$. Then (U, \mathcal{H}, e) is a quadratic Jordan division algebra.*

Proof. By [13, Theorem 2.6] it follows that the map $a \mapsto h_a$ is quadratic. The claim now follows from [13, Theorem 4.6]. \square

The main result of this paper is Theorem 3.10:

THEOREM 1 *Let $\mathbb{M}(U, \tau)$ be a special Moufang set with U abelian and let H be its Hua subgroup. Suppose that there is a subfield \mathbb{k} of $\text{End}_H(U)$ with*

- (i) $\dim_{\mathbb{k}} U < \infty$,
- (ii) $|\mathbb{k} \cap \mathcal{C}| = \infty$,
- (iii) U is generated by the set of squares $\{a^2 \mid a \in U\}$ as a $\text{End}_H(U)$ -module and
- (iv) there is $\lambda \in \mathbb{k}^* \cap \mathcal{C}$ such that $\lambda - 1 \in \mathbb{k}^* \cap \mathcal{C}$ as well.

Then (U, \mathcal{H}, e) is a quadratic Jordan division algebra.

Note that the second condition is automatically satisfied if $\text{char} U = 0$ and the third and fourth if $\text{char} U \neq 2$. In (iv), we can take $\lambda = -1$ if $\text{char} U \neq 2$. Thus we have

Corollary 1.12 *Suppose that $\mathbb{M}(U, \tau)$ is a special Moufang set with U abelian and torsion-free. If U is finite-dimensional over a subfield \mathbb{k} of the centroid of $\mathbb{M}(U, \tau)$, then $\mathbb{M}(U, \tau)$ arises from a quadratic Jordan division algebra.*

As an application, we will discuss Moufang sets whose little projective group can be embedded into a sharply triple transitive group. A sharply triple transitive group whose point stabilisers have normal subgroup acting regularly arises from an algebraic structure called a *KT-nearfield*, named after Karzel and Tits. It is conjectured that every KT-nearfield is actually a commutative field with slightly modified multiplication. In [18] Kerby proved this for nearfields of characteristic different from 2 having dimension 2 over their kernel. We are able to strengthen this result:

THEOREM 2 *Suppose that F is a KT-nearfield with $\text{char} F \neq 2$ such that F has finite dimension over its kernel. Then F is a Dickson nearfield coupled to a commutative field which we also call F , and the corresponding sharply triple group transitive is sandwiched between $\text{PSL}_2(F)$ and $\text{P}\Gamma\text{L}_2(F)$.*

For the proof of our results we will use an identity involving geometric series. However, in order to define an infinite series, we will need an appropriate topology which is a priori not given. In order to get such a topology, we will extend the original Moufang set by an ultrapower. Therefore, we will discuss ultraproducts at first.

Acknowledgement: The author likes to thank Pierre-Emmanuel Caprace, Theo Grundhöfer and Bernhard Mühlherr for their valuable comments.

2 Ultraproducts of Moufang sets

In this section we will introduce ultraproducts of Moufang sets. For an introduction on ultraproducts see [3, §4]

Definition 2.1 Let S be a non-empty set.

- (a) A subset \mathcal{F} of the power set $\mathcal{P}(S)$ of S is called a *filter* on S if

- (i) $S \in \mathcal{F}$ and $\emptyset \notin \mathcal{F}$.
- (ii) For all $A, B \in \mathcal{F}$ we have $A \cap B \in \mathcal{F}$.
- (iii) For all $A \in \mathcal{F}$ and all $A \subseteq B \subseteq S$ we have $B \in \mathcal{F}$.
- (b) A filter \mathcal{F} is called *fixed* if $\bigcap_{A \in \mathcal{F}} A \neq \emptyset$ and *free* otherwise.
- (c) A filter \mathcal{F} is called an *ultrafilter* if for all $A \subseteq S$ either $A \in \mathcal{F}$ or $S \setminus A \in \mathcal{F}$.

Example 2.2 (a) If $\emptyset \neq B \subseteq S$, then $\mathcal{F}_B := \{A \subseteq S \mid B \subseteq A\}$ is a fixed filter on S . It is an ultrafilter if and only if $|B| = 1$.

- (b) If S is infinite, then $\mathcal{F} := \{A \subseteq S \mid |S \setminus A| < \infty\}$ is a free filter on S , but not an ultrafilter.

Theorem 2.3 *Let S be a non-empty set.*

- (a) *If \mathcal{F} a filter on S , then there is an ultrafilter \mathcal{F}^* on S with $\mathcal{F} \subseteq \mathcal{F}^*$.*
- (b) *If S is infinite, then there is a free ultrafilter \mathcal{F} on S .*

Proof.

- (a) This follows by Zorn's Lemma, see for example [3, Corollary 4.1.4] for details.
- (b) This follows by (a) and Example 2.2(b).

□

Now let S be a non-empty set and \mathcal{F} an ultrafilter on S . Suppose that for every $s \in S$ there is a group/ring/etc. X_s . We set

$$I_{\mathcal{F}} := \{(x_s)_{s \in S} \mid \{s \in S \mid x_s = 0\} \in \mathcal{F}\}.$$

Then $I_{\mathcal{F}}$ is normal subgroup/ideal/etc. of $\prod_{s \in S} X_s$. We will write $\prod_{s \in S} X_s / \mathcal{F}$ instead of $\prod_{s \in S} X_s / I_{\mathcal{F}}$ and $[(x_s)_{s \in S}]_{\mathcal{F}}$ or just $[(x_s)_{s \in S}]$ for $(x_s)_{s \in S} + I_{\mathcal{F}}$. Note that if $X_s \neq 0$ for all $s \in S$ and $0 \neq x \in \prod_{s \in S} X_s / \mathcal{F}$, then there is a representative $(x_s)_{s \in S}$ of x such that $x_s \neq 0$ for all $s \in S$. We call $\prod_{s \in S} X_s / \mathcal{F}$ the *ultraproduct* of the X_s with respect to \mathcal{F} . If $X_s = X$ for all $s \in S$, then X^S / \mathcal{F} is called the *ultrapower* of X with respect to \mathcal{F} . In this case, the map $x \mapsto [(x)_{s \in S}]$ is an embedding of X into X^S / \mathcal{F} , thus we may consider X as a substructure of X^S / \mathcal{F} .

Lemma 2.4 *Let \mathcal{F} be an ultrafilter on a set S and let $A \in \mathcal{F}$. Suppose that $A = \dot{\bigcup}_{i=1}^n A_i$ for some $n \geq 1$. Then there is exactly one $1 \leq i \leq n$ with $A_i \in \mathcal{F}$.*

Proof. Induction on n . The claim is trivially true for $n = 1$. For the inductive step suppose that $A = \dot{\bigcup}_{i=1}^{n+1} A_i$. If $A_{n+1} \in \mathcal{F}$, then $A_i \notin \mathcal{F}$ for $1 \leq i \leq n$, since otherwise $\emptyset = A_i \cap A_{n+1} \in \mathcal{F}$. If $A_{n+1} \notin \mathcal{F}$, then $S \setminus A_{n+1} \in \mathcal{F}$, thus $\dot{\bigcup}_{i=1}^n A_i = A \cap (S \setminus A_{n+1}) \in \mathcal{F}$. The claim now follows by inductive hypothesis.

□

Lemma 2.5 Suppose that \mathbb{k}_s is a field for all $s \in S$. Then $\prod_{s \in S} \mathbb{k}_s / \mathcal{F}$ is also field.

Proof. This is well-known, see for example [3, Exercise 4.1.30]. \square

Lemma 2.6 Let S be a non-empty set, \mathcal{F} an ultrafilter on S and \mathbb{k} a field. Set $\tilde{\mathbb{k}} := \mathbb{k}^S / \mathcal{F}$. Then we have

- (a) \mathbb{k} is algebraically closed in $\tilde{\mathbb{k}}$.
- (b) Suppose that $f : S \rightarrow \mathbb{k}$ is a function that is not constant on any subset of S contained in \mathcal{F} . Set $x := f + \mathcal{F}$. Then there is a valuation $v : \tilde{\mathbb{k}} \rightarrow \mathbb{R} \cup \{\infty\}$ such that $v(x) = 1$ and $v(a) = 0$ for all $a \in \mathbb{k}^*$.

Proof.

- (a) Let $p \in \mathbb{k}[t]$ be a non-constant polynomial and $\alpha = [(\alpha_s)_{s \in S}]$ with $p(\alpha) = 0$. Then $S' := \{s \in S \mid p(\alpha_s) = 0\} \in \mathcal{F}$. Let $\alpha_1, \dots, \alpha_n$ be the roots of p in \mathbb{k} . Set $S_i := \{s \in S \mid \alpha_s = \alpha_i\}$ for $i = 1, \dots, n$. Since $S' = \bigcup_{i=1}^n S_i$, by 2.4 there is exactly one $1 \leq i \leq n$ with $S_i \in \mathcal{F}$. Thus $\alpha = \alpha_i \in \mathbb{k}$. Hence the claim follows.
- (b) By assumption $x \notin \mathbb{k}$, thus x is transcendental over \mathbb{k} by (a). Let \mathcal{B} be a transcendental base of $\tilde{\mathbb{k}}$ over $\mathbb{k}(x)$. Then $\mathcal{B} \cup \{x\}$ is a transcendental base of $\tilde{\mathbb{k}}$ over \mathbb{k} and $\{x\}$ is a transcendental base of $\tilde{\mathbb{k}}$ over $\mathbb{K} := \mathbb{k}(\mathcal{B})$. Thus there is a valuation v of $\mathbb{K}(x)$ that is trivial on $\mathbb{K} \supseteq \mathbb{k}$ with $v(x) = 1$. Since $\tilde{\mathbb{k}}$ is algebraic over $\mathbb{K}(x)$, we can extend v to a valuation on $\tilde{\mathbb{k}}$ by [21, XII 4.4]. Thus the claim follows. \square

Remark 2.7 This valuation is not necessarily discrete. For example, if $S = \mathbb{N}$ and $f(n) = a^{n!}$ for some $a \in K^*$ of infinite order, then x is a n -th power for all $n \in \mathbb{N}$, thus the value group contains \mathbb{Q} . The crucial thing is that the value group is Archimedean.

Lemma 2.8 Let S be a non-empty set and \mathcal{F} an ultrafilter on S .

- (a) If R_s is a ring and M_s is an R_s -module for all $s \in S$, then $\prod_{s \in S} M_s / \mathcal{F}$ is a $\prod_{s \in S} R_s / \mathcal{F}$ -module via $[(x_s)_{s \in S}] \cdot [(m_s)_{s \in S}] = [(x_s \cdot m_s)_{s \in S}]$ for all $(x_s)_{s \in S} \in \prod_{s \in S} R_s$ and all $(m_s)_{s \in S} \in \prod_{s \in S} M_s$.
- (b) Suppose that \mathbb{k} is a field, V a \mathbb{k} -vector space of finite dimension and B a \mathbb{k} -basis of V . Set $\tilde{\mathbb{k}} := \mathbb{k}^S / \mathcal{F}$ and $\tilde{V} := V^S / \mathcal{F}$. Then B is $\tilde{\mathbb{k}}$ -basis of \tilde{V} .

Proof.

- (a) This is obvious.
- (b) Since B is finite, V^S is generated by B as a \mathbb{k}^S -module. One easily sees that B is $\tilde{\mathbb{k}}$ -linearly independent. Thus the claim follows. \square

Now suppose that S is a non-empty set, \mathcal{F} an ultrafilter on S and $\mathbb{M}(U_s, \tau_s)$ a Moufang set for all $s \in S$. We suppose that $\tau_s = \mu_{a_s}$ for some $a_s \in U_s^\#$ for all $s \in S$. We set $U := \prod_{s \in S} U_s / \mathcal{F}$. We define the map τ on $U^\#$ as follows: If $a \in U^\#$, choose a representative $(a_s)_{s \in S}$ of a with $a_s \neq 0$ for all $s \in S$ and set $a\tau := [(a_s \tau_s)_{s \in S}]$. This is well-defined. We get

Theorem 2.9 *For all $x = [(x_s)_{s \in S}]$, $a = [(a_s)_{s \in S}] \in U$ with $a_s \neq 0$ for all $s \in S$ we have $xh_a = [(x_s h_{a_s})_{s \in S}]$. In particular $\mathbb{M}(U, \tau)$ is a Moufang set. Moreover, we have*

- (a) *The Moufang set $\mathbb{M}(U, \tau)$ is proper if and only if $\{s \in S \mid (U_s, \tau_s) \text{ is proper}\} \in \mathcal{F}$.*
- (b) *The Moufang set $\mathbb{M}(U, \tau)$ is special if and only if $\{s \in S \mid (U_s, \tau_s) \text{ is special}\} \in \mathcal{F}$.*
- (c) *If $\mathbb{M}(U_s, \tau_s)$ is special for all $s \in S$ and if $T_s \in \text{End}(U_s)$ is in the centroid of $\mathbb{M}(U_s, \tau_s)$ for all $s \in S$, then $[(T_s)_{s \in S}]$ is in the centroid of $\mathbb{M}(U, \tau)$.*

Proof. Let $(x_s)_{s \in S} \in \prod_{s \in S} U_s$ and let $(a_s)_{s \in S} \in \prod_{s \in S} U_s^\#$. We have $S = S_1 \dot{\cup} S_2 \dot{\cup} S_3$ with $S_1 := \{s \in S \mid x_s = 0\}$, $S_2 := \{s \in S \mid x_s = a_s \tau_s^{-1}\}$ and $S_3 := \{s \in S \mid x_s \neq 0, a_s \tau_s^{-1}\}$. Then exactly one of the sets S_1 , S_2 and S_3 are contained in \mathcal{F} by 2.4.

Suppose that $S_1 \in \mathcal{F}$. Then by definition $xh_a = 0$ but also $\{s \in S \mid x_s h_{a_s} = 0\} = S_1 \in \mathcal{F}$, thus the claim follows in this case. If $S_2 \in \mathcal{F}$, then we have $\{s \in S \mid x\tau - a = 0\} \in \mathcal{F}$, thus $(x\tau - a)\tau^{-1} = \infty = (x\tau - a)\tau^{-1} - a\tau^{-1}$, hence $((x\tau - a)\tau^{-1} - a\tau^{-1})\tau = 0$ and so $xh_a = ((x\tau - a)\tau^{-1} - a\tau^{-1})\tau - (-a\tau^{-1})\tau = -(-a\tau^{-1})\tau$. However, for $s \in S_2$ we have $x_s h_{a_s} = a_s \tau_s^{-1} h_{a_s} = -(-a_s \tau_s^{-1})\tau_s$, thus the claim follows.

Now for $s \in S_3$, we have $x_s, x_s \tau_s - a_s, (x_s \tau_s - a_s)\tau_s^{-1} - a\tau_s^{-1} \in U_s^\#$. Thus if $S_3 \in \mathcal{F}$, then $(xh_a)_s = x_s h_{a_s}$ for all $s \in S_3$, hence the claim follows.

Since every map h_{a_s} is an automorphism of U_s for all $s \in S$, it follows that h_a is an automorphism of U . Thus $\mathbb{M}(U, \tau)$ is a Moufang set.

- (a) For every $t \in \{s \in S \mid \mathbb{M}(U_s, \tau_s) \text{ is proper}\}$ there are $x_t, a_t \in U_s^\#$ with $x_t h_{a_t} \neq x_t$. Choose x_s and a_s arbitrarily for the other $s \in S$ and set $x := [(x_s)_{s \in S}]$ and $a := [(a_s)_{s \in S}]$. Then if $\{s \in S \mid \mathbb{M}(U_s, \tau_s) \text{ is proper}\} \in \mathcal{F}$, we have $xh_a \neq x$, thus $\mathbb{M}(U, \tau)$ is proper.

Conversely, for $t \in \{s \in S \mid \mathbb{M}(U_s, \tau_s) \text{ is improper}\}$, we have $x_t h_{a_t} = x_t$ for all $x_t \in U$ and all $a_t \in U_t^\#$. Thus if this set is in S , the Moufang set $\mathbb{M}(U, \tau)$ is improper.

- (b) For all $t \in \{s \in S \mid \mathbb{M}(U_s, \tau_s) \text{ is special}\}$ we have $(-x_t)\tau_t = -x_t\tau_t$ for all $x_t \in U_t^\#$. Thus the claim follows easily.
- (c) This follows also by an easy calculation.

□

Remark 2.10 It is conjectured that the root groups of a proper Moufang set are nilpotent, see [6]. If this conjecture is true, then there is a constant c such that the root groups are nilpotent of class at most c . For if for every natural number n there is a proper Moufang set $\mathbb{M}(U_n, \tau_n)$ such that U_n is of class at least n and if \mathcal{F} is a free ultrafilter on \mathbb{N} , then $\prod_{n \in \mathbb{N}} \mathbb{M}(U_n, \tau_n) / \mathcal{F}$ is a proper Moufang set whose root groups are isomorphic to $\prod_{n \in \mathbb{N}} U_n / \mathcal{F}$ and therefore not nilpotent.

3 Special Moufang sets of finite dimension

Before proving our main result we need some auxiliary results in algebra.

Lemma 3.1 *Let $\tilde{\mathbb{k}}$ be a field with valuation v , \mathcal{O} its valuation ring, \mathbb{k} a subfield of \mathcal{O} , V a vector space over $\tilde{\mathbb{k}}$ and Λ an \mathcal{O} -sublattice of V . Suppose that $v_1, \dots, v_n \in V$ and $\lambda_1, \dots, \lambda_n \in \mathbb{k}$ pairwise distinct such that $\sum_{j=1}^n \lambda_i^{j-1} \cdot v_j \in \Lambda$ for $1 \leq i \leq n$. Then $v_1, \dots, v_n \in \Lambda$.*

Proof. The matrix $A := (\lambda_i^{j-1})_{i,j=1,\dots,n}$ is a Vandermonde matrix with entries in $\mathbb{k} \subseteq \mathcal{O}$. Since the elements λ_i are all distinct and contained in \mathcal{O} , we have $\det(A) := \prod_{i < j} (\lambda_j - \lambda_i) \in \mathcal{O}^*$. Thus if $w_i := \sum_{j=1}^n \lambda_i^{j-1} w_j$ and $A^{-1} = (b_{ij})_{i,j=1,\dots,n} \in \text{GL}_n(\mathcal{O})$, we have $v_i = \sum_{j=1}^n b_{ij} w_j \in \Lambda$. \square

Notation 3.2 Let $\tilde{\mathbb{k}}$ be a field with valuation $v : \tilde{\mathbb{k}} \rightarrow \mathbb{R} \cup \{\infty\}$, \mathcal{O} its valuation ring, V a $\tilde{\mathbb{k}}$ -vector space and B a basis of V . For $v = \sum_{b \in B} \lambda_b \cdot b \in V$ set $\omega(v) := \min\{v(\lambda_b) \mid b \in B\}$. Then the map ω satisfies

- (i) $\omega(v) = \infty \iff v = 0$,
- (ii) $\omega(v + w) \geq \min\{\omega(v), \omega(w)\}$,
- (iii) $\omega(\lambda \cdot v) = \omega(v) + v(\lambda)$

for all $v, w \in V$ and all $\lambda \in \tilde{\mathbb{k}}$. We will call ω the *minimum norm* with respect to v and B . For all $c > 0$ we get an ultrametric d on \tilde{V} by $d(v, w) = c^{\omega(v-w)}$ for $v, w \in \tilde{V}$.

Moreover, we set $\Lambda_r := \{v \in \tilde{V} \mid \omega(v) \geq r\}$ for all $r \in \mathbb{R}$. Then Λ_r is \mathcal{O} -lattice.

Lemma 3.3 *Let $\tilde{\mathbb{k}}$, V and \mathcal{O} as in 3.2. Suppose that $\dim V = n < \infty$ and $B = (b_1, \dots, b_n)$ is a \mathbb{k} -basis of V . Let $f : V \rightarrow V$ be a $\tilde{\mathbb{k}}$ -linear map and $A := (a_{ij})_{i,j=1,\dots,n}$ its transformation matrix with respect to B . Then we have*

- (a) *The map f is Lipschitz-continuous with respect to ω with Lipschitz-constant $L := \min\{v(a_{ij}) \mid 1 \leq i, j \leq n\}$, i.e. $\omega(f(v)) \geq L + \omega(v)$ for all $v \in V$.*
- (b) *Suppose that \mathbb{k} is a subfield of \mathcal{O} . If f is invertible and $a_{ij} \in \mathbb{k}$ for all $1 \leq i, j \leq n$, then f is an isometry with respect to ω .*

Proof. For $v = \sum_{i=1}^n \lambda_i \cdot b_i \in V$ we have

$$\begin{aligned}
\omega(f(v)) &= \omega\left(f\left(\sum_{i=1}^n \lambda_i \cdot b_i\right)\right) = \omega\left(\sum_{i=1}^n \lambda_i \cdot \sum_{j=1}^n a_{ij} \cdot b_j\right) \\
&= \omega\left(\sum_{j=1}^n \sum_{i=1}^n \lambda_i \cdot a_{ij} \cdot b_j\right) \\
&= \min\left\{v\left(\sum_{i=1}^n \lambda_i \cdot a_{ij}\right) \mid 1 \leq j \leq n\right\} \\
&\geq \min\{v(\lambda_i \cdot a_{ij}) \mid 1 \leq i, j \leq n\} \\
&= \min\{v(\lambda_i) + v(a_{ij}) \mid 1 \leq i, j \leq n\} \\
&\geq \min\{v(\lambda_i) \mid 1 \leq i \leq n\} + \min\{v(a_{ij}) \mid 1 \leq i, j \leq n\} \\
&= \omega(v) + L.
\end{aligned}$$

This proves (a). If $a_{ij} \in \mathbb{k}$ for all $1 \leq i, j \leq n$ and f is not the zero map, we have $L = 0$, hence $\omega(f(v)) \geq \omega(v)$ for all $v \in V$. If f is invertible, we also have $\omega(f^{-1}(v)) \geq \omega(v)$ for all $v \in V$. If we replace v by $f(v)$, we get $\omega(v) \geq \omega(f(v))$. Thus the claim follows. \square

From now on we assume that $\mathbb{M}(U, \tau)$ is a special Moufang set with U abelian and $\tau = \mu_e$ for some fixed element $e \in U^\#$. Moreover, \mathbb{k} is a subfield of $\text{End}_H(U)$, where H is the Hua subgroup of $\mathbb{M}(U, \tau)$.

Notation 3.4 For $a \in U$ and $n \in \mathbb{N}$ we define a^n recursively by $a^0 := e$, $a^1 := a$ and $a^{n+2} := a^n h_a$. Moreover, for $a \in U^\#$ we set $a^{-1} := -a\tau$.

Lemma 3.5 For all $a \in U$ and $n, m \in \mathbb{N}$ we have:

- (a) If $2m \geq n$, then $a^n = a^{n-2m} h_a^m$.
- (b) $h_{a^n} = h_a^n$.
- (c) $(a^m)^n = a^{nm}$.
- (d) $a^n h_{e,a} = 2a^{n+1}$.
- (e) $(\lambda a)^n = \lambda^n a^n$ for all $\lambda \in \mathcal{C}$.

Proof. The first statement is very easy to see. We proof (b), (c), (d) and (e) by induction on n .

- (b) For $n = 0$ we have $h_{a^0} = h_e = 1 = h_a^0$, for $n = 1$ we have $h_{a^1} = h_a = h_a^1$. Moreover, we have $h_{a^{n+2}} = h_{a^n h_a} = h_a h_a^n h_a = h_a^{n+2}$. Thus the claim follows.
- (c) The statement is trivially true for $m \in \{0, 1\}$, therefore we may assume $m \geq 2$. We have $(a^m)^0 = e = a^{m \cdot 0}$ and $(a^m)^1 = a^m = a^{m \cdot 1}$. Moreover,

we have

$$\begin{aligned}
(a^m)^{n+2} &= (a^m)^n h_{a^m} = a^{m \cdot n} h_a^m \\
&= a^{m \cdot n - 2 \cdot \lfloor \frac{m \cdot n}{2} \rfloor} h_a^{\lfloor \frac{m \cdot n}{2} \rfloor} h_a^m \\
&= a^{m \cdot n - 2 \cdot \lfloor \frac{m \cdot n}{2} \rfloor + m} \\
&= a^{m \cdot n + 2m} \\
&= a^{m \cdot (n+2)},
\end{aligned}$$

as claimed.

(d) For $n = 0$ and $n = 1$ this is [13, Lemma 3.5 (a) and (c)]. By [4, Proposition 5.8 (3)] the maps h_a and h_{e+a} commute. Hence h_a commutes with $h_{a,e} = h_{a+e} - h_a - h_e = h_{a+e} - h_a - \text{id}_U$. Thus the claim follows by induction.

(e) This is also clear for $n = 0$ and $n = 1$. Suppose the claim is true for n . Since λ is contained in the centroid, we have

$$(\lambda a)^{n+2} = (\lambda a)^n h_{\lambda a} = \lambda^n a^n h_{\lambda a} = \lambda^n \lambda^2 a^n h_a = \lambda^{n+2} a^{n+2}.$$

This proves the claim. □

We will denote elements of the form $a^2 = eh_a$ as *squares*.

Lemma 3.6 *If $\text{char}U \neq 2$, then the group U is generated by squares.*

Proof. We have $2a = eh_{a,e} = eh_{a+e} - eh_a - eh_e = (a+e)^2 - a^2 - e^2$ by 3.5(d). Since U is uniquely 2-divisible, the claim follows. □

Lemma 3.7 *For $a \in U^*$ and $b \in U$ we have $a^{-1}h_{a,b} = 2b$.*

Proof. By [4, Lemma 5.7(2)] we have $a\tau h_{a,b} = -2b$. Thus the claim follows. □

Lemma 3.8 *For $a \neq e$ we have $(e-a)^{-1}h_{a,b} = ((e-a)^{-1} - e)h_{e,b}$.*

Proof. For $a \neq 0$ we have

$$a^{-1}h_{e-a,b} = a^{-1}h_{e,b} + a^{-1}h_{-a,b} = a^{-1}h_{e,b} - 2b = a^{-1}h_{e,b} - eh_{e,b} = (a^{-1} - e)h_{e,b}$$

by [4, 5.10(1)], 3.7 and 3.5(d). Replacing a by $e - a$ yields

$$(e-a)^{-1}h_{a,b} = ((e-a)^{-1} - e)h_{e,b},$$

as desired. □

Proposition 3.9 *Suppose that $v : \mathbb{k} \rightarrow \mathbb{R} \cup \{\infty\}$ is a valuation and that B is a basis of U . For $a = \sum_{b \in B} \lambda_b \cdot b \in U$ set $\omega(a) := \min\{v(\lambda_b) \mid b \in B\}$. Suppose further that all Hua maps are continuous with respect to ω . For all $t \in \mathbb{k} \cap \mathcal{C}$ with $v(t) > 0$ and all $a \in U$ with $\omega(a^n) = 0$ for all $n \in \mathbb{N}$ we have*

$$(e - ta)^{-1} = \sum_{n=0}^{\infty} t^n a^n.$$

Proof. For all $n \in \mathbb{N}$ we have by 3.5(d) and (e).

$$\begin{aligned} ((e - ta)^{-1} - \sum_{k=0}^n t^k a^k) h_{e-ta} &= (e - ta)^{-1} h_{e-ta} - \sum_{k=0}^n t^k a^k h_{e-ta} \\ &= e - ta - \sum_{k=0}^n t^k a^k h_{e-ta} \\ &= e - ta - \sum_{k=0}^n t^k a^k h_e - \sum_{k=0}^n t^k a^k h_{e,-ta} - \sum_{k=0}^n t^k a^k h_{-ta} \\ &= e - ta - \sum_{k=0}^n t^k a^k - \sum_{k=0}^n (ta)^k h_{e,-ta} - \sum_{k=0}^n t^k t^2 a^k h_a \\ &= -2ta - \sum_{k=2}^n t^k a^k + 2 \sum_{k=1}^{n+1} t^k a^k - \sum_{k=2}^{n+2} t^k a^k \\ &= t^{n+1} a^{n+1} - t^{n+2} a^{n+2}. \end{aligned}$$

It follows that $\omega\left(\left((e - ta)^{-1} - \sum_{k=0}^n t^k a^k\right) h_{e-ta}\right) \geq (n+1) \cdot v(t)$. Thus the sequence $\left(\left((e - ta)^{-1} - \sum_{k=0}^n t^k a^k\right) h_{e-ta}\right)_{n \geq 0}$ converges to 0. Since $h_{e-ta}^{-1} = h_{(e-ta)^{-1}}$ is continuous, we also have $\lim_{n \rightarrow \infty} (e - ta)^{-1} - \sum_{k=0}^n t^k a^k = 0$, thus $\sum_{n=0}^{\infty} t^n a^n = (e - ta)^{-1}$. \square

We are now able to prove our main result:

Theorem 3.10 *Suppose that there is a subfield \mathbb{k} of $\text{End}_H(U)$ with*

- (i) $\dim_{\mathbb{k}} U < \infty$,
- (ii) $|\mathbb{k} \cap \mathcal{C}| = \infty$,
- (iii) U is generated by the set of squares $\{a^2 \mid a \in U\}$ as a $\text{End}_H(U)$ -module and
- (iv) there is $\lambda \in \mathbb{k}^* \cap \mathcal{C}$ such that $\lambda - 1 \in \mathbb{k}^* \cap \mathcal{C}$ as well.

Then (U, \mathcal{H}, e) is a quadratic Jordan division algebra.

Proof. We embed $\mathbb{M}(U, \tau)$ in its ultrapower $\mathbb{M}(\tilde{U}, \tilde{\tau})$, where $\tilde{U} := U^{\mathbb{N}}/\mathcal{F}$ for some free ultrafilter \mathcal{F} on \mathbb{N} . Moreover, let $\tilde{\mathbb{k}} := \mathbb{k}^{\mathbb{N}}/\mathcal{F}$. Let $n \mapsto t_n$ be an injective map from \mathbb{N} to $\mathbb{k} \cap \mathcal{C}$ and set $t := [(t_n)_{n \in \mathbb{N}}] \in \tilde{\mathbb{k}}$. Then t is contained in the centroid of $\mathbb{M}(\tilde{U}, \tilde{\tau})$. By 2.6(b) we have a valuation $v : \tilde{\mathbb{k}} \rightarrow \mathbb{R} \cup \{\infty\}$ that is trivial on \mathbb{k} with $v(t) = 1$.

Let B be a \mathbb{k} -basis of U . Then B is also a \mathbb{k} -basis of \tilde{U} . The map ω and the lattices Λ_r are defined as in Notation 3.2.

Let $a, b \in U$. Using Lemma 3.8 for ta in place of a and Proposition 3.9 we have

$$\left(\sum_{n=0}^{\infty} t^n a^n\right) h_{b,ta} = (e - ta)^{-1} h_{b,ta} = ((e - ta)^{-1} - e) h_{b,e} = \left(\sum_{n=1}^{\infty} t^n a^n\right) h_{b,e}.$$

Since the maps $h_{b,ta}$ and $h_{e,b}$ are linear and thus continuous, we get

$$\sum_{n=0}^{\infty} t^n a^n h_{b,ta} = \sum_{n=1}^{\infty} t^n a^n h_{b,e}. \quad (1)$$

For all $n \geq 1$ we have $a^n h_{b,e} \in U$ and thus $\omega(t^n a^n h_{b,e}) \geq n$. Since the series $\sum_{n=0}^{\infty} t^n a^n h_{b,ta}$ converges, the sequence $(t^n a^n h_{b,ta})_{n \geq 0}$ converges to 0. Thus there is $N \in \mathbb{N}$ such that $\omega(t^n a^n h_{b,ta}) \geq 1$ for all $n > N$. Since $\omega(\sum_{n=0}^{\infty} t^n a^n h_{b,ta}) = \omega(\sum_{n=1}^{\infty} t^n a^n h_{b,e}) \geq \min\{\omega(t^n a^n h_{b,e}) \mid n \geq 1\} \geq 1$, we also have $\omega(\sum_{n=0}^N t^n a^n h_{b,ta}) \geq 1$. Therefore we have $\sum_{n=0}^N t^n a^n h_{b,ta} \in \Lambda_1$.

Now let $\lambda \in \mathcal{C} \cap \mathbb{k}$. We have $t^n (\lambda a)^n h_{\lambda ta, \lambda b} = t^n \lambda^{n+2} a^n h_{a,b}$. Replacing a by λa and b by λb in (1), we get

$$\sum_{n=0}^{\infty} t^n \lambda^{n+2} a^n h_{\lambda ta, \lambda b} = \sum_{n=0}^{\infty} t^n (\lambda a)^n h_{\lambda ta, \lambda b} = \sum_{n=1}^{\infty} t^n \lambda^n a^n h_{\lambda b, e}.$$

Since $\omega(t^n \lambda^{n+2} a^n h_{\lambda ta, \lambda b}) = \omega(t^n a^n h_{\lambda ta, \lambda b})$, we conclude that $\sum_{n=0}^N t^n \lambda^{n+2} a^n h_{\lambda ta, \lambda b} \in \Lambda_1$. Dividing by λ^2 yields $\sum_{n=0}^N t^n \lambda^n a^n h_{\lambda ta, \lambda b} \in \Lambda_1$. Since $|k \cap \mathcal{C}| \geq N+1$, Lemma 3.1 yields that $t^n a^n h_{\lambda ta, \lambda b} \in \Lambda_1$ for all $n \geq 0$.

Now let $m \geq 1$. Using the result above for a^{2m+2} instead of a and bh_a^m instead of b for $n = 0$, we get

$$a^{2m} h_{ta^2, b} h_a^m = e h_a^m h_{ta^2, b} h_a^m = e h_{ta^2} h_a^m, b h_a^m = e h_{ta^{m+2}, b} h_a^m \in \Lambda_1,$$

thus $(a^2)^m h_{ta^2, b} = a^{2m} h_{ta^2, b} \in \Lambda_1$ since h_a^m is an isometry. Replacing a by a^2 in (1) yields

$$\sum_{n=0}^{\infty} t^n (a^2)^n h_{b, ta^2} = \sum_{n=1}^{\infty} t^n (a^2)^n h_{b, e}.$$

Since $\omega(t^n (a^2)^n h_{b, ta^2}) \geq 2$ for all $n \geq 1$ and $\omega(t^n (a^2)^n h_{b, e}) \geq 2$ for all $n \geq 2$, we get $e h_{ta^2, b} \equiv ta^2 h_{e, b} \pmod{\Lambda_2}$.

Let $\lambda \in \mathcal{C} \cap \mathbb{k}^*$ such that $\lambda - 1 \in \mathbb{k}^* \cap \mathcal{C}$. Replacing t by λt and b by λb we get

$$\lambda ta^2 h_{e, \lambda b} \equiv e h_{\lambda ta^2, \lambda b} \equiv \lambda^2 e h_{ta^2, b} \equiv \lambda^2 ta^2 h_{e, b} \pmod{\Lambda_2}.$$

Dividing by λt yields

$$a^2 h_{e, \lambda b} \equiv \lambda a^2 h_{e, b} \pmod{\Lambda_1}.$$

Since the map $x \mapsto x + \Lambda_1$ from U to Λ_0/Λ_1 is injective, we $a^2 h_{e, \lambda b} = \lambda a^2 h_{e, b}$. Since U is generated by the squares as a $\text{End}_H(U)$ -module, we get $a h_{e, \lambda b} = \lambda a h_{e, b}$. Now let $0 \neq c \in U$. The same argument for the c -isotope yields that

$ah_c^{-1}h_{c,\lambda b} = \lambda ah_c^{-1}h_{c,b}$. Replacing a by ah_c , we get that $ah_{\lambda b,c} = \lambda ah_{b,c}$ for all $a, b, c \in U$ with $c \neq 0$. This equation is trivially also true for $c = 0$. Thus the claim follows by 1.11. \square

By the classification of quadratic Jordan division algebras in [24, Theorem 15.7], a quadratic Jordan division algebra over a field of characteristic not 2 is a skewfield, a Jordan algebra of Clifford type for an anisotropic quadratic form, an involutory set or an Albert division algebra. The corresponding little projective is a PSL_2 over a skewfield, an orthogonal group, a unitary group or an exceptional algebraic group of type $E_{7,1}^{78}$. Therefore, we can formulate our result in the language of group theory. First, we need two auxiliary results.

Lemma 3.11 *Let G be a doubly transitive permutation group on a set X such that for some $y \in X$ there is a normal subgroup U_y of G_y acting regularly on $X \setminus \{y\}$. For every $x \in X$ choose $g_x \in G$ with $yg_x = x$ and define $U_x := U_y^{g_x}$. Then U_x does not depend of the choice of g_x , and $(X, (U_x)_{x \in X})$ is a Moufang set whose little projective group G^\dagger is normal in G .*

Proof. If h_x is another element with $yh_x = x$, then $h_x g_x^{-1} \in G_y$, so $U_y^{h_x g_x^{-1}} = U_y$, hence $U_y^{h_x} = U_y^{g_x}$. This proves the first part. For $x, z \in X$ and $g \in U_x$ we therefore have $U_z^g = U_y^{g_y g} = U_{y g_y g} = U_{zg}$. This proves that $(X, (U_x)_{x \in X})$ is a Moufang set. Moreover, $\{U_x \mid x \in X\}$ is conjugation class of subgroups in G , hence $G^\dagger = \langle U_x \mid x \in X \rangle$ is normal in G . \square

Lemma 3.12 *Let $\mathbb{M}(U, \tau)$ be a proper Moufang set with U abelian. Let G^\dagger be its little projective group. Suppose that $G^\dagger \trianglelefteq G \leq \mathrm{Sym}(X)$. Then $Z(G_{\infty,0})$ is contained in the centroid of $\mathbb{M}(U, \tau)$.*

Proof. Let $Z := Z(G_{\infty,0})$ and H be the Hua subgroup of $\mathbb{M}(U; \tau)$. Then $Z \subseteq \mathrm{End}_H(U_\infty)$ since $H \leq G_{\infty,0}$. For every $a \in U^\#$, the element μ_a is contained in $G_{\{\infty,0\}}$ and therefore normalises Z . Since $\mu_a^2 = 1$, we have $h^{\mu_a} = h^{-1}$ for all $h \in Z$ or there is $1 \neq z \in Z$ with $z^{\mu_a} = z$. Suppose the latter case holds. Then for all $b \in U^\#$ we have $z = z^{\mu_a \mu_b} = z^{\mu_b}$ since $\mu_a \mu_b \in H \leq G_{\infty,0}$. Therefore we have $\mu_b = \mu_b^h = \mu_{bh}$ for all $b \in U^\#$. Thus we have $bh \in \{b, -b\}$ for all $b \in U$. Therefore $h = 1$ or $b^h = -b$ for all $b \in U$. In both cases we have $h^{-1} = h = h^{\mu_a}$.

Thus we have $h^{\mu_a} = h^{-1}$ for all $h \in Z$ and all $a \in U^\#$. We get

$$h_{ah} = \mu_e \mu_{ah} = \mu_e \mu_a^h = \mu_e h^{-1} \mu_a h = h \mu_e \mu_a h = h h_a h = h^2 h_a$$

\square

Corollary 3.13 *Let G be a group acting doubly transitively on a set X . Suppose that for some $x \in X$ there is an abelian subgroup $U_x \leq G_x$ with*

- (i) U_x acts regularly on $X \setminus \{x\}$,
- (ii) U_x does not contain an involution,

(iii) U_x is finitely generated as a $Z(G_{x,y})$ -module for all $y \in X \setminus \{x\}$.

Let G^\dagger be the subgroup of G generated by $\{U_x^g \mid g \in G\}$. Then $G^\dagger \trianglelefteq G$ and there is a commutative field \mathbb{k} such that one of the following holds:

(I) $\text{char}\mathbb{k} = 2$ and $G^\dagger \cong \text{Aff}(\mathbb{k})$.

(II) $\text{char}\mathbb{k} \neq 2$ and either G^\dagger is a classical group of relative \mathbb{k} -rank one or an exceptional algebraic group of type $E_{7,1}^{78}$.

Proof. By 3.11 $\mathbb{M} := (X, (U_x)_{x \in X})$ is a Moufang set and G^\dagger is normal in G . Suppose that \mathbb{M} is improper. Hence G^\dagger is sharply 2-transitive. Since U_x is abelian, by [28][Remark 2 on page 47/48] there is a commutative field \mathbb{k} such that $G^\dagger \cong \text{Aff}(\mathbb{k})$. Since $\mathbb{k}^* \cong U_x$ does not contain an involution, we have $\text{char}\mathbb{k} = 2$. Hence we are in case (I).

Now suppose that \mathbb{M} is proper. Since U_x is abelian, the Moufang set \mathbb{M} is proper by [26]. By [27] the group $G_{x,y}$ acts irreducibly on U_x , thus $K := \text{End}_{G_{x,y}}(U_x)$ is a skewfield by Schur's Lemma. Let \mathbb{k} be the subfield of K generated by $Z(G_{x,y})$. Then \mathbb{k} is commutative, and since there is no 2-torsion in U_x , we have $\text{char}\mathbb{k} \neq 2$. Suppose that $Z(G_{x,y})$ is finite. Then $Z(G_{x,y})$ is algebraic over the prime field of \mathbb{k} , and so \mathbb{k} is finite-dimensional over its prime field. Moreover, we have $\text{char}\mathbb{k} > 0$, since the map $n^2 \cdot \text{id}_U$ is contained in H for all $n \in \mathbb{N}$ with n relatively prime to the characteristic of U . Therefore \mathbb{k} is finite as well. Since U_x is finitely generated as a $Z(G_{x,y})$ -module, we have $\dim_{\mathbb{k}} U < \infty$, hence U is finite. By [25] we get $G^\dagger \cong \text{PSL}_2(F)$ for some finite field F with $\mathbb{k} \subseteq F$.

Therefore we may assume that $Z(G_{x,y})$ is infinite. By 3.12 and our main theorem we conclude $\mathbb{M}(U, \tau) \cong \mathbb{M}(J)$ for some quadratic Jordan division algebra J . If J is special, then G is a classical group. If J is exceptional, then G^\dagger is an exceptional algebraic group of type $E_{7,1}^{78}$ by [2, Remark after 2.29]. \square

Definition 3.14 (a) Let G be a group acting on a set X . Then G is called *Zassenhaus transitive* on X

- (i) G acts doubly transitively on X ,
- (ii) $G_{x,y} \neq 1$ but $G_{x,y,z} = 1$ for all pairwise distinct $x, y, z \in X$,
- (iii) for all $x \in X$ there is a normal subgroup U_x of G_x acting regularly on $X \setminus \{x\}$ and
- (iv) there is no regular normal subgroup N of G .

- (b) Let $\mathbb{M} = (X, (U_x)_{x \in X})$ be a Moufang set with little projective group G^\dagger . We call \mathbb{M} a *Zassenhaus Moufang set* if G^\dagger is Zassenhaus transitive on X .

Note that if $\mathbb{M}(U, \tau)$ is a Moufang set with Hua subgroup H , then $\mathbb{M}(U, \tau)$ is Zassenhaus if and only if $H \neq 1$ and H acts freely on U . If $\mathbb{M} = \mathbb{M}(J)$ for some quadratic Jordan division algebra (J, Q, e) , this means that if $x, x_1, \dots, x_n \in J^\#$ with $xQ_{x_1} \dots Q_{x_n} = x$, then $yQ_{x_1} \dots Q_{x_n}$ for all $y \in J$.

The following statement is special case of [1, Theorem 4.8]. However, unlike this theorem it does not use the classification of quadratic Jordan division algebras.

Theorem 3.15 *Let (J, Q, e) be a quadratic Jordan division algebra such that $\mathbb{M}(J)$ is Zassenhaus. Then $Q_a Q_b = Q_b Q_a$ for all $a, b \in J$.*

Proof. We will make use of the identities in [14, Chapter 1, Section 2 and 3]. Note that $V_{a,b}$ in our notation corresponds to $V_{b,a}$ in Jacobson's notation. As Jacobson we set $V_a := V_{a,e} = V_{e,a} = Q_{e,a}$ for $a \in J$.

Let $a, b \in J^\#$. By identity QJ4' on [14, p. 18] we have

$$V_{b,a} Q_a = Q_a V_{a,b} = Q_{a,b} Q_a.$$

Applying Q_a^{-1} on the left and right and using (QJ3) we get

$$Q_a^{-1} V_{b,a} = V_{a,b} Q_a^{-1} = Q_a^{-1} Q_{a,b} Q_a Q_a^{-1} = Q_{a^{-1},b}.$$

Thus

$$e Q_{a,b} Q_a^{-1} = e V_{a,b} Q_a^{-1} = e Q_{a^{-1},b}.$$

Therefore we get

$$e Q_{a,b} Q_a^{-1} Q_b^{-1} = e Q_{a^{-1},b^{-1}} = e Q_{a,b} Q_b^{-1} Q_a^{-1}.$$

If $e Q_{a,b} \neq 0$, this implies $Q_a Q_b = Q_b Q_a$.

Now suppose that $e Q_{a,b} = 0$. We will show that $Q_a Q_b = Q_b Q_a$ is also true in this case. Since $e Q_{a,b} = 0 = e Q_{a,-b}$, we have $e Q_{a+b} = e Q_a + e Q_b = e Q_{a-b}$, thus $Q_{a+b} = Q_{a-b}$ and therefore $2Q_{a,b} = 0$. Moreover, we have

$$a^{-1} Q_{a,b} = a V_{b,a^{-1}} = a^{-1} Q_{a^{-1}}^{-1} V_{b,a^{-1}} = a^{-1} V_{a^{-1},b} Q_a = b Q_{a^{-1},a^{-1}} Q_a = 2b Q_a Q_a^{-1} = 2b,$$

therefore we only have to deal with characteristic 2.

By identity QJ18 of [14] we have $a Q_b V_a + a V_b V_{b,a} = a V_{b,a} V_b + a V_a Q_b$. Since $a V_b = 0$, $a V_a = e Q_{a,a} = 2e Q_a = 0$ and $a V_{a,b} = b Q_{a,a} = 2b Q_a = 0$ we have $e Q_{a Q_b, a} = a Q_b V_a = 0$. By identity QJ30 of [14] we have

$$0 = e Q_{e Q_{a,b}} = e Q_a Q_b + e Q_b Q_a + e Q_{a,a} Q_b,$$

thus $e Q_a Q_b = e Q_b Q_a$. Thus the claim follows. \square

Corollary 3.16 *If $\mathbb{M}(J)$ is Zassenhaus, then either J is a commutative field or there is a commutative field \mathbb{k} with $\mathbb{k}^2 \subseteq J \subseteq \mathbb{k}$.*

Proof. This follows by the main theorem of [10]. \square

Corollary 3.17 *Suppose that G acts Zassenhaus transitively on a set X and that for all $x, y \in X$ pairwise distinct*

- (i) *the group U_x is abelian, but not elementary-abelian of exponent 2 and*
- (ii) *U_x is finitely generated as a $Z(G_{x,y})$ -module.*

Then there is a field F with $\text{char} F \neq 2$ and a group $\text{PSL}_2(F) \leq \mathcal{G} \leq \text{P}\Gamma\text{L}_2(F)$ such that (G, X) and $(\mathcal{G}, \text{PG}_1(F))$ are isomorphic as permutation groups, where $\text{PG}_1(F)$ denotes the projective line over F .

Proof. Since G is Zassenhaus transitive on X , there is no regular normal subgroup of G , so case (I) of 3.13 can be excluded. By 3.16 there is a commutative field F with $\text{char} F \neq 2$ such that the corresponding Moufang set is isomorphic to the projective line over F . The little projective group G^\dagger is isomorphic to $\text{PSL}_2(F)$. Thus G is isomorphic to a subgroup of $\text{Aut}(\text{PSL}_2(F)) = \text{P}\Gamma\text{L}_2(F)$.

□

Remark 3.18 By [22], the group \mathcal{G} is contained in $\text{PGL}_2(F)$ if and only if G_{xy} is abelian.

4 KT-nearfields

KT-nearfields are the algebraic structures for Moufang sets whose little projective group is contained in a sharply 3-transitive groups. At first, we recall some notion about nearfields. For an introduction to nearfields see [30].

Definition 4.1 Let F be a set with two binary operations $+$ and \cdot and $0, 1 \in F$ two distinct elements. Then $(F, +, \cdot, 0, 1)$ is called a (*right*) *nearfield* if

- (a) $(F, +, 0)$ is a group.
- (b) $(F^*, \cdot, 1)$ is a group, where $F^* := F \setminus \{0\}$.
- (c) $(a + b) \cdot c = a \cdot c + b \cdot c$ for all $a, b, c \in F$.
- (d) $a \cdot 0 = 0$ for all $a \in F$.

It can be seen that the additive group of a nearfield is necessarily abelian, see [30][2.3].

Definition 4.2 Let F be a nearfield.

- (a) $Z(F) := \{a \in F \mid a \cdot b = b \cdot a \text{ for all } b \in F\}$ is called the *centre* of F .
- (b) $\mathbb{k}(F) := \{a \in F \mid a \cdot (b + c) = a \cdot b + a \cdot c \text{ for all } b, c \in F\}$ is called the *kernel* of F .

One easily sees that $\mathbb{k}(F)$ is a skewfield containing $Z(F)$ and that F is a left vector space over $\mathbb{k}(F)$.

Example 4.3 (a) Let F be a skewfield and $\varphi : F^* \rightarrow \text{Aut}(F)$ a map with $\varphi(a^{\varphi(b)}) = \varphi(a)\varphi(b)$ for all $a, b \in F^*$. Such a map is called a *coupling*. We define a new multiplication \cdot on F by

$$a \cdot b := \begin{cases} a^{\varphi(b)}b, & \text{if } b \in F^*, \\ 0, & \text{if } b = 0 \end{cases}$$

for all $a, b \in F$. Then $F^\varphi := (F, +, \cdot, 0, 1)$ is nearfield, called a *Dickson nearfield*.

- (b) A nearfield that is not a Dickson nearfield is called *wild*. There are up to isomorphism exactly seven finite wild nearfields. Infinite wild nearfields were constructed in [31], [8], [9] and [12].

Definition 4.4 Let F be a nearfield. For $a \in F^*$ and $b \in F$ let $t_{a,b} : F \rightarrow F : x \mapsto ax + b$. Set $\mathcal{T}(F) := \{t_{1,b} \mid b \in F\}$ and $\text{Aff}(F) := \{t_{a,b} \mid a \in F^*, b \in F\}$, the *affine group* of F .

We have

Theorem 4.5 (a) $\text{Aff}(F)$ is a subgroup of $\text{Sym}(F)$ that acts sharply 2-transitively on F .

(b) $\mathcal{T}(F)$ is a normal subgroup of $\text{Aff}(F)$ that acts regularly on F . Moreover, we have $\mathcal{T}(F) \cong (F, +)$.

On the converse, we have

Theorem 4.6 Let X be set, $G \leq \text{Sym}(X)$ a group acting sharply 2-transitively on X and N a normal subgroup of G acting regularly on X . Then there is a nearfield F , such that (G, X) and $(\text{Aff}(F))$ are isomorphic as permutation group. Moreover, if $\varphi : G \rightarrow \text{Aff}(F)$ is an isomorphism, then $\varphi(N) = \mathcal{T}(F)$.

Proofs of these well-known results can for instance be found in [19].

The question whether the sharply 2-transitive action of $\text{Aff}(F)$ on F can be extended to a sharply 3-transitive action leads to the definition of a KT-nearfield.

Definition 4.7 Let F be a nearfield. An involutory automorphism σ of the multiplicative group of F is called a *KT-automorphism* if $(1+a^\sigma)^\sigma = 1 - (1+a)^\sigma$ for all $a \in F^* \setminus \{-1\}$. The pair (F, σ) is called a *KT-nearfield*.

The abbreviation *KT* stands for *Karzel-Tits*.

Example 4.8 Let F be a commutative field, $\varphi : F^* \rightarrow \text{Aut}(F)$ a coupling with $\varphi(a) = \varphi(a^{-1})$ for all $a \in F^*$. Let $a^\sigma := a^{-1}$ (the inverse with respect to the field multiplication) for all $a \in F^*$. Then (F^φ, σ) is a KT-nearfield.

Definition 4.9 Let (F, σ) be a KT-nearfield. Set $X := F \dot{\cup} \{\infty\}$. We embed $\text{Aff}(F)$ to $\text{Sym}(X)$ by setting $\infty g := \infty$ for all $g \in \text{Aff}(F)$. Let $\tau \in \text{Sym}(X)$ be defined by

$$x\tau := \begin{cases} 0, & \text{if } x = \infty, \\ \infty, & \text{if } x = 0, \\ -x^\sigma, & \text{if } x \in F^*. \end{cases}$$

We set $T_3(F) := \langle \tau, \text{Aff}(F) \rangle \leq \text{Sym}(X)$.

Definition 4.10 Let (F, σ) be a KT-nearfield. For $a \in F^*$ let $q_a := a^{-\sigma}a$, the *pseudo-square* associated to a .

Theorem 4.11 Let (F, σ) be a KT-nearfield.

(a) $T_3(F)$ is a sharply 3-transitive permutation group on X with $T_3(F)_\infty = \text{Aff}(F)$ and $T_3(F)_{0,\infty} \cong F^*$.

(b) For all $x \in X$ let U_x be a conjugate of $\mathcal{T}(F)$ in $T_3(F)_x$. Then U_x is the unique conjugate of $\mathcal{T}(F)$ in $T_3(F)_x$ and $(X, (U_x)_{x \in X})$ is a special Moufang set whose little projective group G^\dagger is normal in $T_3(F)$. Moreover, for all $a, x \in F$ with $a \neq 0$ we have $xh_a = x \cdot q_a$.

Proof.

- (a) This follows by [20].
- (b) If U_x and V_x are two conjugates of $\mathcal{T}(F)$ contained in $T_3(F)_x$, then both groups are abelian normal subgroups $T_3(F)_x$ acting regularly on $X \setminus \{x\}$. Since $T_3(F)_x$ acts sharply 2-transitively on $X \setminus \{x\}$, it follows that $U_x = V_x$. Thus for every $x \in X$ there is a unique normal abelian subgroup U_x of $T_3(F)_x$, hence $\{U_x \mid x \in X\}$ is a conjugacy class of subgroups in $T_3(F)$. It follows that $G^\dagger = \langle U_x \mid x \in X \rangle$ is normal in $T_3(F)$.

Since τ interchanges 0 and ∞ , the Moufang sets $(X, (U_x)_{x \in X})$ and $\mathbb{M}(U, \tau)$ are isomorphic Moufang sets. For $a \in U^\#$, we have $(-a)\tau = (-1 \cdot a)\sigma = (-1)\sigma \cdot a\sigma = -a\sigma = -a\tau$. Thus $(X, (U_x)_{x \in X})$ is special, and $U_x \cong (F, +)$ is abelian for all $x \in X$. We have

$$\begin{aligned}
xh_a &= x\tau\alpha_a\alpha_{-a\tau^{-1}}\alpha_{-(-a\tau^{-1})\tau} \\
&= ((x\tau + a)\tau^{-1} - a\tau^{-1})\tau + a \\
&= -(a^\sigma - (a - x^\sigma)^\sigma)^\sigma + a \\
&= ((1 - x^\sigma \cdot a^{-1})^\sigma \cdot a^\sigma - a^\sigma)^\sigma + a \\
&= ((1 - (x \cdot a^{-\sigma})^\sigma)^\sigma - 1)^\sigma \cdot a^{\sigma^2} + a \\
&= (1 - (1 - x \cdot a^{-\sigma})^\sigma - 1) \cdot a + a \\
&= (x \cdot a^{-\sigma} - 1)^{\sigma^2} \cdot a + a \\
&= x \cdot a^{-\sigma} \cdot a - a + a \\
&= x \cdot q_a.
\end{aligned}$$

□

Example 4.12 If F is a commutative field and φ is a coupling as in Example 4.3, then we have $\mathrm{PSL}_2(F) \leq T_3(F^\varphi) \leq \mathrm{P}\Gamma\mathrm{L}_2(F)$. For example, if $F = \mathbb{F}_9$ and φ is the unique non-trivial coupling of F , then $T_3(F^\varphi) = \mathrm{Mat}_{10}$, the Mathieu group of order 10.

Recall that for $n \geq 1$ a permutation group is called n -sharp if the stabiliser of n distinct points is always trivial

Lemma 4.13 *Let G be a 2-sharp permutation group on a set X and A a regular subgroup of G . Suppose that there is $x \in X$ and $h_1, h_2, h_3 \in H := N_G(A) \cap G_x$ such that $a^{h_3} = a^{h_1}a^{h_2}$ for all $a \in A$. Then $Z(G_x) \leq H$.*

Proof. We write the group $A = (A, +, 0)$ additively, identify the set X with A , the element $x \in X$ with $0 \in A$ and the group A with the subgroup $\{\alpha_a \mid a \in A\}$ of $\mathrm{Sym}(A)$, where $b\alpha_a = b + a$ for all $a, b \in A$. Then we have $\alpha_{ah} = \alpha_a^h$ for all $a \in A$ and all $h \in H$. Thus $ah_3 = 0\alpha_{ah_3} = 0\alpha_a^{h_3} = 0\alpha_a^{h_1}\alpha_a^{h_2} = 0\alpha_{ah_1}\alpha_{ah_2} = ah_1 + ah_2$ for all $a \in A$.

Let $t \in Z(G_0)$ and $0 \neq a \in A$. Then $g_a := \alpha_a^t \alpha_{-at} \in G_0$. We have

$$\begin{aligned}
g_{ah_3} \alpha_{ath_1+ath_2} &= g_{ah_3} \alpha_{ath_3} \\
&= g_{ah_3} \alpha_{ah_3 t} \\
&= \alpha_{ah_3}^t \\
&= \alpha_{ah_1+ah_2}^t \\
&= \alpha_{ah_1}^t \alpha_{ah_2}^t \\
&= g_{ah_1} \alpha_{ah_1 t} g_{ah_2} \alpha_{ah_2 t} \\
&= g_{ah_1} \alpha_{ath_1} g_{ah_2} \alpha_{ath_2}.
\end{aligned}$$

Thus

$$ath_1 + ath_2 = 0g_{ah_3} \alpha_{ath_1+ath_2} = 0g_{ah_1} \alpha_{ath_1} g_{ah_2} \alpha_{ath_2} = ath_1 g_{ah_2} + ath_2$$

and hence $ath_1 = ath_1 g_{ah_2}$. Since G is 2-sharp, this means $g_{ah_2} = 1$. Replacing a by ah_2^{-1} yields $g_a = 1$. Therefore we have $\alpha_a^t = \alpha_{at}$ for all $a \in A$, thus $t \in H$.

□

Theorem 4.14 *Let $\mathbb{M}(U, \tau)$ be a special Moufang set whose little projective subgroup G^\dagger is contained in a sharply 3-transitive subgroup G of $\text{Sym}(X)$, where $X := U \dot{\cup} \{\infty\}$. Then $G^\dagger \trianglelefteq G$ and there is a KT -nearfield (F, σ) such that (X, G) and $(F \dot{\cup} \{\infty\}, T_3(F))$ are isomorphic permutation groups.*

Proof. By [6, Corollary 6.4(3)] the group U is abelian.

We claim that U_∞ is normal in G_∞ . Note that G_∞ acts sharply 2-transitively on U . Let J be the set of involutions of G_∞ and $\bar{J} := J \cup \{1\}$. If $\text{char} U = 2$, then $\bar{J} \cap G_\infty$ acts regularly on U by [15, (3.3)]. Since $U_\infty \subseteq \bar{J} \cap G_\infty$ is also regularly on U , we get that $U_\infty = \bar{J}$ is normal in G_∞ .

Now suppose that $\text{char} U = 3$. Set $C := \{x \in G \mid o(x) = 3\}$ and $\bar{C} := C \cup \{1\}$. By [15, (2.1)] is a conjugacy class of G . Therefore every element of C has exactly one fixed point. Thus $C \cap G_\infty$ is a conjugacy class of G_∞ . By [15, (3.1)(b)] the set $\bar{C} \cap G_\infty$ acts regularly on U . Since $U_\infty \subseteq \bar{C} \cap G_\infty$ also acts regularly on U , we have $U_\infty = \bar{C} \cap G_\infty$ and thus $U_\infty \trianglelefteq G_\infty$.

Now suppose that $\text{char} U \neq 2, 3$ and that there is an involution $t \in G_{0,\infty}$ normalising U_∞ . Then t acts freely as an automorphism of U . Thus $at = -a$ for all $a \in U$. Now U is uniquely 2-divisible, hence we have $a = (\frac{1}{2} \cdot a^{-1})^t + \frac{1}{2} \cdot a$ and hence $\alpha_a = \alpha_{-\frac{1}{2} \cdot a}^t \alpha_{\frac{1}{2} \cdot a} = t \alpha_{\frac{1}{2} \cdot a} \in t\bar{J}$. Since $t(J \cap G_\infty)$ acts regularly on U by [15, (3.1)(c)], thus $t(J \cap G_\infty) = U_\infty$. Hence condition (v) and (viii) of [15, (3.7)] apply, thus $U_\infty = (J \cap G_\infty)^2$ is normal in G_∞ by condition (i) of the same theorem. Therefore we show that there is an involution $t \in G_{0,\infty}$ normalising U_∞ .

Let \mathbb{k} be the prime field $\mathbb{M}(U, \tau)$ and let $e \in U^\#$ with $\mu_e = \tau$. For alle $r \in \mathbb{k}^*$ we have $h_{r \cdot e} = r^2 \cdot \text{id}_U$ by [4][Proposition 4.6(6)]. Suppose that $\text{char} U \equiv 1 \pmod{4}$. Then there is an element $r \in \mathbb{k}^*$ with $r^2 = -1$. Thus $h_{r \cdot e} = -\text{id}$ is an involution and we are done.

Now suppose $\text{char}U = 0$ or $\text{char}U > 5$. Since $|(\mathbb{k}^*)^2| \geq 3$, there are $u, v \in \mathbb{k}^*$ such that $u^2 \neq \pm v^2$. Set $\ell := u^2 - v^2$, $m := 2uv$ and $n := u^2 + v^2$. For all $x \in U$ we have $xh_{\ell \cdot e} + xh_{m \cdot e} = \ell^2 \cdot x + m^2 \cdot x = (u^2 - v^2)^2 \cdot x + 4u^2v^2 \cdot x = (u^2 + v^2)^2 \cdot x = xh_{n \cdot e}$. By [15] there is a unique involution t in $G_{0,\infty}$, and by 4.13 we have $t \in N_{G_\infty}(U_\infty)$. Thus $U_\infty \trianglelefteq G_\infty$ in all cases.

Therefore G is a sharply 3-transitive permutation group such that every point stabilizer has an abelian normal subgroup acting regularly on the other points. Thus there is a KT-nearfield (F, σ) such that $G \cong T_3(F)$ by [20]. Since G^\dagger is generated by the root groups, we have $G^\dagger \trianglelefteq G$. \square

Definition 4.15 Let (F, σ) be a KT-nearfield with kernel \mathbb{k} . We set $\mathbb{k}_\sigma := \mathbb{k} \cap (\mathbb{k}^*)^\sigma \cup \{0\}$.

Theorem 4.16 Let (F, σ) be KT-nearfield with kernel \mathbb{k} . Then \mathbb{k}_σ is a commutative subfield of \mathbb{k} such that $x^\sigma = x^{-1}$ for all $x \in \mathbb{k}_\sigma$. If $\text{char}F \neq 2$, then $\mathbb{k} = \mathbb{k}_\sigma$.

Proof. This follows by the main theorem of [16]. \square

We are now ready to prove Theorem 2.

Theorem 4.17 Let (F, σ) be a KT-nearfield with kernel \mathbb{k} such that

- (i) $\text{char}F \neq 2$ and $\dim_{\mathbb{k}} F < \infty$ or
- (ii) $\text{char}F = 2$, $\dim_{\mathbb{k}_\sigma} F < \infty$ and F is generated by the pseudo-squares as a \mathbb{k}_σ -module.

Then F is a Dickson nearfield as in Example 4.8.

Proof. This is well-known if F is finite, so we may assume that F is infinite. Hence \mathbb{k}_σ is infinite as well. Let $\mathbb{M}(F)$ be the Moufang set associated to (F, σ) and H its Hua subgroup. For $a \in \mathbb{k}$ let $\lambda_a : F \rightarrow F : x \mapsto a \cdot x$. Then $\lambda_s \in \text{End}_H(F)$. Moreover, if $a \in \mathbb{k}_\sigma$, then $q_a = a^2 \in Z(F)$ by part (ii) of the main theorem of [16]. Therefore, we have

$$\begin{aligned} xh_b\lambda_a &= xh_{a \cdot b} = x \cdot q_{a \cdot b} \\ &= x \cdot (a \cdot b)^{-\sigma} \cdot a \cdot b = x \cdot b^{-\sigma} \cdot a^2 \cdot b \\ &= x \cdot a^2 \cdot q_b = x\lambda_a^2 h_b. \end{aligned}$$

Therefore λ_a is contained in the centroid. Our main theorem now implies that $\mathbb{M}(F)$ is the Moufang set for a quadratic Jordan division algebra. By 3.15 the pseudo-squares generate an abelian subgroup of F^* . The claim now follows by [17, Theorem 1.3]. \square

The author likes to thank Theo Grundhöfer for pointing out the following corollary. Note that condition (ii) is slightly weaker than condition (ii) in 3.17 since $Z(G_{x,y}) \subseteq \text{End}_{G_{x,y}}(U_x)$.

Corollary 4.18 *Let G be a sharply 3-transitive permutation group on a set X . Assume that*

- (i) *for every $x \in X$ there is a normal abelian subgroup U_x of G_x such that U_x is regular on $X \setminus \{x\}$ and not of exponent 2 and*
- (ii) *for $x, y \in X$ distinct, the group U_x is finitely generated as a module over $\text{End}_{G_{x,y}}(U_x)$.*

Then there is a commutative field F with $\text{char}F \neq 2$ such that $\text{PSL}_2(F) \leq G \leq \text{P}\Gamma\text{L}_2(F)$.

Proof. We have $G = T_3(F)$ for some KT-nearfield (F, σ) . We choose an element $\infty \in X$ and identify F, U_∞ and $X \setminus \{\infty\}$. The group $G_{\infty,0}$ consists of all maps $\rho_a : F \rightarrow F : x \mapsto x \cdot a$ with $a \in F^*$. Let $T \in \text{End}_{G_{\infty,0}}(F)$ and set $a := 1T$. We claim that a is contained in the kernel of F and that $xT = a \cdot x$ for all $x \in F$. Indeed, we have $xT = 1\rho_x T = 1T\rho_x = a\rho_x = a \cdot x$ for all $x \in F$. Moreover, for $x, y \in F$ we have $a \cdot (x + y) = (x + y)T = xT + yT = a \cdot x + a \cdot y$, hence a is in the kernel of F . Conversely, if a is in the kernel of F , then the map $x \mapsto a \cdot x$ is contained in $\text{End}_{G_{\infty,0}}(F)$. Therefore by the previous theorem F is coupled to a commutative field which we also call F and the claim follows. \square

References

- [1] J. Anquela, T. Cortés, H. Petersson, *Commuting U -operators in Jordan algebras*, Trans. Am. Math. Soc. **366**, No. 11 (2014).
- [2] L. Boelaert, T. De Medts, A. Stavrova, *Moufang sets and structurable division algebras*, Mem. Am. Math. Soc. **1245** (2019).
- [3] C. C. Chang, H. J. Keisler, *Model theory* 3d rev. ed., Studies in Logic and the Foundations of Mathematics, **93**, Amsterdam etc.: North-Holland (1990).
- [4] T. De Medts, Y. Segev, *Identities in Moufang sets*, Trans. Amer. Math. Soc. **360**, no. 11, 5831-5852 (2008).
- [5] T. De Medts, Y. Segev, *A course on Moufang sets*, Innov. Incidence Geom. **9**, 79-122 (2009)
- [6] T. De Medts, Y. Segev, K. Tent, *Special Moufang sets, their root groups and their μ -maps*, Proc. London Math. Soc. (3) **96**, 767-791 (2008).
- [7] T. De Medts, R. Weiss, *Moufang sets and Jordan division algebras*, Math. Ann. **335**, no. 2, 415-433 (2006).
- [8] Th. Grundhöfer, *Five wild nearfields*, J. Group Theory **16**, No. 5, 707-718 (2013).
- [9] Th. Grundhöfer, M. Grüniger, *Wild nearfields of dimension 2 over rational function fields*, J. Geom. **107**, No. 2, 317-328 (2016).

- [10] M. Grüninger, *Special Moufang sets with abelian Hua subgroup*, J. Algebra **323**, No. 6, 1797-1801 (2010).
- [11] M. Grüninger, *On the identities defining a quadratic Jordan division algebra*, Arch. Math. **104**, no. 1, 11-24 (2015).
- [12] M. Grüninger, *New examples of wild nearfields via regular matrix groups over skew polynomial rings*, Beitr. Algebra Geom. **57**, No. 4, 789-808 (2016).
- [13] M. Grüninger, *Special Moufang sets coming from quadratic Jordan division algebras*, Trans. Am. Math. Soc. **375**, no. 11, 7831-7852 (2022).
- [14] N. Jacobson, *Lectures on Quadratic Jordan algebras*, Lectures on Mathematics and Physics. Mathematics **45**. Bombay: Tata Institute of Fundamental Research (1969).
- [15] W. Kerby, *On infinite multiply transitive groups*, Hamburger Mathematische Einzelschriften. Neue Folge. Heft 6. Göttingen: Vandenhoeck & Ruprecht (1974).
- [16] W. Kerby, *A structure theorem for KT -nearfields*, Aequationes Math. **31**, 247-252 (1986).
- [17] W. Kerby, *On a Class of Sharply 3-transitive Groups*, Abh. Math. Sem. Hamburg **61**, 1-13 (1991).
- [18] W. Kerby, *KT -nearfields of rank 2*, Aequationes Math. **41**, No. 2-3, 187-191 (1991).
- [19] W. Kerby, H. Wefelscheid, *Bemerkungen über Fastbereiche und scharf zweifach transitive Gruppen*, Abh. Math. Semin. Univ. Hamb. **37**, 20-29 (1972).
- [20] W. Kerby, H. Wefelscheid, *Über eine scharf 3-fach transitiven Gruppe zugeordnete algebraische Struktur*, Abh. Math. Sem. Univ. Hamburg **37**, 225-235 (1972).
- [21] S. Lang, *Algebra*, 3rd revision revision, Undergraduate texts in Mathematics **211**, New York, NY: Springer Verlag (2005)..
- [22] H. Mäurer, *Eine Charakterisierung der zwischen $PS^{-L}(2, K)$ und $PGL(2, K)$ liegenden Permutationsgruppen*, Arch. Math. **40**, 405-411 (1983).
- [23] K. McCrimmon, *A general theory of Jordan rings*, Proc. Natl. Acad. Sci. USA **56**, 1072-1079 (1966).
- [24] K. McCrimmon, E. Zel'manov, *The structure of strongly prime quadratic Jordan algebras*. Adv. Math. **69**, No. 2, 133-222 (1988).
- [25] Y. Segev, *Finite special Moufang sets in odd characteristic*, Commun. Contemp. Math. **10**, No. 3, 455-475 (2008).
- [26] Y. Segev, *Proper Moufang sets with Abelian root groups are special*, J. Am. Math. Soc. **22**, No. 3, 889-908 (2009).

- [27] Y. Segev, R. Weiss, *On the action of the Hua subgroups in special Moufang sets*, Math. Proc. Cambridge Philos. Soc. **144**, no. 1, 77-84 (2008).
- [28] J. Tits, *Généralisations des groupes projectifs basés sur leurs propriétés de transitivité*. Acad. Roy. Belgique. Cl. Sci. Mém. Coll. 16, 18 in 8°, **27**, no. 2, 115 pages. (Collected Works Vol. I, EMS 2013).
- [29] J. Tits, *Twin buildings and groups of Kac-Moody type*, in Groups, combinatorics & geometry (Durham, 1990), 249-286, London Math. Soc. Lecture Note Ser. **165**, Cambridge Univ. Press, Cambridge, 1992.
- [30] H. Wähling, *Theorie der Fastkörper*, Thales Monographs **1**, Essen: Thales Verlag (1987).
- [31] H. Zassenhaus, *On Frobenius groups II: Universal completion of nearfields of finite degree over a field of reference.*, Result. Math. **11**, 317-358 (1987).