## Quantum thermalization of translation-invariant systems at high temperature

Saúl Pilatowsky-Cameo<sup>1</sup> and Soonwon Choi<sup>1</sup>

<sup>1</sup>Center for Theoretical Physics, Massachusetts Institute of Technology, Cambridge, MA 02139, USA

Quantum thermalization describes how closed quantum systems can effectively reach thermal equilibrium, resolving the apparent incongruity between the reversibility of Schrödinger's equation and the irreversible entropy growth dictated by the second law of thermodynamics. Despite its ubiquity and conceptual significance, a complete proof of quantum thermalization has remained elusive for several decades. Here, we prove that quantum thermalization must occur in any qubit system with local interactions satisfying three conditions: (i) high effective temperature, (ii) translation invariance, and (iii) no perfect resonances in the energy spectrum. Specifically, we show that a typical, initially unentangled pure state drawn from any ensemble with maximum entropy becomes locally indistinguishable from a Gibbs state upon unitary evolution. Our proof relies on a recent breakthrough in quantum information theory proving the separability of high-temperature thermal states, as well as on new technical results identifying sufficient conditions for quantum thermalization, whose applicability extends beyond our main result. Our work illustrates that statistical physics can be understood as an emergent phenomenon, explicitly derived from the first principles of quantum mechanics.

Quantum and statistical mechanics are among the most successful theories of modern physics, describing the microscopic dynamics at small scales and the macroscopic emergent phenomena at large scales. However, these theories display an incongruity: the Schrödinger equation always generates reversible time evolution, whereas the second law of thermodynamics prescribes an irreversible entropy growth. This apparent paradox is resolved by the idea of quantum thermalization [1-5], which asserts that even though the time-evolved quantum state has no entropy globally, it locally yields the same measurement outcomes as the maximally-entropic thermal distribution. This nominal resolution, however, brings more questions than answers. What are the physical conditions that lead to quantum thermalization? What is its underlying mechanism? Resolving these is arguably the most important challenge in quantum statistical physics and could provide new insights in related fields ranging from quantum computing [6] to quantum gravity [7].

Proving the emergence of statistical mechanics in quantum systems has remained an open problem for almost a century [8], owing to several challenges. To begin with, we do not even understand under what general conditions quantum thermalization occurs. Known violations include localization [9], Bethe-ansatz integrability [10], quantum many-body scars [11], and Hilbertspace fragmentation [12–14]. Furthermore, the exponentially growing Hilbert-space dimension and computational complexity make it challenging to probe the late-time dynamics of generic systems. These challenges have been attacked by a multitude of complementary approaches, including devising conjectures such as the eigenstate thermalization hypothesis (ETH) [15, 16], performing numerical or quantum simulations [17, 18], and rigorous approaches relying on typicality arguments, added assumptions, and/or symmetries [19–27].

Here, we prove that quantum thermalization must oc-



FIG. 1. Quantum thermalization predicts the local indistinguishability between time-evolved quantum many-body states (left) and the Gibbs states at thermal equilibria (right), establishing the bridge between quantum and statistical physics.

cur in a large class of physical systems at high temperature. Specifically, we consider systems consisting of many qubits (or spin-1/2 particles) on lattices in any dimension, satisfying three conditions: (i) high effective temperature, (ii) translation invariance, and (iii) no perfect resonances in the energy spectrum, so-called nondegenerate spectral gap condition. For all such systems, we prove that typical, initially unentangled states drawn from a maximally entropic pure-state ensemble thermalize. That is, upon unitary time evolution generated by a local Hamiltonian H, the quantum state

$$\left|\psi(t)\right\rangle = e^{-iHt/\hbar}\left|\psi\right\rangle \tag{1}$$

becomes locally indistinguishable from a thermal Gibbs state

$$g_{\beta} = \frac{e^{-\beta H}}{\operatorname{tr}(e^{-\beta H})},\tag{2}$$

at a corresponding inverse temperature  $\beta = 1/(k_B T)$ (Fig. 1). Hereafter, we work with units where the Boltzmann constant  $k_B$  and the Planck constant  $\hbar$  equal 1.

Our proof idea can be summarized as follows. First, the high-temperature condition (i) guarantees that the Gibbs state  $g_{\beta}$  satisfies two key properties: finite correlation length [28] and separability [29]. Separability means that there exists an ensemble of unentangled (product) pure states  $|\psi\rangle$  which averages to the Gibbs state  $\mathbb{E}_{\psi}[|\psi\rangle\langle\psi|] = g_{\beta}$ . Then, by adapting previous results (the weak ETH [25, 30, 31] and quantum equilibration theorems [32]), we develop a new theorem stating that under the nondegenerate gap condition (ii), any typical state in such an ensemble would thermalize as long as the state is formed by the superposition of sufficiently many different energy eigenstates. Crucially, this must be the case: since the energy eigenstates are necessarily translation symmetric (iii), the only way to construct a typical product state is to superpose a large number of eigenstates, as we rigorously quantify below.

Our results bring both conceptual implications and technical innovations to the field of quantum statistical physics. The most prominent is the establishment of thermalization — the foundational principle of statistical mechanics — as an emergent phenomenon, explicitly derived from the first principles of quantum mechanics under certain conditions. In particular, quantum thermalization *must* occur under the three conditions considered here. From a technical point of view, we show this important statement without imposing extra assumptions on energy distributions [24, 26], relaxing the locality constraint of quantum dynamics [33], or relying on empirical approximations such as ETH or other random matrix theories [19–22, 34].

## QUANTUM THERMALIZATION

We begin by formulating the precise statement of quantum thermalization. Let us consider a periodic (Euclidean) lattice of  $N = L^D$  qubits in arbitrary dimension D. The system dynamics is dictated by a local, translation-invariant (TI) Hamiltonian  $H = \sum_j h_j$ , where each term  $h_j$  describes interactions among particles within a finite range, is independent of its position j, and has bounded strength (i.e., has operator norm  $\|h_j\|_{\text{op}} \leq 1$ ). Quantum thermalization states that an initial state  $|\psi\rangle$ , upon sufficiently long time evolution, becomes locally indistinguishable from a thermal state  $g_{\beta}$ . The effective inverse temperature  $\beta$  is determined by matching the conserved global energy  $\langle \psi | H | \psi \rangle =$  $\operatorname{tr}(g_{\beta}H)$ .

The local indistinguishability can be formulated rigorously: for any connected region A of finite size and for any typical late time t, we demand that the evolved state reproduces the thermal expectation values

$$\langle \psi(t)|O_A|\psi(t)\rangle = \operatorname{tr}(g_\beta O_A) \pm \varepsilon(N)$$
 (3)

of any observable  $O_A$  supported in A, where  $\varepsilon(N)$  accounts for any finite-size corrections that must vanish in

the thermodynamic limit. Equivalently,

$$\left\|g_{\beta} - \psi(t)\right\|_{A} \le \varepsilon(N),\tag{4}$$

where we introduced the *local trace norm*  $\|\cdot\|_A$ , which measures the distinguishability between  $g_\beta$  and  $\psi(t) = |\psi(t)\rangle\langle\psi(t)|$  to observables in region A, i.e.,  $\|g_\beta - \psi(t)\|_A \coloneqq \max_{O_A} \frac{1}{\|O_A\|_{op}} |\operatorname{tr}((g_\beta - \psi(t))O_A)|$ . See the Supplemental Material (SM) for more details [35]. We say the initial state  $\psi$  thermalizes under H if Eq. (4) holds.

To formulate a rigorous statement about quantum thermalization, one needs to take into account two important considerations. First, determining whether a given specific state thermalizes under a given Hamiltonian is a computationally undecidable problem, even when the initial state is constrained to be unentangled [36]. Therefore, we focus on establishing a *typicality* statement: among many physically relevant initial states that form a large ensemble  $\mathcal{E}_{\beta}$ , any typical state thermalizes. Second, not all Hamiltonians lead to quantum thermalization; many examples of non-thermalizing systems have been recognized over the past decades [11–14], and the precise conditions that ensure quantum thermalization or the lack thereof have not been completely identified. It turns out that only one extra condition (in addition to TI) suffices to exclude all non-thermalizing Hamiltonians: the energy spectrum  $\{E_0, E_1, E_2, \dots\}$  has nondegenerate gaps, meaning that  $E_i + E_j = E_k + E_l$  is exactly satisfied only if (i, j) = (k, l) or (i, j) = (l, k). This condition has been assumed in many prior works [27, 32, 35] and is believed to hold generically, as we further discuss later. We now state our main result.

**Theorem 1** (Quantum thermalization). Let H be a TI, (geometrically) local Hamiltonian with nondegenerate spectral gaps on a D-dimensional periodic lattice with N qubits. Below a certain threshold inverse temperature  $\beta_* > 0$  which is independent of N, the system exhibits quantum thermalization, in the following sense. For any  $|\beta| < \beta_*$  and any ensemble  $\mathcal{E}_{\beta}$  of pure, unentangled states which maximizes the von Neumann entropy under the global energy constraint  $\mathbb{E}_{\psi}[\operatorname{tr}(\psi H)] = \operatorname{tr}(g_{\beta}H)$ , we have

$$\mathbb{E}_{\psi} \mathbb{E}_{t} \left[ \left\| g_{\beta} - \psi(t) \right\|_{A} \right] \le O \left( N^{-1/2 + \delta} \right)$$
(5)

for any connected region A containing a constant number of qubits and any  $\delta > 0$ .

In Eq. (5), the average  $\mathbb{E}_{\psi}$  is taken over initial states  $\psi$  sampled from the ensemble  $\mathcal{E}_{\beta}$  and  $\mathbb{E}_t[\cdot] := \lim_{T \to \infty} \frac{1}{T} \int_0^T dt [\cdot]$  denotes averaging over evolution times  $t \geq 0$ . The local trace norm being averaged is nonnegative, and therefore, Theorem 1 implies quantum thermalization of all typical initial states in  $\mathcal{E}_{\beta}$  and evolution times up to finite-size corrections. The corrections are expected [37, 38] and vanish in the thermodynamic





FIG. 2. Flowchart for our proof of quantum thermalization in translation-invariant spin systems at high temperature, summarizing logical implications among our results (blue), previously-known results (green), and assumptions (orange).

limit. The maximum-entropy condition ensures that the ensemble  $\mathcal{E}_{\beta}$  is as large as possible, from a physical point of view, and, as we will see shortly, it will also imply that the ensemble averages to the Gibbs state  $\mathbb{E}_{\psi}[\psi] = g_{\beta}$ .

The proof of Theorem 1 is enabled by synthesizing a number of previously-established and newly-developed technical results, summarized by the flowchart in Fig. 2. Among them, three particular results are pivotal. First, we introduce a notion that we call *energy-delocalized* Gibbs ensembles (EDGEs) — these are ensembles of pure states, wherein individual states are typically well spread in the energy eigenbasis (energy delocalized) while their collection averages to the Gibbs state (Gibbs ensemble). We prove in our EDGE Theorem that typical states in any EDGE with finite correlation length thermalize under the nondegenerate spectral gaps condition. Second, we invoke a recent result regarding the separability of Gibbs states: Ref. [29] proves the existence of a Gibbs ensemble consisting of product states at sufficiently high temperature  $|\beta| < \beta_*^{(sep)}$ . Finally, using the translation invariance, we show that any productstate Gibbs ensemble at  $|\beta| < \beta_*^{(corr)}$  — which guarantees a finite correlation length [28] — must also be energy delocalized, as measured by a decaying average inverse participation ratio (IPR), and thus constitute an EDGE. Combined together we obtain Theorem 1 with  $\beta_* = \min(\beta_*^{(sep)}, \beta_*^{(corr)})$ . We now give an overview of these three results.

## EDGE THEOREM

We begin by formally introducing EDGEs.

**Definition 1** (Energy-delocalized Gibbs ensemble). For any N-qubit Hamiltonian H with Gibbs state  $g_{\beta}$ , an ensemble of states  $\mathcal{E}_{\beta}$  is an EDGE if it satisfies the following two conditions:

(a) [Gibbs ensemble] The average of the ensemble is close to  $g_{\beta}$ ,

$$\left\| \mathop{\mathbb{E}}_{\psi}[\psi] - g_{\beta} \right\| \le 2^{-N} \varepsilon_{\rm GE}(N), \tag{6}$$

where  $\|\cdot\|$  is the usual trace norm and  $\psi = |\psi\rangle\langle\psi|$ .

(b) [Energy delocalization] The average inverse participation ratio (IPR) in the energy eigenbasis  $\{|E_j\rangle\}_j$ is upper bounded as

$$\mathbb{E}_{\psi}\left[\sum_{j=1}^{2^{N}}\left|\langle E_{j}|\psi\rangle\right|^{4}\right] \leq \varepsilon_{\mathrm{ED}}(N).$$
(7)

The two functions  $\varepsilon_{\text{ED}}(N)$  and  $\varepsilon_{\text{GE}}(N)$  are nonnegative and vanish in the thermodynamic limit  $N \to \infty$ , but are otherwise arbitrary.

The energy delocalization is measured by the inverse of the participation ratio  $\text{PR} = 1/\sum_j |\langle E_j | \psi \rangle|^4$ , which is also called the effective dimension of  $|\psi\rangle$ , as it estimates the number of eigenstates on which  $|\psi\rangle$  is supported. When the IPR is small,  $|\psi\rangle$  is delocalized in energy.

Our next result shows that typical states in an EDGE thermalize as long as the Gibbs state has finite correlation length, with finite-size corrections bounded by  $\varepsilon_{\text{GE}}(N)$  and  $\varepsilon_{\text{ED}}(N)$ . Finite correlation length means that correlations between any pair of local observables decay at least exponentially fast in their distance [35]. This condition holds below a certain constant inverse temperature  $\beta_*^{(\text{corr})}$  for any local Hamiltonian in any geometry [28, 35]. Our second result reads as follows.

**Theorem 2** (EDGE Theorem). Let H be a TI, (geometrically) local Hamiltonian with nondegenerate spectral gaps on a D-dimensional periodic lattice with N qubits and  $\mathcal{E}_{\beta}$  be an EDGE. Assume that the Gibbs state  $g_{\beta}$  has finite correlation length. Then, for any connected region A with a constant number of qubits  $N_A$ ,

$$\mathbb{E}_{\psi} \mathbb{E}_{t} \left[ \|g_{\beta} - \psi(t)\|_{A} \right] \leq O\left(N^{-1/2+\delta}\right) + \varepsilon_{\mathrm{GE}}(N) + 2^{N_{A}} \varepsilon_{\mathrm{ED}}(N)^{1/2}, \quad (8)$$

for any  $\delta > 0$ .

Our proof of Theorem 2 is inspired by Ref. [27], which studied the quantum thermalization of random product states at infinite temperature ( $\beta = 0$ ). Below we present the key ideas, leaving some technical details for the SM [35]. Proof. Quantum thermalization,  $\|\psi(t) - g_{\beta}\|_A \to 0$ , can be split into two fine-grained statements. First, we show equilibration, which dictates that pure quantum states at late times become locally indistinguishable from their time-averaged mixed state:  $\|\psi(t) - \rho_{\infty}\|_A \to 0$  where  $\rho_{\infty} := \mathbb{E}_{\tau}[\psi(\tau)]$ . Second, we prove a property akin to ergodicity, the local equivalence between temporally and thermally averaged ensembles  $\|\rho_{\infty} - g_{\beta}\|_A \to 0$ . Once these two statements are proven, one obtains the desired result via a triangle inequality.

A sufficient condition for quantum equilibration is well known when assuming nondegenerate spectral gaps: the vanishing of IPR [32], leading to [35]

$$\mathop{\mathbb{E}}_{\psi} \mathop{\mathbb{E}}_{t} \left[ \left\| \psi(t) - \rho_{\infty} \right\|_{A} \right] \leq 2^{N_{A}} \mathop{\mathbb{E}}_{\psi} \left[ \sum_{j=1}^{2^{N}} \left| \langle E_{j} | \psi \rangle \right|^{4} \right]^{1/2},$$

which gives the last term in Eq. (8) upon using property (b) of an EDGE.

To prove the equivalence of ensembles, we utilize a modified version of a seminal result, called the weak eigenstate thermalization hypothesis (weak ETH) [25].

**Proposition 1** (A variant of the weak ETH). Let *H* be a TI, (geometrically) local Hamiltonian on a *D*-dimensional periodic lattice with *N* qubits. Let  $\rho = \sum_{j=1}^{2^N} p_j |E_j\rangle\langle E_j|$  be a diagonal state in the eigenbasis  $\{|E_j\rangle\}_j$  of *H* that has finite correlation length. Consider a connected region *A* with a constant number of qubits. Then, for any  $\delta > 0$ ,

$$\sum_{j=1}^{2^N} p_j \left\| \left| E_j \right\rangle \!\! \left\langle E_j \right| - \rho \left\|_A \le O\left(N^{-1/2+\delta}\right). \tag{9}$$

The proof of Proposition 1 is available in the SM [30, 31, 35, 39]. By applying this result to  $\rho = g_{\beta}$ , we establish the equivalence between thermal and temporal ensembles with bounded error:

$$\begin{split} \mathbb{E}_{\psi} \| \rho_{\infty} - g_{\beta} \|_{A} &= \mathbb{E}_{\psi} \left\| \sum_{j} |\langle E_{j} | \psi \rangle|^{2} (|E_{j} \rangle \langle E_{j} | - g_{\beta}) \right\|_{A} \\ &\leq \sum_{j} \mathbb{E}_{\psi} \left[ \langle E_{j} | \psi | E_{j} \rangle \right] \| |E_{j} \rangle \langle E_{j} | - g_{\beta} \|_{A} \\ &\leq \sum_{j} p_{j} \| |E_{j} \rangle \langle E_{j} | - g_{\beta} \|_{A} + \varepsilon_{\mathrm{GE}}(N), \end{split}$$

where we used  $\rho_{\infty} = \sum_{j} |\langle E_{j} | \psi \rangle|^{2} |E_{j} \rangle \langle E_{j}|$  for a given  $|\psi\rangle$  (owing to the nondegenerate spectral gap condition) in the first line, the triangle inequality in the second line, and the substitution of  $\mathbb{E}_{\psi}[\psi] \approx g_{\beta}$  up to a bounded error  $2^{-N} \varepsilon_{\text{GE}}(N)$  and  $p_{j} \coloneqq \langle E_{j} | g_{\beta} | E_{j} \rangle$  in the third line. This completes the proof.

## PRODUCT-STATE GIBBS ENSEMBLES

While we have shown the thermalization of typical states in EDGEs, we have not yet presented explicit examples or even shown their existence. At infinite temperature, an example of an EDGE has been previously constructed [27]. The random product-state ensemble, where the state at each site is sampled independently and uniformly, constitutes an EDGE and, therefore, thermalizes to the maximally mixed state (or the Gibbs state with  $\beta = 0$ ), which is independent of the Hamiltonian. At finite temperature, however, it is nontrivial to construct an EDGE of low-entanglement states because such ensemble necessarily depends on the details of the Hamiltonian. We address this challenge by using a recent result from quantum information theory.

**Lemma 1** (High-temperature Gibbs states are unentangled, Theorem 1.5 of [29]). Consider a Hamiltonian consisting of Pauli-string interactions on qubits, where each qubit appears in at most  $\mathfrak{d}$  terms and each term contains at most  $\mathfrak{K}$  qubits. For  $|\beta| < \beta_*^{(\text{sep})} := 1/(\gamma \mathfrak{d} \mathfrak{K}^2)$  with a fixed universal constant  $\gamma$ , the corresponding Gibbs state can be expressed as a mixture

$$g_{\beta} = \sum_{|\psi\rangle \in \mathcal{S}} p_{\psi} |\psi\rangle\!\langle\psi|,$$

of product states in  $S = \{ |0\rangle, |1\rangle, |\pm\rangle, |\pm i\rangle \}^{\otimes N}$ .

Lemma 1 guarantees the existence of an ensemble that averages exactly to the thermal state and consists of certain product states at high-enough temperature. Such an ensemble is not unique, and our results are independent of any specific choice. Lemma 1 has an important, general implication. The Gibbs state uniquely maximizes the von Neumann entropy  $S(\rho) = -\operatorname{tr}(\rho \log \rho)$  under the global energy constraint  $\operatorname{tr}(\rho H) = \operatorname{tr}(g_{\beta} H)$  [35], and this maximum is attainable by a certain pure product-state ensemble:

$$S(g_{\beta}) = \max_{\substack{\text{pure product-state } \mathcal{E} \\ \operatorname{tr}(\rho_{\mathcal{E}}H) = \operatorname{tr}(g_{\beta}H)}} S(\rho_{\mathcal{E}}), \qquad (10)$$

with  $\rho_{\mathcal{E}} = \mathbb{E}_{\psi \in \mathcal{E}}[\psi]$ . Thus, any ensemble of pure states with maximal entropy must also describe the Gibbs state, satisfying property (a) of an EDGE with  $\varepsilon_{\text{GE}}(N) = 0$ . It turns out that property (b), energy delocalization, is also satisfied in our setting, as we show next.

## ENERGY DELOCALIZATION FROM TRANSLATION INVARIANCE

We show that any high-temperature product-state Gibbs ensemble is energy-delocalized or, equivalently, has a decaying average IPR in the energy eigenbasis, thus constituting an EDGE. To show this, one possible approach is to utilize Lemma 1 of Ref. [26] which guarantees that any product state with extensive energy variance has an IPR bounded as  $\tilde{O}(N^{-1/2})$  solely due to the locality of the Hamiltonian. Here we provide a (quadratically) stronger bound that does not require an assumption on the energy variance by leveraging the translation invariance.

**Theorem 3** (Bound on average IPR). Let H be a TI, (geometrically) local, Hamiltonian on a D-dimensional periodic lattice with N qubits, and consider an ensemble  $\mathcal{E}_{\beta}$  of product states whose mean value is the Gibbs state  $g_{\beta}$ , at high temperature  $|\beta| < \beta_*^{(\text{corr})}$  (or at any temperature as long as  $g_{\beta}$  has finite correlation length and satisfies the locality of temperature) [35]. Then, the average IPR in a common eigenbasis  $\{|E_j\rangle\}_j$  of the translation operator and the Hamiltonian is bounded

$$\mathbb{E}_{\psi}\left[\sum_{j=1}^{2^{N}} |\langle E_{j} | \psi \rangle|^{4}\right] \le O(N^{-1+\delta}) \tag{11}$$

for any  $\delta > 0$ .

The EDGE Theorem and Theorem 3 together imply our main result, Theorem 1. While the detailed proof is presented in the SM [35], below we illustrate the key intuition behind Theorem 3 by proving a simplified version in 1D.

**Proposition 2.** For any translation invariant Hamiltonian for N qubits on a ring, the average IPR over the uniform ensemble of computational-basis states  $|b\rangle$  with  $b \in \{0,1\}^N$  is upper bounded

$$\mathbb{E}_{b \in \{0,1\}^N} \left[ \sum_j |\langle E_j | b \rangle|^4 \right] \le \frac{1}{N} + O(2^{-N/2}), \qquad (12)$$

where  $\{|E_j\rangle\}_j$  is a common eigenbasis of the translation operator and the Hamiltonian.

Note that the random computational-basis states ensemble describes the Gibbs state at infinite temperature,  $\mathbb{E}_{b \in \{0,1\}^N}[|b\rangle\langle b|] = 1/2^N = g_{\beta=0}$ . Hence, Proposition 2 combined with the EDGE Theorem already proves that a random computational-basis state will typically thermalize to infinite temperature for any TI Hamiltonian with nondegenerate spectral gaps in 1D.

*Proof.* The central idea is to divide the bitstrings  $b \in \{0,1\}^N$  that enumerate the states into two different groups and bound their contributions to the average IPR separately: periodic bitstrings, which are invariant under some nontrivial translation on the circle, and aperiodic ones, which are not equal to any of their nontrivial translations (see Fig. 3a). The former group



FIG. 3. Schematic illustrations of product states on 1D or 2D lattices. **a** Periodic (left) and aperiodic (right) computational basis states in 1D. **b** Approximately periodic product state in 2D: qubit configurations in regions A and B separated by a period are similar, giving rise to long-range correlations.

constitutes a vanishingly small fraction among all  $2^N$  possibilities, hence their contribution becomes small. Specifically, since the period cannot be longer than N/2, there are at most  $O(2^{N/2})$  distinct periodic bitstrings, leading to the second term in Eq. (12). The second, aperiodic group contains the majority of the bitstrings, but their overlaps with the energy eigenstates are necessarily small due to the translation invariance. Specifically, for N distinct bitstrings  $b_1, b_2, \ldots, b_N$  which are translations of each other, their overlaps with an energy eigenstate must be the same up to a global phase, resulting in a bound  $|\langle E_j | b \rangle|^2 \leq 1/N$ . This leads to the first term in Eq. (12).

Theorem 3 is proven by extending the previous argument to arbitrary product states defined on 1D or higherdimensional lattices. This generalization involves solving two nontrivial complications. First, unlike the bitstring case, one cannot sharply differentiate periodic versus aperiodic states because generic product states have a nonzero overlap with their translations. This is overcome by introducing *approximately* periodic states, which are those states whose overlap with some translate is above a certain threshold (Fig. 3b) [35]. Second, at finite temperature, the ensemble  $\mathcal{E}_{\beta}$  is not uniform, hence bounding the contribution to the IPR from (approximately) periodic states requires going beyond simple counting. For this, we utilize the finite correlation length of the Gibbs state: (approximately) periodic states cannot be weighted by high probabilities in the ensemble because otherwise they would induce long-range correlations in  $g_{\beta}$  (see Fig. 3). Here, we additionally utilize the locality of temperature, i.e., that thermal subsystems are well described by local Gibbs states, a property also guaranteed if  $|\beta| < \hat{\beta}_*^{(\text{corr})}$  [28, 35].

## DISCUSSION AND OUTLOOK

Discussion—. Our results showcase that, in certain quantum mechanical systems, statistical mechanics can be understood as an emergent phenomenon. Interestingly, this emergence seems qualitatively different from its classical counterpart. Classically, a single point in the phase space cannot display any subsystem entropy. A nonvanishing amount of entropy must be introduced externally. Only then can the chaotic dynamics effectively increase the (subsystem) entropy over time [40]. This is in stark contrast to quantum systems, where the subsystem entropy is generated by an intrinsically quantum property: entanglement.

Our Theorem 1 establishes a universal statement prescribing conditions under which any system must display quantum thermalization. In practice, one often has access to more fine-grained information, and our other technical results might be useful. For example, if the Gibbs state can be unraveled into an ensemble of pure states each with finite correlation length and extensive energy variance, our EDGE Theorem, compounded with Lemma 1 of Ref. [26], guarantees their typical quantum thermalization.

An important fundamental question is finding the minimum set of conditions needed to make universal statements about quantum thermalization. While our work might be further improved quantitatively, we speculate that completely dropping any of our three conditions —translation invariance, high effective temperature, or nondegenerate spectral gaps- might require qualitatively different tools and insights. Without translation invariance, strongly disordered systems in 1D are largely debated to exhibit many-body localization (MBL) [3, 4, 9], which, if exists, would constitute a robust violation of quantum thermalization. A lower bound on the effective temperature is unavoidable for the thermalization of initial product states. This is simply because states without any entanglement cannot attain arbitrarily low effective temperatures in certain frustrated systems. Finally, the nondegenerate spectral gap condition (or some variant of it) might be inevitable to exclude trivial, noninteracting or otherwise fine-tuned systems which do not thermalize. Note that we can partially relax this condition and allow for few degenerate gaps [41].

Unlike the translation invariance and the high temperature conditions, the nondegenerate spectral gap condition is hidden in the energy spectrum, and hence difficult to check. Nevertheless, it is expected to be a generic condition, in a precise sense. The violations of the nondegenerate spectral gap condition are known to form a zero-measure set in the space of local Hamiltonians without the TI constraint [42], implying that those are rare, fine-tuned exceptions. Even within the space of all local TI Hamiltonians, violations are conjectured to form a zero-measure set [27]. Interesting, if the conjecture were to be false for some systems, then *every* TI local Hamiltonian of the same size would necessarily have at least one gap degeneracy. The latter possibility has been already ruled out by explicit numerical tests for relatively small system sizes [24]. Proving the conjecture for all system sizes is an important open question [35].

*Outlook*—. Our work leaves several questions and generalizations open. It might be possible to broaden the scope of our results by constructing an EDGE at lower temperatures consisting of weakly-entangled states [43, 44] or extending our methods to ensembles with nonthermal averages. It would be important to improve the robustness of our statements (e.g. by allowing for  $\varepsilon_{\rm GE} > 0$  in Theorem 3) and tighten our finitesize bounds, potentially from  $N^{-1/2}$  to  $N^{-1}$ . Another important direction is understanding the finite-time behavior [41] — recent developments have shown that certain TI systems do thermalize, but only after exponentially long timescales [45]. Finally, one should investigate the thermalization of more general physical systems beyond qubits on lattices, such as bosonic particles with unbounded local Hilbert-spaces, those possessing symmetries, those relaxing to generalized Gibbs ensembles [46], and quantum field theories; with the ultimate aim of constructing a universal framework explaining the emergence of statistical mechanics in quantum systems.

#### Acknowledgments

We thank A. M. Alhambra, A. Anshu, A. Bakshi, S. Balasubramanian, F. Brandão, J. Eisert, J. Haferkamp, A. Harrow, W. W. Ho, Y. Huang, I. H. Kim, A. Liu, D. Mark, and D. Ranard for insightful conversations and feedback on earlier versions of this preprint. In particular, we acknowledge A. M. Alhambra for a detailed discussion at the initial stage of this work and Y. Huang for a careful explanation of the techniques used in Ref. [27], some of which are also utilized in the present work.

- [2] J. Eisert, M. Friesdorf, and C. Gogolin, Quantum manybody systems out of equilibrium, Nature Physics 11, 124–130 (2015).
- [3] R. Nandkishore and D. A. Huse, Many-body localization and thermalization in quantum statistical mechanics, Annual Review of Condensed Matter Physics 6, 15–38 (2015).
- [4] D. A. Abanin, E. Altman, I. Bloch, and M. Serbyn, Colloquium: Many-body localization, thermalization, and entanglement, Rev. Mod. Phys. **91**, 021001 (2019).

A. Polkovnikov, K. Sengupta, A. Silva, and M. Vengalattore, Colloquium: Nonequilibrium dynamics of closed interacting quantum systems, Rev. Mod. Phys. 83, 863 (2011).

- [5] A. M. Alhambra, Quantum many-body systems in thermal equilibrium, PRX Quantum 4, 040201 (2023).
- [6] B. J. Brown, D. Loss, J. K. Pachos, C. N. Self, and J. R. Wootton, Quantum memories at finite temperature, Rev. Mod. Phys. 88, 045005 (2016).
- [7] D. Harlow, Jerusalem lectures on black holes and quantum information, Rev. Mod. Phys. 88, 015002 (2016).
- [8] J. von Neumann, Proof of the ergodic theorem and the H-theorem in quantum mechanics: Translation of: Beweis des Ergodensatzes und des H-Theorems in der neuen Mechanik, The European Physical Journal H 35, 201–237 (2010 [original 1929]).
- [9] D. Basko, I. Aleiner, and B. Altshuler, Metal-insulator transition in a weakly interacting many-electron system with localized single-particle states, Annals of Physics **321**, 1126–1205 (2006).
- [10] F. Levkovich-Maslyuk, The Bethe ansatz, Journal of Physics A: Mathematical and Theoretical 49, 323004 (2016).
- [11] M. Serbyn, D. A. Abanin, and Z. Papić, Quantum manybody scars and weak breaking of ergodicity, Nature Physics 17, 675–685 (2021).
- [12] S. Pai, M. Pretko, and R. M. Nandkishore, Localization in fractonic random circuits, Phys. Rev. X 9, 021003 (2019).
- [13] P. Sala, T. Rakovszky, R. Verresen, M. Knap, and F. Pollmann, Ergodicity breaking arising from Hilbert space fragmentation in dipole-conserving hamiltonians, Phys. Rev. X 10, 011047 (2020).
- [14] V. Khemani, M. Hermele, and R. Nandkishore, Localization from Hilbert space shattering: From theory to physical realizations, Phys. Rev. B 101, 174204 (2020).
- [15] J. M. Deutsch, Quantum statistical mechanics in a closed system, Physical Review A 43, 2046–2049 (1991).
- [16] M. Srednicki, Chaos and quantum thermalization, Physical Review E 50, 888–901 (1994).
- [17] M. Rigol, V. Dunjko, and M. Olshanii, Thermalization and its mechanism for generic isolated quantum systems, Nature 452, 854–858 (2008).
- [18] A. M. Kaufman, M. E. Tai, A. Lukin, M. Rispoli, R. Schittko, P. M. Preiss, and M. Greiner, Quantum thermalization through entanglement in an isolated manybody system, Science **353**, 794–800 (2016).
- [19] H. Tasaki, From quantum dynamics to the canonical distribution: General picture and a rigorous example, Phys. Rev. Lett. 80, 1373 (1998).
- [20] S. Goldstein, J. L. Lebowitz, R. Tumulka, and N. Zanghì, Canonical typicality, Phys. Rev. Lett. 96, 050403 (2006).
- [21] S. Popescu, A. J. Short, and A. Winter, Entanglement and the foundations of statistical mechanics, Nature Physics 2, 754–758 (2006).
- [22] M. Cramer, Thermalization under randomized local Hamiltonians, New Journal of Physics 14, 053051 (2012).
- [23] J. P. Keating, N. Linden, and H. J. Wells, Spectra and eigenstates of spin chain Hamiltonians, Communications in Mathematical Physics 338, 81–102 (2015).
- [24] M. P. Müller, E. Adlam, L. Masanes, and N. Wiebe, Thermalization and canonical typicality in translationinvariant quantum lattice systems, Communications in Mathematical Physics **340**, 499–561 (2015).
- [25] T. Mori, Weak eigenstate thermalization with large deviation bound (2016), arXiv:1609.09776 [cond-mat.statmech].
- [26] T. Farrelly, F. G. S. L. Brandão, and M. Cramer, Ther-

malization and return to equilibrium on finite quantum lattice systems, Phys. Rev. Lett. **118**, 140601 (2017).

- [27] Y. Huang and A. W. Harrow, Scrambling and thermalization in translation-invariant systems (2019), 1907.13392 [cond-mat.dis-nn].
- [28] M. Kliesch, C. Gogolin, M. J. Kastoryano, A. Riera, and J. Eisert, Locality of temperature, Phys. Rev. X 4, 031019 (2014).
- [29] A. Bakshi, A. Liu, A. Moitra, and E. Tang, Hightemperature Gibbs states are unentangled and efficiently preparable (2024), arXiv:2403.16850 [quant-ph].
- [30] F. G. S. L. Brandão, E. Crosson, M. B. Şahinoğlu, and J. Bowen, Quantum error correcting codes in eigenstates of translation-invariant spin chains, Phys. Rev. Lett. **123**, 110502 (2019).
- [31] A. M. Alhambra, J. Riddell, and L. P. García-Pintos, Time evolution of correlation functions in quantum many-body systems, Phys. Rev. Lett. **124**, 110605 (2020).
- [32] N. Linden, S. Popescu, A. J. Short, and A. Winter, Quantum mechanical evolution towards thermal equilibrium, Phys. Rev. E 79, 061103 (2009).
- [33] C. Bertoni, C. Wassner, G. Guarnieri, and J. Eisert, Typical thermalization of low-entanglement states (2024), arXiv:2403.18007 [quant-ph].
- [34] A. Dymarsky, N. Lashkari, and H. Liu, Subsystem eigenstate thermalization hypothesis, Phys. Rev. E 97, 012140 (2018).
- [35] See Supplemental Material.
- [36] N. Shiraishi and K. Matsumoto, Undecidability in quantum thermalization, Nature Communications 12, 5084 (2021).
- [37] M. Srednicki, Thermal fluctuations in quantized chaotic systems, Journal of Physics A: Mathematical and General 29, L75–L79 (1996).
- [38] Y. Huang, High-precision simulation of finite-size thermalizing systems at long times (2024), arXiv:2406.05399 [cond-mat.stat-mech].
- [39] A. Anshu, Concentration bounds for quantum states with finite correlation length on quantum spin lattice systems, New Journal of Physics 18, 083011 (2016), we refer to arXiv:1508.07873 which has been updated after publication.
- [40] J. R. Dorfman, An introduction to chaos in nonequilibrium statistical mechanics (Cambridge University Press, Cambridge, England, 1999).
- [41] A. J. Short and T. C. Farrelly, Quantum equilibration in finite time, New Journal of Physics 14, 013063 (2012).
- [42] Y. Huang, Extensive entropy from unitary evolution (2021), 2104.02053 [quant-ph].
- [43] G. Vidal, Efficient classical simulation of slightly entangled quantum computations, Phys. Rev. Lett. 91, 147902 (2003).
- [44] M. Berta, F. G. S. L. Brandão, J. Haegeman, V. B. Scholz, and F. Verstraete, Thermal states as convex combinations of matrix product states, Phys. Rev. B 98, 235154 (2018).
- [45] S. Balasubramanian, S. Gopalakrishnan, A. Khudorozhkov, and E. Lake, Glassy word problems: Ultraslow relaxation, Hilbert space jamming, and computational complexity, Phys. Rev. X 14, 021034 (2024).
- [46] L. Vidmar and M. Rigol, Generalized Gibbs ensemble in integrable lattice models, Journal of Statistical Mechanics: Theory and Experiment **2016**, 064007 (2016).

# Supplemental Material: Quantum thermalization of translation-invariant systems at high temperature

Saúl Pilatowsky-Cameo<sup>1</sup> and Soonwon Choi<sup>1</sup>

<sup>1</sup>Center for Theoretical Physics, Massachusetts Institute of Technology, Cambridge, MA 02139, USA

In this Supplemental Material, we provide detailed information of our results. In Section I, we introduce our notations, the physical setup under consideration, a detailed discussion of the nondegenerate spectral gap condition, and some useful properties of Gibbs states to be used in later sections. Sections II and III are dedicated to formally present and prove our technical Theorems. Specifically, Section II contains the detailed proof of our EDGE Theorem. Finally, Section III contains the detailed proof on our bound on the average inverse participation ratio (IPR) of thermal, product-state ensembles based on translation invariance.

## CONTENTS

I. Preliminaries	1
A. Setting and notations	1
1. Lattice	1
2. Hilbert space, observables and Hamiltonian	2
3. Local trace norm	2
4. Asymptotic notation	2
B. Nondegenerate spectral gaps	2
C. Properties of Gibbs states	3
1. Maximum entropy principle	3
2. Extensive $\log(Z_{\beta})$	3
3. Global purity	4
4. Finite correlation length	4
5. Locality of temperature	5
6. Subsystem purity	5
II. The EDGE theorem	6
A. Equivalence of the temporal and thermal ensembles	6
B. Quantum equilibration	8
III. Bound on average IPR	9
A. Aperiodic states have low IPR	9
B. Approximately periodic states are atypical	10

### I. PRELIMINARIES

#### A. Setting and notations

We begin by introducing our physical setup and general notations.

#### 1. Lattice

We consider systems defined over a *D*-dimensional lattice  $\Lambda = \{0, 1, 2, ..., L-1\}^D$ , consisting of  $N = L^D$  sites. As it will become clear, the specific geometric properties such as the aspect ratios of the lattice are not important for any of our results. We will specialize to periodic boundary conditions, so we will think of  $\Lambda$  as a torus. Given a point  $\boldsymbol{x} \in \Lambda$ , with coordinates  $\boldsymbol{x} = (x_1, \ldots, x_D)$ , we measure its magnitude via the Manhattan norm,

$$\|\boldsymbol{x}\|_{1} = \sum_{j=1}^{D} \min\{x_{j}, L - x_{j}\}.$$
(S1)

A pair of positions in the lattice can be added or subtracted in the conventional way, i.e. with the entrywise modular addition and subtraction. The distance between two sites  $x, y \in \Lambda$  is given by  $||x - y||_1$ .

#### 2. Hilbert space, observables and Hamiltonian

On each lattice site, we place a single qubit, i.e. a spin-1/2 particle, with local Hilbert space  $\mathcal{H}_{\boldsymbol{x}} = \mathbb{C}^2$ . The global Hilbert space is  $\mathcal{H} = \bigotimes_{\boldsymbol{x} \in \Lambda} \mathcal{H}_{\boldsymbol{x}}$ . We say an operator O on  $\mathcal{H}$  acts on a region  $A \subseteq \Lambda$  (or is supported on A) if there is an operator  $O_A$  acting on  $\mathcal{H} = \bigotimes_{\boldsymbol{a} \in A} \mathcal{H}_{\boldsymbol{a}}$  such that  $O = O_A \otimes \bigotimes_{\boldsymbol{x} \notin A} \mathbb{1}_{\boldsymbol{x}}$ . Henceforth, we abuse notation and not distinguish between  $O_A$  and O whenever the latter is supported in A.

For a  $z \in \Lambda$ , the operator  $\mathbb{T}^z$  is the unitary translation operator, which maps a product state  $|\psi\rangle = \bigotimes_{x \in \Lambda} |\psi_x\rangle$  to  $\mathbb{T}^z |\psi\rangle = \bigotimes_{x \in \Lambda} |\psi_{x-z \mod L}\rangle$ . We study (geometrically) k-local, translation-invariant (TI) Hamiltonians H, which means that  $H = \sum_i h_i$ , where each  $h_i$  acts on a connected region containing at most k qubits and has bounded operator norm, and  $H = \mathbb{T}^z H \mathbb{T}^{-z}$  for any  $z \in \Lambda$ . We will often just say H is local, when the number k is not important.

#### 3. Local trace norm

We measure the local indistinguishability of two states with the following notion.

**Definition S1** (Local trace norm). Given a Hermitian operator W, its trace norm ||W|| is equal to the sum of the singular values of W. Given a region  $A \subseteq \Lambda$  with complement  $\overline{A}$ , we denote  $||W||_A = ||\operatorname{tr}_{\overline{A}}(W)||$ .

From the properties of the trace norm, one can see that

$$\|\rho - \sigma\|_A = \max_{\|O_A\|_{\text{op}} \le 1} |\operatorname{tr}(\rho O_A) - \operatorname{tr}(\sigma O_A)|,$$
(S2)

where  $O_A$  runs over all observables supported on A with operator norm  $\|\cdot\|_{op}$  bounded by 1. Technically,  $\|\cdot\|_A$  is not a norm, but only a seminorm as  $\|\rho - \sigma\|_A = 0$  implies only that the states  $\rho$  and  $\sigma$  are locally indistinguishable, meaning that they cannot be discriminated using observables supported on region A.

## 4. Asymptotic notation

We make use of the big-O and related asymptotic notations. When we write  $f(N) \leq O(g(N))$ , we mean that there exist constants C > 0 and  $N_* > 0$  such that  $f(N) \leq Cg(N)$  for all  $N \geq N_*$ . We sometimes also write  $f(N) \leq e^{O(g(N))}$ , which means  $f(N) \leq e^{Cg(N)}$  for all  $N \geq N_*$ . For lower bounds, we write  $f(N) \geq \Omega(g(N))$  which means  $f(N) \geq Cg(N)$  for all  $N \geq N_*$  and  $f(N) \leq e^{-\Omega(g(N))}$  means  $f(N) \leq e^{-Cg(N)}$  for all  $N \geq N_*$ . Finally,  $f(N) = \Theta(g(N))$  means that both the upper bound  $f(N) \leq O(g(N))$  and lower bound  $f(N) \geq \Omega(g(N))$  are satisfied.

We remark that these relations are typically expressed using equality rather than inequality signs. However, we find that the inequalities provide additional clarity.

#### B. Nondegenerate spectral gaps

One of the most important, nontrivial properties that we demand from the spectrum of Hamiltonians is the *non*degenerate spectral gaps condition.

**Definition S2** (Nondegenerate spectral gaps). A Hamiltonian has nondegenerate spectral gaps if its spectrum  $\{E_j\}_j$  has no degeneracies and no gap degeneracies, meaning that

$$E_i + E_j = E_m + E_l \iff (i, j) = (m, l) \text{ or } (i, j) = (l, m).$$

$$(S3)$$

This condition is also called the second no-resonance condition [1]. This condition is believed to hold for generic Hamiltonians. It has been shown that all Hamiltonians violating this condition form a zero measure set in the space of k-local Hamiltonians on N qubits for all N [2]. Even within the space of TI Hamiltonians on N qubits, it is conjectured that all exceptions form a zero measure set. The latter space can be parametrized, for example, as

$$H = \sum_{\boldsymbol{n} \in \Lambda} \mathbb{T}^{\boldsymbol{n}} \sum_{P} J_{P} P \mathbb{T}^{-\boldsymbol{n}}, \tag{S4}$$

where P runs over all Pauli strings which act on some fixed connected region of k qubits, and the coefficients are  $J_P \in [-1, 1]$ .

**Conjecture S1** (Conjecture 2 of Ref. [3]: almost all k-local TI Hamiltonians have nondegenerate gaps). For a fixed  $k \ge 2, N$ , and D, almost all (geometrically) local TI Hamiltonians on the D-dimensional periodic lattice with N qubits have nondegenerate spectral gaps, with respect to the ensemble produced by taking the coefficients  $J_P$  in Eq. (S4) to be uniformly distributed in [-1, 1].

Although proving conjecture S1 remains an open problem, Ref [3] has made some progress by showing the following:

**Theorem S1** (Lemma 8 of [3]). For a fixed  $k \ge 2, N$ , and D one of the following statements holds:

- 1. Almost all k-local TI Hamiltonians on the D-dimensional periodic lattice with N qubits have nondegenerate spectral gaps.
- 2. No k-local TI Hamiltonian on the D-dimensional periodic lattice with N qubits has nondegenerate spectral gaps.

Conjecture S1 asserts that case 1 in the above theorem is the true statement. Since the theorem is already proven, proving the conjecture only requires showing that case 2 does not hold, which reduces to finding a single example of a k-local TI system with nondegenerate spectral gaps. This can be done numerically [4] for accessible system sizes. For all systems sizes N checked so far, case 1 holds as expected. However, for arbitrary system sizes, it appears difficult to attain analytical understanding of the spectrum of a Hamiltonian with nondegenerate spectral gaps. This is an interesting and important problem that we do not address in this work.

## C. Properties of Gibbs states

The Gibbs state of H at inverse temperature  $\beta$  is denoted by

$$g_{\beta} = \frac{e^{-\beta H}}{\operatorname{tr}(e^{-\beta H})}.$$
(S5)

As mentioned in the main text, we always use units where the Boltzmann constant is  $k_B = 1$ . Without loss of generality, we will only consider nonnegative temperatures  $\beta \ge 0$  (a change on the overall sign of the Hamiltonian always ensures this). Below, we introduce several known properties satisfied by Gibbs states under certain conditions that we use throughout our work.

#### 1. Maximum entropy principle

The defining property of the Gibbs state is that it uniquely maximizes the von Neumann entropy  $S(\rho) = -\operatorname{tr}(\rho \log \rho)$ among all states sharing the same energy expectation value  $E_{\beta} = \operatorname{tr}(g_{\beta}H)$ , i.e.,

$$g_{\beta} = \underset{\rho: \operatorname{tr}(\rho H) = E_{\beta}}{\operatorname{argmax}} S(\rho).$$
(S6)

This can be shown by minimizing the relative entropy  $D(\rho||g_{\beta}) = -S(\rho) + \beta \operatorname{tr}(H\rho) + \log(\operatorname{tr}(e^{-\beta H})).$ 

## 2. Extensive $\log(Z_{\beta})$

In a TI Hamiltonian, the quantity  $\log(Z_{\beta})$ , with  $Z_{\beta} = tr(e^{-\beta H})$  is extensive, meaning that the limit

$$p_{\beta} \coloneqq \lim_{N \to \infty} \frac{1}{N} \log(Z_{\beta})$$

exists. From a physical point of view, this statement is trivial, because  $\log(Z_{\beta}) = S(g_{\beta}) - \beta E_{\beta}$ , and both the thermal entropy  $S(g_{\beta})$  and thermal energy  $E_{\beta} = tr(Hg_{\beta})$  are extensive quantities. However, what is nontrivial is that this statement can be rigorously proven (see Ref. [5]). This is true regardless of boundary conditions, i.e., H might be TI except for boundary terms.

## 3. Global purity

The global purity of a Gibbs state at nonzero temperature  $\beta < \infty$  can be proven to decay exponentially in the system size in TI systems. Specifically, if H is a TI invariant local Hamiltonian on a D-dimensional lattice with N qubits (up to boundary conditions, which may be open or periodic). For any finite  $\beta < \infty$ , the purity of the Gibbs state is upper bounded as

$$\operatorname{tr}(g_{\beta}^2) \le e^{-\Omega(N)}.\tag{S7}$$

This follows from computing

$$\operatorname{tr}\left(g_{\beta}^{2}\right) = \frac{Z_{2\beta}}{Z_{\beta}^{2}}$$

By the extensivity  $\log(Z_{\beta}) \to N p_{\beta}$ ,  $\lim_{N\to\infty} \frac{1}{N} \log\left(\operatorname{tr}\left(g_{\beta}^{2}\right)\right)$  converges to  $p_{2\beta} - 2p_{\beta}$ , and consequently we obtain Eq. (S7).

#### 4. Finite correlation length

Intuitively, the correlation length of a state denotes the maximum distance over which correlations, whether classical or quantum, can persist. In many systems, particularly those at high temperatures, the Gibbs state exhibits a finite correlation length, defined as follows.

**Definition S3.** (Finite correlation length) Let  $\rho$  be a state. We define the correlations between A and B by

$$\operatorname{Corr}_{\rho}(A,B) \coloneqq \max_{O_A,O_B} \frac{|\operatorname{tr}(\rho O_A \otimes O_B) - \operatorname{tr}(\rho O_A) \operatorname{tr}(\rho O_B)|}{\|O_A\|_{\operatorname{op}} \|O_B\|_{\operatorname{op}}}$$
(S8)

where  $O_A(O_B)$  runs over all operators with support contained in region A(B). We say  $\rho$  has finite correlation length  $\xi > 0$  if for any regions A and B,

$$\operatorname{Corr}_{\rho}(A,B) \le f_{\operatorname{corr}}(a)e^{-d(A,B)/\xi}$$

where  $f_{corr}(a)$  is a fixed polynomial function, evaluated at  $a = \min\{|\partial A|, |\partial B|\}$ , with  $|\partial A|, (|\partial B|)$  the size of the boundary of A(B), i.e., the number of pairs of contiguous sites such that one site is inside the region and the other outside. The distance between A and B is given by

$$d(A, B) = \inf\{\|x - y\|_1 | x \in A, y \in B\}$$

In one-dimensional systems, Gibbs states of TI local Hamiltonians have finite correlation length at nonzero temperatures, with  $f_{corr}(a) = 1$  independent of a [6, 7]. For higher dimensions, Gibbs states are only guaranteed to have finite correlation length above a certain temperature, as shown by the following result.

**Lemma S1** (High-temperature Gibbs states have finite correlation length. Theorem 2 of [8]). Let  $g_{\beta}$  be the thermal state at inverse temperature  $\beta$  of a Hamiltonian defined over a collection of spins. There exist an inverse temperature  $\beta_*^{(corr)} > 0$  such that for every  $\beta < \beta_*^{(corr)}$ , there is a constant  $\xi'$  such that the Gibbs state  $g_{\beta}$  satisfies  $\operatorname{Corr}_{\rho}(A, B) \leq 4 \operatorname{ac} e^{-d_H(A,B)/\xi'}$  for any regions A, B such that  $d_H(A, B) \geq L_0(a) := \xi' |\log(ac)|$  where  $a = \min\{|\partial^{(H)}A|, |\partial^{(H)}B|\}$  and  $c = (\log(3)(1 - e^{-1/\xi'}))^{-1}$ .

In Lemma S1, the geometry is determined by the interaction terms of the general Hamiltonian, which subsequently determines the distance between regions  $d_H(A, B)$  and their boundaries  $\partial^{(H)}A, \partial^{(H)}B$ . Specifically, Lemma S1 considers the *interaction hypergraph* of H, whose hyperedges are the supports of each of the interactions terms which make up H. The distance between two points in the interaction graph is the shortest path of hyperedges that connect them, and the boundary of a region consists of all hyperedges that contain a site inside the region and a site outside.

This very general statement implies, in our setting, that the Gibbs state has finite correlation length for  $\beta < \beta_*^{(\text{corr})}$ . Specifically, in the case of the *D*-dimensional periodic lattice and *k*-local Hamiltonian we which we study here, the threshold temperature is given by

$$\beta_*^{(\text{corr})} = \frac{1}{2 \max_i \|h_i\|_{\text{op}}} \log\Big(1 + \sqrt{1 + 4/\alpha}\Big),\tag{S9}$$

with  $\alpha \ge ((2R+1)^D - 1)e$  [8], where R is the side-length of a hypercubical region large enough to fit the (translated) support of any of the interaction terms  $h_i$  appearing in the Hamiltonian. The distance in the interaction hypergraph is related to the distance on the lattice by  $d_H(A, B) \ge d(A, B)/k$ . The extra factor k arises because the support of  $h_i$  can have at most diameter k in the lattice. Similarly  $|\partial^{(H)}A| \le b_1|\partial A|$ , (with  $b_1$  an unimportant constant which depends on k and D). Because of the relations above, as long as  $\beta < \beta_*^{(\text{corr})}$ , the Gibbs state has finite correlation length  $\xi \le \xi'/(Dk)$ , with  $f_{\text{corr}}(a) = b_1 4c^2 a^2$ . The extra factor (ca) in  $f_{\text{corr}}$  guarantees that the correlations are bounded for any distance d(A, B), as we demand in Definition S3, even if this distance is smaller than  $L_0$ .

#### 5. Locality of temperature

We expect that temperature is an intensive quantity, i.e., it is independent of the system size and thus can be defined locally. This intuition is captured by the *locality of temperature*, which, in a nutshell, asserts that any reduced density matrix for a localized subsystem of a Gibbs state is well described by the reduced density matrix for the same subsystem obtained from a *reduced* Gibbs state that is defined only for a region slightly larger the subsystem of interest. The formal definition is as follows.

**Definition S4** (Locality of temperature [8]). The locality of temperature at inverse temperature  $\beta$  is satisfied if for any subsystems  $A \subseteq B$ ,

$$\left\|g_{\beta} - g_{\beta}^{B}\right\|_{A} \le f_{\text{LoT}}(|\partial A|, |\partial B|)e^{-d(A,\overline{B})/\nu},\tag{S10}$$

where  $g^B_\beta$  is the Gibbs state of the Hamiltonian  $H|^B$  obtained by removing all terms of H which act nontrivially on  $\overline{B}$ , the complement of the region B,  $f_{\text{LoT}}$  is a polynomial function, and  $\nu$  is a constant.

This property is also called *local indistinguishability* [9–11] and is proven to hold at high temperatures. Specifically, Corollary 2 of Ref. [8] shows that if  $\beta < \beta_*^{(\text{corr})}$  [given by Eq. (S9)], then any local Hamiltonian satisfies the locality of temperature. Note that, as Lemma S1, Corollary 2 of Ref. [8] is stated in terms of the interaction hypergraph of the Hamiltonian and only describes distances  $d_H(A, \overline{B})$  above some  $L_0$  which grows as  $\log(|\partial^{(H)}A|)$ . These differences are resolved by selecting an appropriate function  $f_{\text{LoT}}$  and constant  $\nu$ , much in the same way as we did below Lemma S1.

#### 6. Subsystem purity

The locality of temperature guarantees that not only the global purity of the thermal states decays exponentially, but also the subsystem purity.

**Proposition S1** (Bound on the subsystem purity of the Gibbs state). Let H be a local translation invariant Hamiltonian defined over a D-dimensional periodic lattice and A a hypercubical region of side-length  $\ell = o(L)$  qubits (i.e. such that  $\lim_{L\to\infty} \ell/L = 0$ ). Assuming the locality of temperature, the subsystem purity of the Gibbs state is bounded as

$$\operatorname{tr}((g_{\beta,A})^2) \le e^{-\Omega(\ell)}.\tag{S11}$$

*Proof.* Let  $\ell$  denote the side-length of the hypercubical region A. Consider a bigger hypercube B with length  $\ell_B = \ell + 2d_0$ , which contains A padded around by  $d_0$  qubits. Note that  $d(A, \overline{B}) = d_0$ , so by the locality of temperature, we obtain

$$\left\|g_{\beta} - g_{\beta}^{B}\right\|_{A} \le \operatorname{poly}(\ell, d_{0})e^{-d_{0}/\nu},\tag{S12}$$

because  $|\partial A|$  and  $|\partial B|$  are themselves polynomial functions of  $\ell$  and  $d_0$ . The global and subsystem purities of the Gibbs state  $g^B_\beta$  can be related by

$$\operatorname{tr}\left((g^B_{\beta,A})^2\right) \le 2^{|B \setminus A|} \operatorname{tr}\left((g^B_{\beta})^2\right),\tag{S13}$$

which is a simple consequence of the weak subadditivity of the Rényi-2 entropy [12]. Using that  $|\operatorname{tr}(\rho_A^2) - \operatorname{tr}(\sigma_A^2)| \leq 2\|\rho - \sigma\|_A$  in Eq. (S12), Eq. (S13) with  $|B \setminus A| = (\ell + 2d_0)^D - \ell^D$ , and Eq. (S7) we obtain

$$\operatorname{tr}((g_{\beta,A})^2) \le 2^{(\ell+2d_0)^D - \ell^D - \Omega((\ell+2d_0)^D)} + \operatorname{poly}(\ell, d_0) e^{-d_0/\nu}.$$
(S14)

If we select  $d_0 = c_0 \ell/2$  with  $c_0 > 0$ , the exponent in the first term of the right hand side of Eq. (S14) becomes  $\ell^D((1+c_0)^D - 1 - \Omega(1)) \leq -\Omega(\ell^D)$  if we select sufficiently small  $c_0$ . With this selection, the last term in Eq. (S14) dominates, with a scaling of  $e^{-\Omega(\ell)}$ .

### II. THE EDGE THEOREM

We are now in position to present our first result. We expect that typical product states evolving under generic Hamiltonians will thermalize. Here, we formalize this intuition by considering energy-delocalized Gibbs ensembles (EDGEs) (see Definition 1 from the main text). We prove our EDGE theorem.

**Theorem S2** (EDGE theorem 2 restatement). Let H be a TI, (geometrically) local Hamiltonian with nondegenerate spectral gaps on a D-dimensional periodic lattice with N qubits and  $\mathcal{E}_{\beta}$  be an EDGE. Assume that the Gibbs state  $g_{\beta}$ has finite correlation length. Then, for any connected region A with a constant number of qubits  $N_A$ ,

$$\mathbb{E}_{\psi} \mathbb{E}_{t} \left[ \left\| g_{\beta} - \psi(t) \right\|_{A} \right] \le O\left( N^{-1/2+\delta} \right) + \varepsilon_{\mathrm{GE}}(N) + 2^{N_{A}} \varepsilon_{\mathrm{ED}}(N)^{1/2},$$

for any  $\delta > 0$ .

The rest of this section is devoted to proving Theorem S2. Our proof is inspired by Ref. [3], which studies 1D systems at infinite temperature. We will show that a typical state  $\psi$  in the EDGE satisfies two properties:

(a) Equivalence of ensembles between the temporal  $\rho_{\infty} = \mathbb{E}_t[\psi(t)]$  and the thermal ensemble  $g_{\beta}$ :

$$\|\rho_{\infty} - g_{\beta}\|_A \to 0$$

(b) Equilibration:

$$\mathbb{E}_{t}[\|\rho_{\infty} - \psi(t)\|_{A}] \to 0$$

Taken together, these two properties give us the desired quantum thermalization,

$$\mathbb{E}[\|g_{\beta} - \psi(t)\|_{A}] \to 0.$$

The equivalence of ensembles will follow from property (a) of an EDGE and a variant of the weak ETH [13–15]. We will explain this first. After, we will explain how equilibration follows from property (b) of an EDGE and the seminal results on quantum equilibration from Ref. [16].

#### A. Equivalence of the temporal and thermal ensembles

The weak ETH guarantees that with high likelihood, the expectation value of local observables over the energy eigenstates of a translation invariant system is close to the expectation at equilibrium, with probability exponentially close to 1. It was originally derived in Ref. [13], and then further improved in Refs. [14, 15]. We present a non-probabilistic version whose scaling is not exponential with the system size, but it is independent of spatial dimension. The latter property is favorable for our purposes.

**Proposition S2** (A variant of the weak ETH, Proposition 1 restatement). Let H be a TI, (geometrically) local Hamiltonian on a D-dimensional periodic lattice with N qubits. Let  $\rho = \sum_{j=1}^{2^N} p_j |E_j\rangle\langle E_j|$  be a diagonal state in the eigenbasis  $\{|E_j\rangle\}_j$  of H that has finite correlation length. Consider a connected region A with a constant number of qubits. Then, for any  $\delta > 0$ ,

$$\sum_{j=1}^{2^{N}} p_{j} \| |E_{j} \rangle \langle E_{j}| - \rho \|_{A} \le O\left(N^{-1/2+\delta}\right).$$
(S15)

*Proof.* Our proof initially follows Refs. [14, 15]. Let  $O_A$  be any observable supported on region A with operator norm  $||O_A||_{\text{op}} \leq 1$ . The first step is to utilize the translation invariance of the Hamiltonian to turn the expectation values of  $O_A$  into expectation values of an extensive observable  $\overline{O} = \sum_{n \in \Lambda} \mathbb{T}^n O_A \mathbb{T}^{-n} - N \operatorname{tr}(\rho O_A)$ . Note that

$$\sum_{j=1}^{2^{N}} p_{j} |\langle E_{j} | O_{A} | E_{j} \rangle - \operatorname{tr}(\rho O_{A})| = \frac{1}{N} \sum_{j=1}^{2^{N}} p_{j} |\langle E_{j} | \overline{O} | E_{j} \rangle |$$
$$\leq \frac{1}{N} \sum_{j=1}^{2^{N}} p_{j} \langle E_{j} | |\overline{O} | | E_{j} \rangle = \frac{1}{N} \operatorname{tr}(\rho |\overline{O}|),$$
(S16)

where  $\left|\overline{O}\right| = \sqrt{\overline{O}^2}$ .

The purpose of recasting the inequality in this form is to utilize known concentration bounds for the expectation value of extensive local observables on states with finite correlation length [17]. We diagonalize  $\overline{O} = \sum_{j=1}^{2^N} \lambda_j \Pi_j$ , where  $\Pi_j$  is the projector into the eigensubspace of  $\lambda_j$ . Let  $\gamma > 0$  (we will specify its dependence on N below). Consider the projectors into the bulk (inside a window of size width  $2\gamma$ ) and tails of the spectrum,

$$\Pi_{\text{Bulk}} = \sum_{j:|\lambda_j| < \gamma N} \Pi_j \qquad \qquad \Pi_{\text{Tails}} = \sum_{j:|\lambda_j| \ge \gamma N} \Pi_j. \tag{S17}$$

We can split the trace as

$$\operatorname{tr}(\rho|\overline{O}|) = \operatorname{tr}(\rho\Pi_{\operatorname{Bulk}}|\overline{O}|) + \operatorname{tr}(\rho\Pi_{\operatorname{Tails}}|\overline{O}|).$$
(S18)

We will show that the contribution from the tails is exponentially small in N, while the contribution from the bulk decays as  $N^{-1/2+\delta}$ . The trace on the bulk is bounded as

$$\operatorname{tr}(\rho\Pi_{\operatorname{Bulk}}|\overline{O}|) = \sum_{j:|\lambda_j| < \gamma N} \operatorname{tr}(\rho\Pi_j)|\lambda_j| \le \gamma N$$
(S19)

To bound the trace on the tails, note that

$$\operatorname{tr}(\rho\Pi_{\operatorname{Tails}}|\overline{O}|) \le \left\|\rho\Pi_{\operatorname{Tails}}|\overline{O}|\right\| \le \left\|\rho\Pi_{\operatorname{Tails}}\right\|\left\|\overline{O}\right\|_{\operatorname{op}} = \operatorname{tr}(\rho\Pi_{\operatorname{Tails}})\left\|\overline{O}\right\|_{\operatorname{op}} \le \operatorname{tr}(\rho\Pi_{\operatorname{Tails}})2N$$

where we used Hölder's inequality, the positive semidefiniteness of  $\rho \Pi_{\text{Tails}}$  and  $\|O_A\|_{\text{op}} \leq 1$ . We now use the concentration bound proven in Ref [17].

**Lemma S2** (Th. 4.2 of [17]). If  $\gamma \geq \frac{K_1}{\sqrt{N}}$ , then  $\operatorname{tr}(\Pi_{\operatorname{Tails}}\rho) \leq 2K_2 e^{-K_3(N\gamma^2)^{1/(D+1)}}$ , where the constants  $K_1, K_2$  and  $K_3$  depend on D, |A|,  $\xi$  and  $C = f_{\operatorname{corr}}(|\partial A|)$  (See Ref. [17] for their explicit form).

If we pick  $\gamma = O(N^{-1/2+\delta})$ , by Lemma S2, we obtain

$$\frac{1}{N}\operatorname{tr}\left(\rho\Pi_{\operatorname{Tails}}\left|\overline{O}\right|\right) \le e^{-\Omega(N^{2\delta/(D+1)})}.$$
(S20)

From Eq. (S19) we get

$$\frac{1}{N}\operatorname{tr}\left(\rho\Pi_{\operatorname{Bulk}}\left|\overline{O}\right|\right) \le O(N^{-1/2+\delta}).$$
(S21)

Inserted back into Eqs. (S16) and (S18), the contribution from the bulk dominates so we obtain

$$\sum_{j=1}^{2^N} p_j |\langle E_j | O_A | E_j \rangle - \operatorname{tr}(\rho O_A)| \le O\left(N^{-1/2+\delta}\right).$$

The prefactors on the right-hand side are independent of the choice of observable  $O_A$ , so we can maximize over all  $O_A$  and we obtain Eq. (S15).

**Proposition S3** (Equivalence of temporal and thermal ensembles). Under the assumptions of the EDGE Theorem S2,

$$\mathop{\mathbb{E}}_{\boldsymbol{\phi}\in\mathcal{E}_{\beta}}[\|\boldsymbol{\rho}_{\infty}-\boldsymbol{g}_{\beta}\|_{A}] = O\left(N^{-\frac{1}{2}+\delta}\right) + \varepsilon_{\mathrm{GE}}(N), \tag{S22}$$

where  $\rho_{\infty} = \mathbb{E}_t[\psi(t)]$ , for any  $\delta > 0$ .

*Proof.* By the no-degeneracy condition, the infinite-time-average of any state  $\rho_{\infty}$  is the diagonal ensemble  $\rho_{\infty} = \sum_{j} |\langle \psi | E_{j} \rangle|^{2} |E_{j} \rangle \langle E_{j}|$ . Then

$$\begin{split} & \underset{\psi \in \mathcal{E}_{\beta}}{\mathbb{E}} [\|\rho_{\infty}(\psi) - g_{\beta}\|_{A}] = \underset{\psi \in \mathcal{E}_{\beta}}{\mathbb{E}} \left[ \left\| \sum_{j} \left( |E_{j}\rangle\langle E_{j}| - g_{\beta} \right)|\langle\psi|E_{j}\rangle|^{2} \right\|_{A} \right] \\ & \leq \sum_{j} \||E_{j}\rangle\langle E_{j}| - g_{\beta}\|_{A} \underset{\psi \in \mathcal{E}_{\beta}}{\mathbb{E}} \left[ |\langle\psi|E_{j}\rangle|^{2} \right] \\ & = \sum_{j} \||E_{j}\rangle\langle E_{j}| - g_{\beta}\|_{A} \langle E_{j}| \underset{\psi \in \mathcal{E}_{\beta}}{\mathbb{E}} [\psi] |E_{j}\rangle \\ & \leq \sum_{j} \||E_{j}\rangle\langle E_{j}| - g_{\beta}\|_{A} \left(\langle E_{j}|g_{\beta}|E_{j}\rangle + 2^{-N}\varepsilon_{\mathrm{GE}}(N)\right) \\ & \leq \left(\sum_{j} \||E_{j}\rangle\langle E_{j}| - g_{\beta}\|_{A} \langle E_{j}|g_{\beta}|E_{j}\rangle\right) + \varepsilon_{\mathrm{GE}}(N) \\ & = O\left(N^{-\frac{1}{2}+\delta}\right) + \varepsilon_{\mathrm{GE}}(N), \end{split}$$

where we used property (a) of the EDGE and Proposition S2.

#### B. Quantum equilibration

Now we shift gears to prove equilibration. We utilize the following result from Ref. [16]:

**Lemma S3** (Quantum equilibration, Theorem 1 of Ref. [16]). Let H be a Hamiltonian which has nondegenerate spectral gaps. For any state  $\psi$  with time average  $\rho_{\infty} = \mathbb{E}_t[\psi(t)]$ , and subsystem A of dimension  $d_A$ 

$$\mathbb{E}_{t}[\|\rho_{\infty} - \rho(t)\|_{A}] \le d_{A}\sqrt{\mathrm{IPR}},\tag{S23}$$

where IPR =  $\sum_{j} |\langle E_j | \psi \rangle|^4$  is the inverse participation ratio of  $|\psi\rangle$ .

**Proposition S4** (EDGE equilibration). Under the assumptions of the EDGE Theorem S2,

$$\mathop{\mathbb{E}}_{\psi\in\mathcal{E}_{\beta}}\left[\mathop{\mathbb{E}}_{t}\left[\left\|\rho_{\infty}-\rho(t)\right\|_{A}\right]\right] \leq 2^{N_{A}}\varepsilon_{\mathrm{ED}}(N)^{1/2} \tag{S24}$$

*Proof.* Averaging over all states in the EDGE  $\mathcal{E}_{\beta}$ , using that  $\mathbb{E}[\cdot]^2 \leq \mathbb{E}[(\cdot)^2]$ , and property (b) we obtain

$$\underset{\psi \in \mathcal{E}_{\beta}}{\mathbb{E}}[\underset{t}{\mathbb{E}}[\|\rho_{\infty} - \rho(t)\|_{A}]] \leq \sqrt{\underset{\psi \in \mathcal{E}_{\beta}}{\mathbb{E}}[\underset{t}{\mathbb{E}}[\|\rho_{\infty} - \rho(t)\|_{A}]^{2}]} \leq 2^{N_{A}} \varepsilon_{\mathrm{ED}}(N)^{1/2}.$$

Combining Propositions S3 and S4, by summing Eq. (S22) and Eq. (S24) and applying a triangle inequality, we complete the proof of the EDGE Theorem S2.

Next, will show that for any TI, local Hamiltonian, there exists an EDGE of product states at high enough, but finite temperatures. We invoke a recent result, Theorem 1.5 of Ref. [18], which shows that high-temperature Gibbs states are unentangled for some  $\beta$  below a constant  $\beta_*^{(sep)}$ . Thus, at high temperatures we can construct an ensemble  $\mathcal{E}_{\beta} = \{(p_{\psi}, |\psi\rangle\langle\psi|)\}$  of product states which automatically satisfies property (a) of an EDGE, with  $\varepsilon_{\text{GE}}(N) = 0$ . Moreover, because the thermal state uniquely maximizes the von Neumann entropy  $S(\rho) = -\operatorname{tr}(\rho \log \rho)$ with constrained energy [Eq. (S6)], any product-state ensemble  $\mathcal{E}$  whose mean state  $\rho_{\mathcal{E}} = \mathbb{E}_{\psi \in \mathcal{E}}[\psi]$  also maximizes this quantity

$$\mathcal{E}_{\beta} \in \underset{\operatorname{tr}(\rho_{\mathcal{E}}H) = \operatorname{tr}(g_{\beta}H)}{\operatorname{argmax}} S(\rho_{\mathcal{E}})$$
(S25)

is a Gibbs ensemble. In what follows, we will prove that any Gibbs ensemble of product states is necessarily an EDGE, with  $\varepsilon_{\text{ED}}(N) \leq O(N^{-1+\delta})$ , under certain conditions, which completes the proof of our main result, Theorem 1.

#### III. BOUND ON AVERAGE IPR

In this section, we prove Theorem 3 of the main text.

**Theorem S3** (Bound on average IPR, Theorem 3 restatement). Let H be a TI, (geometrically) local, Hamiltonian on a D-dimensional periodic lattice with N qubits, and consider an ensemble  $\mathcal{E}_{\beta}$  of product states whose mean value is the Gibbs state  $g_{\beta}$ , at high temperature  $\beta < \beta_*^{(corr)}$  (or at any temperature as long as  $g_{\beta}$  has finite correlation length and satisfies the locality of temperature). Then, the average IPR in a common eigenbasis  $\{|E_j\rangle\}_j$  of the translation operator and the Hamiltonian is bounded

$$\mathbb{E}_{\psi}\left[\sum_{j=1}^{2^{N}} |\langle E_{j} | \psi \rangle|^{4}\right] \le O(N^{-1+\delta})$$
(S26)

for any  $\delta > 0$ .

The proof of Theorem S3 below generalizes that of Proposition 2 of the main text. The key idea to recall is that one may upper bound the average IPR by separately upper bounding (i) the overlap of the TI eigenstates with the aperiodic computational-basis states and (ii) the probability of a bitstring being periodic. We generalize this idea to arbitrary product states  $|\psi\rangle = \bigotimes_{i \in \Lambda} |\psi_i\rangle$ . We face one immediate obstacle: For an aperiodic bitstring  $b \in \{0,1\}^N$ the state  $|b\rangle$  is orthogonal to all of its translates  $\langle b|\mathbb{T}^n|b\rangle = 0$  for all 0 < n < N. We relied on this orthogonality in deducing that  $|\langle b|E_j\rangle|^2 \leq 1/N$  for aperiodic b. However, for a general product state  $|\psi\rangle$ , even in 1D, it could be that  $\langle \psi|\mathbb{T}^n|\psi\rangle \neq 0$  even if the sequence  $\{|\psi_1\rangle, |\psi_2\rangle, \dots, |\psi_N\rangle\}$  is not periodic. To circumvent this problem, we establish a robust notion of *approximate* periodicity. We split the expectation value of the IPR into an average over (approximately) periodic and aperiodic states, and prove that (i) the IPR of states which are not approximately periodic is upper bounded by  $\sim 1/N$  and (ii) the approximately periodic states are atypical in any unraveling of the Gibbs state, i.e., they are associated with vanishingly small probabilities.

#### A. Aperiodic states have low IPR

**Definition S5.** (Approximately periodic state). An arbitrary state  $|\psi\rangle$  for N qubits arranged on the D-dimensional lattice  $\Lambda$  is r-approximately periodic if there exists  $\mathbf{n} \in \Lambda$  with  $\mathbf{n} \neq \mathbf{0}$  such that  $|\langle \psi | \mathbb{T}^{\mathbf{n}} | \psi \rangle|^2 \geq 2^{-r}$ . The vector  $\mathbf{n}$  is called a period of  $|\psi\rangle$ .

We begin by generalizing our decomposition of the average IPR from Proposition 2 for approximately periodic states and arbitrary ensembles. We state the result in a general form, which might be useful outside of our current context.

**Lemma S4** (Probabilistic bound on average IPR from symmetry). Consider a set of symmetry operators represented by mutually commuting unitaries  $\{U_1, \ldots, U_M\}$ . Let P be the set of approximately symmetric states, containing all  $\psi$ such that  $|\langle \psi | U_n^{\dagger} U_m | \psi \rangle| \geq \varepsilon$  for some  $n \neq m$ . For any basis  $\{|\alpha\rangle\}_{\alpha}$  of simultaneous eigenstates of all  $U_n$ , the average IPR over any ensemble of states  $\mathcal{E}$  can be upper bounded as

$$\mathop{\mathbb{E}}_{\psi \in \mathcal{E}} \left[ \sum_{\alpha} |\langle \alpha | \psi \rangle|^4 \right] \le \frac{1}{M} + \varepsilon + \mathop{\mathbb{P}}_{\psi \in \mathcal{E}} [\psi \in P], \tag{S27}$$

where  $\mathbb{P}_{\psi \in \mathcal{E}}[\psi \in P]$  denotes the probability that a state drawn from the ensemble  $\mathcal{E}$  is approximately symmetric.

*Proof.* To begin, we split the expectation value conditioning on P,

$$\mathbb{E}_{\psi\in\mathcal{E}}\left[\sum_{\alpha}\left|\langle\alpha|\psi\rangle\right|^{4}\right] = \mathbb{E}_{\psi\in\mathcal{E}}\left[\sum_{\alpha}\left|\langle\alpha|\psi\rangle\right|^{4}\left|\psi\notin P\right]\mathbb{P}_{\psi\in\mathcal{E}}[\psi\notin P] + \mathbb{E}_{\psi\in\mathcal{E}}\left[\sum_{\alpha}\left|\langle\alpha|\psi\rangle\right|^{4}\left|\psi\in P\right]\mathbb{P}_{\psi\in\mathcal{E}}[\psi\in P]\right]$$
(S28)

$$\leq \mathop{\mathbb{E}}_{\psi \in \mathcal{E}} \left[ \sum_{\alpha} \left| \langle \alpha | \psi \rangle \right|^4 \middle| \psi \notin P \right] + \mathop{\mathbb{P}}_{\psi \in \mathcal{E}} [\psi \in P], \tag{S29}$$

where the notation  $\mathbb{E}[\dots | \dots]$  denotes a conditional expectation value. Now we bound the first term, which is the expectation value conditioned on  $\psi \notin P$ , i.e.,  $\psi$  not being approximately symmetric. We denote  $|\psi_n\rangle = U_n |\psi\rangle$ , which

are approximately orthogonal states  $|\langle \psi_n | \psi_m \rangle| < \varepsilon$  if  $\psi \notin P$ . Moreover, observe that  $|\langle \alpha | \psi_n \rangle| = |\langle \alpha | U | \psi \rangle| = |\langle \alpha | \psi \rangle|$ , thus

$$M|\langle \alpha|\psi\rangle|^{2} = \sum_{n=0}^{M-1} \langle \alpha|\psi_{n}\rangle \langle \psi_{n}|\alpha\rangle \leq \langle \alpha|(\tilde{\Pi}-\Pi)|\alpha\rangle + 1,$$
(S30)

where  $\Pi$  is the orthogonal projector into the subspace spanned by  $|\psi_n\rangle$  and  $\Pi := \sum_n |\psi_n\rangle\langle\psi_n|$ . We now show that these two operators are close to one another for  $\psi \notin P$ . Specifically,

$$\langle \alpha | (\tilde{\Pi} - \Pi) | \alpha \rangle^2 \leq \| \tilde{\Pi} - \Pi \|_{\rm op}^2 \leq \| \tilde{\Pi} - \Pi \|_{\rm F}^2 = \operatorname{tr} \left( (\tilde{\Pi} - \Pi)^2 \right) \leq \operatorname{tr} \left( \tilde{\Pi}^2 \right) - M, \tag{S31}$$

where  $||A||_{\rm F} = \sqrt{\operatorname{tr}(AA^{\dagger})}$  is the Frobenius norm, which always upper bounds the operator norm. We may compute,  $\operatorname{tr}(\tilde{\Pi}^2) = \sum_{n,m} |\langle \psi_n | \psi_m \rangle|^2 \leq M(M-1)\varepsilon^2 + M \leq M^2 \varepsilon^2 + M$ , which inserted into Eq. (S31) gives  $\langle \alpha | (\tilde{\Pi} - \Pi) | \alpha \rangle \leq M \varepsilon$ . Thus, from Eq. (S30) we obtain  $|\langle \alpha | \psi \rangle|^2 \leq 1/M + \varepsilon$  for each  $\alpha$ . Summing over all  $\alpha$ , we can bound the IPR of  $\psi \notin P$ as  $\sum_{\alpha} |\langle \alpha | \psi \rangle|^4 \leq \sum_{\alpha} |\langle \alpha | \psi \rangle|^2 (1/M + \varepsilon) = 1/M + \varepsilon$ , where we used the normalization  $\sum_{\alpha} |\langle \alpha | \psi \rangle|^2 = 1$ . Averaging over  $\psi \notin P$ , we obtain that

$$\mathbb{E}_{\psi \in \mathcal{E}} \left[ \sum_{\alpha} \left| \langle \alpha | \psi \rangle \right|^4 \middle| \psi \notin P \right] \le \frac{1}{M} + \varepsilon.$$

Lemma S4 provides a meaningful bound for the average IPR whenever P is atypical in the ensemble  $\mathcal{E}$ , and  $\varepsilon$  is small. We apply Lemma S4 with the unitaries  $U_n$  taken as certain translations on the lattice, which share a basis of eigenstates with the TI Hamiltonian, and we select P to be the set of r-approximately periodic states with periods larger than the translations, and show that  $\mathbb{P}_{\psi \in \mathcal{E}_{\beta}} [\psi \in P]$  is exponentially small.

Let  $\delta > 0$  (smaller than 1). This is the same arbitrary  $\delta$  that appears in the statement of Theorem S3, which will remain fixed henceforth. We consider lattice sites  $x_1, x_2, \ldots, x_M \in \Lambda$  pairwise separated as  $||x_n - x_m||_1 \ge L^{\delta}$  for all  $n \neq m$ . We can select at least

$$M = \left( \left\lfloor \frac{L}{\lceil L^{\delta} \rceil} \right\rfloor - 1 \right)^{D} \ge \Omega(N^{1-\delta})$$

such sites (think of  $\{\boldsymbol{x}_i\}_{i=1}^M$  as approximately forming a coarser sublattice). We take  $U_n = \mathbb{T}^{\boldsymbol{x}_n}$ , so that  $|\langle \psi | U_m^{\dagger} U_n | \psi \rangle| \geq 2^{-r/2}$  for  $n \neq m$  implies that  $\psi$  is r-approximately periodic with a period  $\|\boldsymbol{n}\|_1 = \|\boldsymbol{x}_n - \boldsymbol{x}_m\|_1 \geq L^{\delta}$ , i.e.,

$$P \coloneqq \left\{\psi \left| \exists \boldsymbol{n} \in \Lambda \colon \left\| \boldsymbol{n} \right\|_{1} \ge L^{\delta}, \left| \langle \psi | \mathbb{T}^{\boldsymbol{n}} | \psi \rangle \right|^{2} \ge 2^{-r} 
ight\}.$$

By Lemma S4 we obtain

$$\mathbb{E}_{\psi\in\mathcal{E}_{\beta}}\left[\sum_{j}|\langle E_{j}|\psi\rangle|^{4}\right] \leq O\left(\frac{1}{N^{1-\delta}}\right) + 2^{-r/2} + \mathbb{P}_{\psi\in\mathcal{E}_{\beta}}\left[\psi\in P\right].$$
(S32)

For now, r is arbitrary. We will select an appropriate scaling r(N) at the end of the proof. The reason to pick increasing periods  $\|\boldsymbol{n}\|_1 \ge L^{\delta}$  is that this will guarantee that  $\mathbb{P}_{\psi \in \mathcal{E}_{\beta}}[\psi \in P]$  is exponentially small, as we prove next.

## B. Approximately periodic states are atypical

Intuitively, periodic states with large periods will be long-range correlated, so they cannot have a significant population in  $\mathcal{E}_{\beta}$ , whose average is the Gibbs state with finite correlation length (Lemma S1). Specifically, we bound  $\mathbb{P}_{\psi \in \mathcal{E}_{\beta}} [\psi \in P]$  by separately bounding the probability of a state having fixed period  $\boldsymbol{n}$ , i.e. of being in

$$P_{\boldsymbol{n}} = \left\{ \psi \left| \left| \langle \psi | \mathbb{T}^{\boldsymbol{n}} | \psi \rangle \right|^2 \ge 2^{-r} \right\},$$
(S33)

and then taking the union bound over  $P = \bigcup_{\|n\|_1 > L^{\delta}} P_n$ .

**Proposition S5** (Bound on probability of approximately periodic states). Consider an ensemble of product states  $\mathcal{E}$  for N qubits arranged on the D-dimensional lattice  $\Lambda$ , with mean state  $\rho = \mathbb{E}_{\psi \in \mathcal{E}}[|\psi\rangle\langle\psi|]$  which is translation invariant

and has finite correlation length  $\xi$ . Let  $\mathbf{n} \in \Lambda$  and consider a hypercubical region A with side-length  $\ell < \|\mathbf{n}\|_1/D$  and  $N_A = \ell^D$  qubits. Then,

$$\mathbb{P}_{\boldsymbol{\psi}\in\mathcal{E}}\left[\boldsymbol{\psi}\in P_{\boldsymbol{n}}\right] \le 2^{r} \left(2^{(\ell^{D})} f_{\operatorname{corr}}(2D\ell^{D-1}) e^{-(\|\boldsymbol{n}\|_{1}-D\ell)/\xi} + \operatorname{tr}\left(\rho_{A}^{2}\right)\right),\tag{S34}$$

where  $\rho_A = \operatorname{tr}_{\bar{A}}(\rho)$ , and  $f_{\operatorname{corr}}$  is the polynomial function from Definition S3.

Proof. Let  $O_A = \bigotimes_{i \in A} \sigma_i$  be an arbitrary product of Paulis matrices  $\sigma_i \in \{X, Y, Z, 1\}$  defined over A. Let A + n denote the translation of the region A over the vector n and  $O_{A+n} = \mathbb{T}^n O_A \mathbb{T}^{-n}$ . Note that the distance between A and A + n is at least  $||n||_1 - D\ell$ . Then, because  $\rho$  has finite correlation length  $\xi$  and the number of qubits in the boundary of A is  $|\partial A| = 2D\ell^{D-1}$ ,

$$f_{\rm corr}(2D\ell^{D-1})e^{-(\|\boldsymbol{n}\|_1 - D\ell)/\xi} \ge \operatorname{tr}(\rho O_A \otimes O_{A+\boldsymbol{n}}) - \operatorname{tr}(\rho_A O_A)\operatorname{tr}(\rho_{A+\boldsymbol{n}} O_{A+\boldsymbol{n}})$$
(S35)

$$= \operatorname{tr}\left(\left(\mathbb{E}_{\psi\in\mathcal{E}}\left[\bigotimes_{i\in A}\psi_{i}\otimes\bigotimes_{i\in A}\psi_{i+n}\right] - \rho_{A}\otimes\rho_{A+n}\right)O_{A}\otimes O_{A+n}\right).$$
 (S36)

Summing over all  $4^{(\ell^D)}$  possible Pauli strings over A, and using that  $\sum_{O_A} O_A \otimes O_{A+n} = 2^{(\ell^D)} S_{A,A+n}$ , where  $S_{A,A+n}$  is the swap operator between regions A and A+n, we get

$$2^{(\ell^{D})} f_{\operatorname{corr}}(2D\ell^{D-1}) e^{-(\|\boldsymbol{n}\|_{1} - D\ell)/\xi} \geq \operatorname{tr}\left(\left(\underset{\boldsymbol{\psi}\in\mathcal{E}}{\mathbb{E}}\left[\bigotimes_{\boldsymbol{i}\in A}\psi_{\boldsymbol{i}}\otimes\bigotimes_{\boldsymbol{i}\in A}\psi_{\boldsymbol{i}+\boldsymbol{n}}\right] - \rho_{A}\otimes\rho_{A+\boldsymbol{n}}\right)S_{A,A+\boldsymbol{n}}\right)$$
$$= \underset{\boldsymbol{\psi}\in\mathcal{E}}{\mathbb{E}}\left[\prod_{\boldsymbol{i}\in A}|\langle\psi_{\boldsymbol{i}}|\psi_{\boldsymbol{i}+\boldsymbol{n}}\rangle|^{2}\right] - \operatorname{tr}\left(\rho_{A}^{2}\right)$$
$$\geq \underset{\boldsymbol{\psi}\in\mathcal{E}}{\mathbb{E}}\left[\prod_{\boldsymbol{i}\in A}|\langle\psi_{\boldsymbol{i}}|\psi_{\boldsymbol{i}+\boldsymbol{n}}\rangle|^{2}\Big|\psi\in P_{\boldsymbol{n}}\right]\underset{\boldsymbol{\psi}\in\mathcal{E}}{\mathbb{P}}[\psi\in P_{\boldsymbol{n}}] - \operatorname{tr}\left(\rho_{A}^{2}\right).$$
$$\geq 2^{-r}\underset{\boldsymbol{\psi}\in\mathcal{E}}{\mathbb{P}}[\psi\in P_{\boldsymbol{n}}] - \operatorname{tr}\left(\rho_{A}^{2}\right).$$
(S37)

where we used that  $\operatorname{tr}(S(W \otimes V)) = \operatorname{tr}(WV)$ , the translation invariance  $\rho_A = \rho_{A+n}$  and that for every  $\psi \in P_n$  we have  $\prod_{i \in A} |\langle \psi_i | \psi_{i+n} \rangle|^2 \ge \prod_{i \in \Lambda} |\langle \psi_i | \psi_{i+n} \rangle|^2 \ge 2^{-r}$ .

Performing a union bound over all  $\|\boldsymbol{n}\|_1 \geq L^{\delta}$  in Eq. (S34) and taking  $\rho = g_{\beta}$  and  $\mathcal{E} = \mathcal{E}_{\beta}$ , we obtain

$$\mathbb{P}_{\psi\in\mathcal{E}_{\beta}}\left[\psi\in P\right] \leq N2^{r} \left(2^{(\ell^{D})} f_{\operatorname{corr}}(2D\ell^{D-1})e^{-(L^{\delta}-D\ell)/\xi} + \operatorname{tr}\left((g_{\beta,A})^{2}\right)\right).$$
(S38)

The term  $\operatorname{tr}((g_{\beta,A})^2)$  is the subsystem purity of  $g_{\beta}$ . In Proposition S1, we proved from the locality of temperature that this quantity is exponentially small in  $\ell$ , i.e.,  $\operatorname{tr}((g_{\beta,A})^2) \leq e^{-\Omega(\ell)}$ . Thus, from Eq. (S38) we get the asymptotic bound on the probability of *r*-approximately states

$$\mathbb{P}_{\psi \in \mathcal{E}_{\beta}} \left[ \psi \in P \right] \le N2^{r} \left( \operatorname{poly}(\ell) e^{O(\ell^{D})} e^{-\Omega(L^{\delta})} + e^{-\Omega(\ell)} \right),$$
(S39)

where  $poly(\ell)$  denotes a polynomial on  $\ell$ . Inserting Eq. (S39) into Eq. (S32) gives the bound on the IPR,

$$\mathbb{E}_{\psi \in \mathcal{E}_{\beta}} \left[ \sum_{j} |\langle E_{j} | \psi \rangle|^{4} \right] \leq O\left(\frac{1}{N^{1-\delta}}\right) + N \operatorname{poly}(\ell) e^{O(\ell^{D}) - \Omega(L^{\delta}) + O(r)} + N e^{O(r) - \Omega(\ell)} + e^{-\Omega(r)}.$$
(S40)

We are free to select how  $\ell$  and r scale with  $L = N^{1/D}$ . We select the scaling so the first term on the right-hand side of Eq. (S40) dominates over the other three terms. Specifically, if we pick

$$\ell = \Theta(L^a)$$
 and  $r = \Theta(L^b)$  (S41)

with  $0 < b < a < \delta/D$ , then the last three terms on the right-hand side of Eq. (S40) are upper bounded by  $e^{-\Omega(L^b)}$ , over which the first term dominates. Then we may conclude, as  $e^{-\Omega(L^b)} \leq O(1/N^{1-\delta})$ ,

$$\mathbb{E}_{\psi \in \mathcal{E}_{\beta}} \left[ \sum_{j} |\langle E_{j} | \psi \rangle|^{4} \right] \le O\left(\frac{1}{N^{1-\delta}}\right), \tag{S42}$$

which is the statement of Theorem S3.

### REFERENCES

- D. K. Mark, F. Surace, A. Elben, A. L. Shaw, J. Choi, G. Refael, M. Endres, and S. Choi, A maximum entropy principle in deep thermalization and in Hilbert-space ergodicity (2024), arXiv:2403.11970 [quant-ph].
- [2] Y. Huang, Extensive entropy from unitary evolution (2021), 2104.02053 [quant-ph].
- [3] Y. Huang and A. W. Harrow, Scrambling and thermalization in translation-invariant systems (2019), 1907.13392 [condmat.dis-nn].
- [4] M. P. Müller, E. Adlam, L. Masanes, and N. Wiebe, Thermalization and canonical typicality in translation-invariant quantum lattice systems, Communications in Mathematical Physics 340, 499–561 (2015).
- [5] B. Simon, The Statistical Mechanics of Lattice Gases, Volume I (Princeton University Press, 1993).
- [6] H. Araki, Gibbs states of a one dimensional quantum lattice, Communications in Mathematical Physics 14, 120–157 (1969).
- [7] A. Bluhm, A. Capel, and A. Pérez-Hernández, Exponential decay of mutual information for Gibbs states of local Hamiltonians, Quantum 6, 650 (2022).
- [8] M. Kliesch, C. Gogolin, M. J. Kastoryano, A. Riera, and J. Eisert, Locality of temperature, Phys. Rev. X 4, 031019 (2014).
- [9] F. G. S. L. Brandão and M. J. Kastoryano, Finite correlation length implies efficient preparation of quantum thermal states, Communications in Mathematical Physics 365, 1–16 (2018).
- [10] Ángela Capel, M. Moscolari, S. Teufel, and T. Wessel, From decay of correlations to locality and stability of the Gibbs state (2024), arXiv:2310.09182 [math-ph].
- [11] A. Bluhm, Ángela Capel, and A. Pérez-Hernández, Strong decay of correlations for Gibbs states in any dimension (2024), arXiv:2401.10147 [quant-ph].
- [12] W. van Dam and P. Hayden, Renyi-entropic bounds on quantum communication (2002), arXiv:quant-ph/0204093 [quant-ph].
- [13] T. Mori, Weak eigenstate thermalization with large deviation bound (2016), arXiv:1609.09776 [cond-mat.stat-mech].
- [14] F. G. S. L. Brandão, E. Crosson, M. B. Şahinoğlu, and J. Bowen, Quantum error correcting codes in eigenstates of translation-invariant spin chains, Phys. Rev. Lett. 123, 110502 (2019).
- [15] A. M. Alhambra, J. Riddell, and L. P. García-Pintos, Time evolution of correlation functions in quantum many-body systems, Phys. Rev. Lett. 124, 110605 (2020).
- [16] N. Linden, S. Popescu, A. J. Short, and A. Winter, Quantum mechanical evolution towards thermal equilibrium, Phys. Rev. E 79, 061103 (2009).
- [17] A. Anshu, Concentration bounds for quantum states with finite correlation length on quantum spin lattice systems, New Journal of Physics 18, 083011 (2016), we refer to arXiv:1508.07873 which has been updated after publication.
- [18] A. Bakshi, A. Liu, A. Moitra, and E. Tang, High-temperature Gibbs states are unentangled and efficiently preparable (2024), arXiv:2403.16850 [quant-ph].