Bootstrap Adaptive Lasso Solution Path Unit Root Tests[∗]

Martin C. Arnold†

Thilo Reinschlüssel†‡

martin.arnold@vwl.uni-due.de

thilo.reinschluessel@vwl.uni-due.de

September 12, 2024

Abstract

We propose sieve wild bootstrap analogues to the adaptive Lasso solution path unit root tests of Arnold and Reinschlüssel [\(2024\)](#page-26-0) to improve finite sample properties and extend their applicability to a generalised framework, allowing for non-stationary volatility. Numerical evidence shows the bootstrap to improve the tests' precision for error processes that promote spurious rejections of the unit root null, depending on the detrending procedure. The bootstrap mitigates finite-sample size distortions and restores asymptotically valid inference when the data features time-varying unconditional variance. We apply the bootstrap tests to real residential property prices of the top six Eurozone economies and find evidence of stationarity to be period-specific, supporting the conjecture that exuberance in the housing market characterises the development of Euro-era residential property prices in the recent past.

Keywords: Adaptive Lasso, Autoregressions, Unit root testing, Bootstrap inference, Heteroskedastic errors

JEL classifications: C52, C22, C12, R31

[†]Faculty of Business Administration and Economics, University of Duisburg-Essen, Universitätsstraße 12, 45117 Essen, Germany

[‡]RGS Econ - Ruhr Graduate School in Economics, Hohenzollernstraße 1-3, 45128 Essen, Germany

1 Motivation

In Arnold and Reinschlüssel [\(2024\)](#page-26-0), we propose the *τ* and *τ*˘ adaptive Lasso unit root tests, which exploit distinct stochastic orders of the activation knots of the lagged level regressor on the Lasso solution path in stationary and non-stationary autoregressions. Simulations reveal a decline in the tests' precision when adjusting for deterministic components in the presence of higher-order serial correlation. As is well-documented for established unit root tests (cf. Ng & Perron, [2001\)](#page-28-0), τ and $\tilde{\tau}$ also are vulnerable to large negative moving average (MA) coefficients, which cause significant upward size distortions. Time-varying variance in the error process can exacerbate such distortions. The latter invalidates the tests' homoskedastic limiting null distributions, further promoting spurious rejections.

A well-established strand of the literature on time series regressions addresses inference on model parameters when the generating process affects a test's distribution with nuisance terms. Several contributions to the unit root literature employ the bootstrap for first-order approximation of the null distribution to improve the finite-sample precision and ensure the asymptotic validity of a test. For autoregressive (AR) models, examples are recursive bootstraps in AR(1) models (Ferretti & Romo, [1996;](#page-27-0) Swensen, [2003\)](#page-29-0), sieve bootstraps (Chang & Park, [2003;](#page-27-1) Park, [2003\)](#page-28-1) and the residual-based block bootstrap (Paparoditis & Politis, [2003\)](#page-28-2). Cavaliere and Taylor [\(2008,](#page-26-1) [2009a,](#page-26-2) [2009b\)](#page-26-3) devise wild bootstrap (Liu, [1988\)](#page-28-3) tests for asymptotically valid inference under conditionally heteroskedastic innovations as well as general forms of a time-varying unconditional variance. They propose analogues of the *M* unit root tests suggested by Ng and Perron [\(2001\)](#page-28-0), Perron and Ng [\(1996\)](#page-28-4), and Stock [\(1999\)](#page-29-1), the tests of Phillips [\(1987\)](#page-28-5) and Phillips and Perron [\(1988\)](#page-28-6) and the augmented Dickey-Fuller (ADF) tests of Elliott et al. [\(1996\)](#page-27-2) and Said and Dickey [\(1984\)](#page-29-2).

Bootstraps have also been extensively applied in the recent literature on inference for penalised estimators. Chatterjee and Lahiri [\(2011\)](#page-27-3) utilise a residual-based bootstrap to reliably estimate both the distribution and bias of adaptive Lasso estimators in a cross-sectional setting. Similarly, Audrino and Camponovo [\(2017\)](#page-26-4) use a residual-based block bootstrap for heteroskedasticity-robust inference in adaptively *ℓ*1-penalised time series regressions. Chernozhukov et al. [\(2013\)](#page-27-4) sparked active research by applying the multiplier (wild) bootstrap to penalised estimation in a high-dimensional heteroskedastic setup. Notably, they resample factors that shape the asymptotic distribution, excluding terms that drive asymptotically negligible nuisance parameters by not recalculating the

estimator (the Dantzig selector of Candes & Tao, [2007\)](#page-26-5) during each bootstrap cycle. Hansen and Liao [\(2018\)](#page-27-5) point out that this procedure is computationally efficient to obtain asymptotically valid inference or confidence intervals at the cost of failing to 'capture any finite sample uncertainty introduced in the lasso selection'. Contrary to this minimalistic approach, Dezeure et al. [\(2017\)](#page-27-6) estimate the distribution of a de-sparsified Lasso estimator for high-dimensional regression by computing the Lasso solutions in every iteration of the wild bootstrap. They assess the computational burden as feasible due to the advances in parallel computing.

In this paper, we address the tests' reliability issues outlined above and propose bootstrap analogues to τ and $\tilde{\tau}$, using a wild bootstrap scheme for resampling Lasso solution paths in ADF regressions. The method exploits the computational efficiency of the LARS algorithm to compute the Lasso solution path in every bootstrap iteration, similar to the method of Dezeure et al. [\(2017\)](#page-27-6). Our wild bootstrap algorithm follows Cavaliere et al. [\(2015\)](#page-26-6) and is grounded on the distribution theory, in particular the bootstrap invariance principle, for unconditional heteroskedastic autoregressions of Cavaliere and Taylor [\(2008,](#page-26-1) [2009a\)](#page-26-2). We provide numerical evidence that the wild bootstrap attenuates finite sample size distortions from challenging AR and MA error processes, especially under detrending. Simulations further indicate that—contrary to the unadjusted tests the wild bootstrap permits valid inference when there is non-stationary volatility in the innovations, mirroring the results of Cavaliere and Taylor [\(2008,](#page-26-1) [2009b\)](#page-26-3) for bootstrap variants of established unit root tests.

The remainder of this article is structured as follows. In [Section 2,](#page-3-0) we discuss the theoretical setup and recap the activation knot unit root tests of Arnold and Reinschlüssel [\(2024\)](#page-26-0). [Section 3](#page-6-0) discusses the wild bootstrap algorithm and its implementation. Monte Carlo studies in [Section 4](#page-11-0) investigate finite sample properties of the bootstrap tests. We illustrate applying the methods using real residential property prices for selected OECD countries in [Section 5.](#page-16-0) [Section 6](#page-18-0) concludes and motivates avenues for further research.

We will use the following conventions throughout the manuscript. $\mathbb{I}(\cdot)$ is the indicator function and $\lfloor \cdot \rfloor$ denotes the integer part of its argument. We define $\|\mathbf{x}\|_q$ as the ℓ_q norm of a vector x . Coefficients of the true ADF model have a superscript \star . A superscript $*$ signifies bootstrap quantities conditional on the observed data. $\stackrel{d^*}{\longrightarrow}_p$ means bootstrap weak convergence in probability. Convergence in probability and distribution are denoted by $\stackrel{p}{\to}$ and $\stackrel{d}{\to}$, respectively.

2 Setup and Adaptive Activation Knot Unit Root Tests

2.1 Setup

We consider time series y_t generated by an AR data-generating process (DGP)

$$
y_t = \mathbf{z}'_t \boldsymbol{\theta} + x_t, \qquad x_t = \varrho x_{t-1} + u_t, \qquad t = 0, \dots, T,
$$
 (1)

where $\boldsymbol{z}_t := (1, t, \dots, t^m)'$ is an m^{th} order deterministic component with coefficient vector *θ* and x_t is an AR(1) processes with errors u_t . The stochastic component x_t satisfies $\varrho \in (-1, 1],$ i.e., y_t is stationary when $|\varrho| < 1$ and has a unit root when $\rho = 1$. We make the following assumptions on the error term u_t , cf. Assumption $\mathcal A$ in Cavaliere and Taylor [\(2008\)](#page-26-1).

Assumption 1 (Linear process errors)**.**

- 1. $u_t = \phi(L)\varepsilon_t$ with $\varepsilon_t := \sigma_t \varepsilon_t$. The lag polynomial $\phi(L)$ satisfies $\phi(z) \neq 0$ for all $|z| \leq 1$ *and* $\sum_{j=1}^{\infty} j |\phi_j| < \infty$ *.*
- 2. ϵ_t *is a martingale difference sequence (MDS)* w.r.t. the sigma algebra $\mathcal{F}_t := \{\epsilon_s, s \leq t\}$ *such that* $E(\epsilon_t^2) = \sigma_{\epsilon}^2 = 1$, $T^{-1} \sum_t \epsilon_t^2$ $\frac{p}{\rightarrow} \sigma_{\epsilon}^2$ and $\mathbf{E}|\epsilon_t|^r < K_r$ with $r \geq 4$ and some $K_r < \infty$ *, for all t.*

Assumption 2 (Unconditional homoskedasticity)**.** *The unconditional volatility function* σ_t *satisfies* $\sigma_t = \sigma \in (0, \infty) \ \forall t$ *.*

[Assumption 1](#page-3-1) comprises a set of standard conditions (see Chang & Park, [2002,](#page-26-7) Assumption 1), ensuring that u_t is a weakly-stationary and invertible moving average (MA) process in the ε_t with finite fourth-order moments. Together with [Assumption 2,](#page-3-2) which requires the error process u_t to be unconditionally homoskedastic, [Assumption 1](#page-3-1) imposes the same conditions on *u^t* as we do in Assumption 1 of Arnold and Reinschlüssel [\(2024\)](#page-26-0). We note that the MDS condition in part [2](#page-3-3) allows for conditionally heteroskedastic ε_t , e.g., weakly stationary generalised autoregressive conditionally heteroskedastic (GARCH) and Markov-switching processes, but excludes unconditionally heteroskedastic error processes. The latter is allowed for by the following generalisation of the volatility function in [Assumption 2.](#page-3-2)

Assumption 2' (Unconditional heteroskedasticity). *The unconditional volatility function* σ_t *satisfies* $\sigma_{\lfloor rT \rfloor} = \omega(r)$ *,* $r \in [0,1]$ *with* $\omega(r)$ *a non-stochastic càdlàg function such that* $0 < \sigma_t < \infty$ $\forall t$ *.*

[Assumption 2'](#page-3-4) is a relaxation of [Assumption 2](#page-3-2) and allows σ_t to exhibit non-stochastic time-varying volatility of a quite general form, including (a countable number of) abrupt jumps, polynomial trends and smooth transitions in the unconditional variance. It also generalises the setup of Arnold and Reinschlüssel [\(2024\)](#page-26-0) towards a broader class of empirical processes relevant to macroeconomic applications. We refer to Remarks 1 and 2 in Cavaliere and Taylor [\(2008\)](#page-26-1) and Section 2 in Cavaliere and Taylor [\(2009a\)](#page-26-2) for a discussion of heteroskedastic error processes covered by allowing for MDS innovations ϵ_t as in part [2](#page-3-3) of [Assumption 1](#page-3-1) and deterministic non-stationary volatility implied by [Assumption 2'.](#page-3-4)

Next, we will review the adaptive Lasso activation knot tests of Arnold and Reinschlüssel [\(2024\)](#page-26-0).

2.2 Activation Knot Unit Root Tests

Based on the ADF(∞) representation of [\(1\)](#page-3-5) if $z_t' \theta = 0$,

$$
\Delta y_t = \rho^* y_{t-1} + \sum_{j=1}^{\infty} \delta_j^* \Delta y_{t-j} + \varepsilon_t,
$$
\n(2)

with $\rho^* \in (-2,0], \sum_{j=1}^{\infty} \delta_j^* < 1$, we consider the approximating ADF(*p*) regression model

$$
\Delta y_t = \rho_p y_{t-1} + \sum_{j=1}^p \delta_{p,j} \Delta y_{t-j} + \varepsilon_{p,t}.
$$
\n(3)

The AR lag order *p* in the model [\(3\)](#page-4-0) meets the following assumption.

Assumption 3 (Lag order). *p satisfies* $p = o(T^{1/3})$ *and* $p \to \infty$ *as* $T \to \infty$.

In Arnold and Reinschlüssel [\(2024\)](#page-26-0), we investigate testing for a unit root using the adaptive Lasso solution path to [\(3\)](#page-4-0). Consider the adaptively penalised loss function

$$
\Psi_T(\dot{\rho}, \dot{\delta} | \lambda) := \sum_t \left(\Delta y_t - \dot{\rho} y_{t-1} - \sum_{j=1}^p \dot{\delta}_j \Delta y_{t-j} \right)^2 + 2\lambda \left(w_1^{\gamma_1} |\dot{\rho}| + \sum_{j=1}^p w_{2,j}^{\gamma_2} |\dot{\delta}_j| \right), \qquad (4)
$$

with adaptive weights $w_1 := 1/|\hat{\rho}|, w_{2,j} := 1/\left|\hat{\delta}_j\right|$ determined by the initial OLS estimates

 $\hat{\rho}$ and $\hat{\delta}_j$ in [\(3\)](#page-4-0), and adjustment parameters $\gamma_1, \gamma_2 \in \mathbb{R}^+$. The adaptive Lasso solution path to (4) ,

$$
\mathcal{L} := \left\{ (\widehat{\rho}_{\lambda}, \widehat{\boldsymbol{\delta}}'_{\lambda})' \middle| (\widehat{\rho}_{\lambda}, \widehat{\boldsymbol{\delta}}'_{\lambda})' := \arg\min_{\dot{\rho}, \dot{\delta}} \Psi_T(\dot{\rho}, \dot{\boldsymbol{\delta}} | \lambda), \ \ \lambda \in \mathbb{R}^+ \right\},\tag{5}
$$

is a collection of solutions $\hat{\beta}_{\lambda} := (\hat{\rho}_{\lambda}, \hat{\delta}'_{\lambda})'$ subject to the ℓ_1 penalty parameter λ . A solution path $\mathcal L$ is characterised by knots at which variables are activated or deactivated. Our testing principle leverages that a unit root causes the activation knots of y_{t-1} to be of different stochastic order than under stationarity. We propose a right-sided test of $H_0: \rho^* = 0$ against $H_1: \rho^* \in (-2,0)$ via the statistic $\tau_{\gamma_1} := T^{\gamma_1-1}\lambda_{0,\rho^*}$ with λ_{0,ρ^*} , the first activation knot of y_{t-1} , satisfying

$$
\lambda_{0,\,\rho^{\star}} = w_1^{-\gamma_1} \left| \sum_t y_{t-1} \left(\Delta y_t - \sum_{j=1}^p \widehat{\delta}_{\lambda,\,j} \Delta y_{t-j} \right) \right|.
$$

The knot λ_{0,ρ^*} is a standard output of algorithms for calculating [\(5\)](#page-5-0) such as LARS (Efron et al., [2004\)](#page-27-7), making τ_{γ_1} straightforward to compute.¹

Given consistency of the estimator $\hat{\beta}_{\lambda}$ at $\lambda = \lambda_{0, \rho^* \sim c/T}$, the activation knot of y_{t-1} under local-to-unity roots with non-centrality parameter $c \in (-\infty, 0]$, the (local) distribution of τ_{γ_1} can be identified. Specifically, for $\rho^* = 1 + c/T$ and $\gamma_1 > 1/2$, $\gamma_2 > 0$, Theorem 1 in Arnold and Reinschlüssel [\(2024\)](#page-26-0) states that for unconditionally homoskedastic *u^t* satisfying [Assumption 1,](#page-3-1)

$$
\tau_{\gamma_1} \stackrel{d}{\to} \mathcal{T}_{c,\gamma_1} := \left| \phi(1)^{-1} \frac{W_c(1)^2 - 1}{2 \int_0^1 W_c(r)^2 dr} \right|^{\gamma_1} \left| \frac{1}{2} \sigma^2 \phi(1) \left(W_c(1)^2 - 1 \right) \right|, \tag{6}
$$

where $W_c(r) := \int_0^r \exp[c(r-s)]dW(s)$ is an Ornstein-Uhlenbeck process driven by the standard Wiener process $W(s)$ on $s \in [0,1]$.

A modification of τ_{γ_1} proposed in Arnold and Reinschlüssel [\(2024\)](#page-26-0) derives from enhancing the penalty weight for y_{t-1} with additional information on whether $\rho^* = 0$. This information enrichment of *w*₁ proceeds as $\check{w}_1 := w_1 \cdot J_\alpha$, using the statistic J_α (see Herwartz & Siedenburg, [2010\)](#page-27-8) which exploits different stochastic orders of the OLS estimator in time series regressions when the degree of integration differs. Analogous to τ_{γ_1} , the modified statistic is calculated as $\tilde{\tau}_{\gamma_1} := T^{\gamma_1 - 1} \tilde{\lambda}_{0, \rho^*}$, where $\tilde{\lambda}_{0, \rho^*}$ denotes the first

¹We refer to Section 3.1 in Arnold and Reinschlüssel [\(2024\)](#page-26-0) for the formal definition of an activation knot and a discussion of the stochastic properties of λ_{0, ρ^*} in particular.

activation knot of y_{t-1} on a solution path $\check{\mathcal{L}}$ based on \check{w}_1 .² By Corollary 1 of Arnold and Reinschlüssel [\(2024\)](#page-26-0),

$$
\breve{\tau}_{\gamma_1} \stackrel{d}{\to} \breve{\mathcal{T}}_{c,\,\gamma_1} := \mathcal{T}_{c,\,\gamma_1} / J_{\alpha,\,c}^{\gamma_1},\tag{7}
$$

where $J_{\alpha, c}$ is the *c*-dependent limit of J_{α} .

For implementation we propose the natural choice $\gamma_1 = \gamma_2 = 1$ which avoids an adjustment for $\phi(1) \neq 1$ and yields tests with limit distributions

$$
\tau := \tau_{\gamma_1=1} / \hat{\sigma}^2 \xrightarrow{d} \frac{(W_c(1)^2 - 1)^2}{4 \int_0^1 W_c(r)^2 dr},
$$
\n
$$
\breve{\tau} := \breve{\tau}_{\gamma_1=1} / \hat{\sigma}^2 \xrightarrow{d} \frac{1}{J_{\alpha,c}} \frac{(W_c(1)^2 - 1)^2}{4 \int_0^1 W_c(r)^2 dr},
$$
\n(8)

where $\hat{\sigma}^2$ estimates the error variance based on OLS residuals from the penalty weights ADF regression [\(3\)](#page-4-0).

To accommodate for $z_t \neq 0$, standard detrending ideas can be applied before computing the Lasso solution. In the remainder, we follow Arnold and Reinschlüssel [\(2024\)](#page-26-0) and consider first-difference (FD) detrending (Schmidt & Phillips, [1992\)](#page-29-3) before computing the Lasso solutions and calculating J_α on OLS-adjusted data. Under detrending, the $W_c(r)$ in [\(8\)](#page-6-1) are replaced by the corresponding projection, resulting in limit distributions deviating from the case without adjustment for deterministic components. The *τ*˘ limit is further affected by the distribution of $J_{\alpha,c}$, which is OLS-adjusted. Critical values for *τ* and *τ*˘ under FD adjustment for deterministic components are reported in Table D1 in Arnold and Reinschlüssel [\(2024\)](#page-26-0).

In the next section, we summarise the implications of [Assumption 2'](#page-3-4) and discuss a wild bootstrap correction that addresses the ramifications for the activation knot tests.

3 Wild Bootstrap Activation Knot Tests

While Arnold and Reinschlüssel [\(2024\)](#page-26-0) find that τ and $\tilde{\tau}$ have mostly good size, simulation studies indicate some downward or upward size distortions in small samples for AR error processes with high autocorrelation or processes with MA roots close to −1. These distortions are somewhat more pronounced under detrending.

As we demonstrate in [Section 4,](#page-11-0) heteroskedastic error processes may amplify these

²See Algorithm 1 in Arnold and Reinschlüssel [\(2024\)](#page-26-0) for details on the computation of J_α .

undesirable finite sample properties. Unconditional heteroskedasticity invalidates inference using the homoskedastic null distributions in [\(8\)](#page-6-1). Cavaliere and Taylor [\(2007\)](#page-26-8) show that non-stationary volatility as permitted under [Assumption 2'](#page-3-4) alters the limit distributions of common (unpenalised) regression-based unit root tests in [\(3\)](#page-4-0) as non-stationary volatility introduces nuisance parameters to the tests' homoskedastic distributions in [\(8\)](#page-6-1) that do not vanish asymptotically, invalidating the critical values based on the limits for *c* = 0. For DGP [\(1\)](#page-3-5) in the local-to-unity case $\rho^* = 1 + c/T$ with $-\infty < c \le 0$, this can be traced back to the invariance principle

$$
T^{-1/2}y_{\lfloor rT \rfloor} \stackrel{d}{\to} \overline{\omega}\phi(1)W_{c,\eta}(r), \quad r \in [0,1], \tag{9}
$$

depending on the volatility function $\omega(\cdot)$, cf. the discussion of Theorem 1 in Cavaliere and Taylor [\(2007\)](#page-26-8). Here, $\overline{\omega}^2 := \int_0^1 \omega(s)^2 ds$ is the limit of $T^{-1} \sum_{t=1}^T \sigma_t^2$ and

$$
W_{c,\eta}(r) := \int_0^r \exp[c(r-s)] \, \mathrm{d}W_{\eta}(s), \quad r \in [0,1], \tag{10}
$$

is a diffusion process driven by a time-transformed Wiener process $W_{\eta}(\cdot) := W(\eta(\cdot))$ with directing process $\eta(\cdot)$. A key quantity is the functional

$$
\eta(r) := \overline{\omega}^{-2} \int_0^r \omega(s)^2 \mathrm{d}s, \quad r \in [0, 1], \tag{11}
$$

the so-called *variance profile* of the series, cf. Section 3 in Cavaliere and Taylor [\(2007\)](#page-26-8). A non-constant unconditional volatility function $\omega(\cdot)$ thus alters the tests' (asymptotic) null distributions and local power functions, invalidating the local asymptotic results for *τ* and $\breve{\tau}$ in [\(8\)](#page-6-1). Under constant volatility, $\overline{\omega}^2 = \sigma^2 \in (0, \infty)$. Therefore, $\eta(s) = s, s \in$ $[0,1]$ so that $W_n(\cdot) = W(\cdot)$ and the limiting r.v. in [\(9\)](#page-7-0) reduces to the same (scaled) Ornstein-Uhlenbeck process underlying the limiting functionals in [\(8\)](#page-6-1).

The wild bootstrap proposed in Cavaliere and Taylor [\(2008\)](#page-26-1) samples bootstrap innovations as $\varepsilon_t^* := \xi_t \cdot \breve{\varepsilon}_{p,t}^d$ with $\breve{\varepsilon}_{p,t}^d$ the OLS residuals from the ADF regression [\(3\)](#page-4-0) based on the detrended data and the ξ_t are i.i.d. with $E(\xi_t) = 0$ and $Var(\xi_t) = 1$. The resampled data y_t^* generated by the partial sum process

$$
y_t^* = y_0^* + u_t^*,
$$
 $u_t^* = \sum_{i=1}^t \varepsilon_i^*,$ $t = 1, ..., T,$

initialised at $y_0^* = 0$, then satisfy the bootstrap invariance principle

$$
T^{-1/2}y_{\lfloor rT \rfloor}^* \xrightarrow{d^*} \overline{\omega}W_{0,\eta}(r), \quad r \in [0,1], \tag{12}
$$

cf. Eq. (4) in the proof of Theorem 2 of Cavaliere and Taylor [\(2008\)](#page-26-1).

Since the ξ_t are i.i.d., the device $\varepsilon_t^* = \xi_t \cdot \xi_{q,t}$ anihilates any serial correlation from the original shocks. While this does not impact the asymptotic validity of the bootstrap tests, as follows from the exposition in Cavaliere and Taylor [\(2008,](#page-26-1) [2009a\)](#page-26-2), neglecting correlation in the original shocks may reduce the finite-*T* precision of the bootstrap tests. We thus follow Cavaliere and Taylor [\(2009a\)](#page-26-2) and Smeekes and Taylor [\(2012\)](#page-29-4) and build higher-order stationary dynamics estimated from the data into the bootstrap errors,

$$
u_t^* = \sum_{j=1}^q \check{\delta}_{q,j} u_{t-j}^* + \varepsilon_t^*, \quad t = 1, \dots, T,
$$
\n(13)

and construct the bootstrap sample as

$$
y_t^* := y_0^* + \sum_{i=1}^t u_i^*, \quad 0, \ldots, T.
$$

The $\check{\delta}_{q,j}, j = 1, \ldots, q$ are estimated coefficients of the Δy_{t-j} in a sieve ADF regression [\(3\)](#page-4-0) with lag order *q*, similarly as in the residual-based bootstrap schemes of Ferretti and Romo [\(1996\)](#page-27-0), Chang and Park [\(2003\)](#page-28-1) and Park (2003). Setting $q = p$, the recolouring recursion [\(13\)](#page-8-0) ensures that³

$$
T^{-1/2}y_{\lfloor rT \rfloor}^* = T^{-1/2} \sum_{t=1}^{\lfloor rT \rfloor} u_t^* \xrightarrow{d^*} \overline{\omega} \phi(1) W_{0,\eta}(r), \quad r \in [0,1]. \tag{14}
$$

[Eq. \(14\)](#page-8-1) states that the bootstrap correctly replicates effects from stationary serial corre-lation, cf. [\(9\)](#page-7-0), and thus may better reproduce nuisance terms from adjusting for $\phi(1) \neq 1$ in the finite-*T* null distributions of τ and $\tilde{\tau}$. The invariance principles [\(12\)](#page-8-2) and [\(14\)](#page-8-1), and continuous mapping arguments are the theoretical foundation for applying the (sieve) wild bootstrap to the activation knot tests τ and $\tilde{\tau}$ for improving finite sample precision and valid asymptotic inference in the present heteroskedastic framework.

We next present the wild bootstrap algorithm, which adapts the wild bootstrap algorithms for the popular ADF and *M* tests proposed by Cavaliere and Taylor [\(2008,](#page-26-1)

³Other choices for *q* are discussed in [Remark 1.](#page-10-0) A proof of [\(14\)](#page-8-1) under Assumptions equivalent to ours is given in the proof of Theorem 2 of Smeekes and Taylor [\(2012\)](#page-29-4).

[2009a\)](#page-26-2) and Cavaliere et al. [\(2015\)](#page-26-6) to our activation knot tests.

Algorithm 1 (Wild bootstrap activation knot test)**.**

- 1. Adjust $\{y_t\}_{t=0}^T$ for the determinisitic component $z_t' \theta$ using FD detrending. Denote the resulting series $\{y_t^d\}_{t=0}^T$.
- 2. Select a lag order *q* (see [Remark 1\)](#page-10-0) and obtain the residuals $\{\xi_{q,t}^d\}_{t=1}^T$, where

$$
\check{\varepsilon}_{q,t}^d := \Delta y_t^d - \check{\rho}_q y_{t-1}^d - \sum_{j=1}^q \check{\delta}_{q,j} \Delta y_{t-j},\tag{15}
$$

using the estimates $\check{\rho}_q, \check{\delta}_{q,1}, \ldots, \check{\delta}_{q,q}$, defining $(y_{-1}^d, \ldots, y_{-q}^d)' := \mathbf{0}$. Calculate the adaptive Lasso solution path for the model underlying [\(15\)](#page-9-0) up to $\lambda = \lambda_{0,\rho^*}$, the *first* activation knot of y_{t-1} , and compute τ_{γ_1} .

- 3. Generate wild bootstrap innovations $\{\varepsilon_t^*\}_{t=1}^T$ according to the device $\varepsilon_t^* := \xi_t \cdot \check{\varepsilon}_{q,t}^d$, where the r.v.s ξ_t are i.i.d. and satisfy $E(\xi_t) = 0$ and $Var(\xi_t) = 1$.
- 4. Build the bootstrap error process $\{u_t^*\}_{t=1}^T$ using the recolouring recursion

$$
u_t^* = \sum_{j=1}^q \check{\delta}_{q,j} u_{t-j}^* + \varepsilon_t^*,
$$
\n(16)

with $(u_0^*, \ldots, u_{1-q}^*)' := \mathbf{0}$. Build a bootstrap time series $\{y_t^*\}_{t=0}^T$ via the partial sum process

$$
y_t^* := y_0^* + \sum_{i=1}^t u_i^*,
$$

with $y_0^* = 0$.

5. Adjust the bootstrap sample $\{y_t^*\}_{t=0}^T$ as in Step 1 and compute the bootstrap adaptive Lasso solution path for an ADF regression with lag order p^* (cf. [Remark 1\)](#page-10-0) up to $\lambda = \lambda_{0,\,\rho^*}^*$, with

$$
\lambda_{0,\,\rho^*}^* = \left(\frac{1}{w_1^*}\right)^{\gamma_1} \left| \sum_t y_{t-1}^{*d} \left(\Delta y_t^{*d} - \sum_{j=1}^{p^*} \widehat{\delta}_{\lambda,\,j}^* \Delta y_{t-j}^{*d} \right) \right|,
$$

the *first* activation knot of y_{t-1}^{*d} , and compute the bootstrap test statistic $\tau_{\gamma_1,b}^*$.

6. Obtain bootstrap test statistics $\{\tau_{\gamma_1,b}^*\}_{b=1}^B$ by completing steps 3 to 5 *B* times. Calculate the bootstrap level- α critical value as

$$
CV^*(\alpha) := \max\left\{x : \frac{1}{B} \sum_{b=1}^B \mathbb{I}\left(x < \tau_{\gamma_1, b}^*\right) \le \alpha\right\}.\tag{17}
$$

Reject the unit root null at level α if $CV^*(\alpha) \leq \tau_{\gamma_1}$ with τ_{γ_1} the test statistic computed for lag order p using the detrended data of Step 1.

Remark 1. The lag orders p , q , and p^* must be selected to implement [Algorithm 1.](#page-9-1) Modified information criteria such as the MAIC (Ng $\&$ Perron, [2001\)](#page-28-0) are established procedures for this purpose. The RSMAIC of Cavaliere et al. [\(2015\)](#page-26-6) is a heteroscedasticityrobust variant of the MAIC, which estimates the lag order based on a rescaling of the y_t with a non-parametric estimate of its (assumed sufficiently smooth) variance profile, as suggested by Beare [\(2018\)](#page-26-9). Since the power advantage of using the RSMAIC over the MAIC for sieve wild bootstrap ADF tests under non-stationary volatility is well documented by the simulation results in Cavaliere et al. [\(2015\)](#page-26-6), with negligible effects under homoskedasticity, we apply the RSMAIC throughout.

Although q (and p^*) are not required to diverge for the asymptotic validity of the boot-strap (cf. Cavaliere et al., [2015\)](#page-26-6), selecting p and p^* by the RSMAIC and setting $q = p$ is a convenient choice. Notably, selecting p^* independently from p and q has been documented to help control upward size distortions of sieve bootstrap ADF tests under errors with a large negative $MA(1)$ component, where information criteria yield underspecified models (Richard, [2009\)](#page-29-5).

Remark 2. Obtaining the residuals $\check{\epsilon}_{q,t}$ from the sieve regression underlying [\(15\)](#page-9-0) using OLS is convenient since we require estimates of $(\rho_p, \delta'_p)'$ for computing the adaptive penalty weights of the Lasso estimator anyhow. Cavaliere and Taylor [\(2009a\)](#page-26-2) suggest other asymptotically equivalent strategies to compute the residuals, which we do not consider here for brevity.

If OLS estimation is infeasible (e.g., due to collinearity or $p \geq T$) but zero-consistent initial coefficient estimates⁴ (Huang et al., [2008\)](#page-28-7) are available, ad-hoc estimates for $\check{\varepsilon}_t$ can be obtained by running AL or ALIE and setting $\lambda = \lambda_{\alpha}^{\tau}$ with λ_{α}^{τ} a (upper-tail) quantile of the null distribution of τ . This approach resembles that of Chernozhukov et al. [\(2023\)](#page-27-9) and yields consistent coefficient estimates from conservative model selection, as for AIC-

⁴Zero-consistency ensures the penalty weights to be bounded for relevant variables and to converge to infinity for irrelevant variables. Zero-consistency is the weakest requirement for establishing the oracle property in the literature on the adaptive Lasso to our knowledge.

tuned estimation. However, finite-*T* adaptive Lasso estimates are usually biased and require recentering of the residuals (cf. Chatterjee & Lahiri, [2011\)](#page-27-3). Therefore, we reckon using OLS residuals is more convenient for practitioners if feasible.

Remark 3. The literature on the wild bootstrap (cf. Davidson & Flachaire, [2008;](#page-27-10) Liu, [1988\)](#page-28-3) features several proposals on how to sample the ξ_t in generating the bootstrap errors ε_t^* in step 3. An example is the asymmetric two-point distribution by Mammen [\(1993\)](#page-28-8), designed for higher precision of bootstrap tests under heteroskedastic and non-Gaussian errors. In other applications, e.g., the pooled panel unit root tests of Herwartz and Walle [\(2018\)](#page-28-9), the Rademacher distribution (Davidson & Flachaire, [2008\)](#page-27-10) has been reported to yield better precision. However, and consistent with the results for various wild bootstrap time series unit root tests reported in Cavaliere and Taylor [\(2008,](#page-26-1) [2009b\)](#page-26-3) and Demetrescu and Hanck [\(2016\)](#page-27-11), we find Gaussian ξ_t to yield good performance, with no significant discrepancies to using other candidate distributions. We therefore report results only for Gaussian *ξ^t* . Simulation results for resampling with the Rademacher and Mammen's distribution are available on request.

4 Monte Carlo Evidence

To investigate sample properties of the tests, we generate time series as

$$
y_t = \varrho y_{t-1} + v_t, \quad t = 0, 1, \dots, T,
$$
\n(18)

with starting value $y_0 = 0$ for sample sizes $T \in \{75, 100, 150, 250, 500, 1000\}$. We let $\rho = 1 + c/T$ so that setting $c = 0$ obtains data under the unit root null. As (local) stationary alternatives, we set $c = -7$ when testing based on data adjusted for a constant and $c = -13.5$ for detrending. Pure AR or MA errors are generated with the recursion

$$
v_t = \varphi v_{t-1} + \vartheta \epsilon_{t-1} + \sigma_{T,t} \cdot \epsilon_t, \quad \epsilon_t \stackrel{i.i.d.}{\sim} N(0,1), \tag{19}
$$

with coefficients $\varphi, \vartheta \in \{-8, -4, 0, 4, 8\}.$

Following Cavaliere and Taylor [\(2009b\)](#page-26-3), we model deterministic smooth transitions in the unconditional volatility parameter $\sigma_{T,t}$ between two regimes with variances $s_1^2, s_2^2 > 0$ via a logistic function $\mathbb{S}_{T,t}$,

$$
\sigma_{T,t}^2 := s_1^2 + (s_2^2 - s_1^2) \cdot \mathbb{S}_{T,t}, \quad \mathbb{S}_{T,t} := (1 + \exp(-\tilde{\gamma}_T(t - \lfloor \kappa T \rfloor)))^{-1}, \tag{20}
$$

with transition midpoint $\lbrack \kappa T \rbrack$, $\kappa \in (0,1)$. The parameter $\tilde{\gamma}_T$ determines the transition speed between s_1^2 and s_2^2 , yielding and abrupt regime switch at $t = \lfloor \kappa T \rfloor$ in the sense that $\mathbb{S}_{T,t} \to \mathbb{I}(t \geq \lfloor \kappa T \rfloor)$ as $\tilde{\gamma}_T \to \infty$. We adapt the local drift $\tilde{\gamma}_T := 25/T$ from Cavaliere and Taylor [\(2009b\)](#page-26-3). Setting $s_1^2 = 1$ throughout, we model negative shifts with early transition midpoints ($\kappa = .2, s_2^2 = .25$) as well as positive shifts with late midpoints $(\kappa = .8, s_2^2 = 4).$

The bootstrap variants of τ and $\tilde{\tau}$, denoted τ^* and $\tilde{\tau}^*$, are implemented as detailed in [Section 3](#page-6-0) at the 5% level and computed with $B = 499$ bootstrap replications. The *ξt* in Step 3 of [Algorithm 1](#page-9-1) are standard Gaussian r.v.s, and the lag orders *p, q* and *p* ∗ are estimated using the RSMAIC, if not indicated otherwise. We compute the Lasso solution paths using the implementation of the LARS algorithm (Efron et al., [2004\)](#page-27-7) in the R (2023) package lars (Hastie & Efron, [2022\)](#page-27-12).

We first examine the ability of the wild bootstrap to approximate the tests' finite sample distributions and assess the bootstrap tests' precision and local power in the baseline scenario with uncorrelated homoskedastic errors $(s_2^2 = 1)$. The sieve regression lag order q is zero in this experiment. The results presented in [Table 1](#page-13-0) indicate that *τ*[∗] and *τ*[∗] have excellent precision (top panel), improving on the empirical size of *τ* and $\check{\tau}$, which may be somewhat conservative for small *T*. Furthermore, the bootstrap tests seem to replicate the (size-adjusted) local power function of the standard tests quite well, with only minor deviations for small *T* notable for *τ*˘ (bottom panel).

In a second experiment, we examine the precision of the sieve bootstrap tests under homoskedastic correlated errors. [Table 2](#page-14-0) presents the results. The bootstrap tests perform well for AR errors with $\varphi \in (-.8, .8)$ and under MA errors with $\vartheta = .8$. The most challenging scenario is MA errors with $\vartheta = -.8$, which lead to upward size distortions across *T* that are prohibitive for small samples and only decay slowly with the sample size. Although τ^* and $\tilde{\tau}^*$ are oversized, resampling with recolouring yields significant improvements over τ and $\check{\tau}$. Recolouring is particularly helpful under detrending, where the empirical size of the bootstrap tests converges quickly towards the 5% level, undercutting the rejection rates of the unadjusted tests by over a third in larger samples. Furthermore, comparing the size of the SWB tests with the size of the WB tests with *q* = 0 reported in [Tables A1](#page-13-0) and [A2](#page-14-0) of [Appendix A](#page-20-0) corroborates better precision through the sieve step under MA errors in particular.

Next, we investigate the impact of unconditional heteroscedasticity and its interplay with the effects of autocorrelated errors on the precision of the tests. Our focus is MA processes for which τ and $\breve{\tau}$ show the largest size distortions, and bootstrap inference

		$z_t=1$				$z_t = (1, t)'$				
T	τ^*	$\breve{\tau}^*$	τ	$\breve{\tau}$	τ^*	$\breve{\tau}^*$	τ	$\breve{\tau}$		
				$\rho^* = 0$						
75	.049	.046	.037	.036	.047	.047	.040	.035		
100	.055	.056	.045	.044	.047	.047	.041	.037		
150	.051	.055	.043	.043	.049	.048	.041	.038		
250	.055	.052	.049	.043	.045	.047	.041	.040		
500	.056	.054	.050	.049	.052	.053	.047	.045		
1000	.055	.050	.049	.046	.052	.049	.047	.045		
				$\rho^* = -c/T$						
75	.247	.297	.251	.312	.368	.395	.384	.411		
100	.241	.288	.228	.278	.365	.382	.366	.403		
150	.236	.297	.234	.280	.366	.382	.374	.404		
250	.246	.308	.227	.304	.368	.387	.389	.402		
500	.251	.307	.232	.286	.382	.403	.370	.390		
1000	.310	.359	.296	.351	.391	.416	.382	.418		

Table 1: Rejection rates of the WB activation knot tests for i.i.d. errors

Notes: DGP [\(18\)](#page-11-1) with $\sigma_{T,t} = 1 \forall t$. $B = 499$ wild bootstrap replications with $q = 0$ (no recolouring). The data are adjusted for a constant or a linear time trend using the FD method of Schmidt and Phillips [\(1992\)](#page-29-3). The lag orders *p* and p^* are selected using the RSMAIC. Top panel: unit root model. Bottom panel: local alternative with $c = -7$ if $z_t = 1$ and $c = -13.5$ if $z_t = (1, t)'$. Power estimates of τ and $\breve{\tau}$ are size-adjusted at 5%. 5000 Monte Carlo replications.

with better precision is the most needed. [Table 3](#page-15-0) presents the size of the tests for MA errors with non-stationary volatility. Consistent with the theory, we see (upward) size distortions for τ and $\tilde{\tau}$ under smoothly trending variances across all scenarios. These distortions vary in magnitude with the correlation structure, the type of variance shift and the adjustment for *z^t* . The smallest extent occurs for an early variance reduction and MA coefficient $\vartheta = .8$, where both tests mostly remain below 10%. There are differences between τ and $\check{\tau}$, e.g., for a late variance increase with $\vartheta = .8$, which is likely due to idiosyncracies of the J_{α} statistic under heteroskedasticity. As before, we observe the most pronounced distortions at $\vartheta = -0.8$ for small *T*, which are worse than under homoscedasticity, cf. [Table 2.](#page-14-0) The bootstrap analogues consistently perform better, having size close to the 5% level for positive MA coefficients for upward and downward variance shifts and independently of the adjustment for a deterministic component. For $\vartheta = -.8, \tau^*$ and $\breve{\tau}^*$ tend to be somewhat more oversized than under homoscedasticity and also require large *T* for rejection rates to ameliorate. As in the homoskedastic settings, comparing with additional simulation results for the WB tests with $q = 0$ in [Table A2](#page-14-0) indicates better precision of the SWB tests. [Tables A3](#page-15-0) and [A4](#page-17-0) report qualitatively similar results for heteroskedastic AR errors.

		$z_t = 1$				$z_t = (1, t)'$					
$\cal T$	τ^*	$\breve{\tau}^*$	τ	$\breve{\tau}$	τ^*	$\breve{\tau}^*$	τ	$\breve{\tau}$			
	$\varphi = -.8, \vartheta = 0$										
75	$.047$.042	.054	.049	.026	.033	.059	.068			
100	.044	.044	$.051\,$.047	.030	$.035\,$.051	.054			
150	.041	.040	.045	.042	.044	.047	.060	.060			
$250\,$	$.050\,$.050	$.051\,$.049	$.044$	$.046$.050	$.051\,$			
$500\,$	$.052\,$.049	$.051\,$	$.047$	$.048\,$	$.052\,$.050	.051			
1000	.050	.052	$.051\,$	$.050\,$.047	.047	.048	.045			
	$\varphi = .8, \vartheta = 0$										
75	$.051\,$	$.051\,$.056	.040	$.054\,$.052	.045	$.036\,$			
100	$.047\,$.049	$.047$	$.038\,$	$.046\,$.046	.038	.032			
150	.054	.050	$.052\,$.040	$.046$	$.048\,$.041	.033			
250	.049	.047	$.046\,$.039	.049	.047	.046	.036			
500	$.052\,$.052	$.050\,$.045	.049	$.048\,$.043	.042			
1000	$.053\,$.054	$.053\,$.048	.053	.055	.049	.046			
				$\varphi = 0, \vartheta = -.8$							
$75\,$.169	.173	$.316\,$.325	.277	$.307\,$.500	$.530\,$			
100	$.128\,$.136	$.287\,$.288	.190	$.225\,$.446	.475			
150	.099	.099	.253	$.251\,$	$.124\,$	$.151\,$.400	.402			
250	$.087\,$	$.085\,$.216	.206	$.082\,$	$.091\,$.342	.327			
500	.079	.078	.171	.169	$.069$.074	.287	.259			
1000	.079	.075	.135	.136	.080	.080	.238	.211			
				$\varphi = 0, \vartheta = .8$							
75	.042	.040	.048	.043	.033	.037	.043	.042			
100	.043	$.038\,$	$.051\,$	$.039\,$	$.033\,$	$.037\,$.045	.046			
150	.043	$.043$	$.045$.045	$.036\,$	$.034\,$	$.042$.046			
250	.045	.044	$.049$.049	$.043$	$.044$	$.051$.054			
500	$.055\,$	$.055\,$	$.058\,$	$.056\,$	$.043\,$.044	.047	.050			
1000	.049	$.051\,$	$.050\,$	$.048\,$.050	$.046$	$.052\,$.049			

Table 2: Size of the SWB tests under correlated homoskedastic errors

Notes: DGP [\(18\)](#page-11-1) with $c = 0$ and $\sigma_t = 1 \forall$. The data are adjusted for a constant $(z_t = 1)$ or a linear time trend $(z_t = (1, t)')$ using the FD method of Schmidt and Phillips [\(1992\)](#page-29-3). All lag orders are computed using the RSMAIC. $B = 499$ sieve wild bootstrap replications and 5000 Monte Carlo replications.

		$z_t = 1$			$z_t = (1, t)'$				
T	τ^*	$\breve{\tau}^*$	τ	$\breve{\tau}$	τ^*	$\breve{\tau}^*$	τ	$\breve{\tau}$	
					Early smooth variance reduction, $\varphi = 0, \vartheta = 0$				
$75\,$.060	.061	.067	.079	.056	.058	.081	.079	
100	$.058\,$	$.062\,$	$.063\,$.073	$.052\,$	$.054\,$	$.078\,$.076	
150	$.053\,$	$.056\,$	$.064\,$	$.075\,$	$.053\,$	$.055\,$	$.081\,$.082	
$250\,$	$.050$	$.052\,$	$.063\,$	$.075\,$	$.056\,$	$.054\,$	$.087\,$.086	
500	.049	.046	$.064\,$	$.073\,$	$.053\,$	$.050\,$	$.087\,$.086	
1000	$.048$	$.048$	$.063\,$	$.075\,$	$.053\,$.056	$.083\,$.092	
					Early smooth variance reduction, $\varphi = 0, \vartheta = -.8$				
$75\,$	$.150\,$	$.185\,$.269	.330	$.273\,$	$.338\,$	$.511\,$.605	
100	$.125\,$	$.144\,$	$.255\,$	$.298\,$	$.201\,$	$.261\,$	$.473\,$.561	
150	$.095\,$	$.110\,$	$.219\,$	$.258\,$	$.134\,$	$.175\,$	$.422\,$.483	
$250\,$	$.087\,$	$.090\,$	$.219\,$	$.233\,$	$.094\,$	$.117\,$	$.370\,$.399	
500	$.074\,$	$.072\,$	$.187\,$	$.204\,$	$.084\,$	$.088\,$	$.337\,$	$.334\,$	
1000	$.083\,$	$.076\,$	$.164\,$	$.171\,$	$.076\,$	$.073\,$	$.281\,$.270	
					Early smooth variance reduction, $\varphi = 0, \vartheta = .8$				
$75\,$	$.059\,$.064	.082	$.072\,$	$.056\,$.057	$.108\,$.089	
100	$.047$	$.057\,$	$.075\,$	$.072\,$	$.053\,$	$.060\,$	$.110$.095	
150	$.051\,$	$.058\,$	$.072\,$	$.077\,$	$.048\,$	$.053\,$	$.095\,$.092	
$250\,$	$.054\,$	$.058\,$	$.072\,$	$.082\,$.050	.051	$.094\,$.097	
500	$.051\,$	$.055\,$	$.072\,$	$.088\,$	$.052\,$	$.053\,$	$.098\,$	$.107$	
1000	$.056\,$	$.056\,$	$.068$	$.084\,$	$.047$	$.049\,$	$.085\,$.098	
					Late smooth variance increase, $\varphi = 0, \vartheta = 0$				
$75\,$	$.064\,$	$.056\,$	$.136\,$	$.108\,$	$.055\,$	$.058\,$	$.073\,$.077	
100	$.062\,$	$.062\,$	$.154\,$	$.129\,$	$.048\,$	$.053\,$	$.066\,$.069	
150	$.056\,$	$.055\,$	$.149\,$	$.119\,$	$.051\,$	$.051\,$	$.077\,$	$.081\,$	
$250\,$	$.055\,$	$.051\,$	$.152\,$	$.121\,$	$.050\,$	$.048\,$	$.078\,$.080	
500	$.055\,$	$.051\,$	$.156\,$	$.123\,$	$.052\,$	$.054\,$	$.086\,$.088	
1000	.047	.049	.148	$.125\,$.050	$.057\,$.090	.096	
					Late smooth variance increase, $\varphi = 0, \vartheta = -.8$				
$75\,$.206	.214	.417	.424	$.190\,$.226	$.311\,$.376	
100	$.162\,$	$.175\,$	$.385\,$.389	$.150\,$	$.185\,$	$.288\,$	$.352\,$	
150	$.132\,$	$.137\,$	$.352\,$.366	$.095\,$	$.130\,$.263	.314	
250	$.108\,$.109	.319	.337	.069	.090	.243	.288	
500	.088	.088	.284	.289	.069	.071	.241	.259	
1000	.080	$.083\,$.248	$.258\,$.066	.066	.227	.241	
					Late smooth variance increase, $\varphi = 0, \vartheta = .8$				
$75\,$	$.064\,$.066	$.157\,$	$.132\,$.047	.051	.055	.087	
100	$.065$	$.062\,$	$.167\,$	$.138\,$	$.047$	$.049$	$.059$.090	
150	.060	$.057$	$.154\,$	$.124\,$	$.051\,$	$.054\,$.067	.098	
$250\,$.066	$.062\,$	$.172\,$	$.136\,$	$.050\,$.049	.069	.096	
500	$.059\,$	$.051\,$	$.156\,$	$.127\,$	$.058\,$.055	$.084\,$	$.106$	
1000	$.054\,$	$.058\,$.166	$.134\,$	$.055\,$	$.055\,$	$.079\,$.095	

Table 3: Size of the SWB activation knot tests under heteroskedastic MA errors

Notes: 5% nominal level. DGP [\(18\)](#page-11-1) with $c = 0$ and $\sigma_{T,t}$ as defined in [\(20\)](#page-11-2), with $\kappa = .2$, $s_2^2 = .25$ for early negative shifts and $\kappa = .8$, $s_2^2 = 4$ for late positive shifts. The data are adjusted for a constant sieve wild bootstrap replications. 5000 Monte Carlo replications.

Local power estimates for the SWB tests under heteroskedastic AR and MA processes are provided in [Tables A5](#page-24-0) and [A6.](#page-25-0) With a few exceptions for error processes with negative coefficients and small T , we find that the bootstrap tests effectively approximate the local power functions of the infeasible size-adjusted tests τ and $\tilde{\tau}$. Furthermore, the simulation results show that information enrichment yields power gains even under nonstationary volatility and that the bootstrap retains these gains.

5 Application to Real Residential Property Prices

Figure 1: RRPPI rates and estimated variance profiles of the top six Euro area economies

Notes: Left: Year-on-year quarterly real residential property price inflation rates of selected OECD countries from Q1-1972 to Q2-2023 ($T = 207$). Right: estimated variance profiles, computed as $\hat{\eta}(s) :=$ tries from Q1-1972 to Q2-2023 ($T = 207$). Right: estimated variance profiles, computed as $\hat{\eta}(s) := \left(\sum_{t=1}^{\lfloor sT \rfloor} \hat{u}_t^2 + (sT - \lfloor sT \rfloor) \hat{u}_{\lfloor sT \rfloor + 1} \right) / \sum_{t=1}^T \hat{u}_t^2$, with \hat{u}_t the residuals from an AR(liere and Taylor [\(2007\)](#page-26-8).

The housing price rally sparked by a sustained low-interest environment after the 2008 financial crisis has received much public and scientific attention (see, e.g., Jordà et al., [2016;](#page-28-11) Mian et al., [2015\)](#page-28-12). While monetary policies fueled price dynamics during the COVID-19 pandemic (cf. Francke & Korevaar, [2021\)](#page-27-13), the recent trend reversal due to surges in mortgage and construction interest rates highlights the importance of understanding the dynamics in the market relevant to macroeconomic policy.

We use the tests to assess real residential property price inflation (RRPPI) rates of the six largest Eurozone economies for stochastic trends, covering 207 quarterly obser-

	BE	DE	ES	FR	IT	$\rm NL$
<i>Entire sample: $Q1-1972 - Q2-2023$ (T = 206)</i>						
\hat{k}_i	12	12	5	8	$\overline{5}$	8
τ $\breve{\tau}$	$.007***$ ${<}001***$.230 $.057^{\ast}$	$.016**$ $.002***$	$.012**$ $.008***$	$< .001***$ $<.001***$	$.002***$ $.003***$
τ^* $\breve{\tau}^*$	$.010***$ $< 001***$.298 .107	$.020**$ $.002***$	$.016**$ $.011**$	$< 0.01***$ $< 0.01***$	$.005***$ $.006***$
<i>Pre-Euro:</i> $Q1-1972 - Q4-1998$ $(T = 108)$						
\hat{k}_i	8	4	$\overline{4}$	4	2	4
τ $\breve{\tau}$	$.028**$ $.020**$.378 .170	.109 $.057*$	$.066*$ $.049**$	$< 0.01***$ $< 0.01***$	$.032**$ $.037**$
τ^* $\breve{\tau}^*$	$.044**$ $.015***$.359 .176	$.086*$ $.052*$.100 $.051*$	$< .001***$ $< .001***$	$.027**$ $.031**$
<i>Euro-era:</i> $Q1-1999 - Q2-2023$ (T = 98)						
\hat{k}_i	$\overline{4}$	8	$\overline{4}$	8	$\overline{4}$	4
τ $\breve{\tau}$.569 .356	.108 $.052*$.243 .194	.386 .290	.269 .213	.615 .477
τ^* $\breve{\tau}^*$.589 .471	.197 .121	.257 .214	.439 .406	.275 .275	.673 .610

Table 4: Test outcomes for European residential property price inflation rates

Notes: quarterly year-on-year RRPPI rates from BIS [\(2023\)](#page-26-10). The tests are computed on FD-demeaned data, and the RSMAIC selected all lag orders with maximum lag order $k_{\text{max}} = \lfloor 12 \cdot (100/T)^{25} \rfloor$. k_i is the estimated truncation lag in model (21). The test outcomes are payalues. The supercripts $*$ $**$ and $***$ indicate a rejecti model [\(21\)](#page-18-1). The test outcomes are p-values. The supercripts *, ** and *** indicate a rejection of the unit root null at 10%, 5% and 1%, respectively. The bootstrap tests *τ* and *τ*˘ are computed based on *B* = 4999 bootstrap replications.

vations from Q1-1972 to Q2-2023 for Belgium, France, Germany, Italy, Spain and the Netherlands. The data (BIS, [2023\)](#page-26-10) obtained from CPI deflated price indices summarise all types of new and existing dwellings, except for Germany, where the underlying index considers owner-occupied houses only.

Besides the abovementioned events, the sample period covers further economic turmoil, such as OPEC oil shortages (1973, 1979) and recessions in the 1980s and 1990s, reflecting different variance regimes. We find evidence of such regimes in data displayed in the left panel of [Figure 1,](#page-16-1) showcasing a high degree of co-movement. The estimated volatility profiles in the right panel further indicate the instability of the unconditional variances, showing considerable downward shifts for Belgium (BE), Italy (IT), Spain (ES), and the Netherlands (NL).

We consider ADF regressions with a non-zero mean,

$$
\Delta \text{RRPPI}_{i,t} = \mu_i + \beta_0 \text{RRPPI}_{i,t-1} + \sum_{j=1}^{k_i} \beta_j \Delta \text{RRPPI}_{i,t-j} + e_{i,t},\tag{21}
$$

where the k_i in the baseline regression, as well as lag orders in [Algorithm 1,](#page-9-1) are selected by RSMAIC. We compute the bootstrap tests using $B = 4999$ bootstrap iterations.

[Table 4](#page-17-0) presents the outcomes of the tests for three different periods: the entire data range (Q1-1972–Q2-2023) and subsamples before (Q1-1972–Q4-1998) and for the Euroera (Q1-1999–Q2-2023) which is characterised by macroeconomic convergence, e.g., due to a mutual primary refinancing interest rate applying to the member economies.

Outcomes for the bootstrap tests mostly agree with the standard inference, giving mixed conclusions. For the entire period, the tests indicate mean reversion behaviour for all economies except Germany, where only *τ*˘ rejects at 5%. The pre-Euro subsample shows similar outcomes. Notably, $\tilde{\tau}^*$ tends to have smaller p-values than τ^* , likely due to power gains from information enrichment—a feature of $\tilde{\tau}$ that the simulation outcomes indicate is preserved by the bootstrap. None of the tests rejects the null hypothesis of a stochastic trend for the Euro-era at 5%. The reduced sample size or possible trend changes in the generating process could contribute to the mixed evidence, especially since periods of exuberance in the housing market characterise the Euro-era subsample.

6 Conclusion

In this paper, we assessed the reliability of the adaptive Lasso solution path-based unit root tests τ and $\tilde{\tau}$ proposed in Arnold and Reinschlüssel [\(2024\)](#page-26-0) under weaker assumptions and considered whether resampling offers robust alternatives. Drawing on the theoretical results in Cavaliere and Taylor [\(2008,](#page-26-1) [2009a\)](#page-26-2), we propose the wild bootstrap analogues τ^{*} and τ^{*}, implementing the computation of resampled Lasso solution paths efficiently using the LARS algorithm (Efron et al., [2004;](#page-27-7) Hastie & Efron, [2022\)](#page-27-12). Numerical evidence shows that the bootstrap yields tests with higher precision, allowing more robust inference under correlated error processes of a general form.

Consistent with the theory on heteroskedastic autoregressions in Cavaliere and Taylor [\(2007\)](#page-26-8), *τ* and *τ*˘ do not attain their homoscedastic limits derived in Arnold and Reinschlüssel [\(2024\)](#page-26-0) when the errors are unconditionally heteroscedastic, so that valid inference is not guaranteed, even asymptotically. Our simulations confirm that τ and *τ*˘—like conventional unit root tests—have null distributions and local power functions affected by nuisance parameters for unconditionally heteroskedastic innovations. A consequence is size distortions, with the strength of the effect depending on the adjustment for deterministic components. Correlated errors, such as MA(1) processes with negative coefficients, exacerbate this effect. Our sieve wild bootstrap tests display higher precision than τ and $\check{\tau}$ in these scenarios, indicating their ability to accurately recover the first-order null distributions. In addition, our simulation results show the wild bootstrap variants to approximate the finite-sample (local) power functions of the infeasible size-adjusted implementations of the standard tests under variance shifts.

To illustrate the bootstrap tests, we consider real residential property price inflation rates for selected Eurozone economies. This data set seems representative of our setup, given signs of persistence and non-stationary volatility. Both bootstrap tests yield the same conclusions, pointing to stationarity for the entire period from 1972 to 2023 and the period before the introduction of the Euro. We find no evidence for mean-reversion for the Euro-era subsample.

There are various avenues for further research. Our simulation evidence suggests that the wild bootstrap preserves the benefits of information enrichment, motivating its application to inference for penalised regression when the (asymptotic) distribution is unknown. Given the positive results for τ and $\tilde{\tau}$, expanding the underlying testing principle to other (adaptively) penalised regression estimators, for example, the fused Lasso or the group Lasso seems worthwhile. Good starting points are Qian and Su [\(2016\)](#page-28-13) and Schweikert [\(2021\)](#page-29-6), which employ adaptive variants of the group Lasso and fused Lasso to detect structural breaks in panel and cointegrating regressions.

Our empirical application raises the question of whether the advent of the Euro affected the heterogeneity in the housing price dynamics across the Eurozone countries. To address this question, contemplating additional measures of heterogeneity for information enrichment can improve the discriminatory power of the aforementioned penalised estimators. This topic is closely related to the heterogeneous treatment effect inference literature, which could inspire further extensions.

Another promising direction is inference in high-dimensional regressions for which the double Lasso of Belloni et al. [\(2013\)](#page-26-11) has become a standard method. It would be appealing to investigate whether information enrichment furthermore improves highdimensional inference for which the wild bootstrap has become a cornerstone (cf. Chernozhukov et al., [2023\)](#page-27-9). To this end, one could apply our testing principle to causal inference problems or multivariate time series, potentially using adaptive penalty weights derived using the de-sparsified Lasso. We are currently investigating this approach.

A Additional Simulation Results

		$z_t = 1$			$z_t = (1, t)'$							
$\cal T$	τ^*	$\breve{\tau}^*$	τ	$\breve{\tau}$	τ^*	$\breve{\tau}^*$	τ	$\breve{\tau}$				
	$\varphi = -.8, \vartheta = 0$											
75	.057	.055	.054	.049	.063	.073	.059	.068				
100	.053	$.055\,$	$.051\,$.047	$.054\,$.060	.051	$.054$				
150	.049	.048	.045	$.042\,$	$.061\,$	$.067$.060	.060				
250	.056	.056	$.051\,$.049	$.053\,$	$.057$.050	.051				
500	.053	.053	.051	.047	.052	$.056\,$.050	.051				
1000	$.055\,$.055	$.051\,$	$.050\,$.051	$.053\,$.048	.045				
				$\varphi = .8, \vartheta = 0$								
75	.060	.042	.056	.040	.047	$.039\,$.045	.036				
100	.052	.043	.047	.038	.040	$.038\,$.038	.032				
150	.059	.045	.052	.040	$.045$	$.039\,$.041	.033				
250	$.050$.044	$.046\,$.039	.047	.043	.046	.036				
500	$.054\,$	$.051\,$	$.050\,$.045	$.045$	$.047$.043	.042				
1000	$.057$.055	.053	.048	.051	.052	.049	.046				
				$\varphi = 0, \vartheta = -.8$								
75	.332	$.342\,$	$.316\,$.325	$.514\,$.549	.500	.530				
100	.296	.302	$.287\,$	$.288\,$	$.457\,$	$.493\,$.446	.475				
150	$.261\,$.264	$.253\,$.251	.410	$.420\,$.400	.402				
250	$.221\,$.217	.216	.206	$.353\,$	$.346\,$	$.342\,$.327				
500	.177	$.178\,$.171	.169	.292	.269	.287	.259				
1000	.137	.140	.135	$.136\,$.245	.221	.238	.211				
				$\varphi = 0, \vartheta = .8$								
75	$.059$	$.052\,$	$.048\,$.043	.048	$.051\,$.043	.042				
100	.062	.049	$.051\,$.039	.050	$.059$.045	.046				
150	.052	$.051$.045	.045	.047	.055	.042	.046				
250	.056	$.057\,$	$.049$.049	.055	$.063\,$.051	.054				
$500\,$.063	.062	.058	$.056\,$	$.052\,$	$.059\,$.047	.050				
1000	.053	$.053\,$.050	$.048\,$	$.055\,$	$.055\,$.052	.049				

Table A1: Size of the WB activation knot tests under correlated homoskedastic errors

Notes: DGP [\(18\)](#page-11-1) with $c = 0$ and $\sigma_{T,t}$ as defined in [\(20\)](#page-11-2), with $\kappa = .2$, $s_2^2 = .25$ for early negative shifts and $\kappa = .8$, $s_2^2 = 4$ for late positive shifts. The data are adjusted for a constant $(z_t = 1)$ or a lin the FD method of Schmidt and Phillips [\(1992\)](#page-29-3). All lag orders are computed using the RSMAIC. *B* = 499 wild bootstrap replications with $q = 0$. 5000 Monte Carlo replications.

		$z_t=1$			$z_t = (1, t)'$				
$\cal T$	τ^*	$\breve{\tau}^*$	τ	$\breve{\tau}$	τ^*	$\breve{\tau}^*$	τ	$\breve{\tau}$	
					Early smooth variance reduction, $\varphi = 0, \vartheta = 0$				
75	.065	.064	.067	$.079\,$.059	.059	$.081\,$	$.079\,$	
100	.060	$.061\,$	$.063$	$.073\,$	$.054$	$.055\,$	$.078\,$	$.076\,$	
150	$.056\,$	$.057\,$	$.064\,$	$.075\,$	$.058\,$	$.058\,$	$.081\,$	$.082\,$	
250	$.052\,$	$.054\,$	$.063$	$.075\,$	$.058\,$	$.056\,$	$.087\,$	$.086\,$	
500	.049	$.047$	$.064$	$.073\,$	$.054\,$	$.052\,$	$.087\,$	$.086\,$	
1000	.048	$.048$	$.063\,$	$.075\,$	$.055\,$	$.059\,$	$.083\,$.092	
					Early smooth variance reduction, $\varphi = 0, \vartheta = -.8$				
75	.271	$.322\,$.269	.330	.486	.581	$.511\,$.605	
100	$.253\,$	$.284\,$	$.255\,$	$.298\,$.446	$.530\,$.473	.561	
150	$.217\,$.241	$.219\,$	$.258\,$.388	$.448\,$	$.422\,$.483	
$250\,$	$.213\,$.211	.219	$.233\,$	$.340\,$	$.360\,$	$.370\,$	$.399\,$	
$500\,$.176	.171	$.187\,$	$.204\,$	$.297\,$	$.287\,$.337	$.334\,$	
1000	.150	.141	.164	.171	.237	$.221\,$.281	$.270$	
					Early smooth variance reduction, $\varphi = 0, \vartheta = .8$				
$75\,$	$.075\,$.057	$.082$	$.072\,$.082	.063	.108	.089	
100	.070	$.055\,$	$.075\,$	$.072\,$	$.085\,$	$.067\,$.110	$.095$	
150	.065	$.059\,$	$.072\,$	$.077\,$	$.072\,$	$.067\,$.095	.092	
250	.062	$.061\,$	$.072\,$	$.082\,$	$.066\,$	$.066\,$	$.094\,$.097	
500	.059	$.061\,$	$.072\,$.088	$.063\,$	$.067\,$	$.098\,$.107	
1000	.056	$.059$.068	$.084\,$	$.053\,$	$.058\,$	$.085$.098	
					Late smooth variance increase, $\varphi = 0, \vartheta = 0$				
75	.066	$.058\,$	$.136\,$	$.108\,$	$.062\,$	$.064$	$.073\,$.077	
100	.069	$.066\,$	$.154\,$	$.129\,$	$.054\,$	$.056\,$.066	.069	
150	$.059$	$.055\,$	$.149\,$	$.119\,$	$.054$	$.055\,$	$.077\,$	$.081\,$	
$250\,$	$.057$	$.053\,$	$.152\,$	$.121\,$	$.053\,$	$.052\,$	$.078\,$	$.080$	
$500\,$	$.056\,$	$.051\,$	$.156\,$	$.123\,$.053	$.056\,$	$.086\,$.088	
1000	.047	.049	.148	$.125\,$	$.051\,$	$.057\,$.090	$.096\,$	
					Late smooth variance increase, $\varphi = 0, \vartheta = -0.8$				
75	$.351\,$	$.386\,$.417	$.424\,$	$.303\,$.368	$.311\,$	$.376\,$	
100	.311	$.345\,$.385	$.389\,$.276	$.340\,$	$.288\,$.352	
150	.278	.319	.352	$.366\,$.248	.299	.263	.314	
250	.231	.272	.319	.337	.224	.263	.243	.288	
500	.179	.220	.284	.289	.208	.224	.241	.259	
1000	.141	$.182\,$.248	$.258\,$	$.190\,$	$.200\,$.227	.241	
					Late smooth variance increase, $\varphi = 0, \vartheta = .8$				
$75\,$	$.083\,$.081	$.157\,$	$.132\,$.043	$.075\,$.055	.087	
100	$.081\,$	$.078\,$	$.167\,$	$.138\,$	$.051\,$.076	.059	.090	
150	$.070\,$.073	$.154\,$	$.124\,$	$.050\,$.079	.067	.098	
$250\,$	$.072\,$.073	.172	.136	$.050\,$.071	.069	.096	
500	$.063$.062	.156	$.127\,$.056	.067	.084	.106	
1000	$.057\,$	$.064\,$.166	$.134\,$.054	$.062\,$.079	.095	

Table A2: Size of the WB activation knot tests under heteroskedastic MA errors

Notes: 5% nominal level. DGP [\(18\)](#page-11-1) with $c = 0$ and $\sigma_{T,t}$ as defined in [\(20\)](#page-11-2), with $\kappa = .2$, $s_2^2 = .25$ for early negative shifts and $\kappa = .8$, $s_2^2 = 4$ for late positive shifts. The data are adjusted for a constant wild bootstrap replications with $q = 0$. 5000 Monte Carlo replications.

		$z_t=1$			$z_t = (1, t)'$					
T	τ^*	$\breve{\tau}^*$	τ	$\breve{\tau}$	τ^*	$\breve{\tau}^*$	τ	$\breve{\tau}$		
					Early smooth variance reduction, AR errors: $\varphi = -.8, \vartheta = 0$					
75	.088	$.084\,$.090	.089	.132	.120	.155	$.135\,$		
100	$.087$	$.082\,$	$.091\,$	$.088\,$	$.104\,$	$.106\,$.126	.123		
150	$.076\,$	$.073\,$	$.081\,$.083	.084	.088	$.110$.109		
250	$.071\,$	$.071\,$.079	.090	.074	.071	.099	.096		
500	$.057\,$.060	$.071\,$	$.083\,$.060	$.063$	$.088\,$.092		
1000	$.052\,$	$.055\,$	$.065\,$	$.075\,$.054	$.056\,$	$.085\,$.089		
Early smooth variance reduction, AR errors: $\varphi = .8, \vartheta = 0$										
75	.046	.026	.051	.039	.047	.026	.073	.045		
100	.053	.035	.059	.050	.041	.027	.074	.048		
150	.045	.036	.056	.052	.044	.033	.079	.061		
$250\,$.050	$.047$	$.062\,$.070	.050	.043	$.083\,$.071		
500	.050	.048	$.067\,$	$.072\,$.052	.047	.089	.083		
1000	$.057\,$	$.053\,$	$.071\,$	$.081\,$.050	.044	$.083\,$.083		
					Late smooth variance increase, AR errors: $\varphi = -.8, \vartheta = 0$					
75	.086	.088	.148	.120	.056	.070	.061	.073		
100	.082	.089	.152	.129	.058	.068	$.064$.076		
150	.076	.077	.153	.123	.056	.067	.065	.078		
250	.067	.072	.151	.133	.047	.055	.064	.071		
500	.058	.066	$.159\,$.128	.053	.058	.077	.082		
1000	.054	.055	.147	.125	.051	.054	.081	.089		
					Late smooth variance increase, AR errors: $\varphi = .8, \vartheta = 0$					
75	.088	.073	.158	.132	.044	.076	.056	.095		
100	.077	$.067\,$	$.161\,$.119	.044	$.067$.061	.087		
150	.069	.061	.154	$.123\,$.047	.058	.066	.085		
250	.062	.053	.157	$.117\,$.049	.049	.075	.078		
500	.062	.051	.157	.122	.051	.049	.078	.084		
1000	$.051\,$.050	.152	.125	.048	.047	.082	.081		

Table A3: Size of the WB activation knot tests under heteroskedastic AR errors

Notes: DGP [\(18\)](#page-11-1) with $c = 0$ and $\sigma_{T,t}$ as defined in [\(20\)](#page-11-2), with $\kappa = .2$, $s_2^2 = .25$ for early negative shifts and $\kappa = .8$, $s_2^2 = 4$ for late positive shifts. The data are adjusted for a constant $(z_t = 1)$ or a lin the FD method of Schmidt and Phillips [\(1992\)](#page-29-3). All lag orders are computed using the RSMAIC. $B = 499$ wild bootstrap replications with $q = 0$. 5000 Monte Carlo replications.

		$z_t=1$			$z_t = (1, t)'$					
T	τ^*	$\breve{\tau}^*$	τ	$\breve{\tau}$	τ^*	$\breve{\tau}^*$	τ	$\breve{\tau}$		
					Early smooth variance reduction, AR errors: $\varphi = -.8, \vartheta = 0$					
75	.049	.052	.090	.089	.051	.061	.155	.135		
100	.057	$.056\,$	$.091\,$.088	.047	.055	.126	.123		
150	$.052\,$	$.055\,$	$.081\,$	$.083\,$.043	.052	.110	.109		
250	.056	.057	.079	.090	.047	.048	.099	.096		
500	.050	.053	.071	$.083\,$.048	$.050\,$	$.088\,$.092		
1000	.049	.049	.065	.075	.045	$.050$	$.085\,$.089		
Early smooth variance reduction, AR errors: $\varphi = .8, \vartheta = 0$										
75	.046	$.051\,$.051	.039	$.053$.055	.073	.045		
100	$.054\,$.056	.059	.050	.053	.054	.074	.048		
150	$.052\,$.053	.056	.052	.055	.057	.079	.061		
250	$.054\,$.061	.062	.070	.053	$.059\,$	$.083\,$.071		
500	$.055\,$.056	.067	.072	.053	$.057$.089	.083		
1000	.058	$.057\,$.071	.081	.050	.049	.083	.083		
					Late smooth variance increase, AR errors: $\varphi = -.8, \vartheta = 0$					
75	.057	$.050\,$	$.148\,$.120	.026	.034	.061	.073		
100	.061	$.052\,$	$.152\,$.129	.031	$.038\,$	$.064$.076		
150	.060	.054	.153	.123	.042	.045	.065	.078		
250	.059	.054	.151	.133	.036	.040	.064	.071		
500	$.053\,$.053	.159	.128	.049	.049	.077	.082		
1000	$.052\,$.050	.147	$.125\,$.048	.047	.081	.089		
					Late smooth variance increase, AR errors: $\varphi=.8,\, \vartheta=0$					
75	.067	.059	.158	.132	.056	.061	.056	.095		
100	.058	.054	.161	.119	.056	$.056\,$	$.061\,$.087		
150	$.055\,$.056	.154	.123	$.053\,$.058	.066	.085		
250	.058	.052	.157	.117	.053	$.052\,$.075	.078		
500	.059	.055	.157	.122	.053	.050	.078	.084		
1000	.048	.051	.152	.125	.052	.050	.082	.081		

Table A4: Size of the SWB activation knot tests under heteroskedastic AR errors

Notes: DGP [\(18\)](#page-11-1) with $c = 0$ and $\sigma_{T,t}$ as defined in [\(20\)](#page-11-2), with $\kappa = .2$, $s_2^2 = .25$ for early negative shifts and $\kappa = .8$, $s_2^2 = 4$ for late positive shifts. The data are adjusted for a constant ($z_t = 1$) or a li method of Schmidt and Phillips [\(1992\)](#page-29-3). All lag orders are computed using the RSMAIC. *B* = 499 sieve wild bootstrap replications. 5000 Monte Carlo replications.

		$z_t = 1$			$z_t = (1, t)'$					
T	τ^*	$\breve{\tau}^*$	τ	$\breve{\tau}$	τ^*	$\breve{\tau}^*$	τ	$\breve{\tau}$		
					Early smooth variance reduction, AR errors: $\varphi = -.8, \vartheta = 0$					
75	.100	.113	.105	.113	.128	.188	.125	.176		
100	.099	.109	$.096\,$.100	.153	.189	.174	.183		
150	.099	.111	.099	$.102\,$.148	.186	.176	.192		
250	.108	.116	$.102\,$.104	.165	.196	.174	.207		
500	.126	.130	$.125\,$.126	.192	.214	.205	.226		
1000	$.147$.154	$.152\,$.155	.216	.235	.227	.242		
Early smooth variance reduction, AR errors: $\varphi = .8, \vartheta = 0$										
75	$.113$	$.132\,$	$.107$	$.128\,$	$.133\,$.142	$.113\,$.146		
100	$.100\,$	$.114$	$.086\,$	$.102\,$.135	$.143\,$	$.135\,$.151		
150	$.117$	$.127\,$	$.114\,$	$.125\,$.149	.165	.143	.164		
$250\,$.119	.126	.111	$.113\,$	$.168\,$	$.178\,$	$.158\,$.171		
500	$.130\,$.135	.126	$.126\,$	$.187\,$	$.206\,$	$.176\,$.196		
1000	$.158\,$.169	$.137\,$	$.152\,$.217	.238	$.220\,$.245		
					Late smooth variance increase, AR errors: $\varphi = -.8, \vartheta = 0$					
75	.140	.200	.141	.224	.123	.183	.186	.228		
100	.138	.203	$.127\,$.217	.140	.189	$.205$.245		
150	.132	$.207$	$.116\,$	$.207\,$.167	.210	.209	.242		
$250\,$	$.134\,$.228	$.114\,$	$.230\,$.191	.239	.251	.298		
500	$.150\,$	$.263\,$.144	.269	.235	.278	.255	.286		
1000	$.156\,$.265	.153	.273	.262	.300	$.275\,$.317		
					Late smooth variance increase, AR errors: $\varphi = .8, \vartheta = 0$					
75	.072	.126	.052	$.092\,$.129	.135	.117	.089		
100	.093	.151	.075	$.121\,$.141	.151	.133	.121		
150	$.105\,$.176	$.091\,$.159	.164	.181	$.157\,$.159		
250	.106	.193	.092	$.178\,$.201	$.215\,$	$.197\,$.215		
500	.123	.229	.108	.214	.232	.250	.228	.259		
1000	$.152\,$.265	.165	$.262\,$.266	.294	.272	.307		

Table A5: Local power of the SWB tests under heteroskedastic AR errors

Notes: DGP [\(18\)](#page-11-1) with $c = -7$ for $z_t = 1$ and $c = -13.5$ for $z_t = (1, t)$. $\sigma_{T, t}$ as defined in [\(20\)](#page-11-2), with $\kappa = .2$, $s_2^2 = .25$ for early negative shifts and $\kappa = .8$, $s_2^2 = 4$ for late positive shifts. The data are ad at 5%. All lag orders are computed using the RSMAIC. *B* = 499 sieve wild bootstrap replications. 5000 Monte Carlo replications.

		$z_t = 1$			$z_t = (1, t)'$						
T	τ^*	$\breve{\tau}^*$	τ	$\breve{\tau}$	τ^*	$\breve{\tau}^*$	τ	$\breve{\tau}$			
						Early smooth variance reduction, MA errors: $\varphi = 0, \vartheta = -.8$					
75	.205	.279	.078	.097	.500	.637	.113	.133			
100	.165	.222	$.070\,$.089	.411	.541	.128	.165			
150	$.135\,$	$.177$	$.080\,$.102	.279	.397	.133	.182			
250	.137	.158	$.085$.102	.202	.297	.146	.184			
500	.140	$.154\,$.108	.121	.201	.250	.158	.183			
1000	.166	.174	$.115$	$.120\,$.222	.257	.166	.202			
Early smooth variance reduction, MA errors: $\varphi = 0, \vartheta = .8$											
$75\,$	$.106\,$	$.114\,$	$.092\,$.104	.128	.157	$.124\,$.156			
100	$.107\,$.120	$.110\,$.114	.127	.151	.123	.158			
150	$.107\,$	$.121\,$	$.103\,$	$.110\,$.131	.158	.153	.175			
$250\,$.110	$.126\,$	$.107\,$	$.110$.156	.184	$.162\,$.197			
500	$.124\,$	$.132\,$	$.120\,$.127	.182	.200	.184	.200			
1000	$.157\,$	$.166\,$	$.148\,$.150	.206	.237	.223	.246			
						Late smooth variance increase, MA errors: $\varphi = 0, \vartheta = -.8$					
$75\,$.444	.473	$.186\,$.193	.500	.615	.189	.230			
100	.375	.404	.176	.184	.412	.526	.196	.244			
150	$.314\,$	$.362\,$.191	.201	.275	.383	.191	.233			
$250\,$.301	.343	$.192\,$	$.223\,$.204	.304	.177	.241			
500	$.302\,$.324	.234	.244	.195	.275	.196	.239			
1000	$.309\,$.328	.228	.238	.218	.287	.200	.256			
						Late smooth variance increase, MA errors: $\varphi = 0, \vartheta = .8$					
75	.087	$.157\,$	$.064\,$.134	$.137\,$.161	.163	.145			
100	.089	$.165\,$.067	.160	.137	.166	$.156\,$.168			
150	$.102\,$	$.188\,$	$.083\,$.181	.151	.180	.169	.169			
250	.110	$.206\,$	$.085\,$	$.182\,$.184	.206	.196	.222			
500	.129	.243	.109	.244	$.219\,$.246	.213	.248			
1000	.152	.271	.141	.247	.238	.278	.239	.272			

Table A6: Local power of the SWB tests under heteroskedastic MA errors

Notes: DGP [\(18\)](#page-11-1) with $c = -7$ for $z_t = 1$ and $c = -13.5$ for $z_t = (1, t)$. $\sigma_{T, t}$ as defined in [\(20\)](#page-11-2), with $\kappa = .2$, $s_2^2 = .25$ for early negative shifts and $\kappa = .8$, $s_2^2 = 4$ for late positive shifts. The data are ad at 5%. All lag orders are computed using the RSMAIC, setting $q = p$. $B = 499$ sieve wild bootstrap replications. 5000 Monte Carlo replications.

References

- Arnold, M. C., & Reinschlüssel, T. (2024). Adaptive Unit Root Inference in Autoregressions using the Lasso Solution Path. *<arXiv:2404.06205>* .
- Audrino, F., & Camponovo, L. (2017). Oracle properties, bias correction, and bootstrap inference for adaptive lasso for time series m-estimators. *Journal of Time Series Analysis*, *39* (2), 111–128.
- Beare, B. K. (2018). Unit root testing with unstable volatility. *Journal of Time Series Analysis*, *39* (6), 816–835.
- Belloni, A., Chernozhukov, V., & Hansen, C. (2013). Inference on treatment effects after selection among high-dimensional controls. *Review of Economic Studies*, *81* (2), 608–650.
- BIS. (2023). *Real residential property prices QBER368BIS, QFRR368BIS, QDER368BIS, QITR368BIS, QNLR368BIS, QESR368BIS* [Bank for International Settlements. Retrieved from FRED, Federal Reserve Bank of St. Louis on 3-Dec-2023]. [https:](https://fred.stlouisfed.org/) [//fred.stlouisfed.org/](https://fred.stlouisfed.org/)
- Candes, E., & Tao, T. (2007). The dantzig selector: Statistical estimation when *p* is much larger than *n*. *The Annals of Statistics*, *35* (6).
- Cavaliere, G., Phillips, P. C. B., Smeekes, S., & Taylor, A. M. R. (2015). Lag length selection for unit root tests in the presence of nonstationary volatility. *Econometric Reviews*, *34* (4), 512–536.
- Cavaliere, G., & Taylor, A. M. R. (2007). Testing for unit roots in time series models with non-stationary volatility. *Journal of Econometrics*, $140(2)$, 919–947.
- Cavaliere, G., & Taylor, A. M. R. (2008). Bootstrap unit root tests for time series with nonstationary volatility. *Econometric Theory*, *24* (01).
- Cavaliere, G., & Taylor, A. M. R. (2009a). Bootstrap *M* unit root tests. *Econometric Reviews*, *28* (5), 393–421.
- Cavaliere, G., & Taylor, A. M. R. (2009b). Heteroskedastic time series with a unit root. *Econometric Theory*, *25* (5), 1228–1276.
- Chang, Y., & Park, J. (2002). On the asymptotics of ADF tests for unit roots. *Econometric Reviews*, *21* (4), 431–447.
- Chang, Y., & Park, J. Y. (2003). A sieve bootstrap for the test of a unit root. *Journal of Time Series Analysis*, *24* (4), 379–400.
- Chatterjee, A., & Lahiri, S. N. (2011). Bootstrapping lasso estimators. *Journal of the American Statistical Association*, *106* (494), 608–625.
- Chernozhukov, V., Chetverikov, D., & Kato, K. (2013). Gaussian approximations and multiplier bootstrap for maxima of sums of high-dimensional random vectors. *The Annals of Statistics*, *41* (6).
- Chernozhukov, V., Chetverikov, D., Kato, K., & Koike, Y. (2023). High-dimensional data bootstrap. *Annual Review of Statistics and Its Application*, *10* (1), 427–449.
- Davidson, R., & Flachaire, E. (2008). The wild bootstrap, tamed at last. *Journal of Econometrics*, *146* (1), 162–169.
- Demetrescu, M., & Hanck, C. (2016). Robust inference for near-unit root processes with time-varying error variances. *Econometric Reviews*, *35* (5), 751–781.
- Dezeure, R., Bühlmann, P., & Zhang, C.-H. (2017). High-dimensional simultaneous inference with the bootstrap. *TEST*, *26* (4), 685–719.
- Efron, B., Hastie, T., Johnstone, I., & Tibshirani, R. (2004). Least angle regression. *The Annals of Statistics*, *32* (2), 407–499.
- Elliott, G., Rothenberg, T. J., & Stock, J. H. (1996). Efficient tests for an autoregressive unit root. *Econometrica*, *64* (4), 813–836.
- Ferretti, N. E., & Romo, J. (1996). Bootstrap tests for unit root AR(1) models. *Biometrika*, *83* (4), 849–860.
- Francke, M., & Korevaar, M. (2021). Housing markets in a pandemic: Evidence from historical outbreaks. *The Journal of Real Estate Finance and Economics*.
- Hansen, C., & Liao, Y. (2018). The factor-lasso and k-step bootstrap approach for inference in high-dimensional economic applications. *Econometric Theory*, *35* (03), 465–509.
- Hastie, T., & Efron, B. (2022). *Lars: Least angle regression, lasso and forward stagewise* [R package version 1.3].
- Herwartz, H., & Siedenburg, F. (2010). A new approach to unit root testing. *Computational Economics*, *36* (4), 365–384.
- Herwartz, H., & Walle, Y. M. (2018). A powerful wild bootstrap diagnosis of panel unit roots under linear trends and time-varying volatility. *Computational Statistics*, *33* (1), 379–411.
- Huang, J., Ma, S., & Zhang, C.-H. (2008). Adaptive lasso for sparse high-dimensional regression models. *Statistica Sinica*, *18* (4), 1603–1618.
- Jordà, Ò., Schularick, M., & Taylor, A. (2016). The great mortgaging: Housing finance, crises, and business cycles. *Journal of Economic Perspectives*.
- Liu. (1988). Bootstrap procedures under some non i.i.d. models. *The Annals of Statistics*, *16* (4).
- Mammen, E. (1993). Bootstrap and wild bootstrap for high dimensional linear models. *The Annals of Statistics*, *21* (1).
- Mian, A., Sufi, A., & Trebbi, F. (2015). Foreclosures, house prices, and the real economy. *American Economic Review*.
- Ng, S., & Perron, P. (2001). Lag length selection and the construction of unit root tests with good size and power. *Econometrica*, *69* (6), 1519–1554.
- Paparoditis, E., & Politis, D. N. (2003). Residual-based block bootstrap for unit root testing. *Econometrica*, *52* (1), 139–159.
- Park, J. Y. (2003). Bootstrap unit root tests. *Econometrica*, *71* (6), 1845–1895.
- Perron, P., & Ng, S. (1996). Useful modifications to some unit root tests with dependent errors and their local asymptotic properties. *The Review of Economic Studies*, *63* (3), 435–463.
- Phillips, P. C. B. (1987). Time series regression with a unit root. *Econometrica*, *55* (2), 277–301.
- Phillips, P. C. B., & Perron, P. (1988). Testing for a unit root in time series regression. *Biometrika*, *75* (2), 335–346.
- Qian, J., & Su, L. (2016). Shrinkage estimation of common breaks in panel data models via adaptive group fused lasso. *Journal of Econometrics*, *191* (1), 86–109.
- R Core Team. (2023). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing. Vienna, Austria.
- Richard, P. (2009). Modified fast double sieve bootstraps for adf tests. *Computational Statistics & Data Analysis*, *53* (12), 4490–4499.
- Said, S. E., & Dickey, D. A. (1984). Testing for unit roots in autoregressive-moving average models of unknown order. *Biometrika*, *71* (3), 599–607.
- Schmidt, P., & Phillips, P. C. B. (1992). LM tests for a unit root in the presence of deterministic trends. *Oxford Bulletin of Economics and Statistics*, *54* (3), 257– 287.
- Schweikert, K. (2021). Oracle efficient estimation of structural breaks in cointegrating regressions. *Journal of Time Series Analysis*, *43* (1), 83–104.
- Smeekes, S., & Taylor, A. R. (2012). Bootstrap union tests for unit roots in the presence of nonstationary volatility. *Econometric Theory*, *28* (2), 422–456.
- Stock, J. H. (1999). A class of tests for integration and cointegration. *Cointegration, Causality and Forecasting. A Festschrift in Honour of Clive W.J. Granger*, 135– 167.
- Swensen, A. R. (2003). Bootstrapping unit root tests for integrated processes. *Journal of Time Series Analysis*, *24* (1), 99–126.