

Exotic Hidden-heavy Hadrons and Where to Find Them

Eric Braaten* and Roberto Bruschini†

Department of Physics, The Ohio State University, Columbus, OH 43210, USA

The Born-Oppenheimer potentials for QCD with light quarks include adjoint-hadron potentials that are repulsive at short distances and heavy-hadron-pair potentials that approach thresholds at large distances. The adjoint-hadron potentials must connect smoothly to the heavy-hadron-pair potentials at intermediate distances. We identify exotic hidden-heavy hadrons as bound states and resonances in adjoint-hadron potentials that cross below a heavy-hadron-pair threshold before approaching it. This explains why many exotic hidden-charm and hidden-bottom hadrons have energies near heavy-hadron-pair thresholds. The remarkable properties of some exotic hidden-heavy mesons can be explained by fine tunings of adjoint-meson energies in QCD.

Keywords: Exotic hadrons, heavy quarks, Born-Oppenheimer approximation, hadron spectroscopy.

Introduction. Dozens of exotic heavy hadrons have been discovered since the beginning of the 21st century [1]. Most of them are hidden-heavy hadrons whose constituents include a charm or bottom quark-antiquark ($c\bar{c}$ or $b\bar{b}$) pair. By a recent count, they include as many as 43 $c\bar{c}$ tetraquark mesons, 5 $b\bar{b}$ tetraquark mesons, and 5 $c\bar{c}$ pentaquark baryons [2]. Their discoveries challenge our understanding of quantum chromodynamics (QCD) and other strongly interacting gauge theories.

Most models for exotic heavy hadrons have tenuous connections to the fundamental theory QCD. They have multiple parameters with unknown relations to the QCD coupling constant α_s and the quark masses. Molecular models whose constituents are color-singlet hadrons are motivated by many exotic heavy hadrons having energies near heavy-hadron-pair thresholds. Models whose constituents have color charges predict an explosion in the number of exotic heavy hadrons, making it easy to accommodate new discoveries by overlooking the many predictions of unobserved states.

The Born-Oppenheimer (B-O) approach to multi-heavy hadrons is based firmly on QCD. It has been developed into an effective field theory for multi-heavy hadrons called BOEFT [3–6]. The B-O approach separates the problem into two steps: 1) calculating B-O potentials using lattice QCD, 2) solving the Schrödinger equation for heavy quarks and antiquarks in the B-O potentials. The B-O approach provides a unified framework for describing all multi-heavy hadrons [5–9]. Although theoretically compelling, it has not yet explained the pattern of the observed exotic heavy hadrons or made compelling predictions of new ones before their discoveries.

In this paper, we emphasize that the B-O potentials for hidden-heavy hadrons that approach heavy-hadron-pair thresholds at large distances must, with the exception of the ground-state potential, connect smoothly to repulsive $1/r$ potentials at short distances. This severely reduces the B-O channels that can support bound states and resonances. It explains why many exotic hidden-heavy hadrons have energies near heavy-hadron-pair thresholds. We also identify the fine tunings of QCD responsible for



Figure 1. Exotic hidden-heavy hadrons and other fantastic beasts (image by Jam-Di, [deviantart.com/jam-di](https://www.deviantart.com/jam-di)).

the remarkable properties of some exotic hidden-heavy mesons.

Static Color Sources. The B-O approximation exploits the large mass of a heavy quark compared to the momentum scales of gluons and light quarks. The B-O potentials for hidden-heavy hadrons are discrete energies in the spectrum of QCD with static $\mathbf{3}$ and $\mathbf{3}^*$ color sources separated by a distance r . The sources reduce the rotational symmetry to a cylindrical subgroup with quantum number λ that is integer or half-integer. They reduce the parity and charge-conjugation symmetries with quantum numbers P and C to their product CP . There is also a reflection symmetry through a plane containing the two sources with quantum number R . The B-O symmetries are traditionally denoted Λ_η^ϵ , where $\Lambda = |\lambda|$ (or $\Sigma, \Pi, \Delta, \dots$ if $|\lambda|$ is 0, 1, 2, \dots), $\eta = g, u$ if $CP = +1, -1$, and $\epsilon = +, -$ if $R = +1, -1$. If Λ is not Σ , ϵ is omitted. The B-O potentials are also labeled using approximate light-quark flavor symmetries, such as isospin.

Calculating B-O potentials at large r and small r reduces to the simpler problem of QCD with a single static color source. Its spectrum is determined up to an addi-

tive constant that depends on the color source. The discrete states are labeled by quantum numbers J^{PC} , where J is the angular momentum. They are also labeled using light-quark flavor symmetries. A discrete state bound to a static $\mathbf{3}$ or $\mathbf{3}^*$ color source is called a $\mathbf{3}$ -hadron or $\mathbf{3}^*$ -hadron. A discrete state bound to a static $\mathbf{8}$ color source is called an *adjoint hadron* or an $\mathbf{8}$ -hadron.

As $r \rightarrow \infty$, confinement and the cluster decomposition requires the finite-energy states of QCD with $\mathbf{3}$ and $\mathbf{3}^*$ sources to be pairs of a $\mathbf{3}$ -hadron and a $\mathbf{3}^*$ -hadron. We refer to the corresponding B-O potentials at large r as $(\mathbf{3}+\mathbf{3}^*)$ -potentials. As $r \rightarrow 0$, the $\mathbf{3}$ and $\mathbf{3}^*$ sources reduce to a linear combination of a color-singlet ($\mathbf{1}$) source and an $\mathbf{8}$ source. QCD with a $\mathbf{1}$ source, which is the same as no source, has only one discrete state: the QCD vacuum with B-O quantum numbers Σ_g^+ . The corresponding $\mathbf{1}$ -potential is an attractive $1/r$ potential at small r . We refer to a B-O potential at small r whose QCD state approximates an $\mathbf{8}$ -hadron as an $\mathbf{8}$ -potential. As $r \rightarrow 0$, an $\mathbf{8}$ -potential approaches a repulsive $1/r$ potential shifted by the $\mathbf{8}$ -hadron energy.

The spectrum of QCD with two static color sources must be a smooth function of r . This provides surprisingly strong constraints on the B-O potentials for hidden-heavy hadrons. The ground-state Σ_g^+ potential connects a $(\mathbf{3}+\mathbf{3}^*)$ -potential at large r to the $\mathbf{1}$ -potential at small r . All other $(\mathbf{3}+\mathbf{3}^*)$ -potentials at large r must connect to $\mathbf{8}$ -potentials at small r . For double-heavy hadrons with two heavy quarks, the $(\mathbf{3}+\mathbf{3})$ -potentials at large r in lattice QCD connect at small r to either attractive $\mathbf{3}^*$ -potentials or repulsive $\mathbf{6}$ -potentials [10]. The analogous fact for hidden-heavy hadrons has only been recognized recently [8].

There have been few calculations of the spectrum of $\mathbf{8}$ -hadrons using lattice QCD. The first calculations for gluelumps ($\mathbf{8}$ -mesons that are SU(3)-flavor singlets) by Foster and Michael in 1998 used pure SU(3) gauge theory [11]. The ground-state gluelump is $J^{PC} = 1^{+-}$. The spectrum has recently been calculated more accurately [12]. There are also lattice calculations with 2+1 flavors of light quarks [13]. The spectrum of $\mathbf{8}$ -mesons besides gluelumps has only been calculated using SU(3) gauge theory with light valence quarks [11]. The lowest-energy $\mathbf{8}$ -mesons are 1^{--} and 0^{-+} . Various models for QCD give more detailed predictions for the spectrum of $\mathbf{8}$ -hadrons, as summarized in Ref. [14].

History of B-O Potentials. The first quantitative QCD result from lattice gauge theory was the Σ_g^+ potential in pure SU(3) gauge theory in 1984 [15]. This confining potential is an attractive $1/r$ potential at small r and increases linearly at large r . The ground-state Σ_g^+ potential in QCD with light quarks must cross over at large r to a constant equal to twice the energy of the ground-state $\mathbf{3}$ -meson. Bali *et al.* showed in 1995 that this occurs through a narrow avoided crossing with an excited Σ_g^+ potential that crosses over from approximately constant




Figure 2. An $\mathbf{8}$ -potential at short distances can approach a $(\mathbf{3}+\mathbf{3}^*)$ -potential at long distances by decreasing monotonically (upper curve at small r). It may also decrease below the $(\mathbf{3}+\mathbf{3}^*)$ -threshold before approaching it from below or from above (lower curve at small r), in which case it may support a bound state (solid horizontal line) or a resonance (dashed horizontal line).

to increasing linearly with r [16]. This suggested that QCD with light quarks has additional $(\mathbf{3}+\mathbf{3}^*)$ -potentials at large r for every pair of a $\mathbf{3}$ -hadron and a $\mathbf{3}^*$ -hadron.

The B-O approximation for QCD was pioneered in 1999 by Juge, Kuti, and Morningstar, who applied it to quarkonium hybrid mesons [17]. They showed that pure SU(3) gauge theory also has higher potentials with other B-O quantum numbers [18]. They are confining potentials that increase linearly at large r . The development of pNRQCD revealed that they are $\mathbf{8}$ -potentials at small r that differ by gluelump energy differences [19].

The discoveries of T_{bb1}^+ (10610) and T_{bb1}^+ (10650) (Z_b and Z_b') in 2011 [20] and $T_{c\bar{c}1}^+$ (3900) in 2013 [21, 22] motivated the realization that QCD has additional $\mathbf{8}$ -potentials for every $\mathbf{8}$ -hadron whose flavor is not SU(3)-singlet [23, 24]. At small r , they approach a repulsive $1/r$ potential shifted by the energy of the $\mathbf{8}$ -hadron. Charged quarkonium states were identified as bound states in these potentials, which were assumed to increase linearly at large r . New confining potentials for every $\mathbf{8}$ -hadron imply an explosion in the number of exotic hidden-heavy hadrons.

The continuity constraint that an $\mathbf{8}$ -potential at small r connects to a $(\mathbf{3}+\mathbf{3}^*)$ -potential at large r severely limits the B-O potentials capable of supporting bound states. An $\mathbf{8}$ -potential at small r decreases with increasing r . The simplest possibility is that it decreases monotonically to the $(\mathbf{3}+\mathbf{3}^*)$ -threshold, in which case it cannot support bound states. The second-simplest possibility is that it crosses below the $(\mathbf{3}+\mathbf{3}^*)$ -threshold and then approaches it from below, in which case it can support bound states. The third-simplest possibility is that it crosses below the $(\mathbf{3}+\mathbf{3}^*)$ -threshold, crosses it again, and then approaches it from above, in which case it can also support res-

onances. The simplest and third-simplest possibilities are illustrated in Fig. 2. We identify these bound states and resonances as exotic hidden-heavy hadrons. This explains why the masses of most observed exotic hidden-heavy hadrons are near heavy-hadron-pair thresholds. Since energy splittings between **8**-hadrons are typically larger than those between pairs of a **3**-hadron and a **3***-hadron, most **8**-potentials approach a (**3**+**3***)-threshold from above. Only those associated with the lowest-energy **8**-hadrons cross below the (**3**+**3***)-threshold before approaching it. This avoids an explosion in the number of exotic hidden-heavy hadrons. The thresholds with exotic hidden-heavy hadrons nearby can be predicted most easily by using lattice QCD to calculate the spectrum of **8**-hadrons.

Model B-O potentials. Lattice calculations in SU(3) gauge theory with valence quarks suggest that the lowest-energy **8**-mesons are $J^{PC} = 1^{--}$ and 0^{-+} [11]. The B-O potentials associated with a 1^{--} **8**-meson are $\Sigma_g^{+'}$ and Π_g . The B-O potential associated with a 0^{-+} **8**-meson is Σ_u^- . The lowest-energy multiplets in these B-O potentials are given in Table I of Ref. [8]. For the 1^{--} **8**-meson, the ground-state multiplet is a quartet consisting of a heavy-quark-pair spin-singlet ($S = 0$) 1^{+-} and a spin-triplet ($S = 1$) $(0, 1, 2)^{++}$. For the 0^{-+} **8**-meson, the ground-state multiplet is a doublet consisting of a spin-singlet 0^{++} and a spin-triplet 1^{+-} . Note that 0^{++} and 1^{+-} also appear in the ground-state multiplet for the 1^{--} **8**-meson.

For the purposes of illustration, we use simple models for the $\Sigma_g^{+'}$, Π_g , and Σ_u^- potentials. We take the zero of energy to be the spin-weighted isospin-averaged heavy-meson-pair threshold: $E_{D\bar{D}} = 3946$ MeV for charm mesons and $E_{B\bar{B}} = 10627$ MeV for bottom mesons. Our model for the Λ_η^ϵ potential associated with the J^{PC} **8**-meson with isospin I is

$$V_{\Lambda_\eta^\epsilon}^{(I)}(r) = \begin{cases} \kappa_8/r + E_{J^{PC}}^{(I)} + A_{\Lambda_\eta^\epsilon}^{(I)} r^2 & \text{if } r < R_{\Lambda_\eta^\epsilon}^{(I)}, \\ B_{\Lambda_\eta^\epsilon}^{(I)} e^{-r/d} & \text{if } r > R_{\Lambda_\eta^\epsilon}^{(I)}, \end{cases} \quad (1)$$

where $R_{\Lambda_\eta^\epsilon}^{(I)}$ and $B_{\Lambda_\eta^\epsilon}^{(I)}$ are determined by continuity and smoothness at $r = R_{\Lambda_\eta^\epsilon}^{(I)}$. The strength κ_8 of the color-Coulomb potential is the same for all **8**-potentials. The **8**-meson energy $E_{J^{PC}}^{(I)}$ is the same for all potentials associated with a given **8**-meson. We choose the relaxation length in the large- r potential to be the Sommer scale: $d = r_0 = 0.5$ fm. Parametrizations of 8 of the lowest B-O potentials in pure SU(3) gauge theory were presented in Ref. [25]. We take their values for $\kappa_8 = 0.037$ and for $r_0^3 A_{\Lambda_\eta^\epsilon}$: 0.11, 1.18, and 0.68 for $\Sigma_g^{+'}$, Π_g , and Σ_u^- . We treat the **8**-meson energies $E_{J^{PC}}^{(I)}$ as adjustable parameters.

Schrödinger equation. The Schrödinger equation for a heavy-quark pair in the coupled $\Sigma_g^{+'}$ and Π_g potentials and in the Σ_u^- potential are given in Ref. [8]. We take

the charm and bottom quark masses to be $m_c = 1.48$ GeV and $m_b = 4.89$ GeV. In the isospin-0 channel, we ignore the narrow avoided crossing with the Σ_g^+ quarkonium potential.

The exotic $c\bar{c}$ meson $\chi_{c1}(3872)$ (X_c) discovered in 2003 [26] is remarkable because its mass is within 100 keV of the $D^{*0}\bar{D}^0$ threshold [27]. It has isospin 0 and $J^{PC} = 1^{++}$. We identify X_c with the 1^{++} state in the ground-state multiplet for the isospin-0 1^{--} **8**-meson. The ground-state energy $\varepsilon_1^{(0)}$ depends on $E_{1^{--}}^{(0)}$. There is a critical energy $E_{1^{--}}^{(0)*}$ for which it is exactly at threshold: $\varepsilon_1^{(0)*} = 0$. The critical energy from solving the Schrödinger equation with $m_c = 1.48$ GeV is $E_{1^{--}}^{(0)*} = -157$ MeV.

If a potential has the critical depth for the $c\bar{c}$ system, it must support $b\bar{b}$ bound states. Given $E_{1^{--}}^{(0)*}$, we determine their energies by solving the Schrödinger equation with $m_b = 4.89$ GeV. The ground-state energy is $\varepsilon_1^{(0)} = -17$ MeV. There is one other bound state with energy -4 MeV. Its multiplet is a doublet consisting of a spin-singlet 0^{-+} and a spin-triplet 1^{--} .

The exotic $b\bar{b}$ mesons Z_b and Z'_b are remarkable because their masses are only about 3 MeV above the $B^*\bar{B}$ and $B^*\bar{B}^*$ thresholds. They have isospin 1 and the neutral mesons are 1^{+-} . Two candidates for these two states are the 1^{+-} states in the ground-state multiplets for the isospin-1 1^{--} and 0^{-+} **8**-mesons. The ground-state energies $\varepsilon_1^{(1)}$ and $\varepsilon_0^{(1)}$ depend on $E_{1^{--}}^{(1)}$ and $E_{0^{-+}}^{(1)}$. There are critical energies $E_{1^{--}}^{(1)*}$ and $E_{0^{-+}}^{(1)*}$ for which both ground states are exactly at threshold: $\varepsilon_1^{(1)*} = \varepsilon_0^{(1)*} = 0$. The critical energies from solving the Schrödinger equation with $m_b = 4.89$ GeV are $E_{1^{--}}^{(1)*} = -95$ MeV and $E_{0^{-+}}^{(1)*} = -107$ MeV.

These critical energies suggest that the lowest energy **8**-hadron is an **8**-meson with isospin 0 and $J^{PC} = 1^{--}$. The **8**-mesons with isospin 1 and $J^{PC} = 0^{-+}$ and 1^{--} are higher in energy and close to each other. These predictions present a challenge for lattice QCD.

Spin splittings. Quantitative predictions from the B-O approximation require taking into account the spin and flavor splittings of heavy hadrons. The spin splitting Δ between the 1^- and 0^- heavy mesons M^* and M scales with the heavy quark mass m_Q as $1/m_Q$. The spin splitting is $\Delta_c = 141$ MeV for charm mesons and $\Delta_b = 45$ MeV for bottom mesons. The energies of M and M^* relative to their spin average are $-\frac{3}{4}\Delta$ and $+\frac{1}{4}\Delta$. The thresholds for heavy-meson pairs are $-\frac{3}{2}\Delta$ for MM , $-\frac{1}{2}\Delta$ for $M^*\bar{M}$ and $M\bar{M}^*$, and $+\frac{1}{2}\Delta$ for $M^*\bar{M}^*$.

The B-O potential matrix includes terms of order $1/m_Q$ that break the heavy-quark spin symmetry. They remove the degeneracies between states with different J^{PC} in a B-O multiplet and mix states with the same J^{PC} from different B-O multiplets. At large r , the spin-splitting terms reduce to a constant potential matrix V_{SS}

that depends on the spin vectors \mathbf{S}_1 and \mathbf{S}_2 of the heavy quark and antiquark:

$$V_{\text{SS}}(\mathbf{S}_1, \mathbf{S}_2) = \Delta \mathbf{j}_1 \cdot \mathbf{S}_1 + \Delta \mathbf{j}_2 \cdot \mathbf{S}_2, \quad (2)$$

where \mathbf{j}_1 and \mathbf{j}_2 are the light QCD angular momenta of the well-separated $\mathbf{3}$ -meson and $\mathbf{3}^*$ -meson. To determine its qualitative effects, we will take into account V_{SS} at first order in perturbation theory. This approximation puts the heavy-meson pair thresholds at their correct physical values.

We first consider the ground-state multiplet for a 1^{--} $\mathbf{8}$ -meson. If its energy in the absence of spin splittings is ε_1 , the energies of the four states to order Δ are

$$E_{1^{+-}} = \varepsilon_1, \quad E_{0^{++}} = \varepsilon_1 - \Delta, \quad (3a)$$

$$E_{1^{++}} = \varepsilon_1 - \frac{1}{2}\Delta, \quad E_{2^{++}} = \varepsilon_1 + \frac{1}{2}\Delta. \quad (3b)$$

If ε_1 is tuned to 0, the 1^{++} and 2^{++} states remain near the $M^*\bar{M}$ and $M^*\bar{M}^*$ thresholds. The 0^{++} state is halfway between the $M\bar{M}$ and $M^*\bar{M}$ thresholds. The 1^{+-} state is halfway between the $M^*\bar{M}$ and $M^*\bar{M}^*$ thresholds.

We next consider the ground-state multiplet for a 0^{+-} $\mathbf{8}$ -meson. If its energy in the absence of spin splittings is ε_0 , the energies of the two states to order Δ are

$$E_{1^{+-}} = E_{0^{++}} = \varepsilon_0. \quad (4)$$

There are no corrections at first order in Δ . If ε_0 is tuned to 0, the two states remain degenerate with energies halfway between the $M^*\bar{M}$ and $M^*\bar{M}^*$ thresholds.

If ε_0 and ε_1 are nearly degenerate, it is necessary to diagonalize the 2×2 submatrices of V_{SS} for the 0^{++} and 1^{+-} states in the two multiplets. The resulting pairs of energies can be approximated by

$$E_{0^{++}}^{\pm} = \frac{\varepsilon_1 + \varepsilon_0 - \Delta \pm \sqrt{(\varepsilon_1 - \varepsilon_0 - \Delta)^2 + 3\Delta^2}}{2}, \quad (5a)$$

$$E_{1^{+-}}^{\pm} = \frac{\varepsilon_1 + \varepsilon_0 \pm \sqrt{(\varepsilon_1 - \varepsilon_0)^2 + \Delta^2}}{2}. \quad (5b)$$

If Δ^2 is set to 0, these energies reduce to those in Eqs. (3a) and (4). In the limit $\varepsilon_0 \rightarrow \varepsilon_1$, the energies of the 0^{++} states are $\varepsilon_1 - \frac{3}{2}\Delta$ and $\varepsilon_1 + \frac{1}{2}\Delta$ and those of the 1^{+-} states are $\varepsilon_1 - \frac{1}{2}\Delta$ and $\varepsilon_1 + \frac{1}{2}\Delta$. If ε_1 and ε_0 are both tuned to 0, the two 0^{++} states are at the $M\bar{M}$ and $M^*\bar{M}^*$ thresholds and the two 1^{+-} states are at the $M^*\bar{M}$ and $M^*\bar{M}^*$ thresholds.

The two 0^{++} states with energies in Eq. (5a) and the two 1^{+-} states with energies in Eq. (5b) are both superpositions of heavy-quark-pair spin-singlet and spin-triplet states in the ground-state multiplets of the 0^{+-} and 1^{--} $\mathbf{8}$ -mesons. For the 1^{+-} states, the mixing angle θ_1 satisfies $\tan(2\theta_1) = \Delta/(\varepsilon_1 - \varepsilon_0)$. For the 0^{++} states, the mixing angle θ_0 satisfies $\tan(2\theta_0) = \sqrt{3}\Delta/(\varepsilon_1 - \varepsilon_0 - \Delta)$. In the limit $\varepsilon_0 \rightarrow \varepsilon_1$, the two 1^{+-} states are equal-amplitude superpositions of spin-singlet and spin-triplet.




Figure 3. Energy levels for the multiplet of isospin-0 hidden-charm tetraquark mesons associated with the 1^{--} $\mathbf{8}$ -meson. The mass of $\chi_{c1}(3872)$ was used as an input to determine the energy of the $\mathbf{8}$ -meson. The dashed horizontal lines are the thresholds for charm-meson pairs.

The two 0^{+-} states are superpositions of spin-singlet and spin-triplet with amplitudes that differ by a factor of $\sqrt{3}$.

Note that the limit $\varepsilon_0 \rightarrow \varepsilon_1$ reproduces a molecular model for the tetraquark mesons [28]. In general, the B-O framework reproduces a molecular model in cases where the energy levels closest to the threshold are degenerate for all B-O potentials that approach the same $(\mathbf{3}+\mathbf{3}^*)$ -potential.

X_c and X_b Multiplets. If $E_{1^{--}}^{(0)}$ is tuned to its critical value in the $c\bar{c}$ system, we can predict the masses of X_c and its B-O partners to first order in Δ_c . The mass of X_c is near the $D^*\bar{D}$ threshold at 3.876 GeV. The mass of the 2^{++} partner is near the $D^*\bar{D}^*$ threshold at 4.017 GeV. The masses of the 0^{++} and 1^{+-} partners are near 3.805 GeV and 3.946 GeV. The energy levels in the multiplet including X_c are illustrated in Fig. 3.

If $E_{1^{--}}^{(0)}$ is tuned to its critical value in the $c\bar{c}$ system, we can predict the masses for the ground-state multiplet of the $b\bar{b}$ system to first order in Δ_b . The mass of the 1^{++} state X_b is about 17 MeV below the $B^*\bar{B}$ threshold. The mass of the 2^{++} partner is about 17 MeV below the $B^*\bar{B}^*$ threshold. The masses of the 0^{++} and 1^{+-} partners are near 10.565 GeV and 10.610 GeV.

Z_b and Z'_b Multiplets. Another remarkable feature of Z'_b besides its proximity to the $B^*\bar{B}^*$ threshold is that, although it can decay into $B^*\bar{B}$ and $B\bar{B}^*$ through S -wave channels, these decays are not seen. The suppression of these decays can be explained by a fine-tuning of $E_{1^{--}}^{(1)}$ and $E_{0^{+-}}^{(1)}$ so that $\varepsilon_1^{(1)} = \varepsilon_0^{(1)}$ in the $b\bar{b}$ system. This equality would follow from a *light-quark spin symmetry* proposed by Voloshin [28]. It implies that Z'_b is an equal-amplitude superposition of spin-singlet and spin-triplet, which prevents its decay into $B^*\bar{B}$ or $B\bar{B}^*$. Masses of




Figure 4. Energy levels for the multiplets of isospin-1 hidden-bottom tetraquark mesons associated with the 1^{--} and 0^{++} $\mathbf{8}$ -mesons. The masses of $T_{bb1}^+(10610)$ and $T_{bb1}^+(10650)$ were used as inputs to determine the energies of the $\mathbf{8}$ -mesons. The dashed horizontal lines are the thresholds for bottom-meson pairs.

Z_b and Z'_b near the $B^* \bar{B}$ and $B^* \bar{B}^*$ thresholds can be explained by a further fine-tuning of $E_{1^{--}}^{(1)}$ and $E_{0^{++}}^{(1)}$ so that $\varepsilon_1^{(1)} = \varepsilon_0^{(1)} = 0$. Given these fine tunings, we can predict the masses of Z_b and Z'_b and their B-O partners to first order in Δ_b . The Z_b and Z'_b are near the $B^* \bar{B}$ and $B^* \bar{B}^*$ thresholds. The 1^{++} and 2^{++} partners are also near the $B^* \bar{B}$ and $B^* \bar{B}^*$ thresholds. The two 0^{++} partners are near the $B \bar{B}$ and $B^* \bar{B}$ thresholds. The energy levels in the multiplets including Z_b and Z'_b are illustrated in Fig. 4.

Discussion. A B-O potential for hidden-heavy hadrons that approaches a $\mathbf{8}$ -potential at small r must connect to a $(\mathbf{3}+\mathbf{3}^*)$ -potential at large r . Only those $\mathbf{8}$ -potentials associated with the lowest-energy $\mathbf{8}$ -hadrons cross below the $(\mathbf{3}+\mathbf{3}^*)$ -threshold before approaching it, in which case they can support bound states and/or resonances, which we identify as exotic hidden-heavy hadrons. This explains why most exotic hidden-heavy hadrons have energies near heavy-hadron-pair thresholds.

This also provides insight into the remarkable properties of some of the hidden-heavy tetraquark mesons. The proximity of X_c to the $D^* \bar{D}$ threshold comes from the fine tuning of an isospin-0 $\mathbf{8}$ -meson energy. The suppression of decays of Z'_b into $B^* \bar{B}$ and $B \bar{B}^*$ and the proximity of Z_b and Z'_b to the $B^* \bar{B}$ and $B^* \bar{B}^*$ thresholds come from fine tunings of two isospin-1 $\mathbf{8}$ -meson energies.

We estimated the energies of the B-O partners of X_c , X_b , Z_b , and Z'_b to first order in Δ . Accurate predictions would require solving the coupled-channel Schrödinger equations including the spin-splitting matrix V_{SS} . In the isospin-0 case, it is also necessary to take into account the narrow avoided crossing between the Σ_g^+ and $\Sigma_g^{+'}$

potentials [29].

Similar methods can be applied to hidden-heavy pentaquark baryons. They are bound states or resonances in $\mathbf{8}$ -potentials that cross below the threshold for the pair of a $\mathbf{3}$ -baryon and a $\mathbf{3}^*$ -meson before approaching it. The ground-state pentaquark multiplet could be a triplet associated with a $J^P = \frac{1}{2}^+$ $\mathbf{8}$ -baryon or a quartet associated with a $\frac{3}{2}^+$ $\mathbf{8}$ -baryon [8].

The continuity constraint has important implications for lattice-QCD calculations of B-O potentials. The operators used previously in lattice QCD with light quarks produce states at small r with very small overlaps with $\mathbf{8}$ -hadrons [16, 30–36]. Much larger overlaps would be obtained with operators that connect the $\mathbf{3}$ and $\mathbf{3}^*$ sources by Wilson lines to a junction with a color-octet QCD operator. Until lattice-QCD calculations of the B-O potentials are available, B-O calculations of exotic hidden-heavy hadrons will have to rely on model potentials. Since $\mathbf{8}$ -potentials must connect to $(\mathbf{3}+\mathbf{3}^*)$ -potentials, any model reduces to the interpolation between the large- r and small- r regions. This dramatically reduces the model dependence of the predictions.

Physics beyond the Standard Model may include long-lived heavy particles with $\mathbf{8}$ color charges, such as the gluino in some supersymmetric models. After being created in a collider, a gluino can hadronize before it decays. The ATLAS and CMS collaborations have searched for long-lived gluinos at the Large Hadron Collider [37–44]. The analyses depend on assumptions about the gluino-hadron spectrum [45–49], which is the $\mathbf{8}$ -hadron spectrum up to an additive constant associated with the gluino mass. The exotic hidden-heavy hadrons provide constraints on that spectrum. The masses of X_c and Z_b suggest that the lightest gluino-meson is isospin-0 but isospin-1 is not much heavier. The masses of the $c\bar{c}$ pentaquarks $P_{cc}^+(4312)$ discovered in 2019 [50] and $P_{ccs}^+(4338)$ discovered in 2023 [51] suggest that the lightest gluino-baryon is isospin- $\frac{1}{2}$ but isospin-0 is not much heavier. Quantitative B-O analyses of exotic hidden-heavy hadrons that take into account heavy-hadron spin splittings could determine the lowest-energy states in the $\mathbf{8}$ -hadron spectrum and enable more accurate constraints on long-lived gluinos.

This work was supported in part by the U.S. Department of Energy under grant DE-SC0011726. EB would like to thank A. Mohapatra for illuminating discussions that led to the realization that $\mathbf{8}$ -potentials must connect to $(\mathbf{3}+\mathbf{3}^*)$ -potentials. EB would also like to thank N. Brambilla and A. Vairo for hospitality and discussions during two extended visits to Technical University of Munich. This work contributes to the goals of the US DOE ExoHad Topical Collaboration, Contract DE-SC0023598.

- * braaten.1@osu.edu
† bruschini.1@osu.edu
- [1] N. Brambilla, S. Eidelman, C. Hanhart, A. Nefediev, C.-P. Shen, C. E. Thomas, A. Vairo, and C.-Z. Yuan, The XYZ states: experimental and theoretical status and perspectives, *Phys. Rep.* **873**, 1 (2020), arXiv:1907.07583 [hep-ex].
 - [2] R. F. Lebed, Theory of the (Heavy-Quark) Exotic Hadrons: A Primer for the Flavor Community, *PoS FPCP2023*, 028 (2023), arXiv:2308.00781 [hep-ph].
 - [3] M. Berwein, N. Brambilla, J. Tarrús Castellà, and A. Vairo, Quarkonium Hybrids with Nonrelativistic Effective Field Theories, *Phys. Rev. D* **92**, 114019 (2015), arXiv:1510.04299 [hep-ph].
 - [4] R. Onocala and J. Soto, Heavy Quarkonium Hybrids: Spectrum, Decay and Mixing, *Phys. Rev. D* **96**, 014004 (2017), arXiv:1702.03900 [hep-ph].
 - [5] N. Brambilla, G. Krein, J. Tarrús Castellà, and A. Vairo, Born-Oppenheimer approximation in an effective field theory language, *Phys. Rev. D* **97**, 016016 (2018), arXiv:1707.09647 [hep-ph].
 - [6] J. Soto and J. Tarrús Castellà, Nonrelativistic effective field theory for heavy exotic hadrons, *Phys. Rev. D* **102**, 014012 (2020), arXiv:2005.00552 [hep-ph].
 - [7] E. Braaten and R. Bruschini, Model-independent predictions for decays of hidden-heavy hadrons into pairs of heavy hadrons, *Phys. Rev. D* **109**, 094051 (2024), arXiv:2403.12868 [hep-ph].
 - [8] M. Berwein, N. Brambilla, A. Mohapatra, and A. Vairo, Hybrids, tetraquarks, pentaquarks, doubly heavy baryons, and quarkonia in Born-Oppenheimer effective theory, *Phys. Rev. D* **110**, 094040 (2024), arXiv:2408.04719 [hep-ph].
 - [9] R. Bruschini, Model-independent predictions for decays of double-heavy hadrons into pairs of heavy hadrons, *Phys. Rev. D* **110**, 074033 (2024), arXiv:2408.05150 [hep-ph].
 - [10] P. Bicudo, K. Cichy, A. Peters, and M. Wagner, BB interactions with static bottom quarks from Lattice QCD, *Phys. Rev. D* **93**, 034501 (2016), arXiv:1510.03441 [hep-lat].
 - [11] M. Foster and C. Michael (UKQCD Collaboration), Hadrons with a heavy color adjoint particle, *Phys. Rev. D* **59**, 094509 (1999), arXiv:hep-lat/9811010.
 - [12] J. Herr, C. Schlosser, and M. Wagner, Gluelump masses and mass splittings from SU(3) lattice gauge theory, *Phys. Rev. D* **109**, 034516 (2024), arXiv:2306.09902 [hep-lat].
 - [13] K. Marsh and R. Lewis, A lattice QCD study of generalized gluelumps, *Phys. Rev. D* **89**, 014502 (2014), arXiv:1309.1627 [hep-lat].
 - [14] (ATLAS Collaboration) Generation and Simulation of R -Hadrons in the ATLAS Experiment, ATLAS-CONF-2019-019.
 - [15] J. D. Stack, The Heavy Quark Potential in SU(3) Lattice Gauge Theory, *Phys. Rev. D* **29**, 1213 (1984).
 - [16] G. S. Bali, H. Neff, T. Düssel, T. Lippert, and K. Schilling (SESAM Collaboration), Observation of string breaking in QCD, *Phys. Rev. D* **71**, 114513 (2005), arXiv:hep-lat/0505012.
 - [17] K. J. Juge, J. Kuti, and C. J. Morningstar, Ab Initio Study of Hybrid $\bar{b}gb$ Mesons, *Phys. Rev. Lett.* **82**, 4400 (1999), arXiv:hep-ph/9902336.
 - [18] K. J. Juge, J. Kuti, and C. Morningstar, Fine structure of the QCD string spectrum, *Phys. Rev. Lett.* **90**, 161601 (2003), arXiv:hep-lat/0207004.
 - [19] N. Brambilla, A. Pineda, J. Soto, and A. Vairo, Potential NRQCD: An Effective theory for heavy quarkonium, *Nucl. Phys. B* **566**, 275 (2000), arXiv:hep-ph/9907240.
 - [20] A. Bondar *et al.* (Belle Collaboration), Observation of two charged bottomonium-like resonances in $\Upsilon(5S)$ decays, *Phys. Rev. Lett.* **108**, 122001 (2012), arXiv:1110.2251 [hep-ex].
 - [21] M. Ablikim *et al.* (BESIII Collaboration), Observation of a Charged Charmoniumlike Structure in $e^+e^- \rightarrow \pi^+\pi^-J/\psi$ at $\sqrt{s} = 4.26$ GeV, *Phys. Rev. Lett.* **110**, 252001 (2013), arXiv:1303.5949 [hep-ex].
 - [22] Z. Q. Liu *et al.* (Belle Collaboration), Study of $e^+e^- \rightarrow \pi^+\pi^-J/\psi$ and Observation of a Charged Charmoniumlike State at Belle, *Phys. Rev. Lett.* **110**, 252002 (2013), [Erratum: *Phys.Rev.Lett.* 111, 019901 (2013)], arXiv:1304.0121 [hep-ex].
 - [23] E. Braaten, How the $Z_c(3900)$ Reveals the Spectra of Quarkonium Hybrid and Tetraquark Mesons, *Phys. Rev. Lett.* **111**, 162003 (2013), arXiv:1305.6905 [hep-ph].
 - [24] E. Braaten, C. Langmack, and D. H. Smith, Born-Oppenheimer approximation for the XYZ mesons, *Phys. Rev. D* **90**, 014044 (2014), arXiv:1402.0438 [hep-ph].
 - [25] F. Alasiri, E. Braaten, and A. Mohapatra, Born-Oppenheimer potentials for SU(3) gauge theory, *Phys. Rev. D* **110**, 054029 (2024), arXiv:2406.05123 [hep-ph].
 - [26] S.-K. Choi *et al.* (Belle Collaboration), Observation of a Narrow Charmoniumlike State in Exclusive $B^\pm \rightarrow K^\pm\pi^+\pi^-J/\psi$ Decays, *Phys. Rev. Lett.* **91**, 262001 (2003), arXiv:hep-ex/0309032.
 - [27] S. Navas *et al.* (Particle Data Group), Review of Particle Physics, *Phys. Rev. D* **110**, 030001 (2024).
 - [28] M. B. Voloshin, Light Quark Spin Symmetry in Z_b Resonances?, *Phys. Rev. D* **93**, 074011 (2016), arXiv:1601.02540 [hep-ph].
 - [29] N. Brambilla, A. Mohapatra, T. Scirpa, and A. Vairo, The nature of $\chi_{c1}(3872)$ and $T_{cc}^+(3875)$, arXiv:2411.14306 [hep-ph] (2024).
 - [30] G. S. Bali, B. Bolder, N. Eicker, T. Lippert, B. Orth, P. Ueberholz, K. Schilling, and T. Struckmann (TXL and T(X)L Collaborations), Static potentials and glueball masses from QCD simulations with Wilson sea quarks, *Phys. Rev. D* **62**, 054503 (2000), arXiv:hep-lat/0003012.
 - [31] M. Alberti, G. S. Bali, S. Collins, F. Knechtli, G. Moir, and W. Söldner, Hadroquarkonium from lattice QCD, *Phys. Rev. D* **95**, 074501 (2017), arXiv:1608.06537 [hep-lat].
 - [32] J. Bulava, B. Hörz, F. Knechtli, V. Koch, G. Moir, C. Morningstar, and M. Peardon, String breaking by light and strange quarks in QCD, *Phys. Lett. B* **793**, 493 (2019), arXiv:1902.04006 [hep-lat].
 - [33] P. Bicudo, M. Cardoso, N. Cardoso, and M. Wagner, Bottomonium resonances with $I = 0$ from lattice QCD correlation functions with static and light quarks, *Phys. Rev. D* **101**, 034503 (2020), arXiv:1910.04827 [hep-lat].
 - [34] S. Prelovsek, H. Bahtiyar, and J. Petkovic, Z_b tetraquark channel from lattice QCD and Born-Oppenheimer approximation, *Phys. Lett. B* **805**, 135467 (2020), arXiv:1912.02656 [hep-lat].
 - [35] M. Sadl and S. Prelovsek, Tetraquark systems $\bar{b}b\bar{d}u$ in the

- static limit and lattice QCD, *Phys. Rev. D* **104**, 114503 (2021), [arXiv:2109.08560 \[hep-lat\]](#).
- [36] J. Bulava, F. Knechtli, V. Koch, C. Morningstar, and M. Peardon, The quark-mass dependence of the potential energy between static colour sources in the QCD vacuum with light and strange quarks, *Phys. Lett. B* **854**, 138754 (2024), [arXiv:2403.00754 \[hep-lat\]](#).
- [37] V. Khachatryan *et al.* (CMS Collaboration), Search for Stopped Gluinos in pp collisions at $\sqrt{s} = 7$ TeV, *Phys. Rev. Lett.* **106**, 011801 (2011), [arXiv:1011.5861 \[hep-ex\]](#).
- [38] G. Aad *et al.* (ATLAS Collaboration), Search for stable hadronising squarks and gluinos with the ATLAS experiment at the LHC, *Phys. Lett. B* **701**, 1 (2011), [arXiv:1103.1984 \[hep-ex\]](#).
- [39] G. Aad *et al.* (ATLAS Collaboration), Search for heavy long-lived charged particles with the ATLAS detector in pp collisions at $\sqrt{s} = 7$ TeV, *Phys. Lett. B* **703**, 428 (2011), [arXiv:1106.4495 \[hep-ex\]](#).
- [40] S. Chatrchyan *et al.* (CMS Collaboration), Search for Heavy Long-Lived Charged Particles in pp Collisions at $\sqrt{s} = 7$ TeV, *Phys. Lett. B* **713**, 408 (2012), [arXiv:1205.0272 \[hep-ex\]](#).
- [41] G. Aad *et al.* (ATLAS Collaboration), Searches for heavy long-lived sleptons and R-Hadrons with the ATLAS detector in pp collisions at $\sqrt{s} = 7$ TeV, *Phys. Lett. B* **720**, 277 (2013), [arXiv:1211.1597 \[hep-ex\]](#).
- [42] M. Aaboud *et al.* (ATLAS Collaboration), Search for heavy long-lived charged R -hadrons with the ATLAS detector in 3.2 fb^{-1} of proton–proton collision data at $\sqrt{s} = 13$ TeV, *Phys. Lett. B* **760**, 647 (2016), [arXiv:1606.05129 \[hep-ex\]](#).
- [43] M. Aaboud *et al.* (ATLAS Collaboration), Search for heavy charged long-lived particles in proton-proton collisions at $\sqrt{s} = 13$ TeV using an ionisation measurement with the ATLAS detector, *Phys. Lett. B* **788**, 96 (2019), [arXiv:1808.04095 \[hep-ex\]](#).
- [44] A. M. Sirunyan *et al.* (CMS Collaboration), Search for long-lived particles using nonprompt jets and missing transverse momentum with proton-proton collisions at $\sqrt{s} = 13$ TeV, *Phys. Lett. B* **797**, 134876 (2019), [arXiv:1906.06441 \[hep-ex\]](#).
- [45] H. Baer, K. Cheung, and J. F. Gunion, A Heavy gluino as the lightest supersymmetric particle, *Phys. Rev. D* **59**, 075002 (1999), [arXiv:hep-ph/9806361](#).
- [46] J. L. Hewett, B. Lillie, M. Masip, and T. G. Rizzo, Signatures of long-lived gluinos in split supersymmetry, *JHEP* **09** (2004), 070, [arXiv:hep-ph/0408248](#).
- [47] A. Arvanitaki, S. Dimopoulos, A. Pierce, S. Rajendran, and J. G. Wacker, Stopping gluinos, *Phys. Rev. D* **76**, 055007 (2007), [arXiv:hep-ph/0506242](#).
- [48] G. R. Farrar, R. Mackeprang, D. Milstead, and J. P. Roberts, Limit on the mass of a long-lived or stable gluino, *JHEP* **02** (2011), 018, [arXiv:1011.2964 \[hep-ph\]](#).
- [49] S. Raby, Long-Lived Gluinos and Stable Axinos, *Phys. Rev. Lett.* **115**, 231801 (2015), [arXiv:1508.04373 \[hep-ph\]](#).
- [50] R. Aaij *et al.* (LHCb Collaboration), Observation of a narrow pentaquark state, $P_c(4312)^+$, and of two-peak structure of the $P_c(4450)^+$, *Phys. Rev. Lett.* **122**, 222001 (2019), [arXiv:1904.03947 \[hep-ex\]](#).
- [51] R. Aaij *et al.* (LHCb Collaboration), Observation of a $J/\psi\Lambda$ Resonance Consistent with a Strange Pentaquark Candidate in $B^- \rightarrow J/\psi\Lambda\bar{p}$ Decays, *Phys. Rev. Lett.* **131**, 031901 (2023), [arXiv:2210.10346 \[hep-ex\]](#).