QUANTITATIVE PERIODIC HOMOGENIZATION FOR SYMMETRIC NON-LOCAL STABLE-LIKE OPERATORS

XIN CHEN, ZHEN-QING CHEN, TAKASHI KUMAGAI AND JIAN WANG

ABSTRACT. Homogenization for non-local operators in periodic environments has been studied intensively. So far, these works are mainly devoted to the qualitative results, that is, to determine explicitly the operators in the limit. To the best of authors' knowledge, there is no result concerning the convergence rates of the homogenization for stable-like operators in periodic environments. In this paper, we establish a quantitative homogenization result for symmetric α -stable-like operators on \mathbb{R}^d with periodic coefficients. In particular, we show that the convergence rate for the solutions of associated Dirichlet problems on a bounded domain D is of order

$$\varepsilon^{(2-\alpha)/2} \mathbb{1}_{\{\alpha \in (1,2)\}} + \varepsilon^{\alpha/2} \mathbb{1}_{\{\alpha \in (0,1)\}} + \varepsilon^{1/2} |\log \varepsilon|^2 \mathbb{1}_{\{\alpha=1\}},$$

while, when the solution to the equation in the limit is in $C_c^2(D)$, the convergence rate becomes

 $\varepsilon^{2-\alpha} \mathbb{1}_{\{\alpha \in (1,2)\}} + \varepsilon^{\alpha} \mathbb{1}_{\{\alpha \in (0,1)\}} + \varepsilon |\log \varepsilon|^2 \mathbb{1}_{\{\alpha = 1\}}.$

This indicates that the boundary decay behaviors of the solution to the equation in the limit affects the convergence rate in the homogenization.

Keywords: homogenization; symmetric α -stable-like operator; periodic coefficient; Dirichlet problem MSC 2010: 60G51; 60G52; 60J25; 60J75.

1. INTRODUCTION

This paper is devoted to quantitative results for homogenization of symmetric (non-local) stable-like operators in deterministic periodic media. In line with most of literature (e.g., [12, 29, 30, 41, 43]), we restrict ourselves to the Euclidean space setting. Consider the following symmetric stable-like operator \mathcal{L} on \mathbb{R}^d :

$$\begin{split} \mathcal{L}f(x) &:= \text{p.v.} \int_{\mathbb{R}^d} (f(y) - f(x)) \frac{K(x,y)}{|x-y|^{d+\alpha}} \, dy \\ &:= \lim_{\varepsilon \to 0} \int_{\mathbb{R}^d} \mathbf{1}_{\{|y-x| > \varepsilon\}} (f(y) - f(x)) \frac{K(x,y)}{|x-y|^{d+\alpha}} \, dy \end{split}$$

for $f \in C_c^2(\mathbb{R}^d)$. Here $\alpha \in (0, 2)$, and $K \in C^2(\mathbb{R}^d \times \mathbb{R}^d)$ is multivariate 1-periodic on $\mathbb{R}^d \times \mathbb{R}^d$ (that is, it can be viewed as a bounded C^2 -smooth function defined on $\mathbb{T}^d \times \mathbb{T}^d$, where $\mathbb{T}^d := \mathbb{R}^d / \mathbb{Z}^d$ is the *d*-dimensional torus) so that

$$K(x,y) = K(y,x)$$
 and $\Lambda^{-1} \leq K(x,y) \leq \Lambda$ for $x, y \in \mathbb{R}^d$ (1.1)

with some constant $\Lambda \ge 1$. The symmetric non-local operator \mathcal{L} generates a symmetric bilinear form for $u, v \in C^2_c(\mathbb{R}^d)$:

$$\mathcal{E}(u,v) = \frac{1}{2} \iint_{\mathbb{R}^d \times \mathbb{R}^d} (u(x) - u(y))(v(x) - v(y)) \frac{K(x,y)}{|x - y|^{d + \alpha}} \, dx \, dy. \tag{1.2}$$

Denote by \mathcal{F} the completion of $C^2_c(\mathbb{R}^d)$ under the norm

$$\|u\|_{\alpha/2,2} := \left(\iint_{\mathbb{R}^d \times \mathbb{R}^d} \frac{(u(x) - u(y))^2}{|x - y|^{d + \alpha}} \, dx \, dy + \int_{\mathbb{R}^d} u(x)^2 \, dx\right)^{1/2}.$$
(1.3)

It is easy to check that

$$\mathcal{F} = W^{\alpha/2,2}(\mathbb{R}^d) = \left\{ u \in L^2(\mathbb{R}^d; dx) : \iint_{\mathbb{R}^d \times \mathbb{R}^d} \frac{(u(x) - u(y))^2}{|x - y|^{d + \alpha}} \, dx \, dy < \infty \right\}$$
(1.4)

and $(\mathcal{E}, \mathcal{F})$ is a regular Dirichlet form on $L^2(\mathbb{R}^d; dx)$. Denote by $(\tilde{\mathcal{L}}, \text{Dom}(\tilde{\mathcal{L}}))$ the L^2 -generator of $(\mathcal{E}, \mathcal{F})$. Then $C_c^2(\mathbb{R}^d) \subset \text{Dom}(\tilde{\mathcal{L}})$ and $\tilde{\mathcal{L}}f = \mathcal{L}f$ for $f \in C_c^2(\mathbb{R}^d)$. It is known from [17, 18] that there is a symmetric conservative irreducible Feller process $X := \{X_t\}_{t \ge 0}$ on \mathbb{R}^d associated with the regular Dirichlet form $(\mathcal{E}, \mathcal{F})$ on $L^2(\mathbb{R}^d; dx)$, which has the strong Feller property and can start from any $x \in \mathbb{R}^d$. In the literature, X is called a symmetric α -stable-like process. Since the coefficient K(x, y) is multivariate 1-periodic on $\mathbb{R}^d \times \mathbb{R}^d$, one can regard X as a symmetric \mathbb{T}^d -valued process. It is exponentially ergodic and has the normalized Lebesgue measure on \mathbb{T}^d as its unique invariant probability measure.

For any $\varepsilon > 0$, set

$$X^{\varepsilon} := \{X_t^{\varepsilon}\}_{t \ge 0} := \{\varepsilon X_{\varepsilon^{-\alpha}t}\}_{t \ge 0}$$

Clearly, X^{ε} is a symmetric \mathbb{R}^d -valued Feller process having the strong Feller property with associated Dirichlet form $(\mathcal{E}^{\varepsilon}, W^{\alpha/2,2}(\mathbb{R}^d))$ on $L^2(\mathbb{R}^d; dx)$, where

$$\mathcal{E}^{\varepsilon}(u,v) = \frac{1}{2} \iint_{\mathbb{R}^d \times \mathbb{R}^d} (u(x) - u(y))(v(x) - v(y)) \frac{K(x/\varepsilon, y/\varepsilon)}{|x - y|^{d + \alpha}} \, dx \, dy. \tag{1.5}$$

Denote by $(\mathcal{L}_{\varepsilon}, \operatorname{Dom}(\mathcal{L}_{\varepsilon}))$ the L^2 -generator of $(\mathcal{E}^{\varepsilon}, W^{\alpha/2,2}(\mathbb{R}^d))$. Then $C^2_c(\mathbb{R}^d) \subset \operatorname{Dom}(\mathcal{L}_{\varepsilon})$ and

$$\mathcal{L}_{\varepsilon}f(x) := \text{p.v.} \int_{\mathbb{R}^d} (f(y) - f(x)) \frac{K(x/\varepsilon, y/\varepsilon)}{|x - y|^{d + \alpha}} \, dy, \quad f \in C^2_c(\mathbb{R}^d).$$
(1.6)

Define

$$\bar{K} := \iint_{\mathbb{T}^d \times \mathbb{T}^d} K(x, y) \, dx \, dy. \tag{1.7}$$

Let $\bar{X} := \{\bar{X}_t\}_{t \ge 0}$ be a rotationally symmetric α -stable Lévy process on \mathbb{R}^d with Lévy measure $\bar{K}|z|^{-(d+\alpha)}dz$. Denote the infinitesimal generator of \bar{X} by $(\bar{\mathcal{L}}, \operatorname{Dom}(\bar{\mathcal{L}}))$. Note that $C_c^2(\mathbb{R}^d) \subset \operatorname{Dom}(\bar{\mathcal{L}})$ and

$$\bar{\mathcal{L}}f(x) := \text{p.v.} \int_{\mathbb{R}^d} (f(y) - f(x)) \frac{\bar{K}}{|y - x|^{d + \alpha}} \, dy \quad \text{for } f \in C_c^2(\mathbb{R}^d).$$
(1.8)

One can deduce from [6, Theorem 4.1] that X^{ε} converges weakly to \bar{X} as $\varepsilon \to 0$. Alternatively, this weak convergence can also be established from the uniform heat kernel estimates in [17, Theorem 1.1] for symmetric α -stable-like processes $\{X^{\varepsilon} : \varepsilon \in (0, 1]\}$ and the L^2 -convergence of the resolvents of X^{ε} in [32, Theorem 1.1] as $\varepsilon \to 0$. We thus call $\bar{\mathcal{L}}$ the homogenized operator of $\mathcal{L}_{\varepsilon}$. The reader may refer to [11, 12] and the references therein for more details on homogenization of symmetric stable-like processes in stationary ergodic media and periodic homogenization problem of non-symmetric Lévy-type processes.

Throughout the paper, unless otherwise specified, D is a bounded $C^{1,1}$ domain in \mathbb{R}^d . For any $x \in D$, let $\delta_D(x) := \operatorname{dist}(x, \partial D)$ be the distance from x to the boundary ∂D . For any $\theta \in \mathbb{R}$, $k \in \mathbb{N}$ and $0 < \gamma \notin \mathbb{N}$, define

$$[f]_{k;D}^{(\theta)} := \sup_{x \in D} \left(\delta_D(x)^{k+\theta} |\nabla^k f(x)| \right), \quad [f]_{\gamma;D}^{(\theta)} := \sup_{x,y \in D} \left(\min\left(\delta_D(x), \delta_D(y) \right)^{\gamma+\theta} \frac{|\nabla^{[\gamma]} f(x) - \nabla^{[\gamma]} f(y)|}{|x - y|^{\gamma - [\gamma]}} \right), \\ C_{(\theta)}^{\gamma}(D) := \left\{ f \in C(\mathbb{R}^d) : \|f\|_{\gamma;D}^{(\theta)} := [f]_{0;D}^{(\theta)} + [f]_{\gamma;D}^{(\theta)} < \infty \text{ and } f(x) = 0 \text{ for } x \in D^c \right\}.$$

Here $[\gamma]$ denotes the largest integer not exceeding γ . Let $\mathcal{L}_{\varepsilon}$ and $\overline{\mathcal{L}}$ be the operator defined by (1.6) and (1.8), respectively. Given $h \in C^{\beta}_{(\alpha/2)}(D) \cap C(\overline{D})$ with $\alpha + \beta > 4$ so that neither β nor $\alpha + \beta$ is an integer, consider the following Dirichlet problem

$$\begin{cases} \mathcal{L}_{\varepsilon} u_{\varepsilon}(x) = h(x), & x \in D, \\ u_{\varepsilon}(x) = 0, & x \in D^{c}. \end{cases}$$
(1.9)

Similarly, consider the following Dirichlet problem for the homogenized operator $\overline{\mathcal{L}}$:

$$\begin{cases} \bar{\mathcal{L}}\bar{u}(x) = h(x), & x \in D, \\ \bar{u}(x) = 0, & x \in D^c. \end{cases}$$
(1.10)

According to [40, Proposition 1.4 and Theorem 1.5], the equation (1.10) has the unique solution $\bar{u} \in C^{\alpha+\beta}_{(-\alpha/2)}(D) \cap C(\bar{D})$. On the other hand, according to Lemma 2.3 below, since D is bounded and $h \in C(\bar{D})$, the equation (1.9) has the unique bounded solution $u_{\varepsilon} \in \text{Dom}(\mathcal{L}^{D}_{\varepsilon}) \cap L^{1}(D; dx)$, where $\text{Dom}(\mathcal{L}^{D}_{\varepsilon})$ is the domain of the operator $\mathcal{L}_{\varepsilon}$ with zero Dirichlet condition on D^{c} . Moreover, when the domain D is more

regular, higher order regularity on u_{ε} will be obtained. The reader is referred to [1, Theorem 3.1] and [51, Theorem 1.3] for more details on regularity estimates for solutions to the Dirichlet problem of stable-like operators. The following is the main result of the paper, which establishes an explicit convergence rate of the solution u_{ε} of (1.9) in $L^1(D; dx)$ as $\varepsilon \to 0$.

Theorem 1.1. Suppose that D is a bounded $C^{1,1}$ domain in \mathbb{R}^d . Let u_{ε} and \bar{u} be the unique solutions of (1.9) and (1.10) respectively for some $h \in C^{\beta}_{(\alpha/2)}(D)$ with $\alpha + \beta > 4$, where neither β nor $\alpha + \beta$ is an integer. Then there exists a constant $C_1 > 0$ so that for all $\varepsilon \in (0, 1)$,

$$\|u_{\varepsilon} - \bar{u}\|_{L^{1}(D;dx)} \leqslant C_{1} \begin{cases} \varepsilon^{\alpha/2}, & \alpha \in (0,1), \\ \varepsilon^{1/2}(1+|\log \varepsilon|^{2}), & \alpha = 1, \\ \varepsilon^{(2-\alpha)/2}, & \alpha \in (1,2). \end{cases}$$
(1.11)

As seen from the proof of Theorem 1.1 below, much more efforts are devoted to handling blow up behaviours for $\nabla \bar{u}$ and $\nabla^2 \bar{u}$ near the boundary ∂D , as well as their interactions with the non-local operator $\mathcal{L}_{\varepsilon}$. Indeed, if the solution \bar{u} of the homogenized equation (1.10) is C^2 -smooth with compact support in D (which roughly indicates that we do not need to take care of the boundary behaviours of \bar{u}), then the convergence rates of u_{ε} to \bar{u} will be better than those in (1.11).

Now we consider (1.9) and (1.10) for $h \in C_b^1(\overline{D})$ that does not need to vanish on ∂D but under the assumption that $\overline{u} \in C_c^2(D)$.

Theorem 1.2. Assume that D is a bounded Lipschitz domain in \mathbb{R}^d . Suppose that u_{ε} and \bar{u} are the unique solutions of (1.9) and (1.10), respectively, for some $h \in C_b^1(\bar{D})$. If $\bar{u} \in C_c^2(D)$, then there exists a constant $C_2 > 0$ such that for all $\varepsilon \in (0, 1)$,

$$\|u_{\varepsilon} - \bar{u}\|_{L^{2}(D;dx)} \leqslant C_{2} \begin{cases} \varepsilon^{\alpha}, & \alpha \in (0,1), \\ \varepsilon(1+|\log \varepsilon|^{2}), & \alpha = 1, \\ \varepsilon^{2-\alpha}, & \alpha \in (1,2). \end{cases}$$
(1.12)

The homogenization problem for differential operators with periodic coefficients has been investigated extensively in the literature. The reader is referred to the monographs [4, 7, 31, 44]. Obtaining convergence rate is an important problem in the homogenization theory. The first results in this direction for boundary value problems of elliptic operators with periodic coefficients in L^2 -estimates are due to [7] and [36, 37]. Since then, there are a lot of developments in this topic, see [44] and the references therein for more details. In particular, we want to emphasize that the following sharp estimate for elliptic differential operators (which is comparable with the classical Berry-Esseen bounds for i.i.d. random variables)

$$\|u_{\varepsilon} - \bar{u}\|_{L^2(D;dx)} \leqslant C_3 \varepsilon \tag{1.13}$$

has been established in [7, 24, 33, 34, 35, 45, 48] under different boundary conditions (including the Dirichlet boundary condition and the Neumann boundary condition) and regularity conditions. Here u_{ε} (resp. \bar{u}) in (1.13) is the unique solution to (1.9) (resp. (1.10)) with $\mathcal{L}_{\varepsilon}$ (resp. $\bar{\mathcal{L}}$) being a second-order differential elliptic operator with rapidly oscillating and periodic coefficients (resp. constant coefficients). See also [8, 9, 27, 28, 47, 52, 53] for related results. We note that there is some remarkable progress over the past decade on quantitative stochastic homogenization of elliptic and parabolic equations (for differential operators); see for instance [2, 3, 25, 26].

In recent years, the homogenization for non-local operators in periodic environments has also been studied intensively, see [12, 29, 30, 32, 39, 41, 43, 42, 49]. In all these papers only the qualitative homogenization results for non-local operators are investigated; that is, these papers determine the limit operators and establish the convergence of the processes or the solutions of the non-local partial differential equations to those of the limit operators. The only existing literature we know on quantitative homogenization of nonlocal operators is [38], where the L^2 -convergence rate is obtained by a spectral method for 1-resolvents of a class of non-local operators of convolution type having L^2 integrable jumping kernels. We note that the limit operators in [38] are second order elliptic differential operators, rather than non-local operators, and that the rate of convergence for the associated Dirichlet problem is not investigated in [38]. To the best of our knowledge, there is no result concerning convergence rates of the homogenization for stable-like operators in periodic environments. In this paper we will address this problem for symmetric α -stable-like operators. It is interesting to observe the difference between our results (1.11) and (1.12) for non-local operators and (1.13) for differential operators as well as to the non-local operator investigated in [38] where the jumping kernel has finite second moment. We also want to mention that the interior convergence rate for symmetric α -stable-like operators in i.i.d. environments, where the boundary behavior of the solution was not taken into account, has been obtained in the paper [13].

Below are some additional comments on the results of this paper.

- (1) The interior convergence rate (1.12) seems to be optimal, since it is in line with the Berry-Esseen estimates for i.i.d. random variables having stable laws as limit, see [5, 10, 50] for example. We also believe that the convergence rates in (1.11) for the associated Dirichlet problem are near optimal for the methods used in this paper, due to the blow up behaviors of $\nabla \bar{u}$ and $\nabla^2 \bar{u}$ near the boundary ∂D . These blow ups make the convergence rate slower than (1.12) which can be viewed as the interior convergence rate, even if the domain D is smooth. This is different from the case for the Dirichlet problem of differential operators as limit equations, where \bar{u} are known to decay at the boundary in a linear rate and so ∇u and $\nabla^2 u$ are all bounded, see e.g. [24, 33, 38, 44] for more details.
- (2) We only give the L^1 -convergence rate in (1.11). From (1.11) and Lemma 2.3 below, we could deduce the L^2 -convergence rate from the inequality

$$\|u_{\varepsilon} - \bar{u}\|_{L^{2}(D;dx)} \leq C_{4} \|h\|_{\infty}^{1/2} \|u_{\varepsilon} - \bar{u}\|_{L^{1}(D;dx)}^{1/2}$$

but we do not think the resulting L^2 -convergence rate is optimal. As illustrated in [1, 40, 51], it seems difficult to verify the H^2 -estimate $\int_D |\nabla^2 \bar{u}(x)|^2 dx < \infty$ for the fractional Laplacian operator, which is a key point to establish the L^2 -convergence for differential operators in [24, 33]. Because of the lack of such kind of the H^2 -estimate, it does not seem easy to apply the integration by parts formula to establish a sharper L^2 -convergence rate. On the other hand, ignoring the boundary behaviors of \bar{u} , we can obtain the L^2 -interior convergence rate (1.12). Indeed, in the proof of (1.12) we can use the integration by parts formula to reduce the requirements into the required regularity of \bar{u} up to C^2 , while in our proof of (1.11) higher order regularities on \bar{u} are needed.

(3) Compared with the approach for the quantitative homogenization of differential operators, there are two essential different ingredients in the proof of Theorem 1.1. One is the quantitative estimates (2.41) and (2.44) below for the average in z-variable in the homogenization problem of α -stablelike operators, which represents the jumping size of the associated stable-like process. However, for differential operators only the averaging properties of the position variable x are needed. The other is to give detailed estimates on the interactions between blow up behaviors of $\nabla \bar{u}$ and $\nabla^2 \bar{u}$ near ∂D and the effect of the non-local operator $\mathcal{L}_{\varepsilon}$, see Lemmas 2.4–2.7 below for details. In particular, in the arguments of two ingredients above, it seems natural to use the $C_{(-\alpha/2)}^{\alpha+\beta}$ -norm (with suitable choice of $\beta > 0$) instead of the H^2 -norm for \bar{u} , and the constant C_1 in (1.11) only depends on $\sup_{0 \le k \le 4} \sup_{x \in D} (\delta_D(x)^{k-\alpha/2} |\nabla^k \bar{u}(x)|).$

The rest of this paper is arranged as follows. We present in Section 2 some preliminaries that are needed later in this paper. They include some regularity results for solutions to the Poisson equation associated with symmetric stable-like operators, and several averaging estimates for $\mathcal{L}_{\varepsilon}f$. Their proofs involve some delicate and lengthy estimates for distance functions and cut-off functions. The proofs of Theorems 1.1 and 1.2 are given in Section 3.

2. Preliminaries

In this section we give some preliminaries that are needed for the proofs of our main theorems.

2.1. Solution to the Poisson equation. Since $K \in C^2(\mathbb{R}^d \times \mathbb{R}^d)$ is multivariate 1-periodic on $\mathbb{R}^d \times \mathbb{R}^d$, we can view the corresponding symmetric Feller process $X := \{X_t\}_{t \ge 0}$ on \mathbb{R}^d and its L^2 -generator $(\mathcal{L}, \text{Dom}(\mathcal{L}))$ as a symmetric Feller process $X^{\mathbb{T}^d} := \{X_t^{\mathbb{T}^d}\}_{t \ge 0}$ on the torus \mathbb{T}^d and a generator $(\mathcal{L}, \text{Dom}(\mathcal{L}^{\mathbb{T}^d}))$ on \mathbb{T}^d , respectively.

Lemma 2.1. For every $f \in C(\mathbb{T}^d)$ with $\int_{\mathbb{T}^d} f(x) dx = 0$, there exists a unique $\psi_f \in \text{Dom}(\mathcal{L}^{\mathbb{T}^d}) \cap C(\mathbb{T}^d)$ such that

$$\mathcal{L}\psi_f = -f \quad on \ \mathbb{T}^d \tag{2.1}$$

and $\int_{\mathbb{T}^d} \psi_f(x) dx = 0$. Moreover, the following properties hold.

(1) For any $\alpha \in (0,2)$, there is a constant $C_1 > 0$ so that

$$\|\psi_f\|_{\infty} \leqslant C_1 \|f\|_{\infty}. \tag{2.2}$$

(2) If $\alpha \in (1,2)$, then $\psi_f \in C^1(\mathbb{T}^d)$ and there is a constant $C_2 > 0$ so that

$$\|\psi_f\|_{\infty} + \|\nabla\psi_f\|_{\infty} \leqslant C_2 \|f\|_{\infty}.$$
(2.3)

(3) If $\alpha = 1$, then there exist constants $\theta \in (0, 1)$ and $C_3 > 0$ such that

$$\sup_{x,y\in\mathbb{T}^d}\frac{|\psi_f(x)-\psi_f(y)|}{|x-y|^{\theta}}\leqslant C_3||f||_{\infty}.$$
(2.4)

Proof. Denote by $\{P_t\}_{t\geq 0}$ the transition semigroup of the Feller process $X^{\mathbb{T}^d}$ on \mathbb{T}^d . By [17, Theorem 1.1], it has a strictly positive, jointly continuous transition density function p(t, x, y) with

$$c_0^{-1}\left(t^{-d/\alpha} \wedge \frac{t}{|x-y|^{d+\alpha}}\right) \leqslant p(t,x,y) \leqslant c_0\left(t^{-d/\alpha} \wedge \frac{t}{|x-y|^{d+\alpha}}\right) \quad \text{on } (0,1] \times \mathbb{T}^d \times T^d \tag{2.5}$$

for some $c_0 > 1$. Since \mathbb{T}^d is compact, $X^{\mathbb{T}^d}$ has the normalized Lebesgue measure on \mathbb{T}^d as its equilibrium probability distribution as well as a stationary measure. It is standard to verify that $\{P_t\}_{t\geq 0}$ is uniformly exponentially ergodic (see e.g. [23]), that is,

$$||P_tg||_{\infty} \leqslant c_1 e^{-c_2 t} ||g||_{\infty} \quad \text{for any } g \in C(\mathbb{T}^d) \text{ with } \int_{\mathbb{T}^d} g(x) \, dx = 0.$$

$$(2.6)$$

Since for each t > 0, the operator P_t is Hilbert-Schmidt, it is compact in $L^2(\mathbb{T}^d; dx)$ and thus has discrete L^2 -spectrum $\{e^{-\lambda_k t} : k = 1, 2, \cdots\}$ with $0 = \lambda_0 < \lambda_1 \leq \lambda_2 \leq \cdots$ independent of t > 0. Denote by φ_k the corresponding eigenfunctions so that $\{\varphi_k : k \geq 0\}$ forms an orthonormal basis of $L^2(\mathbb{T}^d; dx)$ with $\varphi_0 = 1/|\mathbb{T}^d|$. Then, for each t > 0,

$$||P_t g||_2 \leq e^{-\lambda_1 t} ||g||_2$$
 for every $g \in L^2(\mathbb{T}^d; dx)$ with $\int_{\mathbb{T}^d} g(x) \, dx = 0.$ (2.7)

Denote by $\langle \cdot, \cdot \rangle_{L^2(\mathbb{T}^d; dx)}$ the inner product on $L^2(\mathbb{T}^d; dx)$. For $f \in C(\mathbb{T}^d)$ with $\int_{\mathbb{T}^d} f(x) dx = 0$, define

$$\psi_f := \int_0^\infty P_t f dt = \sum_{k=1}^\infty \lambda_k^{-1} \langle f, \varphi_k \rangle_{L^2(\mathbb{T}^d; dx)} \varphi_k,$$

which converges both in L^2 and uniformly on \mathbb{T}^d in view of (2.6) and (2.7). Clearly, $\int_{\mathbb{T}^d} \psi_f(x) dx = 0$. Moreover, it follows from the spectral theory that $\psi_f \in \text{Dom}(\mathcal{L}^{\mathbb{T}^d})$ with

$$\mathcal{L}\psi_f = \sum_{k=1}^{\infty} (-\lambda_k) \langle \psi_f, \varphi_k \rangle_{L^2(\mathbb{T}^d; dx)} \varphi_k = -\sum_{k=1}^{\infty} \langle f, \varphi_k \rangle_{L^2(\mathbb{T}^d; dx)} \varphi_k = -f.$$

So ψ_f is a solution to (2.1) with $\int_{\mathbb{T}^d} \psi_f(x) dx = 0$. Such a solution is unique as any $u \in \text{Dom}(\mathcal{L}^{\mathbb{T}^d})$ having $\mathcal{L}u = 0$ is constant on \mathbb{T}^d . It follows from (2.6) that (2.2) holds for every $\alpha \in (0, 2)$.

(i) When $\alpha \in (1,2)$, for $f \in C_c^2(\mathbb{R}^d)$,

$$\mathcal{L}f(x) = \int_{\mathbb{R}^d} (f(x+z) - f(x) - z \cdot \nabla f(x)) \frac{K(x, x+z)}{|z|^{d+\alpha}} dz + b(x) \cdot \nabla f(x),$$

where

$$b(x) = \frac{1}{2} \int_{\mathbb{R}^d} \frac{z(K(x, x+z) - K(x, x-z))}{|z|^{d+\alpha}} \, dz \in C_b(\mathbb{R}^d).$$

In particular, the drift term b(x) belongs to Kato's class in the sense of [22, Definition 1.3]; see [22, Remark 1.4]. If $\alpha \in (1, 2)$, then, by [22, Theorem 1.5(vi)],

$$\|\nabla P_t g\|_{\infty} \leqslant c_3 t^{-1/\alpha} \|g\|_{\infty}, \quad t \in (0,1], \ g \in C(\mathbb{T}^d).$$

Thus,

$$\begin{aligned} \|\nabla\psi_f\|_{\infty} &\leqslant \int_0^1 \|\nabla P_t f\|_{\infty} \, dt + \int_1^\infty \|\nabla P_1 \left(P_{t-1} f\right)\|_{\infty} \, dt \\ &\leqslant c_3 \left(\int_0^1 t^{-1/\alpha} \|f\|_{\infty} \, dt + \int_1^\infty \|P_{t-1} f\|_{\infty} \, dt\right) \\ &\leqslant c_4 \left(\int_0^1 t^{-1/\alpha} \|f\|_{\infty} \, dt + \int_1^\infty e^{-c_5(t-1)} \|f\|_{\infty} \, dt\right) \leqslant c_6 \|f\|_{\infty} \, dt \end{aligned}$$

So (2.3) is proved.

(ii) When $\alpha = 1$, by [17, Theorem 4.14], there exist constants $c_7 > 0$, $\kappa > 1$ and $\beta \in (0, 1)$ such that

$$|P_t g(x) - P_t g(y)| \leq c_7 ||g||_{\infty} t^{-\kappa} |x - y|^{\beta}, \quad t \in (0, 1], \ x, y \in \mathbb{T}^d \text{ and } g \in C(\mathbb{T}^d).$$
(2.8)

Write

$$\begin{aligned} |\psi_f(x) - \psi_f(y)| &\leq \left(\int_0^{|x-y|^{\beta/\kappa}} + \int_{|x-y|^{\beta/\kappa}}^1 + \int_1^\infty \right) |P_t f(x) - P_t f(y)| \, dt \\ &=: I_1(x,y) + I_2(x,y) + I_3(x,y). \end{aligned}$$

It holds that

$$I_1(x,y) \leq 2||f||_{\infty}|x-y|^{\beta/\kappa}.$$

Applying (2.8), we obtain

$$I_2(x,y) \leq c_8 ||f||_{\infty} |x-y|^{\beta(-\kappa+1)/\kappa} |x-y|^{\beta} = c_8 ||f||_{\infty} |x-y|^{\beta/\kappa}.$$

Using (2.6) and (2.8), we deduce

$$I_{3}(x,y) = \int_{1}^{\infty} |P_{1}(P_{t-1}f)(x) - P_{1}(P_{t-1}f)(y)| dt$$

$$\leq c_{7}|x-y|^{\beta} \int_{1}^{\infty} ||P_{t-1}f||_{\infty} dt$$

$$\leq c_{9}|x-y|^{\beta} \int_{1}^{\infty} e^{-c_{10}(t-1)} ||f||_{\infty} dt \leq c_{11} ||f||_{\infty} |x-y|^{\beta}$$

Combining all the estimates for $I_1(x, y)$, $I_2(x, y)$ and $I_3(x, y)$ yields the desired conclusion (2.4) with $\theta = \beta/\kappa$ when $\alpha = 1$.

Recall that $\{X_t^{\varepsilon}\}_{t\geq 0} := \{\varepsilon X_{\varepsilon^{-\alpha}t}\}_{t\geq 0}$, the associated Dirichlet form on $L^2(\mathbb{R}^d; dx)$ is $(\mathcal{E}^{\varepsilon}, W^{\alpha/2,2}(\mathbb{R}^d))$ defined by (1.5), and that $(\mathcal{L}_{\varepsilon}, \operatorname{Dom}(\mathcal{L}_{\varepsilon}))$ is the L^2 -generator of $(\mathcal{E}^{\varepsilon}, W^{\alpha/2,2}(\mathbb{R}^d))$ on $L^2(\mathbb{R}^d; dx)$. For an open subset $D \subset \mathbb{R}^d$, denote by $\{X_t^{\varepsilon,D}\}_{t\geq 0}$ the subprocess of $\{X_t^{\varepsilon}\}_{t\geq 0}$ killed upon leaving D. Its corresponding Dirichlet form $(\mathcal{E}^{\varepsilon}, W_0^{\alpha/2,2}(D))$ is given by

$$\mathcal{E}^{\varepsilon}(f,g) = \frac{1}{2} \iint_{\mathbb{R}^d \times \mathbb{R}^d} (f(x) - f(y))(g(x) - g(y)) \frac{K(x/\varepsilon, y/\varepsilon)}{|x - y|^{d + \alpha}} \, dx \, dy, \quad f,g \in W_0^{\alpha/2,2}(D),$$

$$W_0^{\alpha/2,2}(D) := \overline{C_c^2(D)}^{\|\cdot\|_{\alpha/2,2}}.$$

Denote the infinitesimal generator of $(\mathcal{E}^{\varepsilon}, W_0^{\alpha/2,2}(D))$ by $(\mathcal{L}_{\varepsilon}, \text{Dom}(\mathcal{L}_{\varepsilon}^D))$.

Lemma 2.2. Suppose that D is a bounded open subset of \mathbb{R}^d . Then there exists a constant $C_1 > 0$ such that for every $\varepsilon \in (0, 1)$,

$$\int_{D} |f(x)|^2 dx \leqslant C_1 \mathcal{E}^{\varepsilon}(f, f) \quad \text{for } f \in W_0^{\alpha/2, 2}(D).$$

$$(2.9)$$

Proof. Since the function K(x, y) is bounded between two positive constants, it suffices to show that there is a constant $c_1 > 0$ so that

$$\int_{D} |f(x)|^2 dx \leqslant c_1 \iint_{\mathbb{R}^d \times \mathbb{R}^d} \frac{|f(x) - f(y)|^2}{|x - y|^{d + \alpha}} dx \, dy \quad \text{for } f \in W_0^{\alpha/2, 2}(D).$$
(2.10)

Let $\{\bar{P}_t^D\}_{t\geq 0}$ be the Dirichlet semigroup on D associated with a rotationally symmetric α -stable Lévy process \bar{X} , whose infinitesimal generator is $\bar{\mathcal{L}}$. Since D is bounded, it well known that \bar{P}_t^D is compact on $L^2(D; dx)$ (see e.g. [16] or [21]), and its first eigenvalue is strictly less than 1. This fact implies the inequality (2.10) immediately, and so we can obtain the desired assertion.

Lemma 2.3. Suppose that D is a bounded open subset of \mathbb{R}^d and $f \in L^2(D; dx)$. Then for every $\varepsilon > 0$, there is a unique $\varphi_{\varepsilon} \in \text{Dom}(\mathcal{L}_{\varepsilon}^D)$ so that $\mathcal{L}_{\varepsilon}\varphi_{\varepsilon} = f$ on D, and there exists a constant $C_1 > 0$ such that for all $\varepsilon \in (0, 1)$,

$$\int_{D} |\varphi_{\varepsilon}(x)| \, dx \leqslant C_1 \int_{D} |f(x)| \, dx. \tag{2.11}$$

7

Moreover, if, in addition, f is bounded on D, then there is a constant $C_2 > 0$ such that for all $\varepsilon \in (0, 1)$,

$$\|\varphi_{\varepsilon}\|_{L^{\infty}(D;dx)} \leqslant C_2.$$

Proof. Since D is bounded, $D \subset B(0, R)$ for some R > 0. Suppose $f \in L^2(D; dx)$. By (2.9) and the Riesz representation theorem, there is a unique $\varphi_{\varepsilon} \in W_0^{\alpha/2,2}(D)$ so that

$$\mathcal{E}^{\varepsilon}(\varphi_{\varepsilon},g) = -\int_{D} f(x)g(x) \, dx \quad \text{for every } g \in W_{0}^{\alpha/2,2}(D).$$

It follows that $\varphi_{\varepsilon} \in \text{Dom}(\mathcal{L}_{\varepsilon}^{D})$ with $\mathcal{L}_{\varepsilon}\varphi_{\varepsilon} = f$. In fact, by [15],

$$\varphi_{\varepsilon}(x) = \mathbb{E}_x \left[\int_0^{\tau_D^{\varepsilon}} f(X_s) \, ds \right] = \int_D G_D^{\varepsilon}(x, y) f(y) \, dy, \quad \text{a.e. } x \in D.$$
(2.12)

where $\tau_D^{\varepsilon} := \inf\{t > 0 : X_t^{\varepsilon} \notin D\}$ is the first exit time from D for the process $\{X_t^{\varepsilon}\}_{t \ge 0}$, and $G_D^{\varepsilon}(x, y)$ is the Green function of X^{ε} in D. Note that

$$G_D^{\varepsilon} 1(x) := \int_D G_D^{\varepsilon}(x, y) \, dx = \mathbb{E}_y \left[\tau_D^{\varepsilon} \right], \quad y \in D.$$
(2.13)

By the symmetry of $G_D^{\varepsilon}(x, y)$ in (x, y), we have from (2.12) that

$$\int_{D} |\varphi_{\varepsilon}(x)| \, dx \leqslant \int_{D} |f(x)| G_{D}^{\varepsilon} \mathbb{1}(x) \, dx \leqslant \sup_{y \in D} \mathbb{E}_{y} \left[\tau_{D}^{\varepsilon}\right] \cdot \int_{D} |f(x)| \, dx.$$
(2.14)

By [17, Lemma 5.1], there is a constant $c_1 > 0$ that depends only on the bound $\Lambda \ge 1$ in (1.1) for the function K(x, y) and the dimension d so that

$$\sup_{\varepsilon \in (0,1)} \sup_{y \in D} \mathbb{E}_y \left[\tau_D^{\varepsilon} \right] \leqslant \sup_{\varepsilon \in (0,1)} \sup_{y \in B(0,R)} \mathbb{E}_y \left[\tau_{B(0,R)}^{\varepsilon} \right] \leqslant c_1 R^{\alpha}.$$
(2.15)

This together with (2.14) yields the estimate (2.11).

Furthermore, the last assertion is a direct consequence of (2.12), (2.13) and (2.15) as well as the fact that $G_D^{\varepsilon}(x,y) = G_D^{\varepsilon}(y,x)$ for all $x, y \in D$.

2.2. Distance functions and cut-off functions. It is well known that (see e.g. [46, Theorem 2, p. 171]) for any bounded open set $D \subset \mathbb{R}^d$, there is a C^{∞} -smooth regularizing distance function $\tilde{\delta}_D$ on D so that there are positive constants c_1 , c_2 and c_3 such that

$$c_1\delta_D(x) \leqslant \tilde{\delta}_D(x) \leqslant c_2\delta_D(x)$$
 and $|\nabla \tilde{\delta}_D(x)| \leqslant c_3$ for every $x \in D$.

For any r > 0, we set $D_r := \{x \in D : \widetilde{\delta}_D(x) > r\}.$

Lemma 2.4. Suppose that D is a bounded Lipschitz domain in \mathbb{R}^d . Then the following statements hold.

(1) When $\alpha \in (1,2)$, there is a constant $C_1 > 0$ so that for every $s, \varepsilon \in (0,1)$ and $x \in D$,

$$\int_{\{|z|>s\varepsilon\}} \delta_D(x+z)^{-2+\alpha/2} \mathbb{1}_{\{x+z\in D_{3\varepsilon}\}} \frac{1}{|z|^{d+\alpha-1}} dz$$

$$\leq C_1 \left((s\varepsilon)^{1-\alpha} \delta_D(x)^{-2+\alpha/2} + \varepsilon^{-1+\alpha/2} \delta_D(x)^{-\alpha} \right) \mathbb{1}_{D_{4\varepsilon}}(x) + C_1 s^{1-\alpha} \varepsilon^{-1-\alpha/2} \mathbb{1}_{D\setminus D_{4\varepsilon}}(x)$$

$$(2.16)$$

and

$$\int_{\{|z|>s\varepsilon\}} \mathbb{1}_{\{x+z\in D\setminus D_{3\varepsilon}\}} \frac{1}{|z|^{d+\alpha-1}} dz \leqslant C_1 \varepsilon \delta_D(x)^{-\alpha} \mathbb{1}_{D_{4\varepsilon}}(x) + C_1(s\varepsilon)^{1-\alpha} \mathbb{1}_{D\setminus D_{4\varepsilon}}(x).$$
(2.17)

(2) When $\alpha \in (0,1]$, there is a constant $C_2 > 0$ so that for every $s, \varepsilon \in (0,1)$ and $x \in D$,

$$\int_{\{s\varepsilon < |z| \le s\}} \delta_D(x+z)^{-2+\alpha/2} \mathbb{1}_{\{x+z\in D_{3\varepsilon}\}} \frac{1}{|z|^{d+\alpha-1}} dz
\le C_2 \left(\delta_D(x)^{-2+\alpha/2} \left(\delta_D(x)^{1-\alpha} \mathbb{1}_{\{\alpha\in(0,1)\}} + (1+|\log\varepsilon|) \mathbb{1}_{\{\alpha=1\}} \right) + \varepsilon^{-1+\alpha/2} \delta_D(x)^{-\alpha} \mathbb{1}_{\{s>\delta_D(x)/4\}} \right) \mathbb{1}_{D_{4\varepsilon}}(x)
+ C_2 \varepsilon^{-2+\alpha/2} \left(s^{1-\alpha} \mathbb{1}_{\{\alpha\in(0,1)\}} + (1+|\log\varepsilon|) \mathbb{1}_{\{\alpha=1\}} \right) \mathbb{1}_{D\setminus D_{4\varepsilon}}(x),$$
(2.18)

and

$$\int_{\{s\varepsilon < |z| \leq s\}} \mathbb{1}_{\{x+z\in D\setminus D_{3\varepsilon}\}} \frac{1}{|z|^{d+\alpha-1}} dz$$

$$\leq C_2 \varepsilon \delta_D(x)^{-\alpha} \mathbb{1}_{D_{4\varepsilon}}(x) + C_2 \left(s^{1-\alpha} \mathbb{1}_{\{\alpha\in(0,1)\}} + (1+|\log\varepsilon|) \mathbb{1}_{\{\alpha=1\}}\right) \mathbb{1}_{D\setminus D_{4\varepsilon}}(x),$$
(2.19)

 $as \ well \ as$

$$\int_{\{|z|>s\}} \left(\delta_D(x+z)^{-1+\alpha/2} \mathbb{1}_{\{x+z\in D_{3\varepsilon}\}} + \varepsilon^{-1+\alpha/2} \mathbb{1}_{\{x+z\in D\setminus D_{3\varepsilon}\}} \right) \frac{1}{|z|^{d+\alpha}} \, dz \leqslant C_2 \varepsilon^{-1+\alpha/2} s^{-\alpha}. \tag{2.20}$$

(3) When $\alpha \in (0,1)$, for every $\beta \in (\alpha/2, \alpha)$ there is a constant $C_3 > 0$ so that for all $\varepsilon \in (0,1)$ and $x \in D$,

$$\int_{\{|z|>\varepsilon\}} \delta_D(x+z)^{-1-\beta+\alpha/2} \mathbb{1}_{\{x+z\in D_{3\varepsilon}\}} \frac{1}{|z|^{d+\alpha-\beta}} dz$$

$$\leq C_3 \left(\varepsilon^{-(\beta-\alpha/2)} \delta_D(x)^{-1-(\alpha-\beta)} + \varepsilon^{-(\alpha-\beta)} \delta_D(x)^{-1-\beta+\alpha/2}\right) \mathbb{1}_{D_{4\varepsilon}}(x) + C_3 \varepsilon^{-1-\alpha/2} \mathbb{1}_{D\setminus D_{4\varepsilon}}(x)$$
and
$$(2.21)$$

$$\int_{\{|z|>\varepsilon\}} \mathbb{1}_{\{x+z\in D\setminus D_{3\varepsilon}\}} \frac{1}{|z|^{d+\alpha-\beta}} dz \leqslant C_3 \varepsilon \delta_D(x)^{-1-(\alpha-\beta)} \mathbb{1}_{D_{4\varepsilon}}(x) + C_3 \varepsilon^{-(\alpha-\beta)} \mathbb{1}_{D\setminus D_{4\varepsilon}}(x).$$
(2.22)

Proof. (1) Suppose that $\alpha \in (1,2)$. Let $s, \varepsilon \in (0,1)$. Then, for every $x \in D_{4\varepsilon}$, it holds that

$$\begin{split} &\int_{\{|z|>s\varepsilon\}} \delta_D(x+z)^{-2+\alpha/2} \mathbb{1}_{\{x+z\in D_{3\varepsilon}\}} \frac{1}{|z|^{d+\alpha-1}} \, dz \\ &\leqslant c_1 \left(\delta_D(x)^{-2+\alpha/2} \int_{\{s\varepsilon<|z|\leqslant\delta_D(x)/4\}} \frac{1}{|z|^{d+\alpha-1}} \, dz + \int_{\{x+z\in D:|z|>\delta_D(x)/4,\delta_D(x+z)>3\varepsilon\}} \frac{\delta_D(x+z)^{-2+\alpha/2}}{|z|^{d+\alpha-1}} \, dz \right) \\ &=: I_1(x) + I_2(x), \end{split}$$

where in the inequality above we used

$$\delta_D(x+z) \ge \delta_D(x)/2 \quad \text{for } |z| \le \delta_D(x)/4.$$
 (2.23)

It is clear that

$$I_1(x) \leqslant c_2 \delta_D(x)^{-2+\alpha/2} (s\varepsilon)^{1-\alpha}.$$

On the other hand,

$$I_{2}(x) \leqslant \sum_{k=0}^{\infty} \int_{\{x+z\in D: 2^{k}\delta_{D}(x)/4 < |z| \leq 2^{k+1}\delta_{D}(x)/4, \delta_{D}(x+z) > 3\varepsilon\}} \frac{\delta_{D}(x+z)^{-2+\alpha/2}}{|z|^{d+\alpha-1}} dz$$

$$\leqslant c_{3} \sum_{k=0}^{\infty} (2^{k}\delta_{D}(x))^{-d-\alpha+1} \int_{\{x+z\in D: 2^{k}\delta_{D}(x)/4 < |z| \leq 2^{k+1}\delta_{D}(x)/4, \delta_{D}(x+z) > 3\varepsilon\}} \delta_{D}(x+z)^{-2+\alpha/2} dz \qquad (2.24)$$

$$=: c_{3} \sum_{k=0}^{\infty} (2^{k}\delta_{D}(x))^{-d-\alpha+1} I_{2k}(x).$$

By a change of variable, we have

$$I_{2,k}(x) = \int_{\{z \in D: 2^k \delta_D(x)/4 < |z-x| \le 2^{k+1} \delta_D(x)/4, \delta_D(z) > 3\varepsilon\}} \delta_D(z)^{-2+\alpha/2} \, dz.$$

If $\delta_D(x) > c_4 2^{-k}$ for some $c_4 > 0$ to be determined later, then it holds that

$$I_{2,k}(x) \leqslant \int_{\{z \in D: \delta_D(x) > 3\varepsilon\}} \delta_D(z)^{-2+\alpha/2} dz$$
$$\leqslant c_5 \int_{3\varepsilon}^{\infty} s^{-2+\alpha/2} ds \leqslant c_6 \left(2^{k+1} \delta_D(x)\right)^{d-1} \varepsilon^{-1+\alpha/2},$$

where the last inequality due to the fact $2^k \delta_D(x) \ge c_4$.

When $\delta_D(x) \leq c_4 2^{-k}$, since D is bounded and Lipschitz, we can choose $c_4 > 0$ small enough so that there exists a change of coordinates $\Psi : B(x, 2^{k+1}\delta_D(x)/4) \cap D \to B(x', 2^{k+1}\delta_D(x)) \cap \mathbb{H}_+$ with $\mathbb{H}_+ := \{z \in \mathbb{R}^d : z = (z_1, \cdots, z_d), z_d > 0\}$ which satisfies that

(i) Ψ is a homeomorphism, $\Psi(x) = x'$ and

$$|\Psi(x_1) - \Psi(x_2)| \leq c_7 |x_1 - x_2| \quad \text{for } x_1, x_2 \in B\left(x, 2^{k+1}\delta_D(x)/4\right) \cap D,$$

$$|\Psi^{-1}(z_1) - \Psi^{-1}(z_2)| \leq c_7 |z_1 - z_2| \quad \text{for } z_1, z_2 \in B\left(x', 2^{k+1}\delta_D(x)\right) \cap \mathbb{H}_+;$$

(ii)

$$c_8 \delta_D(z) \leq \delta_{\mathbb{H}_+} (\Psi(z)) \leq c_7 \delta_D(z) \quad \text{for } z \in B\left(x, 2^{k+1} \delta_D(x)/4\right)$$

Here, $\delta_{\mathbb{H}_+}(z) := z_d$, for any $z \in \mathbb{H}_+$, denotes the distance from $z \in \mathbb{H}_+$ to the boundary \mathbb{H}_+ . In particular, we remark that all the constants c_i above are independent of x.

According to these properties above and a change of variable $z \mapsto \Psi(z)$, we derive

$$I_{2,k} \leqslant c_9 \int_{\{z \in \mathbb{H}_+ : |z-x'| \leqslant c_{10}2^{k+1}\delta_D(x), \delta_{\mathbb{H}_+}(z) = z_d > c_{11}\varepsilon\}} \delta_{\mathbb{H}_+}(z)^{-2+\alpha/2} dz$$

$$\leqslant c_9 \int_{\{z \in \mathbb{H}_+ : \sup_{1 \leqslant i \leqslant d-1} |z_i - x'_i| \leqslant c_{10}2^{k+1}\delta_D(x), z_d > c_{11}\varepsilon\}} z_d^{-2+\alpha/2} dz$$

$$\leqslant c_{12} \left(2^{k+1}\delta_D(x)\right)^{d-1} \int_{c_{11}\varepsilon}^{\infty} s^{-2+\alpha/2} ds \leqslant c_{13} \left(2^{k+1}\delta_D(x)\right)^{d-1} \varepsilon^{-1+\alpha/2}.$$

Therefore, summarising all the estimates above together, we arrive at

$$I_{2}(x) \leqslant c_{3} \sum_{k=0}^{\infty} (2^{k} \delta_{D}(x))^{-d-\alpha+1} I_{2,k}(x) \leqslant c_{14} \varepsilon^{-1+\alpha/2} \sum_{k=0}^{\infty} (2^{k} \delta_{D}(x))^{-\alpha} \leqslant c_{15} \varepsilon^{-1+\alpha/2} \delta_{D}(x)^{-\alpha}.$$
(2.25)

When $x \in D \setminus D_{4\varepsilon}$, for any $s, \varepsilon \in (0, 1)$, it holds that

$$\int_{\{|z|>s\varepsilon\}} \left(\delta_D(x+z)^{-2+\alpha/2} \mathbb{1}_{\{x+z\in D_{3\varepsilon}\}} + \varepsilon^{-2+\alpha/2} \mathbb{1}_{\{x+z\in D\setminus D_{3\varepsilon}\}} \right) \frac{1}{|z|^{d+\alpha-1}} dz \\
\leqslant c_{16}\varepsilon^{-2+\alpha/2} \int_{\{|z|>s\varepsilon\}} \frac{1}{|z|^{d+\alpha-1}} dz \leqslant c_{17}s^{1-\alpha}\varepsilon^{-1-\alpha/2}.$$
(2.26)

Combining this with the estimates above for $I_1(x)$ and $I_2(x)$ yields (2.16).

Note that, for any $x \in D_{4\varepsilon}$ and $s, \varepsilon \in (0, 1)$, if $\delta_D(x + z) \leq 3\varepsilon$, then it must hold that $|z| > \delta_D(x)/4$. By this observation, we obtain

$$\int_{\{|z|>s\varepsilon,\delta_D(x+z)\leqslant 3\varepsilon\}} \frac{1}{|z|^{d+\alpha-1}} dz \leqslant \int_{\{|z|>\delta_D(x)/4,\delta_D(x+z)\leqslant 3\varepsilon\}} \frac{1}{|z|^{d+\alpha-1}} dz \\
\leqslant \sum_{k=0}^{\infty} \int_{\{2^k \delta_D(x)/4 < |z|\leqslant 2^{k+1} \delta_D(x)/4,\delta_D(x+z)\leqslant 3\varepsilon\}} \frac{1}{|z|^{d+\alpha-1}} dz \\
\leqslant c_{18} \sum_{k=0}^{\infty} (2^k \delta_D(x))^{-d-\alpha+1} \int_{\{2^k \delta_D(x)/4 < |z|\leqslant 2^{k+1} \delta_D(x)/4,\delta_D(x+z)\leqslant 3\varepsilon\}} dz \\
\leqslant c_{19} \sum_{k=0}^{\infty} (2^k \delta_D(x))^{-d-\alpha+1} \left(2^{k+1} \delta_D(x)\right)^{d-1} \varepsilon \leqslant c_{20}\varepsilon \delta_D(x)^{-\alpha}.$$
(2.27)

Here in the fourth inequality above we have used the fact

$$\int_{\{2^k\delta_D(x)/4 < |z| \le 2^{k+1}\delta_D(x)/4, \delta_D(x+z) \le 3\varepsilon\}} dz \le c_{21}\varepsilon \left(2^{k+1}\delta_D(x)\right)^{d-1}$$

which can be verified by the same argument for the estimate of $I_2(x)$ above. This, along with (2.26), in turn implies (2.17).

(2) For simplicity we only prove (2.18) and (2.19) for the case that $\alpha = 1$, since the case $\alpha \in (0, 1)$ can be proved almost in the same way. Below in Step (2), we let $\alpha = 1$ unless particularly stressed. For every $x \in D_{4\varepsilon}$, it holds for every $s, \varepsilon \in (0, 1)$ that

$$\begin{split} &\int_{\{s\varepsilon<|z|\leqslant s\}} \delta_D(x+z)^{-2+\alpha/2} \mathbb{1}_{\{x+z\in D_{3\varepsilon}\}} \frac{1}{|z|^{d+\alpha-1}} dz \\ &\leqslant c_{22} \left(\delta_D(x)^{-2+\alpha/2} \int_{\{s\varepsilon<|z|\leqslant \min(\delta_D(x)/4,s),\delta_D(x+z)>3\varepsilon\}} \frac{1}{|z|^{d+\alpha-1}} dz \\ &+ \int_{\{\min(\delta_D(x)/4,s)<|z|\leqslant s,\delta_D(x+z)>3\varepsilon\}} \frac{\delta_D(x+z)^{-2+\alpha/2}}{|z|^{d+\alpha-1}} dz \right) \\ &=: J_1(x) + J_2(x), \end{split}$$

where we have used (2.23) in the inequality above for the term $J_1(x)$. It is obvious that

 $J_1(x) \leqslant c_{23}\delta_D(x)^{-2+\alpha/2}(1+|\log\varepsilon|).$

Furthermore, following the argument for the estimate of $I_2(x)$ above, we can obtain that

$$J_2(x) \leqslant c_{24} \mathbb{1}_{\{s > \delta_D(x)/4\}} \int_{\{|z| > \delta_D(x)/4, \delta_D(x+z) > 3\varepsilon\}} \frac{\delta_D(x+z)^{-2+\alpha/2}}{|z|^{d+\alpha-1}} dz \leqslant c_{25} \varepsilon^{-1+\alpha/2} \delta_D(x)^{-\alpha} \mathbb{1}_{\{s > \delta_D(x)/4\}}.$$

When $x \in D \setminus D_{4\varepsilon}$, we have, for any $s, \varepsilon \in (0, 1)$,

$$\int_{\{s\varepsilon < |z| \leqslant s\}} \left(\delta_D(x+z)^{-2+\alpha/2} \mathbb{1}_{\{x+z\in D_{3\varepsilon}\}} + \varepsilon^{-2+\alpha/2} \mathbb{1}_{\{x+z\in D\setminus D_{3\varepsilon}\}} \right) \frac{1}{|z|^{d+\alpha-1}} dz \\
\leqslant c_{26} \varepsilon^{-2+\alpha/2} \int_{\{s\varepsilon < |z| \leqslant s\}} \frac{1}{|z|^{d+\alpha-1}} dz \leqslant c_{27} \varepsilon^{-2+\alpha/2} (1+|\log\varepsilon|).$$
(2.28)

Combining this with the estimates above for $J_1(x)$ and $J_2(x)$ yields (2.18).

According to the argument of (2.27), we deduce that for every $x \in D_{4\varepsilon}$ and $s, \varepsilon \in (0, 1)$,

$$\int_{\{s\varepsilon<|z|\leqslant s,\delta_D(x+z)\leqslant 3\varepsilon\}} \frac{1}{|z|^{d+\alpha-1}} dz \leqslant \int_{\{|z|>\delta_D(x)/4,\delta_D(x+z)\leqslant 3\varepsilon\}} \frac{1}{|z|^{d+\alpha-1}} dz \leqslant c_{28}\varepsilon \delta_D(x)^{-\alpha}.$$

By this and (2.28), we prove (2.19).

For every $\alpha \in (0, 1]$, $x \in D$ and $s, \varepsilon \in (0, 1)$, we have

$$\int_{\{|z|>s\}} \left(\delta_D(x+z)^{-1+\alpha/2} \mathbb{1}_{\{x+z\in D_{3\varepsilon}\}} + \varepsilon^{-1+\alpha/2} \mathbb{1}_{\{x+z\in D\setminus D_{3\varepsilon}\}} \right) \frac{1}{|z|^{d+\alpha}} dz$$
$$\leqslant c_{29}\varepsilon^{-1+\alpha/2} \int_{\{|z|>s\}} \frac{1}{|z|^{d+\alpha}} dz \leqslant c_{30}\varepsilon^{-1+\alpha/2} s^{-\alpha}.$$

So (2.20) is proved.

(3) Suppose that $\alpha \in (0,1)$. For every $\beta \in (\alpha/2, \alpha)$, $\varepsilon \in (0,1)$ and $x \in D_{4\varepsilon}$, following the same argument as that for (2.16), we have

$$\int_{\{|z|>\varepsilon\}} \delta_D(x+z)^{-1-\beta+\alpha/2} \mathbb{1}_{\{x+z\in D_{3\varepsilon}\}} \frac{1}{|z|^{d+\alpha-\beta}} dz
\leq c_{31} \left(\delta_D(x)^{-1-\beta+\alpha/2} \int_{\{\varepsilon<|z|\leqslant\delta_D(x)/4\}} \frac{1}{|z|^{d+\alpha-\beta}} dz + \int_{\{|z|>\delta_D(x)/4,\delta_D(x+z)>3\varepsilon\}} \frac{\delta_D(x+z)^{-1-\beta+\alpha/2}}{|z|^{d+\alpha-\beta}} dz \right)
=: L_1(x) + L_2(x).$$

Furthermore, one can see that

$$L_1(x) \leq c_{32} \delta_D(x)^{-1-\beta+\alpha/2} \varepsilon^{-(\alpha-\beta)}.$$

Similar to the argument for the estimates of $I_2(x)$ above we have

$$L_{2}(x) \leqslant \sum_{k=0}^{\infty} \int_{\{2^{k}\delta_{D}(x)/4 < |z| \leqslant 2^{k+1}\delta_{D}(x)/4, \delta_{D}(x+z) > 3\varepsilon\}} \frac{\delta_{D}(x+z)^{-1-\beta+\alpha/2}}{|z|^{d+\alpha-\beta}} dz$$

$$\leqslant c_{33} \sum_{k=0}^{\infty} (2^{k}\delta_{D}(x))^{-d-\alpha+\beta} \int_{\{2^{k}\delta_{D}(x)/4 < |z| \leqslant 2^{k+1}\delta_{D}(x)/4, \delta_{D}(x+z) > 3\varepsilon\}} \delta_{D}(x+z)^{-1-\beta+\alpha/2} dz$$

$$\leqslant c_{34} \sum_{k=0}^{\infty} (2^{k}\delta_{D}(x))^{-d-\alpha+\beta} \left(2^{k+1}\delta_{D}(x)\right)^{d-1} \int_{3\varepsilon}^{\infty} s^{-1-\beta+\alpha/2} ds$$

$$\leqslant c_{35}\varepsilon^{-(\beta-\alpha/2)}\delta_{D}(x)^{-1-(\alpha-\beta)} \sum_{k=0}^{\infty} 2^{-k(1+\alpha-\beta)} \leqslant c_{36}\varepsilon^{-(\beta-\alpha/2)}\delta_{D}(x)^{-1-(\alpha-\beta)}.$$

When $x \in D \setminus D_{4\varepsilon}$ and $s, \varepsilon \in [0, 1]$, it holds that

$$\int_{\{|z|>\varepsilon\}} \left(\delta_D(x+z)^{-1-\beta+\alpha/2} \mathbb{1}_{\{x+z\in D_{3\varepsilon}\}} + \varepsilon^{-1-\beta+\alpha/2} \mathbb{1}_{\{x+z\in D\setminus D_{3\varepsilon}\}}\right) \frac{1}{|z|^{d+\alpha-\beta}} dz$$

$$\leq c_{37}\varepsilon^{-1-\beta+\alpha/2} \int_{\{|z|>\varepsilon\}} \frac{1}{|z|^{d+\alpha-\beta}} dz \leq c_{38}\varepsilon^{-1-\alpha/2}.$$
(2.29)

Combining this with the estimates above for $L_1(x)$ and $L_2(x)$ yields (2.21).

Following the same argument as that for (2.27), we can show that for every $x \in D_{4\varepsilon}$ and $s, \varepsilon \in [0, 1]$,

$$\int_{\{|z|>\varepsilon,\delta_D(x+z)\leqslant 3\varepsilon\}} \frac{1}{|z|^{d+\alpha-\beta}} dz \leqslant c_{39}\varepsilon \delta_D(x)^{-1-(\alpha-\beta)}.$$

By this and (2.29), we obtain (2.22).

Remark 2.5. In fact, when $\alpha \in (1, 2)$, a direct calculation yields that

$$I_{2}(x) = \int_{\{x+z\in D: |z|>\delta_{D}(x)/4, \delta_{D}(x+z)>3\varepsilon\}} \frac{\delta_{D}(x+z)^{-2+\alpha/2}}{|z|^{d+\alpha-1}} dz$$

$$\leqslant c_{1}\varepsilon^{-2+\alpha/2} \int_{\{z\in D: |z|>\delta_{D}(x)/4\}} \frac{1}{|z|^{d+\alpha-1}} dz \leqslant c_{2}\varepsilon^{-2+\alpha/2} \delta_{D}(x)^{-1+\alpha/2}.$$

However, this is not enough for our application later. So here we need some more effort to obtain a stronger estimate (2.25).

Let $\eta_{\varepsilon} \in C_c^{\infty}(D)$ be a cut-off function near the boundary ∂D (which can be constructed by using the C^{∞} regularizing distance function $\tilde{\delta}_D$) so that $0 \leq \eta_{\varepsilon} \leq 1$ on \mathbb{R}^d ,

$$\eta_{\varepsilon}(x) = \begin{cases} 1, & x \in D_{2\varepsilon}, \\ 0, & x \in \mathbb{R}^d \setminus D_{\varepsilon}, \end{cases}$$
(2.30)

and

$$|\nabla^k \eta_{\varepsilon}(x)| \leqslant c_0 \varepsilon^{-k} \mathbb{1}_{D_{2\varepsilon} \setminus D_{\varepsilon}}(x), \quad k = 1, 2, 3, 4, \ x \in D,$$

$$(2.31)$$

where the constant c_0 is independent of ε and x. We have the following estimate for $\mathcal{L}\eta_{\varepsilon}$, where \mathcal{L} is the fractional Laplacian given by (1.8).

Lemma 2.6. Suppose that D is a bounded Lipschitz domain in \mathbb{R}^d . Then there exists a constant $C_1 > 0$ such that

$$\left|\bar{\mathcal{L}}\eta_{\varepsilon}(x)\right| \leqslant C_1\left(\varepsilon\delta_D(x)^{-1-\alpha}\mathbb{1}_{D_{3\varepsilon}}(x) + \varepsilon^{-\alpha}\mathbb{1}_{D\setminus D_{3\varepsilon}}(x)\right), \quad x \in D.$$
(2.32)

Proof. When $x \in D_{3\varepsilon}$, by (2.30), we know that $\nabla \eta_{\varepsilon}(x) = 0$ and

$$\eta_{\varepsilon}(x+z) - \eta_{\varepsilon}(x) \neq 0$$
 only if $x+z \in D \setminus D_{2\varepsilon}$.

Because $x \in D_{3\varepsilon}$ and $x + z \in D \setminus D_{2\varepsilon}$ imply that $|z| > \delta_D(x)/3$,

$$\begin{aligned} |\bar{\mathcal{L}}\eta_{\varepsilon}(x)| &= \left| \int_{\mathbb{R}^d} \left(\eta_{\varepsilon}(x+z) - \eta_{\varepsilon}(x) - \langle \nabla \eta_{\varepsilon}(x), z \rangle \right) \frac{\bar{K}}{|z|^{d+\alpha}} \, dz \right| \\ &\leq 2 \int_{\{|z| > \delta_D(x)/3, \delta_D(x+z) \leqslant 2\varepsilon\}} \frac{1}{|z|^{d+\alpha}} \, dz \leqslant c_1 \varepsilon \delta_D(x)^{-1-\alpha}, \end{aligned}$$

where in the inequality we have used

$$\int_{\{|z| > \delta_D(x)/3, \delta_D(x+z) \le 2\varepsilon\}} \frac{1}{|z|^{d+\alpha}} dz \le c_2 \varepsilon \delta_D(x)^{-1-\alpha}$$
(2.33)

that can be proved by following the argument for (2.27).

When $x \in D \setminus D_{3\varepsilon}$, it follows from (2.31) that

$$|\bar{\mathcal{L}}\eta_{\varepsilon}(x)| \leqslant \|\nabla^2 \eta_{\varepsilon}\|_{\infty} \int_{\{|z|\leqslant\varepsilon\}} \frac{|z|^2}{|z|^{d+\alpha}} dz + 2 \int_{\{|z|>\varepsilon\}} \frac{1}{|z|^{d+\alpha}} dz \leqslant c_3 \varepsilon^{-\alpha}.$$

$$(32).$$

This establishes (2.32).

Lemma 2.7. Suppose that D is a bounded Lipschitz domain in \mathbb{R}^d . Then, for every $f \in C^1_{(-\alpha/2)}(D)$, there exists a constant $C_1 > 0$ such that for all $\varepsilon \in (0, 1)$ and $x \in D$,

$$\left| \int_{\mathbb{R}^d} \left(f(x+z) - f(x) \right) \left(\eta_{\varepsilon}(x+z) - \eta_{\varepsilon}(x) \right) \frac{1}{|z|^{d+\alpha}} dz \right|$$

$$\leq C_1 \left(\varepsilon \delta_D(x)^{-1-\alpha/2} \mathbb{1}_{D_{4\varepsilon}}(x) + \varepsilon^{-\alpha/2} \mathbb{1}_{D \setminus D_{4\varepsilon}}(x) \right).$$
(2.34)

Proof. Since $f \in C^1_{(-\alpha/2)}(D) \subset C^{\alpha/2}_{(-\alpha/2)}(D)$ (see e.g. [51, Lemma 4.1] for an equivalent characterization of the space $C^{\beta}_{(-\alpha/2)}(D)$) and f(x) = 0 for every $x \in D^c$, it holds that

$$|f(x) - f(x+z)| \leq c_1 |z|^{\alpha/2}, \quad x, z \in \mathbb{R}^d.$$
 (2.35)

and

$$|f(x)| \leqslant c_1 \delta_D(x)^{\alpha/2}, \quad x \in D.$$
(2.36)

When $x \in D_{4\varepsilon}$, due to (2.36) and the fact that

 $\eta_{\varepsilon}(x+z) - \eta_{\varepsilon}(x)$ is possibly non-zero only if when $x+z \in D \setminus D_{2\varepsilon}$,

we obtain

$$\left| \int_{\mathbb{R}^d} \left(f(x+z) - f(x) \right) \left(\eta_{\varepsilon}(x+z) - \eta_{\varepsilon}(x) \right) \frac{1}{|z|^{d+\alpha}} dz \right|$$

$$\leq c_2 \int_{\{|z| > \delta_D(x)/3\}} \left(\delta_D(x+z)^{\alpha/2} + \delta_D(x)^{\alpha/2} \right) \mathbb{1}_{\{x+z \in D \setminus D_{2\varepsilon}\}} \frac{1}{|z|^{d+\alpha}} dz$$
$$\leq c_3 \left(\varepsilon^{\alpha/2} + \delta_D(x)^{\alpha/2} \right) \int_{\{|z| > \delta_D(x)/3, \delta_D(x+z) \leq 2\varepsilon\}} \frac{1}{|z|^{d+\alpha}} dz \leq c_4 \varepsilon \delta_D(x)^{-1-\alpha/2},$$

where in the first inequality we used the fact that $x \in D_{4\varepsilon}$ and $x + z \in D \setminus D_{2\varepsilon}$ imply $|z| > \delta_D(x)/3$, and the last inequality follows from (2.33).

When $x \in D \setminus D_{4\varepsilon}$,

$$\begin{split} \left| \int_{\mathbb{R}^d} \left(f(x+z) - f(x) \right) \left(\eta_{\varepsilon}(x+z) - \eta_{\varepsilon}(x) \right) \frac{1}{|z|^{d+\alpha}} \, dz \right| \\ &\leqslant \left[\int_{\{|z| \leqslant 5\varepsilon\}} + \int_{\{|z| > 5\varepsilon\}} \right] |f(x+z) - f(x)| \left| \eta_{\varepsilon}(x+z) - \eta_{\varepsilon}(x) \right| \frac{1}{|z|^{d+\alpha}} \, dz \\ &=: I_1^{\varepsilon}(x) + I_2^{\varepsilon}(x). \end{split}$$

Using (2.35) and (2.31), we obtain

$$I_1^{\varepsilon}(x) \leqslant c_5 \|\nabla \eta_{\varepsilon}\|_{\infty} \int_{\{|z| \leqslant 5\varepsilon\}} \frac{1}{|z|^{d+\alpha/2-1}} \, dz \leqslant c_6 \varepsilon^{-\alpha/2}$$

and

$$I_2^{\varepsilon}(x) \leqslant c_7 \|\eta_{\varepsilon}\|_{\infty} \int_{\{|z| > 5\varepsilon\}} \frac{1}{|z|^{d+\alpha/2}} \, dz \leqslant c_8 \varepsilon^{-\alpha/2}$$

Therefore, putting all the estimates above together yields (2.34).

2.3. Averaging estimates related to $\mathcal{L}_{\varepsilon}f$. Throughout this subsection, without any mention we always assume that D is a bounded Lipschitz domain in \mathbb{R}^d . For any $f \in C^3_{(-\alpha/2)}(D)$, set $f_{\varepsilon}(x) := f(x)\eta_{\varepsilon}(x)$, where $\eta_{\varepsilon} \in C^{\infty}_{c}(D)$ is a cut-off function satisfying (2.30) and (2.31). By the definition, it holds for $f \in C^3_{(-\alpha/2)}(D)$ that

$$|\nabla^k f(x)| \le \|f\|_{3;D}^{(-\alpha/2)} \delta_D(x)^{-k+\alpha/2}, \quad x \in D, \ k = 0, 1, 2, 3.$$
(2.37)

Combining this with (2.31) yields that

$$|\nabla^k f_{\varepsilon}(x)| \leq c_0 \delta_D(x)^{-k+\alpha/2} \mathbb{1}_{D_{\varepsilon}}(x)$$

$$\leq c_1 \left(\delta_D(x)^{-k+\alpha/2} \mathbb{1}_{D_{3\varepsilon}}(x) + \varepsilon^{-k+\alpha/2} \mathbb{1}_{D \setminus D_{3\varepsilon}}(x) \right), \quad x \in D, \ k = 0, 1, 2, 3.$$

$$(2.38)$$

For every $g \in C_b^2(\mathbb{R}^d)$, define

$$\hat{\mathcal{L}}_{1,\varepsilon}g(x) := \int_{\mathbb{R}^d} \left(g(x+z) - g(x) - \langle \nabla g(x), z \rangle \right) \frac{K\left(x/\varepsilon, (x+z)/\varepsilon\right)}{|z|^{d+\alpha}} dz, \quad \alpha \in (1,2),$$

$$\hat{\mathcal{L}}_{2,\varepsilon}g(x) := \int_{\mathbb{R}^d} \left(g(x+z) - g(x) - \langle \nabla g(x), z \rangle \, \mathbb{1}_{\{|z| \leqslant 1\}} \right) \frac{K\left(x/\varepsilon, (x+z)/\varepsilon\right)}{|z|^{d+\alpha}} dz, \quad \alpha \in (0,1];$$

$$(2.39)$$

and

$$\bar{\mathcal{L}}_{\varepsilon}g(x) := \begin{cases} \int_{\mathbb{R}^d} \left(g(x+z) - g(x) - \langle \nabla g(x), z \rangle \right) \frac{\bar{K}\left(x/\varepsilon\right)}{|z|^{d+\alpha}} dz, & \alpha \in (1,2), \\ \int_{\mathbb{R}^d} \left(g(x+z) - g(x) - \langle \nabla g(x), z \rangle \, \mathbb{1}_{\{|z| \le 1\}} \right) \frac{\bar{K}\left(x/\varepsilon\right)}{|z|^{d+\alpha}} dz, & \alpha \in (0,1], \end{cases}$$
(2.40)

where $\bar{K}(x) := \int_{\mathbb{T}^d} K(x, y) \, dy$. (Note that the constant \bar{K} defined in (1.7) is the value of $\int_{\mathbb{T}^d} \bar{K}(x) \, dx$.) The number of this part is to present extract extraction concerning the extra presenting for $\bar{K}(x)$.

The purpose of this part is to present several estimates concerning the averaging properties for $f \in C^3_{(-\alpha/2)}(D)$. For this, we need to take account into the effect of blow up behaviours for ∇f and $\nabla^2 f$ near the boundary ∂D .

Lemma 2.8. Suppose that $\alpha \in (1,2)$. Then, for every $f \in C^3_{(-\alpha/2)}(D)$, there are constants $C_1, C_2 > 0$ such that for all $\varepsilon \in (0,1)$ and $x \in D$,

$$|\hat{\mathcal{L}}_{1,\varepsilon}f_{\varepsilon}(x) - \bar{\mathcal{L}}_{\varepsilon}f_{\varepsilon}(x)| \leq C_1 \left(\varepsilon^{2-\alpha}\delta_D(x)^{-2+\alpha/2} + \varepsilon^{\alpha/2}\delta_D(x)^{-\alpha}\right) \mathbb{1}_{D_{4\varepsilon}}(x) + C_1\varepsilon^{-\alpha/2}\mathbb{1}_{D\setminus D_{4\varepsilon}}(x)$$
(2.41)

and

$$|\hat{\mathcal{L}}_{1,\varepsilon}\nabla f_{\varepsilon}(x)| \leq C_2 \left(\varepsilon^{1-\alpha}\delta_D(x)^{-2+\alpha/2} + \varepsilon^{-1+\alpha/2}\delta_D(x)^{-\alpha}\right) \mathbb{1}_{D_{4\varepsilon}}(x) + C_2\varepsilon^{-1-\alpha/2}\mathbb{1}_{D\setminus D_{4\varepsilon}}(x).$$
(2.42)

Here $f_{\varepsilon} := f \eta_{\varepsilon}$.

Proof. (1) Set

$$H_{\varepsilon}(x,z) := \left(f_{\varepsilon}(x+z) - f_{\varepsilon}(x) - \langle \nabla f_{\varepsilon}(x), z \rangle\right) |z|^{-(d+\alpha)}$$

For $z = (z_1, \dots, z_d) \in \mathbb{R}^d$, let $|z|_1 := \max_{1 \leq i \leq d} |z_i|$. Write

$$\begin{split} |\hat{\mathcal{L}}_{1,\varepsilon}f_{\varepsilon}(x) - \bar{\mathcal{L}}_{\varepsilon}f_{\varepsilon}(x)| &\leq \int_{\{|z|_{1} \leq 2\varepsilon\}} |H_{\varepsilon}(x,z)| K\left(x/\varepsilon, (x+z)/\varepsilon\right) \, dz + \int_{\{|z|_{1} \leq 2\varepsilon\}} |H_{\varepsilon}(x,z)| \bar{K}\left(x/\varepsilon\right) \, dz \\ &+ \left| \int_{\{|z|_{1} > 2\varepsilon\}} H_{\varepsilon}(x,z) \left(K\left(x/\varepsilon, (x+z)/\varepsilon\right) - \bar{K}\left(x/\varepsilon\right) \right) \, dz \right| \\ &= : I_{1}^{\varepsilon}(x) + I_{2}^{\varepsilon}(x) + I_{3}^{\varepsilon}(x). \end{split}$$

By the mean-value theorem and (2.38), we obtain

$$I_1^{\varepsilon}(x) + I_2^{\varepsilon}(x) \leqslant c_1 \sup_{y:|y-x|_1 \leqslant 2\varepsilon} |\nabla^2 f_{\varepsilon}(y)| \cdot \left(\int_{\{|z|_1 \leqslant 2\varepsilon\}} \frac{|z|^2}{|z|^{d+\alpha}} dz \right)$$
$$\leqslant c_2 \varepsilon^{2-\alpha} \left(\delta_D(x)^{-2+\alpha/2} \mathbb{1}_{D_{4\varepsilon}}(x) + \varepsilon^{-2+\alpha/2} \mathbb{1}_{D \setminus D_{4\varepsilon}}(x) \right).$$

On the other hand, set

$$\begin{split} &\int_{\{|z|_1>2\varepsilon\}} H_{\varepsilon}(x,z) \left(K\left(x/\varepsilon,(x+z)/\varepsilon\right) - \bar{K}\left(x/\varepsilon\right) \right) dz \\ &= \sum_{y \in \varepsilon \mathbb{Z}^d: |y|_1>2\varepsilon} \int_{(y,y+\varepsilon e]} \left(H_{\varepsilon}(x,z) - H_{\varepsilon}(x,y) \right) K\left(x/\varepsilon,(x+z)/\varepsilon\right) dz \\ &+ \sum_{y \in \varepsilon \mathbb{Z}^d: |y|_1>2\varepsilon} \int_{(y,y+\varepsilon e]} \left(H_{\varepsilon}(x,y) - H_{\varepsilon}(x,z) \right) \bar{K}\left(x/\varepsilon\right) dz, \end{split}$$

where $e = (1, \dots, 1), (y, y + \varepsilon e] := \prod_{i=1}^{d} (y_i, y_i + \varepsilon)$ and we have used the following property

$$\int_{(y,y+\varepsilon e]} H_{\varepsilon}(x,y) K\left(x/\varepsilon, (x+z)/\varepsilon\right) dz = H_{\varepsilon}(x,y) \bar{K}\left(x/\varepsilon\right) \varepsilon^d \quad y \in \varepsilon \mathbb{Z}^d \text{ with } |y|_1 > 2\varepsilon,$$

thanks to the fact that $z \mapsto K(x, z)$ is 1-periodic. Hence,

$$I_{3}^{\varepsilon}(x) \leq \sum_{y \in \varepsilon \mathbb{Z}^{d}: |y|_{1} > 2\varepsilon} \left| \int_{(y,y+\varepsilon e]} \left(H_{\varepsilon}(x,z) - H_{\varepsilon}(x,y) \right) K\left(x/\varepsilon, (x+z)/\varepsilon\right) dz \right|$$

+
$$\sum_{y \in \varepsilon \mathbb{Z}^{d}: |y|_{1} > 2\varepsilon} \left| \int_{(y,y+\varepsilon e]} \left(H_{\varepsilon}(x,y) - H_{\varepsilon}(x,z) \right) \bar{K}\left(x/\varepsilon\right) dz \right|$$

$$\leq c_{3} \sum_{y \in \varepsilon \mathbb{Z}^{d}: |y|_{1} > 2\varepsilon} \int_{(y,y+\varepsilon e]} \left| H_{\varepsilon}(x,z) - H_{\varepsilon}(x,y) \right| dz.$$
(2.43)

According to the mean-value theorem, for every $y \in \varepsilon \mathbb{Z}^d$ with $|y|_1 > 2\varepsilon$ and $z \in (y, y + \varepsilon e]$, $|H_{\varepsilon}(x, z) - H_{\varepsilon}(x, y)|$

$$\leq |f_{\varepsilon}(x+z) - f_{\varepsilon}(x) - \langle \nabla f_{\varepsilon}(x), z \rangle| \cdot \left| \frac{1}{|y|^{d+\alpha}} - \frac{1}{|z|^{d+\alpha}} \right|$$

$$+ |(f_{\varepsilon}(x+z) - f_{\varepsilon}(x) - \langle \nabla f_{\varepsilon}(x), z \rangle) - (f_{\varepsilon}(x+y) - f_{\varepsilon}(x) - \langle \nabla f_{\varepsilon}(x), y \rangle)| \cdot \frac{1}{|y|^{d+\alpha}}$$

$$\leq c_{4}\varepsilon \left[\left(\int_{0}^{1} \int_{0}^{1} |\nabla^{2} f_{\varepsilon}(x+stz)| \, ds \, dt \right) \cdot \frac{|z|^{2}}{|z|^{d+\alpha+1}} + \left(\int_{0}^{1} |\nabla f_{\varepsilon}(x+sy+(1-s)z) - \nabla f_{\varepsilon}(x)| \, ds \right) \cdot \frac{1}{|z|^{d+\alpha}} \right]$$

$$\leq c_{5}\varepsilon \left[\left(\int_{0}^{1} \int_{0}^{1} |\nabla^{2} f_{\varepsilon}(x+stz)| \, ds \, dt + \int_{0}^{1} \int_{0}^{1} |\nabla^{2} f_{\varepsilon} \left(x+t(sy+(1-s)z)\right)| \, ds \, dt \right) \cdot \frac{1}{|z|^{d+\alpha-1}} \right]$$

$$\leq c_{6}\varepsilon \left[\int_{0}^{1} \int_{0}^{1} \left(\delta_{D}(x+stz)^{-2+\alpha/2} \mathbb{1}_{\{x+stz\in D_{3}\varepsilon\}} + \delta_{D}(x+tz)^{-2+\alpha/2} \mathbb{1}_{\{x+tz\in D_{3}\varepsilon\}} \right) + \varepsilon^{-2+\alpha/2} \mathbb{1}_{\{x+stz\in D\setminus D_{3}\varepsilon\}} + \varepsilon^{-2+\alpha/2} \mathbb{1}_{\{x+stz\in D\setminus D_{3}\varepsilon\}} \right] ds \, dt = 0$$

where in the last inequality we have used (2.38) and the fact that $y \in \varepsilon \mathbb{Z}^d$ with $|y|_1 > 2\varepsilon$ and $z \in (y, y + \varepsilon e]$. Thus,

$$I_3^{\varepsilon}(x)$$

$$\leqslant c_{7}\varepsilon \sum_{y\in\varepsilon\mathbb{Z}^{d}:|y|_{1}>2\varepsilon} \int_{0}^{1} \int_{0}^{1} \int_{(y,y+\varepsilon e]}^{1} \left(\delta_{D}(x+stz)^{-2+\alpha/2} \mathbb{1}_{\{x+stz\in D\setminus D_{3}\varepsilon\}} + \delta_{D}(x+tz)^{-2+\alpha/2} \mathbb{1}_{\{x+tz\in D_{3}\varepsilon\}} \right) \\ + \varepsilon^{-2+\alpha/2} \mathbb{1}_{\{x+stz\in D\setminus D_{3}\varepsilon\}} + \varepsilon^{-2+\alpha/2} \mathbb{1}_{\{x+tz\in D\setminus D_{3}\varepsilon\}} \right) \frac{1}{|z|^{d+\alpha-1}} dz \, ds \, dt \\ \leqslant c_{8}\varepsilon \int_{0}^{1} \int_{0}^{1} \int_{\{|z|_{1}>\varepsilon\}}^{1} \left(\delta_{D}(x+stz)^{-2+\alpha/2} \mathbb{1}_{\{x+stz\in D_{3}\varepsilon\}} + \delta_{D}(x+tz)^{-2+\alpha/2} \mathbb{1}_{\{x+tz\in D_{3}\varepsilon\}} \right) \\ + \varepsilon^{-2+\alpha/2} \mathbb{1}_{\{x+stz\in D\setminus D_{3}\varepsilon\}} + \varepsilon^{-2+\alpha/2} \mathbb{1}_{\{x+tz\in D\setminus D_{3}\varepsilon\}} \right) \frac{1}{|z|^{d+\alpha-1}} dz \, ds \, dt \\ \leqslant c_{9}\varepsilon \Big(\int_{0}^{1} \int_{0}^{1} (st)^{\alpha-1} \int_{\{|z|_{1}>st\varepsilon\}} \left(\delta_{D}(x+z)^{-2+\alpha/2} \mathbb{1}_{\{x+z\in D_{3}\varepsilon\}} + \varepsilon^{-2+\alpha/2} \mathbb{1}_{\{x+z\in D\setminus D_{3}\varepsilon\}} \right) \frac{1}{|z|^{d+\alpha-1}} dz \, ds \, dt \\ + \int_{0}^{1} \int_{0}^{1} t^{\alpha-1} \int_{\{|z|_{1}>t\varepsilon\}} \left(\delta_{D}(x+z)^{-2+\alpha/2} \mathbb{1}_{\{x+z\in D_{3}\varepsilon\}} + \varepsilon^{-2+\alpha/2} \mathbb{1}_{\{x+z\in D\setminus D_{3}\varepsilon\}} \right) \frac{1}{|z|^{d+\alpha-1}} dz \, ds \, dt$$

where in the last inequality we have used a change of variables. Therefore, according to (2.16) and (2.17), (noting that although here we use the norm $|z|_1$ in place of |z|, it is easy to verify that both (2.16) and (2.17) still hold since $|z|_1$ is equivalent with |z|), we obtain

$$I_3^{\varepsilon}(x) \leqslant c_{10} \left(\varepsilon^{2-\alpha} \delta_D(x)^{-2+\alpha/2} + \varepsilon^{\alpha/2} \delta_D(x)^{-\alpha} \right) \mathbb{1}_{D_{4\varepsilon}}(x) + c_{10} \varepsilon^{-\alpha/2} \mathbb{1}_{D \setminus D_{4\varepsilon}}(x).$$

So, putting all the estimates for $I_1^{\varepsilon}(x)$, $I_2^{\varepsilon}(x)$ and $I_3^{\varepsilon}(x)$ together, we obtain (2.41).

(2) Define

$$G_{\varepsilon}(x,z) := \left(\nabla f_{\varepsilon}(x+z) - \nabla f_{\varepsilon}(x) - \left\langle \nabla^2 f_{\varepsilon}(x), z \right\rangle \right) \frac{1}{|z|^{d+\alpha}}$$

By the mean-value theorem, we have

$$\begin{aligned} |\hat{\mathcal{L}}_{1,\varepsilon}\nabla f_{\varepsilon}(x)| &\leqslant \int_{\{|z|\leqslant 2\varepsilon\}} |G_{\varepsilon}(x,z)| K\left(x/\varepsilon, (x+z)/\varepsilon\right) \, dz + \int_{\{|z|> 2\varepsilon\}} |G_{\varepsilon}(x,z)| K\left(x/\varepsilon, (x+z)/\varepsilon\right) \, dz. \\ &=: J_{1}^{\varepsilon}(x) + J_{2}^{\varepsilon}(x). \end{aligned}$$

According to (2.38),

$$J_1^{\varepsilon}(x) \leqslant c_{11} \left(\sup_{y: |y-x| \leqslant 2\varepsilon} |\nabla^3 f_{\varepsilon}(y)| \right) \cdot \int_{\{|z| \leqslant 2\varepsilon\}} \frac{|z|^2}{|z|^{d+\alpha}} dz$$

XIN CHEN, ZHEN-QING CHEN, TAKASHI KUMAGAI AND JIAN WANG

$$\leq c_{12}\varepsilon^{2-\alpha} \left(\delta_D(x)^{-3+\alpha/2} \mathbb{1}_{D_{4\varepsilon}}(x) + \varepsilon^{-3+\alpha/2} \mathbb{1}_{D\setminus D_{4\varepsilon}}(x) \right).$$

On the other hand, by the mean-value theorem again and (2.38), we derive

$$\begin{split} J_{2}^{\varepsilon}(x) &\leqslant c_{13} \int_{0}^{1} \int_{\{|z|>2\varepsilon\}} \left| \nabla^{2} f_{\varepsilon}(x+sz) \right| \frac{1}{|z|^{d+\alpha-1}} \, dz \, ds + c_{13} |\nabla^{2} f_{\varepsilon}(x)| \cdot \int_{\{|z|>2\varepsilon\}} \frac{1}{|z|^{d+\alpha-1}} \, dz \\ &\leqslant c_{14} \int_{0}^{1} \int_{\{|z|>2\varepsilon\}} \left(\delta_{D}(x+sz)^{-2+\alpha/2} \mathbb{1}_{\{x+sz\in D_{3\varepsilon}\}} + \varepsilon^{-2+\alpha/2} \mathbb{1}_{\{x+sz\in D\setminus D_{3\varepsilon}\}} \right) \frac{1}{|z|^{d+\alpha-1}} \, dz \, ds \\ &+ c_{14} \varepsilon^{1-\alpha} \delta_{D}(x)^{-2+\alpha/2} \mathbb{1}_{D_{4\varepsilon}}(x) + c_{16} \varepsilon^{-1-\alpha/2} \mathbb{1}_{D\setminus D_{4\varepsilon}}(x) \\ &\leqslant c_{15} \int_{0}^{1} \int_{0}^{1} s^{\alpha-1} \int_{\{|z|>s\varepsilon\}} \left(\delta_{D}(x+z)^{-2+\alpha/2} \mathbb{1}_{\{x+z\in D_{3\varepsilon}\}} + \varepsilon^{-2+\alpha/2} \mathbb{1}_{\{x+z\in D\setminus D_{3\varepsilon}\}} \right) \frac{1}{|z|^{d+\alpha-1}} \, dz \, ds \, dt \\ &+ c_{15} \varepsilon^{1-\alpha} \delta_{D}(x)^{-2+\alpha/2} \mathbb{1}_{D_{4\varepsilon}}(x) + c_{17} \varepsilon^{-1-\alpha/2} \mathbb{1}_{D\setminus D_{4\varepsilon}}(x). \end{split}$$

Then by (2.16) and (2.17) we get

$$J_2^{\varepsilon}(x) \leqslant c_{18} \left(\varepsilon^{1-\alpha} \delta_D(x)^{-2+\alpha/2} + \varepsilon^{-1+\alpha/2} \delta_D(x)^{-\alpha} \right) \mathbb{1}_{D_{4\varepsilon}}(x) + c_{18} \varepsilon^{-1-\alpha/2} \mathbb{1}_{D \setminus D_{4\varepsilon}}(x).$$

Putting all the estimates above together, we obtain the second assertion (2.42).

Lemma 2.9. Suppose that $\alpha \in (0,1]$. Then, for every $f \in C^3_{(-\alpha/2)}(D)$, there are positive constants C_1 , C_2 and C_3 such that for all $\varepsilon \in (0,1)$ and $x \in D$,

$$\left| \hat{\mathcal{L}}_{2,\varepsilon} f_{\varepsilon}(x) - \bar{\mathcal{L}}_{\varepsilon} f_{\varepsilon}(x) \right| \leq C_{1} \left(\varepsilon^{\alpha/2} \delta_{D}(x)^{-\alpha} + \varepsilon \delta_{D}(x)^{-3/2} \left(1 + |\log \varepsilon| \mathbb{1}_{\{\alpha=1\}} \right) \mathbb{1}_{D_{4\varepsilon}}(x) + C_{1} \varepsilon^{-1+\alpha/2} \left(1 + |\log \varepsilon| \mathbb{1}_{\{\alpha=1\}} \right) \mathbb{1}_{D \setminus D_{4\varepsilon}}(x).$$

$$(2.44)$$

Here $f_{\varepsilon} = f\eta_{\varepsilon}$. Moreover, when $\alpha = 1$,

$$\left| \hat{\mathcal{L}}_{2,\varepsilon} \nabla f_{\varepsilon}(x) \right| \leq C_2 \left(\delta_D(x)^{-3/2} \left(1 + |\log \varepsilon| \right) + \varepsilon^{-1/2} \delta_D(x)^{-1} \right) \mathbb{1}_{D_{4\varepsilon}}(x) + C_2 \varepsilon^{-3/2} (1 + |\log \varepsilon|) \mathbb{1}_{D \setminus D_{4\varepsilon}}(x),$$

$$(2.45)$$

and, when $\alpha \in (0,1)$, for every $\beta \in (\alpha/2, \alpha)$, it holds that

$$|\mathcal{L}_{\varepsilon}\nabla f_{\varepsilon}(x)| \leq C_{3} \left(\varepsilon^{-(\beta-\alpha/2)} \delta_{D}(x)^{-1-(\alpha-\beta)} + \varepsilon^{-(\alpha-\beta)} \delta_{D}(x)^{-1-\beta+\alpha/2}\right) \mathbb{1}_{D_{4\varepsilon}}(x) + C_{3}\varepsilon^{-1-\alpha/2} \mathbb{1}_{D \setminus D_{4\varepsilon}}(x).$$

$$(2.46)$$

Proof. (i) Set

$$H_{\varepsilon}(x,z) := \left(f_{\varepsilon}(x+z) - f_{\varepsilon}(x) - \langle \nabla f_{\varepsilon}(x), z \rangle \, \mathbb{1}_{\{|z| \leq 1\}} \right) \frac{1}{|z|^{d+\alpha}}$$

Then

$$\begin{split} |\hat{\mathcal{L}}_{2,\varepsilon}f_{\varepsilon}(x) - \bar{\mathcal{L}}_{\varepsilon}f_{\varepsilon}(x)| &\leq \int_{\{|z|_{1} \leq 2\varepsilon\}} |H_{\varepsilon}(x,z)| K\left(x/\varepsilon, (x+z)/\varepsilon\right) \, dz + \int_{\{|z|_{1} \leq 2\varepsilon\}} |H_{\varepsilon}(x,z)| \bar{K}\left(x/\varepsilon\right) \, dz \\ &+ \left| \int_{\{|z|_{1} > 2\varepsilon\}} H_{\varepsilon}(x,z) \left(K\left(x/\varepsilon, (x+z)/\varepsilon\right) - \bar{K}\left(x/\varepsilon\right) \right) \, dz \right| \\ &=: I_{1}^{\varepsilon}(x) + I_{2}^{\varepsilon}(x) + I_{3}^{\varepsilon}(x). \end{split}$$

According to the mean-value theorem and (2.38), we have

$$I_1^{\varepsilon}(x) + I_2^{\varepsilon}(x) \leqslant c_1 \sup_{y:|y-x| \leqslant 2\varepsilon} |\nabla^2 f_{\varepsilon}(y)| \cdot \left(\int_{\{|z| \leqslant 2\varepsilon\}} \frac{|z|^2}{|z|^{d+\alpha}} dz \right)$$
$$\leqslant c_2 \varepsilon^{2-\alpha} \left(\delta_D(x)^{-2+\alpha/2} \mathbb{1}_{D_{4\varepsilon}}(x) + \varepsilon^{-2+\alpha/2} \mathbb{1}_{D \setminus D_{4\varepsilon}}(x) \right).$$

On the other hand, following the same argument as that for (2.43), we obtain

$$I_3^{\varepsilon}(x) \leqslant c_3 \sum_{y \in \varepsilon \mathbb{Z}^d : |y|_1 > 2\varepsilon} \int_{(y, y + \varepsilon e]} |H_{\varepsilon}(x, y) - H_{\varepsilon}(x, z)| \, dz.$$

Applying the mean-value theorem and (2.38) again (by following the same argument as that in the proof of Lemma 2.8), we obtain that for every $x \in D$, $y \in \varepsilon \mathbb{Z}^d$ with $|y|_1 > 2\varepsilon$ and $z \in (y, y + \varepsilon e]$, $|H_{\varepsilon}(x, z) - H_{\varepsilon}(x, y)|$

$$\begin{split} \|I_{\varepsilon}(x,z) - H_{\varepsilon}(x,y)\| \\ &\leqslant |f_{\varepsilon}(x+z) - f_{\varepsilon}(x) - \langle \nabla f_{\varepsilon}(x), z \rangle \mathbb{1}_{\{|z| \leqslant 1-\varepsilon\}} | \left| \frac{1}{|y|^{d+\alpha}} - \frac{1}{|z|^{d+\alpha}} \right| \\ &+ |(f_{\varepsilon}(x+z) - f_{\varepsilon}(x) - \langle \nabla f_{\varepsilon}(x), z \rangle \mathbb{1}_{\{|z| \leqslant 1-\varepsilon\}}) - (f_{\varepsilon}(x+y) - f_{\varepsilon}(x) - \langle \nabla f_{\varepsilon}(x), y \rangle \mathbb{1}_{\{|z| \leqslant 1-\varepsilon\}}) | \frac{1}{|y|^{d+\alpha}} \\ &+ \frac{|\langle \nabla f_{\varepsilon}(x), z \rangle|}{|z|^{d+\alpha}} \mathbb{1}_{\{1-\varepsilon \leqslant |z| \leqslant 1\}} + \frac{|\langle \nabla f_{\varepsilon}(x), y \rangle|}{|y|^{d+\alpha}} \mathbb{1}_{\{1-(1+\sqrt{d})\varepsilon \leqslant |y| \leqslant 1+(1+\sqrt{d})\varepsilon\}} \\ &\leqslant c_{4}\varepsilon \left(\int_{0}^{1} \int_{0}^{1} |\nabla^{2} f_{\varepsilon}(x+stz)| \, ds \, dt + \int_{0}^{1} \int_{0}^{1} |\nabla^{2} f_{\varepsilon}(x+t(sy+(1-s)z))| \, ds \, dt \right) \cdot \frac{1}{|z|^{d+\alpha-1}} \mathbb{1}_{\{|z| \leqslant 1\}} \\ &+ c_{4}\varepsilon \left(\int_{0}^{1} |\nabla f_{\varepsilon}(x+sz)| \, ds + \int_{0}^{1} |\nabla f_{\varepsilon}(x+sy+(1-s)z)| \, ds \right) \cdot \frac{1}{|z|^{d+\alpha}} \mathbb{1}_{\{|z|>1\}} \\ &+ c_{4}|\nabla f_{\varepsilon}(x)| \left(\mathbb{1}_{\{1-\varepsilon \leqslant |z| \leqslant 1\}} + \mathbb{1}_{\{1-(1+\sqrt{d})\varepsilon \leqslant |y| \leqslant 1+(1+\sqrt{d})\varepsilon\}} \right) \\ &\leqslant c_{5}\varepsilon \left[\int_{0}^{1} \int_{0}^{1} \left(\delta_{D}(x+stz)^{-2+\alpha/2} \mathbb{1}_{\{x+stz \in D_{3}\varepsilon\}} + \delta_{D}(x+tz)^{-2+\alpha/2} \mathbb{1}_{\{x+stz \in D_{3}\varepsilon\}} + \varepsilon^{-2+\alpha/2} \mathbb{1}_{\{x+stz \in D_{3}\varepsilon\}} \right) \\ &+ \varepsilon^{-2+\alpha/2} \mathbb{1}_{\{x+stz \in D \setminus D_{3}\varepsilon\}} \right) \, ds \, dt \right] \cdot \frac{1}{|z|^{d+\alpha-1}} \mathbb{1}_{\{|z| \leqslant 1\}} + c_{5}\varepsilon \left[\int_{0}^{1} \left(\delta_{D}(x+sz)^{-1+\alpha/2} \mathbb{1}_{\{x+sz \in D \setminus D_{3}\varepsilon\}} + \delta_{D}(x+z)^{-1+\alpha/2} \mathbb{1}_{\{x+sz \in D \setminus D_{3}\varepsilon\}} \right) \, ds \right] \cdot \frac{1}{|z|^{d+\alpha}} \mathbb{1}_{\{|z|>1\}} \\ &+ c_{5} \left(\delta_{D}(x)^{-1+\alpha/2} \mathbb{1}_{D_{3}\varepsilon}(x) + \varepsilon^{-1+\alpha/2} \mathbb{1}_{D \setminus D_{3}\varepsilon}(x) \right) \left(\mathbb{1}_{\{1-\varepsilon \leqslant |z| \leqslant 1\}} + \mathbb{1}_{\{1-(1+\sqrt{d})\varepsilon \leqslant |y| \leqslant 1+(1+\sqrt{d})\varepsilon\}} \right). \end{split}$$

This, along with the proof of the estimate for $I_3^{\varepsilon}(x)$ in Lemma 2.8 via a change of variables, yields that $I_3^{\varepsilon}(x)$

$$\leq c_{6}\varepsilon \left[\int_{0}^{1} \int_{0}^{1} (st)^{\alpha-1} \int_{\{st\varepsilon < |z|_{1} \leqslant st\}} \left(\delta_{D}(x+z)^{-2+\alpha/2} \mathbb{1}_{\{x+z\in D_{3}\varepsilon\}} + \varepsilon^{-2+\alpha/2} \mathbb{1}_{\{x+z\in D\setminus D_{3}\varepsilon\}} \right) \frac{1}{|z|^{d+\alpha-1}} dz \, ds \, dt \\ + \int_{0}^{1} \int_{0}^{1} s^{\alpha-1} \int_{\{s\varepsilon < |z|_{1} \leqslant s\}} \left(\delta_{D}(x+z)^{-2+\alpha/2} \mathbb{1}_{\{x+z\in D_{3}\varepsilon\}} + \varepsilon^{-2+\alpha/2} \mathbb{1}_{\{x+z\in D\setminus D_{3}\varepsilon\}} \right) \frac{1}{|z|^{d+\alpha-1}} dz \, ds \, dt \right] \\ + c_{6}\varepsilon \left[\int_{0}^{1} s^{\alpha} \int_{\{|z|>s\}} \left(\delta_{D}(x+z)^{-1+\alpha/2} \mathbb{1}_{\{x+z\in D_{3}\varepsilon\}} + \varepsilon^{-1+\alpha/2} \mathbb{1}_{\{x+z\in D\setminus D_{3}\varepsilon\}} \right) \frac{1}{|z|^{d+\alpha}} dz \, ds \\ + \int_{\{|z|>1\}} \left(\delta_{D}(x+z)^{-1+\alpha/2} \mathbb{1}_{\{x+z\in D_{3}\varepsilon\}} + \varepsilon^{-1+\alpha/2} \mathbb{1}_{\{x+z\in D\setminus D_{3}\varepsilon\}} \right) \frac{1}{|z|^{d+\alpha}} dz \right] \\ + c_{6}\varepsilon \left(\delta_{D}(x)^{-1+\alpha/2} \mathbb{1}_{D_{3}\varepsilon}(x) + \varepsilon^{-1+\alpha/2} \mathbb{1}_{D\setminus D_{3}\varepsilon}(x) \right).$$

Now, by (2.18)-(2.20), we have

$$I_{3}^{\varepsilon}(x) \leq c_{7} \left(\varepsilon (\delta_{D}(x)^{-1-\alpha/2} + \delta_{D}(x)^{-2+\alpha/2} (1+|\log\varepsilon|) \mathbb{1}_{\{\alpha=1\}}) + \varepsilon^{\alpha/2} (\delta_{D}(x)^{-\alpha} + 1) \right) \mathbb{1}_{D_{4\varepsilon}}(x)$$

$$+ c_{7} \left(\varepsilon^{-1+\alpha/2} (1+|\log\varepsilon| \mathbb{1}_{\{\alpha=1\}}) + \varepsilon^{\alpha/2} \right) \mathbb{1}_{D \setminus D_{4\varepsilon}}(x)$$

$$\leq c_{8} \left(\varepsilon^{\alpha/2} \delta_{D}(x)^{-\alpha} + \varepsilon \delta_{D}(x)^{-3/2} (1+|\log\varepsilon|) \mathbb{1}_{\{\alpha=1\}} \right) \mathbb{1}_{D_{4\varepsilon}}(x)$$

$$+ c_{8} \varepsilon^{-1+\alpha/2} \left(1+|\log\varepsilon| \mathbb{1}_{\{\alpha=1\}} \right) \mathbb{1}_{D \setminus D_{4\varepsilon}}(x),$$

where in the last inequality we used the fact that $\sup_{x\in D} \delta_D(x) < \infty$ since D is bounded. Putting all the estimates above for $I_1^{\varepsilon}(x)$, $I_2^{\varepsilon}(x)$, $I_3^{\varepsilon}(x)$ together, we arrive at

$$\begin{aligned} &|\hat{\mathcal{L}}_{2,\varepsilon}f_{\varepsilon}(x) - \bar{\mathcal{L}}_{\varepsilon}f_{\varepsilon}(x)| \\ &\leqslant c_9 \left(\varepsilon^{\alpha/2} \delta_D(x)^{-\alpha} + \varepsilon^{2-\alpha} \delta_D(x)^{-2+\alpha/2} + \varepsilon \delta_D(x)^{-3/2} \left(1 + |\log \varepsilon|\right) \mathbb{1}_{\{\alpha=1\}} \right) \mathbb{1}_{D_{4\varepsilon}}(x) \\ &+ c_9 \left(\varepsilon^{-1+\alpha/2} \left(1 + |\log \varepsilon| \mathbb{1}_{\{\alpha=1\}} \right) + \varepsilon^{-\alpha/2} \right) \mathbb{1}_{D \setminus D_{4\varepsilon}}(x) \\ &\leqslant c_{10} \left(\varepsilon^{\alpha/2} \delta_D(x)^{-\alpha} + \varepsilon \delta_D(x)^{-3/2} \left(1 + |\log \varepsilon|\right) \mathbb{1}_{\{\alpha=1\}} \right) \mathbb{1}_{D_{4\varepsilon}}(x) \\ &+ c_{10} \varepsilon^{-1+\alpha/2} \left(1 + |\log \varepsilon| \mathbb{1}_{\{\alpha=1\}} \right) \mathbb{1}_{D \setminus D_{4\varepsilon}}(x). \end{aligned}$$

So (2.44) is proved.

(ii) When $\alpha = 1$, define

$$G_{\varepsilon}(x,z) := \left(\nabla f_{\varepsilon}(x+z) - \nabla f_{\varepsilon}(x) - \left\langle \nabla^2 f_{\varepsilon}(x), z \right\rangle \mathbb{1}_{\{|z| \leq 1\}} \right) \frac{1}{|z|^{d+1}}.$$

Then,

$$\begin{split} |\hat{\mathcal{L}}_{2,\varepsilon}\nabla f_{\varepsilon}(x)| &\leqslant \int_{\{|z|\leqslant 2\varepsilon\}} |G_{\varepsilon}(x,z)| K\left(x/\varepsilon, (x+z)/\varepsilon\right) \, dz + \int_{\{|z|> 2\varepsilon\}} |G_{\varepsilon}(x,z)| K\left(x/\varepsilon, (x+z)/\varepsilon\right) \, dz \\ &=: J_{1}^{\varepsilon}(x) + J_{2}^{\varepsilon}(x). \end{split}$$

Thanks to (2.38), we obtain

$$J_1^{\varepsilon}(x) \leqslant c_{11} \left(\sup_{y:|y-x|\leqslant 2\varepsilon} |\nabla^3 f_{\varepsilon}(y)| \right) \cdot \int_{\{|z|\leqslant 2\varepsilon\}} \frac{|z|^2}{|z|^{d+1}} dz$$

$$\leqslant c_{12}\varepsilon \left(\delta_D(x)^{-5/2} \mathbb{1}_{D_{4\varepsilon}}(x) + \varepsilon^{-5/2} \mathbb{1}_{D\setminus D_{4\varepsilon}}(x) \right).$$

On the other hand, by the mean-value theorem again and (2.38), we derive

$$\begin{split} J_{2}^{\varepsilon}(x) &\leqslant c_{13} \int_{0}^{1} \int_{\{2\varepsilon < |z| \leqslant 1\}} |\nabla^{2} f_{\varepsilon}(x+sz)| \frac{1}{|z|^{d}} dz \, ds + c_{13} |\nabla^{2} f_{\varepsilon}(x)| \cdot \int_{\{2\varepsilon < |z| \leqslant 1\}} \frac{1}{|z|^{d}} dz \\ &+ c_{13} \int_{\{|z|>1\}} |\nabla f_{\varepsilon}(x+z)| \frac{1}{|z|^{d+1}} dz + c_{13} |\nabla f_{\varepsilon}(x)| \cdot \int_{\{|z|>1\}} \frac{1}{|z|^{d+1}} dz \\ &\leqslant c_{14} \int_{0}^{1} \int_{\{2\varepsilon < |z| \leqslant 1\}} \left(\delta_{D}(x+sz)^{-3/2} \mathbb{1}_{\{x+sz \in D_{3}\varepsilon\}} + \varepsilon^{-3/2} \mathbb{1}_{\{x+sz \in D \setminus D_{3}\varepsilon\}} \right) \frac{1}{|z|^{d}} dz \, ds \\ &+ c_{14} \int_{\{|z|>1\}} \left(\delta_{D}(x+z)^{-1/2} \mathbb{1}_{\{x+z \in D_{3}\varepsilon\}} + \varepsilon^{-1/2} \mathbb{1}_{\{x+z \in D \setminus D_{3}\varepsilon\}} \right) \frac{1}{|z|^{d+1}} dz \\ &+ c_{14} \left(1 + |\log \varepsilon| \right) \left(\delta_{D}(x)^{-3/2} \mathbb{1}_{D_{4\varepsilon}}(x) + \varepsilon^{-3/2} \mathbb{1}_{D \setminus D_{4\varepsilon}}(x) \right) \\ &+ c_{14} \left(\delta_{D}(x)^{-1/2} \mathbb{1}_{D_{4\varepsilon}}(x) + \varepsilon^{-1/2} \mathbb{1}_{\{x+z \in D_{3}\varepsilon\}} + \varepsilon^{-3/2} \mathbb{1}_{\{x+z \in D \setminus D_{3}\varepsilon\}} \right) \frac{1}{|z|^{d}} dz \, ds \\ &+ c_{15} \int_{\{|z|>1\}} \left(\delta_{D}(x+z)^{-1/2} \mathbb{1}_{\{x+z \in D_{3}\varepsilon\}} + \varepsilon^{-3/2} \mathbb{1}_{\{x+z \in D \setminus D_{3}\varepsilon\}} \right) \frac{1}{|z|^{d+1}} dz \\ &+ c_{15} \left(\int_{|z|>1} \left(\delta_{D}(x+z)^{-1/2} \mathbb{1}_{\{x+z \in D_{3}\varepsilon\}} + \varepsilon^{-1/2} \mathbb{1}_{\{x+z \in D \setminus D_{3}\varepsilon\}} \right) \frac{1}{|z|^{d+1}} dz \\ &+ c_{15} \left(1 + |\log \varepsilon| \right) \left(\delta_{D}(x)^{-3/2} \mathbb{1}_{D_{4\varepsilon}}(x) + \varepsilon^{-3/2} \mathbb{1}_{D \setminus D_{4\varepsilon}}(x) \right) \\ &+ c_{15} \left(\delta_{D}(x)^{-1/2} \mathbb{1}_{D_{4\varepsilon}}(x) + \varepsilon^{-1/2} \mathbb{1}_{D \setminus D_{4\varepsilon}}(x) \right) . \end{split}$$

Hence, applying (2.18) - (2.20), we get

$$J_{2}^{\varepsilon}(x) \leq c_{16} \left(\delta_{D}(x)^{-3/2} \left(1 + |\log \varepsilon| \right) + \varepsilon^{-1/2} \delta_{D}(x)^{-1} \right) \mathbb{1}_{D_{4\varepsilon}}(x) + c_{16} \varepsilon^{-3/2} \left(1 + |\log \varepsilon| \right) \mathbb{1}_{D \setminus D_{4\varepsilon}}(x).$$

Putting both estimates for $J_1(x)$ and $J_2(x)$ together, we obtain the desired conclusion (2.45).

(iii) Note that, when $\alpha \in (0, 1)$, $f \in C^3_{(-\alpha/2)}(D) \subset C^{1+\beta}_{(-\alpha/2)}(D)$ for every $\beta \in (\alpha/2, \alpha)$. By (2.31) and the definition of f_{ε} , it holds that for all $x, z \in \mathbb{R}^d$ with $|z| \ge 2\varepsilon$

$$\frac{|\nabla f_{\varepsilon}(x) - \nabla f_{\varepsilon}(x+z)|}{|z|^{\beta}} \leqslant c_{17} \Big[\delta_D(x+z)^{-1-\beta+\alpha/2} \mathbb{1}_{\{x+z\in D_{2\varepsilon}\}} + \delta_D(x)^{-1-\beta+\alpha/2} \mathbb{1}_{\{x\in D_{2\varepsilon}\}} + \varepsilon^{-1-\beta+\alpha/2} \Big(\mathbb{1}_{\{x+z\in D\setminus D_{2\varepsilon}\}} + \mathbb{1}_{\{x\in D\setminus D_{2\varepsilon}\}} \Big) \Big].$$

$$(2.47)$$

Here we have also used the fact that for every $|z| \ge 2\varepsilon$, $x + z \in D_{2\varepsilon}$ and $x \in D \setminus D_{2\varepsilon}$

$$\begin{aligned} |\nabla f_{\varepsilon}(x+z) - \nabla f_{\varepsilon}(x)| &\leq |\nabla f_{\varepsilon}(x+z)| + |\nabla f_{\varepsilon}(x)| \\ &\leq c_{18} \left(\delta_D(x+z)^{-1+\alpha/2} \mathbb{1}_{\{x+z\in D_{\varepsilon}\}} + \delta_D(x)^{-1+\alpha/2} \mathbb{1}_{\{x\in D_{\varepsilon}\}} \right) \cdot \left(\frac{|z|}{2\varepsilon}\right)^{\beta} \\ &\leq c_{19}\varepsilon^{-1-\beta+\alpha/2} |z|^{\beta}. \end{aligned}$$

Now we write

$$\begin{aligned} |\mathcal{L}_{\varepsilon} \nabla f_{\varepsilon}(x)| &\leqslant \int_{\{|z| \leqslant 2\varepsilon\}} |\nabla f_{\varepsilon}(x+z) - \nabla f_{\varepsilon}(x)| \frac{K\left(x/\varepsilon, (x+z)/\varepsilon\right)}{|z|^{d+\alpha}} \, dz \\ &+ \int_{\{|z| > 2\varepsilon\}} |\nabla f_{\varepsilon}(x+z) - \nabla f_{\varepsilon}(x)| \frac{K\left(x/\varepsilon, (x+z)/\varepsilon\right)}{|z|^{d+\alpha}} \, dz. \\ &=: L_{1}^{\varepsilon}(x) + L_{2}^{\varepsilon}(x). \end{aligned}$$

It follows from (2.38) that

$$L_1^{\varepsilon}(x) \leq c_{20} \left(\sup_{y:|y-x|\leq 2\varepsilon} |\nabla^2 f_{\varepsilon}(y)| \right) \cdot \int_{\{|z|\leq 2\varepsilon\}} \frac{|z|}{|z|^{d+\alpha}} dz$$

$$\leq c_{21} \varepsilon^{1-\alpha} \left(\delta_D(x)^{-2+\alpha/2} \mathbb{1}_{D_{4\varepsilon}}(x) + \varepsilon^{-2+\alpha/2} \mathbb{1}_{D\setminus D_{4\varepsilon}}(x) \right).$$

On the other hand, according to (2.47),

$$L_{2}^{\varepsilon}(x) \leq c_{22} \int_{\{|z|>2\varepsilon\}} \left(\delta_{D}(x+z)^{-1-\beta+\alpha/2} \mathbb{1}_{\{x+z\in D_{3\varepsilon}\}} + \varepsilon^{-1-\beta+\alpha/2} \mathbb{1}_{\{x+z\in D_{3\varepsilon}\}} \right) \frac{1}{|z|^{d+\alpha-\beta}} dz + c_{22} \left(\delta_{D}(x)^{-1-\beta+\alpha/2} \mathbb{1}_{D_{4\varepsilon}}(x) + \varepsilon^{-1-\beta+\alpha/2} \mathbb{1}_{D\setminus D_{4\varepsilon}}(x) \right) \cdot \int_{\{|z|>2\varepsilon\}} \frac{1}{|z|^{d+\alpha-\beta}} dz \leq c_{23} \left(\varepsilon^{-(\beta-\alpha/2)} \delta_{D}(x)^{-1-(\alpha-\beta)} + \varepsilon^{-(\alpha-\beta)} \delta_{D}(x)^{-1-\beta+\alpha/2} \right) \mathbb{1}_{D_{4\varepsilon}}(x) + c_{23} \varepsilon^{-1-\alpha/2} \mathbb{1}_{D\setminus D_{4\varepsilon}}(x),$$

where in the last inequality we used (2.21), (2.22) and the fact $\beta \in (\alpha/2, \alpha)$.

Putting the estimates for $L_1^{\varepsilon}(x)$ and $L_2^{\varepsilon}(x)$ together, we obtain (2.46).

Lemma 2.10. The following statements hold.

(1) For all $\alpha \in (0,2)$ and $f \in C^3_{(-\alpha/2)}(D)$, there is a constant $C_1 > 0$ such that for all $\varepsilon \in (0,1)$ and $x \in D$,

$$\bar{\mathcal{L}}f_{\varepsilon}(x) - \eta_{\varepsilon}(x)\bar{\mathcal{L}}f(x) \Big| \leqslant C_1 \left(\varepsilon \delta_D(x)^{-1-\alpha/2} \mathbb{1}_{D_{4\varepsilon}}(x) + \varepsilon^{-\alpha/2} \mathbb{1}_{D \setminus D_{4\varepsilon}}(x) \right),$$
(2.48)

where $f_{\varepsilon} := f \eta_{\varepsilon}$.

(2) When $\alpha \in (1,2)$ and $f \in C^4_{(-\alpha/2)}(D)$, there is a constant $C_2 > 0$ such that for all $\varepsilon \in (0,1)$, $x \in D$ and k = 1, 2,

$$\left|\nabla^{k}\bar{\mathcal{L}}f_{\varepsilon}(x)\right| \leq C_{2}(\varepsilon^{1-\alpha}\delta_{D}(x)^{-k-1+\alpha/2} + \varepsilon^{-k+\alpha/2}\delta_{D}(x)^{-\alpha})\mathbb{1}_{D_{4\varepsilon}}(x) + C_{2}\varepsilon^{-k-\alpha/2}\mathbb{1}_{D\setminus D_{4\varepsilon}}(x).$$
(2.49)

(3) When $\alpha \in (0,1]$ and $f \in C^4_{(-\alpha/2)}(D)$, there is a constant $C_3 > 0$ such that for all $\varepsilon \in (0,1)$, $x \in D$ and k = 1, 2,

$$\left|\nabla^{k}\bar{\mathcal{L}}f_{\varepsilon}(x)\right| \leq C_{3}\left(\delta_{D}(x)^{-k-\alpha/2}\mathbb{1}_{\{\alpha\in(0,1)\}} + \delta_{D}(x)^{-k-\alpha/2}\left(1+\left|\log\varepsilon\right|\right)\mathbb{1}_{\{\alpha=1\}}\right)\mathbb{1}_{D_{4\varepsilon}}(x) + C_{3}\varepsilon^{-k+\alpha/2}\delta_{D}(x)^{-\alpha}\mathbb{1}_{D_{4\varepsilon}}(x) + C_{3}\varepsilon^{-k-\alpha/2}\left(1+\left|\log\varepsilon\right|\mathbb{1}_{\{\alpha=1\}}\right)\mathbb{1}_{D\setminus D_{4\varepsilon}}(x).$$

$$(2.50)$$

Proof. (1) By some direct computations, we have

$$\bar{\mathcal{L}}f_{\varepsilon}(x) = \eta_{\varepsilon}(x)\bar{\mathcal{L}}f(x) + f(x)\bar{\mathcal{L}}\eta_{\varepsilon}(x) + \int_{\mathbb{R}^{d}} \left(f(x+z) - f(x)\right)\left(\eta_{\varepsilon}(x+z) - \eta_{\varepsilon}(x)\right)\frac{K}{|z|^{d+\alpha}}dz$$

$$= \eta_{\varepsilon}(x)\bar{\mathcal{L}}f(x) + f(x)\bar{\mathcal{L}}\eta_{\varepsilon}(x) + \int_{\mathbb{R}^{d}} \left(f(x+z) - f(x)\right)\left(\eta_{\varepsilon}(x+z) - \eta_{\varepsilon}(x)\right)\frac{\bar{K}}{|z|^{d+\alpha}}dz$$

$$=: \eta_{\varepsilon}(x)\bar{\mathcal{L}}f(x) + I_{1}^{\varepsilon}(x).$$
(2.51)

Combining (2.32), (2.34) with the fact that $|f(x)| \leq c_1 \delta_D(x)^{\alpha/2}$, we can see

$$|I_1^{\varepsilon}(x)| \leq c_2 \left(\varepsilon \delta_D(x)^{-1-\alpha/2} \mathbb{1}_{D_{4\varepsilon}}(x) + \varepsilon^{-\alpha/2} \mathbb{1}_{D \setminus D_{4\varepsilon}}(x) \right).$$

With this we can verify (2.48) immediately.

(2) For simplicity we only give a proof for (2.49) when k = 1. The k = 2 case can be shown analogously. Suppose that $\alpha \in (1, 2)$. Then, by (2.38), the mean-value theorem and a change of variable as well as (2.16), it holds

$$\begin{split} |\nabla \bar{\mathcal{L}} f_{\varepsilon}(x)| &= |\bar{\mathcal{L}}(\nabla f_{\varepsilon})(x)| \\ &\leqslant \int_{\{|z| \leqslant 2\varepsilon\}} |\nabla f_{\varepsilon}(x+z) - \nabla f_{\varepsilon}(x) - \left\langle \nabla^{2} f_{\varepsilon}(x), z \right\rangle \Big| \frac{\bar{K}}{|z|^{d+\alpha}} dz + \int_{\{|z| > 2\varepsilon\}} |\nabla f_{\varepsilon}(x+z) - \nabla f_{\varepsilon}(x)| \frac{\bar{K}}{|z|^{d+\alpha}} dz \\ &\leqslant c_{3} \sup_{y \in \mathbb{R}^{d}: |y-x| \leqslant 2\varepsilon} |\nabla^{3} f_{\varepsilon}(y)| \cdot \int_{\{|z| \leqslant 2\varepsilon\}} \frac{|z|^{2}}{|z|^{d+\alpha}} dz + c_{3} \int_{0}^{1} \int_{\{|z| > 2\varepsilon\}} |\nabla^{2} f_{\varepsilon}(x+sz)| \frac{1}{|z|^{d+\alpha-1}} dz \, ds \\ &\leqslant c_{4} \varepsilon^{2-\alpha} \delta_{D}(x)^{-3+\alpha/2} \mathbb{1}_{D_{4\varepsilon}}(x) + c_{4} \varepsilon^{-1-\alpha/2} \mathbb{1}_{D \setminus D_{4\varepsilon}}(x) \\ &+ c_{4} \int_{0}^{1} s^{1-\alpha} \int_{\{|z| > 2s\varepsilon\}} \left(\delta_{D}(x+z)^{-2+\alpha/2} \mathbb{1}_{D_{3\varepsilon}}(x+z) + \varepsilon^{-2+\alpha/2} \mathbb{1}_{D \setminus D_{3\varepsilon}}(x+z) \right) \frac{1}{|z|^{d+\alpha-1}} dz \, ds \\ &\leqslant c_{5} \left(\varepsilon^{1-\alpha} \delta_{D}(x)^{-2+\alpha/2} + \varepsilon^{-1+\alpha/2} \delta_{D}(x)^{-\alpha} \right) \mathbb{1}_{D_{4\varepsilon}}(x) + c_{5} \varepsilon^{-1-\alpha/2} \mathbb{1}_{D \setminus D_{4\varepsilon}}(x). \end{split}$$

Hence, (2.49) is established.

(3) Here we only prove (2.50) for k = 1 and $\alpha = 1$, since other cases can be proved by the same way. By (2.38) and a change of variable, it holds

$$\begin{split} |\nabla \bar{\mathcal{L}} f_{\varepsilon}(x)| &= |\bar{\mathcal{L}}(\nabla f_{\varepsilon})(x)| \\ &\leqslant \int_{\{|z| \leq 2\varepsilon\}} \left| \nabla f_{\varepsilon}(x+z) - \nabla f_{\varepsilon}(x) - \left\langle \nabla^{2} f_{\varepsilon}(x), z \right\rangle \right| \frac{\bar{K}}{|z|^{d+1}} dz + \int_{\{2\varepsilon < |z| \leq 1\}} \left| \nabla f_{\varepsilon}(x+z) - \nabla f_{\varepsilon}(x) \right| \frac{\bar{K}}{|z|^{d+1}} dz \\ &+ \int_{\{|z| > 1\}} \left(|\nabla f_{\varepsilon}(x+z)| + |\nabla f_{\varepsilon}(x)| \right) \frac{\bar{K}}{|z|^{d+1}} dz \\ &\leqslant c_{6} \sup_{y \in \mathbb{R}^{d}: |y-x| \leq 2\varepsilon} |\nabla^{3} f_{\varepsilon}(y)| \cdot \int_{\{|z| \leq 2\varepsilon\}} \frac{|z|^{2}}{|z|^{d+1}} dz \\ &+ c_{6} \int_{0}^{1} \int_{\{2\varepsilon < |z| \leq 1\}} |\nabla^{2} f_{\varepsilon}(x+sz)| \frac{1}{|z|^{d}} dz \, ds + c_{6} \int_{\{|z| > 1\}} |\nabla f_{\varepsilon}(x+z)| \frac{1}{|z|^{d+1}} dz + c_{6} |\nabla f_{\varepsilon}(x)| \\ &\leqslant c_{7} \varepsilon \delta_{D}(x)^{-5/2} \mathbbm{1}_{D_{4\varepsilon}}(x) + c_{7} \varepsilon^{-3/2} \mathbbm{1}_{D \setminus D_{4\varepsilon}}(x) \\ &+ c_{7} \int_{0}^{1} \int_{\{2s\varepsilon < |z| \leq 1\}} \left(\delta_{D}(x+z)^{-3/2} \mathbbm{1}_{D_{3\varepsilon}}(x+z) + \varepsilon^{-3/2} \mathbbm{1}_{D \setminus D_{3\varepsilon}}(x+z) \right) \frac{1}{|z|^{d}} dz \, ds \end{split}$$

$$+ c_{7} \int_{\{|z|>1\}} \left(\delta_{D}(x+z)^{-1/2} \mathbb{1}_{D_{3\varepsilon}}(x+z) + \varepsilon^{-1/2} \mathbb{1}_{D \setminus D_{3\varepsilon}}(x+z) \right) \frac{1}{|z|^{d}} dz \\ + c_{7} \left(\delta_{D}(x)^{-1/2} \mathbb{1}_{D_{3\varepsilon}}(x) + \varepsilon^{-1/2} \mathbb{1}_{D \setminus D_{3\varepsilon}}(x) \right) \\ \leqslant c_{8} \left(\varepsilon^{-1/2} \delta_{D}(x)^{-1} + (1+|\log\varepsilon|) \delta_{D}(x)^{-3/2} \right) \mathbb{1}_{D_{4\varepsilon}}(x) + c_{8} \varepsilon^{-3/2} (1+|\log\varepsilon|) \mathbb{1}_{D \setminus D_{4\varepsilon}}(x),$$

where the last step follows from (2.18) and (the proof of) (2.19). This completes the proof.

3. QUANTITATIVE HOMOGENIZATIONS

This section is devoted to the proofs of the main results of this paper, Theorems 1.1 and 1.2. When $\alpha \in (1,2)$, we define

$$F(x) = (F^{(1)}(x), \cdots, F^{(d)}), \quad F^{(i)}(x) := \text{p.v.} \int_{\mathbb{R}^d} z_i \frac{K(x, x+z)}{|z|^{d+\alpha}} \, dz, \quad i = 1, \cdots, d,$$

where z_i is the *i*th coordinate of $z \in \mathbb{R}^d$. By the symmetry and the multivariate 1-periodicity of K(x, y) on $\mathbb{R}^d \times \mathbb{R}^d$, as well as a change of variable, we have

$$\begin{split} \int_{\mathbb{T}^d} F^{(i)}(x) \, dx &= \int_{\mathbb{T}^d} \left(\text{p.v.} \int_{\mathbb{R}^d} z_i \frac{K(x, x+z)}{|z|^{d+\alpha}} \, dz \right) dx = \text{p.v.} \int_{\mathbb{R}^d} z_i \left(\int_{\mathbb{T}^d} \frac{K(x+z, x)}{|z|^{d+\alpha}} \, dx \right) dz \\ &= \text{p.v.} \int_{\mathbb{R}^d} z_i \left(\int_{\mathbb{T}^d} \frac{K(x, x-z)}{|z|^{d+\alpha}} \, dx \right) dz = -\int_{\mathbb{T}^d} \left(\text{p.v.} \int_{\mathbb{R}^d} z_i \frac{K(x, x+z)}{|z|^{d+\alpha}} \, dz \right) dx \\ &= -\int_{\mathbb{T}^d} F^{(i)}(x) \, dx. \end{split}$$

Thus $\int_{\mathbb{T}^d} F^{(i)}(x) dx = 0$. On the other hand, since $K \in C^2(\mathbb{T}^d \times \mathbb{T}^d)$, for every $x \in \mathbb{T}^d$ and $z \in \mathbb{R}^d$ it holds that

$$\begin{aligned} \left| z \frac{K(x, x+z) - K(x, x-z)}{|z|^{d+\alpha}} \right| &\leq 2 \left(\sup_{x, y \in \mathbb{T}^d} |\nabla_y K(x, y)| \right) \frac{1}{|z|^{d+\alpha-2}} \mathbb{1}_{\{|z| \leq 1\}} \\ &+ \left(\sup_{x, y \in \mathbb{T}^d} |K(x, y)| \right) \frac{1}{|z|^{d+\alpha-1}} \mathbb{1}_{\{|z| > 1\}}. \end{aligned}$$

This implies that

$$F(x) = \lim_{\varepsilon \downarrow 0} \frac{1}{2} \int_{\{|z| > \varepsilon\}} z \frac{K(x, x+z) - K(x, x-z)}{|z|^{d+\alpha}} dz$$

is well defined, and that $F \in C(\mathbb{T}^d)$ by the dominated convergence theorem.

Similarly, for $\alpha \in (0, 1]$, define

$$F_{\varepsilon}(x) = (F_{\varepsilon}^{(1)}(x), \cdots, F_{\varepsilon}^{(d)}), \quad F_{\varepsilon}^{(i)}(x) := \text{p.v.} \int_{\{|z| \le 1/\varepsilon\}} z_i \frac{K(x, x+z)}{|z|^{d+\alpha}} dz, \quad i = 1, \cdots, d.$$

We can verify in a similar way as above that, for all $\alpha \in (0,1]$, $\int_{\mathbb{T}^d} F_{\varepsilon}^{(i)}(x) dx = 0$ and $F_{\varepsilon} \in C(\mathbb{T}^d)$ so that

$$||F_{\varepsilon}||_{\infty} \leq c_1(1+\varepsilon^{\alpha-1}\mathbb{1}_{\{\alpha\in(0,1)\}}+|\log\varepsilon|\mathbb{1}_{\{\alpha=1\}}).$$

By Lemma 2.1, there exist $\psi \in \mathscr{D}_{\mathbb{T}^d}(\mathcal{L}), \ \phi = (\phi^{(1)}, \dots, \phi^{(d)}) \in C(\mathbb{T}^d; \mathbb{R}^d)$ with $\phi^{(i)} \in \mathscr{D}_{\mathbb{T}^d}(\mathcal{L})$, and $\phi_{\varepsilon} = (\phi_{\varepsilon}^{(1)}, \dots, \phi_{\varepsilon}^{(d)})$ with $\phi_{\varepsilon}^{(i)} \in \mathscr{D}_{\mathbb{T}^d}(\mathcal{L})$ so that

$$\mathcal{L}\psi(x) = -(\bar{K}(x) - \bar{K}) \quad \text{if } \alpha \in (0, 2),$$

$$\mathcal{L}\phi(x) = -F(x) \quad \text{if } \alpha \in (1, 2),$$

$$\mathcal{L}\phi_{\varepsilon}(x) = -F_{\varepsilon}(x) \quad \text{if } \alpha \in (0, 1],$$

(3.1)

which satisfy that

$$\begin{aligned} \|\psi\|_{\infty} + \|\phi\|_{\infty} + \|\nabla\psi\|_{\infty} + \|\nabla\phi\|_{\infty} &\leq c_2 \quad \text{if } \alpha \in (1,2), \\ \|\psi\|_{\infty} &\leq c_2 \quad \text{if } \alpha \in (0,1] \end{aligned}$$

and

$$\|\phi_{\varepsilon}\|_{\infty} \leq c_2 \left(1 + \varepsilon^{-1+\alpha} \mathbb{1}_{\{\alpha \in (0,1)\}} + |\log \varepsilon| \mathbb{1}_{\{\alpha=1\}}\right).$$

$$(3.2)$$

Moreover, when $\alpha = 1$, there is a constant $\theta \in (0,1)$ so that for all $x, y \in \mathbb{R}^d$,

$$|\psi(x) - \psi(y)| \leqslant c_3 |x - y|^{\theta},$$

$$|\phi_{\varepsilon}(x) - \phi_{\varepsilon}(y)| \leqslant c_3 (1 + |\log \varepsilon|) |x - y|^{\theta}.$$
(3.3)

Proof of Theorem 1.1. Throughout the proof, without any further mention, all the constants c_i are independent of ε . Recall that, by assumption, $\bar{u} \in C^4_{(-\alpha/2)}(D)$. We divide the proof into three cases.

Case 1: $\alpha \in (1, 2)$. Define

$$v_{\varepsilon}(x) := \bar{u}_{\varepsilon}(x) + \varepsilon \left\langle \phi\left(x/\varepsilon\right), \nabla \bar{u}_{\varepsilon}(x) \right\rangle + \varepsilon^{\alpha} \bar{K}^{-1} \psi\left(x/\varepsilon\right) \eta_{\varepsilon}(x) \bar{\mathcal{L}} \bar{u}_{\varepsilon}(x), \quad x \in D.$$

$$(3.4)$$

Here, $\bar{u}_{\varepsilon}(x) := \bar{u}(x)\eta_{\varepsilon}(x)$ with η_{ε} being the cut-off function satisfying (2.30)–(2.31), and ϕ and ψ are those functions given in (3.1).

By the definition (1.6), (2.39) and (2.40) for $\mathcal{L}_{\varepsilon}, \hat{\mathcal{L}}_{1,\varepsilon}$ and $\bar{\mathcal{L}}_{\varepsilon}$ respectively,

$$\mathcal{L}_{\varepsilon}\bar{u}_{\varepsilon}(x) = \hat{\mathcal{L}}_{1,\varepsilon}\bar{u}_{\varepsilon}(x) + \varepsilon^{1-\alpha} \left\langle F\left(x/\varepsilon\right), \nabla\bar{u}_{\varepsilon}(x) \right\rangle = \bar{\mathcal{L}}_{\varepsilon}\bar{u}_{\varepsilon}(x) + \varepsilon^{1-\alpha} \left\langle F\left(x/\varepsilon\right), \nabla\bar{u}_{\varepsilon}(x) \right\rangle + I_{1}^{\varepsilon}(x),$$

where by (2.41), I_1^{ε} satisfies that

$$|I_1^{\varepsilon}(x)| \leq c_1 \left(\varepsilon^{2-\alpha} \delta_D(x)^{-2+\alpha/2} + \varepsilon^{\alpha/2} \delta_D(x)^{-\alpha} \right) \mathbb{1}_{D_{4\varepsilon}}(x) + c_1 \varepsilon^{-\alpha/2} \mathbb{1}_{D \setminus D_{4\varepsilon}}(x)$$

In particular,

$$\int_D |I_1^{\varepsilon}(x)| \, dx \leqslant c_2 \varepsilon^{1-\alpha/2}.$$

It holds that

$$\begin{split} \mathcal{L}_{\varepsilon} \left(\varepsilon \left\langle \phi \left(\varepsilon^{-1} \cdot \right), \nabla \bar{u}_{\varepsilon}(\cdot) \right\rangle \right) (x) \\ &= \varepsilon \left\langle \mathcal{L}_{\varepsilon} \phi \left(\varepsilon^{-1} \cdot \right) (x), \nabla \bar{u}_{\varepsilon}(x) \right\rangle + \varepsilon \left\langle \phi \left(x/\varepsilon \right), \mathcal{L}_{\varepsilon} \left(\nabla \bar{u}_{\varepsilon} \right) (x) \right\rangle \\ &+ \varepsilon \int_{\mathbb{R}^{d}} \left\langle \phi \left((x+z)/\varepsilon \right) - \phi \left(x/\varepsilon \right), \nabla \bar{u}_{\varepsilon}(x+z) - \nabla \bar{u}_{\varepsilon}(x) \right\rangle \frac{K \left(x/\varepsilon, (x+z)/\varepsilon \right)}{|z|^{d+\alpha}} \, dz \\ &= -\varepsilon^{1-\alpha} \left\langle F \left(x/\varepsilon \right), \nabla \bar{u}_{\varepsilon}(x) \right\rangle + \varepsilon \left\langle \phi \left(x/\varepsilon \right), \mathcal{L}_{\varepsilon} \left(\nabla \bar{u}_{\varepsilon} \right) (x) \right\rangle \\ &+ \varepsilon \int_{\mathbb{R}^{d}} \left\langle \phi \left((x+z)/\varepsilon \right) - \phi \left(x/\varepsilon \right), \nabla \bar{u}_{\varepsilon}(x+z) - \nabla \bar{u}_{\varepsilon}(x) \right\rangle \frac{K \left(x/\varepsilon, (x+z)/\varepsilon \right)}{|z|^{d+\alpha}} \, dz \\ &=: -\varepsilon^{1-\alpha} \left\langle F \left(x/\varepsilon \right), \nabla \bar{u}_{\varepsilon}(x) \right\rangle + I_{2}^{\varepsilon}(x) + I_{3}^{\varepsilon}(x), \end{split}$$

where in the second equality we used the equation (3.1).

Since $\bar{u} \in C^3_{(-\alpha/2)}(D)$, by (2.42) and (2.38),

$$|I_{2}^{\varepsilon}(x)| = \left| \left\langle \varepsilon\phi\left(x/\varepsilon\right), \hat{\mathcal{L}}_{1,\varepsilon}(\nabla \bar{u}_{\varepsilon})(x) + \varepsilon^{1-\alpha}\nabla^{2}\bar{u}_{\varepsilon}(x) \cdot F\left(x/\varepsilon\right) \right\rangle \right|$$

$$\leq \varepsilon \|\phi\|_{\infty} \left(\left| \hat{\mathcal{L}}_{1,\varepsilon}\nabla \bar{u}_{\varepsilon}(x) \right| + \varepsilon^{1-\alpha} \|F\|_{\infty} |\nabla^{2}\bar{u}_{\varepsilon}(x)| \right)$$

$$\leq c_{3} \left(\varepsilon^{2-\alpha}\delta_{D}(x)^{-2+\alpha/2} + \varepsilon^{\alpha/2}\delta_{D}(x)^{-\alpha} \right) \mathbb{1}_{D_{4\varepsilon}}(x) + c_{3}\varepsilon^{-\alpha/2} \mathbb{1}_{D\setminus D_{4\varepsilon}}(x)$$

On the other hand, we write

$$I_{3}^{\varepsilon}(x) = \varepsilon^{1-\alpha} \int_{\{|z| \leq 2\}} \left\langle \phi\left(x/\varepsilon + z\right) - \phi\left(x/\varepsilon\right), \nabla \bar{u}_{\varepsilon}(x+\varepsilon z) - \nabla \bar{u}_{\varepsilon}(x) \right\rangle \frac{K\left(x/\varepsilon, x/\varepsilon + z\right)}{|z|^{d+\alpha}} dz + \varepsilon^{1-\alpha} \int_{\{|z| > 2\}} \left\langle \phi\left(x/\varepsilon + z\right) - \phi\left(x/\varepsilon\right), \nabla \bar{u}_{\varepsilon}(x+\varepsilon z) - \nabla \bar{u}_{\varepsilon}(x) \right\rangle \frac{K\left(x/\varepsilon, x/\varepsilon + z\right)}{|z|^{d+\alpha}} dz$$

$$=: I_{31}^{\varepsilon}(x) + I_{32}^{\varepsilon}(x).$$

By the mean-value theorem and (2.38),

$$|I_{31}^{\varepsilon}(x)| \leq c_{4}\varepsilon^{2-\alpha} \sup_{y \in \mathbb{R}^{d}: |y-x| \leq 2\varepsilon} |\nabla^{2}\bar{u}_{\varepsilon}(y)| \cdot \|\nabla\phi\|_{\infty} \cdot \int_{\{|z| \leq 2\}} \frac{|z|^{2}}{|z|^{d+\alpha}} dz$$
$$\leq c_{5}\varepsilon^{2-\alpha}\delta_{D}(x)^{-2+\alpha/2} \mathbb{1}_{D_{4\varepsilon}}(x) + c_{5}\varepsilon^{-\alpha/2} \mathbb{1}_{D \setminus D_{4\varepsilon}}(x).$$

Applying the mean-value theorem again as well as a change of variable, we have

$$\begin{split} |I_{32}^{\varepsilon}(x)| &\leq c_{6}\varepsilon \|\phi\|_{\infty} \cdot \int_{\{|z|>2\varepsilon\}} |\nabla \bar{u}_{\varepsilon}(x+z) - \nabla \bar{u}_{\varepsilon}(x)| \frac{1}{|z|^{d+\alpha}} dz \\ &\leq c_{7}\varepsilon \int_{0}^{1} \int_{\{|z|>2\varepsilon\}} |\nabla^{2}\bar{u}_{\varepsilon}(x+sz)| \frac{1}{|z|^{d+\alpha-1}} dz ds \\ &\leq c_{8}\varepsilon \int_{0}^{1} \int_{\{|z|>2\varepsilon\}} \left(\delta_{D}(x+sz)^{-2+\alpha/2} \mathbb{1}_{\{x+sz\in D_{3}\varepsilon\}} + \varepsilon^{-2+\alpha/2} \mathbb{1}_{\{x+sz\in D\setminus D_{3}\varepsilon\}} \right) \frac{1}{|z|^{d+\alpha-1}} dz ds \\ &\leq c_{9}\varepsilon \int_{0}^{1} s^{1-\alpha} \int_{\{|z|>2s\varepsilon\}} \left(\delta_{D}(x+z)^{-2+\alpha/2} \mathbb{1}_{\{x+z\in D_{3}\varepsilon\}} + \varepsilon^{-2+\alpha/2} \mathbb{1}_{\{x+z\in D\setminus D_{3}\varepsilon\}} \right) \frac{1}{|z|^{d+\alpha-1}} dz ds \\ &\leq c_{10}\varepsilon^{2-\alpha} \delta_{D}(x)^{-2+\alpha/2} \mathbb{1}_{D_{4}\varepsilon}(x) + c_{10}\varepsilon^{\alpha/2} \delta_{D}(x)^{-\alpha} \mathbb{1}_{D_{4}\varepsilon}(x) + c_{10}\varepsilon^{-\alpha/2} \mathbb{1}_{D\setminus D_{4}\varepsilon}(x), \end{split}$$

where in the last inequality we used (2.16) and (2.17). Thus, by all the estimates above for $I_2^{\varepsilon}(x)$, $I_{31}^{\varepsilon}(x)$ and $I_{32}^{\varepsilon}(x)$, we get

$$\int_{D} \left(|I_2^{\varepsilon}(x)| + |I_3^{\varepsilon}(x)| \right) \, dx \leqslant \int_{D} \left(|I_2^{\varepsilon}(x)| + |I_{31}^{\varepsilon}(x)| + |I_{32}^{\varepsilon}(x)| \right) \, dx \leqslant c_{11} \varepsilon^{1-\alpha/2}$$

Moreover, we set

$$\begin{split} \mathcal{L}_{\varepsilon} \left(\varepsilon^{\alpha} \psi \left(\varepsilon^{-1} \cdot \right) \bar{K}^{-1} \eta_{\varepsilon}(\cdot) \bar{\mathcal{L}} \bar{u}_{\varepsilon}(\cdot) \right) (x) \\ &= \varepsilon^{\alpha} \mathcal{L}_{\varepsilon} \psi \left(\varepsilon^{-1} \cdot \right) (x) \cdot \bar{K}^{-1} \eta_{\varepsilon}(x) \bar{\mathcal{L}} \bar{u}_{\varepsilon}(x) + \varepsilon^{\alpha} \bar{K}^{-1} \psi \left(x/\varepsilon \right) \mathcal{L}_{\varepsilon} (\eta_{\varepsilon} \bar{\mathcal{L}} \bar{u}_{\varepsilon}) (x) \\ &+ \varepsilon^{\alpha} \bar{K}^{-1} \int_{\mathbb{R}^{d}} \left(\psi \left((x+z)/\varepsilon \right) - \psi \left(x/\varepsilon \right) \right) \left(\eta_{\varepsilon} (x+z) \bar{\mathcal{L}} \bar{u}_{\varepsilon} (x+z) - \eta_{\varepsilon} (x) \bar{\mathcal{L}} \bar{u}_{\varepsilon} (x) \right) \frac{K \left(\frac{x}{\varepsilon}, \frac{x+z}{\varepsilon} \right)}{|z|^{d+\alpha}} \, dz \\ &=: \eta_{\varepsilon} (x) \left(\bar{\mathcal{L}} \bar{u}_{\varepsilon} (x) - \bar{\mathcal{L}}_{\varepsilon} \bar{u}_{\varepsilon} (x) \right) + I_{4}^{\varepsilon} (x) + I_{5}^{\varepsilon} (x), \end{split}$$

where in the last equality we have used the fact

$$\varepsilon^{\alpha} \mathcal{L}_{\varepsilon} \psi\left(\varepsilon^{-1} \cdot\right)(x) \cdot \bar{K}^{-1} \bar{\mathcal{L}} \bar{u}_{\varepsilon}(x) = \left(-\bar{K}\left(x/\varepsilon\right) + \bar{K}\right) \bar{K}^{-1} \bar{\mathcal{L}} u_{\varepsilon}(x) = \bar{\mathcal{L}} \bar{u}_{\varepsilon}(x) - \bar{\mathcal{L}}_{\varepsilon} \bar{u}_{\varepsilon}(x) \tag{3.5}$$

that can be verified directly by (3.1). We obtain by (2.48) and (1.10) that

$$\bar{\mathcal{L}}\bar{u}_{\varepsilon}(x) = h(x)\eta_{\varepsilon}(x) + I_{6}^{\varepsilon}(x), \qquad (3.6)$$

where

$$|I_6^{\varepsilon}(x)| \leq c_{12} \left(\varepsilon \delta_D(x)^{-1-\alpha/2} \mathbb{1}_{D_{4\varepsilon}}(x) + \varepsilon^{-\alpha/2} \mathbb{1}_{D \setminus D_{4\varepsilon}}(x) \right)$$

This yields that

$$\bar{\mathcal{L}}_{\varepsilon}\bar{u}_{\varepsilon}(x)| = \left|\bar{K}\left(x/\varepsilon\right) \cdot \bar{K}^{-1}\bar{\mathcal{L}}\bar{u}_{\varepsilon}(x)\right| \leqslant c_{13}\varepsilon^{-\alpha/2}, \quad x \in D \setminus D_{4\varepsilon}.$$

$$(3.7)$$

Using (2.48) and (2.49) (by taking $f = \bar{u}$ to get regularity estimates for $\bar{\mathcal{L}}\bar{u}_{\varepsilon}$), based on these regularity estimates for $\bar{\mathcal{L}}\bar{u}_{\varepsilon}$ and following the arguments for $I_2(x)$, $I_{31}^{\varepsilon}(x)$ and $I_{32}^{\varepsilon}(x)$, we can show that

$$|I_4^{\varepsilon}(x)| + |I_5^{\varepsilon}(x)| \leq c_{14} \left(\varepsilon^{2-\alpha} \delta_D(x)^{-2+\alpha/2} + \varepsilon^{\alpha/2} \delta_D(x)^{-\alpha} \right) \mathbb{1}_{D_{4\varepsilon}}(x) + c_{14} \varepsilon^{-\alpha/2} \mathbb{1}_{D \setminus D_{4\varepsilon}}(x),$$

and so

$$\int_D \left(|I_4^{\varepsilon}(x)| + |I_5^{\varepsilon}(x)| \right) \, dx \leqslant c_{15} \varepsilon^{1-\alpha/2}$$

Therefore, putting all the estimates for $I_i^{\varepsilon}(x)$, $i = 1, \dots, 5$ above together, yields that

$$\mathcal{L}_{\varepsilon} v_{\varepsilon}(x) = \eta_{\varepsilon}(x) \bar{\mathcal{L}} \bar{u}_{\varepsilon}(x) + (1 - \eta_{\varepsilon}(x)) \bar{\mathcal{L}}_{\varepsilon} \bar{u}_{\varepsilon}(x) + N_{1}^{\varepsilon}(x), \qquad (3.8)$$

where $N_1^{\varepsilon}(x)$ satisfies

$$\int_D |N_1^{\varepsilon}(x)| \, dx \leqslant c_{16} \varepsilon^{1-\alpha/2}.$$

This, along with (1.9) and (3.6), in turn gives us that

$$\mathcal{L}_{\varepsilon}\left(u_{\varepsilon}-v_{\varepsilon}\right)(x) = h(x)(1-\eta_{\varepsilon}(x)^{2}) - (1-\eta_{\varepsilon}(x))\bar{\mathcal{L}}_{\varepsilon}\bar{u}_{\varepsilon}(x) + N_{2}^{\varepsilon}(x), \quad x \in D,$$

where

$$\int_{D} \left(|h(x)(1 - \eta_{\varepsilon}(x)^{2})| + |(1 - \eta_{\varepsilon}(x))\bar{\mathcal{L}}_{\varepsilon}\bar{u}_{\varepsilon}(x)| + |N_{2}^{\varepsilon}(x)| \right) dx$$

$$\leq \int_{D \setminus D_{2\varepsilon}} \left(|h(x)| + |\bar{\mathcal{L}}_{\varepsilon}\bar{u}_{\varepsilon}(x)| \right) dx + \int_{D} |N_{2}^{\varepsilon}(x)| dx \leq c_{17}\varepsilon^{1 - \alpha/2}.$$
(3.9)

Here we have also used (3.7) and (2.41) in the last inequality.

Note that $\langle \phi(\varepsilon^{-1}\cdot), \nabla \bar{u}_{\varepsilon}(\cdot) \rangle \in \text{Dom}(\mathcal{L}_{\varepsilon}^{D})$ and $\psi(\varepsilon^{-1}\cdot) \eta_{\varepsilon}(\cdot)\mathcal{L}\bar{u}_{\varepsilon}(\cdot) \in \text{Dom}(\mathcal{L}_{\varepsilon}^{D})$, so $u_{\varepsilon} - v_{\varepsilon} \in \text{Dom}(\mathcal{L}_{\varepsilon}^{D})$. Then, according to (2.11) and (3.9), we arrive at

$$\int_{D} |u_{\varepsilon}(x) - v_{\varepsilon}(x)| \, dx \leqslant c_{19} \int_{D} \left(|h(x)(1 - \eta_{\varepsilon}(x)^2)| + |(1 - \eta_{\varepsilon}(x))\bar{\mathcal{L}}_{\varepsilon}\bar{u}_{\varepsilon}(x)| + |N_2^{\varepsilon}(x)| \right) \, dx \leqslant c_{20}\varepsilon^{1 - \alpha/2}.$$

On the other hand, we have by (3.4), (3.6) and (2.38) that

$$\int_{D} |v_{\varepsilon}(x) - \bar{u}(x)| \, dx \leqslant \int_{D} |\bar{u}(x)| |1 - \eta_{\varepsilon}(x)| \, dx + \varepsilon \|\phi\|_{\infty} \int_{D} |\nabla \bar{u}_{\varepsilon}(x)| \, dx + \varepsilon^{\alpha} \|\psi\|_{\infty} \bar{K}^{-1} \int_{D} |\bar{\mathcal{L}}\bar{u}_{\varepsilon}(x)| \, dx \\ \leqslant c_{21}\varepsilon,$$

where in the last inequality we used the fact that D is bounded. Therefore,

$$\int_{D} |u_{\varepsilon}(x) - \bar{u}(x)| \, dx \leq \int_{D} |u_{\varepsilon}(x) - v_{\varepsilon}(x)| \, dx + \int_{D} |v_{\varepsilon}(x) - \bar{u}(x)| \, dx \leq c_{22} \varepsilon^{1 - \alpha/2} + c_{12} \varepsilon^$$

The proof for $\alpha \in (1,2)$ is complete.

Case 2: $\alpha = 1$. In this case, we set

$$v_{\varepsilon}(x) := \bar{u}_{\varepsilon}(x) + \varepsilon \left\langle \phi_{\varepsilon}\left(x/\varepsilon\right), \nabla \bar{u}_{\varepsilon}(x) \right\rangle + \varepsilon \bar{K}^{-1} \psi\left(x/\varepsilon\right) \eta_{\varepsilon}(x) \bar{\mathcal{L}} \bar{u}_{\varepsilon}(x), \quad x \in D,$$

where ϕ_{ε} and ψ are those functions given in (3.1).

According to (2.44), we obtain

$$\mathcal{L}_{\varepsilon}\bar{u}_{\varepsilon}(x) = \hat{\mathcal{L}}_{2,\varepsilon}\bar{u}_{\varepsilon}(x) + \langle F_{\varepsilon}(x/\varepsilon), \nabla\bar{u}_{\varepsilon}(x) \rangle = \bar{\mathcal{L}}_{\varepsilon}\bar{u}_{\varepsilon}(x) + \langle F_{\varepsilon}(x/\varepsilon), \nabla\bar{u}_{\varepsilon}(x) \rangle + J_{1}^{\varepsilon}(x),$$

where $\hat{\mathcal{L}}_{2,\varepsilon}$ is defined by (2.39), and $J_1^{\varepsilon}(x)$ satisfies

$$|J_1^{\varepsilon}(x)| \leq c_1 \left(\varepsilon^{1/2} \delta_D(x)^{-1} + \varepsilon \delta_D(x)^{-3/2} (1+|\log\varepsilon|)\right) \mathbb{1}_{D_{4\varepsilon}}(x) + c_1 \varepsilon^{-1/2} (1+|\log\varepsilon|) \mathbb{1}_{D\setminus D_{4\varepsilon}}(x),$$

which in particular implies that

$$\int_D |J_1^{\varepsilon}(x)| \, dx \leqslant c_2 \varepsilon^{1/2} (1 + |\log \varepsilon|).$$

As in Case 1, by (3.1),

$$\begin{split} \mathcal{L}_{\varepsilon} \left(\varepsilon \left\langle \phi_{\varepsilon} \left(\varepsilon^{-1} \cdot \right), \nabla \bar{u}_{\varepsilon}(\cdot) \right\rangle \right) (x) \\ &= - \left\langle F_{\varepsilon} \left(x/\varepsilon \right), \nabla \bar{u}_{\varepsilon}(x) \right\rangle + \varepsilon \left\langle \phi_{\varepsilon} \left(x/\varepsilon \right), \mathcal{L}_{\varepsilon} \left(\nabla \bar{u}_{\varepsilon} \right) (x) \right\rangle \\ &+ \varepsilon \int_{\mathbb{R}^{d}} \left\langle \phi_{\varepsilon} \left((x+z)/\varepsilon \right) - \phi_{\varepsilon} \left(x/\varepsilon \right), \nabla \bar{u}_{\varepsilon}(x+z) - \nabla \bar{u}_{\varepsilon}(x) \right\rangle \frac{K \left(x/\varepsilon, (x+z)/\varepsilon \right)}{|z|^{d+1}} \, dz \\ &=: - \left\langle F_{\varepsilon} \left(x/\varepsilon \right), \nabla \bar{u}_{\varepsilon}(x) \right\rangle + J_{2}^{\varepsilon}(x) + J_{3}^{\varepsilon}(x). \end{split}$$

Using (2.38), (2.45) and (3.2), we obtain

$$|J_{2}^{\varepsilon}(x)| \leq c_{3}\varepsilon \|\phi_{\varepsilon}\|_{\infty} \left(|\hat{\mathcal{L}}_{2,\varepsilon}(\nabla \bar{u}_{\varepsilon})(x)| + \|F_{\varepsilon}\|_{\infty} |\nabla^{2} \bar{u}_{\varepsilon}(x)| \right)$$

$$\leq c_{4}\varepsilon (1+|\log\varepsilon|) \left(\delta_{D}(x)^{-3/2} (1+|\log\varepsilon|) + \varepsilon^{-1/2} \delta_{D}(x)^{-1} \right) \mathbb{1}_{D_{4\varepsilon}}(x)$$

$$+ c_4 \varepsilon^{-1/2} (1 + |\log \varepsilon|^2) \mathbb{1}_{D \setminus D_{4\varepsilon}}(x).$$

 Set

$$\begin{split} J_{3}^{\varepsilon}(x) &= \int_{\{|z| \leqslant 2\}} \left\langle \phi_{\varepsilon} \left(x/\varepsilon + z \right) - \phi_{\varepsilon} \left(x/\varepsilon \right), \nabla \bar{u}_{\varepsilon}(x + \varepsilon z) - \nabla \bar{u}_{\varepsilon}(x) \right\rangle \frac{K \left(x/\varepsilon, x/\varepsilon + z \right)}{|z|^{d+1}} \, dz \\ &+ \int_{\{|z| > 2\}} \left\langle \phi_{\varepsilon} \left(x/\varepsilon + z \right) - \phi_{\varepsilon} \left(x/\varepsilon \right), \nabla \bar{u}_{\varepsilon}(x + \varepsilon z) - \nabla \bar{u}_{\varepsilon}(x) \right\rangle \frac{K \left(x/\varepsilon, x/\varepsilon + z \right)}{|z|^{d+1}} \, dz \\ &=: J_{31}^{\varepsilon}(x) + J_{32}^{\varepsilon}(x). \end{split}$$

By the mean-value theorem and (3.3), we have

$$\begin{aligned} |J_{31}^{\varepsilon}(x)| &\leq c_{5}\varepsilon(1+|\log\varepsilon|) \sup_{y\in\mathbb{R}^{d}:|y-x|\leq 2\varepsilon} |\nabla^{2}\bar{u}_{\varepsilon}(y)| \cdot \int_{\{|z|\leq 2\}} \frac{|z|^{1+\theta}}{|z|^{d+1}} dz \\ &\leq c_{6}\varepsilon(1+|\log\varepsilon|)\delta_{D}(x)^{-3/2} \mathbb{1}_{D_{4\varepsilon}}(x) + c_{6}\varepsilon^{-1/2}(1+|\log\varepsilon|)\mathbb{1}_{D\setminus D_{4\varepsilon}}(x) \end{aligned}$$

Applying the mean-value theorem again, (3.2) and a change of variable, it holds that $|J_{32}^{\varepsilon}(x)|$

$$\leq c_{7}\varepsilon \|\phi_{\varepsilon}\|_{\infty} \cdot \left(\int_{\{2\varepsilon < |z| \leq 1\}} |\nabla \bar{u}_{\varepsilon}(x+z) - \nabla \bar{u}_{\varepsilon}(x)| \frac{1}{|z|^{d+1}} dz + \int_{\{|z| > 1\}} (|\nabla \bar{u}_{\varepsilon}(x+z)| + |\nabla \bar{u}_{\varepsilon}(x)|) \frac{1}{|z|^{d+1}} dz \right)$$

$$\leq c_{8}\varepsilon (1 + |\log \varepsilon|) \left[\int_{0}^{1} \int_{\{2\varepsilon < |z| \leq 1\}} |\nabla^{2} \bar{u}_{\varepsilon}(x+sz)| \frac{1}{|z|^{d}} dz ds + \left(\delta_{D}(x)^{-1/2} \mathbb{1}_{D_{4\varepsilon}}(x) + \varepsilon^{-1/2} \mathbb{1}_{D \setminus D_{4\varepsilon}}(x) \right) \right.$$

$$+ \int_{\{|z| > 1\}} \left(\delta_{D}(x+z)^{-1/2} \mathbb{1}_{\{x+z \in D_{4\varepsilon}\}} + \varepsilon^{-1/2} \mathbb{1}_{\{x+z \in D \setminus D_{4\varepsilon}\}} \right) \frac{1}{|z|^{d+1}} dz \right]$$

$$\leq c_{9}\varepsilon (1 + |\log \varepsilon|) \left[\int_{0}^{1} \int_{\{2s\varepsilon < |z| \leq s\}} \left(\delta_{D}(x+z)^{-3/2} \mathbb{1}_{\{x+z \in D_{3\varepsilon}\}} + \varepsilon^{-3/2} \mathbb{1}_{\{x+z \in D \setminus D_{3\varepsilon}\}} \right) \frac{1}{|z|^{d}} dz ds$$

$$+ \left(\delta_{D}(x)^{-1/2} \mathbb{1}_{D_{4\varepsilon}}(x) + \varepsilon^{-1/2} \mathbb{1}_{D \setminus D_{4\varepsilon}}(x) \right)$$

$$+ \int_{\{|z| > 1\}} \left(\delta_{D}(x+z)^{-1/2} \mathbb{1}_{\{x+z \in D_{4\varepsilon}\}} + \varepsilon^{-1/2} \mathbb{1}_{\{x+z \in D \setminus D_{4\varepsilon}\}} \right) \frac{1}{|z|^{d+1}} dz \right].$$

Hence, by (2.18)–(2.20) and all the estimates above for $J_2^{\varepsilon}(x)$, $J_{31}^{\varepsilon}(x)$ and $J_{32}^{\varepsilon}(x)$, we get

$$\int_D \left(|J_2^{\varepsilon}(x)| + |J_3^{\varepsilon}(x)| \right) \, dx \leqslant \int_D \left(|J_2^{\varepsilon}(x)| + |J_{31}^{\varepsilon}(x)| + |J_{32}^{\varepsilon}(x)| \right) \, dx \leqslant c_{10} \varepsilon^{1/2} (1 + |\log \varepsilon|^2).$$

Next, we define

$$\begin{split} \mathcal{L}_{\varepsilon} \left(\varepsilon \psi \left(\varepsilon^{-1} \cdot \right) \bar{K}^{-1} \eta_{\varepsilon}(\cdot) \bar{\mathcal{L}} \bar{u}_{\varepsilon}(\cdot) \right) (x) \\ &= \varepsilon \mathcal{L}_{\varepsilon} \psi \left(\varepsilon^{-1} \cdot \right) (x) \cdot \bar{K}^{-1} \eta_{\varepsilon}(x) \bar{\mathcal{L}} \bar{u}_{\varepsilon}(x) + \varepsilon \bar{K}^{-1} \psi \left(x/\varepsilon \right) \mathcal{L}_{\varepsilon} (\eta_{\varepsilon} \bar{\mathcal{L}} \bar{u}_{\varepsilon}) (x) \\ &+ \varepsilon \bar{K}^{-1} \int_{\mathbb{R}^{d}} \left(\psi \left((x+z)/\varepsilon \right) - \psi \left(x/\varepsilon \right) \right) \left(\eta_{\varepsilon} (x+z) \bar{\mathcal{L}} \bar{u}_{\varepsilon} (x+z) - \eta_{\varepsilon} (x) \bar{\mathcal{L}} \bar{u}_{\varepsilon} (x) \right) \frac{K \left(x/\varepsilon, (x+z)/\varepsilon \right)}{|z|^{d+1}} \, dz \\ &=: \eta_{\varepsilon} (x) \left(\bar{\mathcal{L}} \bar{u}_{\varepsilon} (x) - \bar{\mathcal{L}}_{\varepsilon} \bar{u}_{\varepsilon} (x) \right) + J_{4}^{\varepsilon} (x) + J_{5}^{\varepsilon} (x). \end{split}$$

Here we have used the fact (3.5).

By (2.48) and (1.9), we know

$$\bar{\mathcal{L}}\bar{u}_{\varepsilon}(x) = h(x)\eta_{\varepsilon}(x) + J_{6}^{\varepsilon}(x).$$
(3.10)

Here

$$|J_6^{\varepsilon}(x)| \leq c_{11} \left(\varepsilon \delta_D(x)^{-3/2} \mathbb{1}_{D_{4\varepsilon}}(x) + \varepsilon^{-1/2} \mathbb{1}_{D \setminus D_{4\varepsilon}}(x) \right).$$

Using (2.48) and (2.50) (to get regularity estimates for $\bar{\mathcal{L}}\bar{u}_{\varepsilon}$), based on these regularity estimates for $\bar{\mathcal{L}}\bar{u}_{\varepsilon}$ and following the arguments for $J_2^{\varepsilon}(x)$, $J_{31}^{\varepsilon}(x)$ and $J_{32}^{\varepsilon}(x)$, we can show that

$$\int_D \left(|J_4^{\varepsilon}(x)| + |J_5^{\varepsilon}(x)| \right) \, dx \leqslant c_{12} \varepsilon^{1/2} (1 + |\log \varepsilon|^2)$$

Putting all the estimates for $J_i^{\varepsilon}(x)$, $i = 1, \dots, 5$, above together yields that

$$\mathcal{L}_{\varepsilon} v_{\varepsilon}(x) = \eta_{\varepsilon}(x) \bar{\mathcal{L}} \bar{u}_{\varepsilon}(x) + (1 - \eta_{\varepsilon}(x)) \bar{\mathcal{L}}_{\varepsilon} \bar{u}_{\varepsilon}(x) + N_{3}^{\varepsilon}(x), \qquad (3.11)$$

where

$$\int_D |N_3^{\varepsilon}(x)| \, dx \leqslant c_{13} \varepsilon^{1/2} (1+|\log \varepsilon|^2).$$

This, along with (1.9) and (3.10), gives us that

$$\mathcal{L}_{\varepsilon}\left(u_{\varepsilon}-v_{\varepsilon}\right)(x) = h(x)(1-\eta_{\varepsilon}(x)^{2}) - (1-\eta_{\varepsilon}(x))\bar{\mathcal{L}}_{\varepsilon}\bar{u}_{\varepsilon}(x) + N_{4}^{\varepsilon}(x), \quad x \in D,$$

where

$$\int_{D} \left(|h(x)(1 - \eta_{\varepsilon}(x)^2)| + |(1 - \eta_{\varepsilon}(x))\bar{\mathcal{L}}_{\varepsilon}\bar{u}_{\varepsilon}(x)| + |N_4^{\varepsilon}(x)| \right) \, dx \leqslant c_{14}\varepsilon^{1/2}(1 + |\log\varepsilon|^2). \tag{3.12}$$

Here we have also used (3.7) (which still holds for $\alpha \in (0, 1]$ by its proof). As explained in the proof of Case 1, we can apply (3.12) and (2.11) to obtain

$$\int_D |u_{\varepsilon}(x) - v_{\varepsilon}(x)| \, dx \leqslant c_{15} \varepsilon^{1/2} (1 + |\log \varepsilon|^2).$$

Combining (3.4) and (3.10) with (2.38) yields that

$$\int_{D} |v_{\varepsilon}(x) - \bar{u}(x)| \, dx$$

$$\leqslant \int_{D} |\bar{u}(x)| |1 - \eta_{\varepsilon}(x)| \, dx + \varepsilon \|\phi_{\varepsilon}\|_{\infty} \int_{D} |\nabla \bar{u}_{\varepsilon}(x)| \, dx + \varepsilon \|\psi\|_{\infty} \bar{K}^{-1} \int_{D} |\bar{\mathcal{L}}\bar{u}_{\varepsilon}(x)| \, dx$$

$$\leqslant c_{16} \varepsilon^{1/2} (1 + |\log \varepsilon|^{2}).$$

Therefore,

$$\int_{D} |u_{\varepsilon}(x) - \bar{u}(x)| \, dx \leqslant \int_{D} |u_{\varepsilon}(x) - v_{\varepsilon}(x)| \, dx + \int_{D} |v_{\varepsilon}(x) - \bar{u}(x)| \, dx \leqslant c_{17} \varepsilon^{1/2} (1 + |\log \varepsilon|^2).$$

Hence, the proof for the case $\alpha = 1$ is finished.

Case 3: $\alpha \in (0, 1)$. Define

$$v_{\varepsilon}(x) := \bar{u}_{\varepsilon}(x) + \varepsilon \left\langle \phi_{\varepsilon}\left(x/\varepsilon\right), \nabla \bar{u}_{\varepsilon}(x) \right\rangle + \varepsilon^{\alpha} \bar{K}^{-1} \psi\left(x/\varepsilon\right) \eta_{\varepsilon}(x) \bar{\mathcal{L}} \bar{u}_{\varepsilon}(x), \quad x \in D.$$

Here ϕ_{ε} and ψ are those functions in (3.1). Therefore, by (2.44), we obtain

$$\mathcal{L}_{\varepsilon}\bar{u}_{\varepsilon}(x) = \hat{\mathcal{L}}_{2,\varepsilon}\bar{u}_{\varepsilon}(x) + \varepsilon^{1-\alpha} \left\langle F_{\varepsilon}\left(x/\varepsilon\right), \nabla\bar{u}_{\varepsilon}(x) \right\rangle = \bar{\mathcal{L}}_{\varepsilon}\bar{u}_{\varepsilon}(x) + \varepsilon^{1-\alpha} \left\langle F_{\varepsilon}\left(x/\varepsilon\right), \nabla\bar{u}_{\varepsilon}(x) \right\rangle + L_{1}^{\varepsilon}(x),$$

where L_1^{ε} satisfies

$$|L_1^{\varepsilon}(x)| \leq c_1 \varepsilon^{\alpha/2} \delta_D(x)^{-\alpha} \mathbb{1}_{D_{4\varepsilon}}(x) + c_1 \varepsilon^{-1+\alpha/2} \mathbb{1}_{D \setminus D_{4\varepsilon}}(x)$$

that implies

$$\int_D |L_1^{\varepsilon}(x)| \, dx \leqslant c_2 \varepsilon^{\alpha/2}$$

As before, by (3.1), we have

$$\begin{split} \mathcal{L}_{\varepsilon} \left(\varepsilon \left\langle \phi_{\varepsilon} \left(\varepsilon^{-1} \cdot \right), \nabla \bar{u}_{\varepsilon}(\cdot) \right\rangle \right) (x) \\ &= -\varepsilon^{1-\alpha} \left\langle F_{\varepsilon} \left(x/\varepsilon \right), \nabla \bar{u}_{\varepsilon}(x) \right\rangle + \varepsilon \left\langle \phi_{\varepsilon} \left(x/\varepsilon \right), \mathcal{L}_{\varepsilon} \left(\nabla \bar{u}_{\varepsilon} \right) (x) \right\rangle \\ &+ \varepsilon \int_{\mathbb{R}^{d}} \left\langle \phi_{\varepsilon} \left((x+z)/\varepsilon \right) - \phi_{\varepsilon} \left(x/\varepsilon \right), \nabla \bar{u}_{\varepsilon}(x+z) - \nabla \bar{u}_{\varepsilon}(x) \right\rangle \frac{K \left(x/\varepsilon, (x+z)/\varepsilon \right)}{|z|^{d+\alpha}} \, dz \\ &=: -\varepsilon^{1-\alpha} \left\langle F_{\varepsilon} \left(x/\varepsilon \right), \nabla \bar{u}_{\varepsilon}(x) \right\rangle + L_{2}^{\varepsilon}(x) + L_{3}^{\varepsilon}(x). \end{split}$$

According to (2.38), (2.46) and (3.2), we obtain that for every $\beta \in (\alpha/2, \alpha)$,

$$|L_{2}^{\varepsilon}(x)| \leq c_{3}\varepsilon \|\phi_{\varepsilon}\|_{\infty} |\mathcal{L}_{\varepsilon}(\nabla \bar{u}_{\varepsilon})(x)| \leq c_{4}\varepsilon^{\alpha} \left(\varepsilon^{-(\beta-\alpha/2)}\delta_{D}(x)^{-1-(\alpha-\beta)} + \varepsilon^{-(\alpha-\beta)}\delta_{D}(x)^{-1-\beta+\alpha/2}\right) \mathbb{1}_{D_{4\varepsilon}}(x) + c_{4}\varepsilon^{-1+\alpha/2} \mathbb{1}_{D\setminus D_{4\varepsilon}}(x).$$

We set

$$\begin{split} L_{3}^{\varepsilon}(x) &= \varepsilon^{1-\alpha} \int_{\{|z| \leq 2\}} \left\langle \phi_{\varepsilon} \left(x/\varepsilon + z \right) - \phi \left(x/\varepsilon \right), \nabla \bar{u}_{\varepsilon}(x+\varepsilon z) - \nabla \bar{u}_{\varepsilon}(x) \right\rangle \frac{K \left(x/\varepsilon, x/\varepsilon + z \right)}{|z|^{d+\alpha}} \, dz \\ &+ \varepsilon^{1-\alpha} \int_{\{|z| > 2\}} \left\langle \phi_{\varepsilon} \left(x/\varepsilon + z \right) - \phi_{\varepsilon} \left(x/\varepsilon \right), \nabla \bar{u}_{\varepsilon}(x+\varepsilon z) - \nabla \bar{u}_{\varepsilon}(x) \right\rangle \frac{K \left(x/\varepsilon, x/\varepsilon + z \right)}{|z|^{d+\alpha}} \, dz \\ &=: L_{31}^{\varepsilon}(x) + L_{32}^{\varepsilon}(x). \end{split}$$

By the mean-value theorem and (3.2), we have

$$|L_{31}^{\varepsilon}(x)| \leq c_{5}\varepsilon^{2-\alpha} \|\phi_{\varepsilon}\|_{\infty} \cdot \sup_{y \in \mathbb{R}^{d}: |y-x| \leq 2\varepsilon} |\nabla^{2}\bar{u}_{\varepsilon}(y)| \cdot \int_{\{|z| \leq 2\}} \frac{|z|}{|z|^{d+\alpha}} dz$$
$$\leq c_{6}\varepsilon\delta_{D}(x)^{-2+\alpha/2} \mathbb{1}_{D_{4\varepsilon}}(x) + c_{6}\varepsilon^{-1+\alpha/2} \mathbb{1}_{D \setminus D_{4\varepsilon}}(x).$$

Applying (2.47) (with $f = \overline{u}$ in (2.47)) and (3.2), we get

$$\begin{split} |L_{32}^{\varepsilon}(x)| &\leq c_{7}\varepsilon \|\phi_{\varepsilon}\|_{\infty} \cdot \int_{\{|z|>2\varepsilon\}} |\nabla \bar{u}_{\varepsilon}(x+z) - \nabla \bar{u}_{\varepsilon}(x)| \frac{1}{|z|^{d+\alpha}} dz \\ &\leq c_{8}\varepsilon^{\alpha} \int_{\{|z|>2\varepsilon\}} \left(\delta_{D}(x+z)^{-1-\beta+\alpha/2} \mathbb{1}_{\{x+z\in D_{3}\varepsilon\}} + \varepsilon^{-1-\beta+\alpha/2} \mathbb{1}_{\{x+z\in D\setminus D_{3}\varepsilon\}}\right) \frac{1}{|z|^{d+\alpha-\beta}} dz \\ &+ c_{8}\varepsilon^{\alpha} \left(\delta_{D}(x)^{-1-\beta+\alpha/2} \mathbb{1}_{D_{4\varepsilon}}(x) + \varepsilon^{-1-\beta+\alpha/2} \mathbb{1}_{D\setminus D_{4\varepsilon}}(x)\right) \cdot \int_{\{|z|>2\varepsilon\}} \frac{1}{|z|^{d+\alpha-\beta}} dz \\ &\leq c_{9} \left(\varepsilon^{-(\beta-3\alpha/2)} \delta_{D}(x)^{-1-(\alpha-\beta)} + \varepsilon^{\beta} \delta_{D}(x)^{-1-\beta+\alpha/2}\right) \mathbb{1}_{D_{3\varepsilon}}(x) + c_{9}\varepsilon^{-1+\alpha/2} \mathbb{1}_{D\setminus D_{3\varepsilon}}(x), \end{split}$$

where the last inequality follows from (2.21) and (2.22). Hence, by all the estimates above for $L_2^{\varepsilon}(x)$, $L_{31}^{\varepsilon}(x)$ and $L_{32}^{\varepsilon}(x)$ and using the fact $\beta \in (\alpha/2, \alpha)$, we get

$$\int_D \left(|L_2^{\varepsilon}(x)| + |L_3^{\varepsilon}(x)| \right) \, dx \leqslant \int_D \left(|L_2^{\varepsilon}(x)| + |L_{31}^{\varepsilon}(x)| + |L_{32}^{\varepsilon}(x)| \right) \, dx \leqslant c_{10} \varepsilon^{\alpha/2}$$

Define

$$\begin{aligned} \mathcal{L}_{\varepsilon} \left(\varepsilon^{\alpha} \psi \left(\varepsilon^{-1} \cdot \right) \bar{K}^{-1} \eta_{\varepsilon}(\cdot) \bar{\mathcal{L}} \bar{u}_{\varepsilon}(\cdot) \right) (x) \\ &= \varepsilon^{\alpha} \mathcal{L}_{\varepsilon} \psi \left(\varepsilon^{-1} \cdot \right) (x) \cdot \bar{K}^{-1} \eta_{\varepsilon}(x) \bar{\mathcal{L}} \bar{u}_{\varepsilon}(x) + \varepsilon^{\alpha} \bar{K}^{-1} \psi \left(x/\varepsilon \right) \mathcal{L}_{\varepsilon} (\eta_{\varepsilon} \bar{\mathcal{L}} \bar{u}_{\varepsilon}) (x) \\ &+ \varepsilon^{\alpha} \bar{K}^{-1} \int_{\mathbb{R}^{d}} \left(\psi \left((x+z)/\varepsilon \right) - \psi \left(x/\varepsilon \right) \right) \left(\eta_{\varepsilon} (x+z) \bar{\mathcal{L}} \bar{u}_{\varepsilon} (x+z) - \eta_{\varepsilon} (x) \bar{\mathcal{L}} \bar{u}_{\varepsilon} (x) \right) \frac{1}{|z|^{d+\alpha}} dz \\ &=: \eta_{\varepsilon} (x) \left(\bar{\mathcal{L}} \bar{u}_{\varepsilon} (x) - \bar{\mathcal{L}}_{\varepsilon} \bar{u}_{\varepsilon} (x) \right) + L_{4}^{\varepsilon} (x) + L_{5}^{\varepsilon} (x). \end{aligned}$$

Here we have used the property (3.5) again. By (2.48) and (1.9), we know

$$\mathcal{L}\bar{u}_{\varepsilon}(x) = h(x)\eta_{\varepsilon}(x) + L_{6}^{\varepsilon}(x)$$
(3.13)

where

$$|L_6^{\varepsilon}(x)| \leq c_{11} \left(\varepsilon \delta_D(x)^{-1-\alpha/2} \mathbb{1}_{D_{4\varepsilon}}(x) + \varepsilon^{-\alpha/2} \mathbb{1}_{D \setminus D_{4\varepsilon}}(x) \right).$$

Furthermore, we have

$$\begin{aligned} |L_5^{\varepsilon}(x)| &\leq c_{12}\varepsilon^{\alpha} \|\psi\|_{\infty} \left(\int_{\{|z| \leq 2\varepsilon\}} + \int_{\{|z| > 2\varepsilon\}} \right) \left| \eta_{\varepsilon}(x+z)\bar{\mathcal{L}}\bar{u}_{\varepsilon}(x+z) - \eta_{\varepsilon}(x)\bar{\mathcal{L}}\bar{u}_{\varepsilon}(x) \right| \frac{1}{|z|^{d+\alpha}} dz \\ &=: L_{51}^{\varepsilon}(x) + L_{52}^{\varepsilon}(x). \end{aligned}$$

According to (2.50),

$$L_{51}^{\varepsilon}(x) \leqslant c_{13}\varepsilon^{\alpha} \sup_{y:|y-x|\leqslant 2\varepsilon} |\nabla(\eta_{\varepsilon}(y)\bar{\mathcal{L}}\bar{u}_{\varepsilon}(y))| \cdot \int_{\{|z|\leqslant 2\varepsilon\}} \frac{1}{|z|^{d+\alpha-1}} dz$$
$$\leqslant c_{14}\varepsilon \left(\delta_D(x)^{-1-\alpha/2} \mathbb{1}_{D_{4\varepsilon}}(x) + \varepsilon^{-1-\alpha/2} \mathbb{1}_{D\setminus D_{4\varepsilon}}(x)\right)$$
$$= c_{14}\varepsilon \delta_D(x)^{-1-\alpha} \mathbb{1}_{D_{4\varepsilon}}(x) + c_{14}\varepsilon^{-\alpha/2} \mathbb{1}_{D\setminus D_{4\varepsilon}}(x).$$

Using (3.13), we get

$$L_{52}^{\varepsilon}(x) \leq c_{15}\varepsilon^{\alpha} \int_{\{|z|>2\varepsilon\}} \left| h(x+z)\eta_{\varepsilon}^{2}(x+z) - h(x)\eta_{\varepsilon}^{2}(x) \right| \frac{1}{|z|^{d+\alpha}} dz$$
$$+ c_{15}\varepsilon^{\alpha} \int_{\{|z|>2\varepsilon\}} \left(|L_{6}^{\varepsilon}(x)| + |L_{6}^{\varepsilon}(x+z)| \right) \frac{1}{|z|^{d+\alpha}} dz$$
$$=: L_{521}^{\varepsilon}(x) + L_{522}^{\varepsilon}(x).$$

Noting that $h \in C^1_{(\alpha/2)}(D) \subset C^{\beta}_{(\alpha/2)}(D)$ for every $\beta \in (\alpha/2, \alpha)$, by the definition and the argument for (2.47), it holds that for all $x, z \in \mathbb{R}^d$ with $|z| > 2\varepsilon$,

$$\frac{|h(x+z)\eta_{\varepsilon}^{2}(x+z)-h(x)\eta_{\varepsilon}^{2}(x)|}{|z|^{\beta}} \leqslant c_{16} \Big(\delta_{D}(x+z)^{-\beta-\alpha/2} \mathbb{1}_{\{x+z\in D_{3\varepsilon}\}} + \delta_{D}(x)^{-\beta-\alpha/2} \mathbb{1}_{\{x\in D_{3\varepsilon}\}} + \varepsilon^{-\beta-\alpha/2} \Big(\mathbb{1}_{\{x+z\in D\setminus D_{3\varepsilon}\}} + \mathbb{1}_{\{x\in D\setminus D_{3\varepsilon}\}} \Big) \Big).$$

$$(3.14)$$

By (3.14), we deduce that

$$\begin{split} L_{521}^{\varepsilon}(x) \leqslant &c_{17}\varepsilon^{\alpha} \int_{\{|z|>2\varepsilon\}} \left(\delta_D(x+z)^{-\beta-\alpha/2} \mathbb{1}_{\{x+z\in D_{3\varepsilon}\}} + \varepsilon^{-\beta-\alpha/2} \mathbb{1}_{\{x+z\in D\setminus D_{3\varepsilon}\}} \right) \frac{1}{|z|^{d+\alpha-\beta}} dz \\ &+ c_{17}\varepsilon^{\alpha} \left(\delta_D(x)^{-\beta-\alpha/2} \mathbb{1}_{D_{3\varepsilon}}(x) + \varepsilon^{-\beta-\alpha/2} \mathbb{1}_{D\setminus D_{3\varepsilon}}(x) \right) \cdot \int_{\{|z|>2\varepsilon\}} \frac{1}{|z|^{d+\alpha-\beta}} dz \\ &\leqslant &c_{18} \left(\varepsilon^{\beta} \delta_D(x)^{-\beta-\alpha/2} + (\varepsilon^{\alpha} \vee \varepsilon^{1-\beta+\alpha/2}) \cdot \delta_D(x)^{-1-(\alpha-\beta)} \right) \mathbb{1}_{D_{4\varepsilon}}(x) + c_{19}\varepsilon^{-\alpha/2} \mathbb{1}_{D\setminus D_{4\varepsilon}}(x), \end{split}$$

where in the last inequality we have used (2.22) and the fact

$$\int_{\{|z|>2\varepsilon\}} \delta_D(x+z)^{-\beta-\alpha/2} \mathbb{1}_{\{x+z\in D_{3\varepsilon}\}} \frac{1}{|z|^{d+\alpha-\beta}} dz$$

$$\leqslant c_{19} \left(\left(\varepsilon^{1-\beta-\alpha/2} \vee 1\right) \cdot \delta_D(x)^{-1-(\alpha-\beta)} + \varepsilon^{-(\alpha-\beta)} \delta_D(x)^{-\beta-\alpha/2} \right) \mathbb{1}_{D_{4\varepsilon}}(x) + c_{19} \varepsilon^{-3\alpha/2} \mathbb{1}_{D \setminus D_{4\varepsilon}}(x)$$

that can be proved by following the proof of (2.21) line by line. According to (3.13), we obtain

$$\begin{split} L_{522}^{\varepsilon}(x) &\leqslant c_{20}\varepsilon^{1+\alpha} \int_{\{|z|>2\varepsilon\}} \left(\delta_D(x+z)^{-1-\alpha/2} \mathbb{1}_{\{x+z\in D_{3\varepsilon}\}} + \varepsilon^{-1-\alpha/2} \mathbb{1}_{\{x+z\in D\setminus D_{3\varepsilon}\}} \right) \frac{1}{|z|^{d+\alpha}} dz \\ &+ c_{20} \left(\varepsilon \delta_D(x)^{-1-\alpha/2} \mathbb{1}_{D_{4\varepsilon}}(x) + \varepsilon^{-\alpha/2} \mathbb{1}_{D\setminus D_{4\varepsilon}}(x) \right) \\ &\leqslant c_{21} \left(\varepsilon \delta_D(x)^{-1-\alpha/2} \mathbb{1}_{D_{4\varepsilon}}(x) + \varepsilon^{-\alpha/2} \mathbb{1}_{D\setminus D_{4\varepsilon}}(x) \right), \end{split}$$

where the last inequality we have used (2.19) and the fact that

$$\int_{\{|z|>2\varepsilon\}} \delta_D(x+z)^{-1-\alpha/2} \mathbb{1}_{\{x+z\in D_{3\varepsilon}\}} \frac{1}{|z|^{d+\alpha}} dz$$

$$\leqslant c_{22} \left(\varepsilon^{-\alpha} \delta_D(x)^{-1-\alpha/2} + \varepsilon^{-\alpha/2} \delta_D(x)^{-1-\alpha}\right) \mathbb{1}_{D_{4\varepsilon}}(x) + c_{22} \varepsilon^{-3\alpha/2} \mathbb{1}_{D\setminus D_{4\varepsilon}}(x)$$

that can be verified by the same way as that for (2.21).

Using the same procedure (of $I_4^{\varepsilon}(x)$ and $J_4^{\varepsilon}(x)$) to estimate $L_4^{\varepsilon}(x)$ and putting all the estimates above together, as well as taking into account that $\beta > \alpha/2$, yield that

$$\int_D \left(|L_4^{\varepsilon}(x)| + |L_5^{\varepsilon}(x)| \right) \, dx \leqslant c_{23} \varepsilon^{1-\alpha/2} \leqslant c_{23} \varepsilon^{\alpha/2}$$

Combining all the estimates for $L_i^{\varepsilon}(x)$, $i = 1, \dots, 5$, above, we have

$$\mathcal{L}_{\varepsilon} v_{\varepsilon}(x) = \eta_{\varepsilon}(x) \bar{\mathcal{L}} \bar{u}_{\varepsilon}(x) + (1 - \eta_{\varepsilon}(x)) \bar{\mathcal{L}}_{\varepsilon} \bar{u}_{\varepsilon}(x) + N_{5}^{\varepsilon}(x), \qquad (3.15)$$

where

$$\int_D |N_5^{\varepsilon}(x)| \, dx \leqslant c_{24} \varepsilon^{\alpha/2}.$$

According to this, (1.9) and (3.13) yield that

$$\mathcal{L}_{\varepsilon}\left(u_{\varepsilon}-v_{\varepsilon}\right)(x) = h(x)(1-\eta_{\varepsilon}(x)^{2}) - (1-\eta_{\varepsilon}(x))\bar{\mathcal{L}}_{\varepsilon}\bar{u}_{\varepsilon}(x) + N_{6}^{\varepsilon}(x), \quad x \in D,$$

where

$$\int_{D} \left(|h(x)(1 - \eta_{\varepsilon}(x)^2)| + |(1 - \eta_{\varepsilon}(x))\bar{\mathcal{L}}_{\varepsilon}\bar{u}_{\varepsilon}(x)| + |N_6^{\varepsilon}(x)| \right) dx \leqslant c_{25}\varepsilon^{\alpha/2}.$$
(3.16)

Here we have also used (3.7) (by the proof which still holds for $\alpha \in (0, 1]$).

As explained in the proof of Case 1, according to (3.16), we can apply (2.11) to obtain

$$\int_D |u_{\varepsilon}(x) - v_{\varepsilon}(x)| \, dx \leqslant c_{26} \varepsilon^{\alpha/2}.$$

On the other hand, combining (2.38) with (3.13) gives us

$$\begin{split} \int_{D} |v_{\varepsilon}(x) - \bar{u}(x)| \, dx &\leq \int_{D} |\bar{u}(x)| |1 - \eta_{\varepsilon}(x)| \, dx + \varepsilon \|\phi_{\varepsilon}\|_{\infty} \int_{D} |\nabla \bar{u}_{\varepsilon}(x)| \, dx + \varepsilon^{\alpha} \|\psi\|_{\infty} \bar{K}^{-1} \int_{D} |\bar{\mathcal{L}} \bar{u}_{\varepsilon}(x)| \, dx \\ &\leq c_{27} \varepsilon^{\alpha/2}, \end{split}$$

and so

$$\int_{D} |u_{\varepsilon}(x) - \bar{u}(x)| \, dx \leqslant \int_{D} |u_{\varepsilon}(x) - v_{\varepsilon}(x)| \, dx + \int_{D} |v_{\varepsilon}(x) - \bar{u}(x)| \, dx \leqslant c_{28} \varepsilon^{\alpha/2}.$$
have finished the proof for $\alpha \in (0, 1)$.

Therefore, we have finished the proof for $\alpha \in (0, 1)$.

Next, we will present the

Proof of Theorem 1.2. The proof is almost the same as (but simpler than) that of Theorem 1.1, so we only give some crucial different points here. Throughout the proof, all the constants c_i are independent of ε . First, due to the fact $\bar{u} \in C_c^2(D)$, there is a constant $c_1 > 0$ so that for all $x \in D$ and k = 0, 1, 2, d

$$|\nabla^k \bar{u}(x)| \leqslant c_1. \tag{3.17}$$

With this at hand (in particular we do not need to consider blow up behaviors of $|\nabla^k \bar{u}|$ near ∂D), it is easy to verify that there is a constant $c_2 > 0$ so that for all $x \in D$,

$$\begin{aligned} |\hat{\mathcal{L}}_{1,\varepsilon}\bar{u}(x)| + |\hat{\mathcal{L}}_{1,\varepsilon}\nabla\bar{u}(x)| &\leq c_2 \quad \text{if } \alpha \in (1,2), \\ |\hat{\mathcal{L}}_{2,\varepsilon}\bar{u}(x)| + |\hat{\mathcal{L}}_{2,\varepsilon}\nabla\bar{u}(x)| &\leq c_2 \quad \text{if } \alpha \in (0,1], \end{aligned}$$
(3.18)

where $\hat{\mathcal{L}}_{1,\varepsilon}$ and $\hat{\mathcal{L}}_{2,\varepsilon}$ are defined by (2.39).

Using (3.17) in place of (2.38), and following the proofs of Lemmas 2.8 and 2.9 (which are in fact simpler since we do not need to consider the blow up behaviours of $\nabla^k \bar{u}$ near the boundary), we can prove that there is a constant $c_3 > 0$ so that for all $x \in D$,

$$|\hat{\mathcal{L}}_{1,\varepsilon}\bar{u}(x) - \bar{\mathcal{L}}_{\varepsilon}\bar{u}(x)| \leqslant c_3 \varepsilon^{2-\alpha} \quad \text{if } \alpha \in (1,2)$$
(3.19)

and

$$|\hat{\mathcal{L}}_{2,\varepsilon}\bar{u}(x) - \bar{\mathcal{L}}_{\varepsilon}\bar{u}(x)| \leq \begin{cases} c_3\varepsilon(1+|\log\varepsilon|) & \text{if } \alpha = 1, \\ c_3\varepsilon^{\alpha} & \text{if } \alpha \in (0,1), \end{cases}$$
(3.20)

where $\bar{\mathcal{L}}_{\varepsilon}$ is given by (2.40). By assumptions, $\bar{\mathcal{L}}\bar{u}(x) = h(x)$ for every $x \in D$, and $\bar{u} \in C_c^2(D)$. Set $\bar{u}(x) = 0$ for every $x \in D^c$, and then extend h to \mathbb{R}^d by setting $h(x) := \bar{\mathcal{L}}\bar{u}(x)$ for all $x \in \mathbb{R}^d$. Then, it is not difficult to verify that $h \in C_b^1(\mathbb{R}^d)$, and there is a constant $c_4 > 0$ such that for all $x \in \mathbb{R}^d$,

$$|h(x)| + |\nabla h(x)| \leq c_4 (1 + |x|)^{-d-\alpha}.$$
(3.21)

Then, as in the proof of Theorem 1.1 (here we also use the fact that $\mathcal{L}\bar{u}(x) = h(x)$), define for any $x \in D$,

$$v_{\varepsilon}(x) := \begin{cases} \bar{u}(x) + \varepsilon \left\langle \phi\left(x/\varepsilon\right), \nabla \bar{u}(x) \right\rangle + \left(\varepsilon^{\alpha} \bar{K}^{-1} \psi\left(x/\varepsilon\right) h(x) - w_{\varepsilon}(x)\right) & \text{if } \alpha \in (1,2), \\ \bar{u}(x) + \varepsilon \left\langle \phi_{\varepsilon}\left(x/\varepsilon\right), \nabla \bar{u}(x) \right\rangle + \left(\varepsilon^{\alpha} \bar{K}^{-1} \psi\left(x/\varepsilon\right) h(x) - w_{\varepsilon}(x)\right) & \text{if } \alpha \in (0,1]. \end{cases}$$

Here, ϕ , ϕ_{ε} and ψ are those given (3.1), and $w_{\varepsilon} : \mathbb{R}^d \to \mathbb{R}$ is the unique solution to the following Dirichlet exterior condition

$$\begin{cases} \mathcal{L}_{\varepsilon} w_{\varepsilon}(x) = 0, & x \in D, \\ w_{\varepsilon}(x) = \varepsilon^{\alpha} \bar{K}^{-1} \psi(x/\varepsilon) h(x), & x \in D^{c}. \end{cases}$$
(3.22)

The solution to the Dirichlet problem above is given by

$$w_{\varepsilon}(x) = \varepsilon^{\alpha} \bar{K}^{-1} \mathbb{E}_{x} \left[\psi \left(\varepsilon^{-1} X_{\tau_{D}^{\varepsilon}}^{\varepsilon} \right) h \left(X_{\tau_{D}^{\varepsilon}}^{\varepsilon} \right) \right], \quad x \in D$$
(3.23)

where $X^{\varepsilon} := \{X_t^{\varepsilon}\}_{t\geq 0}$ is the Hunt process associated with $\mathcal{L}_{\varepsilon}$, and τ_D^{ε} is the first exit time from D for the process X^{ε} . Indeed, denote by $\tilde{\omega}_{\varepsilon}$ the function given by the right hand side of (3.23). By [14], $\tilde{\omega}_{\varepsilon}$ is locally in $W^{\alpha/2,2}(D)$ and $\mathcal{L}_{\varepsilon}$ -harmonic in D. Since D is Lipschitz and so it satisfies an exterior cone condition, it follows from the heat kernel estimates in [17] for the stable-like process X^{ε} that every boundary point is regular with respect to the process X^{ε} ; see, e.g., [20, Lemma 3.2] for details. Consequently, $\tilde{\omega}_{\varepsilon}$ is continuous up to ∂D and takes values $\varepsilon^{\alpha} \bar{K}^{-1} \psi(x/\varepsilon) h(x)$ on D^{c} . Thus, $\tilde{\omega}_{\varepsilon}$ solves (3.22). By the same argument as that for [20, Theorem 3.8], it can be shown that the weak solution to (3.22) is unique. Consequently $\tilde{\omega}_{\varepsilon} = \omega_{\varepsilon}$ and so (3.23) holds. It follows from (3.23) that there is a constant $c_5 > 0$ so that for all $\varepsilon \in (0, 1)$,

$$\sup_{x \in D} |w_{\varepsilon}(x)| \leqslant c_5 \varepsilon^{\alpha}. \tag{3.24}$$

Now, we split the proof into three cases.

Case 1: $\alpha \in (1,2)$. Using (3.17)–(3.19) and the fact $h \in C_b^1(\mathbb{R}^d)$, and repeating the proof of Theorem 1.1, we can prove that

$$\mathcal{L}_{\varepsilon}\left(\bar{u}(\cdot) + \varepsilon \left\langle \phi\left(\varepsilon^{-1}\cdot\right), \nabla \bar{u}(\cdot) \right\rangle \right)(x) = \bar{K}^{-1}\bar{K}\left(x/\varepsilon\right)h(x) + I_{1}^{\varepsilon}(x) + \varepsilon \left\langle \phi\left(x/\varepsilon\right), \mathcal{L}_{\varepsilon}\left(\nabla \bar{u}\right)(x) \right\rangle, \\ \mathcal{L}_{\varepsilon}\left(\varepsilon^{\alpha}\bar{K}^{-1}\psi\left(\varepsilon^{-1}\cdot\right)h(\cdot)\right)(x) = \left(1 - \bar{K}^{-1}\bar{K}\left(x/\varepsilon\right)\right)h(x) + I_{2}^{\varepsilon}(x) + \varepsilon^{\alpha}\bar{K}^{-1}\psi\left(x/\varepsilon\right)\mathcal{L}_{\varepsilon}h(x),$$

where

$$\int_D (|I_1^{\varepsilon}(x)|^2 + |I_2^{\varepsilon}(x)|^2) \, dx \leqslant c_6 \varepsilon^{4-2\alpha}.$$

Here we remark that the constant c_6 above only depends on $\|\nabla^k \bar{u}\|_{\infty}$ with $k = 0, 1, 2, \|h\|_{\infty}$ and $\|\nabla h\|_{\infty}$. This, along with the equation (3.1) and (3.22), yields that for every $g \in \text{Dom}(\mathcal{L}^D_{\varepsilon})$

$$\int_{D} \mathcal{L}_{\varepsilon} v_{\varepsilon}(x) g(x) \, dx = \int_{D} h(x) g(x) \, dx + \int_{D} (I_{1}^{\varepsilon} + I_{2}^{\varepsilon}(x)) g(x) \, dx + \varepsilon \int_{D} \langle \mathcal{L}_{\varepsilon} \left(\nabla \bar{u} \right) (x) \, dx, \phi \left(x/\varepsilon \right) g(x) \rangle \, dx + \varepsilon^{\alpha} \bar{K}^{-1} \int_{D} \mathcal{L}_{\varepsilon} h(x) \psi \left(x/\varepsilon \right) g(x) \, dx.$$

$$(3.25)$$

Then, by the symmetry of $\mathcal{L}_{\varepsilon}$, it holds that

$$\begin{split} & \left| \varepsilon \int_{D} \left\langle \mathcal{L}_{\varepsilon} \left(\nabla \bar{u} \right)(x), \phi\left(x/\varepsilon \right) g(x) \right\rangle dx \right| \\ &= \frac{\varepsilon}{2} \left| \int_{\mathbb{R}^{d}} \int_{\mathbb{R}^{d}} \left\langle \nabla \bar{u}(x+z) - \nabla \bar{u}(x), \phi\left((x+z)/\varepsilon \right) g(x+z) - \phi\left(x/\varepsilon \right) g(x) \right\rangle \frac{K\left(x/\varepsilon, (x+z)/\varepsilon \right)}{|z|^{d+\alpha}} dz \, dx \right| \\ &= \frac{\varepsilon^{1-\alpha}}{2} \left| \int_{\mathbb{R}^{d}} \int_{\mathbb{R}^{d}} \left\langle \nabla \bar{u}(x+\varepsilon z) - \nabla \bar{u}(x), \phi\left(x/\varepsilon + z \right) g(x+\varepsilon z) - \phi\left(x/\varepsilon \right) g(x) \right\rangle \frac{K\left(x/\varepsilon, x/\varepsilon + z \right)}{|z|^{d+\alpha}} dz \, dx \right| \end{split}$$

$$\leq c_7 \varepsilon^{2-\alpha} \left(\int_D \left(\int_{\mathbb{R}^d} \|\nabla^2 \bar{u}\|_{\infty} |z| \cdot \left(\|\nabla \phi\|_{\infty} |z| \mathbb{1}_{\{|z| \leq 1\}} + \|\phi\|_{\infty} \mathbb{1}_{\{|z| > 1\}} \right) \frac{1}{|z|^{d+\alpha}} dz \right) \cdot |g(x)| dx \right)$$

$$+ c_7 \varepsilon^{1-\alpha} \left(\int_{\mathbb{R}^d} \|\phi\|_{\infty} \cdot \left(\int_{\mathbb{R}^d} |\nabla \bar{u}(x+\varepsilon z) - \nabla \bar{u}(x)| \left|g(x+\varepsilon z) - g(x)\right| \frac{1}{|z|^{d+\alpha}} dz \right) dx \right)$$

$$\leq c_8 \varepsilon^{2-\alpha} \int_D |g(x)| dx + c_8 \varepsilon \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} |\nabla \bar{u}(x+z) - \nabla \bar{u}(x)| \left|g(x+z) - g(x)\right| \frac{1}{|z|^{d+\alpha}} dz dx$$

$$\leq \frac{\varepsilon^{\varepsilon}(g,g)}{8} + c_9 \left(\varepsilon^{4-2\alpha} + \varepsilon^2 \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} |\nabla \bar{u}(x+z) - \nabla \bar{u}(x)|^2 \frac{1}{|z|^{d+\alpha}} dz dx \right)$$

$$\leq \frac{\varepsilon^{\varepsilon}(g,g)}{8} + c_{10} \varepsilon^{4-2\alpha},$$

where in the third inequality we used (2.9), the Young inequality and

$$\mathcal{E}^{\varepsilon}(g,g) := \frac{1}{2} \iint_{\mathbb{R}^d \times \mathbb{R}^d} \left(g(x) - g(y) \right)^2 \frac{K\left(x/\varepsilon, y/\varepsilon \right)}{|x - y|^{d + \alpha}} \, dx \, dy.$$

Using (3.21) and following the arguments above, we can obtain

$$\left|\varepsilon^{\alpha} \int_{D} \mathcal{L}_{\varepsilon} h(x) \psi(x/\varepsilon) g(x) dx\right| \leq \frac{\varepsilon^{\varepsilon}(g,g)}{8} + c_{11} \varepsilon^{4-2\alpha}$$

Putting all the estimates above together and taking $g = u_{\varepsilon} - v_{\varepsilon}$, we have

$$\begin{aligned} \mathcal{E}^{\varepsilon}(u_{\varepsilon} - v_{\varepsilon}, u_{\varepsilon} - v_{\varepsilon}) &= -\int_{D} \mathcal{L}_{\varepsilon}(u_{\varepsilon} - v_{\varepsilon})(x) \cdot (u_{\varepsilon}(x) - v_{\varepsilon}(x)) \, dx \\ &\leqslant \frac{\mathcal{E}^{\varepsilon}(u_{\varepsilon} - v_{\varepsilon}, u_{\varepsilon} - v_{\varepsilon})}{4} + \int_{D} |u_{\varepsilon}(x) - v_{\varepsilon}(x)| (|I_{1}^{\varepsilon}(x)| + |I_{2}^{\varepsilon}(x)|) \, dx + c_{12}\varepsilon^{4-2\alpha} \\ &\leqslant \frac{\mathcal{E}^{\varepsilon}(u_{\varepsilon} - v_{\varepsilon}, u_{\varepsilon} - v_{\varepsilon})}{2} + c_{13}\varepsilon^{4-2\alpha}, \end{aligned}$$

where in the last inequality we also used (2.9) and the Young inequality.

This along with (2.9) again (also by noting that $v_{\varepsilon} \in W_0^{\alpha/2,2}(D)$ due to the definition of w_{ε} on D^c) gives us that

$$\int_{D} |u_{\varepsilon}(x) - v_{\varepsilon}(x)|^2 \, dx \leqslant c_{14} \mathcal{E}^{\varepsilon} (u_{\varepsilon} - v_{\varepsilon}, u_{\varepsilon} - v_{\varepsilon}) \leqslant c_{15} \varepsilon^{4-2\alpha}$$

On the other hand, according to (3.24) and the definition of v_{ε} , it holds that

$$\int_D |\bar{u}(x) - v_{\varepsilon}(x)|^2 dx \leqslant c_{16} \varepsilon^2.$$

Therefore, we get

$$\int_{D} |u_{\varepsilon}(x) - \bar{u}(x)|^2 \, dx \leq 2 \left(\int_{D} |u_{\varepsilon}(x) - v_{\varepsilon}(x)|^2 \, dx + \int_{D} |\bar{u}(x) - v_{\varepsilon}(x)|^2 \, dx \right) \leq c_{17} \varepsilon^{4-2\alpha}.$$

Case 2: $\alpha = 1$. Repeating the arguments for the case $\alpha \in (1,2)$ and using (3.3) to replace $\|\nabla \phi\|_{\infty}$, we can show that

$$\int_D |u_{\varepsilon}(x) - \bar{u}(x)|^2 \, dx \leqslant c_{18} \varepsilon^2 (1 + |\log \varepsilon|)^4.$$

The details are omitted here.

Case 3: $\alpha \in (0,1)$. Using (3.17)–(3.20) and applying the arguments for the case $\alpha \in (1,2)$ (here we also used the approach in the proof of Theorem 1.1), we can prove that for every $g \in \text{Dom}(\mathcal{L}_{\varepsilon}^{D})$,

$$\begin{split} \int_{D} \mathcal{L}_{\varepsilon} v_{\varepsilon}(x) g(x) \, dx &= \int_{D} h(x) g(x) \, dx + \int_{D} (J_{1}^{\varepsilon} + J_{2}^{\varepsilon}(x)) g(x) \, ds + \varepsilon \int_{D} \left\langle \mathcal{L}_{\varepsilon} \left(\nabla \bar{u} \right)(x) \, dx, \phi_{\varepsilon} \left(x/\varepsilon \right) g(x) \right\rangle \, dx \\ &+ \varepsilon^{\alpha} \bar{K}^{-1} \int_{D} \mathcal{L}_{\varepsilon} h(x) \psi \left(x/\varepsilon \right) g(x) \, dx, \end{split}$$

where

$$\int_D (|J_1^{\varepsilon}(x)|^2 + |J_2^{\varepsilon}(x)|^2) \, dx \leqslant c_{19} \varepsilon^{2\alpha}.$$

Then,

$$\begin{split} & \left| \varepsilon \int_{D} \left\langle \mathcal{L}_{\varepsilon} \left(\nabla \bar{u} \right)(x), \phi_{\varepsilon} \left(x/\varepsilon \right) g(x) \right\rangle \, dx \right| \\ &= \frac{\varepsilon}{2} \left| \int_{\mathbb{R}^{d}} \int_{\mathbb{R}^{d}} \left\langle \nabla \bar{u}(x+z) - \nabla \bar{u}(x), \phi_{\varepsilon} \left((x+z)/\varepsilon \right) g(x+z) - \phi_{\varepsilon} \left(x/\varepsilon \right) g(x) \right\rangle \frac{K \left(x/\varepsilon, (x+z)/\varepsilon \right)}{|z|^{d+\alpha}} \, dz \, dx \right| \\ & \leq c_{20} \varepsilon \left(\int_{D} \left(\int_{\mathbb{R}^{d}} \| \phi_{\varepsilon} \|_{\infty} \cdot \left(\| \nabla \bar{u} \|_{\infty} |z| \mathbbm{1}_{\{|z| \leq 1\}} + \| \bar{u} \|_{\infty} \mathbbm{1}_{\{|z| > 1\}} \right) \frac{1}{|z|^{d+\alpha}} \, dz \right) \cdot |g(x)| \, dx \right) \\ & + c_{20} \varepsilon \left(\int_{\mathbb{R}^{d}} \| \phi_{\varepsilon} \|_{\infty} \cdot \left(\int_{\mathbb{R}^{d}} |\nabla \bar{u}(x+z) - \nabla \bar{u}(x)| \, |g(x+z) - g(x)| \frac{1}{|z|^{d+\alpha}} \, dz \right) \, dx \right) \\ & \leq c_{21} \varepsilon^{\alpha} \int_{D} |g(x)| \, dx + c_{21} \varepsilon^{\alpha} \int_{\mathbb{R}^{d}} \int_{\mathbb{R}^{d}} |\nabla \bar{u}(x+z) - \nabla \bar{u}(x)| \, |g(x+z) - g(x)| \frac{1}{|z|^{d+\alpha}} \, dz \, dx \\ & \leq \frac{\varepsilon^{\varepsilon}(g,g)}{8} + c_{22} \left(\varepsilon^{2\alpha} + \varepsilon^{2\alpha} \int_{\mathbb{R}^{d}} \int_{\mathbb{R}^{d}} |\nabla \bar{u}(x+z) - \nabla \bar{u}(x)|^{2} \frac{1}{|z|^{d+\alpha}} \, dz \, dx \right) \\ & \leq \frac{\varepsilon^{\varepsilon}(g,g)}{8} + c_{23} \varepsilon^{2\alpha}. \end{split}$$

Similarly, we can show that

$$\left|\varepsilon^{\alpha} \int_{D} \mathcal{L}_{\varepsilon} h(x) \psi(x/\varepsilon) g(x) dx\right| \leq \frac{\mathcal{E}^{\varepsilon}(g,g)}{8} + c_{24} \varepsilon^{2\alpha}$$

Putting all the estimates above together and taking $g = u_{\varepsilon} - v_{\varepsilon}$, we have

$$\begin{split} \mathcal{E}^{\varepsilon}(u_{\varepsilon} - v_{\varepsilon}, u_{\varepsilon} - v_{\varepsilon}) &= -\int_{D} \mathcal{L}_{\varepsilon}(u_{\varepsilon} - v_{\varepsilon})(x) \cdot (u_{\varepsilon}(x) - v_{\varepsilon}(x)) \, dx \\ &\leqslant \frac{\mathcal{E}^{\varepsilon}(u_{\varepsilon} - v_{\varepsilon}, u_{\varepsilon} - v_{\varepsilon})}{4} + \int_{D} |u_{\varepsilon}(x) - v_{\varepsilon}(x)| (|J_{1}^{\varepsilon}(x)| + |J_{2}^{\varepsilon}(x)|) \, dx + c_{25}\varepsilon^{2\alpha} \\ &\leqslant \frac{\mathcal{E}^{\varepsilon}(u_{\varepsilon} - v_{\varepsilon}, u_{\varepsilon} - v_{\varepsilon})}{2} + c_{26}\varepsilon^{2\alpha}, \end{split}$$

where we used (2.9) and the Young inequality in the last inequality. This, along with (2.9) again, yields that

$$\int_D |u_{\varepsilon}(x) - v_{\varepsilon}(x)|^2 \, dx \leqslant c_{27} \mathcal{E}^{\varepsilon} (u_{\varepsilon} - v_{\varepsilon}, u_{\varepsilon} - v_{\varepsilon}) \leqslant c_{28} \varepsilon^{2\alpha}$$

Hence, it holds that

$$\int_{D} |u_{\varepsilon}(x) - \bar{u}(x)|^2 \, dx \leqslant 2 \left(\int_{D} |u_{\varepsilon}(x) - v_{\varepsilon}(x)|^2 \, dx + \int_{D} |\bar{u}(x) - v_{\varepsilon}(x)|^2 \, dx \right) \leqslant c_{29} \varepsilon^{2\alpha}.$$

Now we have finished the proof.

Acknowledgements. The research of Xin Chen is supported by the National Natural Science Foundation of China (No. 12122111). The research of Zhen-Qing Chen is partially supported by Simons Foundation grant 520542. The research of Takashi Kumagai is supported by JSPS KAKENHI Grant Number JP22H00099 and 23KK0050. The research of Jian Wang is supported by the National Key R&D Program of China (2022YFA1006003) and the National Natural Science Foundation of China (Nos. 12071076 and 12225104).

References

- A. Arapostathis, A. Biswas and L. Caffarelli: The Dirichlet problem for stable-like operators and related probabilistic representations, Comm. Partial Differential Equations, 41 (2016), 1472–1511.
- [2] S. Armstrong, T. Kuusi, and J.-C. Mourrat: The additive structure of elliptic homogenization, *Invent. Math.*, 208 (2017), 999–1154.
- [3] S. Armstrong, T. Kuusi, and J.-C. Mourrat: Quantitative Stochastic Homogenization and Large Scale Regularity, Grundlehren der mathematischen Wissenschaften 352, Springer, Cham, 2019.
- [4] N.S. Bakhvalov and G.P. Panasenko: Homogenisation: Averaging Processes in Periodic Media, Mathematics and Its Applications, vol. 36, Springer, Dordrecht, 1989.
- [5] I.I. Banis: Estimation of the rate of convergence in the metric L_p in the case of a limiting stable law, *Lith. Math.*, **13** (1974), 379–384.
- [6] R. F. Bass, M. Kassmann and T. Kumagai: Symmetric jump processes: localization, heat kernels, and convergence, Ann. Inst. H. Poincaré - Probabilités et Statistiques, 46 (2010), 59–71.
- [7] A. Bensoussan, J.-L. Lions and G.C. Papanicolaou: Asymptotic Analysis for Periodic Structures, Stud. Math. Appl., vol. 5, North-Holland Publishing Co., Amsterdam-New York, 1978.
- [8] M.Sh. Birman and T.A. Suslina: Periodic second-order differential operators. Threshold properties and averaging, St. Petersburg Math. J., 15 (2004), 639–714.
- M.Sh. Birman and T.A. Suslina: Operator error estimates in the homogenization problem for nonstationary periodic equations, St. Petersburg Math. J., 20 (2009), 873–928
- [10] P. Chen, I. Nourdin and L. Xu: Stein's method for asymmetric α-stable distributions, with application to the stable CLT, J. Theoret. Probab., 34 (2021), 1382–1407.
- [11] X. Chen, Z.-Q. Chen, T. Kumagai and J. Wang: Homogenization of symmetric stable-like processes in stationary ergodic media, SIAM J. Math. Anal., 53 (2021), 2957–3001.
- [12] X. Chen, Z.-Q. Chen, T. Kumagai and J. Wang: Periodic homogenization of non-symmetric Lévy-type processes, Ann. Probab., 49 (2021), 2874–2921.
- [13] X. Chen, Z.-Q. Chen, T. Kumagai and J. Wang: Quantitative stochastic homogenization for random conductance models with stable-like jumps, arXiv:2306.15855.
- [14] Z.-Q. Chen: On notions of harmonicity, Proc. Amer. Math. Soc., 137 (2009), 3497–3510.
- [15] Z.-Q. Chen and M. Fukushima. Symmetric Markov Processes, Time Changes, and Boundary Theory, Princeton Univ. Press, 2012.
- [16] Z.-Q. Chen, P. Kim and R. Song: Heat kernel estimates for the Dirichlet fractional Laplacian, J. Eur. Math. Soc., 12 (2010), 1307–1329.
- [17] Z.-Q. Chen and T. Kumagai: Heat kernel estimates for stable-like processes on d-sets, Stoch. Proc. Their Appl., 108 (2003), 27–62.
- [18] Z.-Q. Chen and T. Kumagai: Heat kernel estimates for jump processes of mixed types on metric measure spaces, Probab. Theory Relat. Fields, 140 (2008), 277–317.
- [19] Z.-Q. Chen, T. Kumagai and J. Wang: Stability of heat kernel estimates for symmetric non-local Dirichlet forms, Mem. Amer. Math. Soc., 271 (2021).
- [20] Z.-Q. Chen and J. Peng: Dirichlet boundary value problem for integral-differential operators, in Preparation.
- [21] Z.-Q. Chen and R. Song: Intrinsic ultracontractivity and conditional gauge for symmetric stable processes, J. Funct. Anal., 150 (1997), 204–239.
- [22] Z.-Q. Chen and X. Zhang: Heat kernels for time-dependent non-symmetric stable-like operators, J. Math. Anal. Appl., 465 (2018), 1–21.
- [23] D. Down, S.P. Meyn and R.T. Tweedie: Exponential and uniform ergodicity of Markov processes, Ann. Probab., 23 (1995), 1671–1691.
- [24] J. Geng and Z. Shen: Convergence rates in parabolic homogenization with time-dependent periodic coefficients, J. Funct. Anal., 272 (2017), 2092–2113.
- [25] A. Gloria, S. Neukamm, and F. Otto: A regularity theory for random elliptic operators, Milan J. Math., 88 (2020), 99–170.
- [26] A. Gloria and F. Otto: An optimal variance estimate in stochastic homogenization of discrete elliptic equations, Ann. Probab., 39 (2011), 779–856.
- [27] G. Griso: Error estimate and unfolding for periodic homogenization, Asymptot. Anal., 40 (2004), 269–286.
- [28] G. Griso: Interior error estimate for periodic homogenization Anal. Appl., 4 (2006), 61–79.
- [29] Q. Huang, J.-Q. Duan and R. Song: Homogenization of periodic linear nonlocal partial differential equations, *Bernoulli*, 28 (2022), 1648–1674.
- [30] Q. Huang, J.-Q. Duan and R. Song: Homogenization of non-symmetric jump processes, Adv. Appl. Probab., 56 (2024), 1–33.
- [31] V.V. Jikov, S.M. Kozlov and O.A. Oleinik: Homogenization of Differential Operators and Integral Functionals, Springer-Verlag, Berlin, 1994.
- [32] M. Kassmann, A. Piatnitski and E. Zhizhina: Homogenization of Lévy-type operators with oscillating coefficients, SIAM J. Math. Anal., 51 (2019), 3641–3665.
- [33] C. Kenig, F. Lin and Z. Shen: Convergence rates in L_2 for elliptic homogenization problems, Arch. Ration. Mech. Anal., 203 (2012), 1009–1036.

- [34] C. Kenig, F. Lin and Z. Shen: Homogenization of elliptic systems with Neumann boundary conditions, J. Amer. Math. Soc., 26 (2013), 901–937.
- [35] C. Kenig, F. Lin and Z. Shen: Estimates of eigenvalues and eigenfunctions in periodic homogenization, J. Eur. Math. Soc., 15 (2013), 1901–1925.
- [36] S. Kesavan: Homogenization of elliptic eigenvalue problems: part I, Appl. Math. Optim., 5 (1979), 153–167.
- [37] S. Kesavan: Homogenization of elliptic eigenvalue problems: part II, Appl. Math. Optim., 5 (1979), 197–216.
- [38] A. Piatnitski, V. Sloushch, T. Suslina and E. Zhizhina: On operator estimates in homogenization of nonlocal operators of convolution type, J. Differential Equations, 352 (2023), 153–188.
- [39] A. Piatnitski and E. Zhizhina: Periodic homogenization of nonlocal operators with a convolution-type kernel, SIAM J. Math. Anal., 49 (2017), 64–81.
- [40] X. Ros-Oton and J. Serra: The Dirichlet problem for the fractional Laplacian: regularity up to the boundary, J. Math. Pures Appl., 101 (2014), 275–302.
- [41] N. Sandrić: Homogenization of periodic diffusion with small jumps, J. Math. Anal. Appl., 435 (2016), 551–577.
- [42] R.L. Schilling and T. Uemura: Homogenization of symmetric Lévy processes on R^d, Rev. Roumaine Math. Pures Appl., 66 (2021), 243–253.
- [43] R.W. Schwab: Periodic homogenization for nonliear integro-differential equations, SIAM J. Math. Anal., 42 (2010), 2562– 2680.
- [44] Z. Shen: Periodic Homogenization of Elliptic Systems, Operator Theory: Advances and Applications, vol. 269, Advances in Partial Differential Equations (Basel), Birkhäuser/Springer, Cham, 2018.
- [45] Z. Shen and J. Zhuge: Convergence rates in periodic homogenization of systems of elasticity, Proc. Amer. Math. Soc., 145 (2017), 1187–1202.
- [46] E.M. Stein: Singular Integrals and Differentiability Properties of Functions, Princeton University Press, 1970.
- [47] T.A. Suslina: Homogenization of the elliptic Dirichlet problem: operator error estimates in L_2 , Mathematika, **59** (2013), 463–476.
- [48] T.A. Suslina: Homogenization of the Neumann problem for elliptic systems with periodic coefficients, SIAM J. Math. Anal., 45 (2013), 3453–3493.
- [49] M. Tomisaki: Homogenization of cádlág processes, J. Math. Soc. Japan, 44 (1992), 281–305.
- [50] L. Xu: Approximation of stable law in Wasserstein-1 distance by Stein's method, Ann. Appl. Probab., 29 (2019), 458–504.
- [51] X. Zhang and G. Zhao: Dirichlet problem for supercritical non-local operators, arXiv:1809.05712.
- [52] V.V. Zhikov: On operator estimates in homogenization theory, Dokl. Math., 72 (2005), 534–538.
- [53] V.V. Zhikov and S.E. Pastukhova: On operator estimates for some problems in homogenization theory, Russ. J. Math. Phys., 12 (2005), 515–524.

Xin Chen: School of Mathematical Sciences, Shanghai Jiao Tong University, 200240 Shanghai, P.R. China. chenxin217@sjtu.edu.cn

Zhen-Qing Chen: Department of Mathematics, University of Washington, Seattle, WA 98195, USA. zqchen@uw.edu

Takashi Kumagai: Department of Mathematics, Waseda University, Shinjuku, Tokyo, 169-8555, Japan. t-kumagai@waseda.jp

Jian Wang: School of Mathematics and Statistics & Key Laboratory of Analytical Mathematics and Applications (Ministry of Education) & Fujian Provincial Key Laboratory of Statistics and Artificial Intelligence, Fujian Normal University, 350007 Fuzhou, P.R. China.

jianwang@fjnu.edu.cn