

A RELATION BETWEEN THE DIRICHLET AND THE REGULARITY PROBLEM FOR PARABOLIC EQUATIONS

MARTIN DINDOŠ AND ERIK SÄTTERQVIST

ABSTRACT. We study the relationship between the Dirichlet and Regularity problem for parabolic operators of the form $L = \operatorname{div}(A\nabla\cdot) - \partial_t$ on cylindrical domains $\Omega = \mathcal{O} \times \mathbb{R}$, where the base $\mathcal{O} \subset \mathbb{R}^n$ is a 1-sided chord arc domain (and for one result Lipschitz) in the spatial variables.

In the paper we answer the question when the solvability of the L^p Regularity problem for L (denoted by $(R_L)_p$) can be deduced from the solvability of the $L^{p'}$ Dirichlet problem for the adjoint operator L^* (denoted $(D_L^*)_{p'}$).

We show that this holds if for at least of $q \in (1, \infty)$ the problem $(R_L)_q$ is solvable.

This result is a parabolic equivalent of two elliptic results of Kenig-Pipher (1993) and Shen (2006), the combination of which establishes the elliptic version of our result. For the converse, see Dindoš-Dyer (2019) where it is shown that $(R_L)_p$ implies $(D_L^*)_{p'}$ on parabolic $\operatorname{Lip}(1, 1/2)$ domains.

CONTENTS

1. Introduction	2
2. Basic definitions	4
2.1. Domain	4
2.2. Half time derivative and Parabolic Sobolev Space on \mathbb{R}^n .	6
2.3. Connections to Triebel-Lizorkin and Hajlasz-Sobolev spaces.	7
2.4. Parabolic Regularity problem	12
2.5. Localization	13
3. Parabolic results	16
4. Reverse Hölder inequality	18
5. Proof of <i>Theorem 1.3</i>	20
5.1. Unbounded case	20
5.2. Bounded case	23
6. Interpolation and <i>Theorem 1.4</i> .	28
6.1. Atomic spaces $\dot{H}S_{1,1/2}^{1,(\beta)}(\mathbb{R}^n)$.	28
6.2. Proof of <i>Theorem 1.4</i>	44
References	47

1. INTRODUCTION

In this paper we study the relationship between the parabolic Dirichlet and Regularity problems on parabolic cylinders $\Omega = \mathcal{O} \times \mathbb{R}$ with 1-sided chord arc (CAD) or Lipschitz base $\mathcal{O} \subset \mathbb{R}^n$, for the operators of the form

$$(1.1) \quad L = \operatorname{div}(A\nabla\cdot) - \partial_t,$$

where $A = A(X, t)$ is bounded, measurable and uniformly elliptic.

To be precise, for any $1 < p < \infty$, we wish to show that if the $L^{p'}$ Dirichlet problem is solvable (which we denote by $(D_{L^*})_{p'}$), then the L^p regularity problem is solvable (denoted by $(R_L)_p$) (see Definitions 2.3 and 2.39), provided this problem is solvable for at least one $q \in (1, \infty)$. Note that here $(D_{L^*})_{p'}$ denotes the solvability for the adjoint

$$L^* = \operatorname{div}(A^T\nabla\cdot) + \partial_t,$$

which is a backward in time parabolic operator. Through a change of variables $v(X, t) = u(X, -t)$, and $\tilde{A}(X, t) = A^T(X, -t)$ we can see that the solvability of $(D_{L^*})_{p'}$ is equivalent to solvability of $(D_{\tilde{L}})_{p'}$ where

$$\tilde{L}v := \operatorname{div}(\tilde{A}\nabla v) - \partial_t v = 0,$$

is the usual time-forward parabolic PDE.

Historically, the development of the parabolic theory has been inspired by the elliptic theory, as many elliptic results have their corresponding parabolic counterpart. This is the case of the presented result which was fully formulated by Shen in [37] on bounded Lipschitz domains. It uses substantially an earlier result of Kenig and Pipher [28] where it was been established that $((R_L)_q)$ implies $((R_L)_p)$ for all $1 < p < q + \varepsilon$ for some small $\varepsilon > 0$.

The converse, that $(R_L)_p$ implies $(D_{L^*})_{p'}$ has already been shown in both the elliptic [16, 28] and parabolic [13] settings, with the domain in [13] being a $\operatorname{Lip}(1, 1/2)$ domain.

One major obstacle in generalizing Shen's approach that arises in the parabolic regularity problem, is the fact that our boundary data contains a half-time derivative and thus we also expect to control the non-tangential maximal function of the half time derivative $N(D_t^{1/2}u)$. This is due to the fact that thanks to the natural parabolic scaling ∇u scales like a half derivative time-derivative $D_t^{1/2}u$, since the second derivative $\nabla^2 u$ behaves like $\partial_t u$. However, $D_t^{1/2}u$ is a non-local object which requires some thoughts on how to deal with it and what conditions to impose.

The initial formulation of the Regularity problem is due Fabes and Rivière [18] who formulated it for C^1 cylinders which was then re-formulated to Lipschitz cylinders by Brown [6, 7]. Here by Lipschitz cylinders we understand domains of the form $\Omega = \mathcal{O} \times [0, T]$ where $\mathcal{O} \subset \mathbb{R}^n$ is a bounded Lipschitz domain and time variable runs over a bounded interval $[0, T]$. Subsequent authors such as Mitrea [33], Nyström [36] or Castro-Rodríguez-López-Staubach [9] have dealt with this issue in a similar way.

What Brown has done was to consider solutions on Ω with initial data $u = 0$ at $t = 0$ and naturally extending u by zero for all $t < 0$. Then a definition of half-derivative using fractional integrals I_σ for functions $f \in C^\infty(-\infty, T)$ is used to give meaning to $D_t^{1/2}f$ and $D_t^{1/2}u$.

A variant of this approach (which we present here) is to consider the parabolic PDE on an infinite cylinder $\Omega = \mathcal{O} \times \mathbb{R}$. This can always be achieved by extending the coefficients by setting $A(X, t) = A(X, T)$ for $t > T$ and $A(X, t) = A(X, 0)$ for $t < 0$. Since the parabolic PDE has a defined direction of time the solution on the infinite cylinder $\Omega = \mathcal{O} \times \mathbb{R}$ with boundary data $u|_{\partial\Omega} = 0$ for $t < 0$ will coincide with the solution on the finite cylinder $\mathcal{O} \times [0, T]$ with zero initial data.

Having a well-defined half-derivative the authors [7, 9, 33, 36] then define the parabolic Regularity problem by asking for

$$(1.2) \quad \|N[\nabla u]\|_{L^p(\partial\Omega)} + \|N[D_t^{1/2}u]\|_{L^p(\partial\Omega)} \leq C(\|\nabla_T f\|_{L^p(\partial\Omega)} + \|D_t^{1/2}f\|_{L^p(\partial\Omega)}).$$

Some authors also ask for control of $\|N[D_t^{1/2}H_t u]\|_{L^p}$, where H_t is the one-dimensional Hilbert transform in the t -variable.

This approach is reasonable, if the domain is not time-varying, i.e. of the form $\mathcal{O} \times (t_0, t_1)$ but runs immediately into an obvious issue when this is no longer true, as any reasonable definition of half-derivative requires $u(X, t)$ to be defined for all $t \in \mathbb{R}$ or at least on a half-line.

This is the reason we only consider domains of the form $\mathcal{O} \times \mathbb{R}$ here to avoid further complications connected with dealing with this non-local term.

Our main result is split into two theorems as there are slightly different assumptions on the domain in each case.

Theorem 1.3. *Let \mathcal{O} be a 1-sided chord arc domain and L be a parabolic operator of the form (1.1) on the domain $\Omega = \mathcal{O} \times \mathbb{R}$.*

Suppose that the $L^{p'}$ Dirichlet problem for the adjoint operator, $(D_{L^})_{p'}$, is solvable, and that the regularity problem $(R_L)_q$ is solvable for some $1 < q < p$. Then $(R_L)_p$ is also solvable.*

Theorem 1.4. *Assume that \mathcal{O} is Lipschitz (either bounded or unbounded) and L be a parabolic operator of the form (1.1) on the domain $\Omega = \mathcal{O} \times \mathbb{R}$.*

Suppose that $(R_L)_p$ is solvable for some $1 < p < \infty$. Then $(R_L)_q$ is solvable for $1 < q < p$.

Theorem 1.3 is the parabolic version of the result of [37], while *Theorem 1.4* is an analogue of the elliptic result in [29].

By combining these two theorems we can obtain the following:

Theorem 1.5. *Assume that \mathcal{O} is Lipschitz (either bounded or unbounded) and L be a parabolic operator of the form (1.1) on the domain $\Omega = \mathcal{O} \times \mathbb{R}$.*

Suppose that $(D_{L^})_{p'}$ is solvable for some $p \in (1, \infty)$. Then either $(R_L)_p$ is solvable or $(R_L)_q$ is not solvable for any $1 < q < \infty$.*

In summary, thanks to *Theorem 1.5* and [13] we have the following two implications

$$(R_L)_p \implies (D_{L^*})_{p'} \quad \text{and} \quad (D_{L^*})_{p'} + (R_L)_q \implies (R_L)_p,$$

for parabolic operators on Lipschitz cylinders.

Let us recall that the elliptic versions of these implications played an important role in establishing the main result of the recent paper [15] where solvability of the elliptic Regularity problem for operators satisfying the Carleson condition has been

established. Thus, we expect our results to play a similar role in the parabolic version of an analogous result in the future.

We have left open some further research directions, *Theorem 1.4* has stronger assumption on the domain \mathcal{O} than the other result and thus it might be possible to improve this result further beyond the Lipschitz cylinders. What limits us currently are interpolation results such as *Proposition 6.46* which only works on boundaries that locally look like \mathbb{R}^n .

Another natural direction would be to consider time varying domains (restricting initially to domains that locally look like a graph of $\text{Lip}(1, 1/2)$ functions).

The paper is set up as follows. Section 2 contains definitions needed for the rest of the paper and Section 3 summarises known parabolic results we need for our proofs. *Theorem 1.3* is proven first, motivated by the method from [37]. The two cases $\text{diam } \mathcal{O} = \infty$ and $R := \text{diam } \mathcal{O} < \infty$ are considered separately.

Theorem 1.4 follows by real interpolation against a suitable endpoint space. We construct this endpoint space in section 6. This construction contains some significant novelties, as we need to construct “atoms” that respect different scaling between the space and time. With the interpolation result in hand, the final piece missing is the establishment of an endpoint bound for $\|N[\nabla u]\|_{L^1(\partial\Omega)}$ which closely follows its elliptic version as in [16].

2. BASIC DEFINITIONS

2.1. Domain. In general our domain will be a cylinder $\Omega = \mathcal{O} \times \mathbb{R}$, where the base \mathcal{O} is a 1-sided chord arc domain; that is it is a 1-sided non-tangentially accessible domain (also called uniform domain) with Ahlfors-David regular boundary. As the precise definition does not play any significant role here we omit the details and refer the reader to [12] which has the same settings as here and contains all definitions in great detail (in particular, the existence of interior Harnack chains and corkscrew balls for any boundary point plus the ADR condition).

We denote points in Ω by (X, t) or also (ξ, τ) , where $X, \xi \in \mathcal{O}$ and $t, \tau \in \mathbb{R}$, and points on the boundary by $(P, t) \in \partial\Omega = \partial\mathcal{O} \times \mathbb{R}$. The space Ω is equipped with the norm $\|(\xi, \tau)\| = \rho$, where ρ is the positive solution to the following equation

$$(2.1) \quad \frac{|\xi|^2}{\rho^2} + \frac{\tau^2}{\rho^4} = 1.$$

One can easily show that $\|(\xi, \tau)\| \sim |\xi| + |\tau|^{1/2}$ and that this norm scales correctly according to the parabolic nature of the PDE.

Next we introduce notation for balls, cubes and cylinders

- $B_r(X, t) := \{(Y, s) \in \Omega; \|(Y, s) - (X, t)\| \leq r\}$ (parabolic ball),
- $B_r(X) := \{Y \in \mathcal{O}; |Y - X| \leq r\}$ (ball in space only),
- $C_r(X, t) := B_r(X) \times (t - r^2, t + r^2)$ (parabolic cylinder),
- $Q_r(X, t) := \{(Y, s) \in \Omega; |Y_i - X_i| \leq r, |s - t| \leq r^2\}$ (parabolic cube),
- $\Delta_r(P, t) := \partial\Omega \cap B_r(P, t)$ (boundary ball),
- $T(\Delta_r(P, t)) := \Omega \cap B_r(P, t)$ (Carleson region).

Occasionally, we drop dependence of these sets on the point (X, t) and r and just write B, C, Q respectively. When doing this, the notation $r(B)$ or $r(Q)$ will denote the “radius” r of these sets. We use the similar convention also for Δ and $T(\Delta)$. When writing $2B$ or 5Δ , it means enlarged sets (with same center point but $2\times$ or $5\times$ the radius). Next, given $(X, t) \in \Omega$ we let $\delta(X, t) = \delta(X) = \text{dist}(X, \mathcal{O})$ denotes the parabolic distance to the boundary.

Note that because \mathcal{O} is a CAD we have that for $P \in \mathcal{O}$ and $\text{diam } \mathcal{O} > r > 0$ the existence of a corkscrew point $A_r(P)$. Thus for $\Delta = \Delta(P, t; r) \subset \Omega$ with $r < \text{diam } \mathcal{O}$ we define $V^+(\Delta) = (A_r(P), t + 100r^2)$ and $V^-(\Delta) = (A_r(P), t - 100r^2)$ as the “forward in time” and “backward in time” corkscrew point respectively.

Given an aperture $a > 0$, it is customary to define non-tangential approach regions (“cones”) as follows:

$$\tilde{\Gamma}_a(P, t) := \{(X, s) \in \Omega : \|(X, s) - (P, t)\| < (1 + a)\delta(X)\}.$$

However such cones lack one desired property, namely that the set (which can be thought of as the reverse cone originating from (X, t))

$$\{(P, \tau) \in \partial\Omega : (X, t) \in \tilde{\Gamma}_a(P, \tau)\}$$

always has the surface area comparable to the $\delta(X, t)^{n+1}$. To fix this issue we do as follows. Let

- $\gamma_a(P) := \{X \in \mathcal{O} : |X - P| < (1 + a)\delta(X)\},$
- $S_a(X) := \{P \in \partial\mathcal{O} : X \in \gamma_a(P)\},$
- $\tilde{S}_a(X) := \bigcup_{P \in S_a(X)} \{P' \in \partial\mathcal{O} : |P' - P| < a\delta(X)\},$
- $\tilde{\gamma}_a(P) := \{X \in \Omega : P \in \tilde{S}_a(X)\},$
- $\Gamma_a(P, t) := \{(X, s) \in \Omega : X \in \tilde{\gamma}_a(P), |s - t| < a\delta(X)^2\}.$

Note that $\Gamma_a(P, t)$ defined in the last line is our version of the non-tangential approach region and is equivalent to $\tilde{\Gamma}_a(P, t)$ since for any $a > 0$ there exists $a_1, a_2 > 0$ such that

$$\Gamma_{a_1}(P, t) \subset \tilde{\Gamma}_a(P, t) \subset \Gamma_{a_2}(P, t)$$

and vice versa. However, as the set $\tilde{S}_a(X)$ can be written as a union of boundary balls of radius $a\delta(X)$ it follows that the size of this set is always comparable to $\delta(X, t)^{n+1}$ as desired. For the rest of the paper we will let $a = 1$ and drop the subscript next to Γ .

Definition 2.2. Given a function $u : \Omega \rightarrow \mathbb{R}$, the non-tangential maximal function is given by

$$N_{\text{sup}}[u](P, t) := \sup_{\Gamma(P, t)} |u|.$$

Since ∇u for example, is not defined pointwise it is customary to define a version of non-tangential maximal function using L^2 averages as follows:

$$N[u](P, t) := \sup_{\Gamma(P, t)} \text{Av}^2[u], \quad \text{Av}^2[u](X, s) := \left(\int_{C_{\delta(X)/4}(X, s)} |u|^2 \right)^{1/2}.$$

Occasionally, we will also consider truncated non-tangential cones $\Gamma^r(P, t) := \Gamma(P, t) \cap B_r(P, t)$ and corresponding truncated non-tangential maximal function N^r .

We are now ready to define the Dirichlet problem

Definition 2.3. Let $1 < p < \infty$. We say that the L^p Dirichlet problem for L ($(D_L)_p$) is solvable if for each continuous $f \in L^p(\partial\Omega)$ the solution to

$$\begin{cases} Lu = 0, & \text{in } \Omega \\ u = f, & \text{on } \partial\Omega \end{cases}$$

satisfies

$$\|N[u]\|_{L^p(\partial\Omega)} \lesssim \|f\|_{L^p(\partial\Omega)}.$$

In the definition above it does not matter which version of non-tangential maximal function is used. When it is not explicitly stated, we will always mean N defined using the L^2 averages.

2.2. Half time derivative and Parabolic Sobolev Space on \mathbb{R}^n . Here we consider an appropriate function space for our boundary data on $\mathbb{R}^n = \mathbb{R}^{n-1} \times \mathbb{R}$.

As a rule of thumb, one derivative in time behaves as two derivatives in spatial variables. Thus if we impose boundary data having one spatial derivative, the correct order of the time derivative is $1/2$, as we would like to respect the correct parabolic scaling.

This problem has been studied previously in [24, 25, 36], who have followed [17] in defining the homogeneous parabolic Sobolev space $\dot{L}_{1,1/2}^p$ in the following way.

Definition 2.4. The *homogeneous parabolic Sobolev space* $\dot{L}_{1,1/2}^p(\mathbb{R}^n)$, for $1 < p < \infty$, is defined to consist of an equivalence class of functions f with distributional derivatives satisfying $\|f\|_{\dot{L}_{1,1/2}^p(\mathbb{R}^n)} < \infty$, where

$$(2.5) \quad \|f\|_{\dot{L}_{1,1/2}^p(\mathbb{R}^n)} = \|\mathbb{D}f\|_{L^p(\mathbb{R}^n)}$$

and

$$(2.6) \quad (\mathbb{D}f)^\wedge(\xi, \tau) := \|(\xi, \tau)\| \widehat{f}(\xi, \tau),$$

with $\|(\xi, \tau)\|$ on $\mathbb{R}^{n-1} \times \mathbb{R}$ defined by (2.1).

In addition, following [20], we define a parabolic half-order time derivative by

$$(2.7) \quad (\mathbb{D}_n f)^\wedge(\xi, \tau) := \frac{\tau}{\|(\xi, \tau)\|} \widehat{f}(\xi, \tau).$$

If $0 < \alpha \leq 2$, then for $g \in C_c^\infty(\mathbb{R})$ the *one-dimensional fractional differentiation operators* D_α are defined by

$$(2.8) \quad (D^\alpha g)^\wedge(\tau) := |\tau|^\alpha \widehat{g}(\tau).$$

It is also well known that if $0 < \alpha < 1$ then

$$(2.9) \quad D^\alpha g(s) = c \int_{\mathbb{R}} \frac{g(s) - g(\tau)}{|s - \tau|^{1+\alpha}} d\tau$$

whenever $s \in \mathbb{R}$. If $h(x, t) \in C_c^\infty(\mathbb{R}^n)$ then by $D_t^\alpha h : \mathbb{R}^n \rightarrow \mathbb{R}$ we mean the function $D^\alpha h(x, \cdot)$ defined a.e. for each fixed $x \in \mathbb{R}^{n-1}$. When $\alpha = 1/2$ we have

$$D_t^{1/2} D_t^{1/2} = -H_t \partial_t,$$

where H_t is the Hilbert transform taken in the time variable.

We record the connection between \mathbb{D} , \mathbb{D}_n and $D_t^{1/2}$. By [13, 19, 20] we have that

$$(2.10) \quad \|\mathbb{D}f\|_{L^p(\mathbb{R}^n)} \sim \|\mathbb{D}_n f\|_{L^p(\mathbb{R}^n)} + \|\nabla f\|_{L^p(\mathbb{R}^n)} \sim \|D_t^{1/2} f\|_{L^p(\mathbb{R}^n)} + \|\nabla f\|_{L^p(\mathbb{R}^n)},$$

for all $1 < p < \infty$, as well as at the BMO end-point:

$$(2.11)$$

$$\|\mathbb{D}f\|_{\text{BMO}(\mathbb{R}^n)} \sim \|\mathbb{D}_n f\|_{\text{BMO}(\mathbb{R}^n)} + \|\nabla f\|_{\text{BMO}(\mathbb{R}^n)} \sim \|D_t^{1/2} f\|_{\text{BMO}(\mathbb{R}^n)} + \|\nabla f\|_{\text{BMO}(\mathbb{R}^n)}.$$

Here ∇ denotes the usual gradient in the variables $x \in \mathbb{R}^{n-1}$ and BMO denotes the usual space of bounded mean oscillations (with respect to parabolic balls).

When $p = 1$ we no longer have (2.10) and hence the norm of $\dot{L}_{1,1/2}^1(\mathbb{R}^n)$ might not be equivalent to $\|\nabla f\|_{L^1} + \|D_t^{1/2} f\|_{L^1}$. We therefore define the space

$$(2.12) \quad \dot{W}_{1,1/2}^p(\mathbb{R}^n) = \{f \in L_{loc}^1(\mathbb{R}^n) : \|\nabla f\|_{L^p(\mathbb{R}^n)} + \|D_t^{1/2} f\|_{L^p(\mathbb{R}^n)} < \infty\},$$

with norm $\|\nabla f\|_{L^p} + \|D_t^{1/2} f\|_{L^p}$. Modulo constants when $p \in (1, \infty)$ we have $\dot{W}_{1,1/2}^p(\mathbb{R}^n) = \dot{L}_{1,1/2}^p(\mathbb{R}^n)$ and the norms are comparable, but when $p = 1, \infty$ the spaces might not coincide. These spaces will be useful for interpolation. We will also need a weaker end-point space than $\dot{W}_{1,1/2}^\infty(\mathbb{R}^n)$. We denote by $\dot{W}_{1,1/2}^{\infty, \text{BMO}}(\mathbb{R}^n)$ the space such that $\nabla f \in L^\infty(\mathbb{R}^n)$ and $D_t^{1/2} f \in \text{BMO}(\mathbb{R}^n)$.

2.3. Connections to Triebel-Lizorkin and Hajlasz-Sobolev spaces. Here and in what follows we will use notation $X = (x, t)$ to denote points in \mathbb{R}^n with components $x \in \mathbb{R}^{n-1}$ and $t \in \mathbb{R}$.

Let us recall the definition of Triebel-Lizorkin spaces $F_{p,q}^s$. The fact that our operator \mathbb{D} is defined by (2.6) implies that it fits naturally to the settings of these spaces for $q = 2$, but we need to account for the parabolic scaling.

Definition 2.13. Let $s \in (0, \infty)$ and $p \in (0, \infty]$. Let $\varphi \in \mathcal{S}(\mathbb{R}^n)$ such that

$$\text{supp } \hat{\varphi} \subset \{(\xi, \tau) : 1/2 \leq \|(\xi, \tau)\| \leq 3/2\},$$

and

$$|\hat{\varphi}(\xi, \tau)| \geq c_0 > 0 \text{ for } 3/5 \leq \|(\xi, \tau)\| \leq 5/3.$$

Let $\varphi_\rho(x, t)$ denote the usual parabolic dilation by ρ , that is

$$\varphi_\rho(x, t) = \rho^{-(n+1)} \varphi(x/\rho, t/\rho^2).$$

The homogeneous parabolic Triebel-Lizorkin space $\dot{F}_{p,2}^s(\mathbb{R}^n)$ is defined as the collection of all $f \in \mathcal{S}'(\mathbb{R}^n)$ such that $\|f\|_{\dot{F}_{p,2}^s(\mathbb{R}^n)} < \infty$ where

$$(2.14) \quad \|f\|_{\dot{F}_{p,2}^s(\mathbb{R}^n)} = \left\| \left(\sum_{k \in \mathbb{Z}} 4^{ks} |\varphi_{2^{-k}} * f|^2 \right)^{1/2} \right\|_{L^p(\mathbb{R}^n)},$$

with the usual modification when $p = \infty$.

By the Littlewood-Paley theorem when $M_k f = \varphi_{2^{-k}} * f$ then for $1 < p < \infty$

$$\|f\|_{L^p(\mathbb{R}^n)} \approx \|(M_k f)_{\ell^2(\mathbb{Z})}\|_{L^p(\mathbb{R}^n)} = \|f\|_{\dot{F}_{p,2}^0(\mathbb{R}^n)}.$$

Hence

$$\|f\|_{\dot{L}_{1,1/2}^p(\mathbb{R}^n)} \approx \|(2^k M_k f)_{\ell^2(\mathbb{Z})}\|_{L^p(\mathbb{R}^n)} = \|f\|_{\dot{F}_{p,2}^1(\mathbb{R}^n)}.$$

Note that is is only true when parabolic scaling is applied, in the usual Euclidean metric we have $\dot{L}_{1/2}^p(\mathbb{R}) = \dot{F}_{p,2}^{1/2}(\mathbb{R})$ and $\dot{L}_1^p(\mathbb{R}^{n-1}) = \dot{F}_{p,2}^1(\mathbb{R}^{n-1})$ (see [34]).

The paper [43] gave a different characterisation of these spaces using averages. Again we have to account for parabolic scaling, but the argument adapts easily. Let us first define

$$(2.15) \quad S_{s, \text{disc}}(f)(x, t) = \left(\sum_{k \in \mathbb{Z}} 4^{ks} \left(f(x, t) - \int_{B_{2^k}(x, t)} f \right)^2 \right)^{1/2}.$$

Here $B_\alpha(x, t)$ is the parabolic ball centred at (x, t) with radius α . By Theorem 1.1 of [43] $\|S_{s, \text{disc}}(f)\|_{L^p} \sim \|f\|_{\dot{F}_{p,2}^s}$ for all $1 < p < \infty$. For our further considerations we would prefer continuous version of the square function (2.15). It is not hard to see that by averaging over set of balls of radius $2^{\tau k}$, $1 \leq \tau \leq 2$ we may replace it by

$$(2.16) \quad S_s(f)(x, t) = \left(\int_0^\infty \alpha^{-2s-1} \left(f(x, t) - \int_{B_\alpha(x, t)} f \right)^2 d\alpha \right)^{1/2}.$$

In particular, for $s = 1$ we have $\|f\|_{\dot{L}_{1,1/2}^p(\mathbb{R}^n)} \approx \|S_1(f)\|_{L^p(\mathbb{R}^n)}$ for $1 < p < \infty$.

We develop an interesting connection between this square function and new characterisation of the fact that $\mathbb{D}f \in \text{BMO}(\mathbb{R}^n)$. Recall that [14] (Theorem 2.3) showed the following:

Theorem 2.17. *Let $f : \mathbb{R}^{n-1} \times \mathbb{R} \rightarrow \mathbb{R}$ and $f \in \text{Lip}(1, 1/2)$ then the following conditions are equivalent:*

- $\mathbb{D}f \in \text{BMO}(\mathbb{R}^n)$
-

$$(2.18) \quad \sup_{B_r} \frac{1}{|B_r|} \int_{B_r} \int_{\|(y,s)\| \leq r} \frac{|f(x+y, t+s) - 2f(x, t) + f(x-y, t-s)|^2}{\|(y, s)\|^{n+3}} dy ds dx dt = B_1 < \infty,$$

$$(2.19) \quad \|\mathbb{D}f\|_{\text{BMO}}^2 \sim B_1.$$

Now in the light of (2.16) we need to obtain a new version of the condition (2.18) where the absolute value is moved to the outside of the interior integral. A method how to establish such conditions was developed by Strichartz [40] and our paper [14] adapts it to parabolic settings. We claim the following:

Theorem 2.20. *Let $f : \mathbb{R}^{n-1} \times \mathbb{R} \rightarrow \mathbb{R}$ and $f \in \text{Lip}(1, 1/2)$ then the following conditions are equivalent:*

- $\mathbb{D}f \in \text{BMO}(\mathbb{R}^n)$

$$(2.21) \quad \sup_{B_r} \frac{1}{|B_r|} \int_{B_r} \int_0^r \alpha^{-3} \left| f(x, t) - \fint_{B_\alpha(x, t)} f \right|^2 d\alpha dt dx = B_2 < \infty.$$

Moreover, $\|\mathbb{D}f\|_{\text{BMO}}^2 \sim B_2$ and r is the diameter of the parabolic ball B_r . The statement also holds with parabolic cubes Q_r replacing the parabolic balls B_r .

Perhaps the two conditions (2.18) and (2.21) do not immediately look close, but after writing the interior integral in polar coordinates will reveal that we are taking averages over the shells $\|(y, s)\| = \text{const}$ with the same scaling factor $\|(y, s)\|^{-3}$ as in (2.21). The fact that the absolute value in (2.21) is on the outside of the most interior integral is the innovation [43] has introduced.

Proof. Let us recall the method of Strichartz adapted to the parabolic setting in [14]. Write $F = \mathbb{D}f$ where F is a tempered distribution. Let

$$\varphi = \delta(0, 0) - \chi_{\tilde{Q}_1(0,0)},$$

where $Q_1(0, 0)$ is a cube ball in \mathbb{R}^n . We may take a ball here instead of the cube, it will change the Fourier transform somewhat but not the gist of the argument (we prefer not to deal with Bessel's functions). Then

$$(2.22) \quad \widehat{\varphi}(\xi, \tau) = 1 - \frac{1 - e^{-i\tau}}{i\tau} \prod_{j=1}^{n-1} \frac{1 - e^{-i\xi_j}}{i\xi_j}.$$

with $\widehat{\varphi}(\xi, \tau) \sim |\xi| + |\tau|$ for small ξ and τ . We let

$$\widehat{\psi} = \frac{\widehat{\varphi}(\xi, \tau)}{\|(\xi, \tau)\|},$$

where ψ_ρ denotes the usual parabolic dilation ρ defined above. We may rewrite (2.21) as

$$(2.23) \quad \sup_{Q_r} \frac{1}{|Q_r|} \int_{Q_r} \int_0^r (\psi_\rho * F)^2 \frac{d\rho}{\rho} dx dt \sim B_2.$$

The function ψ satisfy the following conditions for some $\epsilon_i > 0$

$$(2.24) \quad \begin{aligned} \int \psi dx dt &= 0, \\ |\psi(x, t)| &\lesssim \|(x, t)\|^{-n-1-\epsilon_1} \text{ for } \|(x, t)\| \geq a > 0, \\ |\widehat{\psi}(\xi, \tau)| &\lesssim \|(\xi, \tau)\|^{\epsilon_2} \text{ for } \|(\xi, \tau)\| \leq 1, \\ |\widehat{\psi}(\xi, \tau)| &\lesssim \|(\xi, \tau)\|^{-\epsilon_3} \text{ for } \|(\xi, \tau)\| \geq 1. \end{aligned}$$

Therefore if $\mathbb{D}f = F \in \text{BMO}(\mathbb{R}^n)$ then $B_2 \lesssim \|\mathbb{D}\phi\|_*^2$ by [40, Theorem 2.1]; this shows that first condition in [Theorem 2.20](#) implies the second one.

For the converse we proceed via an analogue of the proof of [40, Theorem 2.6]. Consider

$$\widehat{\theta}(\xi, \tau) = \|(\xi, \tau)\| \widehat{\zeta}(\xi, \tau),$$

where $\zeta \in C_0^\infty(\mathbb{R})$. Let H_{00}^1 be the dense subclass of continuous H^1 functions g such that g and all its derivatives decay rapidly, see [39, p. 225]. Via an analogue of [21, Theorem 3], [40, Lemma 2.7] by assuming (2.23) if $g \in H_{00}^1(\mathbb{R}^n)$, then

$$(2.25) \quad \left| \int_0^\infty \iint_{\mathbb{R}^{n-1} \times \mathbb{R}} \psi_\rho * F(x, t) \theta_\rho * g(x, t) dx dt \frac{d\rho}{\rho} \right| \lesssim B_2^{1/2} \|g\|_{H^1}.$$

Let

$$(2.26) \quad m(\xi, \tau) = \int_0^\infty \overline{\hat{\psi}(-\rho\xi, -\rho^2\tau)} \|(\xi, \tau)\| \zeta(\rho \|(\xi, \tau)\|) d\rho.$$

Such function m is homogeneous of degree zero, smooth away from the origin and the associated Fourier multipliers M is a Calderón-Zygmund operator that preserves the class H_{00}^1 and is bounded on H^1 .

The non-degeneracy condition from [8] is the property that $|m(r\xi, r^2\tau)|^2$ does not vanish identically in r for $(\xi, \tau) \neq (0, 0)$. Hence as in [14] we may conclude that $(h, g) = (F, g)$ for all $g \in H_{00}^1$ where h is a BMO function with $\|h\|_*^2 \lesssim B_2$. The identity $\hat{h} = \hat{F}$ does not need to hold at the origin wherefore $\hat{h} - \hat{F}$ may be supported at the origin and hence $F = h + p$ where p is a polynomial. Due to the assumption $\phi \in \text{Lip}(1, 1/2)$, clearly F must be a tempered distribution. Hence as in [40] we may conclude $F = h \in \text{BMO}(\mathbb{R}^n)$. This proves the claim. \square

Now, we shall make connection between $\dot{L}_{1,1/2}^p(\mathbb{R}^n)$ and the Hajłasz-Sobolev space $\dot{M}_d^{1,p}(\mathbb{R}^n)$. In fact, we want to consider a more general setting and work on a set $\partial\mathcal{O} \times \mathbb{R}$, where $\mathcal{O} \subset \mathbb{R}^n$ is a 1-sided chord arc domain and thus $\partial\mathcal{O}$ is Ahlfors-David regular set with respect to the $n - 1$ -dimensional Hausdorff measure. In particular this allows us to talk about the Lebesgues spaces $L^p(\partial\mathcal{O})$. Thus, when $\mathcal{O} = \mathbb{R}_+^n$, we have $\partial\mathcal{O} = \mathbb{R}^{n-1}$ and therefore the discussion below applies to the space $\mathbb{R}^n = \mathbb{R}^{n-1} \times \mathbb{R}$ discussed above.

Definition 2.27. Let d be a complete metric on a measure space (\mathcal{X}, μ) . For $p \geq 1$, $0 < s \leq 1$ we say that a function $f \in L_{loc}^p(\mathcal{X})$ belongs to the Hajłasz-Sobolev space $M^{s,p}(\mathcal{X})$ if there exists a set E of μ -measure zero such that

$$(2.28) \quad |f(X) - f(Y)| \leq d(X, Y)^s [|g(X) + g(Y)|] \quad \forall X, Y \in \mathcal{X} \setminus E.$$

The $M_d^{s,p}(\mathcal{X})$ norm of f is given as the infimum of L^p norms of all such g .

In what follows we shall consider the space $M_d^{1,p}(\partial\mathcal{O} \times \mathbb{R})$ defined with respect to the parabolic distance function

$$(2.29) \quad d(X, X') = d((x, t), (x', t')) = \|(x - x', t - t')\| \sim |x - x'| + |t - t'|^{1/2}.$$

The underlying measure is the n -dimensional Hausdorff measure on $\partial\mathcal{O} \times \mathbb{R} \subset \mathbb{R}^{n+1}$. Here we introduce notation that matches the convention used above, i.e., $X \in \partial\mathcal{O} \times \mathbb{R}$ has the spatial coordinate x and time coordinate t .

Associated with the space $\dot{M}^{1,p}$ is the so-called *Sobolev sharp maximal function* f^\sharp defined by

$$(2.30) \quad f^\sharp(X) = \sup_{B: x \in B} \frac{1}{r(B)} \int_B |f - f_B|.$$

Here B denotes a surface ball on the set $\partial\mathcal{O} \times \mathbb{R}$ with respect to metric d of radius $r(B)$, i.e.,

$$B = B_{r(B)}(Y) = \{Z \in \partial\mathcal{O} \times \mathbb{R} : d(Y, Z) < r(B)\},$$

and f_B denotes the average $f_B = \int_B f(Z) dZ$.

Let us also introduce M which is the usual (non-centred) Hardy-Littlewood parabolic maximal function

$$(2.31) \quad M(f)(X) = \sup_{B: X \in B} \int_B |f| d\mathcal{H}^n,$$

where the supremum is taken over all surface parabolic balls B that contain point X . Denote by

$$M_q(f) = M(|f|^q)^{1/q}.$$

It follows from the usual boundedness of the maximal operator (which is true since $\partial\Omega$ is an ADR set) that M is L^p bounded for $p > 1$ and M_q for $p > q$.

Let us denote by M^t and M^x the variants of the maximal operator when the averages are taken only in t or x direction respectively. That is

$$M^x(f)(x, t) = \sup_{\Delta: x \in \Delta} \int_{\Delta} |f(y, t)| dy,$$

and $\Delta = \Delta(z) = \{y \in \partial\mathcal{O} : |y - z| < r\}$, with M^t defined analogously. We also define M_q^x and M_q^t in analogy to M_q .

As on each fibre for a fixed t we have M^x being bounded on $L^p(\partial\mathcal{O})$ for $p > 1$ we see by Fubini that M^x and M^t are $L^p(\partial\mathcal{O} \times \mathbb{R})$ bounded for $p > 1$ and the same is also true for M_q^x and M_q^t for $p > q$.

On metric spaces with doubling measure μ , that is for some constant $C_s > 1$

$$\mu(B(X, 2r)) \leq C_s \mu(B(X, r)), \quad \text{for all points } X \text{ and } r > 0,$$

we have by [30] for all $p \geq 1$:

$$\|f\|_{\dot{M}_d^{1,p}} \sim \|f^\sharp\|_{L^p}$$

and (2.28) holds with g replaced by f^\sharp . In our case for d defined by (2.29) the volume of surface ball of radius r is proportional to r^{n+1} and hence the doubling constant C_s can be taken to be 2^{n+1} . As in [30] $s = \log_2 C_s$ is called the doubling dimension of our space, thus for $\partial\mathcal{O} \times \mathbb{R}$ equipped with the metric d the doubling dimension is $s = n + 1$.

In particular, we have on our space $\dot{M}_d^{1,p}(\partial\mathcal{O} \times \mathbb{R})$ for a.e. $t \in \mathbb{R}$:

$$(2.32) \quad |f(x, t) - f(y, t)| \leq |x - y| [g(x, t) + g(y, t)] \quad \text{for } \mathcal{H}^{n-1}\text{-almost every } x, y \in \partial\mathcal{O},$$

and for a.e. $x \in \partial\mathcal{O}$

$$(2.33) \quad |f(x, t) - f(x, \tau)| \leq |t - \tau|^{1/2} [g(x, t) + g(x, \tau)] \quad \text{for almost every } t, \tau \in \mathbb{R}.$$

For all $p > 1$, by Corollary 1.2 of [32] when t is fixed we have $\dot{L}_1^p(\mathbb{R}^{n-1}) = \dot{F}_{p,2}^1(\mathbb{R}^{n-1}) = \dot{M}_{|\cdot|}^{1,p}(\mathbb{R}^{n-1})$ and so (2.32) is equivalent to having $\nabla f \in L^p(\mathbb{R}^n)$. Here $\dot{M}_{|\cdot|}^{1,p}$ is defined with respect to the Euclidean distance $|\cdot|$.

When \mathcal{O} is a Lipschitz domain, the Triebel-Lizorkin space $\dot{F}_{p,2}^1(\partial\mathcal{O})$ is still well-defined and we might take it as the definition of the space $\dot{L}_1^p(\partial\mathcal{O})$. Thus in this case we still have $\dot{L}_1^p(\partial\mathcal{O}) = \dot{F}_{p,2}^1(\partial\mathcal{O}) = \dot{M}_{|\cdot|}^{1,p}(\partial\mathcal{O})$ as Corollary 1.2 of [32] applies.

Also since $\dot{L}_{1/2}^p(\mathbb{R}) = \dot{F}_{p,2}^{1/2}(\mathbb{R})$, by Corollary 1.2 of [32] then $\dot{L}_{1/2}^p(\mathbb{R}) = \dot{F}_{p,2}^{1/2}(\mathbb{R}) \subset \dot{F}_{p,\infty}^{1/2}(\mathbb{R}) = \dot{M}_{|\cdot|}^{1/2,p}(\mathbb{R})$ (here the switch to 1/2 derivative comes due to the fact that (2.33) uses the usual Euclidean metric on \mathbb{R}).

Hence for $f \in \dot{L}_{1,1/2}^p(\partial\mathcal{O} \times \mathbb{R})$, $p > 1$ we have that both (2.32) and (2.33) hold. This in turn implies that so does

$$(2.34) \quad |f(x, t) - f(y, \tau)| \leq d((x, t), (y, \tau))[h(x, t) + h(y, \tau)]$$

for almost every pairs of points in $\partial\mathcal{O} \times \mathbb{R}$ with $h = M^t(g) + M^x(g) \in L^p(\partial\mathcal{O} \times \mathbb{R})$. It follows that

$$(2.35) \quad \dot{L}_{1,1/2}^p(\partial\mathcal{O} \times \mathbb{R}) \subset \dot{M}_d^{1,p}(\partial\mathcal{O} \times \mathbb{R}) \quad \text{and} \quad \|f\|_{\dot{M}_d^{1,p}(\partial\mathcal{O} \times \mathbb{R})} \lesssim \|f\|_{\dot{L}_{1,1/2}^p(\partial\mathcal{O} \times \mathbb{R})}, \quad p > 1.$$

Consider any bounded nonnegative function supported in B and let $\chi(B) = \int_B \chi$. Suppose that $\chi(B) \sim |B|$. Consider

$$\begin{aligned} \left| f(X) - \frac{1}{\chi(B)} \int_B f \chi \right| &= \left| \frac{1}{\chi(B)} \int_B (f(X) - f(Z)) \chi(Z) dZ \right| \\ &\leq \sup_{Z \in B} d(X, Z) \frac{1}{\chi(B)} \int_B [f^\sharp(X) + f^\sharp(Z)] \chi(Z) dZ, \end{aligned}$$

which holds by (2.28) and the fact that f^\sharp can be used in place of g . Since $X \in B$ we have $\sup_{Z \in B} d(X, Z) \lesssim r(B)$ and hence

$$(2.36) \quad \frac{1}{r(B)} \left| f(X) - \frac{1}{\chi(B)} \int_B f \chi \right| \lesssim f^\sharp(X) + \int_B f^\sharp,$$

for any X for which $X \in B$.

2.4. Parabolic Regularity problem. The previous section has defined the space $\dot{L}_{1,1/2}(\partial\mathcal{O} \times \mathbb{R})$ in the case when $\mathcal{O} \subset \mathbb{R}^n$ is a bounded or unbounded Lipschitz domain. In particular, our definition implies that when

$$\mathcal{O} = \{(x', x_n) : x_n > \phi(x')\},$$

for some Lipschitz function $\phi : \mathbb{R}^{n-1} \rightarrow \mathbb{R}$, then $f \in \dot{L}_{1,1/2}(\partial\Omega)$ if

$$(2.37) \quad f \circ \pi \in \dot{L}_{1,1/2}(\mathbb{R}^n), \quad \text{where} \quad \pi(x', t) = (x', \phi(x'), t).$$

We would like to extend our definition to the case \mathcal{O} is a 1-sided chord arc domain in a way that is consistent with the previous subsection and so that for example (2.35) still holds.

Motivated by [35] with adaptation to our setting, we may define the spatial gradient on $\partial\mathcal{O}$ as above using the *Hajlasz-Sobolev space*. Namely, For a Borel function f :

$\partial\mathcal{O} \times \mathbb{R} \rightarrow \mathbb{R}$ we say that a Borel function $g : \partial\mathcal{O} \times \mathbb{R} \rightarrow \mathbb{R}$ is a *Hajlasz upper spatial gradient* of f if

$$|f(X, t) - f(Y, t)| \leq |X - Y|(g(X, t) + g(Y, t)) \quad \text{for a.e. } (X, t), (Y, t) \in \partial\mathcal{O} \times \mathbb{R}.$$

We denote the collection of all Hajlasz upper spatial gradients of f as $\mathcal{D}_s(f)$. For a measurable function f on $\partial\Omega = \partial\mathcal{O} \times \mathbb{R}$ we define the Sobolev space $\dot{L}_{1,1/2}^p(\partial\Omega)$ via the norm

$$(2.38) \quad \|f\|_{\dot{L}_{1,1/2}^p(\partial\Omega)} := \inf\{\|g\|_{L^p(\partial\Omega)} : g \in L^p(\partial\Omega) \cap \mathcal{D}_s(f)\} + \|D_{1/2}^t f\|_{L^p(\partial\Omega)}.$$

We can also see that (2.35) still holds as it is a consequence of (2.32)-(2.33) which can be shown for f satisfying (2.38). Also, it is easy to see that this defines the same space as previously for \mathcal{O} Lipschitz (c.f. [35]).

Now we are ready to define the Regularity problem.

Definition 2.39. Let $1 < p < \infty$, and the domain Ω be as above. We say that the L^p Regularity problem for $L(R_L)_p$ is solvable if for each continuous $f \in \dot{L}_{1,1/2}^p(\partial\Omega)$ the solution to

$$\begin{cases} Lu = 0, & \text{in } \Omega, \\ u = f, & \text{on } \partial\Omega, \end{cases}$$

satisfies

$$(2.40) \quad \|N[\nabla u]\|_{L^p(\partial\Omega)} \lesssim \|f\|_{\dot{L}_{1,1/2}^p(\partial\Omega)}.$$

Remark 2.41. One might object that a condition of the type

$$(2.42) \quad \|N[D_{1/2}^t u]\|_{L^p(\partial\Omega)} \lesssim \|f\|_{\dot{L}_{1,1/2}^p(\partial\Omega)},$$

should be satisfied too (c.f. [7, 9, 33, 36]). However, as it turns out, this condition is superfluous. Indeed by [12] we have

$$(2.43) \quad \|N[D_{1/2}^t u]\|_{L^p(\partial\Omega)} \lesssim \|N[\nabla u]\|_{L^p(\partial\Omega)} + \|f\|_{\dot{L}_{1,1/2}^p(\partial\Omega)},$$

and clearly (2.42) follows from (2.43) and (2.40). This simplifies our task, as we only need to establish bounds for $N[\nabla u]$ and the bounds for the half-time derivative will follow.

2.5. Localization. We shall establish the following useful result that will be needed later.

Proposition 2.44. *Let $\Omega = \mathcal{O} \times \mathbb{R}$, where $\mathcal{O} \subset \mathbb{R}^n$ is a 1-sided chord arc domain. Assume that for some $p > 1$ $f \in \dot{L}_{1,1/2}^p(\partial\Omega)$ (with $\dot{L}_{1,1/2}^p$ defined by (2.38)). There exists $h \in L^p(\partial\Omega)$ with $\|h\|_{L^p(\partial\Omega)} \lesssim \|f\|_{\dot{L}_{1,1/2}^p(\partial\Omega)}$ for which the following holds:*

Consider any parabolic ball $B \subset \mathbb{R}^n \times \mathbb{R}$ of radius r centered at the boundary $\partial\Omega$, let $\Delta = B \cap \partial\Omega$ and φ be a $C_0^\infty(\mathbb{R}^n \times \mathbb{R})$ cutoff function such that $\varphi = 1$ on $2B$ and $\varphi = 0$ outside $3B$ with $\|\partial_t \varphi\|_{L^\infty} \lesssim r^{-2}$ and $\|\nabla_x \varphi\|_{L^\infty} \lesssim r^{-1}$. Then we have the following estimate:

$$\|(f - f_\Delta) \varphi\|_{\dot{L}_{1,1/2}^p(\partial\Omega)} \lesssim \|h\|_{L^p(32\Delta)} + \|D_t^{1/2} f\|_{L^p(32\Delta)} \lesssim \|f\|_{\dot{L}_{1,1/2}^p(\partial\Omega)}.$$

Furthermore for all $(x, t) \in 8\Delta$ we have

$$(2.45) \quad |D_t^{1/2}((f - f_\Delta f)\varphi)(x, t) - D_t^{1/2}f(x, t)\varphi(x, t)| \lesssim h(x, t).$$

Proof. Recall that we know that we have for f (2.34) and hence

$$(2.46) \quad |f(x, t) - f(y, \tau)| \leq d((x, t), (y, \tau))[f^\sharp(x, t) + f^\sharp(y, \tau)]$$

for a.e. pair of points $(x, t), (y, \tau) \in \partial\Omega$. We have to be very careful where we use the condition (2.46) as this does not characterise the space of functions for which $\|D_t^{1/2}f\|_{L^p} < \infty$. We have lost a bit of regularity when we have used in the previous section the inclusion $\dot{F}_{p,2}^{1/2} \subset \dot{F}_{p,\infty}^{1/2}$ which is proper. However, we assume $\varphi \in C_0^\infty(\mathbb{R}^n \times \mathbb{R})$ which is better than just $D_t^{1/2}\varphi \in L^p(\mathbb{R}^n \times \mathbb{R})$. This is an extra piece of regularity that will make our argument work. We calculate $D_t^{1/2}((f - f_\Delta f)\varphi)$. Clearly,

$$(2.47) \quad D_t^{1/2}((f - f_\Delta f)\varphi)(x, t) = c \int_{\mathbb{R}} \frac{(f(x, t) - f_\Delta f)\varphi(x, t) - (f(x, s) - f_\Delta f)\varphi(x, s)}{|t - s|^{3/2}} ds$$

$$(2.48) \quad = \varphi(x, t)D_t^{1/2}(f)(x, t) + c \int_{s \in \mathbb{R} \cap \{(x, s) \in 3\Delta\}} \frac{\varphi(x, t) - \varphi(x, s)}{|t - s|^{3/2}} (f(x, s) - f_\Delta f) ds.$$

The second term needs to be considered further. We need to think about when the term $\varphi(x, t) - \varphi(x, s) \neq 0$ as otherwise there is nothing to do. If the point (x, t) is far from support of some enlargement of Δ , say 8Δ , whenever $\varphi(x, t) - \varphi(x, s) \neq 0$ we will have $|t - s| \approx 2^i r^2$ for some $i = 2, 3, \dots$, for all $s \in \mathbb{R}$ such that $(x, s) \in 3\Delta$. We use a crude bound for the numerator $|\varphi(x, t) - \varphi(x, s)| \leq 1$. This allows us to conclude that when $(x, t) \notin 8\Delta$ then the second term will be bounded by

$$(2.49) \quad \left| \int_{\mathbb{R}} \frac{\varphi(x, t) - \varphi(x, s)}{|t - s|^{3/2}} (f(x, s) - f_\Delta f) ds \right| \lesssim (2^{-3i/2})r^{-1} \int_{s \in 3\Delta} |f(x, s) - f_\Delta f| ds$$

$$\lesssim (2^{-3i/2})r^{-1} M^{t < 4r^2} (|f - f_\Delta f|)(x, t'), \quad \text{for some } i \geq 2.$$

Here (x, t') is an arbitrary point in 3Δ and $M^{t < 4r^2}$ is the Hardy-Littlewood maximal function in t -variable only on all balls of size $< 4r^2$.

This leaves us to consider the case $(x, t) \in 8\Delta$. It is then possible that $|t - s|$ is arbitrary small for $s \in \mathbb{R}$ such that $\varphi(x, t) - \varphi(x, s) \neq 0$. We use the fact that φ is Lipschitz in t -variable with Lipschitz constant $\approx r^{-2}$. We also split the integral to sets on which $|s - t| \approx 2^{-j}r^2$ for $j = 0, 1, 2, \dots$. This gives us for the second term of (2.48):

$$(2.50) \quad \left| \int_{\mathbb{R}} \frac{\varphi(x, t) - \varphi(x, s)}{|t - s|^{3/2}} (f(x, s) - f_\Delta f) ds \right| \lesssim \sum_{j=0}^{\infty} r^{-2} \int_{|s-t| \approx 2^{-j}r^2} \frac{|f(x, s) - f_\Delta f|}{|t - s|^{1/2}} ds.$$

By introducing extra term $|t - s|^{1/2}$ into the denominator we again recognise a maximal function in t -variable. Thus continuing our calculation we have

$$\lesssim \sum_{j=0}^{\infty} 2^{-j/2} r^{-1} M^{t < 16r^2} (|f - f_\Delta f|)(x, t) \approx r^{-1} M^{t < 16r^2} (|f - f_\Delta f|)(x, t).$$

We now estimate this maximal function. Let $(x, t) \in 8\Delta$. Using (2.36) for $B = 8\Delta$ we get that

$$(2.51) \quad r^{-1} \left| f(x, t) - \fint_{\Delta} f \right| \lesssim f^{\sharp}(x, t) + \fint_{8\Delta} f^{\sharp},$$

and hence averaging over any time-only ball Δ' that contains (x, t) we get that

$$r^{-1} \left| \fint_{\Delta'} f(x, t) - \fint_{\Delta} f \right| \lesssim \fint_{\Delta'} f^{\sharp}(x, t) + \fint_{8\Delta} f^{\sharp},$$

and after taking sup over all such balls of size $t < 16r^2$ we get that

$$r^{-1} M^{t < 16r^2} (|f - \fint_{\Delta} f|)(x, t) \lesssim M^{t < 16r^2} (f^{\sharp})(x, t) + M^{t < 8r} (f^{\sharp})(x, t) =: h(x, t).$$

We take this as the definition of the function h . Clearly, since M^t and M are L^p bounded for $p > 1$ we have that $\|h\|_{L^p(\partial\Omega)} \lesssim \|f\|_{\dot{L}^p_{1,1/2}(\partial\Omega)}$.

Combing this with (2.48) and (2.50) we obtain for $(x, t) \in 8\Delta$

$$(2.52) \quad |D_t^{1/2}((f - \fint_{\Delta} f)\varphi)(x, t) - D_t^{1/2}f(x, t)\varphi(x, t)| \lesssim h(x, t),$$

thus proving (2.45). After we integrate the inequality above over 8Δ we obtain:

$$\int_{8\Delta} [D_t^{1/2}((f - \fint_{\Delta} f)\varphi)]^p dx dt \lesssim \int_{8\Delta} [|D_t^{1/2}f|^p + h^p] dx dt.$$

Next we estimate the same integral on $\mathbb{R}^n \setminus 8\Delta$. We only need to consider integrating over (x, t) for which there is $s \in \mathbb{R}$ such that $(x, s) \in 8\Delta$. Let us call such $s = s_0$. Let us call B the set of such points x . It follows that

$$\int_{\mathbb{R}^n \setminus 8\Delta} [D_t^{1/2}((f - \fint_{\Delta} f)\varphi)]^p dx dt = \int_{(B \times \mathbb{R}) \setminus 8\Delta} [D_t^{1/2}((f - \fint_{\Delta} f)\varphi)]^p dx dt.$$

Recalling (2.49) and using the estimate in (2.48) we see that after further splitting the integral above into regions such that the distance of points (x, t) to 4Δ is approximately $2^i r^2$ we have that

$$\begin{aligned} & \int_{\mathbb{R}^n \setminus 8\Delta} [D_t^{1/2}((f - \fint_{\Delta} f)\varphi)]^p dx dt \\ & \lesssim \sum_{i=2}^{\infty} \int_{(B \times \{t \in \mathbb{R}: \text{dist}((x, t), 4\Delta) \approx 2^i r^2\})} (2^{-3ip/2}) r^{-p} [M^{t < 4r^2} (|f - \fint_{\Delta} f|)]^p(x, s_0) dx dt \\ & \lesssim \sum_{i=2}^{\infty} 2^{i(1-3p/2)} \int_{32\Delta} h^p dx dt, \end{aligned}$$

which shows that $D_t^{1/2}$ of our function belongs to L^p . For the spatial ‘‘gradient’’ the argument is easier. With h as above we have that

$$\begin{aligned} & |(f - \fint_{\Delta} f)\varphi(x, t) - (f - \fint_{\Delta} f)\varphi(y, t)| \leq |x - y| |f(x, t) - f(y, t)| |\varphi(x, t)| \\ & + |(f(y, t) - \fint_{\Delta} f)| |\varphi(x, t) - \varphi(y, t)| \lesssim |x - y| [f^{\sharp}(x, t) + f^{\sharp}(y, t) + h(y, t)], \end{aligned}$$

where we have used the earlier estimate for $r^{-1} M^{t < 16r^2} (|f - \fint_{\Delta} f|)(y, t)$. This is a good estimate if both $(x, t), (y, t) \in 8\Delta$. If one of the points is not inside 8Δ but $|x - y| \leq r$ we trivially have

$$|(f - \fint_{\Delta} f)\varphi(x, t) - (f - \fint_{\Delta} f)\varphi(y, t)| = 0.$$

and finally if $(x, t) \in 8\Delta$, $(y, t) \notin 8\Delta$ but $|x - y| > r$ we have a trivial estimate

$$|(f - f_\Delta f)\varphi(x, t)| \leq |x - y|r^{-1}M_{t < 16r^2}(|f - f_\Delta f|)(x, t) \lesssim |x - y|h(x, t),$$

which shows that for $G(x, t) = h(x, t)\chi_{8\Delta}$ it always holds that

$$|(f - f_\Delta f)\varphi(x, t) - (f - f_\Delta f)\varphi(y, t)| \lesssim |x - y|[G(x, t) + G(y, x)].$$

Hence G is the Hajlasz–Sobolev spatial gradient of the function $(f - f_\Delta f)\varphi$ and furthermore $\|G\|_{L^p(\partial\Omega)} \lesssim \|h\|_{L^p(8\Delta)} \lesssim \|f\|_{L^p_{1,1/2}(\partial\Omega)}$ as claimed. \square

3. PARABOLIC RESULTS

Below, we will list some known properties of solutions to $Lu = 0$ in Ω . We would like to highlight [Lemma 3.6](#) and [Lemma 3.17](#) that will be key in proving [Theorem 4.1](#). Results here that involve boundary have all been established in the setting we assume, that is $\Omega = \mathcal{O} \times \mathbb{R}$ and hence $\partial\Omega = \partial\mathcal{O} \times \mathbb{R}$ or on even more general boundaries (such as 1-sided NTA domains with time-symmetric ADR boundary c.f. [\[22, 23\]](#)).

Proposition 3.1 (Boundary Poincaré inequality [\[41, 42\]](#)). *Let $\Delta = \Delta_r(P_0)$ be a boundary ball of $\partial\Omega$ and suppose that $u = 0$ on 5Δ . Then*

$$(3.2) \quad \int_{T(\Delta)} \left| \frac{u}{\delta} \right|^2 \lesssim \int_{T(5\Delta)} |\nabla u|^2.$$

Proposition 3.3 (Parabolic Cacciopoli, Lemma 3.3 of Chapter I in [\[26\]](#)). *For a weak solution u in $Q(X, t, 4r) \subset \Omega$ we have*

$$\begin{aligned} r^n \left(\max_{Q(X, t, r/2)} |u| \right)^2 &\lesssim \sup_{s \in (t-r^2, t+r^2)} \int_{Q_X(X, r) \times \{s\}} |u|^2 dZ d\tau \\ &\lesssim \int_{Q(X, t, r)} |\nabla u|^2 dZ d\tau \\ &\lesssim \frac{1}{r^2} \int_{Q(X, t, 2r)} |u|^2 dZ d\tau. \end{aligned}$$

Proposition 3.4 (Boundary Cacciopoli [\[41\]](#)). *Let u be a solution in $T(\Delta(P, s, 2r))$ that vanishes in $\Delta(P, s, 2r)$. Then*

$$\int_{T(\Delta(P, s, r))} |\nabla u|^2 \lesssim \frac{1}{r^2} \int_{T(\Delta(P, s, 2r))} |u|^2.$$

Proposition 3.5. *There exists $p > 2$ such that if Q is a parabolic cube with $2Q \subset \Omega$ then*

$$\left(\int_Q |\nabla u|^p \right)^{1/p} \lesssim \left(\int_{2Q} |\nabla u|^2 \right)^{1/2}$$

Moreover, if $u = 0$ on 4Δ we can replace Q with $T(\Delta)$ in the above inequality.

From this we have the following:

Lemma 3.6. *Let $u = 0$ on 4Δ for some boundary ball Δ . Then*

$$(3.7) \quad \left(\int_{T(\Delta)} |\nabla u|^2 \right)^{1/2} \lesssim \int_{T(2\Delta)} |\nabla u|.$$

This follows from [Proposition 3.5](#) by a standard real-variable argument (c.f. [38], for example).

Proposition 3.8 (Harnack inequality, Lemma 3.5 of Chapter I in [26]). *Let $(Z, \tau), (Y, s) \in Q(X, t, r)$ and u a non-negative weak solution on $Q(X, t, 2r)$. Then for $\tau < s$*

$$u(Z, \tau) \leq u(Y, s) \exp \left[c \left(\frac{|Y - Z|^2}{|s - \tau|} + 1 \right) \right].$$

Proposition 3.9 (Boundary Hölder continuity [22, 41]). *Let u be a solution on $T(\Delta(x, t, 2r))$ that vanishes on continuously on $\Delta(x, t, 2r)$. Then there exists $\alpha > 0$ such that for $(Y, s) \in T(\Delta(x, t, r/2))$*

$$u(Y, s) \lesssim c \left(\frac{\delta(Y, s)}{r} \right)^\alpha \max_{T(\Delta(x, t, r))} |u|.$$

If additionally $u \geq 0$ in $T(\Delta(x, t, 2r))$, then

$$(3.10) \quad u(Y, s) \lesssim \left(\frac{\delta(Y, s)}{r} \right)^\alpha u(V^+(\Delta(x, t, 2r))).$$

By [41] we also have the existence of a Greens function for the operator L and its adjoint L^* with

$$G^*(X, t, Y, s) = G(Y, s, X, t).$$

Moreover, let $\omega(X, t, \Delta)$ denotes the parabolic measure of $\Delta \subset \partial\Omega$ with the pole at $(X, t) \in \Omega$ (and ω^* denotes the parabolic measure of L^*). We have the following comparison result

Proposition 3.11 ([23]). *Let $(X, t), (Y, s) \in \Omega$ with $s + 8r^2 \leq t$, where $r := \delta(Y, s)$. Let Y^* be a point on $\partial\mathcal{O}$ such that $Y \in \tilde{\gamma}(Y^*)$. Then*

$$r^n G(X, t, Y, s) \sim \omega(X, t, \Delta(Y^*, s, r))$$

Similarly, if $s - 8r^2 \geq t$, then

$$r^n G^*(X, t, Y, s) \sim \omega^*(X, t, \Delta(Y^*, s, r))$$

As a consequence, for $d \geq 10r$ we have

$$(3.12) \quad r^n G(V^-(\Delta_r), V^-(\Delta_d)) = r^n G^*(V^-(\Delta_d), V^-(\Delta_r)) \sim \omega^{*V^-(\Delta_d)}(\Delta_r).$$

Furthermore, we have the following decay property for the Green's function:

Lemma 3.13. *There exists $\alpha = \alpha(\lambda, n)$ such that for $(X, t), (Y, s) \in \Omega$ with $t < s$*

$$(3.14) \quad G(X, t, Y, s) \lesssim \delta(X, t)^\alpha (|X - Y|^2 + (s - t))^{-\frac{n+\alpha}{2}}.$$

The proof stems from private communications with Dindoš, Li, and Pipher and can be found as Lemma 2.8.17 in Ulmer thesis ([42]).

Proposition 3.15 (Local comparison Principle [41]). *Let u and v be non-negative solutions to $Lu = 0$ in $T(\Delta_{2r}(Y, s))$ with $u = v = 0$ on $\Delta_{2r}(Y, s)$*

$$\frac{u(X, t)}{v(X, t)} \approx \frac{u(V^+(\Delta_r(Y, s)))}{v(V^-(\Delta_r(Y, s)))}, \quad (X, t) \in T(\Delta_r(Y, s)),$$

and V^+ (V^-) are time forward (backward) corkscrew point relative to the surface ball.

Proposition 3.16 (Lemma 2.15 in [31]). *For solutions to $Lu = 0$ and with $u = 0$ on Δ_{5r}*

$$\sup_{T(\Delta_{2r})} |u| \lesssim \left(\int_{T(\Delta_{4r})} |u|^2 \right)^{1/2}. \quad \square$$

This implies the following:

Lemma 3.17. *Let u be a weak solution such that $u = 0$ on $\Delta_{5r} \subset \Delta_d$, where $d = \text{diam}(\mathcal{O})$. Then for $(X, t) \in T(\Delta_{2r})$*

$$|u(X, t)| \lesssim \frac{G(X, t, V^-(\Delta_d))}{G(V^-(\Delta_r), V^-(\Delta_d))} \left(\int_{T(\Delta_{4r})} |u|^2 \right)^{1/2}.$$

Proof. Let $(X, t) \in T(\Delta_{2r})$ and set $u^+ := \max(u, 0)$ $u^- := \min(-u, 0)$. Next write $u = u_1 - u_2$, where u_1 and u_2 solve

$$\begin{cases} Lu_1 = 0, & \text{in } T(2\Delta), \\ u_1 = u^+, & \text{on } \partial_p T(2\Delta), \end{cases} \quad \begin{cases} Lu_2 = 0, & \text{in } T(2\Delta), \\ u_2 = u^-, & \text{on } \partial_p T(2\Delta), \end{cases}$$

respectively. Here $\partial_p T(2\Delta)$ denotes the parabolic part of the boundary of $T(2\Delta)$. Let $j \in \{1, 2\}$. By [Proposition 3.15](#)

$$u_j(X, t) \lesssim \frac{G(X, t; V^-(\Delta_d))}{G(V^-(\Delta_r); V^-(\Delta_d))} u_j(V^+(\Delta_r)).$$

Next note that since $V^+(\Delta_r) \in T(2\Delta)$ we can apply the maximum principle (see e.g. [1]) followed by [Proposition 3.16](#) to see that

$$u_j(V^+(\Delta_r)) \leq \sup_{\partial T(2\Delta)} |u| \lesssim \left(\int_{T(\Delta_{4r})} |u|^2 \right)^{1/2}.$$

Thus

$$|u(X, t)| \leq u_1(X, t) + u_2(X, t) \lesssim \left(\int_{T(\Delta_{4r})} |u|^2 \right)^{1/2},$$

as desired. □

4. REVERSE HÖLDER INEQUALITY

In this section we establish the Reverse Hölder inequality of Shen [37] in the parabolic case.

Theorem 4.1. *Let $1 \leq p < \infty$ and suppose that $(D_{L^*})_{p'}$ is solvable and suppose that u is a weak solution of $Lu = 0$ with $u = 0$ on $\tilde{\Delta} =: 80\Delta$ where $\Delta = \Delta_r(P_0, \tau_0)$, with $r \lesssim \text{diam}(\mathcal{O})$. Then*

$$\left(\int_{\Delta} N[\nabla u]^p \right)^{1/p} \lesssim \int_{\tilde{\Delta}} N[\nabla u],$$

as well as

$$(4.2) \quad \left(\int_{\Delta} N[\nabla u]^p \right)^{1/p} \lesssim \int_{T(\tilde{\Delta})} |\nabla u|.$$

Proof: Let $(X, t) \in \Gamma(P, \tau)$. If $\delta(X, t) \geq r$, then

$$\tilde{S}(X, t) := \{(P, s) \in \partial\Omega : (X, t) \in \Gamma(P, s)\} = \tilde{S}(X) \times (t - \delta(X)^2, t + \delta(X)^2),$$

i.e., $\tilde{S}(X, t)$, $\tilde{S}(X)$ are backward in space-time (space) cones originating from the point (X, t) defined earlier in Section 2. It follows from the definition of \tilde{S} that $|\tilde{S}(X, t)| \geq 4|\tilde{S}(X)|r^2 \gtrsim r^{n+1}$, and thus

$$\text{Av}^2[\nabla u](X, t) \leq \inf_{\tilde{S}(X, t)} N[u] \leq \int_{\tilde{S}(X, t)} N[u] \lesssim \int_{80\Delta} N[u].$$

Thus we can disregard the points whose distance to the boundary is r or larger and only consider the inequality for truncated non-tangential maximal function at the height r :

$$\left(\int_{\Delta} N^r[\nabla u]^p \right)^{1/p} \lesssim \int_{\tilde{\Delta}} N[\nabla u].$$

By [Proposition 3.3](#) the L^2 average of ∇u over $C_{\delta(X)/4}(X, t)$ can be controlled from above by the L^2 average of $|u/\delta|$ over $C_{\delta(X)/2}(X, t)$ which is further controlled by just the supremum of $|u/\delta|$ over $C_{\delta(X)/2}(X, t)$. It follows that

$$N^r[\nabla u](P, \tau) \lesssim N_{\text{sup}}^r[u/\delta](P, \tau),$$

where

$$N_{\text{sup}}^r[v](P, \tau) = \sup_{\Gamma_{\tilde{a}}^{2r}(P, \tau)} |v|,$$

for some sufficiently large $\tilde{a} > 1$.

Let $d \geq 10r$. By [Lemma 3.17](#)

$$N_{\text{sup}}^r[u/\delta](P, \tau) \lesssim \frac{1}{G(V^-(\Delta), V^-(\Delta_d))} \left(\int_{T(\Delta_{4r})} |u|^2 \right)^{1/2} N_{\text{sup}}^r \left[\frac{G(\cdot, V^-(\Delta_d))}{\delta(\cdot)} \right],$$

and therefore

$$\left(\int_{\Delta} N^r[\nabla u]^p \right)^{1/p} \lesssim \left(\int_{T(\Delta_{4r})} |u|^2 \right)^{1/2} \frac{1}{G(V^-(\Delta), V^-(\Delta_d))} \left(\int_{\Delta} N_{\text{sup}}^r [G(\cdot, V^-(\Delta_d))/\delta(\cdot)]^p \right)^{1/p}.$$

Let $\omega^* = \omega^{*V^-(\Delta_d)}$ be the parabolic measure of the adjoint operator. Observe that by [\(3.12\)](#)

$$\frac{1}{G(V^-(\Delta), V^-(\Delta_d))} \sim \frac{r^n}{\omega^*(\Delta)},$$

It follows that for $\rho := \delta(X, t)$ we have

$$\frac{G(X, t, V^-(\Delta_d))}{\delta(X, t)} \sim \frac{\omega^*(\Delta_\rho(P, \tau + 100\rho))}{\rho^{n+1}} \lesssim_n \frac{\omega^*(\Delta_{11\rho}(P, \tau))}{\rho^{n+1}}.$$

Thus

$$N_{\text{sup}}^r[G(\cdot, V^-(\Delta_d))/\delta(\cdot)] \leq M^r \left[\frac{d\omega^*}{d\sigma} \right] (P, \tau),$$

where as before M^r denotes the usual truncated maximal function which is bounded on L^p for $p > 1$. It follows that

$$\left(\int_{\Delta} N_{\text{sup}}^r [G(\cdot, V^-(\Delta_d))/\delta(\cdot)]^p \right)^{1/p} \lesssim \left(\int_{4\Delta} \left| \frac{d\omega^*}{d\sigma} \right|^p \right)^{1/p} \lesssim r^{-n-1} \omega^*(\Delta),$$

where in the final inequality we have used the fact that $\omega^* \in B_p(d\sigma)$, a fact that follows from the solvability of $(D^*)_{p'}$ and that the measure ω^* is doubling. Hence

$$\begin{aligned} \left(\int_{\Delta} N^r[\nabla u]^p \right)^{1/p} &\lesssim \frac{r^n}{\omega^*(\Delta)} r^{-n-1} \omega^*(\Delta) \left(\int_{T(\Delta_{4r})} |u|^2 \right)^{1/2} = \left(\int_{T(\Delta_{4r})} |u/r|^2 \right)^{1/2} \\ &\lesssim \left(\int_{T(\Delta_{4r})} |u/\delta|^2 \right)^{1/2} \lesssim \left(\int_{T(\Delta_{20r})} |\nabla u|^2 \right)^{1/2} \lesssim \int_{T(\Delta_{40r})} |\nabla u|, \end{aligned}$$

where in the last two inequalities we have used [Proposition 3.1](#) and [Proposition 3.6](#). This gives us (4.2). It remains to show that

$$(4.3) \quad \int_{T(\Delta_{40r})} |\nabla u| \lesssim r \int_{\Delta_{40r}} N[\nabla u],$$

which is a rather straightforward calculation using Fubini's theorem and Whitney decomposition of the set $T(\Delta_{40r})$. We omit the details. \square

5. PROOF OF [Theorem 1.3](#)

5.1. Unbounded case. For this section we assume that $\text{diam}(\mathcal{O}) = \infty$. What this means is that for any boundary point $P \in \partial\mathcal{O}$ there exists interior corkscrew balls of arbitrary large size as the domain is unbounded.

5.1.1. Almost good lambda argument. The bulk of the proof of [Theorem 1.3](#) is proving the ‘‘almost λ -inequality’’ [Lemma 5.1](#). Let $p > q$, and set

$$E(\lambda) := \{M[N[\nabla u]^q] > \lambda\} \quad \text{and} \quad A = A(\varepsilon) := 1/(2\varepsilon)^{q/p}.$$

Let $g \in L^p(\partial\Omega)$ be a Hajlasz upper spatial gradient for f and let h be like in [Proposition 2.44](#). We have the following inequality:

Lemma 5.1. *Let $1 < q < p < \infty$. Suppose that $(R_L)_q$ and $(D_{L^*})_{p'}$ are solvable. There exists ε, γ depending only on p, q, n, L, Ω such that for all $\lambda > 0$*

$$(5.2) \quad |E(A\lambda)| \leq \varepsilon |E(\lambda)| + |\{M[|g|^q + |D_{1/2}^t f|^q + |h|^q] > \gamma\lambda\}|,$$

Proof: Let $\varepsilon \in (0, 1)$ be a small constant to be determined. By [\[10\]](#) there exists a family of ‘‘dyadic cubes’’ such that:

- (1) If $l \geq k$ then either $Q_\beta^l \subset Q_\alpha^k$ or $Q_\beta^l \cap Q_\alpha^k = \emptyset$,
- (2) For each (k, α) and $l < k$ there is a unique β so that $Q_\alpha^k \subset Q_\beta^l$,
- (3) Each Q_α^k contains a ball Δ_α^k , with $r(\Delta) = 2^{-k}$
- (4) There exists a constant $C_0 > 0$ such that $2^{-k} \leq \text{diam}(Q_\alpha^k) \leq C_0 2^{-k}$,

where the diameter is taken in the parabolic metric. The last listed property implies together with the Ahlfors regularity of the surface measure that $\sigma(Q_\alpha^k) \approx 2^{-k(n+1)}$.

Let $E(\lambda) = \bigcup_k Q_k$ be a Whitney decomposition of $E(\lambda)$ into a disjoint collection of maximal such ‘‘dyadic subcubes’’. In this place we use unboundedness of our domain, as on a bounded domain for $\text{diam}(Q_k) \gg \text{diam}(\mathcal{O})$ the cubes become ‘‘misshaped’’

(with long side in t much larger than the size in spatial variables). We claim that we can choose ε, γ, C_0 such that if Q_k is a cube with the property

$$(5.3) \quad Q_k \cap \{M[|D_{1/2}^t f|^q + |g|^q + |h|^q] \leq \gamma\lambda\} \neq \emptyset,$$

then

$$(5.4) \quad |E(A\lambda) \cap Q_k| \leq \varepsilon|Q_k|.$$

The desired estimate (5.2) then follows by summation in k . Let us now fix a cube Q_k and drop the index, writing $Q = Q_k, \Delta = \Delta_k$ and letting $r = r(\Delta)$. To prove the desired implication, first observe that

$$(5.5) \quad M[|N[\nabla u]|^q](P, \tau) \leq \max\{M^{C_0 r}[N[\nabla u]^q](P, \tau), C_1 \lambda\}$$

for any $(P, \tau) \in Q$ and with $C_1 = C_1(\mathcal{O})$. This is because Q is maximal and so $2C_0\Delta \not\subset E(\lambda)$. Assume that $A \geq C_1$. It follows that

$$(5.6) \quad |Q \cap E(A\lambda)| \leq |\{(P, \tau) \in Q : M^{C_0 r}[N[\nabla u]^q](P, \tau) > A\lambda\}|.$$

Next let $\varphi = \varphi_Q$ be a smooth cut-off function on \mathbb{R}^{n+1} such that

$$\varphi = 1 \text{ on } C_2 Q, \quad \varphi = 0, \text{ on } \partial\Omega \setminus 2C_2 Q, \quad |\nabla\varphi| \lesssim \ell(Q)^{-1},$$

where $C_2 \geq 160C_0$. Define

$$f_Q := f - \alpha_Q, \quad \alpha_Q := \int_{80Q} f$$

Now let $v = v_Q$ be the unique weak solution of

$$\begin{cases} Lv = 0, & \text{in } \Omega, \\ v = \varphi f_Q, & \text{on } \partial\Omega, \end{cases}$$

Let $\bar{p} > p$. In view of (5.6), we have

$$\begin{aligned} |Q \cap E(A\lambda)| &\leq |\{P \in Q : M^{C_0 r}[N[\nabla(u-v)]^q](P, \tau) > \frac{A\lambda}{2^q}\}| \\ &\quad + |\{P \in Q : M^{C_0 r}[N[\nabla v]^q](P, \tau) > \frac{A\lambda}{2^q}\}| \\ &\lesssim (A\lambda)^{-\bar{p}/q} \int_{2C_0\Delta} N[\nabla(u-v)]^{\bar{p}} + (A\lambda)^{-1} \int_{2C_0\Delta} N[\nabla v]^q =: J_1 + J_2, \end{aligned}$$

where we have used the weak $(\frac{\bar{p}}{q}, \frac{\bar{p}}{q})$ and $(1, 1)$ estimates for the maximal function.

We start with the term J_2 . Since $(R_L)_q$ is solvable, and applying [Proposition 2.44](#), we have that for some large $C_3 \gg 1$ we have that

$$\begin{aligned} \|N[\nabla v]\|_{L^q(\partial\Omega)} &\lesssim \|\varphi f_Q\|_{L^q_{1,1/2}(\partial\Omega)} \lesssim \|f\|_{L^q_{1,1/2}(C_3\Delta)} + \|h\|_{L^q(C_3\Delta)} \\ &\lesssim \int_{C_3\Delta} |D_t^{1/2} f|^q + |g|^q + |h|^q \lesssim \gamma\lambda|Q| \end{aligned}$$

and hence

$$(5.7) \quad J_2 \lesssim A^{-1}\gamma|Q| \lesssim \gamma\varepsilon^{q/p}|Q|.$$

Now we deal with the first term J_1 . Observe that $(u - \alpha) - v$ is a weak solution whose boundary data $f_Q(1 - \varphi)$ vanishes on $C_2\Delta$. Also note that the solvability of $(D_{L^*})_{p'}$ implies that $(D_{L^*})_{\bar{p}'}$ is solvable for some $\bar{p} > p$ due to properties of the B_p weights. It then follows by [Theorem 4.1](#) that

$$\begin{aligned} \int_{2C_0\Delta} N[\nabla(u - v)]^{\bar{p}} &= |2C_0\Delta| \int_{2Q} N[\nabla(u - v)]^{\bar{p}} \lesssim |Q| \left(\int_{C_2\Delta} N[\nabla(u - v)] \right)^{\bar{p}} \\ &\lesssim |Q| \left[\left(\int_{C_2\Delta} N[\nabla u]^q \right)^{\bar{p}/q} + \left(\int_{C_2\Delta} N[\nabla v]^q \right)^{\bar{p}/q} \right]. \end{aligned}$$

Now, we already know that $\left(\int_{C_2\Delta} \tilde{N}[\nabla v]^q \right)^{1/q} \lesssim \lambda\gamma$ and hence it remains to consider the first term. Note that since $C_2\Delta \not\subset E(\lambda)$ there exists a point $(P', \tau') \in C_2\Delta$ such that $M[N[\nabla u]^q](P', \tau') \leq \lambda$. Let Δ' be a ball obtained by translating Δ to be centered at (P', τ') , then

$$\int_{C_2\Delta} N[\nabla u]^q \lesssim \int_{2C_2\Delta'} N[\nabla u]^q \leq M[N[\nabla u]^q](P', \tau') \leq \lambda.$$

Thus

$$(5.8) \quad J_1 \lesssim \frac{1}{(A\lambda)^{\bar{p}/q}} \lambda^{\bar{p}/q} |Q| = A^{-\bar{p}/q} |Q| = \varepsilon^{\bar{p}/p} |Q| = \varepsilon |Q| \varepsilon^{\bar{p}/p-1}$$

and combining this with [\(5.7\)](#) we get

$$|Q \cap E(A\lambda)| \lesssim J_1 + J_2 \lesssim \varepsilon |Q| (\varepsilon^{\bar{p}/p-1} + \gamma \varepsilon^{-q/p-1}).$$

Hence, first choosing ε small enough that $\varepsilon^{\bar{p}/p-1} \leq 1/2$ and then γ small enough that $\gamma \varepsilon^{q/p-1} \leq 1/2$ we indeed have that

$$|Q \cap E(A\lambda)| \leq \varepsilon |Q|,$$

thus completing our the proof. \square

We are now ready to finish the proof of [Theorem 1.3](#) in the unbounded case, i.e. when $\text{diam}(\mathcal{O}) = \infty$. Like in [\[37\]](#) we leverage [Lemma 5.1](#) for a change of variables to see that

$$\int_{\partial\Omega} N[\nabla u]^p \leq \int_{\partial\Omega} M[N[\nabla u]^q]^{p/q} \sim \lim_{K \rightarrow \infty} \int_0^K |E(\lambda)| \lambda^{p/q-1} d\lambda.$$

For a fixed K the integral on the righthand side is finite as $\lambda^{p/q-1} \leq K^{p/q-1} < \infty$. Thus for a fixed K we have

$$\begin{aligned} \int_0^K |E(\lambda)| \lambda^{p/q-1} d\lambda &\leq \frac{1}{2\varepsilon} \int_0^{K/A} |E(A\lambda)| \lambda^{p/q-1} d\lambda \\ &\leq \frac{1}{2} \int_0^{K/A} |E(\lambda)| \lambda^{p/q-1} d\lambda + \frac{1}{2\varepsilon} \int_0^{K/A} |\{M[|D_{1/2}^t f|^q + |g|^q + |h|^q] > \gamma\lambda\}| \lambda^{p/q-1} d\lambda \\ &\lesssim_\varepsilon \int_0^{K/A} |\{M[|D_{1/2}^t f|^q + |g|^q + |h|^q] > \gamma\lambda\}| \lambda^{p/q-1} d\lambda, \end{aligned}$$

where in the last step since $K/A \leq K$ we have moved the term $\frac{1}{2} \int_0^{K/A} |E(\lambda)| \lambda^{p/q-1} d\lambda$ to the lefthand side. We take limit $K \rightarrow \infty$. Recall, that h is as in [Proposition 2.44](#) and $g \in L^p(\partial\Omega)$ is a Hajlasz upper spatial gradient for f . Thus $\|h\|_{L^p(\partial\Omega)} \lesssim \|D_t^{1/2} f\|_{L^p(\partial\Omega)}$ and $\|g\|_{L^p(\partial\Omega)} \lesssim \|f\|_{\dot{L}_1^p(\partial\Omega)}$ therefore we have that

$$\begin{aligned} \int_{\partial\Omega} N[\nabla u]^p &\lesssim \int_0^\infty |\{M[|D_{1/2}^t f|^q + |g|^q + |h|^q] > \gamma\lambda\}| \lambda^{p/q-1} d\lambda \\ &\lesssim_\gamma \int_{\partial\Omega} M[|D_{1/2}^t f|^q + |g|^q + |h|^q]^{p/q} \\ &\lesssim \int_{\partial\Omega} |D_{1/2}^t f|^p + |g|^p + |h|^p \lesssim \|f\|_{\dot{L}_{1,1/2}^p(\partial\Omega)}^p, \end{aligned}$$

completing the proof. \square

5.2. Bounded case. Now suppose that $\text{diam}(\mathcal{O}) < \infty$, by rescaling we may without loss of generality assume that $\text{diam}(\mathcal{O}) = 1$. The first problem we run into is that the reverse Holder inequality from [Theorem 4.1](#) does not work for boundary cubes Δ with $\text{diam}(\Delta) \gg 1$. For this reason we can prove an analogue of [Lemma 5.1](#) but only for local parabolic cubes of a finite size ~ 1 . Subsequently, we can only obtain local bounds for the L^p norm of $N[\nabla u]$ restricted to boundary parabolic cubes of size ~ 1 . To overcome this we will split our domain into subdomains of finite diameter (in space and time).

Denote by $\Omega_i := \mathcal{O} \times (i, i+1)$ and let $\Delta_i = \partial_p \Omega_i = \partial\mathcal{O} \times (i, i+1)$ for $i \in \mathbb{Z}$. There exists a nonnegative C^1 function $\psi \in C_c^1(\mathbb{R})$, $\text{supp } \psi \subset [-1, 2]$ such that

$$\psi_i(\cdot) = \psi(\cdot + i), \quad |\psi'| \lesssim 1, \quad \int_{\mathbb{R}} \psi = 1, \quad \sum_{i \in \mathbb{Z}} \psi_i \equiv 1.$$

Clearly, $\|D_{1/2}^t \psi\|_{L^r(\mathbb{R})} \lesssim 1$, for each $1 < r < \infty$. Set

$$f_{\Delta_i} := \int_{\Delta_i} f.$$

Finally, let $\Psi_k := \sum_{i \geq k} \psi_i$ and note that $\psi_i = \Psi_{i+1} - \Psi_i$ and $\Psi_k \equiv 1$ for on $[i+2, \infty)$.

Let w_i solve the PDE

$$\begin{cases} Lw_i = 0, & \text{in } \Omega, \\ w_i = (f - f_{\Delta_i})\psi_i & \text{on } \partial\Omega, \end{cases}$$

If $w := \sum_{i \in \mathbb{Z}} w_i$ then $v := u - w$ solves $Lv = 0$ with boundary data

$$\sum_{i \in \mathbb{Z}} f_{\Delta_i} \psi_i = \sum_{i \in \mathbb{Z}} f_{\Delta_i} (\Psi_{i+1} - \Psi_i) = \sum_{i \in \mathbb{Z}} (f_{\Delta_{i-1}} - f_{\Delta_i}) \Psi_i.$$

Thus we can write $v = \sum_{i \in \mathbb{Z}} v_i$ as where v_i solves

$$\begin{cases} Lv_i = 0, & \text{in } \Omega, \\ v_i = (f_{\Delta_{i-1}} - f_{\Delta_i}) \Psi_i & \text{on } \partial\Omega. \end{cases}$$

It follows that we have decomposed the solution u to $Lu = 0$ with $u|_{\partial\Omega} = f$ into

$$u = \sum_i v_i + \sum_i w_i,$$

such that $w_i|_{\partial\Omega}$ is supported only on $\partial\mathcal{O} \times (i-1, i+2)$ and $v_i|_{\partial\Omega} = 0$ for $t < i-1$ and $v_i|_{\partial\Omega}$ is constant for $t > i+2$.

Let us also define \tilde{v}_i , such that \tilde{v}_i solves

$$\begin{cases} L\tilde{v}_i = 0, & \text{in } \Omega, \\ \tilde{v}_i = (f_{\Delta_{i-1}} - f_{\Delta_i})\Psi_i(1 - \Psi_{i+20}) & \text{on } \partial\Omega. \end{cases}$$

Note that by considering the solution \tilde{v}_i instead of v_i , since $v_i|_{\partial\Omega}(P, \tau) = \tilde{v}_i|_{\partial\Omega}(P, \tau)$ for $\tau \leq i+15$ we have $N[\nabla v_i](P, \tau) = N[\nabla \tilde{v}_i](P, \tau)$ for all $\tau \leq i+10$. However, \tilde{v}_i is more similar to w_i as both are now compactly supported and half derivative of $\Psi_i(1 - \Psi_{i+20})$ has faster decay than half derivative of Ψ_i .

Let us study the properties of $N[\nabla v_i]$ and $N[\nabla w_i]$. Without loss of generality, let v be either v_i or w_i for some $i \in \mathbb{Z}$. It follows that $v \equiv 0$ for $t \leq i-1$. Thus $N[\nabla v](P, \tau) \equiv 0$ for all $\tau \leq i-5$.

When $i-5 < \tau < i+10$ we may study $N[\nabla v](P, \tau)$ by considering the $(R_L)_p$ boundary value problem. As we have already observed above, [Lemma 5.1](#) only holds on scales comparable to the diameter of \mathcal{O} . Thus by modifying the calculation for the unbounded case we can obtain an estimate on a bounded (in time) portion of $\partial\Omega$, namely that

$$\int_{\cup_{k=i-5}^{i+9} \Delta_k} N[\nabla v]^p \lesssim \|v\|_{L^p_{1,1/2}(\partial\Omega)}^p.$$

This calculation differs only slightly to the one we gave in the unbounded case, and is actually more similar to what Shen did in [\[37\]](#) as he only considered the case of a bounded domain. Hence we obtain for w_i and v_i (here using \tilde{v}_i as on this portion of the domain they coincide):

$$(5.9) \quad \int_{\cup_{k=i-5}^{i+9} \Delta_k} N[\nabla w_i]^p \lesssim \|(f - f_{\Delta_i})\psi_i\|_{L^p_{1,1/2}(\partial\Omega)}^p, \quad \int_{\cup_{k=i-5}^{i+9} \Delta_k} N[\nabla v_i]^p \lesssim |f_{\Delta_{i-1}} - f_{\Delta_i}|^p.$$

Here, the second term is already simplified, as we use quietly the fact that $\Psi_i(1 - \Psi_{i+20})$ has bounded half time-derivative on $L^p(\partial\Omega)$ for all $p > 1$.

Finally, for $\tau \geq i+10$ we will use the fact that ∇v decays exponentially.

Lemma 5.10. *Let $p \geq 1$ and v be a solution to $Lv = 0$ and suppose that $v = \text{const.}$ on $\partial\mathcal{O} \times (i+2, \infty)$. There exists an $\alpha > 0$ such that for all $j > i+9$ we have that*

$$\|N[\nabla v]\|_{L^p(\Delta_j)} \lesssim e^{-\alpha|i-j|} \|N[\nabla v]\|_{L^p(\Delta_{i+3})}.$$

Hence by [\(5.9\)](#) we have for all $j > i+9$:

$$(5.11) \quad \int_{\Delta_j} N[\nabla w_i]^p \lesssim e^{-\alpha p|i-j|} \|(f - f_{\Delta_i})\psi_i\|_{L^p_{1,1/2}(\partial\Omega)}^p, \quad \int_{\Delta_j} N[\nabla v_i]^p \lesssim e^{-\alpha p|i-j|} |f_{\Delta_{i-1}} - f_{\Delta_i}|^p.$$

In order to establish [Lemma 5.10](#) we start by proving the L^2 decay. Assume first that the coefficients of A as smooth.

Proposition 5.12. *Let u be such that $Lu = 0$ on $\partial\mathcal{O} \times [0, \infty)$ and also Moreover suppose that $u = 0$ on $\partial\mathcal{O} \times [0, \infty)$. Then there exists an $\beta > 0$ such that for $t > 0$*

$$(5.13) \quad \|u(\cdot, t)\|_{L^2(\mathcal{O})} \lesssim e^{-\beta t} \|u(\cdot, 0)\|_{L^2(\mathcal{O})}.$$

Proof. Set $\eta(t) := \|u(\cdot, t)\|_{L^2(\mathcal{O})}$. For $t > 0$

$$2\eta(t)\eta'(t) = \partial_t(\eta(t)^2) = 2 \int_{\Omega_t} u \partial_t u = -2 \int_{\Omega_t} A \nabla u \cdot \nabla u \leq -2\lambda \int_{\Omega_t} |\nabla u|^2,$$

where $\Omega_t = \mathcal{O} \times \{t\}$. Since $u = 0$ on $\partial\Omega_t$, applying the Sobolev inequality on Ω_t we have that

$$- \int_{\Omega_t} |\nabla u|^2 \lesssim - \int_{\Omega_t} |u|^2 = -\eta(t)^2.$$

Hence $\eta'(t) \lesssim -\eta(t)$ and we may apply Gronwall's inequality to deduce [\(5.13\)](#). \square

Observe that this calculation holds on very general domains \mathcal{O} as it only requires the Sobolev inequality. The assumption that A is smooth can be removed by approximation as the estimate we are seeking only depends on the ellipticity constant of the operator A .

Assume now that $\|N[\nabla v]\|_{L^p(\Delta_{i+3})} < \infty$ and that $v|_{\partial\Omega} \equiv 0$ for all times $\geq i + 2$. Clearly, then also

$$\|N[\nabla v]\|_{L^1(\Delta_{i+3})} \lesssim \|N[\nabla v]\|_{L^p(\Delta_{i+3})} < \infty,$$

and thus by [\(4.3\)](#)

$$\int_{\Omega_{i+3}} |\nabla v| \lesssim \|N[\nabla v]\|_{L^1(\Delta_{i+3})} \lesssim \|N[\nabla v]\|_{L^p(\Delta_{i+3})}.$$

By applying [Lemma 3.6](#) as we satisfy $v|_{\partial\Omega}(P, \tau) = 0$ for $\tau > i + 2$ we obtain that for some $\tau \in (i + 3, i + 4)$

$$\left(\int_{\mathcal{O} \times \{\tau\}} |\nabla v|^2 \right)^{1/2} \lesssim \|N[\nabla v]\|_{L^p(\Delta_{i+3})} < \infty,$$

and finally by Sobolev (given that v vanishes on the lateral boundary of Ω_{i+3}):

$$\left(\int_{\mathcal{O} \times \{\tau\}} |v|^2 \right)^{1/2} \lesssim \|N[\nabla v]\|_{L^p(\Delta_{i+3})} < \infty.$$

Fix now some $j > i + 9$. We propagate the L^2 initial data given on $\mathcal{O} \times \{\tau\}$ using [Proposition 5.12](#) to $t \in (j + 7, j + 12)$. It follows that for all such t we have

$$\|v(\cdot, t)\|_{L^2(\mathcal{O})} \lesssim e^{-\beta(j-i-3)} \|v(\cdot, \tau)\|_{L^2(\mathcal{O})} \lesssim e^{-\beta|i-j|} \|N[\nabla v]\|_{L^p(\Delta_{i+3})} < \infty.$$

By integrating over the interval $(j + 7, j + 12)$ and then applying [Proposition 3.16](#)

$$(5.14) \quad \sup_{\Omega_{j+8} \cup \Omega_{j+9} \cup \Omega_{j+10}} |v| \lesssim e^{-\beta|i-j|} \|N[\nabla v]\|_{L^p(\Delta_{i+3})} < \infty.$$

Finally, we repeat the argument we gave in the proof of [Theorem 4.1](#) where we estimated $N^r[\nabla v](P, \tau)$ by $N_{\text{sup}}^r[v/\delta](P, \tau)$ and then using the Green's function eventually we have obtained (for $r = 1$ as the diameter of the domain is 1):

$$\left(\int_{\Delta_{j+9}} N[\nabla v]^p \right)^{1/p} \lesssim \int_{\Omega \cap (j+8.5, j+10.5)} |\nabla v|.$$

The final step is to estimate the above righthand side by the lefthand side of [\(5.14\)](#) which can be done thanks to [Proposition 3.4](#) (boundary Caccioppoli) which finally gives us that

$$\left(\int_{\Delta_{j+9}} N[\nabla v]^p \right)^{1/p} \lesssim e^{-\beta|i-j|} \|N[\nabla v]\|_{L^p(\Delta_{i+3})} < \infty.$$

Form this the claim of [Lemma 5.10](#) follows by applying the above estimate to the function $v - \text{const}$. \square

We are ready to add up the estimates to obtain a bound for the L^p norm of u . Fix $i \in \mathbb{Z}$ and consider $N[\nabla u]$ on Δ_i . Since N is sub-additive, clearly

$$N[\nabla u](P, \tau) \leq \sum_j [N[\nabla v_j] + N[\nabla w_j]](P, \tau),$$

and therefore

$$\begin{aligned} \|N[\nabla u]\|_{L^p(\Delta_i)} &\leq \sum_j [\|N[\nabla v_j]\|_{L^p(\Delta_i)} + \|N[\nabla w_j]\|_{L^p(\Delta_i)}] \\ &\leq \sum_{j=i-9}^{i+5} [\|N[\nabla v_j]\|_{L^p(\Delta_i)} + \|N[\nabla w_j]\|_{L^p(\Delta_i)}] + \sum_{j<i-9} [\|N[\nabla v_j]\|_{L^p(\Delta_i)} + \|N[\nabla w_j]\|_{L^p(\Delta_i)}] \\ &\lesssim \sum_{j=i-9}^{i+5} \left[\|(f - f_{\Delta_j})\psi_j\|_{\dot{L}_{1,1/2}^p} + |f_{\Delta_{j-1}} - f_{\Delta_j}| \right] \\ &\quad + \sum_{j<i-9} e^{-\alpha|i-j|} \left[\|(f - f_{\Delta_j})\psi_j\|_{\dot{L}_{1,1/2}^p} + |f_{\Delta_{j-1}} - f_{\Delta_j}| \right], \end{aligned}$$

using [\(5.9\)](#) and [\(5.11\)](#). We raise both sides to the p -th power. It follows that

$$\begin{aligned} \|N[\nabla u]\|_{L^p(\Delta_i)}^p &\lesssim \sum_{j=i-9}^{i+5} \left[\|(f - f_{\Delta_j})\psi_j\|_{\dot{L}_{1,1/2}^p}^p + |f_{\Delta_{j-1}} - f_{\Delta_j}|^p \right] \\ &\quad + \left(\sum_{j<i-9} e^{-\alpha|i-j|} \left[\|(f - f_{\Delta_j})\psi_j\|_{\dot{L}_{1,1/2}^p} + |f_{\Delta_{j-1}} - f_{\Delta_j}| \right] \right)^p. \end{aligned}$$

For the last term we use Hölder's inequality for series. We split the exponential into two terms

$$e^{-\alpha|i-j|} = e^{-\alpha|i-j|/p'} e^{-\alpha|i-j|/p}$$

to obtain

$$\left(\sum_{j<i-9} e^{-\alpha|i-j|} \left[\|(f - f_{\Delta_j})\psi_j\|_{\dot{L}_{1,1/2}^p} + |f_{\Delta_{j-1}} - f_{\Delta_j}| \right] \right)^p \leq \left(\sum_{j<i-9} \left(e^{-\alpha|i-j|/p'} \right)^{p'} \right)^{p/p'} \times$$

$$\begin{aligned} & \sum_{j < i-9} e^{-\alpha|i-j|} \left[\|(f - f_{\Delta_j})\psi_j\|_{\dot{L}^p_{1,1/2}} + |f_{\Delta_{j-1}} - f_{\Delta_j}| \right]^p \\ & \leq C(p, \alpha) \sum_{j < i-9} e^{-\alpha|i-j|} \left[\|(f - f_{\Delta_j})\psi_j\|_{\dot{L}^p_{1,1/2}}^p + |f_{\Delta_{j-1}} - f_{\Delta_j}|^p \right]. \end{aligned}$$

Now we sum over all indices $i \in \mathbb{Z}$. Since $\|N[\nabla u]\|_{L^p(\partial\Omega)}^p = \sum_i \|N[\nabla u]\|_{L^p(\Delta_i)}^p$ we obtain after rearranging the summation order:

$$\begin{aligned} (5.15) \quad \|N[\nabla u]\|_{L^p(\partial\Omega)}^p & \lesssim \sum_{i \in \mathbb{Z}} \left(1 + \sum_{k > 9} e^{-\alpha k} \right) \left[\|(f - f_{\Delta_i})\psi_i\|_{\dot{L}^p_{1,1/2}(\partial\Omega)}^p + |f_{\Delta_{i-1}} - f_{\Delta_i}|^p \right] \\ & \lesssim \sum_{i \in \mathbb{Z}} \left[\|(f - f_{\Delta_i})\psi_i\|_{\dot{L}^p_{1,1/2}(\partial\Omega)}^p + \sum_{i \in \mathbb{Z}} |f_{\Delta_{i-1}} - f_{\Delta_i}|^p \right]. \end{aligned}$$

We are nearly done. Recall [Proposition 2.44](#) which gives estimate for norm of the term $(f - f_{\Delta_i})\psi_i$. It follows that

$$(5.16) \quad \|N[\nabla u]\|_{L^p(\partial\Omega)}^p \lesssim \sum_{i \in \mathbb{Z}} |f_{\Delta_{i-1}} - f_{\Delta_i}|^p + \|g\|_{L^p(\partial\Omega)}^p + \|h\|_{L^p(\partial\Omega)}^p,$$

where for functions $g \in \mathcal{D}_s(f)$, h we have used the finite overlap of the balls $32\Delta_i$ which allows up to claim that

$$\sum_{i \in \mathbb{Z}} \|g\|_{L^p(32\Delta_i)}^p \lesssim \|g\|_{L^p(\partial\Omega)}^p.$$

For both g and h we have bounds on their L^p norms by $\|f\|_{\dot{L}^p_{1,1/2}(\partial\Omega)}$, provided g is chosen correctly. Thus (5.16) only requires bound on the first term on the righthand side.

We now wish to show

$$(5.17) \quad |f_{\Delta_{i-1}} - f_{\Delta_i}| \lesssim \|f^\#\|_{L^p(\tilde{\Delta}_i)},$$

where $\tilde{\Delta}_i$ is an enlargement of Δ_i and $f^\#$ is the sharp maximal function of f defined by (2.30). Recall that we have established that $\|f^\#\|_{L^p(\partial\Omega)} \leq C\|f\|_{\dot{L}^p_{1,1/2}(\partial\Omega)}$.

We have by (2.35):

$$\left| f(X, t)\chi_{\Delta_{i-1}}(X, t) - \left(\int_{\Delta_i} f \right) \chi_{\Delta_{i-1}}(X, t) \right| \lesssim \left(f^\#(X, t) + \int_B f^\# \right) \chi_{\Delta_{i-1}}(X, t),$$

for $B = \Delta_{i-1} \cup \Delta_i$. Integrating over $\partial\Omega$ yields

$$|f_{\Delta_{i-1}} - f_{\Delta_i}| \leq C \int_B f^\#.$$

Since the volume of B is $O(1)$ the average can be replaced by $\|f^\#\|_{L^1(\tilde{\Delta}_i)}$. From this an estimate analogous to (5.17) follows. In summary, we have for (5.16) that

$$(5.18) \quad \|N[\nabla u]\|_{L^p(\partial\Omega)} \lesssim \|f\|_{\dot{L}^p_{1,1/2}(\partial\Omega)},$$

as desired. Hence the bounded case is also established and [Theorem 1.3](#) holds. \square

6. INTERPOLATION AND *Theorem 1.4*.

6.1. **Atomic spaces** $\dot{HS}_{1,1/2}^{1,(\beta)}(\mathbb{R}^n)$. We plan to define natural end-point spaces for $p = 1$ which interpolates well with the homogeneous parabolic Sobolev spaces $\dot{L}_{1,1/2}^p(\mathbb{R}^n)$ defined above for $p > 1$. It follows that we aim to define spaces that respect parabolic homogeneity. Thus for an “atom” a we expect ∇a to satisfy the usual condition defining Hardy-Sobolev space with one derivative. However in the t -variable we only expect to have a $1/2$ -time derivative. We borrow lots of ideas from [2] (see Definition 2.16 there).

Definition 6.1. Let $1 < \beta < \infty$. We say that a function $b : \mathbb{R}^n \rightarrow \mathbb{R}$ is a homogeneous $(1, 1/2, \beta)$ -atom associated to a parabolic ball $B \subset \mathbb{R}^n$ of radius $r = r(B)$ if

- (i) b is supported in the ball B ,
- (ii) $\|\nabla b\|_{L^\beta(\mathbb{R}^n)} + \|D_t^{1/2}b\|_{L^\beta(\mathbb{R}^n)} \leq |B|^{-1/\beta'}$,
- (iii) $\|b\|_{L^1(\mathbb{R}^n)} \leq r(B)$.

Here $|B|$ denotes the usual n -dimensional Lebesgue measure on \mathbb{R}^n . When $p = \infty$ we modify (ii) and replace it by $\|\nabla b\|_{L^\infty(\mathbb{R}^n)} + \|D_t^{1/2}b\|_{\text{BMO}(\mathbb{R}^n)} \leq |B|^{-1}$.

We say that a locally integrable function f belongs to the homogeneous parabolic atomic Hardy-Sobolev space $\dot{HS}_{1,1/2}^{1,(\beta)}(\mathbb{R}^n)$ if there exists a family of locally integrable homogeneous $(1, 1/2, \beta)$ -atoms $(b_i)_i$ such that $f = \sum_i \lambda_i b_i$, with $\sum_i |\lambda_i| < \infty$. We equip the space $\dot{HS}_{1,1/2}^{1,(\beta)}(\mathbb{R}^n)$ with the norm:

$$\|f\|_{\dot{HS}_{1,1/2}^{1,(\beta)}} = \inf_{(\lambda_i)_i} \sum_i |\lambda_i|.$$

The condition (iii) is not strictly necessary as it follows from the conditions (i) and (ii) and Poincaré’s inequality in spatial gradient ∇b . For each fixed $t \in \mathbb{R}$ let $\Delta_t = B \cap \{t = \text{const}\}$ be the “ball” in spatial variables $x \in \mathbb{R}^{n-1}$. Since b vanishes on $\partial\Delta_t$ we have that

$$\|b\|_{L^1(\cdot, t)} = \int_{\Delta_t} |b| dx \leq r \int_{\Delta_t} |\nabla b| dx.$$

Integrating in t we then get

$$\|b\|_{L^1(\mathbb{R}^n)} \leq r \int_{t \in \mathbb{R}} \int_{\Delta_t} |\nabla b| dx dt = r \int_B |\nabla b| \leq r \|\nabla b\|_{L^\beta} |B|^{1/\beta'},$$

by Hölder. From this using (ii) we see that (iii) follows.

In the definition above the condition (ii) seems to be “global” as the L^β norm is calculated over whole space. However, clearly $\text{supp } \nabla b \subset B$. We also claim that $D_t^{1/2}b$ is “essentially” localised to a ball $2B = B(X, 2r)$ and decaying away from the support of B . We have the following:

Lemma 6.2. For an $(1, 1/2, \beta)$ -atom b supported on $B = B((x, t), r)$ we have for $Y = (y, s)$:

- (i) $D_t^{1/2}b(y, s) = 0$ if $|x - y| \geq r$.
- (ii) If $|x - y| < r$ and $|t - s| > 4r^2$ then

$$|D_t^{1/2}b(y, s)| \leq C|t - s|^{-3/2} \|b\|_{L^1(B \cap \{y = \text{const}\})}.$$

Proof. Using the definition of the half-derivative we have:

$$|D_t^{1/2}b(y, s)| = \left| c_1 \int_{\tau \in \mathbb{R}} \frac{b(y, s) - b(y, \tau)}{|s - \tau|^{3/2}} d\tau \right| \leq c_1 \int_{\tau \in \mathbb{R}} \frac{|b(y, \tau)|}{|s - \tau|^{3/2}} d\tau.$$

Given that b is only supported in B , only $\tau \in (t - r^2, t + r^2)$ contribute to the integral above. Since for such τ : $|s - \tau| \approx |s - t|$ we get that

$$|D_t^{1/2}b(y, s)| \lesssim |t - s|^{-3/2} \|b\|_{L^1(Q \cap \{y = \text{const}\})}.$$

□

Our aim is to prove that if b is a $(1, 1/2, \beta)$ -atom then $b \in \dot{W}_{1,1/2}^1(\mathbb{R}^n)$ with $\dot{W}_{1,1/2}^1(\mathbb{R}^n)$ norm $\lesssim 1$. It then would follow that for every $\beta > 1$ we have that

$$(6.3) \quad \dot{HS}_{1,1/2}^{1,(\beta)}(\mathbb{R}^n) \subset \dot{W}_{1,1/2}^1(\mathbb{R}^n) \quad \text{and} \quad \|f\|_{\dot{W}_{1,1/2}^1(\mathbb{R}^n)} \lesssim \|f\|_{\dot{HS}_{1,1/2}^{1,(\beta)}(\mathbb{R}^n)}.$$

Consider therefore an arbitrary $(1, 1/2, \beta)$ -atom b supported on B centred at (x, t) . By Hölder we have that

$$\begin{aligned} \|\nabla b\|_{L^1(\mathbb{R}^n)} + \|D_t^{1/2}b\|_{L^1(2B)} &= \|\nabla b\|_{L^1(2B)} + \|D_t^{1/2}b\|_{L^1(2B)} \\ &\leq [\|\nabla b\|_{L^\beta(2B)} + \|D_t^{1/2}b\|_{L^\beta(2B)}] |2B|^{1/\beta'}. \end{aligned}$$

Using the fact that we have a doubling measure and condition (ii) in the definition of the atom the claim

$$\|\nabla b\|_{L^1(\mathbb{R}^n)} + \|D_t^{1/2}b\|_{L^1(2B)} \lesssim 1$$

follows. It remains to estimate $D_t^{1/2}b$ away from the ball $2B$. For this we use [Lemma 6.2](#). By (i) and (ii):

$$(6.4) \quad \begin{aligned} \|D_t^{1/2}b\|_{L^1(\mathbb{R}^n \setminus 2B)} &\leq \int_{s \in \mathbb{R}: |t-s| > 4r^2} \int_{|y-x| < r} |D_{1/2}^t b(y, s)| dy ds \\ &\leq \int_{s \in \mathbb{R}: |t-s| > 4r^2} |t-s|^{-3/2} \int_{|y-x| < r} \|b\|_{L^1(B \cap \{y, \cdot\})} dy ds. \end{aligned}$$

The interior integral in y will give us $\|b\|_{L^1(B)}$ thus yielding:

$$(6.5) \quad \|D_t^{1/2}b\|_{L^1(\mathbb{R}^n \setminus 2B)} \leq C \|b\|_{L^1(B)} \int_{4r^2}^{\infty} \tau^{-3/2} d\tau \approx r^{-1} \|b\|_{L^1(B)} \leq 1,$$

where in the final step we have used property (iii) of the definition of an atom b . The calculation of the BMO norm of $D_t^{1/2}b$ is analogous for $\beta = \infty$.

Lemma 6.6. *When $1 < \beta \leq \alpha \leq \infty$ an $(1, 1/2, \alpha)$ -atom b is a multiple of an $(1, 1/2, \beta)$ -atom b and hence*

$$(6.7) \quad \dot{HS}_{1,1/2}^{1,(\infty)}(\mathbb{R}^n) \subset \dot{HS}_{1,1/2}^{1,(\alpha)}(\mathbb{R}^n) \subset \dot{HS}_{1,1/2}^{1,(\beta)}(\mathbb{R}^n).$$

Moreover, for each $\beta > 1$ the space $\dot{HS}_{1,1/2}^{1,(\beta)}(\mathbb{R}^n)$ is a Banach space.

Proof. Only condition (ii) differs for different values of β . A calculation similar to the one we have performed above implies that

$$\|\nabla b\|_{L^\beta(\mathbb{R}^n)} + \|D_t^{1/2}b\|_{L^\beta(2B)} \leq [\|\nabla b\|_{L^\alpha(\mathbb{R}^n)} + \|D_t^{1/2}b\|_{L^\alpha(2B)}] |2B|^{1/\beta-1/\alpha}.$$

This also holds for $\alpha = \infty$ replacing the L^∞ norm by BMO as $\text{BMO} \subset L^\beta$ for $\beta < \infty$. Assuming that b is a $(1, 1/2, \alpha)$ -atom by (ii) of the definition of an atom and using doubling we get that

$$\|\nabla b\|_{L^\beta(\mathbb{R}^n)} + \|D_t^{1/2}b\|_{L^\beta(2B)} \leq C_{n+1}^{1/\beta-1/\alpha} |B|^{1/\beta-1} = C_{n+1}^{1/\beta-1/\alpha} |B|^{-1/\beta'}.$$

Here C_{n+1} is the doubling constant. As before $\|D_t^{1/2}b\|_{L^\beta(\mathbb{R}^n \setminus 2B)}$ requires a separate treatment. We again use [Lemma 6.2](#). By (i) and (ii):

$$(6.8) \quad \begin{aligned} \|D_t^{1/2}b\|_{L^\beta(\mathbb{R}^n \setminus 2B)}^\beta &\leq \int_{s \in \mathbb{R}: |t-s| > 4r^2} \int_{|y-x| < r} |D_{1/2}^t b(y, s)|^\beta dy ds \\ &\leq \int_{s \in \mathbb{R}: |t-s| > 4r^2} |t-s|^{-3\beta/2} \int_{|y-x| < r} \|b\|_{L^1(B \cap \{(y, \cdot)\})}^\beta dy ds. \end{aligned}$$

By Hölder

$$\|b\|_{L^1(B \cap \{(y, \cdot)\})}^\beta \leq \|b\|_{L^\beta(B \cap \{(y, \cdot)\})}^\beta r^{2(\beta-1)}.$$

Hence

$$(6.9) \quad \|D_t^{1/2}b\|_{L^\beta(\mathbb{R}^n \setminus 2B)}^\beta \leq C \|b\|_{L^\beta(B)}^\beta r^{2(\beta-1)} \int_{4r^2}^\infty \tau^{-3\beta/2} d\tau = C \|b\|_{L^\beta(B)}^\beta r^{-\beta}.$$

It follows that

$$\|D_t^{1/2}b\|_{L^\beta(\mathbb{R}^n \setminus 2B)} \leq Cr^{-1} \|b\|_{L^\beta(B)}.$$

As before using the (i) and (ii) we can obtain $\|b\|_{L^\beta} \leq Cr \|\nabla b\|_{L^\beta(B)}$ and hence

$$\|\nabla b\|_{L^\beta(\mathbb{R}^n)} + \|D_t^{1/2}b\|_{L^\beta(\mathbb{R}^n)} \leq C |B|^{-1/\beta'}.$$

From this it follows that $\frac{b}{C}$ is a $(1, 1/2, \beta)$ -atom. The proof that the defined space is Banach is trivial (c.f. [\[2\]](#) for similar proofs). \square

We now consider the real interpolation between the spaces $\dot{H}S_{1,1/2}^{1,(\beta)}(\mathbb{R}^n)$ and $\dot{W}_{1,1/2}^\infty(\mathbb{R}^n)$, where [\(2.12\)](#) defines the space $\dot{W}_{1,1/2}^\infty$. Recall some classical results on K -method of real interpolation from [\[4, 5\]](#).

Let A_0, A_1 be two normed vector spaces embedded in a topological Hausdorff vector space V . For each $a \in A_0 + A_1$ and $t > 0$, we define the K -functional of real interpolation by

$$K(a, t, A_0, A_1) = \inf_{a=a_0+a_1} (\|a_0\|_{A_0} + t\|a_1\|_{A_1}).$$

For $0 < \theta < 1$, $1 \leq q \leq \infty$, we denote by $(A_0, A_1)_{\theta, q}$ the real interpolation space between A_0 and A_1 defined as

$$(A_0, A_1)_{\theta, q} = \left\{ a \in A_0 + A_1 : \|a\|_{\theta, q} = \left(\int_0^\infty (t^{-\theta} K(a, t, A_0, A_1))^q \frac{dt}{t} \right)^{\frac{1}{q}} < \infty \right\}.$$

It is an exact interpolation space of exponent θ between A_0 and A_1 (see [\[5\]](#), Chapter II).

Definition 6.10. Let f be a measurable function on a measure space (X, μ) . The decreasing rearrangement of f is the function f^* defined for every $t \geq 0$ by

$$f^*(t) = \inf \{ \lambda : \mu(\{x : |f(x)| > \lambda\}) \leq t \}.$$

The maximal decreasing rearrangement of f is the function f^{**} defined for every $t > 0$ by

$$f^{**}(t) = \frac{1}{t} \int_0^t f^*(s) ds.$$

From the properties of f^{**} we mention:

1. $(f + g)^{**} \leq f^{**} + g^{**}$.
2. $(Mf)^* \sim f^{**}$.
3. $\mu(\{x; |f(x)| > f^*(t)\}) \leq t$.
4. $\forall 1 < p \leq \infty, \|f^{**}\|_p \sim \|f\|_p$.

We exactly know the functional K for Lebesgue spaces :

Proposition 6.11. Take $0 < p_0 < p_1 \leq \infty$. We have :

$$K(f, t, L^{p_0}, L^{p_1}) \simeq \left(\int_0^{t^\alpha} [f^*(s)]^{p_0} ds \right)^{1/p_0} + t \left(\int_{t^\alpha}^\infty [f^*(s)]^{p_1} ds \right)^{1/p_1},$$

where $\frac{1}{\alpha} = \frac{1}{p_0} - \frac{1}{p_1}$.

When the $p_1 = \infty$ and the L^{p_1} endpoint is replaced by BMO we have by the result [3], (see also [27]) that

$$K(f, t, L^{p_0}, \text{BMO}) \approx t(M_{p_0}^\# f)^*(t^{p_0}),$$

where $M_p^\#$ is a variant of the maximal function that can be bounded by M , the usual maximal function.

We characterize the K -functional of real interpolation in the following theorem:

Proposition 6.12.

1. For all $\beta \in (1, \infty)$, there exists $C_1 > 0$ such that for every $f \in \dot{H}S_{1,1/2}^{1,(\beta)}(\mathbb{R}^n) + \dot{W}_{1,1/2}^{\infty, \text{BMO}}(\mathbb{R}^n)$ and $t > 0$,

$$K(f, t, \dot{H}S_{1,1/2}^{1,(\beta)}, \dot{W}_{1,1/2}^{\infty, \text{BMO}}) \geq C_1 t \left(|\nabla f|^{**} + |M(D_t^{1/2} f)|^{**} \right) (t);$$

2. for $1 < \beta < \infty$, there exists $C_2 > 0$ such that for every $f \in \dot{L}_{1,1/2}^\beta(\mathbb{R}^n)$ and $t > 0$,

$$K(f, t, \dot{H}S_{1,1/2}^{1,(\beta)}, \dot{W}_{1,1/2}^{\infty, \text{BMO}}) \leq C_2 t \left(|\nabla f|^{\beta^{**\frac{1}{\beta}}} + |D_t^{1/2} f|^{\beta^{**\frac{1}{\beta}}} + |S_1(f)|^{\beta^{**\frac{1}{\beta}}} + |f^\#|^{\beta^{**\frac{1}{\beta}}} \right) (t).$$

Proof. The lower bound is trivial and follows from the fact that $\dot{H}S_{1,1/2}^{1,(\beta)} \subset \dot{W}_{1,1/2}^1$ and the characterisation of K between L^1 and BMO.

For the upper bound, the result is based on the following Calderón-Zygmund decomposition lemma.

Proposition 6.13 (Calderón-Zygmund lemma for Sobolev functions). *Let $1 < \beta < \infty$, $f \in L_{1,1/2}^\beta(\mathbb{R}^n)$ and $\alpha > 0$. Then one can find a collection of balls $(B_i)_i$, functions b_i and a function $g \in \dot{W}_{1,1/2}^{\infty, \text{BMO}}(\mathbb{R}^n)$ such that the following properties hold:*

$$(6.14) \quad f = g + \sum_i b_i,$$

$$(6.15) \quad |\nabla g(X)| \leq C\alpha, \quad g^\sharp(X) \leq C\alpha \text{ and } \|D_t^{1/2}g\|_{\text{BMO}(\mathbb{R}^n)} \leq C\alpha \text{ a.e. } x \in \mathbb{R}^n,$$

$$(6.16) \quad \text{supp } b_i \subset B_i, \quad \|b_i\|_{\dot{H}S_{1,1/2}^{1,(\beta)}(\mathbb{R}^n)} \leq C\alpha|B_i|,$$

$$(6.17) \quad \sum_i |B_i| \leq C\alpha^{-\beta} \int (|\nabla f| + |D_t^{1/2}f| + |S_1(f)| + f^\sharp)^\beta,$$

$$(6.18) \quad \sum_i \mathbf{1}_{B_i} \leq N\mathbf{1}_\Omega,$$

$$(6.19) \quad \left\| \sum_i b_i \right\|_{\dot{H}S_{1,1/2}^{1,(\beta)}} \leq \sum_i \|b_i\|_{\dot{H}S_{1,1/2}^{1,(\beta)}} \leq CN\alpha|\Omega|.$$

Here C and N only depend on q , β and on the dimension n . By $\mathbf{1}_A$ we denote the indicator function of the set A and $\Omega = \bigcup_i B_i$.

We postpone the proof and first show how it implies the upper bound in [Proposition 6.12](#).

Take $f \in \dot{L}_{1,1/2}^\beta(\mathbb{R}^n)$ with $\beta > 1$ and $1 < q \leq \beta < \infty$. Let $t > 0$. We consider the Calderón-Zygmund decomposition of [Proposition 6.13](#) for f with $\alpha = \alpha(t) = \left(M(|\nabla f|^\beta + |D_t^{1/2}f|^\beta + |S_1(f)|^\beta + |f^\sharp|^\beta)\right)^{\frac{1}{\beta}}(t)$. We write $f = \sum_i b_i + g = b + g$ where $(b_i)_i$, g satisfy the properties of the proposition. By [\(6.19\)](#)

$$\|b\|_{\dot{H}S_{1,1/2}^{1,(\beta)}} \leq C\alpha(t) \sum_i |B_i| \leq C\alpha(t)\mu(\Omega),$$

where $\Omega = \Omega_t = \bigcup_i B_i$. Moreover, since $(Mf)^* \sim f^{**}$ and $(f+g)^{**} \leq f^{**} + g^{**}$, we get

$$\alpha(t) \lesssim \left(|\nabla f|^{\beta^{**\frac{1}{\beta}}} + |D_t^{1/2}f|^{\beta^{**\frac{1}{\beta}}} + |S_1(f)|^{\beta^{**\frac{1}{\beta}}} + |f^\sharp|^{\beta^{**\frac{1}{\beta}}}\right)(t).$$

Noting that for this choice of $\alpha(t)$, $|\Omega_t| \leq t$ (c.f [\[4\],\[5\]](#)), we deduce that

$$(6.20) \quad \begin{aligned} K(f, t, \dot{H}S_{1,1/2}^{1,(\beta)}, \dot{W}_{1,1/2}^\infty) &\leq \|b\|_{\dot{H}S_{1,1/2}^{1,(\beta)}} + t\|g\|_{\dot{W}_{1,1/2}^\infty} \\ &\lesssim t \left(|\nabla f|^{\beta^{**\frac{1}{\beta}}}(t) + |D_t^{1/2}f|^{\beta^{**\frac{1}{\beta}}}(t) + |S_1(f)|^{\beta^{**\frac{1}{\beta}}}(t) + |f^\sharp|^{\beta^{**\frac{1}{\beta}}}(t)\right), \end{aligned}$$

for all $t > 0$ and obtain the desired inequality for $f \in W^{1,\beta}$, $q \leq \beta < \infty$. \square

Then integrating the K -functional yields

Proposition 6.21. *For all $\beta \in (1, \infty)$ and $p \in [\beta, \infty)$, $\dot{L}_{1,1/2}^p(\mathbb{R}^n)$ is a real interpolation space between $\dot{H}S_{1,1/2}^{1,(\beta)}(\mathbb{R}^n)$ and $\dot{W}_{1,1/2}^{\infty,\text{BMO}}(\mathbb{R}^n)$. More precisely, we have*

$$\left(\dot{H}S_{1,1/2}^{1,(\beta)}(\mathbb{R}^n), \dot{W}_{1,1/2}^{\infty,\text{BMO}}(\mathbb{R}^n) \right)_{1-\frac{1}{p}, p} = \dot{L}_{1,1/2}^p(\mathbb{R}^n).$$

We use the fact that the L^p norm of the functions f^\sharp and $S_1(f)$ are controlled by the norm of f in $\dot{L}_{1,1/2}^p$ for all $p > 1$.

Proof of Proposition 6.13. Let $f \in W^{1,\beta}$, $\alpha > 0$ and consider

$$\Omega = \left\{ X \in \mathbb{R}^n : M(|\nabla f|^\beta + |D_t^{1/2} f|^\beta + |S_1(f)|^\beta + |f^\sharp|^\beta)(X) > \alpha^\beta \right\}.$$

If $\Omega = \emptyset$, then set

$$g = f, \quad b_i = 0 \text{ for all } i$$

so that (6.15) is satisfied according to the Lebesgue differentiation theorem. Otherwise, the maximal theorem yields

$$\begin{aligned} |\Omega| &\leq C\alpha^{-\beta} \| (|\nabla f| + |D_t^{1/2} f| + |S_1(f)|^\beta + |f^\sharp|^\beta) \|_{L^1} \\ (6.22) \quad &\leq C\alpha^{-\beta} \left(\int |\nabla f|^\beta + \int |D_t^{1/2} f|^\beta + \int |S_1(f)|^\beta + \int |f^\sharp|^\beta \right) \\ &< +\infty. \end{aligned}$$

In particular $\Omega \neq \mathbb{R}^n$. Let F be the complement of Ω . Since Ω is an open set distinct of \mathbb{R}^n , let (\underline{B}_i) be a Whitney decomposition of Ω ([11]). That is, the \underline{B}_i are pairwise disjoint, and there exist two constants $C_2 > C_1 > 1$, depending only on the metric, such that

1. $\Omega = \cup_i B_i$ with $B_i = C_1 \underline{B}_i$ and the balls B_i have the bounded overlap property;
2. $r_i = r(B_i) = \frac{1}{2}d(X_i, F)$ and X_i is the center of B_i ;
3. each ball $\overline{B}_i = C_2 \underline{B}_i$ intersects F ($C_2 = 4$ works).

For $X \in \Omega$, denote $I_X = \{i : X \in B_i\}$. Recall that $\sharp I_X \leq N$ and fixing $j \in I_X$, $B_i \subset 7B_j$ for all $i \in I_X$. From this (6.18) follows.

Condition (6.17) is satisfied due to (6.22). Using the doubling property, and the fact that $C_2 B_i$ intersects F we have

$$(6.23) \quad \int_{7B_i} (|\nabla f|^\beta + |D_t^{1/2} f|^\beta + |S_1(f)|^\beta + |f^\sharp|^\beta) d\mu \leq C\alpha^\beta |B_i|.$$

Let us now define the functions b_i . For this, we construct a partition of unity. Let $(\chi_i)_i$ be a partition of unity of Ω subordinated to the covering (B_i) . Each χ_i is a Lipschitz function supported in B_i with $\|\nabla \chi_i\|_\infty \leq Cr_i^{-1}$ and $\|\partial_t \chi_i\|_\infty \leq Cr_i^{-2}$.

We set $b_i = (f - \frac{1}{\chi_i(B_i)} \int_{B_i} f \chi_i) \chi_i$ where $\chi_i(B_i) := \int_{B_i} \chi_i \simeq |B_i|$. We estimate the L^β norms of ∇b_i and $D_t^{1/2} b_i$. We start with the ∇b_i . Clearly this function is supported on B_i and

$$\nabla b_i = (\nabla f) \chi_i + \left(f - \frac{1}{\chi_i(B_i)} \int_{B_i} f \chi_i \right) \nabla \chi_i.$$

Using (2.36) we have that

$$\int_{\mathbb{R}^n} |\nabla b_i|^\beta \leq \int_{B_i} |\nabla f|^\beta + \int_{B_i} |f^\sharp|^\beta.$$

Hence by (6.23) $\|\nabla b_i\|_{L^\beta} \leq C\alpha|B_i|^{1/\beta}$. Moving to $D_t^{1/2}b_i$ we get that

$$\begin{aligned} D_t^{1/2}b_i(x, t) &= [(D_t^{1/2}f)\chi_i](x, t) + c \int_{s \in \mathbb{R}} \frac{\chi_i(x, t) - \chi_i(x, s)}{|s - t|^{3/2}} \left(f - \frac{1}{\chi_i(B_i)} \int_{B_i} f \chi_i \right) (x, s) ds. \\ (6.24) \qquad \qquad &= [(D_t^{1/2}f)\chi_i](x, t) + R_i(x, t). \end{aligned}$$

We first integrate over $2B_i$. The first term enjoys an estimate similar to ∇b_i and again by (6.23) $\|(D_t^{1/2}f)\chi_i\|_{L^\beta(\mathbb{R}^n)} \leq C\alpha|B_i|^{1/\beta}$. For the second term using Lipschitzness of χ_i in the t -variable we may estimate $\frac{|\chi_i(x, t) - \chi_i(x, s)|}{|s - t|^{3/2}}$ by $\frac{C}{r_i^2|s - t|^{1/2}}$. By (2.36) we then have

$$\int_{2B_i} |R_i(x, t)|^\beta \leq \int_{2B_i} r_i^{-\beta} \left(\int_{(x, s) \in B_i} |s - t|^{-1/2} [f^\sharp(x, s) + \int_{B_i} f^\sharp] ds \right)^\beta dt dx.$$

Here the second term containing the average of f^\sharp has an easy estimate by $C \int_{B_i} |f^\sharp|^\beta$. Hence we focus on the first term. Integration over s can be split into integration over the sets where $|s - t| \approx 2^{-i}r_i^2$ for $i = 0, 1, 2, \dots$. Hence we get a bound by

$$\int_{2B_i} \left(\sum_{i=0}^{\infty} 2^{-i/2} \int_{|s-t| \sim 2^{-i}r_i^2} \frac{f^\sharp(x, s)}{2^{-i}r_i^2} \right)^\beta dt dx.$$

We can recognise that the inside integral is bounded by a truncated maximal function $M^t(f^\sharp)(x, t)$ in t -variable. The maximal function is truncated as the sup runs over balls of radius $\leq r_i^2$. It follows that the expression is bounded by

$$\int_{2B_i} [M^t(f^\sharp)]^\beta \leq \int_{3B_i} |f^\sharp|^\beta.$$

We have used boundedness of M^t on L^β and the fact that M^t is truncated version of the maximal function and hence the integral on the righthand side only needs to be taken over an enlargement of $2B_i$. After putting all terms together we get from (6.23) that

$$\|D_t^{1/2}b_i\|_{L^\beta(2B_i)} \leq C\alpha|B_i|^{1/\beta}.$$

It remain to consider the away part when $(x, t) \in \mathbb{R}^n \setminus 2B_i$. There is nothing to calculate unless there exists s for which $(x, s) \in B_i$ as the term $R_i(x, t)$ vanishes otherwise. Hence assuming that, we then have by (2.36)

$$|R_i(x, t)| \leq C \int_{B_i \cap \mathcal{H}^x} \frac{r_i}{|s - t|^{3/2}} [f^\sharp(x, s) + \int_{B_i} f^\sharp] ds.$$

Here \mathcal{H}^x denotes the line $\{(x, s) : s \in \mathbb{R}\}$. Hence

$$|R_i(x, t)| \leq C \left(\frac{r_i}{d((x, t), B_i)} \right)^3 \left[\int_{B_i \cap \mathcal{H}^x} f^\sharp + \int_{B_i} f^\sharp \right].$$

Here $d(x, B_i) = \inf\{d((x, t), Y) : Y \in B_i\}$ measures the parabolic distance of (x, t) to B_i . Hence

$$\int_{\mathbb{R}^n \setminus 2B_i} |R_i|^\beta \lesssim |B_i| \int_{B_i} |f^\#|^\beta.$$

Now (6.23) together with the previous estimates gives us:

$$(6.25) \quad \|b_i\|_{\dot{H}S_{1,1/2}^{1,(\beta)}} = (\|\nabla b_i\|_{L^\beta(\mathbb{R}^n)} + \|D_t^{1/2} b_i\|_{L^\beta(\mathbb{R}^n)}) |B_i|^{1/\beta'} \leq C\alpha |B_i|^{1/\beta+1/\beta'} = C\alpha |B_i|.$$

Hence $a_i := b_i/(C\alpha |B_i|)$ are $(1, 1/2, \beta)$ -atoms and (6.16) is proved. Let $\lambda_i = C\alpha |B_i|$. It follows that

$$b = \sum_i b_i = \sum_i \lambda_i a_i \in \dot{H}S_{1,1/2}^{1,(\beta)}(\mathbb{R}^n) \text{ with norm bounded by: } \sum_i |\lambda_i| = C\alpha \sum_i |B_i|.$$

where in the final step we have used finite overlap of the Whitney balls B_i . Thus $\|b\|_{\dot{H}S_{1,1/2}^{1,(\beta)}} \leq CN\alpha |\Omega|$ by (6.18). From this (6.19) follows.

Set $g = f - \sum_i b_i$. Since the sum is locally finite on Ω , as usual, g is defined almost everywhere on \mathbb{R}^n and $g = f$ on F . Moreover, g is a locally integrable function on \mathbb{R}^n . It remains to prove (6.15). We have

$$\begin{aligned} \nabla g &= \nabla f - \sum_i \nabla b_i \\ &= \nabla f - \left(\sum_i \chi_i \right) \nabla f - \sum_i \left(f - \frac{1}{\chi_i(B_i)} \int_{B_i} f \chi_i \right) \nabla \chi_i \\ &= \mathbf{1}_F(\nabla f) - \sum_i \left(f - \frac{1}{\chi_i(B_i)} \int_{B_i} f \chi_i \right) \nabla \chi_i. \end{aligned}$$

From the definition of F and the Lebesgue differentiation theorem, we have $\mathbf{1}_F |\nabla f| \leq \alpha \mu$ -a.e. We claim that a similar estimate holds for

$$h = \sum_i \left(f - \frac{1}{\chi_i(B_i)} \int_{B_i} f \chi_i \right) \nabla \chi_i,$$

that is $|h(X)| \leq C\alpha$ for all $X \in \mathbb{R}^n$. For this, note first that h vanishes on F and the sum defining h is locally finite on Ω .

Then fix $X \in \Omega$ and fix some $j \in I_X$. Note that $\sum_i \chi_i(X) = 1$ and $\sum_i \nabla \chi_i(X) = 0$ and hence we can rewrite the formula for h as

$$h(X) = \sum_{i \in I_X} \left[\left(\frac{1}{|7B_j|} \int_{7B_j} f \right) - \left(\frac{1}{\chi_i(B_i)} \int_{B_i} f \chi_i \right) \right] \nabla \chi_i(X).$$

We return to the calculation we have done above (2.36). With $B = B_i$, $\chi = \chi_i$ and any $Z \in 7B_j$ we get by it that

$$-Cr_i \left[f^\#(Z) + \int_{B_i} f^\# \right] \leq f(X) - \frac{1}{\chi_i(B_i)} \int_{B_i} f \chi_i \leq Cr_i \left[f^\#(Z) + \int_{B_i} f^\# \right].$$

Hence by averaging over $7B_i$ and using that the measure is doubling we get that

$$(6.26) \quad \left| \int_{7B_j} f - \frac{1}{\chi_i(B_i)} \int_{B_i} f \chi_i \right| \leq Cr_i \int_{7B_j} f^\# \lesssim \alpha r_j,$$

where we have used (6.23) for $7B_j$ in the last step. Hence

$$(6.27) \quad |h(X)| \lesssim \sum_{i \in I_x} \alpha r_j r_j^{-1} \leq CN\alpha,$$

and the bound (6.15) for ∇g holds. Next, we look at $g^\#$. The result for ∇g already proves that

$$|g(x, t) - g(y, t)| \leq C\alpha|x - y|.$$

It follows that we need to prove that

$$(6.28) \quad |g(x, t) - g(x, s)| \leq C\alpha|t - s|^{1/2},$$

since the desired result then follows by triangle inequality $|g(x, t) - g(y, s)| \leq |g(x, t) - g(y, t)| + |g(y, t) - g(y, s)|$. Let us gain a better understanding on the function g . It follows from the definition of g that

$$(6.29) \quad g(x) = \begin{cases} f(x), & \text{when } x \in F, \\ \sum_i (\chi_i(B_i))^{-1} \int_{B_i} f \chi_i, & \text{when } x \in \Omega. \end{cases}$$

Hence if we simplify it, on the good set $g = f$ while on the bad set we take averages of f on B_i and then smoothly glue them together.

It follows that if $(x, t), (x, s) \in F$ there is nothing to do. By (2.34) we have that

$$|g(x, t) - g(x, s)| = |f(x, t) - f(x, s)| = |t - s|^{1/2}[f^\#(x, t) + f^\#(x, s)],$$

and since both points are in the set F , clearly $f^\#(x, t) + f^\#(x, s) \leq 2\alpha$. Next consider the case when both $(x, t), (x, s)$ belong to some neighbouring balls B_i, B_j with $B_i \subset 7B_j$ and $B_j \subset 7B_i$. It follows that $r_i \approx r_j$ and $|t - s| \lesssim r_i^2$. Thus we are in situation where (6.26) holds and we may perform the same calculation for $\partial_t g$ as we have done above for ∇g . It follows that

$$\partial_t g(x, t) = \sum_{i \in I(x, t)} \left[\left(\frac{1}{|7B_j|} \int_{7B_j} f \right) - \left(\frac{1}{\chi_i(B_i)} \int_{B_i} f \chi_i \right) \right] \partial_t \chi_i(x, t),$$

and hence $|\partial_t g| \lesssim \alpha r_i^{-1}$. Same is true at the point (x, s) and in fact any point on the line segment joining these two points by the same argument. Finally by the mean value theorem

$$(6.30) \quad |g(x, t) - g(x, s)| \leq |\partial_t g(x, \tau)| |t - s| \leq C\alpha |t - s|^{1/2} \frac{|t - s|^{1/2}}{r_i}.$$

As $|t - s|^{1/2} \lesssim r_i$ the claim follows. When $(x, t) \in F$ and $(x, s) \in \Omega$ then $(x, s) \in B_i$ for some i and from the construction of the Whitney balls $r_i \lesssim |t - s|^{1/2}$. Let Y_i be the centre of the ball B_i . Previous argument gives us that $|g(Y_i) - g(x, s)| \leq C\alpha r_i \leq C\alpha |t - s|^{1/2}$ and Hence if we prove

$$|g(x, t) - g(Y_i)| = \left| f(x, t) - \frac{1}{\chi_i(B_i)} \int_{B_i} f \chi_i \right| \lesssim C\alpha |t - s|^{1/2},$$

then by triangle inequality we are done. Here we have used the fact that Y_i being the centre of the ball B_i actually belongs to $\overline{B_i}$ and these are non-overlapping and hence $\chi_i(Y_i) = 1$. A calculation identical to the one done above (2.36) yields that

$$(6.31) \quad \left| f(x, t) - \frac{1}{\chi_i(B_i)} \int_{B_i} f \chi_i \right| \leq C \sup_{Z \in B_i} d((x, t), Z) \left[f^\sharp(x, t) + \int_{B_i} f^\sharp \right].$$

Using (6.23) and the fact that $\sup_{Z \in B_i} d((x, t), Z) \approx |s - t|^{1/2}$ we obtain our claim.

The final case is when $(x, t), (x, s) \in \Omega$ but $|t - s|^{1/2} \gg \max\{r_i, r_j\}$, where r_i, r_j are radii of the balls $(x, t) \in B_i, (x, s) \in B_j$ respectively. The argument is similar. We already know that $|g(x, t) - g(Y_i)| \leq C\alpha r_i, |g(x, s) - g(Y_j)| \leq C\alpha r_j$ where Y_i, Y_j are centres of their respective balls. After we average in (6.31) we obtain

$$(6.32) \quad \left| \frac{1}{\chi_i(B_i)} \int_{B_i} f \chi_i - \frac{1}{\chi_j(B_j)} \int_{B_j} f \chi_j \right| \leq C \sup_{Z \in B_i, Z' \in B_j} d(Z, Z') \left[\int_{B_i} f^\sharp + \int_{B_j} f^\sharp \right],$$

and again by (6.23) this is further bounded by $C\alpha|t - s|^{1/2}$. It follows that $g^\sharp \leq C\alpha$ as desired.

Finally, we consider the BMO bound for $D_t^{1/2}g$. This is clearly the hardest estimate as $D_t^{1/2}$ is a non-local operator. We aim to prove that $D_t^{1/2}g \in \text{BMO}(\mathbb{R}^n)$ with norm $\lesssim \alpha$. We want to apply [Theorem 2.20](#) and hence we conclude that $\mathbb{D}g \in \text{BMO}(\mathbb{R}^n)$ instead. But as (2.11) holds and we already know g is Lipschitz in the spatial variables with norm $\lesssim \alpha$ the conclusion about $D_t^{1/2}g$ would follow.

To prove that $\mathbb{D}g \in \text{BMO}(\mathbb{R}^n)$ we use [Theorem 2.20](#) and estimate (2.21) for the function g . Let us fix a parabolic ball B_r . For $(x, t) \in B_r$ let us define a function

$$(6.33) \quad \rho(x, t) = \begin{cases} 0, & \text{for } (x, t) \in F = \mathbb{R}^n \setminus \Omega, \\ \min\{r_i = r(B_i) : (x, t) \in B_i\}, & \text{for } (x, t) \in \Omega. \end{cases}$$

Observe that when $(x, t) \in \Omega$ then due to the way Whitney decomposition of Ω was defined the radii of balls B_i that contain (x, t) are all comparable and hence for simplicity we took the smallest of such radii. It follows that we can write:

$$(6.34) \quad \begin{aligned} & \frac{1}{|B_r|} \int_{B_r} \int_0^r \gamma^{-3} \left| g(x, t) - \int_{B_\gamma(x, t)} g \right|^2 d\gamma dt dx \leq \\ & \frac{1}{|B_r|} \sum_i \int_{B_i \cap B_r} \int_0^{\min\{r, r_i\}} \gamma^{-3} \left| g(x, t) - \int_{B_\gamma(x, t)} g \right|^2 d\gamma dt dx + \\ & \frac{1}{|B_r|} \int_{B_r} \int_{\min\{r, \rho(x, t)\}}^r \gamma^{-3} \left| g(x, t) - \int_{B_\gamma(x, t)} g \right|^2 d\gamma dt dx = I + II. \end{aligned}$$

We estimate terms I and II separately. We start with the term I . Clearly we only need to consider those indices i such that $B_i \cap B_r \neq \emptyset$.

Without loss of generality we may assume that each set B_r is of the form $Q_r \times (t_0 - r^2, t_0 + r^2)$, where Q_r is a ball in spatial variables. Recall again the result of Strichartz, namely [Theorem 3.3](#) of [\[40\]](#). For a fixed $t \in \mathbb{R}$ we have that

$$(6.35) \quad \frac{1}{|Q_r|} \int_{Q_r} \int_{|y| \leq r} \frac{|g(x+y, t) + g(x-y, t) - 2g(x, t)|^2}{|y|^{n+1}} dy dx \lesssim \|\nabla_x g(\cdot, t)\|_{\text{BMO}(\mathbb{R}^{n-1})}^2.$$

Given that we have shown that $|\nabla g(x, t)| \lesssim \alpha$ the quantity above is bounded by $C\alpha^2$.

We claim that therefore we have

$$\begin{aligned} & \frac{1}{|B_r|} \int_{B_r} \int_0^r \gamma^{-3} \left| g(x, t) - \fint_{|y| \leq \gamma} g(x+y, t) dy \right|^2 d\gamma dt dx \\ &= \frac{1}{2} \frac{1}{|B_r|} \int_{B_r} \int_0^r \gamma^{-3} \left| \fint_{|y| \leq \gamma} [g(x+y, t) + g(x-y, t) - 2g(x, t)] dy \right|^2 d\gamma dt dx \\ &\leq \frac{1}{2} \frac{1}{|B_r|} \int_{B_r} \int_0^r \gamma^{-n-2} \int_{|y| \leq \gamma} |g(x+y, t) + g(x-y, t) - 2g(x, t)|^2 dy d\gamma dt dx \\ &\leq \frac{1}{2} \frac{1}{|B_r|} \int_{B_r} \int_{|y| \leq r} \frac{|g(x+y, t) + g(x-y, t) - 2g(x, t)|^2}{|y|^{n+1}} dy dt dx \\ &\leq \sup_t \frac{1}{2} \frac{1}{|Q_r|} \int_{Q_r \times \{t\}} \int_{|y| \leq r} \frac{|g(x+y, t) + g(x-y, t) - 2g(x, t)|^2}{|y|^{n+1}} dy dx \lesssim \alpha^2, \end{aligned}$$

by (6.35). To estimate the term I in (6.34) by $C\alpha^2$ it suffices to establish that

$$\frac{1}{|B_r|} \sum_i \int_{B_i \cap B_r} \int_0^{\min\{r, r_i\}} \gamma^{-3} \left| \fint_{|y| \leq \gamma} g(x+y, t) - \fint_{B_\gamma(x, t)} g \right|^2 d\gamma dt dx \lesssim \alpha^2.$$

Consider any $(x, t) \in B_i \cap B_r$. By the calculations we have done above for $\partial_t g$ on B_i we have that $|\partial_t g(x, t)| \lesssim \alpha r_i^{-1}$, where r_i is the radius of B_i . By the fundamental theorem of calculus (in t -variable) we see that the difference of the two averages in the term above can be bounded by

$$|\partial_t g| \gamma^2 \leq C\alpha r_i^{-1} \gamma^2.$$

It follows that

$$\begin{aligned} & \frac{1}{|B_r|} \sum_i \int_{B_i \cap B_r} \int_0^{\min\{r, r_i\}} \gamma^{-3} \left| \fint_{|y| \leq \gamma} g(x+y, t) - \fint_{B_\gamma(x, t)} g \right|^2 d\gamma dt dx \lesssim \\ & \frac{C\alpha^2}{|B_r|} \sum_i \int_{B_i \cap B_r} r_i^{-2} \int_0^{\min\{r, r_i\}} \gamma d\gamma \leq \frac{C\alpha^2}{|B_r|} \sum_i |B_i \cap B_r| \frac{\min\{r, r_i\}^2}{r_i^2} \lesssim \\ & \alpha^2 \sum_i \frac{|B_i \cap B_r|}{|B_r|} \leq N\alpha^2, \end{aligned}$$

using the finite overlap of the balls B_i .

It remains to consider the term II. We claim that for every $(x, t) \in \mathbb{R}^n$ we have a pointwise bound:

$$(6.36) \quad \int_{\min\{r, \rho(x, t)\}}^r \gamma^{-3} \left| g(x, t) - \fint_{B_\gamma(x, t)} f \right|^2 d\gamma \lesssim \alpha^2.$$

Consider first $(x, t) \in F = \mathbb{R}^n \setminus \Omega$. Then $M(S_1(f)^\beta)(x, t) \leq \alpha^\beta$ and hence in particular $S_1(f)(x, t) \leq \alpha$. Thus by (2.16) the above lefthand side is bounded by $S_1(f)(x, t)^2 \leq \alpha^2$ using the fact that $f(x, t) = g(x, t)$. Hence the claim holds. For $(x, t) \in \Omega$ the definition of the set Ω implies that there exists a point $(y, s) \in F$ whose parabolic distance to (x, t) is at most $4\rho(x, t)$. Again by (2.16) we have that

$$(6.37) \quad \int_{\min\{r, \rho(x, t)\}}^r \gamma^{-3} \left| g(y, s) - \fint_{B_\gamma(y, s)} f \right|^2 d\gamma \leq \alpha^2.$$

We need to compare (6.36) to (6.37). Recall that by (2.28) and the fact that f^\sharp can be used in place of g there, we get that

$$\left| \fint_{B_\gamma(y, s)} f - \fint_{B_\gamma(x, t)} f \right| \leq C\rho(x, t) \left[\fint_{B_\gamma(x, t)} f^\sharp + \fint_{B_\gamma(y, s)} f^\sharp \right].$$

Since $\gamma \geq \rho(x, t)$ we get that $B_\gamma(x, t) \subset B_{4\gamma}(y, s)$. At the same time using how Ω is defined we know that $M((f^\sharp)^\beta)(y, s) \leq \alpha^\beta$. Hence by Hölder inequality we see that

$$\fint_{B_\gamma(x, t)} f^\sharp + \fint_{B_\gamma(y, s)} f^\sharp \leq C \fint_{B_{4\gamma}(y, s)} f^\sharp \leq C \left(\fint_{B_{4\gamma}(y, s)} |f^\sharp|^\beta \right)^{1/\beta} \leq C\alpha.$$

Using the fact that g itself satisfies pointwise bound $|g^\sharp| \leq C\alpha$ we also have that $|g(x, t) - g(y, s)| \leq C\rho(x, t)\alpha$. It follows that

$$(6.38) \quad \int_{\min\{r, \rho(x, t)\}}^r \gamma^{-3} \left| g(x, t) - \fint_{B_\gamma(x, t)} f \right|^2 d\gamma \leq 4 \int_{\min\{r, \rho(x, t)\}}^r \gamma^{-3} \left| g(y, s) - \fint_{B_\gamma(y, s)} f \right|^2 d\gamma + 4C \int_{\min\{r, \rho(x, t)\}}^r \gamma^{-3} \rho(x, t)^2 \alpha^2 d\gamma.$$

The first term on the righthand side enjoys the bound (6.37), while the second integral is bounded by

$$4C\rho(x, t)^2 \alpha^2 [-\gamma^{-2}]_{\rho(x, t)}^\infty \lesssim \alpha^2.$$

Thus (6.36) indeed holds. It follows that to bound the term II of (6.34) by $C\alpha^2$ it suffices to establish that

$$(6.39) \quad \frac{1}{|B_r|} \int_{B_r} \int_{\min\{r, \rho(x, t)\}}^r \gamma^{-3} \left| \fint_{B_\gamma(x, t)} (f - g) \right|^2 d\gamma dt dx \leq C\alpha^2.$$

In order to understand the expression (6.39) recall first that $f = g$ outside Ω . Thus we need to understand the difference of $f - g$ only on the union of the balls B_i . Furthermore, recalling (6.29) we see that on Ω we have that $f = \sum_i f\chi_i$, while $g = \sum_i (\chi_i(B_i))^{-1} \int_{B_i} f\chi_i$ and hence if $B_i \subset B_\gamma(x, t)$ then

$$\fint_{B_\gamma(x, t)} f\chi_i = \fint_{B_\gamma(x, t)} (\chi_i(B_i))^{-1} \int_{B_i} f\chi_i \chi_i,$$

It follows that the i -th terms of the sum defining f and g contribute to the average over $B_\gamma(x, t)$ equally and thus these contributions cancel. Hence, in order to understand $\int_{B_\gamma(x, t)} (f - g)$ we only have to consider contributions of those B_i for which $B_i \cap \partial B_\gamma(x, t) \neq \emptyset$. Recalling (2.36) we see that

$$(6.40) \quad \left| \int_{B_\gamma(x, t)} (f - g) \right| \leq \frac{C}{|B_\gamma|} \sum_{i \in \{j: B_j \cap \partial B_\gamma(x, t)\}} r_i \int_{B_i} f^\# \leq \frac{C}{|B_\gamma|} \sum_{i \in \{j: B_j \cap \partial B_\gamma(x, t)\}} \alpha r_i |B_i|,$$

where in the last step we have used (6.23) and the Hölder's inequality. To further simplify our notation let us introduce the functions $A_i(x, t, \gamma)$ as follows:

$$(6.41) \quad A_i(x, t, \gamma) = \begin{cases} 1, & \text{if } B_i \cap \partial B_\gamma(x, t) \neq \emptyset, \\ 0, & \text{otherwise.} \end{cases}$$

It follows that (6.39) (by using (6.40) and the fact that $|B_\gamma| \approx \gamma^{n+1}$) enjoys the estimate:

$$\begin{aligned} & \frac{1}{|B_r|} \int_{B_r} \int_{\min\{r, \rho(x, t)\}}^r \gamma^{-3} \left| \int_{B_\gamma(x, t)} (f - g) \right|^2 d\gamma dt dx \lesssim \\ & \frac{\alpha^2}{|B_r|} \int_{B_r} \int_{\min\{r, \rho(x, t)\}}^r \gamma^{-(2n+5)} \left(\sum_i A_i r_i |B_i| \right)^2 d\gamma dt dx = \\ & \frac{N^2 \alpha^2}{|B_r|} \sum_i r_i^2 |B_i|^2 \int_{B_r} \int_{\min\{r, \rho(x, t)\}}^r A_i \gamma^{-(2n+5)} d\gamma dt dx + \\ & \frac{\alpha^2}{|B_r|} \sum_i \sum_{j \notin I(i)} r_i r_j |B_i| |B_j| \int_{B_r} \int_{\min\{r, \rho(x, t)\}}^r A_i A_j \gamma^{-(2n+5)} d\gamma dt dx = \\ & = III + IV. \end{aligned}$$

Here we have introduced notation $I(i) = \{j : B_i \cap B_j \neq \emptyset\}$ since we claim that in the last mixed term IV we only need to sum over pair of (i, j) that are separated. This is due to finite overlap property, when a point (x, t) belongs to multiple balls B_i these balls have comparable radius and there are at most N of them. Thus contribution of those terms to $(\sum_i A_i r_i |B_i|)^2$ is at most $N^2 A_i r_i^2 |B_i|^2$ for any such ball $B_i \ni (x, t)$. This also explains why the term III contains the term N^2 .

Let us first estimate the term III . Fix any i . If B_r is such that $r \leq r_i$ there is nothing to do since the integral in variable γ is only over the set where $\gamma \geq \rho(x, t) \geq r_i$ and $\gamma \leq r$. Thus this set is empty unless $r \geq r_i$. Similarly, unless $B_r \cap B_i \neq \emptyset$ there is nothing to do. It follows that

$$\begin{aligned} III & \lesssim \frac{\alpha^2}{|B_r|} \sum_{i: B_r \cap B_i \neq \emptyset} r_i^2 |B_i|^2 \int_{B_r} \int_{\min\{r, \rho(x, t)\}}^r A_i \gamma^{-(2n+5)} d\gamma dt dx \leq \\ & \lesssim \frac{\alpha^2}{|B_r|} \sum_{\{i: B_r \cap B_i \neq \emptyset \text{ \& } r \geq r_i\}} r_i^2 |B_i|^2 \int_{r_i}^\infty \gamma^{-(2n+5)} \left(\int_{B_r} A_i dx dt \right) d\gamma. \end{aligned}$$

We consider the integral $\int_{B_r} A_i dx dt$. Recall that the function A_i is nonzero only for those points (x, t) for which the parabolic distance between the point (x, t) and the

centre of the ball B_i is comparable to $\gamma \pm r_i$. The volume of such set is bounded by

$$|B_{\gamma+r_i}(x, t)| - |B_{\gamma-r_i}(x, t)| \leq C[(\gamma + r_i)^{n+1} - (\gamma - r_i)^{n+1}] \lesssim r_i \gamma^n.$$

Thus III can be further estimated by

$$III \lesssim \frac{\alpha^2}{|B_r|} \sum_{\{i: B_r \cap B_i \neq \emptyset \& r \geq r_i\}} r_i^3 |B_i|^2 \int_{r_i}^{\infty} \gamma^{-(n+5)} d\gamma \lesssim \frac{\alpha^2}{|B_r|} \sum_{\{i: B_r \cap B_i \neq \emptyset \& r \geq r_i\}} |B_i| \frac{r_i^{n+4}}{r_i^{n+4}}.$$

Finally, as the balls B_i have finite overlap and we only count balls intersecting B_r whose radius is less or comparable to B_r we get that $\sum_{\{i: B_r \cap B_i \neq \emptyset \& r \geq r_i\}} |B_i| \lesssim |B_r|$. Hence the estimate $III \lesssim \alpha^2$ follows.

We now look at IV . Fix i, j . As B_i, B_j are well separated we have that the parabolic distance between the centres of the balls is at least $2(r_i + r_j)$. Let us call this distance $2d$. Again we only have to consider pair of indices from the set $S = \{i : B_r \cap B_i \neq \emptyset \& r \geq r_i\}$ for the same reasons as outlined above.

It also follows that unless $\gamma \geq d$ we cannot have in the integral defining IV simultaneously $A_i = A_j = 1$ as the ball $B_\gamma(x, t)$ will fail to intersect at least one of the balls. Thus

$$IV \lesssim \frac{\alpha^2}{|B_r|} \sum_{i, j \in S \& j \notin I(i)} r_i r_j |B_i| |B_j| \int_d^\infty \gamma^{-(2n+5)} \left(\int_{B_r} A_i A_j dx dt \right) d\gamma.$$

We crudely estimate $\int_{B_r} A_i A_j dx dt \lesssim \min\{r_i, r_j\} \gamma^n \leq d \gamma^n$ since $A_i A_j \leq A_i$ and then use the earlier estimate. It follows that

$$IV \lesssim \frac{\alpha^2}{|B_r|} \sum_{i, j \in S \& j \notin I(i)} r_i r_j d |B_i| |B_j| \int_d^\infty \gamma^{-(n+5)} d\gamma \lesssim \frac{\alpha^2}{|B_r|} \sum_{i, j \in S \& d \geq r_i + r_j} \frac{r_i r_j |B_i| |B_j|}{d^{n+3}}.$$

We need to apply some simple geometric considerations to show finiteness of this sum. As now the separation condition is expressed as $d \geq r_i + r_j$ we fix i and then only sum over those balls $j \in S$ for which $r_j \geq r_i$. We further split the sum to $d \approx 2^k r_i$, $k = 0, 1, 2, \dots$. Hence

$$IV \leq \frac{\alpha^2}{|B_r|} \sum_{i \in S} r_i |B_i| \sum_{k=0}^{\infty} \sum_{j \in S, d(B_i, B_j) \approx 2^k r_i, r_j \geq r_i} \frac{r_j |B_j|}{(2^k r_i)^{n+3}}.$$

Actually, the sum in k will be only over those k for which $2^k r_i \leq r$. In the last sum since $r_j \leq d \approx 2^k r_i$ we have that

$$\sum_{j \in S, d(B_i, B_j) \approx 2^k r_i, r_j \geq r_i} \frac{r_j |B_j|}{(2^k r_i)^{n+3}} \leq \sum_{j \in S, d(B_i, B_j) \approx 2^k r_i, r_j \geq r_i} \frac{|B_j|}{(2^k r_i)^{n+2}}.$$

Given the finite overlap of B_j the sum of volumes of those B_j whose distance to B_i is approximately $2^k r_i$ can be at most multiple of the volume of the parabolic ball of radius $2^k r_i$. Hence

$$\sum_{j \in S, d(B_i, B_j) \approx 2^k r_i, r_j \geq r_i} \frac{r_j |B_j|}{(2^k r_i)^{n+3}} \leq \frac{1}{2^k r_i}.$$

It follows that

$$IV \lesssim \frac{\alpha^2}{|B_r|} \sum_{i \in S} r_i |B_i| \sum_{k=0}^{\infty} (2^k r_i)^{-1} \leq \frac{2\alpha^2}{|B_r|} \sum_{i \in S} |B_i| \leq 2N\alpha^2.$$

From this by (6.34) and combining the estimates for I and II we get that

$$\frac{1}{|B_r|} \int_{B_r} \int_0^r \gamma^{-3} \left| g(x, t) - \int_{B_\gamma(x, t)} g \right|^2 d\gamma dt dx \lesssim \alpha^2.$$

Hence $\mathbb{D}g \in \text{BMO}(\mathbb{R}^n)$ and therefore $D_t^{1/2}g \in \text{BMO}(\mathbb{R}^n)$ as desired. \square

Lemma 6.42. *Fix $1 < \beta \leq \infty$. There exists a large constant $M_\beta > 1$ such that any $(1, 1/2, \beta)$ -atom a can be decomposed as*

$$(6.43) \quad a = M_\beta a^\infty + \sum_i \lambda_i a_i,$$

such that a^∞ is an $(1, 1/2, \infty)$ -atom, all a_i are $(1, 1/2, \beta)$ -atoms and $\sum_i |\lambda_i| \leq 1/2$.

Hence for all $k \geq 1$, a can be written as

$$(6.44) \quad a = M_\beta \sum_i \mu_i a_i^\infty + \sum_i \lambda_i a_i,$$

such that all a_i^∞ are $(1, 1/2, \infty)$ -atoms, all a_i are $(1, 1/2, \beta)$ -atoms, $\sum_i |\mu_i| \leq 2 - 2^{-k+1}$ and $\sum_i |\lambda_i| \leq 2^{-k}$.

Proof. Clearly, the second claim follows from the first by induction, since each a_i atom in (6.43) can be again decomposed as a sum of $(1, 1/2, \infty)$ -atom and infinite sum of $(1, 1/2, \beta)$ -atoms. By relabelling the atoms, if necessary, we then obtain (6.44) for $k = 2$ and inductively for any larger k .

When $\beta = \infty$ there is nothing to prove. Suppose therefore that $1 < \beta < \infty$. To prove (6.43) we use the [Proposition 6.13](#) and apply it to the $(1, 1/2, \beta)$ -atom a . Since $a \in L_{1,1/2}^\beta(\mathbb{R}^n)$ we see that for some constant C_1 we have that

$$\begin{aligned} \int (|\nabla a| + |D_t^{1/2}a| + |S_1(a)| + a^\sharp)^\beta &\leq C_1 \|a\|_{L_{1,1/2}^\beta}^\beta = C_1 (\|\nabla a\|_{L^\beta} + \|D_t^{1/2}a\|_{L^\beta})^\beta \\ &\leq C_2 |B|^{-\beta/\beta'} = C_2 |B|^{1-\beta}. \end{aligned}$$

Assume that the atom a is supported in a parabolic ball B or radius $r(B)$. With the constants C and N as in (6.17) and (6.19) we let $\alpha = |B|^{-1} (2C^2 C_2 N)^{-1/(\beta-1)}$ and decompose our atom a as the sum $a = g + \sum b_i$ using [Proposition 6.13](#).

Recall the calculation for each b_i we have done previously in (6.25). It follows from it that

$$b = \sum_i b_i = \sum_i \lambda_i a_i, \quad \text{where each } a_i \text{ is an } (1, 1/2, \beta) \text{ atom and}$$

$$\sum_i |\lambda_i| = C\alpha \sum_i |B_i| \leq C^2 N \alpha^{1-\beta} \int (|\nabla a| + |D_t^{1/2}a| + |S_1(a)| + a^\sharp)^\beta \leq C^2 C_2 N \alpha^{1-\beta} |B|^{1-\beta}.$$

by (6.17) and calculation above for done for a . It follows that a choice of $\alpha = |B|^{-1}(2C^2C_2N)^{-1/(\beta-1)}$ makes the righthand side $\leq 1/2$ as desired. It remains to show that g can be seen as a multiple of a $(1, 1/2, \infty)$ atom.

Recall that it follows from the proof of [Proposition 6.13](#) that $g = a$ on the set where

$$M(|\nabla a|^\beta + |D_t^{1/2}a|^\beta + |S_1(a)|^\beta + |a^\sharp|^\beta)(X) \leq \alpha^\beta.$$

In particular consider $X \in \mathbb{R}^n$ such that the parabolic distance of X to the center of the ball B is at least $C_3r(B)$ for some $C_3 > 1$ to be determined.

Given that a is supported in B and the distance of X to the closest point of B is at least $C_3 - 1$ we see that

$$\begin{aligned} M(|\nabla a|^\beta)(X) &\leq |B_{(C_3-1)r(B)}(X)|^{-1} \int_{\mathbb{R}^n} |\nabla a|^\beta \leq K|B|^{-1}(C_3 - 1)^{-(n+1)}|B|^{-\beta/\beta'} \\ &\leq K(C_3 - 1)^{-(n+1)}|B|^{-\beta} \leq \frac{\alpha^\beta}{4} = |B|^{-\beta}(2C^2C_2N)^{-\beta/(\beta-1)} \end{aligned}$$

will hold, provided C_3 is chosen sufficiently large (and the choice does not depend on B). Similar calculation can be done for $D_t^{1/2}a$ taking into account the decay of the half-derivative away from the support of a as seen in [Lemma 6.2](#), as well as for $S_1(a)$ where by the formula (2.16) we see that

$$S_1(a)(X) \leq \left(\int_{(C_3-1)r(B)}^\infty \alpha^{-2n-5} \|a\|_{L^1}^2 d\alpha \right)^{1/2} \leq K \|a\|_{L^1} r(B)^{-1} |B|^{-1} (C_3 - 1)^{-(n+2)}.$$

Using (iii) of the definition of an atom this is further bounded by $K'|B|^{-1}(C_3 - 1)^{-(n+2)}$. Hence once again for sufficiently large $C_3 > 1$ independent of B we will have $S_1(a)^\beta(X) \leq \frac{\alpha^\beta}{4}$. Analogous calculation works for a^\sharp . Hence we conclude that for sufficiently large $C_3 > 1$ we have $g(X) = a(X) = 0$ for $X \notin C_3B$, where C_3B is the enlargement the original B . Thus g has support in C_3B .

Given this, we need to check the condition (ii) from the definition of an $(1, 1/2, \infty)$ -atom. By [Proposition 6.13](#) we have that $|\nabla g| \leq C\alpha$ and $\|D_t^{1/2}g\|_{\text{BMO}} \leq C\alpha$, and hence

$$\begin{aligned} \|\nabla g\|_{L^\infty(\mathbb{R}^n)} + \|D_t^{1/2}g\|_{\text{BMO}(\mathbb{R}^n)} &\leq C(2C^2C_2N)^{-1/(\beta-1)}|B|^{-1} \\ &\leq C(2C^2C_2N)^{-1/(\beta-1)}C_3^{n+1}|C_3B|^{-1}. \end{aligned}$$

This mean that if we set $M_\beta = C(2C^2C_2N)^{-1/(\beta-1)}C_3^{n+1}$ then $a^\infty = \frac{g}{M_\beta}$ is an $(1, 1/2, \infty)$ -atom with respect to the ball C_3B . From this our claim follows. \square

Lemma above implies that we may write any $(1, 1/2, \beta)$ -atom a as an infinite sum $\sum_i \lambda_i a_i^\infty$ of $(1, 1/2, \infty)$ -atoms a_i^∞ with $\sum_i |\lambda_i| \leq 2M_\beta$, where the equality $a = \sum_i \lambda_i a_i^\infty$ is understood to hold in the space $\dot{W}_{1,1/2}^1(\mathbb{R}^n)$. Hence we have the following as a consequence of [Lemma 6.6](#) and [Lemma 6.42](#).

Proposition 6.45. *For all $1 < \beta \leq \infty$ the Banach spaces $\dot{HS}_{1,1/2}^{1,(\beta)}(\mathbb{R}^n)$ are equal to each other with norms that are equivalent. Thus we may drop the index β and denote this space unambiguously by $\dot{HS}_{1,1/2}^1(\mathbb{R}^n)$.*

Thus as a consequence of this, [Proposition 6.21](#) and re-interpolation theorem [5] we have that

Proposition 6.46. *For all $1 < p < q < \infty$ and $\theta \in (0, 1)$ satisfying $\frac{1}{p} = (1 - \theta) + \frac{\theta}{q}$, $\dot{L}_{1,1/2}^p(\mathbb{R}^n)$ is a real interpolation space between $\dot{H}S_{1,1/2}^1(\mathbb{R}^n)$ and $\dot{L}_{1,1/2}^q(\mathbb{R}^n)$. More precisely, we have*

$$\left(\dot{H}S_{1,1/2}^1(\mathbb{R}^n), \dot{L}_{1,1/2}^q(\mathbb{R}^n) \right)_{\theta,p} = \dot{L}_{1,1/2}^p(\mathbb{R}^n),$$

with equivalent norms.

Corollary 6.47. *Let $\Omega = \mathcal{O} \times \mathbb{R}$, where $\mathcal{O} \subset \mathbb{R}^n$ is a bounded or unbounded Lipschitz domain. Then the conclusions of the above theorem remain to be true, that is for all $1 < p < q < \infty$ and $\theta \in (0, 1)$ satisfying $\frac{1}{p} = (1 - \theta) + \frac{\theta}{q}$, $\dot{L}_{1,1/2}^p(\partial\Omega)$ is a real interpolation space between $\dot{H}S_{1,1/2}^1(\partial\Omega)$ and $\dot{L}_{1,1/2}^q(\partial\Omega)$. More precisely, we have*

$$\left(\dot{H}S_{1,1/2}^1(\partial\Omega), \dot{L}_{1,1/2}^q(\partial\Omega) \right)_{\theta,p} = \dot{L}_{1,1/2}^p(\partial\Omega),$$

with equivalent norms.

This follows in the unbounded case directly from (2.37), and in the bounded case using a partition of unity by splitting $\partial\mathcal{O}$ to local coordinate patches.

6.2. Proof of Theorem 1.4. If we can prove the endpoint bound

$$(6.48) \quad \|N[\nabla u]\|_{L^1(\partial\Omega)} \lesssim \|f\|_{\dot{H}S_{1,1/2}^1(\partial\Omega)},$$

then Theorem 1.4 follows by real interpolation since the operator $f \mapsto N[\nabla u]$ is sub-linear. We follow the same approach as in the elliptic case outlined in [16]. It suffices to prove $\|N[\nabla u]\|_{L^1(\partial\Omega)} \leq C$ for $f = b$, where b is a $(1, 1/2, \infty)$ -atom supported on a boundary ball $\Delta_0 = \Delta_R(P_0, t_0)$. Note that $b = ca$, where a is a $(1, 1/2, p)$ -atom. Hence, by the solvability of $(R_L)_p$ we have that,

$$\begin{aligned} \|N[\nabla u]\|_{L^1(2\Delta_0)} &\leq |\Delta_0|^{1/p'} \|N[\nabla u]\|_{L^p(\partial\Omega)} \\ &\lesssim |\Delta_0|^{1/p'} (\|\nabla a\|_{L^p(\partial\Omega)} + \|D_t^{1/2} a\|_{L^p(\partial\Omega)}) \lesssim 1. \end{aligned}$$

Thus it remains to show that

$$(6.49) \quad \|N[\nabla u]\|_{L^1(\partial\Omega \setminus 2\Delta_0)} \lesssim 1.$$

By the fundamental theorem of calculus

$$|b| \lesssim R \|\nabla b\|_{L^\infty} \lesssim R^{2-n}.$$

Without loss of generality we may assume that $b \geq 0$ since by splitting b into positive and negative part we shall still have the L^∞ bound of both parts by R^{2-n} , which is the only property of b we will use below.

Next, we write $\partial\Omega \setminus 2\Delta_0 = \cup_{j \geq 3} A_j$, where

$$A_j := \{(P, s) \in \partial\Omega : \|(P, s) - (P_0, t_0)\| \approx 2^j R\}.$$

The sets A_j are annuli that can be covered by boundary balls Δ_j^i of finite overlap such that $r(\Delta_j^i) \approx 2^j R$, and on the enlargement $\tilde{\Delta}_j^i$ of Δ_j^i we have $u = 0$. This cover requires a finite fixed number of balls (that only depend on dimension n).

By the proof of [Theorem 4.1](#)

$$(6.50) \quad \int_{\Delta_j^i} N^{2^j R}[\nabla u] \lesssim (2^j R)^{-1} \left(\int_{T(4\Delta_j^i)} u^2 \right)^{1/2} \lesssim (2^j R)^{-1} u(V^+(\Delta_j^i)),$$

where $V^+(\Delta_j^i)$ is time-forward corkscrew point of the ball $4\Delta_j^i$. Give the positions of the balls $4\Delta_j^i$ the $u(V^+(\Delta_j^i))$ can further be controlled by [Proposition 3.8](#) which implies that we have $u(V^+(\Delta_j^i)) \lesssim u(P_0, 2^j R, t_0 + 100(2^j R)^2)$. By [\(3.12\)](#) and [Lemma 3.13](#)

$$\begin{aligned} u(P_0, 2^j R, t_0 + 100(2^j R)^2) &\lesssim R^{2-n} \omega^{(P_0, 2^j R, t_0 + 100(2^j R)^2)}(\Delta_0) \\ &\lesssim G(P_0, 2^j R, t_0 + 100(2^j R)^2, V^+(\Delta_0)) \lesssim R^\alpha (2^j R)^{-(n+\alpha)}. \end{aligned}$$

Using this in [\(6.50\)](#) implies that

$$(6.51) \quad \int_{\Delta_j^i} N^{2^j R}[\nabla u] \lesssim (2^j R)^{-1} R^\alpha (2^j R)^{-(n+\alpha)} \lesssim 2^{-j\alpha} (2^j R)^{-n-1}.$$

Thus

$$\int_{A_j} N^{2^j R}[\nabla u] \leq \sum_i (2^j R)^{n+1} \int_{\Delta_j^i} N^{2^j R}[\nabla u] \lesssim 2^{-j\alpha}.$$

So far we have an estimate for the truncated non-tangential maximal function at the height $2^j R$ for A_j . Consider any point $(X, t) \in \Gamma(S, \tau)$ for $(S, \tau) \in A_j$ and $\delta(X, t) \geq 2^j R$. By Caccioppoli and interior Harnack

$$\begin{aligned} \left(\int_{C_{\delta(X,t)/4}(X,t)} |\nabla u|^2 \right)^{1/2} &\lesssim (2^j R)^{-1} \left(\int_{C_{\delta(X,t)/2}(X,t)} |u|^2 \right)^{1/2} \\ &\lesssim (2^j R)^{-1} u(P_0, \delta(X, t), t_0 + 100\delta(X, t)^2). \end{aligned}$$

This can be further estimated by [\(3.12\)](#) and [Lemma 3.13](#) and taking a supremum over all points with $\delta(X, t) \geq 2^j R$ again yields a bound by $2^{-j\alpha} (2^j R)^{-n-1}$.

Hence by combining these two estimates we obtain that

$$\int_{A_j} N[\nabla u] \leq \int_{A_j} \tilde{N}^{2^j R}[\nabla u] + |A_j| (2^{-j\alpha} (2^j R)^{-n-1}) \lesssim 2^{-j\alpha}.$$

Summing over all $i \geq 3$ yields that

$$\int_{\partial\Omega \setminus 2\Delta_0} N[\nabla u] = \sum_{j \geq 3} \int_{A_j} N[\nabla u] \lesssim \sum_{j \geq 3} 2^{-j\alpha} \lesssim_\alpha 1.$$

In the proof above we have used [Theorem 4.1](#) which implies that the argument given above is fully correct only if the domain \mathcal{O} is unbounded. If \mathcal{O} is bounded we can only consider sets A_j up to the size of comparable to the diameter of \mathcal{O} , that is for $j \in \mathbb{N}$ such that $2^j R \lesssim \text{diam}(\mathcal{O})$. Say $d = \text{diam}(\mathcal{O})$ and let $\Delta_{10d} = \Delta_{10d}(P_0, t_0) \subset \partial\Omega$. Then the calculation above yields the estimate

$$\int_{\Delta_{10d} \setminus 2\Delta_0} N[\nabla u] \lesssim \sum_{j \geq 3}^{[\log_2(R^{-1}d)]+4} 2^{-j\alpha} \lesssim_\alpha 1.$$

The set of boundary points with distance more than $10d$ to (P_0, t_0) requires a separate argument. However, for such points [Lemma 5.10](#) does apply implying an exponential

decay in the L^1 norm of $N[\nabla u]$ and thus obvious summability. Therefore the claim as stated also holds in the case \mathcal{O} is bounded. \square

We record the statement (6.49) more formally as it is interesting on its own right and highlights what is the main obstacle preventing us from establishing solvability of the Regularity problem in the end-point $p = 1$ case. Compare this to Theorem 3.9 of [16] which states a similar result in the elliptic case.

Theorem 6.52. *Assume that $\omega \in A_\infty(d\sigma)$, where ω is the parabolic measure for the operator $L^* = \partial_t + \operatorname{div}(A^T \nabla \cdot)$ on a Lipschitz cylinder $\Omega = \mathcal{O} \times \mathbb{R}$. Then for any $p \in (1, \infty]$ the following holds. There exists $C(p) > 0$ such that for any homogeneous $(1, 1/2, p)$ -atom associated to a parabolic ball $\Delta \subset \partial\Omega$ (c. f. Definition 6.1), the weak solution u of the equation $Lu = -\partial_t u + \operatorname{div}(A \nabla u) = 0$ with boundary datum f satisfies the estimate*

$$(6.53) \quad \|N[\nabla u]\|_{L^1(\partial\Omega \setminus 2\Delta)} \leq C(p).$$

Proof. The proof as given above holds for $(1, 1/2, \infty)$ -atoms. It uses boundedness of such atom and thus does not apply to $(1, 1/2, p)$ -atoms for $p < \infty$. But we may assume that $f \geq 0$ as it can be split as the sum of $f^+ - f^-$ with each part being modulo a constant again an atom. Thanks to Lemma 6.42 we know that any $(1, 1/2, p)$ -atom f , $p > 1$ can be written as as sum

$$f = \sum_i \mu_i a_i^\infty,$$

where a_i^∞ are $(1, 1/2, \infty)$ -atoms and $\sum_i |\mu_i| \leq M_p$ for some large constant $M_p > 1$ independent of f . Furthermore, thinking about the support of the atoms a_i^∞ the construction we gave in the proof of Lemma 6.42 implies that that a_i^∞ will be supported in some enlargement of Δ , say $m\Delta$ for some $m > 1$ which depends of the choice of α in the proof of Lemma 6.42. Thus if we do not make any changes in the argument given we will obtain for each a_i^∞ the estimate (6.49), i.e.:

$$(6.54) \quad \|N[\nabla u_i]\|_{L^1(\partial\Omega \setminus 2m\Delta)} \leq K,$$

making the constant in the estimate explicit. Here u_i solves our PDE with boundary data a_i^∞ . It follows that

$$(6.55) \quad \|N[\nabla u]\|_{L^1(\partial\Omega \setminus 2m\Delta)} \leq \sum_i |\mu_i| \|N[\nabla u_i]\|_{L^1(\partial\Omega \setminus 2m\Delta)} \leq M_p K.$$

This is not quite (6.53) as we have removed a larger set from $\partial\Omega$, namely $2m\Delta$ instead of just 2Δ as claimed. However, this is only a small technical issue. Thinking about the set $2m\Delta \setminus 2\Delta$ we are in the regime where f vanishes, it actually vanishes on an enlargement of this set as well as it clearly vanishes on $4m\Delta \setminus \Delta$.

Hence we get that Proposition 3.4 applies as well as the boundary Hölder continuity (3.10). It follows that u is bounded on the union of non-tangential cones

$$\mathcal{S} := \bigcup_{(q, \tau) \in 3m\Delta \setminus (3/2)\Delta} \Gamma(q, \tau),$$

where the bound for the near part (i.e. points $(X, t) \in \mathcal{S}$ with $\delta(X, t) \lesssim \operatorname{diam}(\Delta) =: R$) is implied by (3.10) which requires bound on value of u at some corkscrew point V^+ of $5m\Delta$ but that bound is implied by The boundary Poincaré inequality (Proposition

3.1) and the bound (6.55) as the point V^+ is in a region covered by the estimates for $N[\nabla u]$ on the set $\partial\Omega \setminus 2m\Delta$. The bound for the away part of the set \mathcal{S} is easier and follows directly from (6.55).

Hence we can claim for all $(X, t) \in \mathcal{S}$ that $|u| \lesssim R(R^{-n-1})\|N[\nabla u]\|_{L^1(\partial\Omega \setminus 2m\Delta)} = R^{-n}M_pK$. Here the normalization factor R^{-n-1} comes from the fact that R^{n+1} is comparable to the surface area of $m\Delta$.

The final piece of the puzzle is Theorem 4.1 which we apply on balls B_i that cover the boundary $2m\Delta \setminus 2\Delta$ of radius comparable to R such that $4B_i \subset 3m\Delta \setminus (3/2)\Delta$ and balls B_i have finite overlap. Thanks to the bound for $|u|$ on \mathcal{S} we have by (4.2) on each B_i

$$\int_{B_i} N[\nabla u] \lesssim \iint_{T(2B_i)} |\nabla u| \lesssim R^{-1} \left(\iint_{T(3B_i)} |u|^2 \right)^{1/2} \lesssim R^{-1}R^{-n}M_pK = R^{-n-1}M_pK,$$

where in the second step we have used the boundary Caccioppoli inequality (Proposition 3.4). Equivalently, $\int_{B_i} N[\nabla u] \lesssim M_pK$ and this after summing over all i we get that

$$\|N[\nabla u]\|_{L^1(2m\Delta \setminus 2\Delta)} \leq CM_pK.$$

From (6.55) and the above estimate (6.53) follows. \square

REFERENCES

- [1] D. G. Aronson, *Non-negative solutions of linear parabolic equations*, Annali della Scuola Normale Superiore di Pisa - Classe di Scienze **22** (1968), no. 4, 607–694. [18](#)
- [2] N. Badr, F. Bernicot, *Abstract Hardy-Sobolev spaces and interpolation*, J. Funct. Anal. 259 (2010), no. 5, 1169–1208. [28](#), [30](#)
- [3] C. Bennett, A. DeVore, R. Sharpley, *Weak-type inequalities for H^p and BMO*, Proc. Symp. Pure Math. XXXV (1) (1979), 201–229. [31](#)
- [4] C. Bennett, R. Sharpley, *Interpolation of Operators*, Academic Press, 1988. [30](#), [32](#)
- [5] J. Bergh, J. Löfström, *Interpolation Spaces, An Introduction*, Springer, Berlin, 1976. [30](#), [32](#), [43](#)
- [6] R. Brown, *Layer Potentials and Boundary Value Problems for the Heat Equation on Lipschitz Cylinders*, Thesis (Ph.D.)—University of Minnesota ProQuest LLC, Ann Arbor, MI, 1987. 94 pp. [2](#)
- [7] R. Brown, *The method of layer potentials for the heat equation in Lipschitz cylinders*, Amer. J. Math. 111 (1989), no. 2, 339–379. [2](#), [3](#), [13](#)
- [8] A. P. Calderón, A. Torchinsky, *Parabolic Maximal Functions Associated with a Distribution I*, Adv. Math., 16(1):1–64, 1975. [10](#)
- [9] A. Castro, S. Rodríguez-López, W. Staubach, *L^2 -solvability of the Dirichlet, Neumann and regularity problems for parabolic equations with time-independent Hölder-continuous coefficients*, Trans. Amer. Math. Soc. 370 (2018), no. 1, 265–319. [2](#), [3](#), [13](#)
- [10] M. Christ, G. Weiss, *A $T(b)$ theorem with remarks on analytic capacity and the Cauchy integral*, Colloquium Mathematicae **60-61** Issue 2 (1977), 601–628. [20](#)
- [11] R. Coifman, G. Weiss, *Extensions of Hardy spaces and their use in analysis*, Bull. Amer. Math. Soc. **83** (1977), 569–645. [33](#)
- [12] M. Dindoš, *On the regularity problem for parabolic operators and the role of half-time derivative*, arXiv:2308.12936. [4](#), [13](#)
- [13] M. Dindoš, L. Dyer, *Parabolic regularity and Dirichlet boundary value problems*, Nonlinear Anal. 185 (2019), 247–263. [2](#), [3](#), [7](#)
- [14] M. Dindoš, L. Dyer, S. Hwang, *Parabolic L^p Dirichlet boundary value problem and VMO-type time-varying domains*, Anal. PDE 13 (2020), no.4, 1221–1268. [8](#), [9](#), [10](#)

- [15] M. Dindoš, S. Hofmann, J. Pipher, *Regularity and Neumann problems for operators with real coefficients satisfying Carleson conditions*, Journal of Functional Analysis 285 (2023), no.6, 110024. [3](#)
- [16] M. Dindoš, J. Kirsch, *The regularity problem for elliptic operators with boundary data in Hardy-Sobolev space HS^1* , Math. Res. Lett.19 (2012), no.3, 699–717. [2](#), [4](#), [44](#), [46](#)
- [17] E. Fabes, M. Jodeit, *L^p boundary value problems for parabolic equations*, Bull. Amer. Math. Soc. 74 (1968), 1098–1102 [6](#)
- [18] E. Fabes, N. Rivière, *Dirichlet and Neumann problems for the heat equation in C^1 -cylinders*, Proc. Sympos. Pure Math., XXXV, Part 2, AMS Providence, RI, 1979, pp. 179–196. [2](#)
- [19] E. Fabes, N. Rivière, *Singular integrals with mixed homogeneity*, Studia Math. 27 (1966), 19–38. [7](#)
- [20] E. Fabes, N. Rivière, *Symbolic calculus of kernels with mixed homogeneity*, Singular Integrals (Proc. Sympos. Pure Math., Chicago, Ill., 1966), pp. 106–127, AMS, Providence, RI, 1967 [6](#), [7](#)
- [21] C. Fefferman, E. M. Stein, *H^p Spaces of Several Variables*, Acta Math., 129(1):137–193, 1972. [10](#)
- [22] A. Genschaw, S. Hofmann, *A Weak Reverse Hölder Inequality for Caloric Measure*, J. Geom. Anal. (2019). [16](#), [17](#)
- [23] A. Genschaw, S. Hofmann, J. Martel, K. Nyström, *Potential Theory in One-Sided Parabolic Chord Arc Domains*, work in progress. [16](#), [17](#)
- [24] S. Hofmann, J. Lewis, *L^2 solvability and representation by caloric layer potentials in time-varying domains*, Ann. of Math. (2) 144 (1996), no. 2, 349–420. [6](#)
- [25] S. Hofmann, J. Lewis, *The L^p regularity problem for the heat equation in non-cylindrical domains*, Hofmann, Steven; Lewis, John L. Illinois J. Math. 43 (1999), no. 4, 752–769. [6](#)
- [26] S. Hofmann, J. Lewis, *Dirichlet Problem for Parabolic Operators with Singular Drift Terms*, Providence American Mathematical Society (2001). [16](#), [17](#)
- [27] B. Jawerth, A. Torchinski, *Local Sharp Maximal Functions*, J. of Approx. Th. 43, 231–270 (1985). [31](#)
- [28] C. Kenig, J. Pipher, *The Neumann problem for elliptic equations with non-smooth coefficients*, Invent. Math. 113 (1993) no. 1, 447–509. [2](#)
- [29] C. Kenig, J. Pipher, *The Neumann problem for elliptic equations with non-smooth coefficients II*, Duke Math. J. 81 (1995), no. 1, 227–250. [3](#)
- [30] J. Kinnunen, H. Tuominen, *Pointwise behaviour of $M^{1,1}$ Sobolev functions*, Math. Z. 257 (2007), no. 3, 613–630. [11](#)
- [31] S. Kim, L. Xu, *Green’s function for second order parabolic equations with singular lower order coefficients*, Arxiv (2021). [18](#)
- [32] P. Koskela, D. Yang, Y. Zhou, *A characterization of Hajlasz-Sobolev and Triebel-Lizorkin spaces via grand Littlewood-Paley functions*, J. Funct. Anal. 258 (2010), no. 8, 2637–2661. [11](#), [12](#)
- [33] M. Mitrea, *The initial Dirichlet boundary value problem for general second order parabolic systems in nonsmooth manifolds*, Comm. Partial Differential Equations 26 (2001), no. 11–12, 1975–2036. [2](#), [3](#), [13](#)
- [34] A. Monguzzi, M. Peloso, M. Salvatori, *Fractional Laplacian, homogeneous Sobolev spaces and their realizations*, Ann. Mat. Pura Appl. (4) 199 (2020), no. 6, 2243–2261. [8](#)
- [35] M. Mourgoglou, B. Poggi, X. Tolsa, *L^p solvability of the Poisson-Dirichlet problem and its applications to the Regularity problem*, arXiv:2207.10554. [12](#), [13](#)
- [36] K. Nyström, *Boundary Value Problems and Duality between L^p Dirichlet and Regularity Problems for Second Order Parabolic Systems in Non-Cylindrical Domains*, Collect. Math.57 (2006), no.1, 93–119. [2](#), [3](#), [6](#), [13](#)
- [37] Z. Shen, *A relationship between the Dirichlet and Regularity Problems for elliptic equations*, Mathematical research letters, 14, 205–213 (2007). [2](#), [3](#), [4](#), [18](#), [22](#), [24](#)
- [38] Z. Shen, *Extrapolation for the L^p Dirichlet Problem in Lipschitz Domains*, Acta Mathematica Sinica, 35, 1074–1084 (2019) no. 6. [17](#)
- [39] E. M. Stein, *Singular Integrals and Differentiability Properties of Functions*, Princeton Univ. Press, Princeton, N.J., 1970. [10](#)

- [40] R. Strichartz, *Bounded Mean Oscillation and Sobolev Spaces*, Indiana Univ. Math. J., 29(4):539–558, 1980. [8](#), [9](#), [10](#), [37](#)
- [41] M. Ulmer. *Perturbation Theory for the Parabolic Regularity Problem*, Preprint arXiv:2408.12529. [16](#), [17](#)
- [42] M. Ulmer, *L^p Boundary Value Problems for Elliptic and Parabolic Operators*, University of Edinburgh, thesis (2024). [16](#), [17](#)
- [43] D. Yang, W. Yuan, Y. Zhou, *A new characterization of Triebel-Lizorkin spaces on \mathbb{R}^n* , Publ. Mat. 57 (2013), no. 1, 57–82. [8](#), [9](#)