

EXISTENCE OF 5 MINIMAL TORI IN 3-SPHERES OF POSITIVE RICCI CURVATURE

ADRIAN CHUN-PONG CHU AND YANGYANG LI

ABSTRACT. In 1989, B. White conjectured that every Riemannian 3-sphere has at least 5 embedded minimal tori. We confirm this conjecture for 3-spheres of positive Ricci curvature. While our proof uses min-max theory, the underlying heuristics are largely inspired by mean curvature flow.

CONTENTS

1. Introduction	1
Part I. Main arguments	12
2. Preliminaries	12
3. Proof of Theorem 1.1	20
Part II. Technical ingredients	38
4. Min-max results I: The Simon-Smith min-max theorem	38
5. The family Ψ	54
6. Perturbing the metric	66
7. Min-max results II: Convergence as currents	71
8. Min-max results III: Interpolation	74
9. Pinch-off process and mean curvature flow	92
10. Topology of genus one cap	94
References	101

1. INTRODUCTION

While enumerative geometry is traditionally a branch of algebraic geometry, there are, in fact, enumerative problems in differential geometry as well.

In 1905, Poincaré conjectured the existence of 3 simple closed geodesic in every smooth, Riemannian 2-sphere [Poi05]. The number 3 is optimal, as is shown in the example of an ellipsoid by Morse [Mor96]. While G. D. Birkhoff [Bir17] showed there is always at least one simple closed geodesic using a min-max argument, a full solution of Poincaré's conjecture was given

Date: September 17, 2024.

by M. Grayson in 1989 using curve shortening flow [Gra89], building upon the previous work of Lyusternik–Schnirelmann [LS29].

In one dimension higher, S.-T. Yau [Yau82] conjectured that there exist at least 4 embedded minimal 2-spheres in every Riemannian 3-sphere. Simon-Smith [Smi83] showed there is always at least one embedded minimal 2-sphere using min-max theory. More recently, Wang-Zhou [WZ24] confirmed Yau’s conjecture for the case of metrics with positive Ricci curvature and bumpy metrics (see also the work of Haslhofer-Ketover [HK19]). Furthermore, B. White [Whi89, Whi91] conjectured that every Riemannian 3-sphere admits at least 5 embedded minimal tori. In fact, he proved that every 3-sphere of positive Ricci curvature has at least 1 embedded minimal torus using a degree theory he developed and Hamilton’s Ricci flow.

In this paper, we confirm B. White’s conjecture for 3-spheres with positive Ricci curvature using min-max theory:

Theorem 1.1. *Every smooth Riemannian 3-sphere of positive Ricci curvature has at least 5 embedded minimal tori.*

Let us briefly explain where the number 5 comes from. The well-known Willmore conjecture [Wil65, Wil71], proved by Marques-Neves [MN14], implies that in the unit 3-sphere (S^3, \bar{g}) , the Clifford torus is the embedded minimal surface with the second smallest area, after the equator. Moreover, the resolution of Lawson’s conjecture on minimal tori [Law70] by S. Brendle [Bre13] indicates that the Clifford torus is the only embedded minimal torus in (S^3, \bar{g}) . Consequently, the space of embedded minimal tori in (S^3, \bar{g}) is exactly the space \mathcal{C} of Clifford tori, which is homeomorphic to $\mathbb{RP}^2 \times \mathbb{RP}^2$. If we now perturb the round metric \bar{g} , the area functional defined on \mathcal{C} changes. On the other hand, the *Lyusternik-Schnirelmann number* of $\mathbb{RP}^2 \times \mathbb{RP}^2$ is 5, meaning every smooth real function on $\mathbb{RP}^2 \times \mathbb{RP}^2$ has at least 5 critical points. This is due to the theorem that the Lyusternik-Schnirelmann number of a space is always strictly greater than its *cup length* in \mathbb{Z}_2 -coefficients (which is the maximal number of first cohomology classes such that their cup product is non-zero), and the topological fact that the cup length of $\mathbb{RP}^2 \times \mathbb{RP}^2$ is 4. Relying on these facts, B. White proved that every 3-sphere equipped with a metric sufficiently close to the round metric has at least 5 embedded minimal tori. It is worth noting that the topological fact that the cup length of $\mathbb{RP}^2 \times \mathbb{RP}^2$ is 4 will play a crucial role in our proof of Theorem 1.1.

For more works on the construction of geodesics or minimal surfaces with controlled topological type¹, see also [GJ86, HK23, Ko23a, Ko23b, Str84, Zho16].

¹During the preparation of this paper, we were informed by another group, Xingzhe Li and Zhichao Wang, that they have also been working on related problems, though we have yet to learn about any content of their work as of this arXiv submission.

1.1. Heuristics for the proof. Let us describe the heuristics behind the proof of Theorem 1.1. Note that while the heuristics rely on mean curvature flow, we would not use mean curvature flow at all in practice: Instead, we primarily use Simon-Smith min-max theory, and also some concepts from Almgren-Pitts min-max theory.

Let (S^3, g_0) be the 3-sphere of positive Ricci curvature given in Theorem 1.1. In their proof of the Willmore conjecture, Marques-Neves defined a 5-parameter family of currents, known as the *canonical family*, associated to each closed embedded surface in the unit 3-sphere (described with more details in §1.2.1). Based on their construction, we define in (S^3, g_0) a 9-parameter family Ψ of surfaces of genus 1 or 0, possibly with singularities.

Now, let us apply mean curvature flow to each member of Ψ . Following the heuristics from gradient flow in the finite-dimensional setting, we assume all such mean curvature flows exist uniquely for all time until the surfaces vanish. Although this assumption is likely false in reality due to the possibility of *fattening* (i.e. non-unique solutions of mean curvature flow), we will ignore this complication in our heuristic argument. Thus, as $t \rightarrow +\infty$, we expect the family Ψ to evolve into some family Ξ that is “optimal”, in the sense that Ξ contains minimal tori or minimal spheres, as well as *eternal or ancient mean curvature flows connecting these minimal surfaces to one another or to the empty set*. These mean curvature flows act as gradient flow lines connecting critical points (see Figure 1). Note that we assume all minimal tori or minimal spheres appearing in Ξ have multiplicity one: This assumption is heuristically justified, as it is conjectured that with the positive Ricci curvature condition, long time limits of mean curvature flow have multiplicity one [CS24, Conjecture 1.3].




FIGURE 1. The figure on the left shows the original family Ψ . The figure on the right shows Ξ , with the minimal surfaces S_1, S_2, S_3 detected, and ancient mean curvature flow lines originating from them. The red part is $T(S_1)$, the green part is $T(S_2)$, and the blue part is $T(S_3)$.

For each minimal surface S detected by Ξ , there are eternal or ancient mean curvature flows originating from S . Let $T(S)$ denote the collection of all time slices in such mean curvature flows. Note that $T(S)$ is a subset

of Ξ , and can be viewed as the “unstable manifold” for S (see Figure 1). Moreover, $\Xi = \cup_S T(S)$ where S ranges over all minimal surfaces (of genus 0 or 1) detected by Ξ , and if S is a minimal sphere, then each element of $T(S)$ should have genus 0 possibly with singularities, as mean curvature flow do not increase genus [BK24].

To prove Theorem 1.1, suppose for the sake of contradiction that there are only 4 minimal tori T_1, T_2, T_3, T_4 detected by Ξ . The case with fewer than 4 minimal tori can be handled similarly. Then, letting $D_0 := \cup_S T(S)$ where S ranges over the minimal *spheres* detected by Ξ , and $D_i := T(T_i)$ for $i = 1, \dots, 4$, we have

$$(1.1) \quad \Xi = D_0 \cup D_1 \cup D_2 \cup D_3 \cup D_4.$$

Now, let us describe a crucial topological fact about Ψ . In fact, ensuring this topological fact holds is the main reason we constructed Ψ using the Marques-Neves canonical family. Recall that the cycle space $\mathcal{Z}_2(S^3; \mathbb{Z}_2)$ is weakly homotopy equivalent to $\mathbb{R}P^\infty$, and thus, the cohomology ring $H^*(\mathcal{Z}_2(S^3; \mathbb{Z}_2); \mathbb{Z}_2)$ can be written as $\mathbb{Z}_2[\bar{\lambda}]$, where $\bar{\lambda}$ denotes the unique non-trivial element in $H^1(\mathcal{Z}_2(S^3; \mathbb{Z}_2); \mathbb{Z}_2)$. Since each member of Ψ can be viewed as a mod 2 cycle, we can view Ψ as a subset of $\mathcal{Z}_2(S^3; \mathbb{Z}_2)$. Let λ be the element in $H^1(\Psi; \mathbb{Z}_2)$ induced by $\bar{\lambda}$. Then, the crucial topological fact is that, there exist $\alpha, \beta \in H^1(\Psi; \mathbb{Z}_2)$ such that

$$(1.2) \quad \lambda^5 \cup \alpha^2 \cup \beta^2 \neq 0.$$

By construction, Ξ and Ψ are homotopic as sweepouts, so we can view λ, α and β as elements of $H^1(\Xi; \mathbb{Z}_2)$ as well. Hence, by an elementary Lyusternik–Schnirelmann argument, (1.1) and (1.2) together immediately imply that one of the following must hold:

- (1) The element $\lambda^5|_{D_0}$ in $H^5(D_0; \mathbb{Z}_2)$ induced by λ^5 is non-zero.
- (2) The element $\alpha|_{D_1}$ in $H^1(D_1; \mathbb{Z}_2)$ induced by α is non-zero.
- (3) The element $\alpha|_{D_2}$ in $H^1(D_2; \mathbb{Z}_2)$ induced by α is non-zero.
- (4) The element $\beta|_{D_3}$ in $H^1(D_3; \mathbb{Z}_2)$ induced by β is non-zero.
- (5) The element $\beta|_{D_4}$ in $H^1(D_4; \mathbb{Z}_2)$ induced by β is non-zero.

And to derive a contradiction, it suffices to show that statements (1) to (5) are all impossible.

To show that (1) is impossible, first note that by construction, D_0 consists entirely of genus 0 surfaces, possibly with singularities. Now, let us consider D_0 as a family in the unit round 3-sphere \mathbb{S}^3 , instead of the original 3-sphere (S^3, g_0) . Note the following two facts:

- If we apply the Simon-Smith min-max theorem to D_0 in \mathbb{S}^3 , then by Wang-Zhou’s multiplicity one theorem [WZ24], we detect the multiplicity-one equatorial 2-sphere, which has area 4π .
- On the other hand, if (1) holds, meaning $\lambda^5|_{D_0} \neq 0$, then by definition, D_0 is a 5-sweepout in the sense of the Almgren-Pitts min-max theory. Since the Simon-Smith width must be greater than or equal to the Almgren-Pitts width, the Simon-Smith width of the family

D_0 (viewed in \mathbb{S}^3) is at least the 5-width of \mathbb{S}^3 , which is $2\pi^2$ by C. Nurser [Nur16].

The above two facts are contradictory, so (1) is impossible.

To show that (2) is impossible, note that by backtracking the mean curvature flow lines, D_1 can be deformation retracted to the single minimal torus $\{T^1\}$. Thus, $H^1(D_1; \mathbb{Z}_2)$ is zero, making (2) is impossible. Similarly, (3) through (5) are also impossible. This concludes the proof of Theorem 1.1.

1.2. Details behind the heuristics. The above heuristic argument contains several details and inaccuracy that need to be addressed.

1.2.1. *The 9-parameter family Ψ .* Let us now explain the construction of the family Ψ . Let \mathcal{C} (resp. $\tilde{\mathcal{C}}$) denote the set of unoriented (resp. oriented) Clifford tori in the unit 3-sphere \mathbb{S}^3 . Note that $\mathcal{C} \cong \mathbb{RP}^2 \times \mathbb{RP}^2$, and $\tilde{\mathcal{C}}$ is a double cover of this space. For each $\Sigma \in \tilde{\mathcal{C}}$, based on Marques-Neves' canonical family, C. Nurser in [Nur16] constructed a 5-sweepout

$$\Phi_5^\Sigma : \mathbb{RP}^5 \rightarrow \mathcal{Z}_2(\mathbb{S}^3; \mathbb{Z}_2).$$

Here is his construction: The set of conformal diffeomorphisms of \mathbb{S}^3 can be parametrized by the open unit 4-ball \mathbb{B}^4 , which we denote by $\{F_v\}_{v \in \mathbb{B}^4}$. For each image $F_v(\Sigma)$, which is oriented, we consider the level surfaces of the *signed* distance function to $F_v(\Sigma)$. This gives us a 5-parameter family of surfaces, possibly with singularities and possibly empty, parametrized by $\mathbb{B}^4 \times [-\pi, \pi]$ (since the set of possible signed distances from $F_v(\Sigma)$ lie in the range $[-\pi, \pi]$).

Marques-Neves [MN14, §5] showed that by ingeniously reparametrizing this family *near* the boundary of its parameter space, one can continuously extend this family to the boundary, obtaining a $\mathbb{B}^4 \times [-\pi, \pi]$ -family. Moreover, any two “antipodal” points on the boundary of this parameter space can be identified, as they represent the same surface (but with opposite orientation), thereby yielding an \mathbb{RP}^5 -family Φ_5^Σ .

Repeating this for every oriented Clifford torus Σ , we obtain an $\mathbb{RP}^5 \times \tilde{\mathcal{C}}$ -family. The subfamilies Ψ_5^Σ and $\Psi_5^{-\Sigma}$ have the same images for each $\Sigma \in \tilde{\mathcal{C}}$, where $-\Sigma$ denotes Σ with an opposite orientation. By identifying all such pairs of subfamilies Ψ_5^Σ and $\Psi_5^{-\Sigma}$, we obtain a family Ψ with parameter space Y , which is an \mathbb{RP}^5 -bundle over $\mathcal{C} \cong \mathbb{RP}^2 \times \mathbb{RP}^2$. This family consists only of genus 0 and genus 1 surfaces, possibly with singularities, and is suitable for running the Simon-Smith min-max process.

1.2.2. *Overcoming the issue of fattening.* In the heuristic argument of §1.1, we proposed using mean curvature flow to obtain an “optimal family” Ξ from Ψ . In practice, the authors were unable to implement this strategy due to the phenomenon of fattening. Indeed, if any member of Ψ fatten along the mean curvature flow, then the flow no longer generates a canonical deformation that is continuous across the entire family Ψ of initial conditions. As a

result, it is unclear how one might obtain the desired family Ξ using this approach.

Instead of using mean curvature flow, we employ a scheme of repeatedly applying the Simon-Smith min-max to Ψ in order to obtain Ξ . Furthermore, we will define a process called *pinch-off process* that resembles mean curvature flow, so that each member $\Psi(x)$ will deform under this pinch-off process to some member $\Xi(x)$. Crucially, this process is genus non-increasing in time. More precisely, a pinch-off process involves a finite number of neck-pinch surgeries in small balls, so that the surface no longer intersects the boundary of the ball, and then shrinking the components inside the ball into a point. Note that this process would not create new conical singularities (which cause the issue of fattening for mean curvature flow). The “topology-decreasing” nature of pinch-off process (which will be discussed in §9) should be compared with the analogous results on mean curvature flow by B. White [Whi95]. At the end, although the family Ξ we obtain may not be as well-behaved as to be considered “optimal” in the sense described in §1.1, it is sufficient for our purposes.

One might ask why we need the pinch-off process. Indeed, solely by the usual Simon-Smith min-max theorem, one already could homotope the initial family Ψ to some other family that is, a certain sense, optimal. The problem is the following. When this optimal family detects a minimal *sphere*, let us consider the “cap” region of this optimal family that is near the minimal sphere. Every member of this cap would only be varifold-close to the minimal sphere, and thus could actually have genus 1, but with some short non-trivial loop. As we will see soon in §1.2.3, this would be problematic (and this deviates from the heuristic picture in §1.1 too). Hence, we need to use the pinch-off process, to pinch all the short non-trivial loops, and make the cap consist of only genus 0 surfaces.

We will explain below in greater details why the pinch-off process is necessary, by actually constructing the family Ξ we use in practice.

1.2.3. Obtaining Ξ through repeated applications of the min-max theorem. First, we assume that the 3-sphere (S^3, g_0) given has only finitely many minimal tori. Furthermore, we show that one can perturb the metric g_0 to assume that it has only finitely many minimal spheres, all of which are non-degenerate and have linearly independent areas over \mathbb{Z} , with no linear combination over \mathbb{Z} of their areas equal to the area of any minimal torus. In the first stage of min-max process, we apply Simon-Smith min-max to Ψ , homotoping it obtain a new pulled-tight “optimal” family $\tilde{\Psi}^1$ that detects finitely many min-max minimal surfaces of genus 0 or 1. For simplicity, we will assume the new parameter space $\text{dmn}(\tilde{\Psi}^1)$ is also Y .

In the new perturbed metric, using D. Ketover’s strengthened genus bound [Ket19] and Wang-Zhou’s multiplicity one theorem [WZ24], which builds on Sarnataro-Stryker’s work [SS23], it is not hard to see that there

are two possible cases about the detected minimal surfaces: (1) each min-max minimal surface is a multiplicity one minimal torus, or (2) only one min-max minimal surface is detected, and it is a multiplicity-one sphere. Note that the Frankel property is needed, which is ensured by the assumption of positive Ricci curvature. For simplicity, in case (1), we assume that there is only one minimal torus.

In both cases, we consider a subset $C^1 \subset \text{dmn}(\tilde{\Psi}^1)$ such that the restriction $\tilde{\Psi}^1|_{C^1}$ is close to the minimal surface V^1 detected. The family $\tilde{\Psi}^1|_{C^1}$ is like a “cap” of large area, such that its complement $\Psi^1 := \tilde{\Psi}^1|_{\overline{Y} \setminus C^1}$ has the property that the maximal area of Ψ^1 is strictly less than the width of Ψ , $\text{area}(V^1)$. Let Y^1 denote the new domain $\text{dmn}(\Psi^1) = \overline{Y} \setminus C^1$.

For case (2) only, we will prove certain interpolation results that allow us, without loss of generality, to assume that $\tilde{\Psi}^1|_{C^1}$ consists entirely of genus 0 surfaces, possibly with singularities: *This is where the pinch-off process described in §1.2.2 comes into place.*

Now, we proceed to the second stage of the min-max process, this time applied to Ψ^1 , regardless of whether case (1) or (2) occurred previously. The width at this stage will be strictly smaller than the previous width, leading to some new minimal surface. We repeat this process iteratively.

Namely, at the k -th stage of the min-max process, we start with a family Ψ^{k-1} with parameter space Y^{k-1} . Applying the Simon-Smith min-max theorem to Ψ^{k-1} would produce a family H^k with parameter space $[0, 1] \times Y^{k-1}$, resembling a homotopy, such that $H^k(0, \cdot) = \Psi^{k-1}$, and the new family $\tilde{\Psi}^k := H^k(1, \cdot)$ (with parameter space Y^{k-1}) is an “optimal” family detecting a multiplicity-one minimal surface V^k of genus 0 or 1.

Let $C^k \subset Y^{k-1}$ be a “cap” such that $\tilde{\Psi}^k|_{C^k}$ has large area and is near V^k . As before, if V^k has genus 0, we can assume, by applying pinch-off process, that $\tilde{\Psi}^k|_{C^k}$ consists entirely of genus 0 surfaces, possibly with singularities. We then consider the new parameter space $Y^k := \overline{Y^{k-1}} \setminus C^k$, and proceed with the $(k + 1)$ -th stage of min-max process, by applying the Simon-Smith min-max theorem to the family $\Psi^k := \tilde{\Psi}^k|_{\overline{Y^{k-1}} \setminus C^k}$. For simplicity, we assume that each Y^k is a 9-dimensional manifold with boundary.

Since we have perturbed the metric g_0 so that all minimal spheres are non-degenerate, there must be only finitely many of them, by Choi-Schoen’s compactness result [CS85]. Consequently, since the width of Ψ^k decreases strictly with each iteration, the repetitive min-max process must terminate. Namely, there exists a $K > 0$ such that at the K -th stage of min-max process, the width of Ψ^{K-1} is zero. At this point, there exists a family H^K (which acts like a homotopy) with parameter space $[0, 1] \times Y^{K-1}$, where $H^K(0, \cdot) = \Psi^{K-1}$, and $\tilde{\Psi}^K := H^K(1, \cdot)$ has a maximal area arbitrarily close to zero.

Finally, let us construct the desired “optimal” family Ξ . Consider the following list of $2K - 1$ Simon-Smith families of genus ≤ 1 . For the sake of visualization, readers may find it helpful to refer to Figure 2.

- (1) $\tilde{\Psi}^1|_{C^1}$
- (2) $H^2|_{[0,1] \times \partial Y^1}$
- (3) $\tilde{\Psi}^2|_{C^2}$
- (4) $H^3|_{[0,1] \times \partial Y^2}$
- ...
- ($2K - 4$) $H^{K-1}|_{[0,1] \times \partial Y^{K-2}}$
- ($2K - 3$) $\tilde{\Psi}^{K-1}|_{C^{K-1}}$
- ($2K - 2$) $H^K|_{[0,1] \times \partial Y^{K-1}}$
- ($2K - 1$) Ψ^K




FIGURE 2. Constructing Ξ .

Immediately from the definition of these $2K - 1$ families, there exists a natural way to glue their boundaries together, as illustrated in Figure 2. Through this process, we obtain a new 9-parameter Simon-Smith family Ξ that, in some sense, is homotopic to the original family Ψ .

1.2.4. *Decomposing Ξ into “unstable manifolds”.* In the heuristic argument in §1.1, we decomposed Ξ into a union of “unstable manifolds” $T(S)$ associated with each minimal surface S detected in Ξ . There are two issues with this approach.

First, the family Ξ we obtained in §1.2.3, when viewed as a subset of the cycle space $\mathcal{Z}_2(S^3; \mathbb{Z}_2)$, may have complicated topology and thus be difficult

to work with from an algebraic topology perspective. Thus, in practice we actually work with its parameter space $\text{dmn}(\Xi)$, and decompose $\text{dmn}(\Xi)$ instead.

Second, our family Ξ no longer consists of mean curvature flow lines. Instead of defining $T(S)$, we are going to define the *trace* $T(C)$ of each cap $C = C^1, C^2, \dots, C^{K-1}$, and each trace $T(C)$ will be a subset of $\text{dmn}(\Xi)$.

Let us fix a cap $C = C^k$, obtained at the k -th stage of the min-max process. Consider the following sets, *which can all be viewed as subsets of $\text{dmn}(\Xi)$* (see Figure (3)):

- $B_k := C$,
- $B_{k+1} := [0, 1] \times \partial Y^k = \text{dmn}(H^{k+1})$,
- $B_{k+2} := [0, 1] \times \left((\{1\} \times \partial Y^k) \cap (\{0\} \times \partial Y^{k+1}) \right) \subset \text{dmn}(H^{k+2})$. Here, the subsets $\{1\} \times \partial Y^k, \{0\} \times \partial Y^{k+1} \subset \text{dmn}(\Xi)$ can both also be viewed as subsets of $\text{dmn}(\tilde{\Psi}^{k+1})$
- $B_{k+3} := [0, 1] \times \left((\{1\} \times \partial Y^{k+1}) \cap (\{0\} \times \partial Y^{k+2}) \right) \subset \text{dmn}(H^{k+3})$.
- ...

Finally, we define the trace of C by

$$T(C) := B_k \cup B_{k+1} \cup \dots \cup B_K \subset \text{dmn}(\Xi).$$

A crucial property of $T(C)$ is that, by definition, $T(C)$ admits a strong deformation retraction back onto the cap C .




FIGURE 3. The trace $T(C)$ in $\text{dmn}(\Xi)$.

In fact, if $C = C^k$ is a cap associated with a genus 0 minimal surface, then by construction $\Xi|_{T(C)}$ consists entirely of genus 0 surfaces, possibly with singularities, because $\tilde{\Psi}^k|_{C^k}$ does and the “homotopies” given by H^{k+1}, H^{k+2}, \dots do not increase genus (this should be compared with the situation in §1.1, where mean curvature flow does not increase genus).

Now, let N be the number of minimal tori we detected throughout the first $K - 1$ stages of the min-max process. To prove Theorem 1.1, we only need to show that $N \geq 5$.

Let us see why $N = 4$ is impossible; the cases for $N = 0, \dots, 3$ are similar. Given a cap $C = C^k$, we say that it has *genus 0* if the minimal surface V^k associated has genus 0, and *genus 1* if V^k has genus 1. For example, when $N = 4$, we have four genus 1 caps, and finitely many genus 0 caps. We can express $\text{dmn}(\Xi)$ as the union of the following five subsets:

- Let D_0 be the union of $\text{dmn}(\Psi^K)$ and $\cup_C T(C)$, where C ranges over all the genus 0 caps.
- For each genus 1 cap C , we consider its trace $T(C)$. There are in total 4 such traces, which we will denote by D_1, \dots, D_4 .

It follows directly from the definition of Ξ that

$$\text{dmn}(\Xi) = D_0 \cup D_1 \cup D_2 \cup D_3 \cup D_4.$$

1.2.5. Topological arguments. With this setup, we can proceed with the remaining steps of the heuristic argument in §1.1. However, instead of considering the cohomology rings of Ψ and Ξ , we should focus on those of Y and $\text{dmn}(\Xi)$. For example, we should instead define λ as an element of $H^1(Y; \mathbb{Z}_2)$, by $\lambda := \Psi^*(\bar{\lambda})$, where Ψ is viewed as a map into $\mathcal{Z}_2(S^3; \mathbb{Z}_2)$. The crucial topological fact is that, there exist $\alpha, \beta \in H^1(Y; \mathbb{Z}_2)$ such that

$$(1.3) \quad \lambda^5 \cup \alpha^2 \cup \beta^2 \neq 0.$$

Moreover, since Y and $\text{dmn}(\Xi)$ are actually homotopy equivalent, λ, α and β can be viewed as elements of $H^1(\text{dmn}(\Xi); \mathbb{Z}_2)$ as well.

Then, as in §1.1, we apply a Lyusternik-Schnirelmann argument to derive a contradiction. The proof that $\lambda^5|_{D_0} = 0$ proceeds similarly to the earlier argument. However, proving $\alpha|_{D_i} = 0$ for $i = 1, \dots, 4$ requires more essential modifications. For instance, unlike in §1.1, D_1 may not be contractible (let us focus on $i = 1$, as the cases of $i = 2, 3, 4$ are the same). Thus, we would instead prove $\alpha|_{D_1} = 0$ by showing that the map

$$H_1(D_1; \mathbb{Z}_2) \rightarrow H_1(\text{dmn}(\Xi); \mathbb{Z}_2)$$

induced by the inclusion $D_1 \hookrightarrow \text{dmn}(\Xi)$ is trivial.

Recall that D_1 admits a strong deformation retraction onto the genus 1 cap C it corresponds to. Therefore, it suffices to show that for any loop $c \subset C$, $[c] = 0$ in $H_1(\text{dmn}(\Xi); \mathbb{Z}_2)$. To achieve this, we show that the cap C can be assumed to be close in the \mathbf{F} -metric for currents to the minimal torus corresponding to C . Consequently, the restriction $\Xi|_c$ actually gives a family of genus 1 surfaces with singularities. Using the homotopy between $\text{dmn}(\Xi)$ and Y , we can deform $\Xi|_c$ back to some subfamily $\Psi|_{c_0}$ of Ψ , for some loop $c_0 \subset Y = \text{dmn}(\Psi)$, while ensuring that throughout the homotopy, *all generalized surfaces are of genus 1*. This is possible because the process of obtain the final family Ξ from the original family Ψ is, in some sense, genus non-increasing (similar to running mean curvature flow). In particular, it suffices to show that the loop $c_0 \subset Y$ is homologically trivial.

Let $Z_0 \subset Y$ be the set of parameters corresponding to (unoriented) Clifford tori under Ψ . From the definition of Ψ and the fact that all members of $\Psi|_{c_0}$ have genus one, it is not hard to show that one can deform c_0 into Z_0 . Thus, it suffices to show that c_0 is homologically trivial in Z_0 . Note that $Z_0 \cong \mathbb{RP}^2 \times \mathbb{RP}^2$, so $H_1(Z_0; \mathbb{Z}_2)$ consists four elements: $(0, 0), (0, 1), (1, 0), (1, 1)$.

First, using the fact that $\Xi|_c$ is close to a single minimal torus, $[c_0]$ cannot be $(0, 1)$ or $(1, 0)$, as both correspond to 1-sweepouts under Ψ . As for $(1, 1)$, one can check that it gives rise to, under Ψ , a loop of *oriented* Clifford tori, $\{\Sigma(\theta)\}_{\theta \in [0, 2\pi]}$, such that the followings hold:

- We can continuously pick an interior region in $\text{in}(\Sigma(\theta))$, bounded by $\Sigma(\theta)$, for each θ .
- While the interior regions in $\text{in}(\Sigma(0))$ and $\text{in}(\Sigma(2\pi))$ coincide, the isomorphism map from $\pi_1(\text{in}(\Sigma(0)))$ to $\pi_1(\text{in}(\Sigma(2\pi)))$ (both groups are \mathbb{Z}) induced by the motion $\{\Sigma(\theta)\}_{\theta \in [0, 2\pi]}$ is not the identity but -1 .

We would show that this is impossible, due to the fact that $\Xi|_c$ is close to a single minimal torus, and that $\Psi|_{c_0}$ was obtained from $\Xi|_c$ through a homotopy which is a family consisting of *genus one* surfaces possibly with singularities.

Hence, $[c_0]$ must be the trivial element $(0, 0)$, as desired. This means $\alpha|_{D_1} = 0$. Similarly, $\alpha|_{D_2}, \beta|_{D_3}, \beta|_{D_4}$ are all trivial. This leads to a contradiction, as shown in §1.1. This completes the proof of Theorem 1.1.

1.3. Open problems. First of all, the full conjecture of B. White is still open. However, in light of the work of [WZ24], we are more optimistic about the existence of at least five embedded minimal tori for a generic metric, though this number might not be sharp for generic metrics, as discussed below.

Recall that the Morse number of a manifold is the minimum number of critical points a Morse function can have. Since $\mathbb{RP}^2 \times \mathbb{RP}^2$ has Morse number 9, B. White [Whi91] conjectured the existence of at least 9 embedded minimal tori in a 3-sphere equipped with a generic metric.

Inspired by the works of [MN16, Li23a, Li23b], and as a further inquiry based on the current work, we conjecture that in every 3-sphere with positive Ricci curvature, there exist at least 5 embedded minimal tori of Morse index at most 9.

It is also natural to raise the question of counting minimal surfaces with higher genus. For instance, let \mathcal{M}_2 denote the space of embedded, unoriented, genus 2 Lawson surfaces in the unit 3-sphere. This space is a closed 6-dimensional manifold. In the spirit of [Whi91], we conjecture that in every Riemannian 3-sphere, the number of closed, embedded, genus 2 minimal surfaces is at least one plus the cup length of \mathcal{M}_2 . This conjecture is, of course, based on the fact that the Lusternik–Schnirelmann number of a manifold is bounded below by one plus the cup length. We also conjecture that in every Riemannian 3-sphere equipped with a generic metric, the number of

closed, embedded, genus 2 minimal surfaces is at least the Morse number of \mathcal{M}_2 .

Finally, we pose the following questions as potential steps towards resolving the fattening issue described in §1.2.2 from a purely mean curvature flow perspective. Given a smooth, embedded, closed surface Σ in a 3-manifold, let \mathfrak{M} denote the union of all time slices of all mean curvature flows with initial data Σ , in whichever appropriate weak sense. Is the set \mathfrak{M} connected? More ambitiously, is \mathfrak{M} contractible?

If \mathfrak{M} is contractible, it may be possible to run mean curvature flow to the whole family Ψ and surpass the time of fattening. (See also the recent work of J. Bernstein, L. Chen, and L. Wang regarding the set of expanders flowing out of a cone [BW22, BCW24].)

1.4. Organization. This paper is divided in two main parts. In part I, we first present the preliminary results in §2, which primarily concern about various min-max theories, and then prove the main theorem in §3. In part II, we prove various theorems and propositions used in part I.

Acknowledgment. We are deeply grateful to Zhihan Wang for his generous help, which has been instrumental in the completion of this project. We also wish to thank André Neves for his constant support and guidance.

This material is based upon work supported by the National Science Foundation under Grant No. DMS-1928930, while the first author was in residence at the Simons Laufer Mathematical Sciences Institute (formerly MSRI) in Berkeley, California, during the Fall semester of 2024. The second author was partially supported by the AMS-Simons travel grant.

Part I. Main arguments

2. PRELIMINARIES

Throughout this paper, all the ambient manifolds M we consider are assumed to be smooth and closed.

2.1. Notations.

- $\mathbf{I}_n(M; \mathbb{Z}_2)$: the set of integral n -dimensional currents in M with \mathbb{Z}_2 -coefficients.
- $\mathcal{Z}_n(M; \mathbb{Z}_2) \subset \mathbf{I}_n(M; \mathbb{Z}_2)$: the subset that consists of elements T such that $T = \partial Q$ for some $Q \in \mathbf{I}_{n+1}(M; \mathbb{Z}_2)$ (such T are also called *flat k -cycles*).
- $\mathcal{Z}_n(M; \nu; \mathbb{Z}_2)$ with $\nu = \mathcal{F}, \mathbf{F}, \mathbf{M}$: the set $\mathcal{Z}_n(M; \mathbb{Z}_2)$ equipped with the three topologies given corresponding respectively to the *flat norm* \mathcal{F} , the *\mathbf{F} -metric*, and the *mass norm* \mathbf{M} (see [Pit81] and the survey paper [MN20]). For the flat norm, there are two definitions that, by the isoperimetric inequality, would induce the same topology:

$$\mathcal{F}(T) := \inf\{\mathbf{M}(P) + \mathbf{M}(Q) : T = P + \partial Q\}$$

and

$$\mathcal{F}(T) := \inf\{\mathbf{M}(Q) : T = \partial Q\}.$$

In this paper, we will use the second definition.

- $\mathcal{V}_n(M)$: the closure, in the varifold weak topology, of the space of k -dimensional rectifiable varifolds in M .
- $\mathcal{C}(M)$: the space of Caccioppoli sets in M , equipped with the metric induced by the Lebesgue measure of the symmetric difference.
- $\partial^*\Omega$: the reduced boundary of $\Omega \in \mathcal{C}(M)$.
- ν_Ω : the inward pointing normal of $\Omega \in \mathcal{C}(M)$.
- $\|V\|$: the Radon measure induced on M by $V \in \mathcal{V}_n(M)$.
- $|T|$: the varifold in $\mathcal{V}_n(M)$ induced by a current $T \in \mathcal{Z}_n(M; \mathbb{Z}_2)$, or a countable n -rectifiable set T . In the same spirit, given a map Φ into $\mathcal{Z}_n(M; \mathbb{Z}_2)$, the associated map into $\mathcal{V}_n(M)$ is denoted by $|\Phi|$.
- $\text{spt}(\cdot)$: the support of a current or a measure.
- $[W]$: the \mathbb{Z}_2 -current induced by W , if W is a countably 2-rectifiable set with $\mathcal{H}^2(W) < \infty$. In the same spirit, given a map f whose images are countably 2-rectifiable sets, the associated map into the space $\mathcal{Z}_n(M; \mathbb{Z}_2)$ is denoted by $[f]$.
- $[\mathcal{W}] := \{[\Sigma_i]\} \subset \mathcal{Z}_n(M; \mathbb{Z}_2)$ for a set \mathcal{W} of varifolds $\{V_i\} \subset \mathcal{V}_n(M)$, each associated with a countably n -rectifiable set Σ_i .
- $\mathbf{B}_\varepsilon^\nu(\cdot)$: the open ε -neighborhood of an element or a subset of the space $\mathcal{Z}_n(M; \nu; \mathbb{Z}_2)$.
- $\mathbf{B}_\varepsilon^{\mathbf{F}}(\cdot)$: the open ε -neighborhood of an element or a subset of $\mathcal{V}_n(M)$ under the \mathbf{F} -metric.
- $\Gamma^\infty(M)$: the set of smooth Riemannian metrics on M .
- $B_r(p)$: the open r -neighborhood of a point p .
- $\mathfrak{g}(S)$: the genus of a surface S .
- $\text{in}(S)$: the inside, open region of an oriented surface S .
- $\text{out}(S)$: the outside, open region of an oriented surface S .

For an m -dimensional cube $I^m = \mathbb{R}^m \cap \{x : 0 \leq x_i \leq 1, i = 1, 2, \dots, m\}$, we can give it *cubical complex structures* as follows.

- $I(1, j)$: the cubical complex on $I := [0, 1]$ whose 1-cells and 0-cells are respectively

$$[0, 1/3^j], [1/3^j, 2/3^j], \dots, [1 - 1/3^j, 1] \quad \text{and} \quad [0], [1/3^j], [2/3^j], \dots, [1].$$

- $I(m, j)$: the cubical complex structure

$$I(m, j) = I(1, j) \otimes \dots \otimes I(1, j) \quad (m \text{ times})$$

on I^m . $\alpha = \alpha_1 \otimes \dots \otimes \alpha_m$ is a q -cell of $I(m, j)$ if and only if each α_i is a cell in $I(1, j)$ and there are exactly q 1-cells. A cell β is a *face* of a cell α if and only if $\beta \subset \alpha$ as sets.

We call $X \subset I(m, j)$ a *cubical subcomplex* of $I(m, j)$ if every face of a cell in X is also a cell in X . For convenience, we also call X a *cubical complex* without referring to the ambient cube. We denote by $|X|$ the *underlying*

space of X . For the sake of simplicity, we will also consider a complex and its underlying space as identical unless there is ambiguity.

Given a cubical subcomplex X of some $I(m, j)$, for $j' > j$, one can *refine* X to a cubical subcomplex

$$X(j') := \{\sigma \in I(m, j') : \sigma \cap |X| \neq \emptyset\}$$

of $I(m, j')$. For the sake of convenience, we will denote the refined cubical subcomplex by X unless there is ambiguity.

2.2. Simon-Smith min-max theory. Let (M, g) be a closed Riemannian 3-manifold.

Definition 2.1 (Generalized surfaces). A closed subset $\Sigma \subset M$ is called a *generalized surface* provided that:

- (1) The 2-dimensional Hausdorff measure $\mathcal{H}^2(\Sigma) \in (0, \infty)$.
- (2) There exists a finite set $P \subset \Sigma$ such that $\Sigma \setminus P$ is an orientable, smooth, embedded surface.
- (3) Let Σ_{iso} be the set of isolated points of Σ . Then $\Sigma \setminus \Sigma_{\text{iso}}$ is the topological boundary of some open region of M .

We denote the space of generalized surfaces in M by $\mathcal{S}(M)$.

For each $\mathfrak{g}_0 \in \mathbb{N}$ (the set of non-negative integers), $\Sigma \in \mathcal{S}(M)$ is called a *generalized surface of genus \mathfrak{g}_0* provided that for a finite set P satisfying (2) above and for a sequence $r_i \rightarrow 0$,

$$\mathfrak{g}(\Phi(x)) := \lim_{r \rightarrow 0} \mathfrak{g}(\Phi(x) \setminus B_{r_i}(P(x))) = \mathfrak{g}_0.$$

Additionally, we denote

$$\mathcal{S}^*(M) = \mathcal{S}(M) \cup \{\Sigma \subset M : \Sigma \text{ is a closed subset, } \mathcal{H}^2(\Sigma) = 0\}.$$

Remark 2.2. For any $\Sigma \in \mathcal{S}(M)$ and a finite set P satisfying (2), by Sard's theorem, there exists a set of full measure $E \subset (0, \infty)$ such that for every $r \in E$, $\Sigma \setminus B_r(P)$ is a smooth surface with boundary.

A surface with smooth boundary can be obtained from a closed surface by removing finitely many disjoint disks. The genus of this surface with boundary is the genus of the original closed surface. Also note that when a surface is disconnected, its genus is defined as the sum of the genus of each of its connected components.

Moreover, for $r_1 > r_2 > 0$ such that both $\Sigma \setminus B_{r_i}(P)$ are smooth surfaces with boundary, by [CM12, Chapter 1 § 2.1 Lemma 1.5], we have the inequality on their genus,

$$\mathfrak{g}(\Phi(x) \setminus B_{r_1}(P(x))) \leq \mathfrak{g}(\Phi(x) \setminus B_{r_2}(P(x))).$$

Hence, the limit

$$(2.1) \quad \lim_{E_x \ni r \rightarrow 0} \mathfrak{g}(\Sigma \setminus B_r(P)) \in \mathbb{N} \cup \{\infty\}.$$

always exists.

In particular, the definition of $\mathfrak{g}(\Sigma)$ is independent of the choice of the sequence r_i . In addition, it is also easy to verify that the definition is independent of the choice of the finite set P satisfying (2).

Definition 2.3 (Simon-Smith family). Let X be a cubical subcomplex of some $I(m, j)$. A map $\Phi : X \rightarrow \mathcal{S}^*(M)$ is called a *Simon-Smith family*, provided that:

- (1) For each $x \in X$ with $\Phi(x) \in \mathcal{S}(M)$, we can choose a finite set $P(x) \subset \Phi(x)$ satisfying Definition 2.1 (2) such that

$$N_P(\Phi) := \sup_{x: \Phi(x) \in \mathcal{S}(M)} |P(x)| < \infty.$$

- (2) The composition map $x \mapsto \mathcal{H}^2 \circ \Phi(x)$ is continuous.
- (3) For any $x_0 \in X$ and any open set $U \supset \Phi(x_0)$, there exists a neighborhood $O \subset X$ of x_0 such that for any $x \in O$, $\Phi(x) \subset U$.
- (4) For any $x_0 \in X$ with $\Phi(x_0) \in \mathcal{S}(M)$, on any open set $U \subset \subset M \setminus P(x_0)$, $\Phi(x) \rightarrow \Phi(x_0)$ smoothly whenever $x \rightarrow x_0$.

In this case, we call X the *parameter space* of Φ .

In addition, for a non-negative integer \mathfrak{g}_0 , we call Φ a *Simon-Smith family of genus $\leq \mathfrak{g}_0$* , if Φ also satisfies:

- (5) For each $x \in X$ with $\Phi(x) \in \mathcal{S}(M)$, $\Phi(x)$ has genus at most \mathfrak{g}_0 .

Remark 2.4. In the literature, there exist various notions of *generalized family of surfaces* in the Simon-Smith min-max setting. Our definition of Simon-Smith family can be viewed as a generalization of those in the following sense:

Our continuity conditions (2), (3) and (4) are essentially the same as those in [DLR18a], but we also allow measure-zero sets in the family. Our genus bound condition (5) generalizes the definitions in [DLP10, CFS22, Fra21] without assuming a dense set of smooth surfaces in the family.

Proposition 2.5 (Simon-Smith families are sweepouts).

- (1) Every element Σ in $\mathcal{S}^*(M)$ is associated with a unique 2-cycle $[\Sigma]$ in $\mathcal{Z}_2(M; \mathbb{Z}_2)$. In particular, if $\mathcal{H}^2(\Sigma) = 0$, then $[\Sigma] = 0$.
- (2) Moreover, for a Simon-Smith family Φ , the induced map

$$[\Phi] : X \rightarrow \mathcal{Z}_2(M; \mathbf{F}; \mathbb{Z}_2)$$

is continuous with respect to the \mathbf{F} -metric.

Remark 2.6. (2) implies that our Simon-Smith family induces a sweepout in the Almgren-Pitts min-max theory.

Proof of Proposition 2.5. By Definition 2.1, for $\Sigma \in \mathcal{S}(M)$, we can choose a finite set $P \subset \Sigma$ such that $\Sigma \setminus P$ is a smooth surface with finite \mathcal{H}^2 measure, so we can define

$$[\Sigma] := [\Sigma \setminus P] \in \mathcal{Z}_2(M; \mathbb{Z}_2).$$

Note that the definition is independent of the choice of P as long as it satisfies the condition (2).

For $\mathcal{H}^2(\Sigma) = 0$, we can simply define

$$[\Sigma] := 0 \in \mathcal{Z}_2(M; \mathbb{Z}_2).$$

For (2), for any $x \in X$ and any sequence $\{y_i\} \subset X$ with $\lim_i y_i = x$, it suffices to prove that $[\Phi(y_i)] \rightarrow [\Phi(x)]$ with respect to the \mathbf{F} -metric. There are two cases: (1) $\mathcal{H}^2(\Phi(x)) = 0$, or (2) $\mathcal{H}^2(\Phi(x)) > 0$.

Case 1: $\mathcal{H}^2(\Phi(x)) = 0$. In this case,

$$\mathbf{M}([\Phi(y_i)]) = \mathcal{H}^2(\Phi(y_i)) \rightarrow \mathcal{H}^2(\Phi(x)) = 0,$$

and thus, $[\Phi(y_i)] \rightarrow 0 = [\Phi(x)]$ with respect to the \mathbf{F} -metric.

Case 2: $\mathcal{H}^2(\Phi(x)) > 0$. In this case, we can assume that every $\Phi(y_i) \in \mathcal{S}(M)$ and choose a finite set P satisfying Condition (2) in Definition 2.1. For any $\varepsilon > 0$, by the fact that $\mathcal{H}^2(\Phi(x)) < \infty$ and Sard's theorem, there exists $r > 0$ such that

$$(2.2) \quad \mathcal{H}^2(\Phi(x) \cap \overline{B}_{2r}(P)) < \frac{\varepsilon}{4},$$

and $\partial B_{2r}(P)$ intersects transversally with $\Phi(x)$ and every $\Phi(y_i)$.

By Definition 2.3 (4), $\Phi(y_i) \rightarrow \Phi(x)$ smoothly outside $\overline{B}_r(P)$. In particular, we have:

- Outside $\overline{B}_{2r}(P)$,

$$(2.3) \quad \lim_i \mathbf{F}([\Phi(y_i) \setminus \overline{B}_{2r}(P)], [\Phi(x) \setminus \overline{B}_{2r}(P)]) = 0.$$

- Inside $\overline{B}_{2r}(P)$,

$$(2.4) \quad \begin{aligned} & \limsup_i \mathcal{H}^2(\Phi(y_i) \cap \overline{B}_{2r}(P)) \\ &= \limsup_i \left(\mathcal{H}^2(\Phi(y_i)) - \mathcal{H}^2(\Phi(y_i) \setminus \overline{B}_{2r}(P)) \right) \\ &= \lim_i \mathcal{H}^2(\Phi(y_i)) - \lim_i \mathcal{H}^2(\Phi(y_i) \setminus \overline{B}_{2r}(P)) \\ &= \mathcal{H}^2(\Phi(x)) - \mathcal{H}^2(\Phi(x) \setminus \overline{B}_{2r}(P)) \\ &= \mathcal{H}^2(\Phi(x) \cap \overline{B}_{2r}(P)) < \frac{\varepsilon}{4}. \end{aligned}$$

Note that by the triangle inequality, we have

$$\begin{aligned} & \mathbf{F}([\Phi(y_i)], [\Phi(x)]) \\ & \leq \mathbf{F}([\Phi(y_i)], [\Phi(y_i) \setminus \overline{B}_{2r}(P)]) + \mathbf{F}([\Phi(x) \setminus \overline{B}_{2r}(P)], [\Phi(x)]) \\ & \quad + \mathbf{F}([\Phi(y_i) \setminus \overline{B}_{2r}(P)], [\Phi(x) \setminus \overline{B}_{2r}(P)]) \\ & = \mathbf{F}([\Phi(y_i) \cap \overline{B}_{2r}(P)], 0) + \mathbf{F}(0, [\Phi(x) \cap \overline{B}_{2r}(P)]) \\ & \quad + \mathbf{F}([\Phi(y_i) \setminus \overline{B}_{2r}(P)], [\Phi(x) \setminus \overline{B}_{2r}(P)]) \\ & \leq 2\mathcal{H}^2(\Phi(y_i) \cap \overline{B}_{2r}(P)) + 2\mathcal{H}^2(\Phi(x) \cap \overline{B}_{2r}(P)) \\ & \quad + \mathbf{F}([\Phi(y_i) \setminus \overline{B}_{2r}(P)], [\Phi(x) \setminus \overline{B}_{2r}(P)]) \end{aligned}$$

where the first identity follows from the definition of \mathbf{F} -metric on $\mathcal{Z}_2(M; \mathbb{Z}_2)$. Applying (2.2), (2.3) and (2.4), for any $\varepsilon > 0$, we obtain

$$\limsup_i \mathbf{F}([\Phi(y_i)], [\Phi(x)]) \leq \varepsilon$$

and thus, $\lim_i \mathbf{F}([\Phi(y_i)], [\Phi(x)]) = 0$. □

Definition 2.7 (Homotopy class). Two Simon-Smith families Φ_0 and Φ_1 parametrized by the same parameter space X are said to be *homotopic* to each other if there exists a continuous map

$$\varphi : [0, 1] \times X \rightarrow \text{Diff}^\infty(M)$$

such that

- $\varphi(0, x) = \text{Id}$ for all $x \in X$,
- $\varphi(1, x)(\Phi_0(x)) = \Phi_1(x)$ for all $x \in X$.

Note that $\text{Diff}^\infty(M)$ denotes the diffeomorphism group of M equipped with the C^∞ topology.

The set of all families homotopic to a Simon-Smith family Φ is called the *homotopy class associated to Φ* , and is denoted by $\Lambda(\Phi)$.

Remark 2.8. The family

$$H : [0, 1] \times X \rightarrow \mathcal{S}^*(M), \quad (t, x) \mapsto \varphi(t, x)(\Phi(x))$$

is also a Simon-Smith family. In particular, for every $t \in [0, 1]$,

$$\Phi_t : X \rightarrow \mathcal{S}^*(M), \quad x \mapsto \varphi(t, x)(\Phi(x))$$

is a Simon-Smith family in $\Lambda(\Phi_0)$.

Furthermore, for any choice of

$$P : \{x \in X : \Phi(x) \in \mathcal{S}(M)\} \rightarrow M$$

for Φ satisfying the conditions (1) and (4) in Definition 2.3, the corresponding homotopy φ induces a choice of P for every $\Phi' \in \Lambda(\Phi)$. Consequently, we may assume that $N_P < \infty$ is a constant within the homotopy class $\Lambda(\Phi)$.

In addition, if Φ is of genus $\leq \mathfrak{g}_0$, then so is every $\Phi' \in \Lambda(\Phi)$.

Definition 2.9 (Simon-Smith width). Given a homotopy class Λ , its *width* is defined by

$$\mathbf{L}(\Lambda) := \inf_{\Phi \in \Lambda} \sup_{x \in X} \mathcal{H}^2(\Phi(x)).$$

Definition 2.10 (Minimizing sequence and min-max sequence). A sequence $\{\Phi_i\}$ in Λ is said to be *minimizing* if

$$\limsup_{i \rightarrow \infty} \sup_{x \in X} \mathcal{H}^2(\Phi_i(x)) = \mathbf{L}(\Lambda).$$

If $\{\Phi_i\}$ is a minimizing sequence in Λ and $\{x_i\} \subset X$ satisfies

$$\lim_{i \rightarrow \infty} \mathcal{H}^2(\Phi_i(x_i)) = \mathbf{L}(\Lambda),$$

then $\{\Phi_i(x_i)\}$ is called a *min-max sequence*. For a minimizing sequence $\{\Phi_i\}$, we define its *critical set* $\mathbf{C}(\{\Phi_i\})$ to be the set of all subsequential varifold-limit of its min-max sequences:

$$\mathbf{C}(\{\Phi_i\}) := \{V = \lim_j |\Phi_{i_j}(x_j)| : x_j \in X, \|V\|(M) = \mathbf{L}(\Lambda)\}.$$

And $\{\Phi_i\}$ is called *pulled-tight* if every varifold in $\mathbf{C}(\{\Phi_i\})$ is stationary.

In a closed Riemannian 3-manifold (M, g) , for $L > 0$, let $\mathcal{W}_L(M, g)$ be the set of all varifolds $W \in \mathcal{V}_2(M)$, with $\|W\|(M) = L$, of the form

$$W = m_1|\Gamma_1| + \cdots + m_l|\Gamma_l|,$$

where $\{m_j\}$ is a set of positive integers and $\{\Gamma_j\}$ is a disjoint set of smooth, connected, embedded minimal surface in (M, g) . For a nonnegative integer $\mathfrak{g}_0 \geq 0$, let $\mathcal{W}_{L, \leq \mathfrak{g}_0}(M, g) \subset \mathcal{W}_L(M, g)$ consist of all the varifolds satisfying the genus bound

$$(2.5) \quad \sum_{j \in I_O} m_j \mathfrak{g}(\Gamma_j) + \frac{1}{2} \sum_{j \in I_N} m_j (\mathfrak{g}(\Gamma_j) - 1) \leq \mathfrak{g}_0,$$

where Γ_j is orientable if $j \in I_O$ and non-orientable if $j \in I_N$. Note that the genus of a non-orientable surface S is the number of cross-caps needed to be attached to a two-sphere in order to obtain a surface homeomorphic to S . For simplicity, when the ambient manifold (M, g) is clear from context, we may use the notations \mathcal{W}_L and $\mathcal{W}_{L, \leq \mathfrak{g}_0}$ without further specification.

Theorem 2.11 (Simon-Smith min-max theorem). *Given an orientable closed Riemannian 3-manifold (M, g) , a cubical subcomplex X of some $I(m, k)$, a Simon-Smith family $\Phi : X \rightarrow \mathcal{S}^*(M)$ of genus $\leq \mathfrak{g}_0$, and a positive real number $r > 0$, if $L := \mathbf{L}(\Lambda(\Phi)) > 0$, then there exists a minimizing sequence $\{\Phi_i\}$ in $\Lambda(\Phi)$ such that:*

(1) *The sequence $\{\Phi_i\}$ is pulled-tight and moreover,*

$$\mathbf{C}(\{\Phi_i\}) \cap \mathcal{W}_{L, \leq \mathfrak{g}_0} \neq \emptyset.$$

(2) *There exists a $W \in \mathcal{W}_{L, \leq \mathfrak{g}_0}$ such that every connected component Γ_j in W satisfies:*

- *if Γ_j is unstable and two-sided, then $m_j = 1$, and*
- *if Γ_j is one-sided, then its connected double cover is stable.*

(3) *Furthermore, there exists $\eta > 0$ such that for all sufficiently large i ,*

$$\mathcal{H}^2(\Phi_i(x)) \geq L - \eta \implies |\Phi_i(x)| \in \mathbf{B}_r^{\mathbf{F}}(\mathcal{W}_{L, \leq \mathfrak{g}_0}).$$

This min-max theorem essentially builds upon the foundational results in [Smi83, DLP10, Ket19, MN21, WZ24]. We postpone the detailed proof of this theorem to §4.

2.3. Relative Simon-Smith min-max theory. In this subsection, we fix a pair of parameter spaces, a cubical subcomplex X and its subcomplex $Z \subset X$. We adopt the concept of relative (X, Z) -homotopy class introduced in [Zho20] to the Simon-Smith min-max theory as follows.

Definition 2.12 (Relative homotopy class). Two Simon-Smith families Φ_0 and Φ_1 parametrized by the same parameter space X with $\Phi_0|_Z = \Phi_1|_Z$ are said to be *homotopic relative to $\Phi_0|_Z$* to each other if there exists a continuous map

$$\varphi : [0, 1] \times X \rightarrow \text{Diff}^\infty(M)$$

such that

- $\varphi(0, x) = \text{Id}$ for all $x \in X$,
- $\varphi(t, z) = \text{Id}$ for all $t \in [0, 1]$ and $z \in Z$,
- $\varphi(1, x)(\Phi_0(x)) = \Phi_1(x)$ for all $x \in X$.

The set of all families homotopic relative to $\Phi|_Z$ to a Simon-Smith family Φ is called the *relative (X, Z) -homotopy class of Φ* , and is denoted by $\Lambda_Z(\Phi)$.

Definition 2.13 (Relative Simon-Smith min-max width). Given a (X, Z) -relative homotopy class Λ_Z , its *width* is defined by

$$\mathbf{L}(\Lambda_Z) := \inf_{\Phi \in \Lambda_Z} \sup_{x \in X} \mathcal{H}^2(\Phi(x)).$$

Definition 2.14 (Minimizing sequence and min-max sequence for Relative Simon-Smith min-max). A sequence $\{\Phi_i\}$ in Λ_Z is said to be *minimizing* if

$$\limsup_{i \rightarrow \infty} \sup_{x \in X} \mathcal{H}^2(\Phi_i(x)) = \mathbf{L}(\Lambda_Z).$$

If $\{\Phi_i\}$ is a minimizing sequence and $\{x_i\} \subset X$ satisfies

$$\lim_{i \rightarrow \infty} \mathcal{H}^2(\Phi_i(x_i)) = \mathbf{L}(\Lambda_Z),$$

then $\{\Phi_i(x_i)\}$ is called a *min-max sequence*. For a minimizing sequence $\{\Phi_i\}$, we define its *critical set* $\mathbf{C}(\{\Phi_i\})$ to be the set of all subsequential varifold-limit of its min-max sequences:

$$\mathbf{C}(\{\Phi_i\}) := \{V = \lim_j |\Phi_{i_j}(x_j)| : x_j \in X, \|V\|(M) = \mathbf{L}(\Lambda_Z)\}.$$

And $\{\Phi_i\}$ is called *pulled-tight* if every varifold in $\mathbf{C}(\{\Phi_i\})$ is stationary.

Lemma 2.15. For a Simon-Smith family $\Phi : X \rightarrow \mathcal{S}^*(M)$ and a subcomplex $Z \subset X$, we have

$$\mathbf{L}(\Lambda(\Phi)) \geq \mathbf{L}(\Lambda_Z(\Phi)).$$

Proof. This follows immediately from the fact that $\Lambda_Z(\Phi) \subset \Lambda(\Phi)$. \square

Theorem 2.16 (Relative Simon-Smith min-max theorem). Given an orientable closed Riemannian 3-manifold (M, g) , a pair of cubical subcomplexes $Z \subset X$ of some $I(m, k)$, a Simon-Smith family $\Phi : X \rightarrow \mathcal{S}^*(M)$ of genus $\leq \mathfrak{g}_0$, and a positive real number $r > 0$, if

$$L = \mathbf{L}(\Lambda_Z(\Phi)) > \sup_{z \in Z} \mathcal{H}^2(\Phi(z)),$$

then there exists a minimizing sequence $\{\Phi_i\}$ in $\Lambda_Z(\Phi)$ such that (1), (2) and (3) in Theorem 2.11 still hold.

This relative min-max theorem can be proved in much the same way as 2.11, carefully following the steps of that proof. A proof is also provided in §4.

2.4. Almgren-Pitts min-max theory. By the Almgren isomorphism theorem [Alm62] (see also [LMN18, §2.5]), when equipped with the flat topology, $\mathcal{Z}_n(M; \mathbb{Z}_2)$ is weakly homotopic equivalent to $\mathbb{R}P^\infty$. Thus we can denote its cohomology ring by $\mathbb{Z}_2[\bar{\lambda}]$, where $\bar{\lambda} \in H^1(\mathcal{Z}_n(M; \mathbb{Z}_2), \mathbb{Z}_2)$ is the generator.

Let \mathcal{P}_p be the set of all \mathbf{F} -continuous maps $\Phi : X \rightarrow \mathcal{Z}_2(M; \mathbb{Z}_2)$, where X is a finite simplicial complex, such that $\Phi^*(\bar{\lambda}^p) \neq 0$. Elements of \mathcal{P}_p are called *p-sweepouts*. Note that every finite cubical complex is homeomorphic to a finite simplicial complex and vice versa (see [BP02, §4]). So in the above notion of *p-sweepouts*, we may as well require X to be a finite cubical complex.

Definition 2.17 (Almgren-Pitts *p*-width). Denoting by $\text{dmn}(\Phi)$ the domain of Φ , the *p*-width of (M, g) is defined by

$$\omega_p(M, g) := \inf_{\Phi \in \mathcal{P}_p} \sup_{x \in \text{dmn}(\Phi)} \mathbf{M}(\Phi(x)).$$

Remark 2.18. There is an equivalent definition of *p*-widths from [MN21, Remark 5.7]: First, an \mathcal{F} -continuous map $\Phi : X \rightarrow \mathcal{Z}_n(M; \mathbb{Z}_2)$ is said to have *no concentration of mass* if

$$\lim_{r \rightarrow 0} \sup_{x \in X, p \in M} \|\Phi(x)\| (B_r(p)) = 0.$$

Then, when defining the *p*-width, instead of using the collection \mathcal{P}_p , we use the collection of all \mathcal{F} -continuous maps Φ with no concentration of mass such that $\Phi^*(\bar{\lambda}^p) \neq 0$.

The interpolation results therein imply that the two definitions yield the same *p*-widths.

As in Simon-Smith min-max theory, we can define the notion of minimizing sequences, critical sets, and min-max sequences for the *p*-widths.

In this paper, the only result we require from the Almgren-Pitts min-max theory is that the 5-width of the unit 3-sphere (S^3, \bar{g}) is greater than 4π . In fact, C. Nurser [Nur16] has proved that

$$\omega_5(S^3, \bar{g}) = 2\pi^2.$$

3. PROOF OF THEOREM 1.1

In this section, we will prove our main theorem, Theorem 1.1: Given a 3-sphere (S^3, g_0) of positive Ricci curvature, we want to show that it has at least 5 embedded minimal tori. The proof consists of five key steps:

- (1) Define a 9-parameter Simon-Smith family Ψ of genus ≤ 1 , building upon the 5-parameter canonical family discovered by Marques-Neves in their proof of the Willmore conjecture [MN14].
- (2) Repeatedly apply Simon-Smith min-max theory to Ψ . Namely, each time we detect a minimal surface, which must be either a minimal sphere or a minimal torus of multiplicity one, according Wang-Zhou's multiplicity one theorem [WZ24], we remove the region in the family near the minimal surface (which is like a cap), resulting in a new family of maximal area strictly less than the area of the minimal surface. We repeat the min-max process on this new family.
- (3) Assemble pieces we obtained in step 2 to get a 9-parameter family Ξ , which is, in a certain sense, homotopic to Ψ .
- (4) Let N be the number of embedded minimal tori obtained via min-max in Step 2. Then we show that the parameter space $\text{dmn}(\Xi)$ of Ξ can be decomposed into a union of subsets D_0, D_1, \dots, D_N , such that the restriction $\Xi|_{D_0}$ is not a 5-sweepout, while for each $i = 1, \dots, N$, the image of $H_1(D_i; \mathbb{Z}_2)$ mapped into $H_1(\text{dmn}(\Xi); \mathbb{Z}_2)$ is trivial.
- (5) Show that $N \geq 5$ by analyzing the cohomology ring of $\text{dmn}(\Xi)$ and applying a Lyusternik-Schnirelmann argument.

3.1. A 9-parameter family. Let us first define a 9-parameter family Ψ of surfaces in the unit 3-sphere. It is worth noting that F. Marques previously considered a similar family to prove that the 8-width of the unit 3-sphere is $2\pi^2$ [Mar23].

For simplicity, throughout §3.1, we denote the unit 3-sphere by \mathbb{S}^3 , and the unit open 4-ball in \mathbb{R}^4 by \mathbb{B}^4 . We fix an unoriented Clifford torus Σ_0 in \mathbb{S}^3 .

Consider for each $v \in \mathbb{B}^4$ the conformal diffeomorphism $F_v : \mathbb{S}^3 \rightarrow \mathbb{S}^3$ given by

$$(3.1) \quad F_v(x) = \frac{1 - |v|^2}{|x - v|^2}(x - v) - v,$$

which pushes everything away from $v/|v|$ and towards $-v/|v|$ whenever $v \neq 0$.

Next, we choose an orientation for Σ_0 . This gives an inward normal direction for Σ_0 , and consequently, for each $F_v(\Sigma_0)$. With this orientation, we can define the signed distance $d_v : \mathbb{S}^3 \rightarrow \mathbb{R}$ to $F_v(\Sigma_0)$, which is positive outside $F_v(\Sigma_0)$ and negative inside.

Then we define the following continuous family of 2-cycles: For each pair $(v, t) \in \mathbb{B}^4 \times [-\pi, \pi]$, let

$$(3.2) \quad \Sigma(v, t) := \partial\{x \in \mathbb{S}^3 : d_v(x) < t\} \in \mathcal{Z}_2(\mathbb{S}^3; \mathbb{Z}_2).$$

Marques-Neves [MN14, §5] showed that by ingeniously reparametrizing this family of cycles near the boundary of the parameter space $\mathbb{B}^4 \times [-\pi, \pi]$, one can actually continuously (in the flat topology) extend this collection to

the boundary. Consequently, we obtain a continuous family of 2-cycles in $\mathcal{Z}_2(\mathbb{S}^3; \mathbb{Z})$ parametrized by $\overline{\mathbb{B}^4} \times [-\pi, \pi]$. Furthermore, any two “antipodal” points (v, t) and $(-v, -t)$ on the boundary would correspond to two 2-cycles that differ exactly by a sign. Then, by identifying the boundary of parameter space via $(v, t) \sim (-v, -t)$ and forgetting the orientations of all cycles, C. Nurser in [Nur16, §3.4.2] obtained an \mathcal{F} -continuous map

$$\Phi_5 : \mathbb{R}\mathbb{P}^5 \rightarrow \mathcal{Z}_2(\mathbb{S}^3; \mathbb{Z}_2),$$

for which he showed to be a 5-sweepout by proving $\Phi_5^*(\bar{\lambda}^5) \neq 0$ and Φ_5 has no concentration of mass ($\bar{\lambda}$ is the non-trivial element of $H^1(\mathcal{Z}_2(\mathbb{S}^3; \mathbb{Z}_2); \mathbb{Z}_2)$). The details of this procedure was carried out carefully in [Nur16, §3.4.2], but for the sake of completeness, we will also provide the details in §5.

Now, for each *oriented* Clifford torus Σ , one can repeat the above procedure, with Σ in place of Σ_0 , to obtain a map

$$\Phi_5^\Sigma : \mathbb{R}\mathbb{P}^5 \rightarrow \mathcal{Z}_2(\mathbb{S}^3; \mathbb{Z}_2).$$

Thus, if we let $\tilde{\mathcal{C}}$ be the space of oriented Clifford tori, we immediately obtain a map

$$(3.3) \quad \mathbb{R}\mathbb{P}^5 \times \tilde{\mathcal{C}} \rightarrow \mathcal{Z}_2(\mathbb{S}^3; \mathbb{Z}_2).$$

However, the topology of the parameter space of this family would not satisfy our need. We would want to identify Clifford tori of opposite orientations. Thus, we wish to identify the subfamily Φ_5^Σ and $\Phi_5^{-\Sigma}$ of the family (3.3), where $-\Sigma$ is Σ with an opposite orientation. We claim that this can be done. More precisely, let us consider instead the space \mathcal{C} of *unoriented* Clifford tori, which is topologically an $\mathbb{R}\mathbb{P}^2 \times \mathbb{R}\mathbb{P}^2$ [Nur16, §3.4.3]. We will prove in §5 the following theorem.

Theorem 3.1. *There exists an $\mathbb{R}\mathbb{P}^5$ -bundle Y over $\mathbb{R}\mathbb{P}^2 \times \mathbb{R}\mathbb{P}^2$, and a Simon-Smith family of genus ≤ 1*

$$\Psi : Y \rightarrow \mathcal{S}^*(S^3)$$

and the following property: If Σ is an oriented Clifford torus and $b \in \mathbb{R}\mathbb{P}^2 \times \mathbb{R}\mathbb{P}^2$ corresponds to $\{\Sigma, -\Sigma\}$, then there exists a homeomorphism $f : Y_b \rightarrow \mathbb{R}\mathbb{P}^5$ such that $\Phi_5^\Sigma \circ f = [\Psi|_{Y_b}]$.

Note that in the above, Y_b denotes the fiber at b , $\Psi|_{Y_b}$ the restriction of Ψ on Y_b , and $[\Psi|_{Y_b}]$ the induced map into $\mathcal{Z}_2(\mathbb{S}^3; \mathbb{Z}_2)$. Also note that since we are in the Simon-Smith setting, Ψ maps into the set $\mathcal{S}^*(S^3)$ of generalized surfaces instead of the cycle space $\mathcal{Z}_2(\mathbb{S}^3; \mathbb{Z}_2)$.

Remark 3.2. In [Nur16, §3.4.3] C. Nurser introduced a 9-parameter family parametrized by $\mathbb{R}\mathbb{P}^5 \times \mathbb{R}\mathbb{P}^2 \times \mathbb{R}\mathbb{P}^2$. However, that family is actually not well-defined. The issue is that to define an $\mathbb{R}\mathbb{P}^5$ -family like above, one needs to specify an orientation for the Clifford torus, but there is not a continuous way to assign an orientation to each unoriented Clifford torus in $\mathbb{R}\mathbb{P}^2 \times \mathbb{R}\mathbb{P}^2$.

3.2. Repetitive min-max. Now, let (S^3, g_0) be the 3-sphere given in our main theorem, Theorem 1.1. To prove the theorem, we can assume (S^3, g_0) has only finitely many embedded minimal tori; otherwise, we would already have at least five embedded minimal tori and the proof would be complete.

In §3.1, we obtained a 9-parameter Simon-Smith family Ψ of genus ≤ 1 on the unit 3-sphere. But this family can also be viewed as a Simon-Smith family in S^3 with any metric.

We will soon run the min-max process repeatedly starting from Ψ . To ensure that the process terminates in finitely many steps and that all the minimal surfaces detected have multiplicity one, we need to perturb g_0 to a new metric g' with the following properties:

Proposition 3.3. *For a Riemannian metric g of positive Ricci curvature on S^3 , if there are finitely many embedded minimal tori in (S^3, g) , then there exists another metric g' of positive Ricci curvature such that:*

- (1) (S^3, g) and (S^3, g') have the same number of embedded minimal tori.
- (2) (S^3, g') has finitely many embedded minimal spheres, each of which is non-degenerate.
- (3) For any embedded minimal sphere S in (S^3, g') and any positive integers m , the value $m \cdot \text{area}_{g'}(S)$ can never equal the g' -area of an embedded minimal torus in (S^3, g') .
- (4) For a pair of distinct embedded minimal spheres S_1, S_2 in (S^3, g') and a pair of positive integers m_1, m_2 , we have

$$m_1 \cdot \text{area}_{g'}(S_1) \neq m_2 \cdot \text{area}_{g'}(S_2).$$

The proof of the proposition above will be postponed to §6. In fact, from the proof, the metric g' can be made arbitrarily close to g in C^∞ , and all the g' -minimal tori coincide with the g -minimal tori. However, we will not need these facts.

Now, let us apply the above proposition to (S^3, g_0) to obtain a modified metric g' satisfying the above properties. To prove Theorem 1.1, it suffices to show that (S^3, g') has at least five embedded minimal tori. Therefore, in what follows, we will focus on (S^3, g') . We will need the following useful lemma.

Lemma 3.4. *In (S^3, g') , for any $L > 0$, suppose that $\mathcal{W}_{L, \leq 1}$ (defined using (2.5)) has a varifold associated with a multiplicity-one minimal surface Σ .*

- (1) *If Σ is a minimal torus, then every varifold in $\mathcal{W}_{L, \leq 1}$ is associated with a multiplicity-one minimal torus.*
- (2) *If Σ is a multiplicity-one minimal sphere, then*

$$\mathcal{W}_{L, \leq 1} = \{[\Sigma]\}.$$

In particular, $\mathcal{W}_{L, \leq 1}$ consists of finitely many varifolds, each associated with a multiplicity-one minimal surface.

Proof. Since (S^3, g') has positive Ricci curvature, the Frankel property holds, i.e., every pair of connected embedded minimal surfaces intersect with each

other. Consequently, every varifold in $\mathcal{W}_{L,\leq 1}$ is associated with a surface Σ , which is either a multiplicity-one minimal torus, or a minimal sphere with multiplicities.

For (1), by (3) of Proposition 3.3, every varifold in $\mathcal{W}_{L,\leq 1}$ is associated with a multiplicity-one minimal torus.

For (2), by (3) of Proposition 3.3 again, every varifold in $\mathcal{W}_{L,\leq 1}$ is associated with a minimal sphere with multiplicities. By (4) of Proposition 3.3, we have $\mathcal{W}_{L,\leq 1} = \{[\Sigma]\}$. \square

We now begin running the min-max process repeatedly, starting with our canonical family Ψ , treated as a family in (S^3, g') . We choose some $d_0 > 0$ to be less than the varifold distance between any two embedded minimal tori in (S^3, g') . (However, in §10, we will need to choose even smaller d_0 , depending on g' only; see the paragraph following the proof of Proposition 10.1.)

3.2.1. First stage. In (S^3, g') , applying Theorem 2.11 to Ψ with $r = d_0$, we obtain a varifold V^1 in $\mathcal{W}^1 := \mathcal{W}_{\mathbf{L}(\Lambda(\Psi)),\leq 1}$, where V^1 is induced by a multiplicity-one minimal surface of genus 0 or 1. By Lemma 3.4, we have the following dichotomy:

Case 1 \mathcal{W}^1 consists finitely many finitely many varifolds, each associated to a multiplicity-one minimal torus.

Case 2 $\mathcal{W}^1 = \{V^1\}$ where $\text{spt } \|V^1\|$ has genus 0.

Let us investigate case 1 first.

Case 1. From Theorem 2.11 (3) it follows immediately that there exists a $\delta > 0$ and some $\Psi' \in \Lambda(\Psi)$ such that

$$\mathbf{M}(\Psi'(x)) \geq \mathbf{L}(\Lambda(\Psi)) - \delta \implies |\Psi'(x)| \in \mathbf{B}_{d_0}^{\mathbf{F}}(\mathcal{W}^1).$$

However, this is not enough for us to prove Theorem 3.16. A stronger result is required.

Theorem 3.5. *Let (M, g) be an orientable closed Riemannian 3-manifold, Λ be a homotopy class of a Simon-Smith families of genus $\leq \mathfrak{g}_0$ and $L = \mathbf{L}(\Lambda) > 0$. Suppose that $\mathcal{W}_{L,\leq \mathfrak{g}_0}$ consists of finitely many varifolds, each associated with a connected multiplicity-one minimal surface. Then for any $r > 0$, there exists $\eta > 0$ and $\Phi \in \Lambda$ such that*

$$\mathbf{M}(\Phi(x)) \geq L - \eta \implies [\Phi(x)] \in \mathbf{B}_r^{\mathbf{F}}([\mathcal{W}_{L,\leq \mathfrak{g}_0}]).$$

The notation $[\mathcal{W}_{L,\leq \mathfrak{g}_0}]$ used here is as introduced in §2, and $\mathbf{B}_r^{\mathbf{F}}([\mathcal{W}_{L,\leq \mathfrak{g}_0}])$ denotes the r -neighborhood of $[\mathcal{W}_{L,\leq \mathfrak{g}_0}]$ in $\mathcal{Z}_2(M; \mathbf{F}; \mathbb{Z}_2)$. We will prove Theorem 3.5 in §7.

Applying the above theorem with $(M, g) = (S^3, g')$, $\Lambda = \Lambda(\Psi)$, $\mathfrak{g}_0 = 1$ and $r := d_0$, we obtain some $\eta > 0$ and $\Phi \in \Lambda(\Psi)$ satisfying the aforementioned property. For later reference, we denote $\delta_1 := \eta$ and $\tilde{\Psi}^1 := \Phi$.

By refining the cubical complex structure of the parameter space Y , we can obtain a *pure* 9-dimensional subcomplex C^1 of Y , which means it is made up of 9-dimensional cubes, such that

$$(3.4) \quad \mathbf{M}(\tilde{\Psi}^1(x)) \geq \mathbf{L}(\Lambda(\Psi)) - \delta_1/2 \implies x \in C^1,$$

and

$$(3.5) \quad x \in C^1 \implies [\tilde{\Psi}^1(x)] \in \mathbf{B}_{d_0}^{\mathbf{F}}([\mathcal{W}^1]).$$

Roughly speaking, C^1 is like the “cap” where $\tilde{\Psi}^1$ has high area, and is close to some embedded minimal tori.

Since (S^3, g') has only finitely many embedded minimal tori, by the definition of d_0 , and by further refining Y , we can assume that C^1 can be decomposed into a *disjoint* union of pure 9-dimensional subcomplexes $C_1^1, \dots, C_{n_1}^1$ such that: For each $j = 1, \dots, n_1$ there is a distinct embedded minimal torus $T_j^1 \in \mathcal{S}^*(S^3)$, with $|T_j^1| \in \mathcal{W}^1$, such that the image

$$[\tilde{\Psi}^1](C_j^1) \subset \mathbf{B}_{d_0}^{\mathbf{F}}([T_j^1]).$$

Finally, we define

$$\Psi^1 := \tilde{\Psi}^1|_{\overline{Y \setminus C^1}}.$$

By (3.4), Ψ^1 satisfies that

$$(3.6) \quad \sup \mathcal{H}^2 \circ \Psi^1 < \mathbf{L}(\Lambda(\Psi)),$$

Later when we run the min-max process for the second time, we will start with Ψ^1 , and the area upper bound guarantees new min-max minimal surfaces will be detected. For notational convenience later, let us denote $\tilde{Y}^1 := \text{dmn}(\tilde{\Psi}^1) = Y$ and $Y^1 = \overline{Y \setminus C^1} = \text{dmn}(\Psi^1)$

Case 2. In this case, the goal is once again to construct a map Ψ^1 that satisfies the bound (3.6). Additionally, we want Ψ^1 , when restricted to the boundary of the removed cap, to form a Simon-Smith family of genus 0. To achieve this, we will need to apply an important interpolation theorem. But first, let us introduce some concepts.

Given two topological spaces X and X' , and a continuous map $f : X' \rightarrow X$, the *mapping cylinder* of f is defined by

$$M_f := (([0, 1] \times X') \sqcup X) / \sim,$$

where $(0, x) \sim f(x)$ for all $x \in X'$.

In the following, we will primarily be interested in the case where X' and X are both finite cubical complexes, and $f : X' \rightarrow X$ is a surjective homotopy equivalence that is a *cubical map*, i.e. every cell of X' is mapped to a cell of X . In this case, given the mapping cylinder $W := M_f$, which can be viewed as a simplicial complex, we define the following subsets of W :

$$\partial_0 W := (\{0\} \times X') / \sim, \quad \partial_1 W := (\{1\} \times X') / \sim.$$




FIGURE 4. The figure on the left shows a map $f : X' \rightarrow X$. The figure on the right shows the mapping cylinder $W := M_f$, where the blue lines describe the deformation retraction of W onto $\partial_0 W$ induced by the map F_W .

Note that $\partial_0 W \cong X$ and $\partial_1 W \cong X'$. Heuristically, W is like a cobordism between $\partial_0 W$ and $\partial_1 W$. Also, we will sometimes by abuse of notation just denote $\partial_0 W$ by X , and $\partial_1 W$ by X' . Now, consider the map

$$[0, 1] \times [0, 1] \times X' \rightarrow [0, 1] \times X'$$

defined by sending each (t, s, x') to (ts, x') . This induces a map

$$(3.7) \quad F_W : [0, 1] \times W \rightarrow W$$

(see Figure 4). The key property of F_W is that the map $(t, w) \mapsto F_W(1-t, w)$, $(t, w) \in [0, 1] \times W$, is a *strong deformation retraction* of W onto $\partial_0 W$: This means for all $t \in [0, 1], w \in W, x \in \partial_0 W$ we have

$$F_W(1, w) = w, \quad F_W(0, w) \in \partial_0 W, \quad F_W(t, x) = x.$$

By [BP02, §4], W can also be viewed as a cubical complex.

Now, we can state the interpolation theorem.

Theorem 3.6. *In a closed 3-dimensional Riemannian manifold (M, g) , let $\Phi : X \rightarrow \mathcal{S}^*(M)$ be a Simon-Smith family of genus ≤ 1 , where X is a cubical subcomplex in I^m . Suppose that $L = \mathbf{L}(\Lambda(\Phi)) > 0$, and the set $\mathcal{W}_{L, \leq 1}$ consists of exactly one varifold associated with a multiplicity-one nondegenerate minimal sphere. Then after refining X , there exists a cubical subcomplex X' in I^{m+1} , a homotopy equivalence $f : X' \rightarrow X$, which is a surjective cubical map, and a Simon-Smith family $\Phi' : X' \rightarrow \mathcal{S}^*(M)$ of genus ≤ 1 with the following properties.*

- (1) *There exists an $\eta > 0$ such that for $x \in X'$,*

$$\mathcal{H}^2(\Phi'(x)) \geq L - \eta \implies \mathbf{g}(\Phi'(x)) = 0.$$

- (2) *Let W be the mapping cylinder $M_f = (([0, 1] \times X') \sqcup X) / \sim$ of f , where $(0, x) \sim f(x)$ for all $x \in X'$. There exists a Simon-Smith family $H : W \rightarrow \mathcal{S}^*(M)$ of genus ≤ 1 such that*

- (a) *$H|_{\partial_0 W} = \Phi$ and $H|_{\partial_1 W} = \Phi'$;*

(b) for all $x \in X'$, $t \mapsto H(t, x)$ is a pinch-off process (see Definition 3.7).

Heuristically, if a torus is sufficiently close to a sphere, one can find a loop within a small ball. Under mean curvature flow, it is expected that the flow would undergo a *neck pinch* along the loop, yielding a genus 0 generalized surface. However, as explained in §1.2.2, we cannot employ mean curvature flow to reduce the genus. Instead, we *pinch off* all components of the torus within the small ball. This process involves a finite number of neck-pinch surgeries, so that the surface no longer intersects the boundary of the small ball, and then shrinking the components inside the ball into a point.

Definition 3.7. In a closed Riemannian manifold (M, g) , a Simon-Smith family $\Phi : [a, b] \rightarrow \mathcal{S}^*(M)$, where the parameter space $[a, b]$ is viewed as a time interval, is called a *pinch-off process* if the following holds. (Note that this definition is purely topological.)

There exist finitely many spacetime points $(t_1, p_1), \dots, (t_n, p_n)$, where $t_i \in [a, b]$ and $p_i \in \Phi(t_i)$, such that:

- (1) For each (t, p) , where $t \in [a, b]$ and $p \in \Phi(t)$, different from any (t_i, p_i) , there exists an open time interval $J \subset [a, b]$ around t , a one-parameter group of diffeomorphisms $\{\varphi_{t'}\}_{t' \in J} \subset \text{Diff}(M)$ and an open ball $U \subset S^3$ around p such that for all $t' \in J$,

$$\varphi_{t'}(\Phi(t)) \cap U = \Phi(t') \cap U.$$

- (2) For each $i = 1, \dots, n$, there exist an open time interval $J \subset [a, b]$ around t_i and a ball $U \subset S^3$ around p_i such that the family

$$\{\Phi(t') \cap U\}_{t' \in J}$$

is of one of the following types:

- A *surgey process*: Pinching a (topological) cylinder at the point p_i , at time t_i , to obtain a (topological) double cone, and then either splitting it into two smooth discs or leaving it intact.
- A *shrinking process*: For each $t' \in J$, $\Phi(t')$ does not intersect ∂U , and $\{\Phi(t') \cap U\}_{t' \in J \cap [a, t_i]}$ is induced by a one-parameter group of diffeomorphisms in U , and $\Phi(t') \cap U = p_i$ for all $t' \in J \cap [t_i, b]$.

Remark 3.8. Let us compare pinch-off process and mean curvature flow. For mean curvature flow, conical singularities may occur, while our pinch-off process would not generate new conical singularities. Nonetheless, as we will prove in §9, a pinch-off process would, in a certain sense, simplify the topology of the initial condition. This should be compared with B. White’s result that mean curvature flow simplify the topology of the surfaces [Whi95].

We will prove the interpolation theorem, Theorem 3.6, in §8. Here is brief outline: First, we show that if a torus is close to a sphere, in the varifold

sense, it must contain a non-trivial loop in a *small* ball. Heuristically, we can then pinch this small loop to obtain a generalized surface with genus 0. The main challenge lies in the fact that the choice of small loops may not be continuous in a Simon-Smith family, causing the pinching process to fail in producing a desired deformation map H . To address this, we will extend Pitts's combinatorial arguments [Pit81, Theorem 4.10], and construct a map H which consists of a union of pinch-off processes.

Note that H itself is not a homotopy between Φ and $\tilde{\Phi}$ in the sense of Simon-Smith min-max theory due to surgeries and shrinking, but the induced maps $[\Phi]$ and $[\tilde{\Phi}]$ are in the same homology class in $\mathcal{Z}_2(M; \mathbf{F}; \mathbb{Z}_2)$, via $[H]$.

Applying Theorem 3.6 to Ψ , we obtain:

- a cubical complex \tilde{Y}^1 ,
- a surjective, cubical, homotopy equivalence $f^1 : \tilde{Y}^1 \rightarrow Y$,
- a mapping cylinder $W^1 := M_{f^1}$ (which can be viewed as a simplicial complex) with

$$Y = \partial_0 W^1, \quad \tilde{Y}^1 = \partial_1 W^1,$$

and the associated map $F^1 := F_{W^1} : [0, 1] \times W^1 \rightarrow W^1$ (see (3.7)),

- a Simon-Smith family of genus ≤ 1

$$\tilde{\Psi}^1 : \tilde{Y}^1 \rightarrow \mathcal{S}^*(S^3),$$

- a Simon-Smith family of genus ≤ 1

$$H^1 : W^1 \rightarrow \mathcal{S}^*(S^3),$$

with $H^1|_{\partial_0 W^1} = \Psi$, $H^1|_{\partial_1 W^1} = \tilde{\Psi}^1$,

- a constant $\delta_1 > 0$ (in place of η),

satisfying the properties listed in the theorem.

Then by refining \tilde{Y}^1 , we can obtain a cubical subcomplex C^1 of \tilde{Y}^1 such that

$$\mathcal{H}^2(\tilde{\Psi}^1(x)) \geq \mathbf{L}(\Lambda(\Psi)) - \delta_1/2 \implies x \in C^1$$

and $\tilde{\Psi}^1|_{C^1}$ is a Simon-Smith family of genus 0. Finally, we let

$$\Psi^1 := \tilde{\Psi}^1|_{\overline{\tilde{Y}^1 \setminus C^1}}.$$

As in case 1, a key property of Ψ^1 is that

$$\sup \mathcal{H}^2 \circ \Psi^1 < \mathbf{L}(\Lambda(\Psi)).$$

Let us denote $Y^1 := \text{dmn}(\Psi^1)$.

In summary, in both case 1 and 2, we obtain a Simon-Smith family Ψ^1 of genus ≤ 1 with parameter space $\overline{\tilde{Y}^1 \setminus C^1}$, and with $\sup \mathcal{H}^2 \circ \Psi^1$ strictly less than the width $\mathbf{L}(\Lambda(\Psi))$. Thus, if we run min-max for the second time using Ψ^1 , all min-max minimal surfaces will have lower area than those we obtained above. And we will keep repeating the min-max process.

3.2.2. *k*-th stage. In this section, we explain in detail what we obtain in the *k*-th stage of min-max for $k \geq 1$. Before running the *k*-th stage of min-max, we would have in our hand a Simon-Smith family of genus ≤ 1

$$\Psi^{k-1} : Y^{k-1} \rightarrow \mathcal{S}^*(S^3),$$

where Y^{k-1} is some cubical complex. If $k = 1$, we would take $\Psi^0 := \Psi$ and $Y^0 := Y$.

Now, let us apply the min-max theorem, Theorem 2.11, to Ψ^{k-1} with $r = d_0$. Like before, we obtain a varifold V^k in $\mathcal{W}^k := \mathcal{W}_{\mathbf{L}(\Lambda(\Psi^{k-1})), \leq 1}$, where V^k is induced by a multiplicity one, smooth, embedded, minimal surface of genus 0 or 1. Again, there are two cases:

- Case 1. each element of \mathcal{W}^k is associated with a multiplicity one torus;
- Case 2. V^k is associated with a smooth, embedded, minimal sphere of multiplicity one and $\mathcal{W}^k = \{V^k\}$.

Remark 3.9. In the following, the superscript k indicates that the object concerned is created from running the min-max process for the *k*-th time.

Case 1. Arguing as before, we obtain the following objects:

- a Simon-Smith family of genus ≤ 1

$$\tilde{\Psi}^k : Y^{k-1} \rightarrow \mathcal{S}^*(S^3)$$

in $\Lambda(\Psi^{k-1})$,

- a “cap” C^k , which is a subcomplex of Y^{k-1} and can be decomposed into a disjoint union of cubical subcomplexes $C_1^k, \dots, C_{n_k}^k$,
- the restriction

$$\Psi^k := \tilde{\Psi}^k|_{\overline{Y^{k-1} \setminus C^k}},$$

- the new domain $Y^k := \text{dmn}(\Psi^k) = \overline{Y^{k-1} \setminus C^k}$,

that have the following properties:

- For each $j = 1, \dots, n_k$, there is a distinct embedded minimal torus $T_j^k \in \mathcal{S}^*(S^3)$, with $|T_j^k| \in \mathcal{W}^k$, such that the image

$$[\tilde{\Psi}^k](C_j^{k-1}) \subset \mathbf{B}_{d_0}^{\mathbf{F}}(|T_j^k|).$$

- $\sup_{Y^k} \mathcal{H}^2 \circ \Psi^k < \mathbf{L}(\Lambda(\Psi^{k-1}))$.

Let us also note that since $\tilde{\Psi}^k$ and Ψ^{k-1} are homotopic in the sense of Simon-Smith min-max theory, there exists a homotopy

$$H^k : [0, 1] \times Y^{k-1} \rightarrow \mathcal{S}^*(S^3)$$

such that

$$H(0, \cdot) = \Psi^{k-1}, \quad H(1, \cdot) = \tilde{\Psi}^k,$$

and for each (t, x) , $H(t, x)$ is obtained from $\Psi^{k-1}(x)$ via a diffeomorphism of S^3 , according to Definition 2.7.

For consistency in both cases 1 and 2, we introduce the following notation, which mirrors the setup in Case 2. Define

$$W^k := [0, 1] \times Y^{k-1}, \quad \partial_0 W^k = \{0\} \times Y^{k-1}, \quad \partial_1 W^k = \{1\} \times Y^{k-1},$$

and a map

$$F^k : [0, 1] \times W^k \rightarrow W^k$$

given by

$$F^k(s, (t, x)) := (st, x).$$

Note that the map $(s, w) \mapsto F^k(1 - s, w)$, with $(s, w) \in [0, 1] \times W^k$, is a strong deformation retraction of W^k onto $\{0\} \times Y^{k-1}$. For convenience, we will identify Y^{k-1} with the subset $\{0\} \times Y^{k-1}$ in W^k , and denote $\{1\} \times Y^{k-1}$ by \tilde{Y}^k .

Remark 3.10. For example, in Case 1, if we have a subset $C \subset Y^{k-1}$, then $(F^k)^{-1}(C)$ and $(F^k(0, \cdot))^{-1}(C)$ are the subset $[0, 1] \times C$ of W^k .

Case 2. In this case, by applying Theorem 3.6 to Ψ^{k-1} with $r = d_0$, we obtain:

- a cubical complex \tilde{Y}^k ,
- a surjective, cubical, homotopy equivalence $f^k : \tilde{Y}^k \rightarrow Y^{k-1}$,
- a mapping cylinder $W^k := M_{f^k}$ (viewed as a simplicial complex) with

$$Y^{k-1} = \partial_0 W^k, \quad \tilde{Y}^k = \partial_1 W^k,$$

and the associated map $F^k := F_{W^k} : [0, 1] \times W^k \rightarrow W^k$ (see (3.7)),

- a Simon-Smith family of genus ≤ 1

$$\tilde{\Psi}^k : \tilde{Y}^k \rightarrow \mathcal{S}^*(S^3),$$

- a Simon-Smith family of genus ≤ 1

$$H^k : W^k \rightarrow \mathcal{S}^*(S^3),$$

with $H^k|_{\partial_0 W^k} = \Psi^{k-1}$, $H^k|_{\partial_1 W^k} = \tilde{\Psi}^k$,

- a ‘‘cap’’ C^k , which is a cubical subcomplex of \tilde{Y}^k ,
- the restriction

$$\Psi^k := \tilde{\Psi}^k|_{\overline{Y^{k-1} \setminus C^k}},$$

- the new domain $Y^k := \text{dmn}(\Psi^k) = \overline{Y^{k-1} \setminus C^k}$,

that satisfy the following:

- The map $(t, w) \mapsto F^k(1 - t, w)$, where $(t, w) \in [0, 1] \times W^k$, is a strong deformation retraction of W^k onto Y^{k-1} .
- For each $x \in \tilde{Y}^k$, the family $t \mapsto H^k(F^k(t, x))$, $t \in [0, 1]$, is a pinch-off process.
- $\tilde{\Psi}^k|_{C^k}$ is a Simon-Smith family of genus 0.
- $\sup \mathcal{H}^2 \circ \Psi^k < \mathbf{L}(\Lambda(\Psi^{k-1}))$.

Note that, in both case 1 and 2, the mass bound

$$(3.8) \quad \sup \mathcal{H}^2 \circ \Psi^k < \mathbf{L}(\Lambda(\Psi^{k-1}))$$

holds. Thus, the width $\mathbf{L}(\Lambda(\Psi^k))$ is strictly decreasing with respect to k . Combined with the assumption that (S^3, g') admits only finitely many embedded minimal spheres and tori, our repetitive min-max process must terminate in finitely many steps. Specifically, for some K , when we run the min-max process for the K -th time, using the family Ψ^{K-1} , the width $\mathbf{L}(\Lambda(\Psi^{K-1}))$ reaches zero.

3.2.3. Last stage. We have shown that the width $\mathbf{L}(\Lambda(\Psi^{K-1}))$ is zero. Thus, there exists some Simon-Smith family $\Psi^K \in \Lambda(\Psi^{K-1})$ and a homotopy in the sense of Simon-Smith min-max,

$$H^K : [0, 1] \times Y^{K-1} \rightarrow \mathcal{S}^*(S^3),$$

such that:

- $H^K(0, \cdot) = \Psi^{K-1}$, $H^K(1, \cdot) = \Psi^K$.
- For each (t, x) , $H^K(t, x)$ is obtained from $\Psi^{K-1}(x)$ via some diffeomorphism of S^3 according to Definition 2.7.
- $\sup_{Y^{K-1}} \mathcal{H}_{\bar{g}}^2 \circ \Psi^K < 1$, where \bar{g} denotes the standard metric on S^3 .

Crucially, note that in the last bullet point, the Hausdorff measure is taken with respect to *the unit 3-sphere* (S^3, \bar{g}) .

We denote

$$W^K := [0, 1] \times Y^{K-1}, \quad \partial_0 W^K = \{0\} \times Y^{K-1}, \quad \partial_1 W^K = \{1\} \times Y^{K-1},$$

and define a map

$$F^K : [0, 1] \times W^K \rightarrow W^K, \quad (s, (t, x)) \mapsto (st, x).$$

Note that the map $(s, w) \mapsto F^K(1-s, w)$, with $(s, w) \in [0, 1] \times W^K$, is a strong deformation retraction of W^K onto $\{0\} \times Y^{K-1}$. As before, we will identify Y^{K-1} with the subset $\{0\} \times Y^{K-1}$ in W^K .

3.3. A new family Ξ . We have already run the min-max process K times already, generating multiple families with from the original family Ψ , each with a different parameter space. In this section, we will use these family to reconstruct a new Simon-Smith family Ξ , whose parameter space is homotopy equivalent to the original parameter space Y (which is an $\mathbb{R}\mathbb{P}^5$ -bundle over $\mathbb{R}\mathbb{P}^2 \times \mathbb{R}\mathbb{P}^2$). This new family remains a Simon-Smith family of genus ≤ 1 .

Consider the following list of $2K - 1$ Simon-Smith families of genus ≤ 1 . For visualization purposes, readers may find it helpful to refer to Figure 5.

- (1) $\tilde{\Psi}^1|_{C^1}$.
- (2) $H^2|_{(F^2)^{-1}(A_1)}$, where $A_1 := C^1 \cap Y^1$. Note that A_1 is simply ∂C^1 if \tilde{Y}^1 is a topological manifold, and $H^2|_{(F^2)^{-1}(A_1)}$ coincides with $H^2|_{[0,1] \times A_1}$ if Case 1 occurred at the second stage of the min-max process; see Remark 3.10.

(3) $\tilde{\Psi}^2|_{C^2}$.

(4) $H^3|_{(F^3)^{-1}(A_2)}$, $A_2 \subset Y^3$ is defined by

$$A_2 := \left((F^2)^{-1}(A_1) \cap Y^2 \right) \cup \left(C^2 \cap Y^2 \right).$$

Note that $H^3|_{(F^3)^{-1}(A_2)}$ coincides with $H^3|_{[0,1] \times A_2}$ if Case 1 occurred at the third stage of min-max.

(5) $\tilde{\Psi}^3|_{C^3}$.

(6) $H^4|_{(F^4)^{-1}(A_3)}$, where $A_3 \subset Y^3$ is defined by

$$A_3 := \left((F^3)^{-1}(A_2) \cap Y^3 \right) \cup \left(C^3 \cap Y^3 \right).$$

...

(2K - 4) ...

(2K - 3) $\tilde{\Psi}^{K-1}|_{C^{K-1}}$.

(2K - 2) $H^K|_{(F^K)^{-1}(A_{K-1})}$, where $A_{K-1} \subset Y^{K-1}$ is defined by

$$A_{K-1} := \left((F^{K-1})^{-1}(A_{K-2}) \cap Y^{K-1} \right) \cup \left(C^{K-1} \cap Y^{K-1} \right).$$

(2K - 1) Ψ^K .




FIGURE 5. Constructing Ξ .

Clearly, by the definition of these $2K - 1$ families, there is a natural way to glue them together. Namely, for each $k = 1, \dots, K - 2$, the following pairs of families share a common subfamily:

- $(2k)$ and $(2k - 1)$
- $(2k)$ and $(2k + 1)$

- $(2k)$ and $(2k + 2)$

Additionally, $(2K - 2)$ and $(2K - 1)$ share a common subfamily as well. The overall gluing scheme is described in the schematic Figure 5.

Now, let the newly obtained family be called Ξ . It is also a Simon-Smith family of genus ≤ 1 .

Proposition 3.11. *There exist:*

- a simplicial complex \widetilde{W} , containing Y and $\text{dmn}(\Xi)$ as subcomplexes,
- a map $\widetilde{F} : [0, 1] \times \widetilde{W} \rightarrow \widetilde{W}$ such that the map $(t, w) \mapsto \widetilde{F}(1 - t, w)$, with $(t, w) \in [0, 1] \times \widetilde{W}$, is a strong deformation retraction of \widetilde{W} onto Y , and $\widetilde{F}(0, \cdot)|_{\text{dmn}(\Xi)}$ is a surjective homotopy equivalence from $\text{dmn}(\Xi)$ onto Y ,
- a Simon-Smith family of genus ≤ 1 ,

$$\widetilde{\Xi} : \widetilde{W} \rightarrow \mathcal{S}^*(S^3),$$

such that:

- (1) $\widetilde{\Xi}|_Y = \Psi$, $\widetilde{\Xi}|_{\text{dmn}(\Xi)} = \Xi$.
- (2) For each $x \in \text{dmn}(\Xi)$, the family $t \mapsto \widetilde{\Xi}(\widetilde{F}(t, x))$, with $t \in [0, 1]$, is a pinch-off process.

Thus, heuristically, the map $(t, w) \mapsto \widetilde{\Xi}(\widetilde{F}(1 - t, w))$, where $(t, w) \in [0, 1] \times \text{dmn}(\Xi)$, can be viewed as a “homotopy” that deforms Ξ back to Ψ .

Proof. By examining the construction of Ξ , there is a natural, canonical way to define the objects mentioned in the proposition. Specifically, we introduce a sequence of “ \mathbf{F} -homotopies” that, in a certain sense, homotope Ξ back to Ψ . Before proceeding, we introduce the following notations:

- The \mathbf{F} -metric in $\mathcal{Z}_2(S^3; \mathbb{Z}_2)$ does not induce a metric on the space $\mathcal{S}^*(S^3)$. However, the \mathbf{F} -metric would induce a pseudometric, and therefore a topology, on $\mathcal{S}^*(S^3)$. With this in mind, the list in the next paragraph would describe a sequence of homotopies, called \mathbf{F} -homotopies under this topology.
- We use “+” to define the addition of chains with \mathbb{Z}_2 -coefficients in $\mathcal{S}^*(S^3)$, where overlapping portions cancel out each other (see, for example, Figure 5).
- We use the numbering $(1), (2), \dots, (2K - 1)$ of the subfamilies of Ξ stated at the beginning of §3.3.

Let us now consider the following the following sequence of \mathbf{F} -homotopies. For visualization purposes, readers may refer to Figure 5.

- The family Ξ can be written as the sum of three subfamilies:

$$\Xi = (2K - 1) + (2K - 2) + \text{remaining part.}$$

We homotope the subfamily $(2K - 1) + (2K - 2)$ back to Ψ^{K-1} (using the family H^K and the strong deformation retraction of W^K onto

Y^{K-1} induced by F^K), while fixing the remaining part. Call this new family Ξ^{K-1} .

- The family Ξ^{K-1} can be written as the sum of three subfamilies:

$$\Xi^{K-1} = \tilde{\Psi}^{K-1} + (2K - 4) + \text{remaining part.}$$

We homotope the subfamily $\tilde{\Psi}^{K-1} + (2K - 4)$ to Ψ^{K-2} (using the family H^{K-1} and the strong deformation retraction of W^{K-1} onto Y^{K-2} induced by F^{K-1}), while fixing the remaining part. Call this new family Ξ^{K-2} .

- ...
- The family Ξ^2 can be written as the sum of three subfamilies:

$$\Xi^2 = \tilde{\Psi}^2 + (2) + \text{remaining part.}$$

We homotope the subfamily $\tilde{\Psi}^2 + (2)$ to Ψ^1 (using the family H^2 and the strong deformation retraction of W^2 onto Y^1 induced by F^2), while fixing the remaining part. Call this new family Ξ^1 , which is actually $\tilde{\Psi}^1$.

- Finally, we homotope $\Xi^1 = \tilde{\Psi}^1$ to Ψ using the family H^1 and the strong deformation retraction of W^1 onto $Y^0 = Y$ induced by F^1 .

From the above chain of \mathbf{F} -homotopies, one easily can construct \tilde{W} , \tilde{F} , and $\tilde{\Xi}$ satisfying the desired conditions. \square

3.4. Decomposing Ξ . Recall that for each $k = 1, \dots, K - 1$, during the k -th min-max process, one of two cases occurred: (1) Some multiplicity one tori $T_1^k, \dots, T_{n_k}^k$ were detected, and (2) a multiplicity one sphere V^k was detected. For each k , if Case 1 occurred, then $C^k \subset \text{dmn}(\tilde{\Psi}^k)$ can be decomposed into a union of $C_1^k, \dots, C_{n_k}^k$, and we will call each C_j^k a *genus 1 cap*. Recall that $[\tilde{\Psi}^k|_{C_j^k}]$ is \mathbf{F} -close to $[T_j^k]$. If instead Case 2 occurred, we call C^k a *genus 0 cap*. Recall that $\tilde{\Psi}^k|_{C^k}$ is a Simon-Smith family of genus 0.

Now, if $C \subset \text{dmn}(\tilde{\Psi}^k)$ is either a genus 0 cap or a genus 1 cap obtained at the k -th stage of min-max, we define the *trace of C* , a subset of $\text{dmn}(\Xi)$, as follows. First, define the following sets (see Figure 6):

- $B_k := C$, a subcomplex of $\text{dmn}(\tilde{\Psi}^k) = \tilde{Y}^k$.
- $B_{k+1} := (F^{k+1})^{-1}(B_k \cap Y^k) \subset W^{k+1}$. Here $B_k \cap Y^k$ can also be viewed as a subset of $\text{dmn}(\tilde{\Psi}^k)$.
- $B_{k+2} := (F^{k+2})^{-1}(B_{k+1} \cap Y^{k+1}) \subset W^{k+2}$. Here $B_{k+1} \cap Y^{k+1}$ can also be viewed as a subset of $\text{dmn}(\tilde{\Psi}^{k+1})$.
- ...

Note that each B_k, B_{k+1}, \dots, B_K is regarded as a subset of $\text{dmn}(\Xi)$.

Finally, we define the trace of C by

$$T(C) := B_k \cup B_{k+1} \cup \dots \cup B_K \subset \text{dmn}(\Xi).$$




FIGURE 6. The trace $T(C)$ in $\text{dmn}(\Xi)$.

Now, let N be the number of all the genus 1 caps we obtained throughout the first $K - 1$ stages of min-max. To prove Theorem 1.1, we just need to show that $N \geq 5$.

Let us decompose $\text{dmn}(\Xi)$ as the union of the following $N + 1$ subcomplexes:

- Let D_0 be the union of $\text{dmn}(\Psi^K)$ and $\cup_C T(C)$, where C ranges over the genus 0 caps.
- For each genus 1 cap C , we consider its trace $T(C)$. There are in total N such traces, which are denoted by D_1, \dots, D_N .

It follows directly from the definition of Ξ that

$$\text{dmn}(\Xi) = D_0 \cup D_1 \cup \dots \cup D_N.$$

Proposition 3.12. *The decompositions above have the following two properties:*

- (1) $\Xi|_{\cup_C T(C)}$, where C ranges over the genus 0 caps, is a Simon-Smith family of genus 0.
- (2) For each genus 0 or genus 1 cap C , there exists a strong deformation retraction of the trace $T(C)$ back to C in $\text{dmn}(\Xi)$.

Proof. For (1), if C is a genus 0 cap obtained at the k -stage of min-max, then $\tilde{\Psi}^k|_{C^k}$ is a Simon-Smith family of genus 0. Then it follows from the definition of the homotopies H^k , namely Theorem 3.6 (2)(b), that $\Xi|_{T(C)}$ is also a Simon-Smith family of genus 0. This proves (1). Item (2) follows directly from the definition of $T(C)$, using the maps F^{k+1}, \dots, F^K . \square

3.5. Topological arguments. Finally in this subsection, we will show that $N \geq 5$.

We first define three elements λ, α, β in $H^1(Y; \mathbb{Z}_2)$:

- Let $\bar{\lambda}$ be the non-trivial element of $H^1(\mathcal{Z}_2(S^3; \mathbb{Z}_2); \mathbb{Z}_2)$. Define λ as the pullback of $\bar{\lambda}$ under $[\Psi]$.

- Let $a = \mathbb{RP}^1 \times \mathbb{RP}^2 \subset \mathbb{RP}^2 \times \mathbb{RP}^2$, and $A \subset Y$ be the \mathbb{RP}^5 -subbundle over a . Define α to be the Poincaré dual $PD(A)$ of A .
- Let $b = \mathbb{RP}^2 \times \mathbb{RP}^1 \subset \mathbb{RP}^2 \times \mathbb{RP}^2$, and $B \subset Y$ be the \mathbb{RP}^5 -subbundle over b . Define β to be the Poincaré dual $PD(B)$ of B .

Here we assumed \mathbb{RP}^2 is built from attaching a 2-disk to the non-trivial loop $\mathbb{RP}^1 \subset \mathbb{RP}^2$.

The following topological fact about Y is crucial and is the main reason we used the family Ψ in the first place.

Theorem 3.13. *In the cohomology ring $H^*(Y; \mathbb{Z}_2)$,*

$$\lambda^5 \cup \alpha^2 \cup \beta^2 \neq 0.$$

We will prove this theorem in §5.

Now, to prove $N \geq 5$, let us suppose by contradiction that $N \leq 4$.

By Proposition 3.11, $\text{dmn}(\Xi)$ and Y are homotopy equivalent, so, by abuse of notation, we will also treat view λ, α, β as elements of $H^1(\text{dmn}(\Xi); \mathbb{Z}_2)$ as well. For each $j = 0, 1, \dots, N$, let i_j be the inclusion map $D_j \hookrightarrow \text{dmn}(\Xi)$. Moreover, given a cohomology class γ of $\text{dmn}(\Xi)$, we denote its pullback under

$$(i_j)^* : H^*(\text{dmn}(\Xi); \mathbb{Z}_2) \rightarrow H^*(D_j; \mathbb{Z}_2)$$

by $\gamma|_{D_j}$. Then by Theorem 3.13 and a Lyusternik-Schnirelmann argument, we obtain:

Lemma 3.14.

- If $N = 4$, then one of $\lambda^5|_{D_0}, \alpha|_{D_1}, \alpha|_{D_2}, \beta|_{D_3}, \beta|_{D_4}$ is non-zero.
- If $N = 3$, then one of $\lambda^5|_{D_0}, \alpha|_{D_1}, \alpha|_{D_2}, \beta^2|_{D_3}$ is non-zero.
- If $N = 2$, then one of $\lambda^5|_{D_0}, \alpha|_{D_1}, (\alpha \cup \beta^2)|_{D_2}$ is non-zero.
- If $N = 1$, then one of $\lambda^5|_{D_0}, (\alpha^2 \cup \beta^2)|_{D_1}$ is non-zero.
- If $N = 0$, i.e. $\text{dmn}(\Xi) = D_0$, then $\lambda^5|_{D_0}$ is non-zero.

By Proposition 3.11, regarding the induced maps $[\Psi]$ and $[\Xi]$ into $\mathcal{Z}_2(S^3; \mathbb{Z}_2)$, we have

$$[\Xi]^*(\bar{\lambda}) = [\Psi]^*(\bar{\lambda}) = \lambda.$$

Hence,

$$\lambda^5|_{D_0} = i_0^*([\Xi]^*(\bar{\lambda})^5) = ([\Xi \circ i_0]^*(\bar{\lambda}))^5 = ([\Xi|_{D_0}]^*(\bar{\lambda}))^5,$$

which is non-zero if and only if $[\Xi|_{D_0}]$ is a 5-sweepout. Consequently, it suffices to prove the following two facts:

Proposition 3.15. $\Xi|_{D_0}$ is not a 5-sweepout.

Theorem 3.16. Suppose $1 \leq N \leq 4$. Then for each $j = 1, \dots, N$, the map

$$(i_j)_* : H_1(D_j; \mathbb{Z}_2) \rightarrow H_1(\text{dmn}(\Xi); \mathbb{Z}_2)$$

is the zero map.

Indeed, Proposition 3.15 implies $\lambda^5|_{D_0} = 0$, and Theorem 3.16, via the universal coefficients theorem, implies the map

$$(i_j)^* : H^1(\text{dmn}(\Xi); \mathbb{Z}_2) \rightarrow H^1(D_j; \mathbb{Z}_2)$$

is trivial for each $j = 1, \dots, N$. This would mean that all the cohomology classes in Lemma 3.14 must be zero, contradicting the assumption that $N \leq 4$. Thus, it remains to prove Proposition 3.15 and Theorem 3.16.

Proof of Proposition 3.15. In this proof, we will treat $\Xi|_{D_0}$ as a Simon-Smith family of genus ≤ 1 in the unit 3-sphere (S^3, \bar{g}) instead of (S^3, g') . Recall that D_0 is the union of $E := \cup_C T(C)$, where C runs over the genus 0 caps, and $\text{dmn}(\Psi^K)$.

By Proposition 3.12, $\Xi|_E$ is a Simon-Smith family of genus 0, while

$$\sup \mathcal{H}_{\bar{g}}^2 \circ \Xi|_{\text{dmn}(\Psi^K)} = \sup \mathcal{H}_{\bar{g}}^2 \circ \Psi^K < 1 \quad (< 4\pi)$$

by definition. Thus, if we apply the *relative* Simon-Smith min-max, Theorem 2.16, on the unit round 3-sphere to $\Xi|_{D_0}$, relative to $\text{dmn}(\Psi^K) \subset D_0$, we will detect the multiplicity one equatorial 2-sphere. This immediately implies that the width $\mathbf{L}(\Lambda(\Xi|_{D_0}))$, in the *non-relative* Simon-Smith min-max setting, is at most 4π .

Now, note the following fact:

Claim 3.17. *Consider a Riemannian 3-manifold M . If Φ is a Simon-Smith family such that $[\Phi]$ is a p -sweepout, then the Simon-Smith min-max width $\mathbf{L}(\Lambda(\Phi))$ satisfies*

$$\mathbf{L}(\Lambda(\Phi)) \geq \omega_p(M).$$

Proof. This is true because it follows from definition that a homotopy in the sense of Simon-Smith min-max is continuous in the flat topology. \square

Thus, whenever $[\Xi|_{D_0}]$ is a p -sweepout,

$$4\pi = \mathbf{L}(\Lambda(\Xi|_{D_0})) \geq \omega_p(S^3, \bar{g}).$$

But by C. Nurser [Nur16], the first four widths of the unit 3-sphere are 4π , while the fifth is $2\pi^2$. Thus, $[\Xi|_{D_0}]$ is not a 5-sweepout. This finishes the proof of Proposition 3.15 \square

We postpone the detailed proof of Theorem 3.16 to §10. For now, let us outline the idea here. Let $1 \leq j \leq N$. By definition, D_j is the trace $T(C)$ of some genus 1 cap $C \subset \text{dmn}(\Xi)$ obtained at the k -th stage of min-max, for some $1 \leq k \leq K - 1$. By Proposition 3.12, there exists a strong deformation retraction of D_j onto C , so it suffices to show that for the inclusion map $i : C \hookrightarrow \text{dmn}(\Xi)$, the map i_* it induces between the first homology groups of C and $\text{dmn}(\Xi)$ is zero. By definition, the entire image of $[\Xi|_C]$ is d_0 -close to a single minimal torus in the \mathbf{F} -metric for currents. We will show in §10 that this means i_* indeed has to be the zero map, thereby proving Theorem 3.16.

Finally, as previously explained, Proposition 3.15 and Theorem 3.16 together contradict Lemma 3.14, thereby proving that the assumption $N \leq 4$ is false. This implies that at least 5 multiplicity one, embedded, minimal tori are detected through the repetitive min-max process. This completes the proof of Theorem 1.1.

Part II. Technical ingredients

4. MIN-MAX RESULTS I: THE SIMON-SMITH MIN-MAX THEOREM

The goal of this section is to prove Theorem 2.11 and subsequently Theorem 2.16.

4.1. Deformation in annuli.

Definition 4.1 ((ε, δ) -deformation). Given $\varepsilon, \delta > 0$, an open set $U \subset M$, and a generalized surface $\Sigma \in \mathcal{S}(M)$, we call an isotopy $\psi \in \mathcal{I}\mathfrak{s}(U)$ an (ε, δ) -deformation of Σ in U if

- (1) $\mathcal{H}^2(\psi(t, \Sigma)) \leq \mathcal{H}^2(\Sigma) + \delta$ for all $t \in [0, 1]$;
- (2) $\mathcal{H}^2(\psi(1, \Sigma)) \leq \mathcal{H}^2(\Sigma) - \varepsilon$.

We define $\mathfrak{a}(U; \varepsilon, \delta)$ to be the set of all generalized surfaces that do not admit (ε, δ) -deformations in U .

Definition 4.2 (Admissible annuli). Given a $K \in \mathbb{N}$ and $p \in M$, a collection of annuli centered in p

$$A(p; s_1, r_1), \dots, A(p; s_K, r_K),$$

is called K -admissible if $2r_{i+1} < s_i$ for all $i = 1, \dots, K-1$.

For $R \in (0, \infty]$, if $\sup_i r_i \leq R$, we will say that these annuli are of *outer radius at most R* .

Here, we adapt Pitts' combinatorial result [Pit81, Theorem 4.10] to fit our setting.

Proposition 4.3. *Given X a finite cubical subcomplex of some cube $I(m, k)$, $D \subset X$ a compact subset (not necessarily a subcomplex) of X , $\Phi : X \rightarrow \mathcal{S}^*(M)$ a Simon-Smith family and $R \in (0, \infty]$ and a pair positive numbers (ε, δ) , we set $K = K(m) := 3^{m3^m}$. Suppose that for every $x \in D$, there exists a K -admissible collection of annuli of outer radius at most R , denoted by $\{A_{x,i}\}_{i=1}^K$, so that*

$$\Phi(x) \notin \bigcup_{i=1}^K \mathfrak{a}(A_{x,i}; 2\varepsilon, \delta/2).$$

Then there exists $\Phi^ \in \Lambda(\Phi)$ in the homotopy class of Φ such that:*

- (1) for every $x \in D$,

$$\mathcal{H}^2(\Phi^*(x)) < \mathcal{H}^2(\Phi(x)) - \varepsilon + (3^m - 1)\delta;$$

(2) for every $x \in X \setminus D$,

$$\mathcal{H}^2(\Phi^*(x)) < \mathcal{H}^2(\Phi(x)) + 3^m \delta;$$

(3) for every $x \in X$, there exist t many points $p_1, \dots, p_t \in M$ such that $t \leq K(m)$ and

$$\Phi^*(x) \setminus (\overline{B}_{2R}(p_1) \cup \dots \cup \overline{B}_{2R}(p_t)) = \Phi(x) \setminus (\overline{B}_{2R}(p_1) \cup \dots \cup \overline{B}_{2R}(p_t)).$$

(4) Moreover, for another compact set $D' \subset X$ with $D' \cap D = \emptyset$, we can choose Φ^* such that $\Phi^*(x) = \Phi(x)$ holds for all $x \in D'$.

Proof. First, by our assumption, for each $x \in D$ and $i \in \{1, 2, \dots, K\}$, there exists an isotopy $\psi_{x,i} \in \mathfrak{Is}(A_{x,i})$ such that

- $\mathcal{H}^2(\psi_{x,i}(t, \Phi(x))) \leq \mathcal{H}^2(\Phi(x)) + \delta/2$ for all $t \in [0, 1]$;
- $\mathcal{H}^2(\psi_{x,i}(1, \Phi(x))) \leq \mathcal{H}^2(\Phi(x)) - 2\varepsilon$.

By Proposition 2.5, $[\Phi]$ is continuous in the \mathbf{F} -metric, and thus, for each $x \in D$, there exists an open neighborhood $O_x \subset X$ of x such that for every $y \in O_x$ and every $i \in \{1, 2, \dots, K\}$, we have

- $\mathcal{H}^2(\psi_{x,i}(t, \Phi(y))) \leq \mathcal{H}^2(\Phi(y)) + \delta$ for all $t \in [0, 1]$;
- $\mathcal{H}^2(\psi_{x,i}(1, \Phi(y))) \leq \mathcal{H}^2(\Phi(y)) - \varepsilon$.

Next, since the set D is compact, we can refine X so that it is a cubical subcomplex of $I(m, k')$ and every cell σ of X with $\sigma \cap D \neq \emptyset$ is contained within some O_{x_σ} where $x_\sigma \in D$. We denote by X_b the smallest cubical subcomplex of X containing

$$\{\sigma \in X : \sigma \cap D \neq \emptyset\}.$$

Moreover, to ensure that conclusion (4) holds, we also require that X'_b the smallest cubical subcomplex of X containing

$$\{\sigma \in X : \sigma \cap D' \neq \emptyset\},$$

satisfies the condition that $X_b \cap X'_b = \emptyset$.

It follows from Proposition 4.9 of [Pit81] that for every cell σ of X_b we can choose an annulus $A_\sigma \in \{A_{x_\sigma, 1}, \dots, A_{x_\sigma, K}\}$ and the corresponding isotopy $\psi_\sigma \in \mathfrak{Is}(A_\sigma)$ with the following property. For every pair of distinct cells σ, τ of X_b , if they are faces of a common cell γ of X_b , then

$$\text{dist}(A_\sigma, A_\tau) > 0.$$

Additionally, for every cell σ of X_b , we can choose a cut-off function

$$\eta_\sigma : X \rightarrow [0, 1],$$

such that:

- $\eta_\sigma(x) \equiv 0$ or every $x \in X$ with $\|x - c_\sigma\|_{\ell_\infty} \geq 3^{-k'}/2$,
- $\eta_\sigma(x) \equiv 1$ for every $x \in X$ with $\|x - c_\sigma\|_{\ell_\infty} \leq 3^{-k'}/3$,

where c_σ is the center of σ . Consequently, we obtain a continuous map

$$\hat{\psi}_\sigma : [0, 1] \times X \rightarrow \text{Diff}^\infty(M), \quad (t, x) \mapsto \psi_\sigma(t\eta_\sigma(x)).$$

Note that $\hat{\psi}_\sigma(t, x) = \text{Id}$ outside A_σ .

Finally, we define a continuous map

$$\varphi : [0, 1] \times X \rightarrow \text{Diff}^\infty(M)$$

by composing all the $\hat{\psi}_\sigma$

$$\varphi(t, x) := \circ_\sigma \text{ is a cell of } X_b \left(\hat{\psi}_\sigma(t, x) \right).$$

Note that no two annuli A_σ and A_τ intersect unless either $\hat{\psi}_\sigma(t, x)$ or $\hat{\psi}_\tau(t, x)$ is an identity map. Consequently, the order of σ 's does not affect the definition of φ . And we can define

$$\Phi^* := \varphi(1, \cdot)(\Phi_i(\cdot)) \in \Lambda(\Phi).$$

Consequently, by the definition of η_σ , for every $x \in X$, we have

- there are at most 3^m cells σ of X_b such that $\eta_\sigma(x) > 0$;
- if two cells σ, τ of X_b has $\eta_\sigma(x), \eta_\tau(x) > 0$, then

$$\text{dist}(A_\sigma, A_\tau) > 0;$$

- if $x \in X_b$, then there exists a cell σ_0 of X_b such that $\eta_{\sigma_0}(x) = 1$.

Therefore, if $x \in X \setminus D$, we have

$$\begin{aligned} \mathcal{H}^2(\Phi^*(x)) &= \sum_{\sigma: \eta_\sigma(x) > 0} \mathcal{H}^2(\Phi^*(x) \cap A_\sigma) + \mathcal{H}^2 \left(\Phi(x) \setminus \bigcup_{\sigma: \eta_\sigma(x) > 0} A_\sigma \right) \\ &\leq \sum_{\sigma: \eta_\sigma(x) > 0} (\mathcal{H}^2(\Phi(x) \cap A_\sigma) + \delta) + \mathcal{H}^2 \left(\Phi(x) \setminus \bigcup_{\sigma: \eta_\sigma(x) > 0} A_\sigma \right) \\ &\leq \mathcal{H}^2(\Phi(x)) + 3^m \delta. \end{aligned}$$

On the other hand, if $x \in D \subset X_b$, let σ_0 of X_b be a cell such that $\eta_{\sigma_0}(x) = 1$, and then we have

$$\begin{aligned} \mathcal{H}^2(\Phi^*(x)) &= \sum_{\sigma: \eta_\sigma(x) > 0} \mathcal{H}^2(\Phi^*(x) \cap A_\sigma) + \mathcal{H}^2 \left(\Phi(x) \setminus \bigcup_{\sigma: \eta_\sigma(x) > 0} A_\sigma \right) \\ &\leq \sum_{\sigma: \sigma \neq \sigma_0, \eta_\sigma(x) > 0} (\mathcal{H}^2(\Phi(x) \cap A_\sigma) + \delta) + (\mathcal{H}^2(\Phi(x) \cap A_{\sigma_0}) - \varepsilon) \\ &\quad + \mathcal{H}^2 \left(\Phi(x) \setminus \bigcup_{\sigma: \eta_\sigma(x) > 0} A_\sigma \right) \\ &\leq \mathcal{H}^2(\Phi(x)) + (3^m - 1)\delta - \varepsilon. \end{aligned}$$

In addition, for $x \in D' \subset X'_b$, we have that $\eta_\sigma(x) = 0$ holds for any cell σ of X_b , and thus,

$$\Phi^*(x) = \Phi(x),$$

which confirms the statement (4).

The statement (3) follows from the construction of the map φ . \square

4.2. Existence and regularity of a smooth minimal surface. From the definition of width, it is clear that one can choose a minimizing sequence $\{\Phi_i\}$ in $\Lambda(\Phi)$.

Recall that any minimizing sequence can be pulled-tight via a family of diffeomorphisms.

Proposition 4.4 (Existence of pulled-tight sequence, [CDL03, Proposition 3.1] [Pit81, Theorem 4.3]). *Given any minimizing sequence $\{\Phi_i\}$ in a homotopy class Λ , there exists a pulled-tight minimizing sequence $\{\Phi_i^*\}$ in Λ such that*

$$\mathbf{C}(\{\Phi_i^*\}) \subset \mathbf{C}(\{\Phi_i\}).$$

Proof. Following Pitts [Pit81, p. 153] (See also [CDL03, MN14]), there exists a continuous map

$$f : [0, 1] \times \{V \in \mathcal{V}_2(M) : \|V\|(M) \leq 2L\} \rightarrow \text{Diff}^\infty(M)$$

such that

- (1) $f(0, \cdot) = \text{id}$;
- (2) $f(t, V) = \text{id}$ for all $t \in [0, 1]$ if V is stationary;
- (3) $\|f(1, V)_\# V\|(M) < \|V\|(M)$ if V is not stationary.

By Proposition 2.5 (2), each $[\Phi_i] : X \rightarrow \mathcal{Z}_2(M; \mathbf{F}; \mathbb{Z}_2)$ is continuous, and thus, $|\Phi_i| : X \rightarrow \mathcal{V}_2(M)$ is continuous. Hence, for each i , we have a continuous map $\varphi_i : [0, 1] \times X \rightarrow \text{Diff}^\infty(M)$ defined by

$$\varphi_i(t, x) := f(t, |\Phi_i(x)|).$$

Let us define Φ_i^* by

$$\Phi_i^*(x) := \varphi_i(1, \|\Phi_i\|(x))(\Phi_i(x)).$$

It follows immediately from the construction that each Φ_i^* is homotopic to Φ_i , $\mathbf{C}(\{\Phi_i^*\}) \subset \mathbf{C}(\{\Phi_i\})$ and every varifold in $\mathbf{C}(\{\Phi_i^*\})$ is stationary. \square

For simplicity, let us assume that the minimizing sequence $\{\Phi_i\}$ is already pulled-tight. Now we aim to show that

$$\mathbf{C}(\{\Phi_i\}) \cap \mathcal{W}_L \neq \emptyset.$$

Indeed, by Proposition 4.8, it suffices to show that at least one varifold in $\mathbf{C}(\{\Phi_i\})$ is *almost minimizing in admissible annuli* as defined in Definition 4.6.

Definition 4.5 (Almost minimizing). For an open subset $U \subset M$ and a sequence of generalized surfaces $\{\Sigma_j\} \subset \mathcal{S}^*(M)$, a varifold $V \in \mathcal{V}_2(M)$ is called *almost minimizing with respect to $\{\Sigma_j\}$* if there exist two sequences of positive real numbers $\varepsilon_j \rightarrow 0$, $\delta_j \rightarrow 0$ such that

- $\mathbf{F}(|\Sigma_j|, V) < \varepsilon_j$;
- $\Sigma_j \in \mathbf{a}(U; \varepsilon_j, \delta_j)$.

Definition 4.6 (Almost minimizing in admissible annuli). Given $R \in (0, \infty]$, a sequence of generalized surfaces $\{\Sigma_j\} \subset \mathcal{S}^*(M)$ and a varifold $V \in \mathcal{V}_2(M)$, we say that a varifold V is *almost minimizing in every K -admissible collection of annuli of outer radius at most R with respect to $\{\Sigma_j\}$* , if there exists $\varepsilon_j \rightarrow 0$, $\delta_j \rightarrow 0$ such that

- $\mathbf{F}(|\Sigma_j|, V) < \varepsilon_j$;
- for any K -admissible collection of annuli $\{A_i\}_{i=1}^K$ each of outer radius at most R , and any Σ_j ,

$$\Sigma_j \in \bigcup_{i=1}^K \mathfrak{a}(A_i; \varepsilon_j, \delta_j).$$

Remark 4.7. Suppose that a varifold V is almost minimizing in every K -admissible collection of annuli of outer radius at most R . Then from the definition, it is clear that for any K -admissible collection of annuli $\{A_i\}_{i=1}^K$ each of radius at most R , V is stable in at least one A_i .

In particular, if $\Sigma = \text{spt } \|V\|$ is a smooth surface, we can say that Σ is *stable in every K -admissible collection of annuli of outer radius at most R* .

Proposition 4.8 (Regularity of varifolds almost minimizing in admissible annuli). *Given a closed Riemannian manifold (M, g) , $K \in \mathbb{N}^+$, $R \in (0, +\infty]$, a sequence $\{\Sigma_j\} \subset \mathcal{S}(M)$, and a stationary varifold $0 \neq V \in \mathcal{V}_2(M)$, suppose that:*

- (1) *We can choose for each $j \in \mathbb{N}^+$ a finite set P_j such that $\Sigma_j \setminus P_j$ is a smooth surface and*

$$P = \lim_{j \rightarrow \infty} P_j$$

in the Hausdorff sense where P is a finite set in M ;

- (2) *V is almost minimizing in every K -admissible collection of annuli of radius at most R with respect to $\{\Sigma_j\}$.*

Then $V \in \mathcal{W}_L$, where $L = \|V\|(M)$.

Proof. Let $\{\varepsilon_j\}$ and $\{\delta_j\}$ be the sequences of positive real numbers for V and $\{\Sigma_j\}$ as in Definition 4.6.

For every fixed $p \in M$, we can choose $R(p) \in (0, R)$ such that

$$B(p, R(p)) \cap (P \setminus \{p\}) = \emptyset.$$

Then by the definition of almost minimizing varifold in admissible annuli, for every $p \in M$ and every subsequence $\{\Sigma_{j_k}\}_{k=1}^\infty$, there exists a radius $r(p) \in (0, R(p))$ such that for every pair $(s, r) \in \mathbb{R}^2$ with $0 < s < r < r(p)$ and a further subsequence $\{\Sigma_{j_{k_l}}\}_{l=1}^\infty$, such that:

- (1) V is almost minimizing in $A(p; s, r)$ with respect to $\{\Sigma_{j_{k_l}}\}$.
- (2) For every $l \in \mathbb{N}^+$, $\Sigma_{j_{k_l}} \cap A(p; s, r)$ is a smooth surface.

Then, following the proof of [CDL03, Theorem 7.1], one can show that V is associated with a disjoint union of minimal surfaces with multiplicities, i.e., $V \in \mathcal{W}_L$. Note that $1/j$ should be replaced by ε_j and $1/(8j)$ by δ_j

therein. It is straightforward to verify that all the replacement arguments hold immediately. \square

The following proposition generalizes [CDL03, Proposition 5.3] to a multi-parameter Simon-Smith family, using Pitts's combinatorial argument [Pit81, Proposition 4.9]. It is worth noting that De Lellis-Ramic has proved a similar multi-parameter generalization in [DLR18a]. However, their homotopy class includes families not derived from isotopies, leading to a slight different definition of almost-minimizing compared to [CDL03] and ours. We provide a proof here for the sake of completeness.

Proposition 4.9 (Existence of varifolds almost minimizing in admissible annuli). *Suppose that X is a finite cubical subcomplex of some cube $I(m, k)$, $\Phi : X \rightarrow \mathcal{S}^*(M)$ is a Simon-Smith family and $\{\Phi_i\}$ is a pulled-tight sequence in $\Lambda(\Phi)$. Then for $K = K(m) := 3^{m3^m}$ and any $R \in (0, \infty]$, there exists an integer and a min-max subsequence $\Sigma_j := \Phi_{i_j}(x_j)$ such that Σ_j converges to some $W \in \mathbf{C}(\{\Phi_i\})$ in the varifold sense. Moreover, W is almost minimizing in every K -admissible collection of annuli of outer radius at most R with respect to $\{\Sigma_j\}$.*

Proof. We set

$$\varepsilon_i = \frac{1}{i}, \quad \delta_i = \frac{\varepsilon_i}{2K(m)}.$$

Then up to a subsequence, we may assume that

$$\sup_{x \in X} \mathcal{H}^2(\Phi_i(x)) < \mathbf{L}(\Lambda(\Phi)) + \delta_i.$$

Suppose for the sake of contradiction that for some large i and for every $\Phi_i(x)$ with $\mathcal{H}^2(\Phi_i(x)) \geq \mathbf{L}(\Lambda(\Phi)) - 2\varepsilon_i$ (and thus, $\Phi_i(x) \in \mathcal{S}(M)$), there exists a $K(m)$ -admissible collection of annuli $\{A_{x,j}\}_{j=1}^K$ of outer radius at most R such that

$$\Phi_i(x) \notin \bigcup_{i=1}^K \mathfrak{a}(A_{x,j}; 2\varepsilon_i, \delta_i/2).$$

It follows from Proposition 4.3 that there exists $\Phi_i^* \in \Lambda(\Phi)$ such that if $x \in X \setminus D$,

$$\begin{aligned} \mathcal{H}^2(\Phi_i^*(x)) &\leq \mathcal{H}^2(\Phi_i(x)) + 3^m \delta_i \\ &\leq (\mathbf{L}(\Lambda(\Phi)) - 2\varepsilon_i) + \varepsilon_i/2 \\ &< \mathbf{L}(\Lambda(\Phi)) - \delta_i. \end{aligned}$$

On the other hand, if $x \in D$,

$$\begin{aligned} \mathcal{H}^2(\Phi_i^*(x)) &\leq \mathcal{H}^2(\Phi_i(x)) + (3^m - 1)\delta_i - \varepsilon_i \\ &< \mathbf{L}(\Lambda(\Phi)) - \delta_i. \end{aligned}$$

In conclusion, we have $\sup_{x \in X} \mathcal{H}^2(\Phi_i^*(x)) < \mathbf{L}(\Lambda(\Phi)) - \delta_i$, a contradiction. This completes the proof. \square

Proof of Theorem 2.11 (1) without genus bound. By Proposition 4.4, there exists a pulled-tight sequence $\{\Phi_i\}$ in $\Lambda(\Phi)$.

Then it follows from Proposition 4.9, there exists an integer $K \in \mathbb{N}^+$ and a varifold $W \in \mathbf{C}(\{\Phi_i\})$ which is almost minimizing in every K -admissible collection of annuli with respect to some $\{\Sigma_j = \Phi_{i_j}(x_j)\}$.

Finally, as discussed in the second paragraph of Remark 2.8, we can choose P_j for each Σ_j such that

$$\sup_j \#P_j < \infty$$

and thus, up to a subsequence, P_j converges to a finite set P in the Hausdorff sense. By Proposition 4.8, we can conclude that

$$W \in \mathbf{C}(\{\Phi_i\}) \cap \mathcal{W}_L.$$

□

4.3. Genus bound. In the previous subsection, we have proved that the existence of a varifold $W \in \mathbf{C}(\{\Phi_i\}) \cap \mathcal{W}_L$ almost minimizing in every K -admissible collection of annuli of outer radius at most R with respect to $\{\Sigma_j\}$. In this and the next subsection, we take $R = +\infty$.

To justify Theorem 2.11 (1), it suffices to show that this W satisfies the genus estimates as follows.

Proposition 4.10 (Genus bound of varifolds almost minimizing in admissible annuli). *Given a closed Riemannian manifold (M, g) , $K \in \mathbb{N}^+$, $\mathfrak{g}_0 \in \mathbb{N}$, $R \in (0, \infty]$, a sequence $\{\Sigma_j\} \subset \mathcal{S}(M)$ and a stationary varifold $V \in \mathcal{V}_2(M)$, suppose that:*

- (1) *We can choose for each $j \in \mathbb{N}^+$ a finite set P_j such that $\Sigma_j \setminus P_j$ is a smooth surface and*

$$P = \lim_{j \rightarrow \infty} P_j$$

in the Hausdorff sense where P is a finite set in M ;

- (2) $\sup_j \mathfrak{g}(\Sigma_j) \leq \mathfrak{g}_0$;
 (3) *V is almost minimizing in every K -admissible collection of annuli of outer radius at most R with respect to $\{\Sigma_j\}$.*

Then $V \in \mathcal{W}_{L, \leq \mathfrak{g}_0}$ for $L = \|V\|(M)$.

The genus bound result essentially relies on the improved lifting lemma with multiplicity proved by Ketover [Ket19, Proposition 2.2], which we have tailored to our setting in the following proposition. More precisely, we do not assume that the limiting sequence consists of smooth surfaces.

Proposition 4.11 (Improved lifting lemma with multiplicity). *Given a closed Riemannian manifold (M, g) , $K \in \mathbb{N}^+$, a sequence $\{\Sigma_j\} \subset \mathcal{S}(M)$ and a stationary varifold $V \in \mathcal{V}_2(M)$, suppose that:*

- (1) *We can choose for each $j \in \mathbb{N}^+$ a finite set P_j such that $\Sigma_j \setminus P_j$ is a smooth surface and*

$$P = \lim_{j \rightarrow \infty} P_j$$

- in the Hausdorff sense where P is a finite set in M ;
- (2) V is almost minimizing in every K -admissible collection of annuli with respect to $\{\Sigma_j\}$.

Thus, by Proposition 4.8, there exists a disjoint set of smooth, connected, embedded minimal surface $\{\Gamma_i\}_{i=1}^l$ and a set of positive integers $\{m_i\}_{i=1}^l$ such that

$$V = m_1|\Gamma_1| + \cdots + m_l|\Gamma_l|.$$

We denote $\Gamma := \bigcup_{i=1}^l \Gamma_i$.

Assume that there exists a collection of points $\{q_i \in \Gamma_i\}_{i=1}^l$ and a collection of simple closed curves $\{\gamma_k\}_{k=1}^t$ contained in $\Gamma \setminus P$ so that for all $k_1 \neq k_2 \in \{1, 2, \dots, t\}$, if $\gamma_{k_1}, \gamma_{k_2} \subset \Gamma_i$ then $\gamma_{k_1} \cap \gamma_{k_2} = \{q_i\}$. Then there exists $\varepsilon_0 > 0$, so that for any $\varepsilon < \varepsilon_0$, there exists curves $\{\tilde{\gamma}_k\}_{k=1}^t$, a subsequence of $\{\Sigma_j\}$ (still labeled as $\{\Sigma_j\}$), and generalized surfaces $\{\tilde{\Sigma}_j\}$ each of which is obtained from Σ_j by finitely many neck-pinch surgeries outside P_j with the following properties.

- (1) Each $\tilde{\gamma}_k$ is homotopic to γ_k in Γ and $\tilde{\gamma}_k \subset T_\varepsilon(\gamma_k)$;
- (2) $|\tilde{\Sigma}_j| \rightarrow V$ as varifolds;
- (3) For each $k \in \{1, 2, \dots, t\}$, if $\tilde{\gamma}_k \subset \Gamma_i$, then exactly one of the following holds:
 - (a) $\pi^{-1}(\tilde{\gamma}_k) \cap T_\varepsilon(\Gamma_i) \cap \tilde{\Sigma}_j$ is a union of m_i closed curves, each of which projects via p onto $\tilde{\gamma}_k$ with degree one.
 - (b) $\pi^{-1}(\tilde{\gamma}_k) \cap T_\varepsilon(\Gamma_i) \cap \tilde{\Sigma}_j$ is a union of $m_i/2$ closed curves, each of which projects via the closest-point projection p onto $\tilde{\gamma}_k$ with degree two. In this case, Γ_i is non-orientable and $\pi^{-1}(\tilde{\gamma}_k)$ is a Mobius band.

Here, $\pi : T_\varepsilon(\Gamma_i) \rightarrow \Gamma_i$ is the closest-point projection onto Γ_i .

Indeed, in [Ket19, Section 4], the proof of lifting lemma, the only point where Ketover essentially relies on the smoothness of the sequence $\{\Sigma_i\}$ is in the application of the following Genus Collapse Lemma by Colding-Minicozzi.

Lemma 4.12 (Genus Collapse, [CM04, Lemma I.0.14]). *Given a 3-manifold (M, g) , a sequence of smooth surfaces $\{\Sigma_j\}$ of genus at most \mathfrak{g}_0 in (M, g) , there exists finitely many points in the manifold $\{x_i\}_{i=1}^m$ with $m \leq \mathfrak{g}_0$ and a subsequence of the surfaces, still denoted $\{\Sigma_j\}$, such that for all $x \notin \{x_i\}_{i=1}^m$, there is a radius $r_x > 0$ such that $\Sigma_j \cap B_{r_x}(x)$ is a union of planar domains, i.e.,*

$$\mathfrak{g}(\Sigma_j \cap B_{r_x}(x)) = 0.$$

Remark 4.13. In the original statement of Colding-Minicozzi, it appears they require each Σ_j is minimal; However, following their proof, as in Ketover's cited statement [Ket19, Lemma 3.4], this minimality condition is unnecessary.

In addition, the requirement by Colding-Minicozzi that the ambient manifold (M, g) be closed is also unnecessary.

Proof of Proposition 4.11. We choose ε_0 as in [Ket19, Set up 4.2] and fix $\varepsilon < \varepsilon_0$.

We choose $s \in (0, \text{dist}(P, \bigcup_k \gamma_k))$ so that for every $p \in P$, $B_s(p) \cap \Gamma$ is either a disk or an empty set. Then we set

$$\widetilde{M} := M \setminus B_s(P).$$

By possibly choosing a subsequence, without loss of generality, we may assume that every $\Sigma_j \cap \widetilde{M}$ is a smooth surface.

Note that all the arguments in [Ket19, Section 4] are confined to a neighborhood of $\bigcup_k \gamma_k$. With Lemma 4.12, we can follow these arguments verbatim to prove our proposition. \square

Proof of Proposition 4.10. In the previous subsection, we prove $V \in \mathcal{W}_L$, i.e.,

$$V = m_1 |\Gamma_1| + \cdots + m_l |\Gamma_l|$$

for a disjoint set of smooth, connected, embedded minimal surface $\{\Gamma_i\}_{i=1}^l$ and a set of positive integers $\{m_i\}_{i=1}^l$.

We follow the proof of [Ket19, Theorem 1.2] to select a set of curves $\{\alpha_k\}_{k=1}^t \subset \Gamma$. Since $\#P < \infty$, we may choose $\{\alpha_k\}_{k=1}^t \subset \Gamma \setminus P$. This allows us to apply our Proposition 4.11 to obtain a sequence of generalized surfaces $\{\widetilde{\Sigma}_j\}$. However, the only issue to continue the arguments from [Ket19] is that our $\widetilde{\Sigma}_j$ is not necessarily smooth everywhere.

To address this issue, for each j , we choose $s_j \rightarrow 0$ such that we can replace $\widetilde{\Sigma}_j \cap B_{s_j}(P_j) = \Sigma_j \cap B_{s_j}(P_j)$ in $\widetilde{\Sigma}_j$ with finitely many disks through surgeries and the removal of connected components, thereby obtaining a sequence of smooth surfaces $\{\widetilde{\Sigma}'_j\}$. For sufficiently large j , we have:

- (1) $B_{s_j}(P_j) \cap \pi^{-1}(\bigcup_{k=1}^t \alpha_k) = \emptyset$;
- (2) $\widetilde{\Sigma}'_j \setminus B_{s_j}(P_j) = \widetilde{\Sigma}_j \setminus B_{s_j}(P_j)$;
- (3) $|\widetilde{\Sigma}'_j| \rightarrow W$ as varifolds;
- (4) $\mathfrak{g}(\widetilde{\Sigma}'_j) \leq \mathfrak{g}(\widetilde{\Sigma}_j) \leq \mathfrak{g}(\Sigma_j) \leq \mathfrak{g}_0$.

Consequently, with the sequence of smooth surfaces $\{\widetilde{\Sigma}'_j\}$, we can follow the proof of [Ket19, Theorem 1.2] verbatim to obtain

$$(4.1) \quad \sum_{i=1}^l m_i \mathfrak{g}(\Gamma_i) \leq \mathfrak{g}_0.$$

\square

4.4. Multiplicity one theorem. The Multiplicity one result is an adaptation of Theorem B in [WZ24], which essentially follows from the PMC min-max theorem, Theorem 2.4 therein.

4.4.1. *PMC Simon-Smith min-max theory.* Let us first recall the concepts of \mathcal{A}^h -functional, \mathcal{VC} -space, $C^{1,1}$ boundary and strong \mathcal{A}^h -stationarity introduced in [WZ24].

Definition 4.14 (Prescribed mean curvature functional). For a function $h \in C^\infty(M)$, we define the *prescribed mean curvature functional associated with h* to be

$$\mathcal{A}^h : \mathcal{V}_2(M) \times \mathcal{C}(M) \rightarrow \mathbb{R}, \quad (V, \Omega) \mapsto \|V\|(M) - \int_{\Omega} h d\mathcal{H}^3.$$

Definition 4.15 (\mathcal{A}^h -stationary pairs). Given an open subset $U \subset M$, a pair $(V, \Omega) \in \mathcal{V}_2(M) \times \mathcal{C}(M)$ is *\mathcal{A}^h -stationary in U* , if for any vector field $X \in \mathcal{X}(U)$ with associated flow φ^t ,

$$\delta \mathcal{A}^h(V, \Omega)(X) := \frac{d}{dt} \Big|_{t=0} \mathcal{A}^h(\varphi_{\#}^t(V, \Omega)) = 0.$$

An \mathcal{A}^h -stationary pair (V, Ω) is *\mathcal{A}^h -stable in U* if for any vector field $X \in \mathcal{X}(U)$ with associated flow φ^t ,

$$\delta^2 \mathcal{A}^h(V, \Omega)(X, X) := \frac{d^2}{dt^2} \Big|_{t=0} \mathcal{A}^h(\varphi_{\#}^t(V, \Omega)) \geq 0.$$

Definition 4.16 (\mathcal{VC} -space). The *\mathcal{VC} -space on M* , denoted by $\mathcal{VC}(M)$, is the space of all pairs $(V, \Omega) \in \mathcal{V}_2(M) \times \mathcal{C}(M)$ such that there is a sequence $\{\Omega_k\} \subset \mathcal{C}(M)$ satisfying

$$|\partial^* \Omega_k| \rightarrow V, \quad \Omega_k \rightarrow \Omega.$$

Given two pairs (V, Ω) and (V', Ω') in $\mathcal{VC}(M)$, the \mathcal{F} -distance between them is

$$\mathcal{F}((V, \Omega), (V', \Omega')) := \mathbf{F}(V, V') + \mathcal{F}(\Omega, \Omega').$$

Definition 4.17 ($C^{1,1}$ almost embedding). Given an open subset $U \subset M$, a $C^{1,1}$ immersed surface $\phi : \Sigma \rightarrow U$ with $\phi(\partial\Sigma) \cap U = \emptyset$ is called a *$C^{1,1}$ almost embedded surface in U* provided that at every point $p \in \phi(\Sigma)$ where ϕ is not an embedding, there exists a neighborhood $W \subset U$ of p , such that

- (1) $\Sigma \cap \phi^{-1}(W)$ is a disjoint union of connected components $\bigsqcup_{i=1}^{\ell} \Gamma^i$;
- (2) $\phi : \Gamma^i \rightarrow W$ is a $C^{1,1}$ embedding for each i ;
- (3) For each pair $i \neq j$, $\phi(\Gamma^j)$ lies on one-side of $\phi(\Gamma^i)$ in W .

We call the set of all such p as the *touching set* of Σ . For simplicity, we will denote $\phi(\Sigma)$ by Σ and $\phi(\Gamma^i)$ by Γ^i .

Definition 4.18 ($C^{1,1}$ boundary). Given an open subset U of M , a $C^{1,1}$ almost embedded surface $\phi : \Sigma \rightarrow U$ and $\Omega \in \mathcal{C}(U)$, (Σ, Ω) is called a *$C^{1,1}$ boundary in U* , provided that Σ is orientable and

$$\phi_{\#}([\Sigma]) = \partial^* \Omega.$$

For a function $h \in C^\infty(M)$, a $C^{1,1}$ boundary (Σ, Ω) in U is called a *$C^{1,1}$ h -boundary* if $(|\Sigma|, \Omega)$ is \mathcal{A}^h -stationary in U .

Definition 4.19 (Strong \mathcal{A}^h -stationarity). Given an open subset U of M , a $C^{1,1}$ h -boundary (Σ, Ω) is said to be *strongly \mathcal{A}^h -stationary in U* provided the following holds.

For every $p \in U$ in the touching set of Σ , there exists a neighborhood $W \subset U$ of p , and a decomposition $\Sigma \cap W = \bigcup_{i=1}^{\ell} \Gamma^i$ into $\ell := \Theta^2(\Sigma, p) \geq 2$ connected disks with a natural ordering

$$\Gamma^1 \leq \Gamma^2 \leq \dots \leq \Gamma^{\ell}.$$

Let W^1 and W^{ℓ} be the bottom and the top components of $W \setminus \Sigma$. For $i = 1$ or ℓ and all $X \in \mathcal{X}(W)$ pointing into W^i along Γ^i , if $W^i \subset \Omega$, then

$$\delta \mathcal{A}^h(\Gamma^i, W^i)(X) \geq 0;$$

otherwise, $W^i \cap \Omega = \emptyset$ and

$$\delta \mathcal{A}^h(\Gamma^i, W \setminus W^i)(X) \geq 0.$$

To adapt their PMC Simon-Smith min-max theory to our setting, note that for every generalized surface $\Sigma \in \mathcal{S}^*(M)$, by definitions, $[\Sigma]$ can be viewed as the reduced boundary of a Caccioppoli set $\Omega \in \mathcal{C}(M)$ and its complement $M \setminus \Omega \in \mathcal{C}(M)$. Therefore, we can extend the definition of \mathcal{E} in [WZ24, Section 2] to

$$\tilde{\mathcal{E}} := \{(\Sigma, \Omega) : \Sigma \in \mathcal{S}^*(M), [\Sigma] = \partial^* \Omega\}.$$

Definition 4.20 (PMC Simon-Smith family). Let X be a cubical subcomplex of some $I(m, j)$. A map $(\Phi, \Omega) : X \rightarrow \tilde{\mathcal{E}}$ is called a *PMC Simon-Smith family*, provided that:

- (1) its first component $\Phi : X \rightarrow \mathcal{S}^*(M)$ is a Simon-Smith family;
- (2) its second component $\Omega : X \rightarrow \mathcal{C}(M)$ is continuous.

Definition 4.21 (Relative PMC homotopy class). Let $Z \subset X$ be cubical subcomplexes of some $I(m, j)$. Two PMC Simon-Smith families (Φ_0, Ω_0) and (Φ_1, Ω_1) parametrized by the same parameter space X with $(\Phi_0, \Omega_0)|_Z = (\Phi_1, \Omega_1)|_Z$ are said to be *homotopic relative to $(\Phi_0, \Omega_0)|_Z$* to each other if there exists a continuous map

$$\varphi : [0, 1] \times X \rightarrow \text{Diff}^{\infty}(M)$$

such that

- $\varphi(0, x) = \text{Id}$ for all $x \in X$,
- $\varphi(t, z) = \text{Id}$ for all $t \in [0, 1]$ and $z \in Z$,
- $(\varphi(1, x)(\Phi_0(x)), \varphi(1, x)(\Omega_0(x))) = (\Phi_1(x), \Omega_1(x))$ for all $x \in X$.

The set of all families homotopic relative to $(\Phi, \Omega)|_Z$ to a PMC Simon-Smith family (Φ, Ω) is called the *relative (X, Z) -homotopy class of (Φ, Ω)* , and is denoted by $\Lambda_Z(\Phi, \Omega)$.

Definition 4.22. Given a (X, Z) -relative homotopy class Λ_Z of a PMC Simon-Smith family and a function $h \in C^{\infty}(M)$, its *h -width* is defined by

$$\mathbf{L}^h(\Lambda_Z) := \inf_{(\Phi, \Omega) \in \Lambda_Z} \sup_{x \in X} \mathcal{A}^h(|\Phi|(x), \Omega(x)), .$$

Definition 4.23. A sequence $\{(\Phi_i, \Omega_i)\}$ in Λ_Z is said to be *minimizing associated with* $h \in C^\infty(M)$ if

$$\limsup_{i \rightarrow \infty} \sup_{x \in X} \mathcal{A}^h(|\Phi|(x), \Omega(x)) = \mathbf{L}^h(\Lambda_Z).$$

If $\{(\Phi_i, \Omega_i)\}$ is a minimizing sequence associated with h and $\{x_i\} \subset X$ satisfies

$$\lim_{i \rightarrow \infty} \mathcal{A}^h(|\Phi_i(x_i)|, \Omega_i(x_i)) = \mathbf{L}^h(\Lambda_Z),$$

then $\{\Phi_i(x_i)\}$ is called a *min-max sequence associated with* h .

For a minimizing sequence $\{(\Phi_i, \Omega_i)\}$, we define its *critical set associated with* h to be the set of all subsequential varifold-limit of its min-max sequences:

$$\begin{aligned} \mathcal{C}^h(\{(\Phi_i, \Omega_i)\}) := \{ & (V, \Omega) = \lim_j (|\Phi_{i_j}(x_j)|, \Omega_{i_j}(x_j)) \in \mathcal{VC}(M) : x_j \in X, \\ & \mathcal{A}^h(V, \Omega) = \mathbf{L}^h(\Lambda_Z)\}. \end{aligned}$$

Definition 4.24 ($(\mathcal{A}^h, \varepsilon, \delta)$ -deformation). Given $\varepsilon, \delta > 0$, an open set $U \subset M$, a function $h \in C^\infty(M)$ and a pair $(\Sigma, \Omega) \in \tilde{\mathcal{E}}$, we call an isotopy $\psi \in \mathfrak{Is}(U)$ an $(\mathcal{A}^h, \varepsilon, \delta)$ -deformation of Σ in U associated with h if

- (1) $\mathcal{A}^h(\psi(t)_\#(|\Sigma|, \Omega)) \leq \mathcal{A}^h(|\Sigma|, \Omega) + \delta$ for all $t \in [0, 1]$;
- (2) $\mathcal{A}^h(\psi(t)_\#(|\Sigma|, \Omega)) \leq \mathcal{A}^h(|\Sigma|, \Omega) - \varepsilon$.

We define $\mathfrak{a}^h(U; \varepsilon, \delta)$ to be the set of all pairs in $\tilde{\mathcal{E}}$ that do not admit $(\mathcal{A}^h, \varepsilon, \delta)$ -deformations in U .

Definition 4.25 (Almost minimizing in admissible annuli). Given a sequence of pairs $\{(\Sigma_j, \Omega_j)\} \subset \tilde{\mathcal{E}}$ and a pair $(V, \Omega) \in \mathcal{VC}(M)$, we say that (V, Ω) is \mathcal{A}^h -almost minimizing in every K -admissible collection of annuli of outer radius at most R with respect to $\{(\Sigma_j, \Omega_j)\}$, if there exists $\varepsilon_j \rightarrow 0$, $\delta_j \rightarrow 0$ such that

- $\mathcal{F}((|\Sigma_j|, \Omega_j), (V, \Omega)) < \varepsilon_j$;
- for any K -admissible collection of annuli $\{A_i\}_{i=1}^K$, and any pair (Σ_j, Ω_j) ,

$$(\Sigma_j, \Omega_j) \in \bigcup_{i=1}^K \mathfrak{a}^h(A_i; \varepsilon_j, \delta_j).$$

Remark 4.26. Similar to Remark 4.7, if $\Sigma = \text{spt } \|V\|$ is a minimal surface, then for every K -admissible collection of annuli, $(|\Sigma|, \Omega)$ is \mathcal{A}^h -stable in at least one annulus.

Theorem 4.27 (Relative PMC min-max theorem, [WZ24, Theorem 2.4]). *Given $Z \subset X$ cubical subcomplexes of some $I(m, j)$, a function $h \in C^\infty(M)$ and a PMC Simon-Smith family (Φ_0, Ω_0) , suppose that*

$$\mathbf{L}^h(\Lambda_Z(\Phi_0, \Omega_0)) > \max\{\sup_{z \in Z} \mathcal{A}^h(|\Phi_0(z)|, \Omega_0(z)), 0\}.$$

Then there exists a minimizing sequence $\{(\Phi_i, \Omega_i)\} \subset \Lambda_Z(\Phi_0, \Omega_0)$, and a strongly \mathcal{A}^h -stationary, $C^{1,1}$ h -boundary (Σ, Ω) with $(|\Sigma|, \Omega) \in \mathbf{C}^h(\{(\Phi_i, \Omega_i)\})$, such that

$$\mathcal{A}^h(\Sigma, \Omega) = \mathbf{L}^h(\Lambda_Z(\Phi_0, \Omega_0)).$$

Moreover, there exists an integer $K = K(m)$ and a min-max subsequence $(\Sigma_j, \Omega_j) := (\Phi_{i_j}(x_j), \Omega_{i_j}(x_j))$ such that $(|\Sigma|, \Omega)$ is \mathcal{A}^h -almost minimizing in every K -admissible collection of annuli with respect to $\{(\Sigma_j, \Omega_j)\}$.

Proof. Firstly, as in the proof of our Proposition 4.9, one can adapt the proof of [WZ24, Theorem 3.8] to obtain a pair $(V_0, \Omega_0) \in \mathcal{VC}(M)$ which is \mathcal{A}^h -almost minimizing in small annuli with respect to $\{(\Sigma_j, \Omega_j) = (\Phi_{i_j}(x_j), \Omega_{i_j}(x_j))\}$.

Then, as discussed in the second paragraph of Remark 2.8, we can choose P_j for each Σ_j such that

$$\sup_j \#P_j < \infty$$

and thus, up to a subsequence, P_j converges to a finite set P in the Hausdorff sense. Since $\Sigma_j \setminus P_j$ is a smooth surface, one can follow verbatim the proof of [WZ24, Theorem 4.7] and conclude that (V_0, Ω_0) is $C^{1,1}$ and strongly \mathcal{A}^h -stationary boundary in $M \setminus P'$ for some finite set $P' \supset P$.

Finally, one can follow their $C^{1,1}$ -version of removal singularity as in the proof of [WZ24, Theorem 2.4] to conclude the theorem. \square

4.4.2. *Proof of Theorem 2.11 (2).* Let $\mathcal{W}_{L, \leq g_0}^m$ be the set of all varifolds $W \in \mathcal{W}_{L, \leq g_0}$, whose support is a smooth embedded minimal surface Σ , such that for every $K(m) := 3^{m3^m}$ -admissible collection of annuli, Σ is stable in at least one annulus. Note that this is just a reformulation of *Property (R')* in [WZ24]. Proposition 4.8 with $R = +\infty$ implies that

$$W \in \mathbf{C}(\{\Phi_i\}) \cap \mathcal{W}_{L, \leq g_0}^m.$$

By Sharp's compactness theorem [Sha17] (See also Theorem [WZ24, Theorem 7.1]), the set $\mathcal{W}_{L, \leq g_0}^m$ is compact. In particular, if g is a bumpy metric on M , then $\mathcal{W}_{L, \leq g_0}^m$ is a finite set.

As in the proof of [WZ24, Theorem 7.3], it suffices to work on bumpy metrics g and the general results follow from a approximation by bumpy metrics.

By Proposition 2.5, for any Simon-Smith family $\Phi : X \rightarrow \mathcal{S}^*(M)$, the map $[\Phi]$ is continuous in the \mathbf{F} -metric, and thus, it can be lifted to a PMC Simon-Smith family

$$(\tilde{\Phi}, \tilde{\Omega}) : \tilde{X} \rightarrow \tilde{\mathcal{E}}$$

where \tilde{X} is a double-cover of X .

Hence, one can follow verbatim the proofs of Theorem 7.2 and Theorem 7.3 of [WZ24] through replacing Theorem 2.4 therein by our Theorem 4.27. Note that by Remark 4.26, Theorem 4.27 also implies the required *Property (R')* as defined in [WZ24, Corollary 3.11]. This finishes the proof of Theorem 2.11 (2).

4.5. Refinement of the critical set by Marques-Neves. Conclusion (3) is adapted from [MN21, Theorem 4.7], which essentially follows from the combinatorial argument of [Pit81, Theorem 4.10]. Note that our Definition 4.1 of (ε, δ) deformation is continuous, so we do not need to appeal to the intricate interpolation lemmas as in [MN21].

Let us first recall a useful lemma proved in Marques-Neves [MN21, Lemma 4.5].

Lemma 4.28. *For every $r, L > 0$ and $m \in \mathbb{N}$, there exists $\tilde{\eta} = \tilde{\eta}(r, L, m) > 0$ so that for every $V, T \in \mathcal{V}_2(M)$ with $\mathbf{M}(V) \leq 2L, \mathbf{M}(T) \leq 2L, V$ stationary,*

$$V \llcorner (M \setminus (\overline{B_{2\tilde{\eta}}}(p_1) \cup \cdots \cup \overline{B_{2\tilde{\eta}}}(p_t))) = T \llcorner (M \setminus (\overline{B_{2\tilde{\eta}}}(p_1) \cup \cdots \cup \overline{B_{2\tilde{\eta}}}(p_t)))$$

for some collection $\{p_1, \dots, p_t\} \subset M, t \leq K(m) := 3^{m3^m}$, and

$$\|V\|(M) - \tilde{\eta} \leq \|T\|(M) \leq \|V\|(M) + \tilde{\eta},$$

then $\mathbf{F}(V, T) < r/4$.

With this lemma, we can fix $\tilde{\eta}$ from Lemma 4.28 using the parameters r, L, m from Theorem 2.11.

As discussed in Remark 4.7, the support of the varifold generated from the min-max theory is also stable in admissible annuli. Let $\mathcal{W}_{L, \leq g_0}^{m, \tilde{\eta}}$ be the set of all varifolds $W \in \mathcal{W}_{L, \leq g_0}$, whose support is a smooth embedded minimal surface Σ , such that for every $K(m) := 3^{m3^m}$ -admissible collection of annuli of outer radius at most $\tilde{\eta}$, Σ is stable in at least one annulus.

Let $\{\Phi_i\}$ be a pulled-tight sequence from Theorem 2.11 (1). In the following, we will deform $\{\Phi_i\}$ and obtain a new pulled-tight sequence $\{\tilde{\Phi}_i\}$ such that there exists $\eta > 0$ such that for all sufficiently large i ,

$$(4.2) \quad \mathcal{H}^2(\Phi_i^*(x)) \geq L - \eta \implies |\Phi_i^*(x)| \in \mathbf{B}_r^{\mathbf{F}}(\mathcal{W}_{L, \leq g_0}^{m, \tilde{\eta}}).$$

Note that

$$\mathcal{W}_{L, \leq g_0}^m \subset \mathcal{W}_{L, \leq g_0}^{m, \tilde{\eta}} \subset \mathcal{W}_{L, \leq g_0}.$$

It follows from Remark 4.7 and Proposition 4.10 (with $R = \tilde{\eta}$) that for each $V \in \mathbf{C}(\{\Phi_i\}) \setminus \mathcal{W}_{L, \leq g_0}^{m, \tilde{\eta}}$, there exists a $K(m) := 3^{m3^m}$ -admissible collection of annuli of outer radius at most $\tilde{\eta}$, denoted by $\{A_{V,j}\}_{j=1}^{K(m)}$, such that V is not almost-minimizing in any $A_{V,j}$ with respect to any min-max subsequence. In particular, for every $V \in \mathbf{C}(\{\Phi_i\}) \setminus \mathcal{W}_{L, \leq g_0}^{m, \tilde{\eta}}$, there exists $\varepsilon_V > 0$ and $N_V \in \mathbb{N}^+$ with the following property: For every $\delta > 0$ and every $\Phi_i(x)$ with $i \geq N_V$, if $\mathbf{F}(V, |\Phi_i(x)|) < \varepsilon_V$, then

$$\Phi_i(x) \notin \bigcup_{j=1}^K \mathbf{a}(A_{V,j}; \varepsilon_V, \delta).$$

Let $\mathcal{K} := \mathbf{C}(\{\Phi_i\}) \setminus \mathbf{B}_{r/2}^{\mathbf{F}}(\mathcal{W}_{L, \leq g_0}^{m, \tilde{\eta}})$. Since \mathcal{K} is compact, we can find a finite set $\{V_k\}_{k=1}^\nu \subset \mathcal{K}$ such that

$$\mathcal{K} \subset \bigcup_{k=1}^\nu \mathbf{B}_{\frac{\varepsilon_k}{4}}^{\mathbf{F}}(V_k),$$

where $\varepsilon_k := \varepsilon_{V_k}$. We set

$$\varepsilon^{(1)} := \frac{\min_k \varepsilon_k}{10}, \quad N^{(1)} := \max_k N_{V_k}.$$

Since $\{\Phi_i\}$ is pulled-tight, we can choose an integer $N^{(2)} \geq N^{(1)}$ and $\varepsilon^{(2)} \in (0, \varepsilon^{(1)})$ such that for each $i \geq N^{(2)}$, and any $x \in X$, if

$$\mathcal{H}^2(\Phi_i(x)) \geq \sup_{y \in X} \mathcal{H}^2(\Phi_i(y)) - 2\varepsilon^{(2)}, \quad \mathbf{F}(|\Phi_i(x)|, \mathcal{W}_{L, \leq \mathfrak{g}_0}^{m, \tilde{\eta}}) \geq \frac{r}{2},$$

then $\mathbf{F}(|\Phi_i(x)|, V_{k_{i,x}}) < \frac{\varepsilon^{k_{i,x}}}{2}$ for some $k_{i,x} \in \{1, 2, \dots, \nu\}$, and thus, for every $\delta > 0$, we have

$$(4.3) \quad \Phi_i(x) \notin \bigcup_{j=1}^K \mathfrak{a}(A_{V_{k_{i,x},j}; \varepsilon_{V_{k_{i,x}}}}, \delta).$$

We denote by D_i the set of all such x (“bad points”) for each $i \geq N^{(2)}$.

We set

$$\varepsilon := \frac{\min \{ \varepsilon^{(1)}, \varepsilon^{(2)}, r, \tilde{\eta} \}}{10}.$$

For each $i \geq N^{(2)}$, we also choose $\delta_i > 0$ such that

$$\lim_i \delta_i = 0.$$

By (4.3), for every $i \geq N^{(2)}$, if $x \in D_i$, then

$$(4.4) \quad \Phi_i(x) \notin \bigcup_{j=1}^K \mathfrak{a}(A_{V_{k_{i,x},j}; 2\varepsilon^{(1)}}, \delta_i/2).$$

Hence, by Proposition 4.3 with the parameters $R = \tilde{\eta}$, $\varepsilon = \varepsilon^{(1)}$, $\delta = \delta_i$ and $D = D_i$, we obtain $\Phi_i^* \in \Lambda(\Phi)$. Moreover, for every $x \in X$,

$$\mathcal{H}^2(\Phi_i^*(x)) \leq \mathcal{H}^2(\Phi_i^*(x)) + 3^m \cdot \delta_i.$$

Since $\delta_i \rightarrow 0$, $\{\Phi_i^*\}$ is also a minimizing sequence.

To show that $\mathbf{C}(\{\Phi_i^*\}) \subset \mathbf{B}_{\frac{3}{4}r}^{\mathbf{F}}(\mathcal{W}_{L, \leq \mathfrak{g}_0}^{m, \tilde{\eta}})$, we need to prove the following claim.

Claim 4.29. *For sufficiently large i , if $\mathcal{H}^2(\Phi_i^*(x)) \geq \sup_{y \in X} \mathcal{H}^2(\Phi_i(y)) - \varepsilon/4$, then*

$$\mathbf{F}(|\Phi_i(x)|, \mathcal{W}_{L, \leq \mathfrak{g}_0}^{m, \tilde{\eta}}) < r/2.$$

Proof. Since $\lim_i \delta_i = 0$, for sufficiently large i , we have

$$\delta_i < \frac{\varepsilon}{1000K(m)}.$$

By the definition of D_i , for $x \in X$, if $\mathbf{F}(|\Phi_i(x)|, \mathcal{W}_{L, \leq \mathfrak{g}_0}^{m, \tilde{\eta}}) \geq r/2$, we have either $\Phi_i(x) \in D_i$ or $\mathcal{H}^2(\Phi_i(x)) < \sup_{y \in X} \mathcal{H}^2(\Phi_i(y)) - 2\varepsilon^{(2)}$.

If $\Phi_i(x) \in D_i$, i.e.,

$$\mathcal{H}^2(\Phi_i(x)) \geq \sup_{y \in X} \mathcal{H}^2(\Phi_i(y)) - 2\varepsilon^{(2)}, \quad \mathbf{F}(|\Phi_i(x)|, \mathcal{W}_{L, \leq g_0}^{m, \tilde{\eta}}) \geq r/2,$$

by Proposition 4.3, we have

$$\mathcal{H}^2(\Phi_i^*(x)) \leq \mathcal{H}^2(\Phi_i(x)) - \varepsilon^{(2)} + (3^m - 1)\delta_i < \sup_y \mathcal{H}^2(\Phi_i(y)) - \varepsilon/4.$$

If $\mathcal{H}^2(\Phi_i(x)) < \sup_{y \in X} \mathcal{H}^2(\Phi_i(y)) - 2\varepsilon^{(2)}$, then by Proposition 4.3 again, we have

$$\mathcal{H}^2(\Phi_i^*(x)) \leq \mathcal{H}^2(\Phi_i(x)) + 3^m \cdot \delta_i < \sup_{y \in X} \mathcal{H}^2(\Phi_i(y)) - 2\varepsilon^{(2)} + 3^m \cdot \delta_i < \mathcal{H}^2(\Phi_i(x)) - \varepsilon/4.$$

This finishes the proof of the claim. \square

Now, every $W \in \mathbf{C}(\{\Phi_i^*\})$ is a varifold limit of some subsequence $\Phi_{i_j}^*(x_{i_j})$ for $j \rightarrow \infty$. After passing to a subsequence, $\Phi_{i_j}(x_{i_j})$ converges to a varifold V . By Proposition 4.3, we know that there exist some points $p_1, \dots, p_t \in M$ ($t \leq K(m)$) such that

$$V \llcorner (M \setminus (\overline{B_{2\tilde{\eta}}}(p_1) \cup \dots \cup \overline{B_{2\tilde{\eta}}}(p_t))) = W \llcorner (M \setminus (\overline{B_{2\tilde{\eta}}}(p_1) \cup \dots \cup \overline{B_{2\tilde{\eta}}}(p_t))).$$

Furthermore, since

$$\begin{aligned} L \geq \|V\|(M) &= \lim_{j \rightarrow \infty} \mathcal{H}^2(\Phi_{i_j}(x_{i_j})) \geq \lim_{j \rightarrow \infty} \mathcal{H}^2(\Phi_{i_j}^*(x_{i_j})) - K(m) \cdot \delta_{i_j} \\ &= \|W\|(M) = L, \end{aligned}$$

by Lemma 4.28, we can conclude that

$$\mathbf{F}(V, W) < r/4.$$

The previous claim implies that $\mathbf{F}(V, \mathcal{W}_{L, \leq g_0}^{m, \tilde{\eta}}) < r/2$, so we have

$$\mathbf{F}(W, \mathcal{W}_{L, \leq g_0}^{m, \tilde{\eta}}) < \frac{3}{4}r.$$

Note that the new minimizing sequence $\{\Phi_i^*\}$ is not necessarily pulled-tight. However, by Proposition 4.4, we can obtain a pulled-tight minimizing sequence $\{\tilde{\Phi}_i\}$ with

$$\mathbf{C}(\{\tilde{\Phi}_i\}) \subset \mathbf{C}(\{\Phi_i^*\}) \subset \mathbf{B}_{\frac{3}{4}r}^{\mathbf{F}}(\mathcal{W}_{L, \leq g_0}^{m, \tilde{\eta}}).$$

The conclusion follows immediately from a suitable choice of η .

4.6. Proof of Theorem 2.16. Since $\mathbf{L}(\Lambda_Z(\Phi)) > \sup_{z \in Z} \mathcal{H}^2(\Phi(z))$, for each $\Phi' \in \Lambda_Z(\Phi)$, one can restrict all the deformations in the previous subsections to occur away from the compact set

$$X' := \left\{ x \in X : \mathcal{H}^2(\Phi(x)) \leq \frac{\mathbf{L}(\Lambda_Z(\Phi)) + \sup_{z \in Z} \mathcal{H}^2(\Phi(z))}{2} \right\}.$$

For example, all deformations can be composed with a cut-off function on X that vanishes on X' .

Consequently, all the results follow immediately.

5. THE FAMILY Ψ

5.1. The space of Clifford Tori. Let us recall parametrization of the space \mathcal{C} of unoriented Clifford tori. A Clifford torus is uniquely determined by an unordered pair of orthogonal unoriented 2-subspaces in \mathbb{R}^4 . Namely, given two orthogonal 2-subspaces P and Q , the Clifford torus they define consists of the points on the unit sphere \mathbb{S}^3 that are equidistant from the equators $P \cap \mathbb{S}^3$ and $Q \cap \mathbb{S}^3$. Note that for every 2-subspace of \mathbb{R}^4 , there exists exactly one orthogonal 2-subspace Q . Hence, to choose an inside direction for a Clifford torus, i.e. to assign an orientation, one simply needs to specify one 2-plane P , so the space of oriented Clifford tori is $G_2(\mathbb{R}^4)$, the space of unoriented 2-planes in \mathbb{R}^4 .

By [HO80, §1], the set $G_2^+(\mathbb{R}^4)$ of *oriented* 2-planes in \mathbb{R}^4 can be identified with the set

$$Q_2 := \{[z_1 : z_2 : z_3 : z_4] : z_1^2 + z_2^2 + z_3^2 + z_4^2 = 0\} \subset \mathbb{C}\mathbb{P}^3.$$

Namely, for an oriented orthonormal basis (u, v) of a 2-plane in \mathbb{R}^4 , we can define

$$(z_1, z_2, z_3, z_4) := u + iv.$$

Moreover, there is a biholomorphism

$$F : \mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1 \rightarrow Q_2,$$

defined in [HO80, §2]:

$$\begin{aligned} F([w_1 : 1], [w_2 : 1]) &= (1 + w_1 w_2, i(1 - w_1 w_2), w_1 - w_2, -i(w_1 + w_2)), \\ F([w_1 : 0], [w_2 : 1]) &= (w_2, -iw_2, 1, -i), \\ F([w_1 : 1], [w_2 : 0]) &= (w_1, -iw_1, -1, -i), \\ F([w_1 : 0], [w_2 : 0]) &= (1, -i, 0, 0). \end{aligned}$$

This gives a homeomorphism between $S^2 \times S^2$ and $G_2^+(\mathbb{R}^4)$. It follows from a straightforward calculation (see [Nur16, §3.4.3]) that $G_2(\mathbb{R}^4)$, or the space of oriented Clifford tori, is homeomorphic to $S^2 \times S^2 / \sim$ with $(x, y) \sim (-x, -y)$. Moreover, one can verify that (x, y) and $(x, -y)$ correspond to two *orthogonal* 2-subspaces (meaning every vector in one plane is perpendicular to every vector in the other plane), so the set \mathcal{C} of unoriented Clifford tori is given by $S^2 \times S^2 / \sim$ with

$$(x, y) \sim (-x, -y) \sim (x, -y) \sim (-x, y),$$

which is an $\mathbb{R}\mathbb{P}^2 \times \mathbb{R}\mathbb{P}^2$ (see also [Whi91, §4]).

5.2. Definition of Ψ . In this section we define the 9-parameter family Ψ . We will reuse many of the notations and presentations in [MN14] by Marques-Neves, [Nur16] by C. Nurser, and an unpublished manuscript [Mar23] by F. Marques, in which he constructed a 9-parameter family (in a slightly different way than us) that is an 8-sweepout to prove that the 8-width of the unit 3-sphere is $2\pi^2$.

Let \mathbb{S}^3 be the unit 3-sphere. Let $Z = \mathbb{RP}^2 \times \mathbb{RP}^2$ parametrize the space of unoriented Clifford tori, and \tilde{Z} denote the space of oriented Clifford tori. Then \tilde{Z} is connected, and there is a natural double-cover map $q : \tilde{Z} \rightarrow Z$ given by forgetting the orientation. Let $\sigma : \tilde{Z} \rightarrow \tilde{Z}$ be the nontrivial deck transformation. Note that $\sigma(\sigma(z)) = z$.

For each $z \in \tilde{Z}$, let $\Sigma(z)$ be the corresponding oriented Clifford torus, and let $A(z)$ and $A^*(z)$ denote the two connected components of $\mathbb{S}^3 \setminus \Sigma(z)$ such that $A(z)$ and $A^*(z)$ vary continuously in z . Let $N(z)$ denote the unit normal vector field on $\Sigma(z)$ that points into $A^*(z)$. Note that for every $z \in \tilde{Z}$, and $p \in \Sigma(z)$,

$$A(z) = A^*(\sigma(z)), \quad N(z)_{-p} = -N(z)_p, \quad N(z) = -N(\sigma(z)).$$

In [Nur16, §3.4.2], C. Nurser defined for each oriented Clifford torus $\Sigma(z)$ a 5-sweepout $\Phi_5^{\Sigma(z)} : \mathbb{RP}^5 \rightarrow \mathcal{Z}_2(\mathbb{S}^3; \mathbb{Z}_2)$. Note that as we will see below, each element of $\Phi_5^{\Sigma(z)}$ can actually be represented as a closed set in \mathbb{S}^3 . Here, we view \mathbb{RP}^5 as the space $\mathbb{B}^4 \times [-\pi, \pi]$ with boundary points (v, t) identified with $(-v, -t)$. Then, we can construct the corresponding family of closed surfaces.

Proposition 5.1. *For each oriented Clifford torus $\Sigma(z)$, with $z \in \tilde{Z}$, there exists a family $\Psi_5^{\Sigma(z)}$ of closed subsets of \mathbb{S}^3 , parametrized by \mathbb{RP}^5 , such that:*

- The maps $[\Psi_5^{\Sigma(z)}]$ and $[\Phi_5^{\Sigma(z)}]$ (defined by C. Nurser) are the same.
- For every $[(v, t)] \in \mathbb{RP}^5$ and $z \in \tilde{Z}$,

$$\Psi_5^{\Sigma(z)}([(v, t)]) = \Psi_5^{\Sigma(\sigma(z))}([(v, -t)]).$$

We will prove this proposition in the following subsections.

Let us assume this proposition for now, and consider the $\mathbb{RP}^5 \times \tilde{Z}$ -family $\tilde{\Psi}$ of closed subsets of \mathbb{S}^3 defined by

$$\tilde{\Psi}([(v, t)], z) := \Psi^{\Sigma(z)}([(v, t)]).$$

Moreover, consider the \mathbb{Z}_2 -action on $\mathbb{RP}^5 \times \tilde{Z}$ where the non-trivial element of \mathbb{Z}_2 acts by

$$(5.1) \quad ([(v, t)], z) \mapsto ([(v, -t)], \sigma(z)),$$

and let Y be the quotient space. Using Proposition 5.1, the family $\tilde{\Psi}$ immediately induces a family Ψ of closed subsets of \mathbb{S}^3 , with parameter space Y . Furthermore, Y is an \mathbb{RP}^5 -bundle over $Z = \mathbb{RP}^2 \times \mathbb{RP}^2$.

5.2.1. *Conformal maps and level surfaces.* For each $v \in \mathbb{B}^4$, we consider the conformal map

$$F_v : \mathbb{S}^3 \rightarrow \mathbb{S}^3, \quad F_v(x) = \frac{(1 - |v|^2)}{|x - v|^2}(x - v) - v.$$

For each $(v, z) \in \mathbb{B}^4 \times \tilde{Z}$, we define $d(v, z) : \mathbb{S}^3 \rightarrow \mathbb{R}$ to be the signed distance to $F_v(\Sigma(z))$, which is positive in $F_v(A^*(z))$ and negative in $F_v(A(z))$. Then for each $v \in \mathbb{B}^4$ and $t \in \mathbb{R}$, we define the open set

$$A(v, t, z) = \{x \in \mathbb{S}^3 : d(v, z)(x) < t\},$$

and the closed set

$$\Sigma(v, t, z) = \partial A(v, t, z).$$

Note that $\Sigma(v, t, z)$ has a natural orientation and induces an element in $\mathcal{Z}(\mathbb{S}^3; \mathbb{Z})$, and

$$A(v, t, \sigma(z)) = \mathbb{S}^3 \setminus (\Sigma(z) \cup A(v, -t, z)).$$

In order to define an $\mathbb{R}\mathbb{P}^5$ -family, we need to reparametrize A by performing blow-up suitably, because A does not give a continuous family near $v \in \Sigma(z)$.

5.2.2. *Reparametrizing A via blow-up.* Define the right-half disk on \mathbb{R}^2 as

$$D_+^2(r) := \{s = (s_1, s_2) \in \mathbb{R}^2 : |s| < r, s_1 \geq 0\}.$$

For $\varepsilon > 0$ sufficiently small, we define the map $\Lambda(z) : \Sigma(z) \times D_+^2(3\varepsilon) \rightarrow \overline{\mathbb{B}^4}$ by

$$\Lambda(z)(p, s) = (1 - s_1)(\cos(s_2)p + \sin(s_2)N(z)_p),$$

which is a diffeomorphism onto a neighborhood of $\Sigma(z)$ in $\overline{\mathbb{B}^4}$. Notice $\Lambda(z)$ maps into \mathbb{S}^3 if $s_1 = 0$.

Let $\Omega_r(z)$ be the image $\Lambda(z)(\Sigma(z) \times D_+^2(r))$ for all $r \leq 3\varepsilon$. Then as s approaches $(0, 3\varepsilon)$ (resp. $(0, -3\varepsilon)$), $\Lambda(z)(p, s)$ approaches $A^*(z) \setminus \Omega_{3\varepsilon}(z)$ (resp. $A(z) \setminus \Omega_{3\varepsilon}(z)$).

Now, we define a continuous map $T(z) : \overline{\mathbb{B}^4} \rightarrow \overline{\mathbb{B}^4}$ which collapses the tubular neighborhood $\Omega_\varepsilon(z)$ of $\Sigma(z)$ onto $\Sigma(z)$:

- $T(z)$ is the identity on $\overline{\mathbb{B}^4} \setminus \Omega_{3\varepsilon}(z)$.
- On $\Omega_{3\varepsilon}(z)$,

$$T(z)(\Lambda(z)(p, s)) = \Lambda(z)(p, \psi(|s|)s),$$

where ψ is smooth, 0 on $[0, \varepsilon]$, strictly increasing on $[\varepsilon, 2\varepsilon]$, and 1 on $[2\varepsilon, 3\varepsilon]$.

Note that $T(z) = T(\sigma(z))$.

For every $p \in \Sigma(z)$ and $k \in [-\infty, +\infty]$, define

$$(5.2) \quad \overline{Q}_{p,k,z} = -\frac{k}{\sqrt{1+k^2}}p - \frac{1}{\sqrt{1+k^2}}N(z)_p \in \mathbb{S}^3.$$

(Note we use the convention that $\frac{k}{\sqrt{1+k^2}} = \pm 1$ and $\frac{1}{\sqrt{1+k^2}} = 0$ for $k = \pm\infty$.)

We define the generalised Gauss map $\overline{Q}(z) : \mathbb{S}^3 \cup \overline{\Omega}_\varepsilon(z) \rightarrow \mathbb{S}^3$ by:

$$(5.3) \quad \overline{Q}(z)(v) = \begin{cases} -T(v) & v \in A^*(z) \setminus \overline{\Omega}_\varepsilon(z) \\ T(v) & v \in A(z) \setminus \overline{\Omega}_\varepsilon(z) \\ \overline{Q}_{p,k(s),z} & v = \Lambda(z)(p, s) \in \overline{\Omega}_\varepsilon(z) \end{cases}$$

where $s = (s_1, s_2) \in D_+^2(\varepsilon)$ and

$$(5.4) \quad k(s) = \frac{s_2}{\sqrt{\varepsilon^2 - s_2^2}} \in [-\infty, +\infty].$$

The key property of \bar{Q} is that, as $v \in \mathbb{S}^3$ moves from $A(z) \setminus \Omega_{3\varepsilon}(z)$, crosses $\Sigma(z)$, and arrives in $A^*(z) \setminus \Omega_{3\varepsilon}(z)$ (so that $v \in \mathbb{S}^3$ crosses from one side of $\Sigma(z)$ into another), $\bar{Q}(z)$ flips from being the identity map to the antipodal map, continuously. And the continuous function $k(s)$ is such that $k(s)$ goes to $+\infty$ (resp. $-\infty$) if $\Lambda(z)(p, s)$ goes towards $A^*(z) \setminus \Omega_\varepsilon(z)$ (resp. $A(z) \setminus \Omega_\varepsilon(z)$).

We define $\bar{r}(z) : \mathbb{S}^3 \cup \bar{\Omega}_\varepsilon(z) \rightarrow [0, \pi]$ by:

$$(5.5) \quad \bar{r}(z)(v) = \begin{cases} 0 & v \in A^*(z) \setminus \bar{\Omega}_\varepsilon(z) \\ \pi & v \in A(z) \setminus \bar{\Omega}_\varepsilon(z) \\ \bar{r}_{k(s)} & v = \Lambda(z)(p, s) \in \bar{\Omega}_\varepsilon(z) \end{cases}$$

where $\bar{r}_k = \pi/2 - \arctan k$. The key point is that $\bar{r}(z)$ changes from 0 to π continuously across the intermediate region $\bar{\Omega}_\varepsilon(z)$.

With the above preparation, we can start defining $\Psi_5^{\Sigma(z)}$.

We know the map $\bar{Q}(z)$ satisfies $\bar{Q}(\sigma(z)) = -\bar{Q}(z)$ from $N(z) = -N(\sigma(z))$, and the function $\bar{r}(z)$ satisfies $r(\sigma(z)) = \pi - \bar{r}(z)$. Notice that the subset $\bar{\Omega}_\varepsilon(z)$ is invariant under the antipodal map, and $\bar{\Omega}_\varepsilon(\sigma(z)) = \bar{\Omega}_\varepsilon(z)$. The function \bar{r} can initially be considered as a continuous function defined on

$$\{(z, v) : z \in \tilde{Z}, v \in \mathbb{S}^3 \cup \bar{\Omega}_\varepsilon(z)\},$$

which is a closed subset of $\tilde{Z} \times \mathbb{B}^4$. We extend it continuously to a function

$$\bar{r} : \tilde{Z} \times \bar{\mathbb{B}}^4 \rightarrow [0, \pi]$$

that still satisfies $\bar{r}(\sigma(z)) = \pi - \bar{r}(z)$. For each $(v, t, z) \in \bar{\mathbb{B}}^4 \times \mathbb{R} \times \tilde{Z}$, we define the set

$$U(v, t, z) = \begin{cases} A(T(z)(v), t, z) & \text{if } v \in \mathbb{B}^4 \setminus \bar{\Omega}_\varepsilon(z), \\ B_{\bar{r}(z)(v)+t}(\bar{Q}(z)(v)) & \text{if } v \in \mathbb{S}^3 \cup \bar{\Omega}_\varepsilon(z). \end{cases}$$

Notice that $U(v, \pi, z) = \mathbb{S}^3$ and $U(v, -\pi, z)$ is empty for every $v \in \mathbb{B}^4$. And for $v \in \mathbb{S}^3 \cup \bar{\Omega}_\varepsilon(z)$, U gives geodesic spheres of varying radius, and the radius always covers the full range $[0, \pi]$ as t ranges over $[-\pi, \pi]$.

Marques-Neves showed that the family ∂U is continuous in the flat topology [MN14, Theorem 5.1].

5.2.3. Closing up the parameter space. Let $\gamma : [0, 1] \rightarrow [0, 1]$ be the continuous function satisfying $\gamma(t) = 0$ if $t \leq \frac{1}{2}$ and $\gamma(t) = 2t - 1$ if $t \geq \frac{1}{2}$. For each

$$(v, t, z) \in \bar{\mathbb{B}}^4 \times [-1, 1] \times \tilde{Z}$$

we define the open set

$$U'(v, t, z) = U \left(v, 2\pi t + \gamma(|v|) \left(\frac{\pi}{2} - \bar{r}(z)(v) \right), z \right).$$

Reparametrizing t as above allows us to close up the boundary of $\overline{\mathbb{B}^4} \times [-1, 1] \times \{z\}$ via the antipodal map, as follows.

For

$$(v, t) \in \partial(\overline{\mathbb{B}^4} \times [-1, 1]) = (\mathbb{S}^3 \times (-1, 1)) \cup (\overline{\mathbb{B}^4} \times \{-1, 1\}),$$

it is straightforward to check that

$$U'(v, t, z) = \begin{cases} B_{\frac{\pi}{2} + 2\pi t}(\overline{Q}(z)(v)) & \text{if } (v, t) \in \mathbb{S}^3 \times (-1, 1), \\ 0 & \text{if } (v, t) \in \overline{\mathbb{B}^4} \times \{-1\}, \\ \mathbb{S}^3 & \text{if } (v, t) \in \overline{\mathbb{B}^4} \times \{1\}. \end{cases}$$

Note that the smallness of ε is used here for the case where $v \in \overline{\Omega}_\varepsilon(z)$. Now, using the fact that $B_{\pi-r}(-p) = \mathbb{S}^3 \setminus \overline{B_r(p)}$, $T(z)(-v) = -T(z)(v)$, and $\overline{Q}(z)(-v) = -\overline{Q}(z)(v)$ by [Nur16, §3.4.3], it follows immediately that for every $(v, t) \in \partial(\overline{\mathbb{B}^4} \times [-\pi, \pi])$,

$$(5.6) \quad U'(-v, -t, z) = \mathbb{S}^3 \setminus \overline{U'(v, t, z)},$$

This allows us to define an \mathbb{RP}^5 -family $\Psi_5^{\Sigma(z)}$ of closed subsets by

$$\Psi_5^{\Sigma(z)}([(v, t)]) = \partial U'(v, t, z).$$

Note that we have disregarded orientation, and a geodesic sphere of radius 0 or π should be both treated as a single point, ensuring $\Psi_5^{\Sigma(z)}$ satisfies Definition 2.3 (3).

5.2.4. *Symmetry of $\Psi_5^{\Sigma(z)}$ in z .* Furthermore, it follows from a straightforward computation that the map U' satisfies the identity

$$U'(v, t, z) = \mathbb{S}^3 \setminus \overline{U'(v, -t, \sigma(z))},$$

for every $(v, t, z) \in \mathbb{B}^4 \times [-1, 1] \times \tilde{Z}$. Thus, we have

$$\Psi_5^{\Sigma(z)}([(v, t)]) = \Psi_5^{\Sigma(\sigma(z))}([(v, -t)]).$$

This finishes the proof of Proposition 5.1.

5.3. Proof of Theorem 3.1. From the above construction of Ψ , it is clear that for each oriented Clifford torus Σ , there exists a homeomorphism $f : Y_{[\Sigma]} \rightarrow \mathbb{RP}^5$ such that $\Phi_5^\Sigma \circ f = [\Psi|_{Y_{[\Sigma]}}]$. So to prove Theorem 3.1, it suffices to show that Ψ is a Simon-Smith family of genus ≤ 1 . We start with studying the topology of the surfaces in Ψ .

5.3.1. *Topology of $\Sigma(v, t, z)$.* Fix a $z_0 \in \tilde{Z}$ which corresponds to the Clifford torus Σ_0 given by

$$X(\alpha, \beta) = \frac{1}{\sqrt{2}}(\cos \alpha, \sin \alpha, \cos \beta, \sin \beta),$$

with $\alpha, \beta \in S^1$ (where S^1 is viewed as $[0, 2\pi]$ with endpoints identified), oriented such that the distance function $d(0, z_0)$ is negative on the “core circle”

$$w(\beta) = \frac{1}{\sqrt{2}}(0, 0, \cos \beta, \sin \beta), \quad \beta \in S^1.$$

Proposition 5.2. *Let $(v, t) \in \mathbb{B}^4 \times [-\pi, \pi]$. We have the following descriptions regarding the closed set*

$$\Sigma(v, t) := \partial\{x \in \mathbb{S}^3 : d(v, z_0)(x) < t\}.$$

If $v = 0$, then $\Sigma(v, t)$ is a smooth torus for $t \in (-\pi/4, \pi/4)$, a great circle for $t = \pm\pi/4$, and empty for other t .

If $v \neq 0$ and $v/|v|$ lies on the circle

$$(5.7) \quad \frac{1}{\sqrt{2}}(\cos \alpha, \sin \alpha, 0, 0), \quad \alpha \in S^1,$$

then there exist $-\pi < t_1 < 0 < t_2 < \pi$, depending on v , such that:

- (1) *For $t_1 < t < t_2$, $\Sigma(v, t)$ is a smooth torus.*
- (2) *For $t = t_1$ and $t = t_2$, $\Sigma(v, t)$ is a great circle.*
- (3) *For $t < t_1$ and $t > t_2$, $\Sigma(v, t)$ is empty.*

If v does not fall into the above two cases, then there exist

$$-\pi < t_1 < t_2 < 0 < t_3 < t_4 < \pi,$$

depending on v , such that:

- (1) *For $t_2 < t < t_3$, $\Sigma(v, t)$ is a smooth torus.*
- (2) *For $t = t_2$ and $t = t_3$, $\Sigma(v, t)$ is topologically a sphere with two points identified such that it is smooth except at a point.*
- (3) *For $t_1 < t < t_2$ and $t_3 < t < t_4$, $\Sigma(v, t)$ is topologically a sphere, smooth except at two points.*
- (4) *For $t = t_1$ and $t = t_4$, $\Sigma(v, t)$ is a point.*
- (5) *For $t < t_1$ and $t > t_4$, $\Sigma(v, t)$ is empty.*

Proof. First, the case $v = 0$ is clear.

Second, in the case where $v \neq 0$ and $v/|v|$ lies on the circle (5.7), note that both the Clifford torus Σ_0 and the map F_v , are symmetric about the antipodal points $\pm v/|v|$. Therefore, the image $F_v(\Sigma_0)$, as well as the level surfaces is parallel to $F_v(\Sigma_0)$, share this symmetry, and the desired result follows immediately.

Finally, consider the case where $v \neq 0$ with $v/|v|$ not lying on the circle (5.7). The Clifford torus Σ_0 is the envelope of the 2-spheres $S_\beta \subset \mathbb{S}^3$ with center $w(\beta)$ and radius $\pi/4$ under the standard metric on \mathbb{S}^3 , where $\beta \in S^1$.

Thus, Σ_0 is a *channel surface*. The sphere S_β is given by the following equations on $x \in \mathbb{R}^4$:

$$|x| = 1, \quad x \cdot w(\beta) = \cos(\pi/4).$$

Now, when we apply the conformal diffeomorphism F_v to S_β , we obtain another 2-sphere $F_v(S_\beta)$ in \mathbb{S}^3 , with a new center $\tilde{w}(\beta)$ and a new radius $\tilde{r}(\beta) \in (0, \pi)$ under the standard metric on \mathbb{S}^3 .

We see that the function $\tilde{r} : S^1 \rightarrow \mathbb{R}$ has the following property.

Claim 5.3. *If $v \in \mathbb{B}^4$ is non-zero and $v/|v|$ does not lie on the circle (5.7), then the function $\tilde{r} : S^1 \rightarrow \mathbb{R}$ has exactly one local maximum and one local minimum.*

We will postpone the proof to the end, but first, let us note that this lemma is sufficient to justify the final case of our proposition.

Without loss of generality, we can focus on the case $t < 0$, as the case $t > 0$ follows from an identical analysis on the Clifford torus $-\Sigma_0$, i.e. the Clifford torus with the opposite orientation to Σ_0 .

Let us consider the closed envelop given by the collection of spheres

$$\{\partial B_{\tilde{r}(\beta)+t}(\tilde{w}(\beta))\}_{\beta \in S^1},$$

where we use the convention that $\partial B_0(p) = \{p\}$ and $\partial B_r = \emptyset$ for $r < 0$. Given that conformal diffeomorphisms preserve channel surfaces, it is straightforward to verify that this envelop is exactly the set $\Sigma(v, t)$.

Now, suppose the function $\tilde{r} : S^1 \rightarrow \mathbb{R}$ has exactly one local maximum and one local minimum. Let us decrease t starting from 0. When $|t|$ is small, the envelop described above forms a torus. And there would be a $t_2 < 0$ such that, when $t = t_2$, the torus neck pinches at exactly one point, corresponding to the β where \tilde{r} achieves a minimum. As we continue decreasing t , the envelop becomes a topological sphere with two singular points, until it eventually vanishes at a point, corresponding to the β where \tilde{r} achieves a maximum. It happens at some time $t_1 < 0$. This finishes the proof of the third case of Proposition 5.2.

Proof of Claim 5.3. Recalling that F_v has a symmetry about the points $\pm v/|v|$, and F_v pushes everything away from $\frac{v}{|v|}$ and towards $-\frac{v}{|v|}$, we know the center $\tilde{w}(\beta)$ of the sphere $F_v(S_\beta)$ takes the form

$$\tilde{w}(\beta) = \frac{-\cos(\theta(\beta))\frac{v}{|v|} + \sin(\theta(\beta))w(\beta)}{\left|-\cos(\theta(\beta))\frac{v}{|v|} + \sin(\theta(\beta))w(\beta)\right|}$$

for some function $\theta(\beta) \in (0, \pi/2)$. For convenience, let us denote the above denominator by $l(\beta)$. Additionally, the new sphere $F_v(S_\beta)$ is characterized by the following equations for $y \in \mathbb{R}^4$:

$$|y| = 1 \quad \text{and} \quad y \cdot \tilde{w}(\beta) = \text{constant},$$

where the constant equals to \cos of the radius of the new sphere.

Thus, let us compute $F_v(x) \cdot \tilde{w}(\beta)$ for $x \in S_\beta$. For simplicity, we omit writing the dependence of l, θ and w on β .

$$\begin{aligned}
 & F_v(x) \cdot \tilde{w}(\beta) \\
 &= \left(\frac{1 - |v|^2}{|x - v|^2} (x - v) - v \right) \cdot \frac{-\cos \theta \frac{v}{|v|} + \sin \theta w}{l} \\
 (5.8) \quad &= \frac{1}{l} \frac{1 - |v|^2}{1 + |v|^2 - 2x \cdot v} \left(-\cos \theta \frac{x \cdot v}{|v|} + \cos \theta |v| + \sin \theta \frac{1}{\sqrt{2}} - \sin \theta w \cdot v \right) \\
 &\quad + \frac{1}{l} (\cos \theta |v| - \sin \theta w \cdot v).
 \end{aligned}$$

Note that in the last equality, we use $w \cdot x = 1/\sqrt{2}$, which holds because $x \in S_\beta$. Hence, for some constant $C = C(v, \beta)$, independent of $x \in S_\beta$, we have

$$-\cos \theta \frac{x \cdot v}{|v|} + \cos \theta |v| + \sin \theta \frac{1}{\sqrt{2}} - \sin \theta w \cdot v = C \left(1 + |v|^2 - 2x \cdot v \right).$$

Now, by setting both the zeroth-order term and the first-order term in x to zero, and canceling C from these two equations, we obtain

$$(5.9) \quad \frac{1 - |v|^2}{2|v|} \cos \theta = \left(\frac{1}{\sqrt{2}} - w \cdot v \right) \sin \theta,$$

which leads to

$$(5.10) \quad \cot \theta = \frac{2|v|}{1 - |v|^2} \left(\frac{1}{\sqrt{2}} - w \cdot v \right).$$

Hence, substituting (5.9) into (5.8), the radius $\tilde{r}(\beta) \in (0, \pi)$ satisfies:

$$\begin{aligned}
 \cos \tilde{r}(\beta) &= F_v(x) \cdot \tilde{w}(\beta) \\
 &= \frac{1}{l} \frac{1 - |v|^2}{|x - v|^2} (x - v) \cos \theta \left(-\frac{x \cdot v}{|v|} + |v| + \frac{1 - |v|^2}{2|v|} \right) \\
 &\quad + \frac{1}{l} (\cos \theta |v| - \sin \theta w \cdot v) \\
 &= \frac{1}{l} \left(\frac{1 - |v|^2}{2|v|} \cos \theta + (\cos \theta |v| - \sin \theta w \cdot v) \right) \\
 &= \frac{1}{l} \left(\frac{1}{|v|} \cos \theta - \frac{1}{\sqrt{2}} \sin \theta \right).
 \end{aligned}$$

Now, by a straightforward computation of l using (5.9), we find that

$$l = \sqrt{\frac{1}{2} \sin^2 \theta + \left(\frac{1}{|v|} \cos \theta - \frac{1}{\sqrt{2}} \sin \theta \right)^2}.$$

It follows straightforward that

$$\cos^2 \tilde{r} = \left(1 + \frac{1}{\left(\frac{\sqrt{2}}{|v|} \cot \theta - 1\right)^2} \right).$$

Since \tilde{r} depends smoothly on θ , the equation can be extended to the case where the denominator is zero. Together with (5.10) one can derive that

$$(5.11) \quad \cot \tilde{r}(\beta) = \frac{\sqrt{2}}{|v|} \cot \theta - 1 = \frac{2(1 - \sqrt{2}w \cdot v)}{1 - |v|^2} - 1.$$

Note that the sign of the right-hand side is positive sign due to the monotonicity of \cot and the geometric behavior of the conformal map F_v : when $w \cdot v$ is larger, \tilde{r} is larger.

Since \cot is a strictly decreasing function with non-zero slope on $(0, \pi)$, by (5.11) and the fact that \tilde{r} depends smoothly on $w \cdot v$, it suffices to show that $w \cdot v(\beta)$, which is a function of $\beta \in S^1$, has exactly one local minimum and one local maximum. This holds true because $w(\beta)$ describe a circles in \mathbb{S}^3 that is not perpendicular to v . This completes the proof of the claim. \square

This finishes the proof of Proposition 5.2. \square

5.3.2. Ψ is a Simon-Smith family of genus ≤ 1 . Let us finish the proof of Theorem 3.1, by showing that Ψ is a Simon-Smith family of genus ≤ 1 .

By the definition of Ψ in the previous section and Proposition 5.2, we can conclude that Ψ consists of the following types of closed subsets:

- (1) A smooth torus: This arises as the nearby level surface of the signed distance function to $F_v(\Sigma(z))$, where $v \notin \Sigma(z)$.
- (2) A smooth sphere: They are geodesic spheres on \mathbb{S}^3 with radius strictly between 0 and π .
- (3) A great circle: This occurs in the second case of v in Proposition 5.2.
- (4) A topological sphere with two points identified, so that it is smooth except at a point: This occurs in the third case of v in Proposition 5.2.
- (5) A topological sphere, smooth except at two points: This occurs in the third case of v in Proposition 5.2.
- (6) A single point: It has several forms. (1) A geodesic sphere on \mathbb{S}^3 with radius 0. (2) A geodesic sphere on \mathbb{S}^3 with radius π . (3) It also occurs in the third case of v in Proposition 5.2.
- (7) The empty set.

Then, it follows easily that each member of Ψ is an element of $\mathcal{S}(S^3)$, and Ψ satisfies Definition 2.3 (1), (2), (3), and also the genus bound (5).

As for Definition 2.3 (4), regarding the local C^∞ -convergence, we know from the definition of Ψ that it suffices to fix a $z = z_0$ and show that $\Psi_5^{\Sigma(z_0)}$

satisfies Definition 2.3 (4). In fact, if we define the family

$$\Psi'_5(v, t) := \partial U(v, t, z_0), \quad (v, t) \in \overline{\mathbb{B}^4} \times [-2\pi, 2\pi]$$

(with $\Psi'_5(v, t)$ treated as a single point if it is a geodesic sphere of radius 0 or π), and notice that $\Psi_5^{\Sigma(z_0)}$ was defined as a subfamily of Ψ'_5 with suitable closing-up of the boundary of the parameter space, it becomes clear that it suffices to show that Ψ'_5 satisfies Definition 2.3 (4).

Let us first show that $\Psi'_5(\cdot, 0)$ satisfies Definition 2.3 (4). We now explicitly describe, for each $\Psi'_5(v, 0)$, what the associated set $P(v, 0)$ of points (according to Definition 2.3) should be. By definition, $\Psi'_5(v, 0)$ takes on one of the following forms, for $v \in \overline{B^4}$.

- (1) If $v \in \overline{\mathbb{B}^4 \setminus \Omega_\varepsilon(z_0)}$, $\Psi'_5(v, 0)$ is a smooth torus.
- (2) If $v \in \overline{\mathbb{S}^3 \setminus \Omega_\varepsilon(z_0)}$, $\Psi'_5(v, 0)$ is empty.
- (3) If $v \in \overline{\mathbb{S}^3 \cap \Omega_\varepsilon(z_0)}$, $\Psi'_5(v, 0)$ is a single point.
- (4) If $v \in \overline{\Omega_\varepsilon(z_0) \setminus \mathbb{S}^3}$, $\Psi'_5(v, 0)$ is a smooth geodesic sphere.

For the first two case, we set $P(v, 0)$ to be empty. For the third case, we take $P(v, 0) = \Psi'_5(v, 0)$. For the last case, we need to be a bit more careful. If v lies in the interior of $\Omega_\varepsilon(z_0)$, we can set $P(v, 0)$ to be empty. However, if $v \in \overline{\mathbb{B}^4 \cap \partial \Omega_\varepsilon(z_0)}$, we should let $P(v, 0)$ consist of the point $-v/|v|$. Indeed, from [MN14, Theorem 5.1], we already know $\Psi'_5(\cdot, 0)$ is continuous in the flat topology near each v . However, the convergence is not smooth when approaching such v from outside $\overline{\Omega_\varepsilon(z_0)}$, as we would see a handle of shrinking size near the point $-v/|v|$. With such definition of $P(v, 0)$ for each v , it is straightforward to verify from the definition that $\Psi'_5(\cdot, 0)$ satisfies Definition 2.3 (4).

Now, consider the case $t \neq 0$. By the definition of U , each $\Psi'_5(v, t)$ is a level surface of the signed distance function to $\Psi'_5(v, 0)$. Thus, using Proposition 5.2 and the fact that $\Psi'_5(\cdot, 0)$ satisfies Definition 2.3 (4), it is easy to check that Ψ'_5 also satisfies Definition 2.3 (4), with $P(v, t)$ be the union of projection of $P(v, 0)$ to $\Psi'_5(v, t)$ and finite singular points as described in Proposition 5.2. This finishes the proof that Ψ'_5 satisfies Definition 2.3 (4).

Finally, to show that the closing-up $\Psi_5^{\Sigma(z_0)}$ satisfies Definition 2.3 (4), one simply need to merge the associated set $P(v, t)$ and $P(v', t')$ if $(v, t) \sim (v', t')$. Since Ψ is induced from $\Psi_5^{\Sigma(z_0)}$ via $SO(4)$ -action and another closing-up, it follows that Ψ also satisfies Definition 2.3 (4).

As a result, Ψ is a Simon-Smith family of genus ≤ 1 .

5.4. A 7-sweepout in Ψ . As part of the setup for the proof of Theorem 3.13, we consider a subset $C \subset Y = \text{dmn}(\Psi)$, where $C \cong \mathbb{RP}^4 \times \mathbb{RP}^2 \times \mathbb{RP}^2$, such that $[\Psi|_C]$ is a 7-sweepout.

First, we define the submanifold

$$\tilde{C} := \{([v, t]), z) \in \mathbb{RP}^5 \times \tilde{Z} : t = 0\}.$$

Evidently, $\tilde{C} \cong \mathbb{RP}^4 \times \tilde{Z}$. Note that \tilde{C} is invariant under the \mathbb{Z}_2 -action on $\mathbb{RP}^5 \times \tilde{Z}$ described in (5.1). Let C be the quotient of \tilde{C} by this \mathbb{Z} -action. Then $C \cong \mathbb{RP}^4 \times Z$, and C can be viewed as a subset of Y . Geometrically, $\Xi|_C$ consists of images of Clifford tori under conformal maps, and geodesic spheres. The fact that C is a product, instead of a non-trivial \mathbb{RP}^4 -bundle over Z , reflects the irrelevance of Clifford tori's orientations in defining $\Xi|_C$, as opposed to the situation over $Y \setminus C$.

Now,

$$H_1(C; \mathbb{Z}_2) \cong H_1(\mathbb{RP}^4; \mathbb{Z}_2) \oplus H_1(\mathbb{RP}^2; \mathbb{Z}_2) \oplus H_1(\mathbb{RP}^2; \mathbb{Z}_2) \cong \mathbb{Z}_2 \oplus \mathbb{Z}_2 \oplus \mathbb{Z}_2.$$

Let $\omega_C, \alpha_C, \beta_C \in H^1(C; \mathbb{Z}_2)$ be respectively the Hom-duals of

$$(5.12) \quad (1, 0, 0), (0, 1, 0), (0, 0, 1) \in H_1(C; \mathbb{Z}_2).$$

By the Künneth formula and the fact that the cohomology ring $H^*(\mathbb{RP}^k; \mathbb{Z}_2)$ is $\mathbb{Z}_2[x]/(x^{k+1})$, we have

$$H^*(C; \mathbb{Z}_2) \cong \mathbb{Z}_2[\omega_C, \alpha_C, \beta_C]/(\omega_C^5, \alpha_C^3, \beta_C^3).$$

As verified in [Nur16, §3.4.3], the three homology classes in (5.12) all correspond to 1-sweepouts under $[\Psi|_C]$, and thus the element

$$\lambda_C := ([\Psi|_C])^*(\bar{\lambda}) \in H^1(C; \mathbb{Z}_2)$$

is equal to $\omega_C + \alpha_C + \beta_C$. Hence, by the Künneth formula,

$$(\lambda_C)^7 = \omega_C^4 \cup \alpha_C^2 \cup \beta_C + \omega_C^4 \cup \alpha_C \cup \beta_C^2 \neq 0$$

in \mathbb{Z}_2 -coefficients. So $[\Psi|_C]$ is a 7-sweepout, as $[\Psi]$ is continuous in the \mathcal{F} -norm by Proposition 2.5.

5.5. Proof of Theorem 3.13. Recall the three elements λ, α, β in $H^1(Y; \mathbb{Z}_2)$ defined by:

- Let $\bar{\lambda}$ be the non-trivial element of $H^1(\mathcal{Z}_2(S^3; \mathbb{Z}_2); \mathbb{Z}_2)$; define $\lambda = [\Psi]^*(\bar{\lambda})$.
- Let $a = \mathbb{RP}^1 \times \mathbb{RP}^2 \subset \mathbb{RP}^2 \times \mathbb{RP}^2$, and $A \subset Y$ be the \mathbb{RP}^5 -subbundle over a ; define α to be the Poincaré dual $PD(A)$ of A .
- Let $b = \mathbb{RP}^2 \times \mathbb{RP}^1 \subset \mathbb{RP}^2 \times \mathbb{RP}^2$, and $B \subset Y$ be the \mathbb{RP}^5 -subbundle over b ; define β to be the Poincaré dual $PD(B)$ of B .

Our goal is to prove $\lambda^5 \cup \alpha^2 \cup \beta^2 \neq 0$.

Let us first write down the generators of $H^*(Y; \mathbb{Z}_2)$. Let $C \subset Y$ be the 8-dimensional subset obtained in the previous subsection such that $[\Psi|_C]$ is a 7-sweepout, and $i_C : C \rightarrow Y$ be the inclusion map. Recall that the element

$$\lambda_C := i_C^*(\lambda) = ([\Psi] \circ i_C)^*(\bar{\lambda}) \in H^1(C; \mathbb{Z}_2)$$

satisfies

$$(5.13) \quad (\lambda_C)^7 \neq 0 \text{ in } H^*(C; \mathbb{Z}_2).$$

Now, set $\gamma := PD(C) \in H^1(Y; \mathbb{Z}_2)$.

Recall that Y is an \mathbb{RP}^5 -bundle over $Z = \mathbb{RP}^2 \times \mathbb{RP}^2$:

$$\mathbb{RP}^5 \rightarrow Y = \text{dmm}(\Psi) \rightarrow Z.$$

Let us denote the projection map by $\pi : Y \rightarrow Z$.

Claim 5.4. *The cohomology ring $H^*(Y; \mathbb{Z}_2)$ is generated by α, β, γ .*

Proof. First, by definition, α, β are the pullbacks of the two generators of $H^*(Z; \mathbb{Z}_2)$ under the projection map $\pi : Y \rightarrow Z$. Moreover, note that for each $z \in Z$, the intersection of C with every fiber $Y_z \cong \mathbb{RP}^5$ of Y is an \mathbb{RP}^4 , whose Poincaré dual in $H^1(Y_z; \mathbb{Z}_2)$ generates

$$H^*(Y_z; \mathbb{Z}_2) \cong H^*(\mathbb{RP}^5; \mathbb{Z}_2).$$

Thus, if we pullback γ to $H^1(Y_z; \mathbb{Z}_2)$ via the inclusion map $Y_z \hookrightarrow Y$, we obtain the generator of $H^*(Y_z; \mathbb{Z}_2)$. Then the claim follows immediately from the Leray–Hirsch Theorem (see [Hat02, Theorem 4D.1]). \square

Let $\pi_1, \pi_2 : Z \rightarrow \mathbb{RP}^2$ be the projection onto the first and second factor respectively. Let $k_1, k_2 \subset \mathbb{RP}^2$ be two transverse isotopic loops in \mathbb{RP}^2 with a single intersection point p . Let $Y_{(p,p)} \subset Y$ be the fiber over (p, p) .

Claim 5.5. $\alpha^2 \cup \beta^2 = PD(Y_{(p,p)})$, and $\alpha^r \cup \beta^s = 0$ whenever $r \geq 3$ or $s \geq 3$.

Proof. By definition, α is the Poincaré dual of $A_i := (\pi_1 \circ \pi)^{-1}(k_i)$ while β is the Poincaré dual of $B_i := (\pi_2 \circ \pi)^{-1}(k_i)$, for $i = 1, 2$. Therefore, $\alpha^2 \cup \beta^2$ is the Poincaré dual of $A_1 \cap A_2 \cap B_1 \cap B_2$, which is the fiber $Y_{(p,p)}$.

If $r \geq 3$ or $s \geq 3$, we just consider three perturbed copies of $k_1 \subset \mathbb{RP}^2$ that do not share a common point, and their preimages under $\pi_1 \circ \pi$ or $\pi_2 \circ \pi$, similarly as above. Then second claim would follow. \square

Claim 5.6. *In $H^1(Y; \mathbb{Z}_2)$, λ is not a linear combination of just α and β .*

Proof. Suppose by contradiction that $\lambda = c_1\alpha + c_2\beta$ for some $c_1, c_2 \in \mathbb{Z}_2$. Then

$$\lambda_C = i_C^*(\lambda) = i_C^*(c_1\alpha + c_2\beta).$$

Recall that $(\lambda_C)^7 \neq 0$ by (5.13). But by Claim 5.5, $(c_1\alpha + c_2\beta)^7 = 0$, which leads to a contradiction. \square

From Claim 5.4 and 5.6, we have $\lambda = \gamma + c_1\alpha + c_2\beta$ for some $c_1, c_2 \in \mathbb{Z}_2$. Theorem 3.13 will follow from the following claim.

Claim 5.7. $\gamma^5 \cup \alpha^2 \cup \beta^2 \neq 0$.

Proof. Since $C \rightarrow Z$ is a trivial \mathbb{RP}^4 -bundle, we can perturb it to obtain C_1, C_2, C_3, C_4, C_5 such that

- Within a trivialization of Y over some open neighborhood $U \subset Z$ of (p, p) , for each i , the restriction of C_i over U is equal to $\hat{C}_i \times U$ for some $\hat{C}_i \subset \mathbb{RP}^5$ homeomorphic to an \mathbb{RP}^4 ;
- All \hat{C}_i 's transversely intersect at a single point.

Thus, $Y_{(p,p)} \cap \bigcap_{i=1}^5 C_i$ is just a single point in Y . Together with Claim 5.5 and the definition $\gamma = PD(C)$, we know $\gamma^5 \cup \alpha^2 \cup \beta^2 \neq 0$. \square

Thus, by the above claim and Claim 5.5, we have

$$\lambda^5 \cup \alpha^2 \cup \beta^2 = (\gamma + (c_1\alpha + c_2\beta))^5 \cup \alpha^2 \cup \beta^2 = \gamma^5 \cup \alpha^2 \cup \beta^2 \neq 0$$

in $H^*(Y; \mathbb{Z}_2)$. This completes the proof of Theorem 3.13.

6. PERTURBING THE METRIC

Note that in a 3-dimensional Riemannian manifold (S^3, g) with positive Ricci curvature, there is no stable minimal surface, and moreover, by [CS85], there exists $L > 0$, such that the area of every minimal surface with genus at most 1 is no greater than $L/2$. Therefore, Proposition 3.3 follows from the following general result.

Proposition 6.1 (Generalization of Proposition 3.3). *Let (S^3, g_0) be a Riemannian 3-sphere with only finitely many embedded minimal tori and no degenerate stable embedded minimal sphere. For any $\varepsilon > 0$ and any L greater than the largest area among all minimal tori in (S^3, g_0) , there exists a metric g' that is ε -close to g_0 in C^∞ such that among all embedded g' -minimal surfaces with area less than L :*

- (1) *the number of embedded g' -minimal tori is the same as that of g_0 -minimal tori;*
- (2) *there are only finitely many embedded minimal spheres $\{S_1, \dots, S_q\}$, and each of them is non-degenerate.*
- (3) *the g' -areas*

$$\text{area}_{g'}(S_1), \dots, \text{area}_{g'}(S_q)$$

are \mathbb{Z} -linearly independent. Moreover, their \mathbb{Z} -linear combination can never achieve the area of any g' -minimal tori with g' -area less than L .

Let us recall a compactness result for minimal surfaces essentially following from [CS85, Page 390, Claim(*)] (cf. [And85, Whi87], [CM11, Proposition 7.14]).

Lemma 6.2. *In S^3 , given a non-negative integer \mathfrak{g}_0 , a sequence of Riemannian metrics $\{g_i\}_{i=1}^\infty$ and a sequence of connected embedded closed surfaces $\{\Sigma_i\}_{i=1}^\infty$, suppose that there exists another Riemannian metrics g_∞ such that $g_i \rightarrow g_\infty$ in C^3 and each Σ_i is a g_i -minimal surface satisfying*

$$\sup_i \text{area}_{g_i}(\Sigma_i) < +\infty, \quad \sup_i \mathfrak{g}(\Sigma_i) \leq \mathfrak{g}_0.$$

Then there exists a subsequence of $\{\Sigma_i\}$, still denoted by $\{\Sigma_i\}$, a connected g_∞ -minimal surface Σ_∞ and a finite number of points $Z = \{x_1, \dots, x_l\}$ such that in the varifold topology,

$$|\Sigma_i| \rightarrow k|\Sigma_\infty|, \quad k\mathfrak{g}(\Sigma_\infty) \leq \mathfrak{g}_0,$$

with some $k \in \mathbb{N}^+$, and for any $r > 0$, $\Sigma_i \rightarrow \Sigma_\infty$ in $S^3 \setminus B_r(Z)$ (multi-)graphically in the C^3 topology.

Furthermore, if $k = 1$, then $Z = \emptyset$, and thus, for sufficiently large i , $\mathfrak{g}(\Sigma_i) = \mathfrak{g}(\Sigma_\infty)$; if $k > 1$, then Σ_∞ is degenerate stable.

Remark 6.3. Note that S^3 does not admit non-orientable closed surfaces, so all Σ_i and Σ_∞ are oriented.

Proof. The first part, except for the genus bound of Σ_∞ , follows from [Whi87, Theorem 3], since a g_i -minimal surface can be viewed as an embedded Φ_i -stationary surface in (S^3, g_∞) for some even elliptic integrand Φ_i , and Φ_i converges to the constant function 1. By the area monotonicity formula for g_i -minimal surfaces, Σ_i also converges to Σ_∞ in the Hausdorff distance sense.

Then there exists $r_0 > 0$ such that $\mathfrak{g}(\Sigma_\infty \setminus B_{r_0}(Z)) = \mathfrak{g}(\Sigma_\infty)$. Since $\Sigma_i \rightarrow \Sigma_\infty$ in $S^3 \setminus B_{r_0}(Z)$ (multi-)graphically in the C^3 topology, for sufficiently large i , $\Sigma_i \setminus B_{r_0}(Z)$ has exactly k connected components $\{\Gamma_i^j\}_{j=1}^k$, and

$$\sum_{j=1}^k \mathfrak{g}(\Gamma_i^j) \leq \mathfrak{g}(\Sigma_i) \leq \mathfrak{g}_0.$$

Hence, for sufficiently large i we have

$$\mathfrak{g}(\Sigma_\infty) = \mathfrak{g}(\Sigma_\infty \setminus B_{r_0}(Z)) = \min_j \mathfrak{g}(\Gamma_i^j) \leq \frac{\mathfrak{g}_0}{k}.$$

When $k = 1$, $Z = \emptyset$ follows immediately from Allard's regularity [All72]. When $k > 1$, it follows from [Sha17, Claim 5 and Claim 6] that there exists a positive Jacobi fields on Σ_∞ , and thus, Σ_∞ is degenerate stable. \square

We prove the following useful lemma.

Lemma 6.4. *Suppose (S^3, g_0) contains only finitely many embedded minimal tori, given by $\mathcal{T} := \{T_1, T_2, \dots, T_l\}$. Then for any $L > 0$, there exists an open subset $U \supset \bigcup_{1 \leq j \leq l} T_j$ in S^3 and an open neighborhood \mathcal{O} of g_0 in C^∞ such that for every $g \in \mathcal{O}$,*

- (a) *Every embedded g -minimal 2-sphere S with g -area less than L in S^3 is not entirely contained in U ;*
- (b) *If we further assume that (S^3, g_0) does not admit any degenerate stable embedded minimal sphere with area less than L , and $g|_U = g_0|_U$. Then every embedded g -minimal torus T with g -area less than L in (S^3, g) is inside \mathcal{T} .*

Proof. We first prove that (a) holds for some U and \mathcal{O} . Suppose for contradiction that there exists an $L > 0$ and a sequence of metrics $\{g_j\}$, which is C^3 -converging to g_0 , as well as a sequence of g_j -minimal spheres $\{S_j\}$ in S^3 , each of which has g_j -area less than L and is contained in $1/j$ -tubular neighborhood of $\bigcup_{1 \leq j \leq l} T_j$.

By Lemma 6.2, up to a subsequence, S_j converges to some minimal sphere S_∞ with multiplicity in $\bigcup_{1 \leq j \leq l} T_j$ as varifolds. However, there is no minimal sphere lying in $\bigcup_{1 \leq j \leq l} T_j$, a contradiction.

Hence, we finish the proof of (a).

Now that we fix L, U as above and argue that for a smaller neighborhood $\mathcal{O}' \subset \mathcal{O}$ of g_0 , (b) is true. To see that, suppose again for contradiction that there exists a sequence of metrics $\{g_j\}_J$ C^3 -converging to g_0 with $g_j|_U = g_0|_U$, as well as a sequence of g_j -minimal torus $\{T_j \notin \mathcal{T}\}$ in S^3 with g_j -area less than L . By Lemma 6.2, $|T_j|$ varifold subconverges to $k|\Sigma|$ for some smooth minimal surface $\Sigma \subset (S^3, g_0)$.

Moreover, if Σ is also a torus, then $\Sigma \in \mathcal{T}$ and $T_j \subset U$ for $j \gg 1$, but since $g_j|_U = g_0|_U$, such T_j is also $g_0|_U$ minimal, which contradicts to the definition of \mathcal{T} . Therefore, Σ is a sphere and the multiplicity $k \geq 2$. Then again by Lemma 6.2, Σ is a degenerate stable minimal sphere in (S^3, g_0) , which contradicts to the assumption in (b).

This completes the proof of (b). \square

Now we adapt White's bumpy metric argument to our setting. In the following, let $3 \leq q \leq \infty$ be an integer (or infinity), $L > \sup_j \{\text{area}_{g_0}(T_j)\}$ be fixed, U, \mathcal{O} be determined from previous Lemma. Let

$$\Gamma^q := \{C^q\text{-metric } g \in \mathcal{O} \text{ conformal to } g_0 : g|_U = g_0|_U\}$$

$$\mathcal{M}^q := \{(g, S) : g \in \Gamma^q; S \text{ is a minimal sphere in } (S^3, g), \text{area}_g(S) < L\};$$

Here we use C^q -topology on Riemannian metrics and $C^{2,\alpha}$ -topology on submanifolds, $\alpha \in (0, 1)$.

Lemma 6.5. *For every integer $3 \leq q < \infty$, \mathcal{M}^q is a separable C^{q-2} Banach manifold modeled on $C^q(S^3)$ and*

$$\Pi : \mathcal{M}^q \rightarrow \Gamma^q, \quad (g, S) \mapsto g$$

is a C^{q-2} Fredholm map with index 0.

Moreover, $g \in \Gamma^q$ is a regular value of Π if and only if for every $(g, S) \in \mathcal{M}^q$, S is non-degenerate.

Proof. For every $(\hat{g}, \hat{S}) \in \mathcal{M}^q$ with unit normal field \hat{v} , we apply [Whi91, Theorem 1.2] with

$$\Gamma := \Gamma^q, \quad X := C^{2,\alpha}(\hat{S}), \quad Y := C^\alpha(\hat{S}), \quad \mathcal{H} := L^2(\hat{S}),$$

$$A : \Gamma \times X \rightarrow \mathbb{R}, \quad (g, w) \mapsto \text{area}_g(\text{graph}_{\hat{S}, \hat{g}}(w)),$$

$$H : \Gamma \times X \rightarrow Y, \quad \text{given by } \langle H(g, w), v \rangle_{L^2} = \left. \frac{d}{dt} \right|_{t=0} A(g, w + tv).$$

where every norm is measured under metric \hat{g} . Note that Γ^q can be identified as an open subset of

$$C_U^q(S^3) := \{f \in C^q(S^3) : f|_U = 0\}.$$

endowed with C^q -topology. This is also a separable Banach space.

Notice that near (\hat{g}, \hat{S}) , $\mathcal{M}^q = \{(g, S) \in \Gamma \times X : H(g, S) = 0\}$. And as is verified in [Whi91, Theorem 1.1],

$$D_2H(\hat{g}, 0) = -(\Delta_{\hat{S}} + |A_{\hat{S}}|^2 + \text{Ric}_{\hat{g}}(\hat{\nu}, \hat{\nu})) =: L_{\hat{S}, \hat{g}} : X \rightarrow Y$$

is a Fredholm map with index 0. Hence to prove the first part of the Lemma, it suffices to verify the condition (C) in [Whi91, Theorem 1.2], namely, for every $0 \neq \kappa \in \ker D_2H(\hat{g}, 0)$, there exists a one parameter family $\{\gamma(s) \in \Gamma^q\}_{s \in (-1, 1)}$ such that $\gamma(0) = \hat{g}$ and

$$(6.1) \quad \left(\frac{\partial^2}{\partial s \partial t} \right)_{s=t=0} A(\gamma(s), t\kappa) \neq 0.$$

To see this, let $\gamma(s) = (1 + sf)\hat{g}$ where $f \in C_U^q(S^3)$. According to the calculation in [Whi91, Theorem 2.1],

$$\left(\frac{\partial^2}{\partial s \partial t} \right)_{s=t=0} A(\gamma(s), t\kappa) = \int_{\hat{S}} \hat{\nu}(f)\kappa \, dx$$

Recall that by Lemma 6.4, $\hat{S} \setminus \bar{U} \neq \emptyset$; And by the unique continuation property, $\kappa|_{S^3 \setminus U}$ is not identically 0. Therefore, (6.1) is true by taking $f \in C_c^q(S^3 \setminus \bar{U})$ such that $\hat{\nu}(f)$ is an L^2 -approximation of $\kappa \cdot \chi_{S^3 \setminus U}$.

To study the regular value of Π , first notice that for every $(g', S') \in \mathcal{M}^q$,

$$\begin{aligned} \text{Tan}_{(g', S')} \mathcal{M}^q &= \{(f, w) : f \in C_U^q(S^3), L_{S', g'} = \hat{\nu}(f) \text{ on } S'\}; \\ d\Pi|_{(g', S')} : \text{Tan}_{(g', S')} \mathcal{M}^q &\rightarrow C_U^q(S^3), (f, w) \mapsto f. \end{aligned}$$

Thus, if $g' \in \Gamma^q$ is a regular value of Π , then for every $(g', S') \in \mathcal{M}^q$ with unit normal field ν' , every $\kappa \in \ker L_{S', g'}$ and every $\phi \in C_c^q(S' \setminus \bar{U})$, we take $f \in C_U^q(S^3)$ such that $\nu'(f) = \phi$, then there exists $w_\phi \in C^{2, \alpha}(S')$ such that $L_{S', g'} w_\phi = \nu'(f) = \phi$. Then

$$0 = \int_{S'} w_\phi \cdot L_{S', g'} \kappa = \int_{S'} \kappa \cdot L_{S', g'} w_\phi = \int_{S'} \kappa \phi.$$

In other words, $\kappa \perp C_c^q(S' \setminus \bar{U})$. Therefore, $\kappa|_{S' \setminus \bar{U}} = 0$. Again by Lemma 6.4 and the unique continuation property, $\kappa \equiv 0$ on S' . Hence, $\ker L_{S', g'} = 0$, that is, S' is non-degenerate. \square

To obtain the third assertion of Proposition 6.1, we need the following lemma in linear algebra.

Lemma 6.6. *Let $k, l \geq 1$. Given $\mathbf{a} \in \mathbb{R}^k, \mathbf{b} \in \mathbb{R}^l$, consider the linear function*

$$f_{\mathbf{a}} : \mathbb{Z}^k \rightarrow \mathbb{R}, \mathbf{u} \mapsto \mathbf{u} \cdot \mathbf{a}; \quad h_{\mathbf{b}} : \mathbb{Z}^l \rightarrow \mathbb{R}, \mathbf{v} \mapsto \mathbf{v} \cdot \mathbf{b}.$$

Then for every $\mathbf{b} \in \mathbb{R}^l$,

$$\mathcal{R} := \{\mathbf{a} \in \mathbb{R}^k : \ker(f_{\mathbf{a}}) = \{0\}, f_{\mathbf{a}}(\mathbb{Z}^k) \cap h_{\mathbf{b}}(\mathbb{Z}^l) = \{0\}\}$$

is a countable intersection of open dense subset of \mathbb{R}^k (and thus dense in \mathbb{R}^k).

Proof. Let $L_k : \mathbb{Z}^k \rightarrow \mathbb{N}$ and $L_l : \mathbb{Z}^l \rightarrow \mathbb{N}$ be bijections, $\Lambda_{k,N} \subset \mathbb{Z}^k$, $\Lambda_{l,N} \subset \mathbb{Z}^l$ be the inverse image of $[0, N]$ under L_k , L_l correspondingly. Clearly, \mathcal{R} is the intersection of

$$\mathcal{R}_N := \left\{ \mathbf{a} \in \mathbb{R}^k : \ker(f_{\mathbf{a}}) \cap \Lambda_{k,N} \subset \{0\}, f_{\mathbf{a}}(\mathbb{Z}^k \cap \Lambda_{k,N}) \cap h_{\mathbf{b}}(\mathbb{Z}^l \cap \Lambda_{l,N}) \subset \{0\} \right\}$$

over $N \in \mathbb{N}$, where each \mathcal{R}_N is open and dense in \mathbb{R}^k . \square

Proof of Proposition 6.1. For every fixed integer $q \geq 3$, $\varepsilon > 0$ and L greater than the largest area among all minimal tori in (S^3, g_0) , let $U \subset S^3$ and $\mathcal{O} \ni g_0$ be determined by Lemma 6.4. By the Sard-Smale Theorem, the regular values of $\Pi : \mathcal{M}^q \rightarrow \Gamma^q$ forms a residual (and hence dense) subset $\Gamma_{bumpy}^q \subset \Gamma^q$. By Lemma 6.2, 6.4, 6.5 and Sharp's compactness theorem [Sha17], the first two assertion of Proposition 6.1 hold for every $g \in \Gamma_{bumpy}^q$. Then by the argument in [Whi17] (See also the proof of [ACS18, Theorem 9]), the first two assertion of Proposition 6.1 also hold for every $g \in \Gamma_{bumpy}^q \cap \Gamma^\infty$, which is dense in Γ^∞ .

Moreover, for each $g'_0 \in \Gamma_{bumpy}^q \cap \Gamma^\infty$, there are only finitely many minimal spheres with area at most L . Therefore, by Lemma 6.2, and Sharp's compactness theorem again, there exists a neighborhood \mathcal{U} of g'_0 in Γ^∞ , such that $\mathcal{U} \subset \Gamma_{bumpy}^q \cap \Gamma^\infty$. In other words, $\Gamma_{bumpy}^q \cap \Gamma^\infty$ is both open and dense in Γ^∞ .

To obtain the third assertion, let $g'_0 \in \Gamma_{bumpy}^q \cap \Gamma^\infty$ be $\varepsilon/2$ - C^∞ close to g_0 and let

$$\{S_1, S_2, \dots, S_k\}$$

be all the embedded minimal 2-spheres in (S^3, g'_0) with area $< L$. Let $r > 0$ and $\{p_j\}_{j=1}^k$ be points in S^3 such that for every j , $p_j \in S_j$ and

$$B_r(p_j) \cap U \cup \bigcup_{i \neq j} S_i = \emptyset.$$

by deforming in $B_r(p_j)$, one can construct an \mathbb{R}^k -parametrized family of metrics $\{g(\mathbf{a})'\}_{\mathbf{a} \in \mathbb{R}^k} \subset \Gamma_{bumpy}^q \cap \Gamma^\infty$ such that $g(0)' = g'_0$, all the embedded $g(\mathbf{a})'$ -minimal spheres $\{S_j(\mathbf{a})'\}_{j=1}^k$ are continuous deformation of $\{S_j\}_{j=1}^k$ and that

$$\mathbf{a} \mapsto \left(\text{area}_{g(\mathbf{a})'}(S_1(\mathbf{a})), \text{area}_{g(\mathbf{a})'}(S_2(\mathbf{a})), \dots, \text{area}_{g(\mathbf{a})'}(S_k(\mathbf{a})) \right)$$

is a local diffeomorphism near 0 onto its image. By Lemma 6.6 with $\mathbf{b} = (\text{area}_{g_0}(T_1), \text{area}_{g_0}(T_2), \dots, \text{area}_{g_0}(T_l))$, in a small neighborhood of 0, those $\mathbf{a} \in \mathbb{R}^k$ such that $g(\mathbf{a})'$ satisfies the third assertion of Proposition 6.1 is residual and hence dense. We choose such a $g' = g(\mathbf{a})'$ within $\varepsilon/2$ -neighborhood of g'_0 in Γ^∞ , which satisfies all the assertion in Proposition 6.1. \square

7. MIN-MAX RESULTS II: CONVERGENCE AS CURRENTS

Theorem 3.5 is analogous to Proposition 4.10 and Theorem 4.11 in [MN21]. However, in our definition of homotopy class (Definition 2.7, two Simon-Smith families must be connected through an isotopy, which precludes the use of the well-known Almgren interpolation scheme. Therefore, the proofs in [MN21] cannot be followed verbatim to justify Theorem 3.5. Instead, we will use results from [Ket19] and [DLP10] to construct appropriate isotopies.

Let $\Phi' : X \rightarrow \mathcal{S}^*(M)$ be a Simon-Smith family of genus $\leq g_0$ in the homotopy class Λ from Theorem 3.5. Suppose that X is a cubical subcomplex of some $I(m, k)$ and $K(m) = 3^{m3^m}$. In addition, by Remark 2.8, we may assume that $N_P \equiv N \in \mathbb{N}$ within the homotopy class Λ . For $L = \mathbf{L}(\Lambda)$, since $\mathcal{W}_{L, \leq g_0}$ consists of finitely many varifolds associated with connected multiplicity-one minimal surfaces, we can enumerate these surfaces as

$$\Gamma_1, \Gamma_2, \dots, \Gamma_q.$$

We fix $r_0 > 0$ such that for each Γ_k , each point $p \in \Gamma_k$, and any $r \in (0, 2r_0)$,

$$B_r(p) \cap \Gamma_k$$

is a topological disk, and moreover, $\partial B_r(p)$ is a mean-convex sphere with transversal intersection with Γ_k :

$$\partial B_r(p) \pitchfork \Gamma_k.$$

Lemma 7.1 ([Pit81, 2.1 (f)]). *Suppose that $T, T_1, T_2, \dots \in \mathcal{Z}_2(M; \mathbb{Z}_2)$, $\lim_i T_i = T$ in the flat topology and $\lim_i |T_i| = V \in \mathcal{V}_2(M)$ in the weak topology. Then*

$$\|T\| \leq \|V\|.$$

Corollary 7.2. *For any $r > 0$, there exists $r_1 > 0$ and $\delta_1 > 0$ such that for every $V \in \mathcal{W}_{L, \leq g_0}$ associated with some Γ_k , the set*

$$\mathcal{D}_k := \{T \in \mathcal{Z}_2(M; \mathbf{F}; \mathbb{Z}_2) : \mathbf{F}(|T|, V) < r_1\}$$

has exactly two connected components $\mathcal{D}_k^0 \cup \mathcal{D}_k^1$ characterized by

$$\begin{aligned} \mathcal{D}_k^0 &= \{T \in \mathcal{D}_k : \mathcal{F}(T, 0) < \delta_1\} \\ \mathcal{D}_k^1 &= \{T \in \mathcal{D}_k : \mathcal{F}(T, 0) > \delta_1\} = \{T \in \mathcal{D}_k : \mathbf{F}(T, [\Gamma_k]) < r\}. \end{aligned}$$

Proof. By Lemma 7.1, every sequence $\{T_i\}_i \subset \mathcal{Z}_2(M; \mathbb{Z}_2)$ satisfying $\lim_i T_i = T$ and $\lim_i |T_i| = V$ has

$$\|T\| \leq \|V\|.$$

Since V is associated with a connected multiplicity-one minimal surface Γ_k and $\partial T = 0$, by the constancy theorem, we have either $T = [\Gamma_k]$ or $T = 0$. Thus, by choosing suitable r_1 and δ_1 , the conclusion follows immediately. \square

Fix r from Theorem 3.5 and then choose r_1 and δ_1 from the previous Corollary. The crucial proposition we need is the following:

Proposition 7.3. *For every $\alpha \in (0, 2r_0)$, there exists $\varepsilon \in (0, r_1)$ such that for any generalized surface $\Sigma \in \mathcal{S}(M)$ and any $\delta > 0$, if for some Γ_k and some $p \in \Gamma_k$,*

- (1) $\mathbf{F}(|\Sigma|, |\Gamma_k|) < \varepsilon$,
- (2) $\Sigma \cap B_\alpha(p)$ is smooth,
- (3) $\mathbf{g}(\Sigma \cap B_\alpha(p)) = 0$,
- (4) $[\Sigma] \in \mathbf{a}(B_\alpha(p); \varepsilon, \delta)$,

then $\mathcal{F}([\Sigma], 0) > \delta_1$.

Proof. Suppose for the sake of contradiction that for $\alpha > 0$, Γ_k and $p \in \Gamma_k$, there exists a sequence of generalized surfaces $\{\Sigma_i\} \subset \mathcal{S}(M)$ and a sequence of positive number $\{\delta^{(i)}\}$ such that for all large i ,

- (1) $\mathbf{F}(|\Sigma_i|, |\Gamma_k|) < \frac{1}{i}$,
- (2) $\Sigma_i \cap B_\alpha(p)$ is smooth,
- (3) $\mathbf{g}(\Sigma_i \cap B_\alpha(p)) = 0$,
- (4) $[\Sigma_i] \in \mathbf{a}(B_\alpha(p); \frac{1}{i}, \delta^{(i)})$,

but $\mathcal{F}([\Sigma_i], 0) \leq \delta_1$. Hence, by Lemma 7.1 and Corollary 7.2, we have $\mathcal{F}([\Sigma_i], 0) < \delta_1$ and

$$\lim_{i \rightarrow \infty} \mathcal{F}([\Sigma_i], 0) = 0.$$

We can choose $\bar{r} \in (\alpha/2, \alpha) \subset (0, 2r_0)$ such that for each i , $\partial B_{\bar{r}}(p) \pitchfork \Sigma_i$. Note that by definition, we also have $[\Sigma_i] \in \mathbf{a}(B_{\bar{r}}(p); \frac{1}{i}, \delta^{(i)})$.

Following [Ket19, Section 3.2], let $\mathfrak{I}\mathfrak{s}_i(\Sigma_i, B_{\bar{r}}(p))$ denote the set of all isotopies ϕ of M supported in $B_{\bar{r}}(p)$ such that for all $t \in [0, 1]$,

$$\mathcal{H}^2(\phi(t, \Sigma_i)) \leq \mathcal{H}^2(\Sigma_i) + \delta^{(i)}.$$

For each $i = 1, 2, 3, \dots$, we can take a *minimizing sequence* $\phi^l(1, \Sigma_i)$ for $\text{Problem}(\Sigma_i, \mathfrak{I}\mathfrak{s}_i(\Sigma_i, B_{\bar{r}}(p)))$, i.e.,

$$\lim_{l \rightarrow \infty} \mathcal{H}^2(\phi^l(1, \Sigma_i)) = \inf_{\phi \in \mathfrak{I}\mathfrak{s}_i(\Sigma_i, B_{\bar{r}}(p))} \mathbf{M}(\phi(1, \Sigma_i)),$$

and let V_i be the varifold limit of $\phi^l(1, \Sigma_i)$.

It follows from [Ket19, Lemma 3.12] that as $i \rightarrow \infty$,

$$V_i \rightarrow |\Gamma_k|$$

in the varifold topology, and the convergence is smooth in compact subsets of $B_{\bar{r}}(p)$. By [DLP10, Proposition 3.2], we also know that $V_i \cap B_{\bar{r}}(p)$ is associated with a smooth minimal surface Δ_i such that $\partial \Delta_i = \partial(\Sigma_i \cap B_{\bar{r}}(p))$.

By [Ket19, Proposition 4.7], we have

$$\mathbf{g}(\Delta_i) \leq \mathbf{g}(\Sigma_i \cap B_{\bar{r}}(p)) = 0,$$

so Δ_i is a minimal disk. In particular, Δ_i separates $B_{\bar{r}}(p)$, and V_i is associated with a cycle $T_i \in \mathcal{Z}_2(M; \mathbb{Z}_2)$.

On one hand, note that

$$T_i \llcorner (M \setminus B_{\bar{r}}(p)) = [\Sigma_i] \llcorner (M \setminus B_{\bar{r}}(p)).$$

In particular,

$$T_i \rightarrow 0$$

outside $B_{2r_0}(p)$.

On the other hand, in $B_{\alpha/2}(p)$, for sufficiently large i , Δ_i is a normal graph over Σ_i , $B_{\alpha/2}(p) \setminus \Delta_i$ has two connected components E_i^1 and E_i^2 and $B_{\alpha/2}(p) \setminus \Gamma_k$ has two connected components E^1 and E^2 satisfying

$$\min\{\text{vol}(E_i^1), \text{vol}(E_i^2)\} \geq \min\{\text{vol}(E^1), \text{vol}(E^2)\}/2 > 0.$$

Therefore, we have

$$\liminf_i \mathcal{F}(T_i, 0) \geq \min\{\text{vol}(E^1), \text{vol}(E^2)\}/2 > 0.$$

Since $|T_i| = V_i \rightarrow |\Gamma_k|$, by Lemma 7.1, we have $\lim T_i = [\Gamma_k]$. This contradicts that $T_i \rightarrow 0$ outside $B_{2r_0}(p)$. \square

Proof of Theorem 3.5. As above, for $r > 0$, we choose r_1 and δ_1 from Corollary 7.2 and also choose ε from Proposition 7.3 with the parameters r_1 and δ_1 .

Let $\alpha = r_0/(16 \cdot 8^{\bar{N}})$ and choose ε from Proposition 7.3. We set $r_3 := \min(r, r_1, \varepsilon)$.

For each Γ_k , choose $p_k \in \Gamma_k$ and for each $i = 1, \dots, \bar{N}$, we set $r_i = \frac{7r_0}{8^i}$ and $s_i = \frac{5r_0}{8^i}$, and we obtain a \bar{N} -admissible collection of annuli, $\{A_i^k := A(p_k; s_i, r_i)\}$. We also choose $\{p_k^1, \dots, p_k^{\bar{N}}\} \subset \Gamma_k$ such that for each $i = 1, \dots, \bar{N}$,

$$\text{dist}(p_k, p_k^i) = \frac{3r_0}{4 \cdot 8^{i-1}}.$$

and thus, $B(p_k^i, \alpha) \subset A_i^k$.

By Theorem 2.11, there exists a pulled-tight sequence $\{\Phi_i\}$ in Λ and some $\eta_3 \in (0, r_3)$ such that for each i and each $x \in X$,

$$\mathcal{H}^2(\Phi_i(x)) \geq L - \eta_3 \implies |\Phi_i(x)| \in \mathbf{B}_{r_3}^{\mathbf{F}}(\mathcal{W}_{L, \leq g_0}).$$

It follows from Corollary 7.2 that the compact set

$$Y_i := \{x \in X : \mathcal{H}^2(\Phi_i(x)) \geq L - \eta_3\}$$

can be decomposed as two disjoint compact sets Y_i^0 and Y_i^1 :

$$Y_i^0 = \{x \in Y_i : \mathcal{F}([\Phi_i(x)], 0) < \delta_1\}$$

$$Y_i^1 = \{x \in Y_i : \mathcal{F}([\Phi_i(x)], 0) > \delta_1\} = \{x \in Y_i : \mathbf{F}([\Phi_i(x)], [\mathcal{W}_{L, \leq g_0}]) < r\}.$$

Now for each $x \in Y_i^0$, we can choose k such that $\mathbf{F}(|\Phi_i(x)|, |\Gamma_k|) < r_3 \leq \varepsilon$. Since $\mathfrak{g}(\Phi_i(x)) \leq \mathfrak{g}_0$ and we can choose a set of at most N points $P(x)$ such that $\Phi_i(x) \setminus P(x)$ is a smooth surface, there are at least $K(m)$ points $\{p_k^{j_l}\}_{j_l}^{K(m)}$ from $\{p_k^j\}_{j=1}^{\bar{N}}$ where for each l ,

- (1) $\Phi_i(x) \cap B_\alpha(p_k^{j_l})$ is smooth,
- (2) $\mathfrak{g}(\Phi_i(x) \cap B_\alpha(p_k^{j_l})) = 0$,

hence, Σ admits an $(\varepsilon, \frac{\eta_3}{1000K(m)})$ deformation in $B_\alpha(p_k^{j_i})$.

By Proposition 4.3, we can obtain Φ_i^* for each Φ_i . Moreover, for sufficiently large i , since $\eta_3 < \varepsilon$, we have the following properties.

For every $x \in Y_i^0$,

$$\begin{aligned} \mathcal{H}^2(\Phi_i^*(x)) &< \mathcal{H}^2(\Phi_i(x)) - \varepsilon + (3^m - 1) \frac{\eta_3}{1000K(m)} \\ &< \mathcal{H}^2(\Phi_i(x)) - \eta_3/2 \\ &< L - \eta_3/2. \end{aligned}$$

For every $x \in X \setminus Y_i$,

$$\begin{aligned} \mathcal{H}^2(\Phi_i^*(x)) &< \mathcal{H}^2(\Phi_i(x)) + 3^m \frac{\varepsilon}{1000K(m)} \\ &< L - \eta_3 + 3^m \frac{\eta_3}{1000K(m)} \\ &< L - \eta_3/2. \end{aligned}$$

Since Y_i^0 and Y_i^1 are disjoint compact set, we also have $\Phi_i^*(x) = \Phi_i(x)$ holds for every $x \in Y_i^1$.

Therefore, for every $x \in X$ with $\mathcal{H}^2(\Phi_i^*(x)) \geq L - \eta_3/2$, then $x \in Y_i^1$ and thus,

$$[\Phi_i(x)] \in \mathbf{B}_{r_3}^{\mathbf{F}}([\mathcal{W}_{L, \leq g_0}]) \subset \mathbf{B}_r^{\mathbf{F}}([\mathcal{W}_{L, \leq g_0}]),$$

which completes the proof with $\eta = \eta_3/2$ and $\Phi = \Phi_i^*$. \square

8. MIN-MAX RESULTS III: INTERPOLATION

The goal of this section is to prove Theorem 3.6. Let (M, g) be a 3-dimensional closed Riemannian manifold.

The key is the following deformation proposition. Intuitively, for a Simon-Smith family of genus ≤ 1 very close to spheres, we show that we can pinch all the small necks and shrink all small connected components in a continuous way to obtain a Simon-Smith family of genus 0.

Proposition 8.1. *Let X be a cubical subcomplex of $I(m, k)$, \mathcal{S} be a compact subset of embeddings of \mathbb{S}^2 into (M, g) (endowed with smooth topology), $q \in \mathbb{N}$ and $\varepsilon \in (0, 1)$. Then there exists $\delta = \delta(\varepsilon, q, m, \mathcal{S}) \in (0, 1)$ with the following property.*

If $\Phi : X \rightarrow \mathcal{S}(M)$ is a Simon-Smith family of genus ≤ 1 , $N_P(\Phi) < q$ and $S : X \rightarrow \mathcal{S}$ is a map such that for every $x \in X$,

$$\mathbf{F}([\Phi(x)], [S(x)]) \leq \delta$$

then there exists a Simon-Smith family $H : [0, 1] \times X \rightarrow \mathcal{S}(M)$ of genus ≤ 1 such that:

- (1) *For every $x \in X$, $H(0, x) = \Phi(x)$, $\mathbf{g}(H(1, x)) = 0$.*
- (2) *For every $x \in X$ and $0 \leq t \leq t' \leq 1$,*

$$\mathcal{H}^2(H(t, x)) \leq \mathcal{H}^2(\Phi(x)) + \varepsilon, \quad \mathbf{g}(H(t', x)) \leq \mathbf{g}(H(t, x)).$$

(3) For each $x \in X$, the Simon-Smith family $\{H(x, t)\}_{t \in [0,1]}$ is a pinch-off process.

We will prove the proposition using the following technical lemmas.

8.1. Nontrivial loop in a small ball. In this subsection, we show that if a (singular) torus is sufficiently close to a sphere, we can find a loop contained within a small ball. In the next subsection, we will explain how to pinch the torus along the boundary of the small ball and then shrink all connected components inside to a point. The small size of the ball helps control the area increase, as required in Proposition 8.1 (2).

Lemma 8.2 (Existence of small nontrivial loops). *Let \mathcal{S} be a compact subset of embeddings of \mathbb{S}^2 into (M, g) (endowed with smooth topology), $\varepsilon \in (0, 1)$. Then there exists $\delta_1(\varepsilon, \mathcal{S}) \in (0, 1)$ with the following property.*

Suppose that $\Sigma \in \mathcal{S}(M)$ has $\mathfrak{g}(\Sigma) = 1$, and there exists a $S \in \mathcal{S}$ such that $\mathbf{F}([\Sigma], [S]) \leq \delta_1(\varepsilon, \mathcal{S})$. Then there exists $p \in S^3$ and an embedded loop $\gamma \subset \Sigma \cap B(p, \varepsilon)$ such that the surface obtained by neckpinch along γ of Σ has genus 0.

Proof. First, we assume that $\mathcal{S} = \{S\}$. The general case will follow by applying a finite covering argument to the compact set \mathcal{S} , which will be addressed at the end.

Suppose that there does not exist such a $\delta_1(\varepsilon, \mathcal{S})$, and then there exists a sequence $\{\Sigma_i\} \subset \mathcal{S}(M)$ with $\mathfrak{g}(\Sigma_i) = 1$ and $\lim_{i \rightarrow \infty} \mathbf{F}([\Sigma_i], [S]) = 0$ but neither Σ_i contains a nontrivial loop γ in any geodesic ball of radius ε .

Step 1. We perform the first surgeries on the singular points of each Σ_i such that they are all smooth surfaces.

By the definition of generalized surface, for each Σ_i , there exists a finite set $P_i \subset \Sigma_i$, such that $\Sigma_i \setminus P_i$ is a smooth surface. By Sard's theorem and the definition of $\mathfrak{g}(\Sigma)$, we can choose an arbitrarily small radius $r_i > 0$ such that

$$B_{r_i}(P_i) \pitchfork (\Sigma_i \setminus P_i)$$

is a finite set of circles, and

$$\mathfrak{g}(\Sigma \setminus B_{r_i}(P_i)) = \mathfrak{g}(\Sigma) = 1.$$

Therefore, by choosing sufficiently small radius r_i and replacing $B_{r_i}(P_i) \cap \Sigma_i$ with a finite number of disks, we obtain a smooth surface $\Sigma_i^{(1)}$ satisfying

$$\mathfrak{g}(\Sigma_i^{(1)}) = 1, \quad \lim_{i \rightarrow \infty} \mathbf{F}([\Sigma_i^{(1)}], [S]) = 0.$$

Moreover, if $\Sigma_i^{(1)}$ contains a nontrivial loop $\gamma_i^{(1)}$ within a geodesic ball of radius $\varepsilon/2$, we can use the simply-connectedness of disks to perturb $\gamma_i^{(1)}$, yielding a nontrivial loop $\gamma \subset \Sigma_i \cap \Sigma_i^{(1)} \subset \Sigma_i$ in a geodesic ball of radius ε , which contradicts to our assumptions.

In the following, we choose $r^{(1)} \in (0, \varepsilon/100)$ depending on S such that there exists a finite covering $\{B_{r^{(1)}}(p_j)\}_{j=1}^K$ of S with the following properties:

- $\{p_j\} \subset S$;
- For each $r \in (0, 2r^{(1)})$, $B_{2r^{(1)}}(p_j) \cap S$ is a disk;
- Every pair $(B_{2r^{(1)}}(p_j), B_{2r^{(1)}}(p_j) \cap S)$ is diffeomorphic to the pair $(\mathbb{B}_{2r^{(1)}}^3, \mathbb{B}_{2r^{(1)}}^3 \cap \{x_3 = 0\})$ in \mathbb{R}^3 with bi-Lipschitz constant $1 + \frac{1}{10000}$;

Then we choose $\varepsilon^{(1)} \in (0, r^{(1)}/100)$ such that the normal exponential map

$$\exp_S^\perp : S \times [-\varepsilon^{(1)}, \varepsilon^{(1)}] \rightarrow T_{\varepsilon^{(1)}}(S)$$

is a diffeomorphism with bi-Lipschitz constant $1 + \frac{1}{10000 \cdot 10^K}$. We denote by $\pi : T_{\varepsilon^{(1)}}(S) \rightarrow S$ the corresponding closest-point projection.

Step 2. We now perform the second surgeries on $\Sigma_i^{(1)}$ such that they are all contained in $T_{\varepsilon^{(1)}}(S)$.

Since $\lim_{i \rightarrow \infty} \mathbf{F}([\Sigma_i^{(1)}], [S]) = 0$, for sufficiently large i , there exists $\eta_i \in (0, \infty)$ such that

$$(8.1) \quad \mathcal{H}^2(\Sigma_i^{(1)} \setminus T_{\varepsilon^{(1)}/2}(S)) < \eta_i, \quad \lim_{i \rightarrow \infty} \eta_i = 0.$$

By the slicing lemma and Sard's lemma, there exists $\sigma \in (\varepsilon^{(1)}/2, \varepsilon^{(1)})$ such that for each sufficiently large i , $\Sigma_i^{(1)} \cap \partial(T_\sigma(S))$ is a finite set of circles and

$$(8.2) \quad \mathcal{H}^1(\Sigma_i^{(1)} \cap \partial(T_\sigma(S))) \leq \frac{4\eta_i}{\varepsilon^{(1)}}.$$

For sufficiently large i , we can perform a surgery on Σ_i in a small neighborhood of $\partial(T_\sigma(S))$ by removing cylinders and gluing appropriate small disks. We will denote the union of all the connected components contained inside $T_\sigma(S)$ by $\Sigma_i^{(2)}$ and the remaining part by $\tilde{\Sigma}_i^{(2)}$. There are three possibilities:

Case 1 $\mathbf{g}(\Sigma_i^{(2)}) = \mathbf{g}(\tilde{\Sigma}_i^{(2)}) = 0$. In this case, the surgery near $\partial(T_\sigma(S))$ detect a nontrivial loop $\gamma^{(1)} \subset \Sigma_i^{(1)}$. For sufficiently large i , by (8.2), the loop $\gamma^{(1)}$ has length no greater than $\varepsilon^{(1)}/100$ and thus, is contained in a geodesic ball of radius $\varepsilon^{(1)}$. This contradicts our assumptions.

Case 2 $\mathbf{g}(\Sigma_i^{(2)}) = 0$ and $\mathbf{g}(\tilde{\Sigma}_i^{(2)}) = 1$. In this case, for sufficiently small η , by the isoperimetric inequality, (8.1) and (8.2) imply that

$$(8.3) \quad \mathcal{H}^2(\tilde{\Sigma}_i^{(2)}) \leq \eta_i + C_2 \cdot \left(\frac{4\eta_i}{\varepsilon^{(1)}}\right)^2$$

for some constant $C_2 = C_2(M, g)$ from the isoperimetric inequality. In (M, g) , we can find a constant $\tilde{c}_2 = \tilde{c}_2(M, g)$ and a triangulation Δ in dependent of i with the following property.

- Every cell transversally intersect with $\tilde{\Sigma}_i^{(2)}$;
- For every 3-cell C in Δ , $\partial C \cap \Sigma_i^{(2)}$ is a finite set of circles;

- Let $\Delta^{(2)}$ be the union of all 2-cells of Δ , and then $\Delta^{(2)} \cap \widetilde{\Sigma}_i^{(2)}$ satisfies

$$(8.4) \quad \mathcal{H}^1(\widetilde{\Sigma}_i^{(2)} \cap \Delta^{(2)}) \leq \frac{\mathcal{H}^2(\widetilde{\Sigma}_i^{(2)})}{\widetilde{c}_2}.$$

Indeed, this can be achieved as follows: we select a triangulation Δ and an isotopy $\{\varphi_t\}_{t \in [0,1]}$ of (M, g) induced by a smooth vector field with only finitely many fixed points outside $\Delta^{(2)}$. Then applying the isotopy $\{\varphi_t\}$ to the given triangulation $\Delta^{(2)}$. At least one of the resulting deformed triangulation $(\varphi_{t_0})_{\#}(\Delta)$ will satisfy the desired transversal property and length estimates.

Then we can perform surgery in a neighborhood of $\Delta^{(2)}$ by replacing cylinders with disks and obtain a union of surfaces, each of which is contained in some 3-cell in Δ . There are two possibilities. Case 2a All new surfaces have genus 0. Then, the surgery near $\Delta^{(2)}$ detects a nontrivial loop γ' of $\widetilde{\Sigma}_i^{(2)}$. Since we only glue disks in all the surgeries, we can perform perturbations to get a nontrivial loop $\gamma^{(1)}$ of $\Sigma_i^{(1)}$. By (8.2), (8.3) and (8.4), $\gamma^{(1)}$ has length

$$\begin{aligned} \mathcal{H}^1(\gamma^{(1)}) &\leq \mathcal{H}^1(\Sigma_i^{(1)} \cap \partial(T_\sigma(S))) + \mathcal{H}^1(\widetilde{\Sigma}_i^{(2)} \cap \Delta^{(2)}) \\ &\leq \eta_i + C_2 \cdot \left(\frac{4\eta_i}{\varepsilon^{(1)}}\right)^2 + \frac{\eta_i + C_2 \cdot \left(\frac{4\eta_i}{\varepsilon^{(1)}}\right)^2}{\widetilde{c}_2}. \end{aligned}$$

For sufficiently large i , $\gamma^{(1)}$ is contained a geodesic ball of radius $\varepsilon^{(1)}$, a contradiction.

Case 2b Some new surface Σ' has genus 1, i.e., it contains a nontrivial loop γ' . Again, we can perform perturbations to get a nontrivial loop $\gamma^{(1)}$ of $\Sigma_i^{(1)}$. As in Case 2a, for sufficiently large i , $\gamma^{(1)}$ is contained a geodesic ball of radius $\varepsilon^{(1)}$, a contradiction.

In summary, Case 2 is impossible.

Case 3 $\mathfrak{g}(\Sigma_i^{(2)}) = 1$ and $\mathfrak{g}(\widetilde{\Sigma}_i^{(2)}) = 0$.

Since Case 1 and Case 2 are impossible, in the following, we assume that Case 3 holds. For simplicity, we will discard $\widetilde{\Sigma}_i^{(2)}$ and focus on $\Sigma_i^{(2)} \subset T_{\varepsilon^{(1)}}(S)$. It is easy to see that we also have

$$(8.5) \quad \lim_i \mathbf{F}([\Sigma_i^{(2)}], [S]) = 0.$$

In particular, for sufficiently large i , S and $\Sigma_i^{(2)}$ are homologous to each other in $T_{\varepsilon^{(1)}}(S)$.

As before, for sufficiently large i , if $\Sigma_i^{(2)}$ contains a nontrivial loop $\gamma^{(2)}$ in a geodesic ball of radius $\varepsilon/2$, there exists a nontrivial loop γ of Σ_i in a geodesic ball of radius ε , which gives us a contradiction.

Step 3. Recall the finite covering $\{B_{r^{(1)}}(p_j)\}$ introduced in the end of Step 1. For sufficiently large i , we claim that a.e. $r \in (r^{(1)}, \frac{7}{4}r^{(1)})$,

$$(8.6) \quad \mathcal{H}^1(\pi^{-1}(\partial(B_r(p_j) \cap S)) \cap \Sigma_i^{(2)}) \geq (1 - \frac{1}{1000 \cdot 10^K}) \mathcal{H}^1(\partial(B_r(p_j) \cap S)).$$

Otherwise, by Sard's theorem we can choose some $r \in (r^{(1)}, \frac{7}{4}r^{(1)})$ such that $\pi^{-1}(\partial(B_r(p_j) \cap S)) \cap \Sigma_i^{(2)}$ is a finite union of circles with

$$\mathcal{H}^1(\pi^{-1}(\partial(B_r(p_j) \cap S)) \cap \Sigma_i^{(2)}) < (1 - \frac{1}{1000 \cdot 10^K}) \mathcal{H}^1(\partial(B_r(p_j) \cap S)).$$

In particular, from the choice of $r^{(1)}$ and $\varepsilon^{(1)}$, the circles are homologous trivial in $\pi^{-1}(\partial(B_r(p_j) \cap S))$, while $\partial(B_r(p_j) \cap S)$ is not. This contradicts that S and $\Sigma_i^{(2)}$ are homologous to each other in $N_{\varepsilon^{(1)}}(S)$.

On the other hand, for sufficiently large i , by the slicing lemma, there exists a subset A of $(r^{(1)}, \frac{7}{4}r^{(1)})$ of positive measure such that for every $r \in A$,

$$(8.7) \quad \mathcal{H}^1(\pi^{-1}(\partial(B_r(p_j) \cap S)) \cap \Sigma_i^{(2)}) \leq (1 + \frac{1}{1000 \cdot 10^K}) \mathcal{H}^1(\partial(B_r(p_j) \cap S)).$$

Combing (8.6) and (8.7), for every p_j , there exists $r_j \in (r^{(1)}, \frac{7}{4}r^{(1)})$ such that $\pi^{-1}(\partial(B_{r_j}(p_j) \cap S)) \cap \Sigma_i^{(2)}$ is a finite union of circles satisfying

$$\frac{\mathcal{H}^1(\pi^{-1}(\partial(B_{r_j}(p_j) \cap S)) \cap \Sigma_i^{(2)})}{\mathcal{H}^1(\partial(B_{r_j}(p_j) \cap S))} \in \left[1 - \frac{1}{1000 \cdot 10^K}, 1 + \frac{1}{1000 \cdot 10^K}\right].$$

Moreover, following the same argument in the proof of (8.7) before, there exists exactly one circle in $\pi^{-1}(\partial(B_{r_j}(p_j) \cap S)) \cap \Sigma_i^{(2)}$ which is homologous nontrivial in the boundary $\pi^{-1}(\partial(B_{r_j}(p_j) \cap S))$ whose length is at least $(1 - \frac{1}{1000 \cdot 10^K}) \mathcal{H}^1(\partial(B_{r_j}(p_j) \cap S))$, referred as the “long circle.” Hence, the sum of the lengths of the remaining circles are at most $\frac{2}{1000 \cdot 10^K} \mathcal{H}^1(\partial(B_{r_j}(p_j) \cap S))$, referred as the “short circles.” Note that short circles are all homologous trivial, so we can perform surgeries on them.

Note that if $\mathfrak{g}(\pi^{-1}(B_{r_j}(p_j) \cap S) \cap \Sigma_i^{(2)}) = 1$ for some j , then there exists a nontrivial loop of $\Sigma_i^{(2)}$ in a geodesic ball of radius $\varepsilon/2$, a contradiction. Therefore, for each j , we have

$$(8.8) \quad \mathfrak{g}(\pi^{-1}(B_{r_j}(p_j) \cap S) \cap \Sigma_i^{(2)}) = 1.$$

Now we would like to inductively perform surgeries on short circles in each neighborhood of $\pi^{-1}(\partial(B_{r_j}(p_j) \cap S))$ from $j = 1$ to K , and obtain a new surface $\Sigma_i^{(3)}$. The issue is that the surgery near the cylinder $\pi^{-1}(\partial(B_{r_j}(p_j) \cap S))$ might change the “short circles” or the “long circle” in another cylinder $\pi^{-1}(\partial(B_{r_{j'}}(p_{j'}) \cap S))$. A priori, some short circle might become homologous nontrivial, and we cannot perform the next surgery. Fortunately, note that every surgery on $\pi^{-1}(\partial(B_{r_j}(p_j) \cap S))$ will only change the lengths of circles in nearby cylinders by at most twice the lengths of the current “short circles”.

Since the initial “short circles” in each cylinder have total length at most $\frac{2}{1000 \cdot 10^K} \mathcal{H}^1(\partial(B_{r_j}(p_j) \cap S))$, the previous argument implies that after performing surgeries on short circles in the neighborhood of $\pi^{-1}(\partial(B_{r_k}(p_k) \cap S))$ for some k , the sum of lengths of short circles in any $\pi^{-1}(\partial(B_{r_j}(p_j) \cap S))$ with $j > k$ is at most

$$\frac{2 \cdot 3^k}{1000 \cdot 10^K} \mathcal{H}^1(\partial(B_{r_j}(p_j) \cap S)) \leq \frac{1}{100} cH^1(\partial(B_{r_j}(p_j) \cap S)).$$

In particular, short circles will always be homologous trivial, and the issue would not occur.

By (8.8) and the fact that our surgery does not increase genus, we know that for each j ,

$$\mathfrak{g}(\pi^{-1}(B_{r_j}(p_j) \cap S) \cap \Sigma_i^{(3)}) = 1.$$

Since $\pi^{-1}(\partial(B_{r_j}(p_j) \cap S)) \cap \Sigma_i^{(3)}$ is a circle, $\pi^{-1}(B_{r_j}(p_j) \cap S) \cap \Sigma_i^{(2)}$ consists of a disk homologous to $B_{r_j}(p_j) \cap S$, and multiple spheres. Therefore, $\Sigma_i^{(3)}$ consists of a large piece isotopic to S and multiple small spheres, i.e.,

$$\mathfrak{g}(\Sigma^{(3)}) = 0.$$

In particular, some surgery along the cylinders $\pi^{-1}(\partial(B_{r_j}(p_j) \cap S))$ detects a nontrivial loop of $\Sigma_i^{(2)}$, which lies in a geodesic ball of radius $\varepsilon/2$. This contradicts to our assumption.

Therefore, in the case where $\mathcal{S} = \{S\}$, there exists a $\delta_1(\varepsilon, \{S\})$ such that the lemma holds.

Finally, for a general compact set \mathcal{S} , each $S \in \mathcal{S}$ has a positive number $\delta_1(\varepsilon, \{S\})$ as above. The compactness of \mathcal{S} implies that there exists a finite subset $\{S_i\}$ such that for every $S' \in \mathcal{S}$, we can find an S_{i_0} with $\mathbf{F}([S'], [S_{i_0}]) \leq \delta_1(\varepsilon, \{S_{i_0}\})/2$.

Thus, we can set

$$\delta_1(\varepsilon, \mathcal{S}) := \min_i \delta_1(\varepsilon, \{S_i\})/2,$$

and the lemma follows immediately. \square

8.2. Local deformations. In the previous subsection, we located nontrivial loops for singular tori within small balls, provided that they are sufficiently close to a sphere. One key reason we do not simply pinch along the loops, which resemble mean curvature flow, is the need to construct the pinching in a continuous manner along a Simon-Smith family. The singular set presents a technical obstruction to this straightforward approach. Therefore, we employ a two-step modification process, Pinching and Shrinking, as detailed in the following lemmas.

Lemma 8.3 (Local Deformation A: Pinching). *There exists a constant C_g depending only on (M, g) with the following property.*

Let $\Lambda > 0$, $0 < r < \text{inrad}(M, g)/4$, $p \in M$. Suppose that $\Phi : X \rightarrow \mathcal{S}^*(M)$ is a Simon-Smith family and $x_0 \in X$ so that $\Phi(x_0) \in \mathcal{S}(M)$ satisfies

$$\mathcal{H}^2(\Phi(x_0) \cap A(p; r, 2r)) \leq \Lambda r^2.$$

Then there exist $s \in (r, 2r)$, $\zeta \in (0, \min\{\Lambda r, s - r, 2r - s\}/5)$, an integer $K \geq 0$, a neighborhood O_{x_0} of x_0 in X , and a Simon-Smith family

$$H : [0, 1] \times O_{x_0} \rightarrow \mathcal{S}(M),$$

with the following properties.

- (1) For every $y \in O_{x_0}$, $\Sigma_y := \Phi(y) \cap A(p; s - 5\zeta, s + 5\zeta)$ is a surface, varying smoothly in y .
- (2) For every $s' \in [s - 4\zeta, s + 4\zeta]$ and every $y \in O_{x_0}$, we have $\Phi(y)$ intersects $\partial B_{s'}(p)$ transversally, and

$$\Phi(y) \cap \partial B_{s'}(p) =: \bigsqcup_{i=1}^K \gamma_i(s', y), \quad \sum_{i=1}^K \mathcal{H}^1(\gamma_i(s', y)) \leq 2\Lambda r.$$

If $K = 0$, then $H(t, y) \equiv \Phi(y)$ for every $y \in O_{x_0}$ and $t \in [0, 1]$; and if $K \geq 1$, then

- (3) For every $y \in O_{x_0}$, $H(0, y) = \Phi(y)$.
- (4) There exists $t_0 = 0 < t_1 < t_2 < \dots < t_K < 1 = t_{K+1}$ such that for every $0 \leq i \leq K$, $t \in (t_i, t_{i+1})$ and $y \in O_{x_0}$,

$$\Sigma_{t,y} := H(t, y) \cap A(p, s - 5\zeta, s + 5\zeta)$$

is a surface, which is diffeomorphic to $\Sigma_y \setminus \bigsqcup_{1 \leq j \leq i} \gamma_j(s, y)$ attached with $2i$ disks. Moreover, $H(1, y) \cap A(p, s - 5\zeta, s + 5\zeta)$ is also a surface and

$$H(1, y) \cap A(p, s - \zeta, s + \zeta) = \emptyset.$$

- (5) For every $i \in \{1, 2, \dots, K\}$ and $y \in O_{x_0}$, near t_i , $H(t, y)$ is a surgery process via pinching the cylinder $\bigcup_{s' \in (s-2\zeta, s+2\zeta)} \gamma_i(s', y)$;
- (6) For every $y \in O_{x_0}$ and $t \in [0, 1]$, we have

$$\begin{aligned} H(t, y) \setminus A(p; s - 3\zeta, s + 3\zeta) &= \Phi(y) \setminus A(p; s - 3\zeta, s + 3\zeta); \\ \mathcal{H}^2(H(t, y)) &\leq \mathcal{H}^2(\Phi(y)) + C_g \Lambda^2 r^2. \end{aligned}$$

Remark 8.4. For every $y \in O_{x_0}$, by (5), we see that $\{H(t, y)\}_{t \in [0, 1]}$ is a pinch-off process.

Proof. Since $\mathcal{H}^2(\Phi(x_0)) \leq \Lambda r^2$ and $\Phi(x_0) \setminus P$ is a smooth surface for some finite set P , by the slicing theorem and Sard's lemma, there exists $s \in (r, 2r)$ such that $\Phi(x_0) \cap \partial B_r(p)$ is a finite union of loops $\{\gamma_i(s, x_0)\}_{i=1}^K$ with

$$\sum_{i=1}^K \mathcal{H}^1(\gamma_i(s, x_0)) \leq \Lambda r.$$

By Definition 2.3 (4), we can choose ζ and O_{x_0} such that (1) and (2) hold. Moreover, one can find a continuous family of $\{\varphi_y\}_{y \in O_{x_0}} \subset \text{Diff}^\infty(M)$ such that

$$\varphi_y(\Phi(x_0)) \cap A(p; s - 5\zeta, s + 5\zeta) = \Sigma_y.$$

Therefore, by possibly choosing a smaller O_{x_0} , it suffices to find a continuous map $\{H(t, x_0)\}_{t \in [0, 1]}$ for Σ supported in $A(p; s - 4\zeta, s + 4\zeta)$ satisfying (3) and

- (4)' There exists $t_0 = 0 < t_1 < t_2 < \dots < t_K < 1 = t_{K+1}$ such that for every $0 \leq i \leq K$, $t \in (t_i, t_{i+1})$ and $x_0 \in O_{x_0}$,

$$\Sigma_{x_0, t} := H(t, x_0) \cap A(p, s - 5\zeta, s + 5\zeta)$$

is a surface, which is diffeomorphic to $\Sigma_{x_0} \setminus \bigsqcup_{1 \leq j \leq i} \gamma_j(s, x_0)$ attached with $2i$ disks. Moreover, $H(1, x_0) \cap A(p, s - 5\zeta, s + 5\zeta)$ is also a surface and

$$H(1, x_0) \cap A(p, s - 1.1\zeta, s + 1.1\zeta) = \emptyset.$$

- (5)' For every $i \in \{1, 2, \dots, K\}$, near t_i , $H(t, x_0)$ is a surgery process via pinching the cylinder $\bigcup_{s' \in (s-1.9\zeta, s+1.9\zeta)} \gamma_i(s', x_0)$;
 (6)' For every $t \in [0, 1]$, we have

$$H(t, x_0) \setminus A(p; s - 3.1\zeta, s + 3.1\zeta) = \Phi(x_0) \setminus A(p; s - 3.1\zeta, s + 3.1\zeta);$$

$$\mathcal{H}^2(H(t, x_0)) \leq \mathcal{H}^2(\Phi(x_0)) + C_g \Lambda^2 r^2 / 1.1.$$

Indeed, the general $H : [0, 1] \times O_x \rightarrow \mathcal{S}(M)$ can be defined by

$$H(t, y) := (\varphi_y(H(t, x_0)) \cap A(p; s - 5\zeta, s + 5\zeta)) \cup (\Phi(y) \setminus A(p; s - 5\zeta, s + 5\zeta)),$$

and one can verify that H satisfies (3) - (6) following the properties of $H(\cdot, x_0)$.

Note that in (M, g) , for every $r \in (0, \text{inrad}(M, g)/4)$ and $p \in S$, the annulus $A(p; r, 2r)$ is diffeomorphic to $\mathbb{A}(\mathbf{0}; r, 2r) \subset \mathbb{R}^3$ with a uniform bi-Lipschitz constant $C_g^{(1)}$ depending only on g . Without loss of generality, we may assume that $A(p; r, 2r)$ is $\mathbb{A}(\mathbf{0}; r, 2r)$.

By the isoperimetric inequality in standard spheres, there exists a dimensional constant $C^{(2)} > 0$ such that for any loop γ in γ in a sphere $\partial\mathbb{B}_s$ for any $s > 0$, γ bounds a disk D in $\partial\mathbb{B}_s$ with

$$\mathcal{H}^2(D) \leq C^{(2)} \mathcal{H}^1(\gamma)^2.$$

Therefore, together with Jordan curve theorem, given a finite set of disjoint loops $\{\gamma_i\}$ in $\partial\mathbb{B}_s$, each of them bounds a disk D_i in $\partial\mathbb{B}_s$ such that

$$\sum_i \mathcal{H}^2(D_i) \leq C^{(2)} \sum_i \mathcal{H}^1(\gamma_i)^2 \leq C^{(2)} \left(\sum_i \mathcal{H}^1(\gamma_i) \right)^2.$$

In addition, for every pair i and j , we have a trichotomy: $D_i \cap D_j = \emptyset$, $D_i \subset D_j$ or $D_j \subset D_i$.

Hence, for $\Phi(x_0)$ with $\Phi(x_0) \cap \partial\mathbb{B}_s(p) = \bigsqcup_{i=1}^K \gamma_i$, we can perform the neckpinch surgery starting from the innermost D_i to the outermost D_i in a the neighborhood $\mathbb{A}(\mathbf{0}; s - 1.9\zeta, s + 1.9\zeta)$ through removing cylinders in

$\mathbb{A}(\mathbf{0}; s - 1.1\zeta, s + 1.1\zeta)$ and gluing disks. Since the total area of all the gluing disks is bounded by

$$3 \sum_i \mathcal{H}^2(D_i) + 6 \left(\sum_i \mathcal{H}^1(\gamma_i) \right) \zeta \leq C^{(3)} \Lambda^2 r^2,$$

for some dimensional constant $C^{(3)}$.

In a general annulus $A(p; r, 2r)$, we only need to replace $C^{(3)}$ by a constant $C(C_g^{(1)}, C^{(2)}, C^{(3)})$, and thus, the statements (3), (4)', (5)' and (6)' above follow immediately. \square

Lemma 8.5 (Local Deformation B: Shrinking). *In a 3-dimensional Riemannian manifold (M, g) , let $p \in M$ and $0 < r^- < r^+ < \text{inrad}(M, g)/4$. Then there exists a smooth one parameter family of maps $\{\mathcal{R}_t : M \rightarrow M\}_{t \in [0, 1]}$ such that*

- (1) $\mathcal{R}_t = \text{id}$ when $t \in [0, 1/4]$; $\mathcal{R}_t(B(p, r^-)) = \{p\}$ when $t \in [3/4, 1]$;
- (2) \mathcal{R}_t is a diffeomorphism when $t \in [0, 3/4]$;
- (3) $\mathcal{R}_t|_{M \setminus B(p, r^+)} = \text{id}$ and $\mathcal{R}_t(B(p, r^\pm)) \subset B(p, r^\pm)$ for every $t \in [0, 1]$;
- (4) $\mathcal{R}_t|_{B(p, r^-)}$ is 1-Lipschitz for every $t \in [0, 1]$.

Proof. Let $\psi \in C^\infty(\mathbb{R})$ be a non-decreasing function such that $\psi = 0$ on $\mathbb{R}_{\leq 1/4}$, $\psi = 1$ on $\mathbb{R}_{\geq 3/4}$, $\psi \in (0, 1)$ on $(1/4, 3/4)$. Let $\eta \in C_c^\infty(\mathbb{R}_{< r^+})$ be a decreasing function such that $\eta = 1$ on $\mathbb{R}_{\leq r^-}$.

Working under normal coordinates of (M, g) at p , it is easy to check that the following \mathcal{R}_t satisfies (1)-(4) in the Lemma:

$$\mathcal{R}_t(q) := \begin{cases} q, & \text{if } q \notin B(p, r^+); \\ (1 - \psi(t)\eta(|q|))q, & \text{if } q \in B(p, r^+). \end{cases}$$

\square

Lemma 8.6. *In (M, g) , for $N \in \mathbb{N}^+$, let $\{A(p_j; r_j^-, r_j^+)\}_{j=1}^N$ be a collection of pairwise disjoint annuli in M , where $r_j^+ < \text{inrad}(M, g)/4$. For each $j \in \{1, 2, \dots, N\}$, let $\mathcal{R}_t^j : M \rightarrow M$ be a map from Lemma 8.5 with p_j, r_j^\pm in place of p, r^\pm . For simplicity, we denote*

$$A_j := A(p_j; r_j^-, r_j^+), \quad B_j^\pm := B(p_j; r_j^\pm).$$

Then the following hold.

- (1) *There exists a permutation $\sigma \in \mathfrak{S}_N$ such that for every $1 \leq i < j \leq N$, either $B_{\sigma(i)}^+ \cap B_{\sigma(j)}^+ = \emptyset$, or $B_{\sigma(i)}^+ \subset B_{\sigma(j)}^-$. In this case, we call such a σ admissible.*
- (2) *For every admissible σ , the map*

$$\mathcal{R}_t^{\sigma(N)} \circ \mathcal{R}_t^{\sigma(N-1)} \circ \dots \circ \mathcal{R}_t^{\sigma(2)} \circ \mathcal{R}_t^{\sigma(1)} : M \rightarrow M$$

is 1-Lipschitz on every connect component of $M \setminus \bigcup_{j=1}^N A_j$; and it is independent of the choice of admissible σ .

Proof. Since $\{A_j\}$ is pairwise disjoint, for every $i \neq j$, we have exactly one of the three possibilities:

- $B_{\sigma(i)}^+ \cap B_{\sigma(j)}^+ = \emptyset$;
- $B_{\sigma(i)}^+ \subset B_{\sigma(j)}^-$;
- $B_{\sigma(j)}^+ \subset B_{\sigma(i)}^-$.

Hence, to achieve (1), we can simply choose a permutation σ such that for every $1 \leq i < j \leq N$, $r_{\sigma(i)}^+ \leq r_{\sigma(j)}^+$.

For (2), the 1-Lipschitz property follows from Lemma 8.5 (3) and (4). The independence on the choice of admissible permutations follows from the fact that if $B_i^+ \cap B_j^+ = \emptyset$, then $\text{spt } \mathcal{R}_t^i \cap \text{spt } \mathcal{R}_t^j = \emptyset$. \square

8.3. Combinatorial arguments. To extend the local deformations from the previous subsection to a global deformation along a Simon-Smith family, we rely on the following two useful combinatorial arguments, inspired by [Pit81, Lemma 4.8, Proposition 4.9] and [DLR18b, Lemma 5.7].

Lemma 8.7 (Combinatorial Argument I). *For any positive integers $m \in \mathbb{N}^+$ and $q \in \mathbb{N}^+$, there exists $N = N(m, q) \in \mathbb{N}^+$ with the following property.*

Given any $k \in \mathbb{N}^+$ and a cubical complex $I(m, k)$, suppose that $\{\mathcal{F}_\sigma\}_{\sigma \in I(m, k)}$ is a family of collections of open sets in M assigned to each cell σ of $I(m, k)$, where each collection \mathcal{F}_σ has the form

$$\mathcal{F}_\sigma = (\mathcal{O}_{\sigma,1}, \mathcal{O}_{\sigma,2}, \dots, \mathcal{O}_{\sigma,q})$$

and each $\mathcal{O}_{\sigma,i} = \{U_{\sigma,i}^1, U_{\sigma,i}^2, \dots, U_{\sigma,i}^N\}$ satisfies

$$(8.9) \quad \text{dist}(U_{\sigma,i}^r, U_{\sigma,i}^s) \geq 2 \min\{\text{diam}(U_{\sigma,i}^r), \text{diam}(U_{\sigma,i}^s)\},$$

for all $\sigma \in I(m, k)$, $i \in \{1, 2, \dots, q\}$ and $r, s \in \{1, 2, \dots, N\}$ with $r \neq s$. Then we can extract a family of open sets

$$\{(U_{\sigma,1}, U_{\sigma,2}, \dots, U_{\sigma,q})\}_{\sigma \in I(m, k)}$$

where $U_{\sigma,i} \in \mathcal{O}_{\sigma,i}$ such that

$$\text{dist}(U_{\sigma,i} \cap U_{\tau,j}) > 0,$$

whenever $\sigma, \tau \in I(m, k)$, $(\sigma, i) \neq (\tau, j)$ and σ, τ are faces of a common cell γ of $I(m, k)$.

Moreover, this is also true if the number of collections in each \mathcal{F}_σ is no greater than q .

Proof. The proof is similar to that of [DLR18b, Lemma 5.7].

Let $\{\mathcal{F}_\sigma\}_{\sigma \in I(m, k)}$ be a family as in the lemma with some $N \in \mathbb{N}^+$ to be determined later.

Note that by our assumption (8.9), for each $U \in \mathcal{O}_{\sigma,i}$ and $V^1, \dots, V^l \in \mathcal{O}_{\tau,j}$ with $\sigma \neq \tau$, if $\text{diam}(U) \leq \text{diam}(V^j)$ for all $j \in \{1, 2, \dots, l\}$, then there exists at most one V^j with

$$\text{dist}(U, V^j) = 0.$$

Here is the algorithm to generate the family of open sets:

- (1) We take an open set $U_{\sigma,i}^r \in \mathcal{O}_{\sigma,i}$ with the smallest diameter among all the remaining open sets $U_{\cdot,\cdot}$ and define $U_{\sigma,i} := U_{\sigma,i}^r$, and then remove all the open sets in $\mathcal{O}_{\sigma,i}$.
- (2) For every $\tau \in I(m,k)$ and every $j \in \{1, 2, \dots, q\}$, where $(\sigma, i) \neq (\tau, j)$ and σ, τ are faces of some cell $\gamma \in I(m, k)$, and we remove all $V \in \mathcal{O}_{\tau,j}$ with $\text{dist}(U_{\sigma,i}, V) = 0$.
- (3) Repeat Step (1) and Step (2), until there is no open set left.

In Step (1), we remove all the open sets in $\mathcal{O}_{\sigma,i}$ after defining $U_{\sigma,i}$, so $U_{\sigma,i}$ can be only defined at most once.

In Step (2), by the argument above, before $U_{\tau,j}$ is defined, the size of $\mathcal{O}_{\tau,j}$ will decrease at most by one if some $U_{\sigma,i}$ is defined where $(\sigma, i) \neq (\tau, j)$ and σ, τ are faces of some cell $\gamma \in I(m, k)$. Since there are at most 5^m many such σ , the size of $\mathcal{O}_{\tau,j}$ will decrease at most by $5^m \cdot q$ in total before $U_{\sigma,i}$ is defined.

Therefore, it suffices to choose

$$N := 5^m \cdot q + 1,$$

and then for every $\sigma \in I(m, k)$ and $i \in \{1, \dots, q\}$, the open set $U_{\sigma,i}$ with required properties is defined before our algorithm halts.

For the general case where $|\mathcal{F}_\sigma| = q' < q$, we can replace each \mathcal{F}_σ by

$$(\mathcal{O}_{\sigma,1}, \mathcal{O}_{\sigma,2}, \dots, \mathcal{O}_{\sigma,q'}, \mathcal{O}_{\sigma,q'}, \dots, \mathcal{O}_{\sigma,q'}).$$

The same arguments apply immediately. \square

Lemma 8.8 (Combinatorial argument II). *Given any integers $q, N \in \mathbb{N}^+$ and any $R_0 \in (0, \text{inrad}(M, g)/4)$, if P is a finite set of at most q points in M , then there exists a radius $R \in (5^{-2Nq^2}R_0, 5^{-2N}R_0)$ such that for every $p \in P$,*

$$A(p; R, 5^{2N}R) \cap P = \emptyset.$$

Proof. For $1 \leq l \leq q^2$, we define

$$s_l = 5^{-2Nl}R_0, \quad r_l = 5^{-2N(l-1)}R_0.$$

For every $p \in P$, since $\#P \leq q$, there are at most $q - 1$ many l such that

$$A(p; s_l, r_l) \cap P \neq \emptyset.$$

Since $q(q - 1) < q^2$, by the pigeon hole principle, there is a l_0 such that $R := 5^{-2Nl_0}R_0$ satisfies the requirement of the lemma. \square

8.4. Proof of Proposition 8.1. The proof will proceed in three steps.

Step 1. Set up. Let C_g be as in Lemma 8.3 and $N := N(m, q)$ as in Lemma 8.7. Without loss of generality, we assume that

$$\varepsilon < 5^{-N-1} \min \left\{ \text{injrads}(M, g), \frac{1}{10000 \cdot 3^m q C_g} \right\},$$

and we set $\varepsilon_1 := 5^{-2N-10}\varepsilon$, and $\varepsilon_2 := 5^{-2Nq^2}\varepsilon_1$.

Since \mathcal{S} is a compact set, we can choose $\delta_0 = \delta_0(\mathcal{S}) > 0$ such that for every $\Sigma \in \mathcal{S}^*(M)$, if $\mathbf{F}([\Sigma], [\mathcal{S}]) < \delta_0$, then for every $r \in (\varepsilon_2, \varepsilon_1)$,

$$(8.10) \quad \sup_{p \in M} \left\{ \mathcal{H}^2(\Sigma \cap A(p, r, 2r)) \right\} \leq 100r^2.$$

We select $\delta_1 := \delta_1(\varepsilon_2, \mathcal{S})$ from Lemma 8.2, and set

$$\delta(\varepsilon, q, m, \mathcal{S}) := \min\{\delta_0, \delta_1\}.$$

For a Simon-Smith family Φ from the proposition, by Lemma 8.2, we see that for every $x \in X$ with $\mathbf{g}(\Phi(x)) = 1$, there exists $p_x \in S^3$ such that $B_{\varepsilon_2}(p_x)$ contains a non-trivial loop in $\Phi(x)$.

For each $x \in X$, we set

$$\tilde{P}(x) := \begin{cases} P(x) \cup \{p_x\}, & \text{if } \mathbf{g}(\Phi(x)) = 1 \\ P(x), & \text{if } \mathbf{g}(\Phi(x)) = 0. \end{cases}$$

By Lemma 8.8, since $\#\tilde{P}(x) \leq q$, for each $x \in X$, take $r_x \in (\varepsilon_2, 5^{-2N}\varepsilon_1)$ such that for every pair $p, p' \in \tilde{P}(x)$, we have

$$p' \notin A(p; r_x, 5^{2N}r_x).$$

For each $l \in \{1, 2, \dots, N\}$, we set

$$r_{x,l} := 5^{2(l-1)}r_x.$$

By (8.10), for every $1 \leq l \leq N$, $x \in X$ and $p \in \tilde{P}(x)$ we have

$$(8.11) \quad \mathcal{H}^2(\Phi(x) \cap A(p; r_{x,l}, 2r_{x,l})) \leq 100r_{x,l}^2.$$

and thus, applying Lemma 8.3 to $\Phi(x)$ in $A(p; r_{x,l}, 2r_{x,l})$ with $\Lambda = 100$, we obtain the following data:

- (1) $s_{x,l,p} \in (r_{x,l}, 2r_{x,l})$,
- (2) $\zeta_{x,l,p} \in (0, \min\{100r_{x,l}, s_{x,l} - r_{x,l}, 2r_{x,l} - s_{x,l}\}/5)$,
- (3) a neighborhood $O_{x,l,p} \subset X$ of x ,
- (4) and a Simon-Smith family

$$H_{x,l,p} : [0, 1] \times O_{x,l,p} \rightarrow \mathcal{S}(M).$$

For each $x \in X$, we set a neighborhood of x

$$O_x \subset \bigcap_{l \in \{1, 2, \dots, N\}, p \in \tilde{P}(x)} O_{x,l,p},$$

such that $y \mapsto \Phi(y) \setminus \bigcup_{p \in \tilde{P}(x)} B(p; r_{x,l,p}, 2r_{x,l,p})$ is continuous in the smooth topology for $y \in O_x$.

Note that by the definition of $\tilde{P}(x)$, $\mathfrak{g}(\Phi(x) \setminus \bigcup_{p \in \tilde{P}(x)} B(p; r_{x,1,p}) = 0$ and for every $y \in O_x$, we also have $\mathfrak{g}(\Phi(y) \setminus \bigcup_{p \in \tilde{P}(x)} B(p; r_{x,1,p}) = 0$. Therefore, if $\mathfrak{g}(\Phi(y)) = 1$, then there exists at least one nontrivial loop contained in $\Phi(y) \cap \bigcup_{p \in \tilde{P}(x)} B(p; r_{x,1,p})$.

Step 2. Refinement. Since X is compact, we can take a finite cover from $\{O_x\}_{x \in X}$. Then we refine X so that for every cell σ of X , we can find an $x_\sigma \in X$ satisfying that every cell τ of X is a subset of O_{x_σ} provided that σ and τ are faces of some cell γ of X . Note that for each σ , the number of such τ is no more than 5^m .

By applying Lemma 8.7 with

$$\mathcal{F}_\sigma = (\{A(p; r_{x_\sigma, i}, 2r_{x_\sigma, i})\}_{i=1}^N)_{p \in \tilde{P}(x_\sigma)},$$

we obtain, associated with each σ , a collection of annuli denoted by

$$(A_{\sigma, p})_{p \in \tilde{P}(x_\sigma)}$$

such that whenever σ and τ are faces of some cell γ of X and for any $p \in \tilde{P}(x_\sigma)$ and $p' \in \tilde{P}(x_\tau)$, we have

$$A_{\sigma, p} \cap A_{\tau, p'} = \emptyset,$$

unless $(\sigma, p) = (\tau, p')$.

For each cell σ and $p \in \tilde{P}(x_\sigma)$, for convenience, we denote the data associated with each $A_{\sigma, p}$, constructed from Lemma 8.3 at the end of Step 1, as follows:

- (1) $s_{\sigma, p} \in (r_{\sigma, p}, 2r_{\sigma, p})$,
- (2) $\zeta_{\sigma, p} \in (0, \min\{s_{\sigma, p} - r_{\sigma, p}, 2r_{\sigma, p} - s_{\sigma, p}\}/5)$,
- (3) a neighborhood $O_\sigma := O_{x_\sigma}$,
- (4) and a Simon-Smith family

$$H_{\sigma, p} : [0, 1] \times O_\sigma \rightarrow \mathcal{S}(M).$$

We also denote

$$r_{\sigma, p}^\pm := s_{\sigma, p} \pm \zeta_{\sigma, p}, \quad B_{\sigma, p}^\pm := B_{r_{\sigma, p}^\pm}(p_{\sigma, p}), \quad \hat{A}_{\sigma, p} := \overline{B_{\sigma, p}^+} \setminus \overline{B_{\sigma, p}^-} \subset A_{\sigma, p}.$$

Step 3. Construction of H . After the refinement in the previous step, suppose that X is a cubical subcomplex of $I(m, k')$. For each cell σ of X and for each $x \in X$, we define

$$\mathbf{d}_\sigma(x) := \min \left\{ \frac{2\|x - c_\sigma\|_{\ell_\infty}}{3^{-k'}}, 1 \right\}.$$

to be the normalized l^∞ distant function to σ . Here, c_σ is the center of σ .

For each $t \in [0, 1/2]$ and each $x \in X$, we define $H(x, t)$ using pinching surgeries as described in Lemma 8.3:

For every cell σ of X and every $p \in \tilde{P}(x_\sigma)$,

$$H(t, x) \cap A_{\sigma, p} = H_{\sigma, p}(\min\{8t(1 - \mathbf{d}_\sigma(x)), 1\}, x) \cap A_{\sigma, p}$$

and that $H(t, x) = \Phi(x)$ outside $\bigcup_{\sigma \in X, p \in \tilde{P}(x_\sigma)} A_{\sigma, p}$. The map has the following properties for every $x \in X$.

- The set

$$Z_x := \{\sigma : \mathbf{d}_\sigma(x) < 1\}$$

has the property that every cell σ in Z_x is a face of the cell γ , which is the smallest cell containing x . By Step 2, the corresponding annuli $\{A_{\sigma, p}\}_{\sigma \in Z_x, p \in \tilde{P}(x_\sigma)}$ are pairwise disjoint, and the number is no greater than $3^m \cdot q$. In particular, the map $H(x, t)$ is well-defined.

- By Lemma 8.3 (4) and (5), $t \mapsto \mathbf{g}(H(x, t))$ is non-increasing and $t \mapsto H(x, t)$ is a pinch-off process.
- By (8.11) and Lemma 8.3 (6), we have for every $t \in [0, 1/2]$,

$$\begin{aligned} \mathcal{H}^2(H(t, x)) &\leq \mathcal{H}^2(\Phi(x)) + (3^m \cdot q) \cdot C_g 100^2 \varepsilon_1^2 \\ &\leq \mathcal{H}^2(\Phi(x)) + \varepsilon. \end{aligned}$$

- If $\mathbf{d}_\sigma(x) \leq 3/4$, then for each $p \in \tilde{P}(x_\sigma)$, in $A_{\sigma, p}$, $H(1/2, x) = H_{\sigma, p}(1, x)$, thus by Lemma 8.3 (4), we know that

$$H(1/2, x) \cap \hat{A}_{\sigma, p} = \emptyset.$$

- If $\mathbf{g}(\Phi(x)) = 1$ and $x \in \sigma$, then by the discussion at the end of Step 1, we know that for some $p \in \tilde{P}(x_\sigma)$, $B_{s_{\sigma, p}}(p)$ contains a nontrivial loop of $\Phi(x)$. Thus, the components of $H(1/2, x)$ outside B_σ^+ always has genus 0.

Finally, for each $t \in [1/2, 1]$ and each $x \in X$, we define $H(x, t)$ using shrinking process as described in Lemma 8.5:

For each cell σ of X and $p \in \tilde{P}(x_\sigma)$, we let $\mathcal{R}_{(\sigma, p), t} : S^3 \rightarrow S^3$ be the one-parameter family of shrinking deformation from Lemma 8.5 with $r^- = r_{\sigma, p}^-$ and $r^+ = r_{\sigma, p}^+$.

Then for each $x \in X$, we label $\{(\sigma, p) \mid \sigma \in Z_x, p \in P'(x_\sigma)\}$ as

$$\{(\sigma_i, p_i)\}_{1 \leq i \leq N'}$$

such that for every $1 \leq i < j \leq N'$, either $B_{\sigma_i, p_i}^+ \cap B_{\sigma_j, p_j}^+ = \emptyset$, or $B_{\sigma_i, p_i}^+ \subset B_{\sigma_j, p_j}^-$. The existence of such labeling follows from Lemma 8.6 (1). Then, we denote for simplicity,

$$\hat{\mathcal{R}}_{t, x}^{(i)} := \mathcal{R}_{(\sigma_i, p_i), (2t-1)(1-\mathbf{d}_\sigma(x))}.$$

Note that

- By Lemma 8.5 (1), if $\mathbf{d}_{\sigma_i}(x) \geq 3/4$ or $t \in [1/2, 5/8]$, then $\hat{\mathcal{R}}_{t, x}^{(i)} = \text{id}$; when $\mathbf{d}_{\sigma_i}(x) < 3/4$, by the construction above, $H(1/2, x) \cap \hat{A}_{\sigma_i, p_i} = \emptyset$.
- if $\mathbf{d}_{\sigma_i}(x) \leq 1/4$, then $\hat{\mathcal{R}}_{1, x}^{(i)}(B_{\sigma_i, p_i}^-) \subset \{p_i\}$.

Now for $t \in [1/2, 1]$ and $x \in X$, we define

$$H(t, x) := \hat{\mathcal{R}}_{t, x}^{(N')} \circ \dots \circ \hat{\mathcal{R}}_{t, x}^{(2)} \circ \hat{\mathcal{R}}_{t, x}^{(1)}(H(x, 1/2)).$$

Intuitively, $\{H(t, x)\}_{t \in [1/2, 1]}$ is obtained by shrinking some connected components of $H(1/2, x)$ to points. Therefore, for every $x \in X$, $t \mapsto \mathfrak{g}(H(t, x))$ is non-increasing, and $\{H(t, x)\}_{t \in [1/2, 1]}$ is a pinch-off process. This confirms statement (3) of the proposition.

By Lemma 8.5 (3), (4) and Lemma 8.6 (2), we see that $\mathcal{H}^2(H(t, x))$ is non-increasing for $t \in [1/2, 1]$. In particular, for every $t \in [1/2, 1]$,

$$\mathcal{H}^2(H(t, x)) \leq \mathcal{H}^2(\Phi(x)) + \varepsilon.$$

This confirms statement (2) of the proposition.

Furthermore, if $\mathfrak{g}(H(1/2, x)) = 1$ and $x \in \sigma$, then by the construction above, $H(1, x)$ consists of some connected components of $H(1/2, x)$ outside $\bigcup_{p \in \tilde{P}(x_\sigma)} B_{s_\sigma, p}(p)$, along with a finite set of points. From the final property of $H(1/2, x)$, it follows that $\mathfrak{g}(H(1, x)) = 0$. This verifies statement (1) of the proposition and completes the proof.

8.5. Proof of Theorem 3.6. Now we use Proposition 8.1 and the following local min-max theorem to prove Theorem 3.6.

Theorem 8.9 (Local min-max theorem [MN21, Theorem 6.1]). *Let Σ be a closed, smooth, embedded non-degenerate minimal surface with Morse index k and multiplicity one, in a closed 3-dimensional manifold (M, g) . For every $\beta > 0$, there exists $\varepsilon_0 > 0$ and a smooth family $\{F_v\}_{v \in \overline{\mathbb{B}}^k} \subset \text{Diff}^\infty(M)$ such that*

- (1) $F_0 = \text{Id}$, $F_v = F_v^{-1}$ for all $v \in \overline{\mathbb{B}}^k$;
- (2) $\|F_v - \text{Id}\|_{C^1} < \beta$ for all $v \in \overline{\mathbb{B}}^k$;
- (3) the function

$$A^\Sigma : \overline{\mathbb{B}}^k \rightarrow [0, \infty], v \mapsto \mathcal{H}^2((F_v)_\# \Sigma),$$

is strictly concave;

- (4) for every $T \in \mathcal{Z}_2(M; \mathbb{Z}_2)$ with $\mathcal{F}(T, [\Sigma]) < \varepsilon_0$, we have

$$\max_{v \in \overline{\mathbb{B}}^k} \mathbf{M}((F_v)_\# T) \geq \mathcal{H}^2(\Sigma)$$

with equality only if $[\Sigma] = (F_v)_\# T$ for some $v \in \overline{\mathbb{B}}^k$.

We proceed the proof in three steps.

Step 1. Set up. Since $\mathcal{W}_{L, \leq g_0}(M, g)$ consists of a varifold associated with a non-degenerate multiplicity-one minimal sphere S , let $\{F_v\}_{v \in \overline{\mathbb{B}}^k}$ be the smooth family associated with S given by Theorem 8.9 such that

- (i) For every $T_1, T_2 \in \mathcal{Z}_n(M; \mathbb{Z}_2)$ and $v \in \overline{\mathbb{B}}^k$,

$$\mathbf{F}((F_v)_\#(T_1), (F_v)_\#(T_2)) \leq 2\mathbf{F}((T_1), (T_2)).$$

We can choose smaller $\varepsilon_0 > 0$ such that

(ii) For every $T \in \mathcal{Z}_2(M; \mathbb{Z}_2)$ with $\mathbf{F}(T, [S]) < \varepsilon_0$, the function

$$A^T : \overline{\mathbb{B}^k} \rightarrow [0, \infty], v \mapsto \mathbf{M}((F_v)_\# T),$$

is strictly concave with a unique maximum in $\mathbb{B}_{1/2}^k$.

(iii) There exists $\varepsilon_1 > 0$ such that for every $T \in \mathcal{Z}_2(M; \mathbb{Z}_2)$ with $\mathbf{F}(T, [S]) < \varepsilon_0$,

$$\varepsilon_1 < \min_{v \in \partial \mathbb{B}^k, T} (\mathbf{M}(S) - \mathbf{M}((F_v)_\# S)).$$

Let $\mathcal{S} := \{(F_v)_\#(S)\}_{v \in \overline{\mathbb{B}^k}}$, which is a compact subset of embedding of \mathbb{S}^2 into (M, g) . Let

$$\varepsilon_1 := \min_{v \in \partial \mathbb{B}^k} (\mathbf{M}(S) - \mathbf{M}((F_v)_\# S)) > 0.$$

Let $\delta(\varepsilon_1/10, N_P(\Phi) + 1, m + 1, \mathcal{S})$ be chosen as in Proposition 8.1.

Step 2. Initial Simon-Smith family. It follows from Theorem 3.5 with $r = \min(\varepsilon_0, \delta(\varepsilon_1/10, \mathcal{S}, m)/10)$ that there exists $\eta > 0$ and $\Phi_1 \in \Lambda(\Phi)$ such that

$$\mathbf{M}(\Phi_1(x)) \geq L - 4\eta \implies [\Phi_1(x)] \in \mathbf{B}_r^{\mathbf{F}}([S]).$$

In particular, there exists a Simon-Smith family

$$H_1 : [0, 1] \times X \rightarrow \mathcal{S}^*(M)$$

with $H_1(0, \cdot) = \Phi$ and $H_1(1, \cdot) = \Phi_1$, which is a pinch-off process, because for every $x \in X$, $H_1(\cdot, x)$ is induced by a one-parameter group of diffeomorphisms.

We can refine the cubical subcomplex X so that, each cell σ of X , exactly one of the following conditions holds:

- There exists a point $x_0 \in \sigma$ such that $\mathbf{M}(\Phi_1(x_0)) \geq L - 2\eta$. In this case, for every $x \in \sigma$, $\mathbf{M}(\Phi_1(x)) \geq L - 4\eta$;
- For every $x \in \sigma$, $\mathbf{M}(\Phi_1(x)) < L - 2\eta$.

Let X_0 be the smallest cubical subcomplex of X containing all cells that satisfy the first condition, and let X_1 be the smallest cubical subcomplex containing all cells that satisfy the second condition. Let

$$Z := X_0 \cap X_1,$$

which is also compact.

Clearly for every $z \in Z$,

$$L - 4\eta \leq \mathcal{H}^2(\Phi_1(z)) \leq L - 2\eta.$$

and thus, $\mathbf{F}([\Phi_1(z)], [S]) < r \leq \varepsilon_0$. For every $z \in Z$, let $A^z : \overline{\mathbb{B}^k} \rightarrow [0, \infty)$ be the function

$$A^z(v) := \mathbf{M}((F_v)_\#([\Phi_1(z)])),$$

By (ii) of Step 1, A^z is strictly concave and has a unique maximum $m(z) \in \mathbb{B}_{1/2}^k$. By Proposition 2.5, $[\Phi_1]$ is continuous in the \mathbf{F} -metric, so the function

$$m : Z \rightarrow \mathbb{B}_{1/2}^k$$

is continuous.

It follows from Theorem 8.9 (4) and $\mathbf{M}([\Phi_1(z)]) = \mathcal{H}^2(\Phi_1(z)) < L$, that $m(z) \neq 0$ for every $z \in Z$. Hence, there exists $\alpha > 0$ such that

$$\alpha \leq |m(z)| < 1/2.$$

Consider the one-parameter flow $\{\phi^z(\cdot, t)\}_{t \geq 0} \subset \text{Diff}(\overline{\mathbb{B}^k})$ generated by

$$v \mapsto -(1 - |v|^2)\nabla A^z(v).$$

For every $v \in \overline{\mathbb{B}^k} \setminus m(y)$, $t \mapsto A^z(\phi^z(v, t))$ is decreasing, and the limit $\lim_{t \rightarrow \infty} \phi^z(v, t) \in \partial \overline{\mathbb{B}^k}$. In particular, by (iii) of Step 1, we have

$$A^z(0) - \lim_{t \rightarrow \infty} A^z(\phi^z(v, t)) \geq \varepsilon_1.$$

By the compactness of Z , we can choose $T_0 > 0$ such that for every $z \in Z$, $t \mapsto A^z(\phi^z(v, t))$ is decreasing along $[0, T_0]$ and

$$A^z(0) - A^z(\phi^z(v, T_0)) \geq \varepsilon_1/2.$$

Now, let us consider a cubical complex X' in I^{m+1} , whose underlying space is

$$X_0 \times \{0\} \cup Z \times [0, 1] \cup X_1 \times \{1\}.$$

From the construction, we also know that the map

$$f : X' \rightarrow X, \quad (x, y) \mapsto x$$

is a homotopy equivalence. Note that f is a surjective cubical map, after refining both X and X' .

We can define a Simon-Smith family $\Phi'_2 : X' \rightarrow \mathcal{S}^*(M)$ by

$$\begin{aligned} \Phi'_2(x, 0) &= \Phi_1(x), \quad \forall x \in X_0 \\ \Phi'_2(x, 1) &= \Phi_1(x), \quad \forall x \in X_1 \\ \Phi'_2(x, y) &= F_{\phi^z(0, 3yT_0)}(\Phi'(x)), \quad \forall x \in Z, y \in [0, 1/3] \\ \Phi'_2(x, y) &= F_{\phi^z(0, T_0)}(\Phi'(x)), \quad \forall x \in Z, y \in [1/3, 2/3] \\ \Phi'_2(x, y) &= F_{\phi^z(0, 3(1-y)T_0)}(\Phi'(x)), \quad \forall x \in Z, y \in [2/3, 1]. \end{aligned}$$

Note that we can define two Simon-Smith family $\Phi' : X' \rightarrow \mathcal{S}^*(M)$ and $\Phi'_1 : X' \rightarrow \mathcal{S}^*(M)$ by extending Φ and Φ_1 to the domain X' :

$$\Phi'(x, y) = \Phi(x), \quad \forall (x, y) \in X'$$

and

$$\Phi'_1(x, y) = \Phi_1(x), \quad \forall (x, y) \in X',$$

Similarly, we also have two deformations $H'_1 : [0, 1] \times X' \rightarrow \mathcal{S}^*(M)$ from Φ' to Φ'_1 and $H'_2 : [0, 1] \times X' \rightarrow \mathcal{S}^*(M)$ from Φ'_1 to Φ'_2 :

$$H'_1(t, x, y) = H_1(t, x), \quad \forall t \in [0, 1], (x, y) \in X'$$

and

$$\begin{aligned}
 H'_2(t, x, 0) &= \Phi_1(x), \quad \forall t \in [0, 1], x \in X_0 \\
 H'_2(t, x, 1) &= \Phi_1(x), \quad \forall t \in [0, 1], x \in X_1 \\
 H'_2(t, x, y) &= F_{\phi^z(0, 3tyT_0)}(\Phi'(x)), \quad \forall t \in [0, 1], x \in Z, y \in [0, 1/3] \\
 H'_2(t, x, y) &= F_{\phi^z(0, tT_0)}(\Phi'(x)), \quad \forall t \in [0, 1], x \in Z, y \in [1/3, 2/3] \\
 H'_2(t, x, y) &= F_{\phi^z(0, 3t(1-y)T_0)}(\Phi'(x)), \quad \forall t \in [0, 1], x \in Z, y \in [2/3, 1].
 \end{aligned}$$

For any $i \in 1, 2$ and any $(x, y) \in X'$, $\{H'_i(t, x, y)\}_{t \in [0, 1]}$ is induced by an isotopy, and thus, a pinch-off process. Furthermore, we have

$$(8.12) \quad N_P(\Phi) = N_P(\Phi_1) = N_P(\Phi') = N_P(\Phi'_1) = N_P(\Phi'_2).$$

Furthermore, Φ'_2 has the following properties: For every $(x, y) \in X'$ with $y \geq 1/3$,

$$(8.13) \quad \mathcal{H}^2(\Phi'_2(x, y)) \leq L - 2\eta.$$

For every $(x, y) \in X'$ with $y \leq 2/3$, by (i) of Step 1,

$$(8.14) \quad \mathbf{F}([\Phi'_2(x, y)], [\mathcal{S}]) \leq 2r < \delta(\varepsilon_1/10, m+1, N_P(\Phi'_2) + 1, \mathcal{S}).$$

For every $(x, y) \in X'$ with $y \in [1/3, 2/3]$,

$$(8.15) \quad \mathcal{H}^2(\Phi'_2(x, y)) \leq L - 2\eta - \varepsilon_1/2.$$

Step 3. Interpolation. Let $Y' := \{(x, y) \in X : y \leq 1/3\}$. By (8.14), for every $x' \in Y'$, $\Phi'_2(x') \in \mathcal{S}(M)$, and we can apply Proposition 8.1 to $\Phi'_2|_{Y'}$ and obtain a deformation

$$H'_{3, Y'} : [0, 1] \times Y' \rightarrow \mathcal{S}(M).$$

Consequently, we can define a Simon-Smith family $H'_3 : [0, 1] \times X' \rightarrow \mathcal{S}^*(M)$ as

$$\begin{aligned}
 H'_3(t, x, y) &= H'_3(t, x, y), \quad \forall t \in [0, 1], (x, y) \in X', y \in [0, 1/3] \\
 H'_3(t, x, y) &= H'_3((2-3y)t, x, 1/3), \quad \forall t \in [0, 1], x \in Z, y \in [1/3, 2/3] \\
 H'_3(t, x, y) &= \Phi'_2(x, y), \quad \forall t \in [0, 1], (x, y) \in X', y \in [2/3, 1].
 \end{aligned}$$

We denote $H'_3(1, \cdot, \cdot) : X' \rightarrow \mathcal{S}^*(M)$ by Φ'_3 .

By Proposition 8.1 (1), for every $(x, y) \in X'$ with $y \in [0, 1/3]$,

$$\mathfrak{g}(\Phi'_3(x, y)) = \mathfrak{g}(H'_3(1, x, y)) = 0.$$

By Proposition 8.1 (2) and (8.15), for every $(x, y) \in X'$ with $y \in [1/3, 2/3]$,

$$\mathcal{H}^2(\Phi'_3(x, y)) = \mathcal{H}^2(H'_3((2-3y)t, x, 1/3)) \leq \mathcal{H}^2(\Phi'_2(x, 1/3)) + \varepsilon_1/10 \leq L - 2\eta.$$

By (8.13), for every $(x, y) \in X'$ with $y \in [2/3, 1]$,

$$\mathcal{H}^2(\Phi'_3(x, y)) = \mathcal{H}^2(\Phi'_2(x, y)) \leq L - 2\eta.$$

Therefore, for every $(x, y) \in X'$,

$$(8.16) \quad \mathcal{H}^2(\Phi'_3(x, y)) \geq L - \eta \implies y \in [0, 1/3] \implies \mathfrak{g}(\Phi'_3(x, y)) = 0.$$

Moreover, by Proposition 8.1 (3), for every $(x, y) \in X'$, $\{H'_3(t, x, y)\}_{t \in [0, 1]}$ is a pinch-off process.

Finally, we can set $\Phi' = \Phi'_3$, which satisfies (1) of the theorem. Let $W := M_f$, and consider the deformation map

$$H' : [0, 1] \times X' \rightarrow \mathcal{S}^*(M)$$

by

$$H'(t, x) = \begin{cases} H_1(3t, x) & t \in [0, 1/3] \\ H_2(3t - 1, x) & t \in [1/3, 2/3] \\ H_3(3t - 2, x) & t \in [2/3, 1] \end{cases}.$$

Since $H'(0, x) = \Phi'(x) = \Phi(f(x))$, H' induces a Simon-Smith family $H : W \rightarrow \mathcal{S}^*(M)$ of genus ≤ 1 satisfying (2)(a) and (2)(b) of the theorem.

This completes the proof.

9. PINCH-OFF PROCESS AND MEAN CURVATURE FLOW

In this section, we show that a pinch-off process (Definition 3.7) would possess certain topological properties similar to those of mean curvature flow. These properties will be used in §10.

Given a level set flow $\{M(t)\}_{t \in [0, \infty)}$ in some $(n+1)$ -Riemannian manifold N , consider the complements of the time-slices,

$$W(t) := \{t\} \times (N \setminus M(t)) \subset [0, \infty) \times N,$$

and

$$W[t_1, t_2] := \bigcup_{t \in [t_1, t_2]} W(t) \subset [0, \infty) \times N.$$

In B. White's work [Whi95], he ingeniously applied the avoidance principle to show:

- For any $T > 0$, any loop in $W[0, T]$ is homotopic to some loop in $W[0]$.
- If N is a complete Riemannian manifold with Ricci curvature bounded from below, then for any $T > 0$, the homomorphism

$$H_{n-1}(W(T); \mathbb{Z}) \rightarrow H_{n-1}(W[0, T]; \mathbb{Z})$$

induced by the inclusion map $W(T) \hookrightarrow W[0, T]$ is an isomorphism.

- The rank of $H_1(W(t); \mathbb{Z})$ is non-increasing in t .

In the following, we prove a similar theorem regarding pinch-off processes. Given a compact Riemannian 3-manifold M , and a pinch-off process $\Phi : [0, T] \rightarrow \mathcal{S}(M)$, consider the complements of the time-slices,

$$W(t) := \{t\} \times (M \setminus \Phi(x)) \subset [0, T] \times M,$$

and

$$W[t_1, t_2] := \bigcup_{t \in [t_1, t_2]} W(t) \subset [0, T] \times M.$$

Proposition 9.1. *If $\Phi : [0, T] \rightarrow \mathcal{S}(M)$ is a pinch-off process, we have:*

- (1) *Any loop in $W(T)$ is homotopic in $W[0, T]$ to a loop in $W(0)$.*
- (2) *If a loop is homologically non-trivial in $W(T)$, then it is homotopic in $W[0, T]$ to a homologically non-trivial loop in $W(0)$.*
- (3) *By Definition 2.1, for each $t \in [0, T]$, $W(t)$ can be written as the disjoint union of two open sets $\text{in}(\Phi(t))$ and $\text{out}(\Phi(t))$, such that both of their reduced boundaries are $\Phi(t) \setminus \Phi(t)_{\text{iso}}$, and they both vary continuously in t (as Caccioppoli sets). Then,*

$$\text{rank}(H_1(\text{in}(\Phi(t)); \mathbb{Z})) \text{ and } \text{rank}(H_1(\text{out}(\Phi(t)); \mathbb{Z}))$$

are both non-increasing in t .

Proof. Let γ be a loop in $W(T)$. We will follow the notation in Definition 3.7. For simplicity, we assume that the “singular times” t_1, \dots, t_n are distinct. It would be clear that the following strategy can still be applied when some of the t_i coincide, by treating the neighborhood of each (t_i, p_i) individually.

To prove (1), in order to homotope γ back to some loop in $W(0)$, let us try to construct a homotopy $\{\gamma_t\}_{t \in [0, T]}$ with $\gamma_T = \gamma$, and $\gamma_t \subset W(t)$ for each t , backwardly from time $t = T$ to $t = 0$. When we decrease t from T to 0, if t is not equal to any of the t_i , then by definition, we can verify that $\Phi(t)$ is moving by isotopy, which yields a homotopy of loops. If $t = t_i$ with (t_i, p_i) corresponding to a shrinking process, letting $J = [s_1, s_2]$ be the time interval of the shrinking process, we can slightly perturb the family of loops $\{\gamma_t\}_{t \in (t_i, s_2]}$ such that the family varies C^2 -continuously and $\inf_{t \in (t_i, s_2]} \text{dist}(\gamma_t, p_i) > 0$. It follows immediately $\gamma_{t_i} := \lim_{t \rightarrow t_i} \gamma_t$ is a loop in $W(t)$. Then, reversing the shrinking process in $[s_1, t_i]$ will yield a homotopy of loops. Finally, if $t = t_i$ with (t_i, p_i) corresponding to a surgery process, let $J = [s_1, s_2]$ be the time interval of the surgery process. By a similar perturbation that avoids the singular point and a reversion of the neck-pinch, we obtain a homotopy of loops for $t \in J$. Thus, the desired homotopy $\{\gamma_t\}_{t \in [0, T]}$ can be constructed, and so (1) is true.

To prove (2), considering the homotopy $\{\gamma_t \subset W(t)\}_{t \in [0, T]}$ of loops obtained previously, it suffices to show that if γ_0 is homologically trivial in $W[0]$ then γ_T is homologically trivial in $W[T]$. Suppose γ_0 bounds a 2-chain in $W[0]$. We are going to show that γ_t bounds some 2-chain Γ_t for each t .

When we increase t from 0 to T , if t is not equal to any of the t_i , then as before, near t , $\Phi(\cdot)$ is moving by isotopy, which yields the desired homotopy of 2-chains. If $t = t_i$ with (t_i, p_i) corresponding to a shrinking process, the shrinking might contract some open subsets of Γ_t to a point at $t = t_i$, but as chains, we still have $\partial \Gamma_{t_i} = \gamma_{t_i}$. If $t = t_i$ with (t_i, p_i) corresponding to a surgery process, let $J = [s_1, s_2]$ be the time interval of the surgery process. In J , the surgery process is also induced by an isotopy except at (t_i, p_i) , and we claim that we can still obtain $\{\Gamma_t\}_{t \in J}$ with $\partial \Gamma_t = \gamma_t$.

Indeed, in this case, from our construction of the family $\{\gamma_t\}_{t \in [0, T]}$, we may, by taking the surgery region U to be sufficiently small, assume that

γ_{s_1} avoids U , even though Γ_{s_1} could intersect U . Now, the boundary the cylinder $\Phi(s_1) \cap U$ consists of two loops. Pick one such loop, and denote by D the disc that it bounds in the sphere ∂U . When we intersect the two chain Γ_{s_1} with D , we can assume that the cross section consists of finitely many loops c_1, \dots, c_m , with each c_j bounding a disc d_j within D . By gluing two copies of each d_j to Γ_{s_1} , and slightly opening up a gap between the two d_j , one can obtain a 2-chain bounded by γ_{s_1} that avoids D . By deforming this 2-chain, we can further assume that it avoids the whole ball U . Let us let Γ_{s_1} denote this new 2-chain instead. Now, when we perform surgery for $t \in [s_1, s_2]$, we can homotope Γ_{s_1} across $t \in [s_1, s_2]$ with no issue. Thus, the desired of two chains Γ_t can be constructed (though $\{\Gamma_t\}_{t \in [0, T]}$ is not a continuous family), and so (2) is true.

As for (3), let us just do the case for $\text{in}(\Phi(t))$, as the case for $\text{out}(\Phi(t))$ is similar. Suppose we have loops $\gamma_T^1, \dots, \gamma_T^k$ in $\text{in}(\Phi(T))$ such that the elements they induce in $H_1(\text{in}(\Phi(T)); \mathbb{Z})$ are linearly independent. From the proof of (1), we can construct for each $j = 1, \dots, k$ a homotopy $\{\gamma_t^j \subset \text{in}(\Phi(t))\}_{t \in [0, T]}$ of loops. To prove (3), it suffices to show that $\gamma_0^1, \dots, \gamma_0^k$ are also linear independent in $H_1(\text{in}(\Phi(0)); \mathbb{Z})$. Suppose by contradiction that $\gamma_0^1 + \dots + \gamma_0^k$ bounds some 2-chain $\Sigma_0 \subset \text{in}(\Phi(0))$. Following the proof of (2), we can similarly construct for each t a 2-chain $\Sigma_t \subset \text{in}(\Phi(t))$ bounded by $\gamma_t^1 + \dots + \gamma_t^k$. Putting $t = T$, contradiction arises. Thus (3) is true too. \square

10. TOPOLOGY OF GENUS ONE CAP

The goal of this section is to prove Theorem 3.16.

As explained at the end of §3, it suffices to show that for each genus 1 cap $C \subset \text{dmn}(\Xi)$, the map

$$i_* : H_1(C; \mathbb{Z}_2) \rightarrow H_1(\text{dmn}(\Xi); \mathbb{Z}_2)$$

induced by the inclusion $i : C \hookrightarrow \text{dmn}(\Xi)$ is trivial.

In particular, for any fixed loop $c \subset C$, we aim to show that

$$[i(c)] = 0 \in H_1(\text{dmn}(\Xi); \mathbb{Z}_2)$$

For simplicity, we may just view c as a subset of $\text{dmn}(\Xi)$, we will show

$$(10.1) \quad [c] = 0 \in H_1(\text{dmn}(\Xi); \mathbb{Z}_2).$$

Note that, by the definition of a genus 1 cap, the image

$$(10.2) \quad [\Xi](c) \subset \mathbf{B}_{d_0}^{\mathbf{F}}([T])$$

for some embedded minimal torus T .

The proof of (10.1) will consist of three main steps. First, we need to understand better the topology of the members of $\Xi|_c$. In the second step, using family $\tilde{\Xi}$ obtained in Proposition 3.11, we, in a certain sense, homotope the family $\Xi|_c$ back to some subfamily $\Phi|_{c_0}$ of Φ , for some loop $c_0 \subset Y$, while keeping track of the topology of the members of this homotopy. We will show that one can assume $\Phi|_{c_0}$ to consist entirely Clifford tori (if Ψ is viewed in

\mathbb{S}^3), and to prove (10.1) it suffices to show that c_0 is homologically trivial in Y . In the third step, we prove that c_0 is homologically trivial by studying the family Ψ .

10.1. **The family $\Xi|_c$.** In Proposition 3.11, we obtained a mapping cone \widetilde{W} containing Y and $\text{dmn}(\Xi)$, a map $\widetilde{F} : [0, 1] \times \widetilde{W} \rightarrow \widetilde{W}$, and a Simon-Smith family $\widetilde{\Xi} : \widetilde{W} \rightarrow \mathcal{S}^*(S^3)$ of genus ≤ 1 . Then we can define a map

$$G_1 : [0, 1] \times S^1 \rightarrow \widetilde{W}$$

such that $G_1(1, \cdot)$ parametrizes a loop $c \subset \widetilde{W}$, and for every $t \in [0, 1]$ and every $\theta \in S^1$,

$$G_1(t, \theta) := \widetilde{F}(t, G_1(1, \theta)).$$

By Proposition 3.11, for each $\theta \in S^1$, the family $\{\widetilde{\Xi} \circ G_1(t, \theta)\}_{t \in [0, 1]}$ is a pinch-off process.

To prove Theorem 3.16, we need more information about $\Xi(x)$ for $x \in C$ beyond the fact that C is a genus 1 cap. More precisely, we need that the two regions, inside and outside, enclosed by each surface resemble solid tori.

Proposition 10.1. *In a Riemannian 3-sphere (S^3, g) , let Σ be a smooth, embedded torus. There exists a constant $\tau(\Sigma) > 0$ with the following property.*

Let $S \in \widetilde{\mathcal{S}}(S^3)$. By definition 2.1, let $\Omega, \Omega' \subset S^3$ be the two open subsets of $S^3 \setminus S$, whose reduced boundaries are both $S \setminus S_{\text{iso}}$. Assume that:

- $H_1(\Omega; \mathbb{Z})$ and $H_1(\Omega'; \mathbb{Z})$ are both either 0 or \mathbb{Z} .
- $\mathbf{F}([S], [\Sigma]) < \tau(\Sigma)$.

Then,

- (1) $H_1(\Omega; \mathbb{Z}) \cong H_1(\Omega'; \mathbb{Z}) \cong \mathbb{Z}$.
- (2) Fixing an inside direction for Σ pointing towards $\text{in}(\Sigma)$, there is a unique choice of an inside direction for S such that

$$\text{vol}(\text{in}(S) \Delta \text{in}(\Sigma)), \text{vol}(\text{out}(S) \Delta \text{out}(\Sigma)) < \tau(\Sigma).$$

Here, $\{\text{in}(S), \text{out}(S)\} = \{\Omega, \Omega'\}$ and $\{\text{in}(\Sigma), \text{out}(\Sigma)\}$ is the connected components of $S^3 \setminus \Sigma$.

- (3) Fixing a generator a_0 for $H_1(\text{in}(\Sigma); \mathbb{Z})$, there is a unique generator a_1 for $H_1(\text{in}(S); \mathbb{Z})$ such that there exists a loop $\gamma_{\text{in}} \subset \text{in}(S) \cap \text{in}(\Sigma)$ that induces both a_0 and a_1 . And an analogous statement holds with $\text{out}(\cdot)$ in place of $\text{in}(\cdot)$.

Remark 10.2. The condition “ $H_1(\Omega; \mathbb{Z})$ and $H_1(\Omega'; \mathbb{Z})$ are both either 0 or \mathbb{Z} ” is different from requiring “ $\mathfrak{g}(S) = 0$ or 1”. Indeed, consider this counterexample: Take three 2-spheres, identify all of their north poles as a point, and similarly, identify all of their three south poles as another point.

Proof. First, choose a constant $d_1 > 0$ small enough such that Σ has a tubular neighborhood of width $2d_1$. Fix an inside direction for Σ . For each $t \in (0, d_1)$, the boundary of the t -neighborhood of Σ consists of two smooth

surfaces, S_t and S_{-t} : We assume that S_t lies outside Σ , while S_{-t} inside. We denote $S_0 := \Sigma$. Now, for sufficiently small $\tau(\Sigma)$, from the fact that

$$\mathcal{F}([S], [\Sigma]) \leq \mathbf{F}([S], [\Sigma]) < \tau(\Sigma),$$

item (2) follows.

Fix a small $\delta = \delta(\Sigma) > 0$ to be determined by Σ only. By $\mathbf{F}(|S|, |\Sigma|) < \tau(\Sigma)$ and by assuming $\tau(\Sigma)$ sufficiently small (depending on δ and Σ), we can assert that: By the coarea formula and Sard's theorem, there exists a $t_0 \in (-d_1, -d_1/2)$ such that:

- The intersection $S \cap S_{t_0}$ is transverse and is a finite union of smooth loops.
- These loops are contained in finitely many balls $\{B_{r_i}(q_i)\}_i$ with

$$\sum_i r_i < \delta.$$

Since Σ is a torus, we can fix generating elements

$$a_0 \in H_1(\text{in}(\Sigma); \mathbb{Z}) \cong \mathbb{Z}, \quad b_0 \in H_1(\text{out}(\Sigma); \mathbb{Z}) \cong \mathbb{Z}$$

such that they have linking number 1 in S^3 . Then, by the bullet points above and $\mathcal{F}([S], [\Sigma]) < \tau(\Sigma)$, we can choose a sufficiently small $\delta = \delta(\Sigma)$, and thus $\tau(\Sigma)$, such that there exists a loop $\gamma_{\text{in}} \subset S_{t_0} \cap \text{in}(S)$ which avoids all loops in $S_{t_0} \cap S$ and satisfies $[\gamma_{\text{in}}] = a_0 \in H_1(\text{in}(\Sigma); \mathbb{Z})$. Arguing similarly, we can choose a loop $\gamma_{\text{out}} \subset \text{out}(\Sigma) \cap \text{out}(S)$ such that $[\gamma_{\text{out}}] = b_0 \in H_1(\text{out}(\Sigma); \mathbb{Z})$. Noting the linking number $\text{link}(\gamma_{\text{in}}, \gamma_{\text{out}})$ is 1, we obtain item (1) of the proposition. Now, set

$$a_1 := [\gamma_{\text{in}}] \in H_1(\text{in}(S); \mathbb{Z}), \quad b_1 := [\gamma_{\text{out}}] \in H_1(\text{out}(S); \mathbb{Z})$$

From $\text{link}(\gamma_{\text{in}}, \gamma_{\text{out}}) = 1$, item (3) also follows easily. \square

Recall that by assumption, (S^3, g') has only finitely many embedded minimal tori. For each embedded minimal torus Σ , the above proposition gives a constant $\tau(\Sigma) > 0$.

Without loss of generality, we now assume that the constant $d_0 > 0$, chosen in the proof of Theorem 1.1 (in the paragraph right before §3.2.1) is smaller than $\tau(\Sigma)$ for every embedded minimal torus Σ . Then for each $x \in c \subset \text{dmn}(\Xi)$, we have the following.

- $\mathbf{F}([\Xi(x)], [T]) < \tau(T)$. In particular, $\Xi(x) \in \mathcal{S}(M)$.
- Since the family $\tilde{\Xi} \circ \tilde{F}(\cdot, x)$ is a pinch-off process (by Proposition 3.11), by considering the shape of the members in original family Ψ (see §5.3.2), we know by Proposition 9.1, that $\Xi(x)$ satisfies the assumptions on S in the first bullet point of Proposition 10.1.

As a result, we can apply Proposition 10.1 to $\tilde{\Xi} \circ G_1(1, \theta)$ for each $\theta \in S^1$ (as $G_1(1, \cdot)$ is a parametrization for the loop c). Thus, fixing an inside direction for the torus T , we have:

(1) For each $\theta \in S^1$,

$$(10.3) \quad H_1(\text{in}(\tilde{\Xi} \circ G_1(1, \theta)); \mathbb{Z}) \cong H_1(\text{out}(\tilde{\Xi} \circ G_1(1, \theta)); \mathbb{Z}) \cong \mathbb{Z}.$$

(2) For each θ , there is a unique way to choose an inside direction for $\tilde{\Xi} \circ G_1(1, \theta)$ such that

$$\text{vol}(\text{in}(T) \Delta \text{in}(\tilde{\Xi} \circ G_1(1, \theta))), \text{vol}(\text{out}(T) \Delta \text{out}(\tilde{\Xi} \circ G_1(1, \theta))) < \tau(T),$$

and this choice is continuous in θ .

(3) Let us fix a generator a_0 for $H_1(\text{in}(T); \mathbb{Z})$. For each θ , there is a unique generator $a(\theta)$ of $H_1(\text{in}(\tilde{\Xi} \circ G_1(1, \theta)); \mathbb{Z})$ such that there exists a loop $\gamma_a(\theta) \subset \text{in}(\tilde{\Xi} \circ G_1(1, \theta)) \cap \text{in}(T)$ that induces both a_0 and $a(\theta)$.

(4) Let us fix a generator b_0 for $H_1(\text{out}(T); \mathbb{Z})$. For each θ , there is a unique generator $b(\theta)$ of $H_1(\text{out}(\tilde{\Xi} \circ G_1(1, \theta)); \mathbb{Z})$ such that there exists a loop $\gamma_b(\theta) \subset \text{out}(\tilde{\Xi} \circ G_1(1, \theta)) \cap \text{out}(T)$ that induces both b_0 and $b(\theta)$.

10.2. Relating $\Xi|_c$ to Ψ . Then, using the item (2) above, we can continuously and uniquely choose an inside direction for $\tilde{\Xi} \circ G_1(t, \theta)$ for each $(t, \theta) \in [0, 1] \times S^1$. Moreover, from the fact that $\tilde{\Xi} \circ G_1(\cdot, \theta)$ is a pinch-off process for each θ , we know by Proposition 9.1 that

$$\text{rank}(H_1(\text{in}(\tilde{\Xi} \circ G_1(t, \theta)); \mathbb{Z})), \quad \text{rank}(H_1(\text{out}(\tilde{\Xi} \circ G_1(t, \theta)); \mathbb{Z}))$$

are both non-increasing in t . Thus, together with (10.3) and the description of the family Ψ given in §5.3.2, we obtain:

- For each $(t, \theta) \in [0, 1] \times S^1$,

$$H_1(\text{in}(\tilde{\Xi} \circ G_1(t, \theta)); \mathbb{Z}) \cong H_1(\text{out}(\tilde{\Xi} \circ G_1(t, \theta)); \mathbb{Z}) \cong \mathbb{Z}.$$

(Recall that the first homology, in \mathbb{Z} -coefficients, of an open subset of S^3 has no torsion.)

- The family $\tilde{\Xi} \circ G_1(0, \cdot)$, which can be viewed as the same as $\Psi|_{\tilde{F}(0, c)}$, consists entirely of smooth tori, where $\tilde{F}(0, c)$ is a loop in Y .

Now, we prove a lemma stating that all smooth tori in Ψ can be deformation retracted to Clifford tori.

Lemma 10.3. *If we view Ψ as a Simon-Smith family in the unit 3-sphere \mathbb{S}^3 , and let $Z_0, Z_1 \subset Y$ denote the sets of parameters corresponding to Clifford tori and smooth tori, respectively, then Z_1 can be deformation retracted onto Z_0 .*

Proof. Proposition 5.2 tells us that for each $v \in \mathbb{B}^4$ and $z \in \tilde{\mathcal{C}}$, we have an interval $I_{v, z} \subset (-\pi, \pi)$ such that for each $t \in I_{v, z}$,

$$\Sigma(v, t, z) := \partial\{x \in S^3 : d(v, z)(x) < t\}$$

is a smooth torus. Thus, by shrinking the interval $(-\pi, \pi)$ into the point $\{0\}$, we can deform $I_{v,z}$ into the point $\{0\}$ for each v and z . This gives us a deformation retraction of Z_1 onto a subset Z_2 of Y . Note that

$$Z_2 \subset \{(v, t, z) \in Y : v \in \mathbb{B}^4, t = 0\}.$$

Thus, Z_2 can be deformation retracted onto the set

$$Z_0 = \{(v, t, z) \in Y : v = 0, t = 0\},$$

which parametrizes the set of Clifford tori under Ψ in \mathbb{S}^3 . \square

Thus, by the above lemma, there exists a homotopy

$$G_2 : [0, 1] \times S^1 \rightarrow Y \subset \widetilde{W}$$

such that $G_2(0, \cdot)$ lies in Z_0 and $G_2(1, \cdot) = G_1(0, \cdot)$. Let c_0 denote the loop in Z_0 parametrized by $G_2(0, \cdot)$. By concatenating the two homotopies G_1 and G_2 , we obtain a homotopy

$$G_3 : [0, 1] \times S^1 \rightarrow \widetilde{W}$$

such that:

- $G_3(0, \cdot)$ parametrizes $c_0 \subset Z_0 \subset \widetilde{W}$
- $G_3(1, \cdot)$ parametrizes $c \subset \text{dmn}(\Xi) \subset \widetilde{W}$.
- For each $\theta \in S^1$, the family $\widetilde{\Xi} \circ G_3(\cdot, \theta)$ is a pinch-off process.
- For each $(t, \theta) \in [0, 1] \times S^1$,

$$H_1(\text{in}(\widetilde{\Xi} \circ G_3(t, \theta)); \mathbb{Z}) \cong H_1(\text{out}(\widetilde{\Xi} \circ G_3(t, \theta)); \mathbb{Z}) \cong \mathbb{Z}.$$

In particular, by the first two bullet points and the fact that \widetilde{F} induces a strong deformation retraction of \widetilde{W} onto Y , to prove that $[c] = 0$ in $H_1(\text{dmn}(\Xi); \mathbb{Z}_2)$, it suffices to show that $[c_0] = 0$ in $H_1(Z_0; \mathbb{Z}_2)$.

Recall that $Z_0 \cong \mathbb{R}\mathbb{P}^2 \times \mathbb{R}\mathbb{P}^2$, so $H_1(Z_0; \mathbb{Z}_2) = \mathbb{Z}^2 \times \mathbb{Z}^2$ has 3 non-trivial elements: $(1, 0), (0, 1), (1, 1)$. Note, on one hand, $(1, 0)$ and $(0, 1)$ (but not $(1, 1)$) both correspond to 1-sweepouts under Ψ as we discussed in §5.4. On the other hand, $\Xi|_c$ is not a 1-sweepout as it is close to a single embedded minimal torus in the flat topology by (10.2), and this implies $\Psi|_{c_0}$, which is homotopic to $\Xi|_c$ in the flat topology via $\widetilde{\Xi} \circ G_3$, is not a 1-sweepout either. Thus, $[c_0]$ cannot be $(1, 0)$ or $(0, 1)$.

It suffices rule out the possibility that $[c_0] = (1, 1)$. Let us extend the inside directions we chose for the family $\widetilde{\Xi} \circ G_1$ to the family $\widetilde{\Xi} \circ G_3$. Note that this extension is continuous and unique. As a result, we can “naturally identify” the groups

$$H_1(\text{in}(\widetilde{\Xi} \circ G_3(t, \theta)); \mathbb{Z}), \quad (t, \theta) \in [0, 1] \times S^1,$$

in the following sense. Let $k \in \mathbb{N}$. For each $i, j = 1, \dots, k$, define the closed rectangle

$$R_{i,j} = \left[\frac{i-1}{k}, \frac{i}{k} \right] \times \left[2\pi \frac{j-1}{k}, 2\pi \frac{j}{k} \right] \subset [0, 1] \times S^1.$$

Proposition 10.4. *For sufficiently large k , there exist for each (i, j) loops $\gamma_{i,j}^{\text{in}}$ and $\gamma_{i,j}^{\text{out}}$ with linking number 1 in S^3 such that the following hold: If $R_{m,n}$ is one of the rectangles that intersect $R_{i,j}$ then for every $p \in R_{m,n}$ we have:*

- (1) $\gamma_{i,j}^{\text{in}}$ lies in $\text{in}(\tilde{\Xi} \circ G_3(p))$ and generates $H_1(\text{in}(\tilde{\Xi} \circ G_3(p)); \mathbb{Z})$.
- (2) $\gamma_{i,j}^{\text{out}}$ lies in $\text{out}(\tilde{\Xi} \circ G_3(p))$ and generates $H_1(\text{out}(\tilde{\Xi} \circ G_3(p)); \mathbb{Z})$.
- (3) $\gamma_{i,j}^{\text{in}}$ and $\gamma_{m,n}^{\text{in}}$ are homologous in $\text{in}(\tilde{\Xi} \circ G_3(p))$.
- (4) $\gamma_{i,j}^{\text{out}}$ and $\gamma_{m,n}^{\text{out}}$ are homologous in $\text{out}(\tilde{\Xi} \circ G_3(p))$.

Proof. For each $(t, \theta) \in [0, 1] \times S^1$, since

$$H_1(\text{in}(\tilde{\Xi} \circ G_3(t, \theta)); \mathbb{Z}) \cong H_1(\text{out}(\tilde{\Xi} \circ G_3(t, \theta)); \mathbb{Z}) \cong \mathbb{Z},$$

there exist $\gamma_{(t,\theta)}^{\text{in}} \subset \text{in}(\tilde{\Xi} \circ G_3(t, \theta))$ and $\gamma_{(t,\theta)}^{\text{out}} \subset \text{out}(\tilde{\Xi} \circ G_3(t, \theta))$ generating these two homology groups, respectively. Moreover, from the fact that $\tilde{\Xi} \circ G_3$ is a Simon-Smith family, there exists a neighborhood $O_{(t,\theta)} \subset [0, 1] \times S^1$ of (t, θ) , such that $\gamma_{(t,\theta)}^{\text{in}} \subset \text{in}(\tilde{\Xi} \circ G_3(p))$ and generates

$$H_1(\text{in}(\tilde{\Xi} \circ G_3(p)); \mathbb{Z}),$$

and $\gamma_{(t,\theta)}^{\text{out}} \subset \text{out}(\tilde{\Xi} \circ G_3(p))$ and generates

$$H_1(\text{out}(\tilde{\Xi} \circ G_3(p)); \mathbb{Z}).$$

Clearly, $\{O_{(t,\theta)}\}_{(t,\theta) \in [0,1] \times S^1}$ is an open covering of $[0, 1] \times S^1$. Since $[0, 1] \times S^1$ is compact, we can find a finite covering, denoted by

$$\{O_l\}_{l=1}^t,$$

together with the corresponding loops $\{\gamma_l^{\text{in}}\}$ and $\{\gamma_l^{\text{out}}\}$. Moreover, we can find a sufficiently large k such that for each $i, j = 1, \dots, k$, there exists some l , such that

$$N(R_{i,j}) \subset O_l$$

where $N(R_{i,j})$ is the union of $R_{i,j}$ and all the rectangles adjacent to $R_{i,j}$.

Let us first work on the rectangles $R_{k,1}, \dots, R_{k,k}$.

For each $R_{k,j}$, it contains some point $(1, \theta)$. Using the property (3) and (4) for the family $\tilde{\Xi} \circ G_3(1, \cdot)$, there exists a loop $\gamma_a(\theta)$ that lies in

$$\text{in}(\tilde{\Xi} \circ G_3(1, \theta)) \cap \text{in}(T).$$

induces both a_0 and $a(\theta)$. Suppose that $R_{k,j} \subset O_l$, and then we can choose $\gamma_{k,j}^{\text{in}}$ to be either γ_l^{in} or $-\gamma_l^{\text{in}}$ such that

$$[\gamma_{k,j}^{\text{in}}] = a_0 \in H_1(\text{in}(T); \mathbb{Z}),$$

and thus,

$$[\gamma_{k,j}^{\text{in}}] = a(\theta) \in H_1(\text{in}(\tilde{\Xi} \circ G_3(1, \theta)); \mathbb{Z}).$$

Similarly, we can choose a $\gamma_{k,j}^{\text{out}}$. This confirms statements (1) and (2) of the proposition for (k, j) . Note that the loops $\gamma_{k,j}^{\text{in}}$ and $\gamma_{k,j}^{\text{out}}$ have linking number

1, and statements (3) and (4) of the proposition follows from the fact that all $\gamma_{k,j}^{\text{in}}$ generates $a_0 \in H_1(\text{in}(T); \mathbb{Z})$ and all $\gamma_{k,j}^{\text{out}}$ generates $b_0 \in H_1(\text{out}(T); \mathbb{Z})$.

Using backward induction, suppose that for some i , $\gamma_{i,j}^{\text{in}}$ and $\gamma_{i,j}^{\text{out}}$ have been chosen for all $j = 1, 2, \dots, k$. For each $(i-1, j)$, as before, suppose that $R_{k,i} \subset O_l$, and then we can choose $\gamma_{k,j}^{\text{in}}$ to be either γ_l^{in} or $-\gamma_l^{\text{in}}$ such that

$$[\gamma_{i-1,j}^{\text{in}}] = [\gamma_{i,j}^{\text{in}}] \in H_1(\text{in}(\tilde{\Xi} \circ G_3(i/k, 2\pi j/k)); \mathbb{Z}).$$

We can similarly choose a $\gamma_{k,j}^{\text{out}}$. This confirms statements (1) and (2) of the proposition for $(i-1, j)$. From the construction and the choice of k , that for each rectangle $R_{m,n}$ with $m \geq i-1$, which intersects $R_{i-1,j}$,

$$\text{link}(\gamma_{m,n}^{\text{in}}, \gamma_{k,j}^{\text{out}}) = 1, \quad \text{link}(\gamma_{m,n}^{\text{out}}, \gamma_{k,j}^{\text{in}}) = 1.$$

This confirms statements (3) and (4) of the proposition in this case.

In conclusion, one can find such a large k and the corresponding loops. \square

10.3. The family $\Phi|_{c_0}$. Focusing on the part in $\{0\} \times S^1 \subset [0, 1] \times S^1$ in Proposition 10.4, one obtains a map $L_0 : S^1 \times S^1 \rightarrow S^3$ such that for each $\theta \in S^1$, the loop $L_0(\theta, \cdot) = \gamma_{0,j}^{\text{in}}$ for some j , which lies in $\text{in}(\tilde{\Xi} \circ G_3(0, \theta))$ and generates

$$H_1(\text{in}(\tilde{\Xi} \circ G_3(0, \theta)); \mathbb{Z}) \cong \mathbb{Z},$$

satisfying conclusions (3) and (4) of the previous proposition.

However, by the topological lemma below, this is impossible if $[c_0] = (1, 1)$.

Lemma 10.5. *Let $d : [0, 2\pi] \rightarrow Z_0$ with $d(0) = d(2\pi)$ parametrize a loop representing $(1, 1) \in H_1(Z_0; \mathbb{Z}_2)$. We continuously choose an inward direction for $\Psi \circ d(\theta)$, for each $\theta \in [0, 2\pi]$. Let*

$$L : [0, 2\pi] \times S^1 \rightarrow S^3$$

be a continuous map such that each loop $L(\theta, \cdot)$ lies in $\text{in}(\Psi \circ d(\theta))$ and generates

$$H_1(\text{in}(\Psi \circ d(\theta)); \mathbb{Z}) \cong \mathbb{Z}.$$

Then $[L(0, \cdot)]$ and $[L(2\pi, \cdot)]$, which both are elements in $H_1(\text{in}(\Psi \circ d(0)); \mathbb{Z})$, differ exactly by a sign.

Proof. Let us view Ψ as a Simon-Smith family in the unit 3-sphere \mathbb{S}^3 , so that each $\Psi \circ d(\theta)$ is an unoriented Clifford torus. Since $Z_0 \cong \mathbb{RP}^2 \times \mathbb{RP}^2$, $\pi_1(Z_0) = \mathbb{Z}_2 \times \mathbb{Z}_2$, so we can also have $[d] = (1, 1)$ in $\pi_1(Z_0)$.

Fix a pair $(x, y) \in S^2 \times S^2$, such that the point $[(x, y)]$ in $S^2 \times S^2 / \sim$ with $(x, y) \sim (-x, -y)$ corresponds to the Clifford torus $\Psi \circ d(0)$ equipped with the chosen inside direction (recall the parametrization of the set of oriented Clifford tori described in §5.1). We consider a path \tilde{d} in $S^2 \times S^2$ joining (x, y) to $(-x, -y)$, which gives us the loop d in $Z_0 \cong \mathbb{RP}^2 \times \mathbb{RP}^2$ under the projection map $S^2 \times S^2 \rightarrow \mathbb{RP}^2 \times \mathbb{RP}^2$.

Consider the set

$$(10.4) \quad \{(\Omega, a_0) : \Omega \subset \mathbb{S}^3 \text{ is a solid torus bounded by some Clifford torus, } \\ a_0 \text{ is one of the two generators of } H_1(\Omega; \mathbb{Z}) \cong \mathbb{Z}\}.$$

Evidently, there is a natural topology that can be equipped on this set, such that it is a double cover of the space

$$\{\Omega \subset \mathbb{S}^3 : \Omega \text{ is a solid torus bounded by some Clifford torus}\}.$$

Note that L induces a map L' from $[0, 2\pi]$ into the space (10.4). To prove the lemma, it suffices to show that, while the solid tori corresponding to $L'(0)$ and $L'(2\pi)$ are the same, the first homology classes of the solid torus given by $L'(0)$ and $L'(2\pi)$ differ by a sign.

Following the discussion in §5.1, we know that the space (10.4) is homeomorphic to $G_2^+(\mathbb{R}^4)$, the space of oriented 2-planes in \mathbb{R}^4 . But $G_2^+(\mathbb{R}^4) \cong S^2 \times S^2$. Thus, to prove the lemma, it suffices to show that the two oriented 2-planes corresponding to $(x, y), (-x, -y) \in S^2 \times S^2$ are the same except differing by an orientation. This was already proved in [Nur16, §3.4.3]. \square

In conclusion, $[c_0] \neq (1, 1)$ in $H_1(Y; \mathbb{Z}_2)$, so $[c_0] = (0, 0)$, as desired. This finishes the proof of Theorem 3.16.

REFERENCES

- [ACS18] Lucas Ambrozio, Alessandro Carlotto, and Ben Sharp. Compactness analysis for free boundary minimal hypersurfaces. *Calc. Var. Partial Differential Equations*, 57(1):Paper No. 22, 39, 2018.
- [All72] William K. Allard. On the first variation of a varifold. *Ann. of Math. (2)*, 95:417–491, 1972.
- [Alm62] Frederick Justin Almgren, Jr. The homotopy groups of the integral cycle groups. *Topology*, 1:257–299, 1962.
- [And85] Michael T. Anderson. Curvature estimates for minimal surfaces in 3-manifolds. *Ann. Sci. École Norm. Sup. (4)*, 18(1):89–105, 1985.
- [BCW24] Jacob Bernstein, Letian Chen, and Lu Wang. Existence of monotone morse flow lines of the expander functional, 2024.
- [Bir17] George D. Birkhoff. Dynamical systems with two degrees of freedom. *Trans. Amer. Math. Soc.*, 18(2):199–300, 1917.
- [BK24] Richard H Bamler and Bruce Kleiner. On the multiplicity one conjecture for mean curvature flows of surfaces, 2024.
- [BP02] Victor M. Buchstaber and Taras E. Panov. *Torus actions and their applications in topology and combinatorics*, volume 24 of *University Lecture Series*. American Mathematical Society, Providence, RI, 2002.
- [Bre13] Simon Brendle. Embedded minimal tori in S^3 and the Lawson conjecture. *Acta Math.*, 211(2):177–190, 2013.
- [BW22] Jacob Bernstein and Lu Wang. A mountain-pass theorem for asymptotically conical self-expanders. *Peking Math. J.*, 5(2):213–278, 2022.
- [CDL03] Tobias H. Colding and Camillo De Lellis. The min-max construction of minimal surfaces. In *Surveys in differential geometry, Vol. VIII (Boston, MA, 2002)*, volume 8 of *Surv. Differ. Geom.*, pages 75–107. Int. Press, Somerville, MA, 2003.

- [CFS22] Alessandro Carlotto, Giada Franz, and Mario B. Schulz. Free boundary minimal surfaces with connected boundary and arbitrary genus. *Camb. J. Math.*, 10(4):835–857, 2022.
- [CM04] Tobias H. Colding and William P. Minicozzi, II. The space of embedded minimal surfaces of fixed genus in a 3-manifold. III. Planar domains. *Ann. of Math. (2)*, 160(2):523–572, 2004.
- [CM11] Tobias Holck Colding and William P. Minicozzi, II. *A course in minimal surfaces*, volume 121 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2011.
- [CM12] Tobias H. Colding and William P. Minicozzi, II. Generic mean curvature flow I: generic singularities. *Ann. of Math. (2)*, 175(2):755–833, 2012.
- [CS85] Hyeong In Choi and Richard Schoen. The space of minimal embeddings of a surface into a three-dimensional manifold of positive Ricci curvature. *Invent. Math.*, 81(3):387–394, 1985.
- [CS24] Jingwen Chen and Ao Sun. Mean curvature flow with multiplicity 2 convergence in closed manifolds, 2024.
- [DLP10] Camillo De Lellis and Filippo Pellandini. Genus bounds for minimal surfaces arising from min-max constructions. *J. Reine Angew. Math.*, 644:47–99, 2010.
- [DLR18a] Camillo De Lellis and Jusuf Ramic. Min-max theory for minimal hypersurfaces with boundary. *Ann. Inst. Fourier (Grenoble)*, 68(5):1909–1986, 2018.
- [DLR18b] Camillo De Lellis and Jusuf Ramic. Min-max theory for minimal hypersurfaces with boundary. *Ann. Inst. Fourier (Grenoble)*, 68(5):1909–1986, 2018.
- [Fra21] Giada Franz. Equivariant index bound for min-max free boundary minimal surfaces. *arXiv preprint arXiv:2110.01020*, 2021.
- [GJ86] M. Grüter and J. Jost. On embedded minimal disks in convex bodies. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 3(5):345–390, 1986.
- [Gra89] Matthew A. Grayson. Shortening embedded curves. *Ann. of Math. (2)*, 129(1):71–111, 1989.
- [Hat02] Allen Hatcher. *Algebraic topology*. Cambridge University Press, Cambridge, 2002.
- [HK19] Robert Haslhofer and Daniel Ketover. Minimal 2-spheres in 3-spheres. *Duke Math. J.*, 168(10):1929–1975, 2019.
- [HK23] Robert Haslhofer and Daniel Ketover. Free boundary minimal disks in convex balls, 2023.
- [HO80] David A. Hoffman and Robert Osserman. The geometry of the generalized Gauss map. *Mem. Amer. Math. Soc.*, 28(236):iii+105, 1980.
- [Ket19] Daniel Ketover. Genus bounds for min-max minimal surfaces. *J. Differential Geom.*, 112(3):555–590, 2019.
- [Ko23a] Dongyeong Ko. Existence and morse index of two free boundary embedded geodesics on riemannian 2-disks with convex boundary, 2023.
- [Ko23b] Dongyeong Ko. Morse index bound of simple closed geodesics on 2-spheres and strong morse inequalities, 2023.
- [Law70] H. Blaine Lawson, Jr. Complete minimal surfaces in S^3 . *Ann. of Math. (2)*, 92:335–374, 1970.
- [Li23a] Yangyang Li. Existence of infinitely many minimal hypersurfaces in higher-dimensional closed manifolds with generic metrics. *J. Differential Geom.*, 124(2):381–395, 2023.
- [Li23b] Yangyang Li. An improved Morse index bound of min-max minimal hypersurfaces. *Calc. Var. Partial Differential Equations*, 62(6):Paper No. 179, 32, 2023.
- [LMN18] Yevgeny Liokumovich, Fernando C. Marques, and André Neves. Weyl law for the volume spectrum. *Ann. of Math. (2)*, 187(3):933–961, 2018.

- [LS29] Lazar Lusternik and Lev Schnirelmann. Sur le problème de trois géodésiques fermées sur les surfaces de genre 0. *CR Acad. Sci. Paris*, 189:269–271, 1929.
- [Mar23] Fernando Codá Marques. Personal communication, 2023.
- [MN14] Fernando C. Marques and André Neves. Min-max theory and the Willmore conjecture. *Ann. of Math. (2)*, 179(2):683–782, 2014.
- [MN16] Fernando C. Marques and André Neves. Morse index and multiplicity of min-max minimal hypersurfaces. *Camb. J. Math.*, 4(4):463–511, 2016.
- [MN20] Fernando C. Marques and André Neves. Applications of min-max methods to geometry. In *Geometric analysis*, volume 2263 of *Lecture Notes in Math.*, pages 41–77. Springer, Cham, [2020] ©2020.
- [MN21] Fernando C. Marques and André Neves. Morse index of multiplicity one min-max minimal hypersurfaces. *Adv. Math.*, 378:Paper No. 107527, 58, 2021.
- [Mor96] Marston Morse. *The calculus of variations in the large*, volume 18 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI, 1996. Reprint of the 1932 original.
- [Nur16] Charles Arthur George Nurser. *Low min-max widths of the round three-sphere*. PhD thesis, Imperial College London, 2016.
- [Pit81] Jon T. Pitts. *Existence and regularity of minimal surfaces on Riemannian manifolds*, volume 27 of *Mathematical Notes*. Princeton University Press, Princeton, NJ; University of Tokyo Press, Tokyo, 1981.
- [Poi05] Henri Poincaré. Sur les lignes géodésiques des surfaces convexes. *Trans. Amer. Math. Soc.*, 6(3):237–274, 1905.
- [Sha17] Ben Sharp. Compactness of minimal hypersurfaces with bounded index. *J. Differential Geom.*, 106(2):317–339, 2017.
- [Smi83] Francis R. Smith. On the existence of embedded minimal 2-spheres in the 3-sphere, endowed with an arbitrary metric. *Bulletin of the Australian Mathematical Society*, 28(1):159–160, 1983.
- [SS23] Lorenzo Sarnataro and Douglas Stryker. Optimal regularity for minimizers of the prescribed mean curvature functional over isotopies, 2023.
- [Str84] M. Struwe. On a free boundary problem for minimal surfaces. *Invent. Math.*, 75(3):547–560, 1984.
- [Whi87] B. White. Curvature estimates and compactness theorems in 3-manifolds for surfaces that are stationary for parametric elliptic functionals. *Invent. Math.*, 88(2):243–256, 1987.
- [Whi89] Brian White. Every three-sphere of positive Ricci curvature contains a minimal embedded torus. *Bull. Amer. Math. Soc. (N.S.)*, 21(1):71–75, 1989.
- [Whi91] Brian White. The space of minimal submanifolds for varying Riemannian metrics. *Indiana Univ. Math. J.*, 40(1):161–200, 1991.
- [Whi95] Brian White. The topology of hypersurfaces moving by mean curvature. *Comm. Anal. Geom.*, 3(1-2):317–333, 1995.
- [Whi17] Brian White. On the bumpy metrics theorem for minimal submanifolds. *Amer. J. Math.*, 139(4):1149–1155, 2017.
- [Wil65] T. J. Willmore. Note on embedded surfaces. *An. Şti. Univ. “Al. I. Cuza” Iaşi Sect. I a Mat. (N.S.)*, 11B:493–496, 1965.
- [Wil71] T. J. Willmore. Mean curvature of Riemannian immersions. *J. London Math. Soc. (2)*, 3:307–310, 1971.
- [WZ24] Zhichao Wang and Xin Zhou. Existence of four minimal spheres in s^3 with a bumpy metric, 2024.
- [Yau82] Shing Tung Yau. Problem section. In *Seminar on Differential Geometry*, volume No. 102 of *Ann. of Math. Stud.*, pages 669–706. Princeton Univ. Press, Princeton, NJ, 1982.
- [Zho16] Xin Zhou. On the free boundary min-max geodesics. *Int. Math. Res. Not. IMRN*, (5):1447–1466, 2016.

- [Zho20] Xin Zhou. On the multiplicity one conjecture in min-max theory. *Ann. of Math.* (2), 192(3):767–820, 2020.

SIMONS LAUFER MATHEMATICAL SCIENCES INSTITUTE, 17 GAUSS WAY, BERKELEY, CA 94720

Email address: acpc@uchicago.edu

THE UNIVERSITY OF CHICAGO, DEPARTMENT OF MATHEMATICS, ECKHART HALL, 5734 S UNIVERSITY AVE, CHICAGO, IL, 60637

Email address: yangyangli@uchicago.edu