Constraining matter bounce scenario from scalar-induced vector perturbations

Mian Zhu^{*a*} Chao Chen^{*b,c*,1}

- ^aFaculty of Physics, Astronomy and Applied Computer Science, Jagiellonian University, 30-348 Krakow, Poland
- ^bDepartment of Physics, School of Science, Jiangsu University of Science and Technology, Zhenjiang, 212003, China

^cJockey Club Institute for Advanced Study, The Hong Kong University of Science and Technology, Hong Kong, China

E-mail: mian.zhu@uj.edu.pl, chaochen012@gmail.com

Abstract. Bouncing cosmologies, while offering a compelling alternative to inflationary models, face challenges from the growth of vector perturbations during the contracting phase. While linear vector instabilities can be avoided with specific initial conditions or the absence of vector degrees of freedom, we demonstrate the significant role of secondary vector perturbations generated by non-linear interactions with scalar fluctuations. Our analysis reveals that in a broad class of single-field matter bounce scenarios, these secondary vector perturbations inevitably get unacceptably large amplitudes, provided the curvature fluctuations are consistent with cosmic microwave background observations. This finding underscores the crucial importance of scalar-induced vector perturbations in bouncing cosmology and highlights the need for further investigation into their potential impact on the viability of these models.

¹Corresponding author

Contents

1	Introduction	1
2	Theoretical setup	2
3	Energy density of SIVP	3
4	Conclusion	5
A	Scalar-induced vector perturbations	6
в	Vector Power Spectrum from Numerical Evaluation	11
С	Analysis on the SIVP Power Spectrum	13

1 Introduction

Inflation [1], the standard paradigm of the early-universe cosmology, provides a natural way to explain the formation of large-scale structures (LSS) and the observation of cosmic microwave background (CMB). Nonetheless, inflationary cosmology may suffer from the initial singularity problem [2–5] and the trans-Planckian problem [6–8]. These challenges motivate us to explore alternative early universe scenarios such as the non-singular bouncing cosmology [9–12], where a contraction phase takes place in prior to the expansion phase. While bouncing cosmology offers an intriguing alternative for the early universe, it faces significant challenges. Conceptual issues [13, 14] and its compatibility with CMB observations [15, 16] remain critical concerns. There are also extensive debate surrounding specific problems of bouncing cosmologies [17–36] and proposed solutions [37–50], a comprehensive review of these challenges is available in the reference [51].

In this paper, we highlight another challenge for bouncing cosmology, the overproduction of vector perturbations, a problem overlooked in the community. Early studies [52, 53] demonstrated that linear vector perturbations scale as $S_i(k) \propto a^{-2}$, leading to its growth that can break down the perturbation theory. Resolving this issue typically requires specific model constructions or finely-tuned initial conditions for vector perturbations. For instance, a single-field bouncing scenario lacks vector degrees of freedom, preventing primordial vector fluctuations from vacuum fluctuations.

However, secondary vector perturbations inevitably arise from non-linear interactions with primordial curvature fluctuations ζ . Those fluctuations cannot be arbitrarily fine-tuned, as the power spectrum of curvature fluctuation \mathcal{P}_{ζ} is determined by CMB observations. In Ref. [54], scalar-induced vector perturbations (SIVP) are investigated in specific collapsing universes with theoretical considerations. For the first time, we in this paper connect the power spectrum \mathcal{P}_{ζ} to CMB observations, establishing a lower bound for the energy density of SIVP. Specifically, we work in matter bounce scenario [55], a simple-yet-significant bouncing scenario where nearly scale-invariant curvature fluctuation is generated in a matterdominated contraction phase (i.e., the effective equation-of-state parameter is zero). Our results demonstrate that the energy density of SIVP becomes comparable to the background energy density at the end of the matter contraction phase, provided the contraction is driven by a k-essence scalar field. This significant back-reaction poses a serious challenge to the viability of the matter bounce scenario.

This paper is organized as follows: Section 2 introduces the theoretical framework. Section 3 presents the calculation of the energy density ratio of SIVP to the background. We conclude in Section 4. Technical details are provided in the appendices. Throughout the paper, we set the Planck mass $M_p = 1$. A dot denotes derivative with respect to cosmic time t, and a prime denotes differentiation with respect to τ , unless otherwise specified.

2 Theoretical setup

We work in a spatially-flat FLRW universe

$$ds^{2} = -dt^{2} + a(t)^{2} dx_{i} dx^{i} = a(\tau)^{2} (-d\tau^{2} + dx_{i} dx^{i}) , \qquad (2.1)$$

where $d\tau = dt/a$ is the conformal time. In matter bounce, the scalar factor scales as $a \propto \tau^2$, and can be parameterized as

$$a(\tau) = (\tau/\tau_0)^2 , \quad \tau < \tau_0 < 0 ,$$
 (2.2)

where τ_0 labels the end of the contraction phase. It will also be useful to define a comoving Hubble parameter $\mathcal{H} \equiv a'/a = 2/\tau < 0$. The background energy density is given by the Friedmann's equation, $\rho_{\rm bg}(\tau) = 3H^2 = 12\tau_0^4/\tau^6$. In the framework of k-essence theory [56],

$$S = \int d^4x \sqrt{-g} \left[\frac{R}{2} + K(\phi, X) \right] , \quad X \equiv -\frac{1}{2} \partial_\mu \phi \partial^\mu \phi , \qquad (2.3)$$

the quadratic action for curvature fluctuation ζ is [57–59]

$$S_{\zeta}^{(2)} = \int d\tau d^3 x \frac{z_s^2}{2} \left[\zeta'^2 - c_s^2 (\partial_i \zeta)^2 \right] , \quad z_s^2 = \frac{3a^2}{c_s^2} , \qquad (2.4)$$

where we used the fact that the effective slow-roll parameter $\epsilon \equiv -\dot{H}/H^2 = 3/2$ in the matter contraction phase, and we regarded the sound speed for curvature perturbations, $c_s \equiv K_{,X}/(K_{,X} + 2XK_{,XX})$, as a constant for simplicity. Working in the Fourier space with a canonical mode function $v_k = z_s \zeta_k$, the dynamical equation for curvature perturbations becomes

$$v_k'' + \left(c_s^2 k^2 - \frac{2}{\tau^2}\right) v_k = 0 .$$
(2.5)

Imposing the vacuum initial condition, we get the expression for curvature fluctuations as

$$\zeta_k(\tau) \equiv \frac{v_k(\tau)}{z_s} = \frac{e^{-ikc_s\tau}c_s}{\sqrt{6c_sk}} \left(1 - \frac{i}{c_sk\tau}\right) \left(\frac{\tau_0}{\tau}\right)^2 \,. \tag{2.6}$$

In contrast to the vanilla slow-roll inflation case, the curvature perturbations grow on superhorizon scales $|k\tau| \ll 1$ in the matter contraction phase (see e.g., Ref. [60]). Hence, one needs to evaluate the curvature power spectrum at the end of the contraction phase:

$$\langle \zeta_{\vec{k}} \zeta_{\vec{p}} \rangle (\tau = \tau_0) = (2\pi)^3 \delta(\vec{k} + \vec{p}) |\zeta_{\vec{k}}|^2 = (2\pi)^3 \delta(\vec{k} + \vec{p}) \frac{c_s}{6k} \left(1 + \frac{1}{k^2 c_s^2 \tau_0^2} \right) .$$
(2.7)

From the definition of the scalar power spectrum,

$$\langle \zeta_{\vec{k}} \zeta_{\vec{p}} \rangle = (2\pi)^3 \delta(\vec{k} + \vec{p}) \frac{2\pi^2}{k^3} \mathcal{P}_{\zeta}(k) , \qquad (2.8)$$

we derive,

$$\mathcal{P}_{\zeta}(k,\tau_0) = \frac{k^2 c_s}{12\pi^2} \left(1 + \frac{1}{k^2 c_s^2 \tau_0^2} \right) \simeq \frac{1}{12\pi^2 c_s \tau_0^2} , \qquad (2.9)$$

which is scale-invariant.

In the FLRW universe, the most general perturbed metric, including only vector perturbation, is given by [61]

$$ds^{2} = a^{2}(\tau) \left[-d\tau^{2} - 2G_{i}d\tau dx^{i} + (\delta_{ij} + F_{ij})dx^{i}dx^{j} \right] , \qquad (2.10)$$

where F_{ij} satisfies $F_{ij} = \partial_i F_j + \partial_j F_i$ and $\partial^i F_i = 0$, and G_i is divergent free, $\partial_i G^i = 0$. Since there is no vector degree of freedom in our setup, F_i and G_i should be regarded as secondorder fluctuations induced by the first-order perturbations. We work in the Newtonian gauge where $F_i = 0$, and the metric involving scalar and vector perturbations can be written as

$$ds^{2} = a^{2}(\tau) \left[-e^{2\Phi} d\tau^{2} - 2G_{i} d\tau dx^{i} + e^{-2\Phi} \delta_{ij} dx^{i} dx^{j} \right] , \qquad (2.11)$$

where the scalar perturbation Φ is related to the curvature fluctuation via [62, 63]

$$\zeta = \Phi + \frac{\mathcal{H}}{\mathcal{H}^2 - \mathcal{H}'} (\Phi' + \mathcal{H}\Phi) . \qquad (2.12)$$

The vector power spectrum, defined as

$$\langle G^{\lambda}(\vec{k})G^{s}(\vec{p})\rangle \equiv (2\pi)^{3}\delta(\vec{k}+\vec{p})\delta^{\lambda s}\frac{2\pi^{2}}{k^{3}}\mathcal{P}_{G}(\tau,\vec{k}) .$$
(2.13)

As discussed above, G_i can be sourced by Φ (or equivalently ζ via Eq. (2.12)) through their nonlinear coupling, its power spectrum is computed as

$$\mathcal{P}_{G}(\tau,k) = \int_{\frac{1}{\sqrt{2}}}^{\infty} \mathrm{d}t \int_{-\frac{1}{\sqrt{2}}}^{\frac{1}{\sqrt{2}}} \mathrm{d}s \frac{(1-2s^{2})(2t^{2}-1)(2st+1)^{2}}{4(t^{2}-s^{2})^{2}} \\ \times \mathcal{P}_{\zeta}\left(\frac{k}{\sqrt{2}}(t-s),\tau\right) \mathcal{P}_{\zeta}\left(\frac{k}{\sqrt{2}}(t+s),\tau\right) \left|\mathcal{I}(t,s,z)\right|^{2} , \qquad (2.14)$$

where $z \equiv k\tau$. We summarize the computational details in App. A.

3 Energy density of SIVP

The energy density of SIVP is given by [64]

$$\rho_V(\tau, \vec{x}) = \frac{1}{4a^2} \partial_i G_j(\tau, \vec{x}) \partial^i G^j(\tau, \vec{x}) . \qquad (3.1)$$

From Eq. (3.1), the energy density of SIVP is related to \mathcal{P}_G as

$$\rho_V(\tau) = \frac{1}{2a^2} \int \mathrm{d}k k \mathcal{P}_G(\tau, \vec{k}) \;. \tag{3.2}$$

The back-reaction is represented by the ratio of the energy density of SIVP against the background one at $\tau = \tau_0$:

$$\delta_V \equiv \frac{\rho_V(\tau_0)}{\rho_{\rm bg}(\tau_0)} = \frac{1}{24} \int (k\tau_0) \mathcal{P}_G(k,\tau_0) d(k\tau_0) \ . \tag{3.3}$$

The \mathcal{P}_{ζ} on super-horizon scales is associated to CMB observations:

$$\mathcal{P}_{\zeta}(\vec{k},\tau_0) \simeq \frac{1}{12\pi^2 c_s \tau_0^2} = A_s , \qquad (3.4)$$

where $A_s = 2.1 \times 10^{-9}$ from Planck collaboration [65]. We denote the scales of fluctuations that "cross the horizon" at the beginning/end of contraction phase to be k_{\min} and k_{\max} :

$$k_{\min} = c_s^{-1} |\mathcal{H}(\tau_{\min})| = -2(c_s \tau_{\min})^{-1} , \ k_{\max} = c_s^{-1} |\mathcal{H}(\tau_0)| = -2(c_s \tau_0)^{-1} , \qquad (3.5)$$

where τ_{ini} labels the initial time of matter contraction phase. The modes with $k_{\text{min}} < k < k_{\text{max}}$ becomes super-horizon during the matter contraction phase, and we adopt a minimal curvature power spectrum for super-horizon perturbations

$$\mathcal{P}_{\zeta} = \begin{cases} A_s , & k_{\min} < k < k_{\max} , \\ 0 , & \text{otherwise} . \end{cases}$$
(3.6)

It is possible that modes with $k < k_{\min}$ or $k > k_{\max}$ become super-horizon before or after the matter contraction phase, hence also give a positive contribution to SIVP. Adopting the ansatz in Eq. (3.6) captures the dominant contribution to SIVP from modes entering the horizon during the contraction phase, providing a lower bound sufficient to analyze the instability.

The scale of scale-invariant curvature fluctuation indicated by CMB observations ranges from $k_C/a_{\text{today}} \simeq 10^{-4} \text{ Mpc}^{-1}$ to $k_L/a_{\text{today}} \simeq 1 \text{ Mpc}^{-1}$. To match CMB data, modes with $k = k_C$ must be well within the horizon ($k_C \gg k_{\min}$), while k_L must be super-horizon at $\tau = \tau_0$ ($k_L \leq k_{\max}$). Introducing a dimensionless scaling factor

$$g \equiv \frac{\tau_{\rm ini}}{\tau_0} = \frac{k_{\rm max}}{k_{\rm min}} , \qquad (3.7)$$

and g must be larger than 10^4 .



Figure 1. Left: The SIVP power spectra (2.14) at the end of matter contraction τ_0 as functions of c_s with $g = 10^4$; Right: The SIVP power spectra (2.14) as functions of g with $c_s = 1$.

The resulting vector power spectrum is determined by the dimensionless parameters c_s and g, as τ_0 is fixed by c_s through Eq. (3.4). We numerically evaluate \mathcal{P}_G and present the result in Fig. 1. In App. B, we demonstrate that \mathcal{P}_G scales as $A_s^2 c_s^{-2} g^4 \log g$. Consequently,

$$\delta_V \propto \frac{A_s^2}{c_s^4} g^4 (\log g + \mathcal{O}(1)) , \qquad (3.8)$$

from our numerical results. We further confirm this scaling through analytical estimations presented in App. C. As a comparison, the energy density of linear curvature perturbations scales as g^2 assuming a scale-invariant \mathcal{P}_{ζ} . As a result, the back-reaction problem associated with SIVP can be more severe than that of curvature fluctuations due to its $g^4 \log g$ scaling.

The value of δ_V is presented in Table 1, revealing that δ_V is less than unity only when $c_s \simeq 1$ and $g \simeq 10^4$. In all other cases, $\delta_V > 1$, indicating either a significant back-reaction on the background evolution or a breakdown of perturbation theory. Considering the consistency relation in the context of k-essence theory during matter contraction, we have $r = 24c_s$ [33], where r is the tensor-to-scalar ratio constrained by $r_{0.002} < 0.044$ [66]. This constraint implies $c_s < 0.02$, inevitably leading to an excessively large δ_V . Even if we artificially set $c_s = 1$, the parameter g is significantly larger than 10^4 in reality because $k_{\min} \gg k_C$. Combining these arguments, we conclude that cosmological models where nearly scale-invariant curvature fluctuations on CMB and LSS scales originate from a matter contraction phase governed by a minimally coupled k-essence field are constrained by the overproduction of SIVP, rendering such models invalid.

Mode	parameters	Outcomes	
c_s	g	$\delta_V(au_0)$	
10^{-2}	10^{4}	9.2×10^6	
10^{-1}	10^{4}	9.3×10^2	
1	10^{4}	$9.3 imes 10^{-2}$	
1	$e^2 \times 10^4$	$3.5 imes 10^2$	
1	$e^4 \times 10^4$	1.2×10^6	
1	$e^6 \times 10^4$	4.3×10^{9}	

Table 1. The values of δ_V in Eq. (3.3) for various values of c_s and g.

4 Conclusion

Vector fluctuations play a crucial role in bouncing cosmologies, particularly during the contraction phase. This study, for the first time, combines the concept of secondary vector fluctuations induced by scalar fluctuations with observational constraints on curvature perturbations, revealing an overproduction of these modes in a matter contraction phase governed by a k-essence scalar field. This finding highlights the importance of vector modes in bouncing cosmologies and motivates further investigation into their impact on various bouncing scenarios, including matter bounces with more complex actions [47], Ekpyrotic scenarios [37], and scenarios where the bouncing phase significantly influences the evolution of curvature fluctuations [67].

Our findings motivate further exploration of perturbation theory within the context of bouncing cosmology. Additionally, our results could be revisited by replacing the cut-off of the curvature power spectrum with regularized primordial fluctuations. This approach could, in principle, yield more accurate results. However, this area currently lacks sufficient research. Additionally, the growth of anisotropic shear during the contraction phase, whose energy density scales as $\rho \propto a^{-6}$, warrants investigation into scalar-induced shear, which could provide additional theoretical constraints on bouncing models. The secondary vector fluctuations in bouncing cosmology in the presence of vector field(s) deserves future investigation due to its potential connection with topics such as primordial magnetogenesis [68].

Acknowledgement

M.Z. thanks Yi Wang for discussions which inspired this work, and Xian Gao, Chunshan Lin and George Zahariade for their critical comments. M.Z. also thanks the correspondence with Yi-Fu Cai and Sabino Matarrese for the discussions on several technical problems. We are grateful to Atsuhisa Ota for his contributions during the initial stage of this work. M.Z. was supported by grant No. UMO 2021/42/E/ST9/00260 from the National Science Centre, Poland. C.C. thanks the support from the Jockey Club Institute for Advanced Study at The Hong Kong University of Science and Technology. C.C. is supported by NSFC (Grants No. 12433002).

A Scalar-induced vector perturbations

This appendix derives the second-order vector fluctuations by solving the second-order Einstein equations, following a similar approach used in the study of scalar-induced gravitational waves (SIGW) (see, for example, the review [69]). Throughout this section, superscripts will be used to denote the order of perturbation. For instance, $\rho^{(2)}$ represents the perturbed energy density at second order. The second-order vector $G_i^{(2)}$ is determined by the *ij* components of the Einstein equations,

$$G_i^{(2)j} = T_i^{(2)j} , \ i \neq j .$$
 (A.1)

Although it is possible to derive the secondary vector fluctuations using momentum constraints (as there are no vector degrees of freedom in our specific scenario), the method of computing with Einstein equations will prove useful in future studies involving vector fields. Examples include primordial magnetogenesis [68] and baryon asymmetry [70]. For detailed explanations of both methods and their equivalence, see Refs. [71, 72].

The computation of geometric quantities based on the metric perturbations in Eq. (2.11) is straightforward. We present some useful expressions below (utilizing the identity $\mathcal{H}' = -\mathcal{H}^2/2$, applicable in the matter bounce scenario, and the notation $\partial^2 \equiv \partial_i \partial^i$),

$$\mathcal{R} = -2\frac{e^{2\Phi}}{a^2} \left[(\partial \Phi)^2 - 2\partial^2 \Phi \right] , \ G_0^{(0)0} = -3\frac{\mathcal{H}^2}{a^2} , \qquad (A.2)$$

$$G_i^{(2)j} = \frac{1}{2a^4} \frac{\mathrm{d}}{\mathrm{d}\tau} [a^2 (\partial_i G^j + \partial^j G_i)] - \frac{1}{a^2} \left[2\partial_i \Phi \partial^j \Phi + \delta_i^j (4\Phi \Phi'' + 5\Phi'^2 - (\partial_l \Phi)^2 + 12\mathcal{H}\Phi\Phi') \right] , \qquad (A.3)$$

where \mathcal{R} is the intrinsic curvature defined for a constant-time hypersurfece associated with the metric (2.11).

For the matter sector, we obtain

$$T^{\nu}_{\mu} = (\rho + P)u_{\mu}u^{\nu} + P\delta^{\nu}_{\mu} + \Sigma^{\nu}_{\mu} , \qquad (A.4)$$

where ρ and P are the energy density and the pressure, respectively; u_{μ} is the four-velocity of the observer; $\Sigma_{\mu\nu}$ is the anisotropic stress subject to the conditions $\Sigma_{00} = \Sigma_{0i} = 0$, $\Sigma_{ij} = \Sigma_{ji}$ and $\delta^{ij}\Sigma_{ij} = 0$. For simplicity, we will set the anisotropic stress to be zero, and leave the study of $\Sigma_{ij} \neq 0$ case in the future work. From (A.4) we get

$$T_i^{(2)j} = (\rho^{(0)} + P^{(0)})u_i^{(1)}u^{(1)j} + P^{(2)}\delta_i^j .$$
(A.5)

The four-velocity by definition is normalized according to $u_{\mu}u_{\nu}g^{\mu\nu} = -1$. Along with the definition $u_{\mu} = g_{\mu\nu}u^{\nu}$, one arrive

$$u^{(0)0} = \left(-g_{00}^{(0)}\right)^{-1/2} , \ u^{(1)0} = \frac{1}{2} \left(-g_{00}^{(0)}\right)^{-3/2} g_{00}^{(1)} , \tag{A.6}$$

and accordingly

$$u_i^{(1)} = -\frac{2a}{3\mathcal{H}^2} (\partial_i \Phi' + \mathcal{H} \partial_i \Phi) , \ u^{(1)i} = -\frac{2}{3a\mathcal{H}^2} (\partial_i \Phi' + \mathcal{H} \partial_i \Phi) .$$
(A.7)

The rest quantities are to be determined by the perturbed Einstein equations $G^{(n)}_{\mu\nu} = T^{(n)}_{\mu\nu}$. For instance,

$$\rho^{(0)} = -T_0^{(0)0} = -G_0^{(0)0} = 3\frac{\mathcal{H}^2}{a^2} , \qquad (A.8)$$

$$P^{(2)} = \frac{1}{3}T_i^{(2)i} - \frac{1}{3}(\rho^{(0)} + P^{(0)})u_i^{(1)}u^{(1)i} = \frac{1}{3}G_i^{(2)i} - \frac{1}{3}(\rho^{(0)} + P^{(0)})u_i^{(1)}u^{(1)i} .$$
(A.9)

We organize some useful expressions as below,

$$\rho^{(0)} = 3 \frac{\mathcal{H}^2}{a^2} , \ P^{(0)} = 0 ,$$
(A.10)

$$P^{(2)} = -\frac{1}{9a^2\mathcal{H}^2} \left[4(\partial_i \Phi')^2 + \mathcal{H}^2(\partial_i \Phi)^2 + 8\mathcal{H}\partial_i \Phi \partial^i \Phi' + 45\mathcal{H}^2 \Phi'^2 \right] .$$
(A.11)

From Eqs. (A.5), (A.3), (A.10), and (A.11), we obtain the following equation,

$$\partial_i G^{j\prime} + 2\mathcal{H}\partial_i G^j - \frac{4\partial_i \Phi' \partial_j \Phi' + 8\mathcal{H}\partial_i \Phi' \partial_j \Phi + 10\mathcal{H}^2 \partial_i \Phi \partial_j \Phi}{3\mathcal{H}^2} + (i \longleftrightarrow j) = 0 .$$
(A.12)

Now we are about to derive the equation for SIVP. The curvature fluctuation is related to the Φ through the relation (2.12). While directly converting Φ to ζ using Eq. (2.12) is challenging, we can leverage the fact that curvature fluctuations grow on super-horizon scales. By adopting the ansatz in Eq. (3.6), we focus on the contributions from modes that are super-horizon at the end of the contraction phase. This allows us to utilize the simplified relationship $\Phi = \frac{3}{2}\zeta$, derived by combining Eqs. (2.12) and (2.6) in the super-horizon regime. We will use this relationship to convert Φ to the curvature fluctuation ζ in the following calculations.

In order to extract the transverse vector modes, we define a projection vector [73],

$$\mathcal{V}_{i}^{kl} \equiv \frac{1}{\nabla^{2}} \partial^{l} \mathcal{T}_{i}^{k} \equiv \frac{1}{\nabla^{2}} \partial^{l} \left(\delta_{i}^{k} - \frac{\partial^{k} \partial_{i}}{\nabla^{2}} \right) , \qquad (A.13)$$

which is able to project a term S_{kl} into a transverse vector G_i such that

$$G_i = \mathcal{V}_i^{kl} S_{kl} , \qquad (A.14)$$

and it is helpful to list the following relationships,

$$\mathcal{V}_{i}^{kl}\delta_{kl} = 0 , \quad \mathcal{V}_{i}^{kl}\partial_{k}\partial_{l}\Phi = 0 , \quad \mathcal{V}_{i}^{kl}\partial_{k}G_{l} = 0 , \quad \mathcal{V}_{i}^{kl}\partial_{l}G_{k} = G_{i} .$$
(A.15)

We then get the equation of G_i as

$$G'_i + 2\mathcal{H}G_i = \mathcal{V}_i^{kl}S_{kl} , \qquad (A.16)$$

where the source term is

$$S_{kl} \equiv 15\partial_k \zeta \partial_l \zeta + \frac{6(\partial_k \zeta' \partial_l \zeta + \partial_k \zeta \partial_l \zeta')}{\mathcal{H}} + \frac{6\partial_k \zeta' \partial_l \zeta'}{\mathcal{H}^2} \,. \tag{A.17}$$

As a final step, we move to the Fourier space. We choose a pair of polarization vector $\{e(\hat{k}), \bar{e}(\hat{k})\}$, which are orthogonal to each other and \vec{k} , satisfying:

$$e_i^{\lambda}(\hat{k})e^{\sigma,i}(\hat{k}) = \delta^{\lambda\sigma} , \ e_i^{\lambda}(\hat{k})k^i = 0 , \ \sum_{\lambda} e^{\lambda,i}(\hat{k})e^{\lambda,j}(\hat{k}) = \delta^{ij} - \frac{k^i k^j}{k^2} .$$
(A.18)

The vector perturbation becomes

$$G_i(\tau, \vec{x}) = \sum_{\lambda} \int \frac{\mathrm{d}^3 \vec{k}}{(2\pi)^3} e^{i\vec{k}\cdot\vec{x}} G^{\lambda}(\tau, \vec{k}) e_i^{\lambda}(\hat{k}) , \qquad (A.19)$$

and we have

$$\mathcal{V}_{i}^{ab}S_{ab}(\tau,\vec{x}) = \int \frac{\mathrm{d}^{3}\vec{k}}{(2\pi)^{3}} \frac{ik^{b}}{k^{2}} \left(\delta_{i}^{a} - \frac{k^{a}k_{i}}{k^{2}}\right) e^{i\vec{k}\cdot\vec{x}}S_{ab}(\tau,\vec{k}) ,$$

$$= \sum_{\lambda} \int \frac{\mathrm{d}^{3}\vec{k}}{(2\pi)^{3}} \frac{ik^{b}}{k^{2}} e_{i}^{\lambda}(\hat{k}) e^{\lambda,a}(\hat{k}) e^{i\vec{k}\cdot\vec{x}}S_{ab}(\tau,\vec{k}) , \qquad (A.20)$$

where $S_{ab}(\tau, \vec{k})$ is the Fourier transform of $S_{ab}(\tau, \vec{x})$. Hence the equation for vector mode becomes

$$G^{\lambda\prime}(\tau,\vec{k}) + 2\mathcal{H}G^{\lambda}(\tau,\vec{k}) = S^{\lambda}(\tau,\vec{k}) , \qquad (A.21)$$

where

$$S^{\lambda}(\tau, \vec{k}) = \frac{ik^{m}}{k^{2}} e^{\lambda, n}(\hat{k}) S_{nm}(\tau, \vec{k})$$

= $-\int \frac{\mathrm{d}^{3}\vec{p}}{(2\pi)^{3}} \frac{ik^{m}}{k^{2}} e^{\lambda, n}(\hat{k}) p_{n} p_{m} f(\vec{p}, \vec{k}, \tau) \zeta_{\vec{p}}(\tau_{0}) \zeta_{\vec{k}-\vec{p}}(\tau_{0}) , \qquad (A.22)$

with the evolution kernel,

$$f(\vec{p}, \vec{k}, \tau) = 15\mathcal{T}(p\tau)\mathcal{T}(|\vec{k} - \vec{p}|\tau) + \frac{6}{\mathcal{H}^2}\mathcal{T}'(p\tau)\mathcal{T}'(|\vec{k} - \vec{p}|\tau) + \frac{6}{\mathcal{H}}[\mathcal{T}'(p\tau)\mathcal{T}(|\vec{k} - \vec{p}|\tau) + \mathcal{T}(p\tau)\mathcal{T}'(|\vec{k} - \vec{p}|\tau)] .$$
(A.23)

Note that the prime in (A.23) denotes derivative with respect to the argument instead of τ . Here, \mathcal{T} is the transfer function defined as

$$\zeta_{\vec{k}}(\tau) = \mathcal{T}(k\tau)\zeta_{\vec{k}}(\tau_0) , \ \tau < \tau_0 < 0 .$$
(A.24)

With the help of Eq. (2.6), we have

$$\mathcal{T}(k\tau) = \frac{\tau_0^3}{\tau^3} e^{ic_s k(\tau_0 - \tau)} \frac{i - c_s k\tau}{i - c_s k\tau_0} \,. \tag{A.25}$$

Now we are about to evaluate the two-point correlation function of G^{λ} , defined as

$$\langle G^{\lambda}(\vec{k})G^{s}(\vec{p})\rangle \equiv (2\pi)^{3}\delta(\vec{k}+\vec{p})\delta_{\lambda s}\frac{2\pi^{2}}{k^{3}}\mathcal{P}_{G}(\tau,\vec{k}) .$$
(A.26)

The general solution of G_i is given by

$$G^{\lambda}(\tau, \vec{k}) = \frac{1}{a(\tau)^2} \int^{\tau} a(\tilde{\tau})^2 S^{\lambda}(\tilde{\tau}, \vec{k}) \mathrm{d}\tilde{\tau} .$$
 (A.27)

Applying the bouncing background (2.2) and specifying the integration range, we have

$$G^{\lambda}(\tau, \vec{k}) = \int_{\tau_{\rm ini}}^{\tau} \mathrm{d}\tilde{\tau} \left(\frac{\tilde{\tau}}{\tau}\right)^4 S^{\lambda}(\tilde{\tau}, \vec{k}) , \qquad (A.28)$$

where τ_{ini} is the conformal time at the beginning of the contraction phase. The two-point correlation function of G^{λ} becomes

$$\langle G^{\lambda}(\tau,\vec{k})G^{s}(\tau,\vec{k}')\rangle = \int_{\tau_{\rm ini}}^{\tau} \mathrm{d}\tilde{\tau}_{1} \int_{\tau_{\rm ini}}^{\tau} d\tilde{\tau}_{2} \left(\frac{\tilde{\tau}_{1}}{\tau}\right)^{4} \left(\frac{\tilde{\tau}_{2}}{\tau}\right)^{4} \langle S^{\lambda}(\tilde{\tau}_{1},\vec{k})S^{s}(\tilde{\tau}_{2},\vec{k}')\rangle$$

$$= -\frac{1}{k^{2}k'^{2}} \int_{\tau_{\rm ini}}^{\tau} \mathrm{d}\tilde{\tau}_{1} \int_{\tau_{\rm ini}}^{\tau} \mathrm{d}\tilde{\tau}_{2} \left(\frac{\tilde{\tau}_{1}}{\tau}\right)^{4} \left(\frac{\tilde{\tau}_{2}}{\tau}\right)^{4} \int \frac{\mathrm{d}^{3}\vec{p}}{(2\pi)^{3}} \langle \zeta_{\vec{p}}\zeta_{\vec{k}-\vec{p}}\zeta_{\vec{q}}\zeta_{\vec{k}'-\vec{q}}\rangle$$

$$\times k^{m} p_{m} p_{n} k'^{i} q_{i} q_{j} e^{\lambda,n}(\hat{k}) e^{s,j}(\hat{k'}) f^{*}(\vec{p},\vec{k},\tilde{\tau}_{1}) f(\vec{q},\vec{k'},\tilde{\tau}_{2}) .$$

$$(A.29)$$

Assuming a Gaussian distribution of curvature fluctuation and with the help of Eq. (2.8), the contraction of the four-point correlator is decomposed as

$$\langle \zeta_{\vec{p}}\zeta_{\vec{k}-\vec{p}}\zeta_{\vec{q}}\zeta_{\vec{k}'-\vec{q}}\rangle = \langle \zeta_{\vec{p}}\zeta_{\vec{q}}\rangle\langle \zeta_{\vec{k}-\vec{p}}\zeta_{\vec{k}'-\vec{q}}\rangle + \langle \zeta_{\vec{p}}\zeta_{\vec{k}'-\vec{q}}\rangle\langle \zeta_{\vec{k}-\vec{p}}\zeta_{\vec{q}}\rangle$$

$$= (2\pi)^{6} \frac{2\pi^{2}}{p^{3}} \frac{2\pi^{2}}{|\vec{k}-\vec{p}|^{3}}\delta(\vec{k}+\vec{k}') \left[\delta(\vec{p}+\vec{q})+\delta(\vec{q}+\vec{k}-\vec{p})\right] \mathcal{P}_{\zeta}(\vec{p})\mathcal{P}_{\zeta}(\vec{k}-\vec{p}) .$$

$$(A.30)$$

Adopting the coordinates of two polarization vectors and \vec{k} as

$$e(\hat{k}) = (1,0,0) , \ \bar{e}(\hat{k}) = (0,1,0) , \ \bar{k} = (0,0,k) , \ \bar{p} = p(\sin\theta\cos\psi,\sin\theta\sin\psi,\cos\theta) , \ (A.31)$$

one can simplify the expressions as

$$\langle G^{\lambda}(\tau,\vec{k})G^{s}(\tau,\vec{k}')\rangle = \frac{8\pi^{5}}{k^{3}}\delta^{\lambda s}\delta(\vec{k}+\vec{k}')\int_{0}^{\infty}\mathrm{d}y\int_{|1-y|}^{1+y}\mathrm{d}x\frac{y^{2}}{x^{2}}\left[1-\left(\frac{1+y^{2}-x^{2}}{2y}\right)^{2}\right] \\ \times \left(\frac{1+y^{2}-x^{2}}{2y}\right)^{2}\mathcal{P}_{\zeta}(ky)\mathcal{P}_{\zeta}(kx)\left|\int_{z_{\mathrm{ini}}}^{z}\mathrm{d}\tilde{z}\frac{\tilde{z}^{4}}{z^{4}}f(x,y,\tilde{z})\right|^{2},$$
(A.32)

with the introduction of auxiliary variables

$$x \equiv \frac{|\vec{k} - \vec{p}|}{k} , \ y \equiv \frac{p}{k} ; \ z \equiv k\tau < 0 , \ z_0 \equiv k\tau_0 < 0 .$$
 (A.33)

Following the convention in the study of induced gravitational waves, we further define

$$s = \frac{y - x}{\sqrt{2}}, \ t = \frac{y + x}{\sqrt{2}},$$
 (A.34)

and the total vector power spectrum is calculated as

$$\mathcal{P}_{G}(\tau, \vec{k}) = \int_{\frac{1}{\sqrt{2}}}^{\infty} dt \int_{-\frac{1}{\sqrt{2}}}^{\frac{1}{\sqrt{2}}} ds \frac{(1-2s^{2})(2t^{2}-1)(2st+1)^{2}}{4(t^{2}-s^{2})^{2}} \\ \times \mathcal{P}_{\zeta}\left(\frac{k}{\sqrt{2}}(t-s), \tau\right) \mathcal{P}_{\zeta}\left(\frac{k}{\sqrt{2}}(t+s), \tau\right) \left|\mathcal{I}(s, t, z)\right|^{2} , \qquad (A.35)$$

where

$$\mathcal{I}(s,t,z) = \int_{z_{\rm ini}}^{z} \mathrm{d}\tilde{z} \frac{\tilde{z}^4}{z^4} f(s,t,\tilde{z}) , \qquad (A.36)$$

is the time integral, z_{ini} corresponds to the value of z at far past, i.e., $z_{ini} = k\tau_{ini}$. Explicitly, we have

$$f(s,t,\tilde{z}) \equiv \frac{3z_0^6 e^{-ic_s(\sqrt{2}t\tilde{z}-2z_0)}}{(t^2-s^2)^3 \tilde{z}^6 (c_s z_0-i)^2} \Big[2\sqrt{2}ic_s t\tilde{z}(14-c_s^2 \tilde{z}^2(t^2-s^2)) + 28-2c_s^2(9t^2-5s^2)\tilde{z}^2 + c_s^4 \tilde{z}^4(t^2-s^2)^2 \Big] .$$
(A.37)

The time integral (A.36) can be integrated out analytically ¹,

$$\mathcal{F}(\tilde{z}) \equiv \int \mathrm{d}\tilde{z} \frac{\tilde{z}^4}{z^4} f(s, t, \tilde{z})$$

= $\frac{3z_0^6 e^{-i(\sqrt{2}c_s t \tilde{z} - 2z_0)}}{\sqrt{2}\tilde{z}t^3 z^4 (t^2 - s^2)^3 (1 + ic_s z_0)^2} \Big[-i\tilde{z}^3 c_s^3 t^2 (t^2 - s^2)^2 + ic_s \tilde{z} (s^4 - 14s^2 t^2 + 21t^4) - \sqrt{2}c_s^2 t \tilde{z}^2 (s^4 - 4s^2 t^2 + 3t^4) + 28\sqrt{2}t^3 \Big].$ (A.38)

Notice that the specific time scale τ_0 appears as we set a reference time scale τ_0 in the definition of transfer function (A.24). Introducing $g \equiv z_{\text{ini}}/z_0$ to label the duration of contraction phase and an auxiliary variable $u \equiv c_s z_0$, we have

$$|\mathcal{I}(z_0)|^2 = \frac{9u^2(1+u^2)^{-2}}{2c_s^2 t^6 (t^2-s^2)^6} \left[\mathcal{I}_1 + \mathcal{I}_2 \cos\left(\sqrt{2}(1-g)tu\right) + \mathcal{I}_3 \sin\left(\sqrt{2}(1-g)tu\right) \right] , \quad (A.39)$$

¹This is a fact from the momentum constraint equation: in the Fourier domain, this Poisson equation simply has a schematic form $G^{\lambda}(k,\tau) = k^2 \mathcal{N}^{\lambda}(\vec{k},\tau)$, in which the right-hand side is evaluated at the specific time τ , so no time integral shall appear in the final result. The triviality of time integral indicates that there's essentially no dynamical vector degree of freedoms, as expected in our scalar-tensor setup.

where

$$\mathcal{I}_{1} = (g^{4} + 1) (t^{2} - s^{2})^{4} t^{4} u^{6} + 8 (g^{2} + 1) (2s^{2} - 3t^{2}) (s^{2} - t^{2})^{2} t^{4} u^{4} + 2 (s^{8} - 28s^{6}t^{2} + 126s^{4}t^{4} - 140s^{2}t^{6} + 105t^{8}) u^{2} + 1568 (1 + g^{-2}) t^{6} ,$$
(A.40)

$$\mathcal{I}_{2} = 2g^{2}t^{2}u^{4} \left(s^{2} - t^{2}\right)^{2} \left[s^{4} - t^{2}u^{2} \left(s^{2} - t^{2}\right)^{2} - 14s^{2}t^{2} + 21t^{4}\right] - 4gt^{2}u^{2} \left(s^{4} - 4s^{2}t^{2} + 3t^{4}\right) \left[u^{2} \left(s^{4} - 4s^{2}t^{2} + 3t^{4}\right) - 28t^{2}\right] + 2u^{2} \left(s^{4} - 14s^{2}t^{2} + 21t^{4}\right) \left[-s^{4} + t^{2}u^{2} \left(t^{2} - s^{2}\right)^{2} + 14s^{2}t^{2} - 21t^{4}\right] + 112g^{-1}t^{4} \left[u^{2} \left(s^{4} - 4s^{2}t^{2} + 3t^{4}\right) - 28t^{2}\right], \qquad (A.41)$$

$$\mathcal{I}_{3} = 2\sqrt{2}g^{2}t^{3}u^{3}(s^{2} - t^{2})^{2} \left[u^{2}\left(s^{4} - 4s^{2}t^{2} + 3t^{4}\right) - 28t^{2}\right] - 2\sqrt{2}gtu^{3}\left(s^{4} - 4s^{2}t^{2} + 3t^{4}\right) \left[-s^{4} + t^{2}u^{2}\left(t^{2} - s^{2}\right)^{2} + 14s^{2}t^{2} - 21t^{4}\right] - 2\sqrt{2}tu\left(s^{4} - 14s^{2}t^{2} + 21t^{4}\right) \left[u^{2}\left(s^{4} - 4s^{2}t^{2} + 3t^{4}\right) - 28t^{2}\right] + 56\sqrt{2}g^{-1}tu\left[t^{4}u^{2}\left(s^{2} - t^{2}\right)^{2} - t^{2}\left(s^{4} - 14s^{2}t^{2} + 21t^{4}\right)\right].$$
(A.42)

B Vector Power Spectrum from Numerical Evaluation

We're interested in the energy density of the induced vector perturbation,

$$\rho_V(\vec{x},\tau) = \frac{1}{4a^2} \partial_i G_j(\vec{x},\tau) \partial^i G^j(\vec{x},\tau) , \qquad (B.1)$$

which is related to the vector power spectrum as

$$\rho_V(\tau) = 2 \times \frac{1}{4a^2} \int k \mathcal{P}_G(\tau, \vec{k}) dk = \frac{1}{2a^2} \int k \mathcal{P}_G(\tau, \vec{k}) dk .$$
(B.2)

The factor of 2 arises from the two polarizations of vector perturbations. The vector energy density, ρ_V , is directly determined by the vector power spectrum, \mathcal{P}_G . Therefore, we will focus on evaluating the vector power spectrum numerically in the following.

First, Fig. 2 demonstrates that the oscillatory terms in Eq. (A.39), namely the \mathcal{I}_2 and \mathcal{I}_3 terms, contribute significantly on larger scales. Conversely, on scales $k \simeq k \max$, the value of Eq. (A.39) is primarily determined by the non-oscillatory term \mathcal{I}_1 . In both cases, the power spectrum rapidly diminishes on scales $k \ge k \max$ due to the cutoff of the curvature power spectrum in Equation (3.4). On scales $k < k_{\max}$, \mathcal{P}_G exhibits a strong blue tilt, which can be parameterized as

$$\mathcal{P}_G(\tau_0, k) = \mathcal{A}_G\left(\frac{k}{k_{\max}}\right)^{n_G} , \ k < k_{\max} , \qquad (B.3)$$

from which we obtain the ratio of ρ_V versus the background energy density value $\rho_{\rm bg}$,

$$\delta_V \equiv \frac{\rho_V(\tau_0)}{\rho_{\rm bg}(\tau_0)} = \frac{1}{24} \int (k\tau_0) \mathcal{P}_G(k,\tau_0) \mathrm{d}(k\tau_0) , \qquad (B.4)$$



Figure 2. Comparison of the vector power spectrum \mathcal{P}_G calculated using Eq. (A.39) (including oscillatory terms) with \mathcal{P}_G evaluated using only the non-oscillatory term \mathcal{I}_1 . The parameters used are $c_s = 1$ and $g = 10^4$.

as

$$\delta_V(\tau_0) = \frac{1}{24} \int (k\tau_0) \mathcal{P}_G(k,\tau_0) \mathrm{d}(k\tau_0)$$

$$< \frac{1}{6c_s^2} \int_0^1 \mathcal{A}_G\left(\frac{k}{k_{\max}}\right)^{n_G+1} \mathrm{d}\left(\frac{k}{k_{\max}}\right) = \frac{\mathcal{A}_G}{6c_s^2(n_G+2)} . \tag{B.5}$$

As Fig. 2 illustrates, the non-oscillatory terms primarily determine the amplitude \mathcal{A}_G of the vector power spectrum, while the oscillatory terms are responsible for its spectral index n_G . However, neglecting the oscillatory terms leads to a difference in the calculated n_G by a factor of order unity. For example, with $c_s = 1$ and $g = 10^4$, we find $n_G = 8.2$ when excluding oscillatory terms and $n_G = 6.0$ when using the full expression in Equation (A.39). This discrepancy can be understood as follows: When $k \simeq k_{\max}$, we have $u \simeq -2$ and $g \gg 1$, making \mathcal{I}_1 the dominant term since $\mathcal{I}_1 = \mathcal{O}(g^4)$ while \mathcal{I}_2 and \mathcal{I}_3 are of order $\mathcal{O}(g^2)$. Conversely, on larger scales where $k \simeq k$ min, we have $u \simeq -2g^{-1}$, resulting in $\mathcal{I}_1 = \mathcal{O}(g^0)$ and \mathcal{I}_2 , $\mathcal{I}_3 = \mathcal{O}(g^{-1})$. In this case, the oscillatory terms become significant. For simplicity, we will proceed with the numerical calculations by evaluating only the contributions from the non-oscillatory term. As we will demonstrate later, the resulting δ_V is significantly larger than unity. Therefore, any potential loss of a factor of order unity in this numerical approximation will not significantly impact our conclusions.

Figure 1 displays the resulting vector power spectrum $\mathcal{P}_G(\tau_0, k)$ for various values of c_s and g. The figure reveals that the amplitude \mathcal{A}_G is proportional to $c_s^{-2}g^4$. More precisely, as we will demonstrate below, \mathcal{A}_G exhibits the following behavior for $g \gg 1$:

$$\mathcal{A}_G \simeq \frac{A_s^2}{c_s^2} g^4(\mathcal{C}_1 \log g + \mathcal{C}_2) + \mathcal{O}(g^3) .$$
 (B.6)

The value of \mathcal{A}_G is shown in Fig. 3. A direct fit of the numerical result gives $\mathcal{C}_1 = 15.45$ and $\mathcal{C}_2 = -14.20$. App. C provides a semi-analytical explanation for Eq. (B.6).

Now we conclude the vector power spectrum for different model parameters in Table 2. The $\log g$ dependence in Eq. (B.6) is implicitly reflected in the running of the spectral indices.



Figure 3. The value of \mathcal{A}_G as a function of g with $c_s = 1$ fixed.

Mode	l parameters	Outcomes			
c_s	g	\mathcal{A}_G	n_G	$\delta_V(au_0)$	
10^{-2}	10^{4}	$5.6 imes 10^4$	8.1	9.2×10^6	
10^{-1}	10^{4}	$5.7 imes 10^2$	8.2	$9.3 imes 10^2$	
1	10^{4}	5.7×10^{0}	8.2	9.3×10^{-2}	
1	$e^2 \times 10^4$	2.1×10^4	8.0	3.5×10^2	
1	$e^4 \times 10^4$	7.4×10^7	8.1	1.2×10^6	
1	$e^6 \times 10^4$	2.6×10^{11}	8.0	4.3×10^9	

Table 2. Vector power spectra (B.3) and the values of δ_V for various values of c_s and g.

As argued in Sec. 3, $g \gg 10^4$, making the log g term in Eq. (B.6) dominant. Therefore, we arrive at the scaling,

$$\mathcal{A}_G \propto \frac{A_s^2}{c_s^2} g^4 \log g \ , \ \delta_V \propto \frac{A_s^2}{c_s^4 (n_G + 1)} g^4 \log g \ . \tag{B.7}$$

As a comparison, the energy density of linear curvature perturbations is estimated as [64]

$$\rho_{\zeta} \equiv -\frac{1}{2} \mathcal{R} \to \rho_{\zeta} \propto \int_{k_{\min}}^{k_{\max}} \mathrm{d}\ln k \frac{k^2}{a^2} \mathcal{P}_{\zeta} , \qquad (B.8)$$

using Eq. (A.2). Notice that ρ_{ζ} scales as g^2 assuming a scale-invariant curvature power spectrum \mathcal{P}_{ζ} . Thus, the energy density of SIVP exhibits a faster growth rate with respect to g, leading to a more severe back-reaction problem during the contraction phase.

C Analysis on the SIVP Power Spectrum

In this appendix, we conduct a semi-analytical study on the vector power spectrum. Using the identity

$$|a^{2}| - 2|ab| + |b^{2}| \le |a - b|^{2} \le |a^{2}| + 2|ab| + |b^{2}|, \qquad (C.1)$$

we get

$$\mathcal{J}_1 - \mathcal{J}_2 + \mathcal{J}_3 \le |\mathcal{F}(z_0) - \mathcal{F}(gz_0)|^2 \le \mathcal{J}_1 + \mathcal{J}_2 + \mathcal{J}_3 , \qquad (C.2)$$



Figure 4. The integration range of momentum integrals. We take g = 4 here for illustrative purposes.

with

$$\mathcal{J}_1 \equiv |\mathcal{F}(z_0)^2| , \ \mathcal{J}_2 \equiv 2|\mathcal{F}(z_0)\mathcal{F}(gz_0)| , \ \mathcal{J}_3 \equiv |\mathcal{F}(gz_0)^2| .$$
(C.3)

Our primary interest lies in determining the value of \mathcal{A}_G , which corresponds to the value of $\mathcal{P}_G(k, \tau_0)$ at k = kmax. For the remainder of this discussion, we will fix k to be equal to kmax. The integration range relevant to the power spectrum is defined by the following identity:

$$k_{\min} \le k_1, k_2 \le k_{\max} ; |k_1 - k_2| \le k_{\max} , k_1 + k_2 \ge k_{\max} ,$$
 (C.4)

which we depict in Fig. 4. In terms of variable s and t, it becomes

$$\int_{\frac{g+1}{\sqrt{2g}}}^{\sqrt{2}} \mathrm{d}t \int_{t-\sqrt{2}}^{\sqrt{2}-t} \mathrm{d}s + \int_{\frac{1}{\sqrt{2g}}}^{\frac{g+1}{\sqrt{2g}}} \mathrm{d}t \int_{\frac{\sqrt{2}}{g}-t}^{t-\frac{\sqrt{2}}{g}} \mathrm{d}s \ . \tag{C.5}$$

One can straightforwardly calculate the integrations. However, the resulting formulae is tediously long and we will use a simpler strategy. We observe that contributions to the integrals mainly comes from the boundary near the points $t = \frac{1}{\sqrt{2}}$, $s = \pm \frac{1}{\sqrt{2}}$, where we can expand the integrand as a series of $(t - \frac{1}{\sqrt{2}})$, which greatly simplify the calculations. Further, we will expand the results around $g \to \infty$, which gives

$$\int \mathrm{d}s \mathrm{d}t \frac{(1-2s^2)(2t^2-1)(2st+1)^2}{4(t^2-s^2)^2} \mathcal{J}_1 \simeq \frac{27g^4 z_0^2 (49+c_s^2 z_0^2)}{56(1+c_s^2 z_0^2)} , \qquad (C.6)$$

$$\int \mathrm{d}s \mathrm{d}t \frac{(1-2s^2)(2t^2-1)(2st+1)^2}{4(t^2-s^2)^2} \mathcal{J}_3 \simeq \frac{3g^4 c_s^2 z_0^4 (45-84c_s^2 z_0^2+140c_s^4 z_0^4 \ln g)}{280(1+c_s^2 z_0^2)^2} , \qquad (C.7)$$

up to $\mathcal{O}(g^3)$ contributions. We are interested in the value of \mathcal{P}_G at $k = k_{\text{max}}$, that is $z_0 c_s = -2$ and

$$\int ds dt \frac{(1-2s^2)(2t^2-1)(2st+1)^2}{4(t^2-s^2)^2} \mathcal{P}_{\zeta}^2 \mathcal{J}_1(k=k_{\max}) \simeq \frac{1431g^4 A_s^2}{350c_s^2} , \qquad (C.8)$$

$$\int \mathrm{d}s \mathrm{d}t \frac{(1-2s^2)(2t^2-1)(2st+1)^2}{4(t^2-s^2)^2} \mathcal{P}_{\zeta}^2 \mathcal{J}_3(k=k_{\mathrm{max}}) \simeq \frac{6g^4 A_s^2}{875c_s^2} (2240\ln g - 291) \ . \tag{C.9}$$

The momentum integral for \mathcal{J}_2 is a bit tricky, since it contains absolute values. For our purpose, we notice that near the pole, the real part of $\mathcal{F}(z_0)$ is always no less than its imaginary part. Thus we take $|\mathcal{F}(z_0)| \leq \sqrt{2} \Re(\mathcal{F}(z_0))$. On the other hand, we have $\mathcal{F}(gz_0)$ to be governed by its g^2 term in the $g \to \infty$ limit. Thus we arrive at

$$\int \mathrm{d}s \mathrm{d}t \frac{(1-2s^2)(2t^2-1)(2st+1)^2}{4(t^2-s^2)^2} \mathcal{P}_{\zeta}^2 \mathcal{J}_2(k=k_{\mathrm{max}}) \simeq \frac{4032g^4 A_s^2}{125c_s^2} \ . \tag{C.10}$$

Combining all pieces, we get

$$15.36 \log g - 30.16 \le \mathcal{A}_G \frac{c_s^2}{A_s^2 g^4} \le 15.36 \log g + 34.35 .$$
 (C.11)

One can see that the semi-analytical result (C.11) agrees with the fitting result (B.6) derived from numerical evaluation.

References

- A. H. Guth, "The Inflationary Universe: A Possible Solution to the Horizon and Flatness Problems," *Phys. Rev. D* 23 (1981) 347–356.
- [2] A. Borde and A. Vilenkin, "Eternal inflation and the initial singularity," *Phys. Rev. Lett.* 72 (1994) 3305–3309, arXiv:gr-qc/9312022.
- [3] A. Borde, A. H. Guth, and A. Vilenkin, "Inflationary space-times are incompletein past directions," *Phys. Rev. Lett.* **90** (2003) 151301, arXiv:gr-qc/0110012.
- [4] J. E. Lesnefsky, D. A. Easson, and P. C. W. Davies, "Past-completeness of inflationary spacetimes," *Phys. Rev. D* 107 no. 4, (2023) 044024, arXiv:2207.00955 [gr-qc].
- [5] G. Geshnizjani, E. Ling, and J. Quintin, "On the initial singularity and extendibility of flat quasi-de Sitter spacetimes," arXiv:2305.01676 [gr-qc].
- [6] J. Martin and R. H. Brandenberger, "The TransPlanckian problem of inflationary cosmology," *Phys. Rev. D* 63 (2001) 123501, arXiv:hep-th/0005209.
- [7] A. Bedroya and C. Vafa, "Trans-Planckian Censorship and the Swampland," JHEP 09 (2020) 123, arXiv:1909.11063 [hep-th].
- [8] Y. Cai and Y.-S. Piao, "Pre-inflation and trans-Planckian censorship," Sci. China Phys. Mech. Astron. 63 no. 11, (2020) 110411, arXiv:1909.12719 [gr-qc].
- M. Novello and S. E. P. Bergliaffa, "Bouncing Cosmologies," *Phys. Rept.* 463 (2008) 127–213, arXiv:0802.1634 [astro-ph].
- [10] J.-L. Lehners, "Ekpyrotic and Cyclic Cosmology," Phys. Rept. 465 (2008) 223-263, arXiv:0806.1245 [astro-ph].
- [11] R. Brandenberger and P. Peter, "Bouncing Cosmologies: Progress and Problems," Found. Phys. 47 no. 6, (2017) 797-850, arXiv:1603.05834 [hep-th].
- [12] Y.-F. Cai, A. Marciano, D.-G. Wang, and E. Wilson-Ewing, "Bouncing cosmologies with dark matter and dark energy," *Universe* 3 no. 1, (2016) 1, arXiv:1610.00938 [astro-ph.CO].
- [13] R. H. Brandenberger, "Alternatives to the inflationary paradigm of structure formation," Int. J. Mod. Phys. Conf. Ser. 01 (2011) 67-79, arXiv:0902.4731 [hep-th].
- [14] A. Ijjas and P. J. Steinhardt, "Bouncing Cosmology made simple," *Class. Quant. Grav.* 35 no. 13, (2018) 135004, arXiv:1803.01961 [astro-ph.CO].
- [15] Y.-F. Cai, J. Quintin, E. N. Saridakis, and E. Wilson-Ewing, "Nonsingular bouncing cosmologies in light of BICEP2," JCAP 07 (2014) 033, arXiv:1404.4364 [astro-ph.CO].
- [16] Y.-F. Cai, "Exploring Bouncing Cosmologies with Cosmological Surveys," Sci. China Phys. Mech. Astron. 57 (2014) 1414-1430, arXiv:1405.1369 [hep-th].
- [17] J. Karouby and R. Brandenberger, "A Radiation Bounce from the Lee-Wick Construction?," *Phys. Rev. D* 82 (2010) 063532, arXiv:1004.4947 [hep-th].
- [18] J. Karouby, T. Qiu, and R. Brandenberger, "On the Instability of the Lee-Wick Bounce," *Phys. Rev. D* 84 (2011) 043505, arXiv:1104.3193 [hep-th].

- [19] K. Bhattacharya, Y.-F. Cai, and S. Das, "Lee-Wick radiation induced bouncing universe models," *Phys. Rev. D* 87 no. 8, (2013) 083511, arXiv:1301.0661 [hep-th].
- [20] Y.-F. Cai, R. Brandenberger, and P. Peter, "Anisotropy in a Nonsingular Bounce," Class. Quant. Grav. 30 (2013) 075019, arXiv:1301.4703 [gr-qc].
- [21] J. Grain and V. Vennin, "Unavoidable shear from quantum fluctuations in contracting cosmologies," *Eur. Phys. J. C* 81 no. 2, (2021) 132, arXiv:2005.04222 [astro-ph.CO].
- [22] J. M. Cline, S. Jeon, and G. D. Moore, "The Phantom menaced: Constraints on low-energy effective ghosts," *Phys. Rev. D* 70 (2004) 043543, arXiv:hep-ph/0311312.
- [23] A. Vikman, "Can dark energy evolve to the phantom?," Phys. Rev. D 71 (2005) 023515, arXiv:astro-ph/0407107.
- [24] J.-Q. Xia, Y.-F. Cai, T.-T. Qiu, G.-B. Zhao, and X. Zhang, "Constraints on the Sound Speed of Dynamical Dark Energy," *Int. J. Mod. Phys. D* 17 (2008) 1229–1243, arXiv:astro-ph/0703202.
- [25] D. A. Easson and A. Vikman, "The Phantom of the New Oscillatory Cosmological Phase," arXiv:1607.00996 [gr-qc].
- [26] M. Libanov, S. Mironov, and V. Rubakov, "Generalized Galileons: instabilities of bouncing and Genesis cosmologies and modified Genesis," JCAP 08 (2016) 037, arXiv:1605.05992 [hep-th].
- [27] T. Kobayashi, "Generic instabilities of nonsingular cosmologies in Horndeski theory: A no-go theorem," *Phys. Rev. D* 94 no. 4, (2016) 043511, arXiv:1606.05831 [hep-th].
- [28] S. Akama and T. Kobayashi, "Generalized multi-Galileons, covariantized new terms, and the no-go theorem for nonsingular cosmologies," *Phys. Rev. D* 95 no. 6, (2017) 064011, arXiv:1701.02926 [hep-th].
- [29] Y.-F. Cai, W. Xue, R. Brandenberger, and X. Zhang, "Non-Gaussianity in a Matter Bounce," JCAP 05 (2009) 011, arXiv:0903.0631 [astro-ph.CO].
- [30] X. Gao, M. Lilley, and P. Peter, "Production of non-gaussianities through a positive spatial curvature bouncing phase," *JCAP* 07 (2014) 010, arXiv:1403.7958 [gr-qc].
- [31] X. Gao, M. Lilley, and P. Peter, "Non-Gaussianity excess problem in classical bouncing cosmologies," *Phys. Rev. D* 91 no. 2, (2015) 023516, arXiv:1406.4119 [gr-qc].
- [32] J. Quintin, Z. Sherkatghanad, Y.-F. Cai, and R. H. Brandenberger, "Evolution of cosmological perturbations and the production of non-Gaussianities through a nonsingular bounce: Indications for a no-go theorem in single field matter bounce cosmologies," *Phys. Rev. D* 92 no. 6, (2015) 063532, arXiv:1508.04141 [hep-th].
- [33] Y.-B. Li, J. Quintin, D.-G. Wang, and Y.-F. Cai, "Matter bounce cosmology with a generalized single field: non-Gaussianity and an extended no-go theorem," *JCAP* 03 (2017) 031, arXiv:1612.02036 [hep-th].
- [34] Y. Ageeva, P. Petrov, and V. Rubakov, "Nonsingular cosmological models with strong gravity in the past," *Phys. Rev. D* **104** no. 6, (2021) 063530, arXiv:2104.13412 [hep-th].
- [35] Y. Ageeva and P. Petrov, "On the strong coupling problem in cosmologies with "strong gravity in the past"," arXiv:2206.10646 [gr-qc].
- [36] S. Akama and S. Hirano, "Primordial non-Gaussianity from Galilean Genesis without strong coupling problem," *Phys. Rev. D* 107 no. 6, (2023) 063504, arXiv:2211.00388 [gr-qc].
- [37] J. Khoury, B. A. Ovrut, P. J. Steinhardt, and N. Turok, "The Ekpyrotic universe: Colliding branes and the origin of the hot big bang," *Phys. Rev. D* 64 (2001) 123522, arXiv:hep-th/0103239.

- [38] J. Middleton and J. D. Barrow, "The Stability of an Isotropic Cosmological Singularity in Higher-Order Gravity," *Phys. Rev. D* 77 (2008) 103523, arXiv:0801.4090 [gr-qc].
- [39] C. Lin, J. Quintin, and R. H. Brandenberger, "Massive gravity and the suppression of anisotropies and gravitational waves in a matter-dominated contracting universe," JCAP 01 (2018) 011, arXiv:1711.10472 [hep-th].
- [40] Y. Cai, Y. Wan, H.-G. Li, T. Qiu, and Y.-S. Piao, "The Effective Field Theory of nonsingular cosmology," JHEP 01 (2017) 090, arXiv:1610.03400 [gr-qc].
- [41] P. Creminelli, D. Pirtskhalava, L. Santoni, and E. Trincherini, "Stability of Geodesically Complete Cosmologies," JCAP 11 (2016) 047, arXiv:1610.04207 [hep-th].
- [42] Y. Cai, H.-G. Li, T. Qiu, and Y.-S. Piao, "The Effective Field Theory of nonsingular cosmology: II," *Eur. Phys. J. C* 77 no. 6, (2017) 369, arXiv:1701.04330 [gr-qc].
- [43] Y. Cai and Y.-S. Piao, "A covariant Lagrangian for stable nonsingular bounce," JHEP 09 (2017) 027, arXiv:1705.03401 [gr-qc].
- [44] R. Kolevatov, S. Mironov, N. Sukhov, and V. Volkova, "Cosmological bounce and Genesis beyond Horndeski," JCAP 08 (2017) 038, arXiv:1705.06626 [hep-th].
- [45] S. Akama and T. Kobayashi, "General theory of cosmological perturbations in open and closed universes from the Horndeski action," *Phys. Rev. D* 99 no. 4, (2019) 043522, arXiv:1810.01863 [gr-qc].
- [46] S. Mironov, V. Rubakov, and V. Volkova, "Genesis with general relativity asymptotics in beyond Horndeski theory," *Phys. Rev. D* 100 no. 8, (2019) 083521, arXiv:1905.06249 [hep-th].
- [47] S. Akama, S. Hirano, and T. Kobayashi, "Primordial non-Gaussianities of scalar and tensor perturbations in general bounce cosmology: Evading the no-go theorem," *Phys. Rev. D* 101 no. 4, (2020) 043529, arXiv:1908.10663 [gr-qc].
- [48] A. Ilyas, M. Zhu, Y. Zheng, Y.-F. Cai, and E. N. Saridakis, "DHOST Bounce," JCAP 09 (2020) 002, arXiv:2002.08269 [gr-qc].
- [49] M. Zhu, A. Ilyas, Y. Zheng, Y.-F. Cai, and E. N. Saridakis, "Scalar and tensor perturbations in DHOST bounce cosmology," *JCAP* 11 no. 11, (2021) 045, arXiv:2108.01339 [gr-qc].
- [50] Y. Cai, J. Xu, S. Zhao, and S. Zhou, "Perturbative unitarity and NEC violation in genesis cosmology," JHEP 10 (2022) 140, arXiv:2207.11772 [gr-qc]. [Erratum: JHEP 11, 063 (2022)].
- [51] D. Battefeld and P. Peter, "A Critical Review of Classical Bouncing Cosmologies," *Phys. Rept.* 571 (2015) 1–66, arXiv:1406.2790 [astro-ph.CO].
- [52] T. J. Battefeld and R. Brandenberger, "Vector perturbations in a contracting universe," *Phys. Rev. D* 70 (2004) 121302, arXiv:hep-th/0406180.
- [53] M. Bojowald and G. M. Hossain, "Cosmological vector modes and quantum gravity effects," *Class. Quant. Grav.* 24 (2007) 4801–4816, arXiv:0709.0872 [gr-qc].
- [54] F. C. Mena, D. J. Mulryne, and R. Tavakol, "Non-linear vector perturbations in a contracting universe," *Class. Quant. Grav.* 24 (2007) 2721–2734, arXiv:gr-qc/0702064.
- [55] R. H. Brandenberger, "The Matter Bounce Alternative to Inflationary Cosmology," arXiv:1206.4196 [astro-ph.CO].
- [56] C. Armendariz-Picon, V. F. Mukhanov, and P. J. Steinhardt, "Essentials of k essence," *Phys. Rev. D* 63 (2001) 103510, arXiv:astro-ph/0006373.
- [57] J. Garriga and V. F. Mukhanov, "Perturbations in k-inflation," Phys. Lett. B 458 (1999) 219-225, arXiv:hep-th/9904176.

- [58] X. Chen, "Primordial Non-Gaussianities from Inflation Models," Adv. Astron. 2010 (2010) 638979, arXiv:1002.1416 [astro-ph.CO].
- [59] X. Gao and D. A. Steer, "Inflation and primordial non-Gaussianities of 'generalized Galileons'," JCAP 12 (2011) 019, arXiv:1107.2642 [astro-ph.CO].
- [60] S. Weinberg, Cosmology. 2008.
- [61] V. F. Mukhanov, H. A. Feldman, and R. H. Brandenberger, "Theory of cosmological perturbations. Part 1. Classical perturbations. Part 2. Quantum theory of perturbations. Part 3. Extensions," *Phys. Rept.* 215 (1992) 203–333.
- [62] J. M. Bardeen, "Gauge Invariant Cosmological Perturbations," Phys. Rev. D 22 (1980) 1882–1905.
- [63] R. H. Brandenberger and R. Kahn, "COSMOLOGICAL PERTURBATIONS IN INFLATIONARY UNIVERSE MODELS," *Phys. Rev. D* 29 (1984) 2172.
- [64] A. Ota, H. J. Macpherson, and W. R. Coulton, "Covariant transverse-traceless projection for secondary gravitational waves," *Phys. Rev. D* 106 no. 6, (2022) 063521, arXiv:2111.09163 [gr-qc].
- [65] Planck Collaboration, N. Aghanim et al., "Planck 2018 results. VI. Cosmological parameters," Astron. Astrophys. 641 (2020) A6, arXiv:1807.06209 [astro-ph.CO]. [Erratum: Astron.Astrophys. 652, C4 (2021)].
- [66] Planck Collaboration, Y. Akrami et al., "Planck 2018 results. X. Constraints on inflation," Astron. Astrophys. 641 (2020) A10, arXiv:1807.06211 [astro-ph.CO].
- [67] Y.-F. Cai, D. A. Easson, and R. Brandenberger, "Towards a Nonsingular Bouncing Cosmology," JCAP 08 (2012) 020, arXiv:1206.2382 [hep-th].
- [68] A. Kandus, K. E. Kunze, and C. G. Tsagas, "Primordial magnetogenesis," *Phys. Rept.* 505 (2011) 1–58, arXiv:1007.3891 [astro-ph.CO].
- [69] G. Domènech, "Scalar Induced Gravitational Waves Review," Universe 7 no. 11, (2021) 398, arXiv:2109.01398 [gr-qc].
- [70] M. Giovannini and M. E. Shaposhnikov, "Primordial magnetic fields, anomalous isocurvature fluctuations and big bang nucleosynthesis," *Phys. Rev. Lett.* 80 (1998) 22–25, arXiv:hep-ph/9708303.
- [71] V. Acquaviva, N. Bartolo, S. Matarrese, and A. Riotto, "Second order cosmological perturbations from inflation," *Nucl. Phys. B* 667 (2003) 119–148, arXiv:astro-ph/0209156.
- [72] N. Bartolo, E. Komatsu, S. Matarrese, and A. Riotto, "Non-Gaussianity from inflation: Theory and observations," *Phys. Rept.* 402 (2004) 103–266, arXiv:astro-ph/0406398.
- [73] Z. Chang, X. Zhang, and J.-Z. Zhou, "The cosmological vector modes from a monochromatic primordial power spectrum," arXiv:2207.01231 [astro-ph.CO].