On the critical finite-size gap scaling for frustration–free Hamiltonians

Marius Lemm^{$*1$} and Angelo Lucia^{$†2,3$}

¹ Department of Mathematics, University of Tübingen, 72076 Tübingen, Germany

 2 Departamento de Análisis Matemático y Matemática Aplicada, Universidad Complutense de Madrid, 28040 Madrid, Spain

³ Instituto de Ciencias Matemáticas, 28049 Madrid, Spain

September 17, 2024

Abstract

We prove that the critical finite-size gap scaling for frustration-free Hamiltonians is of inverse-square type. The novelty of this note is that the result is proved on general graphs and for general finite-range interactions. Therefore, the inverse-square critical gap scaling is a robust, universal property of finite-range frustration-free Hamiltonians. This places further limits on their ability to produce conformal field theories in the continuum limit. Our proof refines the divide–and–conquer strategy of Kastoryano and the second author through the refined Detectability Lemma of Gosset-Huang.

1 Setup and main result

1.1 Assumptions on the graph and the Hamiltonian

We consider graphs Γ that can be embedded in \mathbb{R}^D .

Assumption 1.1 (Assumption on the graph). There exist a dimensional parameter $D \geq 1$, a constant $C_{\Gamma} \geq 1$, and an injective map $\iota : \Gamma \to \mathbb{R}^D$ such that

$$
C_{\Gamma}^{-1}|\iota(i) - \iota(j)| \le d_{\Gamma}(i,j) \le C_{\Gamma}|\iota(i) - \iota(j)|, \qquad \text{for all } i, j \in \Gamma.
$$
 (1)

Assumption [1.1](#page-0-0) is a rather weak assumption that allows many types of graph struc-tures. The prime example satisfying Assumption [1.1](#page-0-0) is the hypercubic lattice \mathbb{Z}^D and many standard lattices are also possible (e.g., honeycomb, triangular, fcc, bcc, hcp).

[∗]marius.lemm@uni-tuebingen.de

[†]anglucia@ucm.es

Assumption [1.1](#page-0-0) implies that the maximal degree of Γ is bounded (see Lemma [3.5](#page-7-0)) for this fact and other consequences of Assumption [1.1\)](#page-0-0). Examples where Assumption [1.1](#page-0-0) is not satisfied include graphs of unbounded degree and the Bethe lattice. While Γ is not required to have infinite cardinality, all relevant examples we have in mind have infinite cardinality.

Let $X \subset \Gamma$ be a finite subset. We place at each site of X a d-dimensional local Hilbert space. Accordingly, the Hilbert space on the region X is set to

$$
\mathcal{H}_X = \bigotimes_{i \in X} \mathbb{C}^d.
$$

We write \mathcal{A}_X for the associated local algebra of bounded linear operators with the natural inclusion $\mathcal{A}_X \subset \mathcal{A}_Y$ for $X \subset Y$ obtained by taking $A_X \otimes \mathrm{Id}_{Y \setminus X} \in \mathcal{A}_Y$. To every finite X, we associate an interaction $\Phi(X) \in \mathcal{A}_X$, a Hermitian bounded linear operator that is subject to the following conditions.

Assumption 1.2 (Assumptions on the interaction).

- (i) Φ has finite interaction range $R > 0$, i.e., $\Phi(X) = 0$ whenever diam $(X) > R$.
- (ii) $\Phi(X) \geq 0$ for all X and it is uniformly bounded from above and away from zero, i.e.,

$$
\Phi_{\text{max}} := \sup_{X \subset \Gamma} \|\Phi(X)\| < \infty,\tag{2}
$$

$$
\Phi_{\min} := \inf_{\substack{X \subset \Gamma: \ \Phi(X) \neq 0}} \text{gap } \Phi(X) > 0,\tag{3}
$$

where we introduced gap $\Phi(X) := \min(\text{spec }\Phi(X) \setminus \{0\}).$

Assumption (ii) is a uniformity assumption that allows us to reduce the Hamiltonian to a sum of local projectors. It holds in most of the practically relevant cases, e.g., in any translation-invariant situation. For finite-range interactions, Assumption (ii) is only violated in rather contrived situations, e.g., when the interactions behave very differently at infinity than in the bulk.

Let $\Lambda \subset \Gamma$ be finite. On the Hilbert space \mathcal{H}_{Λ} , we consider the Hamiltonian

$$
H_{\Lambda} = \sum_{X \subset \Lambda} \Phi(X) \tag{4}
$$

Our main assumption is that the Hamiltonian is frustration-free.

Assumption 1.3 (Assumption on the Hamiltonian). For every $\Lambda \subset \Gamma$, H_{Λ} is frustrationfree, i.e., ker $H_{\Lambda} \neq \{0\}.$

The main quantity of interest is the *spectral gap* gap(Λ) of H_{Λ} ,

$$
gap(H_{\Lambda}) = inf (spec H(\Lambda) \setminus \{0\})
$$
\n(5)

i.e., its smallest non-zero eigenvalue.

Well-known examples of frustration-free Hamiltonians that can be defined on general graphs Γ are the Heisenberg ferromagnet, AKLT Hamiltonians [\[AR+20;](#page-12-0) [Aff+88;](#page-11-0) [LY23;](#page-13-0) [WHR14\]](#page-13-1), and Hamiltonians with low-rank interactions [\[Sat+16;](#page-13-2) [JL22;](#page-12-1) [HJL24\]](#page-12-2). We recall that almost all mathematical investigations of the size of spectral gaps require frustration-freeness — without this assumption even absolutely fundamental questions like the Haldane conjecture [\[Hal83a;](#page-12-3) [Hal83b\]](#page-12-4) (which asserts that any integerspin Heisenberg antiferromagnetic chain is gapped) are completely open.

Our contribution in this work is to show that inverse-square critical gap scaling l^{-2} is a robust, universal property of gapless frustration-free Hamiltonians, in the sense that it holds under the rather weak Assumptions [1.1](#page-0-0) and [1.2](#page-1-0) on the graph and interactions. Our results extend prior ones of this kind [\[Ans20;](#page-12-5) [GM16;](#page-12-6) [LX22\]](#page-13-3) which required more restrictive assumptions. It comes with a mild logarithmic correction, which is unimportant for applications.

In particular, our result applies to any boundary condition and excludes gap closing rates that are of inverse length type l^{-1} , which would be characteristic of edge modes in conformal field theories [\[FMS12;](#page-12-7) [Kit06;](#page-12-8) [LM19\]](#page-12-9). Therefore, the result places further limits on the ability of finite-range frustration-free Hamiltonians to produce conformal field theories in the continuum limit.

1.2 Main result

For simplicity, we identify from now on a graph Γ satisfying Assumption [1.1](#page-0-0) with its Euclidean embedding $\iota(\Gamma) \subset \mathbb{R}^D$. This identification can be made without loss of generality, which can be seen by applying ι^{-1} to the Euclidean rectangles considered below.

We use the following class of domains from [\[KL18\]](#page-12-10) based on a construction originally appearing in [\[Ces01\]](#page-12-11). For each $k \in \mathbb{N}$, we let $l_k = (3/2)^{k/D}$ and introduce the Euclidean rectangle

$$
R(k) = [0, l_{k+1}] \times \cdots \times [0, l_{k+D}] \subset \mathbb{R}^D.
$$
 (6)

We define \mathcal{F}_k to be the collection of subsets of Γ which are contained in $R(k)$ up to translations and permutations of the Euclidean coordinates. We denote

$$
gap(\mathcal{F}_k) = \inf_{\Lambda \in \mathcal{F}_k} gap(H_{\Lambda}), \quad gap(\Gamma) = \inf_{k \ge 1} gap(\mathcal{F}_k). \tag{7}
$$

Theorem 1.4 (Main result). Suppose that

$$
gap(\Gamma) = 0.
$$

Then, for every $\epsilon > 0$,

$$
gap(\mathcal{F}_k) = o\left(\frac{k^{4+\epsilon}}{l_k^2}\right), \qquad \text{as } k \to \infty.
$$
 (8)

That is, if the infinite-volume gap vanishes, then the finite-size gaps gap(\mathcal{F}_k) "cannot close too slowly": they must close at least as the inverse square power of the linear dimension, l_k^{-2} $\frac{-2}{k}$.

We emphasize that the numerator $k^{4+\epsilon}$ is a mild logarithmic correction, since the length parameter $l_k = (3/2)^{k/D}$ grows exponentially in k.

Similar results to Theorem [1.4](#page-2-0) have been proved before using either the divideand-conquer approach [\[KL18\]](#page-12-10) or Knabe-type finite-size criteria [\[GM16;](#page-12-6) [Kna88;](#page-12-12) [LM19;](#page-12-9) [Lem20;](#page-12-13) [LX22\]](#page-13-3). In particular, an inverse-square gap closing rate was obtained in [\[Ans20;](#page-12-5) [GM16;](#page-12-6) [LX22\]](#page-13-3). These works only apply to nearest-neighbor 2-body interactions on hypercubic lattices, as well as the honeycomb and triangular lattices. The novelty of the present result is that it covers general graphs and general interactions. Indeed, it covers all finite-range k-body interactions. (While additional coarse-graining arguments could be applied to expand the scope of the prior works, they would not give a result of this generality.)

To summarize, the result shows that the inverse-square gap closing speed is a universal property of finite-range frustration-free Hamiltonians. We achieve this generality by using the generality of the divide-and-conquer strategy and amplifying it to near-optimal threshold scaling $\sim l^{-2}$ up to logarithmic corrections via the refined Detectability Lemma due to Gosset-Huang [\[GH16\]](#page-12-14); see also [\[AAV16;](#page-11-1) [Ans20;](#page-12-5) [AAG20\]](#page-11-2).

Remark 1.5. The inverse square scaling l^{-2} is well-known to be optimal in view of the Heisenberg ferromagnet, whose spectral gap vanishes as an inverse square on hypercubic lattices in any dimension, which can be easily proved rigorously by constructing explicit spin wave trial states.

2 Proof strategy

2.1 Reduction to sum of projectors

In a preliminary step, we replace each H_{Λ} by the sum of projectors

$$
\tilde{H}_{\Lambda} = \sum_{X \subset \Lambda} h_X \tag{9}
$$

where h_X is the projection onto the range of $\Phi(X)$ (and so $h_X = 0$ if $\Phi(X) = 0$). The fact that $\Phi_{\min}h_X \leq \Phi(X) \leq \Phi_{\max}h_X$ implies that for any finite $\Lambda \subset \Gamma$,

$$
\Phi_{\min} \operatorname{gap}(\tilde{H}_{\Lambda}) \leq \operatorname{gap}(H_{\Lambda}) \leq \Phi_{\max} \operatorname{gap}(\tilde{H}_{\Lambda}).
$$

Therefore, it suffices to prove Theorem [1.4](#page-2-0) for H_{Λ} . Indeed, the assumption $\text{gap}(H(\Lambda)) =$ 0 implies gap $(\tilde{H}_{\Lambda}) = 0$ and the conclusion that $\inf_{\Lambda \in \mathcal{F}_k} \text{gap}(\tilde{H}_{\Lambda}) = o\left(\frac{k^{4+\epsilon}}{l^2}\right)$ $\frac{4+\epsilon}{l_k^2}$ implies k $\inf_{\Lambda \in \mathcal{F}_k} \text{gap}(H_{\Lambda}) = o\left(\frac{k^{4+\epsilon}}{l^2}\right)$ l_k^2 . In the following, we always consider the Hamiltonian [\(9\)](#page-3-0).

2.2 Compatible sequences of domains

Let F denote a family of finite subsets of Γ . We recall a result from [\[KL18\]](#page-12-10), with a minor improvement appearing in [\[LPGPH23\]](#page-13-4).

Denote

$$
gap(\mathcal{F}) = \inf_{\Lambda \in \mathcal{F}} gap(\Lambda), \quad gap(\Lambda) = gap(\tilde{H}(\Lambda)). \tag{10}
$$

Given a finite $A \subset \Gamma$, we write P_A for the projection onto the frustration-free ground state of H_A (which is identical to the frustration-free ground state of the original Hamiltonian H_A).

Theorem 2.1 ([\[LPGPH23,](#page-13-4) Theorem 2.3]). Let F and F' be two families of finite subsets of a graph Γ satisfying Assumption [1.1.](#page-0-0) Suppose that there exists $s \in \mathbb{N}$ and $\delta \in [0,1]$ satisfying the following property: for each $Y \in \mathcal{F}' \setminus \mathcal{F}$, there exists $(A_i, B_i)_{i=1}^s$ pairs of elements in $\mathcal F$ such that:

- 1. $Y = A_i \cup B_i$ for each $i = 1, \ldots, s;$
- 2. $(A_i \cap B_i) \cap (A_j \cap B_j) = \emptyset$ whenever $i \neq j$;
- 3. $||P_{A_i}P_{B_i} P_Y|| \leq \delta$ for every $i = 1, \ldots, s$.

Then

$$
gap(\mathcal{F}') \ge \frac{1-\delta}{1+\frac{1}{s}} \operatorname{gap}(\mathcal{F}).
$$
\n(11)

The point is that elements of \mathcal{F}_k can be nicely decomposed in terms of elements of \mathcal{F}_{k-1} .

Let $\alpha \in \{1, \ldots, D\}$. We write Π_{α} for the coordinate projection

$$
\Pi_{\alpha}(x_1,\ldots,x_D)=(x_1,\ldots,x_{\alpha-1},0,x_{\alpha+1},\ldots,x_D).
$$

Proposition 2.2 ([\[KL18,](#page-12-10) Proposition 1]). Let $s_k \leq \frac{1}{8}$ $\frac{1}{8}l_k$. For each $k \geq 1$, the conditions 1 and 2 of Theorem [2.1](#page-4-0) are satisfied for $\mathcal{F}' = \mathcal{F}_{k+1}$, $\mathcal{F} = \mathcal{F}_k$ and $s = s_k$. Moreover, for each $Y \in \mathcal{F}_k \setminus \mathcal{F}_{k-1}$, it is possible to choose the pairs $(A_i, B_i)_{i=1}^{s_k}$ in such a way that

$$
d_{\Gamma}(A_i \setminus B_i, B_i \setminus A_i) \ge C_{\Gamma}^{-1} \left(\frac{l_k}{8s_k} - 2\right). \tag{12}
$$

Moreover, there exists $\alpha \in \{1, \ldots, D\}$ such that

$$
\Pi_{\alpha} A_i = \Pi_{\alpha} B_i. \tag{13}
$$

Property [\(13\)](#page-4-1) was not explicitly stated in previous works, but it follows directly from the construction. It is important for us here to connect to the refined Detectability Lemma, which requires us to make a 1D reduction to choose a coarse-graining direction.

Iterating Theorem [2.1](#page-4-0) along the sequence $\{\mathcal{F}_k\}_{k\geq 1}$ yields the following gap estimate.

Corollary 2.3. Set

$$
\delta_k := \sup_{(A,B)\in S_k} \|P_A P_B - P_{A\cup B}\|,\tag{14}
$$

where

$$
S_k := \{ (A, B) \in \mathcal{F}_{k-1} \times \mathcal{F}_{k-1} : A \cup B \in \mathcal{F}_k, (12) \text{ and } (13) \text{ hold } \}.
$$
 (15)

Let k_0 be the smallest k such that $\delta_k < 1$ for every $k \geq k_0$.

Then,

$$
gap(\Gamma) \geq \Phi_{\min} \, gap(\mathcal{F}_{k_0}) \prod_{k=k_0}^{\infty} \frac{1 - \delta_k}{1 + \frac{1}{s_k}}.
$$
\n
$$
(16)
$$

Note that the constant on the r.h.s. of [\(16\)](#page-4-3) is strictly positive if and only if $(\delta_k)_k$ and $\sqrt{1}$ $\overline{s_k}$ \setminus are summable sequences. In this case, the "infinite-volume" gap $\inf_{n\geq 1} \text{gap}(\mathcal{F}_n)$ is lower-bounded by a constant and one commonly says that the infinite system is gapped.

2.3 Proof of Theorem [1.4](#page-2-0) assuming Proposition [2.4](#page-5-0)

The proof of Theorem [1.4](#page-2-0) follows from the following Proposition [2.4](#page-5-0) whose proof will occupy us in the rest of the paper.

Proposition 2.4. Let $\lambda_k \in (0,1)$ such that

$$
gap(\mathcal{F}_k) \ge \lambda_k > 0, \quad \forall k. \tag{17}
$$

Suppose that there exists a sequence s_k such that $(\frac{1}{s_k})$ $(\frac{1}{s_k})_k$ is summable and

$$
\liminf_{k \to \infty} \sqrt{\lambda_k} \frac{l_k}{ks_k} > 0. \tag{18}
$$

 \Box

Then gap(Γ) > 0.

Proof of Theorem [1.4.](#page-2-0) As explained in Subsection [2.1,](#page-3-1) it suffices to prove the claim for the Hamiltonians [\(9\)](#page-3-0). We reason by contradiction. Suppose that

$$
gap(\mathcal{F}_k) \ge c \frac{k^{4+\epsilon}}{l_k^2}, \quad \forall k,
$$

for some positive constant c and some $\epsilon > 0$. Set $\lambda_k = c \frac{k^{4+\epsilon}}{l^2}$ $\frac{d^2 k}{l_k^2}$, which is $\lt 1$ for sufficiently large k, and choose $s_k = k^{1+\frac{\epsilon}{2}}$, making $(\frac{1}{s_k})_k$ summable. Moreover

$$
\liminf_{k \to \infty} \sqrt{\lambda_k} \frac{l_k}{ks_k} = \sqrt{c} \liminf_{k \to \infty} \frac{k^{2 + \frac{\epsilon}{2}}}{l_k} \frac{l_k}{ks_k} = \sqrt{c} > 0.
$$

By Proposition [2.4,](#page-5-0) this implies that $\text{gap}(\Gamma) > 0$, leading to a contradiction.

3 Refined overlap bound and proof of Proposition [2.4](#page-5-0)

We derive an effective upper bound on δ_k , defined in [\(14\)](#page-4-4), in terms of the lower bound on the spectral gap, λ_k . Recall S_k defined in [\(15\)](#page-4-5).

Lemma 3.1 (Refined overlap bound). There exist $C_1, C_2 > 0$ such that the following holds. Let $k \ge 1$ and $(A, B) \in S_k$. Suppose that $\lambda_k \in (0, 1)$ satisfies $\text{gap}(\mathcal{F}_k) \ge \lambda_k > 0$. Then,

$$
\delta_k \le C_1 \exp\left(-C_2 \sqrt{\lambda_k} \frac{l_k}{s_k}\right). \tag{19}
$$

This is the key improvement over Theorem 11 in [\[KL18\]](#page-12-10), in which the exponent √ $\overline{\lambda_k}$ behaved as λ_k instead.

We postpone the proof of Lemma [3.1](#page-5-1) for the moment. We first combine it with Corollary [2.3](#page-4-6) to obtain Proposition [2.4.](#page-5-0)

Proof of Proposition [2.4](#page-5-0) assuming Lemma [3.1.](#page-5-1) By Lemma [3.1](#page-5-1) and the definition of δ_k , we have that

$$
\delta_k \le \exp\left(-C\sqrt{\lambda_k} \frac{l_k}{s_k}\right). \tag{20}
$$

We now apply Corollary [2.3,](#page-4-6) and in order to show that the r.h.s. of [\(16\)](#page-4-3) is non-zero, we need to show that $(\delta_k)_k$ is a summable sequence. We will do so by checking that it satisfies the root test:

$$
\limsup_{k \to \infty} \exp \left(-C \sqrt{\lambda_k} \frac{l_k}{s_k} \right)^{1/k} = \exp \left(-C \liminf_{k \to \infty} \sqrt{\lambda_k} \frac{l_k}{ks_k} \right)
$$

which is strictly smaller than 1 by assumption.

In the remainder of this section, we prove Lemma [3.1.](#page-5-1) The proof will follow from a suitable generalization of the refined Detectability Lemma (Lemma [3.6](#page-8-0) below), whose setup we describe next. In contrast to [\[KL18\]](#page-12-10) and rather similarly to [\[Ans20\]](#page-12-5), we first perform a 1D coarse-graining which is a key ingredient in the refined Detectability Lemma. The differences compared to [\[Ans20\]](#page-12-5) are that we work in a more general setup (general number of layers and interaction range) and we need to track how Euclidean versus graph geometry enters in the arguments, as both are relevant.

3.1 Coarse-graining

Let $k \geq 1$ and consider $(A, B) \in S_k$. We denote $\Lambda = A \cup B$. Let $\alpha \in \{1, ..., D\}$ be such that $\Pi_{\alpha}A = \Pi_{\alpha}B$. Without loss of generality, by rotation, we can assume that $\alpha = 1$.

We fix a coarse-graining parameter

$$
t \ge \max\{2, C_{\Gamma}R\}
$$

and we define the width-4t "columns" centered at $j \in \mathbb{Z}$,

$$
\mathcal{C}_j = ([j - 2t + 1, j + 2t - 1] \times \mathbb{R}^{D-1}) \cap \Lambda, \qquad j \in \mathbb{Z}
$$

as well as the associated coarse-grained Hamiltonians

$$
\tilde{H}_j = \sum_{X \subset \mathcal{C}_j} h_X.
$$

We write Q_j for the ground state projector of \tilde{H}_j .

Definition 3.2. Let us define two sets of indices:

$$
\mathcal{I}_e(t) := \{(2+6j) t \mid j \in \mathbb{Z}\}, \quad \mathcal{I}_o(t) := \{(5+6j) t \mid j \in \mathbb{Z}\}.
$$
 (21)

Lemma 3.3. Let $\#\in\{e, o\}$ and let $m, n \in \mathcal{I}_{\#}(t)$ be distinct. Then $[Q_m, Q_n] = 0$.

Proof of Lemma [3.3.](#page-7-1) Without loss of generality, let $\# = e$. Since $m, n \in \mathcal{I}_e(t), Q_m$ is supported on $[6jt+1, 6jt+4t-1] \times \mathbb{R}^{D-1}$ and Q_n is supported on $[6j't+1, 6j't+4t-1]$ $1] \times \mathbb{R}^{D-1}$ for two distinct j, j' ∈ Z. The Euclidean distance between these intervals is at least $2t + 2$. By [\(1\)](#page-0-1), their Γ-distance is at least C_{Γ}^{-1} $\Gamma^{-1}(2t+2)$. Since $t \ge \max\{2, C_{\Gamma}R\},\$ we have C_{Γ}^{-1} $\Gamma^{-1}(2t+2) \geq 2R$. Now $[Q_m, Q_n] = 0$ follows because the interaction range is R. For $\# = o$, the argument is analogous. \Box

Lemma [3.3](#page-7-1) allows us to group the Q_m operators into the even and the odd ones, which commute amongst each other.

Definition 3.4. The *t*-coarse grained detectability lemma operator $DL(t)$ is given by

$$
DL(t) = \prod_{m \in \mathcal{I}_e(t)} Q_m \cdot \prod_{m \in \mathcal{I}_o(t)} Q_m.
$$
 (22)

3.2 Basic properties of the interaction

We note two basic properties of the interaction Φ that follow from Assumption [1.2](#page-1-0) and that are used in the Detectability Lemma.

Lemma 3.5. There exist integers $g, L \geq 1$ such that the following holds.

 \bullet Φ can be decomposed into L layers, i.e., there exists a disjoint decomposition ("L-coloring") of

$$
\{X \subset \Gamma \,:\, \text{diam}(X) \le R\} = \mathcal{X}_1 \sqcup \mathcal{X}_2 \sqcup \ldots \sqcup \mathcal{X}_L \tag{23}
$$

such that for every $X, Y \in \mathcal{X}_i$, it holds that $X \cap Y = \emptyset$.

• For every subset $X \subset \Gamma$ such that $\Phi(X) \neq 0$, there exist at most g subsets $Y \neq X$ such that $[\Phi(X), \Phi(Y)] \neq 0$.

Proof of Lemma [3.5.](#page-7-0) We first observe that Assumption [1.1](#page-0-0) implies that, for any $r > 0$ and $i \in \Gamma$, the ball

$$
B_r^{\Gamma}(i) := \{ j \in \Gamma \ : \ d_{\Gamma}(i,j) \le r \}
$$

has bounded cardinality uniformly in *i*. This follows from the fact that $\iota(B_r^{\Gamma}(i)) \subset \mathbb{R}^D$ is contained in the Euclidean ball $B_{C_{\Gamma}r}(i)$ and all its distinct points are at distance at least $C_{\Gamma}^{-1} > 0$.

Concerning the first bullet point, we consider the hypergraph whose hyperedges are those $X \in \mathcal{P}(\Lambda)$ for which $\Phi(X) \neq 0$. We write $\Delta(\Gamma)$ for the degree of the hypergraph, defined as usual as the maximal number of hyperedges intersecting at a vertex. By the finite-range assumption, given any $i \in \Gamma$, the number of X containing i is bounded by $2^{|B_R(i)|}$ and therefore this number also bounds the degree. As noted above, the bound on $|B_R(i)|$ is uniform in i and so $\Delta(\Gamma)$ is bounded. In this language, L is called the edge-chromatic index of the hypergraph and a simple bound from graph theory is $L \leq \left\lfloor \frac{3\Delta(\Gamma)}{2} \right\rfloor$ $\frac{\Gamma(\Gamma)}{2}$.

Concerning the second bullet point, we note that $[\Phi(X), \Phi(Y)] \neq 0$ implies $X \cap Y \neq 0$ \emptyset . By the finite-range assumption, X and Y must therefore lie in a $B_{2R}(i)$ for some $i \in \Gamma$. Hence, $g \leq 2^{|B_{2R}(i)|} - 1$. As noted above, the bound on $|B_R(i)|$ is uniform in i and so g is bounded. \Box

Of course, the bounds on L and g given in the above proof are far from optimal and much better bounds can usually be read off more simply. Since L and g do not enter in the main results, we do not dwell on this issue and refer readers interested in tighter bounds to the combinatorics literature, e.g., [\[PS89\]](#page-13-5) and the recent survey [\[Kan+23\]](#page-12-15).

3.3 Refined Detectability Lemma

The following refined Detectability Lemma is due to Gosset and Huang [\[GH16\]](#page-12-14); see also [\[AAV16;](#page-11-1) [AAG20\]](#page-11-2).

Lemma 3.6 (Refined Detectability Lemma). Let $k \geq 1$, $(A, B) \in S_k$ and $\text{gap}(\mathcal{F}_k) \geq$ $\lambda_k > 0$. Denote $\Lambda = A \cup B$. For a normalized state $|\phi\rangle$ orthogonal to the ground state sector of H_{Λ} , it holds that

$$
||DL(t)|\phi\rangle|| \leq 2 \exp\left(-\left(\frac{t}{C_{\Gamma}(L-1)R} - 2\right)\sqrt{\frac{\lambda_k}{1+g^2}}\right) \tag{24}
$$

We now prove Lemma [3.6.](#page-8-0) The first step is to show that one can "smuggle" a polynomial F with $F(0) = 1$ and of controlled degree between the products defining $DL(t)$. This polynomial will later be chosen as a suitable Chebyshev polynomial.

Recall the notation [\(23\)](#page-7-2) for the L layers. The standard detectability lemma operator for \tilde{H}_{Λ} from [\(9\)](#page-3-0) is defined as

$$
T := T_L \dots T_1, \qquad T_\beta := \prod_{X \in \mathcal{X}_\beta \cap \Lambda} (1 - h_X), \quad \beta \in \{1, \dots, L\}.
$$
 (25)

The following lemma generalizes Lemma 3.1 in [\[AAG20\]](#page-11-2) to any number of layers, interaction range, and Euclidean-embedded graphs.

Lemma 3.7. Let F be a polynomial of degree at most $\left[\frac{t}{2C_{\Gamma}(L-1)R}-1\right]$, such that $F(0) = 1$. Then

$$
DL(t) = \prod_{m \in \mathcal{I}_e(t)} Q_m \cdot F\left(\mathbb{1} - T^{\dagger}T\right) \cdot \prod_{m \in \mathcal{I}_o(t)} Q_m.
$$
 (26)

Figure 5 in [\[AAG20\]](#page-11-2) is an instructive depiction of the proof idea in one dimension.

Proof. For $\beta \in \{1, ..., L\}$ and $S \subset \Lambda$, define

$$
T_{\beta}^{S} = \prod_{\substack{X \in \mathcal{X}_{\beta} \\ X \cap S \neq \emptyset}} (1 - h_X).
$$

Let

$$
S_0 = \Lambda \setminus \, \mathrm{supp} \left(\prod_{m \in \mathcal{I}_e(t)} Q_m \right)
$$

and define iteratively for $j \geq 1$,

$$
S_j = \{ i \in \Lambda \; : \; d_\Gamma(i, S_{j-1}) \le R \} \, .
$$

Note that, by construction, $S_0 \subset S_1 \subset \cdots \subset S_n$ for a suitable $n \geq 1$.

Because of frustration freeness, we have that

$$
\prod_{m \in \mathcal{I}_e(t)} Q_m \cdot T_1 = \prod_{m \in \mathcal{I}_e(t)} Q_m \cdot T_1^{S_0}.
$$

Since each T_β is a self-adjoint projection, we have that

$$
T^{\dagger}T = T_1T_2 \dots T_{L-1}T_LT_{L-1} \dots L_1,
$$

and so $(T^{\dagger}T)^{q} = (T_{1}T_{2} \ldots T_{L-1}T_{L}T_{L-1} \ldots T_{2})^{q}T_{1}$. Therefore

$$
\prod_{m \in \mathcal{I}_e(t)} Q_m \cdot (T^{\dagger}T)^q = \prod_{m \in \mathcal{I}_e(t)} Q_m \cdot T_1^{S_0} T_2^{S_1} \cdots T_2^{S_{qL(2L-2)-1}} T_1^{S_{q(2L-2)}}.
$$

Now, if $S_{qL(2L-2)-1}$ is contained in the support of $\prod_{m\in\mathcal{I}_o(t)} Q_m$, we have that

$$
T_1^{S_0}T_2^{S_1}\cdots T_2^{S_{qL(2L-2)-1}}T_1^{S_{q(2L-2)}}\cdot \prod_{m\in\mathcal{I}_o(t)}Q_m=\prod_{m\in\mathcal{I}_o(t)}Q_m.
$$

By construction, the set S_0 is at Euclidean distance t from the complement of the support of $\prod_{m\in\mathcal{I}_o(t)}Q_m$ and so its graph distance to the complement of the support of $\prod_{m \in \mathcal{I}_o(t)} Q_m$ is at least C_Γ^{-1} $\int_{\Gamma}^{-1} t$. By induction, S_j is at graph distance C_{Γ}^{-1} $\int_{\Gamma}^{-1} t - jR$ from the complement of the support of $\prod_{m \in \mathcal{I}_o(t)} Q_m$. Therefore, this graph distance is non-zero as long as $j < t/(C_{\Gamma}R)$. Thus by choosing $q \leq \lceil t/(2C_{\Gamma}(L-1)R) - 1 \rceil$, we can show that

$$
\prod_{m \in \mathcal{I}_e(t)} Q_m \cdot (T^{\dagger} T)^q \cdot \prod_{m \in \mathcal{I}_o(t)} Q_m = DL(t).
$$

By linearity, we obtain the stated claim.

The identity [\(26\)](#page-8-1) is used to produce the polynomial and the outside products are then dropped, as summarized in the next corollary. Recall that P_{Λ} denotes the projection onto the ground state of \tilde{H}_{Λ} .

Corollary 3.8. We have

$$
||DL(t)P_{\Lambda}^{\perp}|| \le \inf_{F} F_*(||TP_{\Lambda}^{\perp}||^2), \tag{27}
$$

 \Box

where the infimum runs over polynomials of degree at most $\left\lceil \frac{t}{2C_{\Gamma}(L-1)R} - 1 \right\rceil$ satisfying $F(0) = 1$, and we set

$$
F_*(\epsilon) := \inf_{1-\epsilon \le x \le 1} |F(x)| \,, \quad \forall \epsilon \in [0,1]. \tag{28}
$$

Proof of Corollary [3.8.](#page-9-0) Let F be any of the admissible polynomials. By Lemma [3.7,](#page-8-2)

$$
||DL(t)P_{\Lambda}^{\perp}|| \leq ||F(\mathbb{1} - T^{\dagger}T)P_{\Lambda}^{\perp}||.
$$

Then

$$
(1 - ||TP_{\Lambda}^{\perp}||^2) \mathop{1\!\!\!\!1} \le P_{\Lambda}^{\perp} (1 - T^{\dagger}T) P_{\Lambda}^{\perp} \le 1
$$

yields Corollary [3.8.](#page-9-0)

We recall from [\[Ara+13\]](#page-12-16) a basic fact about Chebyshev polynomials. We let T_q denote the degree-q Chebyshev polynomial.

Lemma 3.9 ([\[Ara+13\]](#page-12-16)). For every integer $q \ge 1$ and number $\gamma \in (0,1)$, let

$$
\operatorname{Step}_{q,\gamma}(x) := \frac{T_q\left(\frac{2(1-x)}{1-\gamma} - 1\right)}{T_q\left(\frac{2}{1-\gamma} - 1\right)}, \qquad x \in \mathbb{R}.
$$

Then

$$
|\text{Step}_{q,\gamma}(x)| \le 2 \exp(-2q\sqrt{\gamma}), \qquad \gamma \le x \le 1. \tag{29}
$$

Proof of Lemma [3.6.](#page-8-0) Take any vector $|\psi^{\perp}\rangle$ orthogonal to the ground state sector of H_{Λ} . By the standard Detectability Lemma [\[AAV16,](#page-11-1) Corollary 1], we have

$$
||T|\psi^{\perp}\rangle||^2 \le \frac{1}{1 + \lambda_k/g^2}
$$

and therefore $||TP_{\Lambda}^{\perp}||^2 \leq \frac{1}{1+\lambda_k/g^2}$.

By applying Corollary [3.8](#page-9-0) and Lemma [3.9](#page-10-0) with $q = \left[\frac{t}{2C_{\Gamma}(L-1)R} - 1\right]$ and $\gamma =$ λ_k $\frac{\lambda_k}{\lambda_k+g^2} \in (0,1)$, we obtain

$$
||DL(t)|\phi\rangle|| \le \inf_{F} F_*(||LP_{\Lambda}^{\perp}||^2) \le 2 \exp\left(-2\left\lceil \frac{t}{2C_{\Gamma}(L-1)R} - 1 \right\rceil \sqrt{\frac{\lambda_k}{\lambda_k + g^2}} \right)
$$

$$
\le 2 \exp\left(-\left(\frac{t}{C_{\Gamma}(L-1)R} - 2\right) \sqrt{\frac{\lambda_k}{1 + g^2}} \right)
$$

where we used $\lambda_k \leq 1$ in the last step. This proves Lemma [3.6.](#page-8-0)

 \Box

3.4 Proof of Lemma [3.1](#page-5-1)

Let $k \geq 1$, $(A, B) \in S_k$, and denote $\Lambda = A \cup B$. We perform the coarse-graining as described in Subsection [3.1.](#page-6-0)

We now define M_A and M_B by reordering the projector Q_m appearing in $DL(t)$ in such a way that $DL(t) = M_A M_B$. We set

$$
M_A = \prod_{\substack{m \in \mathcal{I}_e(t): \\ \text{supp } Q_m \cap A \setminus B \neq \emptyset}} Q_m \prod_{\substack{m \in \mathcal{I}_o(t): \\ \exists m' \in \mathcal{I}_e(t): \\ \text{supp } Q_{m'} \cap A \setminus B \neq \emptyset, \\ \text{supp } Q_m \cap \text{supp } Q_{m'} \neq \emptyset}} Q_m
$$

 \Box

and we write M_B for the remaining Q_m appearing in $DL(t)$, ordered in the same way as they appear in $DL(t)$. Then $DL(t) = M_A M_B$ because, in M_A , the odd-type terms in the second product commute precisely with all even-type terms except for those in the first product. If the Euclidean distance between $A \setminus B$ and $B \setminus A$ exceeds 8t, then M_A is supported on A and M_B is supported on B.

Recall that P_A denotes the projection onto the ground state of H_A . We have that $P_A = P_A M_A$, and therefore $P_A M_B = P_A M_A M_B = P_A D L(t)$. This implies that

$$
||(P_A - M_A)M_B|| = ||DL(t)P_A^{\perp}|| \le ||DL(t)P_{AB}^{\perp}||,
$$

where the inequality follows from the fact that $P_A^{\perp} \leq P_{AB}^{\perp}$, a consequence of frustrationfreeness. Since it also holds that $P_B = P_B M_B$ and $P_B M_A = P_B D L(t)$, the analogous bound with A and B interchanged. This gives that

$$
||P_A P_B - P_{AB}|| = ||P_A P_B - M_A P_B|| + ||M_A P_B - M_A M_B|| + ||M_A M_B - P_{AB}||
$$

\n
$$
\leq ||(P_A - M_A)P_B|| + ||M_A (P_B - M_B)|| + ||DL(t)P_{AB}^{\perp}|| \leq 3||DL(t)P_{AB}^{\perp}||.
$$

By Lemma [3.6,](#page-8-0) we have

$$
||DL(t)P_{AB}^{\perp}|| \leq 2 \exp\left(-\left(\frac{t}{C_{\Gamma}(L-1)R}-2\right)\sqrt{\frac{\lambda_k}{1+g^2}}\right).
$$

By [\(12\)](#page-4-2) and Assumption [1.1,](#page-0-0) we can choose t as a constant multiple of $\frac{l_k}{s_k}$. Defining $C_1, C_2 > 0$ appropriately, this proves Lemma [3.1.](#page-5-1) П

Acknowledgments

The authors thank the organizers of the online Lattice Seminar, during which the idea for this work arose. The research of M.L. is supported by the DFG through the grant TRR 352 – Project-ID 470903074. A. L. acknowledges financial support from grants PID2020-113523GB-I00 and CEX2023-001347-S, funded by MICIU/AEI/ 10.13039/501100011033, and from grant RYC2019-026475-I funded by MICIU/AEI/ 10.13039/501100011033 and "ESF Investing in your future".

References

- [AAG20] Anurag Anshu, Itai Arad, and David Gosset. "Entanglement subvolume law for 2D frustration-free spin systems". Proceedings of the 52nd Annual ACM SIGACT Symposium on Theory of Computing. 2020, pp. 868–874.
- [AAV16] Anurag Anshu, Itai Arad, and Thomas Vidick. "Simple proof of the detectability lemma and spectral gap amplification". Physical Review B 93.20 (2016), p. 205142.
- [Aff+88] Ian Affleck et al. "Valence bond ground states in isotropic quantum antiferromagnets". Communications in Mathematical Physics 115.3 (1988), pp. 477– 528.
- [Ans20] Anurag Anshu. "Improved local spectral gap thresholds for lattices of finite size". Physical Review B 101.16 (Apr. 2020). DOI: [10.1103/physrevb.101.](https://doi.org/10.1103/physrevb.101.165104) [165104](https://doi.org/10.1103/physrevb.101.165104).
- [AR+20] Houssam Abdul-Rahman et al. "A class of two-dimensional AKLT models with a gap". Analytic Trends in Mathematical Physics 741 (2020), pp. 1–21.
- [Ara+13] Itai Arad et al. "An area law and sub-exponential algorithm for 1D systems". arXiv preprint arXiv:1301.1162 (2013).
- [Ces01] Filippo Cesi. "Quasi-factorization of the entropy and logarithmic Sobolev inequalities for Gibbs random fields". Probability Theory and Related Fields 120.4 (Aug. 2001), 569-584. DOI: 10.1007/p100008792.
- [FMS12] Philippe Francesco, Pierre Mathieu, and David Sénéchal. Conformal field theory. Springer Science & Business Media, 2012.
- [GH16] David Gosset and Yichen Huang. "Correlation length versus gap in frustrationfree systems". Physical Review Letters 116.9 (2016), p. 097202.
- [GM16] David Gosset and Evgeny Mozgunov. "Local gap threshold for frustration-free spin systems". Journal of Mathematical Physics 57.9 (2016), p. 091901. DOI: [https://doi.org/10.1063/1.4962337](https://doi.org/https://doi.org/10.1063/1.4962337).
- [Hal83a] F Duncan M Haldane. "Continuum dynamics of the 1-D Heisenberg antiferromagnet: Identification with the $O(3)$ nonlinear sigma model". Physics letters a 93.9 (1983), pp. 464-468. DOI: https://doi.org/10.1016/0375-[9601\(83\)90631-x](https://doi.org/https://doi.org/10.1016/0375-9601(83)90631-x).
- [Hal83b] F Duncan M Haldane. "Nonlinear field theory of large-spin Heisenberg antiferromagnets: semiclassically quantized solitons of the one-dimensional easyaxis Néel state". Physical Review Letters 50.15 (1983), p. 1153. DOI: [https:](https://doi.org/https://doi.org/10.1103/physrevlett.50.1153) [//doi.org/10.1103/physrevlett.50.1153](https://doi.org/https://doi.org/10.1103/physrevlett.50.1153).
- [HJL24] Nicholas Hunter-Jones and Marius Lemm. "Gapped quantum spin systems on general graphs at large local dimension". in preparation (2024).
- [JL22] Ian Jauslin and Marius Lemm. "Random translation-invariant Hamiltonians and their spectral gaps". Quantum 6 (2022), p. 790.
- [Kan+23] Dong Yeap Kang et al. "Graph and hypergraph colouring via nibble methods: A survey". Proceedings of the 8th ECM (2023). DOI: [10.4171/8ECM/11](https://doi.org/10.4171/8ECM/11).
- [Kit06] Alexei Kitaev. "Anyons in an exactly solved model and beyond". Annals of Physics 321.1 (2006), pp. 2–111.
- [KL18] Michael J Kastoryano and Angelo Lucia. "Divide and conquer method for proving gaps of frustration free Hamiltonians". Journal of Statistical Mechanics: Theory and Experiment 2018.3 (Mar. 2018), p. 033105. DOI: [10.1088/1742-](https://doi.org/10.1088/1742-5468/aaa793) [5468/aaa793](https://doi.org/10.1088/1742-5468/aaa793).
- [Kna88] Stefan Knabe. "Energy gaps and elementary excitations for certain VBSquantum antiferromagnets". Journal of statistical physics 52.3 (1988), pp. 627– 638. DOI: [https://doi.org/10.1007/bf01019721](https://doi.org/https://doi.org/10.1007/bf01019721).
- [Lem20] Marius Lemm. "Finite-size criteria for spectral gaps in D-dimensional quantum spin systems". Analytic trends in mathematical physics 741 (2020), p. 121. DOI: [https://doi.org/10.1090/conm/741/14923](https://doi.org/https://doi.org/10.1090/conm/741/14923).
- [LM19] Marius Lemm and Evgeny Mozgunov. "Spectral gaps of frustration-free spin systems with boundary". Journal of Mathematical Physics 60.5 (2019), p. 051901. DOI: [https://doi.org/10.1063/1.5089773](https://doi.org/https://doi.org/10.1063/1.5089773).
- [LPGPH23] Angelo Lucia, David Pérez-García, and Antonio Pérez-Hernández. "Thermalization in Kitaev's quantum double models via tensor network techniques". Forum of Mathematics, Sigma 11 (2023). DOI: [10.1017/fms.2023.98](https://doi.org/10.1017/fms.2023.98).
- [LX22] Marius Lemm and David Xiang. "Quantitatively improved finite-size criteria for spectral gaps". Journal of Physics A: Mathematical and Theoretical 55.29 (2022), p. 295203. DOI: [https://doi.org/10.1088/1751-8121/ac7989](https://doi.org/https://doi.org/10.1088/1751-8121/ac7989).
- [LY23] Angelo Lucia and Amanda Young. "A nonvanishing spectral gap for AKLT models on generalized decorated graphs". Journal of Mathematical Physics 64.4 (2023).
- [PS89] Nicholas Pippenger and Joel Spencer. "Asymptotic behavior of the chromatic index for hypergraphs". Journal of combinatorial theory, Series A 51.1 (1989), pp. 24–42.
- [Sat+16] Or Sattath et al. "When a local Hamiltonian must be frustration-free". Proceedings of the National Academy of Sciences 113.23 (2016), pp. 6433–6437.
- [WHR14] Tzu-Chieh Wei, Poya Haghnegahdar, and Robert Raussendorf. "Hybrid valencebond states for universal quantum computation". Physical Review A 90.4 (2014), p. 042333.