

POLYNOMIAL FUNCTIONS ON A CLASS OF FINITE NON-COMMUTATIVE RINGS

AMR ALI ABDULKADER AL-MAKTRY AND SUSAN F. EL-DEKEN

ABSTRACT

Let R be a finite non-commutative ring with $1 \neq 0$. By a polynomial function on R , we mean a function $F: R \rightarrow R$ induced by a polynomial $f = \sum_{i=0}^n a_i x^i \in R[x]$ via right substitution of the variable x , i.e. $F(a) = f(a) = \sum_{i=0}^n a_i a^i$ for every $a \in R$. In this paper, we study the polynomial functions of the free R -algebra with a central basis $\{1, \beta_1, \dots, \beta_k\}$ ($k \geq 1$) such that $\beta_i \beta_j = 0$ for every $1 \leq i, j \leq k$, $R[\beta_1, \dots, \beta_k]$. Our investigation revolves around assigning a polynomial $\lambda_f(y, z)$ over R in non-commuting variables y and z to each polynomial f in $R[x]$; and describing the polynomial functions on $R[\beta_1, \dots, \beta_k]$ through the polynomial functions induced on R by polynomials in $R[x]$ and by their assigned polynomials in the non-commuting variables y and z . By extending results from the commutative case to the non-commutative scenario, we demonstrate that several properties and theorems in the commutative case can be generalized to the non-commutative setting with appropriate adjustments.

1. INTRODUCTION

Let R denote a finite ring with identity $1 \neq 0$. By a polynomial function on R , we mean a right-polynomial function, which is a function $F: R \rightarrow R$ induced by a polynomial $f = \sum_{i=0}^n a_i x^i \in R[x]$ via right substitution of the variable x , i.e., $F(a) = f(a) = \sum_{i=0}^n a_i a^i$ for every $a \in R$. In particular, if F is a bijection, we call F a *polynomial permutation* and f a *permutation polynomial* on R . We denote the set of polynomial functions on R by $\mathcal{F}(R)$, and by $\mathcal{P}(R)$ we denote the set of all polynomial permutations on R . When, R is a finite commutative ring, the set $\mathcal{F}(R)$ is known to be a monoid under the composition of functions. Further, the group of units of $\mathcal{F}(R)$ is just the set $\mathcal{P}(R)$. While the case that R is a non-commutative is difficult and still open, we however can ensure that the set $\mathcal{F}(R)$ and the set $\mathcal{P}(R)$ generate a monoid and a group, respectively, of functions on R .

Even though polynomial functions on finite fields have been investigated before the 20th century, polynomial functions on finite rings that are not fields were first studied at the beginning of the third decade of the 20th century by Kempner [28]. Kempner intensely studied the polynomial functions of the ring \mathbb{Z}_m of integers modulo m . Perhaps the only defect of his work was a lack of simplicity, but his work nevertheless had a significant impact on other

2010 *Mathematics Subject Classification.* Primary 13F20 16P10; Secondary 05A05, 06B10, 20B35, 11C99.

Key words and phrases. Finite non-commutative rings, dual numbers, polynomials, polynomial functions, polynomial permutations, permutation polynomials, null polynomials, finite polynomial permutation groups.

researchers for decades. For instance, some researchers followed up on his results, derived equivalent relations, and added new ideas to the topic as well [38, 27, 44, 34]. Others generalized his results into Galois rings [11], finite commutative principal rings [35], and the so-called suitable rings [17]. In most of the previously mentioned works, the authors tackled the problems of counting the number of elements in the monoid $\mathcal{F}(R)$ and in the group $\mathcal{P}(R)$, and finding canonical representations of the polynomial functions.

Recently, authors have provided algorithms determining whether or not a given function is a polynomial function and if so, it obtains its polynomial representative on the ring integers modulo m [24] and on any finite commutative rings [13].

To investigate polynomial functions efficiently on finite rings, one needs some essential knowledge of those polynomials inducing the zero function on the ring R . We refer to these polynomials as null polynomials on R . These polynomials are of great interest in this investigation. Because, for example, one can represent all polynomial functions by polynomials of degrees not exceeding $n - 1$ whenever a monic null polynomial of degree n is given. Another reason is the fact that null polynomials form an ideal of the polynomial ring over a quite large class of finite rings including commutative rings, rings of matrices over commutative rings, and local rings (see Section 2). Because of this, some papers mainly considered null polynomials (see for example [21, 6, 42]).

Beyond the significant impact of Kempener's work in the theory, the work of Nöbauer on polynomial permutations and permutation polynomials can not be underestimated. We can not here mention his enormous results in the theory and rather we give a limited brief description of his work and refer the interested reader to his book with Lausch [33]. For instance, he obtained a condition determining whether a given polynomial induces a permutation of the ring \mathbb{Z}_{p^n} of integers modulo a prime power ([37, Hilfssatz 8]), and later he generalized this condition into any finite commutative rings [39]. Also, he expressed the group $\mathcal{P}(\mathbb{Z}_{p^n})$ as the wreath product of two of its subgroups [38]. This structural result was first generalized into finite commutative local rings of nilpotency index 2 by Frisch [17] and finally into any finite local ring by Görcsös, Horváth and Mészáros [22]. Further, some other papers considered the structures of subgroups of the group $\mathcal{P}(R)$ for the case $R = \mathbb{Z}_{p^n}$ [20] or for any finite commutative ring [2].

Despite tackling various aspects related to polynomial functions on finite rings, all rings examined in the previously mentioned references are commutative. Unlikely, polynomial functions on finite non-commutative only arose in special circumstances to answer specified questions. Indeed, most of the theory considered only scalar polynomial functions on the ring of $n \times n$ matrices over a finite field, $M_n(\mathbb{F}_q)$, (see for example [10, 12]), or over a finite commutative local ring [9] for particular considerations. For instance, they considered scalar polynomial functions that permute the entire ring of matrices [9]; or that permute a subset of $M_n(\mathbb{F}_q)$ [48]. Additionally, in [10] and [8] the number of all scalar polynomial functions and the the number of all scalar polynomial permutations, respectively, on the ring $M_n(\mathbb{F}_q)$ have been counted. Meanwhile in [50, 18], the authors focused on null polynomials and their connection with the ring of integer-valued polynomials of certain non-commutative algebras.

To the best of our knowledge, the only reference containing counting formulas for the number of polynomial functions and the number of polynomial permutations on finite non-commutative

rings is [32], where the author counted the elements of $\mathcal{F}(R)$ and $\mathcal{P}(R)$ of any finite non-commutative chain rings (Galois-Eisenstein-Ore ring) of characteristic p .

In this paper, we delve into the realm of polynomial functions within non-commutative rings. we aim namely to explore the polynomial functions on the non-commutative free R -algebra $R_k = R[\beta_1, \dots, \beta_k]$, where β_i lies in the center $C(R_k)$ of R_k and $\beta_i \beta_j = 0$ for $i, j = 1, \dots, k$. Because this ring can be expressed as the quotient ring $R[x_1, \dots, x_k]/T$, where T is the ideal generated by the set $\{x_i x_j \mid i, j = 1, \dots, k\}$, we call it the ring of dual numbers in k variables. When R is commutative, exploring $\mathcal{F}(R_k)$ has been done in two stages. Initially, in [1], only the case $k = 1$ was examined with more consideration on the ring of integers modulo p^n . Then in [3], $\mathcal{F}(R_k)$ was examined for any positive integer k and any finite commutative ring R . In general, these examinations were carried out not only by examining the polynomial functions obtained by polynomials in $R[x]$ but also by examining the polynomial functions obtained by their formal derivatives.

We will show that most of the results of [3] are still true, and losing an important property like commutativity does not affect the results. Furthermore, given a polynomial $f \in R[x]$, we assign a new polynomial $\lambda_f(z, y)$ in non-commuting variables y and z (see Definition 3.5 for detail); and similar to the commutative case, investigating $\mathcal{F}(R_k)$ will depend not on the polynomials from $R[x]$ but also on their assigned polynomials in the non-commutative variables y and z . In other words, conditions derived on f and its formal derivative f' in [3] are revisited here by general conditions on f and $\lambda_f(z, y)$ (compare for example Theorem 4.4 with [3, Theorem 3.4], and Theorem 5.4 with [3, Theorem 4.1]). Furthermore, we also consider the case where R is a chain ring with $Char(R) \neq p$ in some detail.

The influences of polynomial functions can be seen in different topics for example dynamical systems [5], computer science (see for example [46, 47]) and in discrete mathematics [29]. Further, beyond these applications, polynomial functions appear in different contexts of mathematics, for instance, permutation polynomials occur as an R -automorphisms of the polynomial ring (see for example [25, 26]) and as isomorphisms of combinatorial objects [7]. Also, some groups of polynomial permutations occur as subgroups of the permutation groups of cyclic codes [23]. Moreover, null polynomials are efficiently utilized within investigating several rings of integer-valued polynomials (see for example [40, 43]). In our opinion, therefore, exploring polynomial functions on new classes of non-commutative rings opens new gates for applications.

It is worth mentioning that some mathematicians have explored polynomial functions on other algebraic structures such as semi groups [31], monoids [49], groups [30], algebras[16] and Lattices [15].

Here is a brief description of the paper. In Section 2, we review basic definitions and facts about polynomial functions on finite non-commutative rings. Section 3 contains the essential properties of the ring of dual numbers in k variables, R_k , and their polynomials. In Section 4, we describe null polynomials on R_k . In Section 5, we characterize permutation polynomials on R_k . Then in its subsection, we examine permutation polynomials on R_k for the case R is a chain ring. Finally, we devote Section 6 to investigate some groups generated by permutation polynomials on R_k with more consideration on those generated by permutations represented by polynomials over R .

2. BASIC FACTS ON POLYNOMIAL FUNCTIONS

In this section, we expose some basic definitions and facts about polynomial functions on finite non-commutative rings.

Throughout this paper, by g' , we denote the formal derivative of the polynomial g ; and let k denote a positive integer and let \mathbb{F}_q denote the finite field of q elements.

Definition 2.1. Let R be a non-commutative ring, and $g = \sum_{i=0}^n a_i x^i \in R[x]$. Then:

- (1) The polynomial g induces two functions $F_1, F_2: R \rightarrow R$ by right substitution $F_1(a) = f_r(a) = \sum_{i=0}^n a_i a^i$ and left substitution $F_2(a) = f_l(a) = \sum_{i=0}^n a^i a_i$ substitution, for the variable x . We call F_1 a right-polynomial function on R and F_2 a left polynomial function on R .
- (2) By $[g_r]_R$, we denote the right-polynomial function induced by g on R , and by $[g_l]_R$, we denote the left-polynomial function induced by g on R . When the ring is understood, we write $[g_r]$ and $[g_l]$ respectively.
- (3) If $f \in R[x]$ such that f and g induce the same right-function (left-function) on R , i.e. $[g_r] = [f_r]$ ($[g_l] = [f_l]$), then we abbreviate this with $g \triangleq_r f$ ($g \triangleq_l f$) on R .
- (4) We define $\mathcal{F}(R)_r = \{[g_r] \mid g \in R[x]\}$.

$$\mathcal{P}(R)_r = \{[f_r] \mid [f_r] \text{ is a permutation of } R \text{ and } f \in R[x]\}.$$

In similar manner, we define $\mathcal{F}(R)_l$ and $\mathcal{P}(R)_l$.

- (5) If A is a subring of R , and $f \in A[x]$, then f induces two polynomial functions on R and A respectively. To distinguish between them we write $[f]_R$ and $[f]_A$.

It is clear that $\mathcal{F}(R)_r$ ($\mathcal{F}(R)_l$) is an additive group with respect to pointwise addition. However, unlike commutative rings, we can not define pointwise multiplication on the sets $\mathcal{F}(R)_r$ and $\mathcal{F}(R)_l$ since substitution in general is not a homomorphism. Indeed, one may find $f, g \in R[x]$ and $r_1, r_2 \in R$ such that

$$f_r(r_1)g_r(r_1) \neq h_r(r_1) \text{ and } f_l(r_2)g_l(r_2) \neq h_l(r_2),$$

where $h = fg$. In fact, if we write $f(x) = \sum_j a_j x^j$ and $g(x) = \sum_i x^i b_i$, then we can express

h as $h(x) = \sum_j a_j g(x)x^j$ and $h(x) = \sum_i x^i f(x)b_i$. Thus substitutions in h from right and left are respectively given by

$$h_r(c) = \sum_j a_j g_r(c)c^j, \tag{1}$$

and

$$h_l(c) = \sum_i c^i f_l(c)b_i. \tag{2}$$

From now on we only consider the right substitution polynomial functions. By symmetry, all the properties of right-polynomial functions can be drawn into left-polynomial functions with some suitable changes. We will not use the term right and will delete the subscript r indicating it from the notation.

We mention here that if g is central (i.e. in the center of $R[x]$), then $(fg)(r) = f(r)g(r)$ for every $r \in R$. Also, if f is central, then $(fg)(r) = g(r)f(r)$ for every $r \in R$. This can be concluded easily by following definition in which we summarize the relation between the multiplication of two polynomials and its value on R .

Definition 2.2. Let $f, g \in R[x]$. Let $f(x) = \sum_{j=0}^n a_j x^j$. Then

$$(1) (fg)(x) = \sum_{j=0}^n a_j g(x) x^j;$$

$$(2) (fg)(r) = \sum_{j=0}^n a_j g(r) r^j \text{ for every } r \in R.$$

Remark 2.3. The previously defined multiplication of polynomials is the usual multiplication of polynomials which is the same as in the commutative case except that the coefficients are not commutable, and therefore it is associative. But, let us show this by using Definition 2.2.

So, if $f, g, h \in R[x]$, then write $f(x) = \sum_{j=0}^n a_j x^j$ and $g(x) = \sum_{i=0}^m b_i x^i$. Now,

$$(fg)(x) = \sum_{j=0}^n a_j g(x) x^j = \sum_{j=0}^n a_j \sum_{i=0}^m b_i x^i x^j = \sum_{j=0}^n \sum_{i=0}^m a_j b_i x^{i+j}.$$

Therefore

$$((fg)h)(x) = \sum_{j=0}^n \sum_{i=0}^m a_j b_i h(x) x^{i+j} = \sum_{j=0}^n a_j \left(\sum_{i=0}^m b_i h(x) x^i \right) x^j = \sum_{j=0}^n a_j (gh)(x) x^j = (f(gh))(x).$$

This also shows that $((fg)h)(r) = (f(gh))(r)$ for every $r \in R$.

Polynomials that map every element to zero (i.e. induce the zero function) appear widely within investigating polynomial functions, and for this we devote the following definition.

Definition 2.4.

- (1) A polynomial $f \in R[x]$ is called a null polynomial on R if $f(r) = 0$ for every $r \in R$. In this case, we write $f \triangleq 0$ on R .
- (2) We define the set N_R as:
 - (a) $N_R = \{f \in R[x] \mid f \triangleq 0 \text{ on } R\}$.

From Definitions 2.4 and 2.2, one sees without effort the following result.

Corollary 2.5. Let R be a non-commutative ring and let N_R be the set of all null polynomials on R . Then

- (1) N_R is a left ideal of $R[x]$;
- (2) N_R is an ideal of $R[x]$ if and only if N_R is an R -right module.

Remark 2.6. For a ring A , the left ideal N_A defines an equivalence relation “ $\equiv \pmod{N_A}$ ” as the following, for any $f, g \in A[x]$, $f \equiv g \pmod{N_A}$ if and only if $f - g \in N_A$ which is equivalent to $f \triangleq g$ on A . Thus, the relation “ \triangleq on A ” is the same equivalence relation on $A[x]$ defined

by N_A . Also, there is a one-to-correspondence between the equivalence classes of \cong and the polynomial functions on A . Therefore, whenever A is finite, the following equality holds

$$|\mathcal{F}(A)| = [R[x] : N_A]$$

Throughout whenever A is a ring let $J(A)$ denote its Jacobson radical.

Remark 2.7. Unlike the commutative case, we still do not generally know whether N_R is an ideal (i.e. two-sided ideal). However, Werner showed that N_R is an ideal over finite non-commutative rings in which every element can be written as a sum of units (for example semisimple rings and local rings) [50, Theore 3.7]. A result of Stewart [45, Lemm 4.5 and Theorem 4.6] (see also [50, Propostion 3.6]) infers that every element of a finite ring A is a sum of units if only if $\mathbb{F}_2 \times \mathbb{F}_2$ is not a homomorphic image of the ring $A/J(A)$.

Furthermore, Frisch showed that when R is the ring of upper triangular (lower) over commutative ring A , N_R is an ideal [18, Theroem 5.2]. In particular, if R is the non-commutative ring $UT_n(\mathbb{F}_2)$ of all upper triangular matrices with entries in \mathbb{F}_2 of dimension $n \geq 2$, then N_R is an ideal by Frisch result. However, not every element of $UT_n(\mathbb{F}_2)$ is a sum of units. For example, the elementary matrix E_{11} (i.e the matrix with entry 1 in the first position and zero elsewhere) is not a sum of units, because any sum of units in the ring $UT_n(\mathbb{F}_2)$ gives a matrix whose diagonal elements are all equal to zero or all equal to one depends on the number of units in that sum. This shows the previously mentioned condition of Werner is only sufficient but not necessary.

Although we remarked earlier that substitution is not a ring homomorphism from $R[x]$ onto the set of polynomial functions $\mathcal{F}(R)$, as we can not endorse the set $\mathcal{F}(R)$ with pointwise multiplication, we can define multiplication on it utilizing Definition 2.2 whenever N_R is an ideal.

Proposition 2.8. Let R be a finite non-commutative ring. Define an operation “ \cdot ” on $\mathcal{F}(R)$ by letting $F \cdot F_1 = [fg]$, where $f, g \in R[x]$ such that $F = [f]$ and $[g] = F_1$. Then “ \cdot ” is well defined if and only if N_R is a two sided ideal; in this case $\mathcal{F}(R)$ is a ring endowed with multiplication “ \cdot ” and pointwise addition.

Proof. First, we notice by Definition 2.2 that, $[fg] \in \mathcal{F}(R)$ for every $f, g \in R[x]$.

(\Leftarrow) Assume that N_R is an ideal of $R[x]$. Let $F, F_1 \in \mathcal{F}(R)$ and let $f, g \in R[x]$ such that $F = [f]$ and $F_1 = [g]$. To show that $F \cdot F_1$ is well defined, we need to show that $[fg] = [f_1g_1]$ for every $f_1, g_1 \in R[x]$ such that $F = [f_1]$ and $F_1 = [g_1]$. Now, if $f_1, g_1 \in R[x]$ such that $[f] = [f_1]$ and $[g] = [g_1]$, then $f(r) = f_1(r)$ and $g(r) = g_1(r)$ for every $r \in R$. More precisely, the polynomial $h = f - f_1$ is null on R , whence $hg \in N_R$ since N_R is an ideal. Now, writing $f = \sum_{j=0}^n a_j x^j$ and $f_1 = \sum_{j=0}^n b_j x^j$ yields in view of Definition 2.2, for every $r \in R$

$$(fg)(r) - (f_1g_1)(r) = \sum_{j=0}^n a_j g(r) r^j - \sum_{j=0}^n b_j g_1(r) r^j = \sum_{j=0}^n (a_j - b_j) g(r) r^j = (hg)(r) = 0.$$

Thus, $(fg)(r) = (f_1g_1)(r)$ for each r . But, this means $[fg] = [f_1g_1]$.

(\Rightarrow) Assume that “ \cdot ” is well defined. In the light of Corollary 2.5, we need only show that N_R is an R -right module. So, let $f \in N_R$ and let $r \in R$. Then $f \triangleq 0$ (the zero polynomial) on R or equivalently, $[f] = 0$ (the zero function). Thus, if F is the constant polynomial function r , then $[fr] = 0 \cdot F = [0r] = 0$ since “ \cdot ” is well defined. Hence, $fr \triangleq 0$ and $fr \in N_R$. For the other part, one can easily see that $\mathcal{F}(R)$ is an additive group with respect to pointwise addition. We only check that “ \cdot ” is associative and leave other properties to the reader. Let $F_1, F_2, F_3 \in \mathcal{F}(R)$, to show that $(F_1 \cdot F_2) \cdot F_3 = F_1 \cdot (F_2 \cdot F_3)$ it is enough to see show that $(fg)h = f(gh)$, where f, g and h any polynomials induce F_1, F_2 and F_3 respectively. But this what we have already shown that in Remark 2.3. \square

Remark 2.9. Unfortunately, composition of polynomial functions is not compatible with composition of polynomials on non-commutative rings, that is, one can find two polynomials $f, g \in R[x]$ such that $[f] \circ [g] \neq [f \circ g]$, that is, there exists an $r \in R$ such that $(f \circ g)(r) \neq f(g(r))$. This is because, composition depends on multiplication and we already remarked that substitution is not a homomorphism within multiplication of polynomials. For instance, take R to be the ring of 2 by 2 matrices over the field \mathbb{F}_2 , and let $a = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ and $b = \begin{bmatrix} 0 & 1 \\ 0 & 1 \end{bmatrix}$. For $f(x) = x^2$ and $g(x) = ax$, we have the polynomial $h(x) = f \circ g(x) = a^2 x^2 = ax^2$. Then

$$h(b) = ab^2 = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \neq 0 = f(g(b)).$$

Therefore, $[f] \circ [g] \neq [f \circ g]$.

Because of this dis-compatibility of the composition of polynomial functions and the composition of polynomials, we can neither ensure that $\mathcal{F}(R)$ is a monoid nor that $\mathcal{P}(R)$ is a group. However, we will see that $\mathcal{F}(R)$ generates a monoid of functions on R and that $\mathcal{P}(R)$ generates a group of permutations on R . For this we need the following definition.

Definition 2.10. Let A be a non-empty set with associative operation $*$. For every non-empty subset B of A , we define its closure \overline{B} in A by

$$\overline{B} = \{a_1 * a_2 * \cdots * a_n : n \in \mathbb{N} \text{ and } a_i \in B \text{ for } i = 1, \dots, n\}.$$

In view of Definition 2.10, one easily show the following useful fact.

Fact 2.11. Let A be a non-empty set with associative operation $*$ and let B be a non-empty subset of A .

- (1) If $(A, “*”)$ is a monoid and $1_A \in B$, then \overline{B} is a submonoid of A .
- (2) If $(A, “*”)$ is a finite group, then \overline{B} is a subgroup of A .

Now, considering $\mathcal{F}(R)$ as a subset of the monoid of all functions on R and $\mathcal{P}(R)$ as a subset of the symmetric group S_R of all permutations on R together with Fact 2.10 yields the following.

Corollary 2.12. Let R be a finite non-commutative ring and “ \circ ” denote the composition of functions. Then

- (1) $(\overline{\mathcal{F}(R)}, “\circ”)$ is a finite monoid;
- (2) $(\overline{\mathcal{P}(R)}, “\circ”)$ is a finite group.

From now on we call the group (monoid) \overline{B} the closure group (monoid) of B . We conclude this section by indicating that the material in this section is known (see [19, 50]) except Proposition 2.8 which we do not know any reference containing it.

3. DUAL NUMBERS OVER FINITE NON-COMMUTATIVE RINGS

In this section, we expose some elementary properties of the ring of dual numbers in k variables and its polynomial ring. Most, of these properties play an essential role in the arguments of the proven facts through this paper. We start by giving a constructive definition of this ring.

Definition 3.1. Let R be a non-commutative ring and let T be the ideal of the polynomial ring $R[x_1, \dots, x_k]$ generated by the set $\{x_i x_j \mid i, j \in \{1, \dots, k\}\}$. We call the quotient ring $R[x_1, \dots, x_k]/T$ the ring of dual numbers of k variables over R . We write $R[\beta_1, \dots, \beta_k]$ for $R[x_1, \dots, x_k]/T$, where β_i denotes $x_i + T$.

Remark 3.2. Note that every element of $R[x_1, \dots, x_k]/T$ has a unique representation as an R -linear combination of $1, \beta_1, \dots, \beta_k$. That is, $R[\beta_1, \dots, \beta_k]$ is a free R -algebra with basis $\{1, \beta_1, \dots, \beta_k\}$. We call the coefficient of 1 the “constant coefficient”. It follows then that the polynomial ring $R[\beta_1, \dots, \beta_k][x]$ is a free $R[x]$ -algebra with the same basis $\{1, \beta_1, \dots, \beta_k\}$. Also, R is canonically embedded as a subring in $R[\beta_1, \dots, \beta_k]$ by $r \rightarrow r \cdot 1$, and we have

$$R[\beta_1, \dots, \beta_k] = \{r_0 + \sum_{i=1}^k r_i \beta_i \mid r_0, r_i \in R, \text{ with } \beta_i \beta_j = 0 \text{ for } 1 \leq i, j \leq k\}.$$

It follows from this that every polynomial $f \in R[\beta_1, \dots, \beta_k][x]$ has a unique representation $f = f_0 + \sum_{i=1}^k f_i \beta_i$, where $f_0, f_1, \dots, f_k \in R[x]$.

Definition 3.3. Let $f \in R[\beta_1, \dots, \beta_k][x]$ and let $f_0, f_1, \dots, f_k \in R[x]$ be the unique polynomials such that $f = f_0 + \sum_{i=1}^k f_i \beta_i$. Then we call the polynomial f_0 the pure part of f .

The following proposition summarizes some properties of $R[\beta_1, \dots, \beta_k]$ whose proof is immediate from Definition 3.1.

Proposition 3.4. Let R be a non-commutative ring. Then the following statements hold.

(1) For $a_0, \dots, a_k, b_0, \dots, b_k \in R$, we have:

(a) $(a_0 + \sum_{i=1}^k a_i \beta_i)(b_0 + \sum_{i=1}^k b_i \beta_i) = a_0 b_0 + \sum_{i=1}^k (a_0 b_i + a_i b_0) \beta_i;$

(b) $a_0 + \sum_{i=1}^k a_i \beta_i$ is a unit in $R[\beta_1, \dots, \beta_k]$ if and only if a_0 is a unit in R . In this case,

$$(a_0 + \sum_{i=1}^k a_i \beta_i)^{-1} = a_0^{-1} - \sum_{i=1}^k a_0^{-1} a_i a_0^{-1} \beta_i.$$

(2) $R[\beta_1, \dots, \beta_k]$ is a local ring if and only if R is a local ring.

(3) If R is a ring with Jacobson radical $J(R)$ of nilpotency n , then

$$J(R[\beta_1, \dots, \beta_k]) = J(R) + \sum_{i=1}^k \beta_i R \text{ of nilpotency } n + 1.$$

(4) If R is a ring with center $C(R)$, then

- (a) $C(R[\beta_1, \dots, \beta_k]) = C(R) + \sum_{i=1}^k C(R) \beta_i$;
- (b) $C(R)[x] \subseteq C(R[\beta_1, \dots, \beta_k])[x]$.

Unlike the commutative case we can not obtain the same relation as in [3, Lemma 2.6] for the substitution in polynomials. The reason is that the binomial theorem does not hold in non-commutative rings and we need a more general form. To overwhelm this vital obstacle we first introduce the following definition.

Definition 3.5. Let $f = \sum_{j=0}^n a_j x^j \in R[x]$. Then we assign to f a unique polynomial $\lambda_f(y, z)$ in the non-commutative variables y and z defined by

$$\lambda_f(y, z) = \sum_{j=1}^n a_j m_j(y, z), \text{ where } m_j(y, z) = \sum_{r=1}^j y^{r-1} z y^{j-r}.$$

We call λ_f the assigned polynomial of (to) f with respect to two- non-commutative variables, or just the assigned polynomial.

Definition 3.6. Let $f = \sum_{j=0}^n a_j x^j \in R[x]$ and $\lambda_f(y, z)$ be its assigned polynomial in the non-commutative variables y and z . For every $a \in R$, we define the polynomial $\lambda_f(a, z)$ by

$$\lambda_f(a, z) = \sum_{j=1}^n a_j m_j(a, z) = \sum_{j=1}^n \sum_{r=1}^j a_j a^{r-1} z a^{j-r}.$$

Definition 3.7. Let $f = \sum_{j=0}^n a_j x^j \in R[x]$ and $\lambda_f(y, z)$ be its assigned polynomial in the non-commutative variables y and z . Then

- (1) the polynomial λ_f induces (defines) a function $F: R \times R \rightarrow R$ by substituting the variables

$$F(a, b) = \lambda_f(a, b) = \sum_{j=1}^n a_j m_j(a, b) = \sum_{j=1}^n \sum_{r=1}^j a_j a^{r-1} b a^{j-r},$$

which we denote by $[\lambda_f(y, z)]$;

- (2) for every $a \in R$ the polynomial $\lambda_f(a, z)$ defines a function $F_a: R \rightarrow R$ by substituting the variable $F_a(b) = \lambda_f(a, b)$, which we denote by $[\lambda_f(a, z)]$.

Remark 3.8. Let $f = \sum_{j=0}^n a_j x^j \in R[x]$. Then

- (1) the polynomial $m_j(y, z) = \sum_{r=1}^j y^{r-1} z y^{j-r}$ appeared in Definition 3.5 is independent from the polynomial f .
- (2) In similar manner, we can define for every $a \in R$ the polynomial $\lambda_f(y, a)$ by

$$\lambda_f(y, a) = \sum_{j=1}^n a_j m_j(y, a) = \sum_{j=1}^n \sum_{r=1}^j a_j y^{r-1} a y^{j-r}.$$

However, in our text, this type of polynomials is less important than the polynomial $\lambda_f(a, z)$ of Definition 3.6.

(3) For every $a, b \in R$ such that $ab = ba$, one easily sees that

$$\lambda_f(a, b) = \sum_{j=1}^n a_j m_j(a, b) = \sum_{j=1}^n j a_j a^{j-1} b = f'(a)b.$$

Here are straightforward properties of the assigned polynomial which are useful through the coming proofs.

Fact 3.9. Let $r, s, w \in R$. Let f and $g \in R[x]$, and let λ_f and λ_g be their assigned polynomials respectively. Then

- (1) $\lambda_{rf+sg} = \lambda_{rf} + \lambda_{sg}$;
- (2) $\lambda_{fr+gs} = \lambda_{fr} + \lambda_{gs}$;
- (3) $\lambda_f(y, z) = 0$ if and only if f is constant;
- (4) $\lambda_f(y, z) = z$ if and only if $f = x$;
- (5) $\lambda_f(0, z) = a_1 z$ and $\lambda_f(y, 1) = f'(y)$, where $f(x) = \sum_{j=0}^n a_j x^j$;
- (6) $\lambda_f(y, 0) = 0$;
- (7) $\lambda_f(r, s \pm w) = \lambda_f(r, s) \pm \lambda_f(r, w)$.

Now, we are able to state and prove a general form of [3, Lemma 2.6], which will play the same role later in our context.

Lemma 3.10. Let R be a ring and $a, b_1, \dots, b_k \in R$.

(1) If $f \in R[x]$ and λ_f is its assigned polynomial then

$$f\left(a + \sum_{i=1}^k b_i \beta_i\right) = f(a) + \sum_{i=1}^k \lambda_f(a, b_i) \beta_i.$$

(2) If $f \in R[\beta_1, \dots, \beta_k][x]$ and f_0, \dots, f_k are the unique polynomials in $R[x]$ such that

$$f = f_0 + \sum_{i=1}^k f_i \beta_i, \text{ then}$$

$$f\left(a + \sum_{i=1}^k b_i \beta_i\right) = f_0(a) + \sum_{i=1}^k (\lambda_{f_0}(a, b_i) + f_i(a)) \beta_i,$$

where λ_{f_0} is the assigned polynomial to f_0 .

Proof. (1) By induction and the fact that $\beta_i \beta_j = 0$ for $1 \leq i, j \leq k$, we have that for every $a, b_1, \dots, b_k \in R$,

$$\left(a + \sum_{i=1}^k b_i \beta_i\right)^j = a^j + \sum_{i=1}^k \sum_{r=1}^j a^{r-1} b_i a^{j-r} \beta_i = a^j + \sum_{i=1}^k m_j(a, b_i) \beta_i, \quad (3)$$

where $m_j(a, b_i) = \sum_{r=1}^j a^{r-1} b_i a^{j-r}$.

Now, writing $f = \sum_{j=0}^n a_j x^j \in R[x]$ yields, by identity (3) and Definition 3.7,

$$f\left(a + \sum_{i=1}^k b_i \beta_i\right) = \sum_{j=1}^n \left(a + \sum_{i=1}^k b_i \beta_i\right)^j + a_0 = \sum_{j=0}^n a_j a^j + \sum_{j=1}^n a_j \sum_{i=1}^k m_j(a, b_i) \beta_i = f(a) + \sum_{i=1}^k \lambda_f(a, b_i) \beta_i.$$

(2) Follows from (1). \square

Notation 3.11. From now on, let R_k denote the ring $R[\beta_1, \dots, \beta_k]$.

A consequence of Lemma 3.10 is the following result.

Corollary 3.12. Let $F: R_k \rightarrow R_k$ be a polynomial function and let $a, b_1, \dots, b_k \in R$. Then:

- (1) The constant coefficient of $F\left(a + \sum_{i=1}^k a_i \beta_i\right)$ depends only on a ;
- (2) The coefficient of β_i in $F\left(a + \sum_{i=1}^k b_i \beta_i\right)$ depends only on a and b_i .

Remark 3.13. Let σ be a permutation of $1, \dots, k$. Let $a_0, \dots, a_k \in R$ and let $f \in R[x]$ such that $f\left(a_0 + \sum_{i=1}^k a_i \beta_i\right) = b_0 + \sum_{i=1}^k b_i \beta_i$, where $b_0, \dots, b_k \in R$. Then, in view of Lemma 3.10 and Corollary 3.12, we easily see that:

- (1) $f\left(a_0 + \sum_{i=1}^k a_i \beta_{\sigma(i)}\right) = b_0 + \sum_{i=1}^k b_i \beta_{\sigma(i)}$;
- (2) $f\left(a_0 + \sum_{i=1}^k a_{\sigma(i)} \beta_i\right) = b_0 + \sum_{i=1}^k b_{\sigma(i)} \beta_i$;
- (3) $f\left(a_0 + a_i \beta_j\right) = b_0 + b_i \beta_j$ for $i = 1, \dots, k$ and $j \geq 1$.

Now, if $f_1, \dots, f_n \in R[x]$ such that

$[f_n]_{R_k} \circ \dots \circ [f_1]_{R_k}\left(a_0 + \sum_{i=1}^k a_i \beta_i\right) = f_n(\dots(f_1(a_0 + \sum_{i=1}^k a_i \beta_i))\dots) = c_0 + \sum_{i=1}^k c_i \beta_i$, where $c_0, \dots, c_k \in R$. Then inductively, we have that:

- (a) $[f_n]_{R_k} \circ \dots \circ [f_1]_{R_k}\left(a_0 + \sum_{i=1}^k a_i \beta_{\sigma(i)}\right) = c_0 + \sum_{i=1}^k c_i \beta_{\sigma(i)}$;
- (b) $[f_n]_{R_k} \circ \dots \circ [f_1]_{R_k}\left(a_0 + \sum_{i=1}^k a_{\sigma(i)} \beta_i\right) = c_0 + \sum_{i=1}^k c_{\sigma(i)} \beta_i$;
- (c) $[f_n]_{R_m} \circ \dots \circ [f_1]_{R_m}\left(a_0 + a_i \beta_j\right) = c_0 + c_i \beta_j$ ($m = \max(i, j)$) for $i = 1, \dots, k$ and $j \geq 1$.

In the following $F|_A$ stands for the restriction of the function F to A .

Corollary 3.14. Let $k > 1$ and $F, G \in \overline{\mathcal{F}(R_k)}$. Suppose that $F = [f_n]_{R_k} \circ \dots \circ [f_1]$ and $G = [g_m]_{R_k} \circ \dots \circ [g_1]$, where $f_1, \dots, f_n, g_1, \dots, g_m \in R[x]$. Then the following statements are equivalent

- (1) $F = G$;
- (2) $F|_{R_1} = G|_{R_1}$;
- (3) $F|_{R_j} = G|_{R_j}$ for $j = 1, \dots, k-1$.

Proof. The implications (1) \Rightarrow (2), (1) \Rightarrow (2) and (3) \Rightarrow (2) are obvious.

(2) \Rightarrow (1): assume that $F|_{R_1} = G|_{R_1}$. Let $a_0, \dots, a_k \in R$ be arbitrary. We want to show that

$$F(a_0 + \sum_{i=1}^k a_i \beta_i) = G(a_0 + \sum_{i=1}^k a_i \beta_i).$$

Suppose that $F(a_0 + \sum_{i=1}^k a_i \beta_i) = b_0 + \sum_{i=1}^k b_i \beta_i$ and $G(a_0 + \sum_{i=1}^k a_i \beta_i) = c_0 + \sum_{i=1}^k c_i \beta_i$, where $b_0, \dots, b_k, c_0, \dots, c_k \in R$.

By Remark 3.13, $F(a_0 + a_i \beta_1) = b_0 + b_i \beta_1$ and $G(a_0 + a_i \beta_1) = c_0 + c_i \beta_1$ for $i = 1, \dots, k$. But, by assumption, $F|_{R_1} = G|_{R_1}$. Therefore,

$$b_0 + b_i \beta_1 = c_0 + c_i \beta_1 \text{ for } i = 1, \dots, k.$$

Thus, $b_i = c_i$ for $i = 0, \dots, k$, whence $F = G$. \square

4. POLYNOMIAL FUNCTIONS ON R_k

In this section, we characterize null polynomials on the ring R_k . This characterization allows us to determine whether two polynomials are equivalent on R_k , that is whether they represent the same polynomial function on R_k . Finally, we give a counting formula of the polynomial functions of the ring R_k by mean of the indices of the left ideal N_R (see Definition 2.4) and a related left ideal AN_R defined below.

Next, we define a subset of N_R concerning the assigned polynomials of Definition 3.5.

Definition 4.1.

- (1) Let $f \in R[x]$ and let λ_f be its assigned polynomial. We call λ_f is null if $\lambda_f(a, b) = 0$ for every $a, b \in R$. We write $[\lambda_f(y, z)] = 0$.
- (2) We define AN_R as: $AN_R = \{f \in N_R \mid [\lambda_f(y, z)] = 0\}$.
- (3) We define N'_R as: $N'_R = \{f \in N_R \mid f' \in N_R\}$.

Remark 4.2.

- (1) It is obvious that AN_R and N'_R are left ideals of $R[x]$ with $AN_R, N'_R \subseteq N_R$.
- (2) Let $f \in AN_R$. Then the condition $[\lambda_f(y, z)] = 0$ implies that $\lambda_f(a, 1) = f'(a) = 0$ for every $a \in R$ (by Fact 3.9). Hence $f \in N'_R$. Thus, we have the following inclusion

$$AN_R \subseteq N'_R \subseteq N_R.$$

- (3) When R is commutative the condition on λ_f in the definition of AN_R is equivalent to $f' \in N_R$. So, $AN_R = N'_R$ over commutative ring R . However, when R is a non-commutative ring, it can be happened that N'_R contains AN_R properly (see Example 4.13). We will see that the left ideal AN_R plays the same role as N'_R in the context of commutative rings. More precisely, we will show that most of the results of [1, 3] involving N'_R still hold here in the context of non-commutative rings by replacing N'_R by AN_R .

Lemma 4.3. *Let $g \in R[x]$. Then:*

- (1) $g \in N_{R_k}$ if and only if $g \in AN_R$;
- (2) $g \beta_i \in N_{R_k}$ for every $1 \leq i \leq k$ if and only if $g \in N_R$.

Proof. (1) Let $a, b_1, \dots, b_k \in R$. Then by Lemma 3.10,

$$g(a + \sum_{i=1}^k b_i \beta_i) = g(a) + \sum_{i=1}^k \lambda_g(a, b_i) \beta_i.$$

So, g is a null polynomial on R_k is equivalent to

$$g(a + \sum_{i=1}^k b_i \beta_i) = g(a) + \sum_{i=1}^k \lambda_g(a, b_i) \beta_i = 0$$

for every $a, b_1, \dots, b_k \in R$.

A fortiori, this is equivalent to $g(a) = 0$ and $\lambda_g(a, b_i) = 0$ for all $a, b_i \in R$ and $i = 1, \dots, k$. But, this is equivalent to $g \in N_R$ and $[\lambda_g(y, z)] = 0$, whence it is equivalent to $g \in AN_R$.

(2) Straightforward. \square

Theorem 4.4. *Let N_R and AN_R be as in Definition 2.4 and Definition 4.1, respectively. Then*

$$(1) N_{R_k} = AN_R + \sum_{i=1}^k N_R \beta_i;$$

(2) N_{R_k} is an ideal of $R_k[x]$ if and only if AN_R and N_R are ideals of $R[x]$.

Proof. (1) By Lemma 4.3, $AN_R + \sum_{i=1}^k N_R \beta_i \subseteq N_{R_k}$. For the other inclusion let $f \in N_{R_k}$. Then

$f = f_0 + \sum_{i=1}^k f_i \beta_i$, for some $f_0, \dots, f_k \in R[x]$. Let $r \in R$ be arbitrary. Then, by Lemma 3.10 and Fact 3.9 since $f \in N_{R_k}$,

$$0 = f(r) = f_0(r) + \sum_{i=1}^k f_i(r) \beta_i.$$

Thus, since $1, \beta_1, \dots, \beta_k$ is a base of R_k as an R -algebra, $f_i(r) = 0$ ($i = 0, \dots, k$) for every $r \in R$, which implies that $f_0, f_1, \dots, f_k \in N_R$. Now, let $a, b \in R$ be two arbitrary elements. Again by Lemma 3.10 and Fact 3.9, and what we have already shown,

$$0 = f(a + b \beta_1) = \lambda_{f_0}(a, b).$$

Thus, $\lambda_{f_0}(a, b) = 0$ for every $a, b \in R$, whence $[\lambda_{f_0}(a, b)] = 0$. Hence $f_0 \in AN_R$. This finishes the proof of the other inclusion.

(2) Keeping in mind that AN_R, N_R are left ideals of $R[x]$ and N_{R_k} is a left ideal of $R_k[x]$. (\Leftarrow) Assume that N_{R_k} is an ideal of $R_k[x]$ and let $f \in AN_R$ and $g \in N_R$. Then, by part (1), the polynomial $h = f + g \beta_1 \in N_{R_k}$. Hence, $hr = fr + gr \beta_1 \in N_{R_k}$ for every polynomial $r \in R_k[x]$ since N_{R_k} is an ideal, whence in particular, $hr \in N_{R_k}$ for every $r \in R[x]$. Thus, by the first part $fr \in AN_R$ and $gr \in N_R$ for ever $r \in R[x]$. Therefore, AN_R and N_R are ideals of $R[x]$. The other implication is somewhat similar and left to the reader. \square

Proposition 4.5. *Let R be a finite non-commutative ring and R_k be the ring dual numbers of k variables. Then the following statements are equivalent*

- (1) every element of R is a sum of units;
- (2) every element of R_k is a sum of units;
- (3) $R/J(R)$ has no factor ring isomorphic to $\mathbb{F}_2 \times \mathbb{F}_2$;

(4) $R_k/J(R_k)$ has no factor ring isomorphic to $\mathbb{F}_2 \times \mathbb{F}_2$.

Proof. By Remark 2.7, we need to show only (3) \Leftrightarrow (4). By Proposition 3.4, $J(R_k) = J(R) + \sum_{i=1}^k \beta_i R$. Then one easily sees that

$$R_k/J(R_k) = (R + \sum_{i=1}^k \beta_i R)/(J(R) + \sum_{i=1}^k \beta_i R) \cong R/J(R).$$

□

Combining Remark 2.7, Proposition 4.5 and Theorem 4.4 yields the following

Corollary 4.6. Suppose that R (alternatively R_k) satisfies the conditions of Proposition 4.5. Then

- (1) N_{R_k} is an ideal of $R_k[x]$;
- (2) N_R and AN_R are ideals of $R[x]$.

From now on we consider a non-commutative ring R in which N_R and AN_R are ideals of $R[x]$ (equivalently N_{R_k} is an ideal of $R_k[x]$). However, we remark that the results are still true whenever the arguments of the coming proofs involve only the pointwise addition rather than the multiplication “ \cdot ” defined in Proposition 2.8. In such circumstance we deal with $\mathcal{F}(R), \mathcal{F}(R_k), N_R, AN_R$ and N_{R_k} as R left modules.

Because of Lemma 4.3 and the definition of the ideal AN_R , we rephrase the first part of Theorem 4.4 in the following corollary.

Corollary 4.7. Let $f = f_0 + \sum_{i=1}^k f_i \beta_i$, where $f_0, \dots, f_k \in R[x]$. Then the following statements are equivalent

- (1) $f \in N_{R_k}$ (i.e. f is a null polynomial on R_k);
- (2) $f_0, f_i \beta_i \in N_{R_k}$ for $i = 1, \dots, k$;
- (3) $[\lambda_{f_0}(y, z)] = 0$ and $f_i \in N_R$ for $i = 0, \dots, k$.

Another consequence of Theorem 4.4 is the following criterion, which specifies whether two polynomials $f, g \in R_k[x]$ represent the same polynomial function on R_k .

Corollary 4.8. Let $f = f_0 + \sum_{i=1}^k f_i \beta_i$ and $g = g_0 + \sum_{i=1}^k g_i \beta_i$, where $f_0, \dots, f_k, g_0, \dots, g_k \in R[x]$.

Then $f \triangleq g$ on R_k if and only if the following conditions hold:

- (1) $[\lambda_{f_0}(y, z)] = [\lambda_{g_0}(y, z)]$;
- (2) $[f_i]_R = [g_i]_R$ for $i = 0, \dots, k$.

In other words, $f \triangleq g$ on R_k if and only if the following congruences hold:

- (1) $f_0 \equiv g_0 \pmod{AN_R}$;
- (2) $f_i \equiv g_i \pmod{N_R}$ for $i = 1, \dots, k$.

Proof. It is enough to consider the polynomial $h = g - f$ and notice that $g \triangleq f$ on R_k if and only if $h \in N_{R_k}$. □

Concerning AN_R and N_R as subgroups of the additive group $R[x]$ gives the following straightforward result of the Third Isomorphism Theorem of groups.

Lemma 4.9. *Let R be a finite ring. Then $[R[x]: AN_R] = [R[x]: N_R][N_R: AN_R]$.*

Recall from Definition 2.1 that $\mathcal{F}(R_k)$ stands for the set of polynomial functions on R_k . In the following proposition, we find a counting formula for $\mathcal{F}(R_k)$ in terms of the indices of the ideals AN_R, N_R . The proof is quite similar to that of [3, Proposition 3.7] with the difference that we replace here the ideal N'_R with the ideal AN_R .

Proposition 4.10. The number of polynomial functions on R_k is given by

$$|\mathcal{F}(R_k)| = [R[x]: AN_R][R[x]: N_R]^k = [N_R: AN_R]|\mathcal{F}(R)|^{k+1}.$$

Proof. Let $f = f_0 + \sum_{i=1}^k f_i \beta_i$ and $g = g_0 + \sum_{i=1}^k g_i \beta_i$ where $f_0, \dots, f_k, g_0, \dots, g_k \in R[x]$. Then by Corollary 4.8, $[f]_{R_k} = [g]_{R_k}$ if and only if $f_0 \equiv g_0 \pmod{AN_R}$ and $f_i \equiv g_i \pmod{N_R}$ for $i = 1, \dots, k$.

Define $\psi: \bigoplus_{i=0}^k R[x] \longrightarrow \mathcal{F}(R[\beta_1, \dots, \beta_k])$ by $\psi(f_0, \dots, f_k) = [f]_{R_k}$, where $f = f_0 + \sum_{i=1}^k f_i \beta_i$.

Then ψ is a group epimorphism of additive groups with $\ker \psi = AN_R \times \bigoplus_{i=1}^k N_R$ by Theorem 4.4.

Therefore,

$$|\mathcal{F}(R_k)| = \left[\bigoplus_{i=0}^k R[x]: AN_R \times \bigoplus_{i=1}^k N_R \right] = [R[x]: AN_R][R[x]: N_R]^k.$$

The other equality follows by Lemma 4.9 and Remark 2.6. \square

Corollary 4.11. Let $F \in \mathcal{F}(R)$ be fixed.

$$[N_R: AN_R] = |\{[\lambda_f(y, z)] \mid f \in R[x] \text{ such that } [f]_R = F\}|.$$

Proof. Fix an arbitrary $F \in \mathcal{F}(R)$. Then set

$$\mathcal{A} = \{[\lambda_f(y, z)] \mid f \in R[x] \text{ such that } [f]_R = F\},$$

and set

$$\mathcal{B} = \{G \in \mathcal{F}(R_k) \mid G = [g_0 + \sum_{i=1}^k g_i \beta_i]_{R_k} \text{ with } [g_0]_R = F, \text{ where } g_0, g_1, \dots, g_n \in R[x]\}.$$

Then, we claim that $|\mathcal{B}| = |\mathcal{A}| \cdot |\mathcal{F}(R)|^k$. To see this, let $G \in \mathcal{F}(R_k)$ be induced by $g = g_0 + \sum_{i=1}^k g_i \beta_i$ with $[g]_R = F$, where $g_0, g_1, \dots, g_n \in R[x]$. Then, by Corollary 4.8, the number of such different G is corresponding to the number of different $k+1$ tuples $([\lambda_{g_0}(y, z)], [g_1]_R, \dots, [g_k]_R)$ with the condition $[g_0]_R = F$. But, this condition implies that $[\lambda_{g_0}(y, z)]$ can be chosen in $|\mathcal{A}|$ way, whence the claim follows.

Now, we define a map $\phi: \mathcal{F}(R_k) \rightarrow \mathcal{F}(R)$ by letting $\phi(G) = [g_0]_R$. By Corollary 4.8, ϕ is well defined. Also, it is an epimorphism of (additive) groups. Thus, by Proposition 4.10,

$$|\ker \phi| = [N_R: AN_R] \cdot |\mathcal{F}(R)|^k.$$

Thus, $|\phi^{-1}(F)| = [N_R: AN_R] \cdot |\mathcal{F}(R)|^k$. Evidently, $\phi^{-1}(F)$ is the set \mathcal{B} . Hence,

$$[N_R: AN_R] \cdot |\mathcal{F}(R)|^k = |\phi^{-1}(F)| = |\mathcal{B}| = |\mathcal{A}| \cdot |\mathcal{F}(R)|^k.$$

Therefore, $|\mathcal{A}| = [N_R: AN_R]$ □

Remark 4.12.

(1) The division algorithm of polynomials over non-commutative rings can be done from two sides, that is from the left and the right. For, if $f, g \in R[x]$ such that g has a unit leading coefficient, then there exist unique q, q_1, r, r_1 such that $f(x) = q(x)g(x) + r(x)$ with $\deg r < \deg g$ or $r = 0$; and $f(x) = g(x)q_1(x) + r_1(x)$ with $\deg r_1 < \deg g$ or $r_1 = 0$. We will adopt the first one which is the division from the right.

(2) For any finite non-commutative ring A monic null polynomials not only exist but can be also central. Indeed, if $r \in R$, then there exist two positive integers $n_1(r) > n_2(r)$ such that $r^{n_1(r)} = r^{n_2(r)}$. So, $\prod_{r \in A} (x^{n_1(r)} - x^{n_2(r)})$ is a central monic null polynomial (see

for example [50, Proposition 2.2]). In particular, if A is our ring R_k , then the central monic null polynomial on R_k , $\prod_{r \in R_k} (x^{n_1(r)} - x^{n_2(r)})$, is a polynomial with coefficients in

R . Furthermore, we can take the least common multiple of the set of polynomials of the form $(x^{n_1(r)} - x^{n_2(r)})$, where $r \in A$, to get a central monic polynomial of degree $\leq \deg \prod_{r \in A} (x^{n_1(r)} - x^{n_2(r)})$.

Example 4.13. Let $R = UT_2(\mathbb{F}_2)$ (the ring of all upper triangular matrices with entries in \mathbb{F}_2 of dimension 2). Then every $a \in \{0, 1, \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 0 & 1 \end{bmatrix}\}$ is a root of the central

polynomial $x^2 - x$, and every $a \in \{\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}\}$ is a root of the central polynomial $x^4 - x^2$.

Thus, $h(x) = lcm(x^2 - x, x^4 - x^2) = x^4 - x^2$ is a central null polynomial on R . Furthermore, $h'(x) = 4x^3 - 2x = 0 \in N_R$. Hence, $h \in N'_R$. However, by Lemma 4.3, $h \notin AN_R$ since

$$\begin{aligned} h\left(\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} 0 & 1 \\ 0 & 1 \end{bmatrix} \beta\right) &= \left(\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} 0 & 1 \\ 0 & 1 \end{bmatrix} \beta\right)^4 - \left(\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} 0 & 1 \\ 0 & 1 \end{bmatrix} \beta\right)^2 \\ &= \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} - \left(\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} + \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \beta\right) = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \beta \neq 0. \end{aligned}$$

The following proposition shows that to construct a central null polynomial on R_k it is enough to know a central polynomial on R .

Proposition 4.14. Let R be a finite non-commutative ring and let $f \in R[x]$ be a central null polynomial on R . Then f^2 is a central null polynomial on R_k for every $k \geq 1$.

Proof. Since f is a central null polynomial on R , we can write $f = \sum_{i=1}^n a_i x^i$ for some $a_1, \dots, a_n \in C(R)$. Also, f is a central polynomial in the polynomial ring $R_k[x]$ by Proposition 3.4. Thus, f^2 is a central polynomial in $R_k[x]$. Let $r \in R_k$ and put $h = f^2$. Then, by Definition 2.2 since f is central,

$$h(r) = \sum_{i=1}^n a_i f(r) r^i = f(r) f(r) = (f(r))^2.$$

Now, write $r = a + \sum_{i=1}^k b_i \beta_i$, where $a, b_1, \dots, b_k \in R$. Then, by Lemma 3.10 and the fact that f is a null polynomial on R ,

$$h(r) = (f(a + \sum_{i=1}^k b_i \beta_i))^2 = (f(a) + \sum_{i=1}^k \lambda_f(a, b_i) \beta_i)^2 = (\sum_{i=1}^k \lambda_f(a, b_i) \beta_i)^2 = 0.$$

Therefore, h is a null polynomial on R_k on k for every $k \geq 1$. \square

In the following proposition, we obtain an upper bound for the minimal degree of a representative of a polynomial function on R_k .

Proposition 4.15. Let R be a finite non-commutative ring. Let $h_1 \in R_k[x]$ and $h_2 \in R[x]$ be monic null polynomials on R_k and R , respectively, such that $\deg h_1 = d_1$ and $\deg h_2 = d_2$.

Then every polynomial function $F: R_k \rightarrow R_k$ is represented by a polynomial $f = f_0 + \sum_{i=1}^k f_i \beta_i$, where $f_0, \dots, f_k \in R[x]$ such that $\deg f_0 < d_1$ and $\deg f_i < d_2$ for $i = 1, \dots, k$.

Moreover, if F is represented by a polynomial $f \in R[x]$ and $h_1 \in R[x]$ (rather than in $R_k[x]$), then there exists a polynomial $g \in R[x]$ with $\deg g < d_1$, such that $g \triangleq f$ on R and $[\lambda_g(y, z)] = [\lambda_f(y, z)]$.

Proof. Suppose that $h_1 \in R_k[x]$ is a monic null polynomial on R_k of degree d_1 . Let $g \in R_k[x]$ be a polynomial representing F . By the division algorithm, we have $g(x) = q(x)h_1(x) + r(x)$ for some $r, q \in R_k[x]$, where $\deg r \leq d_1 - 1$. Then, since $gh_1 \in N_{R_k}$,

$$g(a) = (qh_1)(a) + r(a) = r(a) \text{ for every } a \in R_k. \quad (4)$$

Thus, $r(x)$ represents F . Then, $r = f_0 + \sum_{i=1}^k r_i \beta_i$ for some $f_0, r_1, \dots, r_k \in R[x]$, and it is obvious that $\deg f_0, \deg r_i \leq d_1 - 1$ for $i = 1, \dots, k$. Now let $h_2 \in N_R$ be a monic polynomial of degree d_2 . Again, by the division algorithm, we have for $i = 1, \dots, k$, $r_i(x) = q_i(x)h_2(x) + f_i(x)$ for some $f_i, q_i \in R[x]$, where $\deg f_i \leq d_2 - 1$. Then by Corollary 4.8, $r_i \beta_i \triangleq f_i \beta_i$ on R_k . Thus $f = f_0 + \sum_{i=1}^k f_i \beta_i$ is the desired polynomial.

For the moreover part, the existence of $g \in R[x]$ with $\deg g < d_1$ such that $[f]_{R_k} = [g]_{R_k}$ follows by the same argument given in the previous part. By Corollary 4.8, $[\lambda_g(y, z)] = [\lambda_f(y, z)]$ and $[g]_R = [f]_R$. \square

Remark 4.16. We here mention that Proposition 4.15 is true in the general case. Indeed, in the proof of Proposition 4.15, Equation (4) is satisfied as well when N_{R_k} is just a left ideal. Also, if we want to argue concerning the left division argument for the same polynomial g and the monic null polynomial h_1 , the same argument works provided that N_{R_k} is an ideal. However, if we suppose that N_{R_k} is only a left ideal but not a right ideal, then we will face some difficulties. Let us illustrate this and suppose that by the left division algorithm, we have that $g(x) = h_1(x)q_1(x) + r_1(x)$. So,

$$g(a) = (h_1q_1)(a) + r_1(a) \text{ for every } a \in R_k.$$

Hence, we can not infer that in this case $f(a) = r_1(a)$ for each a , unless we have $(h_1 q_1)(a) = 0$ for each $a \in R$. Nevertheless, we can solve this obstacle by requiring that the monic null polynomials in that proposition to be central, which they exist by Remark 4.12. Note that in the cases previously discussed, we only consider substituting the variable from the right.

5. PERMUTATION POLYNOMIALS ON R_k

Given a polynomial $f = f_0 + \sum_{i=1}^k f_i \beta_i$, we see in this section that determining whether f is a permutation polynomial on R_k is completely depending on the pure part f_0 of f and its assigned polynomial λ_{f_0} .

Definition 5.1. We call the function $G: R \times R \rightarrow R$ a local permutation in the second coordinate, if for every $a \in R$ the function $G_a: R \rightarrow R$, $r \rightarrow G(a, r)$, is bijective.

When we deal with the function $[\lambda_f(y, z)]$, for some $f \in R[x]$, we use the term " the variable z " instead of " the second coordinate".

Lemma 5.2. Let R be a finite ring and let $f = f_0 + \sum_{i=1}^k f_i \beta_i$, where $f_0, \dots, f_k \in R[x]$. Suppose that f is a permutation polynomial on R_k . Then f_0 is a polynomial permutation on R and $[\lambda_{f_0}(a, z)]$ is a local permutation in the variable z .

Proof. Suppose that f is a permutation polynomial on R_k and let $a, b_1, \dots, b_k \in R$. Then, by Lemma 3.10,

$$f(a + \sum_{i=1}^k a_i \beta_i) = f_0(a) + \sum_{i=1}^k (\lambda_{f_0}(a, b_i) + f_i(a)) \beta_i.$$

So, the constant coefficient of $f(a + \sum_{i=1}^k b_i \beta_i)$ is completely determined by the value of f_0 at a . Hence, f_0 is surjective on R since otherwise f can not be surjective on R_k . Thus f_0 is a permutation on R since R is finite. Now, let $a \in R$ be arbitrary. To show that $[\lambda_{f_0}(a, z)]$ is surjective, let $c \in R$. Then, since f is a permutation polynomial on R_k , there exist $d, b_1, \dots, b_k \in R$ (i.e $d + \sum_{i=1}^k b_i \beta_i \in R_k$) such that

$$f(d + \sum_{i=1}^k b_i \beta_i) = f_0(a) + (f_1(a) + c)\beta_1.$$

But, then by Lemma 3.10,

$$f(d) + \sum_{i=1}^k (\lambda_{f_0}(d, b_i) + f_i(d)) \beta_i = f_0(a) + (f_1(a) + c)\beta_1.$$

Hence, $f_0(d) = f_0(a)$ and $\lambda_{f_0}(d, b_1) + f_1(d) = f_1(a) + c$ since R_k is an R algebra with base $1, \beta_1, \dots, \beta_k$. So, $a = d$ since f_0 is a permutation polynomial on R , whence $\lambda_{f_0}(a, b_1) = c$. Thus, $[\lambda_{f_0}(a, z)]$ is surjective, so it is bijective since R is finite. Thus, $[\lambda_{f_0}(y, z)]$ is a local permutation in z by Definition 5.1. \square

Lemma 5.3. *Let R be a finite ring and let $f = f_0 + \sum_{i=1}^k f_i \beta_i$, where $f_0, \dots, f_k \in R[x]$. Suppose that f_0 is a permutation polynomial on R and $[\lambda_{f_0}(a, z)]$ is a local permutation in the variable z . Then f is a permutation polynomial on R_k .*

Proof. Since R_k is finite, it is enough to show that f is injective on R_k , that is f induces injective function on R_k . Let $a, b_1, \dots, b_k, c, d_1, \dots, d_k \in R$ such that $f(a + \sum_{i=1}^k b_i \beta_i) = f(c + \sum_{i=1}^k d_i \beta_i)$. Then by Lemma 3.10,

$$f(a + \sum_{i=1}^k b_i \beta_i) = f_0(a) + \sum_{i=1}^k (\lambda_{f_0}(d, b_i) + f_i(a)) \beta_i = f_0(c) + \sum_{i=1}^k (\lambda_{f_0}(c, d_i) + f_i(c)) \beta_i.$$

Thus, we have that $f_0(a) = f_0(c)$ and $\lambda_{f_0}(a, b_i) + f_i(a) = \lambda_{f_0}(c, d_i) + f_i(c)$ for $i = 1, \dots, k$. Hence, since f_0 is a permutation polynomial on R , $a = c$, whence $\lambda_{f_0}(a, b_i) = \lambda_{f_0}(a, d_i)$ for $i = 1, \dots, k$. Thus, $b_i = d_i$ for $i = 1, \dots, n$ since $[\lambda_{f_0}(a, z)]$ is injective. Therefore, f is a permutation polynomial on R_k . \square

Now we are in a position to give a characterization of permutation polynomials on the ring R_k .

Theorem 5.4. *Let R be a finite non-commutative ring. Let $f = f_0 + \sum_{i=1}^k f_i \beta_i$, where $f_0, \dots, f_k \in R[x]$, and let λ_{f_0} the assigned polynomial to f_0 in the non-commutative variables y and z . Then the following statements are equivalent:*

- (1) f is a permutation polynomial on R_k ;
- (2) f_0 is a permutation polynomial on R_k ;
- (3) f_0 is a permutation polynomial on R and $[\lambda_{f_0}(a, z)]$ is surjective for every $a \in R$;
- (4) f_0 is a permutation polynomial on R and $[\lambda_{f_0}(a, z)]$ is injective for every $a \in R$;
- (5) f_0 is a permutation polynomial on R and $[\lambda_{f_0}(y, z)]$ is a local permutation in the variable z .

Proof. By Lemma 5.2 and Lemma 5.3, the statements (1), (2), and (5) are equivalent. Since R is finite, the statements (3),(4), and (5) are equivalent. \square

Remark 5.5. It is worth here to mention that for the ring of Matrices of dimension n over a finite local ring R , $M_n(R)$, Brawley [9] characterized scalar permutation polynomials of $M_n(R)$ by putting conditions not only on these polynomials but also on their assigned polynomials. More explicitly he proved the following criteria [9, Theorem 2]:

Let R be a finite commutative local ring with Maximal ideal $M \neq \{0\}$. Let $f \in R[x]$ and let $\bar{f} \in \mathbb{F}_q[x]$ be the image of f in $\mathbb{F}_q[x]$, where $\mathbb{F}_q = R/M$. Then f is a permutation polynomial on $M_n(R)$ if and only if

- (1) \bar{f} is a permutation polynomial on $M_n(\mathbb{F}_q)$, and
- (2) for every matrix $A \in M_n(\mathbb{F}_q)$, the function $[\lambda_{\bar{f}}(A, z)]$ is a permutation of $M_n(\mathbb{F}_q)$.

Theorem 5.4 shows that the criterion to be a permutation polynomial on R_k depends only on f_0 . As a consequence, we have the following corollary.

Corollary 5.6. Let R be a finite non-commutative ring. Let $f = f_0 + \sum_{i=1}^k f_i \beta_i$, where $f_0, \dots, f_k \in R[x]$. Then the following statements are equivalent:

- (1) f is a permutation polynomial on R_k ;
- (2) $f_0 + f_i \beta_i$ is a permutation polynomial on $R[\beta_i]$ for every $i \in \{1, \dots, k\}$;
- (3) f_0 is a permutation polynomial on $R[\beta_i]$ for every $i \in \{1, \dots, k\}$;
- (4) $f_0 + \sum_{i=1}^j f_i \beta_i$ is a permutation polynomial on R_l for every $1 \leq j \leq k$ and $l \geq j$;
- (5) f_0 is a permutation polynomial on R_j for every $j \geq 1$.

Another consequence of Theorem 5.4 is the following.

Corollary 5.7. Let $f = f_0 + \sum_{i=1}^k f_i \beta_i$ be a permutation polynomial on R_k , where $f_0, \dots, f_k \in R[x]$. Then the function $[\lambda_{f_0}(y, z)]: R \times R \rightarrow R$ is surjective.

Remark and Question 5.8.

- (1) If R is a commutative ring, then it will not be hard to see that the condition on $\lambda_{f_0}(y, z)$ ($f_0 \in R[x]$) in Theorem 5.4 is equivalent to f'_0 maps R to its group of units (see also Remark 3.8-(3)). But, then [3, Theorem 4.1] becomes a special case of Theorem 5.4.
- (2) In the special case R is a local commutative that is not a field (hence in the case R is a direct sum of local rings that are not fields), the condition on f'_0 is redundant, that is f_0 is a permutation polynomial on R_k if and only if f_0 is a permutation polynomial on R ([3, Proposition 4.7]). We notice here that over the finite field \mathbb{F}_q there exist $f \in \mathbb{F}_q$ a permutation polynomial on that \mathbb{F}_q is not a permutation polynomial on \mathbb{F}_{q^k} (for example x^q). However, we can always find a polynomial $g \in \mathbb{F}_q$ such that $[g]_{\mathbb{F}_q} = [f]_{\mathbb{F}_q}$ and g is a permutation polynomial on \mathbb{F}_{q^k} (see for instance [1, Lemma 4.9] or [3, Lemma 4.10]).
- (3) The previous point motivates us to ask the following question in the non-commutative case:

Is the condition on $\lambda_{f_0}(y, z)$ in Theorem 5.4 redundant?

Or equivalently:

Given $f_0 \in R[x]$. Is it true that f_0 a permutation polynomial on R if and only if f_0 is a permutation polynomial on R_k ?

In the following, we show that the set of polynomials of the form $x + \sum_{i=1}^k f_i \beta_i$, where $f_i \in R[x]$ for $i = 1, \dots, k$, is an abelian group with respect to composition of polynomials for any ring (not necessarily commutative) with unity $1 \neq 0$. Further, the set of functions induced by these group is an abelian group of permutations on R_j for every $j \geq k$. Moreover, the composition of permutations in this induced group is compatible with the composition of their defining polynomials.

Proposition 5.9. Let R be a ring with $1 \neq 0$ and $k \geq 1$. Set

$$P_{x,k} = \left\{ x + \sum_{i=1}^k f_i \beta_i \mid f_i \in R[x] \text{ for } i = 1, \dots, k \right\}, \text{ and}$$

$$\mathcal{P}_{x,k} = \{[f]_{R_k} \mid f \in P_{x,k}\}.$$

Then

- (1) $P_{x,k}$ is an abelian group with respect to composition of polynomials;
- (2) $P_{x,k} \triangleleft P_{x,j}$ for every $j > k$;
- (3) $\mathcal{P}_{x,k}$ is an abelian group with respect to composition of functions. Further, for every $F_1, F_2 \in \mathcal{P}_{x,k}$ and $f, g \in P_{x,k}$ such that $F_1 = [f]_{R_k}$ and $F_2 = [g]_{R_k}$, we have that

$$F_1 \circ F_2 = [f]_{R_k} \circ [g]_{R_k} = [f \circ g]_{R_k};$$

- (4) $\mathcal{P}_{x,k}$ is embedded normally in $\mathcal{P}_{x,j}$ for every $j > k$;
- (5) $\mathcal{P}_{x,k}$ is in $\overline{\mathcal{P}(R_j)}$ for every $j \geq k$, whenever R is finite. In this case, $|\mathcal{P}_{x,k}| = |\mathcal{F}(R)|^k$.

Proof. (1) First, we show that $P_{x,k}$ is closed with respect to composition. Let $f, g \in P_{x,k}$. Then $f = x + \sum_{i=1}^k f_i \beta_i$ and $g = x + \sum_{i=1}^k g_i \beta_i$, where $f_1, \dots, f_k, g_1, \dots, g_k \in R[x]$. So, we can write $f_i = \sum_{l=0}^{n_i} a_{li} x^l$, where $a_{li} \in R$ for $l = 0, \dots, n_i$ and $i = 1, \dots, k$. Consider,

$$\begin{aligned} f \circ g &= g + \sum_{r=1}^k f_r(g) = g + \sum_{r=1}^k f_r\left(x + \sum_{i=1}^k g_i \beta_i\right) \beta_r \\ &= g + \sum_{r=1}^k \sum_{l=0}^{n_r} a_{lr} \left(x + \sum_{i=1}^k g_i \beta_i\right)^l \beta_r = g + \sum_{r=1}^k \sum_{l=0}^{n_r} a_{lr} x^l \beta_r \quad (\text{since } \beta_i \beta_r = 0) \\ &= x + \sum_{i=1}^k g_i \beta_i + \sum_{r=1}^k f_r \beta_r = x + \sum_{i=1}^k (g_i + f_i) \beta_i \in P_{x,k}. \end{aligned}$$

Similarly, we have that

$$g \circ f = x + \sum_{i=1}^k (f_i + g_i) \beta_i = x + \sum_{i=1}^k (g_i + f_i) \beta_i = f \circ g. \quad (5)$$

Evidently, x is the identity of $P_{x,k}$ and composition of polynomial is associative. Also, if $f = x + \sum_{i=1}^k f_i \beta_i \in P_{x,k}$, then it will not be hard to see that $h = x - \sum_{i=1}^k f_i \beta_i \in P_{x,k}$ is the inverse of f . Therefore, $P_{x,k}$ is an abelian group.

(2) Follows from (1) since $P_{x,k}$ is contained in $P_{x,j}$ for every $j > k$.

(3) First, we show the operation is closed on $\mathcal{P}_{x,k}$. So, let $F_1, F_2 \in \mathcal{P}_{x,k}$. Then $F_1 = [f]_{R_k}$ and $F_2 = [g]_{R_k}$ for some $f, g \in P_{x,j}$ such that $f = x + \sum_{i=1}^k f_i \beta_i$ and $g = x + \sum_{i=1}^k g_i \beta_i$ where

$f_i, g_i \in R[x]$ for $i = 1, \dots, k$. Now, let $a_0, \dots, a_k \in R$ and consider

$$\begin{aligned}
F_1 \circ F_2(a_0 + \sum_{i=1}^k a_i \beta_i) &= F_1(F_2(a_0 + \sum_{i=1}^k a_i \beta_i)) = F_1(g(a_0 + \sum_{i=1}^k a_i \beta_i)) \\
&= F_1(a_0 + \sum_{i=1}^k (a_i + g_i(a_0)) \beta_i) \text{ (by Lemma 3.10)} \\
&= f(a_0 + \sum_{i=1}^k (a_i + g_i(a_0)) \beta_i) = a_0 + \sum_{i=1}^k (a_i + g_i(a_0)) \beta_i + \sum_{i=1}^k f_i(a_0) \beta_i \\
&= a_0 + \sum_{i=1}^k a_i \beta_i + \sum_{i=1}^k (f_i(a_0) + g_i(a_0)) \beta_i \\
&= (f \circ g)(a_0 + \sum_{i=1}^k a_i) \text{ (by Equation (5)).}
\end{aligned}$$

Thus, $F_1 \circ F_2 = [f \circ g]_{R_k} \in \mathcal{P}_{x,k}$ since $f \circ g = x + \sum_{i=1}^k (f_i + g_i) \beta_i \in P_{x,k}$. This also, shows that $[f]_{R_k} \circ [g]_{R_k} = [f \circ g]_{R_k}$, i.e., composition of functions is compatible with composition of their defined polynomials. Further, by Equation (5), we see easily that $F_1 \circ F_2 = F_2 \circ F_1$. Evidently, composition of functions is associative. Further, it is clear that $[x]_{R_k}$ is the identity element of $\mathcal{P}_{x,k}$. Finally, it is not difficult to see that if $F = [x + \sum_{i=1}^k f_i \beta_i]_{R_k}$, then F is invertible and its inverse is the function $[x - \sum_{i=1}^k f_i \beta_i]_{R_k}$.

(4) Let $j > k$ and define a map

$$\alpha : \mathcal{P}_{x,k} \longrightarrow \mathcal{P}_{x,j}, \quad F \mapsto [x + \sum_{i=1}^k f_i \beta_i]_{R_j}, \text{ where } F = [x + \sum_{i=1}^k f_i \beta_i]_{R_k}.$$

By Corollary 4.8, α is well defined. Further, by the compatibility property, we have that α is a homomorphism. Now, if $\alpha(F) = [x + \sum_{i=1}^k f_i \beta_i]_{R_j}$ is the identity function on R_j , then $[x + \sum_{i=1}^k f_i \beta_i]_{R_k} = F$ is the identity on R_k since $[x + \sum_{i=1}^k f_i \beta_i]_{R_k}$ is the restriction of $[x + \sum_{i=1}^k f_i \beta_i]_{R_j}$ to R_k . Thus, $\ker \alpha$ contains only the identity. Therefore, α is a monomorphism, and $\mathcal{P}_{x,k}$ is embedded in $\mathcal{P}_{x,j}$ for every $j > k$. Evidently, $\alpha(\mathcal{P}_{x,k})$ is a normal subgroup of $\mathcal{P}_{x,j}$ (being a subgroup of an abelian group).

(5) Follows from (4) and Corollary 4.8. □

Remark 5.10. We have already seen in the previous proposition that set $\mathcal{P}_{x,k}$ a group of permutations on R_k . So, evidently $\mathcal{P}_{x,k}$ is contained in $\mathcal{P}(R_k)$. Thus, $\mathcal{P}_{x,k}$ is a subgroup of the closure group $\overline{\mathcal{P}(R_k)}$. We will see later in Section 6 in the finite commutative case that $\mathcal{P}_{x,k}$ has a complement in the group $\mathcal{P}(R_k)$ (since in this case $\overline{\mathcal{P}(R_k)} = \mathcal{P}(R_k)$)

In the following, we find the cardinality of the set $\mathcal{P}(R_k)$ in terms of the number of polynomial functions on the ring R and the number of pairs $([g]_R, [\lambda_g(y, z)])$ such that $g \in R[x]$ is a permutation polynomial on R and $\lambda_g(y, z)$ maps $R \times R$ onto R .

Proposition 5.11. Let R be a finite non-commutative ring. Let L designate the number of pairs of functions (F, H) such that

- (1) $F: R \rightarrow R$ is bijective;
- (2) $H: R \times R \rightarrow R$ is a local permutation in the second coordinate;

occurring as $([f]_R, [\lambda_f(y, z)])$ for some $f \in R[x]$.

Then the number of polynomial permutations on R_k is given by

$$|\mathcal{P}(R_k)| = L \cdot |\mathcal{F}(R)|^k.$$

Proof. Given $G \in \mathcal{P}(R_k)$. Then by definition there exist $g_0, \dots, g_k \in R[x]$ such that G is induced by the polynomial $g = g_0 + \sum_{i=1}^k g_i \beta_i$. By Theorem 5.4, $[g_0]_R: R \rightarrow R$ is bijective, $[\lambda_{g_0}(y, z)]$ is a local permutation in z , and $[g_i]_R$ is arbitrary in $\mathcal{F}(R)$ for $i = 1, \dots, k$. Thus the result follows by Corollary 4.8. \square

In the commutative case, the number L is shown to be the numbers of pairs $([f]_R, [f']_R)$ for some permutation $f \in R[x]$ such that $[f']_R$ is a unit-valued polynomial function (i.e. maps R into its units). In particular, for the case the finite field \mathbb{F}_q of q elements, $L = q!(q-1)^q$ (see [3, Proposition 4.11]). In the next section, we will obtain different combinatorial descriptions of the number L of Proposition 5.11.

5.1. Permutation polynomials over finite non-commutative local chain rings. The purpose of this subsection is to answer the question of Remark and Question 5.8 affirmatively for a wide class of finite non-commutative chain rings. For this aim, we recall some facts about finite local rings in general and in particular about finite chain rings, the case of interest.

A finite Ring R is called a local ring if the set M of all zero-divisors of R is an ideal (two-sided ideal) of R . In this case, M is the unique maximal ideal of R ; and there exists a minimal positive integer N such that $M^N = \{0\}$ called the nilpotency index of M . Also, the characteristic of the ring $\text{Char}(R)$ is a power of some prime p , that is $\text{Char}(R) = p^c$ ($1 \leq c \leq N$); and $R/M = \mathbb{F}_q$ where $q = p^w$ ($w \geq 1$). We notice here that if $c = N$, then R is a commutative ring (see for example [41]). Furthermore, if the lattice of left ideals (equivalently of right ideals) is a chain, R is called a chain ring. It follows then that $M^i = t^i R = R t^i$ for some element $t \in M \setminus M^2$ ($i = 0, 1, \dots, N$). In particular, fixing an element $a \in R$ and ($1 \leq i < N$), we have

$$at^i = t^i a_1 \text{ for some } a_1 \in R.$$

Throughout, we use p also to designate the element $\underbrace{1_R + \dots + 1_R}_{p \text{ terms}}$, and by the ramification index

of the finite local chain ring we mean the smallest positive integer e such that $p \in M^e \setminus M^{e+1}$. Also, we mention here that for a finite ring R , being a chain ring is equivalent to being a local principal ideal ring. The above-mentioned properties of finite chain rings can be found in [36] and the references therein, and for more recent results we refer the reader to [4].

As we deal with non-commutative rings, we consider implicitly the case $c < N$, where N is the nilpotency index of the maximal ideal M and $\text{Char}(R) = p^c$.

Recall that a ring A is a semi-commutative ring whenever $a, b \in A$ with $ab = 0$ implies that $aAb = 0$. Then, we have the following lemma which is essential in the forthcoming proofs.

Lemma 5.12. *Let R be a finite chain ring and let $f \in R[x]$. The following statements hold*

- (1) R is semi-commutative;
- (2) $f(a + m) = f(a) + \lambda_f(a, m)$ for every $a, m \in R$ with $m^2 = 0$.

Proof. Let R be a finite chain ring with maximal ideal $M = tR = Rt$.

(1) We want to show that for every $a, b \in R$ such that $ab = 0$, it follows then $adb = 0$ for every $d \in R$. Without losing of generality we may assume that $a, b \neq 0$. So, if $ab = 0$, then $a, b \in M$ since R is local. By the properties of the ideal M , there exist $k_1, k_2 \geq 1$ such that $a \in M^{k_1} \setminus M^{k_1+1}$ and $b \in M^{k_2} \setminus M^{k_2+1}$. So, $a = a_1 t^{k_1}$ and $b = t^{k_2} b_1$ for some $a_1, b_1 \in R$. Hence, a_1, b_1 are units in R since otherwise we will have $a \in M^{k_1+1}$ and $b \in M^{k_2+1}$ which is not possible by our choice of k_1 and k_2 . Now,

$$0 = ab = a_1 t^{k_1} t^{k_2} b_1 = a_1 t^{k_1+k_2} b_1,$$

whence

$$t^{k_1+k_2} = 0.$$

Then, consider an arbitrary element $d \in R$. Since $M^{k_1} = Rt^{k_1} = t^{k_1}R$, there exists $d_1 \in R$ such that $dt^{k_2} = t^{k_2}d_1$. Therefore,

$$adb = a_1 t^{k_1} dt^{k_2} b_1 = a_1 t^{k_1} t^{k_2} d_1 b_1 = a_1 t^{k_1+k_2} d_1 b_1 = 0.$$

(2) Let $m \in R$ with $m^2 = 0$, then by the first assertion, $ma^j m = 0$ for $a \in R$ and $j \geq 1$.

Then it will not be hard to see that, $(a + m)^j = a^j + \sum_{r=1}^j a^{r-1} m a^{j-r} = a^j + m_j(a, m)$, where

$m_j(a, m) = \sum_{r=1}^j a^{r-1} m a^{j-r}$. Now, the rest of the proof is similar to the argument given in the proof of Lemma 3.10, and we leave the details to the reader. \square

We can replace the polynomial λ_f by f' in the second assertion of the previous lemma by requiring that a and m are commutable. The following lemma shows this in general.

Lemma 5.13. *Let R be a finite non-commutative local ring and let $f \in R[x]$. Then $f(a + m) = f(a) + f'(a)m$ for every $a, m \in R$ such that $am = ma$ and $m^2 = 0$.*

Proof. Let $a, m \in R$ such that $am = ma$ and $m^2 = 0$. Then, it follows that for $j \geq 1$, $(a + m)^j = a^j + ja^{j-1}m$. So, if $f = \sum_{j=0}^n a_j x^j$, then

$$f(a + m) = \sum_{j=0}^n a_j (a + m)^j = a_0 + \sum_{j=1}^n a_j a^j + \sum_{j=1}^n j a_j a^{j-1} m = f(a) + f'(a)m.$$

\square

Lemma 5.14. *Let R be a finite chain ring of characteristic p^c with $c > 2$ and maximal ideal M of nilpotency index N , and let $a \in R$. If $pa = 0$, then $a^2 = 0$.*

Proof. Let $M = tR$ and let e be the ramification index of R . Since $pa = 0$, $\text{Char}(R) = p^c$ and $c > 2$, we have that $a \in M$. If e is the ramification index of R , then $p = u_1 t^e$ for some unit u_1 . Further, we can argue as in the proof of Lemma 5.12 to find another three units $u_2, u_3, u_4 \in R$ and a positive integer l such that, $a = t^l u_2$, $a^2 = t^{2l} u_3$ and $p^{c-1} = u_4 t^{ec-e}$. The proof will be finished by showing that $2l \geq N$.

Then $0 = pa = u_1 t^{e+l} u_2$ implies that $t^{e+l} = 0$ and hence

$$l + e \geq N \quad (6)$$

Since $\text{Char}(R) = p^c$, we have $0 \neq p^{c-1} = u_4 t^{ec-e}$ which implies that

$$e + (c-2)e = ec - e < N. \quad (7)$$

Comparing (6) and (7) yields that $l > (c-2)e > e$ since $c > 2$. Therefore, $2l > l + e \geq N$. \square

Through out, by $\bar{f}^{(i)}$ we denote the image of the polynomial $f = \sum_{j=0}^n a_j x^j \in R[x]$, in $R/M^i[x]$, that is $\bar{f}^{(i)} = \sum_{j=0}^n \bar{a}_j^{(i)} x^j$, where $\bar{a}_i^{(i)}$ is the image of a_i in R/M^i .

Lemma 5.15. *Let R be a finite-non commutative local ring of $\text{Char}(R) = p^c$ with $c > 1$, and let $f \in R[x]$ be a permutation polynomial on R . Then $\bar{f}^{(i)}$ is a permutation polynomial on R/M^i for $i = 1, \dots, N$.*

Proof. Fix $1 \leq i \leq N$ and let $\bar{b} \in R/M^i$. Then there exists $b \in R$ such that $b = \bar{b} \pmod{M^i}$. Since f is a permutation polynomial on R there exists $a \in R$ such that $f(a) = b$. But, then we have that, $\bar{f}^{(i)}(\bar{a}) = \bar{b} \pmod{M^i}$, that is $\bar{f}^{(i)}$ is surjective on R/M^i . \square

By definition, a finite local ring is a p -group with respect to addition. Hence, the orders of the elements of R are a power of p which divides $\text{Char}(R)$. When R is a finite chain ring, De Luis [14] obtained a formula for computing the order of the elements of R .

Lemma 5.16. [14, Proposition 2] *Let R be a finite chain ring of $\text{Char}(R) = p^c$ ($c > 1$), and maximal ideal M of nilpotency index N . If $a \in M^i \setminus M^{i+1}$ for some $0 \leq i \leq N-1$, then the order of a is $p^{\lceil \frac{N-i}{e} \rceil}$, where e is the ramification index of R .*

Proposition 5.17. Let R be a finite-non commutative chain ring of $\text{Char}(R) = p^c$ with $c > 1$, and let $f \in R[x]$ be a permutation polynomial on R . Then the following statements hold

- (1) $f'(a) \neq 0 \pmod{M}$ for every $a \in R$;
- (2) $[\lambda_f(y, z)]$ is a local permutation in z .

Proof. (1) Assume to the contrary that $f'(a) = 0 \pmod{M}$ for some $a \in R$. Since $c > 1$, $0 \neq p^{c-1} \in C(R)$ with $p^{2(c-1)} = 0$. Hence, by Lemma 5.13,

$$f(a + p^{c-1}) = f(a) + f'(a)p^{c-1}. \quad (8)$$

Now, if $f'(a)p^{c-1} = 0$ then $f(a + p^{c-1}) = f(a)$ which contradicts the fact that f is a permutation polynomial on R . So, may assume that $f'(a)p^{c-1} = p^{c-1}f'(a) \neq 0$, thus the order of $f'(a)$ with respect to addition is p^c . Apparently, the element $p^{c-1} = p^{c-1}.1_R$ has order p with respect to addition. Hence, if $1 \leq i_1, i_2 < N$ such that $f'(a) \in M^{i_1} \setminus M^{i_1+1}$ and $p^{c-1} \in M^{i_2} \setminus M^{i_2+1}$, then

by Lemma 5.16, $c = \lceil \frac{N-i_1}{e} \rceil$ and $1 = \lceil \frac{N-i_2}{e} \rceil$. Thus, since the least integer function is increasing and $c > 1$, $\frac{N-i_1}{e} > \frac{N-i_2}{e}$, whence $i_1 < i_2$. Therefore, Equation (8) becomes

$$f(a + p^{c-1}) = f(a) \pmod{M^{i_2+1}}.$$

But, $p^{c-1} \neq 0 \pmod{M^{i_2+1}}$, and whence $\bar{f}^{(i_2+1)}$ is not a permutation polynomial on R/M^{i_2+1} which contradicts Lemma 5.15. Therefore, there exists no $a \in R$ such that $f'(a) = 0 \pmod{M}$.

(2) Assume to the contrary that there exists $a \in R$ such that $[\lambda_f(a, z)]$ is not a permutation of R . So, there exist $b_1, b_2 \in R$ such that $\lambda_f(a, b_1) = \lambda_f(a, b_2)$ and $b_1 \neq b_2$. Thus, by fact 3.9

$$\lambda_f(a, b) = 0, \text{ where } b = b_1 - b_2 \neq 0.$$

Now, either $pb \neq 0$ or $pb = 0$. First, consider the case $pb \neq 0$. So, if l is the largest integer such that $p^l b \neq 0$. Then, we can apply Lemma 5.12 by taking $m = p^l b$ to get that, since $p^l \in C(R)$

$$f(a + p^l b) = f(a) + \lambda_f(a, p^l b) = f(a) + p^l \lambda_f(a, b) = f(a),$$

which contradicts the fact that f is a permutation polynomial on R .

Second, consider the case the case $pb = 0$. Here, we discuss two cases on the number $c > 1$. We begin with the case $c > 2$. Then $b^2 = 0$ by Lemma 5.14. So, by applying Lemma 5.12, we see that

$$f(a + b) = f(a) + \lambda_f(a, b) = f(a).$$

Again we reach a contradiction. Finally, assume that $c = 2$. Without losing generality we can suppose that $b^2 \neq 0$, since otherwise we can argue as in the case $c > 2$. So, $b \in M^s \setminus M^{s+1}$ for some $1 \leq s < N - 1$. Thus, $b^2 = 0 \pmod{M^{s+1}}$. Therefore, we can apply Lemma 5.12 in the chain ring R/M^{s+1} to find that,

$$f(a + b) = f(a) + \lambda_f(a, b) \pmod{M^{s+1}} = f(a) \pmod{M^{s+1}}.$$

But, $b \neq 0 \pmod{M^{s+1}}$, and thus $\bar{f}^{(s+1)}$ is not a permutation polynomial on R/M^{s+1} which again conflicts with Lemma 5.15. So, the assumption $[\lambda_f(a, z)]$ is not a permutation of R for some $a \in R$ leads to a contradiction. Therefore, there is no such a and $[\lambda_f(y, z)]$ is a local permutation in z . \square

We are on time to remove the condition on the assigned polynomial in Theorem 5.4 for the class of finite rings with $\text{Char}(R) \neq p$.

Theorem 5.18. *Let R be a finite chain ring of characteristic p^c ($c > 1$). Let $f = f_0 + \sum_{i=1}^k f_i \beta_i$,*

where $f_0, \dots, f_k \in R[x]$. Then the following statements are equivalent:

- (1) f is a permutation polynomial on R_k ;
- (2) f_0 is a permutation polynomial on R_k ;
- (3) f_0 is a permutation polynomial on R .

Proof. We have already seen in Theorem 5.4 that (1) and (2) are equivalent. Also, by Lemma 5.2, (2) implies (3). Now, suppose f_0 is a permutation polynomial on R . Then, by Proposition 5.17, $[\lambda_{f_0}(y, z)]$ is a local permutation in z . Thus, this together with that fact with the fact that f_0 is a permutation polynomial on R implies that f_0 is a permutation polynomial on R_k . \square

Corollary 5.19. Let R be a finite chain ring of characteristic p^c ($c > 1$) and let $f \in R[x]$ be a permutation polynomial on R . Then $[\lambda_{f+g}(y, z)]$ is a local permutation in the variable z for every $g \in N_R$.

Proof. Let $f \in R[x]$ be a permutation polynomial on R and let $g \in N_R$. Then $f+g$ a permutation polynomial on R since $[f+g]_R = [f]_R$. Thus, by Theorem 5.18, $f+g$ is a permutation polynomial on R_k . Therefore, $[\lambda_{f+g}(y, z)]$ is a local permutation in in the variable z by Theorem 5.4. \square

Corollary 5.20. Let R be a finite chain ring of characteristic p^c ($c > 1$), $a \in R$ and $g \in N_R$. Then

- (1) $\sum_{b \in R} \lambda_g(a, b) = 0$;
- (2) $\sum_{a, b \in R} \lambda_g(a, b) = 0$.

Proof. Fix an element $a \in R$ and let $g \in N_R$ be arbitrary. Then applying Corollary 5.19 to the permutation polynomial $f = x$ implies that $[\lambda_f(a, z)]$ and $[\lambda_{f+g}(a, z)]$ are permutations on R . Hence,

$$R = \{\lambda_{f+g}(a, b) \mid b \in R\} = \{\lambda_f(a, b) \mid b \in R\}.$$

Therefore,

$$\sum_{b \in R} b = \sum_{b \in R} \lambda_{f+g}(a, b) = \sum_{b \in R} \lambda_f(a, b),$$

whence

$$\sum_{b \in R} \lambda_f(a, b) + \sum_{b \in R} \lambda_g(a, b) = \sum_{b \in R} \lambda_f(a, b).$$

Thus,

$$\sum_{b \in R} \lambda_g(a, b) = 0.$$

This proves (1). Now (2) follows from (1). \square

In general we have the following result.

Proposition 5.21. Let R be a finite non-commutative ring, $a \in R$ and $g \in N_R$. Suppose that there exists a polynomial $f \in R[x]$ such that f and $f + g$ are permutation polynomials on R_k . Then

- (1) $\sum_{b \in R} \lambda_g(a, b) = 0$;
- (2) $\sum_{a, b \in R} \lambda_g(a, b) = 0$.

Proof. Fix $a \in R$ and fix $g \in N_R$. Suppose the existence of a polynomial $f \in R[x]$ such that f and $f+g$ are permutation polynomials on R_k . Hence, by Theorem 5.4, $[\lambda_f(a, z)]$ and $[\lambda_{f+g}(a, z)]$ are permutations on R . Then the same argument used in the proof of Corollary 5.20 works here as well. \square

6. THE GROUP OF PURE POLYNOMIAL PERMUTATIONS

In this section, we consider a subgroup of $\overline{\mathcal{P}(R_k)}$ whose elements generated by polynomials over R . We call this group the group of pure polynomial permutations. Also, for the commutative case, we show this group has a complement in the group of polynomial permutations. This will allow us to pose a similar question for the non-commutative case. Finally, in a subsection, we consider a subgroup of $\overline{\mathcal{P}(R_k)}$ which maps every element of R to it self.

Recall that from Corollary 2.12 that for a finite non-commutative ring R the closure $\overline{\mathcal{P}(R_k)}$ is a group. Therefore, in view of Fact 2.11, the closure \overline{B} is a subgroup of $\overline{\mathcal{P}(R_k)}$ for every non empty subset $B \subseteq \overline{\mathcal{P}(R_k)}$. We use this fact implicitly in the rest of the paper.

Notation 6.1. For $j \geq 1$ let

$$\mathcal{P}_R(R_j) = \{F \in \mathcal{P}(R_j) \mid F = [f]_{R_j} \text{ for some } f \in R[x]\}.$$

Let $F_1, F_2 \in \mathcal{P}_R(R_j)$ such that $F_1 \neq F_2$. Then, by definition, $F_1 = [f]_{R_k}$ and $F_2 = [g]_{R_k}$ for some $f, g \in R[x]$. Furthermore, by Corollary 4.8, $[f]_R \neq [g]_R$ or $[\lambda_f(y, z)] \neq [\lambda_g(y, z)]$ since $F_1 \neq F_2$. Thus, one can see that the number L defined in Proposition 5.11 is the cardinality of the set $\mathcal{P}_R(R_k)$, and therefore, because of Theorem 5.4, the following result holds.

Corollary 6.2. Let R be a finite non-commutative Ring. Then

$$\begin{aligned} |\mathcal{P}_R(R_k)| &= |\{([f]_R, [\lambda_f(y, z)]) \mid f \in R[x], [f]_R \in \mathcal{P}(R) \text{ and } [\lambda_f(y, z)] \text{ locally permutation in } z\}| \\ &= |\{([f]_R, [\lambda_f(y, z)]) \mid f \in R[x], [f]_{R_k} \in \mathcal{P}(R_k)\}|. \end{aligned}$$

In the case R is a finite chain ring of characteristic p^c with $c > 1$ by means of Theorem 5.18, we have the following

Corollary 6.3. Let R be a finite chain ring with $Char(R) = p^c$ ($c > 1$). Then

$$|\mathcal{P}_R(R_k)| = |\{([f]_R, [\lambda(y, z)]) \mid f \in R[x], [f]_R \in \mathcal{P}(R)\}|.$$

In the following proposition, we show that the group $\overline{\mathcal{P}_R(R_j)}$ is a subgroup of $\overline{\mathcal{P}(R_j)}$, which is independent of the index j , and it is always embedded in $\overline{\mathcal{P}(R_k)}$ for every $k > j$.

Proposition 6.4. Let R be a finite ring and let $k, j \geq 1$. The group $\overline{\mathcal{P}_R(R_k)}$ is a subgroup of $\overline{\mathcal{P}(R_k)}$, and $\overline{\mathcal{P}_R(R_j)} \cong \overline{\mathcal{P}_R(R_k)}$ for every $j \neq k$. In particular, $\overline{\mathcal{P}_R(R_j)}$ is embedded in $\overline{\mathcal{P}(R_k)}$ for every $k > j$.

Proof. Since $\mathcal{P}_R(R_k) \subseteq \mathcal{P}(R_k)$, we have $\overline{\mathcal{P}_R(R_k)} \subseteq \overline{\mathcal{P}(R_k)}$ by Definition 2.10.

Now, let $j \neq k$. Without loss of generality assume that $k > j$ and let $F \in \overline{\mathcal{P}_R(R_j)}$. Then $F = F_1 \circ F_2 \circ \dots \circ F_n$ for some $F_1, \dots, F_n \in \mathcal{P}_R(R_j)$ by Definition 2.10. Then, by the definition of $\mathcal{P}_R(R_j)$, $F_i = [f_i]_{R_j}$ for some $f_i \in R[x]$; $i = 1, \dots, n$. Define

$$\psi: \overline{\mathcal{P}_R(R_j)} \longrightarrow \overline{\mathcal{P}_R(R_k)}, \quad F \mapsto [f_1]_{R_k} \circ \dots \circ [f_n]_{R_k}.$$

We show that ψ is well defined. First assume that there exist $l_1, \dots, l_n \in R[x]$ such that $[l_i]_{R_j} = F_i$ for $i = 1, \dots, n$. Then, by Corollary 4.8, $[l_i]_{R_k} = [f_i]_{R_k}$ for $i = 1, \dots, n$. Thus,

$$\psi(F) = [f_1]_{R_k} \circ \dots \circ [f_n]_{R_k} = [l_1]_{R_k} \circ \dots \circ [l_n]_{R_k}.$$

Also, if there exist $H_1, \dots, H_m \in \mathcal{P}_R(R_j)$ such that $F = H_1 \circ \dots \circ H_m$. Then there exist $h_1, \dots, h_m \in R[x]$ with $H_i = [h_i]_{R_j}$ for $i = 1, \dots, m$. Thus, by Corollary 3.14, $[h_1]_{R_k} \circ \dots \circ [h_m]_{R_k} = [f_1]_{R_k} \circ \dots \circ [f_m]_{R_k} = \psi(F)$. Therefore ψ is well defined. Further, it is a one-to-one homomorphism by Corollary 3.14. Now, let $G \in \overline{\mathcal{P}_R(R_k)}$. Then $G = G_1 \circ \dots \circ G_m$ for some $G_1, \dots, G_m \in \mathcal{P}_R(R_k)$, where $G_i = [g_i]_{R_k}$ for some $g_i \in R[x]$ for $i = 1, \dots, m$. But, then $[g_i]_{R_j} \in \mathcal{P}(R_j)$ by Corollary 5.6, whence $[g_i]_{R_j} \in \mathcal{P}_R(R_j)$ (since $g_i \in R[x]$) for $i = 1, \dots, m$. Thus, $[g_1]_{R_j} \circ \dots \circ [g_m]_{R_j} \in \overline{\mathcal{P}_R(R_j)}$. So, it is not hard to see that

$$\psi([g_1]_{R_j} \circ \dots \circ [g_m]_{R_j}) = [g_1]_{R_k} \circ \dots \circ [g_m]_{R_k}.$$

This shows that ψ is surjective, whence it is an isomorphism. Therefore, $\overline{\mathcal{P}_R(R_j)} \cong \overline{\mathcal{P}_R(R_k)}$. Now, the last statement is obvious since $\overline{\mathcal{P}_R(R_k)}$ is a subgroup of $\overline{\mathcal{P}(R_k)}$. \square

Definition 6.5. We call the group $\overline{\mathcal{P}_R(R_k)}$ the group of pure polynomial permutations.

When R is a finite commutative ring not only the set $\mathcal{P}(R_k)$ is a group but also its subset $\mathcal{P}_R(R_k)$, i.e., $\overline{\mathcal{P}_R(R_k)} = \mathcal{P}_R(R_k)$ (see [3]). In this case, we will show that the group $\mathcal{P}_{x,k}$ defined in Proposition 5.9 is a normal subgroup of $\mathcal{P}(R_k)$ that admits a complement, namely the group $\mathcal{P}_R(R_k)$. Further, we will pose a related question for the non-commutative case involving the group $\mathcal{P}_{x,k}$ and the closure groups $\overline{\mathcal{P}(R_k)}$, $\overline{\mathcal{P}_R(R_k)}$.

Before doing so, we recall the definition of split extensions (see for example [38, Page 760]). The extension

$$1 \rightarrow H \xrightarrow{i} G \xrightarrow{p} N \rightarrow 1$$

is called split if there is a homomorphism $l: N \rightarrow G$ with $p \circ l = id_N$. In such circumstances, the group G is called the semi-direct product of H by N , and we write $G = H \rtimes N$.

Theorem 6.6. *Let R be a finite ring commutative ring. Let $\mathcal{P}(R_k)$ be the group of polynomial permutations on R_k and let $\mathcal{P}_R(R_k)$ and $\mathcal{P}_{x,k}$ as in Notation 6.1 and Proposition 5.9 respectively. Then*

- (1) $\mathcal{P}(R_k) = \mathcal{P}_{x,k} \rtimes \mathcal{P}_R(R_k)$;
- (2) $|\mathcal{P}(R_k)| = |\mathcal{P}_R(R_k)| |\mathcal{F}(R)|^k$.

Proof. (1) Define a map $\varphi: \mathcal{P}(R_k) \rightarrow \mathcal{P}_R(R_k)$ by $\varphi(F) = [f_0]_{R_k}$, where $F = [f]_{R_k}$ with $f = f_0 + \sum_{i=1}^k f_i \beta_i$ and $f_0, \dots, f_k \in R[x]$. Then, by the commutative form of Corollary 4.8 (see Remark 4.2 and see [3, Corollary 3.6]), φ is well defined. Now, let $F_1 \in \mathcal{P}(R_k)$ be induced by $g = g_0 + \sum_{i=1}^k g_i \beta_i$, where $g_0, \dots, g_k \in R[x]$. Then, since composition of polynomial functions and composition of polynomials are compatible over commutative rings, we have

$$\varphi(F \circ F_1) = \varphi([f \circ g]_{R_k}) = [f_0 \circ g_0]_{R_k} = [f_0]_{R_k} \circ [g_0]_{R_k} = \varphi(F) \circ \varphi(F_1).$$

Thus, φ is a homomorphism. Then, it is evident that $\mathcal{P}_{x,k} \subseteq \ker \varphi$. On the other hand if $F \in \ker \varphi$ is induced by $f = f_0 + \sum_{i=1}^k f_i \beta_i$, then $\varphi(F) = [f_0]_{R_k} = id_{R_k} = [x]_{R_k}$. Thus, by the commutative form of Corollary 4.8, $f_0 \equiv x \pmod{N'_R}$, but this implies that (again by

the commutative form of Corollary 4.8) $F = [x + \sum_{i=1}^k f_i \beta_i]_{R_k}$. Thus, $F \in \mathcal{P}_{x,k}$, and therefore $\mathcal{P}_{x,k} = \ker \varphi$. Now, by definition, $\varphi(F) = F$ for every $F \in \mathcal{P}_R(R_k)$, whence φ is an epimorphism. Also, if $i_1: \mathcal{P}_{x,k} \rightarrow \mathcal{P}(R_k)$ and $i_2: \mathcal{P}_R(R_k) \rightarrow \mathcal{P}(R_k)$ are inclusion maps, then $\varphi(i_2(F)) = F$ for every $F \in \mathcal{P}_R(R_k)$ and the following extension is split

$$1 \rightarrow \mathcal{P}_{x,k} \xrightarrow{i_1} \mathcal{P}(R_k) \xrightarrow{\varphi} \mathcal{P}_R(R_k) \rightarrow 1.$$

Therefore, $\mathcal{P}(R_k) = \mathcal{P}_{x,k} \rtimes \mathcal{P}_R(R_k)$. (2) Follows from (1) and Proposition 5.9. □

For the non-commutative case we have the following question.

Question 6.7. Let R be a finite non-commutative ring. Then is it true that

$$\overline{\mathcal{P}(R_k)} = \mathcal{P}_{x,k} \rtimes \overline{\mathcal{P}_R(R_k)}?$$

6.1. The set of stabilizer polynomial permutations of R . In this subsection, we consider a subset of the set $\mathcal{P}(R_k)$ having the property fixing (stabilizing) the elements of the ring R pointwisely. We see set having elements represented by pure polynomials. Also, we use the cardinality of this set to find a counting formula for the number of polynomial permutations on R_k . Further, we consider the closure group of this set.

Similar to the commutative case (see[3, Definition 5.1]), we have the following definition.

Definition 6.8. Let $St_k(R) = \{F \in \mathcal{P}(R_k) \mid F(a) = a \text{ for every } a \in R\}$.

Let R be a finite ring. Then, it is obvious that $St_k(R)$ is a subset of $\mathcal{P}(R_k)$, and therefore $\overline{St_k(R)}$ is a subgroup of $\overline{\mathcal{P}(R_k)}$. Next, we show that the elements of $St_k(R)$ can be obtained by pure polynomials (see Definition 3.3).

Proposition 6.9. Let R be a finite non-commutative ring. Then

$$St_k(R) = \{F \in \mathcal{P}(R_k) \mid F \text{ is induced by } x + h(x), h \in N_R\}.$$

In particular, every element of $St_k(R)$ is induced by a polynomial in $R[x]$.

Proof. It is evident that

$$St_k(R) \supseteq \{G \in \mathcal{P}(R_k) \mid G \text{ is induced by } x + g(x), g \in N_R\}.$$

For the converse, let $G \in \mathcal{P}(R_k)$ such that $G(a) = a$ for every $a \in R$. Then G is represented by $g_0 + \sum_{i=1}^k g_i \beta_i$, where $g_0, \dots, g_k \in R[x]$, and $a = G(a) = g_0(a) + \sum_{i=1}^k g_i(a) \beta_i$ for each $a \in R$. It follows that $g_i(a) = 0$ for each $a \in R$, i.e., g_i is a null polynomial on R for $i = 1, \dots, k$. Thus $g_0 + \sum_{i=1}^k g_i \beta_i \triangleq g_0$ on R_k by Corollary 4.8, that is, G is induced by g_0 . Further, $[g_0]_R = id_R$, where id_R is the identity function on R , i.e. $g_0 \equiv x \pmod{N_R}$ and therefore $g_0(x) = x + h(x)$ for some $h \in N_R$. □

Definition 6.10. We call the group $\overline{St_k(R)}$ the (pure) stabilizer closure group.

Corollary 6.11. Let R be a finite non-commutative ring. Then $\overline{St_k(R)}$ is a subgroup of $\overline{\mathcal{P}_R(R_k)}$.

Next, we show that for all $k \geq 1$ the stabilizer closure groups $\overline{St_k(R)}$ are isomorphic.

Theorem 6.12. Let $k, j \geq 1$. Then $\overline{St_k(R)} \cong \overline{St_j(R)}$.

Proof. Without loss of generality assume that $k > j$ and let $F \in \overline{\mathcal{P}_R(R_j)}$. Then $F = F_1 \circ \cdots \circ F_n$ for some $F_1, \dots, F_n \in \mathcal{P}_R(R_j)$, where $F_i = [f_i]_{R_j}$ and $f_i \in R[x]$ for $i = 1, \dots, n$. Consider the isomorphism

$$\psi: \overline{\mathcal{P}_R(R_j)} \longrightarrow \overline{\mathcal{P}_R(R_k)}, \quad F \mapsto [f_1]_{R_k} \circ \cdots \circ [f_n]_{R_k}$$

defined in the proof of Proposition 6.4. Then $\overline{St_j(R)} \cong \overline{\psi(St_j(R))}$. Also, if $F \in \overline{St_j(R)}$, then $F = [x + h_1(x)]_{R_j} \circ \cdots \circ [x + h_n(x)]_{R_j}$ for some $h_1, \dots, h_n \in N_R$ by Proposition 6.9. Therefore, $\psi(F) = [x + h_1(x)]_{R_k} \circ \cdots \circ [x + h_n(x)]_{R_k}$ since by Proposition 6.9, $[x + h_i(x)]_{R_k} \in St_k(R)$ for $i = 1, \dots, n$. This proves that $\psi(\overline{St_j(R)}) \subseteq \overline{St_k(R)}$. Now, let $G \in \overline{St_k(R)}$. Then, $G = [x + g_1(x)]_{R_k} \circ \cdots \circ [x + g_m(x)]_{R_k}$ for some $g_1, \dots, g_m \in N_R$ by Proposition 6.9. Again by Proposition 6.9, $[x + g_i(x)]_{R_j} \in St_j(R)$ for $i = 1, \dots, m$, and therefore

$$\psi([x + g_1(x)]_{R_j} \circ \cdots \circ [x + g_m(x)]_{R_j}) = [x + g_1(x)]_{R_k} \circ \cdots \circ [x + g_m(x)]_{R_k} = G$$

with $[x + g_1(x)]_{R_j} \circ \cdots \circ [x + g_m(x)]_{R_j} \in \overline{\psi(St_j(R))}$. This proves the other inclusion and ends the proof. \square

By definition, $\overline{St_k(R)}$ stabilizes R pointwisely, that is, $F(a) = a$ for every $a \in R$ and for every $F \in \overline{St_k(R)}$. However, we do not know if every element of $\overline{\mathcal{P}_R(R_k)}$ that stabilizes the elements of R pointwisely is an element of the stabilizer closure group. This motivates to give the following definition.

Definition 6.13. Let $Stb_k(R) = \{F \in \overline{\mathcal{P}_R(R_k)} \mid F(a) = a \text{ for every } a \in R\}$.

It is evident that $Stb_k(R)$ is a subgroup of $\overline{\mathcal{P}_R(R_k)}$ whose elements stabilize the elements of R pointwisely. Because of this we call $Stb_k(R)$ the stabilizer group of R in $\overline{\mathcal{P}_R(R_k)}$. Also, it will not be hard to see that the following inclusions of sets are true

$$St_k(R) \subseteq \overline{St_k(R)} \subseteq Stb_k(R).$$

An interesting question arise from the last relation is do the following equalities hold:

$$St_k(R) = \overline{St_k(R)} = Stb_k(R)?$$

To answer this question affirmatively it would be enough to show that the set of polynomial permutations $\mathcal{P}(R_k)$ is a closed with respect of composition. In particular, in the commutative case these equalities hold.

In the following, we obtain a general form of [3, Theorem 5.7] for finite non-commutative rings. But, we can not ensure that the whole assertion of [3, Theorem 5.7] is valid in the non-commutative case. Because, in general, as we mentioned in Remark 5.8-(3), we do not know if every permutation polynomial $f \in R[x]$ on the finite non-commutative ring R is also a permutation polynomial on the ring R_k with an exceptionally large class of chain rings (see Theorem 5.18).

Proposition 6.14. Let R be a finite ring. Then the stabilizer group $Stb_k(R)$ is a normal subgroup of the group $\overline{\mathcal{P}_R(R_k)}$. Furthermore, if every element of $\mathcal{P}(R)$ is the restriction to R of an element of $\mathcal{P}_R(R_k)$, then

$$\overline{\mathcal{P}_R(R_k)}/Stb_k(R) \cong \overline{\mathcal{P}(R)}.$$

Proof. Let $F \in \overline{\mathcal{P}_R(R_k)}$. By definition, $F = [f_1]_{R_k} \circ \cdots \circ [f_n]_{R_k}$ for some $f_1, \dots, f_n \in R[x]$. Now, define a map $\Psi: \overline{\mathcal{P}_R(R_k)} \rightarrow \overline{\mathcal{P}(R)}$ by $\Psi(F) = [f_1]_R \circ \cdots \circ [f_n]_R$. Then Ψ is well defined by Theorem 5.4 and Corollary 4.8. Evidently, it is a group homomorphism with $\ker \Psi = Stb_k(R)$. Therefore, by the First Isomorphism Theorem,

$$\overline{\mathcal{P}_R(R_k)}/Stb_k(R) \cong \Psi(\overline{\mathcal{P}_R(R_k)}).$$

Furthermore, if the elements of $\mathcal{P}(R)$ obtained by the restriction to R of elements of $\mathcal{P}_R(R_k)$, then it will not be hard to see that $\Psi(\overline{\mathcal{P}_R(R_k)}) = \overline{\mathcal{P}(R)}$. Hence, if $G \in \overline{\mathcal{P}(R)}$ such that $G = G_1 \circ \cdots \circ G_m$ for some $G_1, \dots, G_m \in \mathcal{P}(R)$, then there exist $F_1, \dots, F_m \in \mathcal{P}_R(R_k)$ such that $\Psi(F_i) = G_i$ for $i = 1, \dots, m$. Therefore,

$$G = G_1 \circ \cdots \circ G_m = \Psi(F_1) \circ \cdots \circ \Psi(F_m) = \Psi(F_1 \circ \cdots \circ F_m) \in \Psi(\overline{\mathcal{P}_R(R_k)}).$$

This shows that $\Psi(\overline{\mathcal{P}_R(R_k)}) = \overline{\mathcal{P}(R)}$. □

For chain rings of $Char(R) \neq p$, we can say more.

Theorem 6.15. Let R be a finite chain ring of $Char(R) = p^c$ with $c > 1$. Then:

- (1) each element of $\mathcal{P}(R)$ appears as a restriction on R of some $G \in \mathcal{P}_R(R_k)$;
- (2) $Stb_k(R)$ is a normal subgroup of $\overline{\mathcal{P}_R(R_k)}$ and

$$\overline{\mathcal{P}_R(R_k)}/Stb_k(R) \cong \overline{\mathcal{P}(R)}.$$

Proof. (1) Follows from Theorem 5.18. (2) This a consequence of (1) and Proposition 6.14. □

Remark 6.16.

- (1) Let Ψ be the homomorphism in the proof of Proposition 6.14. Then, since $\Psi^{-1}(id_R) = \ker \Psi = Stb_k(R)$,

$$|\ker \Psi \cap \mathcal{P}_R(R_k)| = |Stb_k(R) \cap \mathcal{P}_R(R_k)| = |St_k(R)|.$$

Therefore,

$$L = |\mathcal{P}_R(R_k)| = |\Psi(\mathcal{P}_R(R_k))| \cdot |St_k(R)|,$$

where where L is as in Proposition 5.11 (see also the paragraph before Corollary 6.2).

- (2) When R is commutative, the homomorphism Ψ in the proof of Proposition 6.14 is a surjection since in this case the restriction condition in the last statement of this proposition valid (see also [3, Theorem 5.7]), and in this case, therefore $\Psi(\mathcal{P}_R(R_k)) = \mathcal{P}(R)$.
- (3) We noticed earlier that conditions on the formal derivatives in the commutative case were replaced by stronger conditions on the assigned polynomials, so, the reader could expect that the results in the commutative case involving $\mathcal{P}(R)$ are still true in the non-commutative case by involving $\Psi(\mathcal{P}_R(R_k))$ instead.

From now on, let $\Psi(\mathcal{P}_R(R_k))$ stands for the subset of $\mathcal{P}(R)$ obtained by restricting the elements of $\mathcal{P}_R(R_k)$ to R . By Remark 6.16, Proposition 5.11 and Theorem 6.15, we have the following.

Corollary 6.17. The number of polynomial permutations on R_k is given by

$$|\mathcal{P}(R_k)| = |\mathcal{F}(R)|^k \cdot |\Psi(\mathcal{P}_R(R_k))| \cdot |St_k(R)|.$$

In particular, when R is a finite chain ring with $Char(R) = p^c$ ($c > 1$),

$$|\mathcal{P}(R_k)| = |\mathcal{F}(R)|^k \cdot |\mathcal{P}(R)| \cdot |St_k(R)|.$$

Similar to the commutative case, for every integer $n \geq 1$ we assign a subset to the ideal N_R and a subset of the ideal AN_R . It is not difficult to see that these subsets are groups concerning the addition of polynomials.

Definition 6.18. For $n \geq 1$, we define

$$N_R(< n) = \{g \in R[x] \mid g \in N_R \text{ with } \deg g < n\},$$

and

$$AN_R(< n) = \{g \in R[x] \mid g \in AN_R \text{ with } \deg g < n\}.$$

In the following proposition, we give another description of the order of the stabilizer set $St_k(R)$. Also, we show this number is bounded by the index of AN_R in N_R .

Proposition 6.19. Let R be a finite non-commutative ring. Then the following hold.

- (1) $|St_k(R)| = |\{[\lambda_g(y, z)] \mid g \in N_R \text{ and } [g + x]_{R_k} \in \mathcal{P}_R(R_k)\}|$.
- (2) If there exists a monic null polynomial on R_k in $R[x]$ of degree n , then:
 - (a) $|St_k(R)| = |\{[\lambda_g(y, z)] \mid g \in N_R \text{ and } [g + x]_{R_k} \in \mathcal{P}_R(R_k) \text{ with } \deg g < n\}|$;
 - (b) $|St_k(R)| \leq [N_R : AN_R] = \frac{|N_R(< n)|}{|AN_R(< n)|}$.

Proof. (1) By Proposition 6.9,

$$\begin{aligned} |St_k(R)| &= |\{[\lambda_{x+g}(y, z)] \mid g \in N_R \text{ and } [g + x]_{R_k} \in St_k(R)\}| \\ &= |\{[\lambda_g(y, z)] \mid g \in N_R \text{ and } [g + x]_{R_k} \in \mathcal{P}_R(R_k)\}|. \end{aligned}$$

Since for every $g \in N_R$, $[x + f]_{R_k} \in St_k(R)$ if and only if $[x + f]_{R_k} \in \mathcal{P}_R(R_k)$.

(2) By Remark 4.12, there is always a monic null polynomial on R_k with coefficients from R .

(2a) If $g \in N_R$, then by Proposition 4.15, there exists $f \in R[x]$ with $\deg f < n$ such that $[\lambda_f(y, z)] = [\lambda_g(y, z)]$ and $[f]_R = [g]_R$. Clearly, $f \in N_R$.

(2b) First, we show the inequality. We have, since every element of $St_k(R)$ induces the identity on R ,

$$\{[\lambda_{x+g}(y, z)] \mid g \in N_R \text{ and } [g + x]_{R_k} \in St_k(R)\} \subseteq \{[\lambda_f(y, z)] \mid f \in R[x] \text{ and } [f]_R = id_R\}.$$

Thus, by (1) and by Corollary 4.11 for the case $F = id_R$,

$$|St_k(R)| \leq |\{[\lambda_f(y, z)] \mid f \in R[x] \text{ and } [f]_R = id_R\}| = [N_R : AN_R]. \quad (9)$$

To compute the ratio, set $\mathcal{B} = \{[\lambda_f(y, z)] \mid f \in R[x]\}$. Then, by Fact 3.9, \mathcal{B} is an additive group with operation defined as the following

$$[\lambda_f(y, z)] + [\lambda_g(y, z)] = [\lambda_{f+g}(y, z)].$$

We leave it to the reader to check the details.

Now define $\phi: N_R \rightarrow \mathcal{B}$ by $\phi(f) = [\lambda_f(y, z)]$.

Then, it is obvious that ϕ is a homomorphism of additive groups. By the definition of AN_R , $\ker \phi = AN_R$, and thus $N_R/AN_R \cong \text{Im}(\phi)$.

So, if ϕ_1 stands to the restriction of ϕ to the subgroup $N_R(< n)$, we have similarly that $\ker \phi_1 = AN_R(< n)$, and $N_R(< n)/AN_R(< n) \cong \text{Im}(\phi_1)$. Therefore, $\frac{|N_R(< n)|}{|AN_R(< n)|} = \text{Im}(\phi_1)$. So to end the proof, we need only show that $\text{Im}(\phi) \subseteq \text{Im}(\phi_1)$ since the other inclusion is valid. Assume that $F \in \text{Im}(\phi)$. Then there exists $g \in N_R$ such that $\phi(g) = [\lambda_g(y, z)] = F$. By part (a), there exists $f \in N_R$ with $\deg f < n$ (i.e. $f \in N_R(< n)$) such that $[\lambda_g(y, z)] = [\lambda_f(y, z)]$ but this mean that $F = [\lambda_f(y, z)] = \phi_1(f) \in \text{Im}(\phi_1)$. \square

Again on the considered class of finite chain rings, one can say more. For instance, the inequality of Proposition 6.19 becomes indeed equality. The following theorem illustrates this.

Theorem 6.20. *Let R be a finite chain ring with $\text{Char}(R) = p^c$ ($c > 1$). Then the following hold.*

- (1) $|St_k(R)| = |\{[\lambda_g(y, z)] \mid g \in N_R\}|$.
- (2) *If there exists a monic null polynomial on R_k in $R[x]$ of degree n , then:*
 - (a) $|St_k(R)| = |\{[\lambda_g(y, z)] \mid g \in N_R \text{ and } \deg g < n\}|$;
 - (b) $|St_k(R)| = [N_R: AN_R] = \frac{|N_R(< n)|}{|AN_R(< n)|}$.

Proof. (1) Follows by Proposition 6.19-(1) since $[f + x]_{R_k} \in \mathcal{P}(R_k)$ for every $f \in N_R$ by Theorem 5.18.

(2a) follows from (1) and Proposition 4.15.

(2b) By Proposition 6.19, we need only show that $|St_k(R)| = [N_R: AN_R]$. Now, let $f \in R[x]$. Then, by Theorem 5.18, $[f]_{R_k} \in St_k(R)$ if and only if $[f]_R = id_R$. Therefore, by Corollary 4.8,

$$|St_k(R)| = |\{[\lambda_f(y, z)] \mid f \in R[x] \text{ and } [f]_R = id_R\}|$$

But, then comparing this relation with Equation (9) in the proof of Proposition 6.19 ends the proof. \square

The previous theorem and Proposition 4.10 imply the following result.

Corollary 6.21. *Let R be a finite chain ring with $\text{Char}(R) = p^c$ ($c > 1$). Then the number of polynomial functions on R_k is given by*

$$|\mathcal{F}(R_k)| = |St_k(R)| |\mathcal{F}(R)|^{k+1}.$$

Acknowledgment. This research was funded in part by the Austrian Science Fund (FWF) [10.55776/P35788]. For open access purposes, the authors have applied a CC BY public copy-right license to any author-accepted manuscript version arising from this submission. The second author is partially supported by the IMU-Simons African Fellowship Program. The second author would like to thank the department of mathematics at the university of Graz for kind hospitality during her stay in Graz.

REFERENCES

- [1] Hasan Al-Ezeh, Amr Ali Al-Maktry, and Sophie Frisch. Polynomial functions on rings of dual numbers over residue class rings of the integers. *Math. Slovaca*, 71(5):1063–1088, 2021.
- [2] Amr Ali Al-Maktry. On the group of unit-valued polynomial functions. *Appl. Algebra Engrg. Comm. Comput.*, 34(3):521–537, 2023.
- [3] Amr Ali Abdulkader Al-Maktry. Polynomial functions over dual numbers of several variables. *J. Algebra Appl.*, 22(11):Paper No. 2350231, 2023.
- [4] Yousef Alkhamees and Sami Alabiad. Classification of chain rings. *AIMS Math.*, 7(4):5106–5116, 2022.
- [5] Vladimir Anashin and Andrei Khrennikov. *Applied algebraic dynamics*, volume 49 of *De Gruyter Expositions in Mathematics*. Walter de Gruyter & Co., Berlin, 2009.
- [6] Andrea Bandini. Functions $f \mathbf{Z}/p^n\mathbf{Z} \rightarrow \mathbf{Z}/p^n\mathbf{Z}$ induced by polynomials of $\mathbf{Z}[X]$. *Ann. Mat. Pura Appl. (4)*, 181(1):95–104, 2002.
- [7] Neal Brand. Isomorphisms of cyclic combinatorial objects. *Discrete Math.*, 78(1-2):73–81, 1989.
- [8] Joel V. Brawley. The number of polynomial functions which permute the matrices over a finite field. *J. Combinatorial Theory Ser. A*, 21(2):147–154, 1976.
- [9] Joel V. Brawley. Polynomials over a ring that permute the matrices over that ring. *J. Algebra*, 38(1):93–99, 1976.
- [10] Joel V. Brawley, Leonard Carlitz, and Jack Levine. Scalar polynomial functions on the $n \times n$ matrices over a finite field. *Linear Algebra Appl.*, 10:199–217, 1975.
- [11] Joel V. Brawley and Gary L. Mullen. Functions and polynomials over Galois rings. *J. Number Theory*, 41(2):156–166, 1992.
- [12] Joel V. Brawley and Gary L. Mullen. Scalar polynomial functions on the nonsingular matrices over a finite field. *Linear Algebra Appl.*, 174:1–12, 1992.
- [13] Balázs Bulyovszky and Gábor Horváth. Polynomial functions over finite commutative rings. *Theoret. Comput. Sci.*, 703:76–86, 2017.
- [14] M. A. De Luis. Group structure computations in finite chain rings. *J. Algebra*, 144(2):399–410, 1991.
- [15] Günther Eigenthaler. On polynomial lattices. In *Proceedings of the Lattice Theory Conference (Ulm, 1975)*, pages 54–61. Univ. Ulm, Ulm, 1975.
- [16] Günther Eigenthaler. On direct products of algebras of polynomials and polynomial functions. In *Contributions to general algebra (Proc. Klagenfurt Conf., Klagenfurt, 1978)*, pages 83–96. Heyn, Klagenfurt, 1979.
- [17] Sophie Frisch. Polynomial functions on finite commutative rings. In *Advances in Commutative Ring Theory (Fez, 1997)*, volume 205 of *Lecture Notes in Pure and Appl. Math.*, pages 323–336. Dekker, New York, 1999.
- [18] Sophie Frisch. Polynomial functions on upper triangular matrix algebras. *Monatsh. Math.*, 184(2):201–215, 2017.
- [19] Sophie Frisch. Polynomial functions on subsets of non-commutative rings — a link between ringsets and null-ideal sets. *ITM Web Conf.*, 20:01003, 2018.
- [20] Sophie Frisch and Daniel Krenn. Sylow p -groups of polynomial permutations on the integers mod p^n . *J. Number Theory*, 133(12):4188–4199, 2013.
- [21] Robert Gilmer. The ideal of polynomials vanishing on a commutative ring. *Proc. Amer. Math. Soc.*, 127(5):1265–1267, 1999.
- [22] Dalma Görcsös, Gábor Horváth, and Anett Mészáros. Permutation polynomials over finite rings. *Finite Fields Appl.*, 49:198–211, 2018.
- [23] Kenza Guenda and T. Aaron Gulliver. On the permutation groups of cyclic codes. *J. Algebraic Combin.*, 38(1):197–208, 2013.
- [24] Ashwin Guha and Ambedkar Dukkipati. A faster algorithm for testing polynomial representability of functions over finite integer rings. *Theoret. Comput. Sci.*, 579:88–99, 2015.
- [25] Jebrel M. Habeb, Mowaffaq Hajja, and William J. Heinzer. Conjugacy classes and invariant subrings of R -automorphisms of $R[x]$. *Comm. Algebra*, 40(4):1496–1524, 2012.
- [26] Mowaffaq Hajja. Algorithms for calculating the inverse of a given R -automorphism of $R[x]$. *Comm. Algebra*, 40(11):4031–4041, 2012.
- [27] Gordon Keller and F. R. Olson. Counting polynomial functions (mod p^n). *Duke Math. J.*, 35:835–838, 1968.

- [28] Aubrey J. Kempner. Polynomials and their residue systems. *Trans. Amer. Math. Soc.*, 22(2):240–266, 267–288, 1921.
- [29] Torleiv Kløve. The modular n -queen problem. *Discrete Math.*, 19(3):289–291, 1977.
- [30] Gerhard Kowol. Polynomial functions over groups: from algebraically closed groups to endomorphism near-rings. In *Contributions to general algebra. 15*, pages 45–62. Heyn, Klagenfurt, 2004.
- [31] Gerhard Kowol and Heinz Mitsch. Polynomial functions over commutative semi-groups. *Semigroup Forum*, 12(2):109–118, 1976.
- [32] O. A. Kozlitin. Polynomial transformations of a GEO-ring of prime characteristic. *Diskret. Mat.*, 16(3):105–117, 2004.
- [33] Hans Lausch and Wilfried Nöbauer. *Algebra of polynomials*, volume Vol. 5 of *North-Holland Mathematical Library*. North-Holland Publishing Co., Amsterdam-London; American Elsevier Publishing Co., Inc., New York, 1973.
- [34] Gary L. Mullen and Harlan Stevens. Polynomial functions (mod m). *Acta Math. Hungar.*, 44(3-4):237–241, 1984.
- [35] Alexander A. Nechaev. Polynomial transformations of finite commutative local rings of principal ideals. 27:425–432, 1980. transl. from 27 (1980) 885–897, 1989.
- [36] Alexandr A. Nechaev. Finite rings with applications. In *Handbook of algebra. Vol. 5*, volume 5 of *Handb. Algebr.*, pages 213–320. Elsevier/North-Holland, Amsterdam, 2008.
- [37] Wilfried Nöbauer. Über Gruppen von Restklassen nach Restpolynomidealen. *Österreich. Akad. Wiss. Math.-Nat. Kl. S.-B. IIa*, 162:207–233, 1953.
- [38] Wilfried Nöbauer. Gruppen von Restpolynomidealrestklassen nach Primzahlpotenzen. *Monatsh. Math.*, 59:194–202, 1955.
- [39] Wilfried Nöbauer. Zur Theorie der Polynomtransformationen und Permutationspolynome. *Math. Ann.*, 157:332–342, 1964.
- [40] Giulio Peruginelli and Nicholas J. Werner. Non-triviality conditions for integer-valued polynomial rings on algebras. *Monatsh. Math.*, 183(1):177–189, 2017.
- [41] R. Raghavendran. Finite associative rings. *Compositio Math.*, 21:195–229, 1969.
- [42] Mark W. Rogers and Cameron Wickham. Polynomials inducing the zero function on chain rings. *J. Algebra Appl.*, 17(8):1850160, 16, 2018.
- [43] Javad Sedighi Hafshejani and Ali Reza Naghipour. Integer-valued polynomials over matrix rings of number fields. *Bull. Iranian Math. Soc.*, 47(6):2005–2013, 2021.
- [44] David Singmaster. On polynomial functions (mod m). *J. Number Theory*, 6:345–352, 1974.
- [45] Ian Stewart. Finite rings with a specified group of units. *Math. Z.*, 126:51–58, 1972.
- [46] Jing Sun and Oscar Y. Takeshita. Interleavers for turbo codes using permutation polynomials over integer rings. *IEEE Trans. Inform. Theory*, 51(1):101–119, 2005.
- [47] Oscar Y. Takeshita. Permutation polynomial interleavers: an algebraic-geometric perspective. *IEEE Trans. Inform. Theory*, 53(6):2116–2132, 2007.
- [48] Timothy C. Teitloff. Permutation polynomials on symmetric matrices. *Linear Algebra Appl.*, 296(1-3):233–243, 1999.
- [49] Robert F. Tichy. Polynomial functions over finite monoids. *Semigroup Forum*, 22(1):83–87, 1981.
- [50] Nicholas J. Werner. Polynomials that kill each element of a finite ring. *J. Algebra Appl.*, 13(3):1350111, 12, 2014.

DEPARTMENT OF ANALYSIS AND NUMBER THEORY (5010), TECHNISCHE UNIVERSITÄT GRAZ, KOPERNIKUS-GASSE 24/II, 8010 GRAZ, AUSTRIA

Email address: `almaktry@math.tugraz.at`

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, HELWAN UNIVERSITY AIN HELWAN, 11790, CAIRO, EGYPT

Email address: `sfdeken@science.helwan.edu.eg`