

Imprints of Dark Photons on Gravitational Wave Polarizations

Kimihiro Nomura[†], Jiro Soda[‡], Kazushige Ueda[§], and Ziwei Wang^{*}

[†] *Department of Physics, Kyoto University, Kyoto 606-8502, Japan*

[‡] *Department of Physics, Kobe University, Kobe 657-8501, Japan*

[§] *National Institute of Technology, Tokuyama College, Gakuendai, Shunan, Yamaguchi
745-8585, Japan*

^{*} *Key Laboratory of Dark Matter and Space Astronomy, Purple Mountain Observatory,
Chinese Academy of Sciences, Nanjing 210023, China*

Abstract

We study conversion processes between gravitons and dark photons and reveal the effects of dark photons on the polarization of gravitational waves. Considering cosmological dark magnetic fields, we investigate the evolution of the intensity and polarization of gravitational waves through the conversion. Specifically, we demonstrate that for minimal coupling between gravitons and dark photons, the intensity, circular polarization, and linear polarization evolve separately. We derive explicit formulas for the statistical mean and variance of the intensity and polarization when the gravitational waves pass through magnetic fields with random orientation. The formulas capture how the initial polarization of dark photons will be imprinted on the observed gravitational wave background.

Contents

1	Introduction	2
2	Cosmological dark magnetic fields and graviton - dark photon conversion	4
2.1	Dark magnetic fields	4
2.2	Graviton - dark photon conversion	6
3	Conversion in a network of magnetic domains	13
3.1	Modeling of a network of magnetic domains	13
3.2	Time evolution of Stokes parameters	13
3.2.1	Class I	18
3.2.2	Class II	18
3.2.3	Class III	19
3.2.4	Class IV	19
4	Imprints on gravitational wave polarizations	20
4.1	Intensity	20
4.1.1	Mean of intensity	20
4.1.2	Variance of intensity	23
4.2	Circular polarization	25
4.2.1	Mean of circular polarization	25
4.2.2	Variance of circular polarization	26
4.3	Linear polarization	27
4.3.1	Mean of linear polarization	27
4.3.2	Variance of linear polarization	27
5	Conclusion and discussion	29

1 Introduction

Astronomy has historically developed by observations of electromagnetic waves. We can gain various information about astrophysical objects from the intensity and the polarization of electromagnetic waves. Remarkably, since the direct discovery of gravitational waves by LIGO in 2015 [1], gravitational wave astronomy has commenced [2, 3]. Now and in the future, the intensity and the polarization of gravitational waves can be valuable tools to explore the universe.

It is known that the universe consists of visible matter, which is described by the Standard Model of particle physics, and invisible matter (dark matter), which can be controlled by physics beyond the Standard Model. Surprisingly, the majority of the matter of the universe belongs to the dark sector including the dark matter. Historically, the presence of dark matter has been suggested by observations of the visible matter affected by the dark matter through gravitational interactions. However, the dark sector can also directly affect the particles in the Standard Model such as photons. For example, an axion field can interact with photons via the Chern-Simons coupling [4]. We should note that magnetic fields exist in the universe on various scales, including stars [5, 6], galaxies [7, 8], intergalactic space [9, 10, 11], and so on [12, 13]. Remarkably, in the presence of magnetic fields, an axion particle can be converted into a photon with a polarization parallel to the magnetic fields [14, 15]. Since the axion can change one of the polarization states, this conversion process changes the polarization pattern of photons. In this way, the polarization pattern of photons carries information about the dark sector. Realistic magnetic fields will have some inhomogeneity. Regarding this point, axion-photon conversion in random magnetic fields has been studied in Refs. [16, 17, 18, 19]. Particularly, in Ref. [19], the asymptotic behavior of Stokes parameters characterizing polarization of photons is investigated. The results show that the equipartition of the degrees of freedom holds.

As mentioned above, on top of electromagnetic waves, gravitational waves can be useful to explore the universe. Furthermore, gravitational waves can convey information about the even earlier universe than electromagnetic waves. Notice that it is natural to expect that the dark sector contains dark gauge fields such as dark photons. Although dark photons

with ultralight masses can be dark matter and generated during inflation [20, 21, 22, 23], we should emphasize that dark photons are not necessarily the main component of dark matter. For instance, they may behave as dark radiation, mediating interactions in mirror dark matter model [24, 25, 26]. Now, we can raise the following question. What polarization pattern of primordial gravitational waves can be expected in the presence of dark photons? More specifically, since dark photons are analogous to the Standard Model photons, it is legitimate to assume cosmological dark magnetic fields [27]. In that case, dark photons can be converted into gravitons and vice versa due to the universal (minimal) coupling [28, 14]. (Here, “gravitons” usually stand for quanta corresponding to gravitational waves, but the conversion itself can also be derived from the classical description.) Hence, the polarization pattern can be affected by dark photons. In this paper, we consider graviton - dark photon conversion in cosmological dark magnetic fields and reveal the evolution of the Stokes parameters of gravitational waves.

This paper is organized as follows. In Section 2, we study the conversion between gravitational waves and dark photons in a domain of a cosmological dark magnetic field where the magnetic field is spatially uniform. In Section 3, after constructing a sequence of domains of magnetic fields with random directions, we derive equations describing the evolution of the Stokes parameters of gravitational waves affected by conversion through the domains. In Section 4, we investigate the asymptotic behavior of the Stokes parameters when gravitational waves pass through many random magnetic field domains. In particular, we examine the statistical mean and variance of the intensity, circular polarization, and linear polarization of gravitational waves. The final section is devoted to the conclusion and discussion. We adopt the natural units, $c = \hbar = \epsilon_0 = 1$ where c is the speed of light, \hbar is the reduced Planck constant, and ϵ_0 is the permittivity in a vacuum. The metric signature is $(-, +, +, +)$.

2 Cosmological dark magnetic fields and graviton - dark photon conversion

In this paper, we consider gravitational waves and waves of a dark $U(1)$ gauge field propagating in the expanding universe accompanied by cosmological dark magnetic fields.

We start with the Einstein–Hilbert action and the action of the $U(1)$ gauge field,

$$S = \int d^4x \sqrt{-g} \left[\frac{1}{2\kappa^2} R - \frac{1}{4} g^{\mu\rho} g^{\nu\sigma} F_{\mu\nu} F_{\rho\sigma} \right]. \quad (2.1)$$

Here, R represents the Ricci scalar associated with the metric $g_{\mu\nu}$, g is the determinant of $g_{\mu\nu}$, $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$ is the field strength tensor of the $U(1)$ gauge field A_μ , and $\kappa^2 \equiv 8\pi G \equiv M_{\text{Pl}}^{-2}$, where G is the Newton constant and M_{Pl} is the reduced Planck mass. The variation of the action with respect to the metric gives the Einstein equation as

$$G^\mu{}_\nu = \kappa^2 \left(F^{\mu\alpha} F_{\nu\alpha} - \frac{1}{4} \delta^\mu_\nu F_{\alpha\beta} F^{\alpha\beta} \right), \quad (2.2)$$

where $G_{\mu\nu}$ is the Einstein tensor. The variation with respect to the gauge field A_μ gives the Maxwell equation as

$$\partial_\mu (\sqrt{-g} F^{\mu\nu}) = 0. \quad (2.3)$$

2.1 Dark magnetic fields

The background spacetime is described by the spatially flat Friedmann-Lemaître-Robertson-Walker metric,

$$\bar{g}_{\mu\nu} dx^\mu dx^\nu = a^2 (-d\eta^2 + \delta_{ij} dx^i dx^j), \quad (2.4)$$

where $\eta (= x^0)$ is the conformal time, x^i with $i = 1, 2, 3$ are the comoving coordinates, and $a = a(\eta)$ is the scale factor.

To consider background magnetic fields and propagating waves of the $U(1)$ gauge field,

we write the total gauge field strength as

$$F_{\mu\nu} = \bar{F}_{\mu\nu} + \partial_\mu A_\nu - \partial_\nu A_\mu. \quad (2.5)$$

Here, $\bar{F}_{\mu\nu}$ denotes the background field strength, and A_μ denotes the propagating field on that background, which is treated as a perturbation. (Note the change in notation: At the beginning of this section, A_μ denoted the total gauge field, but hereafter A_μ stands for the perturbation.) For the gauge field perturbation A_μ , we take the radiation gauge

$$\partial_i A_i = 0, \quad A_0 = 0. \quad (2.6)$$

Since we consider the magnetic configuration as the background, we assume $\bar{F}_{0i} = 0$ in the coordinate system $\{\eta, \mathbf{x}(= x^i)\}$. In this case, the $\mu = i$ component of the Bianchi identity $\epsilon^{\mu\nu\rho\sigma} \partial_\nu \bar{F}_{\rho\sigma} = 0$ tells us that \bar{F}_{ij} is constant in time. Then, we see that the energy density of the background dark magnetic field behaves as

$$\frac{1}{4} \bar{F}^{ij} \bar{F}_{ij} \propto a^{-4}, \quad (2.7)$$

by noting that the spatial indices of \bar{F}^{ij} are raised by the background inverse metric $\bar{g}^{ij} = a^{-2} \delta_{ij}$. Let us introduce the physical dark magnetic field vector $\bar{\mathbf{B}}(a) = (\bar{B}_i(a))$ as

$$\bar{B}_i(a) \equiv \frac{1}{2a^2} \epsilon_{ijk} \bar{F}_{jk}, \quad (2.8)$$

where ϵ_{ijk} is the anti-symmetric symbol normalized as $\epsilon_{123} = 1$. The vector $\bar{\mathbf{B}}(a)$ is defined so that the magnetic field energy density is represented as

$$\frac{1}{4} \bar{F}^{ij} \bar{F}_{ij} = \frac{1}{2} \bar{B}^2(a), \quad (2.9)$$

where $\bar{B}^2(a) \equiv \delta_{ij} \bar{B}_i(a) \bar{B}_j(a)$. Denoting the physical magnetic field magnitude at present by

\bar{B}_0 , the magnitude at the scale factor a is given by

$$\bar{B}(a) = \frac{\bar{B}_0}{a^2}, \quad (2.10)$$

where we normalized the present scale factor to be unity. Throughout this paper, we assume that the magnetic field would be rooted in a primordial origin, and the physical magnitude is only diluted through the cosmic expansion as Eq. (2.10).

If we identify \bar{B}_0 with the present magnetic field made of the Standard Model (SM) photons, various observations tell us a constraint

$$10^{-16} \text{ Gauss} \lesssim \bar{B}_0 \lesssim 10^{-9} \text{ Gauss}. \quad (\text{SM photon}) \quad (2.11)$$

The lower limit is suggested by gamma-ray observations [29, 30, 31], and the upper limit comes from Cosmic Microwave Background (CMB) and matter power spectrum observations [32]. On the other hand, when we consider the magnetic field as made of a $U(1)$ gauge field in a dark sector that does not interact with the SM, the upper bound will be relaxed. In this case, the constraint would be inferred from Big Bang Nucleosynthesis (BBN). We employ the upper limit of the dark magnetic field magnitude obtained in the literature [33],

$$\bar{B}_0 \lesssim 10^{-6} \text{ Gauss}. \quad (\text{dark photon}) \quad (2.12)$$

Below, we assume that the background magnetic field is spatially uniform, $\partial_i \bar{B}_j = 0$, in a certain domain of interest. We expect that the model with a uniform background magnetic field works well if the coherence length of the magnetic field is sufficiently longer than the wavelength of the gravitational and gauge field waves.

2.2 Graviton - dark photon conversion

To incorporate gravitational waves, we write the total metric as

$$g_{\mu\nu} dx^\mu dx^\nu = a^2 \left[-d\eta^2 + (\delta_{ij} + h_{ij}) dx^i dx^j \right], \quad (2.13)$$

where $h_{ij} = h_{ij}(\eta, \mathbf{x})$ represents the metric perturbation propagating as gravitational waves. For h_{ij} , we take the transverse-traceless gauge,

$$h_{ii} = 0, \quad \partial_j h_{ij} = 0. \quad (2.14)$$

Substituting Eqs. (2.5) and (2.13) into the Einstein equation (2.2), and extracting the linear terms with respect to h_{ij} and A_i , we obtain the equation of motion for gravitational waves as

$$(\partial_\eta^2 + 2\mathcal{H}\partial_\eta - \nabla^2)h_{ij} = 2\kappa^2 [\epsilon_{ikl}\bar{B}_l(a)(\partial_j A_k - \partial_k A_j) + \epsilon_{jkl}\bar{B}_l(a)(\partial_i A_k - \partial_k A_i)], \quad (2.15)$$

where $\mathcal{H} \equiv (\partial_\eta a)/a$, $\nabla^2 \equiv \delta_{ij}\partial_i\partial_j$, and the vector $\bar{B}_i(a)$ is defined in Eq. (2.8). Note that we have neglected the terms of the form of $\kappa^2\bar{B}^2h$ on the right-hand side. The contribution of these terms is suppressed by the factor of $M_{\text{Pl}}^{-1}\bar{B}/k$ compared with that of $\kappa^2\bar{B}(\partial A)$ terms, where k abstractly represents the comoving wavenumber of the perturbation. Throughout this paper, we consider the perturbations with large wavenumber k so that we can neglect the $\kappa^2\bar{B}^2h$ terms.

Similarly, from the $\nu = i$ component of the Maxwell equation (2.3), we obtain the linear order equation of motion for the gauge field as

$$(\partial_\eta^2 - \nabla^2)A_i = a^2\epsilon_{jkl}\bar{B}_l(a)\partial_k h_{ij}. \quad (2.16)$$

The gauge field A_i is treated as a massless field so far. However, for the Standard Model (SM) photons, an effective mass is induced by the effects of non-vacuum environments, such as plasma oscillations and quantum corrections in large magnetic fields. For dark photons that are not part of the SM, possibly they have an intrinsic mass. Such (effective) masses for gauge fields generally suppress conversion with gravitational waves. Since our main concern is the cases such that conversion is efficient, hereafter we neglect the (effective) masses for simplicity. For dark photons, this treatment is valid at any frequency if they are decoupled with other particles and are massless.

As an alternative to h_{ij} , it is convenient to introduce a new variable y_{ij} as

$$y_{ij} \equiv (2\kappa)^{-1} a h_{ij}. \quad (2.17)$$

Then, the equations of motion (2.15) and (2.16) are recast to

$$\left(\partial_\eta^2 - \nabla^2 - \frac{\partial_\eta^2 a}{a} \right) y_{ij} = \kappa a [\epsilon_{ikl} \bar{B}_l(a) (\partial_j A_k - \partial_k A_j) + \epsilon_{jkl} \bar{B}_l(a) (\partial_i A_k - \partial_k A_i)], \quad (2.18)$$

$$(\partial_\eta^2 - \nabla^2) A_i = 2 \kappa a \epsilon_{jkl} \bar{B}_l(a) \partial_k y_{ij}. \quad (2.19)$$

We define the Fourier decomposition for the metric and gauge field perturbations as

$$y_{ij}(\mathbf{x}, \eta) = \sum_{P=+, \times} \frac{1}{(2\pi)^{3/2}} \int d^3 \mathbf{k} y_P(\mathbf{k}, \eta) e_{ij}^P(\hat{\mathbf{k}}) e^{i\mathbf{k} \cdot \mathbf{x}}, \quad (2.20)$$

$$A_i(\mathbf{x}, \eta) = \sum_{P=+, \times} \frac{i}{(2\pi)^{3/2}} \int d^3 \mathbf{k} A_P(\mathbf{k}, \eta) e_i^P(\hat{\mathbf{k}}) e^{i\mathbf{k} \cdot \mathbf{x}}, \quad (2.21)$$

where we defined the unit vector $\hat{\mathbf{k}} \equiv \mathbf{k}/k$ with $k \equiv |\mathbf{k}| \equiv \sqrt{\delta_{ij} k^i k^j}$. For each direction $\hat{\mathbf{k}}$, we introduced the polarization vectors $e_i^P(\hat{\mathbf{k}})$ and the polarization tensors $e_{ij}^P(\hat{\mathbf{k}})$ with $P = +, \times$ for the transverse components. The polarization vectors are introduced to be real and satisfy

$$k^i e_i^P(\hat{\mathbf{k}}) = 0, \quad (2.22)$$

$$e_i^P(\hat{\mathbf{k}}) e_i^Q(\hat{\mathbf{k}}) = \delta^{PQ}, \quad (2.23)$$

$$e_i^+(\hat{\mathbf{k}}) = -e_i^+(-\hat{\mathbf{k}}), \quad e_i^\times(\hat{\mathbf{k}}) = e_i^\times(-\hat{\mathbf{k}}). \quad (2.24)$$

Given $e_i^P(\hat{\mathbf{k}})$, we construct the polarization tensors $e_{ij}^P(\hat{\mathbf{k}})$ as

$$e_{ij}^+(\hat{\mathbf{k}}) = \frac{1}{\sqrt{2}} \left[e_i^+(\hat{\mathbf{k}}) e_j^+(\hat{\mathbf{k}}) - e_i^\times(\hat{\mathbf{k}}) e_j^\times(\hat{\mathbf{k}}) \right], \quad (2.25)$$

$$e_{ij}^\times(\hat{\mathbf{k}}) = \frac{1}{\sqrt{2}} \left[e_i^+(\hat{\mathbf{k}}) e_j^\times(\hat{\mathbf{k}}) + e_i^\times(\hat{\mathbf{k}}) e_j^+(\hat{\mathbf{k}}) \right]. \quad (2.26)$$

These tensors satisfy

$$k^i e_{ij}^P(\hat{\mathbf{k}}) = 0, \quad e_{ii}^P(\hat{\mathbf{k}}) = 0, \quad (2.27)$$

$$e_{ij}^P(\hat{\mathbf{k}}) e_{ij}^Q(\hat{\mathbf{k}}) = \delta^{PQ}, \quad (2.28)$$

$$e_{ij}^+(\hat{\mathbf{k}}) = e_{ij}^+(-\hat{\mathbf{k}}), \quad e_{ij}^\times(\hat{\mathbf{k}}) = -e_{ij}^\times(-\hat{\mathbf{k}}). \quad (2.29)$$

Since the fields $h_{ij}(\mathbf{x}, \eta)$ and $A_i(\mathbf{x}, \eta)$ are real (Hermite), the Fourier components $y_P(\mathbf{k}, \eta)$ and $A_P(\mathbf{k}, \eta)$ must satisfy

$$y_+(\mathbf{k}, \eta) = y_+^*(-\mathbf{k}, \eta), \quad y_\times(\mathbf{k}, \eta) = -y_\times^*(-\mathbf{k}, \eta), \quad (2.30)$$

$$A_+(\mathbf{k}, \eta) = A_+^*(-\mathbf{k}, \eta), \quad A_\times(\mathbf{k}, \eta) = -A_\times^*(-\mathbf{k}, \eta), \quad (2.31)$$

where the asterisk (*) denotes the complex conjugate.

We define the polarization vectors so that the magnetic field vector $\bar{\mathbf{B}}(a)$ lies in the plane spanned by $\hat{\mathbf{k}}$ and $\mathbf{e}^\times(\hat{\mathbf{k}})$ without loss of generality. Using $\{\mathbf{e}^+(\hat{\mathbf{k}}), \mathbf{e}^\times(\hat{\mathbf{k}}), \hat{\mathbf{k}}\}$ as the coordinate axes for a given \mathbf{k} and introducing the polar angle θ , we represent the background magnetic field vector $\bar{\mathbf{B}}(a)$ as

$$\bar{\mathbf{B}}(a) = (0, \bar{B}(a) \sin \theta, \bar{B}(a) \cos \theta), \quad (2.32)$$

in that coordinate system. Inserting the Fourier decomposition (2.20) and (2.21) into Eqs. (2.18) and (2.19), and multiplying polarization vectors and tensors, we obtain the equation of motion for each polarization $P = +, \times$ as

$$\left[\partial_\eta^2 + k^2 + \begin{pmatrix} -(\partial_\eta^2 a)/a & 0 \\ 0 & 0 \end{pmatrix} \right] \begin{pmatrix} y_P(\mathbf{k}, \eta) \\ A_P(\mathbf{k}, \eta) \end{pmatrix} = -k \lambda(a) \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} y_P(\mathbf{k}, \eta) \\ A_P(\mathbf{k}, \eta) \end{pmatrix}, \quad (2.33)$$

where we defined

$$\lambda(a) \equiv \sqrt{2} \kappa a \bar{B}(a) \sin \theta = \sqrt{2} \kappa \frac{\bar{B}_0}{a} \sin \theta. \quad (2.34)$$

Equation (2.33) has been derived in Ref. [34], where it was applied to the inflationary universe

and it was shown that tachyonic growth of primordial gravitational waves may appear.

From Eq. (2.33), we see that gravitons of the $+/ \times$ mode have mixing with gauge fields of the $+/ \times$ mode. Let us now investigate conversion between gravitons and gauge fields. To see conversion, it is useful to write the components $y_P(\mathbf{k}, \eta)$ and $A_P(\mathbf{k}, \eta)$ as

$$y_+(\mathbf{k}, \eta) = \tilde{y}_+(\mathbf{k}, \eta) e^{-ik\eta} + \tilde{y}_+^*(-\mathbf{k}, \eta) e^{+ik\eta}, \quad (2.35)$$

$$y_\times(\mathbf{k}, \eta) = \tilde{y}_\times(\mathbf{k}, \eta) e^{-ik\eta} - \tilde{y}_\times^*(-\mathbf{k}, \eta) e^{+ik\eta}, \quad (2.36)$$

$$A_+(\mathbf{k}, \eta) = \tilde{A}_+(\mathbf{k}, \eta) e^{-ik\eta} + \tilde{A}_+^*(-\mathbf{k}, \eta) e^{+ik\eta}, \quad (2.37)$$

$$A_\times(\mathbf{k}, \eta) = \tilde{A}_\times(\mathbf{k}, \eta) e^{-ik\eta} - \tilde{A}_\times^*(-\mathbf{k}, \eta) e^{+ik\eta}. \quad (2.38)$$

(In each equation, the sign of the second term on the right-hand side inherits the properties (2.30) and (2.31).) Inserting Eqs. (2.35)–(2.38) into Eq. (2.33) and assuming $|\partial_\eta^2 \tilde{y}_P| \ll k|\partial_\eta \tilde{y}_P|$ and $|\partial_\eta^2 \tilde{A}_P| \ll k|\partial_\eta \tilde{A}_P|$, we obtain a Schrödinger-like equation

$$i \partial_\eta \vec{\Psi}_P = \mathbf{H}(a) \vec{\Psi}_P \quad (2.39)$$

with

$$\vec{\Psi}_P \equiv \begin{pmatrix} \tilde{y}_P(\mathbf{k}, \eta) \\ \tilde{A}_P(\mathbf{k}, \eta) \end{pmatrix}, \quad \mathbf{H}(a) \equiv \begin{pmatrix} \Delta_h(a) & \Delta_M(a) \\ \Delta_M(a) & 0 \end{pmatrix}, \quad (2.40)$$

where we defined

$$\Delta_M(a) \equiv \frac{\lambda(a)}{2} = \frac{\kappa}{\sqrt{2}} \frac{\bar{B}_0}{a} \sin \theta, \quad (2.41)$$

$$\Delta_h(a) \equiv -\frac{1}{2k} \frac{\partial_\eta^2 a}{a}. \quad (2.42)$$

Since we are interested in cases of efficient conversion, hereafter we assume

$$|\Delta_M(a)| \gg |\Delta_h(a)|, \quad (2.43)$$

so that we neglect the diagonal component of $\mathbf{H}(a)$. Particularly, in the radiation-dominated

epoch in which $a \propto \eta$, the component $\Delta_h(a)$ vanishes. More generally, we can estimate as $|\Delta_h(a)| \sim \mathcal{H}^2/k$ omitting some numerical factor, where $\mathcal{H} = (\partial_\eta a)/a$ is the comoving Hubble scale, and we are interested only in large k so that $|\Delta_h(a)|$ is negligible.

Once the diagonal component is neglected, the matrix $\mathbf{H}(a)$ can be diagonalized as

$$\mathbf{R}^\top \mathbf{H}(a) \mathbf{R} = \begin{pmatrix} +\Delta_M(a) & 0 \\ 0 & -\Delta_M(a) \end{pmatrix}, \quad \mathbf{R} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix}. \quad (2.44)$$

The important notice is that \mathbf{R} is a constant orthogonal matrix. Hence, we can obtain the analytic solution as

$$\vec{\Psi}_P(\eta) = \mathbf{R} \begin{pmatrix} e^{-i \int_{\eta_i}^{\eta} d\eta \Delta_M(a)} & 0 \\ 0 & e^{+i \int_{\eta_i}^{\eta} d\eta \Delta_M(a)} \end{pmatrix} \mathbf{R}^\top \vec{\Psi}_P(\eta_i), \quad (2.45)$$

where $\vec{\Psi}_P(\eta_i)$ is specified by the initial condition at $\eta = \eta_i$. Finally, the time evolution of the variables $\tilde{y}_P(\mathbf{k}, \eta)$ and $\tilde{A}_P(\mathbf{k}, \eta)$ reads

$$\tilde{y}_P(\mathbf{k}, \eta) = \tilde{y}_P(\mathbf{k}, \eta_i) \cos \left(\int_{\eta_i}^{\eta} d\eta \Delta_M(a) \right) - i \tilde{A}_P(\mathbf{k}, \eta_i) \sin \left(\int_{\eta_i}^{\eta} d\eta \Delta_M(a) \right), \quad (2.46)$$

$$\tilde{A}_P(\mathbf{k}, \eta) = \tilde{A}_P(\mathbf{k}, \eta_i) \cos \left(\int_{\eta_i}^{\eta} d\eta \Delta_M(a) \right) - i \tilde{y}_P(\mathbf{k}, \eta_i) \sin \left(\int_{\eta_i}^{\eta} d\eta \Delta_M(a) \right). \quad (2.47)$$

We see that conversion between gravitons and gauge fields occurs on the length scale $\Delta_M^{-1}(a)$, which is numerically given by

$$\Delta_M^{-1}(a) = 1.1 \times 10^6 a \text{ Mpc} \left(\frac{10^{-6} \text{ Gauss}}{\bar{B}_0 \sin \theta} \right). \quad (2.48)$$

In the above derivation, we have assumed $|\partial_\eta^2 \tilde{y}_P| \ll k |\partial_\eta \tilde{y}_P|$ and $|\partial_\eta^2 \tilde{A}_P| \ll k |\partial_\eta \tilde{A}_P|$. For consistency of the solution, the conditions $k \gg |\Delta_M|, |\partial_\eta \Delta_M|/|\Delta_M|$ should be satisfied, which are translated into

$$k \gg |\Delta_M(a)|, \mathcal{H}. \quad (2.49)$$

Considering the Standard Model (SM) photons as the gauge fields, the onset of graviton-

photon conversion should be the decoupling time, $z \sim 1100$. Then, even assuming the observational upper limit of the magnetic field magnitude $\bar{B}_0 \sim 10^{-9}$ Gauss, the conversion length at the decoupling time becomes 10^3 Gpc, far exceeding the present Hubble radius. Therefore, a significant conversion with gravitons is not expected for the SM photons. Indeed, the almost perfect blackbody spectrum of CMB indicates that graviton - SM photon conversion is inefficient [35].

On the other hand, the case of dark photons has a different story. First, the magnitude of the dark magnetic field is observationally less constrained as mentioned in Eq. (2.12). Moreover, as long as the dark photons are decoupled from other particles, the onset of the conversion with gravitons can be earlier. Then, as seen from Eq. (2.48), the length scale of the conversion may be shorter. Hence, for dark photons, conversion with gravitons can occur efficiently and can leave observational signatures on primordial gravitational waves: This is the case we consider in the rest of this paper. In the following sections, we study the imprints of the conversion between gravitational waves and dark photons on the intensity and polarization.

The expression for the conversion length (2.48) implies that conversion can be particularly effective during the radiation-dominated (RD) epoch. Focusing on this epoch, the integrals in Eqs. (2.46) and (2.47) can be performed analytically as follows. From the definition of the conformal time $a d\eta = dt$ and the Hubble parameter $H = (da/dt)/a$, it follows that $d\eta = (a^2 H)^{-1} da$. Additionally, the Friedmann equation in the RD epoch is given by $H = H_0 \sqrt{\Omega_{r0}/a^4}$ where H_0 is the Hubble constant and Ω_{r0} is the present density parameter of radiation. Thus, we obtain

$$\eta = \frac{a}{H_0 \sqrt{\Omega_{r0}}} = 4.6 \times 10^5 a \text{ Mpc}, \quad (\text{RD}) \quad (2.50)$$

where we set the origin of the conformal time to be the Big Bang ($a = 0$), and we employed the observed values, $H_0 = 100 h \text{ km/sec/Mpc}$ and $\Omega_{r0} h^2 = 4.2 \times 10^{-5}$ with $h = 0.68$. Then, the integrals in Eqs. (2.46) and (2.47) are explicitly evaluated as

$$\int_{\eta_i}^{\eta} d\eta \Delta_M(a) = \frac{\kappa}{\sqrt{2}} \frac{\bar{B}_0}{H_0 \sqrt{\Omega_{r0}}} (\sin \theta) \ln \left(\frac{\eta}{\eta_i} \right). \quad (\text{RD}) \quad (2.51)$$

3 Conversion in a network of magnetic domains

3.1 Modeling of a network of magnetic domains

Magnetic fields in the universe have various strengths and coherence lengths at different scales. To model magnetic fields on the cosmological scale, we construct a network of many coherent magnetic field domains as follows (see Fig. 1). First, all domains are assumed to have equal comoving size s , which compares with coherent scale of the field. The background magnetic field is assumed to be uniform within a single domain. For simplicity, we let the magnitude of the magnetic field $|\bar{\mathbf{B}}|$ be equal in all domains on a time slice, but allow the direction of the magnetic field to vary from domain to domain. Specifically, we assume that the direction of the magnetic field is random for each domain. A gravitational wave passes through a network of domains with a fixed wavenumber vector \mathbf{k} . At the n -th domain during propagation, we let θ_n denote the angle between the wavenumber vector \mathbf{k} and the magnetic field vector $\bar{\mathbf{B}}$. When moving from the $(n-1)$ -th domain into the n -th domain, the magnetic field vector rotates by an angle γ_n in the polarization plane (the two-dimensional plane normal to the wavenumber vector \mathbf{k}).

3.2 Time evolution of Stokes parameters

Recall that we have defined the polarization vectors $\mathbf{e}^+(\hat{\mathbf{k}})$ and $\mathbf{e}^\times(\hat{\mathbf{k}})$ so that the magnetic field vector $\bar{\mathbf{B}}$ lies in the $\hat{\mathbf{k}}\text{-}\mathbf{e}^\times(\hat{\mathbf{k}})$ plane. Since the magnetic field vector changes the direction when the domain changes, the definition of the polarization vector must also change. Denoting the polarization vectors in the n -th domain by $\{\mathbf{e}_n^+(\hat{\mathbf{k}}), \mathbf{e}_n^\times(\hat{\mathbf{k}})\}$, the relationship between adjacent domains is given by

$$\begin{pmatrix} \mathbf{e}_n^+(\hat{\mathbf{k}}) \\ \mathbf{e}_n^\times(\hat{\mathbf{k}}) \end{pmatrix} = \begin{pmatrix} \cos \gamma_n & \sin \gamma_n \\ -\sin \gamma_n & \cos \gamma_n \end{pmatrix} \begin{pmatrix} \mathbf{e}_{n-1}^+(\hat{\mathbf{k}}) \\ \mathbf{e}_{n-1}^\times(\hat{\mathbf{k}}) \end{pmatrix}. \quad (3.1)$$




FIG 1. The configuration of the background magnetic fields in our model is illustrated. We assume that gravitational waves and gauge field waves pass through an enormous number of domains with equal comoving sizes. Within each domain, the background magnetic field can be regarded as a uniform vector. The magnitude of the magnetic fields evolves as $|\bar{\mathbf{B}}| \propto 1/a^2$.

This rotation induces the transformation of the polarization tensors as

$$\begin{pmatrix} e_{n,ij}^+(\hat{\mathbf{k}}) \\ e_{n,ij}^\times(\hat{\mathbf{k}}) \end{pmatrix} = \begin{pmatrix} \cos 2\gamma_n & \sin 2\gamma_n \\ -\sin 2\gamma_n & \cos 2\gamma_n \end{pmatrix} \begin{pmatrix} e_{n-1,ij}^+(\hat{\mathbf{k}}) \\ e_{n-1,ij}^\times(\hat{\mathbf{k}}) \end{pmatrix}. \quad (3.2)$$

Let $A_{P,n}$ and $y_{P,n}$ ($P = +, \times$) denote the polarization components defined by means of the polarization vectors and tensors in the n -th domain. According to the rotation (3.1) and (3.2), the relationship between adjacent polarization components reads

$$\begin{pmatrix} A_{+,n}(\eta) \\ A_{\times,n}(\eta) \end{pmatrix} = \begin{pmatrix} \cos \gamma_n & \sin \gamma_n \\ -\sin \gamma_n & \cos \gamma_n \end{pmatrix} \begin{pmatrix} A_{+,n-1}(\eta) \\ A_{\times,n-1}(\eta) \end{pmatrix}, \quad (3.3)$$

$$\begin{pmatrix} y_{+,n}(\eta) \\ y_{\times,n}(\eta) \end{pmatrix} = \begin{pmatrix} \cos 2\gamma_n & \sin 2\gamma_n \\ -\sin 2\gamma_n & \cos 2\gamma_n \end{pmatrix} \begin{pmatrix} y_{+,n-1}(\eta) \\ y_{\times,n-1}(\eta) \end{pmatrix}. \quad (3.4)$$

Here, we omitted the wavenumber vector \mathbf{k} in the argument of $y_{P,n}(\mathbf{k}, \eta)$ and $A_{P,n}(\mathbf{k}, \eta)$ for notational simplicity.

Let η_{n-1} be the conformal time at the end of the passage through the $(n-1)$ -th domain, which also gives the time of entry into the n -th domain. Setting $\eta = \eta_{n-1}$ in Eqs. (3.3) and (3.4) specifies the initial condition for the polarization components defined in the n -th domain $A_{P,n}(\eta_{n-1})$ and $y_{P,n}(\eta_{n-1})$ in terms of the polarization components defined in the previous domain $A_{P,n-1}(\eta_{n-1})$ and $y_{P,n-1}(\eta_{n-1})$. Inserting the initial condition into the general solutions (2.46) and (2.47), we obtain the relationship between the polarization components at the end of the n -th domain and those at the end of the $(n-1)$ -th domain as

$$\varphi_n = \Phi_n \cdot \Gamma_n \cdot \varphi_{n-1}. \quad (3.5)$$

Here, we introduced a four-component vector φ_n as

$$\varphi_n \equiv (\tilde{y}_{+,n}(\eta_n), \tilde{y}_{\times,n}(\eta_n), \tilde{A}_{+,n}(\eta_n), \tilde{A}_{\times,n}(\eta_n))^{\top}. \quad (3.6)$$

We also defined

$$\Phi_n \equiv \begin{pmatrix} \cos \Phi_n & 0 & -i \sin \Phi_n & 0 \\ 0 & \cos \Phi_n & 0 & -i \sin \Phi_n \\ -i \sin \Phi_n & 0 & \cos \Phi_n & 0 \\ 0 & -i \sin \Phi_n & 0 & \cos \Phi_n \end{pmatrix} \quad (3.7)$$

with

$$\Phi_n \equiv \int_{\eta_{n-1}}^{\eta_n} d\eta \Delta_{M,n}(a), \quad \Delta_{M,n}(a) \equiv \frac{\kappa}{\sqrt{2}} \frac{\bar{B}_0}{a} \sin \theta_n, \quad (3.8)$$

and

$$\mathbf{\Gamma}_n \equiv \begin{pmatrix} \cos 2\gamma_n & \sin 2\gamma_n & 0 & 0 \\ -\sin 2\gamma_n & \cos 2\gamma_n & 0 & 0 \\ 0 & 0 & \cos \gamma_n & \sin \gamma_n \\ 0 & 0 & -\sin \gamma_n & \cos \gamma_n \end{pmatrix}. \quad (3.9)$$

To describe polarization, let us introduce Stokes parameters with the vector $\boldsymbol{\varphi}$ defined in Eq. (3.6) as

$$\boldsymbol{\varphi} \boldsymbol{\varphi}^\dagger \equiv \frac{1}{2} \begin{pmatrix} I_h + Q_h & U_h - iV_h & K - iL & M - iN \\ U_h + iV_h & I_h - Q_h & W - iX & Y - iZ \\ K + iL & W + iX & I_\gamma + Q_\gamma & U_\gamma - iV_\gamma \\ M + iN & Y + iZ & U_\gamma + iV_\gamma & I_\gamma - Q_\gamma \end{pmatrix}, \quad (3.10)$$

where the subscripts h and γ represent that the quantities are associated with gravitational waves and gauge fields, respectively. The subscript labeling the number of domains n is omitted here. Specifically, the intensity of gravitational waves and gauge fields are denoted by I_h and I_γ , respectively, and those after passing the n -th domain are given by

$$I_h(\mathbf{k}, \eta_n) = |\tilde{y}_{+,n}(\mathbf{k}, \eta_n)|^2 + |\tilde{y}_{\times,n}(\mathbf{k}, \eta_n)|^2, \quad (3.11)$$

$$I_\gamma(\mathbf{k}, \eta_n) = |\tilde{A}_{+,n}(\mathbf{k}, \eta_n)|^2 + |\tilde{A}_{\times,n}(\mathbf{k}, \eta_n)|^2. \quad (3.12)$$

The variables V_h and V_γ denote the degree of circular polarization of gravitational waves and gauge fields, respectively. Those at the end of the n -th domain are given by

$$V_h(\mathbf{k}, \eta_n) = i[\tilde{y}_{\times,n}^*(\mathbf{k}, \eta_n) \tilde{y}_{+,n}(\mathbf{k}, \eta_n) - \tilde{y}_{+,n}^*(\mathbf{k}, \eta_n) \tilde{y}_{\times,n}(\mathbf{k}, \eta_n)], \quad (3.13)$$

$$V_\gamma(\mathbf{k}, \eta_n) = i[\tilde{A}_{\times,n}^*(\mathbf{k}, \eta_n) \tilde{A}_{+,n}(\mathbf{k}, \eta_n) - \tilde{A}_{+,n}^*(\mathbf{k}, \eta_n) \tilde{A}_{\times,n}(\mathbf{k}, \eta_n)]. \quad (3.14)$$

The variables $Q_{h/\gamma}$ and $U_{h/\gamma}$ denote the degree of linear polarization of gravitational waves/gauge

fields,

$$Q_h(\mathbf{k}, \eta_n) = |\tilde{y}_{+,n}(\mathbf{k}, \eta_n)|^2 - |\tilde{y}_{\times,n}(\mathbf{k}, \eta_n)|^2, \quad (3.15)$$

$$Q_\gamma(\mathbf{k}, \eta_n) = |\tilde{A}_{+,n}(\mathbf{k}, \eta_n)|^2 - |\tilde{A}_{\times,n}(\mathbf{k}, \eta_n)|^2, \quad (3.16)$$

and

$$U_h(\mathbf{k}, \eta_n) = \tilde{y}_{\times,n}^*(\mathbf{k}, \eta_n) \tilde{y}_{+,n}(\mathbf{k}, \eta_n) + \tilde{y}_{+,n}^*(\mathbf{k}, \eta_n) \tilde{y}_{\times,n}(\mathbf{k}, \eta_n), \quad (3.17)$$

$$U_\gamma(\mathbf{k}, \eta_n) = \tilde{A}_{\times,n}^*(\mathbf{k}, \eta_n) \tilde{A}_{+,n}(\mathbf{k}, \eta_n) + \tilde{A}_{+,n}^*(\mathbf{k}, \eta_n) \tilde{A}_{\times,n}(\mathbf{k}, \eta_n). \quad (3.18)$$

Note that $I_{h/\gamma}$ and $V_{h/\gamma}$ are coordinate-independent notions in the sense that they are invariant under rotation of the coordinate axes in the polarization plane, while $Q_{h/\gamma}$ and $U_{h/\gamma}$ are dependent. In Eq. (3.10), we also introduced real parameters K, L, M, N, W, X, Y , and Z , which consist of cross terms of \tilde{y}_P and \tilde{A}_P as follows:

$$K(\mathbf{k}, \eta_n) = \tilde{y}_{+,n}(\mathbf{k}, \eta_n) \tilde{A}_{+,n}^*(\mathbf{k}, \eta_n) + \tilde{y}_{+,n}^*(\mathbf{k}, \eta_n) \tilde{A}_{+,n}(\mathbf{k}, \eta_n), \quad (3.19)$$

$$L(\mathbf{k}, \eta_n) = i[\tilde{y}_{+,n}(\mathbf{k}, \eta_n) \tilde{A}_{+,n}^*(\mathbf{k}, \eta_n) - \tilde{y}_{+,n}^*(\mathbf{k}, \eta_n) \tilde{A}_{+,n}(\mathbf{k}, \eta_n)], \quad (3.20)$$

$$M(\mathbf{k}, \eta_n) = \tilde{y}_{+,n}(\mathbf{k}, \eta_n) \tilde{A}_{\times,n}^*(\mathbf{k}, \eta_n) + \tilde{y}_{+,n}^*(\mathbf{k}, \eta_n) \tilde{A}_{\times,n}(\mathbf{k}, \eta_n), \quad (3.21)$$

$$N(\mathbf{k}, \eta_n) = i[\tilde{y}_{+,n}(\mathbf{k}, \eta_n) \tilde{A}_{\times,n}^*(\mathbf{k}, \eta_n) - \tilde{y}_{+,n}^*(\mathbf{k}, \eta_n) \tilde{A}_{\times,n}(\mathbf{k}, \eta_n)], \quad (3.22)$$

$$W(\mathbf{k}, \eta_n) = \tilde{y}_{\times,n}(\mathbf{k}, \eta_n) \tilde{A}_{+,n}^*(\mathbf{k}, \eta_n) + \tilde{y}_{\times,n}^*(\mathbf{k}, \eta_n) \tilde{A}_{+,n}(\mathbf{k}, \eta_n), \quad (3.23)$$

$$X(\mathbf{k}, \eta_n) = i[\tilde{y}_{\times,n}(\mathbf{k}, \eta_n) \tilde{A}_{+,n}^*(\mathbf{k}, \eta_n) - \tilde{y}_{\times,n}^*(\mathbf{k}, \eta_n) \tilde{A}_{+,n}(\mathbf{k}, \eta_n)], \quad (3.24)$$

$$Y(\mathbf{k}, \eta_n) = \tilde{y}_{\times,n}(\mathbf{k}, \eta_n) \tilde{A}_{\times,n}^*(\mathbf{k}, \eta_n) + \tilde{y}_{\times,n}^*(\mathbf{k}, \eta_n) \tilde{A}_{\times,n}(\mathbf{k}, \eta_n), \quad (3.25)$$

$$Z(\mathbf{k}, \eta_n) = i[\tilde{y}_{\times,n}(\mathbf{k}, \eta_n) \tilde{A}_{\times,n}^*(\mathbf{k}, \eta_n) - \tilde{y}_{\times,n}^*(\mathbf{k}, \eta_n) \tilde{A}_{\times,n}(\mathbf{k}, \eta_n)]. \quad (3.26)$$

To study the evolution of the Stokes parameters, we symbolize the matrix in Eq. (3.10) by $\boldsymbol{\rho}$. Then, at the end of the n -th domain, we obtain the relation

$$\boldsymbol{\rho}_n \equiv \boldsymbol{\varphi}_n \boldsymbol{\varphi}_n^\dagger = (\boldsymbol{\Phi}_n \cdot \boldsymbol{\Gamma}_n) \cdot \boldsymbol{\rho}_{n-1} \cdot (\boldsymbol{\Phi}_n \cdot \boldsymbol{\Gamma}_n)^\dagger, \quad (3.27)$$

where we used Eq. (3.5). The above relation yields recurrence relations between the n -th and

$(n-1)$ -th Stokes parameters. It turns out that not all the Stokes parameters are coupled to each other. Indeed, the recurrence relations are classified into four classes as follows.

3.2.1 Class I

First, let us focus on the part concerning the intensity of gravitational waves and gauge fields, I_h and I_γ . It is found that the variables $\{I_h, I_\gamma, L + Z, X - N\}$ constitute a closed set of equations in the recurrence relation (3.27) as follows:

$$I_h(\eta_n) + I_\gamma(\eta_n) = I_h(\eta_{n-1}) + I_\gamma(\eta_{n-1}), \quad (3.28)$$

$$\begin{aligned} I_h(\eta_n) - I_\gamma(\eta_n) &= \sin 2\Phi_n \{ \cos \gamma_n [L(\eta_{n-1}) + Z(\eta_{n-1})] + \sin \gamma_n [X(\eta_{n-1}) - N(\eta_{n-1})] \} \\ &\quad + \cos 2\Phi_n [I_h(\eta_{n-1}) - I_\gamma(\eta_{n-1})], \end{aligned} \quad (3.29)$$

$$\begin{aligned} L(\eta_n) + Z(\eta_n) &= \cos 2\Phi_n \{ \cos \gamma_n [L(\eta_{n-1}) + Z(\eta_{n-1})] + \sin \gamma_n [X(\eta_{n-1}) - N(\eta_{n-1})] \} \\ &\quad - \sin 2\Phi_n [I_h(\eta_{n-1}) - I_\gamma(\eta_{n-1})], \end{aligned} \quad (3.30)$$

$$X(\eta_n) - N(\eta_n) = \cos \gamma_n [X(\eta_{n-1}) - N(\eta_{n-1})] - \sin \gamma_n [L(\eta_{n-1}) + Z(\eta_{n-1})]. \quad (3.31)$$

3.2.2 Class II

The circular polarizations of gravitational waves and gauge field waves are determined by the following closed set of equations for the variables $\{V_h, V_\gamma, M - W, K + Y\}$:

$$V_h(\eta_n) + V_\gamma(\eta_n) = V_h(\eta_{n-1}) + V_\gamma(\eta_{n-1}), \quad (3.32)$$

$$\begin{aligned} V_h(\eta_n) - V_\gamma(\eta_n) &= -\sin 2\Phi_n \{ \cos \gamma_n [M(\eta_{n-1}) - W(\eta_{n-1})] + \sin \gamma_n [K(\eta_{n-1}) + Y(\eta_{n-1})] \} \\ &\quad + \cos 2\Phi_n [V_h(\eta_{n-1}) - V_\gamma(\eta_{n-1})], \end{aligned} \quad (3.33)$$

$$\begin{aligned} M(\eta_n) - W(\eta_n) &= \cos 2\Phi_n \{ \cos \gamma_n [M(\eta_{n-1}) - W(\eta_{n-1})] + \sin \gamma_n [K(\eta_{n-1}) + Y(\eta_{n-1})] \} \\ &\quad + \sin 2\Phi_n [V_h(\eta_{n-1}) - V_\gamma(\eta_{n-1})], \end{aligned} \quad (3.34)$$

$$K(\eta_n) + Y(\eta_n) = \cos \gamma_n [K(\eta_{n-1}) + Y(\eta_{n-1})] - \sin \gamma_n [M(\eta_{n-1}) - W(\eta_{n-1})]. \quad (3.35)$$

3.2.3 Class III

The linear polarizations of gravitational waves and gauge field waves are determined by the following closed set of equations for the variables $\{Q_h + iU_h, Q_\gamma + iU_\gamma, L - Z, X + N\}$:

$$Q_h(\eta_n) + iU_h(\eta_n) = \cos^2 \Phi_n e^{-4i\gamma_n} [Q_h(\eta_{n-1}) + iU_h(\eta_{n-1})] + \sin^2 \Phi_n e^{-2i\gamma_n} [Q_\gamma(\eta_{n-1}) + iU_\gamma(\eta_{n-1})] + \frac{1}{2} \sin 2\Phi_n e^{-3i\gamma_n} \{ [L(\eta_{n-1}) - Z(\eta_{n-1})] + i [X(\eta_{n-1}) + N(\eta_{n-1})] \}, \quad (3.36)$$

$$Q_\gamma(\eta_n) + iU_\gamma(\eta_n) = \sin^2 \Phi_n e^{-4i\gamma_n} [Q_h(\eta_{n-1}) + iU_h(\eta_{n-1})] + \cos^2 \Phi_n e^{-2i\gamma_n} [Q_\gamma(\eta_{n-1}) + iU_\gamma(\eta_{n-1})] - \frac{1}{2} \sin 2\Phi_n e^{-3i\gamma_n} \{ [L(\eta_{n-1}) - Z(\eta_{n-1})] + i [X(\eta_{n-1}) + N(\eta_{n-1})] \}, \quad (3.37)$$

$$L(\eta_n) - Z(\eta_n) = \cos 2\Phi_n \{ \cos 3\gamma_n [L(\eta_{n-1}) - Z(\eta_{n-1})] + \sin 3\gamma_n [X(\eta_{n-1}) + N(\eta_{n-1})] \} - \frac{1}{2} \sin 2\Phi_n \{ e^{-4i\gamma_n} [Q_h(\eta_{n-1}) + iU_h(\eta_{n-1})] + \text{c.c.} - e^{-2i\gamma_n} [Q_\gamma(\eta_{n-1}) + iU_\gamma(\eta_{n-1})] + \text{c.c.} \}, \quad (3.38)$$

$$X(\eta_n) + N(\eta_n) = \cos 2\Phi_n \{ \cos 3\gamma_n [X(\eta_{n-1}) + N(\eta_{n-1})] - \sin 3\gamma_n [L(\eta_{n-1}) - Z(\eta_{n-1})] \} + \frac{1}{2} \sin 2\Phi_n \{ i e^{-4i\gamma_n} [Q_h(\eta_{n-1}) + iU_h(\eta_{n-1})] + \text{c.c.} - i e^{-2i\gamma_n} [Q_\gamma(\eta_{n-1}) + iU_\gamma(\eta_{n-1})] + \text{c.c.} \}, \quad (3.39)$$

where ‘‘c.c.’’ represents the complex conjugate of the previous term.

3.2.4 Class IV

Additionally, we can find that the variables $\{M + W, K - Y\}$ form a closed set although it is redundant for the purpose of studying the intensity and polarization of gravitational waves:

$$M(\eta_n) + W(\eta_n) = \cos 3\gamma_n [M(\eta_{n-1}) + W(\eta_{n-1})] - \sin 3\gamma_n [K(\eta_{n-1}) - Y(\eta_{n-1})], \quad (3.40)$$

$$K(\eta_n) - Y(\eta_n) = \sin 3\gamma_n [M(\eta_{n-1}) + W(\eta_{n-1})] + \cos 3\gamma_n [K(\eta_{n-1}) - Y(\eta_{n-1})]. \quad (3.41)$$

In summary, we have obtained four classes of recurrence relations. Note that the intensity ($I_{h/\gamma}$), circular polarization ($V_{h/\gamma}$), and linear polarization ($Q_{h/\gamma}, U_{h/\gamma}$) are separated. This

stems from the separation of equations of $+$ and \times modes. Although we have focused here on the minimal coupling, when we take into account non-minimal cases, the situation would be different.

4 Imprints on gravitational wave polarizations

In the previous section, we derived the relationship between the Stokes parameters in adjacent magnetic field domains. In this section, we investigate the asymptotic behavior of the Stokes parameters assuming that the direction of the magnetic field is randomly chosen for each domain. Since we have seen that intensity, circular polarization, and linear polarization are separated in the recurrence relations, we can study each of these classes separately.

4.1 Intensity

First, let us focus on Class I, i.e., Eqs. (3.28)–(3.31), to examine the intensity. Since the magnetic field direction is assumed to be given probabilistically, we study statistical quantities employing an ensemble average. Specifically, the purpose below is to find the asymptotic behavior of the mean (expectation value) of the intensity and the variance of the intensity.

4.1.1 Mean of intensity

Equation (3.28) shows that the sum of I_h and I_γ does not change even when the domain changes. This indicates that the total intensity of the gravitational waves and gauge fields is conserved,

$$I_h(\eta) + I_\gamma(\eta) = I_{h,0} + I_{\gamma,0}, \quad (4.1)$$

where $I_{h,0} = I_h(\eta_0)$ and $I_{\gamma,0} = I_\gamma(\eta_0)$ denote the initial intensity given before entering the first domain.

Next, let us focus on Eq. (3.29). Here we write it down again:

$$I_h(\eta_n) - I_\gamma(\eta_n) = \sin 2\Phi_n \{ \cos \gamma_n [L(\eta_{n-1}) + Z(\eta_{n-1})] + \sin \gamma_n [X(\eta_{n-1}) - N(\eta_{n-1})] \} \\ + \cos 2\Phi_n [I_h(\eta_{n-1}) - I_\gamma(\eta_{n-1})]. \quad (4.2)$$

As mentioned above, the angles θ_n and γ_n are chosen randomly. Then, for the purpose of finding the statistical means (expectation values), we can drop the linear terms concerning $\cos \gamma_n$ or $\sin \gamma_n$ in Eq. (4.2) by averaging over γ_n . On the other hand, the coefficient $\cos 2\Phi_n$ has the θ_n -dependence as is shown in Eq. (3.8), and the average over θ_n is nontrivial in general. Hence, we write the recursion relation for the mean as

$$\langle I_h(\eta_n) - I_\gamma(\eta_n) \rangle = \langle \cos 2\Phi_n \rangle \cdot \langle I_h(\eta_{n-1}) - I_\gamma(\eta_{n-1}) \rangle. \quad (4.3)$$

where the bracket $\langle \dots \rangle$ represents the average over various realizations of the magnetic field directions. It follows from Eq. (4.3) that

$$\langle I_h(\eta_N) - I_\gamma(\eta_N) \rangle = \Pi_N (I_{h,0} - I_{\gamma,0}), \quad (4.4)$$

where we defined

$$\Pi_N \equiv \prod_{n=1}^N \langle \cos 2\Phi_n \rangle. \quad (4.5)$$

Combining Eqs. (4.1) and (4.4), we obtain

$$\langle I_h(\eta_N) \rangle = \frac{1 + \Pi_N}{2} I_{h,0} + \frac{1 - \Pi_N}{2} I_{\gamma,0}, \quad (4.6)$$

$$\langle I_\gamma(\eta_N) \rangle = \frac{1 + \Pi_N}{2} I_{\gamma,0} + \frac{1 - \Pi_N}{2} I_{h,0}. \quad (4.7)$$

The bracket on the right-hand side in Eq. (4.5) consists only of the average over the angle θ_n . Let us rewrite the definition of Φ_n in Eq. (3.8) as

$$\Phi_n = \mathcal{M}_n \sin \theta_n, \quad (4.8)$$

where we defined

$$\mathcal{M}_n \equiv \int_{\eta_{n-1}}^{\eta_n} d\eta \frac{\kappa}{\sqrt{2}} \frac{\bar{B}_0}{a}. \quad (4.9)$$

Then, the average over θ_n yields

$$\begin{aligned} \langle \cos 2\Phi_n \rangle &\equiv \frac{1}{\pi} \int_0^\pi d\theta_n \cos(2\mathcal{M}_n \sin \theta_n) \\ &= J_0(2\mathcal{M}_n), \end{aligned} \quad (4.10)$$

where $J_0(\dots)$ represents the zeroth Bessel function of the first kind. As derived in Eq. (2.51), in the radiation-dominated epoch, the argument \mathcal{M}_n is evaluated as

$$\mathcal{M}_n = \frac{\kappa}{\sqrt{2}} \frac{\bar{B}_0}{H_0 \sqrt{\Omega_{r0}}} \ln \left(\frac{\eta_n}{\eta_{n-1}} \right) = 0.41 \left(\frac{\bar{B}_0}{10^{-6} \text{ Gauss}} \right) \ln \left(\frac{\eta_n}{\eta_{n-1}} \right). \quad (\text{RD}) \quad (4.11)$$

Note that there is a relation $\eta_n = \eta_{n-1} + s$ since we defined the comoving size of each domain as s .

Hereafter, we consider the case that the argument \mathcal{M}_n defined in Eq. (4.9) is large in a single domain, i.e., $\mathcal{M}_n \gg 1$. This essentially corresponds to the case that the length scale of conversion Δ_M^{-1} is much shorter than the domain size so that the conversion occurs efficiently within a domain. More specifically, considering the radiation-dominated epoch (4.11), we see that $\mathcal{M}_n \gg 1$ is realized when $s \gg \eta_{n-1}$, i.e., the value of the conformal time at entry into a domain is sufficiently smaller than the comoving size of the domain. The asymptotic behavior of Eq. (4.10) for large \mathcal{M}_n is given by

$$\langle \cos 2\Phi_n \rangle = J_0(2\mathcal{M}_n) \rightarrow \sqrt{\frac{1}{\pi \mathcal{M}_n}} \cos \left(2\mathcal{M}_n - \frac{\pi}{4} \right). \quad (4.12)$$

Moreover, in the infinite limit $\mathcal{M}_n \rightarrow \infty$, the mean $\langle \cos 2\Phi_n \rangle$ converges to zero and we obtain

$$\langle I_h(\eta) \rangle \rightarrow \frac{1}{2}(I_{h,0} + I_{\gamma,0}), \quad \langle I_\gamma(\eta) \rangle \rightarrow \frac{1}{2}(I_{h,0} + I_{\gamma,0}). \quad (4.13)$$

This result indicates that the equipartition of gravitational waves and gauge fields is realized

irrespective of the initial condition in the limit $\mathcal{M}_n \rightarrow \infty$.

4.1.2 Variance of intensity

Next, we proceed to study the behavior of the statistical variance of the intensity. This is achieved by taking the square of Eqs. (3.28)–(3.31) and taking the ensemble average, i.e., the average over the magnetic field directions. From Eq. (3.28), it follows that

$$\langle [I_h(\eta_n) + I_\gamma(\eta_n)]^2 \rangle = (I_{h,0} + I_{\gamma,0})^2. \quad (4.14)$$

On the other hand, from Eqs. (3.29)–(3.31), we obtain

$$\begin{pmatrix} \langle (I_{h,n} - I_{\gamma,n})^2 \rangle \\ \langle (L_n + Z_n)^2 \rangle \\ \langle (X_n - N_n)^2 \rangle \end{pmatrix} = \begin{pmatrix} \langle \cos^2 2\Phi_n \rangle & \frac{1}{2} \langle \sin^2 2\Phi_n \rangle & \frac{1}{2} \langle \sin^2 2\Phi_n \rangle \\ \langle \sin^2 2\Phi_n \rangle & \frac{1}{2} \langle \cos^2 2\Phi_n \rangle & \frac{1}{2} \langle \cos^2 2\Phi_n \rangle \\ 0 & 1/2 & 1/2 \end{pmatrix} \begin{pmatrix} \langle (I_{h,n-1} - I_{\gamma,n-1})^2 \rangle \\ \langle (L_{n-1} + Z_{n-1})^2 \rangle \\ \langle (X_{n-1} - N_{n-1})^2 \rangle \end{pmatrix}, \quad (4.15)$$

where the subscripts n and $n-1$ attached to the parameters represent that they are evaluated at time η_n and η_{n-1} , respectively. In the right-hand side of Eq. (4.15), the average over the angle γ_n was already performed and the average over θ_n is indicated by the bracket $\langle \dots \rangle$.

As we have done in Eq. (4.13), focusing on the efficient conversion limit, i.e., $\mathcal{M}_n \rightarrow \infty$, we can simplify Eq. (4.15) as

$$\begin{pmatrix} \langle (I_{h,n} - I_{\gamma,n})^2 \rangle \\ \langle (L_n + Z_n)^2 \rangle \\ \langle (X_n - N_n)^2 \rangle \end{pmatrix} = \begin{pmatrix} 1/2 & 1/4 & 1/4 \\ 1/2 & 1/4 & 1/4 \\ 0 & 1/2 & 1/2 \end{pmatrix} \begin{pmatrix} \langle (I_{h,n-1} - I_{\gamma,n-1})^2 \rangle \\ \langle (L_{n-1} + Z_{n-1})^2 \rangle \\ \langle (X_{n-1} - N_{n-1})^2 \rangle \end{pmatrix} \equiv \mathbf{W}_1 \begin{pmatrix} \langle (I_{h,n-1} - I_{\gamma,n-1})^2 \rangle \\ \langle (L_{n-1} + Z_{n-1})^2 \rangle \\ \langle (X_{n-1} - N_{n-1})^2 \rangle \end{pmatrix}. \quad (4.16)$$

The coefficient matrix \mathbf{W}_1 can be diagonalized as

$$\mathbf{P}_1^{-1} \mathbf{W}_1 \mathbf{P}_1 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1/4 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad \mathbf{P}_1 = \begin{pmatrix} 1 & -1/2 & 0 \\ 1 & -1/2 & -1 \\ 1 & 1 & 1 \end{pmatrix}. \quad (4.17)$$

Therefore, the asymptotic values can be derived as

$$\begin{aligned}
\begin{pmatrix} \langle (I_{h,n} - I_{\gamma,n})^2 \rangle \\ \langle (L_n + Z_n)^2 \rangle \\ \langle (X_n - N_n)^2 \rangle \end{pmatrix} &= \mathbf{P}_1 \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1/4 & 0 \\ 0 & 0 & 0 \end{pmatrix}^n \mathbf{P}_1^{-1} \begin{pmatrix} (I_{h,0} - I_{\gamma,0})^2 \\ (L_0 + Z_0)^2 \\ (X_0 - N_0)^2 \end{pmatrix} \\
&\xrightarrow{n \rightarrow \infty} \mathbf{P}_1 \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \mathbf{P}_1^{-1} \begin{pmatrix} (I_{h,0} - I_{\gamma,0})^2 \\ (L_0 + Z_0)^2 \\ (X_0 - N_0)^2 \end{pmatrix} \\
&= \begin{pmatrix} 1/3 & 1/3 & 1/3 \\ 1/3 & 1/3 & 1/3 \\ 1/3 & 1/3 & 1/3 \end{pmatrix} \begin{pmatrix} (I_{h,0} - I_{\gamma,0})^2 \\ (L_0 + Z_0)^2 \\ (X_0 - N_0)^2 \end{pmatrix}. \tag{4.18}
\end{aligned}$$

Combining Eqs. (4.14) and (4.18) yields the asymptotic behavior,

$$\langle I_h^2(\eta) \rangle + \langle I_\gamma^2(\eta) \rangle \rightarrow \frac{2}{3}(I_{h,0}^2 + I_{\gamma,0}^2 + I_{h,0}I_{\gamma,0}) + \frac{1}{6}[(L_0 + Z_0)^2 + (X_0 - N_0)^2]. \tag{4.19}$$

Furthermore, it is legitimate to expect equality between $\langle I_h^2 \rangle$ and $\langle I_\gamma^2 \rangle$ in the case of efficient conversion. In this case, we obtain

$$\langle I_h^2(\eta) \rangle \rightarrow \frac{1}{3}(I_{h,0}^2 + I_{\gamma,0}^2 + I_{h,0}I_{\gamma,0}) + \frac{1}{12}[(L_0 + Z_0)^2 + (X_0 - N_0)^2]. \tag{4.20}$$

This equation gives the asymptotic value of the statistical variance of the gravitational wave intensity in terms of the initial Stokes parameters in the case of efficient conversion through a number of random magnetic field domains.

As a by-product, we can also derive the asymptotic value of the correlation $\langle I_h I_\gamma \rangle$ in the same case as

$$\langle I_h(\eta) I_\gamma(\eta) \rangle \rightarrow \frac{1}{6}(I_{h,0}^2 + I_{\gamma,0}^2) + \frac{2}{3}I_{h,0}I_{\gamma,0} - \frac{1}{12}[(L_0 + Z_0)^2 + (X_0 - N_0)^2]. \tag{4.21}$$

4.2 Circular polarization

Next, we turn to Class II, i.e., Eqs. (3.32)–(3.35), to reveal the asymptotic behavior of the circular polarization denoted by $V_{h/\gamma}$. The analysis is parallel to that of intensity.

4.2.1 Mean of circular polarization

Equation (3.32) shows that, similarly to intensity, the total circular polarization of the gravitational waves and gauge fields is conserved,

$$V_h(\eta) + V_\gamma(\eta) = V_{h,0} + V_{\gamma,0}, \quad (4.22)$$

where $V_{h,0} = V_h(\eta_0)$ and $V_{\gamma,0} = V_\gamma(\eta_0)$ denote the initial circular polarization. The conservation of the total circular polarization follows from the fact that the $+$ and \times modes are separated and evolve in the same manner.

As performed in Eq. (4.3), we take the average of Eq. (3.33) over the magnetic field directions and write the recurrence relation for the mean as

$$\langle V_h(\eta_n) - V_\gamma(\eta_n) \rangle = \langle \cos 2\Phi_n \rangle \cdot \langle V_h(\eta_{n-1}) - V_\gamma(\eta_{n-1}) \rangle. \quad (4.23)$$

Comparing Eqs. (4.3) and (4.23), we see that the mean of the difference of the circular polarization $\langle V_h - V_\gamma \rangle$ has the same structure as that of the intensity $\langle I_h - I_\gamma \rangle$. Combining this with Eq. (4.22), the means of the circular polarization are given by

$$\langle V_h(\eta_N) \rangle = \frac{1 + \Pi_N}{2} V_{h,0} + \frac{1 - \Pi_N}{2} V_{\gamma,0}, \quad (4.24)$$

$$\langle V_\gamma(\eta_N) \rangle = \frac{1 + \Pi_N}{2} V_{\gamma,0} + \frac{1 - \Pi_N}{2} V_{h,0}, \quad (4.25)$$

where Π_N is defined in Eq. (4.5). Furthermore, in the efficient conversion limit $\mathcal{M}_n \rightarrow \infty$, the degree of circular polarization becomes equally distributed between gravitational waves and gauge fields,

$$\langle V_h(\eta) \rangle \rightarrow \frac{1}{2}(V_{h,0} + V_{\gamma,0}), \quad \langle V_\gamma(\eta) \rangle \rightarrow \frac{1}{2}(V_{h,0} + V_{\gamma,0}). \quad (4.26)$$

4.2.2 Variance of circular polarization

The statistical variance of the circular polarization can be found by taking the square of Eqs. (3.32)–(3.35). First, Eq. (3.32) tells us

$$\langle [V_h(\eta_n) + V_\gamma(\eta_n)]^2 \rangle = (V_{h,0} + V_{\gamma,0})^2. \quad (4.27)$$

On the other hand, taking the square of Eqs. (3.33)–(3.35) and averaging them over the angles θ_n and γ_n , we obtain

$$\begin{pmatrix} \langle (V_{h,n} - V_{\gamma,n})^2 \rangle \\ \langle (M_n - W_n)^2 \rangle \\ \langle (K_n + Y_n)^2 \rangle \end{pmatrix} = \begin{pmatrix} \langle \cos^2 2\Phi_n \rangle & \frac{1}{2} \langle \sin^2 2\Phi_n \rangle & \frac{1}{2} \langle \sin^2 2\Phi_n \rangle \\ \langle \sin^2 2\Phi_n \rangle & \frac{1}{2} \langle \cos^2 2\Phi_n \rangle & \frac{1}{2} \langle \cos^2 2\Phi_n \rangle \\ 0 & 1/2 & 1/2 \end{pmatrix} \begin{pmatrix} \langle (V_{h,n-1} - V_{\gamma,n-1})^2 \rangle \\ \langle (M_{n-1} - W_{n-1})^2 \rangle \\ \langle (K_{n-1} + Y_{n-1})^2 \rangle \end{pmatrix}. \quad (4.28)$$

When focusing on the efficient conversion limit $\mathcal{M}_n \rightarrow \infty$, we can simplify Eq. (4.28) as

$$\begin{pmatrix} \langle (V_{h,n} - V_{\gamma,n})^2 \rangle \\ \langle (M_n - W_n)^2 \rangle \\ \langle (K_n + Y_n)^2 \rangle \end{pmatrix} = \begin{pmatrix} 1/2 & 1/4 & 1/4 \\ 1/2 & 1/4 & 1/4 \\ 0 & 1/2 & 1/2 \end{pmatrix} \begin{pmatrix} \langle (V_{h,n-1} - V_{\gamma,n-1})^2 \rangle \\ \langle (M_{n-1} - W_{n-1})^2 \rangle \\ \langle (K_{n-1} + Y_{n-1})^2 \rangle \end{pmatrix} \equiv \mathbf{W}_2 \begin{pmatrix} \langle (V_{h,n-1} - V_{\gamma,n-1})^2 \rangle \\ \langle (M_{n-1} - W_{n-1})^2 \rangle \\ \langle (K_{n-1} + Y_{n-1})^2 \rangle \end{pmatrix}. \quad (4.29)$$

Note that the coefficient matrix \mathbf{W}_2 is the same as \mathbf{W}_1 in Eq. (4.16). Therefore, the asymptotic values are obtained in the same way as Eq. (4.18):

$$\begin{pmatrix} \langle (V_{h,n} - V_{\gamma,n})^2 \rangle \\ \langle (M_n - W_n)^2 \rangle \\ \langle (K_n + Y_n)^2 \rangle \end{pmatrix} \xrightarrow{n \rightarrow \infty} \begin{pmatrix} 1/3 & 1/3 & 1/3 \\ 1/3 & 1/3 & 1/3 \\ 1/3 & 1/3 & 1/3 \end{pmatrix} \begin{pmatrix} (V_{h,0} - V_{\gamma,0})^2 \\ (M_0 - W_0)^2 \\ (K_0 + Y_0)^2 \end{pmatrix}. \quad (4.30)$$

Combining Eqs. (4.27) and (4.30), we obtain the asymptotic behavior,

$$\langle V_h^2(\eta) \rangle + \langle V_\gamma^2(\eta) \rangle \rightarrow \frac{2}{3}(V_{h,0}^2 + V_{\gamma,0}^2 + V_{h,0}V_{\gamma,0}) + \frac{1}{6}[(M_0 - W_0)^2 + (K_0 + Y_0)^2]. \quad (4.31)$$

Furthermore, assuming equality $\langle V_h^2 \rangle = \langle V_\gamma^2 \rangle$, we obtain

$$\langle V_h^2(\eta) \rangle \rightarrow \frac{1}{3}(V_{h,0}^2 + V_{\gamma,0}^2 + V_{h,0}V_{\gamma,0}) + \frac{1}{12}[(M_0 - W_0)^2 + (K_0 + Y_0)^2]. \quad (4.32)$$

This equation represents the asymptotic value of the statistical variance of the circular polarization of gravitational waves in terms of the initial Stokes parameters. The assumptions in derivation were that conversion is efficient in a single domain and that the waves pass through many domains where the direction of the magnetic field is random.

We can also obtain the asymptotic value of the correlation $\langle V_h V_\gamma \rangle$ as

$$\langle V_h(\eta) V_\gamma(\eta) \rangle \rightarrow \frac{1}{6}(V_{h,0}^2 + V_{\gamma,0}^2) + \frac{2}{3}V_{h,0}V_{\gamma,0} - \frac{1}{12}[(M_0 - W_0)^2 + (K_0 + Y_0)^2]. \quad (4.33)$$

4.3 Linear polarization

Finally, we study the asymptotic behavior of the linear polarization denoted by $Q_{h/\gamma}$ and $U_{h/\gamma}$ for which Class III, i.e., the set of Eqs. (3.36)–(3.39), is relevant.

4.3.1 Mean of linear polarization

As we have done for the intensity $I_{h/\gamma}$ and circular polarization $V_{h/\gamma}$, we can express the parameters $Q_{h/\gamma}$ and $U_{h/\gamma}$ in the n -th domain in terms of the Stokes parameters in the $(n - 1)$ -th domain from Eqs. (3.36) and (3.37). In this case, only linear terms with respect to $\cos \gamma_n$, $\cos 2\gamma_n$, and so on, appear. Then, averaging over the angle γ_n leads to

$$\langle Q_h(\eta) \rangle = \langle Q_\gamma(\eta) \rangle = \langle U_h(\eta) \rangle = \langle U_\gamma(\eta) \rangle = 0. \quad (4.34)$$

Hence, the means (expectation values) of linear polarization vanish.

4.3.2 Variance of linear polarization

The statistical variance of linear polarization can be studied by taking the square of Eqs. (3.36)–(3.39) and taking the average over the magnetic field directions characterized by θ_n and γ_n .

Indeed, we obtain the recurrence relation for the average of the square as

$$\begin{pmatrix} \langle Q_{h,n}^2 + U_{h,n}^2 \rangle \\ \langle Q_{\gamma,n}^2 + U_{\gamma,n}^2 \rangle \\ \langle (L_n - Z_n)^2 \rangle \\ \langle (X_n + N_n)^2 \rangle \end{pmatrix} = \begin{pmatrix} \langle \cos^4 \Phi_n \rangle & \langle \sin^4 \Phi_n \rangle & \frac{1}{4} \langle \sin^2 2\Phi_n \rangle & \frac{1}{4} \langle \sin^2 2\Phi_n \rangle \\ \langle \sin^4 \Phi_n \rangle & \langle \cos^4 \Phi_n \rangle & \frac{1}{4} \langle \sin^2 2\Phi_n \rangle & \frac{1}{4} \langle \sin^2 2\Phi_n \rangle \\ \frac{1}{2} \langle \sin^2 2\Phi_n \rangle & \frac{1}{2} \langle \sin^2 2\Phi_n \rangle & \frac{1}{2} \langle \cos^2 2\Phi_n \rangle & \frac{1}{2} \langle \cos^2 2\Phi_n \rangle \\ \frac{1}{2} \langle \sin^2 2\Phi_n \rangle & \frac{1}{2} \langle \sin^2 2\Phi_n \rangle & \frac{1}{2} \langle \cos^2 2\Phi_n \rangle & \frac{1}{2} \langle \cos^2 2\Phi_n \rangle \end{pmatrix} \begin{pmatrix} \langle Q_{h,n-1}^2 + U_{h,n-1}^2 \rangle \\ \langle Q_{\gamma,n-1}^2 + U_{\gamma,n-1}^2 \rangle \\ \langle (L_{n-1} - Z_{n-1})^2 \rangle \\ \langle (X_{n-1} + N_{n-1})^2 \rangle \end{pmatrix}. \quad (4.35)$$

When focusing on the efficient conversion limit $\mathcal{M}_n \rightarrow \infty$, Eq. (4.35) is further simplified:

$$\begin{pmatrix} \langle Q_{h,n}^2 + U_{h,n}^2 \rangle \\ \langle Q_{\gamma,n}^2 + U_{\gamma,n}^2 \rangle \\ \langle (L_n - Z_n)^2 \rangle \\ \langle (X_n + N_n)^2 \rangle \end{pmatrix} = \begin{pmatrix} 3/8 & 3/8 & 1/8 & 1/8 \\ 3/8 & 3/8 & 1/8 & 1/8 \\ 1/4 & 1/4 & 1/4 & 1/4 \\ 1/4 & 1/4 & 1/4 & 1/4 \end{pmatrix} \begin{pmatrix} \langle Q_{h,n-1}^2 + U_{h,n-1}^2 \rangle \\ \langle Q_{\gamma,n-1}^2 + U_{\gamma,n-1}^2 \rangle \\ \langle (L_{n-1} - Z_{n-1})^2 \rangle \\ \langle (X_{n-1} + N_{n-1})^2 \rangle \end{pmatrix} \equiv \mathbf{W}_3 \begin{pmatrix} \langle Q_{h,n-1}^2 + U_{h,n-1}^2 \rangle \\ \langle Q_{\gamma,n-1}^2 + U_{\gamma,n-1}^2 \rangle \\ \langle (L_{n-1} - Z_{n-1})^2 \rangle \\ \langle (X_{n-1} + N_{n-1})^2 \rangle \end{pmatrix}. \quad (4.36)$$

The coefficient matrix \mathbf{W}_3 can be diagonalized as

$$\mathbf{P}_3^{-1} \mathbf{W}_3 \mathbf{P}_3 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1/4 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad \mathbf{P}_3 = \begin{pmatrix} 1 & -1/2 & 0 & -1 \\ 1 & -1/2 & 0 & 1 \\ 1 & 1 & -1 & 0 \\ 1 & 1 & 1 & 0 \end{pmatrix}. \quad (4.37)$$

Hence, we find the asymptotic values through the infinite number of domains $n \rightarrow \infty$ as

$$\begin{aligned}
\begin{pmatrix} \langle Q_{h,n}^2 + U_{h,n}^2 \rangle \\ \langle Q_{\gamma,n}^2 + U_{\gamma,n}^2 \rangle \\ \langle (L_n - Z_n)^2 \rangle \\ \langle (X_n + N_n)^2 \rangle \end{pmatrix} &= \mathbf{P}_3 \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1/4 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}^n \mathbf{P}_3^{-1} \begin{pmatrix} Q_{h,0}^2 + U_{h,0}^2 \\ Q_{\gamma,0}^2 + U_{\gamma,0}^2 \\ (L_0 - Z_0)^2 \\ (X_0 + N_0)^2 \end{pmatrix} \\
&\xrightarrow{n \rightarrow \infty} \mathbf{P}_3 \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \mathbf{P}_3^{-1} \begin{pmatrix} Q_{h,0}^2 + U_{h,0}^2 \\ Q_{\gamma,0}^2 + U_{\gamma,0}^2 \\ (L_0 - Z_0)^2 \\ (X_0 + N_0)^2 \end{pmatrix} \\
&= \begin{pmatrix} 1/3 & 1/3 & 1/6 & 1/6 \\ 1/3 & 1/3 & 1/6 & 1/6 \\ 1/3 & 1/3 & 1/6 & 1/6 \\ 1/3 & 1/3 & 1/6 & 1/6 \end{pmatrix} \begin{pmatrix} Q_{h,0}^2 + U_{h,0}^2 \\ Q_{\gamma,0}^2 + U_{\gamma,0}^2 \\ (L_0 - Z_0)^2 \\ (X_0 + N_0)^2 \end{pmatrix}. \tag{4.38}
\end{aligned}$$

In particular, the first component represents the asymptotic value of the variance of the linear polarization of gravitational waves in terms of the initial Stokes parameters:

$$\langle Q_h^2(\eta) \rangle + \langle U_h^2(\eta) \rangle \rightarrow \frac{1}{3}[(Q_{h,0}^2 + U_{h,0}^2) + (Q_{\gamma,0}^2 + U_{\gamma,0}^2)] + \frac{1}{6}[(L_0 - Z_0)^2 + (X_0 + N_0)^2]. \tag{4.39}$$

5 Conclusion and discussion

Gravitational waves will offer a novel and powerful tool for exploring the early universe. Furthermore, since gravitational waves have universal coupling with particles in the dark sector, they can be useful in probing the dark sector. In this paper, we considered the conversion processes between gravitons and dark photons in cosmological dark magnetic fields and investigated their imprints on the gravitational waves. Specifically, we considered a sequence of magnetic domains in which the direction of the magnetic field is randomly given. Then, we studied the asymptotic behavior of the intensity and polarization of gravitational waves when they pass through many domains.

As indicated in Sec. 3.2, the evolution equations for gravitational wave intensity, circular polarization, and linear polarization are separated in the framework of minimal coupling between gravitons and dark photons. This is clearly different from the case of axion-photon conversion [18, 19], where axions are converted only to the polarization component of photons parallel to the magnetic field.

We explicitly showed that the statistical means (expectation values) of the intensity and circular polarization become equally distributed between gravitational waves and dark photons as Eqs. (4.13) and (4.26) when the conversion is efficient. Besides, the mean of the linear polarization vanishes as Eq. (4.34), which is due to the randomness of the magnetic field directions. Going further, we obtained asymptotic values of the statistical variance of the intensity and polarization of gravitational waves in terms of the initial Stokes parameters as Eqs. (4.20), (4.32), and (4.39). We should emphasize that the obtained results can be used to infer the initial polarization of dark photons. For example, if the primordial circular polarization of dark photons exists, it can be transferred to that of gravitational waves as Eq. (4.24). Hence, we can get information on primordial dark photons through the observations of gravitational waves. Furthermore, observational confirmation of the predicted variance (4.32) will corroborate that the conversion occurred through random magnetic fields.

It will be important to study to what extent the results in this paper depend on the setup. In particular, non-minimal coupling between gravitons and dark photons and peculiar magnetic field configurations may alter the results. We leave these issues for future work.

Acknowledgments

We appreciate Hidetoshi Omiya and Ann Mukuno for their useful comments and daily discussions of related topics. K. N. was supported by JSPS KAKENHI Grant Number JP24KJ0117. J. S. was in part supported by JSPS KAKENHI Grant Numbers JP23K22491, and JP24K21548. K. U. was supported by JSPS KAKENHI Grant Number JP24K17050. Z. W. appreciates Kobe University for Hospitality when the project is initiated. Z. W. was supported from Ministry of Science and Technology of China by a special grant.

References

- [1] B. P. Abbott et al. “Observation of Gravitational Waves from a Binary Black Hole Merger”. In: *Phys. Rev. Lett.* 116.6 (2016), p. 061102. arXiv: 1602.03837 [gr-qc].
- [2] Alberto Sesana. “Prospects for Multiband Gravitational-Wave Astronomy after GW150914”. In: *Phys. Rev. Lett.* 116.23 (2016), p. 231102. arXiv: 1602.06951 [gr-qc].
- [3] Misao Sasaki et al. “Primordial black holes—perspectives in gravitational wave astronomy”. In: *Class. Quant. Grav.* 35.6 (2018), p. 063001. arXiv: 1801.05235 [astro-ph.CO].
- [4] Jihn E. Kim. “Light Pseudoscalars, Particle Physics and Cosmology”. In: *Phys. Rept.* 150 (1987), pp. 1–177.
- [5] S. Akiyama et al. “Interplanetary Magnetic Flux Rope Observed at Ground Level by HAWC”. In: *Astrophys. J.* 905.1 (2020), p. 73. arXiv: 2101.03243 [astro-ph.SR].
- [6] Andrei P. Igoshev, Sergei B. Popov, and Rainer Hollerbach. “Evolution of Neutron Star Magnetic Fields”. In: *Universe* 7.9 (2021), p. 351. arXiv: 2109.05584 [astro-ph.HE].
- [7] Lawrence M. Widrow. “Origin of galactic and extragalactic magnetic fields”. In: *Rev. Mod. Phys.* 74 (2002), pp. 775–823. arXiv: astro-ph/0207240.
- [8] Ronnie Jansson and Glennys R. Farrar. “A New Model of the Galactic Magnetic Field”. In: *Astrophys. J.* 757 (2012), p. 14. arXiv: 1204.3662 [astro-ph.GA].
- [9] F. Aharonian et al. “Constraints on the Intergalactic Magnetic Field Using Fermi-LAT and H.E.S.S. Blazar Observations”. In: *Astrophys. J. Lett.* 950.2 (2023), p. L16. arXiv: 2306.05132 [astro-ph.HE].
- [10] J. Tjemsland, M. Meyer, and F. Vazza. “Constraining the astrophysical origin of intergalactic magnetic fields”. In: (Nov. 2023). arXiv: 2311.04273 [astro-ph.HE].
- [11] V. A. Acciari et al. “A lower bound on intergalactic magnetic fields from time variability of 1ES 0229+200 from MAGIC and Fermi/LAT observations”. In: *Astron. Astrophys.* 670 (2023), A145. arXiv: 2210.03321 [astro-ph.HE].
- [12] Dario Grasso and Hector R. Rubinstein. “Magnetic fields in the early universe”. In: *Phys. Rept.* 348 (2001), pp. 163–266. arXiv: astro-ph/0009061.

- [13] Ruth Durrer and Andrii Neronov. “Cosmological Magnetic Fields: Their Generation, Evolution and Observation”. In: *Astron. Astrophys. Rev.* 21 (2013), p. 62. arXiv: 1303.7121 [astro-ph.CO].
- [14] Georg Raffelt and Leo Stodolsky. “Mixing of the Photon with Low Mass Particles”. In: *Phys. Rev. D* 37 (1988), p. 1237.
- [15] L. Maiani, R. Petronzio, and E. Zavattini. “Effects of Nearly Massless, Spin Zero Particles on Light Propagation in a Magnetic Field”. In: *Phys. Lett. B* 175 (1986), pp. 359–363.
- [16] Yuval Grossman, Sourov Roy, and Jure Zupan. “Effects of initial axion production and photon axion oscillation on type Ia supernova dimming”. In: *Phys. Lett. B* 543 (2002), pp. 23–28. arXiv: hep-ph/0204216.
- [17] Alessandro Mirizzi and Daniele Montanino. “Stochastic conversions of TeV photons into axion-like particles in extragalactic magnetic fields”. In: *JCAP* 12 (2009), p. 004. arXiv: 0911.0015 [astro-ph.HE].
- [18] Nicola Bassan, Alessandro Mirizzi, and Marco Roncadelli. “Axion-like particle effects on the polarization of cosmic high-energy gamma sources”. In: *JCAP* 05 (2010), p. 010. arXiv: 1001.5267 [astro-ph.HE].
- [19] Emi Masaki, Arata Aoki, and Jiro Soda. “Photon-Axion Conversion, Magnetic Field Configuration, and Polarization of Photons”. In: *Phys. Rev. D* 96.4 (2017), p. 043519. arXiv: 1702.08843 [astro-ph.CO].
- [20] Konstantinos Dimopoulos. “Can a vector field be responsible for the curvature perturbation in the Universe?” In: *Phys. Rev. D* 74 (2006), p. 083502. arXiv: hep-ph/0607229.
- [21] Kazunori Nakayama. “Vector Coherent Oscillation Dark Matter”. In: *JCAP* 10 (2019), p. 019. arXiv: 1907.06243 [hep-ph].
- [22] Kazunori Nakayama. “Constraint on Vector Coherent Oscillation Dark Matter with Kinetic Function”. In: *JCAP* 08 (2020), p. 033. arXiv: 2004.10036 [hep-ph].

- [23] Yuichiro Nakai, Ryo Namba, and Ziwei Wang. “Light Dark Photon Dark Matter from Inflation”. In: *JHEP* 12 (2020), p. 170. arXiv: 2004.10743 [hep-ph].
- [24] Robert Foot and Xiao-Gang He. “Comment on Z Z-prime mixing in extended gauge theories”. In: *Phys. Lett. B* 267 (1991), pp. 509–512.
- [25] Robert Foot. “Mirror matter-type dark matter”. In: *Int. J. Mod. Phys. D* 13 (2004), pp. 2161–2192. arXiv: astro-ph/0407623.
- [26] R. Foot. “Mirror dark matter: Cosmology, galaxy structure and direct detection”. In: *Int. J. Mod. Phys. A* 29 (2014), p. 1430013. arXiv: 1401.3965 [astro-ph.CO].
- [27] Emi Masaki and Jiro Soda. “Conversion of Gravitons into Dark Photons in Cosmological Dark Magnetic Fields”. In: *Phys. Rev. D* 98.2 (2018), p. 023540. arXiv: 1804.00458 [astro-ph.CO].
- [28] M.E. Gertsenshtein. “Wave Resonance of Light and Gravitational Waves”. In: *J.Exp.Theor.Phys.* 14 (1962), p. 84.
- [29] A. Neronov and I. Vovk. “Evidence for strong extragalactic magnetic fields from Fermi observations of TeV blazars”. In: *Science* 328 (2010), pp. 73–75. arXiv: 1006.3504 [astro-ph.HE].
- [30] A. M. Taylor, I. Vovk, and A. Neronov. “Extragalactic magnetic fields constraints from simultaneous GeV-TeV observations of blazars”. In: *Astron. Astrophys.* 529 (2011), A144. arXiv: 1101.0932 [astro-ph.HE].
- [31] Zi-Qing Xia et al. “A delayed 400 GeV photon from GRB 221009A and implication on the intergalactic magnetic field”. In: *Nature Commun.* 15.1 (2024), p. 4280. arXiv: 2210.13052 [astro-ph.HE].
- [32] Dai G. Yamazaki et al. “The Search for a Primordial Magnetic Field”. In: *Phys. Rept.* 517 (2012), pp. 141–167. arXiv: 1204.3669 [astro-ph.CO].
- [33] Masahiro Kawasaki and Motohiko Kusakabe. “Updated constraint on a primordial magnetic field during big bang nucleosynthesis and a formulation of field effects”. In: *Phys. Rev. D* 86 (2012), p. 063003. arXiv: 1204.6164 [astro-ph.CO].

- [34] Sugumi Kanno et al. “A peak in the power spectrum of primordial gravitational waves induced by primordial dark magnetic fields”. In: *JCAP* 05 (2023), p. 052. arXiv: 2301.13540 [hep-th].
- [35] Pisin Chen and Teruaki Suyama. “Constraining Primordial Magnetic Fields by CMB Photon-Graviton Conversion”. In: *Phys. Rev. D* 88.12 (2013), p. 123521. arXiv: 1309.0537 [astro-ph.CO].