

Quantized axial charge of staggered fermions and the chiral anomaly

Arkya Chatterjee[✉], Salvatore D. Pace[✉], Shu-Heng Shao^{✉✉}

[✉]*Department of Physics, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA*

[✉]*Yang Institute for Theoretical Physics, Stony Brook University, Stony Brook 11794, USA*

In the 1+1D ultra-local lattice Hamiltonian for staggered fermions with a finite-dimensional Hilbert space, there are two conserved, integer-valued charges that flow in the continuum limit to the vector and axial charges of a massless Dirac fermion with a perturbative anomaly. Each of the two lattice charges generates an ordinary U(1) global symmetry that acts locally on operators and can be gauged individually. Interestingly, they do not commute on a finite lattice, but their commutator goes to zero in the continuum limit. This non-abelian algebra presents an alternative lattice realization of the vector and axial symmetries, which is consistent with the Nielsen-Ninomiya theorem. We further prove that the presence of these two conserved lattice charges forces the low-energy phase to be gapless, reminiscent of the consequence from perturbative anomalies of continuous global symmetries in continuum field theory. Upon bosonization, these two charges lead to two exact U(1) symmetries in the XY model that flow to the momentum and winding symmetries in the free boson conformal field theory.

CONTENTS

I. Introduction	1
II. Symmetries and anomalies in the continuum	2
III. Axial charges of staggered fermions	3
A. Quantized axial charge	3
B. Unquantized axial charge	4
C. From the lattice to the continuum	5
D. The non-abelian algebra	5
E. Comparison with other constructions	6
IV. Coupling to gauge fields	6
A. Gauging the vector symmetry	6
B. Gauging the axial symmetry	6
V. A lattice anomaly as an obstruction to gapped phases	7
VI. Exact winding symmetry in the XY model	7
VII. Conclusion	8
Acknowledgments	8
References	8
A. Continuum limit of $[Q^V, Q^A]$	10
B. Deformations preserving both Q^V and Q^A	11

I. INTRODUCTION

The realization of chiral symmetries on the lattice has been a long-standing problem in lattice field theory [1]. One prominent difficulty is that the global symmetries of interest typically have anomalies, in the sense that

they cannot be coupled to dynamical gauge fields in a consistent manner.¹

Anomalies are traditionally linked to divergences in continuum field theory and are often considered challenging, if not impossible, to realize on a lattice with finite lattice spacing. However, this piece of lore is not true. Numerous examples of anomalies for discrete global symmetries are realized on the lattice. See, for example, Ref. [6] for a recent survey of this topic.

A more refined piece of lore is that perturbative anomalies of continuous global symmetries cannot be realized on a lattice. By perturbative or local anomalies, we mean those captured by Feynman diagrams and local operator product expansions. This piece of lore also turns out to be incorrect: perturbative anomalies can be realized in Villain-type lattice models [6–11]. The local Hilbert spaces of these models involve continuous variables and are infinite-dimensional. See Ref. [11] for discussions of chiral fermion symmetry in 1+1D QED in this setup, and Ref. [12] for a Euclidean lattice realization of mixed gravitational anomalies with global symmetries.

Perhaps a more interesting question is: *can perturbative anomalies be realized in a lattice model whose local Hilbert space is finite-dimensional (e.g., qubits)?*

¹ Anomalies of global symmetries are often called 't Hooft anomalies, which indicate that even though the global symmetries are conserved and exact at the quantum level, there is an obstruction to gauging them. This contrasts with the Schwinger anomaly [2] and the Adler-Bell-Jackiw (ABJ) anomaly [3, 4], where a classical symmetry fails to be a true global symmetry at the quantum level. See, for example, Ref. [5] for this terminology. These two notions of anomalies are not unrelated. For example, the Schwinger anomaly of the axial symmetry in 1+1D QED arises from the 't Hooft anomaly involving both the axial and vector symmetries of a free Dirac fermion. The violation of the axial symmetry when the vector symmetry is gauged implies an obstruction to gauging both global symmetries simultaneously.

Figure 1. In an ultra-local Hamiltonian lattice model, we find two conserved, quantized charges Q^V, Q^A , which become the vector and axial charges $\mathcal{Q}^V, \mathcal{Q}^A$ with a mixed anomaly in the continuum limit. (We use calligraphic and ordinary fonts for operators in the continuum and on the lattice, respectively.)

There is a simple no-go argument against this hope. This can be seen most readily for $U(1)$ symmetries in 1+1D, where perturbative anomalies are encoded in the equal-time commutator of the conserved current operators $[j^0(t, x), j^x(t, x')] \sim i\partial_x \delta(x - x')$, known as the Schwinger term [13, 14].² Taking the trace of this relation immediately shows that the Schwinger term cannot be realized verbatim on a finite-dimensional Hilbert space since the trace of a commutator in a finite-dimensional vector space necessarily vanishes. This is analogous to the fact that $[X, P] = i\hbar$ cannot be realized in a finite-dimensional Hilbert space in quantum mechanics. The Schwinger term, however, has recently been realized exactly on a lattice model with an infinite-dimensional Hilbert space [9].

These difficulties are rigorously formulated in the Nielsen-Ninomiya theorem [15–17]. The theorem, most precisely formulated in [18], asserts that in any quadratic lattice fermion Hamiltonian with certain locality properties in odd spatial dimensions, there must be an equal number of left- and right-moving fermions within each irreducible representation of the global symmetries.³ In particular, this theorem prohibits the existence of an axial charge that (i) has quantized, integer eigenvalues and (ii) commutes with the vector $U(1)$ symmetry. If such an axial charge existed, one could restrict to the fixed charge sector of both the vector and axial symmetries and would find a single left-moving fermion.

Given the above, what fingerprints of anomalies can we hope for on a lattice with a finite-dimensional Hilbert space? In this letter, we focus on the anomaly between the vector and axial symmetries of a 1+1D massless Dirac fermion, which is the oldest and arguably the simplest anomaly of all. In a microscopic Hamiltonian lattice realization, we discuss two conserved, quantized charges that become the vector and axial charges in the continuum limit. While each of the two charges generates an ordinary $U(1)$ symmetry satisfying all the locality properties, they do *not* commute on a finite-size lattice. Together they form a non-abelian algebra, which is consistent with the Nielsen-Ninomiya theorem and can be

viewed as the lattice avatar of the continuum anomaly (see Figure 1).

II. SYMMETRIES AND ANOMALIES IN THE CONTINUUM

We start with a review of the symmetries and anomalies of the massless Dirac fermion field theory in 1+1D. We denote the left- and right-moving (complex, one-component) Weyl fermions as $\Psi_L(t, x)$ and $\Psi_R(t, x)$, respectively. The action for a free, massless Dirac fermion $\Psi = (\Psi_L \ \Psi_R)^T$ is

$$S = i \int dt dx \bar{\Psi} \Gamma^\mu \partial_\mu \Psi = i \int dt dx \left[\Psi_L^\dagger (\partial_t - \partial_x) \Psi_L + \Psi_R^\dagger (\partial_t + \partial_x) \Psi_R \right], \quad (1)$$

where our conventions for the Dirac matrices are $\Gamma^0 = \sigma^x$, $\Gamma^1 = -i\sigma^y$ and $\bar{\Psi} = \Psi^\dagger \Gamma^0$. The internal global symmetry for the right-moving Weyl fermion is $O(2)^R = U(1)^R \rtimes \mathbb{Z}_2^{\mathcal{C}^R}$, where $U(1)^R$ and the charge conjugation symmetry $\mathbb{Z}_2^{\mathcal{C}^R}$ act on the fermion as $e^{i\varphi \mathcal{Q}^R} \Psi_R e^{-i\varphi \mathcal{Q}^R} = e^{-i\varphi} \Psi_R$ and $\mathcal{C}^R \Psi_R (\mathcal{C}^R)^{-1} = \Psi_R^\dagger$, respectively. Here \mathcal{Q}^R is the quantized charge for the right movers, which obeys $\mathcal{C}^R \mathcal{Q}^R (\mathcal{C}^R)^{-1} = -\mathcal{Q}^R$. A similar global $O(2)^L$ symmetry applies to the left movers, making the total internal global symmetry $O(2)^L \times O(2)^R$.

The (quantized) vector and axial charges are defined as $\mathcal{Q}^V = \mathcal{Q}^L + \mathcal{Q}^R$ and $\mathcal{Q}^A = \mathcal{Q}^L - \mathcal{Q}^R$ and act on the fermions as

$$\begin{aligned} [\mathcal{Q}^V, \Psi_L^\dagger] &= \Psi_L^\dagger, & [\mathcal{Q}^V, \Psi_R^\dagger] &= \Psi_R^\dagger, \\ [\mathcal{Q}^A, \Psi_L^\dagger] &= \Psi_L^\dagger, & [\mathcal{Q}^A, \Psi_R^\dagger] &= -\Psi_R^\dagger. \end{aligned} \quad (2)$$

The \mathbb{Z}_2 subgroups of the vector and axial $U(1)$ symmetries act identically on the fermions as a fermion parity, so the global form of these symmetries is $[U(1)^V \times U(1)^A] / \mathbb{Z}_2$. The vector and axial charges are related by:

$$\mathcal{Q}^A = \mathcal{C}^R \mathcal{Q}^V (\mathcal{C}^R)^{-1}. \quad (3)$$

The vector and axial $U(1)$ symmetries are separately anomaly-free in the sense that there is no obstruction to gauging either one of the two. However, there is a mixed anomaly between them, which implies that when the vector symmetry is gauged (*i.e.*, in QED), the axial symmetry is broken, and vice versa [2, 19]. This can be seen from the anomalous conservation equation of the axial current $j_\mu^A = \bar{\Psi} \Gamma_\mu \Gamma^5 \Psi$ (with $\Gamma^5 = \Gamma^0 \Gamma^1$):

$$\partial^\mu j_\mu^A = \frac{1}{\pi} E, \quad (4)$$

where E is the electric field.

² We thank Tom Banks and Nathan Seiberg for this discussion.

³ This statement is only meaningful once a specific charge sector of the global symmetries is fixed. Otherwise, for instance, complex conjugation changes the chirality and global symmetry charge of a Weyl fermion field in 3+1D.

III. AXIAL CHARGES OF STAGGERED FERMIONS

Consider the 1+1D staggered fermion Hamiltonian [20–22]

$$H = -i \sum_{j=1}^L \left(c_j^\dagger c_{j+1} + c_j c_{j+1}^\dagger \right), \quad (5)$$

where there is a single-component complex fermion c_j at every site j , satisfying $\{c_j, c_{j'}\} = \{c_j^\dagger, c_{j'}^\dagger\} = 0$ and $\{c_j, c_{j'}^\dagger\} = \delta_{j,j'}$. This Hamiltonian is ultra-local, with only nearest-neighbor couplings.⁴ The continuum limit is a single, free, massless Dirac fermion (1). We consider both periodic and anti-periodic boundary conditions on a closed chain with L sites, *i.e.*, $c_{j+L} = (-1)^\nu c_j$ with $\nu = 0, 1$. We assume L to be even for simplicity. The Hilbert space is 2^L dimensional.

It will be important for the following discussion to decompose the complex fermion into two real fermions as $c_j = \frac{1}{2}(a_j + ib_j)$, where $a_j = a_j^\dagger$ and $b_j = b_j^\dagger$ are decoupled Majorana fermions satisfying $\{a_j, a_{j'}\} = \{b_j, b_{j'}\} = 2\delta_{j,j'}$. In terms of these Majorana fermions, the Hamiltonian (5) becomes

$$H = -\frac{i}{2} \sum_{j=1}^L (a_j a_{j+1} + b_j b_{j+1}). \quad (6)$$

The Hamiltonian (5) has a manifest $U(1)^V$ fermion-number symmetry whose quantized charge is⁵

$$Q^V = \sum_{j=1}^L \left(c_j^\dagger c_j - \frac{1}{2} \right) = \frac{i}{2} \sum_{j=1}^L a_j b_j \equiv \sum_{j=1}^L q_j^V. \quad (7)$$

It flows to the vector charge Q^V of the Dirac fermion field theory in the continuum limit.⁶ We will hence refer to it as the vector charge or the fermion number. The $U(1)^V$ symmetry acts on the lattice fermions as

$e^{i\varphi Q^V} c_j e^{-i\varphi Q^V} = e^{-i\varphi} c_j$, or equivalently, as

$$\begin{aligned} e^{i\varphi Q^V} a_j e^{-i\varphi Q^V} &= \cos \varphi a_j + \sin \varphi b_j, \\ e^{i\varphi Q^V} b_j e^{-i\varphi Q^V} &= \cos \varphi b_j - \sin \varphi a_j. \end{aligned} \quad (8)$$

A. Quantized axial charge

Is the axial symmetry exact on the lattice? We claim that the lattice operator

$$\begin{aligned} Q^A &= \frac{1}{2} \sum_{j=1}^L (c_j + c_j^\dagger) (c_{j+1} - c_{j+1}^\dagger) \\ &= \frac{i}{2} \sum_{j=1}^L a_j b_{j+1} \equiv \sum_{j=1}^L q_{j+\frac{1}{2}}^A, \end{aligned} \quad (9)$$

obeys the following properties: (a) It is quantized with integer eigenvalues. (b) It commutes with the ultra-local Hamiltonian (5). (c) It is a sum of local charge density operators $q_{j+\frac{1}{2}}^A$. (d) It is bilinear in the fermions. (e) It becomes the continuum axial charge Q^A in the continuum limit. Property (a) follows from the fact that the local factors $q_{j+\frac{1}{2}}^A$, which is the lattice version of the (time component of the) Noether current, commute with each other and can be simultaneously diagonalized. Properties (b), (c), and (d) are straightforward to verify. Below, we will argue for the last property.

To find a lattice axial charge, we note that since the vector charge is manifest on the lattice, using (3), it suffices to find a lattice realization of the right-moving charge conjugation C^R . The latter is a chiral fermion parity that flips the sign of a single Majorana-Weyl fermion. Specifically, we decompose the Weyl fermion into two Majorana-Weyl fermions as $\Psi_R = \lambda_R + i\chi_R$, then C^R only flips the sign of χ_R . It is known that such a chiral fermion parity is realized as a lattice translation on the lattice, which we review below.

We focus on the Majorana fermion b_j in the lattice Hamiltonian (6), which flows to the left and right Majorana-Weyl fermions χ_L and χ_R in the continuum. Up to a constant, the Hamiltonian for b_j in momentum space is $\sum_{0 < k < L/2} \sin \frac{2\pi k}{L} \beta_{-k} \beta_k$, where $\beta_k = \frac{1}{\sqrt{L}} \sum_{j=1}^L e^{-\frac{2\pi i k j}{L}} b_j$ with $k \in \mathbb{Z} + \nu/2$ and $k \sim k + L$. The ground state(s) $|\Omega\rangle$ obeys $\beta_k |\Omega\rangle = 0$ for all $0 < k < \frac{L}{2}$. The left- and right-moving modes in the continuum field theory created by χ_L and χ_R arise from the lattice modes created by β_{-k} with k close to $k = 0$ and $k = \frac{L}{2}$, respectively.

The key symmetry in the lattice model (6) is the translation operator T_b that shifts the b_j fermion by one site but leaves a_j invariant:

$$T_b a_j T_b^{-1} = a_j, \quad T_b b_j T_b^{-1} = b_{j+1}. \quad (10)$$

⁴ A Hamiltonian is called “ultra-local” if every term only involves operators in a finite neighborhood whose size does not scale with the overall system size. In some literature, this property is simply referred to as “local”, while in others a “local” Hamiltonian can have long-range interaction which decays exponentially in the distance.

⁵ We choose the constant in Q^V so that $C Q^V C^{-1} = -Q^V$. Here, C is charge conjugation, which acts on the fermion as $C c_j C^{-1} = c_j^\dagger$. It becomes the non-chiral charge conjugation $C^L C^R$ in the continuum limit. The axial charge Q^A defined later in (9) also obeys $C Q^A C^{-1} = -Q^A$.

⁶ By continuum limit, we mean we focus on finitely many excited states of the lattice model and then send $L \rightarrow \infty$.

In momentum space, $T_b \beta_k T_b^{-1} = e^{\frac{2\pi i k}{L}} \beta_k$, hence T_b acts with a relative minus sign between the modes around $k=0$ and $k=\frac{L}{2}$ in the limit $L \rightarrow \infty$. We conclude that on the low-lying states, T_b acts as the right-moving charge conjugation times the continuum translation operator [23]:

$$T_b = \mathcal{C}^R e^{\frac{2\pi i \mathcal{P}}{L}} \quad (11)$$

where \mathcal{P} is the continuum momentum operator acting only on the Majorana fermion χ (but not λ).⁷

Having identified the lattice origin of the right-moving charge conjugation \mathcal{C}^R , we follow (3) to define the lattice axial charge:

$$Q^A = T_b Q^V T_b^{-1}, \quad (12)$$

which gives (9). On the low-lying states in the $L \rightarrow \infty$ limit, $e^{2\pi i \mathcal{P}/L}$ becomes 1 and $T_b \sim \mathcal{C}^R$, so from (3) we see that Q^A becomes the continuum axial charge \mathcal{Q}^A . From this expression, it is clear that Q^A has integer eigenvalues since it is unitarily equivalent to Q^V . It is also clear that it commutes with the Hamiltonian since Q^V and T_b do.

Since Q^A is quantized, we can exponentiate it to find an exact $U(1)^A$ axial symmetry on the lattice, which acts locally on the fermions as:

$$\begin{aligned} e^{i\varphi Q^A} a_j e^{-i\varphi Q^A} &= \cos \varphi a_j + \sin \varphi b_{j+1}, \\ e^{i\varphi Q^A} b_j e^{-i\varphi Q^A} &= \cos \varphi b_j - \sin \varphi a_{j-1}, \end{aligned} \quad (13)$$

or equivalently, as

$$\begin{aligned} e^{i\varphi Q^A} c_j e^{-i\varphi Q^A} &= \cos \varphi c_j \\ &- i \sin \varphi \frac{1}{2} \left(c_{j-1}^\dagger + c_{j-1} - c_{j+1}^\dagger + c_{j+1} \right). \end{aligned} \quad (14)$$

This quantized axial charge was first identified in [25, 26] from the connection to integrability. Here, we provide an alternative derivation using lattice translation.

While each of Q^V and Q^A generates an ordinary $U(1)$ global symmetry, interestingly, these two lattice charges do *not* commute:

$$[Q^V, Q^A] = - \sum_{j=1}^L \left(c_j c_{j+1} + c_j^\dagger c_{j+1}^\dagger \right). \quad (15)$$

This is to be contrasted with the continuum where $[Q^V, \mathcal{Q}^A] = 0$. In Appendix A, we show that the non-vanishing lattice commutator goes to zero on the low-lying states in the $L \rightarrow \infty$ limit. Note that

$(-1)^F \equiv e^{i\pi Q^V} = e^{i\pi Q^A}$ is the (non-anomalous) fermion parity that flips the sign of all the fermions. It commutes with both $U(1)$'s, *i.e.*, $[(-1)^F, Q^A] = [(-1)^F, Q^V] = 0$.

B. Unquantized axial charge

Although $[Q^A, Q^V] \neq 0$, there is another conserved operator $\tilde{Q}^A = \frac{1}{2}(Q^A + e^{\frac{i\pi}{2} Q^V} Q^A e^{-\frac{i\pi}{2} Q^V})$ that obeys $[\tilde{Q}^A, Q^V] = 0$. This operator, which is the most straightforward lattice regularization of the axial charge [21, 22], can also be written as a sum of local terms:

$$\begin{aligned} \tilde{Q}^A &= \frac{1}{2} \sum_{j=1}^L \left(c_j^\dagger c_{j+1} - c_j c_{j+1}^\dagger \right) \\ &= \frac{i}{4} \sum_{j=1}^L (a_j b_{j+1} - b_j a_{j+1}) \equiv \sum_{j=1}^L \tilde{q}_{j+\frac{1}{2}}^A. \end{aligned} \quad (16)$$

In the continuum limit, Q^V and Q^A commute, and \tilde{Q}^A flows to the same continuum axial charge \mathcal{Q}^A . However, the price we pay is that the eigenvalues of \tilde{Q}^A are generally irrational and are not quantized.⁸

The commutator between the local charge densities q_j^V and $\tilde{q}_{j+\frac{1}{2}}^A$ is [27]⁹

$$[q_j^V, \tilde{q}_{j+\frac{1}{2}}^A] = \frac{i}{2} (\delta_{j,j'} - \delta_{j-1,j'}) h_{j+\frac{1}{2}}, \quad (17)$$

where $h_{j+\frac{1}{2}} = -\frac{i}{2}(a_j a_{j+1} + b_j b_{j+1})$ is the Hamiltonian density. This is the lattice avatar of the Schwinger term, which encodes the mixed anomaly of $U(1)^V$ and $U(1)^A$ in the continuum.

The simultaneous realization of both vector and axial $U(1)$ symmetries seems to conflict with the well-known Nielsen-Ninomiya theorem. However, neither of the conserved lattice operators Q^A and \tilde{Q}^A satisfies both conditions (i) and (ii) mentioned in the Introduction: Q^A is quantized but does not commute with Q^V , while \tilde{Q}^A commutes with Q^V but is not quantized. (See Table I.) We, therefore, cannot define a notion of lattice chirality using them. The fact that these two conditions cannot be met simultaneously is in harmony with the Nielsen-Ninomiya theorem. Relatedly, we do not have $U(1)^L$ or $U(1)^R$ symmetries on the lattice since $\frac{1}{2}(Q^V \pm Q^A)$ do not have quantized eigenvalues.

⁷ This is to be contrasted with the lattice translation $T = T_a T_b$ that shifts both a_j and b_j by one site. In the continuum limit, T flips the signs of both λ_R, χ_R [21, 22], while T_b only flips the sign of χ_R . See Refs. [23, 24] for recent discussions of chiral fermion parities from the lattice.

⁸ Up to an overall constant, \tilde{Q}^A is equivalent to the Hamiltonian (5) for $L=0 \pmod{4}$, under the unitary transformation $c_j \mapsto i^j c_j$.

⁹ We thank Tom Banks and Nathan Seiberg for discussions on the Schwinger term and collaboration on another project.

lattice operators	Q^A	\tilde{Q}^A
quantized eigenvalues?	✓	
$[\bullet, Q^V] = 0?$		✓
$[\bullet, H] = 0?$	✓	✓
sum of local terms?	✓	✓

Table I. Properties of the two lattice operators, both flowing to the same axial charge of a free Dirac fermion in the continuum limit.

C. From the lattice to the continuum

We now present a more detailed analysis in momentum space to relate the lattice and continuum symmetries. Let us define $\gamma_k = \frac{1}{\sqrt{L}} \sum_{j=1}^L e^{-\frac{2\pi i k j}{L}} c_j$, which satisfies $\{\gamma_k, \gamma_{k'}\} = \{\gamma_k^\dagger, \gamma_{k'}^\dagger\} = 0$ and $\{\gamma_k, \gamma_{k'}^\dagger\} = \delta_{k, k'}$.¹⁰ The Hamiltonian (5) in momentum space is $H = \sum_{-\frac{L}{2} < k \leq \frac{L}{2}} 2 \sin \frac{2\pi k}{L} \gamma_k^\dagger \gamma_k$. The ground state(s) is(are) annihilated by γ_k with $0 < k < \frac{L}{2}$, and by γ_k^\dagger with $-\frac{L}{2} < k < 0$. For periodic boundary condition $\nu = 0$, there are four degenerate ground states due to the zero modes at $k = 0$ and $\frac{L}{2}$. For anti-periodic boundary condition $\nu = 1$, there is a unique ground state.

The conserved operators in momentum space are

$$\begin{aligned}
Q^V &= \sum_k \left(\gamma_k^\dagger \gamma_k - \frac{1}{2} \right), \\
\tilde{Q}^A &= \sum_k \cos \frac{2\pi k}{L} \gamma_k^\dagger \gamma_k \\
Q^A &= \tilde{Q}^A + \frac{1}{2} \sum_k \left(e^{-\frac{2\pi i}{L} k} \gamma_k \gamma_{-k} + \text{h.c.} \right)
\end{aligned} \tag{18}$$

where all the sums are over $-\frac{L}{2} < k \leq \frac{L}{2}$. The middle equation shows that the eigenvalues of \tilde{Q}^A are not quantized, as mentioned earlier.

In the continuum limit, we only retain the Fourier modes γ_k^\dagger , γ_{-k} , $\gamma_{\frac{L}{2}-k}^\dagger$, and $\gamma_{-\frac{L}{2}+k}$ with $|k| \ll L$, which become the creation operators for Ψ_L^\dagger , Ψ_L , Ψ_R^\dagger , and Ψ_R , respectively. We now compute the commutators of the unquantized \tilde{Q}^A and quantized Q^A axial charges with the creation operators:

$$\begin{aligned}
[\tilde{Q}^A, \gamma_k^\dagger] &= \cos \frac{2\pi k}{L} \gamma_k^\dagger, \\
[Q^A, \gamma_k^\dagger] &= \cos \frac{2\pi k}{L} \gamma_k^\dagger + i \sin \frac{2\pi k}{L} \gamma_{-k},
\end{aligned} \tag{19}$$

¹⁰ They are related to the Fourier modes α_k, β_k of a_j, b_j by $\gamma_k = \frac{1}{2}(\alpha_k + i\beta_k)$, $\gamma_k^\dagger = \frac{1}{2}(\alpha_{-k} - i\beta_{-k})$.

lattice states	Q^V	\tilde{Q}^A	$\langle Q^A \rangle$	cont. op.
$ \Omega\rangle$	0	0	0	1
$\gamma_{\frac{L}{2}}^\dagger \Omega\rangle$	+1	$\cos \frac{\pi}{L}$	$\cos \frac{\pi}{L}$	Ψ_L^\dagger
$\gamma_{-\frac{L}{2}} \Omega\rangle$	-1	$-\cos \frac{\pi}{L}$	$-\cos \frac{\pi}{L}$	Ψ_L
$\gamma_{\frac{L}{2}-\frac{1}{2}}^\dagger \Omega\rangle$	+1	$-\cos \frac{\pi}{L}$	$-\cos \frac{\pi}{L}$	Ψ_R^\dagger
$\gamma_{-\frac{L}{2}+\frac{1}{2}} \Omega\rangle$	-1	$\cos \frac{\pi}{L}$	$\cos \frac{\pi}{L}$	Ψ_R

Table II. The eigenvalues of the vector charge Q^V and the unquantized axial charge \tilde{Q}^A for the low-lying energy eigenstates of the Hamiltonian (5) with anti-periodic boundary condition. Since the quantized axial charge Q^A does not commute with Q^V or \tilde{Q}^A , it cannot be simultaneously diagonalized; instead, we record its expectation values. The rightmost column shows the local operators that create these states in the continuum limit.

while those with γ_k can be obtained similarly. Hence, for finite k and $L \rightarrow \infty$,

$$\begin{aligned}
\lim_{L \rightarrow \infty} [Q^A, \gamma_k^\dagger] &= \lim_{L \rightarrow \infty} [\tilde{Q}^A, \gamma_k^\dagger] = \gamma_k^\dagger, \\
\lim_{L \rightarrow \infty} [Q^A, \gamma_{\frac{L}{2}-k}^\dagger] &= \lim_{L \rightarrow \infty} [\tilde{Q}^A, \gamma_{\frac{L}{2}-k}^\dagger] = -\gamma_{\frac{L}{2}-k}^\dagger.
\end{aligned} \tag{20}$$

Comparing with (2), this proves that both Q^A and \tilde{Q}^A flow to the continuum axial charge Q^A in the limit. As a demonstration, we compute the eigenvalues and expectation values of these conserved charges on the low-lying lattice states in Table II.

D. The non-abelian algebra

The charges Q^V and Q^A (as well as \tilde{Q}^A) are part of a larger algebra of conserved operators. Let us define $A_n = \frac{i}{2} \sum_j a_j a_{j+n}$, $B_n = -\frac{i}{2} \sum_j b_j b_{j+n}$ and $Q_n = \frac{i}{2} \sum_j a_j b_{j+n}$ with $n \in \mathbb{Z}$. We have $A_{n+L} = (-1)^\nu A_n$, and similarly for B_n and Q_n . These operators all commute with the Hamiltonian and obey the following closed algebra:

$$\begin{aligned}
[Q_n, Q_m] &= i(B_{m-n} + A_{m-n}), \\
[Q_n, A_m] &= [Q_n, B_m] = i(Q_{n-m} - Q_{n+m}),
\end{aligned} \tag{21}$$

while all the other commutators vanish. In particular, $Q^V = Q_0$, $Q^A = Q_1$, $\tilde{Q}^A = \frac{1}{2}(Q_1 + Q_{-1})$ and $H = A_{-1} + B_1$. While $Q_n = T_b^n Q_0 T_b^{-n}$ have integer eigenvalues, A_n and B_n do not.

For a fixed finite n , these operators are sum of local operators, and using (11), we find

$$\lim_{L \rightarrow \infty} Q_n = \begin{cases} Q^V & \text{for } n \text{ even,} \\ Q^A & \text{for } n \text{ odd.} \end{cases} \tag{22}$$

Therefore, we find an infinite tower of conserved lattice operators, with Q^A and \tilde{Q}^A special cases of them, that flow to the same axial charge in the continuum limit.

Since the model (5) is integrable, it is expected to have many conserved quantities (in particular, our algebra contains the $sl(2)$ loop algebra of Ref. [28] as a subalgebra). However, the majority of these conserved quantities are non-local, meaning they cannot be expressed as a sum of operators supported in a finite region, nor do they map local operators to other local operators. The interesting point, however, is that among these conserved quantities, Q^A (along with Q_n for small, odd n) is local and flows to the axial charge in the continuum, which has an anomaly with the vector charge.

E. Comparison with other constructions

Many of the discussions of lattice chiral symmetries are in the context of Euclidean lattice models, while ours concerns the Hamiltonian lattice models. We will not attempt to give a comprehensive comparison with all the existing constructions, but refer the readers to Ref. [1] for a review.

The Euclidean version of the staggered fermion, which is reviewed in Ref. [29], has two $U(1)$ symmetries. The first is the manifest vector, or fermion number, symmetry. The second is sometimes referred to as the $U(1)_\epsilon$ symmetry [30, 31]. In contrast to the symmetries in the Hamiltonian model, these two $U(1)$ symmetries commute in the Euclidean model. However, $U(1)_\epsilon$ acts with an alternating sign in the Euclidean time direction, and therefore does not lead to an ordinary global symmetry in the Hamiltonian formalism. It would be interesting to further explore the connection between $U(1)_\epsilon$ and our non-abelian algebra, especially along the lines of Ref. [32].

Next, we compare this discussion with other Hamiltonian lattice models in the literature. Refs. [33–35] discuss an axial charge that is both quantized and commutes with the vector charge, but their Hamiltonian is not local, and the Nielsen-Ninomiya theorem does not apply. The quantized axial charge was pointed out in Refs. [25, 26], and was referred to as the “arrow charge” in the latter reference in connection with the integrability literature. In this letter, we provided an alternative derivation of this quantized axial charge, from which we found a family of other lattice realizations and identified the non-abelian algebra they generate. The unquantized axial charge was first discussed in Refs. [21, 22], and was later elaborated in Ref. [26, 36] in the context of the Hamiltonian version of the overlap formulation [37–39] and the Ginsparg-Wilson relation [40]. Finally, the Hamiltonian in (5) is in 1+1D with nearest-neighbor couplings, contrasting with the domain wall fermion approach where the model is effectively 2+1D [41]. (See also Refs. [42, 43] for recent progress.)

IV. COUPLING TO GAUGE FIELDS

A. Gauging the vector symmetry

Let us now discuss the fate of the two axial charges after we gauge $U(1)^V$. The gauged Hamiltonian is a lattice regularization of 1+1D QED, *i.e.*, the Schwinger model [2]. We introduce a $U(1)$ -valued gauge field $U_{j+\frac{1}{2}}$ and an integer-valued electric field operator $L_{j+\frac{1}{2}}$ on each link. They satisfy $[L_{j+\frac{1}{2}}, U_{j'+\frac{1}{2}}] = \delta_{j,j'} U_{j'+\frac{1}{2}}$. The gauged Hamiltonian is [21, 44]

$$H_V = -i \sum_j \left(c_j^\dagger U_{j+\frac{1}{2}} c_{j+1} + c_j U_{j+\frac{1}{2}}^\dagger c_{j+1}^\dagger \right) + K \sum_j L_{j+\frac{1}{2}}^2. \quad (23)$$

Furthermore, we impose the Gauss constraint, $L_{j+\frac{1}{2}} - L_{j-\frac{1}{2}} = q_{j+\frac{1}{2}}^V + \frac{(-1)^j}{2}$ [45, 46]. The Gauss law restricts the Hilbert space to a subsector of fixed Q^V charge. Since the quantized axial charge Q^A does not commute with Q^V , it does not act within the gauge-invariant subspace. Q^A is therefore no longer a symmetry when we couple to the dynamical gauge field for $U(1)^V$. This is analogous to gauging a $U(1)$ subgroup of $SU(2)$: if the S^z symmetry is gauged, the S^x and S^y charges cannot be made gauge-invariant and are, therefore, explicitly broken.

On the other hand, the unquantized axial charge \tilde{Q}^A can be made gauge-invariant as $\tilde{Q}^A(U) = \frac{1}{2} \sum_j (c_j^\dagger U_{j+\frac{1}{2}} c_{j+1} - c_j U_{j+\frac{1}{2}}^\dagger c_{j+1}^\dagger)$.¹¹ Even though it is a gauge-invariant operator acting in the gauged theory, it fails to commute with the gauged Hamiltonian,

$$-i [\tilde{Q}^A(U), H_V] = -\frac{K}{2} \sum_j \{L_{j+\frac{1}{2}}, h_{j+\frac{1}{2}}\}, \quad (24)$$

where $h_{j+\frac{1}{2}}(U) = -i (c_j^\dagger U_{j+\frac{1}{2}} c_{j+1} + c_j U_{j+\frac{1}{2}}^\dagger c_{j+1}^\dagger)$ is the Hamiltonian density for the fermions. This is the lattice avatar of the Schwinger anomaly of the continuum axial charge $\frac{d}{dt} Q^A = -i [Q^A, H] = \frac{1}{\pi} \int dx E$ [48], which is the integrated form of (4). We conclude that both the quantized Q^A and the unquantized \tilde{Q}^A axial symmetries are broken as we gauge the vector $U(1)^V$ symmetry.

B. Gauging the axial symmetry

The main advantage of the quantized axial charge Q^A , compared to the unquantized one \tilde{Q}^A , is that it can be

¹¹ The continuum counterpart is the following: in the point-splitting regularization of the axial current j^A , an infinitesimal Wilson line must be introduced to ensure gauge invariance [47].

coupled to dynamical (compact) $U(1)$ gauge fields. The gauged Hamiltonian is obtained by conjugating H_V with T_b :

$$H_A = \sum_j \left[-\frac{i}{4}(a_j - ib_{j+1})U_{j+\frac{1}{2}}(a_{j+1} + ib_{j+2}) + \text{h.c.} \right] + K \sum_j L_{j+\frac{1}{2}}^2. \quad (25)$$

The Gauss law is obtained similarly, and found to be $L_{j+\frac{1}{2}} - L_{j-\frac{1}{2}} = q_{j+\frac{1}{2}}^A + \frac{(-1)^j}{2}$. We conclude that Q^V and Q^A can individually be gauged on the lattice and are free of self-anomalies.

V. A LATTICE ANOMALY AS AN OBSTRUCTION TO GAPPED PHASES

Do the lattice axial and vector symmetries have an anomaly? Conventionally, the anomaly of a global symmetry is defined as the obstruction to gauging the symmetry. However, the lattice charges Q^A and Q^V generate the non-abelian algebra (21), which includes highly non-local charges, so it is not clear if there is a sensible prescription for gauging it. On the other hand, it has been advocated in Refs. [49–52] that a global symmetry should be called anomalous if there does not exist a trivially gapped phase (*i.e.*, a gapped phase with a non-degenerate ground state and no long-range entanglement) preserving the symmetry. This definition is inspired by the 't Hooft anomaly matching argument and avoids the need to discuss the gauging of said global symmetry. Below, we will show that the lattice axial and vector symmetries together are anomalous in this sense. See Refs. [53–55] for the relation between these two definitions of anomalies.

Concretely, we prove in Appendix B that local deformations of the Hamiltonian (5) preserving both Q^V and Q^A are necessarily quadratic in the fermions. In fact, the only Q^V and Q^A symmetric local deformations are of the form ($n \geq 2$)

$$\delta H_n = \sum_{j=1}^L -i(c_j^\dagger c_{j+n} + c_j c_{j+n}^\dagger), \quad (26)$$

which flow to irrelevant deformations of the Dirac CFT in the IR. For a small deformation strength, such terms only renormalize the velocities of the left- and right-moving fermions in the continuum limit.

In particular, the quartic Thirring coupling $c_j^\dagger c_j c_{j+1}^\dagger c_{j+1}$ preserves Q^V but breaks Q^A . This is to be contrasted with its continuum limit; the continuum Thirring coupling $\frac{1}{4}(\bar{\Psi}\Gamma^\mu\Psi)(\bar{\Psi}\Gamma_\mu\Psi) = \Psi_L^\dagger\Psi_L\Psi_R^\dagger\Psi_R$ preserves both the vector and axial symmetries Q^V, Q^A , and leads to an exactly marginal deformation of the Dirac fermion CFT.

One interesting corollary is that *any Hamiltonian that commutes with both Q^V and Q^A must be gapless*. This is reminiscent of the constraint from perturbative anomalies in continuum field theory. In continuum field theory, perturbative anomalies of continuous global symmetries are encoded in the local operator product expansion and, therefore, cannot be matched by a gapped phase. Even when the symmetry is spontaneously broken, there are gapless Goldstone boson modes. The symmetries Q^V and Q^A present an example of such a gapless constraint on a lattice with a finite-dimensional local Hilbert space.

This constraint is in fact much stronger than the typical discrete anomalies or the Lieb-Schultz-Mattis constraints [56–58], where the low-energy phase is constrained to be either gapless, *or* gapped with some non-trivial features, such as degenerate ground states and/or long-range entanglement described by a topological quantum field theory. It is also stronger than the “symmetry-enforced gaplessness” discussed in Refs. [59–65], where a gapped phase with spontaneous broken discrete symmetries (*e.g.*, time-reversal symmetry) remains a possibility. The charges Q^V and Q^A entirely exclude any gapped phases, leaving the gapless phase as the only possibility.

While the preservation of both Q^V and Q^A imposes a nontrivial gapless constraint, each of these two conserved charges can individually be coupled to respective gauge fields (as in (23) and (25)) and is free of anomalies. Moreover, it is possible to deform the gapless Hamiltonian (5) to a trivially gapped phase with a unique vacuum while preserving either Q^V or Q^A (but not both). If we choose to preserve Q^V , while violating Q^A , we can add the deformation $\delta H = \sum_j (-1)^j c_j^\dagger c_j$ to open an energy gap. In the continuum, this corresponds to deforming the Lagrangian density by the mass term $\Psi_R^\dagger\Psi_L + \Psi_L^\dagger\Psi_R$ [21, 22]. See Ref. [46] for the precise relation between this lattice deformation and the continuum mass term. On the other hand, the deformation $T_b \delta H T_b^{-1} = \sum_j (-1)^j q_{j+\frac{1}{2}}^A$ preserves Q^A but violates Q^V and drives the system to a trivially gapped phase. In the continuum, it corresponds to $\Psi_R\Psi_L + \Psi_L^\dagger\Psi_R^\dagger$.

Finally, we note that the charge Q^V is on-site in the sense that the local terms q_j^V involve fermions only on site j , and it maps a local operator at site j to another at the same location. On the other hand, Q^A is not on-site since it smears a local operator at site j to its nearest neighbors. The unitary transformation by T_b^{-1} renders Q^A on-site, but this comes at the cost of Q^V no longer being on-site. However, there does not seem to be a unitary transformation that makes both Q^V and Q^A on-site simultaneously, hinting at a mixed anomaly between them. See Refs. [66, 67] for related discussions.

VI. EXACT WINDING SYMMETRY IN THE XY MODEL

It is well-known that upon bosonization, the fermionic Hamiltonian (5) becomes the XY model,

$H = \sum_j X_j X_{j+1} + Y_j Y_{j+1}$ [56]. The vector charge Q^V becomes $\frac{1}{2} \sum_j Z_j$, the charge of the manifest $U(1)$ spin rotation symmetry of the XY Hamiltonian. In the continuum, this flows to the momentum $U(1)$ symmetry of the $c = 1$ compact boson CFT at radius $R = \sqrt{2}$.¹² On the other hand, the axial charge Q^A is mapped to a second $U(1)$ symmetry of the XY model, which flows to the winding $U(1)$ symmetry in the continuum; its explicit form on a chain with even L is¹³

$$\frac{1}{4} \sum_{j=1}^{L/2} (Y_{2j-1} X_{2j} - X_{2j} Y_{2j+1}). \quad (27)$$

The global aspects of bosonization, the exact winding symmetry, and the relation to a non-invertible symmetry of the XY model will be discussed in Ref. [73].

VII. CONCLUSION

While a lattice model with a finite-dimensional Hilbert space cannot host the exact chiral anomaly, the symmetry generators for the axial and vector symmetries – which in the continuum limit have a mixed anomaly – exist exactly in the staggered fermion lattice model. The quantized charges Q^V, Q^A [25, 26] resemble their continuum counterparts closely: each one of them generates a (compact) $U(1)$ global symmetry and can be gauged. However, they do not commute with each other on the lattice. By contrast, the unquantized axial charge \tilde{Q}^A [21, 22, 26, 36] generates an \mathbb{R} global symmetry, and it is not clear how to couple it to $U(1)$ gauge fields.

It would be interesting to investigate the fate of these charges for interacting fermions *e.g.*, in 1+1D QED with multiple flavors of fermions and in QCD, where gauge interactions make the models non-integrable. It would also be natural to analyze chiral global symmetries in other models, such as the 3450 model. Another generalization is to explore quantized axial charges in 3+1D stag-

gered fermions, which is more phenomenologically relevant. One qualitative difference compared to the 1+1D case is that the axial symmetry in 3+1D not only has a mixed anomaly with the vector symmetry but it is also anomalous by itself.

The phenomenon discussed here is qualitatively similar to the emanant symmetry of Ref. [9] and the non-invertible Kramers-Wannier symmetry in Refs. [23, 74–76]. In all these cases, we start with an internal global symmetry G_{IR} in a continuum field theory in the IR and seek a UV symmetry G_{UV} in a microscopic lattice realization. While the symmetry operators for G_{IR} exist exactly on the lattice (and therefore G_{IR} is not emergent), they obey a different algebra G_{UV} compared to the IR one. In Refs. [23, 74–77], the UV symmetries mix with lattice translations. In the current letter, the IR symmetry $G_{\text{IR}} = [U(1)^V \times U(1)^A]/\mathbb{Z}_2$, which has an anomaly, emanates [9] from the non-abelian UV symmetry G_{UV} (21).

ACKNOWLEDGMENTS

We would like to thank Ömer Aksoy, Tom Banks, Will Detmold, Igor Klebanov, Ho Tat Lam, Michael Levin, Laurens Lootens, Nathan Seiberg, Hersh Singh, Senthil Todadri, Frank Verstraete, and Yichen Xu for interesting discussions. SHS is especially grateful to Tom Banks and Nathan Seiberg for numerous enlightening discussions on lattice fermions. We would also like to express our special gratitude to Hersh Singh for his valuable insights and guidance toward relevant literature. We thank Simon Catterall, Aleksey Cherman, Theo Jacobson, Igor Klebanov, and Hersh Singh for helpful comments on a draft. AC is supported by NSF DMR-2022428 and by the Simons Collaboration on Ultra-Quantum Matter, which is a grant from the Simons Foundation (651446). SDP is supported by the National Science Foundation Graduate Research Fellowship under Grant No. 2141064. SHS is supported in part by NSF grant PHY-2210182.

¹² As standard in the string theory literature, we refer to the two $U(1)$ symmetries of the compact boson CFT as the “momentum” and “winding” symmetries. In some communities, they are respectively referred to as the “charge” and “vortex” symmetries. Here “momentum” refers to that in the target space. In the language of the Luttinger liquid, these two $U(1)$ symmetries respectively shift the two scalar fields. Our convention for the target space radius R is so that $R = 1$ is the self-dual point under T-duality, which enjoys an enhanced $su(2)$ chiral algebra.

¹³ In fact, when $L = 0 \pmod{4}$, the XY model is unitarily equivalent to the Levin-Gu spin chain [68, 69], and this lattice winding charge becomes the $U(1)$ charge $\frac{1}{4} \sum_j Z_j Z_{j+1}$ of Ref. [70]. On the other hand, the Levin-Gu spin chain is known to be equiva-

-
- [1] D. B. Kaplan, in *Les Houches Summer School: Session 93: Modern perspectives in lattice QCD: Quantum field theory and high performance computing* (2009) pp. 223–272, [arXiv:0912.2560 \[hep-lat\]](#).
 - [2] J. Schwinger, *Phys. Rev.* **128**, 2425 (1962).
 - [3] S. L. Adler, *Phys. Rev.* **177**, 2426 (1969).
 - [4] J. S. Bell and R. Jackiw, *Nuovo Cim. A* **60**, 47 (1969).
 - [5] A. Kapustin and R. Thorngren, *Phys. Rev. Lett.* **112**, 231602 (2014), [arXiv:1403.0617 \[hep-th\]](#).

lent to the XY model coupled to a \mathbb{Z}_2 gauge field. Therefore, the XY model is self-dual under gauging, leading to a non-invertible symmetry. The latter is the lattice realization of the symmetry discussed in Refs. [52, 71, 72].

- [6] S. Seifnashri, *SciPost Phys.* **16**, 098 (2024), arXiv:2308.05151 [cond-mat.str-el].
- [7] T. Sulejmanpasic and C. Gatteringer, *Nucl. Phys. B* **943**, 114616 (2019), arXiv:1901.02637 [hep-lat].
- [8] P. Gorantla, H. T. Lam, N. Seiberg, and S.-H. Shao, *J. Math. Phys.* **62**, 102301 (2021), arXiv:2103.01257 [cond-mat.str-el].
- [9] M. Cheng and N. Seiberg, *SciPost Phys.* **15**, 051 (2023), arXiv:2211.12543 [cond-mat.str-el].
- [10] L. Fazza and T. Sulejmanpasic, *JHEP* **05**, 017 (2023), arXiv:2211.13047 [hep-th].
- [11] E. Berkowitz, A. Cherman, and T. Jacobson, *Phys. Rev. D* **110**, 014510 (2024), arXiv:2310.17539 [hep-lat].
- [12] S. Catterall, J. Laiho, and J. Unmuth-Yockey, *JHEP* **10**, 013 (2018), arXiv:1806.07845 [hep-lat].
- [13] J. S. Schwinger, *Phys. Rev. Lett.* **3**, 296 (1959).
- [14] S. B. Treiman, E. Witten, R. Jackiw, and B. Zumino, *Current Algebra and Anomalies* (Princeton University Press, 2014).
- [15] H. B. Nielsen and M. Ninomiya, *Nucl. Phys. B* **185**, 20 (1981), [Erratum: *Nucl.Phys.B* 195, 541 (1982)].
- [16] H. B. Nielsen and M. Ninomiya, *Phys. Lett. B* **105**, 219 (1981).
- [17] H. B. Nielsen and M. Ninomiya, *Nucl. Phys. B* **193**, 173 (1981).
- [18] D. Friedan, *Commun. Math. Phys.* **85**, 481 (1982).
- [19] K. Johnson, *Physics Letters* **5**, 253 (1963).
- [20] J. B. Kogut and L. Susskind, *Phys. Rev. D* **11**, 395 (1975).
- [21] T. Banks, L. Susskind, and J. B. Kogut, *Phys. Rev. D* **13**, 1043 (1976).
- [22] L. Susskind, *Phys. Rev. D* **16**, 3031 (1977).
- [23] N. Seiberg and S.-H. Shao, *SciPost Phys.* **16**, 064 (2024), arXiv:2307.02534 [cond-mat.str-el].
- [24] M. Clancy, D. B. Kaplan, and H. Singh, *Phys. Rev. D* **109**, 014502 (2024), arXiv:2309.08542 [hep-lat].
- [25] H. B. Thacker, *Nucl. Phys. B Proc. Suppl.* **42**, 642 (1995), arXiv:hep-lat/9412019.
- [26] I. Horvath and H. B. Thacker, *Nucl. Phys. B Proc. Suppl.* **73**, 682 (1999), arXiv:hep-lat/9809108.
- [27] T. Banks, N. Seiberg, and S.-H. Shao, unpublished.
- [28] T. Deguchi, K. Fabricius, and B. M. McCoy, *J. Statist. Phys.* **102**, 701 (2001), arXiv:cond-mat/9912141.
- [29] M. Golterman, “Staggered fermions,” (2024), arXiv:2406.02906 [hep-lat].
- [30] L. H. Karsten and J. Smit, *Nucl. Phys. B* **183**, 103 (1981).
- [31] M. F. L. Golterman and J. Smit, *Nucl. Phys. B* **245**, 61 (1984).
- [32] S. Catterall, *Phys. Rev. D* **107**, 014501 (2023), arXiv:2209.03828 [hep-lat].
- [33] S. D. Drell, M. Weinstein, and S. Yankielowicz, *Phys. Rev. D* **14**, 1627 (1976).
- [34] L. H. Karsten and J. Smit, *Nucl. Phys. B* **144**, 536 (1978).
- [35] M. Weinstein, *Phys. Rev. D* **26**, 839 (1982).
- [36] M. Creutz, I. Horvath, and H. Neuberger, *Nucl. Phys. B Proc. Suppl.* **106**, 760 (2002), arXiv:hep-lat/0110009.
- [37] R. Narayanan and H. Neuberger, *Nucl. Phys. B* **443**, 305 (1995), arXiv:hep-th/9411108.
- [38] H. Neuberger, *Phys. Lett. B* **417**, 141 (1998), arXiv:hep-lat/9707022.
- [39] H. Neuberger, *Phys. Lett. B* **427**, 353 (1998), arXiv:hep-lat/9801031.
- [40] P. H. Ginsparg and K. G. Wilson, *Phys. Rev. D* **25**, 2649 (1982).
- [41] D. B. Kaplan, *Phys. Lett. B* **288**, 342 (1992), arXiv:hep-lat/9206013.
- [42] D. B. Kaplan and S. Sen, *Phys. Rev. Lett.* **132**, 141604 (2024), arXiv:2312.04012 [hep-lat].
- [43] D. B. Kaplan, *Phys. Rev. Lett.* **132**, 141603 (2024), arXiv:2312.01494 [hep-lat].
- [44] A. Carroll, J. B. Kogut, D. K. Sinclair, and L. Susskind, *Phys. Rev. D* **13**, 2270 (1976), [Erratum: *Phys.Rev.D* 14, 1729 (1976)].
- [45] C. J. Hamer, W.-h. Zheng, and J. Oitmaa, *Phys. Rev. D* **56**, 55 (1997), arXiv:hep-lat/9701015.
- [46] R. Dempsey, I. R. Klebanov, S. S. Pufu, and B. Zan, *Phys. Rev. Res.* **4**, 043133 (2022), arXiv:2206.05308 [hep-th].
- [47] M. Shifman, *Physics Reports* **209**, 341 (1991).
- [48] N. S. Manton, *Annals Phys.* **159**, 220 (1985).
- [49] C.-M. Chang, Y.-H. Lin, S.-H. Shao, Y. Wang, and X. Yin, *JHEP* **01**, 026 (2019), arXiv:1802.04445 [hep-th].
- [50] X.-G. Wen, *Phys. Rev. B* **99**, 205139 (2019), arXiv:1812.02517 [cond-mat.str-el].
- [51] R. Thorngren and Y. Wang, *JHEP* **04**, 132 (2024), arXiv:1912.02817 [hep-th].
- [52] Y. Choi, C. Cordova, P.-S. Hsin, H. T. Lam, and S.-H. Shao, *Phys. Rev. D* **105**, 125016 (2022), arXiv:2111.01139 [hep-th].
- [53] Z. Komargodski, K. Ohmori, K. Roumpedakis, and S. Seifnashri, *JHEP* **03**, 103 (2021), arXiv:2008.07567 [hep-th].
- [54] Y. Choi, B. C. Rayhaun, Y. Sanghavi, and S.-H. Shao, *Phys. Rev. D* **108**, 125005 (2023), arXiv:2305.09713 [hep-th].
- [55] Y. Choi, D.-C. Lu, and Z. Sun, *JHEP* **01**, 142 (2024), arXiv:2310.19867 [hep-th].
- [56] E. Lieb, T. Schultz, and D. Mattis, *Annals of Physics* **16**, 407 (1961).
- [57] M. Oshikawa, *Phys. Rev. Lett.* **84**, 3370 (2000), arXiv:cond-mat/0002392 [cond-mat.str-el].
- [58] M. B. Hastings, *Phys. Rev. B* **69**, 104431 (2004), arXiv:cond-mat/0305505.
- [59] C. Wang and T. Senthil, *Phys. Rev. B* **89**, 195124 (2014), arXiv:1401.1142 [cond-mat.str-el].
- [60] C. Wang and T. Senthil, *Phys. Rev. B* **94**, 245107 (2016), arXiv:1604.06807 [cond-mat.str-el].
- [61] I. Sodemann, I. Kimchi, C. Wang, and T. Senthil, *Phys. Rev. B* **95**, 085135 (2017), arXiv:1609.08616 [cond-mat.str-el].
- [62] C. Wang, A. Nahum, M. A. Metlitski, C. Xu, and T. Senthil, *Phys. Rev. X* **7**, 031051 (2017), arXiv:1703.02426 [cond-mat.str-el].
- [63] C. Córdova and K. Ohmori, “Anomaly Obstructions to Symmetry Preserving Gapped Phases,” (2019), arXiv:1910.04962 [hep-th].
- [64] C. Córdova and K. Ohmori, *Phys. Rev. D* **102**, 025011 (2020), arXiv:1912.13069 [hep-th].
- [65] A. Apte, C. Cordova, and H. T. Lam, *Phys. Rev. B* **108**, 045134 (2023), arXiv:2212.14605 [hep-th].
- [66] J. Wang and X.-G. Wen, *Phys. Rev. B* **107**, 014311 (2023), arXiv:1307.7480 [hep-lat].
- [67] J. Wang and X.-G. Wen, *Phys. Rev. D* **99**, 111501 (2018), arXiv:1807.05998 [hep-lat].
- [68] X. Chen, Y.-M. Lu, and A. Vishwanath, *Nat Commun* **5**, 4507 (2014), arXiv:1303.4301.
- [69] L. Li, M. Oshikawa, and Y. Zheng, *SciPost Phys.* **17**, 013 (2024), arXiv:2204.03131.

- [70] M. Levin and Z.-C. Gu, *Physical Review B* **86**, 115109 (2012), [arXiv:1202.3120](#).
- [71] W. Ji, S.-H. Shao, and X.-G. Wen, *Phys. Rev. Res.* **2**, 033317 (2020), [arXiv:1909.01425 \[cond-mat.str-el\]](#).
- [72] R. Thorngren and Y. Wang, *JHEP* **07**, 051 (2024), [arXiv:2106.12577 \[hep-th\]](#).
- [73] A. Chatterjee, S. D. Pace, and S.-H. Shao, in preparation.
- [74] N. Seiberg, S. Seifnashri, and S.-H. Shao, *SciPost Phys.* **16**, 154 (2024), [arXiv:2401.12281 \[cond-mat.str-el\]](#).
- [75] P. Gorantla, S.-H. Shao, and N. Tantivasadakarn, “Tensor networks for non-invertible symmetries in 3+1d and beyond,” (2024), [arXiv:2406.12978 \[quant-ph\]](#).
- [76] S. D. Pace, G. Delfino, H. T. Lam, and Ö. M. Aksoy, “Gauging modulated symmetries: Kramers-Wannier dualities and non-invertible reflections,” (2024), [arXiv:2406.12962 \[cond-mat.str-el\]](#).
- [77] M. A. Metlitski and R. Thorngren, *Phys. Rev. B* **98**, 085140 (2018), [arXiv:1707.07686 \[cond-mat.str-el\]](#).

Appendix A: Continuum limit of $[Q^V, Q^A]$

In the main text, we claim that the quantized lattice vector and axial charges Q^V and Q^A , given by Eqs. (7) and (9) respectively, commute in the continuum limit of the staggered fermion Hamiltonian. In this appendix, we prove this statement. More precisely, we will show that the matrix elements of $[Q^V, Q^A]$ between any two low-energy states vanish in the limit thermodynamic limit $L \rightarrow \infty$.

Low-energy states of the staggered fermion Hamiltonian model are most naturally understood in momentum space, where they reside near momenta $k = 0$ and $k = L/2$. Recall that in terms of the momentum space operators $\gamma_k = \frac{1}{\sqrt{L}} \sum_{j=1}^L e^{-\frac{2\pi i k j}{L}} c_j$, the Hamiltonian is re-written as

$$H = \sum_{-\frac{L}{2} < k \leq \frac{L}{2}} 2 \sin \frac{2\pi k}{L} \gamma_k^\dagger \gamma_k. \quad (\text{A.1})$$

When the fermions obey periodic boundary conditions, there are zero modes at $k = 0$ and $k = L/2$, which cause the ground states $|\Omega; J\rangle$ to be 4-fold degenerate ($J = 1, 2, 3, 4$). For anti-periodic boundary conditions, $k = 0$ and $k = L/2$ are not allowed values of momenta, and there is a single ground state $|\Omega; J\rangle$ with $J = 1$. For both periodic and anti-periodic boundary conditions, the ground state(s) satisfy

$$\gamma_k^\dagger \gamma_k |\Omega; J\rangle = \begin{cases} |\Omega; J\rangle & -\frac{L}{2} < k < 0, \\ 0 & 0 < k < \frac{L}{2}, \end{cases} \quad (\text{A.2})$$

and the low-energy states are created by acting γ_k and γ_k^\dagger on $|\Omega; J\rangle$ with k near 0 or $L/2$. Therefore, it is convenient to rewrite the commutator (15),

$$Q_{\text{com}} \equiv [Q^V, Q^A] = - \sum_{j=1}^L \left(c_j c_{j+1} + c_j^\dagger c_{j+1}^\dagger \right) \quad (\text{A.3})$$

in momentum space as

$$Q_{\text{com}} = \sum_k \left(e^{\frac{2\pi i}{L} k} \gamma_{-k}^\dagger \gamma_k^\dagger - e^{-\frac{2\pi i}{L} k} \gamma_k \gamma_{-k} \right). \quad (\text{A.4})$$

Consider two arbitrary low-energy states $|\phi_1\rangle = \gamma_{k_1}^\dagger \dots \gamma_{k_n}^\dagger \gamma_{q_1} \dots \gamma_{q_m} |\Omega, J\rangle$ and $|\phi_2\rangle = \gamma_{k'_1}^\dagger \dots \gamma_{k'_n}^\dagger \gamma_{q'_1} \dots \gamma_{q'_m} |\Omega, J'\rangle$ and the related matrix element of the commutator

$$\langle \phi_2 | Q_{\text{com}} | \phi_1 \rangle = \langle \Omega, J' | \gamma_{q'_m}^\dagger \dots \gamma_{q'_1}^\dagger \gamma_{k'_n} \dots \gamma_{k'_1} Q_{\text{com}} \gamma_{k_1}^\dagger \dots \gamma_{k_n}^\dagger \gamma_{q_1} \dots \gamma_{q_m} |\Omega, J\rangle. \quad (\text{A.5})$$

In the continuum limit, this matrix element simplifies. Indeed, using (A.4), it follows that

$$[Q_{\text{com}}, \gamma_k^\dagger] = -2i \sin \frac{2\pi k}{L} \gamma_{-k}, \quad [Q_{\text{com}}, \gamma_k] = 2i \sin \frac{2\pi k}{L} \gamma_{-k}^\dagger. \quad (\text{A.6})$$

For k close to 0 or $L/2$, these commutators vanish in the $L \rightarrow \infty$ limit. Therefore, the commutator Q_{com} commutes with all low-energy operators in the continuum limit. Using this, we can rewrite the matrix elements (A.5) in the

continuum limit as

$$\lim_{L \rightarrow \infty} \langle \phi_2 | Q_{\text{com}} | \phi_1 \rangle = \langle \Omega, J' | \gamma_{q'_m}^\dagger \cdots \gamma_{q'_1}^\dagger \gamma_{k'_n} \cdots \gamma_{k'_1} \gamma_{k_1}^\dagger \cdots \gamma_{k_n}^\dagger \gamma_{q_1} \cdots \gamma_{q_m} | \Omega, J \rangle. \quad (\text{A.7})$$

However, using Eqs. (A.2) and (A.4), the commutator Q_{com} satisfies $Q_{\text{com}} |\Omega, J\rangle = 0$. This is because for any $k \neq 0, L/2$, either γ_k or γ_{-k} annihilates $|\Omega, J\rangle$, and so $\gamma_k \gamma_{-k}$ necessarily annihilates $|\Omega, J\rangle$. Furthermore, this is true for both periodic and anti-periodic boundary conditions. Indeed, for the zero modes $k = 0, L/2$ occurring from periodic boundary conditions, we have $\gamma_k \gamma_{-k} = \gamma_k \gamma_k = 0$ and so these values of k do not contribute to Q_{com} . Following this argument, the matrix element (A.7) simplifies to

$$\lim_{L \rightarrow \infty} \langle \phi_2 | Q_{\text{com}} | \phi_1 \rangle = 0. \quad (\text{A.8})$$

Since $|\phi_1\rangle$ and $|\phi_2\rangle$ are arbitrary low-energy states, we conclude that all matrix elements of $Q_{\text{com}} \equiv [Q^V, Q^A]$ involving low-energy states vanish in the continuum limit, and Q^V and Q^A commute in the continuum limit.

Appendix B: Deformations preserving both Q^V and Q^A

In the main text, we discussed how the lattice vector and axial symmetries are anomalous in the sense that there does not exist a non-degenerated gapped Hamiltonian with $U(1)^V$ and $U(1)^A$ symmetries. In fact, we claimed that all local Hamiltonians that are $U(1)^V$ and $U(1)^A$ symmetric take the form

$$H = \sum_{n=1}^N \left(-i g_n \sum_{j=1}^L (c_j^\dagger c_{j+n} + c_j c_{j+n}^\dagger) \right), \quad (\text{B.1})$$

with $\lim_{L \rightarrow \infty} N/L = 0$. This is a gapless Hamiltonian for all couplings g_n with dispersion $\omega_k = 2 \sum_{n=1}^N g_n \sin(2\pi n k/L)$. In this appendix, we prove this statement. In particular, we will now prove that all local deformations of the staggered fermion Hamiltonian that commute with the $U(1)^V$ symmetry operator $e^{i\varphi Q^V}$ and the $\mathbb{Z}_4^A \subset U(1)^A$ symmetry operator $e^{-i\frac{\pi}{2} Q^A}$ are quadratic and of the type (B.1), which has the full $U(1)^A$ symmetry.

We start by considering the product of symmetry operators $e^{-i\frac{\pi}{2} Q^A} e^{i\frac{\pi}{2} Q^V}$, which satisfies $e^{-i\frac{\pi}{2} Q^A} e^{i\frac{\pi}{2} Q^V} = T_b T_a^{-1}$. Indeed, this identity can be seen by noting that $e^{-i\frac{\pi}{2} Q^A} e^{i\frac{\pi}{2} Q^V}$ act on a_j and b_j as

$$\begin{aligned} e^{-i\frac{\pi}{2} Q^A} e^{i\frac{\pi}{2} Q^V} a_j e^{-i\frac{\pi}{2} Q^V} e^{i\frac{\pi}{2} Q^A} &= e^{-i\frac{\pi}{2} Q^A} b_j e^{i\frac{\pi}{2} Q^A} = a_{j-1}, \\ e^{-i\frac{\pi}{2} Q^A} e^{i\frac{\pi}{2} Q^V} b_j e^{-i\frac{\pi}{2} Q^V} e^{i\frac{\pi}{2} Q^A} &= e^{-i\frac{\pi}{2} Q^A} (-a_j) e^{i\frac{\pi}{2} Q^A} = b_{j+1}, \end{aligned} \quad (\text{B.2})$$

which is the same action as $T_b T_a^{-1}$. Then, because a_j and b_j form a faithful basis for the algebra of operators acting on this tensor product Hilbert space, the operator identity $e^{-i\frac{\pi}{2} Q^A} e^{i\frac{\pi}{2} Q^V} = T_b T_a^{-1}$ holds. Imposing the symmetry generated by $T_b T_a^{-1}$ forbids deformations to the staggered fermion Hamiltonian that couple the a and b Majorana fermions. This is because repeated actions of $T_b T_a^{-1}$ increase the distance between the a and b Majorana operators, so that any local product of them will become increasingly non-local. Therefore, by imposing that all deformations commute with $T_b T_a^{-1}$, we need only consider deformations whose individual terms consist of only the a_j Majorana fermions or only the b_j Majorana fermions. Furthermore, by locality of the Hamiltonian, this implies that T_a and T_b individually commute as well, and therefore lattice translations $T = T_a T_b$ too.

Having found the effect of requiring $T_b T_a^{-1}$ invariance, we next require all deformations to commute with $e^{i\varphi Q^V}$. Let us first consider the $T_b T_a^{-1}$ -symmetric Majorana bi-linears

$$O_2^a(j, j') = a_j a_{j'}, \quad O_2^b(j, j') = b_j b_{j'}, \quad (\text{B.3})$$

where we assume $j < j'$ without loss of generality. An infinitesimal Q^V action transforms these terms by

$$\begin{aligned} \delta O_2^a(j, j') &= \varphi (b_j a_{j'} + a_j b_{j'}) + \mathcal{O}(\varphi^2), \\ \delta O_2^b(j, j') &= -\varphi (a_j b_{j'} + b_j a_{j'}) + \mathcal{O}(\varphi^2). \end{aligned} \quad (\text{B.4})$$

Therefore, the only combinations of these bi-linears that are $U(1)^V$ symmetric take the form

$$O_2^a(j, j') + O_2^b(j, j') = a_j a_{j'} + b_j b_{j'} = 2(c_j^\dagger c_{j'} + c_j c_{j'}^\dagger), \quad (\text{B.5})$$

and the most general $T_b T_a^{-1}$ and $U(1)^V$ symmetric, local deformation using fermion bi-linears is

$$\delta H = -\frac{i}{2} \sum_{j=1}^L \sum_{n=1}^N g_n (a_j a_{j+n} + b_j b_{j+n}), \quad (\text{B.6})$$

where $\lim_{L \rightarrow \infty} N/L = 0$. Note that the original staggered fermion Hamiltonian (5) is of such type for which $n = 1$. Furthermore, because δH commutes with T_b and Q^V , in light of (12), it also commutes with Q^A too.

Having considered symmetric bi-linear operators, we now consider $T_b T_a^{-1}$ -symmetric $2n$ -body terms with $n \geq 2$, which take the form

$$O_{2n}^a(j_1, \dots, j_{2n}) = \prod_{\ell=1}^{2n} a_{j_\ell}, \quad O_{2n}^b(j_1, \dots, j_{2n}) = \prod_{\ell=1}^{2n} b_{j_\ell}. \quad (\text{B.7})$$

An infinitesimal Q^V action transforms these operators by

$$\begin{aligned} \delta O_{2n}^a(\{j_\ell\}) &= \varphi \sum_{\ell'=1}^{2n} a_{j_1} \cdots b_{j_{\ell'}} \cdots a_{j_{2n}} + \mathcal{O}(\varphi^2), \\ \delta O_{2n}^b(\{j_\ell\}) &= -\varphi \sum_{\ell'=1}^{2n} b_{j_1} \cdots a_{j_{\ell'}} \cdots b_{j_{2n}} + \mathcal{O}(\varphi^2). \end{aligned} \quad (\text{B.8})$$

Because each term in δO_{2n}^a consists of $2n - 1$ a_j operators and 1 b_j operator, while each term in δO_{2n}^b contains $2n - 1$ b_j operators and 1 a_j operator, δO_{2n}^a and δO_{2n}^b are linearly independent when $n \geq 2$. Therefore, there is no linear combination of $O_{2n}^a(j_1, \dots, j_{2n})$ and $O_{2n}^b(j_1, \dots, j_{2n})$ for $n \geq 2$ that is symmetric under the entire $U(1)^V$ symmetry, and the only allowed deformations of the staggered fermion model (6) are quadratic terms of the form (B.6).