

The loop equation for the Burgers–KdV hierarchy

Di Yang, Chunhui Zhou

Abstract

The Burgers–KdV hierarchy was introduced towards understanding intersection numbers on the moduli space of Riemann surfaces with boundaries [9, 23]. The goal of this paper is to derive the Dubrovin–Zhang type loop equation for the topological tau-function of the Burgers–KdV hierarchy. As an application, we provide some relations for open intersection numbers.

1 Introduction

Let w, ρ be indeterminates, and let $(\mathcal{A} = \mathbb{C}[w, \rho, w_x, \rho_x, \dots], \partial = \partial_x)$ be the differential algebra. Recall that the Burgers–KdV hierarchy is the following integrable hierarchy introduced by Buryak [9] (cf. [23]):

$$\frac{\partial w}{\partial t_n} = \partial_x K_n, \quad \frac{\partial \rho}{\partial t_n} = \partial_x R_n, \quad n \geq 0, \quad (1.1)$$

$$\frac{\partial w}{\partial s_n} = 0, \quad \frac{\partial \rho}{\partial s_n} = \partial_x Q_n, \quad n \geq 0, \quad (1.2)$$

where $K_n, R_n, Q_n \in \mathcal{A}[\epsilon]$ are given by

$$K_0 = w, \quad R_0 = \rho, \quad Q_0 = w + \frac{\rho^2}{2} + \epsilon \frac{\rho_x}{2}, \quad (1.3)$$

$$\partial_x K_n = \frac{2}{2n+1} \left(w \partial_x + \frac{1}{2} w_x + \frac{\epsilon^2}{8} \partial_x^3 \right) K_{n-1}, \quad (1.4)$$

$$R_n = \frac{2}{2n+1} \left(\left(w + \frac{\rho^2}{2} + \epsilon \left(\frac{\rho_x}{2} + \rho \partial_x \right) + \frac{\epsilon^2}{2} \partial_x^2 \right) R_{n-1} + \left(\frac{\rho}{2} + \frac{3}{4} \epsilon \partial_x \right) K_{n-1} \right), \quad (1.5)$$

$$Q_n = \frac{1}{n+1} \left(w + \frac{\rho^2}{2} + \epsilon \left(\frac{\rho_x}{2} + \rho \partial_x \right) + \frac{\epsilon^2}{2} \partial_x^2 \right) Q_{n-1}, \quad n \geq 1. \quad (1.6)$$

Here the integration constants in (1.4) are chosen as zero. Equations (1.1) with $n = 0$ read

$$\frac{\partial w}{\partial t_0} = \frac{\partial w}{\partial x}, \quad \frac{\partial \rho}{\partial t_0} = \frac{\partial \rho}{\partial x}, \quad (1.7)$$

so we identify t_0 with x .

For an arbitrary solution (w, ρ) in $\mathbb{C}[[\mathbf{t}; \epsilon]] \times \mathbb{C}[[\mathbf{t}, \mathbf{s}; \epsilon]]$ to the Burgers–KdV hierarchy, there exists a tuple $(\tau_1, \tau_2) \in \mathbb{C}((\epsilon))[[\mathbf{t}]] \times \mathbb{C}((\epsilon))[[\mathbf{t}, \mathbf{s}]]$ satisfying

$$w = \epsilon^2 \partial_x^2 \log \tau_1, \quad \rho = \epsilon \partial_x \log \tau_2 \quad (1.8)$$

and a sequence of similar conditions (see Section 2 for the details), called [26] the tau-tuple of the solution (w, ρ) . We call the product $\tau = \tau_1 \tau_2$ the tau-function of the solution.

We are particularly interested in the solution $(w_{\text{top}}, \rho_{\text{top}}) \in \mathbb{C}[[\mathbf{t}; \epsilon]] \times \mathbb{C}[[\mathbf{t}, \mathbf{s}; \epsilon]]$ specified by

$$(w_{\text{top}}, \rho_{\text{top}})|_{t_n=x\delta_{n,0}, s_n=0, n \geq 0} = (x, 0). \quad (1.9)$$

We call $(w_{\text{top}}, \rho_{\text{top}})$ the *topological solution* to the Burgers–KdV hierarchy. The tau-function $\tau_{\text{top}} = \tau_{1,\text{top}} \tau_{2,\text{top}}$ of this solution can [9] be further chosen to satisfy the equation

$$\sum_{n \geq 1} \left(\tilde{t}_n \frac{\partial \tau_{\text{top}}}{\partial t_{n-1}} + s_n \frac{\partial \tau_{\text{top}}}{\partial s_{n-1}} \right) + \frac{t_0^2}{2\epsilon^2} + \frac{s_0}{\epsilon} = 0, \quad (1.10)$$

where $\tilde{t}_n = t_n - \delta_{n,1}$. In this way, τ_{top} is unique up to a constant factor. We call τ_{top} the *topological tau-function* of the Burgers–KdV hierarchy.

According to the Witten–Kontsevich theorem [21, 25], the first component $\tau_{1,\text{top}}$ has the form

$$\tau_{1,\text{top}}(\mathbf{t}; \epsilon) = e^{\sum_{g \geq 0} \epsilon^{2g-2} \mathcal{F}_g^c(\mathbf{t})}, \quad (1.11)$$

where $\mathcal{F}_g^c(\mathbf{t})$ has the expression

$$\mathcal{F}_g^c(\mathbf{t}) = \sum_{n \geq 0} \sum_{k_1, \dots, k_n \geq 0} \frac{\langle \tau_{k_1} \cdots \tau_{k_n} \rangle_g^c}{n!} t_{k_1} \cdots t_{k_n}, \quad \langle \tau_{k_1} \cdots \tau_{k_n} \rangle_g^c = \int_{\overline{\mathcal{M}}_{g,n}} \psi_1^{k_1} \cdots \psi_n^{k_n}, \quad (1.12)$$

where $\overline{\mathcal{M}}_{g,n}$ denotes the moduli space of smooth complex algebraic curves of genus g with n distinct marked points, and ψ_i denotes the first Chern class of the tautological line bundle on $\overline{\mathcal{M}}_{g,n}$. We call $\mathcal{F}_g^c(\mathbf{t})$ the genus g topological free energy for the KdV hierarchy. It is well-known that $\tau_{1,\text{top}}$ satisfies the following Virasoro constraints:

$$\mathcal{L}_m \tau_{1,\text{top}} = 0, \quad m \geq -1. \quad (1.13)$$

where \mathcal{L}_m are operators as follows:

$$\mathcal{L}_{-1} = \sum_{n \geq 1} \tilde{t}_n \frac{\partial}{\partial t_{n-1}} + \frac{t_0^2}{2\epsilon^2}, \quad (1.14)$$

$$\begin{aligned} \mathcal{L}_m &= \sum_{n \geq 0} \frac{(2n+2m+1)!!}{2^{m+1}(2n-1)!!} \tilde{t}_n \frac{\partial}{\partial t_{n+m}} + \frac{1}{16} \delta_{m,0} \\ &\quad + \frac{\epsilon^2}{2} \sum_{n=0}^{m-1} \frac{(2n+1)!!(2m-2n-1)!!}{2^{m+1}} \frac{\partial^2}{\partial t_n \partial t_{m-n-1}}, \quad m \geq 0. \end{aligned} \quad (1.15)$$

The second component has the form

$$\tau_{2,\text{top}}(\mathbf{t}, \mathbf{s}; \epsilon) = e^{\sum_{p \geq 0} \epsilon^{p-1} \mathcal{F}_p^o(\mathbf{t}, \mathbf{s})}, \quad (1.16)$$

where $\mathcal{F}_p^o(\mathbf{t}, \mathbf{s})$ has the expression

$$\mathcal{F}_p^o(\mathbf{t}, \mathbf{s}) = \sum_{m, n \geq 0} \sum_{\substack{k_1, \dots, k_m \geq 0, \\ \ell_1, \dots, \ell_n \geq 0}} \frac{\langle \tau_{k_1} \cdots \tau_{k_m} \sigma_{\ell_1} \cdots \sigma_{\ell_n} \rangle_p^o}{m!n!} t_{k_1} \cdots t_{k_m} s_{\ell_1} \cdots s_{\ell_n}, \quad (1.17)$$

where the correlators $\langle \tau_{k_1} \cdots \tau_{k_m} \sigma_{\ell_1} \cdots \sigma_{\ell_n} \rangle_p^o$ are (cf. [8, 9, 23]) related with intersection numbers on moduli space of Riemann surfaces with boundaries. We call $\mathcal{F}_p^o(\mathbf{t}, \mathbf{s})$ the genus p topological open free energy for the Burgers–KdV hierarchy.

Recall from e.g. [16, 17, 18] that for every $g \geq 1$ there exists a function of $(3g - 2)$ variables

$$F_g^c = F_g^c(v_1, v_2, \dots, v_{3g-2})$$

such that

$$\mathcal{F}_g^c(\mathbf{t}) = F_g^c\left(\frac{\partial v_{\text{top}}(\mathbf{t})}{\partial t_0}, \dots, \frac{\partial^{3g-2} v_{\text{top}}(\mathbf{t})}{\partial t_0^{3g-2}}\right), \quad (1.18)$$

where $v_{\text{top}}(\mathbf{t}) := \partial_{t_0}^2 \mathcal{F}_0^c(\mathbf{t})$. For example,

$$F_1^c = \frac{1}{24} \log v_1, \quad F_2^c = \frac{v_4}{1152v_1^2} - \frac{7v_2v_3}{1920v_1^3} + \frac{v_2^3}{360v_1^4}. \quad (1.19)$$

By using a method from [17] (see also [22]), we will show in Section 2 the following theorem, which gives the jet-variable representations for the topological open free energies \mathcal{F}_p^o , $p \geq 1$.

Theorem 1.1 *There exists a sequence of functions*

$$F_1^o = \frac{1}{2} \log(v_1 + rr_1), \quad (1.20)$$

$$\begin{aligned} F_p^o &= F_p^o(r, v_1, r_1, \dots, v_{2p-1}, r_{2p-1}) \\ &\in v_1^{-(5p-2)}(v_1 + rr_1)^{-(5p-2)} \mathbb{C}[r, v_1, r_1, v_2, r_2, \dots, v_{2p-1}, r_{2p-1}], \quad p \geq 2 \end{aligned} \quad (1.21)$$

satisfying the homogeneous conditions

$$\sum_{k \geq 1} kv_k \frac{\partial F_p^o}{\partial v_k} + \sum_{k \geq 0} kr_k \frac{\partial F_p^o}{\partial r_k} = (p-1)F_p^o, \quad p \geq 2 \quad (1.22)$$

$$\sum_{k \geq 1} (k-1)v_k \frac{\partial F_p^o}{\partial v_k} + \sum_{k \geq 0} \left(k - \frac{1}{2}\right) r_k \frac{\partial F_p^o}{\partial r_k} = \frac{3}{2}(p-1)F_p^o, \quad p \geq 2 \quad (1.23)$$

such that for $p \geq 1$,

$$\mathcal{F}_p^o(\mathbf{t}, \mathbf{s}) = F_p^o\left(r_{\text{top}}(\mathbf{t}, \mathbf{s}), \frac{\partial v_{\text{top}}(\mathbf{t})}{\partial t_0}, \frac{\partial r_{\text{top}}(\mathbf{t}, \mathbf{s})}{\partial t_0}, \dots, \frac{\partial^{2p-1} v_{\text{top}}(\mathbf{t})}{\partial t_0^{2p-1}}, \frac{\partial^{2p-1} r_{\text{top}}(\mathbf{t}, \mathbf{s})}{\partial t_0^{2p-1}}\right), \quad (1.24)$$

where $r_{\text{top}}(\mathbf{t}, \mathbf{s}) = \partial_{t_0} \mathcal{F}_0^o(\mathbf{t}, \mathbf{s})$.

It was shown in [9] that the topological tau-function τ_{top} satisfies the following Virasoro constraints:

$$\mathcal{L}_m^{\text{ext}} \tau_{\text{top}} = 0, \quad m \geq -1, \quad (1.25)$$

where $\mathcal{L}_m^{\text{ext}}$ are operators given by

$$\mathcal{L}_{-1}^{\text{ext}} = \mathcal{L}_{-1} + \sum_{n \geq 1} s_n \frac{\partial}{\partial s_{n-1}} + \frac{s_0}{\epsilon}, \quad (1.26)$$

$$\mathcal{L}_m^{\text{ext}} = \mathcal{L}_m + \sum_{n \geq 0} \frac{(n+m+1)!}{n!} s_n \frac{\partial}{\partial s_{n+m}} + \frac{3(m+1)!}{4} \epsilon \frac{\partial}{\partial s_{m-1}} + \frac{3}{4} \delta_{m,0} \quad m \geq 0. \quad (1.27)$$

Introduce the following notations:

$$A = \frac{1}{\lambda - v}, \quad B = \frac{1}{\lambda - v - \frac{r^2}{2}}, \quad \partial = \sum_{k \geq 0} \left(v_{k+1} \frac{\partial}{\partial v_k} + r_{k+1} \frac{\partial}{\partial r_k} \right). \quad (1.28)$$

By using the Virasoro constraints (1.26)–(1.27) we will show in Section 3 the following theorem.

Theorem 1.2 *The series $\Delta F = \sum_{g \geq 1} \epsilon^{2g-2} F_g^c + \sum_{p \geq 1} \epsilon^{p-1} F_p^o$ is the unique solution to the following equation:*

$$\begin{aligned} & \sum_{k \geq 0} \frac{\partial \Delta F}{\partial v_k} \left(\partial^k(A) + \sum_{j=1}^k \binom{k}{j} \partial^{j-1}(A^{\frac{1}{2}}) \partial^{k+1-j}(A^{\frac{1}{2}}) \right) \\ & + \frac{1}{2} \sum_{k \geq 0} \frac{\partial \Delta F}{\partial r_k} \left(\partial^k(rAB) + \sum_{j=1}^k \binom{k}{j} \partial^{j-1}(A^{\frac{1}{2}}) \partial^{k+1-j}(rA^{\frac{1}{2}}B) \right) \\ & = \frac{\epsilon^2}{2} \sum_{k, \ell \geq 0} \left(\frac{\partial^2 \Delta F}{\partial v_k \partial v_\ell} + \frac{\partial \Delta F}{\partial v_k} \frac{\partial \Delta F}{\partial v_\ell} \right) \partial^{k+1}(A^{\frac{1}{2}}) \partial^{\ell+1}(A^{\frac{1}{2}}) \\ & + \frac{\epsilon^2}{2} \sum_{k, \ell \geq 0} \left(\frac{\partial^2 \Delta F}{\partial v_k \partial r_\ell} + \frac{\partial \Delta F}{\partial v_k} \frac{\partial \Delta F}{\partial r_\ell} \right) \partial^{k+1}(A^{\frac{1}{2}}) \partial^{\ell+1}(rA^{\frac{1}{2}}B) \\ & + \frac{\epsilon^2}{8} \sum_{k, \ell \geq 0} \left(\frac{\partial^2 \Delta F}{\partial r_k \partial r_\ell} + \frac{\partial \Delta F}{\partial r_k} \frac{\partial \Delta F}{\partial r_\ell} \right) \partial^{k+1}(rA^{\frac{1}{2}}B) \partial^{\ell+1}(rA^{\frac{1}{2}}B) \\ & + \sum_{k \geq 0} \frac{\partial \Delta F}{\partial v_k} \left(\frac{\epsilon^2}{16} \partial^{k+2}(A^2) + \epsilon r A^{\frac{1}{2}} B \partial^{k+1}(A^{\frac{1}{2}}) \right) \\ & + \sum_{k \geq 0} \frac{\partial \Delta F}{\partial r_k} \left(\epsilon^2 \partial^{k+1} \left(\frac{A}{16r} \partial(r^2 AB) + \frac{r}{12} \partial(B^3) \right) + \epsilon \left(\frac{1}{4} r A^{\frac{1}{2}} B \partial^{k+1}(rA^{\frac{1}{2}}B) + \frac{3}{4} \partial^{k+1}(B^2) \right) \right) \\ & + \epsilon \left(\frac{A}{16r} \partial(r^2 AB) + \frac{r}{12} \partial(B^3) \right) + \frac{A^2}{16} + B^2 - \frac{1}{4} AB. \end{aligned} \quad (1.29)$$

Example 1.3 Denote $u := v + \frac{r^2}{2}$ and $u_k := \partial^k(u)$ for $k \geq 0$. From Theorem 1.2 and the formulae (1.19), we obtain the following expressions:

$$\begin{aligned} F_1^o &= \frac{1}{2} \log(v_1 + rr_1) = \frac{1}{2} \log u_1, \\ F_2^o &= \frac{1}{24v_1 (v_1 + rr_1)^3} \left(-r^2 v_2 r_1^3 + 3r^2 r_3 v_1^2 - 4r^3 v_1 r_2^2 + 4v_1^3 r_2 - 3rv_1 r_1^4 + v_1^2 r_1^3 + 3r^3 r_3 v_1 r_1 \right. \\ &\quad \left. + 6r^2 v_1 r_1^2 r_2 + 18rv_1^2 r_1 r_2 + 3v_3 r^2 v_1 r_1 + 3v_3 r v_1^2 - 8r^2 v_1 v_2 r_2 - 9rv_1 v_2 r_1^2 - 4rv_1 v_2^2 \right) \\ &= -\frac{ru_2^2}{6u_1^3} + \frac{r_1 u_2 + 3ru_3}{24u_1^2} + \frac{r_1(r_1^2 - u_2)}{24(u_1 - rr_1)u_1} + \frac{r_2}{8u_1} - \frac{r_2}{24(u_1 - rr_1)}. \end{aligned}$$

Analogous to the work of Itzykson–Zuber [20], we introduce the following variables:

$$I_k := \sum_{n \geq 0} t_{n+k} \frac{v_{\text{top}}^n}{n!}, \quad (1.30)$$

$$J_k := \sum_{n \geq 0} t_{n+k+1} \sum_{i=0}^n \frac{v_{\text{top}}^i r_{\text{top}}^{2n-2i+1}}{i!(2n-2i+1)!!} + \sum_{n \geq 0} \frac{s_{n+k}}{n!} \left(v_{\text{top}} + \frac{r_{\text{top}}^2}{2} \right)^n. \quad (1.31)$$

We will show in Section 2 that the transformation from \mathbf{t}, \mathbf{s} to $I_k, J_k, k \geq 0$ is invertible. By using Theorem 1.1, we will prove in Section 4 that for $p \geq 2$, the genus p open free energy \mathcal{F}_p^o can be expressed as an element in $\mathbb{C}[(1-I_1)^{-1}, (1-I_1-J_0J_1)^{-1}, J_0, J_1, I_2, J_2, \dots, I_{2p-1}, J_{2p-1}]$. As an application of these expressions, we will prove in Section 4 the following corollary.

Corollary 1.4 *Let $m, n \geq 0$, $k_1, \dots, k_m \geq 2$, $\ell_1, \dots, \ell_n \geq 2$ and $d \geq 0$, $p \geq 2$. If $m + n + d > 13p - 7$, the following identities hold true:*

$$\sum_{q \geq 0} \frac{(-1)^q}{(d-q)!q!(7p-4-q)!} \Lambda_{k_1, \dots, k_m; 1^{d-q}, \ell_1, \dots, \ell_n}(\mathcal{F}_p^o)|_{s_0=1, t_{\geq 0}=s_{\geq 1}=0} = 0. \quad (1.32)$$

Here $\Lambda_{k_1, \dots, k_m; \ell_1, \dots, \ell_n}$ are operators defined as follows:

$$\Lambda_{k_1, \dots, k_m; \ell_1, \dots, \ell_n} := \prod_{i=1}^m \left(\frac{\partial}{\partial t_{k_i}} - \sum_{j=0}^{k_i-1} \frac{(-1)^j}{2^j(2j+1)j!} \frac{\partial}{\partial s_{k_i-j-1}} \right) \prod_{i=1}^n \left(\sum_{j=0}^{\ell_i} \frac{(-1)^j}{2^j j!} \frac{\partial}{\partial s_{\ell_i-j}} \right). \quad (1.33)$$

We also have the following conjectural statement: for $m, n \geq 0$, $k_1, \dots, k_m \geq 2$, $\ell_1, \dots, \ell_n \geq 2$ and $d \geq 0$, $p \geq 2$, the identities

$$\sum_{q \geq 0} \frac{(-1)^q}{(d-q)!q!(3p-3-q)!} \Lambda_{k_1, \dots, k_m; 1^{d-q}, \ell_1, \dots, \ell_n}(\mathcal{F}_p^o)|_{s_0=1, t_{\geq 0}=s_{\geq 1}=0} = 0 \quad (1.34)$$

hold true when $m + n + d > 6p - 6$. This is stronger than (1.32).

Organization of the paper. In Section 2, we prove Theorem 1.1. In Section 3, we prove Theorem 1.2. In Section 4, we present some applications.

Acknowledgements. We are grateful to Professor Youjin Zhang his advice and for suggesting this project. We also thank Professor Si-Qi Liu for helpful discussions. The work is partially supported by NSFC grants (No. 12371254, No. 12322111), by National Key R and D Program of China 2020YFA0713100, and by CAS No. YSBR-032.

2 Jet-variable representation

In this section, we will prove Theorem 1.1.

Let $\Omega_{m,n}$, $m, n \geq 0$, be the two-point correlation functions of the KdV hierarchy (cf. [4, 17]). Recall that, for an arbitrary solution (w, ρ) to the Burgers-KdV hierarchy, the defining equations of the associated tau-tuple (τ_1, τ_2) [26] (cf. [4, 9, 16, 17]) are given by

$$\epsilon^2 \frac{\partial^2 \log \tau_1}{\partial t_m \partial t_n} = \Omega_{m,n}, \quad \epsilon \frac{\partial \log \tau_2}{\partial t_n} = R_n, \quad \epsilon \frac{\partial \log \tau_2}{\partial s_n} = Q_n, \quad m, n \geq 0. \quad (2.1)$$

We denote

$$\langle\langle \tau_{k_1} \cdots \tau_{k_m} \rangle\rangle^c := \epsilon \frac{\partial^m \log \tau_1}{\partial t_{k_1} \cdots \partial t_{k_m}}, \quad (2.2)$$

$$\langle\langle \tau_{k_1} \cdots \tau_{k_m} \sigma_{\ell_1} \cdots \sigma_{\ell_n} \rangle\rangle^o := \epsilon \frac{\partial^{m+n} \log \tau_2}{\partial t_{k_1} \cdots \partial t_{k_m} \partial s_{\ell_1} \cdots \partial s_{\ell_n}}. \quad (2.3)$$

It is well known that $\langle\langle \tau_{k_1} \cdots \tau_{k_m} \rangle\rangle^c$ are differential polynomials of w . By the definition (2.1) and the hierarchy (1.1)–(1.2), we know that $\langle\langle \tau_{k_1} \cdots \tau_{k_m} \sigma_{\ell_1} \cdots \sigma_{\ell_n} \rangle\rangle^o$ are also differential polynomials in $\mathcal{A}[\epsilon]$.

It follows from equation (1.10) that the topological solution $(w_{\text{top}}, \rho_{\text{top}})$ satisfies the following Euler–Lagrange equations:

$$\sum_{n \geq 0} \tilde{t}_{n+1} K_n(\underline{w}; \epsilon) + t_0 = 0, \quad (2.4)$$

$$\sum_{n \geq 0} (\tilde{t}_{n+1} R_n(\underline{w}, \underline{\rho}; \epsilon) + s_{n+1} Q_n(\underline{w}, \underline{\rho}; \epsilon)) + s_0 = 0. \quad (2.5)$$

Here, $\underline{w} = (w, w_x, w_{xx}, \dots)$ and $\underline{\rho} = (\rho, \rho_x, \rho_{xx}, \dots)$, and for simplifying the notations we use (w, ρ) to denote $(w_{\text{top}}, \rho_{\text{top}})$.

By taking $\epsilon = 0$ in the system of equations (2.4) and (2.5), we have that the pair

$$(v, r) := (w, \rho)|_{\epsilon=0}$$

satisfies the following genus zero Euler–Lagrange equations:

$$\sum_{n \geq 0} \tilde{t}_n \frac{v^n}{n!} = 0, \quad (2.6)$$

$$\sum_{n \geq 0} \tilde{t}_{n+1} \sum_{i=0}^n \frac{v^i r^{2n-2i+1}}{i!(2n-2i+1)!!} + \sum_{n \geq 0} \frac{s_n}{n!} \left(v + \frac{r^2}{2}\right)^n = 0. \quad (2.7)$$

Then it is easy to see from the definitions (1.30), (1.31) that

$$I_0 = v, \quad J_0 = r, \quad (2.8)$$

$$I_k = \frac{\partial_x(I_{k-1}) - \delta_{k,1}}{v_1}, \quad J_k = \frac{\partial_x(J_{k-1}) - r_1 I_k}{v_1 + rr_1}, \quad (2.9)$$

where $k \geq 1$. Straightforwardly, we have the following lemma.

Lemma 2.1 *The functions $I_k, J_k, k \geq 1$, have the form*

$$I_k = \sum_{\lambda \in \mathcal{P}_{k-1}} c'_{k;\lambda} v_1^{-k-\ell(\lambda)} v_{\lambda+1}, \quad (2.10)$$

$$J_k = \sum_{1 \leq d \leq k} \sum_{\substack{\lambda, \mu \in \mathcal{P} \\ |\lambda| + |\mu| = k-d}} c''_{k;\lambda;\mu}(v_1^{-1}, u_1^{-1}) r_d v_{\lambda+1} u_{\mu+1}, \quad (2.11)$$

where $c'_{k;\lambda}$ are rational numbers, and $c''_{k;\lambda;\mu}(z_1, w_1)$ are homogeneous polynomials of z_1, w_1 with degree $k + \ell(\lambda) + \ell(\mu) + 1$ and with rational coefficients. Here we denote by \mathcal{P} the set of partitions

$$\mathcal{P} = \{(\lambda_1, \lambda_2, \dots) | \lambda_1 \geq \lambda_2 \geq \dots \geq 1, \sum_i \lambda_i < \infty\},$$

by $\ell(\lambda)$ the length of λ , by $|\lambda|$ the summation of the components of λ , and by \mathcal{P}_k the set of partitions λ with $|\lambda| = k$. We also denote $a_\lambda := a_{\lambda_1} \cdots a_{\lambda_{\ell(\lambda)}}$.

For example,

$$I_1 = 1 - \frac{1}{v_1}, \quad I_2 = \frac{v_2}{v_1^3}, \quad J_1 = \frac{r_1}{v_1 u_1}, \quad J_2 = \frac{r_2}{v_1 u_1^2} - r_1 \left(\frac{u_2}{v_1 u_1^3} + \frac{v_2}{v_1^3 u_1} + \frac{v_2}{v_1^2 u_1^2} \right),$$

and conversely,

$$v_1 = \frac{1}{U_1}, \quad r_1 = \frac{J_1}{U_1 U_2}, \quad v_2 = \frac{I_2}{U_1^3}, \quad r_2 = \frac{J_2}{U_2^3} + \frac{J_1^3}{U_1^2 U_2^3} + J_1 J_2 \left(\frac{1}{U_1 U_2^3} + \frac{1}{U_1^2 U_2^2} + \frac{1}{U_1^3 U_2} \right),$$

where we denote

$$U_1 := 1 - I_1, \quad U_2 := 1 - I_1 - J_0 J_1. \quad (2.12)$$

The following lemma gives the inverse of (1.30)–(1.31), which generalizes the one in [16, 27].

Lemma 2.2 *For $k \geq 0$, we have*

$$t_k = \sum_{n \geq 0} \frac{(-1)^n}{n!} I_0^n I_{n+k}, \quad (2.13)$$

$$s_k = \sum_{n \geq 0} \sum_{i=0}^n \frac{(-1)^{n+1} I_0^i J_0^{2n-2i+1}}{2^{n-i} i! (n-i)! (2n-2i+1)} I_{n+k+1} + \sum_{n \geq 0} \frac{(-1)^n}{n!} \left(I_0 + \frac{1}{2} J_0^2 \right)^n J_{n+k}. \quad (2.14)$$

Proof See in [27] the proof for (2.13). To prove (2.14), let us denote

$$\alpha_n := \sum_{i=0}^n \frac{(-1)^{n+1} I_0^i J_0^{2n-2i+1}}{2^{n-i} i! (n-i)! (2n-2i+1)}, \quad \beta_n := \frac{(-1)^n}{n!} \left(I_0 + \frac{1}{2} J_0^2 \right)^n. \quad (2.15)$$

Substitute (2.13) and (2.14) into (1.31) and then we need to verify

$$\begin{aligned} J_k &= \sum_{n,m \geq 0} \frac{(-1)^m I_0^m}{m!} I_{n+m+k+1} \sum_{i=0}^n \frac{I_0^i J_0^{2n-2i+1}}{i! (2n-2i+1)!!} \\ &\quad + \sum_{n,m \geq 0} \frac{1}{n!} \left(I_0 + \frac{J_0^2}{2} \right)^n (\alpha_m I_{n+m+k+1} + \beta_m J_{n+m+k}), \quad k \geq 0. \end{aligned} \quad (2.16)$$

Equivalently, it suffices to check the following identities:

$$\sum_{n=0}^k \frac{1}{n!} \left(I_0 + \frac{1}{2} J_0^2 \right)^n \alpha_{k-n} = -\frac{J_0^{2k+1}}{(2k+1)!!} = -\sum_{n=0}^k \frac{(-1)^{k-n} I_0^{k-n}}{(k-n)!} \sum_{i=0}^n \frac{I_0^i J_0^{2n-2i+1}}{i! (2n-2i+1)!!}, \quad (2.17)$$

$$\sum_{n=0}^k \frac{1}{n!} \left(I_0 + \frac{1}{2} J_0^2 \right)^n \beta_{k-n} = \delta_{k,0}. \quad (2.18)$$

The lemma is proved. \square

It follows from Lemma 2.2 and the relations (2.8)–(2.9) that

$$t_k = v_k - \delta_{k,1} + \text{higher-order terms in } \mathbb{C}[[v, v_1 - 1, v_2, \dots]], \quad (2.19)$$

$$s_k = r_k + \text{higher-order terms in } \mathbb{C}[[v, r, v_1 - 1, r_1, v_2, r_2, \dots]]. \quad (2.20)$$

By applying this transformation, the free energies \mathcal{F}_p^o can be expressed as an element

$$F_p^o \in \mathbb{C}[[v, r, v_1 - 1, r_1, v_2, r_2, \dots]]. \quad (2.21)$$

Lemma 2.3 For $n \geq 0$, the differential polynomials R_n and Q_n can be expanded into the form

$$R_n = \sum_{i=0}^n \left(a_i(\underline{w}, \underline{\rho}; \epsilon) \sum_{j=0}^{n-i} \frac{w^j \rho^{2n-2i-2j+1}}{j!(2n-2i-2j+1)!} + b_i(\underline{w}, \underline{\rho}; \epsilon) \frac{w^{n-i+1}}{(n-i+1)!} \right), \quad (2.22)$$

$$Q_n = \sum_{i=0}^{n+1} a_i(\underline{w}, \underline{\rho}; \epsilon) \frac{(w + \frac{1}{2}\rho^2)^{n-i+1}}{(n-i+1)!}, \quad (2.23)$$

where the coefficients a_i and b_i are certain elements in $\mathcal{A}[\epsilon]$, which do not explicitly contain w . Moreover, a_i 's and b_i 's have the following form:

$$a_i(\underline{w}, \underline{\rho}; \epsilon) = \sum_{j=\lceil \frac{i+1}{2} \rceil}^{2i-1} \epsilon^j a_{i,j}(\underline{w}, \underline{\rho}), \quad b_i(\underline{w}, \underline{\rho}; \epsilon) = \sum_{j=\lceil \frac{i+1}{2} \rceil}^{2i-2} \epsilon^j b_{i,j}(\underline{w}, \underline{\rho}). \quad (2.24)$$

Proof By using formula (1.4), one can obtain

$$K_n(\underline{w}; \epsilon) = \sum_{i \geq 0} c_i(\underline{w}; \epsilon) \frac{w^{n-i+1}}{(n-i+1)!}. \quad (2.25)$$

where the coefficients $c_i = c_i(\underline{w}; \epsilon)$ satisfy

$$\begin{aligned} \left(i - \frac{1}{2} \right) \partial_x c_i &= \left(\left(-i + \frac{1}{2} \right) w_x + \frac{1}{8} \epsilon^2 \partial_x^3 \right) c_{i-1} + \frac{1}{8} \epsilon^2 \left(w_{xxx} + 3w_{xx} \partial_x + 3w_x \partial_x^2 \right) c_{i-2} \\ &\quad + \frac{3}{8} \epsilon^2 \left(w_x w_{xx} + w_x^2 \partial_x \right) c_{i-3} + \frac{1}{8} \epsilon^2 w_x^3 c_{i-4}. \end{aligned} \quad (2.26)$$

By substituting (2.22)–(2.23) into (1.5)–(1.6) respectively, we have

$$\begin{aligned} ia_i &= \left(\frac{1}{2} \epsilon \rho_x + \epsilon \rho \partial_x + \frac{1}{2} \epsilon^2 \partial_x^2 \right) a_{i-1} + \left(\epsilon \rho u_x + \frac{1}{2} (w_x + \rho \rho_x)_x + \epsilon^2 (w_x + \rho \rho_x) \partial_x \right) a_{i-2} \\ &\quad + \frac{1}{2} \epsilon^2 (w_x + \rho \rho_x)^2 a_{i-3}, \end{aligned} \quad (2.27)$$

$$\begin{aligned} \left(i - \frac{1}{2} \right) b_i &= \left(\frac{1}{2} \rho^2 + \frac{1}{2} \epsilon \rho_x + \frac{1}{2} \epsilon^2 \partial_x^2 \right) b_{i-1} + \left(\epsilon \rho w_x + \frac{1}{2} w_{xx} + \epsilon^2 w_x \partial_x \right) b_{i-2} \\ &\quad + \frac{1}{2} \epsilon^2 w_x^2 b_{i-3} - \frac{1}{2} \rho a_{i-1} + \left(\epsilon \rho \rho_x + \frac{1}{2} \epsilon^2 \rho_{xx} + \epsilon^2 \rho_x \partial_x \right) a_{i-2} \\ &\quad + \frac{1}{2} \epsilon^2 \rho_x (2w_x + \rho \rho_x) a_{i-3} + \left(\frac{1}{2} \rho + \frac{3}{4} \epsilon \partial_x \right) c_{i-1} + \frac{3}{4} \epsilon w_x c_{i-2}. \end{aligned} \quad (2.28)$$

The lemma can then be proved by induction. \square

Example 2.4 The first several coefficients a_i and b_i are as follows:

$$\begin{aligned} a_0 &= 1, \quad a_1 = \frac{1}{2}\epsilon\rho_x, \\ a_2 &= \frac{1}{2}\epsilon(\rho w_x + \rho^2\rho_x) + \frac{1}{8}\epsilon^2(3\rho_x^2 + 2w_{xx} + 4\rho\rho_{xx}) + \frac{1}{8}\epsilon^3\rho_{xxx}. \\ b_0 &= 0, \quad b_1 = \frac{1}{2}\epsilon(w_x + \rho\rho_x) + \frac{1}{3}\epsilon^2\rho_{xx}, \\ b_2 &= \frac{1}{24}\epsilon^2(12w_x\rho_x + 15\rho\rho_x^2 + 4\rho w_{xx} + 4\rho^2\rho_{xx}) + \frac{1}{24}\epsilon^3(16\rho_x\rho_{xx} + 3w_{xxx} + 5\rho\rho_{xxx}) + \frac{1}{15}\epsilon^4\rho^{(4)}. \end{aligned}$$

Lemma 2.5 Let $\Delta\underline{v} = (\Delta v_0, \Delta v_1, \dots)$ and $\Delta\underline{r} = (\Delta r_0, \Delta r_1, \dots)$. Then we have

$$\begin{aligned} &\sum_{n \geq 0} (t_{n+1}R_n(\underline{v}(\mathbf{t}) + \Delta\underline{v}, \underline{r}(\mathbf{t}, \mathbf{s}) + \Delta\underline{r}; \epsilon) + s_{n+1}Q_n(\underline{v}(\mathbf{t}) + \Delta\underline{v}, \underline{r}(\mathbf{t}, \mathbf{s}) + \Delta\underline{r}; \epsilon)) + s_0 \\ &= \sum_{m,n \geq 0} \sum_{\substack{k_1, \dots, k_m; \ell_1, \dots, \ell_n \\ \ell_1, \dots, \ell_n \geq 0}} \mathcal{V}_{k_1, \dots, k_m; \ell_1, \dots, \ell_n}(\mathbf{t}, \mathbf{s}; \epsilon) \frac{\Delta v_{k_1} \cdots \Delta v_{k_m} \Delta r_{\ell_1} \cdots \Delta r_{\ell_n}}{m!n!}, \end{aligned} \quad (2.29)$$

where the coefficients $\mathcal{V}_{k_1, \dots, k_m; \ell_1, \dots, \ell_n}$ have the following expressions:

$$\mathcal{V}_{k_1, \dots, k_m; \ell_1, \dots, \ell_n}(\mathbf{t}, \mathbf{s}; \epsilon) = \sum_{p \geq 0} \epsilon^p V_{k_1, \dots, k_m; \ell_1, \dots, \ell_n}^{[p]}(\underline{v}(\mathbf{t}), \underline{r}(\mathbf{t}, \mathbf{s}))$$

with

$$V_{k_1, \dots, k_m; \ell_1, \dots, \ell_n}^{[p]}(\underline{v}, \underline{r}) \in \sum_{\substack{i,j \geq 0 \\ i+j \leq 4p+2c_0-1}} v_1^{-i}(v_1 + rr_1)_1^{-j} \mathbb{C}[r, v_1, r_1, \dots, v_{2p+c_0}, r_{2p+c_0}]. \quad (2.30)$$

Here c_0 counts the number of 0 in $(k_1, \dots, k_m, \ell_1, \dots, \ell_n)$.

Proof By using Lemma 2.3, we can express $\mathcal{V}_{k_1, \dots, k_m; \ell_1, \dots, \ell_n}(\mathbf{t}, \mathbf{s}; \epsilon)$ in terms of $a_{k,i}, b_{k,i}$ and I_k, J_k as follows:

$$\begin{aligned} \mathcal{V}_{\emptyset; \emptyset} &= \sum_{n \geq 0} \left(t_n \sum_{k \geq 0} a_k(\underline{v}, \underline{r}; \epsilon) \sum_{j=0}^{n-k-1} \frac{v^j r^{2n-2k-2j-1}}{j!(2n-2k-2j-1)!!} + t_n \sum_{k \geq 0} b_k(\underline{v}, \underline{r}; \epsilon) \frac{v^{n-k-1}}{(n-k-1)!} \right. \\ &\quad \left. + s_n \sum_{k \geq 0} a_k(\underline{v}, \underline{r}; \epsilon) \frac{(v + \frac{1}{2}r^2)^{n-k}}{(n-k)!} \right) \\ &= \sum_{k \geq 0} \sum_{j=\lceil \frac{k+1}{2} \rceil}^{2k-1} \epsilon^j a_{k,j}(\underline{v}, \underline{r}) J_k + \sum_{k \geq 0} \sum_{j=\lceil \frac{k+1}{2} \rceil}^{2k-2} \epsilon^j b_{k,j}(\underline{v}, \underline{r}) I_{k+1}, \end{aligned} \quad (2.31)$$

$$\mathcal{V}_{0; \emptyset} = \sum_{k \geq 0} \left(\sum_{j=\lceil \frac{k+1}{2} \rceil}^{2k-1} \epsilon^j a_{k,j}(\underline{v}, \underline{r}) J_{k+1} + \sum_{j=\lceil \frac{k+1}{2} \rceil}^{2k-2} \epsilon^j b_{k,j}(\underline{v}, \underline{r}) I_{k+2} \right), \quad (2.32)$$

$$\mathcal{V}_{\emptyset; 0} = \sum_{k \geq 0} \left(\sum_{j=\lceil \frac{k+1}{2} \rceil}^{2k-1} \epsilon^j \frac{\partial a_{k,j}(\underline{v}, \underline{r})}{\partial r} J_k + r \sum_{j=\lceil \frac{k+1}{2} \rceil}^{2k-1} \epsilon^j a_{k,j}(\underline{v}, \underline{r}) J_{k+1} \right)$$

$$+ \left(\sum_{j=\lceil \frac{k+1}{2} \rceil}^{2k-1} \epsilon^j a_{k,j}(\underline{v}, \underline{r}) + \sum_{j=\lceil \frac{k+1}{2} \rceil}^{2k-2} \epsilon^j \frac{\partial b_{k,j}(\underline{v}, \underline{r})}{\partial r} \right) I_{k+1} \right), \\ \dots \quad \dots \quad (2.33)$$

Then by using Lemma 2.1, the lemma is proved. \square

Now we are ready to prove Theorem 1.1.

Proof of Theorem 1.1 Using Lemma 2.5 and taking

$$\Delta v_k = \sum_{g \geq 1} \epsilon^{2g} \partial_x^{k+2} \mathcal{F}_g^c, \quad \Delta r_k = \sum_{p \geq 1} \epsilon^p \partial_x^{k+1} \mathcal{F}_p^o, \quad (2.34)$$

we have

$$\sum_{m,n \geq 0} \sum_{\substack{k_1, \dots, k_m \geq 0, \\ \ell_1, \dots, \ell_n \geq 0}} \frac{\mathcal{V}_{k_1, \dots, k_m; \ell_1, \dots, \ell_n}}{m!n!} \Delta v_{k_1} \cdots \Delta v_{k_m} \Delta r_{\ell_1} \cdots \Delta r_{\ell_n} - r_0 - \Delta r_0 = 0. \quad (2.35)$$

Then by taking the coefficient of ϵ^p of (2.35), we have the following recursion:

$$-\frac{1}{v_1 + rr_1} \partial_x F_p^o = \widetilde{\sum} \frac{V_{k_1, \dots, k_m; \ell_1, \dots, \ell_n}^{[q]}}{m!n!} \partial_x^{k_1+2} F_{g_1}^c \cdots \partial_x^{k_m+2} F_{g_m}^c \partial_x^{\ell_1+1} F_{p_1}^o \cdots \partial_x^{\ell_n+1} F_{p_n}^o, \quad (2.36)$$

where the summation $\widetilde{\sum}$ is over

$$\begin{aligned} k_1, \dots, k_m, \ell_1, \dots, \ell_n &\geq 0, \quad g_1, \dots, g_m, p_1, \dots, p_n \geq 1, \\ k_1 + \cdots + k_m + \ell_1 + \cdots + \ell_n &\leq q, \quad q + 2g_1 + \cdots + 2g_m + p_1 + \cdots + p_n = p, \end{aligned}$$

and we used the result

$$V_{\emptyset;0}^{[0]} = I_1 + J_0 J_1 = 1 + \frac{1}{v_1 + rr_1}. \quad (2.37)$$

By using formulae (2.30), (2.36) and the fact that

$$\partial_x^2 F_p^c \in v_1^{-(4p-4)} \mathbb{C}[v_1, v_2, \dots, v_{3p}], \quad p \geq 1, \quad (2.38)$$

we can obtain recursively from (2.36) that

$$\partial_x F_p^o \in v_1^{-(5p-1)} (v_1 + rr_1)^{-(5p-1)} \mathbb{C}[r, v_1, r_1, \dots, v_{2p}, r_{2p}]. \quad (2.39)$$

It follows from (2.21) and (2.39) that

$$F_p^o \in \mathbb{C}[\log v_1, \log(v_1 + rr_1), v_1^{-1}, (v_1 + rr_1)^{-1}, r, v_1, r_1, v_2, r_2, \dots, v_{2p-1}, r_{2p-1}]. \quad (2.40)$$

For $p = 1$, formula (2.36) reads

$$\partial_x F_1^o = -(v_1 + rr_1) V_{\emptyset;\emptyset}^{[1]} = \frac{1}{2} \partial_x \log(v_1 + rr_1). \quad (2.41)$$

Hence (1.20) is proved.

Now let us consider $p \geq 2$. In this case, we will use two scaling invariant properties. From the equations (1.13) and (1.25), we obtain

$$\sum_{n \geq 0} \frac{2n+1}{2} \tilde{t}_n \frac{\partial \mathcal{F}_p^o}{\partial t_n} + \sum_{n \geq 0} (n+1)s_n \frac{\partial \mathcal{F}_p^o}{\partial s_n} = 0. \quad (2.42)$$

From (1.1)–(1.6), we find that when $m+n \geq 1$, $\langle\langle \tau_{k_1} \cdots \tau_{k_m} \sigma_{\ell_1} \cdots \sigma_{\ell_n} \rangle\rangle^o \in \mathcal{A}[\epsilon]$ satisfy

$$\begin{aligned} & \sum_{k \geq 0} \left(kw_{kx} \frac{\partial}{\partial w_{kx}} + k\rho_{kx} \frac{\partial}{\partial \rho_{kx}} - \epsilon \frac{\partial}{\partial \epsilon} \right) \langle\langle \tau_{k_1} \cdots \tau_{k_m} \sigma_{\ell_1} \cdots \sigma_{\ell_n} \rangle\rangle^o \\ &= (m+n-1) \langle\langle \tau_{k_1} \cdots \tau_{k_m} \sigma_{\ell_1} \cdots \sigma_{\ell_n} \rangle\rangle^o, \end{aligned} \quad (2.43)$$

$$\begin{aligned} & \sum_{k \geq 0} \left((k-1)w_{kx} \frac{\partial}{\partial w_{kx}} + \left(k-\frac{1}{2}\right)\rho_{kx} \frac{\partial}{\partial \rho_{kx}} - \frac{3}{2}\epsilon \frac{\partial}{\partial \epsilon} \right) \langle\langle \tau_{k_1} \cdots \tau_{k_m} \sigma_{\ell_1} \cdots \sigma_{\ell_n} \rangle\rangle^o \\ &= -\left(k_1 + \cdots + k_m + \ell_1 + \cdots + \ell_n - m - \frac{n}{2} + \frac{3}{2}\right) \langle\langle \tau_{k_1} \cdots \tau_{k_m} \sigma_{\ell_1} \cdots \sigma_{\ell_n} \rangle\rangle^o. \end{aligned} \quad (2.44)$$

Note that

$$\partial_x^k w_{\text{top}} \Big|_{\mathbf{t}=0} = \delta_{k,1}, \quad \partial_x^k \rho_{\text{top}} \Big|_{\mathbf{t}=\mathbf{s}=0} = 0. \quad (2.45)$$

Then by taking $\mathbf{t} = \mathbf{s} = 0$ in (2.3) and comparing the coefficients of each power of ϵ in the both sides, we obtain that

$$\langle\langle \tau_{k_1} \cdots \tau_{k_m} \sigma_{\ell_1} \cdots \sigma_{\ell_n} \rangle\rangle_p^o \neq 0, \quad (2.46)$$

only if

$$k_1 + \cdots + k_m + \ell_1 + \cdots + \ell_n - m - \frac{n}{2} = \frac{3}{2}(p-1). \quad (2.47)$$

Equivalently, we have

$$\sum_{k \geq 0} \left((k-1)t_k \frac{\partial \mathcal{F}_p^o}{\partial t_k} + \left(k-\frac{1}{2}\right)s_k \frac{\partial \mathcal{F}_p^o}{\partial s_k} \right) = \frac{3}{2}(p-1)\mathcal{F}_p^o. \quad (2.48)$$

Together with (2.42), we arrive at

$$\sum_{k \geq 1} \left(\tilde{t}_k \frac{\partial \mathcal{F}_p^o}{\partial t_k} + s_k \frac{\partial \mathcal{F}_p^o}{\partial s_k} \right) = -(p-1)\mathcal{F}_p^o. \quad (2.49)$$

The conditions (1.22) and (1.23) follow from (2.49) and (2.48) respectively. Then (1.21) follows from (1.22) and (2.40). The theorem is proved. \square

We also have the conjectural form for F_p^o with $p \geq 2$:

$$F_p^o = \frac{v_1^{4p-4}}{(v_1 + rr_1)^{3p-3}} \sum_{i=1}^{3p-3} \sum_{\substack{\lambda, \mu \in \mathcal{P} \\ \lambda_i \leq 2p-2 \\ \mu_i \leq 2p-1}} \sum_{|\lambda|+|\mu|=} C_{\lambda; \mu}^{[i]} r^i \frac{v_{\lambda+1} r_\mu}{v_1^{|\lambda|+\ell(\lambda)+|\mu|}}, \quad (2.50)$$

where $C_{\lambda; \mu}^{[i]}$ are certain rational numbers.

3 Loop equation

In this section, we derive the Dubrovin–Zhang type loop equation for the topological tau-function of the Burgers–KdV hierarchy using the Virasoro constraints.

Consider the generating series

$$\mathcal{L}(\lambda) := \sum_{m \geq -1} \frac{\mathcal{L}_m^{\text{ext}}}{\lambda^{m+2}}, \quad (3.1)$$

where $\mathcal{L}_m^{\text{ext}}$ are the operators given by (1.27). It is easy to verify that the series $\mathcal{L}(\lambda)$ can be equivalently written as

$$\mathcal{L}(\lambda) = [\mathcal{I}_1(\lambda)\mathcal{J}_1(\lambda) + \mathcal{I}_2(\lambda)\mathcal{J}_2(\lambda)]_- + \frac{\epsilon^2}{2}\mathcal{J}_1(\lambda)^2 - \frac{3\epsilon}{4}\partial_\lambda\mathcal{J}_2(\lambda) + \frac{1}{\lambda} \left(\frac{t_0^2}{2\epsilon^2} + \frac{s_0}{\epsilon} \right) + \frac{5}{8\lambda^2}, \quad (3.2)$$

where $[]_-$ means taking the terms with negative powers of λ , and

$$\mathcal{I}_1(\lambda) := \sum_{n \geq 0} \frac{2^n \lambda^{n-\frac{1}{2}}}{(2n-1)!!} \tilde{t}_n, \quad \mathcal{I}_2(\lambda) := \sum_{n \geq 0} \frac{\lambda^n}{n!} s_n, \quad (3.3)$$

$$\mathcal{J}_1(\lambda) := \sum_{n \geq 0} \frac{(2n+1)!!}{2^{n+1} \lambda^{n+\frac{3}{2}}} \frac{\partial}{\partial t_n}, \quad \mathcal{J}_2(\lambda) := \sum_{n \geq 0} \frac{(n+1)!}{\lambda^{n+2}} \frac{\partial}{\partial s_n}. \quad (3.4)$$

We have the following lemma.

Lemma 3.1 *The constraints (1.25) are equivalent to the following equations:*

$$[(\mathcal{I}_1(\lambda)\mathcal{J}_1(\lambda) + \mathcal{I}_2(\lambda)\mathcal{J}_2(\lambda))(\mathcal{F}_0^c)]_- + \frac{1}{2}(\mathcal{J}_1(\lambda)(\mathcal{F}_0^c))^2 + \frac{t_0^2}{2\lambda} = 0, \quad (3.5)$$

$$[(\mathcal{I}_1(\lambda)\mathcal{J}_1(\lambda) + \mathcal{I}_2(\lambda)\mathcal{J}_2(\lambda))(\mathcal{F}_0^o)]_- + \mathcal{J}_1(\lambda)(\mathcal{F}_0^c)\mathcal{J}_1(\lambda)(\mathcal{F}_0^o) + \frac{s_0}{\lambda} = 0, \quad (3.6)$$

$$\begin{aligned} &[(\mathcal{I}_1(\lambda)\mathcal{J}_1(\lambda) + \mathcal{I}_2(\lambda)\mathcal{J}_2(\lambda))(\Delta\mathcal{F})]_- + \mathcal{J}_1(\lambda)(\mathcal{F}_0^c)\mathcal{J}_1(\lambda)(\Delta\mathcal{F}) \\ &+ \frac{\epsilon^2}{2} \left(\mathcal{J}_1(\lambda)^2(\Delta\mathcal{F}) + (\mathcal{J}_1(\lambda)(\Delta\mathcal{F}))^2 \right) + \epsilon \left(\mathcal{J}_1(\lambda)(\mathcal{F}_0^o)\mathcal{J}_1(\lambda)(\Delta\mathcal{F}) - \frac{3}{4}\partial_\lambda\mathcal{J}_2(\lambda)(\Delta\mathcal{F}) \right) \\ &+ \frac{1}{2}\mathcal{J}_1(\lambda)^2(\mathcal{F}_0^c) + \frac{\epsilon}{2}\mathcal{J}_1(\lambda)^2(\mathcal{F}_0^o) + \frac{1}{2}\mathcal{J}_1(\lambda)(\mathcal{F}_0^o)^2 - \frac{3}{4}\partial_\lambda\mathcal{J}_2(\lambda)(\mathcal{F}_0^o) + \frac{5}{8\lambda^2} = 0, \end{aligned} \quad (3.7)$$

where $\Delta\mathcal{F}$ is defined as

$$\Delta\mathcal{F} := \sum_{g \geq 1} \epsilon^{2g-2} \mathcal{F}_g^c + \sum_{p \geq 1} \epsilon^{p-1} \mathcal{F}_p^o. \quad (3.8)$$

Proof By using (1.25) and (3.1), we have

$$\frac{1}{\tau_{\text{top}}} \mathcal{L}(\lambda)(\tau_{\text{top}}) = 0. \quad (3.9)$$

By substituting (3.2) into the above (3.9), and then taking the coefficients of ϵ^{-2} , ϵ^{-1} and the remaining part, we get (3.5), (3.6) and (3.7) respectively. Hence the lemma is proved. \square

Now we are ready to prove Theorem 1.2.

Proof of Theorem 1.2 Let us denote

$$\mathcal{D}(\lambda) := -[(\mathcal{I}_1(\lambda)\mathcal{J}_1(\lambda) + \mathcal{I}_2(\lambda)\mathcal{J}_2(\lambda))]_- - \mathcal{J}_1(\lambda)(\mathcal{F}_0^c)\mathcal{J}_1(\lambda). \quad (3.10)$$

Then equation (3.7) can be written as follows:

$$\begin{aligned} \mathcal{D}(\lambda)(\Delta\mathcal{F}) &= \frac{\epsilon^2}{2} \left(\mathcal{J}_1(\lambda)^2(\Delta\mathcal{F}) + (\mathcal{J}_1(\lambda)(\Delta\mathcal{F}))^2 \right) + \epsilon \left(\mathcal{J}_1(\lambda)(\mathcal{F}_0^c)\mathcal{J}_1(\lambda)(\Delta\mathcal{F}) - \frac{3}{4}\partial_\lambda\mathcal{J}_2(\lambda)(\Delta\mathcal{F}) \right) \\ &\quad + \frac{1}{2}\mathcal{J}_1(\lambda)^2(\mathcal{F}_0^c) + \frac{\epsilon}{2}\mathcal{J}_1(\lambda)^2(\mathcal{F}_0^o) + \frac{1}{2}\mathcal{J}_1(\lambda)(\mathcal{F}_0^o)^2 - \frac{3}{4}\partial_\lambda\mathcal{J}_2(\lambda)(\mathcal{F}_0^o) + \frac{5}{8\lambda^2}. \end{aligned} \quad (3.11)$$

Recall that $v = \partial_x^2\mathcal{F}_0^c$ and $r = \partial_x\mathcal{F}_0^o$. It follows from the genus-zero part of equations (1.1)–(1.2) that for $k \geq 0$,

$$\frac{\partial v}{\partial t_k} = \frac{v^k}{k!}v_x, \quad \frac{\partial r}{\partial t_k} = \sum_{i=0}^{k-1} \frac{v^i r^{2k-2i+1}}{i!(2k-2i+1)!!}(v_x + rr_x) + \frac{v^k}{k!}r_x, \quad (3.12)$$

$$\frac{\partial v}{\partial s_k} = 0, \quad \frac{\partial r}{\partial s_k} = \frac{1}{k!} \left(v + \frac{r^2}{2} \right)^k (v_x + rr_x). \quad (3.13)$$

Then we have

$$\mathcal{J}_1(\lambda)(\partial_x\mathcal{F}_0^c) = \sum_{i \geq 0} \frac{(2i+1)!!}{2^{i+1}} \lambda^{-i-\frac{3}{2}} \frac{v^{i+1}}{(i+1)!} = A^{\frac{1}{2}} - \lambda^{-\frac{1}{2}}, \quad (3.14)$$

$$\mathcal{J}_1(\lambda)^2(\mathcal{F}_0^c) = \sum_{i,j \geq 0} \frac{(2i+1)!!(2j+1)!!}{2^{i+j+2}i!j!(i+j+1)} \lambda^{-i-j-3} v^{i+j+1} = \frac{1}{8}A^2 - \frac{1}{8}\lambda^{-2}, \quad (3.15)$$

$$\partial_\lambda J_2(\lambda)(\mathcal{F}_0^o) = - \sum_{i \geq 0} (i+2)\lambda^{-i-3} \left(v + \frac{1}{2}r^2 \right)^{i+1} = \lambda^{-2} - B^2, \quad (3.16)$$

$$\mathcal{J}_1(\lambda)(\mathcal{F}_0^o) = \sum_{i \geq 0} \sum_{j=0}^i \frac{(2i+1)!!}{2^{i+1}} \lambda^{-i-\frac{3}{2}} \frac{v^j r^{2i-2j+1}}{j!(2i-2j+1)!!} = \frac{r}{2}A^{\frac{1}{2}}B, \quad (3.17)$$

$$\mathcal{J}_1(\lambda)^2(\mathcal{F}_0^o) = \sum_{i \geq 0} \frac{(2i+1)!!}{2^{i+1}} \lambda^{-i-\frac{3}{2}} \frac{\partial}{\partial t_i} \mathcal{J}_1(\lambda)(\mathcal{F}_0^o) = \frac{A}{8r}\partial_x(r^2AB) + \frac{r}{6}\partial_x(B^3). \quad (3.18)$$

Here we recall that A and B are given by (1.28). It follows from (3.5) and (3.6) that

$$\mathcal{D}(\lambda)(v) = A, \quad \mathcal{D}(\lambda)(r) = \frac{1}{2}rAB. \quad (3.19)$$

Note that $[\mathcal{D}(\lambda), \partial_x] = A^{\frac{1}{2}}\mathcal{J}_1(\lambda)$. Then we obtain by induction that

$$\mathcal{D}(\lambda)(v_i) = \partial_x^i(A) + \sum_{j=1}^i \binom{i}{j} \partial_x^{j-1}(A^{\frac{1}{2}}) \partial_x^{i+1-j}(A^{\frac{1}{2}}), \quad (3.20)$$

$$\mathcal{D}(\lambda)(r_i) = \frac{1}{2}\partial_x^i(rAB) + \frac{1}{2} \sum_{j=1}^i \binom{i}{j} \partial_x^{j-1}(A^{\frac{1}{2}}) \partial_x^{i+1-j}(rA^{\frac{1}{2}}B). \quad (3.21)$$

Using (3.14)–(3.18), (3.20), (3.21) and (3.11), we arrive at (1.29). Here we note that equation (1.29) holds identically in λ .

It remains to show the uniqueness of the solution to equation (1.29). By using Theorem 1.1, we can write ΔF into the form

$$\Delta F =: \sum_{p \geq 0} \epsilon^p X_p(r, v_1, r_1, \dots, v_{2p+1}, r_{2p+1}). \quad (3.22)$$

Hence, by taking the coefficient of ϵ^p in (1.29), we have

$$\begin{aligned} & \sum_{i=0}^{2p+1} \frac{\partial X_p}{\partial v_i} \left(\partial^i(A) + \sum_{j=1}^i \binom{i}{j} \partial^{j-1}(A^{\frac{1}{2}}) \partial^{i+1-j}(A^{\frac{1}{2}}) \right) \\ & + \frac{1}{2} \sum_{i=0}^{2p+1} \frac{\partial X_p}{\partial r_i} \left(\partial^i(rAB) + \sum_{j=1}^i \binom{i}{j} \partial^{j-1}(A^{\frac{1}{2}}) \partial^{i+1-j}(rA^{\frac{1}{2}}B) \right) \\ & = \frac{1}{2} \sum_{i=0}^{2p-3} \left(\frac{\partial X_{p-2}}{\partial v_i} \partial^{i+2}\left(\frac{A^2}{8}\right) + \frac{\partial X_{p-2}}{\partial r_i} \partial^{i+1}\left(\frac{A}{8r} \partial(r^2AB) + \frac{r}{6} \partial(B^3)\right) \right) \\ & + \frac{1}{2} \sum_{i,j=0}^{2g-3} \left[\left(\frac{\partial^2 X_{p-2}}{\partial v_i \partial v_j} + \sum_{k=0}^{p-2} \frac{\partial X_k}{\partial v_i} \frac{\partial X_{p-k-2}}{\partial v_j} \right) \partial^{i+1}(A^{\frac{1}{2}}) \partial^{j+1}(A^{\frac{1}{2}}) \right. \\ & + \frac{1}{4} \left(\frac{\partial^2 X_{p-2}}{\partial r_i \partial r_j} + \sum_{k=0}^{p-2} \frac{\partial X_k}{\partial r_i} \frac{\partial X_{p-k-2}}{\partial r_j} \right) \partial^{i+1}(rA^{\frac{1}{2}}B) \partial^{j+1}(rA^{\frac{1}{2}}B) \\ & \left. + \left(\frac{\partial^2 X_{p-2}}{\partial v_i \partial r_j} + \sum_{k=2}^{p-2} \frac{\partial X_k}{\partial v_i} \frac{\partial X_{p-k-2}}{\partial r_j} \right) \partial^{i+1}(A^{\frac{1}{2}}) \partial^{j+1}(rA^{\frac{1}{2}}B) \right] \\ & + \sum_{i=0}^{2p-1} \left(\frac{1}{2} \frac{\partial X_{p-1}}{\partial v_i} r A^{\frac{1}{2}} B \partial^{i+1}(A^{\frac{1}{2}}) + \frac{\partial X_{p-1}}{\partial r_i} \left(\frac{1}{4} r A^{\frac{1}{2}} B \partial^{i+1}(rA^{\frac{1}{2}}B) + \frac{3}{4} \partial^{i+1}(B^2) \right) \right) \\ & + \delta_{p,1} \left(\frac{A}{16r} \partial(r^2AB) + \frac{r}{12} \partial(B^3) \right) + \delta_{p,0} \left(\frac{A^2}{16} + B^2 - \frac{1}{4} AB \right) \end{aligned} \quad (3.23)$$

By writing it into partial fraction with respect to λ , and then taking the coefficients of the powers of A and B , we arrive at a system of equations in the following form:

$$M_p \left(\frac{\partial X_p}{\partial v}, \frac{\partial X_p}{\partial r}, \dots, \frac{\partial X_p}{\partial v_{2p+1}}, \frac{\partial X_p}{\partial r_{2p+1}} \right)^T = C_p, \quad (3.24)$$

where M_p is an upper triangular matrix with diagonal elements being

$$1, \frac{1}{r}, \dots, \frac{(2i+1)!!}{2^i} v_1^i, \frac{(i+1)!}{r} (v_1 + rr_1)^i, \dots, \frac{(4p+3)!!}{2^{2p+1}} v_1^{2p+1}, \frac{(2p+2)!}{r} (v_1 + rr_1)^{2p+1}, \quad (3.25)$$

and C_p is a column vector determined by X_0, \dots, X_{p-1} . Then the gradient of X_p can be uniquely determined by the system of linear equations (3.24). The theorem is proved. \square

Using the loop equation, we obtain the expressions in Example 1.3 as well as

$$F_3^o = \frac{1}{v_1^4 (v_1 + rr_1)^6} \left(\frac{1}{8} r_2^2 v_1^8 + \frac{5}{24} r_1 r_3 v_1^8 + \frac{73 v_4 v_1^8}{1152} + \frac{1}{12} r r_4 v_1^8 - \frac{1}{8} r r_1 r_2^2 v_1^7 - \frac{9}{16} r_1^3 r_2 v_1^7 \right)$$

$$\begin{aligned}
& -\frac{7}{12}r_1v_2r_2v_1^7 - \frac{3}{16}r_1^2v_3v_1^7 - \frac{149}{640}v_2v_3v_1^7 - \frac{13}{48}rr_2v_3v_1^7 + \frac{35}{48}rr_1^2r_3v_1^7 - \frac{7}{24}rv_2r_3v_1^7 \\
& - \frac{1}{8}r^2r_2r_3v_1^7 + \frac{19}{64}rr_1v_4v_1^7 + \frac{23}{48}r^2r_1r_4v_1^7 + \frac{1}{48}r^2v_5v_1^7 + \frac{1}{48}r^3r_5v_1^7 + \frac{1}{8}r_1^6v_1^6 + \frac{61}{360}v_2^3v_1^6 \\
& - \frac{1}{4}r^3r_2^3v_1^6 + \frac{5}{12}r_1^2v_2^2v_1^6 - \frac{23}{8}r^2r_1^2r_2^2v_1^6 + \frac{1}{6}r^2v_2r_2^2v_1^6 - \frac{1}{8}r^2v_3^2v_1^6 - \frac{1}{8}r^4r_3^2v_1^6 + \frac{19}{48}r_1^4v_2v_1^6 \\
& - \frac{31}{16}rr_1^4r_2v_1^6 + \frac{7}{12}rv_2^2r_2v_1^6 - \frac{67}{48}rr_1^2v_2r_2v_1^6 - \frac{19}{24}rr_1^3v_3v_1^6 - \frac{307}{320}rr_1v_2v_3v_1^6 - \frac{29}{16}r^2r_1r_2v_3v_1^6 \\
& + \frac{1}{48}r^2r_1^3r_3v_1^6 - \frac{43}{24}r^2r_1v_2r_3v_1^6 - \frac{49}{24}r^3r_1r_2r_3v_1^6 - \frac{1}{4}r^3v_3r_3v_1^6 + \frac{133}{384}r^2r_1^2v_4v_1^6 - \frac{1}{6}r^2v_2v_4v_1^6 \\
& - \frac{1}{6}r^3r_2v_4v_1^6 + \frac{37}{48}r^3r_1^2r_4v_1^6 - \frac{1}{6}r^3v_2r_4v_1^6 - \frac{1}{6}r^4r_2r_4v_1^6 + \frac{1}{16}r^3r_1v_5v_1^6 + \frac{1}{16}r^4r_1r_5v_1^6 \\
& + \frac{1}{2}rr_1^7v_1^5 + \frac{3}{5}rr_1v_2^3v_1^5 + 2r^4r_1r_3^2v_1^5 + \frac{73}{48}rr_1^3v_2^2v_1^5 + \frac{3}{8}r^3r_1^3r_2^2v_1^5 + \frac{67}{12}r^3r_1v_2r_2^2v_1^5 - \frac{1}{4}r^3r_1v_3^2v_1^5 \\
& - \frac{1}{4}r^5r_1r_3^2v_1^5 + \frac{73}{48}rr_1^5v_2v_1^5 + \frac{13}{16}r^2r_1^5r_2v_1^5 + \frac{25}{6}r^2r_1v_2^2r_2v_1^5 + \frac{239}{48}r^2r_1^3v_2r_2v_1^5 - \frac{1}{4}r^2r_1^4v_3v_1^5 \\
& + \frac{3}{4}r^2v_2^2v_3v_1^5 + \frac{3}{4}r^4r_2^2v_3v_1^5 + \frac{33}{128}r^2r_1^2v_2v_3v_1^5 - \frac{21}{16}r^3r_1^2r_2v_3v_1^5 + \frac{3}{2}r^3v_2r_2v_3v_1^5 - \frac{9}{16}r^3r_1^4r_3v_1^5 \\
& + \frac{3}{4}r^3v_2^2r_3v_1^5 + \frac{3}{4}r^5r_2^2r_3v_1^5 - \frac{29}{24}r^3r_1^2v_2r_3v_1^5 - \frac{53}{24}r^4r_1^2r_2r_3v_1^5 + \frac{3}{2}r^4v_2r_2r_3v_1^5 - \frac{1}{2}r^4r_1v_3r_3v_1^5 \\
& + \frac{17}{288}r^3r_1^3v_4v_1^5 - \frac{1}{3}r^3r_1v_2v_4v_1^5 - \frac{1}{3}r^4r_1r_2v_4v_1^5 + \frac{7}{16}r^4r_1^3r_4v_1^5 - \frac{1}{3}r^4r_1v_2r_4v_1^5 - \frac{1}{3}r^5r_1r_2r_4v_1^5 \\
& + \frac{1}{16}r^4r_1^2v_5v_1^5 + \frac{1}{16}r^5r_1^2r_5v_1^5 - \frac{1}{8}r^2r_1^8v_1^4 - \frac{1}{2}r^2v_2^4v_1^4 - \frac{1}{2}r^6r_2^4v_1^4 - \frac{37}{24}r^2r_1^2v_2^3v_1^4 + \frac{1}{4}r^5r_1^2r_2^3v_1^4 \\
& - 2r^5v_2r_2^3v_1^4 - \frac{95}{48}r^2r_1^4v_2^2v_1^4 - 3r^4v_2^2r_2^2v_1^4 - \frac{7}{12}r^4r_1^2v_2r_2^2v_1^4 - \frac{1}{8}r^4r_1^2v_3^2v_1^4 - \frac{1}{8}r^6r_1^2r_3^2v_1^4 \\
& - \frac{43}{48}r^2r_1^6v_2v_1^4 + \frac{3}{16}r^3r_1^6r_2v_1^4 - 2r^3v_2^3r_2v_1^4 - \frac{29}{12}r^3r_1^2v_2^2r_2v_1^4 - \frac{3}{16}r^3r_1^4v_2r_2v_1^4 + \frac{3}{8}r^3r_1^5v_3v_1^4 \\
& + \frac{3}{4}r^3r_1v_2^2v_3v_1^4 + \frac{3}{4}r^5r_1r_2^2v_3v_1^4 + \frac{91}{96}r^3r_1^3v_2v_3v_1^4 + \frac{11}{48}r^4r_1^3r_2v_3v_1^4 + \frac{3}{2}r^4r_1v_2r_2v_3v_1^4 - \frac{1}{16}r^4r_1^5r_3v_1^4 \\
& + \frac{3}{4}r^4r_1v_2^2r_3v_1^4 + \frac{3}{4}r^6r_1r_2^2r_3v_1^4 + \frac{7}{24}r^4r_1^3v_2r_3v_1^4 - \frac{7}{24}r^5r_1^3r_2r_3v_1^4 + \frac{3}{2}r^5r_1v_2r_2r_3v_1^4 - \frac{1}{4}r^5r_1^2v_3r_3v_1^4 \\
& - \frac{19}{384}r^4r_1^4v_4v_1^4 - \frac{1}{6}r^4r_1^2v_2v_4v_1^4 - \frac{1}{6}r^5r_1^2r_2v_4v_1^4 + \frac{1}{16}r^5r_1^4r_4v_1^4 - \frac{1}{6}r^5r_1^2v_2r_4v_1^4 - \frac{1}{6}r^6r_1^2r_2r_4v_1^4 \\
& + \frac{1}{48}r^5r_1^3v_5v_1^4 + \frac{1}{48}r^6r_1^3r_5v_1^4 + \frac{1}{18}r^3r_1^3v_2^3v_1^3 - \frac{5}{48}r^3r_1^5v_2^2v_1^3 - \frac{1}{48}r^3r_1^7v_2v_1^3 + \frac{1}{48}r^4r_1^5v_2r_2v_1^3 \\
& + \frac{1}{48}r^4r_1^6v_3v_1^3 - \frac{7}{128}r^4r_1^4v_2v_3v_1^3 + \frac{1}{192}r^5r_1^5v_4v_1^3 + \frac{1}{24}r^4r_1^4v_2^3v_1^2 - \frac{1}{48}r^4r_1^6v_2^2v_1^2 - \frac{7}{320}r^5r_1^5v_2v_3v_1^2 \\
& + \frac{r^6r_1^6v_4v_1^2}{1152} + \frac{1}{60}r^5r_1^5v_2^3v_1 - \frac{7r^6r_1^6v_2v_3v_1}{1920} + \frac{1}{360}r^6r_1^6v_2^3 \\
& = -\frac{r^2u_2^4}{u_1^6} + \frac{ru_2^2(r_1u_2 + 3ru_3)}{4u_1^5} - \frac{r_1^2u_2^2 - 4u_2^3 - 2rr_2u_2^2 + 6rr_1u_2u_3 + 3r^2u_3^2 + 4r^2u_2u_4}{24u_1^4} \\
& + \frac{r_1^2(r_1^2u_2 - u_2^2)}{48v_1u_1^3} - \frac{r_1^2(r_1^4 - 2r_1u_2 + u_2^2 + rr_2u_2)}{48v_1^2u_1^2} - \frac{3r_1^3r_2 - r_1^2u_3}{48v_1u_1^2} - \frac{2r_1^3r_2 + 3r_1r_2u_2}{48v_1^2u_1} \\
& + \frac{2r_1r_2u_2 + r_2^2u_3 - 11u_2u_3 - 2rr_2u_3 - 3rr_3u_2 + 2rr_1u_4 + r^2u_5}{48u_1^3}
\end{aligned}$$

$$-\frac{2r_2^2 + r_1r_3 - 3u_4 - rr_4}{48u_1^2} - \frac{r_1r_3}{48v_1u_1} - \frac{r_2^2}{48v_1^2}.$$

We observe that the expressions of F_p^o in terms of r, u_k, r_k are much shorter than those in terms of r, v_k, r_k .

4 Applications

In this section, we give some applications of Theorems 1.1 and 1.2. Up to now, by solving the loop equation (3.23) and by using the expressions (1.19), we have obtained the expressions of the first two open free energies F_p^o (see Example 1.3). Next we will further study how to relate our results and observations with the Taylor coefficients in the power series \mathcal{F}_p^o .

Let us consider the change of the independent variables in $F_p^o, p \geq 1$ from the jet variables to the Itzykson-Zuber type variables, as the latter have more direct relations with the correlators. From the recursion relations (2.8)–(2.9), we obtain

$$\sum_{k \geq 0} kv_k \frac{\partial I_n}{\partial v_k} = \delta_{n,1} - I_n, \quad \sum_{k \geq 0} (k-1)v_k \frac{\partial I_n}{\partial v_k} = (n-1)I_n, \quad (4.1)$$

$$\sum_{k \geq 0} \left(kv_k \frac{\partial J_n}{\partial v_k} + kr_k \frac{\partial J_n}{\partial r_k} \right) = -J_n, \quad \sum_{k \geq 0} \left((k-1)v_k \frac{\partial J_n}{\partial v_k} + \left(k - \frac{1}{2} \right) r_k \frac{\partial J_n}{\partial r_k} \right) = \left(n - \frac{1}{2} \right) J_n, \quad (4.2)$$

and by using the formula

$$\frac{\partial}{\partial x} = \frac{1}{1-I_1} \frac{\partial}{\partial I_0} + \sum_{k \geq 1} \frac{I_{k+1}}{1-I_1} \frac{\partial}{\partial I_k} + \sum_{k \geq 0} \frac{J_{k+1} + \frac{J_1}{1-I_1} I_{k+1}}{1-I_1 - J_0 J_1} \frac{\partial}{\partial J_k}, \quad (4.3)$$

we have the form:

$$v_k = \sum_{\lambda \in \mathcal{P}_{k-1}} C'_{k;\lambda} \frac{I_{\lambda+1}}{U_1^{k+\ell(\lambda)}}, \quad (4.4)$$

$$r_k = \sum_{d=0}^{k+1} \sum_{\substack{\lambda, \mu \in \mathcal{P} \\ |\lambda| + |\mu| = k - \frac{-\ell(\mu) - d + 1}{2}}} C''_{k;d;\lambda;\mu}(U_1^{-1}, U_2^{-1}) J_0^d I_{\lambda+1} J_\mu, \quad (4.5)$$

where $C'_{k;\lambda}$ are certain rational numbers, and $C''_{k;d;\lambda;\mu}(U_1^{-1}, U_2^{-1})$ are homogeneous polynomials in U_1^{-1} and U_2^{-1} of degree $m + n + k$ with rational coefficients. Then by using Theorem 1.1, we have that

$$F_1^o = -\frac{1}{2} \log(1 - I_1 - J_0 J_1), \quad (4.6)$$

and the function $F_p^o, p \geq 2$ can be expressed as rational functions in $J_0, I_1, J_1, \dots, I_{2p-1}, J_{2p-1}$ satisfying the conditions

$$-\sum_{k \geq 1} \left((I_k - \delta_{k,1}) \frac{\partial F_p^o}{\partial I_k} + J_k \frac{\partial F_p^o}{\partial J_k} \right) = (p-1)F_p^o, \quad (4.7)$$

$$\sum_{k \geq 1} \left((k-1)I_k \frac{\partial F_p^o}{\partial I_k} + \left(k - \frac{1}{2} \right) J_k \frac{\partial F_p^o}{\partial J_k} \right) = \frac{3}{2}(p-1)F_p^o. \quad (4.8)$$

For example,

$$\begin{aligned} F_2^o &= \frac{1}{(1-I_1)^2 (1-I_1-J_0J_1)^3} \left(\frac{1}{24} (3J_3J_0^2 + 3I_3J_0 + 4J_2) (I_1-1)^3 \right. \\ &\quad + \frac{1}{24} (-5J_2^2J_0^3 + 3J_1J_3J_0^3 + 3I_3J_1J_0^2 - 10I_2J_2J_0^2 - 5I_2^2J_0 - 6J_1J_2J_0 - 12I_2J_1) (I_1-1)^2 \\ &\quad \left. + \left(\frac{5}{24}J_1^3 - \frac{1}{8}J_0J_1^2I_2 \right) (I_1-1) - \frac{1}{24}J_0^2J_1^3I_2 \right). \end{aligned} \quad (4.9)$$

In general, F_p^o , $p \geq 2$ can be written into the form

$$F_p^o = \frac{\sum_{m,n \geq 0, m+n \leq 13p-7} \sum_{2 \leq k_1, \dots, k_m \leq 2p-1, 1 \leq \ell_1, \dots, \ell_n \leq 2p-1} \frac{B_{k_1, \dots, k_m; \ell_1, \dots, \ell_n}}{m!n!} \prod_{i=1}^m \frac{J_0^{2k_i-2} I_{k_i}}{1-I_1} \prod_{i=1}^n \frac{J_0^{2\ell_i-1} J_{\ell_i}}{1-I_1}}{J_0^{3p-3} (1-I_1)^{p-1} (1-\frac{J_0J_1}{1-I_1})^{7p-4}}. \quad (4.10)$$

Similar to [16], we consider the restriction of $\mathcal{F}_p^o(\mathbf{t}, \mathbf{s})$, $p \geq 1$ at

$$t_0 = t_1 = 0, \quad \sum_{k \geq 2} \frac{t_k}{(2k+1)!} + \sum_{k \geq 0} \frac{s_k}{2^k k!} = 1. \quad (4.11)$$

According to the formulae (2.6), (2.7) and Lemma 2.2, this is equivalent to consider the restriction of F_p^o , $p \geq 1$ at $I_0 = I_1 = 0$, $J_0 = 1$ as follows:

$$F_1^{o,*} := F_1^o|_{I_1=0, J_0=1} = -\frac{1}{2} \log(1-J_1), \quad (4.12)$$

$$F_p^{o,*} := F_p^o|_{I_1=0, J_0=1} \in (1-J_1)^{-(7p-4)} \mathbb{C}[I_2, \dots, I_{2p-1}, J_1, \dots, J_{2p-1}], \quad p \geq 2. \quad (4.13)$$

(Here since F_p^o , $p \geq 1$ are independent of I_0 , we do not write $I_0 = 0$ in the above formulae.) The expression of $F_p^{o,*}$ is simpler than F_p^o . For $p \geq 2$, we can recover all the information of F_p^o from $F_p^{o,*}$ by

$$F_p^o = \frac{1}{J_0^{3p-3} (1-I_1)^{p-1}} F_p^{o,*}|_{I_k \rightarrow (1-I_1)^{-1} J_0^{2k-2} I_k, J_k \rightarrow (1-I_1)^{-1} J_0^{2k-1} J_k}. \quad (4.14)$$

In other words, the restriction of $\mathcal{F}_p^o(\mathbf{t}, \mathbf{s})$, $p \geq 2$ at (4.11) has no loss of information. Below we will use the expressions of $F_p^{o,*}$ to give relations among the correlators. Indeed, we have, by using Lemma 2.2, that

$$F_p^{o,*} = \mathcal{F}_p^o(t_0^*, s_0^*, t_1^*, s_1^*, \dots), \quad p \geq 2 \quad (4.15)$$

where

$$t_0^* = 0, \quad s_0^* = 1 + \sum_{n \geq 1} \frac{(-1)^n}{2^n n!} \left(J_n - \frac{1}{2n+1} I_{n+1} \right), \quad (4.16)$$

$$t_k^* = (1 - \delta_{k,1}) I_k, \quad s_k^* = \sum_{n \geq 0} \frac{(-1)^n}{2^n n!} \left(J_{k+n} - \frac{1}{2n+1} I_{n+k+1} \right), \quad k \geq 1. \quad (4.17)$$

It is obvious from (4.10) that $F_p^{o,*}$ has the form

$$F_p^{o,*} = \frac{1}{(1 - J_1)^{7p-4}} \sum_{\substack{m,n \geq 0, \\ m+n \leq 13p-7}} \frac{B_{k_1, \dots, k_m; \ell_1, \dots, \ell_n}}{m!n!} I_{k_1} \cdots I_{k_m} J_{\ell_1} \cdots J_{\ell_n}. \quad (4.18)$$

For example,

$$\begin{aligned} F_2^{o,*} = & \frac{1}{24}(-I_2 - 5) - \frac{5}{8(J_1 - 1)} + \frac{-3I_2 + I_3 - 2J_2 + J_3 - 5}{8(J_1 - 1)^2} \\ & - \frac{5(I_2^2 + 2J_2I_2 + 2I_2 + J_2^2 + 2J_2 + 1)}{24(J_1 - 1)^3}. \end{aligned} \quad (4.19)$$

Proof of Corollary 1.4 By using Lemma 2.2, we know that

$$t_0|_{I_0=I_1=0, J_0=1} = 0, \quad t_k|_{I_0=I_1=0, J_0=1} = (1 - \delta_{k,1})I_k, \quad (4.20)$$

$$s_0|_{I_0=I_1=0, J_0=1} = 1 + \sum_{n \geq 1} \frac{(-1)^n}{2^n n!} \left(J_n - \frac{1}{2n+1} I_{n+1} \right), \quad (4.21)$$

$$s_k|_{I_0=I_1=0, J_0=1} = \sum_{n \geq 0} \frac{(-1)^n}{2^n n!} \left(J_{k+n} - \frac{1}{2n+1} I_{n+k+1} \right), \quad (4.22)$$

where $k \geq 1$. Hence by taking the derivatives of (4.15), we obtain

$$\frac{\partial^{m+n} F_p^{o,*}}{\partial I_{k_1} \cdots \partial I_{k_m} \partial J_{\ell_1} \cdots \partial J_{\ell_n}} \Big|_{I_{\geq 2}=J_{\geq 1}=0} = \Lambda_{k_1, \dots, k_m; \ell_1, \dots, \ell_n}(\mathcal{F}_p)|_{s_0=1, t_{\geq 0}=s_{\geq 1}=0}, \quad (4.23)$$

where $k_1, \dots, k_m \geq 2$ and $\ell_1, \dots, \ell_n \geq 1$ and the operator $\Lambda_{k_1, \dots, k_m; \ell_1, \dots, \ell_n}$ is defined by (1.33). Then the corollary is proved. \square

The property (2.48) implies that

$$\langle \tau_{k_1} \cdots \tau_{k_m} \sigma_{\ell_1} \cdots \sigma_{\ell_n} \rangle_p^o \neq 0, \quad (4.24)$$

only if

$$k_1 + \cdots + k_m + \ell_1 + \cdots + \ell_n = \frac{3p-3}{2} + m + \frac{n}{2}. \quad (4.25)$$

It follows that

$$\frac{\partial^{m+n} \mathcal{F}_p^o}{\partial t_{k_1} \cdots \partial t_{k_m} \partial s_{\ell_1} \cdots \partial s_{\ell_n}} \Big|_{s_0=1, t_{\geq 0}=s_{\geq 1}=0} = \frac{\langle \sigma_0^{3-3p} \prod_{i=1}^m (\sigma_0^{2k_i-2} \tau_{k_i}) \prod_{j=1}^n (\sigma_0^{2\ell_j-1} \sigma_{\ell_j}) \rangle_p^o}{(2 \sum_{i=1}^m k_m + 2 \sum_{j=1}^n \ell_j - 3(p-1) - 2m - n)!} \quad (4.26)$$

if

$$k_1 + \cdots + k_m + \ell_1 + \cdots + \ell_n \geq \frac{3}{2}(p-1) + m + \frac{n}{2}, \quad (4.27)$$

and vanishes otherwise. Hence the left-hand side of (1.32) is just a linear combination of the correlators.

Example 4.1 It follows from Corollary 1.4 together with the formulae (4.23) and (4.26) that when n is sufficiently large, we have

$$\begin{aligned} B_{\emptyset;1^n}^{[p]} &= (7p-4)! \sum_{m=0}^n \sum_{k=3p-3}^{n-m} \binom{n}{m} \binom{n-m}{k} \frac{(-1)^{n-k}}{2^{n-m-k}(7p-4-m)!} \frac{\langle \sigma_0^{k-3p+3} \sigma_1^k \rangle_2^o}{(2k-n+m-3p+3)!} = 0, \\ B_{2;1^n}^{[p]} &= (7p-4)! \sum_{m=0}^n \sum_{k=3p-3}^{n-m} \binom{n}{m} \binom{n-m}{k} \frac{(-1)^{n-k}}{2^{n-m-k}(7p-4-m)!} \\ &\quad \times \left(\frac{\langle \tau_2 \sigma_0^{k-3p+5} \sigma_1^k \rangle_p^o}{(2k-n+m-3p+6)!} - \frac{\langle \sigma_0^{k-3p+4} \sigma_1^{k+1} \rangle_p^o}{(2k-n+m-3p+4)!} + \frac{1}{6} \frac{\langle \sigma_0^{k-3p+3} \sigma_1^k \rangle_p^o}{(2k-n+m-3p+2)!} \right) = 0. \end{aligned}$$

We also have a conjectural form for F_p^o , $p \geq 2$:

$$F_p^o = \frac{\sum_{\substack{m,n \geq 0 \\ 0 \leq m+n \leq 6p-6}} \sum_{\substack{2 \leq k_1, \dots, k_m \leq 2p-1 \\ 1 \leq \ell_1, \dots, \ell_n \leq 2p-1}} \frac{\tilde{B}_{k_1, \dots, k_m; \ell_1, \dots, \ell_n}^{[p]}}{m! n!} \prod_{i=1}^m \frac{J_0^{2k_i-2} I_{k_i}}{1-I_1} \prod_{i=1}^n \frac{J_0^{2\ell_i-1} J_{\ell_i}}{1-I_1}}{J_0^{3p-3} (1-I_1)^{p-1} (1 - \frac{J_0 J_1}{1-I_1})^{3p-3}}. \quad (4.28)$$

It would be interesting if the two conjectural forms (2.50) and (4.28) could be proved to be equivalent using the transformation between v_k, r_k and I_k, J_k . Under the restriction $I_1 = 0, J_0 = 1$, the conjectural formula (4.28) becomes

$$F_p^{o,*} = \frac{1}{(1-J_1)^{3p-3}} \sum_{0 \leq m+n \leq 6p-6} \tilde{B}_{k_1, \dots, k_m; \ell_1, \dots, \ell_n}^{[p]} I_{k_1} \cdots I_{k_m} J_{\ell_1} \cdots J_{\ell_n}, \quad (4.29)$$

which is equivalent to (1.34).

Example 4.2 For $p = 2$, we have the explicit expression (4.19) of $F_2^{o,*}$, which implies that the conjectural relations (1.34) hold true for $p = 2$. For examples,

$$\tilde{B}_{\emptyset;1^n}^{[2]} = 6 \sum_{m=0}^3 \sum_{k=3}^{n-m} \binom{n}{m} \binom{n-m}{k} \frac{(-1)^{n-k}}{2^{n-m-k}(3-m)!} \frac{\langle \sigma_0^{k-3} \sigma_1^k \rangle_2^o}{(2k-n+m-3)!} = 0$$

for $n \geq 7$, and

$$\begin{aligned} \tilde{B}_{2;1^n}^{[2]} &= 6 \sum_{m=0}^3 \sum_{k=3}^{n-m} \binom{n}{m} \binom{n-m}{k} \frac{(-1)^{n-k}}{2^{n-m-k}(3-m)!} \\ &\quad \times \left(\frac{\langle \tau_2 \sigma_0^{k-1} \sigma_1^k \rangle_2^o}{(2k-n+m-1)!} - \frac{\langle \sigma_0^{k-2} \sigma_1^{k+1} \rangle_2^o}{(2k-n+m-2)!} + \frac{1}{6} \frac{\langle \sigma_0^{k-3} \sigma_1^k \rangle_2^o}{(2k-n+m-4)!} \right) = 0 \end{aligned}$$

for $n \geq 6$. But in fact, from the expression (4.19) we obtain more vanishing relations, namely, the above two identities hold for $n \geq 4$.

Since the explicit expression (4.19) of $F_2^{o,*}$ is obtained, we have even more relations:

$$\tilde{B}_{2,2;\emptyset}^{[2]} = -2 \langle \tau_2 \sigma_1 \rangle_2^o + \langle \tau_2^2 \sigma_0 \rangle_2^o = \frac{5}{12},$$

$$\begin{aligned}
\tilde{B}_{\emptyset;1,2}^{[2]} &= -3\langle \sigma_2 \rangle_2^o + \langle \sigma_0 \sigma_1 \sigma_2 \rangle_2^o = \frac{1}{4}, \\
\tilde{B}_{3;\emptyset}^{[2]} &= -\langle \sigma_2 \rangle_2^o + \langle \tau_3 \sigma_0 \rangle_2^o = \frac{1}{8}, \\
\tilde{B}_{\emptyset;3}^{[2]} &= -\frac{1}{2}\langle \sigma_2 \rangle_2^o + \frac{1}{2}\langle \sigma_0^2 \sigma_3 \rangle_2^o = \frac{1}{8}, \\
\tilde{B}_{2;1,1}^{[2]} &= \langle \tau_2 \sigma_0 \sigma_1^2 \rangle_2^o - \langle \sigma_1^3 \rangle_2^o - 6\langle \tau_2 \tau_1 \rangle_2^o = -\frac{1}{4}, \\
\tilde{B}_{\emptyset;2,2}^{[2]} &= -\langle \sigma_0 \sigma_1 \sigma_2 \rangle_2^o + \frac{1}{6}\langle \sigma_0^3 \sigma_2^2 \rangle_2^o = \frac{5}{12}, \\
\tilde{B}_{2;1,1,1}^{[2]} &= 9\langle \sigma_1^3 \rangle_2^o - \langle \sigma_0 \sigma_1^4 \rangle_2^o + 18\langle \tau_2 \sigma_1 \rangle_2^o - \frac{21}{2}\langle \tau_2 \sigma_0 \sigma_1^2 \rangle_2^o + \frac{1}{2}\langle \tau_2 \sigma_0^2 \sigma_1^3 \rangle_2^o = \frac{1}{4}, \\
\tilde{B}_{2;2}^{[2]} &= -\langle \sigma_0 \sigma_1 \sigma_2 \rangle_2^o - \frac{1}{2}\langle \tau_2 \sigma_1 \rangle_2^o + \frac{1}{2}\langle \tau_2 \sigma_0^2 \sigma_2 \rangle_2^o = \frac{5}{12}, \\
\tilde{B}_{3;1}^{[2]} &= 3\langle \sigma_2 \rangle_2^o - \langle \sigma_0 \sigma_1 \sigma_2 \rangle_2^o - \frac{7}{2}\langle \tau_3 \sigma_0 \rangle_2^o + \frac{1}{2}\langle \tau_3 \sigma_0^2 \sigma_1 \rangle_2^o = -\frac{1}{8}, \\
\tilde{B}_{\emptyset;1,3}^{[2]} &= \frac{3}{2}\langle \sigma_2 \rangle_2^o - \frac{1}{2}\langle \sigma_0 \sigma_1 \sigma_2 \rangle_2^o - 2\langle \sigma_0^2 \sigma_3 \rangle_2^o + \frac{1}{6}\langle \sigma_0^3 \sigma_1 \sigma_3 \rangle_2^o = -\frac{1}{8}.
\end{aligned}$$

5 Conclusions

The genus zero part of the topological tau-function for the Burgers–KdV hierarchy evaluated at $t_1 = t_2 = \dots = 0, s_1 = s_2 = \dots = 0$ gives rise to

$$F = \frac{1}{6}(v^1)^3, \quad F^o = v^1 v^2 + \frac{1}{6}(v^2)^3, \quad (5.1)$$

which satisfy the open WDVV equations [19, 23]. In general, starting with an n -dimensional Frobenius manifold, one can impose the open WDVV equations [19] to obtain an $(n+1)$ -dimensional F -manifold (due to an observation from Alcolado¹ and P. Rossi). A reconstruction of higher genus theory was given in [2] for flat F -manifolds. It is also possible to reconstruct higher genus parts by using the Dubrovin–Zhang approach together with the results in [3] similarly to what we do in this paper. We plan to do general cases including extended r -spin [10, 11] and the open analogue of a semisimple Frobenius manifold in future publications.

References

- [1] A. Alcolado. Extended Frobenius manifolds and the open WDVV equations. PhD Thesis. McGill University (Canada), 2017.
- [2] A. Arsie, A. Buryak, P. Lorenzoni, P. Rossi, Semisimple flat F -manifolds in higher genus. Comm. Math. Phys. **397** (2023), 141–197.
- [3] A. Basalaev, A. Buryak, Open WDVV equations and Virasoro constraints. Arnold Math. J. **5** (2019), 145–186.
- [4] M. Bertola, B. Dubrovin, D. Yang, Correlation functions of the KdV hierarchy and applications to intersection numbers over $\overline{\mathcal{M}}_{g,n}$. Phys. D **327** (2016), 30–57.

¹One of the authors D.Y. would like to thank Xujia Chen who kindly pointed out the interesting reference [1].

- [5] M. Bertola, G. Ruzza, The Kontsevich-Penner matrix integral, isomonodromic tau functions and open intersection numbers. *Ann. Henri Poincaré* **20** (2019), 393–443.
- [6] M. Bertola, D. Yang, The partition function of the extended r -reduced Kadomtsev-Petviashvili hierarchy. *J. Phys. A* **48** (2015), Paper No. 195205, 20 pp.
- [7] G. Borot, V. Bouchard, N. Chidambaram, T. Creutzig, D. Noshchenko, Higher Airy structures, \mathcal{W} -algebras and topological recursion. *Mem. Amer. Math. Soc.* **296** (2024), No. **1476**, v+108 pp.
- [8] A. Buryak, Equivalence of the open KdV and the open Virasoro equations for the moduli space of Riemann surfaces with boundary. *Lett. Math. Phys.* **105** (2015), 1427–1448.
- [9] A. Buryak, Open intersection numbers and the wave function of the KdV hierarchy. *Mosc. Math. J.* **16** (2016), 27–44.
- [10] A. Buryak, E. Clader, R. J. Tessler, Closed extended r -spin theory and the Gelfand-Dickey wave function. *J. Geom. Phys.* **137** (2019), 132–153.
- [11] A. Buryak, E. Clader, R. J. Tessler, Open r -spin theory III: A prediction for higher genus. *J. Geom. Phys.* **192** (2023), 12 pp.
- [12] A. Buryak, R. J. Tessler, R. J. Tessler, Matrix models and a proof of the open analog of Witten’s conjecture. *Comm. Math. Phys.* **353** (2017), 1299–1328.
- [13] P. Deligne, D. Mumford, The irreducibility of the space of curves of given genus. *Publ. Math. I.H.E.S.* **36** (1969), 75–109.
- [14] R. Dijkgraaf, E. Witten, Developments in topological gravity. *Internat. J. Modern Phys. A* **33** (2018), 63 pp.
- [15] B. Dubrovin, Geometry of 2D topological field theories. In: Francaviglia, M., Greco, S. (eds.) “Integrable Systems and Quantum Groups” (Montecatini Terme, 1993), Lecture Notes in Math., Vol. **1620**, Springer, Berlin, 1996, pp. 120–348.
- [16] B. Dubrovin, D. Yang, Remarks on intersection numbers and integrable hierarchies. I. Quasi-triviality. *Adv. Theor. Math. Phys.* **24** (2020), 1055–1085.
- [17] B. Dubrovin, Y. Zhang, Normal forms of hierarchies of integrable PDEs, Frobenius manifolds and Gromov–Witten invariants. arXiv:math/0108160.
- [18] T. Eguchi, Y. Yamada, S.-K. Yang, On the genus expansion in the topological string theory. *Rev. Maht. Phys.* **7** (1995), 279–309
- [19] A. Horev, J. P. Solomon, The open Gromov–Witten–Welschinger theory of blow-ups of the projective plane. arXiv:12104043.
- [20] C. Itzykson, J.B. Zuber, Combinatorics of the modular group II the Kontsevich integrals. *Internat. J. Modern Phys. A*, **23** (1992), 5661–5705.
- [21] M. Kontsevich, Intersection theory on the moduli space of curves and the matrix Airy function. *Comm. Math. Phys.* **147** (1992), 1–23.

- [22] S.-Q. Liu, D. Yang, Y. Zhang, C. Zhou, The Hodge-FVH correspondence. *J. Reine Angew. Math.* **775** (2021), 259–300.
- [23] R. Pandharipande, J. P. Solomon, R. J. Tessler, Intersection theory on moduli of disks, open KdV and Virasoro. arXiv:1409.2191.
- [24] K. Okuyama, K. Sakai: Genus expansion of open free energy in 2d topological gravity. *J. High Energy Phys.* **2021** (2021), Paper No. 217, 21 pp.
- [25] E. Witten, Two-dimensional gravity and intersection theory on moduli space. In "Surveys in differential geometry (Cambridge, MA, 1990)", pp. 243–310. Lehigh Univ., Bethlehem, PA, 1991.
- [26] D. Yang, C. Zhou, On an extension of the generalized BGW tau-function. *Lett. Math. Phys.* **111** (2021), 23 pp.
- [27] J. Zhou, On topological 1D gravity. I. arXiv:1412.1604.

Di Yang, School of Mathematical Sciences, University of Science and Technology of China,
Hefei 230026, P.R. China
e-mail: diyang@ustc.edu.cn

Chunhui Zhou, Institute of Geometry and Physics, University of Science and Technology of China,
Hefei 230026, P.R. China
e-mail: zhouch@ustc.edu.cn