Geometric, Variational, and Bracket Descriptions of Fluid Motion with Open Boundaries

Christopher Eldred celdred@sandia.gov Computer Science Research Institute Sandia National Laboratories

Francois Gay-Balmaz francois.gb@ntu.edu.sg Division of Mathematical Sciences Nanyang Technological University

Meng Wu

meng.wu@lmd.ipsl.fr Laboratoire de Météorologie Dynamique Ecole Normale Supérieure de Paris & Division of Mathematical Sciences Nanyang Technological University

Abstract

We develop a Lie group geometric framework for the motion of fluids with permeable boundaries that extends Arnold's geometric description of fluid in closed domains. Our setting is based on the classical Hamilton principle applied to fluid trajectories, appropriately amended to incorporate bulk and boundary forces via a Lagrange-d'Alembert approach, and to take into account only the fluid particles present in the fluid domain at a given time. By applying a reduction by relabelling symmetries, we deduce a variational formulation in the Eulerian description that extends the Euler-Poincar´e framework to open fluids. On the Hamiltonian side, our approach yields a bracket formulation that consistently extends the Lie-Poisson bracket of fluid dynamics and contains as a particular case a bulk+boundary bracket formulation proposed earlier. We illustrate the geometric framework with several examples, and also consider the extension of this setting to include multicomponent fluids, the consideration of general advected quantities, the analysis of higher-order fluids, and the incorporation of boundary stresses. Open fluids are found in a wide range of physical systems, including geophysical fluid dynamics, porous media, incompressible flow and magnetohydrodynamics. This new formulation permits a description based on geometric mechanics that can be applied to a broad class of models.

Contents

1 Introduction

Geometric mechanics formulations (ex. variational [\[1](#page-47-0), [2](#page-47-1), [3\]](#page-47-2), Hamiltonian, [\[4,](#page-47-3) [5,](#page-47-4) [6](#page-47-5), [7,](#page-47-6) [8\]](#page-47-7), metriplectic/GENERIC [\[9,](#page-47-8) [10,](#page-47-9) [11](#page-47-10), [12](#page-47-11), [13](#page-47-12), [14\]](#page-47-13)) have proven to be a powerful tool in the development of physically based models for fluid systems, with deep connections to fundamental features such as conservation laws [\[15](#page-47-14), [16,](#page-47-15) [17](#page-48-0), [18\]](#page-48-1) and involution constraints. However, existing formulations are largely restricted to the case of fixed^{[1](#page-1-1)}, material^{[2](#page-1-2)} and adiabatic^{[3](#page-1-3)} boundaries. Real physical systems often have open boundaries where mass, momentum, energy, entropy and other quantities are exchanged with an environment; this exchange can extend even to the bulk of the fluid. For example, in geophysical fluid dynamics the atmosphere has surface exchanges of moisture and heat with the ocean or land surface through precipitation and evaporation, radiative interactions in the bulk, surfaces stresses due to wave-atmosphere interaction and land-atmosphere interaction; with similar processes occuring in the ocean. Plasma systems [\[20](#page-48-2), [21](#page-48-3)] might have external magnetic fields such as those found in tokamaks and stellarators, and leakage or injection of charged material across (computational) boundaries. The general setting we are interested in is a fluid in a fixed^{[4](#page-1-4)} domain Ω , but which can escape the boundary. The particular nature of the boundary exchange is left open, and might include no-flux or material, open and/or prescribed inflow or outflow boundary conditions.

Existing geometric mechanics formulations are incapable of directly handling such situations, especially the general case of arbitrary boundary conditions, both Lagrangian and Eulerian

¹No exchange of momentum, since there is no movement of the boundary, although see [\[19\]](#page-48-4).

²No exchange of mass, also known as no-flux.

³No exchange of heat, and therefore energy.

⁴ In this work we will restrict ourselves to the case of fixed boundaries, with an extension to moving boundaries [\[19\]](#page-48-4) the subject of future work.

coordinates, and a full treatment of both variational and bracket-based approaches. Some work has been done in the context of Eulerian bracket-based formulations using GENERIC [\[22](#page-48-5), [23](#page-48-6)], but it is limited to open boundary conditions, only certain fluid models, does not treat the variational case and does not handle prescribed boundary conditions (such as those needed for inflow). There is also significant work in this area that has occurred in the port-Hamiltonian literature [\[24](#page-48-7), [25,](#page-48-8) [26,](#page-48-9) [27](#page-48-10), [28](#page-48-11), [29,](#page-48-12) [30,](#page-49-0) [31](#page-49-1)], mainly in the context of single component incompressible and compressible fluids. This work is based on Stokes-Dirac structures using Eulerian coordinates, and is disconnected from first principles such as Hamilton's or Lagranged'Alembert. Related to this, the variational case is not treated and it is limited to certain fluid models.

This paper addresses these concerns by developing a geometric theory of fluid dynamics in open domains, applicable to both Lagrangian and Eulerian coordinates, starting from first principles and closely following the existing theory for closed domains. Furthermore, it is shown that several important extensions can be naturally incorporated within this geometric framework, such as the treatment of multicomponent fluids, general advected quantities, higher-order fluids, and boundary stresses. This is done through an application of the classical Hamilton's principle applied to Lagrangian fluid trajectories, appropriately amended to incorporate bulk and boundary exchanges via a Lagrange-d'Alembert approach, and to take into account only the fluid particles present in the fluid domain. By applying a reduction by relabelling symmetries, we deduce a variational formulation in the Eulerian description that extends the Euler-Poincaré framework to open fluids. On the Hamiltonian side, our approach yields a bracket formulation that consistently extends the Lie-Poisson bracket of fluid dynamics and contains as a particular case the GENERIC-based bulk+boundary bracket formulation proposed earlier [\[22,](#page-48-5) [23](#page-48-6)]. This new theory is applicable to a wide range of fluid models with arbitrary advected quantities, and can handle arbitrary boundary conditions.

Beyond providing a unifying geometric framework, our approach offers significant insights into the treatment of boundary conditions for open fluids, which we will elaborate on later. Specifically, it identifies non-trivial terms that arise when the relationship between (absolute) fluid momentum $\frac{\partial l}{\partial u}$ and the fluid velocity u is no longer the usual $\frac{\partial l}{\partial u} = \rho u$. Furthermore, this approach sheds light on the handling of fluids with tensorial advected quantities and higherorder fluids, where additional nontrivial contributions to the boundary conditions emerge in the context of open fluid systems.

The two diagrams below provide a schematic overview of our geometric framework, highlighting how the setting for open fluids naturally extends the existing framework for closed fluids, from both the Lagrangian and Hamiltonian perspectives, in both the material and Eulerian descriptions. All the notations used will be explained in details later.

(i) Overview of the case of closed fluids:

Hamilton's principle	$L(\varphi, \dot{\varphi}, \varrho_0, S_0)$	Legendre trans.	$H(\varphi, p, \varrho_0, S_0)$	canonical bracket
$(\S1.1)$	$= \int_{\Omega} \mathfrak{L}(\varphi, \dot{\varphi}, \nabla \varphi, \varrho_0, S_0) dX$	$= \int_{\Omega} \mathfrak{H}(\varphi, p, \nabla \varphi, \varrho_0, S_0) dX$	$\dot{F} = \{F, H\}_{\text{can}}$	
Lagrangian-to-Eulerian	Lagrangian-to-Eulerian			
Euler-Poincaré principle	$\ell(u, \rho, s)$	Legendre trans.	$h(m, \rho, s)$	Lie-Poisson bracket
$(\S1.1)$	$= \int_{\Omega} I(u, \rho, s) dx$	$= \int_{\Omega} \mathfrak{h}(m, \rho, s) dx$	$\dot{f} = \{f, h\}_{\text{LP}}$	

Figure 1.1: Closed fluids: On the left side of the diagram, one starts with a formulation based on a Lagrangian $L : T \text{Diff}(\Omega) \to \mathbb{R} = L(\varphi, \dot{\varphi}, \varrho_0, S_0)$ using the classical Hamilton principle in material (configuration-velocity, unfortunately also known as Lagrangian) variables, from which the variational principle in *Eulerian* variables is derived through the use of *relabelling symmetries* (*Euler-*Poincaré reduction, left arrow pointing down, also known as the Lagrangian-to-Eulerian map). Alternatively, one starts on the right side of the diagram with a formulation based on a Hamiltonian $H: T^*$ Diff $(\Omega) \to \mathbb{R} = H(\varphi, p, \varrho_0, S_0)$ in material (configuration-momentum) variables. These dynamics are governed by the canonical symplectic form, with the associated canonical Poisson structure $\{\cdot,\cdot\}_{\text{can}}$. The *(non-canonical) Lie-Poisson structure* $\{f,h\}_{\text{LP}}$ in the *Eulerian* formulation is then derived through relabelling symmetries (Lie-Poisson reduction, right arrow pointing down, again also known as the Lagrangian-to-Eulerian map). Under the appropriate regularity conditions, the Lagrangian and Hamiltonian approaches can be related through the Legendre transform, at the level of either material (top arrow) or Eulerian (bottom arrow) variables. For example, if the Lagrangian L comes from a regular Lagrangian density \mathfrak{L} , we can define the associated Hamiltonian function H.

(ii) Overview of the case of open fluids:

Figure 1.2: Open fluid: On the left side of the diagram, one again starts with a formulation based on a Lagrangian $L : T$ Diff $(\Omega) \to \mathbb{R} = L(\varphi, \dot{\varphi}, \varrho_0, S_0)$ using now the Lagrange-d'Alembert principle in material variables, incorporating bulk and boundary sources of momentum, mass, and entropy. From this, the corresponding variational principle in Eulerian variables is derived through relabelling symmetries (Euler-Poincaré-d'Alembert reduction for the group $\text{Diff}(\mathbb{R}^n)$ (not $\text{Diff}(\Omega)$), left arrow pointing down). Alternatively, one again starts on the right side of the diagram with a formulation based on a *Hamiltonian* $H: T^* \text{Diff}(\Omega) \to \mathbb{R} = H(\varphi, p, \varrho_0, S_0)$ in material variables. These dynamics are now governed by an *extension* of the canonical Poisson structure $\{\cdot,\cdot\}_{\text{can}}$ which incorporates the bulk and boundary contributions. From this, an extended Lie-Poisson formulation in the Eulerian picture is derived by relabelling symmetries, mimicking the Lie-Poisson reduction process (right arrow pointing down). Again, under the appropriate regularity conditions, the Lagrangian and Hamiltonian approaches can be related through the Legendre transform, at the level of either material (top arrow) or Eulerian (bottom arrow) variables.

In summary, our geometric approach for open fluids meets the following criteria:

- (i) The fluid's evolution equations can be derived equivalently from either a variational perspective or a bracket (extended Poisson) perspective.
- (ii) It consistently extends the geometric Lie group formulation of closed fluids on both the Lagrangian and Hamiltonian sides, in both the material and Eulerian descriptions.
- (iii) It incorporates the process of reduction by symmetries by the relabelling group of diffeomorphisms.
- (iv) It coherently extends the fundamental variational and geometric formulations of finite dimensional mechanics, including the Hamilton and Lagrange-d'Alembert principles, as well as the canonical symplectic and Poisson structures.

Plan of the paper. The remainder of this paper is structured as follows. This section closes with a review of the existing general theory for closed fluids. Section [2](#page-10-0) discusses the Lagrangian variational principle for open fluids, in both Lagrangian and Eulerian coordinates. Section [3](#page-26-0) discusses the Hamiltonian variational principle and associated brackets for open fluids, in both Lagrangian and Eulerian coordinates. Section [4](#page-34-0) introduces extensions of the theory to handle multicomponent fluids, general advected quantities, higher-order fluids and boundary stresses. Finally, conclusions are discussed in Section [5.](#page-43-0) Appendix [A](#page-44-0) demonstrates the equivalent between the Lagrangian and Eulerian coordinate versions of the theory.

Hamilton and Lagrange-d'Alembert principles. Let us recall that the critical action principle of classical mechanics for a system with configuration manifold Q and Lagrangian $L: TQ \to \mathbb{R}$ is given by the *Hamilton principle*

$$
\delta \int_0^T L(q, \dot{q}) dt = 0. \tag{1.1}
$$

It seeks for critical curves of the Lagrangian action functional, among curves with fixed endpoints, leading to $\delta q(0) = \delta q(T) = 0$. Here TQ denotes the tangent bundle, or velocity space, of the manifold Q. When the system is subject to an external force F, then the *Lagranged'Alembert* principle must be used, in which the critical condition is amended by the virtual work of the force, giving

$$
\delta \int_0^T L(q, \dot{q}) dt + \int_0^T \langle F(q, \dot{q}), \delta q \rangle dt = 0.
$$
 (1.2)

In order to derive the equations and boundary conditions for open fluids, we shall use a continuum version of this principle. As we shall see, in this continuum extension, the configuration manifold Q of the system is a group of diffeomorphisms and the virtual force term has both distributed and boundary contributions.

We review below the geometric variational description for closed fluids. It takes its origin in the geometric formulation due to $[32]$ of the ideal fluid equations as geodesics on the group of volume preserving diffeomorphisms of the fluid domain. For compressible fluids, one can more generally formulate the equations of motion as Euler-Lagrange equations on the group of all diffeomorphisms of the fluid domain. In the material picture, the equations thus follows from the standard Hamilton variational principle associated to the fluid Lagrangian given by its kintic minus potential energy. The induced variational principle in the Eulerian picture is best understood by using the theory of Euler-Poincaré reduction for systems on Lie groups $[1]$. This geometric setting is also useful to understand the Hamiltonian side and to systematically explain the occurrence of Lie-Poisson bracket structures in the Eulerian description, emerging as reduction by symmetry of canonical Poisson bracket in the material description, see [\[7](#page-47-6), [8\]](#page-47-7). These geometric developments only hold for closed fluids, and it is our purpose to extend them to fluids with open boundaries.

1.1 Review of the case of closed fluids

The material description and Lagrangian densities. We consider the motion of a fluid in a bounded domain $\Omega \subset \mathbb{R}^n$, $n = 2, 3$. If we assume that the motion is smooth and the

fluid cannot escape the domain, then its evolution is fully described by a time dependent diffeomorphism of this domain, namely $\varphi(t, \cdot) \in \text{Diff}(\Omega)$. Here $\text{Diff}(\Omega)$ is the infinite dimensional Lie group of smooth diffeomorphisms of Ω , and it plays the role of the configuration manifold for the fluid system. Note in particular that the domain boundary is kept invariant during such motion, i.e. $\varphi(t, \partial \Omega) \subset \partial \Omega$. We shall use the standard continuum mechanics notation $x = \varphi(t, X)$ to describe the position $x \in \Omega$ of a fluid particle with label $X \in \Omega$, which is referred to as the material or Lagrangian description of the fluid, see Remark [1.1.](#page-7-0)

In addition to the fluid motion, for a full compressible fluid, one also needs to describe the evolution of its mass and entropy densities. For closed fluids and in absence of irreversible processes, these are described by time independent fields $\rho_0(X)$ and $S_0(X)$ in the material description.

As recalled earlier, following the classical Lagrangian mechanics setting, the Lagrangian function is defined on the tangent bundle TQ of the configuration manifold Q . For compressible fluids $Q = \text{Diff}(\Omega)$ and for each fixed ϱ_0 , S_0 , we have a Lagrangian function

$$
L(\cdot, \cdot, \varrho_0, S_0) : T \text{Diff}(\Omega) \to \mathbb{R}, \qquad (\varphi, \dot{\varphi}) \mapsto L(\varphi, \dot{\varphi}, \varrho_0, S_0).
$$

Here $\dot{\varphi} \in T_{\varphi}$ Diff(Ω) denotes an arbitrary tangent vector to the manifold Diff(Ω) at φ . It is given as a vector field on Ω along φ parallel to the boundary, i.e., $\dot{\varphi}: \Omega \to T\Omega$ is such that $\dot{\varphi}(X) \in T_{\varphi(X)}\Omega$, for all $X \in \Omega$ and $\dot{\varphi}(X)\|\partial\Omega$ for all $X \in \partial\Omega$. In particular for $\varphi = id$ the identity map this tangent space becomes the Lie algebra of $Diff(\Omega)$ given by the space of vector fields on Ω tangent to the boundary: T_{id} Diff(Ω) = $\mathfrak{X}_{\parallel}(\Omega)$.

In the main part of the paper, we shall assume that the Lagrangian function is given in terms of a Lagrangian density \mathfrak{L} as

$$
L(\varphi, \dot{\varphi}, \varrho_0, S_0) = \int_{\Omega} \mathfrak{L}(\varphi(X), \dot{\varphi}(X), \nabla_X \varphi(X), \varrho_0(X), S_0(X)) \mathrm{d}X,\tag{1.3}
$$

which is the case for the main fluid models. We shall briefly consider a more general case in §[4.3.](#page-42-0)

From now on we will not explicitly write the X-dependence of the fields in expressions such as (1.3) .

Hamilton's principle. For fluid motion in a domain Ω , Hamilton's principle [\(1.1\)](#page-5-1) reads, for each fixed ϱ_0 and S_0 ,

$$
\delta \int_0^T L(\varphi, \dot{\varphi}, \varrho_0, S_0) dt = 0 \tag{1.4}
$$

for arbitrary variations $\delta\varphi$ with $\delta\varphi|_{t=0,T} = 0$.

For L given in terms of a density $\mathfrak L$ as in [\(1.3\)](#page-6-0), Hamilton's principle [\(1.4\)](#page-6-1) gives the Euler-Lagrange equations and boundary conditions

$$
\frac{d}{dt}\frac{\partial \mathcal{L}}{\partial \dot{\varphi}} = \frac{\partial \mathcal{L}}{\partial \varphi} - \text{DIV} \frac{\partial \mathcal{L}}{\partial \nabla \varphi} \quad \text{and} \quad \frac{\partial \mathcal{L}}{\partial \nabla \varphi}(N, \delta \varphi) = 0, \ \forall \delta \varphi \parallel \partial \Omega,
$$
\n(1.5)

where N is the outward pointing unit normal vector field to Ω , seen as the domain of labels, see Remark [1.1,](#page-7-0) and DIV denotes the divergence operator with respect to the material variable X.

Using the local coordinate notations X^A for the labels X and x^a for the spatial points x, the equations above have the local form

$$
\frac{d}{dt}\frac{\partial \mathfrak{L}}{\partial \dot{\varphi}^a} = \frac{\partial \mathfrak{L}}{\partial \varphi^a} - \partial_A \frac{\partial \mathfrak{L}}{\partial \varphi^a_{,A}} \quad \text{and} \quad \frac{\partial \mathfrak{L}}{\partial \varphi^a_{,A}} N_A \delta \varphi^a = 0, \ \forall \delta \varphi \parallel \partial \Omega,
$$

where $\varphi^a_{,A} = \partial \varphi^a / \partial X^A$. The local notation makes clear the meaning of $\frac{\partial \mathfrak{L}}{\partial \nabla \varphi} (N, \delta \varphi)$ in [\(1.5\)](#page-6-2), namely, the evaluation of the 2-point tensor $\frac{\partial \mathfrak{L}}{\partial \nabla \varphi}$ along the material and spatial vectors N and δϕ.

The boundary condition emerging from Hamilton's principle indicates that the traction on the boundary can only be normal to the boundary, i.e. $\frac{\partial \mathfrak{L}}{\partial \varphi_{,A}^a} N_A = \lambda n_a$ for some λ and with n the outward pointing unit normal vector field to Ω seen as the spatial domain, see Remark [1.1.](#page-7-0) As we will see later, when $\mathfrak L$ satisfies the relabelling symmetry, this boundary condition becomes empty. For more general classes of Lagrangian than (1.3) , such as those used in elasticity, such a boundary condition is not empty even under relabelling symmetry, see [\[19\]](#page-48-4).

Note that the writing of these equations needs the introduction of a metric, chosen here as the Euclidean metric on Ω induced from the one on \mathbb{R}^n , see also Remark [1.2.](#page-8-1)

Remark 1.1 (Double role of Ω). In the case of a closed fluid with fixed boundary, it is possible to identify the domain of labels with the spatial domain of the fluid, which we have done here by choosing them both equal to Ω . In general however they have to be thought of as being distinct [\[33](#page-49-3)]. For instance the metrics on the space of labels and the spatial domain do not necessarily coincide and have totally different meanings. Even though in our case we choose in both cases the Euclidean metric, we introduce two different notations for the outward pointing unit normal vector field, N and n , according if we look at it on the domain of labels or the spatial domain. Such confusion can not arise for open fluids as these domains are distinct.

Relabelling symmetries. By the relabelling symmetry (or material covariance) of fluid dynamics, the Lagrangian density \mathfrak{L} must satisfy

$$
\mathfrak{L}(\varphi \circ \psi, \dot{\varphi} \circ \psi, \nabla_X(\varphi \circ \psi), (\varrho_0 \circ \psi)J\psi, (S_0 \circ \psi)J\psi) = \mathfrak{L}(\varphi, \dot{\varphi}, \nabla_X \varphi, \varrho_0, S_0) \circ \psi J\psi, \qquad (1.6)
$$

for all $\psi \in \text{Diff}(\Omega)$ with $J\psi$ the Jacobian of ψ , see [\[34\]](#page-49-4). This property turns out to be equivalent to the existence of a Lagrangian density $I(u, \rho, s)$ in the Eulerian description, such that

$$
\mathfrak{L}(\varphi, \dot{\varphi}, \nabla_X \varphi, \varrho_0, S_0) \circ \varphi^{-1} J \varphi^{-1} = \mathfrak{l}(u, \rho, s)
$$
\n(1.7)

for the Eulerian velocity, mass density, and entropy density

$$
u = \dot{\varphi} \circ \varphi^{-1}, \quad \rho = (\varrho_0 \circ \varphi^{-1}) J \varphi^{-1}, \quad s = (S_0 \circ \varphi^{-1}) J \varphi^{-1},
$$

see [\[34\]](#page-49-4) for details.

In terms of the associated Lagrangian function L in (1.3) , property (1.6) corresponds to its right $Diff(\Omega)$ -invariance, namely

$$
L(\varphi \circ \psi, \dot{\varphi} \circ \psi, \varrho_0 \circ \psi J\psi, S_0 \circ \psi J\psi) = L(\varphi, \dot{\varphi}, \varrho_0, S_0)
$$

for all $\psi \in \text{Diff}(\Omega)$, and (1.7) yields

$$
L(\varphi, \dot{\varphi}, S_0, \varrho_0) = \ell(u, \rho, s),
$$

for the reduced Lagrangian function

$$
\ell(u,\rho,s) = \int_{\Omega} \mathfrak{l}(u,\rho,s) dx.
$$

The Eulerian variational principle. As shown in $\boxed{1}$ via Lagrange reduction by symmetry, the Hamilton principle [\(1.4\)](#page-6-1) induces the following variational principle in the Eulerian picture

$$
\delta \int_0^T \ell(u,\rho,s) \mathrm{d}t = 0 \tag{1.8}
$$

for variations

$$
\delta u = \partial_t \zeta + \mathcal{L}_u \zeta, \qquad \delta \rho = -\operatorname{div}(\rho \zeta), \qquad \delta s = -\operatorname{div}(s \zeta), \tag{1.9}
$$

where ζ is an arbitrary time dependent vector field parallel to the boundary, vanishing at $t = 0, T$. The associated equations of motion read

$$
\begin{cases}\n\partial_t \frac{\partial \mathfrak{l}}{\partial u} + \mathcal{L}_u \frac{\partial \mathfrak{l}}{\partial u} = \rho \nabla \frac{\partial \mathfrak{l}}{\partial \rho} + s \nabla \frac{\partial \mathfrak{l}}{\partial s} \\
\partial_t \rho + \text{div}(\rho u) = 0, \qquad \partial_t s + \text{div}(su) = 0 \\
u \cdot n = 0 \quad \text{on} \quad \partial \Omega,\n\end{cases} (1.10)
$$

with

$$
\pounds_u \frac{\partial \mathfrak{l}}{\partial u} = u \cdot \nabla \frac{\partial \mathfrak{l}}{\partial u} + \nabla u^{\mathsf{T}} \frac{\partial \mathfrak{l}}{\partial u} + \frac{\partial \mathfrak{l}}{\partial u} \operatorname{div} u
$$

the Lie derivative of the fluid momentum density along u. We note that (1.8) – (1.9) only give the first equation. The advection equations follow from the definition $\rho = (\varrho_0 \circ \varphi^{-1})J\varphi^{-1}$ and $s = (S_0 \circ \varphi^{-1})J\varphi^{-1}$, while the boundary condition $u \cdot n = 0$ follow since $u = \dot{\varphi} \circ \varphi^{-1}$ and $\varphi \in \text{Diff}(\Omega)$.

Remark 1.2 (Equations on manifolds). In this paper we assume for simplicity that Ω is a bounded domain in the Euclidean space \mathbb{R}^n , however our approach can be extended to domains given by compact manifolds with boundary by following [\[33,](#page-49-3) [19\]](#page-48-4). In particular, an intrinsic formulation which makes explicit the dependence on Riemannian metrics on Ω is crucial to discuss the notion of covariances. These aspects will be pursued somewhere else.

1.2 Statement of an equivalent principle for closed fluids

While in the previous approach we fix the fields ϱ_0 and S_0 a priori and compute the critical condition by varying φ only, for our subsequent developments it is useful to reformulate the Hamilton principle [\(1.4\)](#page-6-1) as a critical action principle in which all the fields can be varied independently. We will utilize an extension of Hamilton's action functional recently developed in nonequilibrium thermodynamic, $\left[3, 35\right]$, which is reminiscent of a time integration by parts of a Clebsch-type variational principle, [\[36](#page-49-6)].

Material description. We consider the augmented Hamilton principle

$$
\delta \int_0^T \left[L(\varphi, \dot{\varphi}, \varrho, S) + \varrho \dot{W} + S\dot{\Gamma} \right] dt = 0 \qquad (1.11)
$$

for arbitrary variations $\delta\varphi$, δW , $\delta\Gamma$, $\delta\varrho$, δS , vanishing at $t = 0, T$.

While the criticality condition with respect to $\delta\varphi$ yields the same equations [\(1.5\)](#page-6-2) as above, the other variations give, respectively,

$$
\frac{d}{dt}\rho = 0, \quad \frac{d}{dt}S = 0, \quad \frac{d}{dt}W = -\frac{\partial \mathfrak{L}}{\partial \rho}, \quad \frac{d}{dt}\Gamma = -\frac{\partial \mathfrak{L}}{\partial S}.
$$
\n(1.12)

The first two equations recover the fact that the mass and entropy densities are constant in time, while the last two equations govern the dynamics of the additional fields W and Γ which do not play a role in the dynamics. In this approach, ρ and S are seen as independent variables, not predetermined. The actual mass and entropy densities of the fluid $\varrho_0(X)$ and $S_0(X)$ are inserted as initial conditions for $\rho(t, X)$ and $S(t, X)$. From the last two equations in [\(1.12\)](#page-9-0), the fields W and Γ acquire the meaning of thermodynamic displacements, as defined in [\[3,](#page-47-2) [37](#page-49-7)], which are relevant for the variational formulation of thermodynamic systems experiencing irreversible processes associated with heat and matter exchanges.

Eulerian description. The Eulerian version of [\(1.11\)](#page-8-4) will be derived in details later in the more general case of open fluids. For now we shall just notice that by defining the Eulerian thermodynamic displacments $w = W \circ \varphi^{-1}$ and $\gamma = \Gamma \circ \varphi^{-1}$, the variational principle [\(1.11\)](#page-8-4) yields

$$
\delta \int_0^T \left[\ell(u,\rho,s) + \rho D_t w + s D_t \gamma \right] dt = 0 \tag{1.13}
$$

for variations $\delta u = \partial_t \zeta + \mathcal{L}_u \zeta$ and for free variations $\delta \rho$, δs , δw , and $\delta \gamma$, vanishing at $t = 0, T$ and where, as earlier, ζ is an arbitrary time dependent vector field parallel to the boundary, vanishing at $t = 0, T$. We have denoted by D_t the Eulerian time derivative of scalar fields, i.e.,

$$
D_t w = \partial_t w + u \cdot \nabla w, \qquad D_t \gamma = \partial_t \gamma + u \cdot \nabla \gamma.
$$

Now the following Eulerian version of (1.12) emerges

$$
\bar{D}_t \rho = 0,
$$
 $\bar{D}_t s = 0,$ $D_t w = -\frac{\partial \mathfrak{l}}{\partial \rho},$ $D_t \gamma = -\frac{\partial \mathfrak{l}}{\partial s}$

from the variations δw , $\delta \gamma$, $\delta \rho$, δs , while the variations proportional to ζ yield the fluid mo-mentum equation, i.e., the first equation in [\(1.10\)](#page-8-5). Above we have introduced the notation \bar{D}_t for the Eulerian time derivative of density fields i.e.,

$$
\bar{D}_t \rho = \partial_t \rho + \text{div}(\rho u), \qquad \bar{D}_t s = \partial_t s + \text{div}(s u).
$$

In conclusion, in a similar way with its Lagrangian version (1.11) earlier, (1.13) is an alternative Lagrangian variational derivation of the system [\(1.10\)](#page-8-5) for closed fluids, which does not a priori assume the advection equations for the mass and entropy density, but derives them as critical conditions.

The principles [\(1.11\)](#page-8-4) and [\(1.13\)](#page-9-1) will be appropriately extended to handle the case of an open fluid below.

2 Lagrangian variational formulation for open fluids

2.1 Material description and the Lagrange-d'Alembert principle

Lagrangian and configuration group for open fluids. We now consider as before a domain Ω, but we allow the fluid to enter or leave the domain. In this context, while we remain focused on solving for the fluid motion within Ω , the fluid configuration map is no longer a diffeomorphism of Ω . Not only is the boundary $\partial\Omega$ not preserved by the fluid motion, but the set of fluid labels corresponding to particles currently within the domain Ω also depends on the fluid configuration.

This fact is made precise by considering as configuration group the group $\text{Diff}(\mathbb{R}^n)$ of diffeomorphisms of the whole Euclidean space and making the following definition. As before, we choose a Lagrangian density \mathfrak{L} , and ρ , S denote the mass and entropy densities in the material description.

Definition 2.1. *The Lagrangian function for an open fluid with domain* Ω *and Lagrangian density* L *is*

$$
L(\cdot, \cdot, \varrho, S) : T \operatorname{Diff}(\mathbb{R}^n) \to \mathbb{R}
$$

defined by:

$$
L(\varphi, \dot{\varphi}, \varrho, S) = \int_{\varphi^{-1}(\Omega)} \mathfrak{L}(\varphi, \dot{\varphi}, \nabla \varphi, \varrho, S) dX.
$$

We note the nontrivial dependence of L on φ which arises from the fact that we integrate only on the domain $\varphi^{-1}(\Omega) \subset \mathbb{R}^n$ which contains the labels X such that $x = \varphi(X)$ belongs to Ω . A similar Lagrangian appears in the treatment of fluid conveying tubes, see [\[38\]](#page-49-8), which is playing a key role in modeling the effect of boundary forces.

Remark 2.2 (Case with open and closed boundaries). In practical situations, the boundary can be made of several pieces some with closed and other with open boundary conditions. We shall see later how such cases can be treated, while still using $\text{Diff}(\mathbb{R}^n)$ as the configuration Lie group.

Remark 2.3 (Case of manifolds). This setting can be extended to the case when Ω is a compact manifold with smooth boundary. For this extension we use the fact that any such manifold Ω can be properly embedded into a smooth manifold without boundary Ω , which is given by a copy of its interior Int Ω , see [\[39\]](#page-49-9). In this case one uses Diff($\tilde{\Omega}$) instead of Diff(\mathbb{R}^n) in Definition [2.1.](#page-10-2)

Lagrange-d'Alembert principle for open fluids. We shall now insert bulk and boundary external flows by using a continuum version of the Lagrange-d'Alembert principle [\(1.2\)](#page-5-2). We shall describe these bulk and boundary flows as being associated to the fluid momentum, the mass density, and the entropy density. Following the Lagrange-d'Alembert approach, these effects can be naturally included by considering virtual works associated to the variation $\delta\varphi$, δW, and δΓ, both at the interior and at the boundary. In order to be consistently paired with these variations, the distributed/bulk and boundary sources must be given, for each $\varphi \in \text{Diff}(\Omega)$, as the following maps:

• A distributed source of momentum $\mathfrak{B}dX = \mathfrak{B}(\varphi, \dot{\varphi}, \nabla \varphi, \varrho, S)dX$, given as a map

$$
\mathfrak{B}dX : \varphi^{-1}(\Omega) \to T^*\mathbb{R}^n \otimes \Lambda^n(\varphi^{-1}(\Omega)); \tag{2.1}
$$

• A boundary source of momentum $\mathfrak{J}dX = \mathfrak{J}(\varphi, \dot{\varphi}, \nabla \varphi, \varrho, S)dX$, given as a map

$$
\mathfrak{J}dA : \varphi^{-1}(\partial \Omega) \to T^* \mathbb{R}^n \otimes \Lambda^{n-1}(\varphi^{-1}(\partial \Omega)); \tag{2.2}
$$

• Distributed sources of mass $\Theta_{\varrho}dX = \Theta_{\varrho}(\varphi,\dot{\varphi},\nabla\varphi,\varrho,S)dX$ and of entropy $\Theta_S dX =$ $\Theta_S(\varphi, \dot{\varphi}, \nabla \varphi, \varrho, S) dX$, given as maps

$$
\Theta_{\varrho}dX, \ \Theta_{S}dX : \varphi^{-1}(\Omega) \to \Lambda^{n}(\varphi^{-1}(\Omega)); \tag{2.3}
$$

• Boundary sources of mass $j_{\varrho}(\varphi, \dot{\varphi}, \nabla \varphi, \varrho, S)dA = j_{\varrho}dA$ and of entropy $j_{S}dA = j_{S}(\varphi, \dot{\varphi}, \nabla \varphi, \varrho, S)dA$, given as maps

$$
j_{\varrho}dA, \ \ j_{S}dA: \varphi^{-1}(\partial\Omega) \to \Lambda^{n-1}(\varphi^{-1}(\partial\Omega)). \tag{2.4}
$$

Here dA denotes the area element induced by dX on $\varphi^{-1}(\Omega)$ and $\Lambda^k(\varphi^{-1}(\Omega))$, resp., $\Lambda^k(\varphi^{-1}(\partial\Omega))$, denotes the bundle of k-forms on $\varphi^{-1}(\partial\Omega)$, resp., $\varphi^{-1}(\Omega)$.

More precisely, for each $X \in \varphi^{-1}(\Omega)$, we have

$$
\mathfrak{B}(X)\mathrm{d}X \in T_{\varphi(X)}^* \mathbb{R}^n \otimes \Lambda_X^n(\varphi^{-1}(\Omega))
$$

\n
$$
\Theta_{\varrho}(X)\mathrm{d}X \in \Lambda_X^{n-1}(\varphi^{-1}(\Omega))
$$

\n
$$
\Theta_S(X)\mathrm{d}X \in \Lambda_X^{n-1}(\varphi^{-1}(\Omega))
$$

and, for each $X \in \varphi^{-1}(\partial\Omega)$, we have

$$
\mathfrak{J}(X)\mathrm{d}A \in T_{\varphi(X)}^* \mathbb{R}^n \otimes \Lambda_X^{n-1}(\varphi^{-1}(\partial \Omega))
$$

$$
j_{\varrho}(X)\mathrm{d}A \in \Lambda_X^{n-1}(\varphi^{-1}(\partial \Omega))
$$

$$
j_S(X)\mathrm{d}A \in \Lambda_X^{n-1}(\varphi^{-1}(\partial \Omega)),
$$

with $\Lambda_X^k(\varphi^{-1}(\Omega))$ and $\Lambda_X^k(\varphi^{-1}(\partial\Omega))$ the vector fibers of the bundles at X.

In each case, the above clarification of the spaces in which these objects live will be useful to understand their transformation to the Eulerian frame. For this step it is important to have the dependence on $\varphi \in \text{Diff}(\mathbb{R}^n)$ explicitly written. Note that we have allowed these objects to have the same dependences as the Lagrangian density, i.e., on $(\varphi, \dot{\varphi}, \nabla \varphi, \varrho, S)$.

Based on the preceding considerations regarding

- (i) The general statement of the Lagrange-d'Alembert principle [\(1.2\)](#page-5-2) in classical mechanics;
- (ii) The equivalent formulation of Hamilton's principle given in $\S1.2$;
- (iii) The definition of the Lagrangian for open fluids in Definition [2.1;](#page-10-2)

we propose the following *Lagrange-d'Alembert principle for open fluids*:

$$
\frac{d}{d\varepsilon}\Big|_{\varepsilon=0} \int_0^T \int_{\varphi_\varepsilon^{-1}(\Omega)} \left[\mathfrak{L}(\varphi_\varepsilon, \dot{\varphi}_\varepsilon, \nabla \varphi_\varepsilon, \varrho_\varepsilon, S_\varepsilon) + \varrho_\varepsilon \dot{W}_\varepsilon + S_\varepsilon \dot{\Gamma}_\varepsilon \right] dX dt \n+ \int_0^T \int_{\varphi^{-1}(\Omega)} \left[\mathfrak{B} \cdot \delta \varphi + \Theta_\varrho \delta W + \Theta_S \delta \Gamma \right] dX dt \quad \leftarrow \text{bulk contribution} \n+ \int_0^T \int_{\varphi^{-1}(\partial \Omega)} \left[\mathfrak{J} \cdot \delta \varphi + \mathfrak{j}_\varrho \delta W + \mathfrak{j}_S \delta \Gamma \right] dA dt = 0 \quad \leftarrow \text{boundary contribution},
$$
\n(2.5)

where $\delta(\cdot) = \frac{d}{d\varepsilon}\big|_{\varepsilon=0}$ and we assume $\delta\varphi$, δW , and $\delta\Gamma$ vanish at $t=0,T$. In particular, we have applied the Lagrange-d'Alembert principle [\(1.2\)](#page-5-2) in which the action functional term $\int_0^T L(q, \dot{q}) dt$ is given by the one in [\(1.11\)](#page-8-4) and the virtual force term $\int_0^T \langle F(q, \dot{q}), \delta q \rangle dt$ is given by the sum of the contribution of all the distributed and boundary forces described above. Note that in this treatment the configuration variable q in (1.2) is now represented by the list of variables $(\varphi, \rho, S, W, \Gamma)$. Note also that while the application of this type of principle implies that all variations vanish at $t = 0, T$, we didn't assume that δS and $\delta \varrho$ vanish. This is because, in our case, the action functional does not involve \dot{S} and $\dot{\rho}$.

Proposition 2.4. *The variational principle* [\(2.5\)](#page-12-0) *gives the following open Euler-Lagrange equations and boundary conditions:*

$$
\begin{cases}\n\frac{d}{dt}\frac{\partial \mathfrak{L}}{\partial \dot{\varphi}} + \text{DIV} \frac{\partial \mathfrak{L}}{\partial \nabla \varphi} - \frac{\partial \mathfrak{L}}{\partial \varphi} = \mathfrak{B} \\
\frac{d}{dt}\varrho = \Theta_{\varrho}, \qquad \frac{d}{dt}S = \Theta_{S} \\
N \cdot \left(\frac{\partial \mathfrak{L}}{\partial \nabla \varphi} + (\nabla_{x}\varphi^{-1} \circ \varphi) \cdot \dot{\varphi} \frac{\partial \mathfrak{L}}{\partial \dot{\varphi}} + \left(\varrho \frac{\partial \mathfrak{L}}{\partial \varrho} + S \frac{\partial \mathfrak{L}}{\partial S} - \mathfrak{L}\right) \nabla_{x}\varphi^{-1} \circ \varphi\right) = -\mathfrak{J} \quad on \quad \partial \Omega \\
N \cdot \left((\nabla_{x}\varphi^{-1} \circ \varphi) \cdot \dot{\varphi} \varrho \right) = -\mathfrak{j}_{\varrho}, \qquad N \cdot \left((\nabla_{x}\varphi^{-1} \circ \varphi) \cdot \dot{\varphi} S \right) = -\mathfrak{j}_{S} \quad on \quad \partial \Omega,\n\end{cases}
$$
\n(2.6)

where N *is the outward pointing unit normal vector field to the boundary* $\partial \varphi^{-1}(\Omega)$ *.*

Proof. The treatment of this variational principle is not trivial since we take variations of a Lagrangian which is integrated on a domain that is ε -dependent: $\varphi_{\varepsilon}^{-1}(\Omega)$. The following formula will be useful

$$
\frac{d}{d\varepsilon}\bigg|_{\varepsilon=0} \int_{\varphi_{\varepsilon}^{-1}(\Omega)} f_{\varepsilon} dX = \int_{\varphi^{-1}(\partial\Omega)} f(\delta \varphi^{-1} \circ \varphi) \cdot N dA + \int_{\varphi^{-1}(\Omega)} \delta f dX, \tag{2.7}
$$

for f_{ε} and φ_{ε} families of functions and diffeomorphisms, both smoothly depending on ε , with $f_{\varepsilon=0} = f$ and $\varphi_{\varepsilon=0} = \varphi$.

In the following computation of (2.5) , we omit the dependence on S since it is identical to

the treatment of ϱ for simplicity. We compute

$$
\frac{d}{d\varepsilon}\Big|_{\varepsilon=0} \int_{0}^{T} \int_{\varphi_{\varepsilon}^{-1}(\Omega)} \mathfrak{L}(\varphi_{\varepsilon}, \dot{\varphi}_{\varepsilon}, \nabla_{X}\varphi_{\varepsilon}, \varrho_{\varepsilon}) dX dt \n= \int_{0}^{T} \int_{\varphi^{-1}(\Omega)} \left[\frac{\partial \mathfrak{L}}{\partial \varphi} \cdot \delta \varphi + \frac{\partial \mathfrak{L}}{\partial \dot{\varphi}} \cdot \delta \dot{\varphi} + \frac{\partial \mathfrak{L}}{\partial \nabla \varphi} \cdot \nabla_{X} \delta \varphi + \frac{\partial \mathfrak{L}}{\partial \varrho} \delta \varrho \right] dX dt \n+ \int_{0}^{T} \int_{\varphi^{-1}(\partial \Omega)} \left[\left[\frac{\partial \mathfrak{L}}{\partial \varphi} - \frac{d}{dt} \frac{\partial \mathfrak{L}}{\partial \dot{\varphi}} - \mathrm{DIV} \frac{\partial \mathfrak{L}}{\partial \nabla \varphi} \right] \cdot \delta \varphi + \frac{\partial \mathfrak{L}}{\partial \varrho} \delta \varrho \right] dX dt \n+ \int_{0}^{T} \frac{d}{dt} \int_{\varphi^{-1}(\Omega)} \left[\frac{\partial \mathfrak{L}}{\partial \varphi} \cdot \delta \varphi dX dt \n+ \int_{0}^{T} \int_{\varphi^{-1}(\partial \Omega)} \left[\frac{\partial \mathfrak{L}}{\partial \nabla \varphi} \cdot \delta \varphi + \mathfrak{L} \delta \varphi^{-1} \circ \varphi - \left(\frac{\partial \mathfrak{L}}{\partial \dot{\varphi}} \cdot \delta \varphi \right) \frac{d}{dt} \varphi^{-1} \circ \varphi \right] \cdot N dA dt \n+ \int_{0}^{T} \int_{\varphi^{-1}(\partial \Omega)} \left[\left[\frac{\partial \mathfrak{L}}{\partial \varphi} - \frac{d}{dt} \frac{\partial \mathfrak{L}}{\partial \dot{\varphi}} - \mathrm{DIV} \frac{\partial \mathfrak{L}}{\partial \nabla \varphi} \right] \cdot \delta \varphi + \frac{\partial \mathfrak{L}}{\partial \varrho} \delta \varrho \right] dX dt \n+ \int_{0}^{T} \int_{\varphi^{-1}(\partial \Omega
$$

where we used $\delta\varphi^{-1} = -\nabla_x\varphi^{-1} \cdot \delta\varphi \circ \varphi^{-1}$ as well as $\delta\varphi_0 = \delta\varphi_T = 0$. We also have

$$
\frac{d}{d\varepsilon}\Big|_{\varepsilon=0} \int_0^T \int_{\varphi_\varepsilon^{-1}(\Omega)} \varrho_\varepsilon \dot{W}_\varepsilon dX dt \n= \int_0^T \int_{\varphi^{-1}(\Omega)} \left[\dot{W} \delta \varrho + \frac{d}{dt} (\varrho \delta W) - \dot{\varrho} \delta W \right] dX dt + \int_0^T \int_{\varphi^{-1}(\partial \Omega)} \varrho \dot{W} \delta \varphi^{-1} \circ \varphi \cdot N dA dt \n= \int_0^T \int_{\varphi^{-1}(\Omega)} \left[\dot{W} \delta \varrho - \dot{\varrho} \delta W \right] dX dt + \int_0^T \int_{\varphi^{-1}(\partial \Omega)} \left[\varrho \dot{W} \delta \varphi^{-1} \circ \varphi - \varrho \delta W \frac{d}{dt} \varphi^{-1} \circ \varphi \right] \cdot N dA dt \n= \int_0^T \int_{\varphi^{-1}(\Omega)} \left[\dot{W} \delta \varrho - \dot{\varrho} \delta W \right] dX dt \n+ \int_0^T \int_{\varphi^{-1}(\partial \Omega)} \left[-\varrho \dot{W} (\nabla_x \varphi^{-1} \circ \varphi) \cdot \delta \varphi + (\nabla_x \varphi^{-1} \circ \varphi) \cdot \dot{\varphi} \varrho \delta W \right] \cdot N dA dt.
$$

From the variations $\delta \varrho$, we get the condition

$$
\dot{W} = -\frac{\partial \mathfrak{L}}{\partial \varrho}.
$$

Then, by using this condition, and collecting the terms proportional to the variations $\delta\varphi$ and δW at the interior and at the boundary, we get the system (2.6) .

Remark 2.5 (Local expression). For further understanding of the notations, it is useful to rewrite the equations [\(2.6\)](#page-12-1) locally. Using as before X^A and x^a for the material and spatial

coordinates, we have

$$
\begin{cases}\n\frac{d}{dt}\frac{\partial \mathfrak{L}}{\partial \dot{\varphi}^{a}} + \partial_{A}\frac{\partial \mathfrak{L}}{\partial \varphi^{a}_{,A}} - \frac{\partial \mathfrak{L}}{\partial \varphi^{a}} = \mathfrak{B}_{a} \\
\frac{d}{dt}\varrho = \Theta_{\varrho}, \qquad \frac{d}{dt}S = \Theta_{S} \\
N_{A}\left(\frac{\partial \mathfrak{L}}{\partial \varphi^{a}_{,A}} + \left((\varphi^{-1})^{A}_{,b}\circ\varphi\right)\dot{\varphi}^{b}\frac{\partial \mathfrak{L}}{\partial \dot{\varphi}^{a}} + \left(\varrho\frac{\partial \mathfrak{L}}{\partial \varrho} + S\frac{\partial \mathfrak{L}}{\partial S} - \mathfrak{L}\right)(\varphi^{-1})^{A}_{,a}\circ\varphi\right) = -\mathfrak{J}_{a} \text{ on } \partial\Omega \\
N_{A}\left((\varphi^{-1})^{A}_{,b}\circ\varphi\right)\dot{\varphi}^{b}\varrho = -\mathfrak{j}_{\varrho} \quad \text{on} \quad \partial\Omega, \qquad N_{A}\left((\varphi^{-1})^{A}_{,b}\circ\varphi\right)\dot{\varphi}^{b}S = -\mathfrak{j}_{S} \quad \text{on} \quad \partial\Omega.\n\end{cases}
$$
\n(2.8)

Example: the compressible Euler fluid with open boundaries. The Lagrangian density is given as

$$
\mathfrak{L}(\varphi, \dot{\varphi}, \nabla \varphi, \varrho, S) = \frac{1}{2} \varrho |\dot{\varphi}|^2 - \varepsilon \left(\frac{\varrho}{J\varphi}, \frac{S}{J\varphi} \right) J\varphi,
$$
\n(2.9)

with ε the energy density of the fluid, expressed in terms of the Eulerian mass and entropy densities ρ and s. We shall denote by $\frac{\partial \varepsilon}{\partial \rho}$ and $\frac{\partial \varepsilon}{\partial s}$ the partial derivative, even though ρ and s do not appear in the material description (recall that $\rho = (\varrho \circ \varphi)J\varphi$ and $s = (S \circ \varphi)J\varphi$). The partial derivatives are:

$$
\frac{\partial \mathfrak{L}}{\partial \varphi} = 0, \quad \frac{\partial \mathfrak{L}}{\partial \dot{\varphi}} = \varrho \dot{\varphi}, \quad \frac{\partial \mathfrak{L}}{\partial \varrho} = \frac{1}{2} |\dot{\varphi}|^2 - \frac{\partial \varepsilon}{\partial \rho}, \quad \frac{\partial \mathfrak{L}}{\partial S} = -\frac{\partial \varepsilon}{\partial s}
$$
\n
$$
\frac{\partial \mathfrak{L}}{\partial \nabla \varphi} = \left(\frac{\partial \varepsilon}{\partial \rho} \frac{\varrho}{J \varphi} + \frac{\partial \varepsilon}{\partial s} \frac{S}{J \varphi} - \varepsilon \right) J \varphi \left(\nabla_x \varphi^{-1} \circ \varphi \right) = (p \circ \varphi) J \varphi \left(\nabla_x \varphi^{-1} \circ \varphi \right). \tag{2.10}
$$

In the last equality, we used the definition of the pressure in terms of the internal energy density as

$$
p = \frac{\partial \varepsilon}{\partial \rho}(\rho, s)\rho + \frac{\partial \varepsilon}{\partial s}(\rho, s)s - \varepsilon(\rho, s)
$$

which, in material coordinates, results in

$$
p \circ \varphi = \frac{\partial \varepsilon}{\partial \rho} \left(\frac{\varrho}{J\varphi}, \frac{S}{J\varphi} \right) \frac{\varrho}{J\varphi} + \frac{\partial \varepsilon}{\partial s} \left(\frac{\varrho}{J\varphi}, \frac{S}{J\varphi} \right) \frac{S}{J\varphi} - \varepsilon \left(\frac{\varrho}{J\varphi}, \frac{S}{J\varphi} \right).
$$

Then, we have

$$
\text{DIV} \left(p \circ \varphi \left(\nabla \varphi^{-1} \circ \varphi \right) J\varphi \right) = \nabla_X (p \circ \varphi) \cdot \left(\nabla \varphi^{-1} \circ \varphi \right) J\varphi = (\nabla_x p \circ \varphi) J\varphi,
$$

by using the Piola identity $\text{DIV}(J\varphi(\nabla_x\varphi^{-1} \circ \varphi)) = 0$. With this, the first equation of [\(2.6\)](#page-12-1) reduces to

$$
\frac{d}{dt}(\varrho\dot{\varphi}) + (\nabla_x p \circ \varphi)J\varphi = \mathfrak{B},
$$

which is the fluid momentum equation in the material description.

By using (2.10) remarkable cancellations occur in the third equation of (2.6) yielding simply

$$
N \cdot \left[\left(\nabla_x \varphi^{-1} \circ \varphi \right) \cdot \dot{\varphi} \right] \varrho \dot{\varphi} = -\mathfrak{J}.
$$

We shall mention later the origin of these cancellations, see Remark [2.8.](#page-17-0) We have thus obtained the following statement.

Proposition 2.6. *The equation for an open compressible Euler fluid in material coordinates are given by*

$$
\begin{cases}\n\frac{d}{dt}(\varrho\dot{\varphi}) + (\nabla_x p \circ \varphi)J\varphi = \mathfrak{B} \\
\frac{d}{dt}\varrho = \Theta_{\varrho}, \qquad \frac{d}{dt}S = \Theta_S \\
N \cdot [(\nabla_x \varphi^{-1} \circ \varphi) \cdot \dot{\varphi}] \varrho\dot{\varphi} = -\mathfrak{J} \quad on \quad \partial\Omega \\
N \cdot [(\nabla_x \varphi^{-1} \circ \varphi) \cdot \dot{\varphi}] \varrho = -\mathfrak{j}_{\varrho}, \quad N \cdot [(\nabla_x \varphi^{-1} \circ \varphi) \cdot \dot{\varphi}] S = -\mathfrak{j}_{S} \quad on \quad \partial\Omega.\n\end{cases}
$$
\n(2.11)

Balance laws in the material description. The seemingly involved expressions of the boundary conditions in (2.6) in the material description, see also (2.11) , is due to the fact that the domain of labels is time dependent. These expressions naturally appear in the boundary integrals when computing the balance laws.

For instance, for the balance of mass, one notes

$$
\frac{d}{dt} \int_{\varphi^{-1}(\Omega)} \varrho \, dX = \int_{\varphi^{-1}(\Omega)} \frac{d}{dt} \varrho \, dX + \int_{\varphi^{-1}(\partial \Omega)} \frac{d}{dt} (\varphi^{-1}) \circ \varphi \cdot N \, dA
$$
\n
$$
= \int_{\varphi^{-1}(\Omega)} \frac{d}{dt} \varrho \, dX - \int_{\varphi^{-1}(\partial \Omega)} N \cdot \left[(\nabla_x \varphi^{-1} \circ \varphi) \cdot \dot{\varphi} \right] \varrho \, dA
$$
\n
$$
= \int_{\varphi^{-1}(\Omega)} \Theta_{\varrho} \, dX + \int_{\varphi^{-1}(\partial \Omega)} j_{\varrho} dA,
$$

which used the open boundary condition for the mass density in (2.6) in the last equality. Similarly, for the entropy density, one has

$$
\frac{d}{dt} \int_{\varphi^{-1}(\Omega)} S \, dX = \int_{\varphi^{-1}(\Omega)} \Theta_S \, dX + \int_{\varphi^{-1}(\partial \Omega)} j_S dA.
$$

This is consistent with the interpretation of Θ_{ϱ} , Θ_{S} as the distributed/bulk source of mass and entropy, and j_{ρ} and j_{S} as the boundary sources of mass and entropy.

Regarding the energy balance, we recall that the total energy density in the material description, for a given Lagrangian density \mathfrak{L} is found as

$$
\mathfrak{E} = \frac{\partial \mathfrak{L}}{\partial \dot{\varphi}} \cdot \dot{\varphi} - \mathfrak{L}.
$$

From [\(2.6\)](#page-12-1), we get the local energy balance

$$
\frac{d}{dt}\mathfrak{E} = -\operatorname{DIV}\left(\frac{\partial \mathfrak{L}}{\partial \nabla \varphi} \cdot \dot{\varphi}\right) + \dot{\varphi} \cdot \mathfrak{B} - \frac{\partial \mathfrak{L}}{\partial \varrho} \Theta_{\rho} - \frac{\partial \mathfrak{L}}{\partial S} \Theta_{S}.
$$
\n(2.12)

Hence, the total energy balance follows from the three boundary conditions in [\(2.6\)](#page-12-1) as

$$
\frac{d}{dt} \int_{\varphi^{-1}(\Omega)} \mathfrak{E} dX = \int_{\varphi^{-1}(\Omega)} \left[\dot{\varphi} \cdot \mathfrak{B} - \Theta_{\varrho} \frac{\partial \mathfrak{L}}{\partial \varrho} - \Theta_{S} \frac{\partial \mathfrak{L}}{\partial S} \right] dX - \int_{\varphi^{-1}(\partial \Omega)} \left[\dot{\varphi} \cdot \mathfrak{J} - \frac{\partial \mathfrak{L}}{\partial \varrho} \mathfrak{j}_{\rho} - \frac{\partial \mathfrak{L}}{\partial S} \mathfrak{j}_{S} \right] dA,
$$

which show the change of energy due to bulk (first integral) and boundary (second integral) contributions. In particular, the somehow involved expression of the first boundary condition in (2.6) is used to express the boundary integral arising from the divergence term in (2.12) thereby yielding the simple final form.

2.2 Eulerian variational principle

As recalled in §[1.1](#page-5-0) for the case of closed fluids, the variational formulation in the Eulerian frame is best explained via the process of Lagrangian reduction by symmetry. In our case, the fluid Lagrangian is a function

$$
L: T\operatorname{Diff}(\mathbb{R}^n) \times \Omega^n(\mathbb{R}^n) \times \Omega^n(\mathbb{R}^n) \to \mathbb{R},
$$

$$
L(\varphi, \dot{\varphi}, \varrho, S) = \int_{\varphi^{-1}(\Omega)} \mathfrak{L}(\varphi(X), \dot{\varphi}(X), \nabla_X \varphi(X), \varrho(X), S(X)) \,dX
$$

and the relabelling symmetry is given as the $\text{Diff}(\mathbb{R}^n)$ -invariance

$$
L(\varphi \circ \psi, \dot{\varphi} \circ \psi, (\varrho \circ \psi)J\psi, (S \circ \psi)J\psi) = L(\varphi, \dot{\varphi}, \varrho, S), \text{ for all } \psi \in \text{Diff}(\mathbb{R}^n). \tag{2.13}
$$

From this, L induces a unique reduced Lagrangian

$$
\ell : \mathfrak{X}(\mathbb{R}^n) \times \Omega^n(\mathbb{R}^n) \times \Omega^n(\mathbb{R}^n) \to \mathbb{R}
$$
\n
$$
(2.14)
$$

defined by $\ell(u, \rho, s) = L(\varphi, \dot{\varphi}, \rho, S)$, where the variables on the right hand side are such that $u = \dot{\varphi} \circ \varphi^{-1}, \, \rho = (\varrho \circ \varphi^{-1})J\varphi^{-1}, \text{ and } S = (S \circ \varphi^{-1})J\varphi^{-1}.$ Let us emphasize that even though the Lagrangian is not integrated over the whole space \mathbb{R}^n , it is still Diff(\mathbb{R}^n)-invariant, so that the ideas of Lagrangian reduction still apply in the open case. We refer to Remark [2.10](#page-20-0) for the link with the theory of Lagrangian reduction theory by symmetry. See Remark [3.3](#page-29-0) for similar comments on the Hamiltonian side.

Relabelling symmetries. For our development it is useful to put the emphasis on the symmetries of the Lagrangian density rather than those of the Lagrangian function. We will say that the Lagrangian density $\mathfrak L$ is materially covariant if it satisfies

$$
\mathfrak{L}(\varphi \circ \psi, \dot{\varphi} \circ \psi, \nabla_X(\varphi \circ \psi), (\varrho \circ \psi)J\psi, (S \circ \psi)J\psi) = \mathfrak{L}(\varphi, \dot{\varphi}, \nabla_X \varphi, \varrho, S) \circ \psi J\psi, \quad (2.15)
$$

for all $\psi \in \text{Diff}(\mathbb{R}^n)$, see [\[34\]](#page-49-4). This property is the natural extension of the material covariance property of stored energy functions in elasticity [\[33,](#page-49-3) [19](#page-48-4)] to the case of general Lagrangian density. It is easy to see that it implies the Diff(\mathbb{R}^n)-invariance [\(2.13\)](#page-16-1) of the associated Lagrangian function L. Also, the symmetry (2.15) is equivalent to the existence of a Lagrangian density $l(u, \rho, s)$ in the Eulerian description, such that

$$
\mathfrak{L}(\varphi, \dot{\varphi}, \nabla_X \varphi, \varrho, S) \circ \varphi^{-1} J \varphi^{-1} = \mathfrak{l}(u, \rho, s).
$$
\n(2.16)

where $u = \dot{\varphi} \circ \varphi^{-1}$, $\rho = (\varrho \circ \varphi^{-1})J\varphi^{-1}$, $s = (S \circ \varphi^{-1})J\varphi^{-1}$. From this, the reduced Lagrangian ℓ in (2.14) associated to L is given in terms of l

$$
\ell: \mathfrak{X}(\mathbb{R}^n) \times \Omega^n(\mathbb{R}^n) \times \Omega^n(\mathbb{R}^n) \to \mathbb{R}, \quad \ell(u,\rho,s) = \int_{\Omega} \mathfrak{l}(u(x),\rho(x),s(x))dx.
$$

It will be important to distinguish between the Lagrangian function ℓ and the Lagrangian density l. For instance, we would like to emphasize that the Lagrangian ℓ is defined on the whole Lie algebra $\mathfrak{X}(\mathbb{R}^n)$, but its expression effectively depends only on the restriction $u|_{\Omega}$ of $u \in \mathfrak{X}(\mathbb{R}^n)$ to Ω . Similarly for the variables ρ and s.

Remark 2.7 (Properties of \mathfrak{L}). While the properties [\(2.15\)](#page-16-2) and [\(2.16\)](#page-16-4) are written, for convenience, in terms of fields, such as φ , ϱ , and u , these properties only depends on the point values of such fields, as a careful examination shows, so that they are really properties of the densities \mathfrak{L} and **l**, see [\[34\]](#page-49-4) for details.

Remark 2.8 (Property of materially covariant Lagrangian densities). We noticed earlier a drastic simplification of one of the boundary condition in the case of the compressible Euler fluid. It is not associated to this particular example, but follows from the material covariance [\(2.15\)](#page-16-2) of the Lagrangian density. Indeed, the infinitesimal property associated to the material covariance (2.15) yields the following relation between the partial derivatives of \mathfrak{L} :

$$
\frac{\partial \mathfrak{L}}{\partial \nabla \varphi} + \left(\frac{\partial \mathfrak{L}}{\partial \varrho} \varrho + \frac{\partial \mathfrak{L}}{\partial S} S - \mathfrak{L} \right) \nabla_x \varphi^{-1} \circ \varphi = 0,
$$

see, e.g., $[34]$. In this case, the first boundary condition in (2.6) reduces to just

$$
\left(N \cdot (\nabla_x \varphi^{-1} \circ \varphi) \cdot \dot{\varphi}\right) \frac{\partial \mathfrak{L}}{\partial \dot{\varphi}} = -\mathfrak{J}, \quad \text{i.e.,} \quad N_A\big((\varphi^{-1})^A_{,b} \circ \varphi\big) \dot{\varphi}^b \frac{\partial \mathfrak{L}}{\partial \dot{\varphi}^a} = -\mathfrak{J}_a.
$$

In a similar way with the Lagrangian density, the bulk and boundary fluxes must also satisfy symmetries in order to admit a purely Eulerian formulation. They are given by

$$
\mathfrak{B}(\varphi \circ \psi, \dot{\varphi} \circ \psi, \nabla_X(\varphi \circ \psi), (\varrho \circ \psi)J\psi, (S \circ \psi)J\psi) = \mathfrak{B}(\varphi, \dot{\varphi}, \nabla \varphi, \varrho, S) \circ \psi J\psi \n\Theta_{\varrho}(\varphi \circ \psi, \dot{\varphi} \circ \psi, \nabla_X(\varphi \circ \psi), (\varrho \circ \psi)J\psi, (S \circ \psi)J\psi) = \Theta_{\varrho}(\varphi, \dot{\varphi}, \nabla \varphi, \varrho, S) \circ \psi J\psi,
$$
\n(2.17)

similarly for Θ_s , and by

$$
\mathfrak{J}(\varphi \circ \psi, \dot{\varphi} \circ \psi, \nabla_X(\varphi \circ \psi), (\varrho \circ \psi)J\psi, (S \circ \psi)J\psi) = \mathfrak{J}(\varphi, \dot{\varphi}, \nabla \varphi, \varrho, S) \circ \psi_{\partial}J\psi_{\partial}
$$
\n
$$
\mathfrak{j}_{\varrho}(\varphi \circ \psi, \dot{\varphi} \circ \psi, \nabla_X(\varphi \circ \psi), (\varrho \circ \psi)J\psi, (S \circ \psi)J\psi) = \mathfrak{j}_{\varrho}(\varphi, \dot{\varphi}, \nabla \varphi, \varrho, S) \circ \psi_{\partial}J\psi_{\partial},
$$
\n(2.18)

similarly for js, for all $\psi \in \text{Diff}(\mathbb{R}^n)$. We used the notation ψ_{∂} for the restriction of the diffeomorhism ψ to the boundary, and $J\psi_{\partial}$ for the Jacobian of ψ_{∂} . These properties are equivalent to the existence of the following Eulerian objects:

• a distributed source of momentum

$$
b(u, \rho, s)dx : \Omega \to T^*\mathbb{R}^n \otimes \Lambda^n(\Omega);
$$

• a boundary source of momentum

$$
j(u, \rho, s)da : \partial\Omega \to T^*\mathbb{R}^n \otimes \Lambda^{n-1}(\partial\Omega);
$$

• distributed sources of mass and entropy

$$
\theta_{\rho}(u,\rho,s)dx, \ \theta_{s}(u,\rho,s)dx:\Omega\to\Lambda^{n}(\Omega);
$$

• boundary sources of mass and entropy

$$
j_{\rho}(u,\rho,s)da, j_s(u,\rho,s)da : \partial\Omega \to \Lambda^{n-1}(\partial\Omega);
$$

such that

$$
\mathfrak{B}(\varphi, \ldots) \circ \varphi^{-1} J \varphi^{-1} = b(u, \rho, s) \n\Theta_{\varrho}(\varphi, \ldots) \circ \varphi^{-1} J \varphi^{-1} = \theta_{\rho}(u, \rho, s), \qquad \Theta_{S}(\varphi, \ldots) \circ \varphi^{-1} J \varphi^{-1} = \theta_{s}(u, \rho, s)
$$
\n(2.19)

and

$$
\mathfrak{J}(\varphi, \ldots) \circ \varphi_{\partial}^{-1} J \varphi_{\partial}^{-1} = J(u, \rho, s), \n\mathfrak{j}_{\varrho}(\varphi, \ldots) \circ \varphi_{\partial}^{-1} J \varphi_{\partial}^{-1} = j_{\rho}(u, \rho, s), \qquad \mathfrak{j}_{S}(\varphi, \ldots) \circ \varphi_{\partial}^{-1} J \varphi_{\partial}^{-1} = j_{s}(u, \rho, s),
$$
\n(2.20)

where $u = \dot{\varphi} \circ \varphi^{-1}$, $\rho = (\varrho \circ \varphi^{-1})J\varphi^{-1}$, and $S = (S \circ \varphi^{-1})J\varphi^{-1}$. Above, we have used the abbreviation $(\varphi, \ldots) = (\varphi, \dot{\varphi}, \varrho, S).$

Note that the transformations appearing in (2.17) – (2.20) are the natural ones associated to the geometric type of these objects, see (2.1) – (2.4) .

Eulerian variational principle for open fluids. Under the relabelling symmetries described above, we can state the following variational formulation of open fluids in the Eulerian frame:

Proposition 2.9 (Eulerian variational principle for open fluids). *Assume that the Lagrangian density and the distributed and boundary sources of momentum, mass and entropy satisfy the relabelling symmetries. Then the following hold:*

(i) *The Lagrange-d'Alembert principle* [\(2.5\)](#page-12-0) *yields the following variational formulation in the Eulerian frame (Euler-Poincaré-d'Alembert):*

$$
\delta \int_0^T \int_{\Omega} \left[I(u, \rho, s) + \rho D_t w + s D_t \gamma \right] dx dt
$$

+
$$
\int_0^T \int_{\Omega} \left[b \cdot \zeta + \theta_{\rho} D_{\delta} w + \theta_s D_{\delta} \gamma \right] dx dt \leftarrow bulk contribution +
$$
\int_0^T \int_{\partial \Omega} \left[J \cdot \zeta + j_{\rho} D_{\delta} w + j_s D_{\delta} \gamma \right] d\alpha dt = 0 \leftarrow boundary contribution
$$
 (2.21)
$$

with respect to variations $\delta u = \partial_t \zeta + [\zeta, u]$ *and free variations* $\delta \rho$ *,* δs *,* δw *,* $\delta \gamma$ *, with* δw *and* $\delta \gamma$ *vanishing at* $t = 0, T$ *and* ζ *an arbitrary time dependent vector field vanishing at* $t = 0, T$.

The relation with the material variables used in [\(2.5\)](#page-12-0) *is given by*

$$
u = \dot{\varphi} \circ \varphi^{-1}, \quad \rho = (\varrho \circ \varphi^{-1})J\varphi^{-1}, \quad s = (S \circ \varphi^{-1})J\varphi^{-1}, \quad w = W \circ \varphi^{-1}, \quad \gamma = \Gamma \circ \varphi^{-1}.
$$

(ii) *This principle yields the following equations and boundary conditions for open fluids:*

$$
\begin{cases}\n\partial_t \frac{\partial \mathfrak{l}}{\partial u} + \mathcal{L}_u \frac{\partial \mathfrak{l}}{\partial u} - \rho \nabla \frac{\partial \mathfrak{l}}{\partial \rho} - s \nabla \frac{\partial \mathfrak{l}}{\partial s} = b \\
\partial_t \rho + \operatorname{div}(\rho u) = \theta_\rho, \qquad \partial_t s + \operatorname{div}(s u) = \theta_s \\
(u \cdot n) \frac{\partial \mathfrak{l}}{\partial u} = -J \quad on \quad \partial \Omega \\
\rho u \cdot n = -j_\rho, \qquad su \cdot n = -j_s \quad on \quad \partial \Omega.\n\end{cases} \tag{2.22}
$$

Proof. (i) By using the relations [\(2.19\)](#page-18-1), [\(2.20\)](#page-18-0), as well as $\delta W = D_{\delta}w \circ \varphi$ and $\delta \Gamma = D_{\delta} \gamma \circ \varphi$, the bulk and boundary source terms in (2.5) yield the ones in (2.21) . From the relabelling symmetry of \mathfrak{L} , we have from (2.16)

$$
\frac{d}{d\varepsilon}\Big|_{\varepsilon=0} \int_{\varphi_{\varepsilon}^{-1}(\Omega)} \mathfrak{L}(\varphi_{\varepsilon}, \dot{\varphi}_{\varepsilon}, \nabla_X \varphi_{\varepsilon}, \varrho_{\varepsilon}, S_{\varepsilon}) dX = \frac{d}{d\varepsilon}\Big|_{\varepsilon=0} \int_{\Omega} \mathfrak{L}(\varphi_{\varepsilon}, \dot{\varphi}_{\varepsilon}, \nabla_X \varphi_{\varepsilon}, \varrho_{\varepsilon}, S_{\varepsilon}) \circ \varphi_{\varepsilon}^{-1} J \varphi_{\varepsilon}^{-1} dx
$$

\n
$$
= \int_{\Omega} \frac{d}{d\varepsilon}\Big|_{\varepsilon=0} \mathfrak{L}(\varphi_{\varepsilon}, \dot{\varphi}_{\varepsilon}, \nabla_X \varphi_{\varepsilon}, \varrho_{\varepsilon}, S_{\varepsilon}) \circ \varphi_{\varepsilon}^{-1} J \varphi_{\varepsilon}^{-1} dx
$$

\n
$$
= \int_{\Omega} \frac{d}{d\varepsilon}\Big|_{\varepsilon=0} \mathfrak{l}(u_{\varepsilon}, \rho_{\varepsilon}, s_{\varepsilon}) dx,
$$

where $u_{\varepsilon} = \dot{\varphi}_{\varepsilon} \circ \varphi^{-1}$ and $\rho_{\varepsilon} = \varrho_{\varepsilon} \circ \varphi_{\varepsilon}^{-1} J \varphi_{\varepsilon}^{-1}$, $s_{\varepsilon} = S_{\varepsilon} \circ \varphi_{\varepsilon}^{-1} J \varphi_{\varepsilon}^{-1}$. From these relations, we have δ $u = \partial_t \zeta + \pounds_u \zeta$

while $\delta \rho$ and δs are free since $\delta \rho$ and δS are free. We also have, by using $w = W \circ \varphi^{-1}$,

$$
\int_{\varphi^{-1}(\Omega)} \varrho \dot{W} \mathrm{d}X = \int_{\Omega} \varrho \circ \varphi^{-1} J \varphi^{-1} \dot{W} \circ \varphi^{-1} \mathrm{d}x = \int_{\Omega} \rho D_t w \mathrm{d}x.
$$

Similarly we have $\int_{\varphi^{-1}(\Omega)} S\dot{\Gamma} dX = \int_{\Omega} sD_t \gamma dx$. In conclusion, the principle [\(2.5\)](#page-12-0) yields [\(2.21\)](#page-18-2).

(ii) The treatment of (s, γ) is similar to that of (ρ, w) , so it is enough to present the computation for the variation of $\int_0^T \int_{\Omega} [(u,\rho) + \rho D_t w] \, \mathrm{d}x \mathrm{d}t$. The variation with respect to ρ gives the condition

$$
D_t w = -\frac{\partial \mathfrak{l}}{\partial \rho}.\tag{2.23}
$$

The variation of $\int_0^T \int_{\Omega} [I(u,\rho) + \rho D_t w] \, \mathrm{d}x \mathrm{d}t$ with respect to u and w is computed as follows:

$$
= \int_{0}^{T} \int_{\Omega} \left[\frac{\partial l}{\partial u} \cdot (\partial_{t} \zeta + \mathcal{L}_{u} \zeta) + \rho D_{t} \delta w + (\partial_{t} \zeta + \mathcal{L}_{u} \zeta) \cdot \rho \nabla w \right] dxdt
$$

\n
$$
= \int_{0}^{T} \int_{\Omega} \left[\left(-\partial_{t} \frac{\partial l}{\partial u} - \mathcal{L}_{u} \frac{\partial l}{\partial u} \right) \cdot \zeta + \text{div} \left(\left(\frac{\partial l}{\partial u} \cdot \zeta \right) u \right) + \bar{D}_{t} (\rho \delta w) - \bar{D}_{t} \rho \delta w + \bar{D}_{t} (\rho \zeta \cdot \nabla w)
$$

\n
$$
- \left(\partial_{t} (\rho \nabla w) + \mathcal{L}_{u} (\rho \nabla w) \right) \cdot \zeta \right] dxdt
$$

\n
$$
= \int_{0}^{T} \int_{\Omega} \left[\left(-\partial_{t} \frac{\partial l}{\partial u} - \mathcal{L}_{u} \frac{\partial l}{\partial u} \right) \cdot \zeta + \text{div} \left(\left(\frac{\partial l}{\partial u} \cdot \zeta + \rho D_{\delta} w \right) u \right) - \bar{D}_{t} \rho D_{\delta} w - \rho \zeta \cdot \nabla D_{t} w \right] dxdt
$$

\n
$$
= \int_{0}^{T} \int_{\Omega} \left[\left(-\partial_{t} \frac{\partial l}{\partial u} - \mathcal{L}_{u} \frac{\partial l}{\partial u} + \rho \nabla \frac{\partial l}{\partial \rho} \right) \cdot \zeta - \bar{D}_{t} \rho D_{\delta} w \right] dxdt
$$

\n
$$
+ \int_{0}^{T} \int_{\partial \Omega} \left[\left(\frac{\partial l}{\partial u} \cdot \zeta \right) (u \cdot n) + \rho u \cdot n D_{\delta} w \right] dadt,
$$

where we used [\(2.23\)](#page-19-0), as well as ζ and δw vanishing at $t = 0, T$. Hence, when [\(2.21\)](#page-18-2) is considered, and by collecting the terms proportional to ζ , $D_{\delta}w$, and $D_{\delta}\gamma$, both at the interior and at the boundary, and using the equalities $D_t w = -\frac{\partial l}{\partial \rho}$, $D_t \gamma = -\frac{\partial l}{\partial s}$ arising from the variations $\delta \rho$ and δs , we get the six equations [\(2.22\)](#page-18-3).

Remark 2.10 (Lagrangian reduction by symmetry). Let us explain how the variational setting employed above fits in the Lagrangian reduction framework. In general, given a configuration manifold Q and a Lagrangian $L: TQ \to \mathbb{R}$, we assume that the Lagrangian is invariant under the (tangent lifted) action of a Lie group action of G on Q . In the case when the quotient $(TQ)/G$ is a smooth manifold, L induces a reduced Lagrangian $\ell : (TQ)/G \to \mathbb{R}$ and the Hamilton principle for the action functional of L induces a variational principle for the action functional of ℓ . The so called reduced principle is in general more involved than the Hamilton principle since it uses constrained variations, namely, it considers only the variations of the curves in $(TQ)/G$ that are induced by the free variations of the curves in Q. The same idea extends easily to the case with forcing, i.e., the Lagrange-d'Alembert principle, when the force is equivariant with respect to G . For the case of open fluid, the configuration manifold is originally $\text{Diff}(\mathbb{R}^n) \times \Omega^n(\mathbb{R}^n) \times \Omega^n(\mathbb{R}^n) \ni (\varphi, \varrho, S)$, but we extend it to

$$
Q = \text{Diff}(\mathbb{R}^n) \times \Omega^n(\mathbb{R}^n) \times \Omega^n(\mathbb{R}^n) \times \Omega^0(\mathbb{R}^n) \times \Omega^0(\mathbb{R}^n) \ni q = (\varphi, \varrho, S, W, \Gamma)
$$

in order to include the thermodynamic displacements W and Γ. The Lagrangian $L(q, \dot{q})$ is

$$
L(q, \dot{q}) = \int_{\varphi^{-1}(\Omega)} \left[\mathfrak{L}(\varphi, \dot{\varphi}, \nabla \varphi, \varrho) + \varrho \dot{W} + S\dot{\Gamma} \right] dX dt
$$

and the force $F(q, \dot{q})$ is given by

$$
\langle F(q,\dot{q}),\delta q\rangle = \int_{\varphi^{-1}(\Omega)} \left[\mathfrak{B} \cdot \delta \varphi + \Theta_{\varrho} \delta W + \Theta_S \delta \Gamma \right] dX + \int_{\varphi^{-1}(\partial \Omega)} \left[\mathfrak{J} \cdot \delta \varphi + \mathfrak{j}_{\varrho} \delta W + \mathfrak{j}_S \delta \Gamma \right] dA.
$$

This identification explicitly shows how (2.5) emerges from (1.2) . Now the symmetry Lie group is $G = \text{Diff}(\mathbb{R}^n)$ and it acts on Q by the right action

$$
q = (\varphi, \varrho, S, W, \Gamma) \mapsto q \cdot \psi = (\varphi \circ \psi, (\varrho \circ \psi) J\psi, (S \circ \psi) J\psi, W \circ \psi, \Gamma \circ \psi),
$$

thereby leading to the Eulerian variables we have used above.

Balance laws in the Eulerian description. From the local balance of mass and entropy and the last two boundary conditions in [\(2.22\)](#page-18-3), the corresponding global balance equations are

$$
\frac{d}{dt} \int_{\Omega} \rho dx = \int_{\Omega} \theta_{\rho} dx + \int_{\partial \Omega} j_{\rho} da \quad \text{and} \quad \frac{d}{dt} \int_{\Omega} s dx = \int_{\Omega} \theta_s dx + \int_{\partial \Omega} j_s da, \quad (2.24)
$$

showing the transfer of mass and entropy via bulk and boundary contributions.

We now consider the total energy density, defined for a general Lagrangian density l as

$$
\mathbf{e} = \frac{\partial \mathbf{I}}{\partial u} \cdot u - \mathbf{I}(u, \rho, s) \tag{2.25}
$$

and we compute the local energy balance

$$
\partial_t \mathbf{e} = \partial_t \frac{\partial \mathbf{I}}{\partial u} \cdot u - \frac{\partial \mathbf{I}}{\partial \rho} \cdot \partial_t \rho - \frac{\partial \mathbf{I}}{\partial s} \cdot \partial_t s \n= -\mathcal{L}_u \frac{\partial \mathbf{I}}{\partial u} \cdot u + \rho u \cdot \nabla \frac{\partial \mathbf{I}}{\partial \rho} + su \cdot \nabla \frac{\partial \mathbf{I}}{\partial s} + b \cdot u + \frac{\partial \mathbf{I}}{\partial \rho} \operatorname{div}(\rho u) - \theta_\rho \frac{\partial \mathbf{I}}{\partial \rho} + \frac{\partial \mathbf{I}}{\partial s} \operatorname{div}(su) - \theta_s \frac{\partial \mathbf{I}}{\partial s} \n= \operatorname{div} \left(-\left(\frac{\partial \mathbf{I}}{\partial u} \cdot u \right) u + \rho \frac{\partial \mathbf{I}}{\partial \rho} u + s \frac{\partial \mathbf{I}}{\partial s} u \right) + b \cdot u - \theta_\rho \frac{\partial \mathbf{I}}{\partial \rho} - \theta_s \frac{\partial \mathbf{I}}{\partial s}.
$$
\n(2.26)

From this, using all three boundary conditions in [\(2.22\)](#page-18-3), the global energy balance follows as

$$
\frac{d}{dt} \int_{\Omega} \mathbf{e} \, dx = \int_{\Omega} \left[b \cdot u - \theta_{\rho} \frac{\partial \mathbf{I}}{\partial \rho} - \theta_{s} \frac{\partial \mathbf{I}}{\partial s} \right] dx + \int_{\partial \Omega} \left[J \cdot u - j_{\rho} \frac{\partial \mathbf{I}}{\partial \rho} - j_{s} \frac{\partial \mathbf{I}}{\partial s} \right] da,
$$
\n(2.27)

showing the energy transfer between the fluid and its exterior via both bulk and boundary contributions. We note that from the boundary conditions in (2.22) we have the following relations at the boundary

$$
j_s = \frac{s}{\rho} j_\rho
$$
 and $J = j_\rho \frac{1}{\rho} \frac{\partial \mathfrak{l}}{\partial u} = j_s \frac{1}{s} \frac{\partial \mathfrak{l}}{\partial u}$ on $\partial \Omega$,

hence the boundary contribution to the energy balance in [\(2.27\)](#page-21-0) can be written in either of the following forms

$$
\int_{\partial\Omega} \left[j_{\rho} \left(\frac{1}{\rho} \frac{\partial \mathfrak{l}}{\partial u} \cdot u - \frac{\partial \mathfrak{l}}{\partial \rho} \right) - j_{s} \frac{\partial \mathfrak{l}}{\partial s} \right] da = \int_{\partial\Omega} j_{\rho} \left(\frac{1}{\rho} \frac{\partial \mathfrak{l}}{\partial u} \cdot u - \frac{\partial \mathfrak{l}}{\partial \rho} - \frac{s}{\rho} \frac{\partial \mathfrak{l}}{\partial s} \right) da
$$
\n
$$
= \int_{\partial\Omega} \left(\rho \frac{\partial \mathfrak{l}}{\partial \rho} + s \frac{\partial \mathfrak{l}}{\partial s} - u \cdot \frac{\partial \mathfrak{l}}{\partial u} \right) (u \cdot n) da. \tag{2.28}
$$

The compressible Euler fluid with open boundaries. Let us consider the Lagrangian density of compressible Euler fluid as given in [\(2.9\)](#page-14-1). One can readily checks that it satisfies the material covariance assumption (2.15) . In the Eulerian description the Lagrangian density takes the standard form

$$
\mathfrak{l}(u,\rho,s) = \frac{1}{2}\rho|u|^2 - \varepsilon(\rho,s),\tag{2.29}
$$

as it is seen from (2.16) . In this case the system (2.22) becomes

$$
\begin{cases}\n\partial_t(\rho u) + \operatorname{div}(\rho u \otimes u) = -\nabla p + b \\
\partial_t \rho + \operatorname{div}(\rho u) = \theta_\rho, \qquad \partial_t s + \operatorname{div}(s u) = \theta_s \\
(u \cdot n)\rho u = -J \quad \text{on} \quad \partial\Omega \\
\rho u \cdot n = -j_\rho, \qquad su \cdot n = -j_s \quad \text{on} \quad \partial\Omega.\n\end{cases}
$$
\n(2.30)

We note that when written in terms of the velocity, the momentum equation acquires an additional term due to the presence of the distributed source of mass θ_{ρ} , namely,

$$
\rho(\partial_t u + u \cdot \nabla u) = -\nabla p + b - \theta_\rho u. \tag{2.31}
$$

By using $\frac{\partial \mathfrak{l}}{\partial \rho} = \frac{1}{2}$ $\frac{1}{2}|u|^2-g$ and $\frac{\partial \ell}{\partial s}=-T$ with $g=\frac{\partial \varepsilon}{\partial \epsilon}=(\varepsilon+p-sT)/\rho$ the Gibbs potential and $T = \frac{\partial \varepsilon}{\partial s}$ the temperature, the local energy balance [\(2.26\)](#page-20-1) becomes

$$
\partial_t \mathfrak{e} + \mathrm{div}((\mathfrak{e} + p)u) = b \cdot u + \theta_\rho \Big(g - \frac{1}{2}|u|^2\Big) + \theta_s T,
$$

giving the total energy balance [\(2.27\)](#page-21-0) in the form.

$$
\frac{d}{dt} \int_{\Omega} \mathbf{e} \, dx = \int_{\Omega} \left[b \cdot u + \theta_{\rho} \left(g - \frac{1}{2} |u|^2 \right) + \theta_s T \right] dx + \int_{\partial \Omega} \left[J \cdot u + j_{\rho} \left(g - \frac{1}{2} |u|^2 \right) + j_s T \right] da. \tag{2.32}
$$

Following [\(2.28\)](#page-21-1), other useful expressions for the boundary contribution to energy change are given by

$$
\int_{\partial\Omega} \Big[j_{\rho} \Big(\frac{1}{2} |u|^2 + g \Big) + j_s T \Big] da = \int_{\partial\Omega} j_{\rho} \Big(\frac{1}{2} |u|^2 + h \Big) da = - \int_{\partial\Omega} (\mathfrak{e} + p) u \cdot n da,
$$

with $h = (\varepsilon + p)/\rho$ the enthalpy. One of these expressions may be useful depending on whether it is desirable to have the boundary sources j_{ρ} and j_{s} explicitly appear or not.

Apart from the overall energy balance, in the case of Euler fluid, we can calculate the balance law for the kinetic energy density $\mathfrak{k} = \frac{1}{2}$ $\frac{1}{2}\rho |u|^2$ and the internal energy density ε . Using the equations, we get

$$
\partial_t \mathbf{\hat{t}} + \text{div}(\mathbf{\hat{t}}u) = -u \cdot \nabla p + b \cdot u - \theta_\rho \frac{1}{2} |u|^2
$$

$$
\partial_t \varepsilon + \text{div}(\varepsilon u) = -p \text{div } u + \theta_\rho g + \theta_s T.
$$

Then, from the boundary conditions the global balance equations become

$$
\frac{d}{dt} \int_{\Omega} \mathbf{t} \, dx = \int_{\Omega} \Big[-u \cdot \nabla p + b \cdot u - \theta_{\rho} \frac{1}{2} |u|^2 \Big] dx + \int_{\partial \Omega} j_{\rho} \frac{1}{2} |u|^2 da
$$

$$
\frac{d}{dt} \int_{\Omega} \varepsilon \, dx = \int_{\Omega} \Big[u \cdot \nabla p + \theta_{\rho} g + \theta_{s} T \Big] dx + \int_{\partial \Omega} \Big[j_{\rho} g + j_{s} T \Big] da
$$

$$
= \int_{\Omega} \Big[u \cdot \nabla p + \theta_{\rho} g + \theta_{s} T \Big] dx + \int_{\partial \Omega} j_{\rho} h da.
$$

Adding these two equations up, we recover exactly the total energy balance law [\(2.32\)](#page-21-2), however these detailed balance equations show that in the interior of the fluid domain, the source term b contributes uniquely to the kinetic energy and the source term θ_s completely to the internal energy, while the source θ_{ρ} splits in two parts $-\theta_{\rho} \frac{1}{2}$ $\frac{1}{2}|u|^2$ for the kinetic energy contribution and $\theta_{\rho}g$ for the internal energy contribution. A similar situation occurs on the boundary. The term $\int_{\Omega} u \cdot \nabla p \,dx$ determines the exchange rate of energy between the kinetic and the internal energy.

Effect of rotation and gravity. In the case of a rotating fluid subject to gravity, the Lagrangian density is

$$
I(u, \rho, s) = \frac{1}{2}\rho |u|^2 + \rho R \cdot u - \varepsilon(\rho, s) - \rho \phi.
$$

Here $R(x)$ is the vector potential for the Coriolis parameter, i.e., curl $R = 2\omega$ with ω the angular velocity of the Earth, and $\phi(x)$ is the gravitational potential. In this case, system (2.22) becomes

$$
\begin{cases}\n\partial_t(\rho u) + \operatorname{div}(\rho u \otimes u) + 2\omega \times \rho u = -\nabla p - \rho \nabla \phi + b - \theta_\rho R \\
\partial_t \rho + \operatorname{div}(\rho u) = \theta_\rho, \qquad \partial_t s + \operatorname{div}(s u) = \theta_s \\
(u \cdot n)\rho(u + R) = -J \quad \text{on} \quad \partial\Omega \\
\rho u \cdot n = -j_\rho, \qquad su \cdot n = -j_s \quad \text{on} \quad \partial\Omega.\n\end{cases}
$$

We note the appearance of the vector potential R in the fluid momentum equation, due to the bulk source of mass θ_{ρ} and in the boundary condition for the fluid momentum, compare with the non rotating case (2.30) . The fluid momentum equation can be rewritten in a similar way with (2.31) as

$$
\rho(\partial_t u + u \cdot \nabla u + 2\omega \times u) = -\nabla p - \rho \nabla \phi + b - \theta_\rho(u + R), \tag{2.33}
$$

which also show the occurrence of R.

We note that the total energy density as defined in (2.25) becomes

$$
\mathfrak{e} = \frac{1}{2}\rho |u|^2 + \varepsilon(\rho, s) + \rho \phi,
$$

in which the contribution of rotation as been eliminated. In order to further show the impact of rotation and gravity we now list all the global and local balance equations for kinetic, internal, potential, and total energy densities:

$$
\partial_t \mathbf{\ell} + \text{div}(\mathbf{\ell}u) = -u \cdot \nabla p - \rho u \cdot \nabla \phi + b \cdot u - \theta_\rho \left(\frac{1}{2}|u|^2 + u \cdot R\right)
$$

$$
\partial_t \varepsilon + \text{div}(\varepsilon u) = -p \text{div } u + \theta_\rho g + \theta_s T
$$

$$
\partial_t (\rho \phi) + \text{div}(\rho \phi u) = \theta_\rho \phi + \rho u \cdot \nabla \phi
$$

$$
\partial_t \mathbf{\ell} + \text{div}((\mathbf{\ell} + p)u) = b \cdot u + \theta_\rho \left(g + \phi - u \cdot R - \frac{1}{2}|u|^2\right) + \theta_s T,
$$

and

$$
\frac{d}{dt} \int_{\Omega} \mathbf{t} \, d\mathbf{t} = \int_{\Omega} \Big[-u \cdot \nabla p - \rho u \cdot \nabla \phi + b \cdot u - \theta_{\rho} \Big(\frac{1}{2} |u|^2 + u \cdot R \Big) \Big] dx + \int_{\partial \Omega} j_{\rho} \frac{1}{2} |u|^2 da
$$

$$
\frac{d}{dt} \int_{\Omega} \varepsilon \, dx = \int_{\Omega} \Big[u \cdot \nabla p + \theta_{\rho} g + \theta_{s} T \Big] dx + \int_{\partial \Omega} j_{\rho} h da
$$

$$
\frac{d}{dt} \int_{\Omega} \rho \phi \, dx = \int_{\Omega} \Big[\theta_{\rho} \phi + \rho u \cdot \nabla \phi \Big] dx + \int_{\partial \Omega} j_{\rho} \phi da
$$

$$
\frac{d}{dt} \int_{\Omega} \mathbf{t} \, dx = \int_{\Omega} \Big[b \cdot u - \theta_{\rho} \Big(\frac{1}{2} |u|^2 + u \cdot R - g - \phi \Big) + \theta_{s} T \Big] dx + \int_{\partial \Omega} j_{\rho} \Big(\frac{1}{2} |u|^2 + h + \phi \Big) da.
$$

Remark 2.11 (Absolute Momentum vs Mass Flux ρu). A major insight of our variational approach is the clear distinction between the absolute fluid momentum $\frac{\partial l}{\partial u}$, and the mass flux ρu . We see that both of them appear in the expression of the boundary conditions for the general open fluid equations in [\(2.22\)](#page-18-3), namely:

$$
(u \cdot n)\frac{\partial \mathfrak{l}}{\partial u} = -J
$$
 and $\rho u \cdot n = -j_{\rho}$,

(instead of $\frac{\partial l}{\partial u} \cdot n = -j_\rho$ for instance for the second one). These quantities coincide if l depends on u only through the usual expression $\frac{1}{2}\rho |u|^2$ which is the case for nonrotating Euler, amongst other equations, but is not true in the general case. When they do coincide we have the relation j ^{μ} = J on the boundary, which is not true in general, for example the rotating case giving $j_{\rho}(u+R)=J.$

Remark 2.12 (Choice of the rotational vector potential). For open fluids, the choice of the rotational vector potential R such that curl $R = 2\omega$ impacts both the choice of the distributed

momentum source b and mass source j_{ρ} , since it changes the relationship between velocity u and absolute momentum $\frac{\partial l}{\partial u}$ by a factor of ρR . Therefore, when using $R' = R + \nabla f$ for some function f , one needs to accordingly make the changes

$$
b \to b' = b - \theta_{\rho} \nabla f
$$
 and $J \to J' = J - j_{\rho} \nabla f$

in the distributed and boundary source of fluid momentum, to have the physics unchanged.

The rotating shallow water equations. The Lagrangian density for the rotating shallow water equations is

$$
I(u, h) = \frac{1}{2}h|u|^2 + hR \cdot u - \frac{1}{2}g(h+Z)^2
$$

with $Z(x)$ the bottom topography and h the water depth, so that the surface level is $h + Z$. The associated open fluid equations found from [\(2.22\)](#page-18-3) are

$$
\begin{cases}\n\partial_t(hu) + \operatorname{div}(hu \otimes u) + 2\omega \times hu = -gh\nabla(h+Z) + b - \theta_h R \\
\partial_t h + \operatorname{div}(hu) = \theta_h \\
h(u \cdot n)(u+R) = -J \quad \text{on} \quad \partial\Omega \\
hu \cdot n = -j_h \quad \text{on} \quad \partial\Omega.\n\end{cases}
$$

The fluid momentum equation can be written as

$$
h(\partial_t u + u \cdot \nabla u + 2\omega \times u) = -gh\nabla(h+Z) - \theta_h(u+R) + b,
$$

showing again the role of the source term θ_h . Distributed source terms θ_h in the continuity equation of shallow water models appear, for instance, in the modeling of moist shallow water equations, as discussed in [\[40\]](#page-49-10).

The total energy density as defined in [\(2.25\)](#page-20-2) is now

$$
\mathfrak{e} = \frac{1}{2}h|u|^2 + \frac{1}{2}g(h+Z)^2,
$$

in which the contribution of rotation as been eliminated. The list of global and local balance equations for kinetic, potential $p = \frac{1}{2}$ $\frac{1}{2}g(h+Z)^2$, and total energy densities reads as follows:

$$
\partial_t \mathbf{\ell} + \text{div}(\mathbf{\ell}u) = -ghu \cdot \nabla(h+Z) + b \cdot u - \theta_h \left(\frac{1}{2}|u|^2 + u \cdot R\right)
$$

$$
\partial_t \mathbf{p} + \text{div}(\mathbf{p}u) = \text{div}\left(\frac{1}{2}g(Z^2 - h^2)u\right) + ghu \cdot \nabla(h+Z) + g(h+Z)\theta_h
$$

$$
\partial_t \mathbf{e} + \text{div}\left(\mathbf{e}u + \frac{1}{2}g(h^2 - Z^2)u\right) = b \cdot u + \theta_h \left(g(h+Z) - u \cdot R - \frac{1}{2}|u|^2\right)
$$

,

and

$$
\frac{d}{dt} \int_{\Omega} \mathbf{t} \, dx = \int_{\Omega} \Big[-ghu \cdot \nabla (h+Z) + b \cdot u - \theta_h \Big(\frac{1}{2} |u|^2 + R \cdot u \Big) \Big] dx + \int_{\partial \Omega} j_h \frac{1}{2} |u|^2 da
$$

$$
\frac{d}{dt} \int_{\Omega} \mathbf{p} \, dx = \int_{\Omega} \Big[ghu \cdot \nabla (h+Z) + \theta_h g(h+Z) \Big] dx + \int_{\partial \Omega} j_h g(h+Z) da
$$

$$
\frac{d}{dt} \int_{\Omega} \mathbf{e} \, dx = \int_{\Omega} \Big[b \cdot u - \theta_h \Big(\frac{1}{2} |u|^2 + R \cdot u - g(h+Z) \Big) \Big] dx + \int_{\partial \Omega} j_h \Big(\frac{1}{2} |u|^2 + g(h+Z) \Big) da.
$$

On the choice of the fluxes and inflow/outflow boundary conditions. We now focus on the non rotating case and analyse how our setting allows the treatment of inflow and outflow boundary conditions. For this discussion we consider the Lagrangian [\(2.29\)](#page-21-5) and we assume $\rho > 0$ everywhere. The boundary conditions take the following form:

$$
(u \cdot n)\rho u = -J, \quad \rho u \cdot n = -j_\rho, \quad su \cdot n = -j_s.
$$

(i) If we first assume $j_{\rho} = 0$ then we have $u \cdot n = 0$ on $\partial \Omega$ which corresponds to an impermeable boundary. In this case, the relations above impose the choice $j_s = 0$ and $J = 0$ and there are no boundary conditions for $\rho|_{\partial\Omega}$ and $s|_{\partial\Omega}$, consistently with the treatment of fluids with impermeable boundaries. It is interesting to note that, while our geometric setting is primarily designed for open fluid motion, it allows the treatment of closed fluids by appropriate choice of J, j_ρ, j_s .

(ii) If $j_{\rho} \neq 0$, then $u \cdot n \neq 0$ and J must satisfy $J \cdot n \neq 0$. We get the following boundary *relations*

$$
u|_{\partial\Omega} = \frac{J}{j_{\rho}}, \quad \rho|_{\partial\Omega} = -\frac{j_{\rho}^2}{J \cdot n}, \quad s|_{\partial\Omega} = -\frac{j_{\rho}j_s}{J \cdot n}.
$$

In particular, we note the relations $j_s = \frac{\rho}{s}$ $\frac{\rho}{s}j_{\rho}$ and $J \cdot n < 0$ on the boundary. Let us show that these relations do yield appropriate boundary *conditions* for inflow and outflow.

(ii.a) The case $j_{\rho} > 0$ corresponds to inflow: $u \cdot n < 0$. In this situation J, j_{ρ} and j_{s} can be freely chosen, independently on the current fluid motion, so that $u|_{\partial\Omega}$, $\rho|_{\partial\Omega}$, $s|_{\partial\Omega}$ are all prescribed on the boundary, consistently with the treatment of inflow. Equivalently, $u|_{\partial\Omega}$, $\rho|_{\partial\Omega}$, $T|_{\partial\Omega}$ are all prescribed, with T the temperature.

(ii.b) The case $j_{\rho} < 0$ corresponds to outflow: $u \cdot n > 0$. In this situation, we choose

$$
j_{\rho} = -\rho u_0 \cdot n \quad \text{and} \quad J = -(u_0 \cdot n)\rho u_0,
$$

where we recall that a dependence of j_ρ on the current fluid variables (here ρ) is allowed in our approach, see §[2.2,](#page-16-0) and where u_0 is a prescribed velocity field on $\partial\Omega$ with $u_0 \cdot n > 0$. In this case the first two boundary relations just give one boundary condition

$$
u|_{\partial\Omega}=u_0,
$$

consistently with the treatment of outflow boundary conditions for which there is no prescription on $\rho|_{\partial\Omega}$. The temperature can be prescribed at the outflow boundary by choosing

$$
j_s = -s(\rho, T_0)u_0 \cdot n,
$$

where the state equation is used to write s in terms of ρ and T_0 . With these choices, we see that the three boundary relations above yield the two boundary conditions $u|_{\partial\Omega} = u_0$ and $T|_{\partial\Omega} = T_0$, consistently with what is assumed for outflow in the viscous case, see, e.g., [\[41](#page-49-11)]. In the inviscid case, we choose

$$
j_{\rho} = -\rho \nu_0
$$
, $j_s = -s\nu_0$, and $J = -\nu_0 \rho u$,

for $\nu_0 > 0$. With these choices, we see that the three boundary relations above yield the single boundary condition

$$
u\cdot n|_{\partial\Omega}=\nu_0,
$$

consistently with what is assumed for outflow in the non-viscous case, see, e.g., [\[41](#page-49-11)].

While the boundary conditions as detailed above are used for well-posedness questions, [\[41](#page-49-11)], in practice the number of boundary conditions at the inflow and outflow depend on the (subsonic/supersonic) fluid regime. Appropriate choices of J , j_{ρ} , and j_{s} can also handle these cases.

3 Hamiltonian variational and bracket formulations.

In this section we present the Hamiltonian counterpart of the approach presented in §[2.](#page-10-0) This approach is useful for deriving the appropriate extension of the Lie-Poisson bracket in fluid dynamics that can accommodate open boundary conditions. As in the previous section, we start with the material description as it is in this description that the variational principle is simpler. The Eulerian version is then derived through reduction. We do not include detailed computations here, as they can be carried out in parallel with those on the Lagrangian side.

We are using the word Hamiltonian in this section, since the setting is based on using Hamiltonian functions and densities, rather than Lagrangian ones. However, the systems under consideration, being open, are *not Hamiltonian systems* and, consequently, we do not aim to find a Poisson bracket for this system, but to describe a natural extension of the Lie-Poisson bracket that accounts for the bulk and boundary fluxes.

3.1 Material formulation

Start by assuming the existing of a Hamiltonian function $H: T^*Diff(\mathbb{R}^n) \times \Omega^n(\mathbb{R}^n) \times \Omega^n(\mathbb{R}^n) \to$ R and Hamiltonian density $\mathfrak{H}(\varphi, M, \nabla \varphi, \varrho, S)$, related by

$$
H(\varphi, M, \varrho, S) := \int_{\varphi^{-1}(\Omega)} \mathfrak{H}(\varphi, M, \nabla \varphi, \varrho, S) \, dX. \tag{3.1}
$$

The next result is the Hamiltonian analog to [\(2.5\)](#page-12-0) and Proposition [2.4.](#page-12-2)

Proposition 3.1. *The Hamilton-d'Alembert phase-space principle*

$$
\frac{d}{d\varepsilon}\Big|_{\varepsilon=0} \int_0^T \int_{\varphi_{\varepsilon}^{-1}(\Omega)} \Big[M_{\varepsilon} \cdot \dot{\varphi}_{\varepsilon} + \varrho_{\varepsilon} \dot{W}_{\varepsilon} + S_{\varepsilon} \dot{\Gamma}_{\varepsilon} - \mathfrak{H}(\varphi_{\varepsilon}, M_{\varepsilon}, \nabla \varphi_{\varepsilon}, \varrho_{\varepsilon}, S_{\varepsilon}) \Big] dX dt + \int_0^T \int_{\varphi^{-1}(\Omega)} (\mathfrak{B} \cdot \delta \varphi + \Theta_{\varrho} \delta W + \Theta_S \delta \Gamma) dX dt \leftarrow bulk \iota contribution + \int_0^T \int_{\partial(\varphi^{-1}(\Omega))} (\mathfrak{J} \cdot \delta \varphi + \mathfrak{j}_{\varrho} \delta W + \mathfrak{j}_S \delta \Gamma) dA dt = 0 \leftarrow boundary \iota contribution
$$
\n(3.2)

with free variations δM , $\delta \varphi$, $\delta \varrho$, δW , δS , $\delta \Gamma$, such that $\delta \varphi$, δW , and $\delta \Gamma$ vanish at $t = 0, T$, *gives the following equations:*

$$
\begin{cases}\n\frac{d}{dt}\varphi = \frac{\partial \mathfrak{H}}{\partial M}, & \frac{d}{dt}M - \text{DIV} \frac{\partial \mathfrak{H}}{\partial \nabla \varphi} + \frac{\partial \mathfrak{H}}{\partial \varphi} = \mathfrak{B} \\
\frac{d}{dt}\varrho = \Theta_{\varrho}, & \frac{d}{dt}S = \Theta_{S} \\
N \cdot \left(-\frac{\partial \mathfrak{H}}{\partial \nabla \varphi} + (\nabla_{x}\varphi^{-1} \circ \varphi) \cdot \frac{\partial \mathfrak{H}}{\partial M} M + \left(\mathfrak{H} - M \cdot \frac{\partial \mathfrak{H}}{\partial M} - \varrho \frac{\partial \mathfrak{H}}{\partial \varrho} - S \frac{\partial \mathfrak{H}}{\partial S} \right) \nabla_{x}\varphi^{-1} \circ \varphi \right) = -\mathfrak{J} \quad on \quad \partial \Omega \\
N \cdot \left((\nabla_{x}\varphi^{-1} \circ \varphi) \cdot \frac{\partial \mathfrak{H}}{\partial M} \varrho \right) = -\mathfrak{j}_{\varrho}, \quad N \cdot \left((\nabla_{x}\varphi^{-1} \circ \varphi) \cdot \frac{\partial \mathfrak{H}}{\partial M} S \right) = -\mathfrak{j}_{S} \quad on \quad \partial \Omega,\n\end{cases} \tag{3.3}
$$

Remark 3.2 (Hamilton-d'Alembert phase space principle). The variational principle [\(3.2\)](#page-26-2) is an instance of the Hamilton-d'Alembert phase space principle, given in general as

$$
\delta \int_0^T \left[\langle p, \dot{q} \rangle - H(q, p) \right] dt + \int_0^T \langle F(q, \dot{q}), \delta q \rangle dt = 0,
$$

for a Hamiltonian $H: T^*Q \to \mathbb{R}$ and a force $F: TQ \to T^*Q$. It is the Hamiltonian counterpart of the Lagrange-d'Alembert principle [\(1.2\)](#page-5-2).

3.1.1 Relating Hamiltonian and Lagrangian formulations

Given a Lagrangian density $\mathfrak{L}(\varphi, \dot{\varphi}, \nabla \varphi, S)$, the corresponding Hamiltonian density is obtained by the *Legendre transform*, assuming that the map

$$
\dot{\varphi} \mapsto M = \frac{\partial \mathfrak{L}}{\partial \dot{\varphi}} (\varphi, \dot{\varphi}, \nabla \varphi, \varrho, S) \tag{3.4}
$$

is a diffeomorphism for all $\varphi, \dot{\varphi}, \varrho, S$ with $\varrho > 0^5$ $\varrho > 0^5$. In this case we can define the Hamiltonian density \mathfrak{H} in the following way:

$$
\mathfrak{H}(\varphi, M, \nabla \varphi, \varrho, S) := M \cdot \dot{\varphi} - \mathfrak{L}(\varphi, \dot{\varphi}, \nabla \varphi, \varrho, S),
$$

where $\dot{\varphi}$ in the formula above is the inverse image of M by [\(3.4\)](#page-27-2). Alternatively, one could start with a Hamiltonian and obtain a Lagrangian, assuming that the *inverse Legendre transform* is a diffeomorphism. This is obviously the case for all the fluid models treated above, although there are fluid models where this is not the case and there is either a Lagrangian or a Hamiltonian formulation, but not both. For example, the hydrostatic primitive equations in Eulerian coordinates have only a Lagrangian representation $[42]$. We note the relations

$$
\frac{\partial \mathfrak{H}}{\partial M} = \dot{\varphi}, \quad \frac{\partial \mathfrak{H}}{\partial \cdot \cdot} = -\frac{\partial \mathfrak{L}}{\partial \cdot \cdot} \text{ for } \varphi, \nabla \varphi, \varrho, S.
$$

Example: the compressible Euler fluid with open boundaries. By taking the Legendre transform of the Lagrangian density [\(2.9\)](#page-14-1) of the compressible Euler fluid, we get

$$
\mathfrak{H}(\varphi, M, \nabla \varphi, \varrho, S) = \frac{|M|^2}{2\varrho} + \varepsilon \left(\frac{\varrho}{J\varphi}, \frac{S}{J\varphi} \right) J\varphi.
$$

The partial derivatives are

$$
\frac{\partial \mathfrak{H}}{\partial \varphi} = 0, \quad \frac{\partial \mathfrak{H}}{\partial M} = \frac{M}{\varrho}, \quad \frac{\partial \mathfrak{H}}{\partial \varrho} = \frac{\partial \varepsilon}{\partial \rho} - \frac{|M|^2}{2\varrho^2}, \quad \frac{\partial \mathfrak{H}}{\partial S} = \frac{\partial \varepsilon}{\partial s}
$$
\n
$$
\frac{\partial \mathfrak{L}}{\partial \nabla \varphi} = \left(\varepsilon - \frac{\partial \varepsilon}{\partial \rho} \frac{\varrho}{J\varphi} - \frac{\partial \varepsilon}{\partial s} \frac{S}{J\varphi}\right) J\varphi \left(\nabla_x \varphi^{-1} \circ \varphi\right) = -(p \circ \varphi) J\varphi \left(\nabla_x \varphi^{-1} \circ \varphi\right)
$$
\n(3.5)

 5 In other words, the Lagrangian is non-degenerate and therefore the Legendre transform is invertible.

hence the system [\(3.3\)](#page-26-3) becomes

$$
\begin{cases}\n\frac{d}{dt}\varphi = \frac{M}{\varrho}, & \frac{d}{dt}M + (\nabla_x p \circ \varphi)J\varphi = \mathfrak{B} \\
\frac{d}{dt}\varrho = \Theta_{\varrho}, & \frac{d}{dt}S = \Theta_{S} \\
N \cdot \left[(\nabla_x \varphi^{-1} \circ \varphi) \cdot \frac{M}{\varrho} \right] M = -\mathfrak{J} \quad \text{on} \quad \partial\Omega \\
N \cdot \left[(\nabla_x \varphi^{-1} \circ \varphi) \cdot \frac{M}{\varrho} \right] \varrho = -\mathfrak{j}_{\varrho}, & N \cdot \left[(\nabla_x \varphi^{-1} \circ \varphi) \cdot \frac{M}{\varrho} \right] S = -\mathfrak{j}_{S} \quad \text{on} \quad \partial\Omega.\n\end{cases}
$$
\n(3.6)

This system of equations and boundary conditions for the open compressible fluid in the material descriptions is readily seen to be equivalent to the one derived in the Lagrangian side in [\(2.11\)](#page-15-0) by writing $M = \varrho \dot{\varphi}$.

3.2 Eulerian formulation

We now assume relabelling symmetries and deduce from it the variational principle in the Eulerian description, induced by the Hamilton-d'Alembert phase space principle [\(3.3\)](#page-26-3).

Relabelling symmetries. We say that a Hamiltonian density is materially covariant it it satisfies

$$
\mathfrak{H}(\varphi \circ \psi, (M \circ \psi)J\psi, \nabla_X(\varphi \circ \psi), (\varrho \circ \psi)J\psi, (S \circ \psi)J\psi) = \mathfrak{H}(\varphi, M, \nabla_X \varphi, \varrho, S) \circ \psi J\psi, \quad (3.7)
$$

for all $\psi \in \text{Diff}(\mathbb{R}^n)$. It is clear that if a Lagrangian density $\mathfrak L$ is materially covariant, see [\(2.15\)](#page-16-2), then the associated Hamiltonian density obtained via the Legendre transform is also. The symmetry [\(3.7\)](#page-28-1) is equivalent to the existence of a Hamiltonian density $\mathfrak{h}(m, \rho, s)$ in the Eulerian description, such that

$$
\mathfrak{H}(\varphi, M, \nabla_X \varphi, \varrho, S) \circ \varphi^{-1} J \varphi^{-1} = \mathfrak{h}(m, \rho, s).
$$
\n(3.8)

where $m = (M \circ \varphi^{-1})J\varphi^{-1}, \ \rho = (\varrho \circ \varphi^{-1})J\varphi^{-1}, \ s = (S \circ \varphi^{-1})J\varphi^{-1}.$ Under the assumption [\(3.7\)](#page-28-1), the associated Hamiltonian function $H: T^* \text{Diff}(\mathbb{R}^n) \times \Omega^n(\mathbb{R}^n) \times \Omega^n(\mathbb{R}^n) \to \mathbb{R}$ is $\text{Diff}(\mathbb{R}^n)$ invariant, namely

$$
H(\varphi \circ \psi, (M \circ \psi)J\psi, (\varrho \circ \psi)J\psi, (S \circ \psi)J\psi) = H(\varphi, M, \varrho, S), \text{ for all } \psi \in \text{Diff}(\mathbb{R}^n). \tag{3.9}
$$

Therefore, it induces a unique reduced Hamiltonian h , which turns out to be associated to the Hamiltonian density in [\(3.8\)](#page-28-2)

$$
h: (\Omega^1(\mathbb{R}^n) \otimes \Omega^n(\mathbb{R}^n)) \times \Omega^n(\mathbb{R}^n) \times \Omega^n(\mathbb{R}^n) \to \mathbb{R}, \quad h(m,\rho,s) = \int_{\Omega} \mathfrak{h}(m(x),\rho(x),s(x))dx. \tag{3.10}
$$

Again, we emphasize the distinction between the function h, which is defined on forms on \mathbb{R}^n , and the density h depending on the point values of these forms on Ω only.

If the Legendre transform is invertible, the reduced Hamiltonian density can be equivalently defined in the following way :

$$
\mathfrak{h}(m,\rho,s)=m\cdot u-\mathfrak{l}(u,\rho,s)
$$

where $\mathfrak l$ is the reduced Lagrangian density and u is the inverse image of m via the Legendre transform $u \mapsto \frac{\partial \ell}{\partial u}$.

Remark 3.3 (Reduction of the phase space). We note that the passing from the Hamiltonian $\text{function } H: T^* \text{Diff}(\mathbb{R}^n) \times \Omega^n(\mathbb{R}^n) \times \Omega^n(\mathbb{R}^n) \to \mathbb{R} \text{ to its reduced version } h: (\Omega^1(\mathbb{R}^n) \otimes \Omega^n(\mathbb{R}^n)) \times$ $\Omega^n(\mathbb{R}^n) \times \Omega^n(\mathbb{R}^n) \to \mathbb{R}$, is simply understood as taking the function h induced by H on the quotient by the symmetry group $\text{Diff}(\mathbb{R}^n)$:

$$
\left[T^*\,\mathrm{Diff}(\mathbb{R}^n)\times\Omega^n(\mathbb{R}^n)\times\Omega^n(\mathbb{R}^n)\right]/\,\mathrm{Diff}(\mathbb{R}^n)\simeq\left(\Omega^1(\mathbb{R}^n)\otimes\Omega^n(\mathbb{R}^n)\right)\times\Omega^n(\mathbb{R}^n)\times\Omega^n(\mathbb{R}^n).
$$

While this reduction process is formally the same as that for a closed fluid, the key difference is that it is applied to Hamiltonians in which the domain of the fluid enter in a nontrivial way, see [\(3.1\)](#page-26-4), being found by integrating the density on $\varphi^{-1}(\Omega)$. We stress again that the relabelling symmetry group is not Diff(Ω), with Ω the fluid domain, but is the group Diff(\mathbb{R}^n).

Based on these developments, we can state the following result which is the Hamiltonian counterpart of Proposition [2.9.](#page-18-4)

Proposition 3.4 (Hamiltonian variational principle for open fluids). *Assume that the Hamiltonian density and the distributed and boundary sources of momentum, mass and entropy satisfy the relabelling symmetries. Then the following hold:*

(i) *The Hamilton-d'Alembert phase space principle* [\(3.2\)](#page-26-2) *yields the following Hamltonian variational formulation in the Eulerian frame:*

$$
\delta \int_0^T \int_{\Omega} \left[m \cdot u + \rho D_t w + s D_t \gamma - \mathfrak{h}(m, \rho, s) \right] dx dt + \int_0^T \int_{\Omega} \left[b \cdot \zeta + \theta_{\rho} D_{\delta} w + \theta_s D_{\delta} \gamma \right] dx dt \leftarrow bulk contribution + \int_0^T \int_{\partial \Omega} \left[J \cdot \zeta + j_{\rho} D_{\delta} w + j_s D_{\delta} \gamma \right] da dt = 0 \leftarrow boundary contribution
$$
\n(3.11)

with respect to variations $\delta u = \partial_t \zeta + [\zeta, u]$ *and free variations* δm *,* $\delta \rho$ *,* δs *,* δw *,* $\delta \gamma$ *, with* δw and δ $γ$ vanishing at $t = 0, T$ and $ζ$ an arbitrary time dependent vector field vanishing $at t = 0, T$.

The relation with the material variables used in [\(3.2\)](#page-26-2) *is given by*

$$
m = (M \circ \varphi^{-1})J\varphi^{-1}, \quad u = \dot{\varphi} \circ \varphi^{-1},
$$

$$
\rho = (\varrho \circ \varphi^{-1})J\varphi^{-1}, \quad s = (S \circ \varphi^{-1})J\varphi^{-1}, \quad w = W \circ \varphi^{-1}, \quad \gamma = \Gamma \circ \varphi^{-1}.
$$

(ii) *This principle yields the following equations and boundary conditions for open fluids:*

$$
\begin{cases}\n\partial_t m + \mathcal{L}_{\frac{\partial \mathfrak{h}}{\partial m}} m + \rho \nabla \frac{\partial \mathfrak{h}}{\partial \rho} + s \nabla \frac{\partial \mathfrak{h}}{\partial s} = b \\
\partial_t \rho + \text{div} \left(\rho \frac{\partial \mathfrak{h}}{\partial m} \right) = \theta_{\rho}, \quad \partial_t s + \text{div} \left(s \frac{\partial \mathfrak{h}}{\partial m} \right) = \theta_s \\
\left(\frac{\partial \mathfrak{h}}{\partial m} \cdot n \right) m = -J \quad on \quad \partial \Omega \\
\rho \frac{\partial \mathfrak{h}}{\partial m} \cdot n = -j_{\rho}, \quad s \frac{\partial \mathfrak{h}}{\partial m} \cdot n = -j_s \quad on \quad \partial \Omega.\n\end{cases} (3.12)
$$

The compressible Euler fluid with open boundaries. The reduced Hamiltonian density has the form:

$$
\mathfrak{h}(m,\rho,s)=\frac{|m|^2}{2\rho}+\varepsilon(\rho,s),
$$

from which the system [\(3.12\)](#page-29-1) gives the equations and boundary condition for the compressible fluid as follows

$$
\begin{cases}\n\partial_t m + \operatorname{div}(m \otimes \frac{m}{\rho}) = -\nabla p + b \\
\partial_t \rho + \operatorname{div}(m) = \theta_\rho, \qquad \partial_t s + \operatorname{div}(\frac{s m}{\rho}) = \theta_s \\
(\frac{m}{\rho} \cdot n)\rho u = -J \quad \text{on} \quad \partial \Omega \\
m \cdot n = -j_\rho, \qquad \frac{s m}{\rho} \cdot n = -j_s \quad \text{on} \quad \partial \Omega.\n\end{cases} (3.13)
$$

Rotating compressible fluid and shallow water equations. They can be treated in a similar way by noting the the Hamiltonian densities are given by

$$
\mathfrak{h}(m,\rho,s) = \frac{1}{2\rho^2}|m-\rho R|^2 + \varepsilon(\rho,s) + \rho\phi
$$

$$
\mathfrak{h}(m,h) = \frac{1}{2h^2}|m-hR|^2 + \frac{1}{2}g(h+Z)^2.
$$

3.3 Extended Lie-Poisson bracket formulation

In this paragraph we derive a bracket formulation for open fluids based on the above description. This formulation extends the Lie-Poisson bracket formulation of fluid dynamics([\[4](#page-47-3), [6,](#page-47-5) [5,](#page-47-4) [7,](#page-47-6) [8\]](#page-47-7)) to fluids with open boundaries.

In the context of Hamiltonian systems, the most efficient way to derive Lie-Poisson brackets is to obtain them via reduction by symmetry of a canonical Poisson bracket on the phase space of the system, given by the cotangent bundle of the configuration manifold. For fluid dynamics, this corresponds to passing from the material to the Eulerian description. This is the approach taken in $\lvert \mathcal{S} \rvert$ for compressible fluids, in which the canonical Poisson bracket on T^* Diff(Ω) corresponding to the material description, is reduced by the group of relabelling symmetries to give, in the Eulerian description, a Lie-Poisson bracket associated to a semidirect product Lie algebra, see $[7]$ for the incompressible fluid. We refer to $[37, 43]$ $[37, 43]$ $[37, 43]$ for the derivation of bracket formalisms from a variational perspective for irreversible processes.

We shall take the same approach here, by deriving the bracket formulation associated to the material and Eulerian descriptions described in §[3.1](#page-26-1) and §[3.2](#page-28-0) via reduction by symmetries. This results in a suitable extension of the Lie-Poisson bracket. As a particular case of our bracket formulation, we derive derives the "bulk+boundary" bracket formulation proposed in $|22|$, see also $|23|$.

Bracket formulation in the material description. In order to get the bracket formulation in the material description, we shall focus on functionals having the same form as the Hamiltonian for open fluids, namely $F: T^* \text{Diff}(\mathbb{R}^n) \times \Omega^n(\mathbb{R}^n) \times \Omega^n(\mathbb{R}) \to \mathbb{R}$ given in terms of

a density as follows

$$
F(\varphi, M, \varrho, S) = \int_{\varphi^{-1}(\Omega)} \mathfrak{F}(\varphi, M, \nabla \varphi, \varrho, S) dX,
$$
\n(3.14)

for some fixed domain $\Omega \subset \mathbb{R}^n$. In a similar way with the Poisson formulation $\dot{F} = \{F, H\}$ of Hamiltonian systems, our goal is to find an expression for the time derivative of any functional F of the form [\(3.14\)](#page-31-0) along a curve in T^* Diff(\mathbb{R}^n) $\times \Omega^n(\mathbb{R}^n) \times \Omega^n(\mathbb{R})$, which extends $\{F, H\}$ and completely characterizes the evolution equations and boundary conditions of the system [\(3.3\)](#page-26-3). In order to get this expression, we compute the evolution of such functionals along the solution of [\(3.3\)](#page-26-3), thereby giving

$$
\frac{d}{dt}F = \int_{\varphi^{-1}(\Omega)} \left[\left(\frac{\partial \mathfrak{F}}{\partial \varphi} - \text{DIV} \frac{\partial \mathfrak{F}}{\partial \nabla \varphi} \right) \cdot \frac{\partial \mathfrak{H}}{\partial M} - \left(\frac{\partial \mathfrak{H}}{\partial \varphi} - \text{DIV} \frac{\partial \mathfrak{H}}{\partial \nabla \varphi} \right) \cdot \frac{\partial \mathfrak{F}}{\partial M} \right] dX \tag{3.15}
$$

$$
+\int_{\varphi^{-1}(\Omega)} \left[\frac{\partial \mathfrak{F}}{\partial M} \cdot \mathfrak{B} + \frac{\partial \mathfrak{F}}{\partial \varrho} \cdot \Theta_{\varrho} + \frac{\partial \mathfrak{F}}{\partial S} \cdot \Theta_{S}\right] dX\tag{3.16}
$$

$$
+\int_{\varphi^{-1}(\partial\Omega)}\frac{\partial\mathfrak{H}}{\partial M}\cdot\left(\frac{\partial\mathfrak{F}}{\partial\nabla\varphi}-\mathfrak{F}\left(\nabla_x\varphi^{-1}\circ\varphi\right)\right)\cdot N\mathrm{d}A.\tag{3.17}
$$

We now multiply the three boundary conditions in [\(3.3\)](#page-26-3) by, respectively, $\frac{\partial \mathfrak{F}}{\partial M}$, $\frac{\partial \mathfrak{F}}{\partial \varrho}$, and $\frac{\partial \mathfrak{F}}{\partial S}$, integrate them on $\varphi^{-1}(\partial\Omega)$ and add them to the above expression. The boundary integral [\(3.17\)](#page-31-1) then remarkably combines with these added terms to yield the expression

$$
\frac{d}{dt}F = \int_{\varphi^{-1}(\Omega)} \left[\left(\frac{\partial \mathfrak{F}}{\partial \varphi} - \text{DIV} \frac{\partial \mathfrak{F}}{\partial \nabla \varphi} \right) \cdot \frac{\partial \mathfrak{H}}{\partial M} - \left(\frac{\partial \mathfrak{H}}{\partial \varphi} - \text{DIV} \frac{\partial \mathfrak{H}}{\partial \nabla \varphi} \right) \cdot \frac{\partial \mathfrak{F}}{\partial M} \right] dX \tag{3.18}
$$

$$
+ \int_{\varphi^{-1}(\partial\Omega)} \frac{\partial \mathfrak{H}}{\partial M} \cdot \left(\frac{\partial \mathfrak{F}}{\partial \nabla \varphi} + \left(\frac{\partial \mathfrak{F}}{\partial M} \cdot M + \frac{\partial \mathfrak{F}}{\partial \varrho} \varrho + \frac{\partial \mathfrak{F}}{\partial S} S - \mathfrak{F} \right) \nabla_x \varphi^{-1} \circ \varphi \right) \cdot N \mathrm{d}A \quad (3.19)
$$

$$
- \int_{\varphi^{-1}(\partial\Omega)} \frac{\partial \mathfrak{F}}{\partial M} \cdot \left(\frac{\partial \mathfrak{H}}{\partial \nabla \varphi} + \left(\frac{\partial \mathfrak{H}}{\partial M} \cdot M + \frac{\partial \mathfrak{H}}{\partial \varrho} \varrho + \frac{\partial \mathfrak{H}}{\partial S} S - \mathfrak{H} \right) \nabla_x \varphi^{-1} \circ \varphi \right) \cdot N \mathrm{d}A \quad (3.20)
$$

$$
+\underbrace{\int_{\varphi^{-1}(\Omega)} \left[\frac{\partial \mathfrak{F}}{\partial M} \cdot \mathfrak{B} + \frac{\partial \mathfrak{F}}{\partial \varrho} \cdot \Theta_{\varrho} + \frac{\partial \mathfrak{F}}{\partial S} \cdot \Theta_{S}\right] dX}_{\tag{3.21}
$$

bulk contribution
+
$$
\underbrace{\int_{\varphi^{-1}(\partial\Omega)} \left[\frac{\partial \mathfrak{F}}{\partial M} \cdot \mathfrak{J} + \frac{\partial \mathfrak{F}}{\partial \varrho} j_{\varrho} + \frac{\partial \mathfrak{F}}{\partial S} j_{S} \right] dA}_{\text{boundary contribution}}
$$
(3.22)

A direct verification then shows that the previous equality holding for all F is equivalent to the system (3.3) . The two terms (3.21) and (3.22) contain the contribution of each of the bulk and boundary source terms, which are paired with the corresponding partial derivative of \mathfrak{F} on the domain and on the boundary. The nontrivial appearance of the boundary terms [\(3.19\)](#page-31-4) and [\(3.20\)](#page-31-5) is due to the domain of integration being a variable. Remarkably these two terms vanish for materially covariant densities \mathfrak{F} and \mathfrak{H} , see Remark [3.5,](#page-32-0) which is always the case for fluid dynamics, thereby yielding

$$
\frac{d}{dt}F = (3.18) + (3.21) + (3.22)
$$

for materially covariant densities. While the bracket (3.18) – (3.20) takes a somehow involved form which deserves further investigations of its geometric properties, it takes the usual Lie-Poisson form in the Eulerian description, as we shall show.

Remark 3.5 (Property of materially covariant Hamiltonian densities). When a Hamiltonian density $\mathfrak H$ is materially covariant, see [\(3.7\)](#page-28-1), then its partial derivatives satisfy:

$$
\frac{\partial \mathfrak{H}}{\partial \nabla \varphi} + \left(\frac{\partial \mathfrak{H}}{\partial \varrho} \varrho + \frac{\partial \mathfrak{H}}{\partial S} S + \frac{\partial \mathfrak{H}}{\partial M} \cdot M - \mathfrak{H} \right) \nabla_x \varphi^{-1} \circ \varphi = 0.
$$

This property is similar with the one mentioned in Remark [2.8](#page-17-0) for Lagrangian densities, and is responsible for the cancellation of (3.19) and (3.20) in the materially covariant case.

Bracket formulation in the Eulerian description. As we have seen earlier for the Hamil-tonian, see [\(3.8\)](#page-28-2) and [\(3.9\)](#page-28-3), when the density \mathfrak{F} in [\(3.14\)](#page-31-0) is materially covariant, the functional F is Diff(\mathbb{R}^n)-invariant, and induces a functional $f: (\Omega^1(\mathbb{R}^n) \otimes \Omega^n(\mathbb{R}^n)) \times \Omega^n(\mathbb{R}^n) \times \Omega^n(\mathbb{R}^n) \to \mathbb{R}$ of the form

$$
f(m, \rho, s) = \int_{\Omega} \mathfrak{f}(m(x), \rho(x), s(x)) \mathrm{d}x,
$$

see Remark [3.3.](#page-29-0) By taking the time derivative of such functionals on the solutions of system [\(3.12\)](#page-29-1), we get the following result.

Proposition 3.6. *The system* [\(3.12\)](#page-29-1) *can be equivalently written in the extended Lie-Poisson form*

$$
\frac{d}{dt}f = \{f, h\}_{\text{LP}} + \underbrace{\int_{\Omega} \left(b \cdot \frac{\partial f}{\partial m} + \theta_{\rho} \frac{\partial f}{\partial \rho} + \theta_{s} \frac{\partial f}{\partial s} \right) dx}_{\text{bulk contribution}} + \underbrace{\int_{\partial \Omega} \left(J \cdot \frac{\partial f}{\partial m} + j_{\rho} \frac{\partial f}{\partial \rho} + j_{s} \frac{\partial f}{\partial s} \right) da}_{\text{boundary contribution}}, \quad \text{for all } f, \tag{3.23}
$$

where

$$
\{f, h\}_{\text{LP}} = \int_{\Omega} m \cdot \left[\frac{\partial f}{\partial m}, \frac{\partial h}{\partial m} \right] dx + \int_{\Omega} \rho \left(\frac{\partial h}{\partial m} \cdot \nabla \frac{\partial f}{\partial \rho} - \frac{\partial f}{\partial m} \cdot \nabla \frac{\partial h}{\partial \rho} \right) dx + \int_{\Omega} s \left(\frac{\partial h}{\partial m} \cdot \nabla \frac{\partial f}{\partial s} - \frac{\partial f}{\partial m} \cdot \nabla \frac{\partial h}{\partial s} \right) dx
$$

is the Lie-Poisson bracket.

In particular, taking $f = h$ gives the energy balance of the open fluid system (3.12) as

$$
\frac{d}{dt}h = \underbrace{\int_{\Omega} \left(b \cdot \frac{\partial \mathfrak{h}}{\partial m} + \theta_{\rho} \frac{\partial \mathfrak{h}}{\partial \rho} + \theta_{s} \frac{\partial \mathfrak{h}}{\partial s} \right) dx}_{\text{bulk source}} + \underbrace{\int_{\partial \Omega} \left(J \cdot \frac{\partial \mathfrak{h}}{\partial m} + j_{\rho} \frac{\partial \mathfrak{h}}{\partial \rho} + j_{s} \frac{\partial \mathfrak{h}}{\partial s} \right) da}_{\text{boundary source}},
$$

while the balance of momentum, mass, and entropy follow by appropriate choice for f , namely

$$
\frac{d}{dt}m_i = \int_{\Omega} \partial_i \left(\mathfrak{h} - m \cdot \frac{\partial \mathfrak{h}}{\partial m} - \rho \frac{\partial \mathfrak{h}}{\partial \rho} - s \frac{\partial \mathfrak{h}}{\partial s} \right) dx + \int_{\Omega} b_i dx + \int_{\partial \Omega} J_i da
$$

$$
\frac{d}{dt} \rho = \int_{\Omega} \theta_\rho dx + \int_{\partial \Omega} j_\rho da
$$

$$
\frac{d}{dt} s = \int_{\Omega} \theta_s dx + \int_{\partial \Omega} j_s da.
$$

Remark 3.7 (On the Lie-Poisson bracket). The Lie-Poisson bracket appearing above can be seen as the Lie-Poisson associated to the Lie algebra X(Ω) of *all* smooth vector fields on Ω , i.e., without any conditions on the boundary. We have obtained it via reduction by Lie group symmetry applied to a specific class of functionals appearing for open fluid systems, namely, of the form (3.14) . It should be noted that it *does not* come from the group $Diff(\Omega)$ of diffeomorphism Ω . First Diff(Ω) is not the Lie group configuration of our system (it is the configuration Lie group of a closed fluid in Ω), and, second, its Lie algebra is the space $\mathfrak{X}_{\parallel}(\Omega)$ of vector fields parallel to the boundary, and not the Lie algebra $\mathfrak{X}(\Omega)$ we are considering here.

Particular cases. In order to present particular cases and compare our bracket formulation with earlier works, it is useful to split the Lie-Poisson bracket in two terms, obtained by integration by parts in a way to isolate the partial derivatives $\frac{\partial f}{\partial m}$, $\frac{\partial f}{\partial \rho}$, and $\frac{\partial f}{\partial s}$. This results in the expression

$$
\{f, h\}_{\text{LP}} = -\int_{\Omega} \left[\left(\nabla \frac{\partial h}{\partial m} m + \left(\nabla \frac{\partial h}{\partial m} \right)^{\mathsf{T}} \cdot m + m \operatorname{div} \frac{\partial h}{\partial m} + \rho \nabla \frac{\partial h}{\partial \rho} + s \nabla \frac{\partial h}{\partial s} \right) \cdot \frac{\partial f}{\partial m} \right. \\
\left. - \operatorname{div} \left(\rho \frac{\partial h}{\partial m} \right) \frac{\partial f}{\partial \rho} - \operatorname{div} \left(s \frac{\partial h}{\partial m} \right) \frac{\partial f}{\partial s} \right] dx \\
+ \int_{\partial \Omega} \left[\left(\frac{\partial h}{\partial m} \cdot m \right) \left(m \cdot \frac{\partial f}{\partial m} \right) + \rho \left(\frac{\partial h}{\partial m} \cdot n \right) \frac{\partial f}{\partial \rho} + s \left(\frac{\partial h}{\partial m} \cdot n \right) \frac{\partial f}{\partial s} \right] da \\
=: \{f, h\}_{\text{bulk}} + \{f, h\}_{\text{boundary}},
$$

with $\{f, h\}_{\text{bulk}}$, resp., $\{f, h\}_{\text{boundary}}$, defined as the integral over Ω , resp., $\partial\Omega$. With this notation, our bracket formulation [\(3.23\)](#page-32-1) reads

$$
\dot{f} = \{f, h\}_{\text{bulk}} + \underbrace{\int_{\partial\Omega} \left[m \left(\frac{\partial \mathfrak{h}}{\partial m} \cdot n \right) + J \right] \cdot \frac{\partial \mathfrak{f}}{\partial m} + \left[\rho \left(\frac{\partial \mathfrak{h}}{\partial m} \cdot n \right) + j_{\rho} \right] \frac{\partial \mathfrak{f}}{\partial \rho} + \left[s \left(\frac{\partial \mathfrak{h}}{\partial m} \cdot n \right) + j_{s} \right] \frac{\partial \mathfrak{f}}{\partial s} \mathrm{d}a}_{\text{= {\{f, h\}_{\text{boundary}}+\text{boundary contribution}} \atop \text{interior contribution}}} \tag{3.24}
$$

which shows how the boundary term $\{\cdot,\cdot\}$ boundary can be combined with the term including the boundary fluxes J, j_{ρ} , and j_{ρ} . We now consider several particular cases of the extended Lie-Poisson formulation. For this discussion we set

$$
b = 0
$$
 and $\theta_{\rho} = \theta_s = 0$.

(A) If the fluxes are chosen as $J = 0$ and $j_{\rho} = j_s = 0$, then the bracket formulation is

$$
\dot{f} = \{f, h\}_{\text{LP}} = \{f, h\}_{\text{bulk}} + \{f, h\}_{\text{boundary}}
$$

which yields the boundary conditions:

$$
\left(\frac{\partial \mathfrak{h}}{\partial m} \cdot n\right) m = 0, \quad \rho \frac{\partial \mathfrak{h}}{\partial m} \cdot n = 0 \quad \text{and} \quad s \frac{\partial \mathfrak{h}}{\partial m} \cdot n = 0 \quad \text{on} \quad \partial \Omega,
$$
 (3.25)

the fluid is closed. It formally recovers the usual Lie-Poisson bracket from semidirect product theory. There is however a major difference in the formulation, which is due to the fact that we are considering the Lie-Poisson structure on the whole Lie algebra $\mathfrak{X}(\Omega)$, not $\mathfrak{X}_{\parallel}(\Omega)$. In our case the Lie-Poisson formulation $f = \{f, h\}_{\text{LP}}$ for all f, not only produces the fluid equations of motion, but also imposes the boundary conditions [\(3.25\)](#page-33-0).

(B) If the boundary fluxes J, j_{ρ} , and j_s are chosen as

$$
J:=-m\left(\frac{\partial h}{\partial m}\cdot m\right), \quad j_{\rho}:=\rho\left(\frac{\partial h}{\partial m}\cdot n\right) \quad \text{and} \quad j_s:=s\left(\frac{\partial h}{\partial m}\cdot n\right),
$$

(recall that our approach allows them to depend on m, ρ, s), then the bracket formulation [\(3.23\)](#page-32-1) reduces to

$$
\dot{f} = \{f, h\}_{\text{bulk}}
$$

and the boundary conditions just vanish, and the fluid is freely open. It is important to remember that $\{f, h\}_{\text{bulk}}$ is not a Poisson bracket. We can write it in terms of the Lie-Poisson bracket as $\{f, h\}_{\text{bulk}} = \{f, h\}_{\text{LP}} - \{f, h\}_{\text{boundary}}$. This setting recovers the bracket formulation for open fluids in [\[22\]](#page-48-5), see also [\[23\]](#page-48-6).

(C) The other cases are when J and j are prescribed, i.e., the fluid is open, but its behavior through the boundary is prescribed.

It is important to note that the boundary can be divided into non-intersecting pieces on which (A) , (B) or (C) hold. For example there could be inflow along one part of the domain, no-flux along another, and open flow along a third part.

4 Extensions

We extend the previously developed framework to address several key aspects, including the treatment of multicomponent fluids, the consideration of general advected quantities, the analysis of higher-order fluids, and the incorporation of boundary stresses. As we shall see, some of these extensions lead to nontrivial modifications of the boundary conditions for open fluids.

4.1 Multicomponent fluids

To treat multicomponent fluids, one considers Lagrangian densities of the form $\mathfrak{l}(u, \rho_k, s)$, i.e., depending on the mass density of each component $k = 1, ..., N$. These single velocity, single temperature, multiple component models are ubiquitous in geophysical fluid dynamics [\[44](#page-50-0), [45](#page-50-1), [46,](#page-50-2) [47](#page-50-3), [43](#page-49-13), [48,](#page-50-4) [49](#page-50-5)] and porous media [\[50](#page-50-6), [51,](#page-50-7) [52,](#page-50-8) [53](#page-50-9), [54\]](#page-50-10). The extension of Proposition [2.9](#page-18-4) to this case is straightforward. In the material description, we replace the terms ϱW , $\Theta_{\rho} \delta W$ and $j_{\rho}\delta W$ by the terms $\sum_k \varrho_k \dot{W}_k$, $\sum_k \Theta_{\rho_k} \delta W_k$ and $\sum_k j_{\rho_k} \delta W_k$ in the Lagrange-d'Alembert principle [\(2.5\)](#page-12-0). In the Eulerian version this gives the variational principle

$$
\delta \int_0^T \int_{\Omega} \left[\mathbf{I}(u, \rho, s) + \sum_k \rho_k D_t w_k + s D_t \gamma \right] \mathrm{d}x \mathrm{d}t \n+ \int_0^T \int_{\Omega} \left[b \cdot \zeta + \sum_k \theta_{\rho_k} D_{\delta} w_k + \theta_s D_{\delta} \gamma \right] \mathrm{d}x \mathrm{d}t \n+ \int_0^T \int_{\partial \Omega} \left[J \cdot \zeta + \sum_k j_{\rho_k} D_{\delta} w_k + j_s D_{\delta} \gamma \right] \mathrm{d}a \mathrm{d}t = 0
$$
\n(4.1)

with respect to variations $\delta u = \partial_t \zeta + [\zeta, u]$ and free variations $\delta \rho_k$, δs , δw_k , $\delta \gamma$, with δw_k and $\delta\gamma$ vanishing at $t = 0, T$ and ζ an arbitrary time dependent vector field vanishing at $t = 0, T$. From this, the following system is obtained

$$
\begin{cases}\n\partial_t \frac{\partial \mathfrak{l}}{\partial u} + \mathcal{L}_u \frac{\partial \mathfrak{l}}{\partial u} - \sum_k \rho_k \nabla \frac{\partial \mathfrak{l}}{\partial \rho_k} - s \nabla \frac{\partial \mathfrak{l}}{\partial s} = b \\
\partial_t \rho_k + \operatorname{div}(\rho_k u) = \theta_{\rho_k}, \text{ for all } k, \qquad \partial_t s + \operatorname{div}(s u) = \theta_s \\
(u \cdot n) \frac{\partial \mathfrak{l}}{\partial u} = -J \quad \text{on } \partial \Omega \\
\rho_k u \cdot n = -j_{\rho_k}, \text{ for all } k, \qquad su \cdot n = -j_s \quad \text{on } \partial \Omega.\n\end{cases} (4.2)
$$

From this point, the variational formulation on the Hamiltonian side, as well as the extension of the Lie-Poisson bracket can be derived in a straightforward way.

4.2 General advected quantities

In this section we consider the case in which the fluid depends or other quantities than the densities ρ and s. We treat the general case of advected tensor fields and tensor field densities. These sorts of fluid models occur in the complex fluids literature [\[55](#page-50-11), [56\]](#page-50-12) and the plasma physics literature [\[57,](#page-50-13) [58,](#page-50-14) [59,](#page-51-0) [60](#page-51-1)], amongst others. For example, magnetohydrodynamics [\[5](#page-47-4)] can be understood as a compressible fluid with an advected magnetic field 2-form.

Case of tensor field densities. Let us assume that the Lagrangian density depends on a (p, q) tensor field density, denoted Π dX in the material description. Following the approach described in §[2.1](#page-10-1) for the mass and entropy densities, we consider a "distributed source" $\Theta_{\Pi} dX =$ $\Theta_{\Pi}(\varphi, \dot{\varphi}, \nabla \varphi, \varrho, K)dX$ of the tensorial quantity, given as a map

$$
\Theta_{\Pi} dX : \varphi^{-1}(\Omega) \to T_q^p(\varphi^{-1}(\Omega)) \otimes \Lambda^n(\varphi^{-1}(\Omega))
$$

with $T_q^p(\varphi^{-1}(\Omega))$ the bundle of (p, q) tensors. We also consider a "boundary source" $\mathfrak{J}_\Pi dA =$ $\mathfrak{J}_{\Pi}(\varphi,\dot{\varphi},\nabla\varphi,\varrho,\Pi)dA$ of this tensorial quantity, given as a map

$$
\mathfrak{J}_{\Pi} dA : \varphi^{-1}(\partial \Omega) \to T_q^p(\varphi^{-1}(\partial \Omega)) \otimes \Lambda^{n-1}(\varphi^{-1}(\partial \Omega)).
$$

The corresponding virtual displacement to be considered in this case, is a (q, p) tensor field Z. Denoting by $\Pi:Z$ the full contraction of tensors, the Lagrange-d'Alembert principle [\(2.5\)](#page-12-0) adapted to this case becomes

$$
\frac{d}{d\varepsilon}\Big|_{\varepsilon=0} \int_0^T \int_{\varphi_\varepsilon^{-1}(\Omega)} \left[\mathfrak{L}(\varphi_\varepsilon, \dot{\varphi}_\varepsilon, \nabla \varphi_\varepsilon, \varrho_\varepsilon, \Pi_\varepsilon) + \varrho_\varepsilon \dot{W}_\varepsilon + \Pi_\varepsilon : \dot{Z}_\varepsilon \right] dX dt \n+ \int_0^T \int_{\varphi^{-1}(\Omega)} \left[\mathfrak{B} \cdot \delta \varphi + \Theta_\varrho \delta W + \Theta_\Pi : \delta Z \right] dX dt \leftarrow \text{bulk contribution} \n+ \int_0^T \int_{\varphi^{-1}(\partial \Omega)} \left[\mathfrak{J} \cdot \delta \varphi + \mathfrak{j}_\varrho \delta W + \mathfrak{j}_\Pi : \delta Z \right] dA dt = 0, \leftarrow \text{boundary contribution}
$$
\n(4.3)

where we assume $\delta\varphi$, δW , and δZ vanish at $t = 0, T$. We leave the derivation of equations in the material description to the reader and focus below on the Eulerian description.

In a similar way with [\(2.17\)](#page-17-1) and [\(2.18\)](#page-17-2), the relabelling symmetry of the tensorial source terms are

$$
\Theta_{\Pi}(\varphi \circ \psi, \dot{\varphi} \circ \psi, \nabla_{X}(\varphi \circ \psi), (\varrho \circ \psi)J\psi, \psi^{*}\Pi J\psi) = \psi^{*}(\Theta_{\Pi}(\varphi, \dot{\varphi}, \nabla \varphi, \varrho, \Pi))J\psi \n\mathfrak{j}_{\Pi}(\varphi \circ \psi, \dot{\varphi} \circ \psi, \nabla_{X}(\varphi \circ \psi), (\varrho \circ \psi)J\psi, \psi^{*}\Pi J\psi) = \psi_{\partial}^{*}(\mathfrak{j}_{\varrho}(\varphi, \dot{\varphi}, \nabla \varphi, \varrho, \Pi))J\psi_{\partial},
$$
\n(4.4)

for all $\psi \in \text{Diff}(\mathbb{R}^n)$, with ψ^* and ψ^*_{∂} denoting the pull-back of tensor fields. From this, one gets the corresponding Eulerian source terms

$$
\theta_{\pi}(u,\rho,\pi)dx:\Omega \to T_q^p(\Omega) \otimes \Lambda^n(\Omega)
$$

$$
j_{\pi}(u,\rho,\pi)da:\partial\Omega \to T_q^p(\Omega)\Lambda^{n-1}(\partial\Omega),
$$

related to Θ_{Π} and \mathfrak{J}_{Π} via

$$
\varphi_*\big(\Theta_\Pi(\varphi,\ldots)\big)J\varphi^{-1} = \theta_\pi(u,\rho,\pi)
$$

$$
(\varphi_\partial)_*\big(\mathfrak{j}_\pi(\varphi,\ldots)\big)J\varphi_\partial^{-1} = \mathfrak{j}_\pi(u,\rho,\pi),
$$
\n(4.5)

with $u = \dot{\varphi} \circ \varphi^{-1}$ and $\rho = (\varrho \circ \varphi^{-1})J\varphi^{-1}$ as earlier, and with the Eulerian version of Π defined by $\pi = (\varphi_* \Pi) J \varphi^{-1}$. Above, we have used the abbreviation $(\varphi, \ldots) = (\varphi, \dot{\varphi}, \varrho, \Pi)$.

We extend to tensor fields the notations $\bar{D}_t \rho$ and $D_t w$ used earlier for the Eulerian time derivatives of densities and functions: given a tensor field density π and a tensor field κ , we use

$$
\bar{D}_t \pi := \partial_t \pi + \pounds_u \pi \quad \text{and} \quad D_t \kappa := \partial_t \kappa + \pounds_u \kappa,
$$

with $\mathcal{L}_u\kappa$ and $\mathcal{L}_u\pi$ the Lie derivative of the tensor field κ and the tensor field density π . We recall their coordinate expressions

$$
\begin{split}\n(\pounds_{u}\kappa)^{a_{1...a_{p}}}_{b_{1...b_{q}}} &= u^{c}\partial_{c}\kappa^{a_{1...a_{p}}}_{b_{1...b_{q}}}-\kappa^{a_{1...a_{r-1}ca_{r+1}...a_{p}}}_{b_{1...b_{q}}}\partial_{c}u^{a_{r}}+\kappa^{a_{1...a_{p}}}_{b_{1...b_{r-1}cb_{r+1}...b_{q}}}\partial_{b_{r}}u^{c} \\
&= u^{c}\partial_{c}\kappa^{a_{1...a_{p}}}_{b_{1...b_{q}}}+\widehat{\kappa}^{a_{1...a_{p}}}_{b_{1...b_{q}}d}\partial_{c}u^{d} \\
(\pounds_{u}\pi)^{a_{1...a_{p}}}_{b_{1...b_{q}}}&= \partial_{c}(u^{c}\pi^{a_{1...a_{p}}}_{b_{1...b_{q}}})-\pi^{a_{1...a_{r-1}ca_{r+1}...a_{p}}}_{b_{1...b_{q}}}\partial_{c}u^{a_{r}}+\pi^{a_{1...a_{p}}}_{b_{1...b_{r-1}cb_{r+1}...b_{q}}}\partial_{b_{r}}u^{c} \\
&= \partial_{c}(u^{c}\pi^{a_{1}...a_{p}}_{b_{1...b_{q}}})+\widehat{\pi}^{a_{1...a_{p}}}_{b_{1...b_{q}}d}\partial_{c}u^{d},\n\end{split}
$$

where we introduced the $(p+1, q+1)$ tensor field defined by

$$
\widehat{\kappa}_{b_1...b_q d}^{a_1...a_p c} = \sum_r \left(\kappa_{b_1...b_{r-1}db_{r+1}...b_q}^{a_1...a_p} \delta_{b_r}^c - \kappa_{b_1...b_q}^{a_1...a_{r-1}ca_{r+1}...a_p} \delta_{d}^{a_r} \right), \tag{4.6}
$$

similarly for $\hat{\pi}$. We can hence write these Lie derivatives as

$$
\pounds_u \kappa = \nabla_u \kappa + \hat{\kappa} : \nabla u
$$
 and $\pounds_u \pi = \text{div}(u\pi) + \hat{\pi} : \nabla u$.

These notations are convenient to write explicitly the equations of motion below. We shall also use the following technical result, see [\[34\]](#page-49-4).

Lemma 4.1. Let κ be a (p, q) tensor field, π a (q, p) tensor field density, and u a vector field. *Then*

$$
\pounds_u \kappa : \pi = (u \cdot \nabla \kappa) : \pi - \text{div}(\pi \cdot \hat{\kappa}) \cdot u + \text{div}((\pi \cdot \hat{\kappa}) \cdot u)
$$

and

$$
\pounds_u \pi : \kappa = -\pi : (u \cdot \nabla \kappa) - \text{div}(\kappa : . \widehat{\pi}) \cdot u + \text{div} ((\pi : \kappa)u + (\kappa : . \widehat{\pi}) \cdot u),
$$

where π ∴ $\hat{\kappa}$ *is the* (1, 1) *tensor field density obtained by contracting all the respective indices of* $\hat{\kappa}$ *and* π *except the last covariant and contravariant indices of* $\hat{\kappa}$ *, similarly for* κ ∴ $\hat{\pi}$ *. Also, we have* π ∴ $\hat{\kappa} + \kappa$ ∴ $\hat{\pi} = 0$ *and* $\pounds_u \kappa : \pi + \pounds_u \pi : \kappa = \text{div}((\pi : \kappa)u)$ *.*

We can now state the extension of Proposition [2.9](#page-18-4) to the tensorial case.

Proposition 4.2 (Eulerian variational principle for open fluids with tensorial advection I). *Assume that the Lagrangian density and the distributed and boundary sources satisfy the relabelling symmetries. Then the following hold:*

(i) *The Lagrange-d'Alembert principle* [\(4.3\)](#page-36-0) *yields the following variational formulation in the Eulerian frame:*

$$
\delta \int_0^T \int_{\Omega} \left[I(u, \rho, \pi) + \rho D_t w + \pi \, :D_t z \right] \mathrm{d}x \mathrm{d}t \n+ \int_0^T \int_{\Omega} \left[b \cdot \zeta + \theta_{\rho} D_{\delta} w + \theta_{\pi} \, :D_{\delta} z \right] \mathrm{d}x \mathrm{d}t \leftarrow bulk \, contribution \n+ \int_0^T \int_{\partial \Omega} \left[J \cdot \zeta + j_{\rho} D_{\delta} w + j_{\pi} \, :D_{\delta} z \right] \mathrm{d}a \mathrm{d}t = 0 \leftarrow boundary \, contribution
$$
\n(4.7)

with respect to variations $\delta u = \partial_t \zeta + [\zeta, u]$ *and free variations* $\delta \rho$, $\delta \pi$, δw , δz , with δw *and* δz *vanishing at* $t = 0, T$ *and* ζ *an arbitrary time dependent vector field vanishing at* $t = 0, T$.

The relation with the material variables used in [\(4.3\)](#page-36-0) *is given by*

$$
u = \dot{\varphi} \circ \varphi^{-1}, \quad \rho = (\varrho \circ \varphi^{-1})J\varphi^{-1}, \quad \pi = \varphi_* \Pi J\varphi^{-1}, \quad w = W \circ \varphi^{-1}, \quad z = \varphi_* Z.
$$

(ii) *This principle yields the following equations and boundary conditions for open fluids:*

$$
\begin{cases}\n\partial_t \frac{\partial \mathfrak{l}}{\partial u} + \mathcal{L}_u \frac{\partial \mathfrak{l}}{\partial u} - \rho \nabla \frac{\partial \mathfrak{l}}{\partial \rho} - \pi : \nabla \frac{\partial \mathfrak{l}}{\partial \pi} + \text{div} \left(\pi : \frac{\partial \mathfrak{l}}{\partial \pi} \right) = b \\
\partial_t \rho + \text{div}(\rho u) = \theta_\rho, \qquad \partial_t \pi + \mathcal{L}_u \pi = \theta_\pi \\
(u \cdot n) \frac{\partial \mathfrak{l}}{\partial u} + \left(\pi : \frac{\partial \mathfrak{l}}{\partial \pi} \right) \cdot n = -J \quad on \quad \partial \Omega \\
\rho u \cdot n = -j_\rho, \qquad \pi u \cdot n = -j_\pi \quad on \quad \partial \Omega.\n\end{cases} \tag{4.8}
$$

Proof. The proof of Part (i) is similar to that in Proposition [2.9.](#page-18-4) For (ii), the treatment of ρ has already been done earlier, so it is enough to present the computation for the variation of $\int_0^T \int_{\Omega} [(u, \pi) + \pi : D_t z] dx dt$. The variation with respect to π gives the condition

$$
D_t z = -\frac{\partial \mathfrak{l}}{\partial \pi}.\tag{4.9}
$$

The variation with respect to u and z is computed as follows:

$$
= \int_{0}^{T} \int_{\Omega} \left[\frac{\partial I}{\partial u} \cdot (\partial_{t} \zeta + \mathcal{L}_{u} \zeta) + \pi \, :D_{t} \delta z + \pi \, : \mathcal{L}_{\partial_{t} \zeta + \mathcal{L}_{u} \zeta z} \right] dxdt
$$

\n
$$
= \int_{0}^{T} \int_{\Omega} \left[\left(-\partial_{t} \frac{\partial I}{\partial u} - \mathcal{L}_{u} \frac{\partial I}{\partial u} \right) \cdot \zeta + \text{div} \left(\left(\frac{\partial I}{\partial u} \cdot \zeta \right) u \right) \right.
$$

\n
$$
+ \pi \, :D_{t} D_{\delta} z - \pi \, :D_{t} \mathcal{L}_{\zeta} z + \pi \, : \mathcal{L}_{\partial_{t} \zeta + \mathcal{L}_{u} \zeta z} \right] dxdt
$$

\n
$$
= \int_{0}^{T} \int_{\Omega} \left[\left(-\partial_{t} \frac{\partial I}{\partial u} - \mathcal{L}_{u} \frac{\partial I}{\partial u} \right) \cdot \zeta + \text{div} \left(\left(\frac{\partial I}{\partial u} \cdot \zeta \right) u \right) \right.
$$

\n
$$
+ \bar{D}_{t} (\pi \, : D_{\delta} z) - \bar{D}_{t} \pi \, : D_{\delta} z - \pi \, : \mathcal{L}_{\zeta} D_{t} z \right] dxdt
$$

\n
$$
= \int_{0}^{T} \int_{\Omega} \left[\left(-\partial_{t} \frac{\partial I}{\partial u} - \mathcal{L}_{u} \frac{\partial I}{\partial u} \right) \cdot \zeta + \text{div} \left(\left(\frac{\partial I}{\partial u} \cdot \zeta + \pi \, : D_{\delta} z \right) u \right) - \bar{D}_{t} \pi \, : D_{\delta} z + \pi \, : \mathcal{L}_{\zeta} \frac{\partial I}{\partial \pi} \right] dxdt
$$

\n
$$
= \int_{0}^{T} \int_{\Omega} \left[\left(-\partial_{t} \frac{\partial I}{\partial u} - \mathcal{L}_{u} \frac{\partial I}{\partial u} + \pi \, : \nabla \frac{\partial I}{\partial \pi} -
$$

where we used [\(4.9\)](#page-38-0), we used ζ , δw , and δz vanishing at $t = 0, T$, and, in the last equality we used Lemma [4.1.](#page-37-0)

Hence, when [\(4.7\)](#page-37-1) is considered, and by collecting the terms proportional to ζ , $D_{\delta}w$, and $D_{\delta}z$, both at the interior and at the boundary, and using the equalities $D_t w = -\frac{\partial l}{\partial \rho}$, $D_t z = -\frac{\partial l}{\partial \sigma}$ ∂π arising from the variations $\delta \rho$ and $\delta \pi$, we get the six equations [\(4.8\)](#page-37-2).

The local and global energy balance equations are found as

$$
\partial_t \mathfrak{e} = \text{div}\left(-\left(\frac{\partial \mathfrak{l}}{\partial u} \cdot u\right)u + \rho \frac{\partial \mathfrak{l}}{\partial \rho} u + \pi \cdot \frac{\partial \mathfrak{l}}{\partial \pi} u + \left(\frac{\partial \mathfrak{l}}{\partial \pi} \cdot \hat{\pi}\right) \cdot u\right) + b \cdot u - \frac{\partial \mathfrak{l}}{\partial \rho} \theta_\rho - \frac{\partial \mathfrak{l}}{\partial \pi} \cdot \theta_\pi
$$

and

$$
\frac{d}{dt} \int_{\Omega} \mathbf{e} dx = \int_{\Omega} \left[b \cdot u - \theta_{\rho} \frac{\partial \mathbf{l}}{\partial \rho} - \theta_{\pi} \, ; \frac{\partial \mathbf{l}}{\partial \pi} \right] dx + \int_{\partial \Omega} \left[J \cdot u - j_{\rho} \frac{\partial \mathbf{l}}{\partial \rho} - j_{\pi} \, ; \frac{\partial \mathbf{l}}{\partial \pi} \right] da.
$$

Remark 4.3 (Additional stress and examples). The dependence of the fluid Lagrangian on the tensor density π induces the additional stress

$$
\sigma_\pi:=\pi\mathrel{\mathop:} \frac{\partial \mathfrak{l}}{\partial \pi}
$$

appearing both in the fluid momentum equation and its boundary condition in [\(4.8\)](#page-37-2). It can be explicitly computed depending on the nature of the tensor π , by using the general formula [\(4.6\)](#page-36-1). For example, we get the following cases

π vector field density
\n
$$
(\sigma_{\pi})_{d}^{c} = \pi^{c} \frac{\partial I}{\partial \pi^{d}}
$$
\n
$$
(\sigma_{\pi})_{d}^{c} = -\pi_{d} \frac{\partial I}{\partial \pi_{c}}
$$
\n
$$
\pi
$$
 2-contravariant symmetric density
\n
$$
(\sigma_{\pi})_{d}^{c} = 2\pi^{ac} \frac{\partial I}{\partial \pi^{ad}}
$$
\n
$$
\pi
$$
 2-covariant symmetric density
\n
$$
(\sigma_{\pi})_{d}^{c} = -2\pi_{ad} \frac{\partial I}{\partial \pi_{ac}}
$$
\n
$$
(\sigma_{\pi})_{d}^{c} = -2\pi_{ad} \frac{\partial I}{\partial \pi_{ac}}
$$

We note that σ_{π} are (1,1) tensor field densities, hence the question of its symmetry needs to explicitly use a metric (here the Euclidean metric) for rising or lowering one of the indices. From the Lagrangian variational point of view we have here, this symmetry is related to the spatial (as opposed to material) covariance of the Lagrangian density. We refer to [\[19,](#page-48-4) [34](#page-49-4)] for these questions.

.

Remark 4.4 (Formulation on manifolds and bundle valued forms). Although we have assumed a bounded domain in \mathbb{R}^n , the developments in this section can be extended to arbitrary Riemannian manifolds. Our treatment can also naturally be extended to (vector) bundle-valued differential forms [\[61](#page-51-2), [30](#page-49-0), [31,](#page-49-1) [62,](#page-51-3) [63](#page-51-4)], of which κ and π are special cases (tensor-valued 0-forms and n-forms, respectively). This will be the subject of future work.

Case of tensor fields. Let us now assume that the Lagrangian density depends on a (p,q) tensor field (not tensor field density), denoted K in the material description. The sources are hence given as $\Theta_K = \Theta_K(\varphi, \dot{\varphi}, \nabla \varphi, \varrho, K)$ with

$$
\Theta_K : \varphi^{-1}(\Omega) \to T_q^p(\varphi^{-1}(\Omega))
$$

and $\mathfrak{J}_K = \mathfrak{J}_K(\varphi, \dot{\varphi}, \nabla \varphi, \varrho, K)$ with

$$
\mathfrak{J}_K: \varphi^{-1}(\partial\Omega) \to T^p_q(\varphi^{-1}(\partial\Omega)).
$$

The corresponding virtual displacement to be considered in this case, is a (q, p) tensor field density Z dX and the Lagrange-d'Alembert principle [\(2.5\)](#page-12-0) adapted to this case becomes

$$
\frac{d}{d\varepsilon}\Big|_{\varepsilon=0} \int_0^T \int_{\varphi_\varepsilon^{-1}(\Omega)} \Big[\mathfrak{L}(\varphi_\varepsilon, \dot{\varphi}_\varepsilon, \nabla \varphi_\varepsilon, \varrho_\varepsilon, K_\varepsilon) + \varrho_\varepsilon \dot{W}_\varepsilon + K_\varepsilon : \dot{Z}_\varepsilon \Big] dX dt \n+ \int_0^T \int_{\varphi^{-1}(\Omega)} \Big[\mathfrak{B} \cdot \delta \varphi + \Theta_\varrho \delta W + \Theta_K : \delta Z \Big] dX dt \quad \leftarrow \text{bulk contribution} \n+ \int_0^T \int_{\varphi^{-1}(\partial \Omega)} \Big[\mathfrak{J} \cdot \delta \varphi + \mathfrak{j}_\varrho \delta W + \mathfrak{j}_K : \delta Z \Big] dA dt = 0, \quad \leftarrow \text{boundary contribution}
$$
\n(4.10)

where we assume $\delta\varphi$, δW , and $\delta\Gamma$ vanish at $t = 0, T$.

Remark 4.5. We note that the geometric setting in this case is different from that in [\(4.3\)](#page-36-0) regarding the construction of the boundary integral. In [\(4.3\)](#page-36-0) the boundary source term $\mathfrak{J}_{\Pi}dA$ is given as a tensor field density on the boundary, which can be contracted, on the boundary $\varphi^{-1}(\partial\Omega)$, with the tensor field δZ to give the density on the boundary to be integrated. In the present case $\delta Z dX$ is a tensor field density on the domain $\varphi^{-1}(\Omega)$ and one needs to induce from it a tensor field density on the boundary, that we denoted as δZdA in [\(4.3\)](#page-36-0), to be contracted with the tensor field \mathfrak{J}_K on the boundary. This step of passing from $\delta Z dX$ to $\delta Z dA$ needs the explicit use of a metric, here the Euclidean one, which was not necessary in (4.3) . This issue appears more clearly when formulating the framework on Riemannian manifolds, which will be pursued elsewhere.

Proposition 4.6 (Eulerian variational principle for open fluids with tensorial advection II). *Assume that the Lagrangian density and the distributed and boundary sources satisfy the relabelling symmetries. Then the following hold:*

(i) *The Lagrange-d'Alembert principle* [\(4.10\)](#page-39-0) *yields the following variational formulation in the Eulerian frame:*

$$
\delta \int_0^T \int_{\Omega} \left[I(u, \rho, \kappa) + \rho D_t w + \kappa : \bar{D}_t z \right] dx dt \n+ \int_0^T \int_{\Omega} \left[b \cdot \zeta + \theta_{\rho} D_{\delta} w + \theta_{\kappa} : \bar{D}_{\delta} z \right] dx dt \leftarrow bulk contribution \n+ \int_0^T \int_{\partial \Omega} \left[J \cdot \zeta + j_{\rho} D_{\delta} w + j_{\kappa} : \bar{D}_{\delta} z \right] da dt = 0 \leftarrow boundary contribution
$$
\n(4.11)

with respect to variations $\delta u = \partial_t \zeta + [\zeta, u]$ *and free variations* $\delta \rho$ *,* $\delta \kappa$ *,* δw *,* δz *, with* δw *and* δz *vanishing* at $t = 0, T$ *and* ζ *an arbitrary time dependent vector field vanishing at* $t = 0, T$.

The relation with the material variables used in [\(4.3\)](#page-36-0) *is given by*

$$
u = \dot{\varphi} \circ \varphi^{-1}, \quad \rho = (\varrho \circ \varphi^{-1})J\varphi^{-1}, \quad \kappa = \varphi_*K, \quad w = W \circ \varphi^{-1}, \quad z = \psi_*ZJ\varphi^{-1}.
$$

(ii) *This principle yields the following equations and boundary conditions for open fluids:*

$$
\begin{cases}\n\partial_t \frac{\partial \mathfrak{l}}{\partial u} + \mathcal{L}_u \frac{\partial \mathfrak{l}}{\partial u} - \rho \nabla \frac{\partial \mathfrak{l}}{\partial \rho} - \kappa : \nabla \frac{\partial \mathfrak{l}}{\partial \kappa} + \text{div} \left(\left(\frac{\partial \mathfrak{l}}{\partial \kappa} : \kappa \right) \delta + \kappa : \frac{\partial \mathfrak{l}}{\partial \kappa} \right) = b \\
\partial_t \rho + \text{div}(\rho u) = \theta_\rho, \qquad \partial_t \kappa + \mathcal{L}_u \kappa = \theta_\kappa \\
(u \cdot n) \frac{\partial \mathfrak{l}}{\partial u} + \left(\left(\frac{\partial \mathfrak{l}}{\partial \kappa} : \kappa \right) \delta + \kappa : \frac{\partial \mathfrak{l}}{\partial \kappa} \right) \cdot n = -J \quad on \quad \partial \Omega \\
\rho u \cdot n = -j_\rho, \qquad \kappa u \cdot n = -j_\kappa \quad on \quad \partial \Omega.\n\end{cases} \tag{4.12}
$$

The local and global energy balance equations are found as

$$
\partial_t \mathfrak{e} = \text{div}\left(-\left(\frac{\partial \mathfrak{l}}{\partial u} \cdot u\right)u + \rho \frac{\partial \mathfrak{l}}{\partial \rho} u + \left(\frac{\partial \mathfrak{l}}{\partial \kappa} \cdot \widehat{\kappa}\right) \cdot u\right) + b \cdot u - \frac{\partial \mathfrak{l}}{\partial \rho} \theta_\rho - \frac{\partial \mathfrak{l}}{\partial \kappa} \cdot \theta_\kappa
$$

and

$$
\frac{d}{dt} \int_{\Omega} \mathbf{e} dx = \int_{\Omega} \left[b \cdot u - \theta_{\rho} \frac{\partial \mathbf{l}}{\partial \rho} - \theta_{\kappa} \cdot \frac{\partial \mathbf{l}}{\partial \kappa} \right] dx + \int_{\partial \Omega} \left[J \cdot u - j_{\rho} \frac{\partial \mathbf{l}}{\partial \rho} - j_{\kappa} \cdot \frac{\partial \mathbf{l}}{\partial \kappa} \right] da.
$$

Examples. The additional stress due to the dependence on the tensor κ is now given by

$$
\sigma_{\kappa} := \left(\frac{\partial \mathfrak{l}}{\partial \kappa} : \kappa\right) \delta + \kappa \therefore \frac{\partial \mathfrak{l}}{\partial \kappa},
$$

which also appears both in the fluid momentum equation and its boundary condition in (4.12) . It can be explicitly computed depending on the nature of the tensor κ , by using the general formula [\(4.6\)](#page-36-1). For example, we get the following cases

$$
\kappa \text{ vector field} \qquad (\sigma_{\kappa})_d^c = \frac{\partial \mathfrak{l}}{\partial \kappa^a} \kappa^a \delta_d^c + \kappa^c \frac{\partial \mathfrak{l}}{\partial \kappa^d}
$$

$$
\kappa \text{ 1-form} \qquad (\sigma_{\kappa})_d^c = \frac{\partial \mathfrak{l}}{\partial \kappa_a} \kappa_a \delta_d^c - \kappa_d \frac{\partial \mathfrak{l}}{\partial \kappa_c}
$$

$$
\kappa \text{ 2-form} \qquad (\sigma_{\kappa})_d^c = \frac{\partial \mathfrak{l}}{\partial \kappa_{ac}} \kappa_{ac} \delta_d^c - 2\kappa_{ad} \frac{\partial \mathfrak{l}}{\partial \kappa_{ac}}.
$$

Magnetohydrodynamics with open boundaries. In particular, for κ a closed 2-form, which we identify with a divergence free vector field B , this is the setting of magnetohydrodynamics. In this case, the equations [\(4.12\)](#page-40-0) become

$$
\begin{cases}\n\partial_t \frac{\partial \mathfrak{l}}{\partial u} + \mathcal{L}_u \frac{\partial \mathfrak{l}}{\partial u} - \rho \nabla \frac{\partial \mathfrak{l}}{\partial \rho} - B \cdot \nabla \frac{\partial \mathfrak{l}}{\partial B} + \text{div} \left(B \otimes \frac{\partial \mathfrak{l}}{\partial B} \right) = b \\
\partial_t \rho + \text{div}(\rho u) = \theta_\rho, \qquad \partial_t B + \text{curl}(B \times u) = \theta_B \\
(u \cdot n) \frac{\partial \mathfrak{l}}{\partial u} + \left(B \otimes \frac{\partial \mathfrak{l}}{\partial B} \right) \cdot n = -J \quad \text{on} \quad \partial \Omega \\
\rho u \cdot n = -j_\rho, \qquad Bu \cdot n = -j_B \quad \text{on} \quad \partial \Omega.\n\end{cases} \tag{4.13}
$$

The local and global energy balance equations are found as

$$
\partial_t \mathfrak{e} = \text{div} \left(- \left(\frac{\partial \mathfrak{l}}{\partial u} \cdot u \right) u + \rho \frac{\partial \mathfrak{l}}{\partial \rho} u + (B \times u) \times \frac{\partial \mathfrak{l}}{\partial B} \right) + b \cdot u - \frac{\partial \mathfrak{l}}{\partial \rho} \theta_\rho - \frac{\partial \mathfrak{l}}{\partial B} \cdot \theta_B
$$

and

$$
\frac{d}{dt} \int_{\Omega} \mathbf{e} dx = \int_{\Omega} \left[b \cdot u - \frac{\partial \mathbf{I}}{\partial \rho} \theta_{\rho} - \frac{\partial \mathbf{I}}{\partial B} \cdot \theta_{B} \right] dx + \int_{\partial \Omega} \left[J \cdot u - j_{\rho} \frac{\partial \mathbf{I}}{\partial \rho} - j_{B} \cdot \frac{\partial \mathbf{I}}{\partial B} \right] da.
$$

In particular, it is interesting to note how the term associated to B in the local energy balance, i.e., $(B \times u) \times \frac{\partial I}{\partial B}$, contributes to both the momentum, $J \cdot u$, and magnetic, $j_B \cdot \frac{\partial I}{\partial B}$, effect in the boundary term of the global energy balance. Using

$$
(B \times u) \times \frac{\partial \mathfrak{l}}{\partial B} = \left(\frac{\partial \mathfrak{l}}{\partial B} \cdot B\right)u - \left(\frac{\partial \mathfrak{l}}{\partial B} \cdot u\right)B\tag{4.14}
$$

we get

$$
\int_{\Omega} \operatorname{div} \left(-\left(\frac{\partial \mathfrak{l}}{\partial u} \cdot u \right) u + (B \times u) \times \frac{\partial \mathfrak{l}}{\partial B} \right) \mathrm{d}x
$$
\n
$$
= \int_{\partial \Omega} \left[-\left(\frac{\partial \mathfrak{l}}{\partial u} \cdot u \right) u \cdot n - \left(\frac{\partial \mathfrak{l}}{\partial B} \cdot u \right) B \cdot n + \left(\frac{\partial \mathfrak{l}}{\partial B} \cdot B \right) u \cdot n \right] \mathrm{d}a
$$
\n
$$
= \int_{\partial \Omega} \left[J \cdot u - j_{B} \cdot \frac{\partial \mathfrak{l}}{\partial B} \right] \mathrm{d}a,
$$

showing the different role played by both terms in [\(4.14\)](#page-41-0). It also explains the occurrence of the magnetic field contribution $(B \otimes \frac{\partial R}{\partial P})$ $\frac{\partial \tilde{l}}{\partial B}$ *n* in the open boundary condition for the fluid momentum.

4.3 Higher order fluids

We consider the case of a Lagrangian density depending also on the first order derivative of ρ . An example of a fluid model with this feature is the Euler-Kortweg equations for multiphase flow [\[64](#page-51-5)]. As we shall see this induces the appearance of an additional boundary condition. Given $\mathfrak{l}(u, \rho, \nabla \rho)$, we consider the same Lagrange-d'Alembert principle as in (2.5) , in which we do not include the entropy for simplicity. We thus have

$$
\delta \int_0^T \int_{\Omega} \left[\mathfrak{l}(u, \rho, \nabla \rho) + \rho D_t w \right] dx dt + \int_0^T \int_{\Omega} \left[b \cdot \zeta + \theta_{\rho} D_{\delta} w \right] dx dt + \int_0^T \int_{\partial \Omega} \left[J \cdot \zeta + j_{\rho} D_{\delta} w \right] d\alpha dt = 0
$$
\n(4.15)

with respect to variations $\delta u = \partial_t \zeta + [\zeta, u]$ and free variations $\delta \rho$, δw , with δw vanishing at $t = 0, T$ and ζ an arbitrary time dependent vector field vanishing at $t = 0, T$.

This principle yields the following equations and boundary conditions for open fluids:

$$
\begin{cases}\n\partial_t \frac{\partial \mathfrak{l}}{\partial u} + \mathcal{L}_u \frac{\partial \mathfrak{l}}{\partial u} - \rho \nabla \Big(\frac{\partial \mathfrak{l}}{\partial \rho} - \text{div } \frac{\partial \mathfrak{l}}{\partial \nabla \rho} \Big) = b \\
\partial_t \rho + \text{div}(\rho u) = \theta_\rho \\
(u \cdot n) \frac{\partial \mathfrak{l}}{\partial u} = -J \quad \text{on} \quad \partial \Omega \\
\rho u \cdot n = -j_\rho, \qquad \frac{\partial \mathfrak{l}}{\partial \nabla \rho} \cdot n = 0 \quad \text{on} \quad \partial \Omega.\n\end{cases} (4.16)
$$

We note the occurrence of the last boundary condition on $\frac{\partial \mathcal{I}}{\partial \nabla \rho}$. Its role in the energy balance is illustrated as follows. System [\(4.16\)](#page-42-2) implies the following local energy balance for the total energy density $\mathfrak{e}(u, \rho, \nabla \rho) = \frac{\partial \mathfrak{l}}{\partial u} \cdot u - \mathfrak{l}(u, \rho, \nabla \rho)$ defined from \mathfrak{l} :

$$
\partial_t \mathfrak{e} = \text{div} \left(- \left(\frac{\partial \mathfrak{l}}{\partial u} \cdot u \right) u + \rho \left(\frac{\partial \mathfrak{l}}{\partial \rho} - \text{div} \frac{\partial \mathfrak{l}}{\partial \nabla \rho} \right) u + \frac{\partial \mathfrak{l}}{\partial \nabla \rho} (\text{div}(\rho u) - \theta_\rho) \right) + b \cdot u - \theta_\rho \left(\frac{\partial \mathfrak{l}}{\partial \rho} - \text{div} \frac{\partial \mathfrak{l}}{\partial \nabla \rho} \right).
$$

This should be compared to (2.26) . Now, by integrating it over the domain, we use the four boundary conditions in [\(4.16\)](#page-42-2) to finally get

$$
\frac{d}{dt} \int_{\Omega} \mathfrak{e} \, dx = \int_{\Omega} \left[b \cdot u - \theta_{\rho} \left(\frac{\partial \mathfrak{l}}{\partial \rho} - \text{div} \frac{\partial \mathfrak{l}}{\partial \nabla \rho} \right) \right] dx + \int_{\partial \Omega} \left[J \cdot u - j_{\rho} \left(\frac{\partial \mathfrak{l}}{\partial \rho} - \text{div} \frac{\partial \mathfrak{l}}{\partial \nabla \rho} \right) \right] da.
$$

We note in particular, how the additional condition on $\frac{\partial \ell}{\partial \nabla \rho}$ is used so that the term $\frac{\partial \ell}{\partial \nabla \rho}(\text{div}(\rho u) \theta_{\rho}$) has no boundary flow contribution.

4.4 Boundary stresses

As a final extension, we add the contribution of a (bulk) external stress tensor σ . This is a common feature in geophysical fluid dynamics models, used to model surface stresses due to the wind in ocean models and the land or waves in atmospheric models [\[65\]](#page-51-6). By appropriately adding the stress contribution in the Euler-Poincaré-d'Alembert principle (2.21) , we get:

$$
\delta \int_0^T \int_{\Omega} \left[I(u,\rho) + \rho D_t w \right] dx dt + \int_0^T \int_{\Omega} \left[b \cdot \zeta - \sigma \cdot \nabla \zeta + \theta_{\rho} D_{\delta} w \right] dx dt + \int_0^T \int_{\partial \Omega} \left[J \cdot \zeta + j_{\rho} D_{\delta} w \right] d\alpha dt = 0.
$$
\n(4.17)

This gives

$$
\begin{cases}\n\partial_t \frac{\partial \mathfrak{l}}{\partial u} + \mathcal{L}_u \frac{\partial \mathfrak{l}}{\partial u} - \rho \nabla \frac{\partial \mathfrak{l}}{\partial \rho} = b + \operatorname{div} \sigma \\
\partial_t \rho + \operatorname{div}(\rho u) = \theta_\rho \\
(u \cdot n) \frac{\partial \mathfrak{l}}{\partial u} = -J + \sigma \cdot n \quad \text{on} \quad \partial \Omega \\
\rho u \cdot n = -j_\rho \quad \text{on} \quad \partial \Omega\n\end{cases}
$$
\n(4.18)

in which we note the appearance of the external stress tensor in the boundary condition for the fluid momentum.

The local and balance equations are found

$$
\partial_t \mathfrak{e} = \text{div} \left(- \left(\frac{\partial \mathfrak{l}}{\partial u} \cdot u \right) u + \rho \frac{\partial \mathfrak{l}}{\partial \rho} u \right) + b \cdot u + \text{div} \, \sigma \cdot u - \theta_\rho \frac{\partial \mathfrak{l}}{\partial \rho}
$$

and

$$
\frac{d}{dt} \int_{\Omega} \mathfrak{e} \, dx = \int_{\Omega} \left[b \cdot u - \theta_{\rho} \frac{\partial \mathfrak{l}}{\partial \rho} - \sigma : \nabla u \right] dx + \int_{\partial \Omega} \left[J \cdot u - j_{\rho} \frac{\partial \mathfrak{l}}{\partial \rho} \right] da.
$$

5 Conclusions

We have developed a geometric framework based on Lie groups for the motion of fluids with fixed, permeable boundaries, that extends the usual geometric desciption in closed domains. Our setting is based on the classic approach with appropriate extensions to incorporate bulk and boundary forces via a Lagrange-d'Alembert approach, taking into account only the fluid parcels present in the domain. Using this, we were able to develop both a variational formulation in the Eulerian description that extends the Euler-Poincaré framework to open fluids; and a bracket-based formulations that extends the Lie-Poisson bracket. It is capable of reproducing, as particular cases, existing results from the GENERIC literature [\[22,](#page-48-5) [23](#page-48-6)]. A comparison with related port-Hamiltonian results [\[24](#page-48-7), [25](#page-48-8), [26](#page-48-9), [27](#page-48-10), [28](#page-48-11), [29,](#page-48-12) [30,](#page-49-0) [31\]](#page-49-1) will be the subject of future work. This general framework was illustrated with several examples (shallow water and compressible Euler), and extensions to multicomponent fluids, general tensor advected quantities, higherorder fluids and boundary stresses were demonstrated. In future work we will consider the extension of these ideas to moving boundaries, following the approach in [\[19](#page-48-4)]; and also an extension to interactive coupling of multiple fluid domains.

A Equivalence between Lagrangian and Eulerian open fluid equations

We have seen in Proposition [2.9](#page-18-4) that the variational principles in regard to the action integrals in the Lagrangian perspective (2.5) and the Eulerian perspective (2.21) are equivalent. We can also show directly that the Euler-Poincaré equations derived from the two variational principles are equivalent by change of perspective. To see this, we first calculate the relationship between the derivatives of reduced and non-reduced Lagrangians.

In fact, from the relation

$$
\mathfrak{L}(\varphi, \dot{\varphi}, \nabla \varphi, \varrho, S)(X) = \mathfrak{l}(u, \rho, s) \circ \varphi(X) J \varphi(X)
$$

we have

$$
\mathfrak{L}(\varphi(X),\dot{\varphi}(X),\nabla\varphi(X),\varrho(X))=\mathfrak{l}\left(\dot{\varphi}(X),\frac{\varrho(X)}{J\varphi(X)},\frac{S(X)}{J\varphi(X)}\right)J\varphi(X)
$$

by definition of u , ρ and s . From this equation, we have the following equalities:

$$
\frac{\partial \mathfrak{L}}{\partial \varphi} = 0
$$

\n
$$
\frac{\partial \mathfrak{L}}{\partial \dot{\varphi}} = \frac{\partial \mathfrak{l}}{\partial u} \circ \varphi J \varphi
$$

\n
$$
\frac{\partial \mathfrak{L}}{\partial \nabla x \varphi} = \left(\left(\mathfrak{l} - \rho \frac{\partial \mathfrak{l}}{\partial \rho} - s \frac{\partial \mathfrak{l}}{\partial s} \right) \circ \varphi \right) \left(\nabla_x \varphi^{-1} \circ \varphi \right) J \varphi
$$

\n
$$
\frac{\partial \mathfrak{L}}{\partial \varrho} = \frac{\partial \mathfrak{l}}{\partial \rho} \circ \varphi
$$

\n
$$
\frac{\partial \mathfrak{L}}{\partial S} = \frac{\partial \mathfrak{l}}{\partial s} \circ \varphi.
$$

(1) We first prove the equivalence of the following two equations:

$$
D_t \frac{\partial \mathfrak{L}}{\partial \dot{\varphi}} + \text{DIV } \frac{\partial \mathfrak{L}}{\partial \nabla \varphi} - \frac{\partial \mathfrak{L}}{\partial \varphi} = \mathfrak{B}
$$

$$
\partial_t \frac{\partial \mathfrak{l}}{\partial u} + \mathcal{L}_u \frac{\partial \mathfrak{l}}{\partial u} - \rho \nabla \frac{\partial \mathfrak{l}}{\partial \rho} - s \nabla \frac{\partial \mathfrak{l}}{\partial s} = b.
$$

First of all, taking the material time derivative on both sides of the relation $\frac{\partial \mathcal{L}}{\partial \dot{\varphi}} = \frac{\partial \mathcal{L}}{\partial u} \circ \varphi J\varphi$, we have that:

$$
D_t \frac{\partial \mathfrak{L}}{\partial \dot{\varphi}} = \partial_t \frac{\partial \mathfrak{l}}{\partial u} \circ \varphi J\varphi + \nabla \frac{\partial \mathfrak{l}}{\partial u} \circ \varphi \cdot \dot{\varphi} J\varphi + \left(\frac{\partial \mathfrak{l}}{\partial u} \text{div} u\right) \circ \varphi J\varphi.
$$

Then, we have that:

$$
\text{DIV} \frac{\partial \mathcal{L}}{\partial \nabla \varphi} = \nabla_X \left(\left(\ell - \rho \frac{\partial \ell}{\partial \rho} - s \frac{\partial \ell}{\partial s} \right) \circ \varphi \right) \cdot \left(\nabla_x \varphi^{-1} \circ \varphi \right) J \varphi.
$$

In local coordinates, it is:

$$
\left(\frac{\partial \mathfrak{L}}{\partial \nabla \varphi^i_{,A}}\right)_{,A}=\left(\left(\ell-\rho\frac{\partial \ell}{\partial \rho}-s\frac{\partial \ell}{\partial s}\right)\circ \varphi\right)_{,A}(\varphi^{-1})^A_{,i}\circ \varphi J\varphi,
$$

where we have used Piola's identity:

$$
\text{DIV}\left(\left(\nabla_x \varphi^{-1} \circ \varphi\right) J\varphi\right) = 0.
$$

Next we have that:

$$
\begin{split}\n&\left(\left(\ell-\rho\frac{\partial\ell}{\partial\rho}-s\frac{\partial\ell}{\partial s}\right)\circ\varphi\right)_{,A}(\varphi^{-1})_{,i}^{A}\circ\varphi J\varphi=\left(\left(\ell-\rho\frac{\partial\ell}{\partial\rho}-s\frac{\partial\ell}{\partial s}\right)_{,k}\circ\varphi\right)\varphi_{,A}^{k}(\varphi^{-1})_{,i}^{A}\circ\varphi J\varphi \\
&=\left(\left(\ell-\rho\frac{\partial\ell}{\partial\rho}-s\frac{\partial\ell}{\partial s}\right)_{k}\circ\varphi\right)e_{ki}J\varphi=\left(\left(\ell-\rho\frac{\partial\ell}{\partial\rho}-s\frac{\partial\ell}{\partial s}\right)_{,i}\circ\varphi\right)J\varphi \\
&=\left(\left(\frac{\partial\ell}{\partial u^{j}}u_{,i}^{j}+\frac{\partial\ell}{\partial\rho}\rho_{,i}+\frac{\partial\ell}{\partial s}s_{,i}-\frac{\partial\ell}{\partial\rho}\rho_{,i}-\rho\left(\frac{\partial\ell}{\partial\rho}\right)_{,i}-\frac{\partial\ell}{\partial s}s_{,i}-s\left(\frac{\partial\ell}{\partial s}\right)_{,i}\right)\circ\varphi\right)J\varphi \\
&=\left(\left(\frac{\partial\ell}{\partial u^{j}}u_{,i}^{j}-\rho\left(\frac{\partial\ell}{\partial\rho}\right)_{,i}-s\left(\frac{\partial\ell}{\partial s}\right)_{,i}\right)\circ\varphi\right)J\varphi.\n\end{split}
$$

Thus

$$
\text{DIV} \frac{\partial \mathfrak{L}}{\partial \nabla \varphi} = \left(\left((\nabla u)^{\mathsf{T}} \cdot \frac{\partial \ell}{\partial u} - \rho \nabla \frac{\partial \ell}{\partial \rho} - s \nabla \frac{\partial \ell}{\partial s} \right) \circ \varphi \right) J\varphi.
$$

Hence

$$
\partial_t \frac{\partial \mathfrak{L}}{\partial \dot{\varphi}} + \text{DIV} \frac{\partial \mathfrak{L}}{\partial \nabla \varphi} - \frac{\partial \mathfrak{L}}{\partial \varphi} = \left(\partial_t \frac{\partial \mathfrak{l}}{\partial u} + \left(\frac{\partial \mathfrak{l}}{\partial u} \text{div} u \right) + (\nabla u)^{\mathsf{T}} \cdot \frac{\partial \mathfrak{l}}{\partial u} + u \cdot \nabla \frac{\partial \mathfrak{l}}{\partial u} - \rho \nabla \frac{\partial \mathfrak{l}}{\partial \rho} - s \nabla \frac{\partial \mathfrak{l}}{\partial s} \right) \circ \varphi J \varphi
$$

$$
= \left(\partial_t \frac{\partial \mathfrak{l}}{\partial u} + \mathfrak{L}_u \frac{\partial \mathfrak{l}}{\partial u} - \rho \nabla \frac{\partial \mathfrak{l}}{\partial \rho} - s \nabla \frac{\partial \mathfrak{l}}{\partial s} \right) \circ \varphi J \varphi.
$$

On the other hand, by definition,

$$
b\circ\varphi J\varphi=\mathfrak{B}.
$$

We therefore have the equivalence.

$$
D_t \frac{\partial \mathfrak{L}}{\partial \dot{\varphi}} + \text{DIV} \frac{\partial \mathfrak{L}}{\partial \nabla \varphi} - \frac{\partial \mathfrak{L}}{\partial \varphi} = \mathfrak{B} \quad \Leftrightarrow \quad \partial_t \frac{\partial \mathfrak{l}}{\partial u} + \pounds_u \frac{\partial \mathfrak{l}}{\partial u} - \rho \nabla \frac{\partial \mathfrak{l}}{\partial \rho} - s \nabla \frac{\partial \mathfrak{l}}{\partial s} = b.
$$

(2) Regarding the balance of mass, we have

$$
D_t \varrho = D_t(\rho \circ \varphi J \varphi) = \partial_t \rho \circ \varphi J \varphi + \text{div}(\rho u) \circ \varphi J \varphi.
$$

On the other hand

$$
\theta_{\varrho} \circ \varphi J \varphi = \Theta_{\rho}.
$$

Thus we have

$$
D_t \varrho = \Theta_\varrho \quad \Leftrightarrow \quad \partial_t \rho + \text{div}(\rho u) = \theta_\rho.
$$

A similar proof shows the equivalence

$$
D_t S = \Theta_S \quad \Leftrightarrow \quad \partial_t s + \text{div}(s u) = \theta_s.
$$

(3) On the boundary $\partial(\varphi^{-1}(\Omega))$, we have that for any vector field $w(x)$:

$$
N \cdot ((\nabla_x \varphi^{-1} \cdot w) \circ \varphi) J\varphi = (n \cdot w) \circ \varphi_{\partial} J\varphi_{\partial}, \tag{A.1}
$$

where $N(X)$ at $X \in \partial(\varphi^{-1}(\Omega))$ and $n(x)$ at $x = \varphi(X) \in \partial(\Omega)$ are the normal vectors on the boundary pointing outward. In fact, due to Piola's identity,

$$
\text{DIV}\left((\nabla_x \varphi^{-1} \cdot w) \circ \varphi)J\varphi\right) = \text{div}(w) \circ \varphi J\varphi.
$$

Applying the divergence theorem on both sides and a change of variables, we have:

$$
\int_{\partial(\varphi^{-1}(\Omega))} N \cdot ((\nabla_x \varphi^{-1} \cdot w) \circ \varphi) J \varphi dA = \int_{\partial \Omega} n \cdot w da.
$$

Using the change of variables $\varphi_{\partial}: X \mapsto x$ on the boundary integral, we also have:

$$
\int_{\partial\Omega} n \cdot w \mathrm{d}a = \int_{\partial(\varphi^{-1}(\Omega))} (n \cdot w) \circ \varphi_{\partial} J \varphi_{\partial} \mathrm{d}A
$$

so

$$
\int_{\partial(\varphi^{-1}(\Omega))} N \cdot ((\nabla_x \varphi^{-1} \cdot w) \circ \varphi) J \varphi dA = \int_{\partial(\varphi^{-1}(\Omega))} (n \cdot w) \circ \varphi_{\partial} J \varphi_{\partial} dA.
$$

Since w is an arbitrary vector field, we can multiply w with test functions that converge to the Dirac delta function, and the identity remains true. We can thus remove the integral and get the identity [A.1.](#page-45-0)

Taking $w = \rho u$, we have that $w \circ \varphi J\varphi = \varrho \dot{\varphi}$. Piola's identity gives that:

$$
N \cdot (\varrho(\nabla_x \varphi^{-1} \circ \varphi \cdot \dot{\varphi})) = (n \cdot (\rho u)) \circ \varphi_{\partial} J \varphi_{\partial}.
$$

On the other hand, by definition

$$
\mathfrak{j}_{\varrho} = j_{\rho} \circ \varphi_{\partial} J \varphi_{\partial}.
$$

We conclude that on the boundary

$$
((\nabla_x \varphi^{-1} \circ \varphi \cdot \dot{\varphi}) \varrho) \cdot N = -\mathfrak{j}_{\varrho} \iff \rho u \cdot n = -\mathfrak{j}_{\rho}.
$$

The same proof shows

$$
((\nabla \varphi_x^{-1} \circ \varphi \cdot \dot{\varphi})S) \cdot N = -\mathfrak{j}_S \iff su \cdot n = -\mathfrak{j}_s.
$$

(4) Finally, note that

$$
\left(\rho \frac{\partial \mathfrak{L}}{\partial \rho} + S \frac{\partial \mathfrak{L}}{\partial S} - \mathfrak{L}\right) \nabla \varphi^{-1} \circ \varphi = \left(\rho \frac{\partial \mathfrak{l}}{\partial \rho} + s \frac{\partial \mathfrak{l}}{\partial s} - \mathfrak{l}\right) \circ \varphi J \varphi \nabla \varphi^{-1} \circ \varphi = -\frac{\partial \mathfrak{L}}{\partial \nabla \varphi}
$$

and that

$$
N \cdot (\nabla \varphi^{-1} \circ \varphi \cdot \dot{\varphi}) \frac{\partial \mathfrak{L}}{\partial \dot{\varphi}} = N \cdot (\nabla \varphi^{-1} \circ \varphi \cdot \dot{\varphi}) \frac{\partial \mathfrak{l}}{\partial u} \circ \varphi J \varphi = (n \cdot u) \frac{\partial \mathfrak{l}}{\partial u} \circ \varphi_{\partial} J \varphi_{\partial}
$$

which is equation [A.1](#page-45-0) with $w = \frac{\partial l}{\partial u}u$. By definition

$$
\mathfrak{J}=J\circ\varphi_{\partial}J\varphi_{\partial},
$$

we conclude that on the boundary

$$
N \cdot \left(\frac{\partial \mathfrak{L}}{\partial \nabla \varphi} + (\nabla \varphi^{-1} \circ \varphi \cdot \dot{\varphi}) \frac{\partial \mathfrak{L}}{\partial \dot{\varphi}} + \left(\varrho \frac{\partial \mathfrak{L}}{\partial \varrho} + S \frac{\partial \mathfrak{L}}{\partial S} - \mathfrak{L}\right) \nabla \varphi^{-1} \circ \varphi\right) = -\mathfrak{J}
$$

$$
\Leftrightarrow
$$

$$
(u \cdot n) \frac{\partial \mathfrak{l}}{\partial u} = -J.
$$

References

- [1] D. D. Holm, J. E. Marsden, and T. S. Ratiu. The Euler–Poincar´e equations and semidirect products with applications to continuum theories. *Advances in Mathematics*, 137(1):1–81, 1998.
- [2] F. Gay-Balmaz and H. Yoshimura. A Lagrangian variational formalism for nonequilibrium thermodynamics. Part I: discrete systems. *J. Geom. Phys.*, 111:169–193, 2017.
- [3] F. Gay-Balmaz and H. Yoshimura. A Lagrangian variational formalism for nonequilibrium thermodynamics. Part II: continuum systems. *J. Geom. Phys.*, 111:194–212, 2017.
- [4] P. Morrison and J. Greene. Noncanonical Hamiltonian density formulation of hydrodynamics and ideal magnetohydrodynamics. *Phys. Rev. Letters*, 45:790–794, 1980.
- [5] D. D. Holm and B. A. Kupershmidt. Noncanonical Hamiltonian formulation of ideal magnetohydrodynamics. *Physica D: Nonlinear Phenomena*, 7(1-3):330–333, 1983.
- [6] I. E. Dzyaloshinskii and G. E. Volovick. Poisson brackets in condensed matter physics. *Ann. Phys.*, 125:67–97, 1980.
- [7] J E. Marsden and A. Weinstein. Coadjoint orbits, vortices, and Clebsch variables for incompressible fluids. *Physica D: Nonlinear Phenomena*, 7(1):305–323, 1983.
- [8] J. E. Marsden, T. S. Ratiu, and A. Weinstein. Semidirect product and reduction in mechanics. *Trans. Amer. Math. Soc.*, 281:147–177, 1984.
- [9] M. Grmela. Bracket formulation of dissipative fluid mechanics equations. *Phys. Lett. A*, 102:355–358, 1984.
- [10] A. Kaufman. Dissipative Hamiltonian systems: A unifying principle. *Phys. Lett. A*, 100: 419–422, 1984.
- [11] P. Morrison. Bracket formulation for irreversible classical fields. *Phys. Lett. A*, 100:423–427, 1984.
- [12] P. Morrison. Some observations regarding brackets and dissipation. Technical report, University of California, Berkeley, 1984.
- [13] M. Grmela and H.-C. Ottinger. Dynamics and thermodynamics of complex fluids. I. Development of a general formalism. *Phys. Rev. E*, 56:6620–6632, 1997.
- [14] H.-C. Ottinger and M. Grmela. Dynamics and thermodynamics of complex fluids. II. Illustrations of a general formalism. *Phys. Rev. E*, 56:6633–6655, 1997.
- [15] D. D. Holm, J. E. Marsden, T. S. Ratiu, and A. Weinstein. Nonlinear stability of fluid and plasma equilibria. *Physics Reports*, 123(1-2):1–116, 1985.
- [16] G. M. Webb and R. L. Mace. Noether's theorems and fluid relabelling symmetries in magnetohydrodynamics and gas dynamics. *J. Phys. A Math. Theoret. article No. JPHYSA-101-057, available at http://arxiv. org/abs/1403.3133 (submitted on March 10)*, 2014.
- [17] G. M. Webb, B. Dasgupta, J. F. McKenzie, Q. Hu, and G. P. Zank. Local and nonlocal advected invariants and helicities in magnetohydrodynamics and gas dynamics I: Lie dragging approach. *Journal of Physics A: Mathematical and Theoretical*, 47(9):095501, 2014.
- [18] G. M. Webb, B. Dasgupta, J. F. McKenzie, Q. Hu, and G. P. Zank. Local and nonlocal advected invariants and helicities in magnetohydrodynamics and gas dynamics II: Noether's theorems and Casimirs. *Journal of Physics A: Mathematical and Theoretical*, 47(9):095502, 2014.
- [19] F. Gay-Balmaz, J. E. Marsden, and T. S. Ratiu. Reduced variational formulations in free boundary continuum mechanics. *J. Nonlin. Sci.*, 22(4):463–497, 2012.
- [20] P. Makaremi-Esfarjani and A. Najafi-Yazdi. Characteristic boundary conditions for magnetohydrodynamic equations. *Computers & Fluids*, 241:105461, 2022. ISSN 0045-7930. doi: https://doi.org/10.1016/j.compfluid.2022.105461. URL <https://www.sciencedirect.com/science/article/pii/S0045793022001116>.
- [21] E. T. Meier, A. H. Glasser, V. S. Lukin, and U. Shumlak. Modeling open boundaries in dissipative mhd simulation. *Journal of Computational Physics*, 231(7):2963– 2976, 2012. ISSN 0021-9991. doi: https://doi.org/10.1016/j.jcp.2012.01.003. URL <https://www.sciencedirect.com/science/article/pii/S0021999112000137>.
- [22] H. C. Ottinger. Nonequilibrium thermodynamics for open systems. ¨ *Physical Review E—Statistical, Nonlinear, and Soft Matter Physics*, 73(3):036126, 2006.
- [23] A. Moses Badlyan, B. Maschke, C. Beattie, and V. Mehrmann. Open physical systems: from GENERIC to port-Hamiltonian systems, 2018. URL <https://arxiv.org/abs/1804.04064>.
- [24] R. Rashad, F. Califano, F. P. Schuller, and S. Stramigioli. Port-Hamiltonian modeling of ideal fluid flow: Part I. Foundations and kinetic energy. *Journal of Geometry and Physics*, 164:104201, 2021.
- [25] R. Rashad, F. Califano, F. P. Schuller, and S. Stramigioli. Port-Hamiltonian modeling of ideal fluid flow: Part II. Compressible and incompressible flow. *Journal of Geometry and Physics*, 164:104199, 2021.
- [26] M. Lohmayer and S. Leyendecker. Exergetic port-Hamiltonian systems: Navier-Stokes-Fourier fluid. *IFAC-PapersOnLine*, 55(18):74–80, 2022.
- [27] L. A. Mora, Y. Le Gorrec, D. Matignon, and H. Ramirez. Irreversible port-Hamiltonian modelling of 3D compressible fluids. *IFAC-PapersOnLine*, 56(2):6394–6399, 2023.
- [28] F. L. Cardoso-Ribeiro, D. Matignon, and L. Lefèvre. Dissipative shallow water equations: A port-Hamiltonian formulation. *IFAC-PapersOnLine*, 54(19):167–172, 2021.
- [29] G. Haine and D. Matignon. Incompressible Navier-Stokes equation as port-Hamiltonian systems: velocity formulation versus vorticity formulation. *IFAC-PapersOnLine*, 54(19): 161–166, 2021.
- [30] R. Rashad, F. Califano, A. Brugnoli, F. P. Schuller, and S. Stramigioli. Exterior and vector calculus views of incompressible Navier-Stokes port-Hamiltonian models. *IFACpapersonline*, 54(19):173–179, 2021.
- [31] F. Califano, R. Rashad, F. P. Schuller, and S. Stramigioli. Geometric and energy-aware decomposition of the Navier–Stokes equations: A port-Hamiltonian approach. *Physics of fluids*, 33(4), 2021.
- [32] V. Arnold. Sur la géométrie différentielle des groupes de Lie de dimension infinie et ses applications `a l'hydrodynamique des fluides parfaits. In *Annales de l'institut Fourier*, volume 16, pages 319–361, 1966.
- [33] J. E. Marsden and T. J. R. Hughes. *Mathematical foundations of elasticity*. Courier Corporation, 2012.
- [34] F. Gay-Balmaz. General relativistic Lagrangian continuum theories part I: reduced variational principles and junction conditions for hydrodynamics and elasticity. *Journal of Nonlinear Science*, 34(3):46, 2024.
- [35] F. Gay-Balmaz and H. Yoshimura. From Lagrangian mechanics to nonequilibrium thermodynamics: a variational perspective. *Entropy*, 21:8, 2019.
- [36] R. L. Seliger and F. R. S. Whitham. Variational principles in continuum mechanics. *Proc. Roy. Soc. A*, 305:1–25, 1968.
- [37] F. Gay-Balmaz and H. Yoshimura. From variational to bracket formulations in nonequilibrium thermodynamics of simple systems. *J. Geom. Phys.*, 158:103812, 2020.
- [38] F. Gay-Balmaz and V. Putkaradze. Geometric theory of flexible and expandable tubes conveying fluid: equations, solutions, and shock waves. *Journal of Nonlinear Science*, 29 (2):377–414, 2019.
- [39] J. M. Lee. *Smooth manifolds*. Springer, 2012.
- [40] F. Bouchut, J. Lambaerts, G. Lapeyre, and V. Zeitlin. Fronts and nonlinear waves in a simplified shallow-water model of the atmosphere with moisture and convection. *Physics of Fluids*, 21(11), 2009.
- [41] J. Serrin. On the uniqueness of compressible fluid motions. *Archive for Rational Mechanics and Analysis*, 3:271–288, 1959.
- [42] T. Dubos and M. Tort. Equations of atmospheric motion in non-eulerian vertical coordinates: Vector-invariant form and quasi-hamiltonian formulation. *Monthly Weather Review*, 142(10):3860–3880, 2014.
- [43] C. Eldred and F. Gay-Balmaz. Single and double generator bracket formulations of multicomponent fluids with irreversible processes. *Journal of Physics A: Mathematical and Theoretical*, 53(39):395701, 2020.
- [44] F. Gay-Balmaz. A variational derivation of the thermodynamics of a moist atmosphere with rain process and its pseudoincompressible approximation. *Geophysical & Astrophysical Fluid Dynamics*, 113(5-6):428–465, 2019.
- [45] A. M. Makarieva, V. G. Gorshkov, A. V. Nefiodov, D. Sheil, A. D. Nobre, P. Bunyard, P. Nobre, and B. Li. The equations of motion for moist atmospheric air. *Journal of Geophysical Research: Atmospheres*, 122(14):7300–7307, 2017.
- [46] A. Bott. Theoretical considerations on the mass and energy consistent treatment of precipitation in cloudy atmospheres. *Atmospheric Research*, 89(3):262–269, 2008.
- [47] B. Catry, J.-F. Geleyn, M. Tudor, P. Bénard, and A. Trojáakováa. Flux-conservative thermodynamic equations in a mass-weighted framework. *Tellus A: Dynamic Meteorology and Oceanography*, 59(1):71–79, 2007.
- [48] U. Wacker and F. Herbert. Continuity equations as expressions for local balances of masses in cloudy air. *Tellus A: Dynamic Meteorology and Oceanography*, 55(3):247–254, 2003.
- [49] P. R. Bannon. Theoretical foundations for models of moist convection. *Journal of the Atmospheric Sciences*, 59(12):1967–1982, 2002.
- [50] F. G. Helfferich. Theory of multicomponent, multiphase displacement in porous media. *Society of Petroleum Engineers Journal*, 21(01):51–62, 1981.
- [51] M. Quintard, L. Bletzacker, D. Chenu, and S. Whitaker. Nonlinear, multicomponent, mass transport in porous media. *Chemical Engineering Science*, 61(8):2643–2669, 2006.
- [52] R. Chella, D. Lasseux, and M. Quintard. Multiphase, multicomponent fluid flow in homogeneous and heterogeneous porous media. *Revue de l'Institut Français du Pétrole*, 53(3): 335–346, 1998.
- [53] T. Farkhutdinov, F. Gay-Balmaz, and V. Putkaradze. Geometric variational approach to the dynamics of porous media filled with incompressible fluid. *Acta Mechanica*, 431(9): 3897–3924, 2020.
- [54] F. Gay-Balmaz and V. Putkaradze. Variational geometric approach to the thermodynamics of porous media. *Zeitschrift f¨ur Angewandte Mathematik und Mechanik*, 102(11), 2022.
- [55] F. Gay-Balmaz and T. S. Ratiu. The geometric structure of complex fluids. *Advances in Applied Mathematics*, 42(2):176–275, 2009.
- [56] D. D. Holm. Euler-Poincar´e dynamics of perfect complex fluids. In *Geometry, mechanics, and dynamics*, pages 169–180. Springer, 2002.
- [57] D. D. Holm and B. A. Kupershmidt. Poisson brackets and Clebsch representations for magnetohydrodynamics, multifluid plasmas, and elasticity. *Physica D: Nonlinear Phenomena*, 6(3):347–363, 1983.
- [58] D. D. Holm and B. A. Kupershmidt. Hydrodynamics and electrohydrodynamics of adiabatic multiphase fluids and plasmas. *International journal of multiphase flow*, 12(4): 667–680, 1986.
- [59] D. D. Holm. Hamiltonian dynamics of a charged fluid, including electro-and magnetohydrodynamics. *Physics Letters A*, 114(3):137–141, 1986.
- [60] P. J. Morrison. Hamiltonian and action principle formulations of plasma physics. *Physics of plasmas*, 12(5), 2005.
- [61] E. Kanso, M. Arroyo, Y. Tong, A. Yavari, J. E. Marsden, and M. Desbrun. On the geometric character of stress in continuum mechanics. Zeitschrift für Angewandte Mathematik *und Mechanik*, 58:1–14, 2007.
- [62] A. D. Gilbert and J. Vanneste. A geometric look at momentum flux and stress in fluid mechanics. *Journal of Nonlinear Science*, 33(2):31, 2023.
- [63] R. Rashad, A. Brugnoli, F. Califano, E. Luesink, and S. Stramigioli. Intrinsic nonlinear elasticity: An exterior calculus formulation. *Journal of Nonlinear Science*, 33(5):84, 2023.
- [64] D. Bresch, M. Gisclon, and I. Lacroix-Violet. On Navier–Stokes–Korteweg and Euler– Korteweg systems: application to quantum fluids models. *Archive for Rational Mechanics and Analysis*, 233:975–1025, 2019.
- [65] E. B. Kraus and J. A. Businger. *Atmosphere-Ocean Interaction*, volume 27. Oxford University Press, 1994.