ON REES ALGEBRAS OF IDEALS AND MODULES WITH WEAK RESIDUAL CONDITIONS

ALESSANDRA COSTANTINI, EDWARD F. PRICE III, AND MATTHEW WEAVER

ABSTRACT. Let E be a module of projective dimension one over $R = k[x_1, \ldots, x_d]$. If E is presented by a matrix φ with linear entries and the number of generators of E is bounded locally up to codimension d-1, the Rees ring $\mathcal{R}(E)$ is well understood. In this paper, we study $\mathcal{R}(E)$ when this generation condition holds only up to codimension s-1, for some s < d. Moreover, we provide a generating set for the ideal defining this algebra by employing a method of successive approximations of the Rees ring. Although we employ techniques regarding Rees rings of modules, our findings recover and extend known results for Rees algebras of perfect ideals with grade two in the case that rank E = 1.

1. INTRODUCTION

For $I = (f_1, \ldots, f_n)$ an ideal in a Noetherian ring R, the Rees ring of I is the subring $\mathcal{R}(I) = R[It] = R[f_1t, \ldots, f_nt]$ of the polynomial ring R[t], for t an indeterminate. As a graded algebra, one also has that $\mathcal{R}(I) = R \oplus It \oplus I^2 t^2 \oplus \cdots$. From this latter description, one sees that $\mathcal{R}(I)$ carries information on every power I^j and their asymptotic behavior for large exponents $j \gg 0$. As such, the Rees ring and its associated algebras have proven to be indispensable within the study of reductions and various multiplicities. In the geometric setting, the Rees ring $\mathcal{R}(I)$ is often called the *blowup algebra*, as $\operatorname{Proj}(\mathcal{R}(I))$ is precisely the blowup of $\operatorname{Spec}(R)$ along the subscheme V(I). Alternatively, if $R = k[x_1, \ldots, x_d]$ and the polynomials f_1, \ldots, f_n are homogeneous of a common degree, the Rees algebra serves as the coordinate ring of graph(Φ) where $\Phi : \mathbb{P}_k^{d-1} \longrightarrow \mathbb{P}_k^{n-1}$ is the rational map defined by f_1, \ldots, f_n . We may also extend this notion to Rees algebras of modules in order to treat the case of repeated or successive blowups, or also compositions of rational maps between projective varieties. Indeed, the successive blowup of an affine scheme along disjoint subschemes V(I) and V(J) corresponds to the Rees ring $\mathcal{R}(I \oplus J)$.

For an *R*-module *E*, the Rees ring $\mathcal{R}(E)$ is defined as $\mathcal{R}(E) = \mathcal{S}(E)/\tau(\mathcal{S}(E))$ where $\mathcal{S}(E)$ is the symmetric algebra of *E* and $\tau(\mathcal{S}(E))$ is its *R*-torsion submodule. Although seemingly different, this recovers the previous notion when *E* is actually an *R*-ideal. Since identifying the torsion of $\mathcal{S}(E)$ is seldom a simple task, it is typically more advantageous to describe $\mathcal{R}(E)$ as a quotient of a polynomial ring $R[T_1, \ldots, T_n]$. The ideal \mathcal{J} defining this quotient is aptly called the *defining ideal* of $\mathcal{R}(E)$ and its generators are called the *defining equations* of $\mathcal{R}(E)$.

In general, determining the equations of \mathcal{J} is an arduous task, however there has been notable success for Rees rings of perfect ideals with small codimension [3, 10, 12, 21, 22, 25, 26, 27, 28, 37, 38] and Rees algebras of modules with small projective dimension [8, 10, 24, 31, 37, 39] in a multitude of settings. Many of these results share common assumptions, most notably the *residual condition* G_d , where $d = \dim R$. This condition was introduced by Artin and Nagata in [1], to study residual intersections of an ideal (see Section 2). For E a module with rank e, one says that E satisfies the condition G_s if $\mu(E_p) \leq \dim R_p + e - 1$ for every $\mathfrak{p} \in \operatorname{Spec}(R)$ with $1 \leq \dim R_p \leq s - 1$. Additionally, E satisfies G_{∞} if E satisfies G_s for all s. Here $\mu(E_p)$ denotes the minimal number of generators of E_p .

Although seemingly unassuming, the condition G_d where $d = \dim R$ is quite powerful within the study of Rees algebras, as it often dictates the prime ideals of R at which $\mathcal{R}(E)$ and $\mathcal{S}(E)$ coincide locally. As such, this assumption can seldom be weakened when comparing these two algebras. However, there are many classes of ideals and modules that do not satisfy this condition, yet have notable Rees rings. For example, let $\Omega_{R/k}$ denote the module of Kähler differentials of a complete intersection ring $R = k[x_1, \ldots, x_n]/I$ with dim $R = d \ge 2$. The Rees algebra $\mathcal{R}(\Omega_{R/k})$ and its related rings are called *tangent algebras* due to their connections with tangential varieties arising in algebraic geometry, and have been well studied [32]. However, the module $\Omega_{R/k}$ often fails to satisfy G_d , for instance if R is not a normal ring [39, Sec. 6]. Additionally, classes of examples of grade-two perfect ideals that do not satisfy G_d can easily be found, for instance in [9, 12].

While the condition G_d is essential for most arguments, there has been success in determining the defining ideal of Rees rings of perfect ideals with grade two satisfying G_{d-1} [12, 27, 28]. Much of this work was extended more generally to modules of projective dimension one satisfying G_{d-1} in our recent paper [10]. In each of these instances, the weaker residual condition is supplemented with a strict rank condition on a presentation matrix φ , which appears difficult to relax. The objective of this paper is to further extend this previous work to modules of projective dimension one satisfying G_s for any s < d. Our main results Theorem 4.5 and Theorem 5.3 are summarized as follows.

Theorem A. Let $R = k[x_1, \ldots, x_d]$ be a polynomial ring with $d \ge 3$ over a field k, and let E be a R-module of projective dimension one and rank E = e. Assume that E is minimally generated by $\mu(E) = n \ge d + e$ elements and E satisfies G_s , but not G_{s+1} , for some integer $2 \le s \le d - 1$. Assume furthermore that E has a presentation matrix φ consisting of linear entries in R with $I_1(\varphi) = (x_1, \ldots, x_d)$ and that, after possibly a change of coordinates, modulo (x_1, \ldots, x_s) the matrix $\overline{\varphi}$ has rank 1.

(a) If the nonzero entries of $\overline{\varphi}$ are in one column, the Rees algebra of E is $\mathcal{R}(E) \cong \mathbb{R}[T_1, \ldots, T_n]/\mathcal{J}$ with

$$\mathcal{J} = (\ell_1, \dots, \ell_{n-e}) : (x_1, \dots, x_s) = (\ell_1, \dots, \ell_{n-e}) + I_s(M)$$

where $\ell_1, \ldots, \ell_{n-e}$ are linear forms such that $[\ell_1 \ldots \ell_{n-e}] = [T_1 \ldots T_n] \cdot \varphi$ and M is an $s \times (n-e-1)$ matrix with entries in $k[T_1, \ldots, T_n]$ such that $[\ell_1 \ldots \ell_{n-e-1}] = [x_1 \ldots x_s] \cdot M$.

(b) If the nonzero entries of $\overline{\varphi}$ are in one row, the Rees algebra of E is $\mathcal{R}(E) \cong R[T_1, \ldots, T_n]/\mathcal{J}$ with

$$\mathcal{J} = (\ell_1, \dots, \ell_{n-e}) : (x_1, \dots, x_s) = (\ell_1, \dots, \ell_{n-e}) + I_s(M) + I_{s+1}(C)$$

where $\ell_1, \ldots, \ell_{n-e}$ are linear forms such that $[\ell_1 \ldots \ell_{n-e}] = [T_1 \ldots T_n] \cdot \varphi$. Moreover, M is an $s \times (n-e-d+s)$ matrix with entries in $k[T_1, \ldots, T_n]$ such that $[\ell_1 \ldots \ell_{n-e-d+s}] = [x_1 \ldots x_s] \cdot M$ and C is an $(s+1) \times (n-e)$ matrix such that $[\ell_1 \ldots \ell_{n-e}] = [x_1 \ldots x_s \ T_n] \cdot C$.

Moreover, in either case, the Rees algebra $\mathcal{R}(E)$ is a Cohen-Macaulay domain of dimension d + e.

With the condition that rank $\overline{\varphi} = 1$, the nonzero entries of $\overline{\varphi}$ are concentrated in either a single row or a single column. Interestingly, $\mathcal{R}(E)$ is Cohen-Macaulay in either case, but the shape of the defining ideal differs in the two settings. However, in either case, the matrices M and C which provide the nontrivial equations are obtained from submatrices of the Jacobian dual matrix of the presentation φ (see Section 2).

Not only does the Rees ring $\mathcal{R}(E)$ differ in each of the cases above, but the *special fiber ring* $\mathcal{F}(E) := \mathcal{R}(E) \otimes_R k$ does as well. In particular, the *analytic spread* of E, i.e. the Krull dimension $\ell(E) := \dim \mathcal{F}(E)$ differs in each case. Indeed, as a consequence of Theorem A, we obtain the result below, which is a compilation of Corollary 4.6 and Corollary 5.12.

Theorem B. With the assumptions of Theorem A, we have the following.

- (a) If the nonzero entries of $\overline{\varphi}$ are in one column, the special fiber ring is $\mathcal{F}(E) \cong k[T_1, \ldots, T_n]/I_s(M)$. Moreover, $\mathcal{F}(E)$ is a Cohen-Macaulay domain of dimension $\ell(E) = s + e$.
- (b) If the nonzero entries of $\overline{\varphi}$ are in one row, the special fiber ring is $\mathcal{F}(E) \cong k[T_1, \ldots, T_n]/I_s(M)$. Moreover, $\mathcal{F}(E)$ is a Cohen-Macaulay domain of dimension $\ell(E) = d + e - 1$.

We note that the Cohen-Macaulay property of $\mathcal{F}(E)$ does not depend on the two cases of Theorem B, but the analytic spread of E does. However, the two phenomena do coincide when s = d - 1, recovering our findings in [10]. Additionally, when rank E = 1, the module E is isomorphic to an R-ideal I, hence Theorem A and Theorem B are applicable to Rees algebras of perfect ideals of grade two. In particular, when e = 1 and s = d - 1, Theorem A and Theorem B recover the main results of [27, 28]. Moreover, to the best of our knowledge, our findings provide the first known results for Rees algebras of ideals and modules satisfying G_s for $s \leq d - 2$.

In [10], the primary technique to study the Rees ring was to produce a generic Bourbaki ideal I of E, reducing the study of $\mathcal{R}(E)$ to the study of $\mathcal{R}(I)$, where information is more accessible. However, in the present paper, we forgo this technique and instead employ a method of successive approximations of the

Rees algebra $\mathcal{R}(E)$, as the essential ingredient to the proofs of Theorem A and Theorem B. From a free presentation $\mathbb{R}^m \xrightarrow{\varphi} \mathbb{R}^n \to E \to 0$, one factors the map $\mathbb{R}^n \to E$ into a sequence of epimorphisms

$$R^n = E_m \twoheadrightarrow E_{m-1} \twoheadrightarrow \cdots \twoheadrightarrow E_0 = E$$

which induces a sequence of epimorphisms

$$R[T_1, \ldots, T_n] = \mathcal{R}(E_m) \twoheadrightarrow \mathcal{R}(E_{m-1}) \twoheadrightarrow \cdots \longrightarrow \mathcal{R}(E_0) = \mathcal{R}(E)$$

factoring the natural map $\pi : R[T_1, \ldots, T_n] \twoheadrightarrow \mathcal{R}(E)$. If one is able to understand these intermediate algebras, and their defining ideals in particular, then one has a much better chance of determining the defining ideal of $\mathcal{R}(E)$. We remark that this is not an entirely novel approach, and this technique originates in [24]. Moreover, we note that variations of this technique have been previously applied in [3, 21, 37, 38]. However, our main contribution is to show that the construction of these approximation modules is compatible with the residual condition G_s , when E has projective dimension one. Moreover, we describe how the shape of the defining ideal of an intermediate ring above can be deduced from the previous algebra.

We now briefly describe how this paper is organized. In Section 2, we review the necessary preliminary material on Rees rings of modules and their associated algebras, as well as residual intersections of ideals. We also introduce the method of successive approximations of the Rees algebra $\mathcal{R}(E)$ and how to implement this construction in the study of the defining ideal. In Section 3, we discuss the primary setting for the paper and identify the two cases that must be considered, stemming from the rank condition in Theorem A. The remainder of the section is spent discussing the similarities between the two cases, while the proceeding sections are spent analyzing their differences. In Sections 4 and 5, we determine the defining ideal \mathcal{J} of the Rees ring $\mathcal{R}(E)$ in each of the two cases. With this, the Cohen-Macaulay and fiber-type property are explored in each case. Finally, in Section 6 we include several examples showing that the conclusion of Theorem A may not necessarily hold if we weaken any of the assumptions.

2. Preliminaries

In this section, we recall background information on modules with a rank and their Rees algebras, needed throughout the article. In particular, we include crucial observations regarding modules of projective dimension one in Section 2.2. For the present setting, assume that R is a Noetherian ring and E is a finitely generated R-module. We note that an ideal of positive grade is isomorphic to a torsion-free module of rank one, hence all statements discussed here apply to Rees rings of ideals when rank E = 1.

2.1. Modules and their Rees algebras. Recall that E is said to have a rank if $E \otimes_R Quot(R) \cong Quot(R)^e$ for some integer e, where Quot(R) denotes the total ring of quotients of R; in this case, we write rank(M) = e. In particular, free modules have a rank, hence modules of finite projective dimension do as well, since rank is additive along exact sequences.

Given a generating set $E = Ra_1 + \cdots + Ra_n$, such a module admits a corresponding free presentation

$$R^m \xrightarrow{\varphi} R^n \longrightarrow E \longrightarrow 0. \tag{2.1}$$

Recall that, for $0 \leq i \leq n$, the *i*th *Fitting ideal* of *E* is the ideal $\text{Fitt}_i(E) \coloneqq I_{n-i}(\varphi)$, i.e. the ideal generated by all $(n-i) \times (n-i)$ minors of φ . We remark that these ideals are particularly useful invariants of the module *E*.

Proposition 2.1 ([14, 20.4–20.6]). With a module E as above, we have the following.

- (a) Fitt_i(E) depends only on the module E and the index i. In particular, it is independent of the choice of presentation.
- (b) If R is local, then $\operatorname{Fitt}_i(E) = R$ if and only if $\mu(E) \leq i$.
- (c) $V(\operatorname{Fitt}_i(E)) = \{ \mathfrak{p} \in \operatorname{Spec}(R) \mid \mu(E_{\mathfrak{p}}) \ge i+1 \}.$

Condition (c) above is particularly useful as it easily relates to the condition G_s from the introduction.

Remark 2.2. A module E with rank E = e satisfies G_s if and only if $\operatorname{ht} \operatorname{Fitt}_i(E) \ge i - e + 2$ for all $e \le i \le s + e - 2$. As before, we say that E satisfies G_∞ if E satisfies G_s for all s.

In addition to the Fitting invariants, a free presentation of E also provides information on the symmetric algebra of E, S(E). More precisely, if $E = Ra_1 + \ldots + Ra_n$ and (2.1) is a presentation corresponding to this generating set, there is a natural homogeneous epimorphism of graded R-algebras

$$\eta \colon R[T_1, \dots, T_n] \longrightarrow \mathcal{S}(E) \tag{2.2}$$

defined by mapping each T_i to $a_i \in [\mathcal{S}(E)]_1$, and extending *R*-linearly. Moreover, the kernel $\mathcal{L} = \ker(\eta)$ is generated by linear forms ℓ_1, \ldots, ℓ_m such that $[T_1 \ldots T_n] \cdot \varphi = [\ell_1 \ldots \ell_m]$. Thus, there is an induced isomorphism $\mathcal{S}(E) \cong R[T_1, \ldots, T_n]/\mathcal{L}$. We remark that this description is independent of the choice of presentation (2.1).

As noted in the introduction, the *Rees algebra* of E is $\mathcal{R}(E) = \mathcal{S}(E)/\tau(\mathcal{S}(E))$, where $\tau(\mathcal{S}(E))$ is the R-torsion submodule of $\mathcal{S}(E)$. If $E \cong I$ for an R-ideal I, the Rees algebra $\mathcal{R}(E)$ is isomorphic to the subalgebra $\mathcal{R}(I) = R[It] \subseteq R[t]$ defined in the introduction. If E is an R-module with rank e, the Krull dimension of $\mathcal{R}(E)$ is dim $\mathcal{R}(E) = d + e$, where $d = \dim R$ (see, e.g., [31, 2.2]).

Composing the map η in (2.2) with the natural map factoring $\tau(\mathcal{S}(E))$, one obtains a second homogeneous epimorphism

$$\pi \colon R[T_1, \dots, T_n] \longrightarrow \mathcal{R}(E) \tag{2.3}$$

mapping $T_i \mapsto a_i \in [\mathcal{R}(E)]_1$. There is an induced isomorphism $\mathcal{R}(E) \cong R[T_1, \ldots, T_n]/\mathcal{J}$, where $\mathcal{J} \coloneqq \ker(\pi)$ is the *defining ideal* of $\mathcal{R}(E)$. By construction, it is clear that $\mathcal{L} \subseteq \mathcal{J}$. Moreover, \mathcal{L} is actually the degree-one component of \mathcal{J} , $\mathcal{L} = [\mathcal{J}]_1$ [36]. One says that E is of *linear type* if $\mathcal{L} = \mathcal{J}$, as in this case all generators of \mathcal{J} are linear forms.

If E is not of linear type, the higher-degree generators of \mathcal{J} can often be detected by means of a Jacobian dual matrix of the presentation φ of E, or via the special fiber ring $\mathcal{F}(E)$ of E. We next briefly recall both of these notions.

Definition 2.3. Let *E* be a module with presentation (2.1) and let ℓ_1, \ldots, ℓ_m be the generators of \mathcal{L} as before. There exists an $r \times m$ matrix $B(\varphi)$ with linear entries in $R[T_1, \ldots, T_n]$ such that

$$[T_1 \dots T_n] \cdot \varphi = [\ell_1 \dots \ell_m] = [x_1 \dots x_r] \cdot B(\varphi), \qquad (2.4)$$

where (x_1, \ldots, x_r) is an ideal containing $I_1(\varphi)$. The matrix $B(\varphi)$ is a Jacobian dual of φ , with respect to the sequence x_1, \ldots, x_r .

We remark that $B(\varphi)$ is not unique in general. However, if $R = k[x_1, \ldots, x_d]$, then there is a unique Jacobian dual matrix $B(\varphi)$, with respect to x_1, \ldots, x_d , if and only if the entries of φ are linear [29, p. 47]. In this case, the entries of $B(\varphi)$ belong to the subring $k[T_1, \ldots, T_n]$, a fact which we will exploit in several proofs in later sections of this article.

Definition 2.4. Assume that R is a local ring with maximal ideal \mathfrak{m} and residue field k. Let E be a finitely generated R-module as above. The *special fiber ring* of E is

$$\mathcal{F}(E) \coloneqq \mathcal{R}(E) \otimes_R k \cong \mathcal{R}(E)/\mathfrak{m}\mathcal{R}(E).$$

The Krull dimension of $\mathcal{F}(E)$ is called the *analytic spread* of E and is denoted by $\ell(E)$.

This notion may also be adapted to the graded setting when R is a standard graded polynomial ring $R = k[x_1, \ldots, x_d]$, over a field k. In this setting, the special fiber ring is similarly defined using the unique homogeneous maximal ideal $\mathfrak{m} = (x_1, \ldots, x_d)$.

The natural map π of (2.3) induces an epimorphism of graded k-algebras

$$\psi \colon R[T_1, \dots, T_n] \otimes_R k \cong k[T_1, \dots, T_n] \longrightarrow \mathcal{F}(E) \cong \mathcal{R}(E)/\mathfrak{m}\mathcal{R}(E)$$

such that $\psi(T_i) = a_i + \mathfrak{mR}(E) \in [\mathcal{F}(E)]_1$. In particular, $\ker(\psi)R[T_1, \ldots, T_n] + \mathcal{L} \subseteq \mathcal{J}$ and we say that E is of *fiber type* if this containment is an equality.

2.2. Successive approximations of Rees algebras. In this subsection, we discuss a useful technique used to approximate Rees algebras of modules of projective dimension one, expanding on the method introduced in [24]. Our setting throughout is as follows.

Setting 2.5. Let R be a Noetherian ring, and let E be an R-module of rank e having projective dimension one and minimally generated by $\mu(E) = n$ forms. With this, we may write

$$0 \longrightarrow R^{n-e} \xrightarrow{\varphi} R^n \longrightarrow E \longrightarrow 0 \tag{2.5}$$

to denote a minimal resolution of E. Let φ_i denote the submatrix of φ obtained by deleting the last i columns of φ and let $E_i = \operatorname{coker} \varphi_i$.

Notice that there is a chain of epimorphisms

$$R^n = E_{n-e} \twoheadrightarrow E_{n-e-1} \twoheadrightarrow \dots \twoheadrightarrow E_0 = E.$$

$$(2.6)$$

factoring the map $\mathbb{R}^n \twoheadrightarrow \mathbb{E}$ in (2.5). With the sequence of modules above, we make the following observation.

Lemma 2.6. With the assumptions of Setting 2.5 and the modules in (2.6), we have the following.

- (a) For $1 \le i \le n e 1$, each E_i is an *R*-module of projective dimension one with rank $E_i = e + i$.
- (b) If E_i satisfies G_s , then E_j satisfies G_s as well, for all $j \ge i$.

Proof. To verify (a) it suffices to show that φ_i is an injective map for each *i* in the given range. However, this is clear from linear algebra, as φ is injective. Now that φ_i is seen to be injective, the short exact sequence

$$0 \longrightarrow R^{n-e-i} \xrightarrow{\varphi_i} R^n \longrightarrow E_i \longrightarrow 0$$
(2.7)

shows that E_i has projective dimension one and rank $E_i = e + i$, by additivity of rank.

To prove (b), it suffices to show that if $E = E_0$ satisfies G_s , then E_1 does as well. The claim then follows by induction after reindexing. Since E satisfies G_s and $\operatorname{Fitt}_i(E) \subseteq \operatorname{Fitt}_{i+1}(E_1)$, one has that ht $\operatorname{Fitt}_{i+1}(E_1) \ge i - e + 2$ for all $e \le i \le s + e - 2$. After adjusting the indices, we see that ht $\operatorname{Fitt}_i(E_1) \ge i - (e+1) + 2$ for all $e + 1 \le i \le s + (e+1) - 2$. Since $\operatorname{rank} E_1 = e + 1$, it follows that E_1 satisfies G_s .

Remark 2.7. From Lemma 2.6 it follows that if the module E satisfies G_s , then each of the modules in (2.6) satisfy G_s as well. However, it is possible that some of the modules satisfy G_t for t > s. For instance, $E_{n-e} = \mathbb{R}^n$ is free and hence satisfies G_{∞} .

Notice that the sequence of epimorphisms in (2.6) induces a sequence of homogeneous epimorphisms of symmetric algebras:

$$R[T_1, \dots, T_n] = \mathcal{S}(E_{n-e}) \twoheadrightarrow \mathcal{S}(E_{n-e-1}) \twoheadrightarrow \dots \twoheadrightarrow \mathcal{S}(E_0) = \mathcal{S}(E), \qquad (2.8)$$

which factors the natural map in (2.2). Moreover, this induces a sequence of homogeneous epimorphisms of Rees algebras:

$$R[T_1, \dots, T_n] = \mathcal{R}(E_{n-e}) \twoheadrightarrow \mathcal{R}(E_{n-e-1}) \twoheadrightarrow \dots \twoheadrightarrow \mathcal{R}(E_0) = \mathcal{R}(E)$$
(2.9)

by further factoring *R*-torsion, which also factors the natural map in (2.3). In particular, the ideals defining the intermediate algebras of (2.8) and (2.9) relate to each other.

Notation 2.8. Write \mathcal{L}_i to denote the defining ideal of $\mathcal{S}(E_i)$, i.e. the kernel of the epimorphism

$$R[T_1,\ldots,T_n] \longrightarrow \mathcal{S}(E_i),$$

and write \mathcal{J}_i to denote the defining ideal of $\mathcal{R}(E_i)$, i.e. the kernel of the epimorphism

$$R[T_1,\ldots,T_n] \longrightarrow \mathcal{R}(E_i).$$

Writing \mathcal{L} to denote the defining ideal of $\mathcal{S}(E)$, with the presentation of E in (2.5) note that $\mathcal{L} = (\ell_1, \ldots, \ell_{n-e})$ where $[\ell_1 \ldots \ell_{n-e}] = [T_1 \ldots T_n] \cdot \varphi$. Moreover, the defining ideal of $\mathcal{S}(E_i)$ is precisely $\mathcal{L}_i = (\ell_1, \ldots, \ell_{n-e-i})$ as E_i is presented by (2.7) and φ_i is a submatrix of φ . Hence there is a chain of inclusions

$$0 = \mathcal{L}_{n-e} \subsetneq \mathcal{L}_{n-e-1} \subsetneq \cdots \cdots \subsetneq \mathcal{L}_1 \subsetneq \mathcal{L}_0 = \mathcal{L}_0$$

where the generators of the ideals \mathcal{L}_i are well understood.

Unfortunately, the ideals \mathcal{J}_i are not as simple. However, we have a chain of inclusions

$$0 = \mathcal{J}_{n-e} \subsetneq \mathcal{J}_{n-e-1} \subsetneq \cdots \cdots \subsetneq \mathcal{J}_1 \subsetneq \mathcal{J}_0 = \mathcal{J},$$

and we note that, if R is a domain, for each i the Rees ring $\mathcal{R}(E_i)$ is a domain of dimension d + e + i, hence each \mathcal{J}_i is a prime ideal of height n - e - i. Moreover, each successive quotient $\mathcal{J}_i/\mathcal{J}_{i+1}$ fits into the short exact sequence

$$0 \longrightarrow \mathcal{J}_i/\mathcal{J}_{i+1} \longrightarrow \mathcal{R}(E_{i+1}) \longrightarrow \mathcal{R}(E_i) \longrightarrow 0.$$

The quotients $\mathcal{J}_i/\mathcal{J}_{i+1}$ have the advantage of being simpler than the entire ideal \mathcal{J} , as they are prime $\mathcal{R}(E_{i+1})$ -ideals of height 1, whenever R is catenary. Moreover, if generating sets can be found for each $\mathcal{J}_i/\mathcal{J}_{i+1}$, then they can be lifted and combined to produce a generating set of \mathcal{J} .

2.3. Residual intersections. Lastly, we very briefly recall the notion of residual intersections of an ideal. This notion goes back to the work of Artin and Nagata [1], who informally also introduced the G_s condition; indeed, if an ideal I satisfies G_s , one can easily identify an optimal generating set for a residual intersection of I (see, e.g., [33, 1.6]). In the present paper, we aim to describe the structure of certain quotients $\mathcal{J}_i/\mathcal{J}_{i+1}$ in terms of residual intersections, for which we require very few technical aspects of the subject. Hence, we omit much of the unnecessary background material, and refer the curious reader to [19, 20] for a more rigorous treatment of the matter.

Definition 2.9 ([19, 1.1]). Let R be a Cohen-Macaulay local ring and I an R-ideal. For s an integer with $s \ge \operatorname{ht} I$, a proper ideal $J = \mathfrak{a} : I$ is an s-residual intersection of I if $\mathfrak{a} = (a_1, \ldots, a_s) \subseteq I$ and $\operatorname{ht} J \ge s$.

In particular, we restrict our attention to residual intersections of ideals generated by regular sequences. Not only do these ideals possess strong properties, their generation and resolutions are also well known [5].

Theorem 2.10. Let R be a Cohen-Macaulay local ring and let $I = (x_1, \ldots, x_g)$ be an ideal with x_1, \ldots, x_g an R-regular sequence. Let $J = \mathfrak{a} : I$ be an s-residual intersection of I for some $\mathfrak{a} = (a_1, \ldots, a_s) \subseteq I$.

- (a) ([20, 1.4 and 1.5]) R/J is a Cohen-Macaulay ring.
- (b) ([20, 1.5 and 1.8]) If B is a $g \times s$ matrix with $[a_1 \dots a_s] = [x_1 \dots x_g] \cdot B$, then $J = \mathfrak{a} + I_g(B)$.

3. Modules with weak residual conditions

We now begin our treatment of the Rees algebra of an R-module E with projective dimension one satisfying G_s for $s < d = \dim R$. We remark that we supplement the weakening of the condition G_d with a strict rank condition on a presentation matrix φ of E, similar to the settings of [10, 28]. We also note that for s < d-1, this rank assumption gives rise to two cases which must be treated separately. For the duration of the paper, we adopt the following setting.

Setting 3.1. Let $R = k[x_1, \ldots, x_d]$ be a standard-graded polynomial ring over a field k, with $\mathfrak{m} = (x_1, \ldots, x_d)$ and $d \ge 3$. Let s be an integer with $2 \le s \le d-1$. Let E be a R-module of projective dimension one and rank E = e minimally generated by $\mu(E) = n \ge d + e$ elements, satisfying the following assumptions:

- (i) The module E has a presentation matrix φ consisting of linear entries with $I_1(\varphi) = \mathfrak{m}$.
- (ii) After a possible change of coordinates, the matrix φ has rank 1 modulo $\mathfrak{p} = (x_1, \ldots, x_s)$.
- (iii) The module E satisfies G_s , but not G_{s+1} .

Condition (ii) is the aforementioned rank condition and is certainly the most strict of the assumptions above. However, we note that this condition holds automatically if s = d - 1 and n = d + e [10, 4.4]. Additionally, notice that as E satisfies G_2 , it is torsion-free. As a consequence, modules of rank one as in Setting 3.1 are isomorphic to notable ideals.

Remark 3.2. Under the assumptions of Setting 3.1, E has rank e = 1 if and only if E is isomorphic to a perfect ideal of grade two.

Proof. Let E be a module as in Setting 3.1 with rank e = 1. Since E satisfies G_s for some $s \ge 2$, it is torsion-free, and hence isomorphic to an R-ideal I. Moreover, as $E \cong I$ has projective dimension one, from (3.1) it follows that I is presented by a $n \times (n-1)$ matrix φ . Moreover, as I satisfies G_2 we have that $\operatorname{ht} I_{n-1}(\varphi) = \operatorname{ht} \operatorname{Fitt}_1(I) \ge 2$ and the claim follows from the Hilbert-Burch theorem [14, 20.15]. \Box

In particular, any result obtained in the present setting for Rees algebras of modules with projective dimension one can be applied to Rees algebras of perfect ideals of grade two.

The rank condition (ii) above has a profound implication on the structure of the matrix φ , as discussed in the following crucial remark.

Remark 3.3. With the assumptions of Setting 3.1, the module E has minimal free resolution of the form

$$0 \longrightarrow R^{n-e} \xrightarrow{\varphi} R^n \longrightarrow E \longrightarrow 0.$$
(3.1)

Moreover, with conditions (i) and (ii) above, after possible row and column operations, the matrix $\overline{\varphi}$ obtained from φ modulo (x_1, \ldots, x_s) has nonzero entries x_{s+1}, \ldots, x_d which are concentrated all in one row or all in one column. Hence, there are two possible shapes of the matrix φ . Namely, after row and column operations and a possible change of coordinates, the matrix φ may be taken to have either the form $\varphi = \varphi_C$ or $\varphi = \varphi_R$ where

In particular, the entries of the $(n - d + s) \times (n - e - 1)$ submatrix φ'_C and the $(n - 1) \times (n - e - d + s)$ submatrix φ'_R , as well as all the * entries, belong to the subring $A = k[x_1, \ldots, x_s] \subset R$.

As the indeterminates x_{s+1}, \ldots, x_d are either concentrated in a single column or a single row of φ , for the remainder of this paper we refer to these two cases as the *column setting* and the *row setting*, respectively. The next two sections are dedicated to a more thorough treatment of each case, highlighting substantial differences in the structure of the Rees algebra. Nevertheless, the two cases share several common features, which we discuss in the remainder of this section.

Remark 3.4. Notice that if s = d - 1, then the two possible shapes for the matrix φ in (3.2) coincide and Setting 3.1 recovers the setting of [10, sec. 4]. Moreover, following Remark 3.2, if s = d - 1 and additionally e = 1 and d = 3, then Setting 3.1 reduces to the setting of [28]. Likewise, when s = d - 1 and also e = 1and n = d + 1, one recovers the setting of [27].

In particular, in our study of the Rees ring $\mathcal{R}(E)$ our main interest will be in the case when s < d - 1, which will produce novel results, even in the rank-one case. We begin by identifying the *non-linear type locus* of the module E, i.e. the set of prime ideals $q \in \text{Spec}(R)$ for which E_q is not of linear type. We note that the following proposition holds more generally, outside of the assumptions of Setting 3.1.

Proposition 3.5. Let R be a Noetherian ring and E a finitely generated R-module of rank e and projective dimension one. If E satisfies G_s but not G_{s+1} , then $E_{\mathfrak{q}}$ is of linear type for all primes $\mathfrak{q} \in \operatorname{Spec}(R)$ with $\mathfrak{q} \notin V(\operatorname{Fitt}_{s+e-1}(E))$.

Proof. Notice that $V(\operatorname{Fitt}_{s+e-1}(E)) = \{ \mathfrak{q} \in \operatorname{Spec}(R) \mid \mu(E_{\mathfrak{q}}) \geq s+e \}$ by Proposition 2.1. Thus for any prime ideal \mathfrak{q} not contained in this set, we have that $\mu(E_{\mathfrak{q}}) \leq s+e-1$. Since E satisfies G_s , by transitivity of localization, for any $\mathfrak{p} \subseteq \mathfrak{q}$ it follows that $\mu((E_{\mathfrak{q}})_{\mathfrak{p}R_{\mathfrak{q}}}) = \mu(E_{\mathfrak{p}}) \leq \min\{\operatorname{ht}\mathfrak{p}, s\} + e - 1$, so $E_{\mathfrak{q}}$ satisfies G_{∞} and is hence of linear type by [2, Prop. 3 and 4].

Thus, we are reduced to investigate the minimal primes of $\operatorname{Fitt}_{s+e-1}(E)$. To this end, we recall a short lemma to assist us.

Lemma 3.6 ([10, 2.4]). Let R be a Noetherian ring and E a finitely generated R-module of rank e satisfying G_s but not G_{s+1} . Then ht Fitt_{s+e-2}(E) = ht Fitt_{s+e-1}(E) = s.

Now returning to the assumptions of Setting 3.1, we show that $\operatorname{Fitt}_{s+e-1}(E)$ has a *unique* minimal prime. This fact is crucial for the following arguments and, unlike the previous items, we note that this does require the strength of Setting 3.1. We present some examples in Section 6 of the behavior of $\mathcal{R}(E)$ outside of this setting, when this phenomenon does not occur. **Proposition 3.7.** With the assumptions of Setting 3.1, $\mathfrak{p} = (x_1, \ldots, x_s)$ is the unique minimal prime of Fitt_{s+e-1}(E) = $I_{n-s-e+1}(\varphi)$.

Proof. Since E satisfies G_s but not G_{s+1} , we have ht $I_{n-s-e+2}(\varphi) = \operatorname{ht} I_{n-s-e+1}(\varphi) = s$ by Lemma 3.6. We now consider two cases, depending on the two possible shapes of the matrix φ in (3.2).

For the column setting, let $\varphi = \varphi_C$ as in (3.2). Following the notation of Section 2, let φ_1 denote the $n \times (n - e - 1)$ submatrix of φ obtained by deleting the last column of φ . Then, the entries of φ_1 are linear forms in $A = k[x_1, \ldots, x_s]$, and also we have $I_{n-s-e+2}(\varphi) \subseteq I_{n-s-e+1}(\varphi_1) \subseteq (x_1, \ldots, x_s) = \mathfrak{p}$. Since ht $I_{n-s-e+2}(\varphi) = s = \operatorname{ht} \mathfrak{p}$, it follows that ht $I_{n-s-e+1}(\varphi_1) = s$. Thus, $I_{n-s-e+1}(\varphi_1)$ is a \mathfrak{p} -primary ideal in A, and so \mathfrak{p} is its only minimal prime in A and hence in R. On the other hand, since $I_{n-s-e+1}(\varphi_1) \subseteq I_{n-s-e+1}(\varphi) \subseteq \mathfrak{p}$ and ht $I_{n-s-e+1}(\varphi) = s$, it follows that \mathfrak{p} is the unique minimal prime of $I_{n-s-e+1}(\varphi)$ too.

For the row setting, let $\varphi = \varphi_R$ as in (3.2). A similar argument as above yields the same result, by considering the submatrix obtained by deleting the last row of φ .

With the non-linear type locus of E determined in Proposition 3.5, we may now introduce our first description of the defining ideal \mathcal{J} of the Rees ring $\mathcal{R}(E)$.

Proposition 3.8. With the assumptions of Setting 3.1, \mathcal{J} is a prime ideal of height n - e. Moreover, we have that $\mathcal{J} = \mathcal{L} : \mathfrak{p}^{\infty}$ where $\mathfrak{p} = (x_1, \ldots, x_s)$ as before.

Proof. The initial statement is clear as $\mathcal{R}(E) \cong R[T_1, \ldots, T_n]/\mathcal{J}$ and $\mathcal{R}(E)$ is a domain of dimension d + e. For the second statement, notice that $\mathcal{L} : (x_1, \ldots, x_s)^{\infty} \subseteq \mathcal{J}$ as \mathcal{J} is prime, $\mathcal{L} \subseteq \mathcal{J}$ and $(x_1, \ldots, x_s) \notin \mathcal{J}$. To prove the reverse containment, consider the quotient $\mathcal{A} = \mathcal{J}/\mathcal{L}$. It suffices to show that \mathcal{A} is annihilated by some power of $\mathfrak{p} = (x_1, \ldots, x_s)$. From Proposition 3.5 and Proposition 3.7, it follows that \mathcal{A} is supported only at primes containing \mathfrak{p} , hence \mathfrak{p} is the only minimal prime of ann \mathcal{A} . Thus $\mathfrak{p} = \sqrt{\operatorname{ann} \mathcal{A}}$, and so indeed a power of \mathfrak{p} annihilates \mathcal{A} .

The description of \mathcal{J} in Proposition 3.8 will be crucial to our study. Our search for an explicit generating set of \mathcal{J} now begins by extracting information from a Jacobian dual matrix $B(\varphi)$ of φ . We note, however, as there are two possible forms of the presentation φ in (3.2), there are two possible shapes of its Jacobian dual matrix, with respect to x_1, \ldots, x_d . Namely $B(\varphi)$ coincides with either $B(\varphi_R)$ or $B(\varphi_C)$ where

$$B(\varphi_C) = \begin{pmatrix} \bullet & & \\ B'_C & \vdots & \\ \hline 0 & \cdots & 0 & T_{n-d+s+1} \\ \vdots & \vdots & \vdots & \\ 0 & \cdots & 0 & T_n \end{pmatrix}, \qquad B(\varphi_R) = \begin{pmatrix} B'_R & \psi & \\ \hline 0 & \cdots & 0 & T_n & \\ \vdots & \vdots & \ddots & \\ 0 & \cdots & 0 & T_n \end{pmatrix}$$
(3.3)

which can be seen from (2.4) and the two possible shapes of φ in (3.2).

Here B'_C is an $s \times (n-e-1)$ matrix with linear entries in $k[T_1, \ldots, T_n]$, and the \bullet entries of $B(\varphi_C)$ belong to $k[T_1, \ldots, T_{n-d+s}]$. Moreover, B'_R is an $s \times (n-e-d+s)$ matrix and ψ is an $s \times (d-s)$ matrix, both consisting of linear entries in $k[T_1, \ldots, T_n]$. Additionally, the bottom right block of $B(\varphi_R)$ is $T_n \cdot I_{d-s}$ where I_{d-s} is the $(d-s) \times (d-s)$ identity matrix.

Remark 3.9. Notice that if s = d - 1, then the two possible shapes for $B(\varphi)$ in (3.3) coincide with

$$B(\varphi) = \begin{pmatrix} \bullet \\ B' & \vdots \\ \hline 0 & \cdots & 0 & T_n \end{pmatrix}$$

and one has $[x_1 \dots x_d] \cdot B' = [\ell_1 \dots \ell_{n-e-1}]$, as observed in [10]. Moreover, in this case, the matrix B' completely determines the nonlinear equations of the Rees algebra, which coincide with the *fiber equations* of $\mathcal{R}(E)$ [10, 4.11, 4.12]. In the proceeding sections, we will see that this is not always true when s < d - 1.

As we will see in Section 4 and Section 5, the two possible shapes of φ and $B(\varphi)$ produce two very different generating sets for the defining ideal \mathcal{J} of $\mathcal{R}(E)$ (compare Theorem 4.5 and Theorem 5.3), yielding very different behaviors in terms of the fiber type property and the analytic spread of E (compare Corollary 4.6 and Corollary 5.12). To understand the differences between the two cases, it will be useful to relate the submatrices B'_C and B'_R of (3.3) to the submatrices φ_i and the ideals \mathcal{L}_i introduced in Setting 2.5 and Notation 2.8. Notice that B'_C satisfies the equation

$$[T_1 \dots T_n] \cdot \varphi_1 = [\ell_1 \dots \ell_{n-e-1}] = [x_1 \dots x_s] \cdot B'_C$$

$$(3.4)$$

and so B'_C is precisely the Jacobian dual of the matrix φ_1 , with respect to the sequence x_1, \ldots, x_s . Similarly, with B'_R we have

$$[T_1 \dots T_n] \cdot \varphi_{d-s} = [\ell_1 \dots \ell_{n-e-(d-s)}] = [x_1 \dots x_s] \cdot B'_R$$

$$(3.5)$$

and so B'_R is the Jacobian dual of the matrix φ_{d-s} with respect to the sequence x_1, \ldots, x_s .

Recall that $\mathcal{L} = (\ell_1, \ldots, \ell_{n-e})$, where $[\ell_1 \ldots \ell_{n-e}] = [T_1 \ldots T_n] \cdot \varphi$, is the defining ideal of $\mathcal{S}(E)$. Notice that with (3.4) and (3.5), depending on whether $\varphi = \varphi_C$ or $\varphi = \varphi_R$, we have $\mathcal{L} + I_s(B'_C) \subseteq \mathcal{J}$ or $\mathcal{L} + I_s(B'_R) \subseteq \mathcal{J}$ due to Cramer's rule, since \mathcal{J} is a prime ideal. With this, we introduce the following result, which will be a crucial technical step towards determining the defining ideal \mathcal{J} of $\mathcal{R}(E)$.

Proposition 3.10. With the assumptions of Setting 3.1 and B'_C and B'_R as in (3.3), the following hold.

- (a) If $\varphi = \varphi_C$, then $I_s(B'_C)$ is a Cohen-Macaulay prime ideal of height n s e.
- (b) If $\varphi = \varphi_R$, then $I_s(B'_R)$ is a Cohen-Macaulay prime ideal of height n e d + 1.

Proof. By (3.3), B'_C is an $s \times (n-e-1)$ matrix with linear entries in $k[T_1, \ldots, T_n]$. Moreover, $[\ell_1 \ldots \ell_{n-e-1}] = [x_1 \ldots x_s] \cdot B'_C$ by (3.4). Now, as $\mathcal{L} : (x_1, \ldots, x_s)^{\infty}$ is a prime ideal of height n-e by Proposition 3.8, from [3, 2.2] it follows that $(\ell_1, \ldots, \ell_{n-e-1}) : (x_1, \ldots, x_s)^{\infty}$ is a prime ideal of height n-e-1. Hence, [3, 2.4] implies that $I_s(B'_C)$ is a prime ideal with ht $I_s(B'_C) = (n-e-1) - s + 1 = n-e-s$. Since this is the maximal possible height by [13, Thm. 1], the Cohen-Macaulayness of $I_s(B'_C)$ then follows from [14, A2.13]. This proves (a).

For (b), notice that according to (3.3) and (3.5), B'_R is an $s \times (n-e-d+s)$ matrix with linear entries in $k[T_1, \ldots, T_n]$ and $[\ell_1 \ldots \ell_{n-e-d+s}] = [x_1 \ldots x_s] \cdot B'_R$. As $\mathcal{L} : (x_1, \ldots, x_s)^{\infty}$ is a prime ideal of height n-e, by successively applying [3, 2.2] d-s times, we obtain that $(\ell_1, \ldots, \ell_{n-e-d+s}) : (x_1, \ldots, x_s)^{\infty}$ is a prime ideal of height n-e-d+s. Thus, $I_s(B'_R)$ is a prime ideal with ht $I_s(B'_R) = (n-e-d+s)-s+1 = n-e-d+1$. Again, this is the maximal possible height by [13, Thm. 1], hence $I_s(B'_R)$ is Cohen-Macaulay by [14, A2.13].

In the next sections, we will combine the height calculation of Proposition 3.10 with an argument allowing us to determine the defining ideal of $\mathcal{R}(E)$ from that of $\mathcal{R}(E_1)$, where $E = \operatorname{coker} \varphi_1$ as in Setting 2.5. The following proposition guarantees that the assumptions of Setting 3.1 are preserved when passing from E to E_1 , in either case of Remark 3.3.

Proposition 3.11. The module E_1 is an R-module of projective dimension one satisfying G_s , but not G_{s+1} .

Proof. From Lemma 2.6, we know that E_1 satisfies G_s , hence we only need to show that E_1 does not satisfy G_{s+1} . As E_1 has rank e+1, it suffices to show that ht $\operatorname{Fitt}_{s+e}(E_1) \leq s$. We distinguish two cases, depending on the two possible shapes for the matrix φ as in (3.2).

In the column setting, let $\varphi = \varphi_C$ as in (3.2) and write $\operatorname{Fitt}_{s+e}(E_1) = I_{n-s-e}(\varphi_1)$, with φ_1 as in Setting 2.5. Notice that $n-s-e \ge d-s \ge 1$, as $n \ge d+e$ and $s \ge d-1$ by assumption. Thus, $I_{n-s-e}(\varphi_1)$ is not the unit ideal and is hence contained in (x_1, \ldots, x_s) , as the entries of φ_1 belong to $A = k[x_1, \ldots, x_s]$. Thus, ht $\operatorname{Fitt}_{s+e}(E_1) \le s$ and the claim follows.

In the row setting, let $\varphi = \varphi_R$ as in (3.2). It is enough to show that E_{d-s} does not satisfy G_{s+1} , since then neither does E_1 , by Lemma 2.6. To prove this, notice that E_{d-s} is an *R*-module of rank e + d - sand $\operatorname{Fitt}_{d+e-1}(E_{d-s}) = I_{n-d-e+1}(\varphi_{d-s})$. Similarly to the previous case, it follows that $I_{n-d-e+1}(\varphi_{d-s}) \subseteq$ (x_1, \ldots, x_s) , as the entries of φ_{d-s} are in *A* and this is not the unit ideal. Thus, ht $\operatorname{Fitt}_{d+e-1}(E_{d-s}) \leq s$ and so E_{d-s} does not satisfy G_{s+1} , as required.

Remark 3.12. The proof of Proposition 3.11 shows that, in the row setting when $\varphi = \varphi_R$ as in (3.2), actually all modules E_i for $0 \le i \le d-s$ satisfy G_s but not G_{s+1} . That is, the maximal index s such that E_i satisfies G_s is constant in this range.

4. The column case

Throughout this section, we let R and E be as in Setting 3.1 and we assume that the presentation matrix φ of E is in column form, $\varphi = \varphi_C$ as in (3.2). Our primary task is to determine the defining ideal \mathcal{J} of the Rees ring $\mathcal{R}(E)$ through means of successive approximations of this algebra, as described in Section 2.2. As we will see, the fact that a subset of variables is concentrated in one column of the presentation matrix φ makes this setting quite amenable to this approximation method. Indeed, the first algebra $\mathcal{R}(E_1)$ alone will provide enough information to determine \mathcal{J} . Our main setting throughout is as follows.

Setting 4.1. Adopt the assumptions of Setting 3.1 and assume that, after the appropriate change of coordinates and row and column operations, the presentation matrix φ has the shape $\varphi = \varphi_C$ in (3.2).

We omit the subscript in φ_C and instead write φ throughout to simplify the notation. With this, the matrix φ and its Jacobian dual, with respect to x_1, \ldots, x_d , are

$$\varphi = \begin{pmatrix} & & * \\ & & \ddots \\ & & * \\ \hline * & \cdots & * & x_{s+1} \\ \vdots & & \vdots & \vdots \\ * & \cdots & * & x_d \end{pmatrix} \quad \text{and} \quad B(\varphi) = \begin{pmatrix} & & \bullet \\ & B' & \vdots \\ & & \bullet \\ \hline 0 & \cdots & 0 & T_{n-d+s+1} \\ \vdots & & \vdots & \vdots \\ 0 & \cdots & 0 & T_n \end{pmatrix}$$
(4.1)

where the entries of φ' and the * entries belong to the subring $A = k[x_1, \ldots, x_s]$, and the entries of $B(\varphi)$ belong to $k[T_1, \ldots, T_n]$. We also recall that

$$[T_1 \dots T_n] \cdot \varphi = [\ell_1 \dots \ell_{n-e}] = [x_1 \dots x_d] \cdot B(\varphi)$$

$$(4.2)$$

where $\mathcal{L} = (\ell_1, \dots, \ell_{n-e})$ is the defining ideal of the symmetric algebra $\mathcal{S}(E)$, as in Section 2.

As noted, we aim to approximate, in a sense, the Rees ring $\mathcal{R}(E)$ with a well-understood algebra surjecting onto it. By employing the technique of successive approximations, we take this algebra to be the Rees ring of E_1 , where E_1 is the *R*-module defined in Section 2.2. We begin by making some observations on the module E_1 , and then we will consider its Rees ring.

Proposition 4.2. With the assumptions of Setting 4.1, let φ_1 denote the submatrix of φ as in Section 2.2, and let E_1 denote the cohernel of φ_1 . The module E_1 has projective dimension one and satisfies G_s , but not G_{s+1} , both as an R-module and as an A-module, where $A = k[x_1, \ldots, x_s]$.

Proof. The first claim follows from Proposition 3.11. Notice that since φ has the shape in (4.1), the entries of φ_1 belong to the subring A, hence E_1 is an A-module of projective dimension one. Moreover, since $\operatorname{Spec}(A) \subset \operatorname{Spec}(R), E_1$ satisfies G_s , but not G_{s+1} , as an A-module as well. \Box

With this, we may describe the Rees ring of E_1 as both an A-module and an R-module.

Proposition 4.3. The Rees algebra of E_1 , as an R-module, is $\mathcal{R}(E_1) \cong \mathbb{R}[T_1, \ldots, T_n]/\mathcal{J}_1$ where

$$\mathcal{J}_1 = \mathcal{L}_1 : (x_1, \dots, x_s) = \mathcal{L}_1 + I_s(B')$$

and $\mathcal{L}_1 = (\ell_1, \ldots, \ell_{n-e-1})$. Moreover, $\mathcal{R}(E_1)$ is Cohen-Macaulay, both as an R-module and as an A-module.

Proof. Notice that the submatrix B' is precisely the Jacobian dual of φ_1 with respect to x_1, \ldots, x_s , as observed in (3.4). Moreover, by Proposition 4.2, E_1 satisfies G_s as an A-module, where $A = k[x_1, \ldots, x_s]$. Hence, from [31, 4.11] it follows that the Rees ring of E_1 , as an A-module, is $\mathcal{R}_A(E_1) \cong A[T_1, \ldots, T_n]/\mathcal{J}_1$ with $\mathcal{J}_1 = \mathcal{L}_1 : (x_1, \ldots, x_s) = \mathcal{L}_1 + I_s(B')$ and moreover, this ring is Cohen-Macaulay.

Viewing now E_1 as an *R*-module, it then follows that $\mathcal{R}(E_1) \cong R[T_1, \ldots, T_n]/\mathcal{J}_1$, with \mathcal{J}_1 extended to $R[T_1, \ldots, T_n]$. Moreover, $\mathcal{R}(E_1)$ is Cohen-Macaulay, as x_{s+1}, \ldots, x_d is a regular sequence modulo \mathcal{J}_1 . \Box

We remark that, although in the previous proof we include a description of the Rees algebra of E_1 over the simpler ring A, the description of $\mathcal{R}(E_1)$ with E_1 viewed as an R-module is required for our purposes. Indeed, we require a map of R-modules $E_1 \to E$ to induce the map of R-algebras $\mathcal{R}(E_1) \to \mathcal{R}(E)$, as in Section 2.2. Recall that the kernel of this map is $\mathcal{J}/\mathcal{J}_1$, and this is a prime ideal of height one. **Lemma 4.4.** With the isomorphism $\mathcal{R}(E_1) \cong R[T_1, \ldots, T_n]/\mathcal{J}_1$ in Proposition 4.3, write $\overline{\cdot}$ to denote images modulo \mathcal{J}_1 . We have the following.

- (a) Letting $\mathbf{p} = (x_1, \ldots, x_s)$, $\overline{\mathbf{p}}$ is a Cohen-Macaulay prime $\mathcal{R}(E_1)$ -ideal of height one.
- (b) The $\mathcal{R}(E_1)$ -ideal $\mathcal{L} + I_s(B')$ is Cohen-Macaulay of height one.

Proof. Recall that $\mathcal{R}_A(E_1) \cong A[T_1, \ldots, T_n]/\mathcal{J}_1$, where $A = k[x_1, \ldots, x_s]$. As $\mathfrak{p} = (x_1, \ldots, x_s)$ is the homogeneous maximal ideal of A, the special fiber ring of E_1 , as an A-module, is

$$\mathcal{F}_A(E_1) \cong \mathcal{R}_A(E_1)/\mathfrak{p}\mathcal{R}_A(E_1) \cong k[T_1,\ldots,T_n]/I_s(B')$$

by [31, 4.11], noting that $\mathcal{L}_1 \subseteq \mathfrak{p}$. Moreover, this is a Cohen-Macaulay domain of dimension s + e, following Proposition 3.10(a). On the other hand, notice that $\mathcal{R}(E_1) \cong \mathcal{R}_A(E_1)[x_{s+1}, \ldots, x_d]$ and so

$$\mathcal{R}(E_1)/\mathfrak{p}\mathcal{R}(E_1) \cong \mathcal{F}_A(E_1)[x_{s+1},\ldots,x_d].$$

Hence this quotient is a Cohen-Macaulay domain of dimension s + e + (d - s) = d + e. Now since E_1 has rank e + 1 by Lemma 2.6(a), it follows that dim $\mathcal{R}(E_1) = d + e + 1$. Thus $\overline{\mathfrak{p}}$ is a prime ideal with ht $\overline{\mathfrak{p}} = 1$, which shows (a).

For (b), notice that $\overline{\mathcal{L} + I_s(B')} = \overline{(\ell_{n-e})}$, which follows immediately from the expression of \mathcal{J}_1 in Proposition 4.3. Since $\ell_{n-e} \notin \mathcal{J}_1$, it follows that $\overline{\ell_{n-e}} \neq 0$. As $\mathcal{R}(E_1)$ is a domain, the ideal $\overline{(\ell_{n-e})}$ is thus generated by a regular element. Hence it is Cohen-Macaulay of height one, as claimed.

In [10, 4.11] it was shown that when s = d - 1, the defining ideal of $\mathcal{R}(E)$ is $\mathcal{J} = \mathcal{L}: (x_1, \ldots, x_{d-1}) = \mathcal{L} + I_{d-1}(B')$. We now prove that the same property holds for every $s \leq d - 1$, in the present setting.

Theorem 4.5. With φ and E as in Setting 4.1, the defining ideal \mathcal{J} of $\mathcal{R}(E)$ is

$$\mathcal{J} = \mathcal{L} : \mathfrak{p} = \mathcal{L} + I_s(B')$$

where $\mathfrak{p} = (x_1, \ldots, x_s)$. Moreover, $\mathcal{R}(E)$ is Cohen-Macaulay.

Proof. Recall from Proposition 3.8 that $\mathcal{J} = \mathcal{L} : \mathfrak{p}^{\infty}$. Hence, by Cramer's rule one has the containments $\mathcal{L} + I_s(B') \subseteq \mathcal{L} : \mathfrak{p} \subseteq \mathcal{J}$, and so we only need to show that $\mathcal{J} = \mathcal{L} + I_s(B')$. Moreover, since $\mathcal{J}_1 \subseteq \mathcal{L} + I_s(B') \subseteq \mathcal{J}$, it suffices to prove that $\overline{\mathcal{J}} = \overline{\mathcal{L}} + I_s(B') = \overline{(\ell_{n-e})}$.

Since clearly $\overline{(\ell_{n-e})} \subseteq \overline{\mathcal{J}}$, it is enough to show that $\overline{(\ell_{n-e})}$ and $\overline{\mathcal{J}}$ agree locally at the associated primes of $\overline{(\ell_{n-e})}$. Recall from Lemma 4.4(b) that this ideal is Cohen-Macaulay of height one, hence all of its associated primes are minimal of height one. Thus it is enough to prove the stronger statement that $\overline{(\ell_{n-e})}_{\mathfrak{q}} = \overline{\mathcal{J}}_{\mathfrak{q}}$ for any prime $\mathcal{R}(E_1)$ -ideal \mathfrak{q} with height one. Recall that $\overline{\mathfrak{p}}$ is one such prime, due to Lemma 4.4(a).

First, notice that if $q \neq \overline{p}$, we have

$$\overline{\mathcal{J}}_{\mathfrak{q}} = \overline{\mathcal{L}}_{\mathfrak{q}} : \overline{\mathfrak{p}}_{\mathfrak{q}}^{\infty} = \overline{\mathcal{L}}_{\mathfrak{q}} : \mathcal{R}(E_1)_{\mathfrak{q}} = \overline{\mathcal{L}}_{\mathfrak{q}} = \overline{(\ell_{n-e})}_{\mathfrak{q}},$$

since $\overline{\mathcal{L}} = \overline{(\ell_{n-e})}$. If instead $\mathfrak{q} = \overline{\mathfrak{p}}$, note that $\overline{\ell_{n-e}} \notin \overline{\mathfrak{p}}$. Indeed, one has that $\overline{(\ell_{n-e})} + \overline{\mathfrak{p}} = (\overline{\gamma}) + \overline{\mathfrak{p}}$ where $\gamma = x_{s+1}T_{n-d+s+1} + \cdots + x_dT_n$ following (4.2). Since $\mathfrak{p} = (x_1, \ldots, x_s)$ and x_1, \ldots, x_s, γ is a regular sequence, it follows that $\overline{(\ell_{n-e})} + \overline{\mathfrak{p}} \neq \overline{\mathfrak{p}}$, and so $\overline{\ell_{n-e}} \notin \overline{\mathfrak{p}}$ as claimed. Therefore, $\overline{(\ell_{n-e})}_{\overline{\mathfrak{p}}} = \mathcal{R}(E_1)_{\overline{\mathfrak{p}}}$. Moreover, since $\overline{(\ell_{n-e})} \subseteq \overline{\mathcal{J}}$, it follows that $\overline{\mathcal{J}}_{\overline{\mathfrak{p}}} = \mathcal{R}(E_1)_{\overline{\mathfrak{p}}}$ as well. Thus $\overline{\mathcal{J}} = \overline{(\ell_{n-e})}$, and so $\mathcal{J} = \mathcal{L} + I_s(B')$, as claimed.

As for the Cohen-Macaulayness of $\mathcal{R}(E)$, notice that we have shown that $\mathcal{J} = \mathcal{J}_1 + (\ell_{n-e})$. As noted in Lemma 4.4, the element ℓ_{n-e} is regular modulo \mathcal{J}_1 . Moreover, by Proposition 4.3 we know that $\mathcal{R}(E_1)$ is Cohen-Macaulay, hence $\mathcal{R}(E) \cong \mathcal{R}(E_1)/(\overline{\ell_{n-e}})$ is Cohen-Macaulay as well.

With this, we may easily describe the ideal defining the special fiber ring $\mathcal{F}(E)$.

Corollary 4.6. With the assumptions of Setting 4.1, the special fiber ring of E is

$$\mathcal{F}(E) \cong k[T_1, \dots, T_n]/I_s(B')$$

In particular, $\mathcal{F}(E)$ is a Cohen-Macaulay domain of dimension s + e, i.e. E has analytic spread $\ell(E) = s + e$.

Proof. Recall that the entries of B' belong to $k[T_1, \ldots, T_n]$. From Theorem 4.5, it follows that $\mathcal{J} + (x_1, \ldots, x_d) = I_s(B') + (x_1, \ldots, x_d)$, and so $\mathcal{F}(E) \cong k[T_1, \ldots, T_n]/I_s(B')$. Moreover, by Proposition 3.10(a) we have that $I_s(B')$ is a Cohen-Macaulay prime ideal of $k[T_1, \ldots, T_n]$ with ht $I_s(B') = n - s - e$. Hence it follows that $\mathcal{F}(E)$ is a Cohen-Macaulay domain with $\ell(E) = \dim \mathcal{F}(E) = s + e$.

Remark 4.7. With the shape of the defining ideal \mathcal{J} in Theorem 4.5, we note that the module E is of *fiber type* by Corollary 4.6. In other words, aside from the generators of \mathcal{L} , the equations of \mathcal{J} belong to the subring $k[T_1, \ldots, T_n]$.

As another consequence of Theorem 4.5, we may also describe the defining ideal \mathcal{J} as a residual intersection.

Proposition 4.8. With the assumptions of Setting 4.1, the defining ideal \mathcal{J} of $\mathcal{R}(E)$ may be realized as

$$\mathcal{J} = \mathcal{L} : (x_1, \dots, x_s, \gamma)$$

where $\gamma = x_{s+1}T_{n-d+s+1} + \cdots + x_dT_n$, as in the proof of Theorem 4.5. Moreover, this is an (n-e)-residual intersection of $(x_1, \ldots, x_s, \gamma)$.

Proof. Recall from the proof of Theorem 4.5 that $\mathcal{L} \subseteq (x_1, \ldots, x_s, \gamma)$. Moreover, notice that we have the containment

$$\mathcal{L}: (x_1, \dots, x_s, \gamma) \subseteq \mathcal{L}: (x_1, \dots, x_s) = \mathcal{J}$$

$$(4.3)$$

where the equality follows from Theorem 4.5. Cramer's rule then implies that $I_s(B') = I_{s+1}(B'') \subseteq \mathcal{L}$: $(x_1, \ldots, x_s, \gamma)$, where B'' is the matrix

$$B'' = \begin{pmatrix} & & \bullet \\ & B' & \vdots \\ & & \bullet \\ \hline 0 & \cdots & 0 & 1 \end{pmatrix}$$
(4.4)

satisfying the matrix equation

$$[\ell_1 \dots \ell_{n-e}] = [x_1 \dots x_s \ \gamma] \cdot B''. \tag{4.5}$$

Thus, $\mathcal{L} + I_s(B') \subseteq \mathcal{L} : (x_1, \ldots, x_s, \gamma)$, hence by Theorem 4.5 we have $\mathcal{J} = \mathcal{L} : (x_1, \ldots, x_s, \gamma)$. Moreover, as ht $\mathcal{J} = n - e$ and $\mathcal{L} = (\ell_1, \ldots, \ell_{n-e}) \subseteq (x_1, \ldots, x_s, \gamma)$, we conclude that \mathcal{J} is an (n - e)-residual intersection of $(x_1, \ldots, x_s, \gamma)$.

Remark 4.9. As noted in the proof of Theorem 4.5, we have that x_1, \ldots, x_s, γ is a regular sequence. Hence by Proposition 4.8, \mathcal{J} is a residual intersection of a complete intersection ideal. The proof of the main result of [10] relied on a similar observation, and then used Theorem 2.10 to obtain a generating set of \mathcal{J} . We remark that a similar path could be followed to give an alternative proof of Theorem 4.5, however the method of successive approximations of $\mathcal{R}(E)$ allows for a much quicker proof, and gives that \mathcal{J} is a residual intersection as a byproduct. Whereas the methods of [10] could have been applied in the present setting, we will see in the proceeding section that they fail for the row case; see Remark 5.14.

As noted in Remark 3.4, the column setting and the row setting of Remark 3.3 coincide when s = d - 1, in which case one has the main result of [10], which shows that the defining ideal of $\mathcal{R}(E)$ is $\mathcal{J} = \mathcal{L} + I_{d-1}(B')$, similarly to Theorem 4.5. In the next section, we show that in the row setting, i.e. when $\varphi = \varphi_R$ in (3.2), the ideal $\mathcal{L} + I_s(B')$ will be *strictly* contained in \mathcal{J} , when s < d - 1. Moreover, unlike in Theorem 4.5, the module E will not be of fiber type.

5. The row case

In this section, we describe the defining ideal \mathcal{J} of $\mathcal{R}(E)$ with E as in Setting 3.1, assuming that the presentation matrix φ of E is in row form $\varphi = \varphi_R$ as in (3.2). As before, we employ the method of successive approximations to study $\mathcal{R}(E)$. The key difference here, opposed to the case of Section 4, is that this method becomes part of an induction argument to determine the defining ideal of $\mathcal{R}(E)$. Our setting throughout is as follows.

Setting 5.1. Adopt the assumptions of Setting 3.1 and assume that, after the appropriate change of coordinates, and row and column operations, φ has the shape $\varphi = \varphi_R$ in (3.2).

As in the previous section, we omit the subscript and write φ for φ_R throughout to simplify the notation. With this, the matrix φ and its Jacobian dual, with respect to x_1, \ldots, x_d , are

$$\varphi = \begin{pmatrix} & & \ast & \cdots & \ast \\ & & \vdots & & \vdots \\ & & \ast & \cdots & \ast \\ \hline \ast & \cdots & \ast & x_{s+1} & \cdots & x_d \end{pmatrix} \quad \text{and} \quad B(\varphi) = \begin{pmatrix} & B' & & \psi \\ & & & & \\ \hline 0 & \cdots & 0 & T_n \\ \vdots & & \vdots & & \ddots \\ 0 & \cdots & 0 & & & T_n \end{pmatrix}$$
(5.1)

as in (3.2) and (3.3). Along with the Jacobian dual $B(\varphi)$, consider the matrix

$$C = \begin{pmatrix} B' & \psi \\ \hline 0 & \cdots & 0 & x_{s+1} & \cdots & x_d \end{pmatrix}$$
(5.2)

where B' and ψ are the submatrices of $B(\varphi)$ in (5.1).

Remark 5.2. As noted prior to Proposition 3.10, recall that $B' = B(\varphi_{d-s})$, the Jacobian dual of the submatrix φ_{d-s} with respect to x_1, \ldots, x_s , adopting the notation of Section 2.2. Moreover, notice that the matrix C is a transition matrix as well. Indeed, from (3.5) and the shape of C above, we see that

$$\left[\ell_1 \dots \ell_{n-e-(d-s)}\right] = \left[x_1 \dots x_s\right] \cdot B' \quad \text{and} \quad \left[\ell_1 \dots \ell_{n-e}\right] = \left[x_1 \dots x_s \ T_n\right] \cdot C. \tag{5.3}$$

where $[\ell_1 \dots \ell_{n-e}] = [T_1 \dots T_n] \cdot \varphi$ are the equations defining the symmetric algebra $\mathcal{S}(E)$.

Notice that, by Cramer's rule, and since \mathcal{J} is a prime ideal, we have that $\mathcal{L} + I_s(B') + I_{s+1}(C) \subseteq \mathcal{J}$. The following theorem, which we prove by induction in Section 5.1, shows that this containment is actually an equality.

Theorem 5.3. With the assumptions of Setting 5.1 and φ the presentation of E as in (5.1), the Rees algebra of E is $\mathcal{R}(E) \cong R[T_1, \ldots, T_n]/\mathcal{J}$ with

$$\mathcal{J} = \mathcal{L} : (x_1, \dots, x_s) = \mathcal{L} + I_s(B') + I_{s+1}(C),$$

where $\mathcal{L} = (\ell_1, \dots, \ell_{n-e})$, and B' and C are the matrices in (5.1) and (5.2), satisfying the equations of (5.3). Moreover, $\mathcal{R}(E)$ is Cohen-Macaulay.

5.1. **Proof by Induction.** We devote this subsection to the proof of Theorem 5.3. As such, we provide the induction setting below.

Setting 5.4. With the assumptions of Setting 5.1, we proceed by induction on the difference d-s in order to prove Theorem 5.3. We note that the initial case d-s = 1 has been shown in [10, 3.9], observing that $I_{s+1}(C) \subseteq I_s(B')$ in this setting. (Alternatively, this follows from Theorem 4.5, due to Remark 3.4.) Hence we assume that $d-s \ge 2$ and the claim holds up to d-s-1.

As in the previous section, we employ the technique of successive approximations to produce a ring surjecting onto $\mathcal{R}(E)$ so that the kernel of this map is a prime ideal of height one. As before, we take such a ring to be the Rees ring of $E_1 = \operatorname{coker} \varphi_1$, where φ_1 and E_1 are as in Section 2.2. Notice that from the shape of φ in (5.1), the entries of φ_1 belong to the subring $R' = k[x_1, \ldots, x_{d-1}]$. In particular, E_1 may be viewed as a module over this subring.

Proposition 5.5. The Rees algebra of E_1 is, as an *R*-module, $\mathcal{R}(E_1) \cong \mathbb{R}[T_1, \ldots, T_n]/\mathcal{J}_1$ with

$$\mathcal{J}_1 = \mathcal{L}_1 : (x_1, \dots, x_s) = \mathcal{L}_1 + I_s(B') + I_{s+1}(C_1)$$

where $\mathcal{L}_1 = (\ell_1, \ldots, \ell_{n-e-1})$, B' is the submatrix of $B(\varphi)$ in (5.1), and C_1 is the submatrix of C in (5.2) obtained by deleting its last column. Moreover, $\mathcal{R}(E_1)$ is a Cohen-Macaulay domain of dimension d + e + 1.

Proof. Recall that by Proposition 3.11, E_1 satisfies G_s , but not G_{s+1} , as an R-module. Moreover, from the shape of φ and its submatrix φ_1 in (5.1), an argument similar to the one in Proposition 4.2 shows that E_1 satisfies G_s , but not G_{s+1} , as an R'-module as well, where $R' = k[x_1, \ldots, x_{d-1}]$. Hence E_1 satisfies the assumptions of Setting 5.1 as an R'-module. Moreover, as $R' = k[x_1, \ldots, x_{d-1}]$, we see that dim R' - s = d - s - 1. Hence by the induction hypothesis in Setting 5.4, the Rees ring of E_1 , as an R'-module, is

$$\mathcal{R}_{R'}(E_1) \cong R'[T_1, \dots, T_n]/\mathcal{J}_1$$

where

$$\mathcal{J}_1 = \mathcal{L}_1 + I_s(B') + I_{s+1}(C_1)$$

with $\mathcal{L}_1 = (\ell_1, \ldots, \ell_{n-e-1})$ and $[\ell_1 \ldots \ell_{n-e-1}] = [T_1 \ldots T_n] \cdot \varphi_1$. Indeed, observe that B' and C_1 satisfy the matrix equations

$$[\ell_1 \dots \ell_{n-(e+1)-(d-s-1)}] = [x_1 \dots x_s] \cdot B'$$
 and $[\ell_1 \dots \ell_{n-e-1}] = [x_1 \dots x_s \ T_n] \cdot C_1$

similar to those in (5.3), noting that rank $E_1 = e + 1$ and dim R' = d - 1. Hence n - (e + 1) - (d - s - 1) = n - e - (d - s) and the equations above agree with (5.3). Alternatively, again noting that dim R' - s = d - s - 1, from Remark 5.2 the first matrix required here may be realized as the Jacobian dual of $(\varphi_1)_{d-s-1} = \varphi_{d-s}$, which is again B' by Remark 5.2.

As the ideal \mathcal{J}_1 above is the ideal defining the Rees ring of E_1 as an R'-module, its extension to $R[T_1, \ldots, T_n]$ is the defining ideal of the Rees ring of E_1 as an R-module. In other words, viewing \mathcal{J}_1 as an ideal of $R[T_1, \ldots, T_n]$, we have

$$\mathcal{R}(E_1) \cong R[T_1, \ldots, T_n]/\mathcal{J}_1.$$

Finally, since $\mathcal{R}(E_1) \cong \mathcal{R}_{R'}(E_1)[x_d]$ and $\mathcal{R}_{R'}(E_1)$ is Cohen-Macaulay by the induction hypothesis, it follows that $\mathcal{R}(E_1)$ is a Cohen-Macaulay domain of dimension d + e + 1.

Similarly to the previous section, we regard E_1 as an *R*-module so as to consider the epimorphism of *R*-algebras $\mathcal{R}(E_1) \to \mathcal{R}(E)$, as outlined in Section 2. Notice that the kernel of this map is precisely $\mathcal{J}/\mathcal{J}_1$, which is a prime $\mathcal{R}(E_1)$ -ideal of height one. Inspired by analogous constructions used in [3, 21, 37, 38], we now introduce certain ideals in $\mathcal{R}(E_1)$ to study this kernel. To this end, the following notation will be useful.

Notation 5.6. Let $\overline{\cdot}$ denote images modulo \mathcal{J}_1 in $\mathcal{R}(E_1)$. Let C' denote the $s \times (n-e-1)$ submatrix of C_1 obtained by deleting its last row (equivalently, C' is obtained from C by deleting its last row and column). Consider the $R[T_1, \ldots, T_n]$ -ideal $\mathcal{K} \coloneqq (\ell_1, \ldots, \ell_{n-e-1}) + I_s(C') + (T_n)$.

Proposition 5.7. The $\mathcal{R}(E_1)$ -ideal $\overline{\mathcal{K}}$ is a Cohen-Macaulay ideal of height one.

Proof. From the description of \mathcal{J}_1 in Proposition 5.5, we have the containment $\mathcal{J}_1 \subseteq \mathcal{K}$, noting that $I_s(B') \subseteq I_s(C')$. Thus ht $\mathcal{K} \ge n - e$, as \mathcal{J}_1 is a prime ideal of height n - e - 1 and $T_n \in \mathcal{K} \setminus \mathcal{J}_1$. We next show that ht $\mathcal{K} = n - e$. To this end, notice that, since $T_n \in \mathcal{K}$, there exists an $s \times (n - e - 1)$ matrix $\widetilde{C'}$ with entries in $k[T_1, \ldots, T_{n-1}]$ and elements $\tilde{\ell}_1, \ldots, \tilde{\ell}_{n-e-1} \in R[T_1, \ldots, T_{n-1}]$ such that

$$\mathcal{K} = (\widetilde{\ell}_1, \dots, \widetilde{\ell}_{n-e-1}) + I_s(\widetilde{C'}) + (T_n)$$
(5.4)

and $[\tilde{\ell}_1 \dots \tilde{\ell}_{n-e-1}] = [x_1 \dots x_s] \cdot \widetilde{C'}$. Since ht $\mathcal{K} \ge n-e$, from (5.4) above it follows that ht $((\tilde{\ell}_1, \dots, \tilde{\ell}_{n-e-1}) + I_s(\widetilde{C'})) \ge n-e-1$, as this is an ideal of the subring $R[T_1, \dots, T_{n-1}]$. However, since

$$(\widetilde{\ell}_1, \dots, \widetilde{\ell}_{n-e-1}) + I_s(\widetilde{C}') \subseteq (\widetilde{\ell}_1, \dots, \widetilde{\ell}_{n-e-1}) : (x_1, \dots, x_s)$$

it follows that $\operatorname{ht}\left((\ell_1,\ldots,\ell_{n-e-1}):(x_1,\ldots,x_s)\right) \ge n-e-1$. Thus this ideal is an (n-e-1)-residual intersection of (x_1,\ldots,x_s) , and so by Theorem 2.10 it follows that

$$(\tilde{\ell}_1, \dots, \tilde{\ell}_{n-e-1}) + I_s(\widetilde{C}') = (\tilde{\ell}_1, \dots, \tilde{\ell}_{n-e-1}) : (x_1, \dots, x_s)$$
(5.5)

and this is a Cohen-Macaulay ideal of height exactly n - e - 1. As T_n is regular modulo this ideal, we then have that \mathcal{K} is Cohen-Macaulay of height n - e, as claimed. Thus $\overline{\mathcal{K}}$ is a Cohen-Macaulay $\mathcal{R}(E_1)$ -ideal with ht $\overline{\mathcal{K}} = 1$.

With the Cohen-Macaulayness of $\overline{\mathcal{K}}$ established, we now show that this ideal is *linked*, in the sense of [17], to a particular $\mathcal{R}(E_1)$ -ideal. We begin with the following lemma.

Lemma 5.8. Writing $\mathcal{K} = (\tilde{\ell}_1, \ldots, \tilde{\ell}_{n-e-1}) + I_s(\widetilde{C'}) + (T_n)$ as in the proof of Proposition 5.7, we have $I_s(\widetilde{C'}) \neq 0$, modulo \mathcal{J}_1 .

Proof. First note that, by degree considerations, if $I_s(\widetilde{C'}) \subseteq \mathcal{J}_1$, then it must be that $I_s(\widetilde{C'}) \subseteq I_s(B')$. We show that this is impossible. Recall from the proof of Proposition 5.7 that the ideal $(\tilde{\ell}_1, \ldots, \tilde{\ell}_{n-e-1}) + I_s(\widetilde{C'})$ has height n - e - 1. As $(\tilde{\ell}_1, \ldots, \tilde{\ell}_{n-e-1}) \subseteq (x_1, \ldots, x_s)$, it follows that ht $((x_1, \ldots, x_s) + I_s(\widetilde{C'})) \ge n - e - 1$. Since $I_s(\widetilde{C'})$ is an ideal of the subring $k[T_1, \ldots, T_{n-1}]$, this implies that ht $I_s(\widetilde{C'}) \ge n - e - s - 1$. Thus if $I_s(\widetilde{C'}) \subseteq I_s(B')$, then by Proposition 3.10 we have that $n - e - s - 1 \le n - e - d + 1$, hence $s \ge d - 2$. However, from Setting 5.4 we have that $s \le d - 2$, so we only need to consider the case that s = d - 2.

So, suppose that $I_s(\widetilde{C'}) \subseteq I_s(B')$ where s = d - 2, and note that then $d \ge 4$, as $s \ge 2$ by assumption. Write $\widetilde{B'}$ for the submatrix of $\widetilde{C'}$ consisting of the first n - e - 2 columns, which coincides with the matrix obtained from B' by extracting its terms containing T_n . Notice that

$$I_{d-2}(\widetilde{C}') \subseteq I_{d-2}(B') \subseteq I_{d-2}(B') + (T_n) = I_{d-2}(\widetilde{B}') + (T_n)$$

hence it follows that $I_{d-2}(\widetilde{C'}) \subseteq I_{d-2}(\widetilde{B'})$ as both matrices consist of entries in $k[T_1, \ldots, T_{n-1}]$. Thus we have that $I_{d-2}(\widetilde{C'}) = I_{d-2}(\widetilde{B'})$, as $\widetilde{B'}$ is a submatrix of $\widetilde{C'}$.

With this, we next show that $\tilde{\ell}_{n-e-1}$ is a regular element modulo $(\tilde{\ell}_1, \ldots, \tilde{\ell}_{n-e-2}) + I_{d-2}(\tilde{B}')$. Recall from the proof of Proposition 3.10(b) that $(\ell_1, \ldots, \ell_{n-e-2}) : (x_1, \ldots, x_{d-2})^{\infty}$ is a prime ideal of height n-e-2. Hence by [3, 2.4] and (5.3), we see that

$$(\ell_1, \dots, \ell_{n-e-2}) : (x_1, \dots, x_{d-2})^{\infty} = (\ell_1, \dots, \ell_{n-e-2}) : (x_1, \dots, x_{d-2}) = (\ell_1, \dots, \ell_{n-e-2}) + I_{d-2}(B').$$

As this is a prime ideal not containing T_n , it follows that

$$(\ell_1, \dots, \ell_{n-e-2}) + I_{d-2}(B') + (T_n) = (\tilde{\ell}_1, \dots, \tilde{\ell}_{n-e-2}) + I_{d-2}(\widetilde{B'}) + (T_n)$$

and this ideal has height n - e - 1. Thus we deduce that ht $((\tilde{\ell}_1, \ldots, \tilde{\ell}_{n-e-2}) + I_{d-2}(\widetilde{B'})) = n - e - 2$ as this ideal belongs to $R[T_1, \ldots, T_{n-1}]$. As $(\tilde{\ell}_1, \ldots, \tilde{\ell}_{n-e-2}) + I_{d-2}(\widetilde{B'}) \subseteq (\tilde{\ell}_1, \ldots, \tilde{\ell}_{n-e-2}) : (x_1, \ldots, x_{d-2})$, by height considerations and Theorem 2.10, it follows that

$$(\widetilde{\ell}_1,\ldots,\widetilde{\ell}_{n-e-2}):(x_1,\ldots,x_{d-2})=(\widetilde{\ell}_1,\ldots,\widetilde{\ell}_{n-e-2})+I_{d-2}(\widetilde{B'})$$

with height exactly n - e - 2, and this ideal is Cohen-Macaulay. Now, since $I_{d-2}(\widetilde{C}') = I_{d-2}(\widetilde{B}')$ and $(\widetilde{\ell}_1, \ldots, \widetilde{\ell}_{n-e-1}) + I_{d-2}(\widetilde{C}')$ is a Cohen-Macaulay ideal of height n - e - 1, the height calculation above implies that $\widetilde{\ell}_{n-e-1}$ is a regular element modulo $(\widetilde{\ell}_1, \ldots, \widetilde{\ell}_{n-e-2}) + I_{d-2}(\widetilde{B}')$, as claimed.

Having shown that $\tilde{\ell}_{n-e-1}$ is regular modulo $(\tilde{\ell}_1, \ldots, \tilde{\ell}_{n-e-2}) + I_{d-2}(\widetilde{B'})$, we now use this to reach the desired contradiction. Observe that, as $I_{d-2}(\widetilde{B'}) = I_{d-2}(\widetilde{C'}) \neq 0$, there exists a nonzero minor of $\widetilde{B'}$ and, without loss of generality, we may assume this is the minor consisting of the first d-2 columns. Hence, there is a nonzero $(d-3) \times (d-3)$ minor δ within the first d-3 columns (recall that $d \ge 4$). Assume δ is obtained by deleting row i, for some $1 \le i \le d-2$. Let Δ denote the minor of $\widetilde{C'}$ consisting of the first d-3 columns of $\widetilde{B'}$ and the last column of $\widetilde{C'}$.

Notice that $x_i \Delta \in (x_1, \ldots, x_{d-2}) I_{d-2}(\widetilde{C}') = (x_1, \ldots, x_{d-2}) I_{d-2}(\widetilde{B}') \subseteq (\tilde{\ell}_1, \ldots, \tilde{\ell}_{n-e-2})$, recalling that $I_{d-2}(\widetilde{C}') = I_{d-2}(\widetilde{B}')$. Moreover, by Cramer's rule, modulo $(\tilde{\ell}_1, \ldots, \tilde{\ell}_{d-3})$ we have $x_i \Delta \equiv (-1)^{i+d-2} \tilde{\ell}_{n-e-1} \delta$ and so it follows that

$$\widetilde{\ell}_{n-e-1}\delta \subseteq (\widetilde{\ell}_1, \dots, \widetilde{\ell}_{n-e-2}) + I_{d-2}(\widetilde{B'}).$$

However, recall that $\tilde{\ell}_{n-e-1}$ is regular modulo $(\tilde{\ell}_1, \ldots, \tilde{\ell}_{n-e-2}) + I_{d-2}(\widetilde{B'})$. Hence it must be that $\delta \in (\tilde{\ell}_1, \ldots, \tilde{\ell}_{n-e-2}) + I_{d-2}(\widetilde{B'})$. But this is impossible by degree considerations, and so this is a contradiction. \Box

Proposition 5.9. With \mathcal{K} as in Notation 5.6, we have that $\overline{\mathcal{K}}$ and $\overline{(x_1, \ldots, x_s, T_n)}$ are linked through the regular element $\overline{T_n}$, namely $\overline{(x_1, \ldots, x_s, T_n)} = \overline{(T_n)} : \overline{\mathcal{K}}$ and $\overline{\mathcal{K}} = \overline{(T_n)} : \overline{(x_1, \ldots, x_s, T_n)}$. In particular, the $\mathcal{R}(E_1)$ -ideal $\overline{(x_1, \ldots, x_s, T_n)}$ is unmixed of height one.

<u>Proof.</u> From the construction of \mathcal{K} , it is clear that we have the containment $\overline{(x_1, \ldots, x_s, T_n)} \overline{\mathcal{K}} \subseteq \overline{(T_n)}$, hence $\overline{(x_1, \ldots, x_s, T_n)} \subseteq \overline{(T_n)} : \overline{\mathcal{K}}$ and also $\overline{\mathcal{K}} \subseteq \overline{(T_n)} : \overline{(x_1, \ldots, x_s, T_n)}$. In order to conclude these containments are

actually equalities, it suffices to show they agree locally at the associated primes of the smaller ideal. To this end, it suffices to show that $\overline{\mathcal{K}}$ and $\overline{(x_1, \ldots, x_s, T_n)}$ have no common associated prime.

Suppose that there is some $\mathbf{q} \in \operatorname{Spec}(\mathcal{R}(E_1))$ with $\mathbf{q} \in \operatorname{Ass}(\overline{\mathcal{K}}) \cap \operatorname{Ass}(\overline{(x_1, \ldots, x_s, T_n)})$. From Proposition 5.7, we see that $\mathbf{q} \in \operatorname{Ass}\overline{\mathcal{K}} = \min\overline{\mathcal{K}}$, hence ht $\mathbf{q} = 1$. Moreover, as $\mathbf{q} \supseteq \overline{(x_1, \ldots, x_s, T_n)} \neq 0$, it follows that \mathbf{q} is a minimal prime of this ideal and so ht $\overline{(x_1, \ldots, x_s, T_n)} = 1$ as well. Thus \mathbf{q} is an ideal of height one which contains $\overline{\mathcal{K}} + \overline{(x_1, \ldots, x_s, T_n)} = \overline{I_s(\widetilde{C'})} + \overline{(x_1, \ldots, x_s, T_n)}$. However, from (5.5) and Lemma 5.8 it follows that, modulo $\overline{I_s(\widetilde{C'})} \neq 0$, $\overline{(x_1, \ldots, x_s, T_n)}$ contains a regular element. Hence ht $(\overline{\mathcal{K}} + \overline{(x_1, \ldots, x_s, T_n)}) \ge 2$, noting that $\mathcal{R}(E_1)$ is a Cohen-Macaulay domain by Proposition 5.5. However, this is a contradiction.

Now that $\overline{\mathcal{K}}$ and $\overline{(x_1,\ldots,x_s,T_n)}$ have been shown to be linked, the last statement follows from [17, 0.1].

With the $\mathcal{R}(E_1)$ -ideal $\overline{\mathcal{K}}$ and Proposition 5.9, we now introduce a divisorial ideal that will ultimately pave the path to the defining ideal \mathcal{J} of $\mathcal{R}(E)$. As previously noted, similar constructions and techniques have been applied in [3, 21, 37, 38].

Notation 5.10. Consider the divisorial ideal $\mathcal{D} = \frac{\overline{\ell_{n-e} \mathcal{K}}}{\overline{T_n}}$ and note that, since $\ell_{n-e} \in (x_1, \ldots, x_s, T_n)$, this is actually an $\mathcal{R}(E_1)$ -ideal, by Proposition 5.9.

Notice that, since $\mathcal{R}(E_1)$ is a domain and both $\overline{\ell_{n-e}} \neq 0$ and $\overline{T_n} \neq 0$, the ideals $\overline{\mathcal{K}}$ and \mathcal{D} are isomorphic. In particular, \mathcal{D} is a Cohen-Macaulay $\mathcal{R}(E_1)$ -ideal of height one following Proposition 5.7. Also notice that $\mathcal{D} \subseteq \overline{\mathcal{J}}$ since $\mathcal{D}(\overline{T_n}) \subseteq (\overline{\ell_{n-e}}) \subseteq \overline{\mathcal{J}}$, the ideal $\overline{\mathcal{J}}$ is prime, and $\overline{T_n} \notin \mathcal{J}$.

Proposition 5.11. With B' and C the matrices in (5.2) and $\mathcal{L} = (\ell_1, \ldots, \ell_{n-e})$, we have the equality

$$\overline{\mathcal{L} + I_s(B') + I_{s+1}(C)} = \mathcal{D}$$

in $\mathcal{R}(E_1)$.

Proof. Notice that, by Proposition 5.5, we have $\overline{\mathcal{L} + I_s(B') + I_{s+1}(C)} = (\overline{\ell_{n-e}}) + \overline{I_{s+1}(C)}$ and the only nonzero minors of $\overline{I_{s+1}(C)}$ are those involving the last column. With this, recall that $T_n \in \mathcal{K}$, hence

$$\overline{\ell_{n-e}} = \frac{\overline{\ell_{n-e}} \overline{T_n}}{\overline{T_n}}$$
(5.6)

and so $\overline{\ell_{n-e}} \in \mathcal{D}$. Moreover, to show that $\overline{I_{s+1}(C)} \subseteq \mathcal{D}$, we need only show that every $(s+1) \times (s+1)$ minor of \overline{C} involving the last column is contained in \mathcal{D} , as the other minors vanish. Without loss of generality, let M denote the submatrix consisting of the first s columns and the last column of C. By the matrix equation in (5.3) and Cramer's rule, in $\mathcal{R}(E_1)$ one has the equality

$$\overline{\det M} \cdot \overline{T_n} = \overline{\ell_{n-e}} \cdot \overline{\det M'} \tag{5.7}$$

where M' is the submatrix of C' consisting of its first s columns. Thus $\overline{\det M} = \frac{\overline{I_{n-e}} \overline{\det M'}}{\overline{T_n}} \in \mathcal{D}$ as well, and so it follows that $\overline{\mathcal{L} + I_s(B') + I_{s+1}(C)} \subseteq \mathcal{D}$.

To prove that the reverse containment holds, notice that $\overline{\mathcal{K}} = \overline{I_s(C')} + (\overline{T_n})$. Now, (5.6) shows that $\frac{\overline{\ell_{n-e}}T_n}{\overline{T_n}} = \overline{\ell_{n-e}} \in \overline{\mathcal{L}} + I_s(B') + I_{s+1}(C)$. Moreover, a similar argument using Cramer's rule and producing an equation similar to (5.7) proves that $\frac{\overline{\ell_{n-e}}I_s(C')}{(T_n)} \subseteq \overline{\mathcal{L}} + I_s(B') + I_{s+1}(C)$ as well.

Notice that Proposition 5.11 proves that the ideal \mathcal{D} agrees with the candidate for the defining ideal in Theorem 5.3, modulo \mathcal{J}_1 in $\mathcal{R}(E_1)$. Thus the induction argument will be complete once it has been shown that \mathcal{D} also agrees with $\overline{\mathcal{J}}$, and that this is a Cohen-Macaulay ideal. With this, we are ready to prove Theorem 5.3.

Proof of Theorem 5.3. Notice that $\mathcal{J}_1 \subseteq \mathcal{L} + I_s(B') + I_{s+1}(C) \subseteq \mathcal{J}$, where the first containment is a consequence of Proposition 5.5 and the second follows from Cramer's rule and the transition equations in (5.3). Thus it suffices to show that $\mathcal{D} = \overline{\mathcal{J}}$ in $\mathcal{R}(E_1)$, by Proposition 5.11. Moreover, with this containment, it is enough to prove that \mathcal{D} and $\overline{\mathcal{J}}$ agree locally at the associated primes of \mathcal{D} . As noted, \mathcal{D} and $\overline{\mathcal{K}}$ are isomorphic, hence by Proposition 5.7 we only to show that $\mathcal{D}_{\mathfrak{q}} = \overline{\mathcal{J}}_{\mathfrak{q}}$ for any prime \mathfrak{q} of $\mathcal{R}(E_1)$ with ht $\mathfrak{q} = 1$. We consider the following cases.

First, suppose that \mathfrak{q} is a height one prime ideal in $\mathcal{R}(E_1)$ and $\overline{(x_1,\ldots,x_s)} \notin \mathfrak{q}$. Recall from Proposition 3.8 that $\mathcal{J} = \mathcal{L} : (x_1,\ldots,x_s)^{\infty}$, hence $\overline{\mathcal{J}}_{\mathfrak{q}} = \overline{\mathcal{L}}_{\mathfrak{q}} = (\overline{\ell_{n-e}})_{\mathfrak{q}}$. Moreover, since $\overline{(x_1,\ldots,x_s,T_n)} \notin \mathfrak{q}$ as well, it follows from Proposition 5.9 that $\mathcal{R}(E_1)_{\mathfrak{q}} = \overline{(T_n)_{\mathfrak{q}}} : \overline{\mathcal{K}}_{\mathfrak{q}}$, hence $\overline{\mathcal{K}}_{\mathfrak{q}} \subseteq \overline{(T_n)}_{\mathfrak{q}}$. However, recall that $T_n \in \mathcal{K}$, and so $\overline{\mathcal{K}}_{\mathfrak{q}} = \overline{(T_n)}_{\mathfrak{q}}$. Thus $\mathcal{D}_{\mathfrak{q}} = \frac{(\overline{\ell_{n-e}})_{\mathfrak{q}}\overline{\mathcal{K}}_{\mathfrak{q}}}{(T_n)_{\mathfrak{q}}} = (\overline{\ell_{n-e}})_{\mathfrak{q}}$ as well.

Before we consider the case that $(x_1, \ldots, x_s) \subseteq \mathfrak{q}$, we observe that from the shape of φ in (5.1) and the description of \mathcal{J}_1 in Proposition 5.5 it follows that

$$\mathcal{J}_1 + (x_1, \dots, x_s) = (x_{s+1}T_n, \dots, x_{d-1}T_n) + I_s(B') + I_s(C_1) + (x_1, \dots, x_s).$$

Thus any prime ideal of $\mathcal{R}(E_1)$ containing $\overline{(x_1, \ldots, x_s)}$ contains either $\overline{(T_n)}$ or $\overline{(x_{s+1}, \ldots, x_{d-1})}$. In particular, we deduce that the set of minimal primes of $\overline{(x_1, \ldots, x_s)}$ is

$$\operatorname{Min}\left(\overline{x_1,\ldots,x_s}\right) = \left\{\overline{(x_1,\ldots,x_{d-1})}\right\} \bigcup \operatorname{Min}\left(\overline{x_1,\ldots,x_s,T_n}\right),\tag{5.8}$$

noting that $(\overline{x_1, \ldots, x_{d-1}})$ is a prime $\mathcal{R}(E_1)$ -ideal of height one. Indeed, this ideal defines the special fiber ring, a domain, of E_1 viewed as a module over $R' = k[x_1, \ldots, x_{d-1}]$, as in the proof of Proposition 5.5. With this, suppose that \mathfrak{q} is a prime of $\mathcal{R}(E_1)$ with height one such that $(\overline{x_1, \ldots, x_s}) \subseteq \mathfrak{q}$, and consider the following two cases of (5.8) above.

First, suppose that $\mathbf{q} = \overline{(x_1, \ldots, x_{d-1})}$, and notice that then $\overline{(x_1, \ldots, x_s, T_n)} \not\subseteq \mathbf{q}$. Thus by Proposition 5.9 it follows that $\overline{\mathcal{K}}_{\mathbf{q}} = \overline{(T_n)}_{\mathbf{q}}$. On the other hand, observe that $\overline{\ell_{n-e}} \notin \mathbf{q}$. Indeed, we have that $(\overline{\ell_{n-e}}) + \mathbf{q} = (\overline{x_d} \overline{T_n}) + \mathbf{q}$ and neither $\overline{T_n}$ nor $\overline{x_d}$ is contained in \mathbf{q} . Hence $\mathcal{D}_{\mathbf{q}} = \frac{(\overline{\ell_{n-e}})_{\mathbf{q}} \overline{\mathcal{K}}_{\mathbf{q}}}{(T_n)_{\mathbf{q}}} = (\overline{\ell_{n-e}})_{\mathbf{q}} = \mathcal{R}(E_1)_{\mathbf{q}}$. Moreover, as $\mathcal{D}_{\mathbf{q}} \subseteq \overline{\mathcal{J}}_{\mathbf{q}}$, it follows that $\overline{\mathcal{J}}_{\mathbf{q}} = \mathcal{R}(E_1)_{\mathbf{q}}$ as well.

Lastly, assume that $\mathbf{q} \in \operatorname{Min}(\overline{x_1, \ldots, x_s, T_n})$. Notice that $\overline{\mathcal{K}} \not \subseteq \mathbf{q}$ as $\overline{\mathcal{K}}$ and $\overline{(x_1, \ldots, x_s, T_n)}$ have no common associated prime, as noted in the proof of Proposition 5.9. Thus $\overline{\mathcal{K}}_{\mathbf{q}} = \mathcal{R}(E_1)_{\mathbf{q}}$ and so Proposition 5.9 implies that $\overline{(x_1, \ldots, x_s, T_n)}_{\mathbf{q}} = \overline{(T_n)}_{\mathbf{q}}$. Moreover, observe that $\ell_{n-e} \in (x_1, \ldots, x_s, T_n)$, whence $(\overline{\ell_{n-e}})_{\mathbf{q}} \subseteq \overline{(x_1, \ldots, x_s, T_n)}_{\mathbf{q}} = \overline{(T_n)}_{\mathbf{q}}$. We prove that this containment is an equality. Recall that $\mathcal{J}_1 \subset R'[T_1, \ldots, T_n]$ where $R' = k[x_1, \ldots, x_{d-1}]$, thus it follows that $\overline{x_d} \notin \mathbf{q}$, as ht $\mathbf{q} = 1$. Hence, in the localization at \mathbf{q} , it becomes a unit. Moreover, since $\ell_{n-e} \in (x_1, \ldots, x_s, x_d T_n)$, it follows that $\overline{\ell_{n-e}}$ is a unit multiple of $\overline{T_n}$ locally, hence $(\overline{\ell_{n-e})}_{\mathbf{q}} = \overline{(T_n)}_{\mathbf{q}}$. Thus $\mathcal{D}_{\mathbf{q}} = \frac{(\ell_{n-e})_{\mathbf{q}}\overline{\mathcal{K}}_{\mathbf{q}}}{(T_n)_{\mathbf{q}}} = \overline{\mathcal{K}}_{\mathbf{q}} = \mathcal{R}(E_1)_{\mathbf{q}}$. Again noting that $\mathcal{D}_{\mathbf{q}} \subseteq \overline{\mathcal{J}}_{\mathbf{q}}$, it follows that $\overline{\mathcal{J}}_{\mathbf{q}} = \mathcal{R}(E_1)_{\mathbf{q}}$ as well.

Now that it has been shown that $\mathcal{D} = \overline{\mathcal{J}}$, recall that this ideal is isomorphic to $\overline{\mathcal{K}}$, which is Cohen-Macaulay by Proposition 5.7. Thus $\overline{\mathcal{J}}$ is a Cohen-Macaulay $\mathcal{R}(E_1)$ -ideal, and so $\mathcal{R}(E) \cong \mathcal{R}(E_1)/\overline{\mathcal{J}}$ is Cohen-Macaulay.

Now that the defining ideal of Rees ring $\mathcal{R}(E)$ is understood, in the next subsection we proceed to analyze the special fiber ring $\mathcal{F}(E)$.

5.2. The fiber ring. With the assumptions of Setting 5.1, recall that the defining ideal of $\mathcal{R}(E)$ is identified in Theorem 5.3. From this, we can also determine the ideal defining the special fiber ring $\mathcal{F}(E)$.

Corollary 5.12. The special fiber ring of E is $\mathcal{F}(E) \cong k[T_1, \ldots, T_n]/I_s(B')$. Moreover, $\mathcal{F}(E)$ is a Cohen-Macaulay domain with dimension $\ell(E) = d + e - 1$. In particular, E has maximal analytic spread.

Proof. With the defining ideal \mathcal{J} in Theorem 5.3, we see that $\mathcal{J} + (x_1, \ldots, x_d) = I_s(B') + (x_1, \ldots, x_d)$ and the initial claim follows, noting that B' consists of entries in $k[T_1, \ldots, T_n]$. The second assertion then follows from Proposition 3.10.

Notice that from Corollary 4.6 and Corollary 5.12, the analytic spread of E differs in the row setting and the column setting; however, they coincide when s = d - 1, as expected from Remark 3.4.

Like in the column case (see Proposition 4.8), in addition to the description provided in Theorem 5.3, the defining ideal \mathcal{J} of $\mathcal{R}(E)$ may also be realized as a residual intersection.

Proposition 5.13. With the assumptions of Setting 5.1, the defining ideal \mathcal{J} of $\mathcal{R}(E)$ may be realized as

$$\mathcal{J} = \mathcal{L} : (x_1, \dots, x_s, x_{s+1}T_n, \dots, x_dT_n)$$

Moreover, this is an (n-e)-residual intersection.

Proof. Notice that we have the containments

$$\mathcal{L}: (x_1, \dots, x_s, x_{s+1}T_n, \dots, x_dT_n) \subseteq \mathcal{L}: (x_1, \dots, x_s) = \mathcal{J}$$
(5.9)

where the equality follows from Theorem 5.3. From Cramer's rule and (5.3), we have $I_{s+1}(C) \subseteq \mathcal{L}$: $(x_1, \ldots, x_s, T_n) \subseteq \mathcal{L}: (x_1, \ldots, x_s, x_{s+1}T_n, \ldots, x_dT_n)$. Moreover, using Cramer's rule again, we see $I_s(B') = I_d(B'') \subseteq \mathcal{L}: (x_1, \ldots, x_s, x_{s+1}T_n, \ldots, x_dT_n)$, where B'' is the matrix

$$B'' = \begin{pmatrix} B' & \psi \\ 0 & \cdots & 0 & 1 \\ \vdots & \vdots & \ddots \\ 0 & \cdots & 0 & & 1 \end{pmatrix}$$
(5.10)

obtained by replacing the T_n entries of the lower diagonal block of $B(\varphi)$ in (5.1) with 1s. Notice that this matrix satisfies the transition equation

$$[\ell_1 \dots \ell_{n-e}] = [x_1 \dots x_s \ x_{s+1} T_n \ \dots \ x_d T_n] \cdot B''.$$
(5.11)

Thus $\mathcal{L} + I_s(B') + I_{s+1}(C) \subseteq \mathcal{L} : (x_1, \dots, x_s, x_{s+1}T_n, \dots, x_dT_n)$, and so by Theorem 5.3 we have $\mathcal{J} = \mathcal{L} : (x_1, \dots, x_s, x_{s+1}T_n, \dots, x_dT_n)$. Moreover, as ht $\mathcal{J} = n - e$ and $\mathcal{L} = (\ell_1, \dots, \ell_{n-e})$ it follows that this is an (n-e)-residual intersection.

Remark 5.14. Although the defining ideal may be realized as a residual intersection, this description of \mathcal{J} in not very useful in order to describe the properties of \mathcal{J} . Indeed, it is particularly ill-behaved, as the ideal $I = (x_1, \ldots, x_s, x_{s+1}T_n, \ldots, x_dT_n)$ is not unmixed unless s = d - 1, in which case I is a complete intersection and a generating set of \mathcal{J} may be obtained from Theorem 2.10. Moreover, in this case the residual intersection of Proposition 5.13 coincides with the residual intersection of Proposition 4.8, and the main result of [10] is recovered. If however $s \leq d-2$, then I is not a complete intersection, and the techniques used in [10] fail in the present setting, so the method of successive approximations must be applied instead; compare with Remark 4.9 in the column setting.

With the bigrading on $R[T_1, \ldots, T_n]$ given by bideg $x_i = (1, 0)$ and bideg $T_i = (0, 1)$, observe that the equations of $I_{s+1}(C)$ in Theorem 5.3 have bidegree (1, s). Hence it follows that the module E is not of fiber type, once it has been shown that the equations of $I_{s+1}(C)$ are minimal generators of \mathcal{J} when $s \leq d-2$. We prove this in the following proposition, by exploiting the residual intersection property of \mathcal{J} .

Proposition 5.15. With the assumptions of Setting 5.1, the module E is of fiber type if and only if s = d-1.

Proof. Recall from Theorem 5.3 that the defining ideal of $\mathcal{R}(E)$ is $\mathcal{J} = \mathcal{L} + I_s(B') + I_{s+1}(C)$. From bidegree considerations, it follows that the fiber type property is equivalent to the containment $I_{s+1}(C) \subseteq \mathcal{L} + I_s(B')$ or rather $\mathcal{J} = \mathcal{L} + I_s(B')$. Thus we have already seen that E is of fiber type if s = d - 1 within the initial case of the induction proof (see also [10]), hence we only need to show the converse.

Suppose that E is of fiber type, and so $\mathcal{J} = \mathcal{L} + I_s(B')$. Writing $I = (x_1, \ldots, x_s, x_{s+1}T_n, \ldots, x_dT_n)$, recall from Proposition 5.13 that \mathcal{J} may also be realized as the (n - e)-residual intersection $\mathcal{J} = \mathcal{L} : I$. We prove that the ideals $\mathcal{L} : I$ and $\mathcal{L} + I_s(B')$ can only coincide when s = d - 1, by showing that otherwise these two descriptions of \mathcal{J} provide two distinct values for the type of $\mathcal{R}(E)$, denoted $r(\mathcal{R}(E))$ [4]. We will crucially use the fact that $r(\mathcal{R}(E))$ coincides with the minimal number of generators of the canonical module $\omega_{\mathcal{R}(E)}$ [4, 3.3.11].

Recall that B' is the submatrix of the Jacobian dual $B(\varphi)$ satisfying the transition equation in (5.3). Thus, B' is the Jacobian dual of the matrix φ_{d-s} with respect to the sequence x_1, \ldots, x_s . Also, φ_{d-s} presents E_{d-s} , and by Lemma 2.6 E_{d-s} is an A-module satisfying G_s , where $A = k[x_1, \ldots, x_s]$. Hence by [31, 4.11], the defining ideal of $\mathcal{R}(E_{d-s})$ is $\mathcal{L}_{d-s} : (x_1, \ldots, x_s) = \mathcal{L}_{d-s} + I_s(B')$, where $\mathcal{L}_{d-s} = (\ell_1, \ldots, \ell_{n-e-(d-s)})$. Moreover, this is a Cohen-Macaulay prime ideal of height n - e - (d - s). With this, we obtain that $\mathcal{J} = \mathcal{L} + I_s(B') = \mathcal{J}_{d-s} + (\ell_{n-e-(d-s)+1}, \ldots, \ell_{n-e})$. Since \mathcal{J} is a prime ideal of height n - e, it then follows that $\ell_{n-e-(d-s)+1}, \ldots, \ell_{n-e}$ is a regular sequence modulo \mathcal{J}_{d-s} . As $\mathcal{J}_{d-s} = (\ell_1, \ldots, \ell_{n-e-(d-s)}) + I_s(B')$ is Cohen-Macaulay of height n-e-(d-s), the transition equation (5.3) and Proposition 3.10 show that the complex $\mathbb{T} = \mathbb{T}(\mathcal{L}_{d-s}, B')$ of [5] is a minimal free resolution of $\mathcal{R}(E_{d-s})$. In particular, from [5, 3.5], the last Betti number of \mathbb{T} is $\binom{n-e-d+s}{s-1}$ and we note that this is precisely the type of $\mathcal{R}(E_{d-s})$ [4, 3.3.9]. Moreover, as the type of a ring is unchanged modulo a regular sequence [4, 1.2.19], from the previous observation, it follows that $\mu(\omega_{\mathcal{R}(E)}) = r(\mathcal{R}(E)) = \binom{n-e-d+s}{s-1}$ as well.

On the other hand, as $\mathcal{J} = \mathcal{L} : I$ is a residual intersection, by [19, 5.1] it follows that the calculated module $\omega_{\mathcal{R}(E)}$ can be realized as the symmetric power $\omega_{\mathcal{R}(E)} \cong S_B^{n-e-s}(I/\mathcal{L})$, where $B = R[T_1, \ldots, T_n]$. Notice that, due to the shape of φ in (5.1), we have that $\mu(I/\mathcal{L}) = s$. Indeed, modulo \mathcal{L} the images of $x_{s+1}T_n, \ldots, x_dT_n$ are contained in (x_1, \ldots, x_s) . Therefore, we obtain that

$$\mu(\omega_{\mathcal{R}(E)}) = \dim_k \mathcal{S}_B^{n-e-s}(I/\mathcal{L}) \otimes k = \dim_k \mathcal{S}_k^{n-e-s}(k^s) = \binom{n-e-1}{s-1}.$$

Thus, if $\mathcal{J} = \mathcal{L} : I = \mathcal{L} + I_s(B')$ we must have that $\binom{n-e-d+s}{s-1} = \binom{n-e-1}{s-1}$, from which it follows that s = d-1.

6. EXAMPLES AND FUTURE DIRECTIONS

In this section, we explore various questions related to modifying the conditions of Setting 3.1. When these conditions are changed, do the results obtained in this paper remain the same? Additionally, what are some potential next steps that could be taken to further explore the study of defining equations of Rees algebras when a module or ideal only satisfies the condition G_s but not G_{s+1} ?

As noted in Proposition 3.7 and Proposition 3.8, the fact that $\operatorname{Fitt}_{s+e-1}(E) = I_{n-s-e+1}(\varphi)$ has a unique minimal prime in Setting 3.1 is crucial to the proofs of Theorem 4.5 and Theorem 5.3. Whereas Proposition 3.7 shows that the assumptions of Setting 3.1 are sufficient to ensure this phenomenon, one might ask if the assumptions of Setting 3.1 are necessary. Moreover, one might ask what the structure of the defining ideal \mathcal{J} is when $\operatorname{Fitt}_{s+e-1}(E)$ has more than one minimal prime.

We present some examples here illustrating interesting phenomena related to these questions. Recall from Remark 3.2 that if the rank of E is e = 1, then E is isomorphic to a perfect R-ideal of grade two. Hence, for computational purposes, we provide examples in the case of these ideals, making use of the Hilbert-Burch theorem [14, 20.15] throughout. Each of the examples presented here was produced and verified through Macaulay2 [23].

Example 6.1. Consider the following 6×5 matrix with entries in $R = \mathbb{Q}[x_1, x_2, x_3, x_4]$:

$$\varphi = \begin{pmatrix} 0 & 0 & 0 & 0 & x_2 \\ x_2 & x_1 + x_2 & 0 & x_1 + x_2 & x_1 \\ 0 & 0 & x_3 & x_3 & x_4 \\ 0 & x_2 & x_1 + x_2 & 0 & x_1 + x_2 \\ x_4 & x_3 + x_4 & 0 & 0 & x_3 \\ 0 & 0 & x_4 & 0 & x_1 \end{pmatrix}$$

and consider the ideal $I = I_5(\varphi)$. One has that ht I = 2, hence I is perfect of height two. Moreover, the ideal I satisfies G_2 but not G_3 . In this example, $\operatorname{Fitt}_2(I) = I_4(\varphi)$ has two minimal prime ideals, (x_1, x_2) and (x_3, x_4) . As I satisfies assumptions (i) and (iii) of Setting 3.1, it follows from Proposition 3.7 that modulo any set of two linear forms, φ has rank at least 2.

With the bigrading on $R[T_1, \ldots, T_5]$ given by bideg $x_i = (1, 0)$ and bideg $T_i = (0, 1)$, the defining ideal \mathcal{J} of $\mathcal{R}(I)$ consists of the equations of $\mathcal{L} = ([T_1 \ldots T_5] \cdot \varphi)$ in bidegree (1, 1), one equation of bidegree (1, 3), two equations of bidegree (2, 2), and fiber equations with one of bidegree (0, 3), and four of bidegree (0, 4). Hence the shape of \mathcal{J} does not agree with either of the forms in Theorem 4.5 or Theorem 5.3, as it is minimally generated in different bidegrees.

Whereas, in the example above, the ideal $\operatorname{Fitt}_2(I)$ has two minimal primes, they are both complete intersections generated by s = 2 many linear forms; this is comparable to the behavior discussed in Proposition 3.7. However, even this behavior is not guaranteed and, as the next example shows, this Fitting ideal may have multiple minimal primes of differing codimensions.

Example 6.2. Consider the following 5×4 matrix with entries in $R = \mathbb{Q}[x_1, x_2, x_3, x_4]$:

$$\varphi = \begin{pmatrix} x_1 - x_2 & x_2 & x_2 & x_1 \\ x_2 & 0 & x_2 & x_1 \\ x_1 + x_2 & 0 & x_2 & x_1 \\ x_4 & x_1 & x_3 & 0 \\ x_1 & x_3 & x_1 & x_4 \end{pmatrix}$$

and consider the ideal $I = I_4(\varphi)$. One has that ht I = 2, hence I is perfect of height two. Moreover, I satisfies G_2 but not G_3 . Here, $\operatorname{Fitt}_2(I) = I_3(\varphi)$ has two minimal primes of different heights, namely (x_1, x_2) and (x_1, x_3, x_4) . As I satisfies assumptions (i) and (iii) of Setting 3.1, it follows from Proposition 3.7 that modulo any set of two linear forms, φ has rank at least 2. For instance, modulo (x_1, x_2) , φ has rank 2.

With the bigrading on $R[T_1, \ldots, T_5]$ given by bideg $x_i = (1, 0)$ and bideg $T_i = (0, 1)$, the defining ideal \mathcal{J} of $\mathcal{R}(I)$ consists of the equations of $\mathcal{L} = ([T_1 \ldots T_5] \cdot \varphi)$ in bidegree (1, 1), one equation of bidegree (2, 2), and one fiber equation with bidegree (0, 4). Hence the shape of \mathcal{J} does not agree with either of the forms in Theorem 4.5 or Theorem 5.3, as there is a minimal generator with bidegree (2, 2).

In light of Example 6.1 and Example 6.2, we can see that the results obtained in this article are not generally true in the absence of condition (ii) of Setting 3.1. However, there are still examples for a module E satisfying G_s , but not G_{s+1} , where $\operatorname{Fitt}_{s+e-1}(E)$ has $\mathfrak{p} = (x_1, \ldots, x_s)$ as its unique minimal prime, as in Proposition 3.7, even if φ has rank greater than 1 modulo this ideal.

Example 6.3. Consider the following 5×4 matrix with entries in $R = \mathbb{Q}[x_1, x_2, x_3, x_4]$:

$$\varphi = \begin{pmatrix} x_2 & 0 & x_2 & 0 \\ x_2 & x_1 & x_4 & x_2 \\ 0 & x_1 & x_2 & x_3 \\ 0 & x_2 & x_3 & x_1 \\ x_1 & x_2 & x_1 & x_4 \end{pmatrix}$$

and consider the ideal $I = I_4(\varphi)$. This ideal is perfect of grade two and satisfies G_2 but not G_3 . The ideal I satisfies conditions (i) and (iii) of Setting 3.1, and the ideal Fitt₂(I) has (x_1, x_2) as its only minimal prime. However, the rank of φ is 2 modulo (x_1, x_2) .

Moreover, Proposition 3.5 and the proof of Proposition 3.8 imply that the defining ideal of $\mathcal{R}(I)$ is $\mathcal{J} = \mathcal{L} : \mathfrak{p}^{\infty}$ where $\mathcal{L} = ([T_1 \dots T_5] \cdot \varphi)$ and $\mathfrak{p} = (x_1, x_2)$. Computations through *Macaulay2* [23] show that actually $\mathcal{J} = \mathcal{L} : \mathfrak{p}$ and that \mathcal{J} is minimally generated by the equations of \mathcal{L} and one fiber equation of bidegree (0, 2), which may be taken as the determinant of a 2 × 2 submatrix of $B(\varphi)$. In particular, \mathcal{J} has the shape of the ideal in Theorem 4.5 without φ or $B(\varphi)$ having the form in (4.1), after a change of coordinates.

As noted, the condition that $\operatorname{Fitt}_{s+e-1}(E)$ has a unique minimal prime allows the defining ideal \mathcal{J} of $\mathcal{R}(E)$ to be written as a saturation. Moreover, since the defining ideal \mathcal{J} in Example 6.3 resembles the behavior observed in Theorem 4.5, one might ask the following question.

Question 6.4. Can one characterize when $\operatorname{Fitt}_{s+e-1}(E)$ has a unique minimal prime if E is a module satisfying all of the assumptions of Setting 3.1 except for condition (ii)?

An answer to this question could possibly allow for a description of \mathcal{J} in a more general setting than the assumptions of Setting 3.1. Additionally, Example 6.1 and Example 6.2 lead to the following more general question.

Question 6.5. What conditions can be placed, to allow one to determine the minimal primes of the Fitting ideals of E, for a module E satisfying all conditions of Setting 3.1 except for the rank condition (ii)?

In [12], the notion of the *chaos invariant* was introduced to relate the minimal primes of Fitting ideals and the defining equations of the Rees ring, without relying on an assumption like condition (ii) of Setting 3.1. This invariant was introduced in the setting where $R = k[x_1, x_2, x_3]$ and I is a linearly presented perfect ideal of height 2 satisfying G_2 but not G_3 . Perhaps a generalization of the chaos invariant to $R = k[x_1, \dots, x_d]$ and a linearly presented R-module E satisfying G_s but not G_{s+1} could assist in answering Question 6.4 and Question 6.5. In addition to condition (ii) of Setting 3.1, we also remark that condition (i) is restrictive as well. The assumption that the module E has a presentation matrix φ consisting of linear entries was crucial for the arguments presented here and, aside from [3, 11, 8, 21], this condition has seldom been weakened in the context of modules of projective dimension one. However, it is curious if the techniques presented here can be combined with the techniques presented in [3], to study the case when condition (i) is relaxed.

Question 6.6. Can the equations defining $\mathcal{R}(E)$ be determined for a module E satisfying the assumptions of Setting 3.1 except for condition (i)?

In particular, as a natural extension of the work in [3, 8], one might consider ideals and modules of projective dimension one not satisfying G_d , that have *almost* linear presentation; that is, their presentation matrix φ consists of linear entries, except for one column of entries of a higher degree.

Example 6.7. Consider the following 5×4 matrix with entries in $R = \mathbb{Q}[x_1, x_2, x_3, x_4]$:

$$\varphi = \begin{pmatrix} x_1^2 & x_1 & x_2 & 0\\ 0 & 0 & x_1 & x_1\\ x_2^2 & x_2 & x_1 & 0\\ 0 & x_1 & x_2 & x_2\\ x_2^2 & x_1 & x_3 & x_4 \end{pmatrix}$$

and let $I = I_4(\varphi)$. The ideal I is perfect of height two and satisfies G_2 but not G_3 . Modulo (x_1, x_2) , the matrix φ has rank 1, and moreover $\operatorname{Fitt}_2(I)$ has a unique minimal prime, namely (x_1, x_2) .

By Proposition 3.5 and by repeating the proof of Proposition 3.8, it follows that the defining ideal \mathcal{J} of $\mathcal{R}(E)$ is $\mathcal{L} : \mathfrak{p}^{\infty}$ where $\mathcal{L} = ([T_1 \dots T_5] \cdot \varphi)$ and $\mathfrak{p} = (x_1, x_2)$. However, computations through *Macaulay2* [23] show that actually $\mathcal{J} = \mathcal{L} : \mathfrak{p}^2 \neq \mathcal{L} : \mathfrak{p}$, differing from the behavior in Theorem 4.5 or Theorem 5.3, but similar to the behavior in [3, 3.6], as the entries in the nonlinear column of φ have degree 2.

References

- [1] M. Artin and M. Nagata, Residual intersections in Cohen-Macaulay rings, J. Math. Kyoto Univ. 12 (1972), 307-323.
- [2] L. Avramov, Complete intersections and symmetric algebras, J. Algebra 73 (1981), 248–263.
- [3] J. A. Boswell and V. Mukundan, Rees algebras of almost linearly presented ideals, J. Algebra 460 (2016), 102–127.
- [4] W. Bruns and J. Herzog, Cohen-Macaulay rings, Cambridge University Press, Cambridge, 1993.
- [5] W. Bruns, A.R. Kustin, and M. Miller, the resolution of the generic residual intersection of a complete intersection, J. Algebra 128 (1990), 214–239.
- [6] M. Cooper and E. F. Price, Bounding the degrees of the defining equations of Rees rings for certain determinantal and Pfaffian ideals, J. Algebra 606 (2022), 613–653.
- [7] A. Corso, C. Polini and B. Ulrich, The structure of the core of ideals, Math. Ann. 321 (2001), 89–105.
- [8] A. Costantini, Cohen-Macaulay fiber cones and defining ideal of Rees algebras of modules, Women in Commutative Algebra – Proceedings of the 2019 WICA Workshop, Association for Women in Mathematics Series, vol. 29, Springer (2022).
- [9] A. Costantini, B. Drabkin and L. Guerrieri, The Rees algebra of ideals of star configurations, Linear Algebra Appl. 645 (2022), 91–122.
- [10] A. Costantini, E. F. Price, and M. Weaver, On Rees algebras of linearly presented ideals and modules, to appear in Collect. Math. https://doi.org/10.1007/s13348-024-00440-0, arxiv:2308.16010.
- [11] D. Cox, J. W. Hoffman, and H. Wang, Syzygies and the Rees algebra, J. Pure Appl. Algebra 212 (2008), 1787–1796.
- [12] A. Doria, Z. Ramos and A. Simis, Linearly presented perfect ideals of codimension 2 in three variables, J. Algebra 512 (2018), 216–251.
- [13] J. A. Eagon and D. G. Northcott, Ideals defined by matrices and a certain complex associated with them, Proc. Roy. Soc. Ser. A 269 (1962), 188–204.
- [14] D. Eisenbud, Commutative Algebra With a View Towards Algebraic Geometry, Springer, 1995.
- [15] L. Fouli, J. Montaño, C. Polini and B. Ulrich, The core of monomial ideals, to appear in Algebra Number Theory, arxiv:2303.10486.
- [16] J. Herzog, A. Simis and W. Vasconcelos, Koszul homology and blowing-up rings, in : Commutative Algebra, Proceedings, Trento, 1981, in: Lect. Notes Pure Appl. Math., vol. 84, Marcel Dekker, 1983, pp.79–169.
- [17] C. Huneke, Linkage and the Koszul homology of ideals, Amer. J. Math. 104 (1982), 1043–1062.
- [18] C. Huneke and M. Rossi, The dimension and components of symmetric algebras, J. Algebra 98 (1986), 200-210.
- [19] C. Huneke and B. Ulrich, Residual intersections, J. Reine Angew Math. 390 (1988), 1-20.
- [20] C. Huneke and B. Ulrich, Generic residual intersections, Lecture Notes in Math. 1430, Springer-Verlag, Berlin, 1990, 47–60.
- [21] A. Kustin, C. Polini and B. Ulrich, Rational normal scrolls and the defining equations of Rees algebras, J. Reine Angew. Math. 650 (2011), 23–65.

- [22] A. Kustin, C. Polini and B. Ulrich, The equations defining blowup algebras of height three Gorenstein ideals, Algebra Number Theory 11 (2017), 1489–1525.
- [23] D. R. Grayson and M. E. Stillman, Macaulay2, a software system for research in algebraic geometry. Available at http://www.math.uiuc.edu/Macaulay2/
- [24] J. Madsen, Equations of Rees algebras of ideals in two variables, (2015) arxiv:1511.04073.
- [25] S. Morey, Equations of blowups of ideals of codimension two and three, J. Pure Appl. Algebra 109 (1996), 197–211.
- [26] S. Morey and B. Ulrich, Rees algebras of ideals of low codimension, Proc. Amer. Math. Soc. 124 (1996), 3653–3661.
- [27] P. H. L. Nguyen, On Rees algebras of linearly presented ideals, J. Algebra 420 (2014), 186–200.
- [28] P. H. L. Nguyen, On Rees algebras of linearly presented ideals in three variables, J. Pure Appl. Algebra 221 (2017), 2180—2198.
- [29] A. Simis, B. Ulrich and W. Vasconcelos, Jacobian dual fibrations, Amer. J. Math. 115 (1993), 47–75.
- [30] A. Simis, B. Ulrich, and W. Vasconcelos, Cohen-Macaulay Rees algebras and degrees of polynomial relations, Math. Ann. 301 (1995), 421–444.
- [31] A. Simis, B. Ulrich, and W. Vasconcelos, Rees algebras of modules, Proc. London Math. Soc. 87 (2003), 610-646.
- [32] A. Simis, B. Ulrich, and W. V. Vasconcelos, Tangent algebras, Trans. Amer. Math. Soc. 364 (2012), 571–594.
- [33] B. Ulrich, Artin-Nagata properties and reductions of ideals, Contemp. Math. 159 (1994), 373–400.
- [34] B. Ulrich, Ideals having the expected reduction number, Amer. J. Math 118 (1996), 17–38.
- [35] B. Ulrich and W. V. Vasconcelos, The equations of Rees algebras of ideals with linear presentation, Math. Z. 214 (1993), 79–92.
- [36] W. V. Vasconcelos, On the equations of Rees algebras, J. Reine Angew. Math. 418 (1991), 189–218.
- [37] M. Weaver, On Rees algebras of ideals and modules over hypersurface rings, J. Algebra 636 (2023), 417-454.
- [38] M. Weaver, The equations of Rees algebras of height three Gorenstein ideals in hypersurface rings, J. Commut. Algebra 16 (2024), 123–149.
- [39] M. Weaver, Duality and the equations of Rees rings and tangent algebras, arxiv:2406.06766.

DEPARTMENT OF MATHEMATICS, TULANE UNIVERSITY, GIBSON HALL, NEW ORLEANS LA 70118 Email address: acostantini@tulane.edu

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, COLORADO COLLEGE, TUTT SCIENCE CENTER, COLORADO SPRINGS CO 80903

Email address: eprice@coloradocollege.edu

School of Mathematical and Statistical Sciences, Arizona State University, Wexler Hall, Tempe AZ 85281 *Email address:* matthew.j.weaver@asu.edu