8th International Conference on Computational and Mathematical Biomedical Engineering – CMBE2024 24-26 June 2024, USA P. Nithiarasu and R. Löhner (Eds.)

Multi-Scale Fiber Remodeling in HCM Using a Stress-Based Fiber Reorientation Law

Mohammad Mehri¹, Hossein Sharifi¹, Kenneth S. Campbell ², and Jonathan F. Wenk ^{1,3}

¹Department of Mechanical and Aerospace Engineering, University of Kentucky, Lexington, Kentucky, USA, {<u>m.mehri@uky.edu</u>, <u>hossein.sharifi@uky.edu</u>, <u>jonathan.wenk@uky.edu</u>}

² Division of Cardiovascular Medicine and Department of Physiology, University of Kentucky, Lexington, KY, USA, <u>k.s.campbell@uky.edu</u>

³Department of Surgery, University of Kentucky, Lexington, Kentucky, USA

SUMMARY

Quantifying fiber disarray, which is a prominent maladaptation associated with hypertrophic cardiomyopathy, remains critical to understanding the disease's complex pathophysiology. This study investigates the role of heterogeneous impairment of fiber contractility and fibrosis in the induction of disarray and their subsequent impact on cardiac pumping function. Fiber disarray is modeled via a stress-based fiber reorientation law within a multiscale finite element cardiac modeling framework called MyoFE. Using multiscale modeling capabilities, this study quantifies the distinct impacts of hypocontractility, hypercontractility and fibrosis on the development of fiber disarray and quantifies how their contributions affect the functional characteristics of the heart.

Keywords: *Multiscale modeling, Hypertrophic Cardiomyopathy (HCM), Finite element modeling, Cardiac mechanics, Fiber remodeling*

1 INTRODUCTION

Myocardial fiber disarray, a hallmark feature of hypertrophic cardiomyopathy (HCM), is defined as the irregular arrangement and disorganization of myocardial fibers. Heart muscle in HCM patients exhibits impaired contractility, fibrosis, and fiber disarray. The emergence of fiber disarray is commonly associated with contractile imbalance and fibrosis, yet the precise mechanisms are not fully understood. It has been reported that some HCM mutations result in hypercontractility, a condition previously observed to be caused by the mutation-induced unlocking of myosin heads from a state known as the super-relaxed state (SRX) [1]. HCM mutations do not all result in hypercontractility; different mutations alter the force generation mechanism in different ways, possibly including the inhibition of SRX [2]. The imbalance of contractile function (as a result of unequal fractions of hyper-hypo-contractile mutant proteins and wildtype proteins) will lead to cardiomyocyte disarray over time [3, 4]. In addition, replacement fibrosis, the most prevalent form of myocardial fibrosis in HCM [5], can induce heterogeneous changes in the material properties of muscle fibers [6] and is hypothesized to contribute to fiber disarray as well.

While previous modeling studies have incorporated the effects of fiber disarray on cardiac performance [7, 8], this study uniquely investigates the underlying mechanisms that trigger fiber reorientation and lead to myocardial fiber disarray. This study uses the MyoFE framework to identify the distinct contributions of contractile imbalance and fibrosis to fiber disarray. Moreover, this study aims to quantify the specific impact of microscopic abnormalities on global pump function. By elucidating how cell level alternation impacts organ level performance, this study aims to provide a deeper understanding of HCM disease progression.

2 METHODOLOGY

The current study presents a comprehensive modeling approach to investigate cardiac remodeling using a multiscale finite element (FE) framework, called MyoFE. This framework integrates various characteristics of cardiovascular function, including calcium handling, myofiber active contraction, and circulatory hemodynamics, in a closed loop model of the left ventricle (LV) and systemic circulation. In the present study, a fiber reorientation module is implemented within the MyoFE framework. At each time step, a novel stress-based reorientation law is used to compute fiber reorientation at each integration point in the mesh.



Figure 1: Stress based fiber reorientation. The unit vector f_0 represents the initial direction of myofiber and collagen. f_0 reorients toward the local traction vector Sf_0 , which is associated with the cross-sectional face of the fiber. S is the total stress tensor, encompassing both passive and active stresses.

As shown in Figure 1, f_0 represents the unit vector along the fiber direction in the reference configuration, which is shared by the myofibers and collagen fibers. The traction vector (Sf_0) is associated with the cross-sectional face of the fibers at a point in the LV. Our stress-based reorientation law incrementally reorients the fibers at each integration point in the LV mesh to be aligned with the traction vector as below:

$$\frac{df_0}{dt} = \frac{1}{\kappa} \left(\frac{Sf_0}{\|Sf_0\|} - f_0 \right) \tag{1}$$

where S is the 2nd Piola-Kirchhoff stress and κ is a time constant that is employed to enforce the separation of time scales. More specifically, changes in fiber orientation occur over an extended period of time (e.g., days to weeks), whereas the current study achieves comparable fiber reorientation within several heartbeats. Through a gradual reorientation process, the difference between the fiber direction and the direction of traction experienced by the fibers is minimized. To account for the reorientation that occurs in both the myofibers and collagen, it is necessary to employ the total stress tensor, which encompasses both active and passive stress components.

Based on previous observations in myocardium with HCM, we implemented the following perturbations to our LV models. In some HCM mutations, a substantial variance in force generation is observed within cardiomyocytes at physiological calcium concentrations [3, 4]. In addition, disruption of the SRX state, which modulates the contraction of the heart, is also reported [3, 9]. Accordingly, for hypercontractile models we incorporated an increased (up to 100%) K-SRX value for 30% of cardiomyocytes distributed heterogeneously within the myocardium, where K-SRX is the transition rate of SRX in our contraction model called MyoSim [1, 9]. For the hypocontractile models, K-SRX is reduced with a similar percentage in cardiomyocytes with impaired contractility. In order to model the effects of heterogeneous replacement fibrosis in the myocardium, 30% of the cardiomyocytes are assigned with a 100% increase in stiffness, as well as being isotropic [5, 6, 8]. To facilitate head-to-head comparisons, a specific random distribution of abnormal cells (based on the number and location of cells) is assigned to all cases. For all models, the initial

fiber configuration was assigned to the LV using a rule-based approach. Then, fiber reorientation was simulated until each case-specific fiber configuration reached steady state.

3 RESULTS AND CONCLUSIONS

As shown in Figure 2, all perturbed cases displayed considerable fiber disarray in contrast to the baseline model. As seen from the endocardial view, fiber disarray is more pronounced in hypocontractile and fibrous models whereas in the epicardial view, hypercontractility exhibits a greater degree of fiber disarray.



Figure 2: Final fiber configuration in LV models. A: Baseline LV model, B: LV model with heterogenous hypocontractility, C: LV model with heterogenous hypercontractility, D: LV model with heterogenous fibrosis. Top: Endocardial side of the myocardium. Bottom: Epicardial side of the myocardium.

In order to assess the sensitivity of fiber reorientation to the disruption of the SRX state, we implemented different levels of hypercontractility in order to evaluate the effect of K_{-SRX} on the induction of fiber disarray. By increasing K_{-SRX}, the average value of fiber orientation within the LV was observed to increase proportionally as shown in Figure 3.



Figure 3: The average value of fiber reorientation in LV models with different increased values of K_{-SRX} in hypercontractile cardiomyocytes.

As shown in Figure 4, pressure volume (PV) loops are compared to evaluate the effect of cell level perturbation on LV pumping performance. In the hypercontractile case, the end systolic volume (ESV) is reduced, and the PV loop is shifted to the left. Both hypocontractile and fibrous models result in a shrinkage of the PV loop and a significant increase in ESV due to reduced contractile abilities.



Figure 4: Pressure-volume (PV) loops of baseline and perturbed LV models.

In conclusion, the current study demonstrates the effectiveness of a stress-based law, within the multiscale cardiac modeling framework MyoFE, to predict fiber reorientation in the LV. These findings hold potential to enhance our understanding of pathological conditions and potentially aid in the development of therapeutic strategies to mitigate adverse remodeling effects due to HCM.

REFERENCES

- Vander Roest, A.S., et al., Hypertrophic cardiomyopathy beta-cardiac myosin mutation (P710R) leads to hypercontractility by disrupting super relaxed state. Proc Natl Acad Sci U S A, 2021. 118(24).
- [2] Vikhorev, P.G. and N.N. Vikhoreva, *Cardiomyopathies and Related Changes in Contractility of Human Heart Muscle*. International Journal of Molecular Sciences, 2018. 19(8).
- [3] Kraft, T. and J. Montag, *Altered force generation and cell-to-cell contractile imbalance in hypertrophic cardiomyopathy*. Pflugers Arch, 2019. 471(5): p. 719-733.
- [4] Montag, J., et al., Burst-Like Transcription of Mutant and Wildtype Alleles as Possible Origin of Cellto-Cell Contractile Imbalance in Hypertrophic Cardiomyopathy. Frontiers in Physiology, 2018. 9.
- [5] Liu, J., et al., *Patterns of Replacement Fibrosis in Hypertrophic Cardiomyopathy*. Radiology, 2022. 302(2): p. 298-306.
- [6] Sewanan, L.R., et al., *Extracellular Matrix From Hypertrophic Myocardium Provokes Impaired Twitch Dynamics in Healthy Cardiomyocytes*. JACC Basic Transl Sci, 2019. 4(4): p. 495-505.
- [7] Kovacheva, E., et al., *Causes of altered ventricular mechanics in hypertrophic cardiomyopathy: an in-silico study.* Biomed Eng Online, 2021. 20(1): p. 69.
- [8] Mojumder, J., et al., *Computational analysis of ventricular mechanics in hypertrophic cardiomyopathy patients.* Sci Rep, 2023. 13(1): p. 958.
- [9] Campbell, K.S., *Dynamic coupling of regulated binding sites and cycling myosin heads in striated muscle.* J Gen Physiol, 2014. 143(3): p. 387-99.