

THE WEYL TRANSFORM OF A COMPACTLY SUPPORTED DISTRIBUTION

MANSI MISHRA AND M. K. VEMURI

ABSTRACT. If T is a compactly supported distribution on \mathbb{R}^{2n} , then the Fourier transform of T is p -th power integrable if and only if the Weyl transform of T is p -th power traceable, and the Fourier transform of T vanishes at infinity if and only if the Weyl transform of T is a compact operator.

Let $\mathcal{H} = L^2(\mathbb{R}^n)$. Let $S^p(\mathcal{H})$ denote the p^{th} Schatten class of \mathcal{H} , $1 \leq p \leq \infty$ with the convention that $S^\infty(\mathcal{H})$ is the set of bounded operators on \mathcal{H} . Define $\rho : \mathbb{R}^{2n} \rightarrow S^\infty(\mathcal{H})$ by

$$(\rho(x, y)\varphi)(t) = e^{\pi i(x \cdot y + 2y \cdot t)}\varphi(t + x), \quad \varphi \in \mathcal{H}.$$

If $X \in S^1(\mathcal{H})$, the *Fourier-Wigner transform* of X is the function on \mathbb{R}^{2n} defined by

$$\alpha(X)(x, y) = \text{tr}(\rho(x, y)X).$$

Let $\mathcal{S}(\mathbb{R}^n)$ and $\mathcal{S}'(\mathbb{R}^n)$ denote the Schwartz class and the space of tempered distributions on \mathbb{R}^n respectively. For $\varphi, \psi \in \mathcal{S}(\mathbb{R}^n)$, let $\varphi \otimes \psi$ denote the operator whose kernel is $\varphi(x)\psi(y)$. If T is a tempered distribution on \mathbb{R}^{2n} , the *Weyl Transform* of T is the operator $W(T) : \mathcal{S}(\mathbb{R}^n) \rightarrow \mathcal{S}'(\mathbb{R}^n)$ defined by

$$(W(T)\varphi)(\psi) = T(\alpha(\varphi \otimes \psi)^\sim),$$

where $f^\sim(x, y) = f(-x, -y)$ (see [2, §5.3.2], but note that in that work, our Fourier-Wigner transform is called the Weyl transform, and our Weyl transform is called the inverse Weyl transform).

Theorem 1. *Let T be a compactly supported distribution on \mathbb{R}^{2n} . Let \hat{T} denote the Fourier transform of T . Then $W(T) \in S^p(\mathcal{H})$ if and only if $\hat{T} \in L^p(\mathbb{R}^{2n})$ for $1 \leq p \leq \infty$. Moreover, if K is a compact set in \mathbb{R}^{2n} , then there exists a constant C_K such that*

$$C_K^{-1} \left\| \hat{T} \right\|_p \leq \|W(T)\|_{S^p} \leq C_K \left\| \hat{T} \right\|_p,$$

whenever $\text{supp}(T) \subseteq K$. Furthermore, $\hat{T} \in \mathcal{C}_0(\mathbb{R}^{2n})$ if and only if $W(T)$ is compact.

When T is a Radon measure (i.e., when the order of T is zero), this theorem was proved by Luef and Samuelsen (see [3, Theorem 1.4]). Here we give a short proof of Theorem 1

2010 *Mathematics Subject Classification.* 42B10, 43A05, 47B10.

Key words and phrases. Symplectic Fourier transform; Schatten class.

based on Tim Steger's observation that Fourier-Wigner transforms of trace-class operators are locally the same as symplectic Fourier transforms of integrable functions (cf [6]).

Let $e : \mathbb{R}^{2n} \times \mathbb{R}^{2n} \rightarrow \mathbb{C}$ be the function defined by

$$e((\xi, \eta), (x, y)) = e^{2\pi i(\xi \cdot y - \eta \cdot x)}.$$

Recall that the symplectic Fourier transform of a function $f \in L^1(\mathbb{R}^{2n})$ is the function on \mathbb{R}^{2n} defined by

$$\check{f}(\xi, \eta) = \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} e((\xi, \eta), (x, y)) f(x, y) dx dy.$$

More generally (see [4]), if T is a tempered distribution on \mathbb{R}^{2n} , the symplectic Fourier transform of T is given by

$$\check{T}(\varphi) = T((\check{\varphi})^\sim), \quad \varphi \in \mathcal{S}(\mathbb{R}^{2n}).$$

Since the symplectic Fourier transform is a rotation of the Fourier transform, it follows that $\hat{T} \in L^p(\mathbb{R}^{2n})$ if and only if $\check{T} \in L^p(\mathbb{R}^{2n})$ and $\|\check{T}\|_p = \|\hat{T}\|_p$, and $\hat{T} \in \mathcal{C}_0(\mathbb{R}^{2n})$ if and only if $\check{T} \in \mathcal{C}_0(\mathbb{R}^{2n})$.

Let $g(x, y) = e^{-\frac{\pi}{2}(|x|^2 + |y|^2)}$, $(x, y) \in \mathbb{R}^{2n}$. Let $X \in S^1(\mathcal{H})$. Let $\beta(X)$ be the function on \mathbb{R}^{2n} defined by

$$\beta(X)(x, y) = g(x, y)\alpha(X)(x, y).$$

Let $X \in S^1(\mathcal{H})$, and let $\check{\beta}(X)$ denote the symplectic Fourier transform of $\beta(X)$. According to [7, Theorem 4.3], $\|\check{\beta}(X)\|_1 \leq \|X\|_{S^1}$. By Plancherel theorems for the Fourier transform and the Weyl transform (see e.g., [1]), it follows that

$$\|\check{\beta}(X)\|_2 = \|\beta(X)\|_2 \leq \|g\|_\infty \|\alpha(X)\|_2 = \|\alpha(X)\|_2 = \|X\|_{S^2}.$$

By the Calderon-Lions interpolation theorem ([5, Theorem 9.20, Example 1, Proposition 8]), it follows that if $1 \leq p \leq 2$, then

$$(1) \quad \|\check{\beta}(X)\|_p \leq \|X\|_{S^p}.$$

Let $f \in L^1(\mathbb{R}^{2n})$, and let $\Gamma(f) = W(g\check{f})$. Observe that $W(g) = \varphi \otimes \varphi$, where $\varphi(t) = 2^{-1/4}e^{-\pi|t|^2}$, $t \in \mathbb{R}^n$. It follows that $\|W(g)\|_{S^1} = 1$. By [7, Lemma 3.10], $\|\Gamma f\|_{S^1} \leq \|f\|_1$. If $f \in L^2(\mathbb{R}^{2n})$, then by Plancherel theorems for the Fourier transform and the Weyl transform,

$$\|\Gamma f\|_{S^2} = \|g\check{f}\|_2 \leq \|g\|_\infty \|\check{f}\|_2 = \|\check{f}\|_2 = \|f\|_2.$$

By the Calderon-Lions interpolation theorem, it follows that if $1 \leq p \leq 2$, then

$$(2) \quad \|\Gamma f\|_{S^p} \leq \|f\|_p.$$

Observe that $\check{\beta}$ is the formal adjoint of the operator Γ . By a duality argument, we conclude that Equations (1) and (2) hold for $2 \leq p \leq \infty$.

Let K be a compact set in \mathbb{R}^{2n} . Suppose T is supported in K , and $\check{T} \in L^q(\mathbb{R}^{2n})$, $1 \leq q \leq \infty$. Let f be a compactly supported smooth function which is identically one on K , and put $h(x, y) = e^{\frac{\alpha}{2}(|x|^2 + |y|^2)} f(x, y)$. Then \check{h} is integrable; put $C_K = \|\check{h}\|_1$. Let $T_1 = hT$. By Young's inequality, $\check{T}_1 = \check{h} * \check{T} \in L^q(\mathbb{R}^{2n})$.

Let p be the conjugate of q , i.e., $\frac{1}{p} + \frac{1}{q} = 1$. Define $L_T : S^p(\mathcal{H}) \rightarrow \mathbb{C}$ by

$$L_T(X) = \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \check{T}_1(x, y) \check{\beta}(X)(x, y) dx dy.$$

Then

$$|L_T(X)| \leq \|\check{\beta}(X)\|_p \|\check{T}_1\|_q \leq \|\check{\beta}(X)\|_p \|\check{h}\|_1 \|\check{T}\|_q \leq C_K \|X\|_{S^p} \|\check{T}\|_q.$$

Therefore L_T is a bounded linear functional on $S^p(\mathcal{H})$ and $\|L_T\|_{\text{op}} \leq C_K \|\check{T}\|_q$. Therefore, there exists $Y \in S^q(\mathcal{H})$ with $\|Y\|_{S^q} \leq C_K \|\check{T}\|_q$ such that $L_T(X) = \text{tr}(YX)$ ([5, Proposition 7]). However, if $\varphi, \psi \in \mathcal{S}(\mathbb{R}^n)$, then

$$\begin{aligned} (W(T)\varphi)(\psi) &= T(\alpha(\varphi \otimes \psi)^\sim) \\ &= T(h\beta(\varphi \otimes \psi)^\sim) \\ &= T_1(\beta(\varphi \otimes \psi)^\sim) \\ &= \check{T}_1(\check{\beta}(\varphi \otimes \psi)) \\ &= \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \check{T}_1(x, y) \check{\beta}(\varphi \otimes \psi)(x, y) dx dy \\ &= L_T(\varphi \otimes \psi) \\ &= \text{tr}(Y(\varphi \otimes \psi)) \\ &= \int_{\mathbb{R}^n} (Y\varphi)(t)\psi(t) dt \\ &= (Y\varphi)(\psi). \end{aligned}$$

Therefore $W(T) = Y$, and hence $W(T) \in S^q(\mathcal{H})$ and

$$(3) \quad \|W(T)\|_{S^q} \leq C_K \|\check{T}\|_q.$$

Conversely, assume that T is supported in K , and $W(T) \in S^q(\mathcal{H})$, $1 \leq q \leq \infty$. Let $Z = \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \check{h}(x, y) \rho(x, y) W(T) \rho(x, y)^{-1} dx dy$. Then $Z \in S^q(\mathcal{H})$, and $\|Z\|_{S^q} \leq C_K \|W(T)\|_{S^q}$ (cf the proof of [7, Theorem 3.7]). Define $M_T : L^p(\mathbb{R}^{2n}) \rightarrow \mathbb{C}$ by

$$M_T(\varphi) = \text{tr}(\Gamma(\varphi)Z), \quad \varphi \in L^p(\mathbb{R}^{2n}).$$

Then M_T is a bounded linear functional on $L^p(\mathbb{R}^{2n})$ and $\|M_T\|_{\text{op}} \leq C_K \|W(T)\|_{S^q}$. Therefore there exists $\psi \in L^q(\mathbb{R}^{2n})$ with $\|\psi\|_q \leq C_K \|W(T)\|_{S^q}$ such that $M_T(\varphi) = \int \varphi \psi$. It

follows like before that $\psi = \check{T}$. Therefore, $\check{T} \in L^q(\mathbb{R}^{2n})$, and

$$C_K^{-1} \left\| \check{T} \right\|_q \leq \|W(T)\|_{S^q}.$$

Now suppose $\check{T} \in \mathcal{C}_0(\mathbb{R}^{2n})$. Let $\varepsilon > 0$. Let B denote the closed unit ball centered at 0 in \mathbb{R}^{2n} and put $K = \text{supp}(T) + B$ (Minkowski sum). Then K is compact. Let ρ be a non-negative smooth function supported in B with $\int \rho = 1$. For each $r > 1$, let ρ_r be defined by $\rho_r(x, y) = r^{2n} \rho(rx, ry)$. Observe that for each $r > 0$, $\check{\rho}_r \in L^1(\mathbb{R}^{2n})$, $\|\check{\rho}_r\|_\infty = 1$, and $\lim_{r \rightarrow \infty} \check{\rho}_r = 1$ uniformly on compact sets. It follows that for sufficiently large r , $\left\| \check{\rho}_r \check{T} - \check{T} \right\|_\infty < \varepsilon / C_K$. Put $Y = W(\rho_r * T)$. Then $Y \in S^1(\mathcal{H})$. Since $\rho_r * T - T$ is supported in K , it follows from Equation (3) that

$$\|Y - W(T)\|_{\text{op}} = \|W(\rho_r * T - T)\|_{\text{op}} \leq C_K \left\| \check{\rho}_r \check{T} - \check{T} \right\|_\infty < \varepsilon.$$

Therefore $W(T)$ is compact. An analogous argument where convolution is replaced by twisted convolution shows that if $W(T)$ is compact, then $\check{T} \in \mathcal{C}_0(\mathbb{R}^{2n})$.

REFERENCES

1. Gerald B. Folland, *Harmonic analysis in phase space*, Annals of Mathematics Studies, vol. 122, Princeton University Press, Princeton, NJ, 1989. MR 983366
2. M. Keyl, J. Kiukas, and R. F. Werner, *Schwartz operators*, Rev. Math. Phys. **28** (2016), no. 3, 1630001, 60. MR 3499017
3. Franz Luef and Helge Jørgen Samuelsen, *Fourier restriction for Schatten class operators and functions on phase space*, arXiv:2407.16259 (2024).
4. J.-M. Maillard, *On the twisted convolution product and the Weyl transformation of tempered distributions*, J. Geom. Phys. **3** (1986), no. 2, 230–261. MR 879367
5. Michael Reed and Barry Simon, *Methods of modern mathematical physics. II. Fourier analysis, self-adjointness*, Academic Press [Harcourt Brace Jovanovich, Publishers], New York-London, 1975. MR 493420
6. M. K. Vemuri, *Realizations of the canonical representation*, ProQuest LLC, Ann Arbor, MI, 1997, Thesis (Ph.D.)—The University of Chicago. MR 2716708
7. ———, *Realizations of the canonical representation*, Proc. Indian Acad. Sci. Math. Sci. **118** (2008), no. 1, 115–131. MR 2434197

DEPARTMENT OF MATHEMATICAL SCIENCES, IIT(BHU), VARANASI 221005.