

# R<sub>II</sub> TYPE THREE TERM RELATIONS FOR BIVARIATE POLYNOMIALS ORTHOGONAL WITH RESPECT TO VARYING WEIGHTS

CLEONICE F. BRACCIALI, ANTONIA M. DELGADO, LIDIA FERNÁNDEZ, AND  
TERESA E. PÉREZ

ABSTRACT. Given a bivariate weight function defined on the positive quadrant of  $\mathbb{R}^2$ , we study polynomials in two variables orthogonal with respect to varying measures obtained by special modifications of this weight function. In particular, the varying weight functions are given by the multiplication of  $x_1^{-n}x_2^{-n}$  times the original weight function. Apart from the question of the existence and construction of such kind of orthogonal polynomials, we show that the systems of bivariate polynomials orthogonal with respect to this kind of varying weights satisfy R<sub>II</sub> type three term relations, one for every variable. A method to construct bivariate orthogonal systems with respect to varying weights based in the Koornwinder's method is developed. Finally, several examples and particular cases have been analysed.

## 1. INTRODUCTION

Consider a positive measure  $\psi(x)$  defined on the real interval  $(a, b)$  and the sequence of monic orthogonal polynomials,  $\{P_n(x)\}_{n \geq 0}$  that satisfy the orthogonality conditions

$$\int_a^b x^k P_n(x) d\psi(x) = 0, \quad 0 \leq k \leq n-1.$$

It is well known that the polynomials,  $P_n(x)$ , satisfy the three term recurrence relation

$$P_{n+1}(x) = (x - c_{n+1})P_n(x) - \lambda_{n+1}P_{n-1}(x), \quad n \geq 0,$$

with  $P_{-1}(x) = 0$ ,  $P_0(x) = 1$ ,  $\lambda_{n+1} \neq 0$ , and that these polynomials are connected with continued fractions known as  $J$ -fractions, see Chihara [4].

Within the scope of the connection between orthogonal polynomials and continued fractions Ismail and Masson in [7] investigated what they call continued fractions of type R<sub>I</sub> and of type R<sub>II</sub>. The continued fractions of

---

*Date:* September 26, 2024.

*2020 Mathematics Subject Classification.* Primary: 42C05; 33C47.

*Key words and phrases.* Bivariate orthogonal polynomials; varying weights; R<sub>II</sub> type three term relations.

type  $R_I$  are associated with sequences of polynomials that satisfy the  $R_I$  type three term recurrence relation in the form

$$P_{n+1}(x) = (x - c_{n+1})P_n(x) - \lambda_{n+1}(x - a_{n+1})P_{n-1}(x), \quad n \geq 0,$$

with  $P_{-1}(x) = 0$ ,  $P_0(x) = 1$ , and  $\lambda_{n+1} \neq 0$ . If  $P_n(a_k) \neq 0$ , for  $n \geq 1$ ,  $k = 1, 2, \dots, n$ , then there is a linear functional  $\mathcal{L}$  such that the rational functions  $P_n(x)/\prod_{j=1}^n(x - a_j)$  satisfy the orthogonality conditions

$$\mathcal{L} \left[ x^k \frac{P_n(x)}{\prod_{j=1}^n(x - a_j)} \right] = 0, \quad 0 \leq k \leq n - 1.$$

The special case of  $R_I$  type three term recurrence relations when  $a_{n+1} = 0$  appeared in [16], associated with polynomials that satisfy

$$\int_a^b x^k P_n(x) \frac{d\psi(x)}{x^n} = 0, \quad 0 \leq k \leq n - 1, \quad (1.1)$$

where  $0 \leq a < b \leq \infty$  and  $\psi(x)$  is a positive measure such that

$$\int_a^b x^k d\psi(x) < \infty, \quad k = 0, \pm 1, \pm 2, \dots$$

Similar polynomials have first appeared in [9], related to orthogonal Laurent polynomials (see also [6]). In [15], another special case of  $R_I$  type three term recurrence relation have been studied with applications to orthogonal polynomials on the unit circle.

On the other hand, continued fractions of type  $R_{II}$  are associated with sequences of polynomials that satisfy the  $R_{II}$  type three term recurrence relation

$$P_{n+1}(x) = (x - c_{n+1})P_n(x) - \lambda_{n+1}(x - a_{n+1})(x - b_{n+1})P_{n-1}(x), \quad n \geq 0,$$

with  $P_{-1}(x) = 0$ ,  $P_0(x) = 1$ , and  $\lambda_{n+1} \neq 0$ . If  $P_n(a_k) \neq 0$  and  $P_n(b_k) \neq 0$ , for  $n \geq 1$  and  $k = 1, 2, \dots, n$ , there exists a linear functional  $\mathcal{L}$  such that the rational functions  $P_n(x)/\prod_{j=1}^n(x - a_j)(x - b_j)$  satisfy the orthogonality conditions

$$\mathcal{L} \left[ x^k \frac{P_n(x)}{\prod_{j=1}^n(x - a_j)(x - b_j)} \right] = 0, \quad 0 \leq k \leq n - 1.$$

Zhedanov in [18] studied the connection of these sequences of polynomials with biorthogonal rational functions and the generalized eigenvalue problem. The special case of  $R_{II}$  type three term recurrence relation

$$P_{n+1}(x) = (x - c_{n+1})P_n(x) - \lambda_{n+1}(x^2 + 1)P_{n-1}(x), \quad n \geq 0,$$

was studied by Ismail and Sri Ranga [8] in connection with orthogonal polynomials on the unit circle. In [8] the authors have given some conditions for coefficients  $c_{n+1}$  and  $\lambda_{n+1}$  such that these polynomials satisfy

$$\int_{-\infty}^{\infty} x^k P_n(x) \frac{d\psi(x)}{(x^2 + 1)^n} = 0, \quad 0 \leq k \leq n - 1, \quad (1.2)$$

where  $\psi(x)$  is a positive measure.

Polynomials that satisfy (1.1) or (1.2) can be seen as orthogonal polynomials with respect to varying measures. In the literature we can find many references on orthogonal polynomials with respect to varying measures, see [2, 3, 17], and specially on varying measures defined on the unit circle, see [1, 13, 14].

The theory of orthogonal polynomials in one variable has been extended to several variables with many applications (see, for instance, [5] and the references therein). Hence, it seems reasonable to consider if it is possible to extend the R<sub>I</sub> or the R<sub>II</sub> type three term relation to several variables. This paper addresses the second relation for the bivariate case, see Theorem 4.2.

The goal of this paper is to extend results and properties about R<sub>II</sub> type three term relations for systems of bivariate polynomials that satisfy orthogonality conditions with respect to varying weights in two variables. In particular, we work with varying weight functions of two variables in the form

$$W_n(x_1, x_2) = x_1^{-n} x_2^{-n} W(x_1, x_2), \quad n \geq 0,$$

where  $W(x_1, x_2)$  is a weight function defined on a domain  $\Omega$  contained on the first quadrant of  $\mathbb{R}^2$ . We define and prove the existence of bivariate orthogonal polynomials systems with respect to a varying weight. As main result, we found R<sub>II</sub> type three term relations satisfied by these polynomials. We provide methods to construct the polynomials: one method is based on the moment matrices, another method is inspired by the well-known Koornwinder's method to construct bivariate orthogonal polynomials. Examples are also presented.

The paper is structured as follows. Section 2 brings the basic facts about bivariate polynomial systems. In Section 3 we define varying weight functions of two variables and bivariate orthogonal systems with respect to varying weights. Besides, we prove the existence of these polynomials and how to construct them. In Section 4 we obtain the R<sub>II</sub> type three term relations satisfied by the polynomials, as well as R<sub>II</sub> type three term recurrence relations. In Section 5 a method to construct bivariate orthogonal systems with respect to varying weights based in the Koornwinder's method is developed. The last section brings several examples, in the first one the varying weight function in two variables is given by a product of one variable ones. In this case, the associated polynomials are given by product of orthogonal polynomials with respect to the varying weight functions in one variable. Second example uses the construction method developed on Section 5, where the domain  $\Omega$  is a triangle on the first quadrant of  $\mathbb{R}^2$ . In the last example we analyse a varying weight function defined in a shifted simplex on  $\mathbb{R}^2$ .

## 2. BASIC TOOLS AND FIRST RESULTS

We collect the basic tools and first results that we will need along this work. For more information, see [5].

A bivariate polynomial of total degree  $n \geq 0$  with real coefficients is a linear combination of the monomials in two variables as

$$p(x_1, x_2) = \sum_{\substack{i+j=0 \\ i,j \geq 0}}^n c_{i,j} x_1^i x_2^j, \quad c_{i,j} \in \mathbb{R}.$$

When there exists only one term of highest degree in the form

$$p(x_1, x_2) = c_{n-k,k} x_1^{n-k} x_2^k + \sum_{\substack{i+j=0 \\ i,j \geq 0}}^{n-1} c_{i,j} x_1^i x_2^j,$$

we say that it is *monomial*, and when  $c_{n-k,k} = 1$ , the polynomial is called *monic*.

Let us denote by  $\Pi_n$  the linear space of bivariate polynomials in two variables of total degree less than or equal to  $n$  with real coefficients. Then

$$\Pi_n = \text{span}\{1, x_1, x_2, x_1^2, x_1 x_2, x_2^2, \dots, x_1^n, x_1^{n-1} x_2, \dots, x_2^n\},$$

and

$$\dim \Pi_n = (n+2)(n+1)/2 = t_n. \quad (2.1)$$

Let  $\Pi = \cup_{n \geq 0} \Pi_n$ . The canonical basis for  $\Pi$  can be written as a sequence  $\{\mathbb{X}_n\}_{n \geq 0}$  of vectors of increasing size  $n+1$  such that  $\mathbb{X}_n$  contains the  $n+1$  different monomials of total degree  $n$  arranged in reverse lexicographical order, this is

$$\mathbb{X}_n = \begin{pmatrix} x_1^n \\ x_1^{n-1} x_2 \\ \vdots \\ x_1 x_2^{n-1} \\ x_2^n \end{pmatrix}, \quad n \geq 0. \quad (2.2)$$

Observe that the set of entries of  $\{\mathbb{X}_m\}_{m=0}^n$  forms a basis of  $\Pi_n$ .

Now, we define the vectors for negative indexes.

**Definition 2.1.** For  $n \geq 1$ , we define

$$\mathbb{X}_{-n} = x_1^{-n} x_2^{-n} \mathbb{X}_n = \begin{pmatrix} x_2^{-n} \\ x_1^{-1} x_2^{-n+1} \\ \vdots \\ x_1^{-n+1} x_2^{-1} \\ x_1^{-n} \end{pmatrix}.$$

Consider  $L_{n,1}$  and  $L_{n,2}$  matrices of size  $(n + 1) \times (n + 2)$  given by

$$L_{n,1} = \left( \begin{array}{ccc|c} 1 & \circ & 0 \\ \cdot & \cdot & \vdots \\ \circ & & 1 & 0 \end{array} \right) \quad \text{and} \quad L_{n,2} = \left( \begin{array}{c|cc} 0 & 1 & \circ \\ \vdots & & \cdot \\ 0 & \circ & 1 \end{array} \right). \quad (2.3)$$

Observe that  $L_{n,i}$  are full rank matrices, such that  $L_{n,i} L_{n,i}^T = I_{n+1}$ . For  $n \geq 0$ , we get

$$x_1 \mathbb{X}_n = x_1 \begin{pmatrix} x_1^n \\ x_1^{n-1} x_2 \\ \vdots \\ x_1 x_2^{n-1} \\ x_2^n \end{pmatrix} = \begin{pmatrix} x_1^{n+1} \\ x_1^n x_2 \\ \vdots \\ x_1^2 x_2^{n-1} \\ x_1 x_2^n \end{pmatrix} = L_{n,1} \mathbb{X}_{n+1},$$

analogously,  $x_2 \mathbb{X}_n = L_{n,2} \mathbb{X}_{n+1}$ . In addition, for  $n \geq 1$ ,

$$x_1^{-1} \mathbb{X}_{-n} = x_1^{-1} \begin{pmatrix} x_2^{-n} \\ x_1^{-1} x_2^{-n+1} \\ \vdots \\ x_1^{-n+1} x_2^{-1} \\ x_1^{-n} \end{pmatrix} = \begin{pmatrix} x_1^{-1} x_2^{-n} \\ x_1^{-2} x_2^{-n+1} \\ \vdots \\ x_1^{-n} x_2^{-1} \\ x_1^{-n-1} \end{pmatrix} = L_{n,2} \mathbb{X}_{-n-1},$$

and analogously,  $x_2^{-1} \mathbb{X}_{-n} = L_{n,1} \mathbb{X}_{-n-1}$ .

We can abbreviate above expressions in the form

$$\text{for } n \geq 1, \quad x_i^{-1} \mathbb{X}_{-n} = L_{n,j} \mathbb{X}_{-(n+1)}, \quad i, j = 1, 2 \quad \text{and} \quad i \neq j.$$

also,

$$\text{for } n \geq 0, \quad x_i \mathbb{X}_n = L_{n,i} \mathbb{X}_{n+1}, \quad i = 1, 2. \quad (2.4)$$

A basis of  $\Pi$  can be organized as sequences of polynomial vectors of increasing size whose entries are independent polynomials of the same total degree.

**Definition 2.2** ([11, 12]). A polynomial system (PS) is a sequence of column polynomial vectors  $\{\mathbb{P}_n\}_{n \geq 0}$  of increasing size  $n + 1$  in the form

$$\mathbb{P}_n = (P_{n,0}(x_1, x_2), P_{n-1,1}(x_1, x_2), \dots, P_{0,n}(x_1, x_2))^T,$$

such that every entry  $P_{n-k,k}(x_1, x_2)$ ,  $k = 0, 1, \dots, n$ , is a bivariate polynomial of total degree  $n$ , and  $\{P_{n,0}(x_1, x_2), P_{n-1,1}(x_1, x_2), \dots, P_{0,n}(x_1, x_2)\}$  is a linearly independent set.

Observe that the set of entries of  $\{\mathbb{P}_m\}_{m=0}^n$  is a basis of  $\Pi_n$ , and  $\{\mathbb{X}_n\}_{n \geq 0}$  defined as in (2.2) is a PS. Therefore, every polynomials vector,  $\mathbb{P}_n$ , can be expressed as

$$\mathbb{P}_n = G_n \mathbb{X}_n + G_{n,n-1} \mathbb{X}_{n-1} + G_{n,n-2} \mathbb{X}_{n-2} + \dots + G_{n,1} \mathbb{X}_1 + G_{n,0} \mathbb{X}_0, \quad (2.5)$$

where  $G_{n,j}$  are matrices of size  $(n+1) \times (j+1)$ , and the square matrix  $G_n = G_{n,n}$  is non-singular, and it is called the *matrix leading coefficient* of  $\mathbb{P}_n$ .

Moreover, a PS is called *monic* if every involved polynomial is monic as

$$P_{n-k,k}(x_1, x_2) = x_1^{n-k} x_2^k + \sum_{\substack{i+j=0 \\ i,j \geq 0}}^{n-1} c_{i,j} x_1^i x_2^j, \quad c_{i,j} \in \mathbb{R}.$$

In that case, the matrix leading coefficient is the identity,  $G_n = I_{n+1}$ .

When computing a monic PS, every entry as in (2) has  $t_{n-1} = n + (n-1) + \dots + 2 + 1$  coefficients  $c_{i,j} \in \mathbb{R}$ , and then, the number of real coefficients in the monic polynomial vector  $\mathbb{P}_n$  is

$$t_{n-1} \times (n+1) = \frac{n(n+1)^2}{2}. \quad (2.6)$$

### 3. BIVARIATE VARYING ORTHOGONALITY

Let  $\Omega$  be a domain on the first quadrant of  $\mathbb{R}^2$ , i.e.,  $\Omega \subset \mathbb{R}_+^2 = \{(x_1, x_2) \in \mathbb{R}^2 : x_1, x_2 > 0\}$ , and let  $W(x_1, x_2)$  be a weight function defined on  $\Omega$  such that

$$\mu_{k,m} = \iint_{\Omega} x_1^k x_2^m W(x_1, x_2) dx_1 dx_2 < \infty,$$

for  $k, m = 0, \pm 1, \pm 2, \dots$ . For every two polynomials  $f, g \in \Pi$ , we define the inner product

$$\langle f, g \rangle = \iint_{\Omega} f(x_1, x_2) g(x_1, x_2) W(x_1, x_2) dx_1 dx_2.$$

For a given weight function  $W(x_1, x_2)$ , we are going to work with varying weight functions in the form

$$W_n(x_1, x_2) = x_1^{-n} x_2^{-n} W(x_1, x_2), \quad (3.1)$$

for  $n \geq 0$ , that are positive on  $\Omega \subset \mathbb{R}_+^2$ . To this end, we start studying this kind of varying orthogonality.

Given a natural number  $n \geq 0$ , we define

$$\begin{aligned} \mu_{k,m}^{(n)} &= \iint_{\Omega} x_1^k x_2^m W_n(x_1, x_2) dx_1 dx_2 \\ &= \iint_{\Omega} x_1^{-n+k} x_2^{-n+m} W(x_1, x_2) dx_1 dx_2 = \mu_{-n+k, -n+m}, \end{aligned}$$

for  $k, m = 0, \pm 1, \pm 2, \dots$ . Observe that we have defined the moments for the varying weight (3.1), and they are well defined.

Associated with the varying weight  $W_n(x_1, x_2)$ , we define the  $n$ th varying inner product, in short *VIP*,

$$\langle f, g \rangle_n = \iint_{\Omega} f(x_1, x_2) g(x_1, x_2) W_n(x_1, x_2) dx_1 dx_2 \quad (3.2)$$

$$= \iint_{\Omega} f(x_1, x_2) g(x_1, x_2) x_1^{-n} x_2^{-n} W(x_1, x_2) dx_1 dx_2.$$

Obviously,  $\langle \cdot, \cdot \rangle = \langle \cdot, \cdot \rangle_0$ .

**3.1. Moment matrices.** For  $n, r, s \geq 0$ , we define the matrix  $M_{r,s}^{(n)}$  of size  $(r+1) \times (s+1)$ , as

$$\begin{aligned} M_{r,s}^{(n)} &= \langle \mathbb{X}_r, \mathbb{X}_s^T \rangle_n = \iint_{\Omega} \mathbb{X}_r \mathbb{X}_s^T W_n(x_1, x_2) dx_1 dx_2 \\ &= \begin{pmatrix} \mu_{r+s,0}^{(n)} & \mu_{r+s-1,1}^{(n)} & \cdots & \mu_{r,s}^{(n)} \\ \mu_{r+s-1,1}^{(n)} & \mu_{r+s-2,2}^{(n)} & \cdots & \mu_{r-1,s+1}^{(n)} \\ \vdots & \vdots & & \vdots \\ \mu_{s,r}^{(n)} & \mu_{s-1,r+1}^{(n)} & \cdots & \mu_{0,r+s}^{(n)} \end{pmatrix}. \end{aligned} \quad (3.3)$$

For  $0 \leq k \leq s$ , we denote each column of  $M_{r,s}^{(n)}$  by  $\mathbf{m}_{r,s,k}^{(n)}$ , hence

$$M_{r,s}^{(n)} = \begin{pmatrix} \mathbf{m}_{r,s,0}^{(n)} & \mathbf{m}_{r,s,1}^{(n)} & \cdots & \mathbf{m}_{r,s,s}^{(n)} \end{pmatrix},$$

with

$$\begin{aligned} \mathbf{m}_{r,s,k}^{(n)} &= \iint_{\Omega} \mathbb{X}_r x_1^{s-k} x_2^k W_n(x_1, x_2) dx_1 dx_2 \\ &= \left( \mu_{r+s-k,k}^{(n)} \quad \mu_{r+s-k-1,k+1}^{(n)} \quad \mu_{r+s-k-2,k+2}^{(n)} \quad \cdots \quad \mu_{s-k,k+r}^{(n)} \right)^T \\ &= \langle \mathbb{X}_r, x_1^{s-k} x_2^k \rangle_n. \end{aligned} \quad (3.4)$$

Observe that, from the definition, the matrices  $M_{r,s}^{(n)}$  satisfy

$$M_{r,s}^{(n)} = (M_{s,r}^{(n)})^T.$$

Now we define the moment matrix  $\mathcal{M}_n$  of size  $t_n \times t_n$ , constructed by blocks  $M_{r,s}^{(n)}$  for  $0 \leq r, s \leq n$ , in the form

$$\mathcal{M}_n = \begin{pmatrix} M_{0,0}^{(n)} & M_{0,1}^{(n)} & \cdots & M_{0,n}^{(n)} \\ M_{1,0}^{(n)} & M_{1,1}^{(n)} & \cdots & M_{1,n}^{(n)} \\ \vdots & \vdots & & \vdots \\ M_{n,0}^{(n)} & M_{n,1}^{(n)} & \cdots & M_{n,n}^{(n)} \end{pmatrix}, \quad (3.5)$$

where  $t_n$  was defined in (2.1).

Some examples of  $\mathcal{M}_n$ ,

$$\mathcal{M}_0 = (M_{0,0}^{(0)}) = (\mu_{0,0}) \quad \text{of size } t_0 \times t_0 = 1 \times 1,$$

$$\mathcal{M}_1 = \begin{pmatrix} M_{0,0}^{(1)} & M_{0,1}^{(1)} \\ M_{1,0}^{(1)} & M_{1,1}^{(1)} \end{pmatrix} = \left( \begin{array}{c|cc} \mu_{0,0}^{(1)} & \mu_{1,0}^{(1)} & \mu_{0,1}^{(1)} \\ \hline \mu_{1,0}^{(1)} & \mu_{2,0}^{(1)} & \mu_{1,1}^{(1)} \\ \mu_{0,1}^{(1)} & \mu_{1,1}^{(1)} & \mu_{0,2}^{(1)} \end{array} \right)$$

of size  $t_1 \times t_1 = 3 \times 3$ , and  $\mathcal{M}_2$  of size  $t_2 \times t_2 = 6 \times 6$  is

$$\mathcal{M}_2 = \begin{pmatrix} M_{0,0}^{(2)} & M_{0,1}^{(2)} & M_{0,2}^{(2)} \\ M_{1,0}^{(2)} & M_{1,1}^{(2)} & M_{1,2}^{(2)} \\ M_{2,0}^{(2)} & M_{2,1}^{(2)} & M_{2,2}^{(2)} \end{pmatrix} = \left( \begin{array}{c|ccc|ccc} \mu_{0,0}^{(2)} & \mu_{1,0}^{(2)} & \mu_{0,1}^{(2)} & \mu_{2,0}^{(2)} & \mu_{1,1}^{(2)} & \mu_{0,2}^{(2)} \\ \hline \mu_{1,0}^{(2)} & \mu_{2,0}^{(2)} & \mu_{1,1}^{(2)} & \mu_{3,0}^{(2)} & \mu_{2,1}^{(2)} & \mu_{1,2}^{(2)} \\ \mu_{0,1}^{(2)} & \mu_{1,1}^{(2)} & \mu_{0,2}^{(2)} & \mu_{2,1}^{(2)} & \mu_{1,2}^{(2)} & \mu_{0,3}^{(2)} \\ \hline \mu_{2,0}^{(2)} & \mu_{3,0}^{(2)} & \mu_{2,1}^{(2)} & \mu_{4,0}^{(2)} & \mu_{3,1}^{(2)} & \mu_{2,2}^{(2)} \\ \mu_{1,1}^{(2)} & \mu_{2,1}^{(2)} & \mu_{1,2}^{(2)} & \mu_{3,1}^{(2)} & \mu_{2,2}^{(2)} & \mu_{1,3}^{(2)} \\ \mu_{0,2}^{(2)} & \mu_{1,2}^{(2)} & \mu_{0,3}^{(2)} & \mu_{2,2}^{(2)} & \mu_{1,3}^{(2)} & \mu_{0,4}^{(2)} \end{array} \right).$$

From  $\mathcal{M}_n$ , we define the matrix  $\widehat{\mathcal{M}}_n$  of size  $t_{n-1} \times t_{n-1}$  by deleting the last block row and column

$$\widehat{\mathcal{M}}_n = \begin{pmatrix} M_{0,0}^{(n)} & M_{0,1}^{(n)} & \cdots & M_{0,n-1}^{(n)} \\ M_{1,0}^{(n)} & M_{1,1}^{(n)} & \cdots & M_{1,n-1}^{(n)} \\ \vdots & \vdots & & \vdots \\ M_{n-1,0}^{(n)} & M_{n-1,1}^{(n)} & \cdots & M_{n-1,n-1}^{(n)} \end{pmatrix}.$$

We observe that  $\mathcal{M}_n$  is the  $n$ th block moment matrix, and  $\widehat{\mathcal{M}}_n$  in the  $(n-1)$ th block moment matrix associated with the weight function  $W_n(x_1, x_2) = x_1^{-n} x_2^{-n} W(x_1, x_2)$ . Consequently, they are symmetric and positive definite matrices ([5, p. 64]).

Hence

$$\mathcal{M}_n = \left( \begin{array}{cccc|c} & & & & M_{0,n}^{(n)} \\ & & & & M_{1,n}^{(n)} \\ & & & & \vdots \\ & & & & M_{n-1,n}^{(n)} \\ \hline M_{n,0}^{(n)} & M_{n,1}^{(n)} & \cdots & M_{n,n-1}^{(n)} & M_{n,n}^{(n)} \end{array} \right).$$

### 3.2. Orthogonal polynomial systems with respect to varying weights.

Our objective is to construct a PS  $\{\mathbb{P}_n\}_{n \geq 0}$  satisfying orthogonality conditions by using varying weights. Let  $n \geq 0$ , then for  $k = 0, 1, \dots, n-1$ ,

$$\begin{aligned} \langle \mathbb{X}_k, \mathbb{P}_n^T \rangle_n &= \iint_{\Omega} \mathbb{X}_k \mathbb{P}_n^T W_n(x_1, x_2) dx_1 dx_2 \\ &= \iint_{\Omega} \mathbb{X}_k \mathbb{P}_n^T x_1^{-n} x_2^{-n} W(x_1, x_2) dx_1 dx_2 = 0_{(k+1) \times (n+1)}, \end{aligned} \quad (3.6)$$



where  $\mathbf{0}_{(k+1) \times (n+1)}$  denotes a zero matrix of size  $(k+1) \times (n+1)$ , and

$$\langle \mathbb{X}_n, \mathbb{P}_n^T \rangle_n = \Lambda_n, \quad (3.7)$$

where  $\Lambda_n$  is a non-singular real matrix of size  $(n+1) \times (n+1)$ .

**Definition 3.1.** A PS satisfying the varying orthogonality conditions (3.6)-(3.7) will be called a polynomial system orthogonal with respect to the varying weight  $W_n(x_1, x_2)$ , in short VOPS.

**3.3. Existence of VOPS.** From (2.5), we get

$$\mathbb{P}_n^T = \mathbb{X}_0^T G_{n,0}^T + \mathbb{X}_1^T G_{n,1}^T + \cdots + \mathbb{X}_{n-1}^T G_{n,n-1}^T + \mathbb{X}_n^T G_n^T = \sum_{i=0}^n \mathbb{X}_i^T G_{n,i}^T,$$

and, using the orthogonality conditions (3.6) and (3.7), making  $\langle \mathbb{X}_k, \mathbb{P}_n^T \rangle_n$  for  $k = 0, 1, \dots, n-1, n$ , we can write

$$\begin{aligned} \sum_{i=0}^n \langle \mathbb{X}_0, \mathbb{X}_i^T \rangle_n G_{n,i}^T &= \mathbf{0}_{1 \times (n+1)} \\ \sum_{i=0}^n \langle \mathbb{X}_1, \mathbb{X}_i^T \rangle_n G_{n,i}^T &= \mathbf{0}_{2 \times (n+1)} \\ &\vdots \\ \sum_{i=0}^n \langle \mathbb{X}_{n-1}, \mathbb{X}_i^T \rangle_n G_{n,i}^T &= \mathbf{0}_{n \times (n+1)} \\ \sum_{i=0}^n \langle \mathbb{X}_n, \mathbb{X}_i^T \rangle_n G_{n,i}^T &= \Lambda_n. \end{aligned}$$

Using (3.5) we get the matrix system

$$\begin{pmatrix} M_{0,0}^{(n)} & M_{0,1}^{(n)} & \cdots & M_{0,n}^{(n)} \\ M_{1,0}^{(n)} & M_{1,1}^{(n)} & \cdots & M_{1,n}^{(n)} \\ \vdots & \vdots & & \vdots \\ M_{n,0}^{(n)} & M_{n,1}^{(n)} & \cdots & M_{n,n}^{(n)} \end{pmatrix} \begin{pmatrix} G_{n,0}^T \\ G_{n,1}^T \\ \vdots \\ G_n^T \end{pmatrix} = \begin{pmatrix} \mathbf{0}_{1 \times (n+1)} \\ \mathbf{0}_{2 \times (n+1)} \\ \vdots \\ \Lambda_n \end{pmatrix},$$

whose coefficient matrix is  $\mathcal{M}_n$ , the moment matrix. Observe that  $G_{n,k}^T$ , for  $k = 0, 1, \dots, n$  are unknown matrices of size  $(k+1) \times (n+1)$ . Then we have unique solution since the coefficient matrix is non-singular.

Next result shows how to construct the monic VOPS.

**Theorem 3.2.** For  $n \geq 0$  and  $k = 0, 1, \dots, n$ , the entries of every vector  $\mathbb{P}_n$  are given by

$$P_{n-k,k}(x_1, x_2) = \frac{1}{\det(\widehat{\mathcal{M}}_n)} \left| \begin{array}{cccc|c} & & & & \mathbf{m}_{0,n,k}^{(n)} \\ & & & & \mathbf{m}_{1,n,k}^{(n)} \\ & & & & \vdots \\ & & & & \mathbf{m}_{n-1,n,k}^{(n)} \\ \hline \mathbb{X}_0^T & \mathbb{X}_1^T & \dots & \mathbb{X}_{n-1}^T & x_1^{n-k} x_2^k \end{array} \right|,$$

where  $\mathbf{m}_{r,n,k}^{(n)}$  was defined in (3.4).

*Proof.* Obviously,  $P_{n-k,k}(x_1, x_2)$  is a monic polynomial whose highest degree term is  $x_1^{n-k} x_2^k$ . Computing  $\langle x_1^{m-j} x_2^j, P_{n-k,k} \rangle_n$ , for  $m < n$ , and  $0 \leq j \leq m$ , the last row of the the determinant coincides with another row above, therefore the determinant is zero, and the orthogonality holds.  $\square$

#### 4. $R_{II}$ TYPE THREE TERM RELATIONS

We can now prove that the VOPS  $\{\mathbb{P}_n\}_{n \geq 0}$  that satisfy the orthogonality conditions (3.6)-(3.7), also satisfy three term relations.

**Theorem 4.1.** Let  $\{\mathbb{P}_n\}_{n \geq 0}$  be a VOPS associated with the VIP (3.2). For  $n \geq 0$  and  $i = 1, 2$ , there exist matrices  $A_{n,i}$ ,  $B_{n,i}$ ,  $C_{n,i}$ , of respective sizes  $(n+1) \times (n+2)$ ,  $(n+1) \times (n+1)$  and  $(n+1) \times n$ , such that the following  $R_{II}$  type three term relations hold

$$x_i \mathbb{P}_n = A_{n,i} \mathbb{P}_{n+1} + B_{n,i} \mathbb{P}_n + x_1 x_2 C_{n,i} \mathbb{P}_{n-1}, \quad i = 1, 2, \quad n \geq 0, \quad (4.1)$$

with  $\mathbb{P}_{-1} = 0$ ,  $\mathbb{P}_0 = (1)$ , if and only if, for  $n \geq 1$ ,

$$A_{n,i} G_{n+1} = G_n L_{n,i} - C_{n,i} G_{n-1} L_{n-1,1} L_{n,2}, \quad (4.2)$$

$$L_{n-1,1} L_{n,2} \Lambda_{n+1} A_{n,i}^T = L_{n-1,i} \Lambda_n - \Lambda_{n-1} C_{n,i}^T, \quad (4.3)$$

where  $G_n$  is the leading coefficient matrix of  $\mathbb{P}_n$ ,  $L_{n,i}$  and  $\Lambda_n$  are defined in (2.3) and (3.6), respectively.

Moreover, in this case, we have that

$$B_{n,i} \Upsilon_n^T = -C_{n,i} \Upsilon_{n-1}^T, \quad n \geq 0, \quad (4.4)$$

for  $i = 1, 2$ , where

$$\Upsilon_n = \langle \mathbb{X}_{-1}, \mathbb{P}_n^T \rangle_n, \quad n \geq 0.$$

*Proof.* For  $n \geq 0$  and  $i = 1, 2$ , let  $A_{n,i}$  and  $C_{n,i}$  be  $(n+1) \times (n+2)$  and  $(n+1) \times n$  matrices, respectively, satisfying (4.2). We define the  $(n+1) \times 1$  polynomial vector

$$\mathbb{T}_i = A_{n,i} \mathbb{P}_{n+1} - x_i \mathbb{P}_n + x_1 x_2 C_{n,i} \mathbb{P}_{n-1},$$

of degree  $n + 1$ . From the explicit expression (2.5), we can write

$$\mathbb{T}_i = A_{n,i} \sum_{k=0}^{n+1} G_{n+1,k} \mathbb{X}_k - x_i \sum_{k=0}^n G_{n,k} \mathbb{X}_k + x_1 x_2 C_{n,i} \sum_{k=0}^{n-1} G_{n-1,k} \mathbb{X}_k.$$

Using (2.4), we get

$$\begin{aligned} \mathbb{T}_i &= A_{n,i} \sum_{k=0}^{n+1} G_{n+1,k} \mathbb{X}_k - \sum_{k=0}^n G_{n,k} L_{k,i} \mathbb{X}_{k+1} + C_{n,i} \sum_{k=0}^{n-1} G_{n-1,k} L_{k,1} L_{k+1,2} \mathbb{X}_{k+2} \\ &= [A_{n,i} G_{n+1} - G_n L_{n,i} + C_{n,i} G_{n-1} L_{n-1,1} L_{n,2}] \mathbb{X}_{n+1} \\ &\quad + A_{n,i} \sum_{k=0}^n G_{n+1,k} \mathbb{X}_k - \sum_{k=0}^{n-1} G_{n,k} L_{k,i} \mathbb{X}_{k+1} + C_{n,i} \sum_{k=0}^{n-2} G_{n-1,k} L_{k,1} L_{k+1,2} \mathbb{X}_{k+2}. \end{aligned}$$

Then, from (4.2),  $\mathbb{T}_i$  has degree  $n$ . Therefore, since  $\{\mathbb{P}_k\}_{k=0}^n$  is a basis of  $\Pi_n$ , there exist matrices  $F_{k,i}$  of size  $(n+1) \times (k+1)$ ,  $i = 1, 2$ , such that

$$\mathbb{T}_i = \sum_{k=0}^n F_{k,i} \mathbb{P}_k.$$

Now we denote  $F_{n,i} = -B_{n,i}$ , hence

$$\mathbb{T}_i = A_{n,i} \mathbb{P}_{n+1} - x_i \mathbb{P}_n + x_1 x_2 C_{n,i} \mathbb{P}_{n-1} = -B_{n,i} \mathbb{P}_n + \sum_{k=0}^{n-1} F_{k,i} \mathbb{P}_k,$$

and therefore we can write

$$A_{n,i} \mathbb{P}_{n+1} - x_i \mathbb{P}_n + B_{n,i} \mathbb{P}_n + x_1 x_2 C_{n,i} \mathbb{P}_{n-1} = \mathbb{Q}_{n-1,i} \quad (4.5)$$

where

$$\mathbb{Q}_{n-1,i} = \sum_{k=0}^{n-1} F_{k,i} \mathbb{P}_k$$

is a polynomial vector of degree  $n - 1$  and of size  $(n + 1) \times 1$ . Our objective is to prove that  $\mathbb{Q}_{n-1,i}$  vanishes.

Using the transpose of equation (4.5), multiplying it by  $\mathbb{X}_k$ , for  $k = 0, 1, \dots, n - 1$ , and applying the  $n$ th VIP, we get

$$\begin{aligned} \langle \mathbb{X}_k, \mathbb{P}_{n+1}^T \rangle_n A_{n,i}^T - \langle \mathbb{X}_k, x_i \mathbb{P}_n^T \rangle_n + \langle \mathbb{X}_k, \mathbb{P}_n^T \rangle_n B_{n,i}^T \\ + \langle \mathbb{X}_k, x_1 x_2 \mathbb{P}_{n-1}^T \rangle_n C_{n,i}^T = \langle \mathbb{X}_k, \mathbb{Q}_{n-1,i}^T \rangle_n. \end{aligned}$$

Taking into account that  $x_1^{-1} x_2^{-1} W_n = W_{n+1}$ , and  $x_1 x_2 W_n = W_{n-1}$ , we have

$$\begin{aligned} \langle x_1 x_2 \mathbb{X}_k, \mathbb{P}_{n+1}^T \rangle_{n+1} A_{n,i}^T - \langle x_i \mathbb{X}_k, \mathbb{P}_n^T \rangle_n + \langle \mathbb{X}_k, \mathbb{P}_n^T \rangle_n B_{n,i}^T \\ + \langle \mathbb{X}_k, \mathbb{P}_{n-1}^T \rangle_{n-1} C_{n,i}^T = \langle \mathbb{X}_k, \mathbb{Q}_{n-1,i}^T \rangle_n, \end{aligned}$$

and from (2.4) we can write

$$\begin{aligned} L_{k,1} L_{k+1,2} \langle \mathbb{X}_{k+2}, \mathbb{P}_{n+1}^T \rangle_{n+1} A_{n,i}^T - L_{k,i} \langle \mathbb{X}_{k+1}, \mathbb{P}_n^T \rangle_n \\ + \langle \mathbb{X}_k, \mathbb{P}_n^T \rangle_n B_{n,i}^T + \langle \mathbb{X}_k, \mathbb{P}_{n-1}^T \rangle_{n-1} C_{n,i}^T = \langle \mathbb{X}_k, \mathbb{Q}_{n-1,i}^T \rangle_n. \end{aligned} \quad (4.6)$$

Using (3.6), we observe that all terms in the left hand side vanish for  $k = 0, 1, \dots, n-2$ , and then

$$\langle \mathbb{X}_k, \mathbb{Q}_{n-1,i}^T \rangle_n = \mathbf{0}_{(k+1) \times (n+1)}.$$

Substituting  $k = n-1$  in (4.6) and using (3.6) and (3.7), we obtain

$$L_{n-1,1} L_{n,2} \Lambda_{n+1} A_{n,i}^T - L_{n-1,i} \Lambda_n + \Lambda_{n-1} C_{n,i}^T = \langle \mathbb{X}_{n-1}, \mathbb{Q}_{n-1,i}^T \rangle_n.$$

Therefore, (4.3) for  $i = 1, 2$ , provides the square homogeneous linear system of size  $(n+1)t_{n-1}$

$$\langle \mathbb{X}_k, \mathbb{Q}_{n-1,i}^T \rangle_n = \mathbf{0}_{(k+1) \times (n+1)}, \quad k = 0, 1, 2, \dots, n-1. \quad (4.7)$$

Now we look at the matrix of the homogeneous system (4.7). Setting

$$\mathbb{Q}_{n-1,i} = \sum_{r=0}^{n-1} E_{r,i} \mathbb{X}_r \quad \text{and} \quad \mathbb{Q}_{n-1,i}^T = \sum_{r=0}^{n-1} \mathbb{X}_r^T E_{r,i}^T,$$

where  $E_{r,i}$  are constant  $(n+1) \times (r+1)$  matrices, we write (4.7) as

$$\sum_{r=0}^{n-1} \langle \mathbb{X}_k, \mathbb{X}_r^T \rangle_n E_{r,i}^T = \mathbf{0}_{(k+1) \times (n+1)}, \quad k = 0, 1, \dots, n-1.$$

Using the definition of  $M_{r,s}^{(n)}$  given in (3.3), we can write the above linear system of equations as

$$\begin{pmatrix} M_{0,0}^{(n)} & M_{0,1}^{(n)} & \cdots & M_{0,n-1}^{(n)} \\ M_{1,0}^{(n)} & M_{1,1}^{(n)} & \cdots & M_{1,n-1}^{(n)} \\ \vdots & \vdots & & \vdots \\ M_{n-1,0}^{(n)} & M_{n-1,1}^{(n)} & \cdots & M_{n-1,n-1}^{(n)} \end{pmatrix} \begin{pmatrix} E_{0,i}^T \\ E_{1,i}^T \\ \vdots \\ E_{n-1,i}^T \end{pmatrix} = \begin{pmatrix} \mathbf{0}_{n \times (n+1)} \\ \mathbf{0}_{(n-1) \times (n+1)} \\ \vdots \\ \mathbf{0}_{1 \times (n+1)} \end{pmatrix}.$$

The coefficient matrix in the above homogeneous system is  $\widehat{\mathcal{M}}_n$  defined in (3.5), and, since  $\det \widehat{\mathcal{M}}_n > 0$ , the unique solution of the system is zero and  $\mathbb{Q}_{n-1,i} \equiv \mathbf{0}$ . Therefore, the  $R_{II}$  type three term relations (4.1) hold.

The reciprocal can be showed in an analogous way.

Finally, we can take  $k = -1$  in (4.6). Since all integrals are well defined for  $n \geq 0$ , we deduce

$$\langle \mathbb{X}_{-1}, \mathbb{P}_n^T \rangle_n B_{n,i}^T + \langle \mathbb{X}_{-1}, \mathbb{P}_{n-1}^T \rangle_{n-1} C_{n,i}^T = \mathbf{0}_{2 \times (n+1)},$$

and (4.4) holds.  $\square$

**4.1. R<sub>II</sub> type three term recurrence relations.** Observe that the R<sub>II</sub> type three term relations given by (4.1) do not play a rule of recurrence relation, since the matrices  $A_{n,i}$ ,  $i = 1, 2$  are non squared and they can not be invertible. Next result brings R<sub>II</sub> type three term recurrence relations for bivariate polynomials. Given  $M_1, M_2$ , two matrices of the same size, we define the *joint matrix* as

$$M = \begin{pmatrix} M_1 \\ M_2 \end{pmatrix}.$$

**Theorem 4.2.** *Let  $\{\mathbb{P}_n\}_{n \geq 0}$  be a VOPS satisfying the three term relations (4.1). If the matrices  $A_{n,i}$ ,  $i = 1, 2$ , and the joint matrix  $A_n$ , have full rank, then*

$$\mathbb{P}_{n+1} = x_1 D_{n,1}^T \mathbb{P}_n + x_2 D_{n,2}^T \mathbb{P}_n + E_n \mathbb{P}_n + x_1 x_2 F_n \mathbb{P}_{n-1}, \quad n \geq 0, \quad (4.8)$$

with  $\mathbb{P}_{-1} = 0$ ,  $\mathbb{P}_0 = (1)$ , where  $D_n^T = (D_{n,1}^T \ D_{n,2}^T)$ , such that  $D_{n,1}^T A_{n,1} + D_{n,2}^T A_{n,2} = I_{n+2}$ , (it is the pseudo inverse of  $A_n$ ) and

$$E_n = - \sum_{i=1}^2 D_{n,i}^T B_{n,i}, \quad \text{and} \quad F_n = - \sum_{i=1}^2 D_{n,i}^T C_{n,i}.$$

*Proof.* Since the joint matrix  $A_n$  of size  $2(n+1) \times (n+2)$  has full rank, then there exists a (not unique) full rank matrix  $D_n^T = (D_{n,1}^T \ D_{n,2}^T)$  of size  $(n+2) \times 2(n+1)$  such that

$$D_n^T A_n = D_{n,1}^T A_{n,1} + D_{n,2}^T A_{n,2} = I_{n+2}.$$

Therefore, multiplying (4.1) for  $i = 1$  by  $D_{n,1}^T$ , and for  $i = 2$  by  $D_{n,2}^T$  by the left hand side, and sum, we get

$$\sum_{i=1}^2 x_i D_{n,i}^T \mathbb{P}_n = \mathbb{P}_{n+1} + \sum_{i=1}^2 D_{n,i}^T B_{n,i} \mathbb{P}_n + x_1 x_2 \sum_{i=1}^2 D_{n,i}^T C_{n,i} \mathbb{P}_{n-1},$$

and (4.8) holds. □

## 5. A METHOD FOR GENERATING VOPS

Inspired by the well known Koornwinder's method (see Dunkl and Xu [5] and Koornwinder [10]) used to obtain bivariate orthogonal polynomials from univariate orthogonal polynomials, we develop a similar construction to obtain VOPS.

Let us consider two weight functions in one variable,  $w_1(x_1)$  defined in  $(a, b) \subset \mathbb{R}^+$  and  $w_2(x_2)$  in  $(c, d) \subset \mathbb{R}^+$ , and let  $\rho$  be a positive polynomial of degree one in  $(a, b)$ . We can construct a weight function in two variables as

$$W(x_1, x_2) = w_1(x_1) w_2 \left( \frac{x_2}{\rho(x_1)} \right),$$

defined on

$$\Omega = \{(x_1, x_2) : a < x_1 < b, c\rho(x_1) < x_2 < d\rho(x_1)\} \subset \mathbb{R}_+^2.$$

For a fixed  $n$  and a fixed  $k$ , we consider the sequences of monic orthogonal polynomials,  $\{p_m^{(n,k)}\}_{m \geq 0}$  and  $\{q_m^{(n)}\}_{m \geq 0}$ , with respect to the weight functions

$$\frac{\rho(x_1)^{2k+1}w_1(x_1)}{\rho(x_1)^n x_1^n} \quad \text{and} \quad \frac{w_2(x_2)}{x_2^n},$$

respectively. We can write VOPS associated with  $W(x_1, x_2)$

$$\mathbb{P}_n = (P_{n,0}(x_1, x_2), P_{n-1,1}(x_1, x_2), \dots, P_{0,n}(x_1, x_2))^T$$

as

$$P_{n-k,k}(x_1, x_2) = p_{n-k}^{(n,k)}(x_1) \rho(x_1)^k q_k^{(n)}\left(\frac{x_2}{\rho(x_1)}\right).$$

To prove the orthogonality (3.6), let  $m = 0, 1, \dots, n-1$  and  $j \leq m$ , then

$$\begin{aligned} I &= \iint_{\Omega} x_1^{m-j} x_2^j P_{n-k,k}(x_1, x_2) W_n(x_1, x_2) dx_1 dx_2 \\ &= \iint_{\Omega} x_1^{m-j} x_2^j p_{n-k}^{(n,k)}(x_1) \rho(x_1)^k q_k^{(n)}\left(\frac{x_2}{\rho(x_1)}\right) \frac{w_1(x_1)}{x_1^n} \frac{w_2\left(\frac{x_2}{\rho(x_1)}\right)}{x_2^n} dx_1 dx_2, \end{aligned}$$

Setting  $t = x_2/\rho(x_1)$ , since  $c\rho(x_1) < x_2 < d\rho(x_1)$ , then  $c < t < d$  and the integral can be expressed as

$$I = \int_c^d \int_a^b x_1^{m-j} t^j \rho(x_1)^{j+k+1} p_{n-k}^{(n,k)}(x_1) q_k^{(n)}(t) \frac{w_1(x_1)}{\rho(x_1)^n x_1^n} \frac{w_2(t)}{t^n} dx_1 dt,$$

and we can split it into two integrals

$$I = \int_a^b x_1^{m-j} p_{n-k}^{(n,k)}(x_1) \frac{\rho(x_1)^{j+k+1} w_1(x_1)}{\rho(x_1)^n x_1^n} dx_1 \int_c^d t^j q_k^{(n)}(t) \frac{w_2(t)}{t^n} dt.$$

When  $j < k$ , since the second integral in  $I$  vanishes, then  $I = 0$ . In the case  $j = k$ , we get

$$\int_c^d t^k q_k^{(n)}(t) \frac{w_2(t)}{t^n} dt = C_k \neq 0,$$

and we can write

$$I = C_k \int_a^b x_1^{m-k} p_{n-k}^{(n,k)}(x_1) \frac{\rho(x_1)^{2k+1} w_1(x_1)}{\rho(x_1)^n x_1^n} dx_1.$$

Using the orthogonality of  $p_{n-k}^{(n,k)}(x_1)$ , since  $m-k < n-k$ , we conclude that also  $I = 0$ .

When  $j > k$ , we can write

$$I = \int_a^b x_1^{m-j} p_{n-k}^{(n,k)}(x_1) \rho(x_1)^{j-k} \frac{\rho(x_1)^{2k+1} w_1(x_1)}{\rho(x_1)^n x_1^n} dx_1 \int_c^d t^j q_k^{(n)}(t) \frac{w_2(t)}{t^n} dt,$$

we observe that the first integral vanishes, since  $x_1^{m-j} \rho(x_1)^{j-k}$  is a polynomial of degree  $m-k < n-k$ .

6. EXAMPLES

6.1. **VOPS in a rectangle.** We consider the weight function in one variable

$$w(t) = \frac{1}{\sqrt{t-a}\sqrt{b-t}}$$

defined in  $(a, b)$ , where  $0 < a < b < \infty$ . Setting  $\beta = \sqrt{ab}$ , the moments associated with  $w$  satisfy  $\mu_k = \beta^{2k+1}\mu_{-(k+1)}$  (see [16]).

Then, considering two weight functions  $w_1(x_1)$  and  $w_2(x_2)$  defined on positive intervals  $(a, b)$  and  $(c, d)$ , respectively, we can construct the weight function in two variables as

$$W(x_1, x_2) = w_1(x_1)w_2(x_2) = \frac{1}{\sqrt{x_1-a}\sqrt{b-x_1}} \frac{1}{\sqrt{x_2-c}\sqrt{d-x_2}}$$

defined on the rectangular region

$$\Omega = \{(x_1, x_2) : 0 < a < x_1 < b < \infty, 0 < c < x_2 < d < \infty\}.$$

The Figure 6.1 shows the region  $\Omega$ .




FIGURE 6.1. Rectangular region.

Considering  $\beta_1 = \sqrt{ab}$  and  $\beta_2 = \sqrt{cd}$ , then the moments of  $w_1(x_1)$  and  $w_2(x_2)$  satisfy  $\mu_r^{(1)} = \beta_1^{2r+1}\mu_{-(r+1)}^{(1)}$  and  $\mu_s^{(2)} = \beta_2^{2s+1}\mu_{-(s+1)}^{(2)}$ , respectively. Hence, we observe that the moments associated with  $W(x_1, x_2)$  satisfy the identity  $\mu_{r,s} = \beta_1^{2r+1}\beta_2^{2s+1}\mu_{-(r+1),-(s+1)}$ .

For a fixed  $n$ , we consider the sequences of monic orthogonal polynomials,  $\{p_m^{(n)}\}_{m \geq 0}$  and  $\{q_m^{(n)}\}_{m \geq 0}$ , with respect to the weight functions

$$\frac{w_1(x_1)}{x_1^n} \quad \text{and} \quad \frac{w_2(x_2)}{x_2^n},$$

respectively. Hence we can write the monic VOPS associated with  $W(x_1, x_2)$ ,

$$\mathbb{P}_n = (P_{n,0}(x_1, x_2), P_{n-1,1}(x_1, x_2), \dots, P_{0,n}(x_1, x_2))^T,$$

as

$$P_{n-k,k}(x_1, x_2) = p_{n-k}^{(n)}(x_1)q_k^{(n)}(x_2),$$

since

$$\int_c^d \int_a^b x_1^j x_2^h p_{n-k}^{(n)}(x_1) q_k^{(n)}(x_2) \frac{w(x_1)}{x_1^n} \frac{w(x_2)}{x_2^n} dx_1 dx_2 = 0,$$

for  $0 \leq j < n - k$  and  $0 \leq h < k$ .

In order to give some numerical computations, let us choose specific values of the parameter, as  $a = 1/4$ ,  $b = 4$ ,  $c = 4/9$ , and  $d = 9$ . Then the weight function becomes

$$W(x_1, x_2) = \frac{1}{\sqrt{x_1 - \frac{1}{4}\sqrt{4 - x_1}}} \frac{1}{\sqrt{x_2 - \frac{4}{9}\sqrt{9 - x_2}}}, \quad (6.1)$$

defined on  $\Omega = \{(x_1, x_2) : 1/4 < x_1 < 4, 4/9 < x_2 < 9\}$ .

Table 6.1 shows the sequence of monic orthogonal polynomials in one variable  $\{p_m^{(n)}\}_{m \geq 0}$  and  $\{q_m^{(n)}\}_{m \geq 0}$ , with respect to the weight functions  $1/(\sqrt{x_1 - 1/4}\sqrt{4 - x_1}x_1^n)$  and  $1/(\sqrt{x_2 - 4/9}\sqrt{9 - x_2}x_2^n)$ , respectively.

The entries of the first VOPS with respect the weight function (6.1) are given in Table 6.2.

$k$	0	1	2	3
$p_k^{(1)}(x_1)$	1	$x_1 - 1$		
$q_k^{(1)}(x_2)$	1	$x_2 - 2$		
$p_k^{(2)}(x_1)$	1	$x_1 - \frac{8}{17}$	$x_1^2 - \frac{25}{8}x_1 + 1$	
$q_k^{(2)}(x_2)$	1	$x_2 - \frac{72}{85}$	$x_2^2 - \frac{121}{18}x_2 + 4$	
$p_k^{(3)}(x_1)$	1	$x_1 - \frac{272}{803}$	$x_1^2 - \frac{59}{25}x_1 + \frac{16}{25}$	$x_1^3 - \frac{75}{16}x_1^2 + \frac{75}{16}x_1 - 1$
$q_k^{(3)}(x_2)$	1	$x_2 - \frac{4080}{6793}$	$x_2^2 - \frac{582}{121}x_2 + \frac{288}{121}$	$x_2^3 - \frac{121}{12}x_2^2 + \frac{121}{6}x_2 - 8$

TABLE 6.1. Table of the monic orthogonal polynomials  $p_k^{(n)}(x_1)$  and  $q_k^{(n)}(x_2)$ , for  $1 \leq n \leq 3$  and degree  $0 \leq k \leq n$ .

In this case the matrices in (3.7),  $\Lambda_n = \langle \mathbb{X}_n, \mathbb{P}_n^T \rangle_n$  are diagonal, for example for  $n = 3$  we get

$$\Lambda_3 = \langle \mathbb{X}_3, \mathbb{P}_3^T \rangle_3 = \begin{pmatrix} \frac{183411}{524288} & 0 & 0 & 0 \\ 0 & \frac{160083}{2717200} & 0 & 0 \\ 0 & 0 & \frac{60025}{194326} & 0 \\ 0 & 0 & 0 & \frac{94472147}{2985984} \end{pmatrix}.$$



$P_{0,0}$	1
$P_{1,0}$	$x_1 - 1$
$P_{0,1}$	$x_2 - 2$
$P_{2,0}$	$x_1^2 - \frac{25}{8}x_1 + 1$
$P_{1,1}$	$x_1x_2 - \frac{72}{85}x_1 - \frac{8}{17}x_2 + \frac{576}{1445}$
$P_{0,2}$	$x_2^2 - \frac{121}{18}x_2 + 4$
$P_{3,0}$	$x_1^3 - \frac{75}{16}x_1^2 + \frac{75}{16}x_1 - 1$
$P_{2,1}$	$x_1^2x_2 - \frac{4080}{6793}x_1^2 - \frac{59}{25}x_1x_2 + \frac{48144}{33965}x_1 + \frac{16}{25}x_2 - \frac{13056}{33965}$
$P_{1,2}$	$x_1x_2^2 - \frac{582}{121}x_1x_2 - \frac{272}{803}x_2^2 + \frac{288}{121}x_1 + \frac{158304}{97163}x_2 - \frac{78336}{97163}$
$P_{0,3}$	$x_2^3 - \frac{121}{12}x_2^2 + \frac{121}{6}x_2 - 8$

TABLE 6.2. Entries  $P_{n-k,k}$  of the first VOPS with respect the weight function (6.1) for degree  $0 \leq n \leq 3$  and for  $0 \leq k \leq n$ .

6.2. **VOPS in a triangle.** Using the construction described in Section 5, if we choose

$$w_1(x_1) = (x_1 - a)^\alpha(b - x_1)^{\beta+\gamma} \quad \text{and} \quad w_2(x_2) = (x_2 - c)^\beta(d - x_2)^\gamma,$$

defined in  $(a, b)$  and  $(c, d)$ , respectively, and  $\rho(x) = b - x_1$ , then the weight function in two variables takes the form

$$\begin{aligned} W(x_1, x_2) &= (x_1 - a)^\alpha(b - x_1)^{\beta+\gamma} \left( \frac{x_2}{b - x_1} - c \right)^\beta \left( d - \frac{x_2}{b - x_1} \right)^\gamma \\ &= (x_1 - a)^\alpha (x_2 - c(b - x_1))^\beta (d(b - x_1) - x_2)^\gamma, \end{aligned}$$

defined in the triangular region

$$\Omega = \{(x_1, x_2) : a < x_1 < b, c(b - x_1) < x_2 < d(b - x_1)\},$$

as in Figure 6.2.

Choosing  $\alpha = 1, \beta = 2, \gamma = 1, a = 1, b = 2, c = 3$  and  $d = 5$ , the weight functions become

$$\begin{aligned} w_1(x_1) &= (x_1 - 1)(2 - x_1)^3, \quad x_1 \in (1, 2), \\ w_2(x_2) &= (x_2 - 3)^2(5 - x_2), \quad x_2 \in (3, 5), \end{aligned}$$

and

$$W(x_1, x_2) = \tau(x_1 - 1)(x_2 + 3x_1 - 6)^2(10 - 5x_1 - x_2), \quad (6.2)$$




FIGURE 6.2. Triangular region.

defined in  $\Omega = \{(x_1, x_2) : 1 < x_1 < 2, 6 - 3x_1 < x_2 < 10 - 5x_1\}$ . Here  $\tau = 45/2$  is chosen so that  $\mu_{0,0} = 1$ .

Since

$$P_{n-k,k}(x_1, x_2) = p_{n-k,k}^{(n)}(x_1) \rho(x_1)^k q_k^{(n)}\left(\frac{x_2}{\rho(x_1)}\right),$$

for fixed  $n$  and  $k$ , we need to construct the orthogonal polynomials,  $p_{n-k,k}^{(n)}$  and  $q_k^{(n)}$ , with respect to the weight functions

$$\frac{(x_1 - 1)(2 - x_1)^{3+2k+1-n}}{x_1^n} \quad \text{and} \quad \frac{(x_2 - 3)^2(5 - x_2)}{x_2^n},$$

respectively.

Table 6.3 shows the entries of the VOPS with respect the weight function given in (6.2).

In this case the matrix in (3.7) for  $n = 2$ , is given by

$$\Lambda_2 = \begin{pmatrix} 0.000144303270 & 0 & 0 \\ -0.00059443234 & 0.000138244373 & 0 \\ 0.00247304236 & -0.00113123217 & 0.000759945541 \end{pmatrix}.$$

**6.3. VOPS in a shifted simplex.** We consider the family of VOPS on the shifted simplex associated with the weight function

$$W(x_1, x_2) = (x_1 - a)^\alpha (x_2 - a)^\beta (a + b - x_1 - x_2)^\gamma, \quad (6.3)$$

where  $\alpha, \beta, \gamma > -1$ , defined on the region

$$\Omega = \{(x_1, x_2) \in \mathbb{R}^2 : x_1 \geq a, x_2 \geq a, a + b - x_1 - x_2 \geq 0\},$$

with  $0 < a < b < \infty$ . Shifted simplex is represented in Figure 6.3.

$P_{0,0}$	1
$P_{1,0}$	$x_1 - 1.31041371$
$P_{0,1}$	$x_2 + 4.16034291x_1 - 8.32068582$
$P_{2,0}$	$x_1^2 - 2.78649820x_1 + 1.90208670$
$P_{1,1}$	$x_1x_2 + 4.11932686x_1^2 - 13.3840858x_1 - 1.24909536x_2 + 10.2908642$
$P_{0,2}$	$x_2^2 + 8.18284423x_1x_2 + 16.5699978x_1^2 - 66.2799910x_1 - 16.3656885x_2 + 66.2799910$
$P_{3,0}$	$x_1^3 - 4.37745730x_1^2 + 6.28003391x_1 - 2.95184419$
$P_{2,1}$	$x_1^2x_2 + 4.07720565x_1^3 - 18.9879650x_1^2 - 2.65710257x_1x_2 + 28.7424743x_1 + 1.73534712x_2 - 14.1507342$
$P_{1,2}$	$x_1x_2^2 + 8.13057120x_1^2x_2 + 16.3546681x_1^3 - 85.1393062x_1^2 - 26.0650722x_1x_2 - 1.20581071x_2^2 + 144.301208x_1 + 19.6078596x_2 - 78.8825355$
$P_{0,3}$	$x_2^3 + 48.9437310x_1^2x_2 + 12.1645735x_1x_2^2 + 65.1198605x_1^3 - 390.719163x_1^2 - 195.774924x_1x_2 - 24.3291470x_2^2 + 781.4383265x_1 + 195.774924x_2 - 520.958883$

TABLE 6.3. Entries  $P_{n-k,k}$  of the first VOPS with respect to the weight function (6.2) for degree  $0 \leq n \leq 3$  and for  $0 \leq k \leq n$ .




FIGURE 6.3. Shifted simplex.

Choosing  $\alpha = 3, \beta = 2, \gamma = 1, a = 1,$  and  $b = 2,$  the weight function

$$W(x_1, x_2) = (x_1 - 1)^3(x_2 - 1)^2(3 - x_1 - x_2)$$

is defined in  $\Omega = \{(x_1, x_2) \in \mathbb{R}^2 : x_1 \geq 1, x_2 \geq 1, 3 - x_1 - x_2 \geq 0\}.$

Using Theorem 3.2 we can calculate the monic VOPS associated with  $W(x_1, x_2)$  given by (6.3), see Table 6.4.

$P_{0,0}$	1
$P_{1,0}$	$x_1 - 1.43796769$
$P_{0,1}$	$x_2 - 1.32696025$
$P_{2,0}$	$x_1^2 - 2.88799337x_1 + 0.00263767x_2 + 2.05644459$
$P_{1,1}$	$x_1x_2 - 1.26099644x_1 - 1.35144600x_2 + 1.71293494$
$P_{0,2}$	$x_2^2 + 0.00131771x_1 - 2.70824586x_2 + 1.80864659$
$P_{3,0}$	$x_1^3 - 4.34366084x_1^2 + 0.00992183x_1x_2 + 0.00014521x_2^2 + 6.21350091x_1$ $- 0.01365970x_2 - 2.92734391$
$P_{2,1}$	$x_1^2x_2 - 1.21564853x_1^2 - 2.74130572x_1x_2 + 0.00177590x_2^2 + 3.34491332x_1$ $+ 1.85415301x_2 - 2.27334201$
$P_{1,2}$	$x_1x_2^2 + 0.00091564x_1^2 - 2.58880086x_1x_2 - 1.29133163x_2^2 + 1.65554061x_1$ $+ 3.35648957x_2 - 2.15629266$
$P_{0,3}$	$x_2^3 + 0.00006145x_1^2 + 0.00572113x_1x_2 - 4.11853295x_2^2 - 0.00732316x_1$ $+ 5.58718398x_2 - 2.49732589$

TABLE 6.4. Entries  $P_{n-k,k}$  of the first VOPS with respect the weight function (6.3) for degree  $0 \leq n \leq 3$  and for  $0 \leq k \leq n$ .

Also,

$$\Lambda_2 = \langle \mathbb{X}_2, \mathbb{P}_2^T \rangle_2 = \begin{pmatrix} 0.000263986510 & 0.000143544887 & 0.000089632056 \\ 0.000143544887 & 0.000157172682 & 0.000132333688 \\ 0.000089632057 & 0.000132333688 & 0.000214040951 \end{pmatrix}$$

and

$$\Upsilon_2 = \langle \mathbb{X}_{-1}, \mathbb{P}_2^T \rangle_2 = \begin{pmatrix} 0.000090452349 & 0.000049064200 & 0.000030587667 \\ 0.000036048468 & 0.000053277399 & 0.000086301383 \end{pmatrix}.$$

#### ACKNOWLEDGEMENTS

C.F.B. thanks the support by the grant 2022/09575-5 from FAPESP, Sao Paulo, Brazil, and the support by the grant 88887.310463/2018-00 from the program CAPES/PrInt of Brazil.

A.M.D., L.F., and T.E.P. thank grant CEX 2020-001105-M funded by MCIN/AEI/10.13039/501100011033 and Research Group Goya-384, Spain.

T.E.P. also thanks grant 88887.911704/2023-00 from CAPES/PrInt of Brazil.

## REFERENCES

- [1] M. P. Alfaro, M. Bello, J. M. Montaner, J. L. Varona, Some asymptotic properties for orthogonal polynomials with respect to varying measures, *J. Approx. Theory* 135 (2005) 22-34.
- [2] V. S. Buyarov, J. S. Dehesa, A. Martínez-Finkelshtein, E. B. Saff, Asymptotics of the information entropy for Jacobi and Laguerre polynomials with varying weights, *J. Approx. Theory* 99 (1999) 153-166.
- [3] B. de la Calle Ysern, G. López Lagomasino, Strong asymptotics of orthogonal polynomials with varying measures and Hermite-Padé approximants, *J. Comput. Appl. Math.* 99 (1998) 91-103.
- [4] T.S. Chihara, *An Introduction to Orthogonal Polynomials*, Mathematics and Its Applications Series, Gordon and Breach, 1978.
- [5] C. F. Dunkl, Y. Xu, *Orthogonal Polynomials of Several Variables*, 2nd edition, Encyclopedia of Mathematics and its Applications, vol. 155, Cambridge Univ. Press, 2014.
- [6] E. Hendriksen, H. van Rossum, Orthogonal Laurent polynomials, *Indag. Math. (Ser. A)* 48 (1986) 17-36.
- [7] M. E. H. Ismail, D. R. Masson, Generalized orthogonality and continued fractions, *J. Approx. Theory* 83 (1995) 1-40.
- [8] M. E. H. Ismail, A. Sri Ranga, R<sub>II</sub> type recurrence, generalized eigenvalue problem and orthogonal polynomials on the unit circle, *Linear Algebra Appl.* 562 (2019) 63-90.
- [9] W. B. Jones, W. J. Thron, H. Waadeland, A strong Stieltjes moment problem. *Trans. Amer. Math. Soc.* 261 (1980) 503-528.
- [10] T. H. Koornwinder, Two variable analogues of the classical orthogonal polynomials, in Askey R (ed) *Theory and application of special functions*. Academic Press, New York, pp. 435-495, 1975.
- [11] M. A. Kowalski, The recursion formulas for orthogonal polynomials in  $n$  variables, *SIAM J. Math. Anal.* 13 (1982) 309-315.
- [12] M. A. Kowalski, Orthogonality and recursion formulas for polynomials in  $n$  variables, *SIAM J. Math. Anal.* 13 (1982) 316-323.
- [13] G. López Lagomasino, Asymptotics of polynomials orthogonal with respect to varying measures, *Constr. Approx.* 5 (1989) 199-219.
- [14] K. Pan, On orthogonal polynomials with respect to varying measures on the unit circle, *Trans. Amer. Math. Soc.* 346 (1994) 331-340.
- [15] A. P. Silva, A. Sri Ranga, Polynomials generated by a three term recurrence relation: bounds for complex zeros, *Linear Algebra Appl.* 397 (2005) 299-324
- [16] A. Sri Ranga, Another quadrature rule of highest algebraic degree of precision, *Numer. Math.* 68 (1994) 283-294.
- [17] V. Totik, Orthogonal polynomials with respect to varying weights, *J. Comput. Appl. Math.* 99 (1998) 373-385.
- [18] A. Zhedanov, Biorthogonal rational functions and generalized eigenvalue problem, *J. Approx. Theory* 101 (1999) 303-329.

(C. F. Bracciali) DEPARTAMENTO DE MATEMÁTICA, IBILCE, UNESP - UNIVERSIDADE ESTADUAL PAULISTA, CAMPUS AT SÃO JOSÉ DO RIO PRETO, SP (BRAZIL)  
*Email address:* cleonice.bracciali@unesp.br

(A. M. Delgado) INSTITUTO DE MATEMÁTICAS IMAG & DEPARTAMENTO DE MATEMÁTICA APLICADA, FACULTAD DE CIENCIAS. UNIVERSIDAD DE GRANADA (SPAIN)  
*Email address:* amdelgado@ugr.es

(L. Fernández) INSTITUTO DE MATEMÁTICAS IMAG & DEPARTAMENTO DE MATEMÁTICA APLICADA, FACULTAD DE CIENCIAS. UNIVERSIDAD DE GRANADA (SPAIN)

*Email address:* [lidiafr@ugr.es](mailto:lidiafr@ugr.es)

(T. E. Pérez) INSTITUTO DE MATEMÁTICAS IMAG & DEPARTAMENTO DE MATEMÁTICA APLICADA, FACULTAD DE CIENCIAS. UNIVERSIDAD DE GRANADA (SPAIN)

*Email address:* [tperez@ugr.es](mailto:tperez@ugr.es)