

**( $\lambda, \lambda$ )-EIGENFUNCTIONS ON COMPACT MANIFOLDS**

THOMAS JACK MUNN AND OSKAR RIEDLER

ABSTRACT. In this note we study  $(\lambda, \mu)$ -eigenfamilies on compact Riemannian manifolds when  $\lambda = \mu$ . We show that any compact manifold admitting a  $(\lambda, \lambda)$ -eigenfunction is a mapping torus and that any  $(\lambda, \lambda)$ -eigenfamily is one dimensional. Additionally, we consider generalised eigenfamilies, which can have higher dimension, and relate these to harmonic Riemannian submersions to a torus.

## 1. INTRODUCTION

Let  $(M, g)$  be a Riemannian manifold,  $\mathcal{F} = \{\phi_i : M \rightarrow \mathbb{C} \mid i \in I\}$  a family of complex-valued functions, and  $\lambda, \mu \in \mathbb{C}$ . We say that  $\mathcal{F}$  is a  $(\lambda, \mu)$ -*eigenfamily*, if for every  $i, j \in I$  we have that

$$\tau(\phi_i) = \lambda \cdot \phi_i, \quad g(\nabla \phi_i, \nabla \phi_j) = \mu \cdot \phi_i \phi_j,$$

where  $\tau$  is the Laplace-Beltrami operator and  $g$  denotes the complex bilinear extension of the metric to the complexified tangent bundle. Elements of  $(\lambda, \mu)$ -eigenfamilies are called  $(\lambda, \mu)$ -*eigenfunctions*.

Eigenfamilies were introduced by Gudmundsson and Sakovich [7] in order to produce harmonic morphisms. Since then they have been utilised to generate other geometrically interesting data, such as  $r$ -harmonic functions [10], and minimal submanifolds of codimension two in [6] by Gudmundsson and the first named author.

Eigenfamilies have been shown to exist on all the classical Riemannian symmetric spaces, see [7, 8, 9, 3, 4]. The well studied case of  $(\lambda, \mu)$ -eigenfunctions when  $(\lambda, \mu) = (0, 0)$  is that of *harmonic morphisms with codomain  $\mathbb{C}$* . For the general theory of harmonic morphisms between Riemannian manifolds we refer to the excellent book [1] and the regularly updated online bibliography [5].

Recently, there has been interest in classifying eigenfamilies.  $(\lambda, \mu)$ -eigenfamilies on  $m$ -spheres are shown to correspond to  $(0, 0)$ -eigenfamilies of homogeneous polynomials on the ambient space  $\mathbb{R}^{m+1}$  in [12] by the second named author. When  $M$  is *compact* many results describing the spectra, existence, and topological properties of eigenfunctions are derived in [13] by Siffert and the second named author. In particular, they show that eigenfamilies on compact manifolds have  $\lambda \leq \mu < 0$  and that the case  $\lambda = \mu$  is rather singular: the image degenerates to a circle and this case must be excluded from certain constructions.

---

2020 *Mathematics Subject Classification*. 53C35, 53C43, 58E20.

*Key words and phrases*. Eigenfunctions, Eigenfamilies, Harmonic morphisms.

OR gratefully acknowledges the support of the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) - Project-ID 427320536 - SFB 1442, as well as of Germany's Excellence Strategy EXC 2044 390685587, Mathematics Münster: Dynamics-Geometry-Structure. TM gratefully acknowledges the hospitality of Mathematics Münster during his visit.

We begin this note by providing a complete characterisation of  $(\lambda, \lambda)$ -eigenfamilies on compact manifolds. The situation for a single eigenfunction is as follows:

**Theorem 1.1.** *Let  $(M, g)$  be a compact and connected Riemannian manifold,  $\phi : M \rightarrow \mathbb{C}$  a non-constant smooth map,  $\lambda < 0$ . The following are equivalent:*

- (i)  $\phi$  is a  $(\lambda, \lambda)$ -eigenfunction.
- (ii) For any  $x_0 \in M$  the map  $\pi : (M, g) \rightarrow (S^1, \frac{1}{|\lambda|} dt^2)$ ,  $x \mapsto \frac{\phi(x)}{|\phi(x_0)|}$  is a well-defined harmonic Riemannian submersion.

Harmonic Riemannian submersions to  $S^1$  are well understood. The only Riemannian manifolds admitting Riemannian submersions to a circle are mapping tori, i.e. manifolds which may be written as

$$(M, g) \cong \left( \frac{M_0 \times [0, 2\pi]}{(x, 0) \sim (\eta(x), 2\pi)}, g(t) + \frac{1}{|\lambda|} dt^2 \right)$$

where  $M_0$  is compact,  $g(t)$  a Riemannian metric on  $M_0$  for all  $t$ , and  $\eta : M_0 \rightarrow M_0$  is a diffeomorphism with  $\eta^*g(2\pi) = g(0)$ . The Riemannian submersion  $[x, t] \mapsto e^{it}$  is then harmonic if and only if the fibres are minimal, i.e. if and only if the volume density of  $g(t)$  on  $M_0$  is constant in  $t$ .

The above remarks can be considered to be a version of Tischler's Theorem [14] in the Riemannian setting. We formulate and prove this as Theorem 3.1 in Section 3 below.

Beyond this we show that any  $(\lambda, \lambda)$ -eigenfamily on a compact and connected  $(M, g)$  must necessarily be spanned by a single function, whence the previous theorem already gives the complete picture for  $(\lambda, \lambda)$ -eigenfamilies:

**Proposition 1.2.** *Let  $(M, g)$  be a compact and connected Riemannian manifold,  $\lambda \in \mathbb{C}$  and  $\mathcal{F}$  a  $(\lambda, \lambda)$ -eigenfamily on  $M$ . Then  $\dim(\text{span}_{\mathbb{C}}(\mathcal{F})) \leq 1$ .*

In certain situations a generalised notion of a  $(\lambda, \mu)$ -eigenfamily appears, where one allows different values of  $\lambda$  for each function, and the number  $\mu$  is replaced by a matrix. To be precise:

**Definition 1.3.** Let  $\mathcal{F} = \{\phi_1, \dots, \phi_k\}$  be a finite family of functions  $M \rightarrow \mathbb{C}$  and  $\lambda_i, A_{ij} \in \mathbb{C}$  for  $i, j \in \{1, \dots, k\}$ . We say that  $\mathcal{F}$  is  $(\lambda_i, A_{ij})$ -eigenfamily if

$$\tau(\phi_i) = \lambda_i \phi_i, \quad g(\nabla \phi_i, \nabla \phi_j) = A_{ij} \phi_i \phi_j$$

for all  $i, j \in \{1, \dots, k\}$ .

Examples of these families appear on Sasaki manifolds, where one can build an embedding

$$(\phi_1, \dots, \phi_k) : (M, g) \rightarrow \mathbb{C}^k$$

for which  $\{\phi_1, \dots, \phi_k\}$  is a  $(\lambda_i, -\sqrt{\mu_i \mu_j})$ -eigenfamily for appropriate values  $\lambda_i, \mu_i$ . The construction of these families is carried out in Theorem 3 of [13].

**Example 1.4.** Let  $\mathbf{w} = (w_1, \dots, w_n) \in \mathbb{R}_{>0}^n$ , we denote the associated weighted Sasakian sphere by  $S_{\mathbf{w}}^{2n-1}$ . As a Riemannian manifold this is the unit sphere  $S^{2n-1} \subset \mathbb{C}^n$  equipped with the metric

$$g_{\mathbf{w}}(v, v) = \frac{1}{\eta(\xi_{\mathbf{w}})} \left( \|v\|^2 - 2\langle \xi_{\mathbf{w}}, v \rangle \eta_{\mathbf{w}}(\xi_{\mathbf{w}}) \eta_{\mathbf{w}}(v) \right) + \left( 1 + \frac{\|\xi_{\mathbf{w}}\|^2}{\eta(\xi_{\mathbf{w}})} \right) \eta_{\mathbf{w}}(v)^2$$

where  $v \in TS^{2n-1}$  and  $\xi_{\mathbf{w}} = \sum_i w_i(x_i \partial_{y_i} - y_i \partial_{x_i})$ ,  $\eta = \sum_i (x_i dy_i - y_i dx_i)$ ,  $\eta_{\mathbf{w}} = \frac{1}{\eta(\xi_{\mathbf{w}})} \eta$  for  $(x_1 + iy_1, \dots, x_n + iy_n)$  the standard complex coordinates on  $\mathbb{C}^n$ . Here  $\langle \cdot, \cdot \rangle$ ,  $\|\cdot\|$  denote the usual Euclidean scalar product and norm. Then the family  $\{\phi_1 = (x_1 + iy_1), \dots, \phi_n = (x_n + iy_n)\}$  is a  $(\lambda_i, A_{ij})$ -eigenfamily on  $S_{\mathbf{w}}^{2n-1}$  for

$$\lambda_i = -w_i^2 - w_i(2n - 2), \quad A_{ij} = -w_i w_j.$$

These kinds of families can still be used to construct local harmonic morphisms:

**Theorem 1.5.** *Let  $(M, g)$  be a Riemannian manifold,  $\lambda_i \in \mathbb{C}$ ,  $A_{ij} \in \mathbb{C}$  for  $i, j \in \{1, \dots, k\}$ , and  $\{\phi_1, \dots, \phi_k\}$  a  $(\lambda_i, A_{ij})$ -eigenfamily on  $(M, g)$ . For  $U \subseteq \mathbb{C}^k$  open and  $F : U \rightarrow \mathbb{C}$  holomorphic with:*

$$F(t_1 z_1, \dots, t_k z_k) = \left( \prod_i t_i^{d_i} \right) F(z_1, \dots, z_k)$$

for all  $(z_1, \dots, z_k) \in U$  and  $t_i \in \mathbb{R}$  close enough to 1. Then

$$(\phi_1, \dots, \phi_k)^{-1}(U) \rightarrow \mathbb{C}, \quad x \mapsto F(\phi_1(x), \dots, \phi_k(x))$$

is a  $(\tilde{\lambda}, \tilde{\mu})$ -eigenfunction for  $\tilde{\lambda} = \sum_i d_i(\lambda_i - A_{ii}) + \sum_{ij} d_i d_j A_{ij}$ ,  $\tilde{\mu} = \sum_{ij} d_i d_j A_{ij}$ . In particular if

$$\sum_{ij} d_i d_j A_{ij} = 0, \quad \sum_i d_i(\lambda_i - A_{ii}) = 0,$$

then the map is a harmonic morphism.

**Example 1.6.** Let  $F : \mathbb{C}^k \rightarrow \mathbb{C}$ ,  $(z_1, \dots, z_k) \mapsto z_1^{d_1} \cdots z_k^{d_k}$  be a monomial map. Then  $F$  satisfies the conditions of Theorem 1.5, and so  $F \circ (\phi_1, \dots, \phi_k)$  gives a new globally defined eigenfunction for any  $(\lambda_i, A_{ij})$ -eigenfamily on a manifold  $M$ .

In Section 4 we prove Theorem 4.5, which states that a generalised eigenfamily  $\mathcal{F}$  consisting of  $(-A_{ii}, -A_{ij})$ -eigenfunctions on  $M$  is equivalent to the existence of a harmonic Riemannian submersion  $\pi : (M, g) \rightarrow (T^k, A^{-1})$ . After this, we conclude the paper in Section 5 by applying the characterisations of Theorems 1.1 and 4.5 in order to derive existence and non-existence results for such families from curvature conditions.

## 2. EIGENFUNCTIONS AND EIGENFAMILIES

In this section we remind the reader of some important definitions and results that we will use later. Let  $(M, g)$  be an  $m$ -dimensional Riemannian manifold and  $T^{\mathbb{C}}M$  be the complexification of the tangent bundle  $TM$  of  $M$ . We extend the metric  $g$  to a complex bilinear form on  $T^{\mathbb{C}}M$ . Then the gradient  $\nabla\phi$  of a complex-valued function  $\phi : (M, g) \rightarrow \mathbb{C}$  is a section of  $T^{\mathbb{C}}M$ . In this situation, the well-known complex linear *Laplace-Beltrami operator* (alt. *tension field*)  $\tau$  on  $(M, g)$  acts locally on  $\phi$  as follows

$$\tau(\phi) = \operatorname{div}(\nabla\phi) = \sum_{i,j=1}^m \frac{1}{\sqrt{|g|}} \frac{\partial}{\partial x_j} \left( g^{ij} \sqrt{|g|} \frac{\partial \phi}{\partial x_i} \right).$$

For two complex-valued functions  $\phi, \psi : (M, g) \rightarrow \mathbb{C}$  we have the following well-known fundamental relation

$$\tau(\phi \psi) = \tau(\phi) \psi + 2 \kappa(\phi, \psi) + \phi \tau(\psi),$$

where the complex bilinear *conformality operator*  $\kappa$  is given by

$$\kappa(\phi, \psi) = g(\nabla\phi, \nabla\psi).$$

In local coordinates it takes the form:

$$\kappa(\phi, \psi) = \sum_{i,j=1}^m g^{ij} \frac{\partial\phi}{\partial x_i} \frac{\partial\psi}{\partial x_j}.$$

**Definition 2.1.** [7] Let  $(M, g)$  be a Riemannian manifold,  $\lambda, \mu \in \mathbb{C}$ . Then a complex-valued function  $\phi : M \rightarrow \mathbb{C}$  is said to be a  $(\lambda, \mu)$ -*eigenfunction* if it is eigen both with respect to the Laplace-Beltrami operator  $\tau$  and the conformality operator  $\kappa$  with respective eigenvalues  $\lambda, \mu$ , i.e.

$$\tau(\phi) = \lambda \cdot \phi \quad \text{and} \quad \kappa(\phi, \phi) = \mu \cdot \phi^2.$$

A set  $\mathcal{E}$  of complex-valued functions on  $M$  is said to be a  $(\lambda, \mu)$ -*eigenfamily* on  $M$  if for all  $\phi, \psi \in \mathcal{E}$  we have

$$\tau(\phi) = \lambda \cdot \phi \quad \text{and} \quad \kappa(\phi, \psi) = \mu \cdot \phi \psi.$$

Eigenfamilies can be used to produce complex-valued harmonic morphisms:

**Theorem 2.2** ([7]). *Let  $(M, g)$  be a semi-Riemannian manifold,  $\lambda, \mu \in \mathbb{C}$ , and*

$$\mathcal{F} = \{\phi_1, \dots, \phi_n\}$$

*a  $(\lambda, \mu)$ -eigenfamily. If  $P, Q : \mathbb{C}^n \rightarrow \mathbb{C}$  are linearly independent homogeneous polynomials of the same positive degree then the quotient*

$$\frac{P(\phi_1, \dots, \phi_n)}{Q(\phi_1, \dots, \phi_n)}$$

*is a non-constant harmonic morphism on the open and dense subset*

$$\{p \in M \mid Q(\phi_1(p), \dots, \phi_n(p)) \neq 0\}.$$

Away from its zeros, one can write a complex function in polar form, in this case the condition of being an eigenfunction becomes as follows:

**Lemma 2.3** ([13]). *Let  $(U, g)$  be a Riemannian manifold, not necessarily compact or complete, and let  $\phi : U \rightarrow \mathbb{C}$  be a  $(\lambda, \mu)$ -eigenfunction with  $\lambda, \mu$  both real and  $\phi(x) \neq 0$  for all  $x \in U$ . Suppose  $\phi(x) = e^{i\vartheta(x)}|\phi(x)|$  for some smooth function  $\vartheta : U \rightarrow \mathbb{R}$ . Then:*

- (i)  $\tau(\vartheta) = 0$ ;
- (ii)  $\tau(\ln|\phi|) = \lambda - \mu$ ;
- (iii)  $\kappa(\vartheta, |\phi|) = 0$ ;
- (iv)  $\kappa(\ln|\phi|, \ln|\phi|) = \kappa(\vartheta, \vartheta) + \mu$ .

The case of a  $(\lambda, \lambda)$ -eigenfunction is special. One has the following characterisation:

**Proposition 2.4** ([13]). *Let  $(M, g)$  be a compact connected Riemannian manifold and  $\phi : M \rightarrow \mathbb{C}$  a non-constant  $(\lambda, \mu)$ -eigenfunction. The following are equivalent:*

- (i)  $\lambda = \mu$ .
- (ii)  $|\phi|^2$  is constant.
- (iii)  $\phi(x) \neq 0$  for all  $x \in M$ .

### 3. $(\lambda, \lambda)$ -EIGENFUNCTIONS

Theorem 1.1 is a special case of Theorem 4.5, which will be proven in the next section. We begin this section by proving Proposition 1.2, showing that all  $(\lambda, \lambda)$ -eigenfamilies on compact connected manifolds are necessarily one-dimensional:

*Proof of Proposition 1.2.* Suppose that  $\phi, \psi$  are non-constant and in the same  $(\lambda, \lambda)$ -eigenfamily  $\mathcal{F}$ . Then it follows from Theorem 2.2 that the quotient

$$\frac{\phi}{\psi} : M \setminus \psi^{-1}(\{0\}) \rightarrow \mathbb{C}$$

is a harmonic morphism, in particular it is a harmonic map. Since  $\psi(x) \neq 0$  for all  $x \in M$  by Proposition 2.4, the domain of  $\frac{\phi}{\psi}$  is all of  $M$ . By compactness of  $M$  it must then be constant, i.e.  $\psi$  and  $\phi$  are linearly dependent.  $\square$

The following result characterises compact manifolds which admit  $(\lambda, \lambda)$ -eigenfunctions.

**Theorem 3.1.** *Let  $(M, g)$  be compact and connected,  $\lambda < 0$ , and  $\pi : (M, g) \rightarrow (S^1, \frac{1}{|\lambda|} dt^2)$  a smooth map. The following are equivalent:*

- (i) *The map  $\pi$  is a harmonic Riemannian submersion.*
- (ii)  *$M$  is a mapping torus*

$$M_0 \times_{\eta} [0, 2\pi] = \frac{M_0 \times [0, 2\pi]}{(x, 0) \sim (\eta(x), 2\pi)}$$

with metric

$$g = g(t) + \frac{1}{|\lambda|} dt^2$$

and monodromy map  $\eta : M_0 \rightarrow M_0$  with  $\eta^*g(2\pi) = g(0)$ ,  $M_0$  is compact, the volume density of  $g(t)$  is constant in  $t$ , and  $\pi([x, t]) = e^{it}$ .

*Proof.* (i)  $\Rightarrow$  (ii). Since  $\pi : M \rightarrow S^1$  is a surjective submersion from a compact manifold it is a proper map, and thus by Ehresmann's Theorem it is a fibre bundle over  $S^1$ . We let  $M_0$  denote a typical fibre, and  $X = \nabla \hat{\pi}$ , where  $\hat{\pi}$  is the locally defined lift of  $\pi$  with co-domain  $\mathbb{R}$ .

Note that  $X$  is a horizontal vector field, it is divergence free since  $\hat{\pi}$  is harmonic. Additionally from  $\hat{\pi}$  being a Riemannian submersion:

$$d\hat{\pi}(X) = g(X, X) = \frac{d\hat{\pi}(X)^2}{|\lambda|},$$

whence  $d\hat{\pi}(X) = \|X\|^2 = -\lambda$  is constant. Letting  $\eta_t$  denote the flow of  $X$  one then sees that

$$\partial_t(\hat{\pi}\eta_t(x)) = d\hat{\pi}(X) = -\lambda,$$

and so  $\hat{\pi}(\eta_t(x)) = -\lambda \cdot t + \hat{\pi}(x)$ , i.e.  $\pi(\eta_t(x)) = e^{it}\pi(x)$ . It follows that  $M$  is a mapping torus

$$M \cong \frac{M_0 \times [0, 2\pi]}{(x, 0) \sim (\eta(x), 2\pi)}$$

with monodromy map  $\eta = \eta_{2\pi}$ , and  $\pi([x, t]) = e^{it}$ . Under the above diffeomorphism  $X \equiv \partial_t$ , since  $X$  was horizontal one finds that the metric on  $M$  is of the form

$$g = g(t) + \frac{1}{|\lambda|} dt^2$$

where  $\eta^*g(2\pi) = g(0)$  and  $g(t)$  is a family of metrics on the fibre  $M_0$ .

We now show that each  $g(t)$  induces the same volume density. Assume first that  $M$  is oriented, let  $\omega(t)$  denote the volume form of  $g(t)$  so that  $\omega(t) \wedge dt$  is the volume form of  $M$  at  $[x, t]$ . Then

$$(\partial_t \omega(t)) \wedge dt = (\mathcal{L}_X \omega(t)) \wedge dt = \mathcal{L}_X(\omega(t) \wedge dt) = \operatorname{div}(X) \omega(t) \wedge dt = 0,$$

where we used the formula  $\mathcal{L}_X d\operatorname{Vol} = \operatorname{div}(X)$  and  $\operatorname{div}(X) = 0$ . Since the above argument is entirely local it also shows that the volume density induced by  $g(t)$  is constant in  $t$  for the non-orientable case.

(ii)  $\Rightarrow$  (i). Note that the map

$$\hat{\pi} : (M_0 \times \mathbb{R}, g(t) + dt^2) \rightarrow (\mathbb{R}, \cdot), \quad (x, t) \mapsto t$$

is clearly a Riemannian submersion. We now verify that it is harmonic. Let  $(x_0, t_0) \in M_0 \times \mathbb{R}$  and let  $x_1, \dots, x_m$  be a coordinate system of  $M_0$  so that  $\partial_{x_1}, \dots, \partial_{x_m}$  is an orthonormal basis of  $T_{x_0} M_0$  with respect to  $g(t_0)$ . We denote  $g_{ij}(t) = g(t)(\partial_{x_i}, \partial_{x_j})$ , note that  $g(t_0)_{ij} = \delta_{ij}$ . One checks that  $\nabla \hat{\pi} = \partial_t$ , from which it follows that

$$\begin{aligned} \Delta \hat{\pi} &= \sum_{i=1}^m g(t_0) (\nabla_{\partial_{x_i}} \partial_t, \partial_{x_i}) = \sum_{i=0}^m g(t_0) (\nabla_{\partial_t} \partial_{x_i}, \partial_{x_i}) = \frac{1}{2} \frac{d}{dt} \sum_{i=0}^m g_{ii}(t)|_{t=t_0} \\ &= \frac{1}{2} \frac{d}{dt} \operatorname{trace}((g_{ij})_{1 \leq i, j \leq m})|_{t=0} = \frac{1}{2} \frac{d}{dt} \ln(\det((g_{ij}(t))_{1 \leq i, j \leq m}))|_{t=t_0}. \end{aligned}$$

This last expression is 0, since the volume form  $\sqrt{\det((g_{ij})_{1 \leq i, j \leq m})} dx_1 \wedge \dots \wedge dx_m$  is constant in the variable  $t$  by assumption, and so  $\hat{\pi}$  is harmonic.

Now the diagram

$$\begin{array}{ccc} (M_0 \times \mathbb{R}, g(t) + dt^2) & \xrightarrow{\hat{\pi}} & (\mathbb{R}, \cdot) \\ \downarrow / \mathbb{Z} & & \downarrow / \mathbb{Z} \\ (M_0 \times_{\eta} I, g(t) + dt^2) & \xrightarrow{\pi} & (S^1, dt^2) \end{array}$$

commutes by construction, here the vertical arrows are the natural covering maps. Since the vertical maps are also local isometries it follows that  $\pi$  is a harmonic Riemannian submersion.  $\square$

#### 4. GENERALISED EIGENFAMILIES

The previous section shows that the notion of a  $(\lambda, \mu)$ -eigenfamily becomes degenerate in the case  $\lambda = \mu$ , since these are then always spanned by a single function. By relaxing some of the conditions on  $(\lambda, \mu)$ -eigenfamilies the  $(\lambda, \lambda)$ -eigenfunctions can be part of a family:

**Definition 4.1.** Let  $\mathcal{F} = \{\phi_1, \dots, \phi_k\}$  be a finite family of functions  $M \rightarrow \mathbb{C}$  and  $\lambda_i \in \mathbb{C}$ ,  $1 \leq i \leq k$  a vector in  $\mathbb{C}^k$  and  $A_{ij}$ ,  $1 \leq i, j \leq k$  a symmetric complex  $k \times k$  matrix.

(i) We call  $\mathcal{F}$  an  $(\lambda_i, A_{ij})$ -*eigenfamily* if for all  $\phi_i, \phi_j \in \mathcal{F}$ :

$$\kappa(\phi_i, \phi_j) = A_{ij}\phi_i\phi_j, \quad \tau(\phi_i) = \lambda_i\phi_i.$$

- (ii) We say that the family is  $\lambda$ -*diagonal* if additionally  $\lambda_i = A_{ii}$  for all  $i$ .  
 (iii)  $\mathcal{F}$  is said to be *reduced* if  $A_{ij}$  is a non-degenerate matrix.

The following is the prototypical example of a  $\lambda$ -diagonal  $(\lambda_i, A_{ij})$ -eigenfamily.

**Example 4.2** ([13]). Let  $\Gamma$  be a lattice in  $\mathbb{R}^n$  and  $M = \mathbb{R}^n/\Gamma$  be the corresponding flat torus. Then consider the dual lattice

$$\Gamma^* = \{k \in \mathbb{R}^n \mid \langle k, k' \rangle \in \mathbb{Z}, \text{ for all } k' \in \Gamma\}.$$

For any subset  $K \subset \Gamma^*$  we define the family

$$\mathcal{F} = \{f_k(x) = e^{2\pi i \langle k, x \rangle} \mid k \in K\}.$$

A direct computation shows

$$\tau(f_k) = -4\pi^2 \langle k, k \rangle \cdot f_k, \quad \kappa(f_k, f_{k'}) = -4\pi^2 \langle k, k' \rangle \cdot f_k f_{k'}$$

and thus  $\mathcal{F}$  is a  $\lambda$ -diagonal eigenfamily. Furthermore the set  $K$  is linearly independent if and only if  $\mathcal{F}$  is reduced.

*Proof of Theorem 1.5.* Let  $F : U \rightarrow \mathbb{C}$  as in the statement of the theorem. The homogeneity condition means that  $z_i \cdot \partial_{z_i} F(z_1, \dots, z_k) = d_i F$  for all  $i$ . Writing  $\wedge$  to denote precomposition with  $(\phi_1, \dots, \phi_k)$  one then finds:

$$\kappa(\hat{F}, \hat{F}) = \sum_{ij} (\partial_{z_i} F)^\wedge (\partial_{z_j} F)^\wedge \kappa(\phi_i, \phi_j) = \sum_{ij} A_{ij} (z_i \partial_{z_i} F \ z_j \partial_{z_j} F)^\wedge = \sum_{ij} A_{ij} d_i d_j (\hat{F})^2.$$

Similarly:

$$\Delta \hat{F} = \sum_i (\partial_{z_i} F)^\wedge \Delta \phi_i + \sum_{ij} (\partial_{z_i} \partial_{z_j} F)^\wedge \cdot \kappa(\phi_i, \phi_j) = \sum_i d_i (\lambda_i - A_{ii}) \hat{F} + \sum_{ij} A_{ij} d_i d_j \hat{F}.$$

Completing the proof. □

We now turn to  $\lambda$ -diagonal eigenfamilies:

**Proposition 4.3.** *Let  $(M, g)$  be compact and connected,  $A \in M_{k \times k}(\mathbb{C})$ , and  $\mathcal{F} = \{\phi_1, \dots, \phi_k\}$  a  $(A_{ii}, A_{ij})$ -eigenfamily on  $M$ , in particular  $\mathcal{F}$  is  $\lambda$ -diagonal. Then:*

- (i) *A is a negative semi-definite real matrix.*  
 (ii)  *$\mathcal{F}$  is not reduced if and only if there exist  $\alpha_i \in \mathbb{R}$  not all vanishing so that  $\prod_{i=1}^k (\phi_i)^{\alpha_i}$  is constant.*

*Proof.* Since  $\mathcal{F}$  is  $\lambda$ -diagonal  $|\phi_i(x)|$  is constant on  $M$  for all  $i$ . Then the locally defined functions  $\ln \phi_i$  are purely imaginary and the vector fields  $\nabla \vartheta_i := \nabla \text{Im}(\ln \phi_i) = \frac{\nabla \phi_i}{i \phi_i}$  satisfy:

$$g(\nabla \vartheta_i, \nabla \vartheta_j) = -\frac{A_{ij} \phi_i \phi_j}{\phi_i \phi_j} = -A_{ij},$$

whence  $A_{ij}$  is negative semi-definite. Further it is degenerate if and only if the  $\nabla\vartheta_i$  are linearly dependent, i.e. iff  $\sum_i \alpha_i \nabla\vartheta_i = 0$  for some  $\alpha_i \in \mathbb{R}$  not all vanishing. This is equivalent to

$$0 = \sum_i \alpha_i \nabla \ln(\phi_i) = \nabla \ln \prod_i (\phi_i)^{\alpha_i},$$

which completes the proof.  $\square$

**Definition 4.4.** For  $A$  a positive definite  $k \times k$  matrix, let  $(T^k, A^{-1})$  denote the flat torus  $T^k = (S^1)^k$  equipped with metric  $A^{-1}$ .

**Theorem 4.5.** Let  $(M, g)$  be a compact Riemannian manifold,  $A$  a symmetric real  $k \times k$  matrix, and  $\mathcal{F} = \{\phi_1, \dots, \phi_k\}$  a family of functions  $M \rightarrow \mathbb{C}$ . The following are equivalent:

- (i)  $\mathcal{F}$  is a reduced  $(-A_{ii}, -A_{ij})$ -eigenfamily.
- (ii)  $A$  is positive definite and for any  $x_0 \in M$  the map

$$\pi : (M, g) \rightarrow (T^k, A^{-1}), \quad x \mapsto \left( \frac{\phi_1(x)}{|\phi_1(x_0)|}, \dots, \frac{\phi_k(x)}{|\phi_k(x_0)|} \right)$$

is a well-defined and harmonic Riemannian submersion.

*Proof.* (i)  $\implies$  (ii): Since  $\phi_i$  is a  $(-A_{ii}, -A_{ij})$ -eigenfunction for all  $i$  one has by Proposition 2.4 that  $|\phi_i(x)|$  is constant and non-zero for all  $i$ , and so  $\pi : M \rightarrow T^k, x \mapsto \left( \frac{\phi_1(x)}{|\phi_1(x_0)|}, \dots, \frac{\phi_k(x)}{|\phi_k(x_0)|} \right)$  is well defined.

Additionally by Lemma 2.3 one finds around each point in  $M$  a neighbourhood  $U$  and maps  $\vartheta_i : U \rightarrow \mathbb{R}$  so that

$$\left( \frac{\phi_1(x)}{|\phi_1(x_0)|}, \dots, \frac{\phi_k(x)}{|\phi_k(x_0)|} \right) = \left( e^{i\vartheta_1(x)}, \dots, e^{i\vartheta_k(x)} \right).$$

One verifies:

$$g(\nabla\vartheta_i, \nabla\vartheta_j) = -g(\nabla \ln \phi_i, \nabla \ln \phi_j) = A_{ij}, \quad \tau(\vartheta_i) = 0. \quad (4.1)$$

Since  $\mathcal{F}$  is reduced Proposition 4.3 implies that  $A$  is a positive definite matrix. Equation (4.1) then implies that the map  $\tilde{\pi} : (U, g) \rightarrow (\mathbb{R}^k, A^{-1}), x \mapsto (\vartheta_1(x), \dots, \vartheta_k(x))$  is a harmonic Riemannian submersion to its image.

Letting  $L : (\mathbb{R}^k, A^{-1}) \rightarrow (T^k, A^{-1}), (t_1, \dots, t_k) \mapsto (e^{it_1}, \dots, e^{it_k})$  denote the standard isometric covering map, one has by construction that  $\pi|_U = L \circ \tilde{\pi}$ , whence  $\pi$  is a harmonic Riemannian submersion.

(ii)  $\implies$  (i): Since we assume that  $\pi : M \rightarrow T^k, x \mapsto \left( \frac{\phi_1(x)}{|\phi_1(x_0)|}, \dots, \frac{\phi_k(x)}{|\phi_k(x_0)|} \right)$  is well defined it follows that  $|\phi_i(x)|$  is constant non-zero for all  $i$ . We then find locally defined functions  $\vartheta_i := \text{Im}(\ln \phi_i)$  so that

$$\pi(x) = (e^{i\vartheta_1(x)}, \dots, e^{i\vartheta_k(x)}) = (L \circ \tilde{\pi})(x).$$

Here we let  $L : (\mathbb{R}^k, A^{-1}) \rightarrow (T^k, A^{-1}), (t_1, \dots, t_k) \mapsto (e^{it_1}, \dots, e^{it_k})$  denote the standard isometric covering map. The map  $\tilde{\pi}(x) = (\vartheta_1(x), \dots, \vartheta_k(x))$  can be defined locally around any point.

Note that:

$$D\tilde{\pi}(\nabla\vartheta_i) = \sum_j d\vartheta_j(\nabla\vartheta_i) \partial_{t_j} = \sum_j g(\nabla\vartheta_i, \nabla\vartheta_j) \partial_{t_j}.$$



Since  $\pi$  is Riemannian submersion so is  $\tilde{\pi}$ , and it follows that:

$$g(\nabla\vartheta_i, \nabla\vartheta_j) = A^{-1}(D\tilde{\pi}\nabla\vartheta_i, D\tilde{\pi}\nabla\vartheta_j) = \sum_{k,\ell} g(\nabla\vartheta_i, \nabla\vartheta_k) A_{k\ell}^{-1} g(\nabla\vartheta_\ell, \nabla\vartheta_j),$$

which implies

$$g(\nabla\phi_i, \phi_j) = -\phi_i\phi_j g(\nabla\vartheta_i, \nabla\vartheta_j) = -A_{ij}\phi_i\phi_j.$$

Additionally  $\tilde{\pi}$  is harmonic since  $\pi$  is, and so

$$\tau(\phi_i) = -\phi_i g(\nabla\vartheta_i, \nabla\vartheta_i) + i\phi_i\tau(\vartheta_i) = -A_{ii}\phi_i.$$

It follows that  $\mathcal{F}$  is a  $(-A_{ii}, -A_{ij})$ -eigenfamily, since  $A$  is positive definite it is reduced by Proposition 4.3.  $\square$

**Remark 4.6.** Theorem 4.5 shows that any compact manifold  $(M, g)$  admitting a reduced  $\lambda$ -diagonal  $(-A_{ii}, -A_{ij})$ -eigenfamily is a fibre bundle over the torus  $T^k$ . If, additionally, the horizontal distribution is integrable then one can proceed as in Theorem 3.1 to show that  $M$  is a quotient of

$$M_0 \times [0, 2\pi]^k$$

with metric  $g(t_1, \dots, t_{n-k}) + A^{-1}$  such that  $g(t_1, \dots, t_{n-k})$  induces the a constant volume density on  $M_0$ .

## 5. EXISTENCE AND NON-EXISTENCE FROM CURVATURE

In this section we apply our characterisations of  $(\lambda, \lambda)$ -eigenfunctions and  $\lambda$ -diagonal eigenfunctions by harmonic Riemannian submersions in order to derive existence and non-existence results.

We begin by remarking that harmonic maps  $\pi : M \rightarrow S^1$  correspond, up to rotations of  $S^1$ , with harmonic 1-forms on  $M$  with integral periods (see e.g. [1], [11]). The condition that the resulting map is submersive is equivalent to the form being *nowhere vanishing*. Theorems 1.1 and 4.5 then imply:

**Corollary 5.1.** *Let  $(M, g)$  be a compact and connected Riemannian manifold.*

- (i). *Suppose  $\text{Ric} \geq 0$  and first Betti number  $\beta_1(M) = k$ . Then there exists a reduced  $\lambda$ -diagonal eigenfamily  $\mathcal{F} = \{\phi_1, \dots, \phi_k\}$ .*
- (ii). *Suppose the sectional curvature of  $(M, g)$  is strictly negative, then  $(M, g)$  does not admit any non-constant  $(\lambda, \lambda)$ -eigenfunctions for any value of  $\lambda$ .*

*Proof.* We prove (i) first. Since  $M$  is compact and  $\beta_1(M) = k$ , there exist linearly independent integral harmonic 1-forms  $\omega_1, \dots, \omega_k$ . Let  $\phi_1, \dots, \phi_k$  denote the corresponding harmonic maps into  $S^1$ .

Since  $\text{Ric} \geq 0$  and  $M$  is compact, we have that the  $\omega_i$  are parallel by Bochner [2] and thus

$$g(\omega_i, \omega_j) = g(d\phi_i, d\phi_j)$$

is constant and so the resulting map  $(\phi_1, \dots, \phi_k)$  is a harmonic Riemannian submersion to a torus.

Point (ii) follows from the fact, due to Tsagas [15], that every closed 1-form on a compact manifold with strictly negative curvature vanishes at at least one point.  $\square$

## REFERENCES

- [1] P. Baird and J. C. Wood, *Harmonic morphisms between Riemannian manifolds*, London Math. Soc. Monogr. No. **29**, Oxford Univ. Press (2003).
- [2] S. Bochner, *Curvature and Betti Numbers*, Ann. of Math. **49**, 379-390 (1948), <https://doi.org/10.2307/1969287>.
- [3] E. Ghandour, S. Gudmundsson, *Explicit  $p$ -harmonic functions and harmonic morphisms on the real Grassmannians*, Adv. Geom. (2023).  
<https://doi.org/10.1515/advgeom-2023-0015>.
- [4] E. Ghandour, S. Gudmundsson, *Explicit harmonic morphisms and  $p$ -harmonic functions from the complex and quaternionic Grassmannians*, Ann. Global Anal. Geom. **64** (2023), 15.
- [5] S. Gudmundsson, *The Bibliography of Harmonic Morphisms*, [www.matematik.lu.se/matematiklu/personal/sigma/harmonic/bibliography.html](http://www.matematik.lu.se/matematiklu/personal/sigma/harmonic/bibliography.html).
- [6] S. Gudmundsson, T. J. Munn, *Minimal submanifolds via complex-valued eigenfunctions*, J. Geom. Anal. **34** (2024), 190.
- [7] S. Gudmundsson, A. Sakovich, *Harmonic morphisms from the classical compact semisimple Lie groups*, Ann. Global Anal. Geom. **33** (2008), 343-356.
- [8] S. Gudmundsson, A. Sakovich, *Harmonic morphisms from the classical non-compact semisimple Lie groups*, Differential Geom. Appl. **27** (2009), 47-63.
- [9] S. Gudmundsson, A. Siffert, M. Sobak, *Explicit proper  $p$ -harmonic functions on the Riemannian symmetric spaces  $\mathbf{SU}(n)/\mathbf{SO}(n)$ ,  $\mathbf{Sp}(n)/\mathbf{U}(n)$ ,  $\mathbf{SO}(2n)/\mathbf{U}(n)$ ,  $\mathbf{SU}(2n)/\mathbf{Sp}(n)$* , J. Geom. Anal. **32**, 147 (2022).
- [10] S. Gudmundsson, M. Sobak, *Proper  $r$ -harmonic functions from Riemannian manifolds*, Ann. Global Anal. Geom. **57** (2020), 217-223.
- [11] F. Hélein, J. C. Wood, *Harmonic maps*, Handbook of global analysis, 417-491 (2008).
- [12] O. Riedler, *Polynomial harmonic morphisms and eigenfamilies on spheres*, arXiv preprint arXiv:2310.19565 (2023).
- [13] O. Riedler, A. Siffert, *Global eigenfamilies on compact manifolds*, arXiv preprint arXiv:2401.17750 (2024).
- [14] D. Tischler, *On fibering certain foliated manifolds over  $S^{1*}$* , Topology **9**, 153-154, Springer (1970).
- [15] Gr. Tsagas, *Properties of closed 1-forms on special Riemannian manifold*, Proc. Amer. Math. Soc. **81**, 104-106 (1981).

MATHEMATICS, FACULTY OF SCIENCE, UNIVERSITY OF LUND, BOX 118, LUND 221 00, SWEDEN  
*Email address:* `Thomas.Munn@math.lu.se`

UNIVERSITÄT MÜNSTER, MATHEMATISCHES INSTITUT, EINSTEINSTR. 62, 48149 MÜNSTER, GERMANY  
*Email address:* `oskar.riedler@uni-muenster.de`