

Lie–Hamilton systems associated with the symplectic Lie algebra $\mathfrak{sp}(6, \mathbb{R})$

OSCAR CARBALLAL^{†1}, RUTWIG CAMPOAMOR-STURSBURG^{†,*} AND FRANCISCO J. HERRANZ[‡]

[†] Departamento de Álgebra, Geometría y Topología, Facultad de Ciencias Matemáticas, Universidad Complutense de Madrid, Plaza de Ciencias 3, E-28040 Madrid, Spain

^{*} Instituto de Matemática Interdisciplinar, Universidad Complutense de Madrid, E-28040 Madrid, Spain

[‡] Departamento de Física, Universidad de Burgos, E-09001 Burgos, Spain

E-mail: oscarbal@ucm.es, rutwig@ucm.es, fjherranz@ubu.es

Abstract

New classes of Lie–Hamilton systems are obtained from the six-dimensional fundamental representation of the symplectic Lie algebra $\mathfrak{sp}(6, \mathbb{R})$. The ansatz is based on a recently proposed procedure for constructing higher-dimensional Lie–Hamilton systems through the representation theory of Lie algebras. As applications of the procedure, we study a time-dependent electromagnetic field and several types of coupled oscillators. The irreducible embedding of the special unitary Lie algebra $\mathfrak{su}(3)$ into $\mathfrak{sp}(6, \mathbb{R})$ is also considered, yielding Lie–Hamilton systems arising from the sum of the quark and antiquark three-dimensional representations of $\mathfrak{su}(3)$, which are applied in the construction of t -dependent coupled systems. In addition, t -independent constants of the motion are obtained explicitly for all these Lie–Hamilton systems, which allows the derivation of a nonlinear superposition rule

Keywords: constants of the motion, coupled oscillators, Lie systems, Minkowski, nonlinear differential equations, nonlinear superposition rules, special unitary Lie algebra, symplectic Lie algebra

MSC 2020 codes: 17B10, 34A26, 34C14, 58A30

¹Based on the contribution presented at the “XXIVth International Conference on Geometry, Integrability and Quantization” held in Varna, Bulgaria, June 6–13, 2024

Contents

1	Introduction	2
2	Basics on Lie and Lie–Hamilton systems	3
3	The symplectic Lie algebra $\mathfrak{sp}(6, \mathbb{R})$	4
3.1	Constants of the motion and superposition rule	7
3.2	Applications	8
3.2.1	A time-dependent electromagnetic field	8
3.2.2	Time-dependent coupled oscillators	9
4	The special unitary Lie algebra $\mathfrak{su}(3)$	11
4.1	Applications to coupled systems	14
5	Final remarks	16
	Acknowledgements	16
	References	16

1 Introduction

Group theory has been shown to be a very effective method within the theory of differential equations, helping to identify symmetry schemes and linearization properties, as well as effective criteria to derive (nonlinear) superposition rules [3, 17, 21]. The combination of Lie groups with additional geometric structures, on the other hand, has enlarged the number of techniques to analyze the existence and computation of superposition rules for differential equations, hence enlarging the original approach of S. Lie [13]. In this context, the theory of Lie–Hamilton systems (LH in short) has been shown to be an effective approach, with wide applications to both classical and quantum systems (see e.g. [15] and references therein). In this context, genuinely indecomposable LH systems in higher dimension, i.e., systems that cannot be decoupled into systems in lower dimension, have recently been introduced and analyzed in connection with the representation theory of (semisimple) Lie algebras [6], leading to the construction of new intrinsic hierarchies of higher-dimensional LH systems. In this paper, we generalize the construction done in [6] for the symplectic Lie algebra $\mathfrak{sp}(4, \mathbb{R})$ to the rank-three case, i.e., the Lie algebra $\mathfrak{sp}(6, \mathbb{R})$, hence deriving new three-dimensional LH systems related to coupled oscillators.

This paper is structured as follows. After presenting some generalities on Lie and LH systems in Section 2, we extend in Section 3 the formalism developed in [6] based on the Lie algebra $\mathfrak{sp}(4, \mathbb{R})$ to the case of the Lie algebra $\mathfrak{sp}(6, \mathbb{R})$, leading to novel LH systems on $\mathbb{T}^*\mathbb{R}^3$. Furthermore, in Subsection 3.1 their corresponding t -independent constants of the motion are obtained in an explicit form following the coalgebra formalism introduced in [1, 2]. These can then be used to deduce a nonlinear superposition rule which is also indicated. As applications of such

$\mathfrak{sp}(6, \mathbb{R})$ -LH systems, we construct a t -dependent electromagnetic field and several coupled oscillators in Subsection 3.2, which generalize the systems obtained in [6]. In Section 4 we study LH systems based on the regular subalgebra $\mathfrak{su}(3) \subset \mathfrak{sp}(6, \mathbb{R})$, which allow the construction of coupled systems presented in Subsection 4.1. In particular, we obtain an $\mathfrak{su}(3)$ -LH system that can be interpreted as a coupling of two LH systems, each of them being a coupled oscillator on the Minkowski plane $\mathbb{T}^*\mathbf{M}^{1+1}$. The paper finishes with some concluding remarks and future work perspectives.

2 Basics on Lie and Lie–Hamilton systems

The solutions of a non-autonomous first-order system of ordinary differential equations (ODEs) on a smooth manifold M in normal form

$$\frac{dx^j}{dt} = \psi^j(t, \mathbf{x}), \quad \mathbf{x} \in M, \quad 1 \leq j \leq n = \dim M \quad (1)$$

are given by the integral curves of a t -dependent vector field $\mathbf{X} : \mathbb{R} \times M \rightarrow \mathbb{T}M$, the local expression of which is given by

$$\mathbf{X}(t, x) := \psi^j(t, \mathbf{x}) \frac{\partial}{\partial x^j}.$$

This allows us to univocally identify the system (1) with the t -dependent vector field \mathbf{X} . We say that (1) is a *Lie system* if its general solution $\mathbf{x}(t)$ can be written in terms of a t -independent function Ψ , a *superposition rule*, as

$$\mathbf{x}(t) = \Psi(\mathbf{x}_1(1), \dots, \mathbf{x}_r(t); k_1, \dots, k_n)$$

where $\mathbf{x}_1(t), \dots, \mathbf{x}_r(t)$ denotes a generic finite family of particular solutions and $(k_1, \dots, k_n) \in M$ a certain point related to the initial conditions (for general aspects of the theory of Lie systems, see e.g. [8, 15] and references therein).

The fundamental Lie–Scheffers Theorem states that a system (1) is a Lie system if and only if the t -dependent vector field \mathbf{X} can be decomposed as

$$\mathbf{X}(t, \mathbf{x}) = \sum_{i=1}^{\ell} b_i(t) \mathbf{X}_i(x)$$

for some functions $b_1(t), \dots, b_{\ell}(t)$ and vector fields $\mathbf{X}_1, \dots, \mathbf{X}_{\ell}$ on M spanning an ℓ -dimensional real Lie algebra V^X of vector fields, the so-called *Vessiot–Guldberg (VG) Lie algebra* [13]. This allows to regard a Lie system as a curve taking values values in a finite-dimensional Lie algebra V^X of vector fields.

One of the main features of Lie systems is that they can be made compatible with a lot of supplementary geometric structures (Jacobi, Poisson, symplectic, multi-symplectic, Dirac, contact, etc.), a fact that provides additional properties of notable interest for both the quantitative and qualitative analysis of the associated systems of ODEs and, in some cases, facilitates the deduction of an explicit superposition rule.

For the particular case of symplectic geometry, a Lie system \mathbf{X} on a symplectic manifold (M, ω) is called a *LH system* if it admits a VG Lie algebra V^X formed by Hamiltonian vector fields with respect to the symplectic form ω . Moreover, if $\mathbf{X}_1, \dots, \mathbf{X}_\ell$ span V^X , their associated Hamiltonian functions $h_1, \dots, h_\ell \in C^\infty(M)$, determined by the inner product condition

$$\iota_{\mathbf{X}_i} \omega = dh_i, \quad 1 \leq i \leq \ell, \quad (2)$$

span a finite-dimensional (functional) real Lie algebra \mathcal{H}_ω with respect to the Poisson bracket $\{\cdot, \cdot\}_\omega$ induced by the symplectic form ω , known as a *LH algebra* of \mathbf{X} . At this point, we recall that the symmetric algebra $S(\mathcal{H}_\omega)$ of \mathcal{H}_ω can be endowed with a Poisson–Hopf algebra structure which allows an algorithmic computation of t -independent constants of the motion, hence eventually simplifying the deduction of a superposition rule [1, 2].

3 The symplectic Lie algebra $\mathfrak{sp}(6, \mathbb{R})$

The real symplectic Lie algebra $\mathfrak{sp}(6, \mathbb{R})$ is defined by the matrix constraint

$$\mathfrak{sp}(6, \mathbb{R}) := \{X \in \mathfrak{gl}(6, \mathbb{R}); X^T J + JX = 0\}$$

where J is the matrix $J := \begin{pmatrix} 0 & \text{Id}_3 \\ -\text{Id}_3 & 0 \end{pmatrix}$. Instead of using the usual basis of $\mathfrak{sp}(6, \mathbb{R})$ described through elementary matrices, we consider the so-called boson basis [11, 14], as done in [6] for the construction dealing with the Lie algebra $\mathfrak{sp}(4, \mathbb{R})$.

In terms of the creation a_i and annihilation a_i^\dagger operators ($1 \leq i \leq 3$), the Lie algebra $\mathfrak{sp}(6, \mathbb{R})$ is spanned by the operators $a_i^\dagger a_j$, $a_i^\dagger a_j^\dagger$ and $a_i a_j$. We label this boson basis as

$$X_{i,j} := a_i^\dagger a_j, \quad X_{-i,j} := a_i^\dagger a_j^\dagger, \quad X_{i,-j} := a_i a_j$$

in such a way that the constraint

$$X_{i,j} + \varepsilon_i \varepsilon_j X_{-j,-i} = 0, \quad \varepsilon_i := \text{sign}(i), \quad \varepsilon_j := \text{sign}(j), \quad -3 \leq i, j \leq 3$$

is satisfied. The commutation relations of $\mathfrak{sp}(6, \mathbb{R})$ over this basis are

$$[X_{i,j}, X_{k,\ell}] = \delta_{j,k} X_{i,\ell} - \delta_{i,\ell} X_{k,j} + \varepsilon_i \varepsilon_j \delta_{j,-\ell} X_{k,-i} - \varepsilon_i \varepsilon_j \delta_{i,-k} X_{-j,\ell}$$

where $-3 \leq i, j, k, \ell \leq 3$.

Let us consider the six-dimensional fundamental (defining) representation $\Gamma_{\omega_1} : \mathfrak{sp}(6, \mathbb{R}) \rightarrow \mathfrak{gl}(6, \mathbb{R})$ of $\mathfrak{sp}(6, \mathbb{R})$ given in matrix form by

$$A_{\Gamma_{\omega_1}} = \begin{pmatrix} X_{1,1} & X_{1,2} & X_{1,3} & -X_{-1,1} & -X_{-1,2} & -X_{-1,3} \\ X_{2,1} & X_{2,2} & X_{2,3} & -X_{-1,2} & -X_{-2,2} & -X_{-2,3} \\ X_{3,1} & X_{3,2} & X_{3,3} & -X_{-1,3} & -X_{-2,3} & -X_{-3,3} \\ X_{1,-1} & X_{1,-2} & X_{1,-3} & -X_{1,1} & -X_{2,1} & -X_{3,1} \\ X_{1,-2} & X_{2,-2} & X_{2,-3} & -X_{1,2} & -X_{2,2} & -X_{3,2} \\ X_{1,-3} & X_{2,-3} & X_{3,-3} & -X_{1,3} & -X_{2,3} & -X_{3,3} \end{pmatrix}.$$

As $\mathfrak{sp}(6, \mathbb{R})$ is a simple Lie algebra of rank 3, it possesses 3 functionally independent invariants C_2 , C_4 and C_6 of degrees 2, 4 and 6, respectively, which can be obtained by the standard determinantal methods [10, 19]. Considering the characteristic equation $|A_{\Gamma_{\omega_1}} - \lambda \text{Id}_6|$ of the matrix, the expansion gives (see [4])

$$|A_{\Gamma_{\omega_1}} - \lambda \text{Id}_6| = \lambda^6 + C_2 \lambda^4 + C_4 \lambda^2 + C_6.$$

The quadratic Casimir operator C_2 , corresponding to the Killing form, has the following symmetric expression

$$\begin{aligned} C_2 = & X_{1,1}^2 + X_{2,2}^2 + X_{3,3}^2 + X_{1,2}X_{2,1} + X_{2,1}X_{1,2} + X_{1,3}X_{3,1} + X_{3,1}X_{1,3} \\ & + X_{2,3}X_{3,2} + X_{3,2}X_{2,3} - X_{-1,2}X_{1,-2} - X_{1,-2}X_{-1,2} - X_{-1,3}X_{1,-3} \\ & - X_{1,-3}X_{-1,3} - X_{-2,3}X_{2,-3} - X_{2,-3}X_{-2,3} - 2X_{-1,1}X_{1,-1} \\ & - 2X_{1,-1}X_{-1,1} - 2X_{-2,2}X_{2,-2} - 2X_{2,-2}X_{-2,2} - 2X_{-3,3}X_{3,-3} \\ & - X_{3,-3}X_{-3,3}. \end{aligned} \quad (3)$$

For the purposes of this work, the invariant C_2 is sufficient. The explicit expressions for C_4 and C_6 , that are obtained along the same lines, are omitted due to their length.

The realization $\Phi : \mathfrak{sp}(6, \mathbb{R}) \rightarrow \mathfrak{X}(\mathbb{T}^*\mathbb{R}^3)$ induced by Γ_{ω_1} is spanned by the following 21 vector fields expressed in global coordinates $(\mathbf{q}, \mathbf{p}) = (q_1, q_2, q_3, p_1, p_2, p_3)$ of $\mathbb{T}^*\mathbb{R}^3$:

$$\begin{aligned} \mathbf{X}_1 &:= \Phi(X_{1,1}) = q_1 \frac{\partial}{\partial q_1} - p_1 \frac{\partial}{\partial p_1}, & \mathbf{X}_2 &:= \Phi(X_{1,2}) = q_1 \frac{\partial}{\partial q_2} - p_2 \frac{\partial}{\partial p_1} \\ \mathbf{X}_3 &:= \Phi(X_{1,3}) = q_1 \frac{\partial}{\partial q_3} - p_3 \frac{\partial}{\partial p_1}, & \mathbf{X}_4 &:= \Phi(X_{2,1}) = q_2 \frac{\partial}{\partial q_1} - p_1 \frac{\partial}{\partial p_2} \\ \mathbf{X}_5 &:= \Phi(X_{2,2}) = q_2 \frac{\partial}{\partial q_2} - p_2 \frac{\partial}{\partial p_2}, & \mathbf{X}_6 &:= \Phi(X_{2,3}) = q_2 \frac{\partial}{\partial q_3} - p_3 \frac{\partial}{\partial p_2} \\ \mathbf{X}_7 &:= \Phi(X_{3,1}) = q_3 \frac{\partial}{\partial q_1} - p_1 \frac{\partial}{\partial p_3}, & \mathbf{X}_8 &:= \Phi(X_{3,2}) = q_3 \frac{\partial}{\partial q_2} - p_2 \frac{\partial}{\partial p_3} \\ \mathbf{X}_9 &:= \Phi(X_{3,3}) = q_3 \frac{\partial}{\partial q_3} - p_3 \frac{\partial}{\partial p_3}, & \mathbf{X}_{10} &:= \Phi(X_{-1,1}) = -q_1 \frac{\partial}{\partial p_1} \\ \mathbf{X}_{11} &:= \Phi(X_{-1,2}) = -q_2 \frac{\partial}{\partial p_1} - q_1 \frac{\partial}{\partial p_2}, & \mathbf{X}_{12} &:= \Phi(X_{-1,3}) = -q_3 \frac{\partial}{\partial p_1} - q_1 \frac{\partial}{\partial p_3} \\ \mathbf{X}_{13} &:= \Phi(X_{-2,2}) = -q_2 \frac{\partial}{\partial p_2}, & \mathbf{X}_{14} &:= \Phi(X_{-2,3}) = -q_3 \frac{\partial}{\partial p_2} - q_2 \frac{\partial}{\partial p_3} \\ \mathbf{X}_{15} &:= \Phi(X_{-3,3}) = -q_3 \frac{\partial}{\partial p_3}, & \mathbf{X}_{16} &:= \Phi(X_{1,-1}) = p_1 \frac{\partial}{\partial q_1} \\ \mathbf{X}_{17} &:= \Phi(X_{1,-2}) = p_2 \frac{\partial}{\partial q_1} + p_1 \frac{\partial}{\partial q_2}, & \mathbf{X}_{18} &:= \Phi(X_{1,-3}) = p_3 \frac{\partial}{\partial q_1} + p_1 \frac{\partial}{\partial q_3} \\ \mathbf{X}_{19} &:= \Phi(X_{2,-2}) = p_2 \frac{\partial}{\partial q_2}, & \mathbf{X}_{20} &:= \Phi(X_{2,-3}) = p_3 \frac{\partial}{\partial q_2} + p_2 \frac{\partial}{\partial q_3} \\ \mathbf{X}_{21} &:= \Phi(X_{3,-3}) = p_3 \frac{\partial}{\partial q_3}. \end{aligned} \quad (4)$$

As these vector fields are naturally associated to the fundamental representation Γ_{ω_1} of $\mathfrak{sp}(6, \mathbb{R})$, they are Hamiltonian vector fields with respect to the canonical symplectic form

$$\omega := dq_1 \wedge dp_1 + dq_2 \wedge dp_2 + dq_3 \wedge dp_3 \quad (5)$$

of $\mathbb{T}^*\mathbb{R}^3$. The corresponding Hamiltonian functions, determined by means of the inner product condition (2), have the following expressions:

$$\begin{aligned} h_1 &= q_1 p_1, & h_2 &= q_1 p_2, & h_3 &= q_1 p_3, & h_4 &= q_2 p_1 \\ h_5 &= q_2 p_2, & h_6 &= q_2 p_3, & h_7 &= q_3 p_1, & h_8 &= q_3 p_2 \\ h_9 &= q_3 p_3, & h_{10} &= \frac{1}{2} q_1^2, & h_{11} &= q_1 q_2, & h_{12} &= q_1 q_3 \\ h_{13} &= \frac{1}{2} q_2^2, & h_{14} &= q_2 q_3, & h_{15} &= \frac{1}{2} q_3^2, & h_{16} &= \frac{1}{2} p_1^2 \\ h_{17} &= p_1 p_2, & h_{18} &= p_1 p_3, & h_{19} &= \frac{1}{2} p_2^2, & h_{20} &= p_2 p_3 \\ h_{21} &= \frac{1}{2} p_3^2. \end{aligned} \quad (6)$$

A routine computation shows that they span a Lie algebra isomorphic to $\mathfrak{sp}(6, \mathbb{R})$ with respect to the Poisson bracket $\{\cdot, \cdot\}_\omega$ induced on $C^\infty(\mathbb{T}^*\mathbb{R}^3)$ by the canonical symplectic form ω .

By the procedure described in [6], the t -dependent vector field

$$\mathbf{X} := \sum_{i=1}^{21} b_i(t) \mathbf{X}_i \quad (7)$$

where the $b_i \in C^\infty(\mathbb{R})$ are arbitrary t -dependent functions ($1 \leq i \leq 21$), determines a LH system on $\mathbb{T}^*\mathbb{R}^3$. Its VG Lie algebra V^X and LH algebra \mathcal{H}_ω , respectively spanned by the vector fields (4) and the Hamiltonian functions (6), are both isomorphic to $\mathfrak{sp}(6, \mathbb{R})$. The first-order system of ODEs on $\mathbb{T}^*\mathbb{R}^3$ associated to \mathbf{X} is given by

$$\begin{pmatrix} \dot{q}_1 \\ \dot{q}_2 \\ \dot{q}_3 \\ \dot{p}_1 \\ \dot{p}_2 \\ \dot{p}_3 \end{pmatrix} = \begin{pmatrix} b_1(t) & b_4(t) & b_7(t) & b_{16}(t) & b_{17}(t) & b_{18}(t) \\ b_2(t) & b_5(t) & b_8(t) & b_{17}(t) & b_{19}(t) & b_{20}(t) \\ b_3(t) & b_6(t) & b_9(t) & b_{18}(t) & b_{20}(t) & b_{21}(t) \\ -b_{10}(t) & -b_{11}(t) & -b_{12}(t) & -b_1(t) & -b_2(t) & -b_3(t) \\ -b_{11}(t) & -b_{13}(t) & -b_{14}(t) & -b_4(t) & -b_5(t) & -b_6(t) \\ -b_{12}(t) & -b_{14}(t) & -b_{15}(t) & -b_7(t) & -b_8(t) & -b_9(t) \end{pmatrix} \begin{pmatrix} q_1 \\ q_2 \\ q_3 \\ p_1 \\ p_2 \\ p_3 \end{pmatrix} \quad (8)$$

where it is easily seen that the coefficient matrix itself is symplectic. We observe that this Lie system \mathbf{X} cannot be reduced by invariants to a lower-dimensional Lie system, as the maximum rank of the generalized distribution associated to \mathbf{X} is 6 (see [6] for details on the reduction by invariants of Lie systems).

3.1 Constants of the motion and superposition rule

We now obtain t -independent constants of the motion for the $\mathfrak{sp}(6, \mathbb{R})$ -LH systems (8) by applying the so-called coalgebra formalism [1, 2], which, in turn, will allow us to deduce a (nonlinear) superposition rule.

In terms of the Hamiltonian functions (6), spanning the $\mathfrak{sp}(6, \mathbb{R})$ -LH algebra \mathcal{H}_ω , the quadratic Casimir invariant C_2 in (3) reads

$$C_2 = h_1^2 + h_5^2 + h_9^2 + 2h_2h_4 + 2h_3h_7 + 2h_6h_8 - 2h_{11}h_{17} - 2h_{12}h_{18} - 2h_{14}h_{20} - 4h_{10}h_{16} - 4h_{13}h_{19} - 4h_{15}h_{21}. \quad (9)$$

The smallest integer s such that the diagonal prolongations of the vector fields (4) are linearly independent at a generic point of the product manifold

$$(\mathbb{T}^*\mathbb{R}^3)^s := \mathbb{T}^*\mathbb{R}^3 \times \dots \times \mathbb{T}^*\mathbb{R}^3$$

is $s = 6$. Thus, we consider the Hamiltonian functions $h_i^{(k)} \in C^\infty((\mathbb{T}^*\mathbb{R}^3)^k)$, for $1 \leq k \leq 7$, given by

$$h_i^{(k)} = \sum_{\ell=1}^k h_i(\mathbf{q}^{(\ell)}, \mathbf{p}^{(\ell)}), \quad 1 \leq i \leq 21$$

where $(\mathbf{q}^{(\ell)}, \mathbf{p}^{(\ell)})$ denote the canonical global coordinates in the ℓ^{th} -copy of $\mathbb{T}^*\mathbb{R}^3$ within the product manifold. Using the Casimir invariant (9), we obtain the constants of the motion $F^{(k)} := C_2(h_1^{(k)}, \dots, h_{21}^{(k)})$ for the diagonal prolongation $\tilde{\mathbf{X}}^7$ of \mathbf{X} to $(\mathbb{T}^*\mathbb{R}^3)^7$. Concretely, we find that $F^{(1)} = 0$, while

$$F^{(2)} = - \left(p_1^{(2)} q_1^{(1)} - p_1^{(1)} q_1^{(2)} + p_2^{(2)} q_2^{(1)} - p_2^{(1)} q_2^{(2)} + p_3^{(2)} q_3^{(1)} - p_3^{(1)} q_3^{(2)} \right)^2.$$

This second-order constant of the motion further gives rise to the following 10 constants of the motion by permutation S_{ij} of the variables $(\mathbf{q}^{(i)}, \mathbf{p}^{(i)}) \leftrightarrow (\mathbf{q}^{(j)}, \mathbf{p}^{(j)})$:

$$\begin{aligned} F_{13}^{(2)} &= S_{13}(F^{(2)}), & F_{14}^{(2)} &= S_{14}(F^{(2)}), & F_{15}^{(2)} &= S_{15}(F^{(2)}) \\ F_{16}^{(2)} &= S_{16}(F^{(2)}), & F_{17}^{(2)} &= S_{17}(F^{(2)}), & F_{23}^{(2)} &= S_{23}(F^{(2)}) \\ F_{24}^{(2)} &= S_{24}(F^{(2)}), & F_{25}^{(2)} &= S_{25}(F^{(2)}), & F_{26}^{(2)} &= S_{26}(F^{(2)}) \\ F_{27}^{(2)} &= S_{27}(F^{(2)}). \end{aligned}$$

Note that the third-order constant of the motion $F^{(3)}$ is expressed in terms of the quadratic constants of the motion $F^{(2)}$, $F_{13}^{(2)}$ and $F_{23}^{(2)}$ as

$$F^{(3)} = F^{(2)} + F_{13}^{(2)} + F_{23}^{(2)}.$$

In this situation, a superposition rule for the $\mathfrak{sp}(6, \mathbb{R})$ -LH system (8) in terms of 6 significative constants k_ℓ and 6 particular solutions $(\mathbf{q}^{(\ell)}, \mathbf{p}^{(\ell)})$, with $1 \leq \ell \leq 6$, can be derived by solving the system of algebraic equations

$$\begin{aligned} F^{(2)} &= -k_1^2, & F_{23}^{(2)} &= -k_2^2, & F_{24}^{(2)} &= -k_3^2, & F_{25}^{(2)} &= -k_4^2 \\ F_{26}^{(2)} &= -k_5^2, & F_{27}^{(2)} &= -k_6^2 \end{aligned}$$

provided that the general solution corresponds to $(\mathbf{q}^{(7)}(t), \mathbf{p}^{(7)}(t))$. The explicit expression, even with the use of symbolic computation packages, is computationally cumbersome and quite long, for which reason we omit its detailed derivation.

3.2 Applications

The generic t -dependent Hamiltonian h obtained from the Hamiltonian functions (6) is characterized by

$$h = \sum_{i=1}^{21} b_i(t) h_i$$

in such a way that the system (8) corresponds to the Hamilton equations on $\mathbb{T}^*\mathbb{R}^3$ with respect to the canonical symplectic form (5). In order to find some interesting and manageable physical applications, some restrictions on the generators are required, i.e., considering sets of generators \mathcal{S} of the LH algebra $\mathcal{H}_\omega \simeq \mathfrak{sp}(6, \mathbb{R})$ such that $\text{card}(\mathcal{S}) < 21$.

Considering for example the subset

$$\mathcal{S}_I := \{h_2, h_4, h_9, h_{10}, h_{13}, h_{15}, h_{16}, h_{19}, h_{21}\}$$

formed by nine generators of the LH algebra \mathcal{H}_ω , it follows that the t -dependent Hamiltonian h^I is given by

$$\begin{aligned} h^I = & b_2(t)q_1p_2 + b_4(t)q_2p_1 + b_9(t)q_3p_3 + \frac{b_{10}(t)}{2}q_1^2 + \frac{b_{13}(t)}{2}q_2^2 + \frac{b_{15}(t)}{2}q_3^2 \\ & + \frac{b_{16}(t)}{2}p_1^2 + \frac{b_{19}(t)}{2}p_2^2 + \frac{b_{21}(t)}{2}p_3^2. \end{aligned} \quad (10)$$

An alternative choice that involves the same number of generators can be taken as

$$\mathcal{S}_{II} := \{h_2 - h_4, h_3 - h_7, h_6 - h_8, h_{10}, h_{13}, h_{15}, h_{16}, h_{19}, h_{21}\}$$

where the corresponding t -dependent Hamiltonian h^{II} has the form

$$\begin{aligned} h^{II} = & \frac{b_{16}(t)}{2}p_1^2 + \frac{b_{10}(t)}{2}q_1^2 + \frac{b_{19}(t)}{2}p_2^2 + \frac{b_{13}(t)}{2}q_2^2 + \frac{b_{21}(t)}{2}p_3^2 + \frac{b_{15}(t)}{2}q_3^2 \\ & + b_2(t)(q_1p_2 - q_2p_1) + b_3(t)(q_1p_3 - q_3p_1) + b_6(t)(q_2p_3 - q_3p_2) \end{aligned} \quad (11)$$

provided that $b_4(t) = -b_2(t)$, $b_7(t) = -b_3(t)$ and $b_8(t) = -b_6(t)$. In the latter expression, the mechanical meaning of each of the intervening terms is more transparent, suggesting that the Hamiltonian can be used to describe (coupled) systems in either Euclidean or non-Euclidean spaces.

3.2.1 A time-dependent electromagnetic field

Let us now consider t -dependent functions $m_i(t), e_i(t), \gamma(t) \in C^\infty(\mathbb{R})$, with $1 \leq i \leq 3$, such that

$$m_i(t) > 0 \quad (1 \leq i \leq 3), \quad \ddot{\gamma}(t) \neq 0, \quad t \in \mathbb{R}.$$

Define now the t -dependent vector field \mathbf{A} on \mathbb{R}^3 given by

$$A_1 := -\frac{1}{2}q_2\gamma(t), \quad A_2 := \frac{1}{2}q_1\gamma(t), \quad A_3 := \frac{1}{2}q_3\gamma(t)$$

as well as the scalar oscillator potential

$$\phi := \phi_1 + \phi_2 + \phi_3, \quad \phi_i := \frac{1}{2}q_i^2 \quad (1 \leq i \leq 3).$$

Consider the following choices for the t -dependent functions $b_i(t)$ appearing in the t -dependent Hamiltonian (10):

$$\begin{aligned} b_2(t) &= -\frac{\gamma(t)e_2(t)}{2m_2(t)}, & b_{10}(t) &= e_1(t) + \frac{\gamma^2(t)e_2^2(t)}{4m_2(t)}, & b_{16}(t) &= \frac{1}{m_1(t)} \\ b_4(t) &= \frac{\gamma(t)e_1(t)}{2m_1(t)}, & b_{13}(t) &= e_2(t) + \frac{\gamma^2(t)e_1^2(t)}{4m_1(t)}, & b_{19}(t) &= \frac{1}{m_2(t)} \\ b_9(t) &= -\frac{\gamma(t)e_3(t)}{2m_3(t)}, & b_{15}(t) &= e_3(t) + \frac{\gamma^2(t)e_3^2(t)}{4m_3(t)}, & b_{21}(t) &= \frac{1}{m_3(t)}. \end{aligned}$$

The t -dependent Hamiltonian h^E obtained from (10) with these choices turns out to be

$$h^E = \sum_{i=1}^3 \left(\frac{1}{2m_i(t)} (p_i - e_i(t)A_i)^2 + e_i(t)\phi_i \right).$$

This describes the motion on \mathbb{R}^3 of three particles of time-dependent masses $m_i(t)$ and time-dependent electric charges $e_i(t)$ ($1 \leq i \leq 3$) under the action of a time-dependent electromagnetic field. The magnetic field is given by

$$\mathbf{B} = \nabla \times \mathbf{A} = (0, 0, \gamma(t))$$

while the electric field reads

$$\mathbf{E} = -\nabla\phi - \frac{\partial\mathbf{A}}{\partial t} = -\frac{1}{2}(2q_1 - q_2\dot{\gamma}(t), 2q_2 + q_1\dot{\gamma}(t), 2q_3 + q_3\dot{\gamma}(t)).$$

As we are assuming that $\dot{\gamma}(t) \neq 0$ for all $t \in \mathbb{R}$, the electric field \mathbf{E} is also time-dependent.

3.2.2 Time-dependent coupled oscillators

We next show that the Hamiltonian (11) can be considered as a coupled system, using a similar ansatz to that proposed in [6]. To this extent, we first consider the Whitney sum $\mathbb{T}^*\mathbb{R}^3 = \mathbb{T}^*\mathbb{R} \oplus \mathbb{T}^*\mathbb{R} \oplus \mathbb{T}^*\mathbb{R}$ together with the three canonical projections pr_k ($1 \leq k \leq 3$) onto its factors:

$$\text{pr}_k : \mathbb{T}^*\mathbb{R}^3 \rightarrow \mathbb{T}^*\mathbb{R}, \quad (\mathbf{q}, \mathbf{p}) \mapsto (q_k, p_k), \quad 1 \leq k \leq 3.$$

The t -dependent Hamiltonian $h_{i,j}^{1D}$ on $\mathbb{T}^*\mathbb{R}$ given by

$$h_{i,j}^{1D} := \frac{b_i(t)}{2}p^2 + \frac{b_j(t)}{2}q^2$$

determines a LH system on $\mathbb{T}^*\mathbb{R}$ such that the Hamiltonian (11) can be rewritten as

$$h = \text{pr}_1^*(h_{16,10}^{\text{1D}}) + \text{pr}_2^*(h_{19,13}^{\text{1D}}) + \text{pr}_3^*(h_{21,15}^{\text{1D}}) + b_2(t)(q_1p_2 - q_2p_1) \\ + b_3(t)(q_1p_3 - q_3p_1) + b_6(t)(q_2p_3 - q_3p_2).$$

This shows that (11) describes a system formed by three 1D t -dependent Hamiltonians (namely, $h_{16,10}^{\text{1D}}$, $h_{19,13}^{\text{1D}}$ and $h_{21,15}^{\text{1D}}$) coupled through the angular momentum terms $q_1p_2 - q_2p_1$, $q_1p_3 - q_3p_1$ and $q_2p_3 - q_3p_2$.

Several interesting choices can be made for the t -dependent functions taking part in the t -dependent Hamiltonian (11), yielding generalized coupled oscillators with different properties. In order to illustrate the situation, let us consider the positive functions $m_i(t), k_i(t), \gamma_i(t) \in C^\infty(\mathbb{R})$ for $1 \leq i \leq 3$.

- *A time-dependent Hamiltonian of coupled harmonic oscillators.* Suppose that

$$k_i(t) > \frac{\gamma_i^2(t)}{4m_i(t)}, \quad t \in \mathbb{R}, \quad 1 \leq i \leq 3.$$

Define now

$$\Omega_i(t) := \sqrt{\frac{1}{m_i(t)} \left(k_i(t) - \frac{\gamma_i^2(t)}{4m_i(t)} \right)}, \quad t \in \mathbb{R}, \quad 1 \leq i \leq 3.$$

Denote by h^{CHO} the t -dependent Hamiltonian obtained from (11) after the following choices:

$$b_{16}(t) = \frac{1}{m_1(t)}, \quad b_{10}(t) = m_1(t)\Omega_1^2(t), \quad b_{19}(t) = \frac{1}{m_2(t)} \\ b_{13}(t) = m_2(t)\Omega_2^2(t), \quad b_{21}(t) = \frac{1}{m_3(t)}, \quad b_{15}(t) = m_3(t)\Omega_3^2(t).$$

Then, h^{CHO} reads as

$$h^{\text{CHO}} = \sum_{i=1}^3 \left(\frac{1}{2m_i(t)} p_i^2 + \frac{1}{2} m_i(t) \Omega_i^2(t) q_i^2 \right) + b_2(t)(q_1p_2 - q_2p_1) \\ + b_3(t)(q_1p_3 - q_3p_1) + b_6(t)(q_2p_3 - q_3p_2).$$

In this case, the three 1D systems $h_{16,10}^{\text{1D}}$, $h_{19,13}^{\text{1D}}$ and $h_{21,15}^{\text{1D}}$ turn out to be time-dependent harmonic oscillators with time-dependent masses $m_i(t)$, time-dependent damping ‘‘constants’’ $\gamma_i(t)$, time-dependent spring ‘‘constants’’ $k_i(t)$ and time-dependent frequencies $\Omega_i(t)$ ($1 \leq i \leq 3$). Additional interesting choices can be considered for the t -dependent coupling constants $b_2(t)$, $b_3(t)$ and $b_6(t)$, as they can be taken to depend on the time-dependent masses $m_i(t)$, Hooke ‘‘constants’’ $k_i(t)$ or friction ‘‘constants’’ $\gamma_i(t)$.

- *Coupled Caldirola–Kanai Hamiltonian.* Let us now define

$$\lambda_i(t) := \frac{\gamma_i(t)}{m_i(t)}, \quad \Omega_i(t) := \sqrt{\frac{k_i(t)}{m_i(t)}}, \quad t \in \mathbb{R}, \quad 1 \leq i \leq 3$$

and denote by h^{CCK} the t -dependent Hamiltonian obtained from (11) after the choices

$$\begin{aligned} b_{16}(t) &= \frac{1}{m_1(t)} e^{-2 \int_0^t \lambda_1(s) ds}, & b_{10}(t) &= m_1(t) \Omega_1^2(t) e^{2 \int_0^t \lambda_1(s) ds} \\ b_{19}(t) &= \frac{1}{m_2(t)} e^{-2 \int_0^t \lambda_2(s) ds}, & b_{13}(t) &= m_2(t) \Omega_2^2(t) e^{2 \int_0^t \lambda_2(s) ds} \\ b_{21}(t) &= \frac{1}{m_3(t)} e^{-2 \int_0^t \lambda_3(s) ds}, & b_{15}(t) &= m_3(t) \Omega_3^2(t) e^{2 \int_0^t \lambda_3(s) ds}. \end{aligned}$$

In this case, h^{CCK} adopts the form

$$\begin{aligned} h^{\text{CCK}} &= \sum_{i=1}^3 \left(\frac{1}{2m_i(t)} e^{-2 \int_0^t \lambda_i(s) ds} p_i^2 + \frac{1}{2} m_i(t) \Omega_i^2(t) e^{2 \int_0^t \lambda_i(s) ds} q_i^2 \right) \\ &\quad + b_2(t)(q_1 p_2 - q_2 p_1) + b_3(t)(q_1 p_3 - q_3 p_1) + b_6(t)(q_2 p_3 - q_3 p_2). \end{aligned}$$

It is straightforward to verify that the 1D systems $h_{16,10}^{\text{1D}}$, $h_{19,13}^{\text{1D}}$ and $h_{21,15}^{\text{1D}}$ correspond to the so-called Caldirola–Kanai Hamiltonians [7, 12]. They describe the motion of a particle of time-dependent mass $m_i(t)$ attached to a spring with time-dependent Hooke “constant” given by $k_i(t) := m_i(t) \Omega_i^2(t)$, with $\Omega_i(t)$ being the time-dependent frequency, and subjected to a frictional force with a time-dependent friction “constant” $\gamma_i(t) := m_i(t) \lambda_i(t)$ for $1 \leq i \leq 3$.

4 The special unitary Lie algebra $\mathfrak{su}(3)$

The relevance of the Lie algebra $\mathfrak{su}(3)$ is widely recognized due to its importance in particle physics (see [9] and references therein). From the algebraic point of view, recall that $\mathfrak{su}(3)$ is the compact real form of the (complex) Lie algebra $\mathfrak{sl}(3, \mathbb{C})$. Let us now consider the $\mathfrak{su}(3)$ -subalgebra of $\mathfrak{sp}(6, \mathbb{R})$ spanned by

$$\begin{aligned} E_1^+ &:= \frac{1}{2}(X_{1,2} - X_{2,1} - X_{-1,2} + X_{1,-2}) \\ E_2^+ &:= \frac{1}{2}(X_{2,3} - X_{3,2} - X_{-2,3} + X_{2,-3}) \\ E_1^- &:= \frac{1}{2}(-X_{1,2} + X_{2,1} - X_{-1,2} + X_{1,-2}) \\ E_2^- &:= \frac{1}{2}(-X_{2,3} + X_{3,2} - X_{-2,3} + X_{2,-3}). \end{aligned} \tag{12}$$

The elements E_1^+, E_2^+ correspond to the two positive simple roots of $\mathfrak{su}(3)$, while E_1^- and E_2^- are associated with the negative roots. The generators of the Cartan subalgebra are

$$\begin{aligned} H_1 &:= [E_1^+, E_1^-] = -X_{-1,1} + X_{-2,2} + X_{1,-1} - X_{2,2} \\ H_2 &:= [E_2^+, E_2^-] = -X_{-2,2} + X_{-3,3} + X_{2,-2} - X_{3,-3}. \end{aligned} \tag{13}$$

The Cartan–Weyl basis of $\mathfrak{su}(3)$ is obtained by from (12) and (13) together with the non-simple roots

$$\begin{aligned} E_3^+ &:= [E_1^+, E_2^+] = \frac{1}{2}(X_{1,3} - X_{3,1} - X_{-1,3} + X_{1,-3}) \\ E_3^- &:= -[E_1^-, E_2^-] = \frac{1}{2}(-X_{1,3} + X_{3,1} - X_{-1,3} + X_{1,-3}). \end{aligned}$$

The nonvanishing commutation relations of $\mathfrak{su}(3)$ over this basis, up to skew-symmetry, are given by

$$\begin{aligned} [E_1^+, E_2^+] &= E_3^+, & [E_1^+, E_1^-] &= H_1, & [E_1^+, E_3^-] &= -E_2^- \\ [E_1^+, H_1] &= -2E_1^+, & [E_1^+, H_2] &= E_1^+, & [E_2^+, E_2^-] &= H_2 \\ [E_2^+, E_3^-] &= E_1^-, & [E_2^+, H_1] &= E_2^+, & [E_2^+, H_2] &= -2E_2^+ \\ [E_3^+, E_1^-] &= -E_2^+, & [E_3^+, E_2^-] &= E_1^+, & [E_3^+, E_3^-] &= H_1 + H_2 \\ [E_3^+, H_1] &= -E_3^+, & [E_3^+, H_2] &= -E_3^+, & [E_1^-, E_2^-] &= -E_3^- \\ [E_1^-, H_1] &= 2E_1^-, & [E_1^-, H_2] &= -E_1^-, & [E_2^-, H_1] &= -E_2^- \\ [E_2^-, H_2] &= 2E_2^-, & [E_3^-, H_1] &= E_3^-, & [E_3^-, H_2] &= E_3^-. \end{aligned}$$

The fundamental representation Γ_{ω_1} of $\mathfrak{sp}(6, \mathbb{R})$, when restricted to the subalgebra $\mathfrak{su}(3)$, gives rise to the branching rule $\mathfrak{su}(3)$ -representation $\Gamma_{\omega_1|_{\mathfrak{su}(3)}} : \mathfrak{su}(3) \rightarrow \mathfrak{gl}(6, \mathbb{R})$ determined by the matrix condition

$$A_{\Gamma_{\omega_1|_{\mathfrak{su}(3)}}} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \otimes B_1 + \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \otimes B_2$$

where B_1 and B_2 are the 3×3 real matrices given by

$$\begin{aligned} B_1 &:= \frac{1}{2} \begin{pmatrix} 0 & E_1^+ - E_1^- & E_3^+ - E_3^- \\ -E_1^+ + E_1^- & 0 & E_2^+ - E_2^- \\ -E_3^+ + E_3^- & -E_2^+ + E_2^- & 0 \end{pmatrix} \\ B_2 &:= \frac{1}{2} \begin{pmatrix} 2H_1 & E_1^+ + E_1^- & E_3^+ + E_3^- \\ E_1^+ + E_1^- & -2H_1 + 2H_2 & E_2^+ + E_2^- \\ E_3^+ + E_3^- & E_2^+ + E_2^- & -2H_2 \end{pmatrix}. \end{aligned}$$

In particular, note that the restriction of the fundamental representation Γ_{ω_1} of $\mathfrak{sp}(4, \mathbb{R})$ to $\mathfrak{su}(3)$ gives rise to the branching rule (see [16])

$$(1, 0, 0) \downarrow (1, 0) \oplus (0, 1)$$

where $(1, 0, 0)$ is highest weight of the representation Γ_{ω_1} , while $(1, 0)$ and $(0, 1)$ are the highest weights of the three-dimensional quark and antiquark representations of $\mathfrak{su}(3)$, respectively².

²The representation $(1, 0) \oplus (0, 1)$ is irreducible as a real representation of $\mathfrak{su}(3)$, but reducible as a complex one.

The associated realization $\Psi : \mathfrak{su}(3) \rightarrow \mathfrak{X}(\mathbb{T}^*\mathbb{R}^3)$ induced by the representation $\Gamma_{\omega_1}|_{\mathfrak{su}(3)}$ is thus spanned by the eight vector fields

$$\begin{aligned} \mathbf{Y}_1 &:= \Psi(E_1^+), & \mathbf{Y}_2 &:= \Psi(E_2^+), & \mathbf{Y}_3 &:= \Psi(E_3^+) \\ \mathbf{Y}_4 &:= \Psi(E_1^-), & \mathbf{Y}_5 &:= \Psi(E_2^-), & \mathbf{Y}_6 &:= \Psi(E_3^-) \\ \mathbf{Y}_7 &:= \Psi(H_1), & \mathbf{Y}_8 &:= \Psi(H_2). \end{aligned}$$

In terms of the global coordinates (\mathbf{q}, \mathbf{p}) of $\mathbb{T}^*\mathbb{R}^3$, these vector fields are explicitly given by

$$\begin{aligned} \mathbf{Y}_1 &= \frac{1}{2} \left((-q_2 + p_2) \frac{\partial}{\partial q_1} + (q_1 + p_1) \frac{\partial}{\partial q_2} + (q_2 - p_2) \frac{\partial}{\partial p_1} + (q_1 + p_1) \frac{\partial}{\partial p_2} \right) \\ \mathbf{Y}_2 &= \frac{1}{2} \left((-q_3 + p_3) \frac{\partial}{\partial q_2} + (q_2 + p_2) \frac{\partial}{\partial q_3} + (q_3 - p_3) \frac{\partial}{\partial p_2} + (q_2 + p_2) \frac{\partial}{\partial p_3} \right) \\ \mathbf{Y}_3 &= \frac{1}{2} \left((-q_3 + p_3) \frac{\partial}{\partial q_1} + (q_1 + p_1) \frac{\partial}{\partial q_3} + (q_3 - p_3) \frac{\partial}{\partial p_1} + (q_1 + p_1) \frac{\partial}{\partial p_3} \right) \\ \mathbf{Y}_4 &= \frac{1}{2} \left((q_2 + p_2) \frac{\partial}{\partial q_1} + (-q_1 + p_1) \frac{\partial}{\partial q_2} + (q_2 + p_2) \frac{\partial}{\partial p_1} + (q_1 - p_1) \frac{\partial}{\partial p_2} \right) \\ \mathbf{Y}_5 &= \frac{1}{2} \left((q_3 + p_3) \frac{\partial}{\partial q_2} + (-q_2 + p_2) \frac{\partial}{\partial q_3} + (q_3 + p_3) \frac{\partial}{\partial p_2} + (q_2 - p_2) \frac{\partial}{\partial p_3} \right) \\ \mathbf{Y}_6 &= \frac{1}{2} \left((q_3 + p_3) \frac{\partial}{\partial q_1} + (-q_1 + p_1) \frac{\partial}{\partial q_3} + (q_3 + p_3) \frac{\partial}{\partial p_1} + (q_1 - p_1) \frac{\partial}{\partial p_3} \right) \\ \mathbf{Y}_7 &= p_1 \frac{\partial}{\partial q_1} - p_2 \frac{\partial}{\partial q_2} + q_1 \frac{\partial}{\partial p_1} - q_2 \frac{\partial}{\partial p_2} \\ \mathbf{Y}_8 &= p_2 \frac{\partial}{\partial q_2} - p_3 \frac{\partial}{\partial q_3} + q_2 \frac{\partial}{\partial p_2} - q_3 \frac{\partial}{\partial p_3}. \end{aligned} \tag{14}$$

As expected, they are Hamiltonian vector fields with respect to the canonical symplectic form (5), as they have been obtained by restriction of the fundamental representation Γ_{ω_1} of $\mathfrak{sp}(6, \mathbb{R})$. The corresponding Hamiltonian functions h'_i , fulfilling the inner product condition (2), are easily verified to be

$$\begin{aligned} h'_1 &= \frac{1}{2}(q_1 p_2 - q_2 p_1 + p_1 p_2 - q_1 q_2), & h'_2 &= \frac{1}{2}(q_2 p_3 - q_3 p_2 + p_2 p_3 - q_2 q_3) \\ h'_3 &= \frac{1}{2}(q_1 p_3 - q_3 p_1 + p_1 p_3 - q_1 q_3), & h'_4 &= \frac{1}{2}(-q_1 p_2 + q_2 p_1 + p_1 p_2 - q_1 q_2) \\ h'_5 &= \frac{1}{2}(-q_2 p_3 + q_3 p_2 + p_2 p_3 - q_2 q_3), & h'_6 &= \frac{1}{2}(-q_1 p_3 + q_3 p_1 + p_1 p_3 - q_1 q_3) \\ h'_7 &= \frac{1}{2}(-q_1^2 + q_2^2 + p_1^2 - p_2^2), & h'_8 &= \frac{1}{2}(-q_2^2 + q_3^2 + p_2^2 - p_3^2). \end{aligned} \tag{15}$$

A routine computation shows that they span a Lie algebra isomorphic to $\mathfrak{su}(3)$. It follows that the t -dependent vector field

$$\mathbf{Y} := \sum_{i=1}^8 a_i(t) \mathbf{Y}_i \tag{16}$$

where $a_i \in C^\infty(\mathbb{R})$ is an arbitrary t -dependent function ($1 \leq i \leq 8$), determines a LH system on $\mathbb{T}^*\mathbb{R}^3$. Indeed, its VG Lie algebra V^Y , spanned by the vector fields (14), and its LH algebra \mathcal{H}'_ω , spanned by the Hamiltonian functions (15), are both isomorphic to $\mathfrak{su}(3)$.

It is worthy to be observed that the $\mathfrak{su}(3)$ -Lie system \mathbf{Y} cannot be reduced by invariants to a lower-dimensional system, as the subalgebra $\mathfrak{su}(3) \subset \mathfrak{sp}(6, \mathbb{R})$ is irreducibly embedded and the $\mathfrak{sp}(6, \mathbb{R})$ -Lie system \mathbf{X} (7) cannot itself be reduced by invariants, as pointed out before (see [5] for more details). It should be observed that a superposition rule for the $\mathfrak{su}(3)$ -LH system (16) can be derived following the same procedure described in Subsection 3.1 for the $\mathfrak{sp}(6, \mathbb{R})$ -LH system (8).

4.1 Applications to coupled systems

Following an approach similar to that of Subsection 3.2, we first look for a minimal set of generators of the $\mathfrak{su}(3)$ -LH algebra \mathcal{H}'_ω . A simple inspection shows that h'_1, h'_2, h'_4 and h'_5 span \mathcal{H}'_ω , as they are associated to the two simple roots of the subalgebra $\mathfrak{su}(3) \subset \mathfrak{sp}(6, \mathbb{R})$. Among all possible linear combinations of these generators, let us consider the following new set of generators of \mathcal{H}'_ω :

$$\begin{aligned}\tilde{h}_1 &:= h'_1 + h'_4 = p_1 p_2 - q_1 q_2, & \tilde{h}_2 &:= h'_1 - h'_4 = q_1 p_2 - q_2 p_1 \\ \tilde{h}_3 &:= h'_2 + h'_5 = p_2 p_3 - q_2 q_3, & \tilde{h}_4 &:= h'_2 - h'_5 = q_2 p_3 - q_3 p_2.\end{aligned}$$

The t -dependent Hamiltonian \tilde{h} corresponding to this choice of generators reads

$$\begin{aligned}\tilde{h} &= \tilde{a}_1(t)(p_1 p_2 - q_1 q_2) + \tilde{a}_2(t)(q_1 p_2 - q_2 p_1) \\ &\quad + \tilde{a}_3(t)(p_2 p_3 - q_2 q_3) + \tilde{a}_4(t)(q_2 p_3 - q_3 p_2).\end{aligned}\tag{17}$$

Alternatively, \tilde{h} can also be obtained from the t -dependent Hamiltonian $h' := \sum_{i=1}^8 a_i(t) h'_i$ with the following identifications of the t -dependent functions a_i :

$$\begin{aligned}a_1 &= \tilde{a}_1 + \tilde{a}_2, & a_2 &= \tilde{a}_3 + \tilde{a}_4, & a_4 &= \tilde{a}_1 - \tilde{a}_2, & a_5 &= \tilde{a}_3 - \tilde{a}_4 \\ a_3 &= a_6 = a_7 = a_8 = 0.\end{aligned}$$

The Hamilton equations associated to \tilde{h} allow the following matrix form

$$\begin{pmatrix} \dot{q}_1 \\ \dot{q}_2 \\ \dot{q}_3 \\ \dot{p}_1 \\ \dot{p}_2 \\ \dot{p}_3 \end{pmatrix} = \begin{pmatrix} 0 & -\tilde{a}_2(t) & 0 & 0 & \tilde{a}_1(t) & 0 \\ \tilde{a}_2(t) & 0 & -\tilde{a}_4(t) & \tilde{a}_1(t) & 0 & \tilde{a}_3(t) \\ 0 & \tilde{a}_4(t) & 0 & 0 & \tilde{a}_3(t) & 0 \\ 0 & \tilde{a}_1(t) & 0 & 0 & -\tilde{a}_2(t) & 0 \\ \tilde{a}_1(t) & 0 & \tilde{a}_3(t) & \tilde{a}_2(t) & 0 & -\tilde{a}_4(t) \\ 0 & \tilde{a}_3(t) & 0 & 0 & \tilde{a}_4(t) & 0 \end{pmatrix} \begin{pmatrix} q_1 \\ q_2 \\ q_3 \\ p_1 \\ p_2 \\ p_3 \end{pmatrix}.\tag{18}$$

A remarkable fact is that \tilde{h} can be interpreted as a non-trivially coupled system formed by two subsystems defined on the Minkowskian plane \mathbf{M}^{1+1} . Recall that \mathbf{M}^{1+1} is the real plane equipped with the Lorentzian metric $ds^2 = dq_x^2 - dq_y^2$, where (q_x, q_y) are global coordinates on \mathbf{M}^{1+1} . Using the standard light-cone coordinates

$$q_+ = q_x + q_y, \quad q_- = q_x - q_y$$

the Minkowskian metric reads as $ds^2 = dq_+dq_-$. The cotangent bundle $\mathbb{T}^*\mathbf{M}^{1+1}$ is naturally equipped with the canonical global coordinates (q_\pm, p_\pm) , where p_\pm are the conjugate momentum associated to q_\pm , that is,

$$p_+ = \frac{1}{2}(p_x + p_y), \quad p_- = \frac{1}{2}(p_x - p_y).$$

It follows that the kinetic energy associated to the Minkowskian metric in the light-cone coordinates (q_\pm, p_\pm) is $2p_+p_-$, while q_+q_- is just the potential of the isotropic oscillator on \mathbf{M}^{1+1} , which is one of the widely studied Drach potentials [18, 20]. As in Subsection 3.2.2, we now describe some coupled LH systems associated to the compact Lie algebra $\mathfrak{su}(3)$. Let us first consider the projections

$$\begin{aligned} \text{pr}_{12} : \mathbb{T}^*\mathbb{R}^3 &\rightarrow \mathbb{T}^*\mathbf{M}^{1+1}, & (\mathbf{q}, \mathbf{p}) &\mapsto (q_+ = q_1, q_- = q_2, p_+ = p_1, p_- = p_2) \\ \text{pr}_{23} : \mathbb{T}^*\mathbb{R}^3 &\rightarrow \mathbb{T}^*\mathbf{M}^{1+1}, & (\mathbf{q}, \mathbf{p}) &\mapsto (q_+ = q_3, q_- = q_2, p_+ = p_3, p_- = p_2). \end{aligned}$$

Then, the t -dependent Hamiltonian h_{ij}^{2D} on $\mathbb{T}^*\mathbf{M}^{1+1}$ given by

$$h_{ij}^{2D} := \tilde{a}_i(t)(p_+p_- - q_+q_-) + \tilde{a}_j(t)(q_+p_- - q_-p_+) \quad (19)$$

is a LH system on $\mathbb{T}^*\mathbf{M}^{1+1}$ with respect to the canonical symplectic form $\omega = dq_+ \wedge dp_+ + dq_- \wedge dp_-$ and such that the Hamiltonian \tilde{h} (17) associated to the $\mathfrak{su}(3)$ -LH system (18) can be expressed as

$$\tilde{h} = \text{pr}_{12}^*(h_{12}^{2D}) + \text{pr}_{23}^*(h_{34}^{2D})$$

thus corresponding to a coupling of two LH systems defined on $\mathbb{T}^*\mathbf{M}^{1+1}$.

The Hamiltonian h_{ij}^{2D} (19) is expressed in the usual canonical coordinates (q_x, q_y, p_x, p_y) of $\mathbb{T}^*\mathbf{M}^{1+1}$ as

$$h_{ij}^{2D} = \frac{\tilde{a}_i(t)}{4}(p_x^2 - p_y^2) - \tilde{a}_i(t)(q_x^2 - q_y^2) - \tilde{a}_j(t)(q_x p_y - q_y p_x).$$

This shows that h_{ij}^{2D} is a particular case of the $\mathfrak{so}(1, 3)$ -LH systems on $\mathbb{T}^*\mathbf{M}^{1+1}$ studied in [6], where it was shown that h_{ij}^{2D} actually corresponds to a coupling of two oscillator LH systems defined on $\mathbb{T}^*\mathbb{R}$ through the angular momentum term $q_x p_y - q_y p_x$. More precisely, h_{ij}^{2D} (19) can be written as

$$h_{ij}^{2D} = \text{pr}_1^*(h_i^{1D}) - \text{pr}_2^*(h_i^{1D}) - \tilde{a}_j(t)(q_x p_y - q_y p_x) \quad (20)$$

where $\text{pr}_1 : \mathbb{T}^*\mathbf{M}^{1+1} \rightarrow \mathbb{T}^*\mathbb{R}$ and $\text{pr}_2 : \mathbb{T}^*\mathbf{M}^{1+1} \rightarrow \mathbb{T}^*\mathbb{R}$ are the projections of the Whitney sum $\mathbb{T}^*\mathbf{M}^{1+1} = \mathbb{T}^*\mathbb{R} \oplus \mathbb{T}^*\mathbb{R}$ onto the first and second factors, respectively, while h_i^{1D} is the t -dependent Hamiltonian on $\mathbb{T}^*\mathbb{R}$ given by

$$h_i^{1D} = \frac{\tilde{a}_i(t)}{4}(p^2 - 4q^2). \quad (21)$$

Thus, combining the decompositions (15) and (20), we conclude that the $\mathfrak{su}(3)$ -LH system (18) on $\mathbb{T}^*\mathbb{R}^3$ can be interpreted as a coupling of two LH systems on $\mathbb{T}^*\mathbf{M}^{1+1}$ given by the Hamiltonians h_{12}^{2D} and h_{34}^{2D} in (19), with each one of these corresponding to a coupling of two LH systems on $\mathbb{T}^*\mathbb{R}$ with t -dependent Hamiltonians h_1^{1D} , h_2^{1D} and h_3^{1D} , h_4^{1D} obtained from (21).

5 Final remarks

It has been shown that the formal construction of LH systems based on representation theoretical grounds and with underlying VG algebra isomorphic to $\mathfrak{sp}(4, \mathbb{R})$ developed in [6] can be naturally extended to higher rank. In particular, several of the two-dimensional LH systems obtained in that reference can be obtained by restriction of the three-dimensional systems associated to $\mathfrak{sp}(6, \mathbb{R})$, as follows from the canonical embedding chain of symplectic Lie algebras. In particular, the three-dimensional version of time-dependent electromagnetic fields and coupled oscillators, as coupled harmonic oscillators and coupled Caldirola–Kanai systems [7, 12], have been obtained. On the other hand, restricted LH systems related to the subalgebra embedding $\mathfrak{su}(3) \subset \mathfrak{sp}(6, \mathbb{R})$ have been considered, illustrating how the branching rules of (semisimple) Lie algebras can be used to determine new systems associated to distinctive subalgebras. Formally, there is no obstruction to generalize these results to the generic symplectic algebra $\mathfrak{sp}(2n, \mathbb{R})$ for $n > 3$, as well as to relevant semisimple subalgebras of the latter, although the explicit derivation of the corresponding superposition rules is expected to be computationally demanding. Another question to be analyzed in the future is to find suitable conditions on the t -dependent parameters that allow us to determine an explicit solution of the LH system, specially in connection with physically relevant cases, such as the coupled oscillators or systems with electromagnetic fields. Work in these directions are currently in progress.

Acknowledgements

This work has been supported by Agencia Estatal de Investigación (Spain) under grant PID2023-148373NB-I00 funded by MCIN /AEI /10.13039/501100011033 / FEDER, UE. O.C. acknowledges a fellowship (grant C15/23) supported by Universidad Complutense de Madrid and Banco de Santander. F.J.H. acknowledges support by the Q-CAYLE Project funded by the Regional Government of Castilla y León (Junta de Castilla y León, Spain) and by the Spanish Ministry of Science and Innovation (MCIN) through the European Union funds NextGenerationEU (PRTR C17.I1). The authors also acknowledge the contribution of RED2022-134301-T funded by MCIN/AEI/10.13039/501100011033 (Spain).

References

- [1] A. Ballesteros, R. Campoamor-Stursberg, E. Fernández-Saiz, F. J. Herranz and J. de Lucas. Poisson–Hopf deformations of Lie–Hamilton systems revisited: deformed superposition rules and applications to the oscillator algebra. *J. Phys. A: Math. Theor.* **54** (2021) 205202. DOI:[10.1088/1751-8121/abf1db](https://doi.org/10.1088/1751-8121/abf1db)
- [2] A. Ballesteros, J. F. Cariñena, F. J. Herranz, J. de Lucas and C. Sardón. From constants of motion to superposition rules for Lie–Hamilton systems. *J. Phys. A: Math. Theor.* **46** (2013) 285203. DOI:[10.1088/1751-8113/46/28/285203](https://doi.org/10.1088/1751-8113/46/28/285203)
- [3] G. Bluman and J. Cole. *Similarity Methods for Differential Equations*. (New York: Springer Verlag) 1974. DOI:[10.1007/978-1-4612-6394-4](https://doi.org/10.1007/978-1-4612-6394-4)

- [4] R. Campoamor-Stursberg. A new matrix method for the Casimir operators of the Lie algebras $w\mathfrak{sp}(N, \mathbb{R})$ and $I\mathfrak{sp}(2N, \mathbb{R})$. *J. Phys. A: Math. Gen.* **38** (2005) 4187–4208. DOI:[10.1088/0305-4470/38/19/009](https://doi.org/10.1088/0305-4470/38/19/009)
- [5] R. Campoamor-Stursberg. Invariant functions of vector field realizations of Lie algebras and some applications to representation theory and dynamical systems. *J. Phys.: Conf. Ser.* **1071** (2018) 012005. DOI:[10.1088/1742-6596/1071/1/012005](https://doi.org/10.1088/1742-6596/1071/1/012005)
- [6] R. Campoamor-Stursberg, O. Carballal and F. J. Herranz. A representation-theoretical approach to higher-dimensional Lie–Hamilton systems: The symplectic Lie algebra $\mathfrak{sp}(4, \mathbb{R})$. [arXiv:2406.17479](https://arxiv.org/abs/2406.17479) [math-ph].
- [7] P. Caldirola. Forze non conservative nella meccanica quantistica. *Nuovo Cim.* **18** (1941) 393–400. DOI:[10.1007/BF02960144](https://doi.org/10.1007/BF02960144)
- [8] J. F. Cariñena and A. Ramos. Applications of Lie systems in quantum mechanics and control theory. *Banach Cent. Publ.* **59** (2003) 143–162. DOI:[10.4064/bc59-0-7](https://doi.org/10.4064/bc59-0-7)
- [9] H. Georgi. *Lie Algebras in Particle Physics*. (Reading MA: The Benjamin/Cummings Publ. Co., Inc.) 1999.
- [10] B. Gruber and L. O’Raifeartaigh. S Theorem and Construction of the Invariants of the Semisimple Compact Lie Algebras. *J. Math. Phys.* **5** (1964) 1796–1804. DOI:[10.1063/1.1704102](https://doi.org/10.1063/1.1704102)
- [11] M. Havlíček and W. Lassner. Canonical realization of the Lie algebras $\mathfrak{sp}(2n, \mathbb{R})$. *Int. J. Theor. Phys.* **15** (1976) 867–876. DOI:[10.1007/BF01807449](https://doi.org/10.1007/BF01807449)
- [12] E. Kanai. On the Quantization of the Dissipative Systems. *Prog. Theor. Phys.* **3** (1948) 440–442. DOI:[10.1143/ptp/3.4.440](https://doi.org/10.1143/ptp/3.4.440)
- [13] S. Lie. *Vorlesungen über Differentialgleichungen mit Bekannten Infinitesimalen Transformationen*. (Leipzig: B. G. Teubner) 1891.
- [14] H. Lipkin. *Lie groups for pedestrians*. (Amsterdam: North-Holland) 1965.
- [15] J. de Lucas and C. Sardón. *A Guide to Lie Systems with Compatible Geometric Structures*. (Singapore: World Scientific Singapore) 2020. DOI:[10.1142/q0208](https://doi.org/10.1142/q0208)
- [16] W. G. McKay and J. Patera. *Tables of Dimensions, Indices, and Branching Rules for Representations of Simple Lie Algebras*. (New York: Marcel Dekker) 1981.
- [17] P. J. Olver. *Applications of Lie Groups to Differential Equations*. (New York: Springer Verlag) 1985. DOI:[10.1007/978-1-4684-0274-2](https://doi.org/10.1007/978-1-4684-0274-2)
- [18] A. M. Perelomov. *Integrable Systems of Classical Mechanics and Lie Algebras*. (Bassel: Birkhäuser Verlag) 1990. DOI:[10.1007/978-3-0348-9257-5](https://doi.org/10.1007/978-3-0348-9257-5)
- [19] A. M. Perelomov A. and V. S. Popov. Casimir Operators for Semisimple Lie Groups. *Izv. Akad. Nauk SSSR* **32** (1968) 1368–1390. DOI:[10.1070/IM1968v002n06ABEH000731](https://doi.org/10.1070/IM1968v002n06ABEH000731)

- [20] M. F. Rañada and M. Santander. Complex Euclidean Super-Integrable Potentials, Potentials of Drach, and Potentials of Holt. *Phys. Lett. A* **278** (2001) 271–279. DOI:[10.1016/S0375-9601\(00\)00782-9](https://doi.org/10.1016/S0375-9601(00)00782-9)
- [21] P. Winternitz. *Lie groups and solutions of nonlinear differential equations*. In *Nonlinear Phenomena*, K. B. Wolf (Ed.), Lect. Notes Phys. **189**, Springer, Berlin 1983, pp. 263–331. DOI:[10.1007/3-540-12730-5_12](https://doi.org/10.1007/3-540-12730-5_12)