

EXPLICIT CONSTRUCTION OF RECURRENT NEURAL NETWORKS EFFECTIVELY APPROXIMATING DISCRETE DYNAMICAL SYSTEMS

CHIKARA NAKAYAMA AND TSUYOSHI YONEDA

ABSTRACT. We consider arbitrary bounded discrete time series originating from dynamical system with recursivity. More precisely, we provide an explicit construction of recurrent neural networks which effectively approximate the corresponding discrete dynamical systems.

1. INTRODUCTION

In this paper we provide an explicit construction of recurrent neural networks (RNNs) which effectively approximate discrete dynamical systems. In general, Lyapunov indexes for reconstructed and original systems have a gap (see Dechert-Gençay [2] and Berry-Das [1]), and our explicit construction of RNNs at least minimize the maximal Lyapunov exponent. As far as the authors are aware, none of the numerous works to date attempted to explicitly construct RNNs in terms of an elementary algebraic approach, and in this paper we initiate it (for investigating the universality of periodic points based on the similar algebraic approach, see [4]).

First we define the dynamical system in a delay coordinate. For the dimension of the dynamical system $L \in \mathbb{Z}_{\geq 1}$, let $\Phi : [-1, 1]^L \rightarrow [-1, 1]^L$ $((w_1, w_2, \dots, w_L)^T \mapsto (x_1, x_2, \dots, x_L)^T)$ be such that

$$(1) \quad w_1 = x_2, \quad w_2 = x_3, \dots, w_{L-1} = x_L$$

(i.e. discrete dynamical system in a delay coordinate, see [5], and see also [3]). Let us assume that a time series $y : \mathbb{Z} \rightarrow [-1, 1]$ satisfies

$$Y_{t+1} = \Phi(Y_t) \quad \text{for } t \in \mathbb{Z},$$

where

$$Y_t := (y(t), y(t-1), \dots, y(t-L+1))^T.$$

From this Φ , we define $\lambda \in \mathbb{R}$ to be the maximum Lyapunov exponent, and assume it is finite, as follows:

$$e^\lambda := \sup_{\substack{w, w' \in [-1, 1]^L, \\ w \neq w'}} \frac{|\Phi(w) - \Phi(w')|}{|w - w'|} < \infty.$$

Also we assume recursivity to y : For any $\epsilon > 0$, there is $\{t_j\}_{j=1}^\infty \subset \mathbb{Z}$ ($t_j \rightarrow -\infty$ for $j \rightarrow \infty$) such that

$$|Y_0 - Y_{t_j}| < \epsilon.$$

2020 *Mathematics Subject Classification.* Primary 37N30; Secondary 11B50.

Key words and phrases. recurrent neural networks, discrete dynamical systems, key-value pairs, topological ring.

Now we define the conventional RNNs. For initial hidden state vector $r(0) \in \mathbb{R}^N$, initial data $\hat{y}(0) \in [-1, 1]$, recurrent weight matrix $W \in \mathbb{R}^{N \times N}$, input weight matrix (vector) $W^{in} \in \mathbb{R}^{N \times 1}$, output weight matrix (vector) $W^{out} \in \mathbb{R}^{1 \times N}$ and activate function $h : \mathbb{R} \rightarrow [-1, 1]$, the RNNs can be expressed as follows:

$$(2) \quad \begin{cases} r(t+1) & := h(Wr(t) + W^{in}\hat{y}(t)), \\ \hat{y}(t+1) & := W^{out}r(t+1) \quad \text{for } t \geq 0. \end{cases}$$

The main theorem is as follows:

Theorem 1. *Let $y : \mathbb{Z} \rightarrow [-1, 1]$ be a time series generated by the discrete dynamical system in a delay coordinate $\Phi : [-1, 1]^L \rightarrow [-1, 1]^L$ ($Y_{t-1} \mapsto Y_t$) with recursivity and a finite $\lambda \geq 0$. Then for any large $K \in \mathbb{Z}_{\geq 1}$ and for any constant C which is slightly larger than 1, there exist $N \leq K^L$, h , explicit W , W^{in} and W^{out} which are composed of $y(t)$ ($t \in \mathbb{Z}_{\leq 0}$), an initial hidden state vector $r(0)$, and initial data $\hat{y}(0)$ such that the following holds:*

$$(3) \quad |\hat{y}(t) - y(t)| \leq (2t+1)e^{\lambda t} \frac{\sqrt{LC}}{K} \quad \text{for } t \geq 0.$$

This is only on the order of log worse (since $t = e^{\log t}$). Thus, this RNNs construction seems effective for re-expressing the original system.

2. PROOF OF MAIN THEOREM

First let us discretize the range $[-1, 1]$ as follows: For $K \in \mathbb{Z}_{\geq 1}$, we choose $\{a_k^K\}_{k=1}^K \subset [-1, 1]$ such that

- $-1 < a_1^K < a_2^K < \dots < a_K^K < 1$,
- $a_i^K \neq 0$ for any $1 \leq i \leq K$ and
- $\sum_{\ell=1}^L \frac{a_{k_\ell}^{k'_\ell}}{a_{k_\ell}^K} \neq L$ for each $(k_1, \dots, k_L) \neq (k'_1, \dots, k'_L)$,
- $\sup \left\{ \frac{a_{k+1}^K - a_{k-1}^K}{2} \ (1 < k < K), \frac{a_1^K + a_2^K}{2} + 1, 1 - \frac{a_{K-1}^K + a_K^K}{2} \right\} \leq C/K$
- For any t with $-L+1 \leq t \leq 0$ and for any k with $1 \leq k < K$,

$$(4) \quad y(t) \neq \frac{a_k^K + a_{k+1}^K}{2}.$$

We need the last condition for appropriately separating the time series y into training and inference phases. Note that the above second condition is almost always satisfied in the following sense.

Proposition 2. *There is a closed set Z of $[-1, 1]^K$ whose Lebesgue measure is zero such that the above second condition is satisfied as soon as (a_1^K, \dots, a_K^K) is not in Z .*

Proof. We drop the superscript K for simplicity. Regard a_i ($1 \leq i \leq K$) as indeterminates. Clearly the measure of the set where $a_i = 0$ is zero. Under the assumption that any a_i is not zero, the condition $\sum_{\ell=1}^L \frac{a_{k'_\ell}}{a_{k_\ell}} \neq L$ is equivalent to $\sum_{\ell} a_{k_1} \dots a_{k'_\ell} \dots a_{k_L} - L \prod_{\ell} a_{k_\ell} \neq 0$, the left-hand-side of which is a nonzero polynomial whenever $k_\ell \neq k'_\ell$ for some ℓ (then the term $\prod_{\ell} a_{k_\ell}$ does not vanish). Hence by the following simple fact, the non-equality holds for almost all values of a_i :

- Let f be a nonzero polynomial in M variables. Then the Lebesgue measure of the zero point set $\{x \in \mathbb{R}^M : f(x) = 0\}$ is zero.

□

We now discretize y (in the training phase) as follows:

$$(5) \quad \bar{y}_K(t) := \arg \min_{a \in \{a_k^K\}_{k=1}^K} |y(t) - (a - 0)| \quad \text{for } t \in \mathbb{Z}_{\leq 0},$$

where $a - 0 := a - \varepsilon$ for any sufficiently small $\varepsilon > 0$. From this discretization, we define “dictionary”, in other word, “key-value pairs”. Let σ_n ($n = 1, 2, \dots, N$) be a permutation operator, namely, a map

$$\sigma_n^K : \{1, 2, \dots, L\} \rightarrow \{a_1^K, a_2^K, \dots, a_{K-1}^K, a_K^K\}$$

($\ell \mapsto \sigma_n^K(\ell)$) with $\sigma_n^K \neq \sigma_{n'}^K$ ($n \neq n'$), and we impose the following two conditions for determining N :

$$(6) \quad \begin{cases} \text{For any } t \in \mathbb{Z}_{\leq 0}, \text{ there is } n \in \{1, \dots, N\} \text{ such that} \\ \quad \sigma_n^K(\ell) = \bar{y}_K(t - \ell) \text{ for } \ell = 1, 2, \dots, L, \\ \text{For any } n \in \{1, \dots, N\} \text{ there is } t \in \mathbb{Z}_{\leq 0} \text{ such that} \\ \quad \sigma_n^K(\ell) = \bar{y}_K(t - \ell) \text{ for } \ell = 1, 2, \dots, L. \end{cases}$$

Note that $N \leq K^L$ due to the sequence with repetition. We now define the dictionary $\{(\sigma_n, a_{k(n)})\}_{n=1}^N$, in other word, key-value pairs (i.e. σ_n is the “key” and $a_{k(n)}$ is the corresponding “value”). By (6), for any key σ_n^K , there exists a $t' \in \mathbb{Z}_{\leq 0}$ and then we choose such a t' and the corresponding $k(n) \in \{1, 2, \dots, K\}$ such that

$$(7) \quad \text{key: } \sigma_n^K(\ell) = \bar{y}_K(t' - \ell) \quad (\ell = 1, 2, \dots, L) \quad \text{and} \quad \text{value: } a_{k(n)}^K := \bar{y}_K(t').$$

Now, by using these key-value pairs, we first construct a time series $y^*(t)$ ($t \geq 0$), and then show an estimation analogous to (3), that is, the inequality obtained by replacing \hat{y} in (3) by y^* . After that, we give $h, W, W^{in}, W^{out}, r(0)$ and $\hat{y}(0)$ such that $\hat{y} = y^*$, which completes the proof.

We define $y^*(t)$ by induction. Let $y^*(t) = \bar{y}_K(t)$ for $t = 0, -1, \dots, -L$. Then the permutation operator $\ell \mapsto y^*(1 - \ell)$ is in the dictionary by recursivity and (4). Let $t_0 > 0$. Assume that we have defined $y^*(t)$ for $-L \leq t < t_0$ and assume that we can find the permutation operator $\ell \mapsto y^*(t_0 - \ell)$ is in the dictionary, that is, it is σ_n for some n . Define $y^*(t_0) := a_{k(n)}$. Then we see that the permutation operator $\ell \mapsto y^*(t_0 - \ell + 1)$ also in the dictionary as follows. Recall that for some $t' \leq 0$, $\sigma_n(\ell) = \bar{y}_K(t' - \ell)$ ($\ell = 1, 2, \dots, L$) and $a_{k(n)} = \bar{y}_K(t')$. Hence the permutation operator concerned coincides with $\ell \mapsto \bar{y}_K(t' + 1 - \ell)$. If $t' < 0$, there is \tilde{n} such that $\bar{y}_K(t' - \ell + 1) = \sigma_{\tilde{n}}(\ell)$ ($\ell = 1, 2, \dots, L$) by (6). If $t' = 0$, we just apply recursivity and (4). Thus the induction goes and we have defined y^* .

We estimate the difference of $y^*(t)$ and $y(t)$, which eventually implies (3). The case $t = 0$ is by the third condition in the beginning of this section. Let

$$\begin{aligned} \bar{Y}_t &:= (\bar{y}_K(t), \bar{y}_K(t-1), \dots, \bar{y}_K(t-L+1))^T \quad (t \leq 0), \\ Y_t^* &:= (y^*(t), y^*(t-1), \dots, y^*(t-L+1))^T \quad (t \geq 0). \end{aligned}$$

Then we see that, for any $t \in \mathbb{Z}_{\geq 1}$, there exists $t' \in \mathbb{Z}_{\leq 0}$ such that

$$(8) \quad \begin{cases} \bar{Y}_{t'} = Y_t^*, & |Y_{t'} - Y_t^*| \leq \frac{\sqrt{LC}}{K}, \\ Y_{t'} = \Phi(Y_{t'-1}), \\ \bar{Y}_{t'-1} = Y_{t-1}^*, & |Y_{t'-1} - Y_{t-1}^*| \leq \frac{\sqrt{LC}}{K}. \end{cases}$$

Recall $e^\lambda \geq 1$, and then we have

$$\begin{aligned} |y^*(t) - y(t)| &\leq |Y_{t'}^* - Y_t| \\ &\leq |Y_{t'} - Y_t| + \frac{\sqrt{LC}}{K} \\ &\leq e^\lambda |Y_{t'-1} - Y_{t-1}| + \frac{\sqrt{LC}}{K} \\ &\leq e^\lambda |Y_{t-1}^* - Y_{t-1}| + e^\lambda \frac{\sqrt{LC}}{K} + \frac{\sqrt{LC}}{K} \\ &\leq e^\lambda |Y_{t-1}^* - Y_{t-1}| + 2e^\lambda \frac{\sqrt{LC}}{K} \leq \dots \\ &\leq e^{\lambda t} |Y_0^* - Y_0| + 2te^{\lambda t} \frac{\sqrt{LC}}{K} \\ &\leq (2t+1)e^{\lambda t} \frac{\sqrt{LC}}{K}. \end{aligned}$$

Next we explicitly construct effective recurrent neural networks. Let σ_n^* ($n = 1, 2, \dots, N$) be an adjoint type of permutation operator, more precisely, let

$$(9) \quad \sigma_n^*(\ell - 1) := \frac{1}{\sigma_n(\ell)}$$

for $\ell \in \{1, \dots, L\}$ and $n \in \{1, \dots, N\}$. The definition of the following $N \times N$ matrix X is the crucial to prove the main theorem:

$$X := \left[h \left(\sum_{\ell=1}^L \sigma_i^*(\ell - 1) \sigma_j(\ell) \right) \right]_{i,j}$$

for some $h : \mathbb{R} \rightarrow [-1, 1]$. Also let G be such that

$$G := \left\{ \sum_{\ell=1}^L \sigma_i^*(\ell - 1) \sigma_j(\ell) : i, j \in \{1, 2, \dots, N\} \right\}.$$

Note that $L \in G$.

Lemma 3. *X is a regular matrix for almost all $h : \mathbb{R} \rightarrow [-1, 1]$ in the following sense. Let $\gamma_1, \dots, \gamma_M$ be the all elements of G . Then there is a closed set Z of \mathbb{R}^M whose Lebesgue measure is zero such that X is regular as soon as $(h(\gamma_1), \dots, h(\gamma_M))$ is not in Z .*

Proof. Take a bijection $x : \gamma_m \mapsto x_m$ from G to the set of indeterminates $\{x_1, \dots, x_M\}$. Consider the determinant D of the matrix $[x(\sum_{\ell=1}^L \sigma_i^*(\ell - 1) \sigma_j(\ell))]_{i,j \in N}$ with polynomial coefficients. Then D is a nonzero polynomial because it is a monic polynomial of degree N with respect to the indeterminate $x(L)$ (the leading term $x(L)^N$ comes from the diagonal). Now we apply the simple fact in the proof of Proposition 2 to

D . Let Z be the zero point set of D . Assume that $(h(\gamma_1), \dots, h(\gamma_M))$ is not in Z . Then $\det X$ defined by h is not zero so that X is regular. \square

With the aid of the inverse of X , we can provide the $N \times N$ recurrent weight matrix W as follows:

$$(10) \quad W := YX^{-1} \quad \text{with} \quad Y := \left[\sum_{\ell=1}^{L-1} \sigma_i^*(\ell) \sigma_j(\ell) \right]_{i,j}.$$

Remark 1. The rank of W is less than $L - 1$, because Y is the product of the $N \times (L - 1)$ -matrix: $[\sigma_i^*(\ell - 1)]_{i,\ell}$ and $(L - 1) \times N$ -matrix: $[\sigma_j(\ell)]_{\ell,j}$.

We now define the input weight vector W^{in} and the the output weight matrix (vector) W^{out} as follows:

$$\begin{aligned} W^{in} &:= \sigma^*(0), \\ W^{out} &:= (a_{k(1)}, a_{k(2)}, \dots, a_{k(N)})X^{-1}, \end{aligned}$$

where $\sigma^*(\ell) := (\sigma_1^*(\ell), \sigma_2^*(\ell), \dots, \sigma_N^*(\ell))^T$ for $0 \leq \ell \leq L - 1$. Let us set the initial hidden state vector $r(0)$ and the initial data $\hat{y}(0)$ as follows:

$$\begin{aligned} r(0) &= \left[h \left(\sum_{\ell=1}^L \sigma_i^*(\ell - 1) \bar{y}_K(-\ell) \right) \right]_i, \\ \hat{y}(0) &:= \bar{y}_K(0), \end{aligned}$$

then we have the desired RNNs. Let \hat{y} be the time series generated by these initial data. Now we certify $\hat{y} = y^*$, which completes the proof of Theorem. For $t \geq 0$, the permutation operator $\ell \mapsto y^*(t - \ell)$ is σ_n for some n . Let n_t be this n . Then we define the column vector

$$e_{n_t} := \underbrace{(0, 0, \dots, 0, 1, 0 \dots, 0)}_{N}^T.$$

We prove $\hat{y}(t) = y^*(t)$ for $t \geq 0$ together with $r(t) = X e_{n_t}$ by induction. The case $t = 0$ is by the definition. Assume that $t \geq 0$ and they hold for $0, 1, \dots, t$. Then we have

$$\text{key: } y^*(t - \ell) = \sigma_{n_t}(\ell) \quad \text{for } \ell = 1, 2, \dots, L.$$

Thus we have

$$\begin{aligned} W r(t) &= W X e_{n_t} = Y e_{n_t} = \sum_{\ell=1}^{L-1} \sigma^*(\ell) y^*(t - \ell), \\ r(t + 1) &= h(Y e_{n_t} + \sigma^*(0) y^*(t)) = X e_{n_{t+1}} \quad \text{and} \end{aligned}$$

$$\hat{y}(t + 1) = W^{out} r(t + 1) = (a_{k(1)}, \dots, a_{k(N)}) e_{n_{t+1}} = a_{k(n_{t+1})} = y^*(t + 1).$$

Hence they hold also for $t + 1$.

Acknowledgments. The first author thanks J. C. for leading him to join this work. Research of TY was partly supported by the JSPS Grants-in-Aid for Scientific Research 24H00186. Research of CN was partly supported by the JSPS Grants-in-Aid for Scientific Research 21K03199.

REFERENCES

- [1] T. Berry and S. Das, *Learning theory for dynamical systems*, SIAM J. Appl. Dyn. Syst., **22**, (2023) 10.1137/22M1516865.
- [2] W. Dechert and R. Gençay, *The topological invariance of lyapunov exponents in embedded dynamics*, Phys. D, **90** (1996) 40–55.
- [3] K. Nakai and Y. Saiki, *Machine-learning construction of a model for a macroscopic fluid variable using the delay-coordinate of a scalar observable*, Discr. Conti. Dyn. Sys.-S, **14**, (2021) 1079-1092.
- [4] C. Nakayama and T. Yoneda, *Universality of almost periodicity in bounded discrete time series*, arXiv:2310.00290.
- [5] F. Takens, *Detecting strange attractors in turbulence*, Dynamical Systems of Turbulence, Lecture Notes in Mathematics, **898**, Springer-Verlag, Berlin, (1981) 366-381.

GRADUATE SCHOOL OF ECONOMICS, HITOTSUBASHI UNIVERSITY, 2-1 NAKA, KUNITACHI, TOKYO
186-8601, JAPAN

Email address: `c.nakayama@r.hit-u.ac.jp`

GRADUATE SCHOOL OF ECONOMICS, HITOTSUBASHI UNIVERSITY, 2-1 NAKA, KUNITACHI, TOKYO
186-8601, JAPAN

Email address: `t.yoneda@r.hit-u.ac.jp`