ISOMETRIC ACTIONS OF COMPACT QUANTUM GROUPS ON GRAPH *C* ∗ **-ALGEBRAS**

SOUMALYA JOARDAR AND JITENDER SHARMA

ABSTRACT. For a strongly connected directed graph Λ , it is shown that the quantum automorphism group of Λ acts isometrically in the sense of D. Goswami et al. ([\[3\]](#page-18-0)) on the spectral triple $(C^*(\Lambda), L^2(\Lambda^\infty, M), D)$ constructed by Farsi et al.. It is also shown that the natural action of U_n^+ is not isometric on the Cuntz algebra \mathcal{O}_n .

1. INTRODUCTION

Given a mathematical structure, to understand its symmetry is a fundamental problem. It often helps in reducing the complexity of a system exhibiting a given mathematical structure. Classically such symmetries are modelled by groups. For example, given a Riemannian manifold *M*, a symmetry of the manifold should be given by some group which acts on *M* preserving the Riemannian metric. In the realm of noncommutative geometry or topology, a 'compact' space is typically replaced by a unital C^{*}-algebra or von Neumann algebra. Then it becomes very important to understand 'generalized symmetry' of such algebras. By now it has become standard to replace groups by what are known as compact quantum groups (CQG's in short). Since Wang in his seminal paper (see [\[25\]](#page-20-0)) showed the existence of a genuine quantum symmetry for a space consisting of finite points (at least 4 points), CQG's have fulfilled its role as quantum symmetry object in the realm of noncommutative geometry very successfully. To study more geometric properties, A. Connes introduced the concept of a spectral triple on *C* ∗ -algebras. A spectral triple is a triple (A, \mathcal{H}, D) where typically $\mathcal A$ is a C^* -algebra represented faithfully on a Hilbert space $\mathcal H$ and D is some unbounded operator, called the Dirac operator, capturing the 'geometric' data. Quite naturally one seeks for a notion of an 'isometric' action of CQG's on such spectral triples which should be a generalization of classical isometric group actions on Riemannian manifolds. This is done in a series of papers by D. Goswami and his collaborators (see [\[3\]](#page-18-0), [\[10\]](#page-19-0), [\[11\]](#page-19-1) for example). In these papers, the authors develop a notion of a 'quantum isometric' action of CQG's on spectral triples both in terms of the Dirac operator of the spectral triple and in terms of a suitable Laplacian operator derived from the spectral triple data. A very important and tractable class of *C* ∗ -algebras are given by what are known as graph C^* -algebras. As the name hints, these C^* -algebras are built from directed graphs. Due to the underlying combinatorial structure of the graph, this class has

²⁰²⁰ *Mathematics Subject Classification.* 46L89, 81R50, 81R60.

Key words and phrases. Compact qauntum group, quantum isometric action, spectral triple, graph *C*[∗] -algebras.

2 JOARDAR AND SHARMA

become very well studied class of C^* -algebras (see [\[23\]](#page-19-2),[\[18\]](#page-19-3)). This underlying combinatorial data has also been useful in studying quantum symmetry of the graph C^{*}-algebras. For example, the quantum symmetry of the underlying graph naturally acts on the corresponding graph C^* -algebra (see [\[24\]](#page-19-4), [\[14\]](#page-19-5), [\[15\]](#page-19-6), [\[16\]](#page-19-7), [\[17\]](#page-19-8), [\[2\]](#page-18-1) for quantum group actions on graph C^* -algebras). Recently, a spectral triple has been introduced on the graph C^* -algebras (see [\[5\]](#page-19-9), [\[7\]](#page-19-10)). In these spectral triples, the graph C^* -algebras corresponding to strongly connected finite directed graphs are faithfully represented on an L^2 -space of infinite path spaces completed with respect to a measure derived from its unique KMS-state. So it becomes quite natural to understand 'quantum isometric' actions on such spectral triples. In this paper, we show that given a strongly connected finite graph Λ without multiple edge, loop or source and a spectral triple on the corresponding graph *C* ∗ -algebra, the natural action of the quantum automorphism group of the underlying graph (to be denoted by Aut⁺(Λ)) is 'isometric' in the sense of Goswami et al.([\[3\]](#page-18-0)). The essential step is to construct a unitary co-representation \tilde{U} of the CQG on the L^2 space of infinite paths that can implement the given action. To prove the unitarity of \bar{U} , the invariance of the unique KMS-state under the action of the $\text{Aut}^+(\Lambda)$ becomes important. That leads us naturally to consider the universal object in the category of CQG's acting linearly and preserving the KMS state on a graph C^* -algebra. In case of the Cuntz algebra with *n*-generators, it is well known that the quantum unitary group U_n^+ is the universal object in that category. We show that the action of U_n^+ is not isometric however. This makes $Aut^+(\Lambda)$ a natural general class of quantum groups that can act 'quantum isometrically' on the corresponding graph C ^{*}-algebras. Note that although isometric group action has been studied on Cuntz algebras before (see $[6, 19]$ $[6, 19]$), the study of quantum isometries for a general class of graph C^* -algebras is certainly new and we hope that this work will give impetus to understand quantum isometries of graph C^* -algebras.

Notations: In this paper, we will use some standard notations. Sp{*A*} will stand for the linear span of elements of *A* in a given vector space; $\mathcal{H}\overline{\otimes}\mathcal{A}$ will refer to the external tensor product of Hilbert space with a C^* -algebra to form a Hilbert- $\mathcal A$ module. We shall use $\mathcal{B}(\mathcal{H}), \mathcal{K}(\mathcal{H})$ to denote the set of bounded and compact linear operators on a Hilbert space H respectively. If $\mathcal E$ is a Hilbert $\mathcal A$ -module then $\mathcal L(\mathcal E)$ is the *C* ∗ -algebra of adjointable A-linear maps. ⊗ will denote the minimal tensor product between C^* -algebras as well as tensor product of Hilbert spaces whereas ⊗_{alg} will stand for algebraic tensor product. We use \langle , \rangle for both the *C*^{*}-algebra valued inner product on a Hilbert module and a C-valued inner product on a Hilbert space. For a C^* -algebra $A, M(A)$ will denote its multiplier algebra.

2. Preliminaries

2.1. **Compact quantum groups.** In this section we shall recall the preliminaries on compact quantum groups and their actions on C^* -algebras. The reader is referred to any of [\[26\]](#page-20-1), [\[20\]](#page-19-13) for a thorough exposition on compact quantum groups. For generalities on actions of CQG on C^* -algebras the reader can consult [\[4\]](#page-18-2).

Definition 2.2. A compact quantum group \mathbb{G} is a pair $(C(\mathbb{G}), \Delta)$ where $C(\mathbb{G})$ is a unital C^* -algebra and $\Delta : C(\mathbb{G}) \to C(\mathbb{G}) \otimes C(\mathbb{G})$ is a unital C^* -homomorphism which satisfies the following conditions:

- (I) $(id \otimes \Delta) \circ \Delta = (\Delta \otimes id) \circ \Delta$
- (II) $\text{Sp}\{\Delta(C(\mathbb{G}))(1 \otimes C(\mathbb{G}))\}$ and $\text{Sp}\{\Delta(C(\mathbb{G}))(C(\mathbb{G}) \otimes 1)\}\$ are dense in $C(\mathbb{G}) \otimes$ $C(\mathbb{G}).$

Given a CQG \mathbb{G} , there is a canonical dense Hopf $*$ -algebra $C(\mathbb{G})_0$ of $C(\mathbb{G})$ where the coproduct is algebraic i.e. $\Delta(C(\mathbb{G})_0) \subset C(\mathbb{G})_0 \otimes_{\text{alg}} C(\mathbb{G})_0$. Given any CQG \mathbb{G} , $C(\mathbb{G})_0$ will denote this canonical dense Hopf \ast -algebra.

Example 2.3 (Quantum permutation group). Consider the universal C^* -algebra generated by a magic unitary of order *n*, denoted by $C(S_n^+)$ i.e.

$$
C(S_n^+) := C^* \left(q_{ij}, 1 \le i, j \le n \mid q_{ij}^2 = q_{ij} = q_{ij}^* \text{ and } \sum_k q_{ik} = \sum_k q_{kj} = 1 \right).
$$

With the coproduct map on the generators of $C(S_n^+)$ given by $\Delta(q_{ij}) = \sum_k q_{ik} \otimes q_{kj}$, $(C(S_n^+), ∆)$ becomes a compact quantum group. It is denoted by S_n^+ and called the quantum permutation group on *n*-points.

Example 2.4 (Quantum unitary group). Consider the universal C^* -algebra generated by unitary of order *n*, denoted by $C(U_n^+)$, that is,

$$
C(U_n^+) := C^* \left(q_{ij}, 1 \le i, j \le n | q = (q_{ij}) \text{ and } \overline{q} = (q_{ij}^*) \text{ are unitaries} \right).
$$

The co-product map on the generators given by $\Delta(q_{ij}) = \sum_{k} q_{ik} \otimes q_{kj}$. Then $(C(U_n^+), \Delta)$ is a compact quantum group. This is called the quantum unitary group and is denoted by U_n^+ .

Example 2.5 (Qaut of a finite directed graph). (see [\[1\]](#page-18-3)) Let $\Lambda = (V, E, r, s)$ be a finite directed graph without loops or multiple edges, where *V* and *E* are the sets of vertices and edges respectively and *r, s* are the standard range and source maps. The quantum automorphism group of Λ is the pair $(C(\text{Aut}^+(\Lambda)), \Delta)$ where $C(\text{Aut}^+(\Lambda))$ is the universal *C*^{*}-algebra generated by the elements $\{q_{ij}\}_{i,j\in V}$ such that the matrix $U = ((q_{ij}))_{i,j \in V}$ is a magic unitary of order |*V*| and $UA = AU$ where *A* is the adjacency matrix of Λ . The co-product Δ is given on the generators by $\Delta(q_{ij}) = \sum_k q_{ik} \otimes q_{kj}$. One can write down all the relations between the generators explicitly. In particular, we note down the following relations for future purpose (see [\[9\]](#page-19-14))

(2.1)
$$
q_{s(\gamma)i}q_{r(\gamma)k} = q_{r(\gamma)k}q_{s(\gamma)i} = 0, \ \gamma \in E, \ (i,k) \notin E
$$

$$
q_{is(\gamma)}q_{kr(\gamma)} = q_{kr(\gamma)}q_{is(\gamma)} = 0, \ \gamma \in E, \ (i,k) \notin E.
$$

Definition 2.6. Given a unital C^{*}-algebra A, a compact quantum group **G** is said to act on A if there is a unital C^* -homomorphism α : $\mathcal{A} \to \mathcal{A} \otimes C(\mathbb{G})$ such that

- (I) $(id \otimes \Delta) \circ \alpha = (\alpha \otimes id) \circ \alpha$.
- (II) $\text{Sp}\{\alpha(\mathcal{A})(1 \otimes C(\mathbb{G}))\}$ is dense in $\mathcal{A} \otimes C(\mathbb{G})$.

Moreover, a state ϕ on a unital *C*^{*}-algebra *A* is said to be *invariant* under α if $(\phi \otimes id) \circ \alpha(a) = \phi(a)$. for all $a \in \mathcal{A}$.

Now let us recall the rudiments of graph C^* -algebras and in particular the actions of the quantum automorphism groups of the underlying graphs on them. This is going to be the central object of this paper. The reader is referred to [\[23\]](#page-19-2) and [\[14,](#page-19-5) [24\]](#page-19-4) for more details on this topic. Let $\Lambda = (V, E, r, s)$ be a strongly connected finite directed graph without multiple edges, loops and sources. Recall that by a strongly connected graph, we mean a graph where any two vertices can be joined by a path of finite length.

We remark that if $\Lambda = (V, E, r, s)$ is a finite directed graph without loops and multiple edges, we say two edges *e*, f are composable if $s(e) = r(f)$ and write it as *ef*. More generally, for two paths λ, μ , we say λ, μ are composable if $s(\lambda) = r(\mu)$ and write the composed path as $\lambda \mu$ (see [\[7\]](#page-19-10)). Also note that the use of the term 'loop' is different from that in [\[7\]](#page-19-10) in the context of higher rank graph. In [\[7\]](#page-19-10), a loop is a path λ of finite length such that $r(\lambda) = s(\lambda)$. However, we use the term 'loop' to represent an edge *e* such that $r(e) = s(e)$. In this paper we call a path λ of length greater than 1 such that $r(\lambda) = s(\lambda)$ a closed path. Thus for us a closed path of length 1 is a loop. Given a finite path λ , we denote the length of the path by $d(\lambda)$.

Definition 2.7. Let $\Lambda = (V, E, r, s)$ be a row finite directed graph consisting of a countable vertex set *V*, countable edge set *E* and the maps $r, s : E \to V$ identifying the range and source of each edge, then $C^*(\Lambda)$ is defined as the universal C^* algebra generated by mutually orthogonal projections $\{p_v, v \in V\}$ and partial isometries ${S_e, e \in E}$ that satisfy,

(I) $S_e^* S_e = p_{s(e)}$ for all $e \in E$; and

(II)
$$
p_v = \sum_{\{e \in E, r(e) = v\}} S_e S_e^*
$$
.

It follows immediately that for a finite graph Λ , $\sum_{v} p_v = 1$ and therefore the graph C^* -algebra $C^*(\Lambda)$ is unital. In this paper Λ will always be finite. For a finite directed graph $\Lambda = (V, E, r, s)$ without loops and multiple edges and sources, let $V = \{1, 2, \dots, n\}$ and $E = \{e_1, e_2, \dots, e_m\}$. For the proof of the following proposition, the reader is referred to [\[14,](#page-19-5) [24\]](#page-19-4).

Proposition 2.8. *Let* $\Lambda = (V, E, r, s)$ *be a finite directed graph without loops or multiple edges or sources and* $C^*(\Lambda)$ *be the corresponding graph* C^* *algebra. Then the quantum automorphism group* $Aut^+(\Lambda)$ *acts on* $C^*(\Lambda)$ *. The action* $\alpha: C^*(\Lambda) \to$ $C^*(\Lambda) \otimes C(Aut^+(\Lambda))$ *is given by,*

$$
\alpha(p_i) = \sum_{k=1}^n p_k \otimes q_{ki} \quad 1 \le i \le n
$$

and

$$
\alpha(S_{e_j}) = \sum_{l=1}^m S_{e_l} \otimes q_{r(e_l)r(e_j)}q_{s(e_l)s(e_j)} \quad 1 \leq j \leq m.
$$

Remark 2.9*.* The above action is slightly different form the action defined by Schmidt and Weber. The difference lies in the coefficients of S_{e_l} . In [\[24\]](#page-19-4), the coefficients are taken as ${}^{\alpha}q_{s(e_j)s(e_l)}q_{r(e_j)r(e_l)}$. This difference arises from the fact that we are using slightly different convention for composability of paths.

There is a very important class of states on graph C^* -algebras known as the KMS states. We are not going to recall the basics of KMS states. The reader is referred to [\[13\]](#page-19-15) for details on KMS states on graph *C* ∗ -algebras. It is well known that for a strongly connected graph Λ , the graph C^* -algebra $C^*(\Lambda)$ has a unique KMS state at the critical inverse temperature. If we denote the unique KMS state by ϕ , then ϕ is given on the vertex projections by $\phi(p_v) = x_v^{\Lambda}$, where $\{x_v^{\Lambda}\}_{v \in V}$ is the normalized Perron-Frobenius vector of the vertex matrix (see [\[8\]](#page-19-16)). Moreover ϕ is invariant under the action of $\text{Aut}^+(\Lambda)$ (Theorem 3.1 of [\[15\]](#page-19-6)).

2.10. **Spectral triples and isometric actions.** We start this subsection by recalling the definition of a spectral triple.

Definition 2.11. A spectral triple (A, H, D) consists of $*$ -algebra A with a faithful representation

 $\pi : A \rightarrow B(H)$

where H is a Hilbert space and a densely defined operator $D(p$ ossibly unbounded) called the *Dirac operator* which satisfies the following conditions:

- (I) *D* is self-adjoint;
- (II) For all $c \in \mathbb{R}$ in the resolvent, $(D c)^{-1} \in \mathcal{K}(\mathcal{H})$;
- (III) For all $a \in \mathcal{A}, [D, a]$ is a bounded operator on \mathcal{H} .

Now let us recall the spectral triple introduced in [\[7\]](#page-19-10) for a strongly connected finite *k*-graph very briskly. We are going to specialize the construction for $k = 1$. To that end, let $\Lambda = (V, E, r, s)$ be a finite directed graph without sources. We remark that for the spectral triple discussion we don't need to restrict the class of graphs without multiple edges or loops. Let $\Lambda^* = \bigcup_{n\geq 0} E^n$ be the collection of all finite paths of Λ with range and source maps extended accordingly. This along with the usual length function $d: E^* \to \mathbb{N}$, forms a *k*-graph with $k = 1$. We shall consider the set of all infinite paths,

 $\Lambda^{\infty} = {\lambda = (\lambda_1 \lambda_2 \cdots) | r(\lambda_{i+1}) = s(\lambda_i), i = 1, 2, \cdots; \lambda_1 \lambda_2 \cdots$ is an infinite sequence}

This infinite path space Λ^{∞} being a subset of $\prod E$ has the natural subspace topology from product space for which the *cylinder sets* ,

 $[\lambda] = \{x \in \Lambda^{\infty} | \lambda = x_1 x_2 \cdots x_{d(\lambda)}\}$

parametrised by finite length paths λ form a basis of open set in the subspace topology.

Remark 2.12 ([\[22\]](#page-19-17))*.* Generally speaking, infinite path space for higher rank graph is given by the set of all degree-preserving morphisms from Ω_k to Λ , where Λ is a *k*-graph. However, for a 1-graph, the situation is simpler. Using the *unique* *factorisation property*, given a degree preserving morphism $x : \Omega_1 \to \Lambda$, we can uniquely identify *x* with an infinite path

$$
\lambda = x(0,1)x(1,2)x(2,3)x(3,4)\cdots
$$

vice versa. As a result, the typical endless path and the infinite path provided by the morphism for the 1-graph may be identified with each other.

For the following proposition, see [\[7\]](#page-19-10).

Proposition 2.13. *Let* Λ *be a strongly connected finite* 1*-graph with the vertex matrix A. There is a unique Borel probability measure M on* Λ^{∞} *, given on the cylinder sets by*

$$
M([\lambda]) = \rho(\Lambda)^{-d(\lambda)} x_{s(\lambda)}^{\Lambda}
$$

Where $\rho(\lambda)$ *is the spectral radius of the vertex matrix A and* $(x_v^{\Lambda})_{v \in V}$ *is the unique normalized Perron-Frobenious eigenvector of* Λ*.*

Now we are ready to introduce the spectral triple given by Farsi et al. in [\[7\]](#page-19-10). Recall that a spectral triple is a data comprising of a C^* -algebra, a representation of the algebra on a Hilbert space and a Dirac operator. We are going to describe the data next. To that end, let $\Lambda = (V, E, r, s)$ be a strongly connected finite graph without multiple edges, loops and sources. Let the corresponding 1-graph be Λ .

The Hilbert space: The Hilbert space of the spectral triple is taken to be $L^2(\Lambda^\infty, M)$. There is a very useful dense subspace of the Hilbert space given by the linear span of the characteristic functions of the cylinder sets parametrized by finite length paths.

The representation of $C^*(\Lambda)$: Consider the seperable Hilbert space $\mathcal{H} = L^2(\Lambda^\infty, M)$. Using a semi-branching function system on the measure space (Λ^{∞}, M) , one can construct a representation $\pi: C^*(\Lambda) \to B(\mathcal{H})$ which is given by the following:

(2.2)
$$
\pi(S_{\lambda})(\chi_{[\eta]}) = \rho(\Lambda)^{d(\lambda)/2} \chi_{[\lambda \eta]},
$$

where λ , η are finite paths and $\chi_{[\eta]}$ is a characteristic function on the cylinder set $[\eta]$ as usual. To define $\pi(S^*_{\lambda})$, we consider two cases:

Case I $d(\lambda) \geq d(\eta)$: There are two subcases. Firstly, let $\lambda = \eta \beta$ for some β . Then

(2.3)
$$
(\pi(S_{\lambda}))^*(\chi_{[\eta]}) = \rho(\Lambda)^{-d(\lambda)/2} \chi_{[s(\lambda)]}
$$

Otherwise

(2.4)
$$
(\pi(S_{\lambda}^*))(\chi_{[\eta]}) = 0.
$$

Case 2: $d(\lambda) < d(\eta)$: Again there are two subcases. Firstly, let $\lambda\beta = \eta$ for some β . Then

(2.5)
$$
(\pi(S_{\lambda}^{\ast}))(\chi_{[\eta]}) = \rho(\Lambda)^{-d(\lambda)/2} \chi_{[\beta]}
$$

Otherwise as before,

(2.6)
$$
(\pi(S^*_{\lambda}))(\chi_{[\eta]}) = 0
$$

The Dirac Operator: Define R−¹ to be the linear subspace of *L* 2 (Λ∞*, M*) spanned by constant function on Λ^{∞} . For $k \in \mathbb{N}$, define $\mathcal{R}_k \subset L^2(\Lambda^{\infty}, M)$ by,

$$
\mathcal{R}_k = \mathrm{Sp}\{\chi_{[\eta]} \mid \eta \in \Lambda^*, d(\eta) \le k\}
$$

Now let us note down the following proposition which is going to be important in proving the main results of this paper.

Proposition 2.14. For a fixed $k > 1$, consider the subset $\mathcal{C}_k = \{ \chi_{[\eta]} | d(\eta) = k \} \subset$ \mathcal{R}_k *. Then elements of* \mathcal{C}_k *forms a linear basis for the space* \mathcal{R}_k *. Consequently, if the graph has m number of edges,* dim $(\mathcal{R}_k) \leq m^k$.

Proof. First we can observe that for any $\lambda \in \Lambda^*$ and $n \in \mathbb{N}$,

$$
[\lambda]=\bigcup_{\substack{\mu\in\Lambda^*,d(\mu)=n\\s(\mu)=r(\lambda)}}[\mu\lambda]
$$

For $\eta_1, \eta_2 \in \Lambda^*$ such that $\eta_1 \neq \eta_2$ and $d(\eta_1) = d(\eta_2)$, it is easy to see that $[\eta_1] \cap [\eta_2] =$ *≬*. Thus for $\mu_1 \neq \mu_2$ with $d(\mu_1) = d(\mu_2)$, $[\mu_1 \lambda] ∩ [\mu_2 \lambda] = ∅$. This implies that any $x \in [\lambda]$ lies in a unique cylinder set $[\mu\lambda]$ for a unique finite path μ with $d(\mu) = n$. Therefore,

$$
\chi_{[\lambda]} = \sum_{\substack{\mu, d(\mu) = n \\ s(\mu) = r(\lambda)}} \chi_{[\mu\lambda]}
$$

Now if we consider any $\lambda \in \Lambda^*$ such that $d(\lambda) < k$ then we have,

$$
\chi_{[\lambda]} = \sum_{\substack{\mu, d(\mu)=k-d(\lambda) \\ s(\mu)=r(\lambda)}} \chi_{[\mu\lambda]}
$$

As $d(\mu\lambda) = k$ in the above formula, every element of \mathcal{R}_k is a linear combination of elements in \mathcal{C}_k . Now let

$$
\sum_{\eta,d(\eta)=k} c_{\eta} \chi_{[\eta]} = 0,
$$

for some scalars *cη*. As the intersection of any two cylinder sets is empty for different paths of same degree, by choosing one of the infinite paths for each η we get $c_n = 0$. Hence these vectors are linearly independent in \mathcal{R}_k . There can be at most m^k paths of length *k* if number of edges in the graph is *m* giving the desired bound on the dimension of \mathcal{R}_k . \square

Let Ξ_k be the orthogonal projection in $L^2(\Lambda^\infty, M)$ onto \mathcal{R}_k . For a pair $(k, l) \in$ $\mathbb{N} \times (\mathbb{N} \cup \{-1\})$ with $k > l$, let

$$
\widehat{\Xi}_{k,l} = \Xi_k - \Xi_l.
$$

As $\mathcal{R}_l \subset \mathcal{R}_k$ and hence $\Xi_{k,l}$ is an orthogonal projection onto $\mathcal{R}_k \cap \mathcal{R}_l^{\perp}$. Given an increasing sequence of $\alpha = (\alpha_q)$ of positive real number with $\lim_{q\to\infty} \alpha_q = \infty$ such that there exists a $C > 0$ for which $| \alpha_{q+1} - \alpha_q | < C$ for all q, define an unbounded operator *D* on $L^2(\Lambda^\infty, M)$ by,

(2.7)
$$
D := \sum_{q \in \mathbb{N}} \alpha_q \widehat{\Xi}_{q,q-1}.
$$

For the proof of the following proposition, the reader is referred to [\[7\]](#page-19-10).

Proposition 2.15. *Let* $\Lambda = (V, E, r, s)$ *be a finite strongly connected directed graph without multiple edges and loops and the corresponding* 1*-graph be* Λ*. Let* A^Λ *be the dense* *-subalgebra of $C^*(\Lambda)$ spanned by $S_{\mu}S_{\eta}^*$ and D be the Dirac operator *on* $\mathcal{H} = L^2(\Lambda^\infty, M)$ *given by the formula [2.7.](#page-6-0) Then the data* $(\mathcal{A}_\Lambda, \mathcal{H}, D)$ *gives a spectral triple for* $C^*(\Lambda)$ *.*

Now we prove the following proposition which is a bit of digression. This proposition is not going to be used in proving the main results in this paper.

Proposition 2.16. With the choice $\alpha_q = q^{\frac{1}{2} + \epsilon}$ for some $0 < \epsilon < \frac{1}{2}$ in the formula [2.7](#page-6-0) of the Dirac operator, $(C^*(\Lambda), L^2(\Lambda^\infty, M), D)$ is a Θ -summable spectral triple.

Proof. It is easy to see that $| \alpha_{q+1} - \alpha_q | \leq 1$ for all *q* so that $(C^*(\Lambda), L^2(\Lambda^\infty, M), \mathcal{D})$ is a spectral triple. As e^{-tD^2} is a positive operator for all $t > 0$, to prove that e^{-tD^2} is trace class, it is enough to show that $\text{Tr}(e^{-tD^2}) < \infty$ for all $t > 0$. If $\dim(\mathcal{R}_q \cap \mathcal{R}_{q-1}^{\perp})$ is n_q , then fixing an orthonormal basis $\{\xi_1, \ldots, \xi_{n_q}\}$ of $(\mathcal{R}_q \cap \mathcal{R}_{q-1}^{\perp})$, we get

$$
\text{Tr}(e^{-tD^2}) = \sum_{q \in \mathbb{N}, j=1}^{j=n_q} \langle e^{-tD^2} \xi_j, \xi_j \rangle = \sum_{q \in \mathbb{N}} e^{-tq^{1+2\epsilon}} n_q.
$$

Now if we assume that the number of edges of the graph is *m*, then by Proposition [2.14,](#page-6-1) $n_q \leq m^q$. Then the above series is dominated by the series $\sum_q e^{-tq^{1+2\epsilon}} m^q$. If we denote the *q*-th term of the later by a_q , then $a_q > 0$ for all *q* and

$$
\text{lim}_q (a_q)^{1/q} = \text{lim}_q m (e^{-tq^{1+2\epsilon}})^{\frac{1}{q}} = \text{lim}_q m / (e^{tq^{1+2\epsilon}})^{\frac{1}{q}}.
$$

Now the denominator goes to infinity for all $t > 0$ as $q \to \infty$. This can be seen by taking logarithm of the denominator. Therefore, $\lim_{q} (a_q)^{1/q} = 0$. Hence by the root test, $\text{Tr}(e^{-tD^2}) < \infty$ for all $t > 0$.

2.16.1. *Isometric Action.* Now we are going to recall the notion of 'orientation preserving quantum isometric action'. We shall drop the term orientation preserving in the sequel. The main ideas are taken from [\[3\]](#page-18-0). We do slight modifications according to our needs. We start with the standard definition of unitary co-representation of a CQG on a Hilbert space.

Definition 2.17. (see [\[20,](#page-19-13)[21\]](#page-19-18)) A *unitary co-representation* of a CQG G on a Hilbert space H (*possibly infinite dimensional*) is a unitary element \tilde{U} of $\mathcal{M}(\mathcal{K}(\mathcal{H})\otimes C(\mathbb{G}))$ such that

$$
(\mathrm{id} \otimes \Delta) \circ \widetilde{U} = \widetilde{U}_{(12)} \widetilde{U}_{(13)},
$$

There is a standard equivalent picture of a unitary co-representations which we recall now. It is going to be useful in the sequel. Let *U* be a C-linear map from a Hilbert space H to the Hilbert $C(\mathbb{G})$ -module $H \otimes C(\mathbb{G})$ which satisfies

- (I) $\langle U(\xi), U(\eta) \rangle = \langle \xi, \eta \rangle$.1 for all $\xi, \eta \in \mathcal{H}$;
- (II) $(U \otimes id) \circ U = (id \otimes \Delta) \circ U$ on some dense subspace $\mathcal{K} \subset \mathcal{H}$ where U is algebraic i.e. $U(\mathcal{K}) \subset \mathcal{K} \otimes_{\text{alg}} C(\mathbb{G})_0;$

(III) $\text{Sp}\{U(\xi), q \mid \xi \in \mathcal{H}, q \in C(\mathbb{G})\}$ is dense in $\mathcal{H} \overline{\otimes} C(\mathbb{G})$.

Given such a $\mathbb{C}\text{-linear map}$, we have a unitary adjointable map on the Hilbert module $\mathcal{H} \overline{\otimes} C(\mathbb{G})$ given by $\overline{U}(\xi \otimes q) = U(\xi) . q$. Therefore, we have $\overline{U} \in \mathcal{L}(\mathcal{H} \overline{\otimes} C(\mathbb{G}))$ = $\mathcal{M}(\mathcal{K}(\mathcal{H})\otimes C(\mathbb{G}))$. Moreover \widetilde{U} satisfies (id $\otimes \Delta$) $\circ \widetilde{U} = \widetilde{U}_{(12)}\widetilde{U}_{(13)}$ making \widetilde{U} a unitary co-representation of $\mathbb G$ on $\mathcal H$. Conversely, given a unitary co-representation \tilde{U} , one can construct the C-linear map *U* by the formula $U(\xi) := \tilde{U}(\xi \otimes 1)$. Then it is easy to check that this *U* satisfies (I), (II) and (III).

Now let $(A, \mathcal{H}, \mathcal{D})$ be a spectral triple on a C^* -algebra and $\alpha : A \to A \otimes C(\mathbb{G})$ be an action of a CQG G on A. Also let τ be a faithful state on $C(\mathbb{G})$ with $L^2(\mathbb{G})$ being the GNS-space with respect to τ . If we denote the representation of $\mathcal A$ on *H* by *π* and the GNS representation of the C^* -algebra $C(\mathbb{G})$ on $L^2(\mathbb{G})$ by [.], then there is a representation of the *C*^{*}-algebra $\mathcal{A} \otimes C(\mathbb{G})$ on $\mathcal{B}(\mathcal{H} \otimes L^2(\mathbb{G}))$. We denote the representation by $(\pi \otimes [.)$. We say that the action α is implemented on the spectral data by some unitary co-representation \tilde{U} of \mathbb{G} on H if

$$
(\pi \otimes [.])\alpha(a) = \widetilde{U}(\pi(a) \otimes [1])\widetilde{U}^* \in \mathcal{B}(\mathcal{H} \otimes L^2(\mathbb{G})) \ \forall \ a \in \mathcal{A},
$$

where 1 is the identity element of $C(\mathbb{G})$. Note that $\mathcal{M}(\mathcal{K}(\mathcal{H})\otimes C(\mathbb{G}))$ is a subalgebra of $\mathcal{B}(\mathcal{H} \otimes L^2(\mathbb{G}))$ and therefore $\widetilde{U} \in \mathcal{B}(\mathcal{H} \otimes L^2(\mathbb{G}))$. We shall denote the map $\widetilde{U}(\pi(.)\otimes[.])\widetilde{U}^*$ from *A* to $B(\mathcal{H}\otimes L^2(\mathbb{G}))$ by ad_{\widetilde{U}}. Clearly ad_{\widetilde{U}} is a *C*^{*}-homomorphism.
The reader can compare the following definition with Definition 2.6 of [3] (also see The reader can compare the following definition with Definition 2.6 of [\[3\]](#page-18-0) (also see the discussion in the second paragraph on page no. 1803 of [\[6\]](#page-19-11)).

Definition 2.18. Let (A, H, D) be a spectral triple where A is a unital C^* -algebra. An action α of a CQG \mathbb{G} on \mathcal{A} is said to be 'quantum isomteric' (or simply isometric) if

- (I) α is implemented on the spectral data by some unitary co-representation \tilde{U} .
- (II) $\tilde{U} \circ (D \otimes 1) = (D \otimes 1) \circ \tilde{U}$ on appropriate domains.

Remark 2.19. Note that although in the definition of a spectral triple (A, H, D) , it is in general assumed that the representation π : $\mathcal{A} \to \mathcal{B}(\mathcal{H})$ is faithful, our consideration of quantum isometric action does not really require π to be faithful. In this context we would like to remind the reader that for the spectral triple $(C^*(\Lambda), L^2(\Lambda^\infty, M), \mathcal{D})$, the representation π is faithful if and only if the corresponding 1-graph is aperiodic if and only if every closed cycle has an exit.

3. Main section

In this section we are going to prove the following main theorem:

Theorem 3.1. *Let* Λ *be a strongly connected* 1*-graph without multiple edge, loop or source and* $(C^*(\Lambda), L^2(\Lambda^\infty, M), \mathcal{D})$ *be the spectral triple as given in Proposition* [2.15.](#page-7-0) Then the action of $Aut^+(\Lambda)$ on $C^*(\Lambda)$ given in Proposition [2.8](#page-3-0) is isometric *in the sense of Definition [2.18.](#page-8-0)*

We start with constructing a unitary co-representation \tilde{U} of $Aut^+(\Lambda)$ on the Hilbert space $\mathcal{H} = L^2(\Lambda^\infty, M)$. We are going to denote the Hilbert space $L^2(\Lambda^\infty, M)$ by H throughout this section. Let us denote the dense subspace say of H given by the

10 JOARDAR AND SHARMA

increasing union $\cup_k \mathcal{R}_k$ by K. The idea is to define an isometric C-linear map U from K to $\mathcal{H}\overline{\otimes}C(\mathbb{G})$ and extending it to the whole of H. The important observation is that since we are seeking for a unitary co-representation that would implement the given action α , once the representation is defined on \mathcal{R}_0 , it is essentially determined by the action α on the whole of $\mathcal H$ which leads to the natural formula for *U* on the eigenspaces \mathcal{R}_k (see Formula [3.1](#page-9-0) below). The challenging part is to prove that the formulae gel together to produce a well-defined unitary $\tilde{U} \in \mathcal{M}(\mathcal{K}(\mathcal{H}) \otimes C(\mathbb{G}))$. With this heuristics we start with the following theorem.

Theorem 3.2. *There exists a well-defined linear* \mathbb{C} *-linear map* $U : \mathcal{K} \to \mathcal{K} \otimes_{\text{alg}}$ $C(\text{Aut}^+(\Lambda))_0$ *given on the characteristic function of a cylinder set* $[\lambda]$ *for some path* $\lambda = \lambda_1 \lambda_2 \ldots \lambda_k$ *of length k by the following formula:*

$$
(3.1) \tU(\chi_{[\lambda]}) = \sum_{\eta} \chi_{[\eta]} \otimes q_{r(\eta_1)r(\lambda_1)} q_{s(\eta_1)s(\lambda_1)} \cdots q_{r(\eta_k)r(\lambda_k)} q_{s(\eta_k)s(\lambda_k)}
$$

where the summation varies over all the paths $\eta = \eta_1 \eta_2 \dots \eta_k$ *of degree k. Moreover,* $(U \otimes id) \circ U = (id \otimes \Delta) \circ U$ *on* K .

Proof. Fix $k \in \mathbb{N}$ and let \mathcal{R}_k be the finite-dimensional subspace of $L^2(\Lambda^\infty, M)$ for which \mathcal{C}_k forms a basis of \mathcal{R}_k as proved in proposition [2.14.](#page-6-1) So we have a welldefined C-linear map $U_k : \mathcal{R}_k \to \mathcal{K} \otimes_{\text{alg}} C(\text{Aut}^+(\Lambda))_0$, given on the basis elements by formula [3.1.](#page-9-0) Note that for $k = 0$, \mathcal{R}_0 is the vector space $\text{Sp}\{\chi_{[i]} : i = 1, 2, \ldots, n\},\$ where the vertices of Λ are $\{1, 2, \ldots, n\}$. Then the map U_0 on \mathcal{R}_0 has the following formula:

(3.2)
$$
U_0(\chi_{[i]}) = \sum_j \chi_{[j]} \otimes q_{ji}.
$$

As \mathcal{R}_k forms an increasing chain of finite-dimensional subspaces, to get a well defined linear map on \mathcal{K} , we just need to show that for $l < k$ the maps satisfy, $U_k|_{\mathcal{R}_l} = U_l$. To show this we need to show equality only for the basis set \mathcal{C}_l of \mathcal{R}_l . Let $\chi_{[\lambda]} \in \mathcal{C}_l$. Then an element in \mathcal{R}_k can written as a linear combination of basis element of \mathcal{R}_k . In particular,

$$
\chi_{[\lambda]} = \sum_{\substack{\mu,d(\mu)=k-l \\ s(\mu)=r(\lambda)}} \chi_{[\mu\lambda]}
$$

where μ varies over all the paths of length $m = k - l$. So we need to prove,

$$
U_l(\chi_{[\lambda]}) = \sum_{\substack{\mu, d(\mu) = m \\ s(\mu) = r(\lambda)}} U_k(\chi_{[\mu\lambda]})
$$

Expanding LHS by definition we get,

$$
U(\chi_{[\lambda]}) = \sum_{\substack{\xi \in \Lambda^* \\ d(\xi)=l}} \chi_{[\xi]} \otimes q_{r(\xi_{m+1})r(\lambda_{m+1})} q_{s(\xi_{m+1})s(\lambda_{m+1})} \cdots q_{r(\xi_{m+l})r(\lambda_{m+l})} q_{s(\xi_{m+l})s(\lambda_{m+l})}
$$

where we are writing $\xi = \xi_{m+1}\xi_{m+2}\cdots\xi_{m+l}$ and $\lambda = \lambda_{m+1}\lambda_{m+2}\cdots\lambda_{m+l}$. On the other hand by expanding RHS we get,

$$
\sum_{\substack{\mu \in \Lambda^* \\ d(\mu)=m, \\ s(\mu)=r(\lambda)}} U(\chi_{[\mu\lambda]}) = \sum_{\substack{\mu \in \Lambda^* \\ d(\mu)=m}} \sum_{\substack{\delta \in \Lambda^* \\ d(\mu)=m}} \chi_{[\delta]} \otimes q_{r(\delta_1)r(\mu_1)} q_{s(\delta_1)s(\mu_1)} \cdots q_{r(\delta_m)r(\mu_m)} q_{s(\delta_m)s(\mu_m)}
$$
\n
$$
q_{r(\delta_{m+1})r(\lambda_{m+1})} q_{s(\delta_{m+1})s(\lambda_{m+1})} \cdots \cdots
$$
\n
$$
\cdots \cdots q_{r(\delta_{m+l})r(\lambda_{m+l})} q_{s(\delta_{m+l})s(\lambda_{m+l})}
$$

Where $\delta = \delta_1 \cdots \delta_m \delta_{m+1} \cdots \delta_{m+l}$ and $\mu = \mu_1 \mu_2 \cdots \mu_m$.

Since every path δ of degree $k=m+l$ has a unique decomposition into a path ζ,ξ of degree *m* and *l*, where we write $\xi = \xi_{m+1} \cdots \xi_{m+l}, \zeta = \zeta_1 \zeta_2 \cdots \zeta_m$ and the above summation becomes,

$$
= \sum_{\mu \in \Lambda^*, d(\mu) = m} \sum_{\substack{\delta = \zeta \xi \in \Lambda^* \\ s(\mu) = r(\lambda)}} \chi_{[\zeta \xi]} \otimes q_{r(\zeta_1)r(\mu_1)} q_{s(\zeta_1)s(\mu_1)} \cdots q_{r(\zeta_m)r(\mu_m)} q_{s(\zeta_m)s(\mu_m)}
$$

\n
$$
= \sum_{\mu \in \Lambda^*} \sum_{\substack{\xi \in \Lambda^* \\ s(\mu) = r(\lambda)}} \sum_{\substack{\delta \in \Lambda^* \\ d(\zeta) = m}} \sum_{\substack{\chi(\zeta \xi) \\ \chi(\zeta \xi) = m}} \chi_{[\zeta \xi]} \otimes q_{r(\zeta_1)r(\mu_1)} q_{s(\zeta_1)s(\mu_1)} \cdots q_{r(\zeta_m)r(\mu_m)} q_{s(\zeta_m)s(\mu_m)}
$$

\n
$$
= \sum_{\substack{\mu \in \Lambda^* \\ s(\mu) = r(\lambda)}} \sum_{\substack{\delta \in \Lambda^*, d(\zeta) = m \\ d(\zeta) = r(\xi)}} \chi_{[\zeta \xi]} \otimes q_{r(\zeta_1)r(\mu_1)} q_{s(\zeta_1)s(\mu_1)} \cdots q_{r(\zeta_m)r(\mu_m)} q_{s(\zeta_m)s(\mu_m)}
$$

\n
$$
\cdots \cdots q_{r(\xi_{m+1})r(\lambda_{m+1})} q_{s(\xi_{m+1})s(\lambda_{m+1})} \cdots \cdots
$$

We shall prove now that the coefficients for a fixed ξ and for any ζ with $s(\zeta) = r(\xi)$ are the same. Now in the summands for fix ξ , ζ such that $s(\zeta) = r(\xi)$, the coefficients of $\chi_{\left[\zeta\xi\right]}$ is

$$
= \sum_{\substack{\mu,d(\mu)=m \ g(\mu)=r(\lambda)}} q_{r(\zeta_1)r(\mu_1)}q_{s(\zeta_1)s(\mu_1)}\cdots q_{r(\zeta_m)r(\mu_m)}q_{s(\zeta_m)s(\mu_m)}q_{r(\xi_{m+1})r(\lambda_{m+1})}q_{s(\xi_{m+1})s(\lambda_{m+1})}
$$

Now note that in the above summation, $(r(\mu_1), s(\mu_1))$ can be replaced by $(i, j) \in$ *V* × *V*. This can be proved by the following argument: Whenever $(i, j) \notin E$, $q_{r(\zeta_1)i}q_{s(\zeta_1),j} = 0$ by [2.1.](#page-2-0) As for any $(i,j) \in E$, there are two possibilities: Either (i, j) is the first edge of a *k*-length path in which case it is already included in the summation, or (i, j) fails to be the first edge of a *k*-length path. In this case, as $\zeta_1 \cdots \zeta_m \zeta_{m+1} \cdots \zeta_{m+l}$ forms a *k*-length path,

$$
q_{r(\zeta_1)i}q_{s(\zeta_1)j}\cdots q_{r(\zeta_m)r(\mu_m)}q_{s(\zeta_m)s(\mu_m)}q_{r(\xi_{m+1})r(\lambda_{m+1})}\cdots q_{s(\xi_{m+1})s(\lambda_{m+l})}=0,
$$

which enables us to replace $(r(\mu_1), s(\mu_1))$ by $(i, j) \in V$. Therefore the summand becomes equal to

$$
\sum_{\mu,d(\mu)=m-1} \sum_{i,j \in V} q_{r(\zeta_1)i} q_{s(\zeta_1)j} \cdots q_{r(\zeta_m)r(\mu_m)} q_{s(\zeta_m)s(\mu_m)} q_{r(\xi_{m+1})r(\lambda_{m+1})} q_{s(\xi_{m+1})s(\lambda_{m+1})}
$$
\n
$$
= \sum_{\mu,d(\mu)=m-1} q_{r(\zeta_2)r(\mu_2)} q_{s(\zeta_2)s(\mu_2)} \cdots q_{r(\zeta_m)r(\mu_m)} q_{s(\zeta_m)s(\mu_m)} q_{r(\xi_{m+1})r(\lambda_{m+1})} q_{s(\xi_{m+1})s(\lambda_{m+1})}
$$
\n
$$
= \sum_{\substack{\mu,d(\mu)=m-1 \ g(\mu)=r(\lambda_m) \\ s(\mu)=r(\lambda_m)}} q_{r(\zeta_2)r(\mu_2)} q_{s(\zeta_2)s(\mu_2)} \cdots q_{r(\zeta_m)r(\mu_m)} q_{s(\zeta_m)s(\mu_m)} q_{r(\xi_{m+1})r(\lambda_{m+1})} q_{s(\xi_{m+1})s(\lambda_{m+1})}
$$

By repeating the same argument, we can get rid of the first *m* indices and the summand reduces to

$$
q_{r(\xi_{m+1})r(\lambda_{m+1})}q_{s(\xi_{m+1})s(\lambda_{m+1})}\cdots q_{r(\xi_{m+1})r(\lambda_{m+1})}q_{s(\xi_{m+1})s(\lambda_{m+1})}
$$

Therefore the term $\mu \in \Lambda^*$, $\overline{d}(\mu) = m$ $s(\mu)=r(\lambda)$ $U(\chi_{\lbrack \mu\lambda \rbrack})$ becomes

$$
= \sum_{\substack{\xi \in \Lambda^*, d(\xi) = l \\ \xi = \xi_{m+1} \cdots \xi_{m+1}}} \sum_{\substack{\zeta \in \Lambda^*, d(\zeta) = m \\ \zeta = \zeta_1 \zeta_2 \cdots \zeta_m, \\ s(\zeta) = r(\xi)}} \chi_{[\zeta \xi]} \otimes \sum_{\substack{\mu \in \Lambda^* \\ d(\mu) = m}} q_{r(\zeta_1)r(\mu_1)} q_{s(\zeta_1)s(\mu_1)} \cdots q_{r(\zeta_m)r(\mu_m)} q_{s(\zeta_m)s(\mu_m)}
$$

which by the above calculations equals to

$$
\sum_{\xi \in \Lambda, d(\xi)=l} \sum_{\zeta, d(\zeta)=m} \chi_{[\zeta \xi]} \otimes q_{r(\xi_{m+1})r(\lambda_{m+1})} q_{s(\xi_{m+1})s(\lambda_{m+1})} q_{r(\xi_{m+2})r(\lambda_{m+2})}
$$
\n
$$
\xi = \xi_{m+1} \cdots \xi_{m+l} \zeta_{\zeta} \zeta_{\zeta} \cdots \zeta_{m}, \qquad q_{s(\xi_{m+2})s(\lambda_{m+2})} \cdots \cdots q_{r(\xi_{m+l})r(\lambda_{m+l})} q_{s(\xi_{m+l})s(\lambda_{m+l})}
$$
\n
$$
= \sum_{\xi \in \Lambda, d(\xi)=l} \left(\sum_{\zeta, d(\zeta)=m} \chi_{[\zeta \xi]} \right) \otimes q_{r(\xi_{m+1})r(\lambda_{m+1})} q_{s(\xi_{m+1})s(\lambda_{m+1})} \cdots \cdots \zeta_{\zeta = \xi_{m+1} \cdots \xi_{m+l}} \zeta_{\zeta = \zeta_1 \zeta_2 \cdots \zeta_m, s(\zeta) = r(\xi)} \cdots q_{r(\xi_{m+l})r(\lambda_{m+l})} q_{s(\xi_{m+l})s(\lambda_{m+l})}
$$
\nAs we know that from the proof of proposition 2.14, $\chi_{[\xi]} = \sum \chi_{[\zeta \xi]}$ and

therefore the summation becomes

$$
\sum_{\substack{\xi \in \Lambda^* \\ d(\xi)=l}} \chi_{[\xi]} \otimes q_{r(\xi_{m+1})r(\lambda_{m+1})} q_{s(\xi_{m+1})s(\lambda_{m+1})} \cdots q_{r(\xi_{m+l})r(\lambda_{m+l})} q_{s(\xi_{m+l})s(\lambda_{m+l})}
$$

$$
\xi = \xi_{m+1} \cdots \xi_{m+l}
$$

ζ,d(*ζ*)=*m ζ*=*ζ*1*ζ*2···*ζ^m* $s(\zeta)=r(\xi)$

which is nothing but $U(\chi_{[\lambda]})$. In conclusion we have,

$$
U_l(\chi_{[\lambda]}) = \sum_{\substack{\mu, d(\mu) = m \\ s(\mu) = r(\lambda)}} U_k(\chi_{[\mu\lambda]})
$$

and we have a well defined C-linear map $U : \mathcal{K} \to \mathcal{K} \otimes_{\text{alg}} C(\text{Aut}^+(\Lambda))_0$ where $\mathcal{K} = \bigcup_k \mathcal{R}_k$ is dense in $L^2(\Lambda^\infty, M)$. The formula $(U \otimes id) \circ U = (id \otimes \Delta) \circ U$ follows from the well known fact that the action α of $\text{Aut}^+(\Lambda)$ on $C(V)$ extends as representation of $Aut^+(\Lambda)$ on all path spaces of finite lengths. \square

Proposition 3.3. *Let* $\lambda, \eta \in \Lambda^*$ *such that* $d(\lambda) = d(\eta)$ *. Then*

(3.3)
$$
\langle U(\chi_{[\lambda]}), U(\chi_{[\eta]}) \rangle = \langle \chi_{[\lambda]}, \chi_{[\eta]} \rangle.
$$

Proof. Fix some $k > 0$ and two elements $\lambda, \eta \in \Lambda^*$ such that $d(\lambda) = d(\eta) = k$. Write λ and η as $\lambda_1 \lambda_2 \cdots \lambda_k$ and $\eta_1 \eta_2 \cdots \eta_k$ for appropriately composable edges λ_i 's and η_i 's respectively. Then

$$
\langle U(\chi_{[\lambda]}), U(\chi_{[\eta]})\rangle = \left\langle \sum_{\zeta; d(\zeta) = k; \zeta = \zeta_1 \cdots \zeta_k} \chi_{[\zeta]} \otimes q_{r(\zeta_1)r(\lambda_1)} q_{s(\zeta_1)s(\lambda_1)} \cdots q_{r(\zeta_n)r(\lambda_k)} q_{s(\zeta_k)s(\lambda_k)},\right.\
$$

$$
\sum_{\xi; d(\xi) = k; \xi = \xi_1 \cdots \xi_k} \chi_{[\xi]} \otimes q_{r(\xi_1)r(\eta_1)} q_{s(\xi_1)s(\eta_1)} \cdots q_{r(\xi_k)r(\eta_k)} q_{s(\xi_k)s(\eta_k)} \right\rangle
$$

$$
= \sum_{\zeta,\xi} \langle \chi_{[\zeta]}, \chi_{[\xi]} \rangle \otimes q_{s(\zeta_k)s(\lambda_k)} q_{r(\zeta_k)r(\lambda_k)} \cdots q_{s(\zeta_1)s(\lambda_1)} q_{r(\zeta_1)r(\lambda_1)} q_{r(\xi_1)r(\eta_1)} q_{s(\xi_1)s(\eta_1)} q_{r(\xi_2)r(\eta_2)} q_{s(\xi_2)s(\eta_2)} \cdots q_{r(\xi_k)r(\eta_k)} q_{s(\xi_k)s(\eta_k)}
$$

As $d(\zeta) = d(\xi), \langle \chi_{[\zeta]}, \chi_{[\xi]} \rangle = 0$ for $\zeta \neq \xi$, the last expression becomes

$$
= \sum_{\zeta} x_{s(\zeta)}^{\Lambda} q_{s(\zeta_k)s(\lambda_k)} q_{r(\zeta_k)r(\lambda_k)} \cdots q_{s(\zeta_1)s(\lambda_1)} q_{r(\zeta_1)r(\lambda_1)} q_{r(\zeta_1)r(\eta_1)} q_{s(\zeta_1)s(\eta_1)} \cdots
$$

$$
\cdots q_{r(\zeta_k)r(\eta_k)} q_{s(\zeta_k)s(\eta_k)}
$$

Therefore, to prove [3.3,](#page-11-0) one needs to show that

$$
\delta_{\lambda,\eta} x_{s(\lambda)}^{\Lambda} = \delta_{\lambda,\eta} \sum_{\zeta} x_{s(\zeta)}^{\Lambda} q_{s(\zeta_k)s(\lambda_k)} q_{r(\zeta_k)r(\lambda_k)} \cdots q_{s(\zeta_1)s(\lambda_1)} q_{r(\zeta_1)r(\lambda_1)} q_{r(\zeta_1)r(\eta_1)} q_{s(\zeta_1)s(\eta_1)}
$$

$$
q_{r(\zeta_2)r(\eta_2)} q_{s(\zeta_2)s(\eta_2)} \cdots q_{r(\zeta_k)r(\eta_k)} q_{s(\zeta_k)s(\eta_k)}.
$$

Now the above equation follows from the invariance of the unique KMS state under the action α from Proposition [2.8.](#page-3-0) The reader can compare with the relevant expression from the proof of Proposition 2.31 in [\[15\]](#page-19-6). \Box

Lemma 3.4. *The* $\mathbb C$ *linear map* $U : \mathcal K \to \mathcal K \otimes_{alg} C(Aut^+(\Lambda))_0$ *in theorem* [3.2](#page-9-1) *is inner product preserving on* K*.*

Proof. Consider paths λ , η of length k , l respectively. Then $\chi_{[\lambda]}, \chi_{[\eta]} \in \mathcal{R}_{\max\{k,l\}}$. Therefore, we can write $\chi_{[\lambda]}, \chi_{[\eta]}$ as linear combinations of basis elements in $\mathcal{R}_{\max\{k,l\}}$. But a basis for $\mathcal{R}_{\max\{k,l\}}$ is given by $\mathcal{C}_{\max\{k,l\}} = {\chi_{[\zeta]} : d(\zeta) = \max\{k,l\}}$. As *U* is inner product preserving on $\mathcal{C}_{\max\{k,l\}}$ by Proposition [3.3,](#page-11-0) $\langle U(\chi_{[\lambda]}), U(\chi_{[\eta]}) \rangle =$ $\langle \chi_{[\lambda]}, \chi_{[\eta]} \rangle$ 1. Therefore *U* preserves the inner product on \mathcal{K} . \Box

As K is dense in H and $U : \mathcal{K} \to \mathcal{K} \otimes_{alg} C(\text{Aut}^+(\Lambda))_0$ is inner product preserving, U extends to an inner product preserving $\mathbb{C}\text{-linear map from } \mathcal{H}$ to the Hilbert module $\mathcal{H} \overline{\otimes} C(\mathbb{G})$. We continue to denote the extension by *U*. Now we move forward to prove the density condition (III) of Definition [2.17.](#page-7-1)

Proposition 3.5. *For the inner product preserving map* $U: \mathcal{H} \to \mathcal{H} \overline{\otimes} C(Aut^{+}(\Lambda)),$ $\text{Sp}\lbrace U(\xi), q \mid \xi \in \mathcal{H}, q \in C(Aut^+(\Lambda)) \rbrace$ *is dense in* $\mathcal{H} \overline{\otimes} C(Aut^+(\Lambda))$.

Proof. Note that as the linear span of elements $\{\chi_{[\lambda]} : \lambda \in \Lambda^*\}$ is dense in H, it is enough to approximate the elements $\chi_{[\lambda]} \otimes 1$ for all $\lambda \in \Lambda^*$. When length of λ is 0 or 1, this can be shown along the lines of [\[24\]](#page-19-4). Let us illustrate the proof for a path *λ* of length two. For a path of any finite length the argument can be adapted. To that end let $\lambda = \gamma \beta$ be a two length path for edges γ , β with $r(\beta) = s(\gamma)$. Then

$$
Sp{U(\xi).q} \ni \sum_{i \in V} \sum_{\substack{\zeta, \xi, r(\xi)=s(\zeta) \\ r(\zeta\xi)=i \\ d(\xi)=d(\zeta)=1}} U(\chi_{[\zeta\xi]})(q_{s(\beta)s(\xi)}q_{r(\beta)r(\xi)}q_{s(\gamma)s(\zeta)}q_{r(\gamma)r(\zeta)})
$$

Upon expanding, the last summation becomes

$$
\sum_{i \in V} \sum_{\substack{\zeta, \xi \in \Lambda^* \\ r(\zeta \xi) = i}} \sum_{\substack{\eta, \delta \in \Lambda^* \\ r(\delta) = s(\eta)}} \left(\chi_{[\eta \delta]} \otimes q_{r(\eta)r(\zeta)} q_{s(\eta)s(\zeta)} q_{r(\delta)r(\xi)} q_{s(\delta)s(\zeta)} \right)
$$

$$
= \sum_{\substack{i \in V \\ \zeta, \xi \in \Lambda^*}} \sum_{\eta, \delta} \chi_{[\eta \delta]} \otimes q_{r(\eta)r(\zeta)} q_{s(\eta)s(\zeta)} q_{r(\delta)r(\xi)} q_{s(\delta)s(\xi)} q_{s(\beta)s(\xi)} q_{r(\beta)r(\xi)} q_{s(\gamma)s(\zeta)} q_{r(\gamma)r(\zeta)}
$$

For $s(\delta) \neq s(\beta)$, the corresponding coefficients in the above summation are zero and therefore the summation is equal to

$$
\sum_{i \in V} \sum_{\substack{\zeta \\ r(\zeta) = i}} \sum_{\substack{\eta, \delta, \xi \\ s(\eta) = r(\delta) \\ s(\zeta) = r(\xi)}} \chi_{[\eta\delta]} \otimes q_{r(\eta)r(\zeta)} q_{s(\eta)s(\zeta)} q_{r(\delta)r(\xi)} q_{s(\beta)s(\xi)} q_{r(\beta)r(\xi)} q_{s(\gamma)s(\zeta)} q_{r(\gamma)r(\zeta)}
$$

Now in the above summation the term $q_{r(\delta)r(\xi)}q_{s(\beta)s(\xi)}q_{r(\beta)r(\xi)}$ can be replaced by $q_{r(\delta)k}q_{s(\beta)l}q_{r(\beta)k}$ for $(k, l) \in V \times V$. This is because if (k, l) in not an edge, $q_{r(\delta)k}q_{s(\beta)l} = 0$ by [2.1](#page-2-0) as $s(\beta) = s(\delta)$. All the edges already appear in the summation. Therefore using the fact that $\sum_l q_{s(\beta)l} = 1$, the summation reduces to

$$
\sum_{i \in V} \sum_{\substack{\zeta \\ r(\zeta) = i}} \sum_{\substack{\eta, \delta, \xi \\ s(\delta) = s(\zeta) \\ s(\eta) = r(\delta)}} \chi_{[\eta\delta]} \otimes q_{r(\eta)r(\zeta)} q_{s(\eta)s(\zeta)} \left(\sum_{k \in V} q_{r(\delta)k} q_{r(\beta)k} \right) q_{s(\gamma)s(\zeta)} q_{r(\gamma)r(\zeta)}.
$$

Again for $r(\delta) \neq r(\beta)$, $q_{r(\delta)k}q_{r(\beta)k} = 0$ for all $k \in V$. Therefore using the fact that $\sum_{k} q_{r(\beta)k} = 1$ and the graph has no multiple edges, the last summation further reduces to

$$
\sum_{i\in V}\sum_{\zeta,r(\zeta)=i}\sum_{\eta}\chi_{[\eta\beta]}\otimes q_{r(\eta)r(\zeta)}q_{s(\eta)s(\zeta)}q_{s(\gamma)s(\zeta)}q_{r(\gamma)r(\zeta)}.
$$

Now applying the same argument for a single edge proved in [\[24\]](#page-19-4), we have

$$
\sum_{i \in V} \sum_{\zeta, \xi, r(\xi) = s(\zeta)} U(\chi_{[\zeta\xi]})(1 \otimes q_{s(\beta)s(\xi)} q_{r(\beta)r(\xi)} q_{s(\gamma)s(\zeta)} q_{r(\gamma)r(\zeta)}) = \chi_{[\gamma\beta]} \otimes 1,
$$

i.e. $\chi_{[\lambda]} \otimes 1 \in \mathrm{Sp}\{U(\xi), q : \xi \in \mathcal{H}, q \in C(\mathrm{Aut}^+(\Lambda)\} \square$

Hence the $\mathbb C$ -linear map U satisfies the conditions (I), (II) and (III) following the Definition [2.17](#page-7-1) and therefore, thanks to the discussion following Definition [2.17,](#page-7-1) $\widetilde{U} \in \mathcal{L}(\mathcal{H} \overline{\otimes} C(\mathrm{Aut}^+(\Lambda))) = \mathcal{M}(\mathcal{K}(\mathcal{H}) \otimes C(\mathrm{Aut}^+(\Lambda)))$ is a unitary co-representation of Aut⁺(Λ) on the Hilbert space H.

Proposition 3.6. The unitary co-representation \tilde{U} commutes with the Dirac op*erator D i.e.* $\widetilde{U} \circ (D \otimes 1) = (D \otimes 1) \circ \widetilde{U}$ *on appropriate domains.*

Proof. Note that it is enough to show that \tilde{U} commutes with the projections on the eigen spaces of *D* viz. $\Xi_{q,q-1}$ for all *q* (see the discussion after Proposition [2.14](#page-6-1) for the notations). It is clear from the formula [3.1](#page-9-0) that *U* preserves the closed subspaces \mathcal{R}_q for all *q*. Therefore \tilde{U} commutes with the projections Ξ_q for all *q*. Hence *U* being a unitary co-representation, commutes with Ξ_{q-1}^{\perp} for all *q*. Then *U* commutes with $\Xi_q \wedge \Xi_{q-1}^{\perp}$ which is by definition $\Xi_{q,q-1}$. The last claim follows from the fact that $\Xi_q \wedge \Xi_{q-1}^{\perp} = \text{SOT} - \lim_n (\Xi_q \Xi_{q-1}^{\perp} \Xi_q)^n$. \square

Proposition 3.7. *The action* α *of* $\text{Aut}^+(\Lambda)$ *on* $C^*(\Lambda)$ *as given in Proposition* [2.8](#page-3-0) *is implemented on the spectral data* $(C^*(\Lambda), \mathcal{H}, D)$ *by the unitary co-representation* \tilde{U} .

Proof. We start by showing that for any $\lambda, \eta \in \Lambda^*$,

(3.4)
$$
(\pi \otimes [.)\alpha(S^*_{\lambda})U(\chi_{[\eta]}) = U(\pi(S^*_{\lambda})\chi_{[\eta]})
$$

Case1: Consider the case $d(\lambda) \geq d(\eta)$. Then there are two subcases: Firstly let $\lambda = \eta \beta$ for some path β of finite length. Let's write $\eta = \eta_1 \eta_2 \dots \eta_m$ and $\lambda = \lambda_1 \lambda_2 \cdots \lambda_n$. Then by the unique factorization of paths, λ can be written as $\eta_1 \eta_2 \dots \eta_m \lambda_{m+1} \dots \lambda_n$ where η_i and λ_i 's are appropriate composable edges. Note that $s(\lambda) = s(\lambda_n)$. Then by [2.3,](#page-5-0)

$$
U(\pi(S_{\lambda}^*)\chi_{[\eta]}) = \rho(\Lambda)^{-d(\lambda)/2} U(\chi_{[s(\lambda)]}) = \rho(\Lambda)^{-d(\lambda)/2} \sum_{i \in V} \chi_{[i]} \otimes q_{is(\lambda)}
$$

By the Formula [3.1,](#page-9-0)

$$
U(\chi_{[\eta]}) = \left(\sum_{\zeta: \zeta = \zeta_1 \zeta_2 \dots \zeta_m} \chi_{[\zeta]} \otimes q_{r(\zeta_1)r(\eta_1)} q_{s(\zeta_1)s(\eta_1)} \cdots q_{r(\zeta_m)r(\eta_m)} q_{s(\zeta_m)s(\eta_m)}\right)
$$

Therefore,

$$
(\pi \otimes [.])\alpha(S_{\lambda}^{*})U(\chi_{[\eta]}) = (\pi \otimes [.])(\alpha(S_{\lambda_{n}})^{*} \cdots \alpha(S_{\lambda_{1}})^{*})U(\chi_{[\eta]})
$$

\n
$$
= (\pi \otimes [.])\left(\left(\sum_{\xi_{n}} S_{\xi_{n}}^{*} \otimes q_{s(\xi_{n})s(\lambda_{n})} q_{r(\xi_{n})r(\lambda_{n})}\right) \cdots \left(\sum_{\xi_{1}} S_{\xi_{1}}^{*} \otimes q_{s(\xi_{1})s(\lambda_{1})} q_{r(\xi_{1})r(\lambda_{1})}\right)\right)
$$

\n
$$
\left(\sum_{\zeta} \chi_{[\zeta]} \otimes q_{r(\zeta_{1})r(\eta_{1})} q_{s(\zeta_{1})s(\eta_{1})} \cdots q_{r(\zeta_{m})r(\eta_{m})} q_{s(\zeta_{m})s(\eta_{m})}\right)
$$

\n
$$
= (\pi \otimes [.])\left(\sum_{\xi = \xi_{1} \cdots \xi_{n}} S_{\xi}^{*} \otimes q_{s(\xi_{n})s(\lambda_{n})} q_{r(\xi_{n})r(\lambda_{n})} \cdots q_{s(\xi_{1})s(\lambda_{1})} q_{r(\xi_{1})r(\lambda_{1})}\right)
$$

\n
$$
\left(\sum_{\zeta} \chi_{[\zeta]} \otimes q_{r(\zeta_{1})r(\eta_{1})} q_{s(\zeta_{1})s(\eta_{1})} \cdots q_{r(\zeta_{m})r(\eta_{m})} q_{s(\zeta_{m})s(\eta_{m})}\right)
$$

\n
$$
= \left(\sum_{\xi = \xi_{1} \cdots \xi_{n}} \pi(S_{\xi})^{*} \otimes [q_{s(\xi_{n})s(\lambda_{n})} q_{r(\xi_{n})r(\lambda_{n})} \cdots q_{s(\xi_{1})s(\lambda_{1})} q_{r(\xi_{1})r(\lambda_{1})}]\right)
$$

\n
$$
= \sum_{\zeta, \zeta, \zeta} \chi_{[\zeta]} \otimes q_{r(\zeta_{1})r(\eta_{1})} q_{s(\zeta_{1})s(\eta_{1})} \cdots q_{r(\zeta_{m})r(\eta_{m})} q_{s(\zeta_{m})s(\eta_{m})}\right)
$$

\n
$$
= \sum_{\zeta, \zeta
$$

Note that $d(\xi) \geq d(\zeta)$ as the action α and the map *U* are both degree preserving. Now for any path ξ, ζ with $d(\xi) \geq d(\zeta)$ such that $\xi = \zeta \gamma$ for some path γ , $\pi(S_{\xi}^{*})(\chi_{\{\zeta\}}) = \chi_{[s(\xi)]}$. If there is no path γ of length $(n - m)$ such that $\xi = \zeta \gamma$, then $\pi(S_{\xi}^{*})(\chi_{[\zeta]}) = 0$. For $\xi = \zeta \gamma$, using the unique factorisation of paths, we write

16 JOARDAR AND SHARMA

 $\xi = \zeta_1 \zeta_2 \cdots \zeta_m \xi_{m+1} \cdots \xi_n$, $\gamma = \xi_{m+1} \xi_{m+2} \cdots \xi_n$. Note that $s(\xi) = s(\xi_n)$. Therefore, using the expression $\lambda = \eta_1 \cdots \eta_m \lambda_{m+1} \cdots \lambda_n$, the last summation becomes

$$
\rho(\Lambda)^{-d(\xi)/2} \sum_{\substack{\xi,\zeta\\d(\xi)\geq d(\zeta)}} \chi_{[s(\xi)]} \otimes q_{s(\xi_n)s(\lambda_n)} q_{r(\xi_n)r(\lambda_n)} \cdots q_{s(\xi_{m+1})s(\lambda_{m+1})} q_{r(\xi_{m+1})r(\lambda_{m+1})}
$$

\n
$$
q_{s(\zeta_m)s(\eta_m)} q_{r(\zeta_m)r(\eta_m)} \cdots q_{s(\zeta_1)s(\eta_1)} q_{r(\zeta_1)r(\eta_1)}
$$

\n
$$
q_{r(\zeta_1)r(\eta_1)} q_{s(\zeta_1)s(\eta_1)} \cdots q_{r(\zeta_m)r(\eta_m)} q_{s(\zeta_m)s(\eta_m)}.
$$

As $d(\xi) = d(\lambda)$, ignoring the common scalar factor $\rho(\Lambda)^{-d/2}$, we shall show that for a fixed $i \in V$, the coefficient of $\chi_{[i]}$ in the above summation is $q_{is(\lambda)}$ proving [3.4](#page-14-0) in this subcase. The coefficient is given by

$$
\sum_{\substack{\xi,\zeta\\n=d(\xi)\geq d(\zeta)=m}}q_{s(\xi_n)s(\lambda_n)}q_{r(\xi_n)r(\lambda_n)}\cdots q_{s(\xi_{m+1})s(\lambda_{m+1})}q_{r(\xi_{m+1})r(\lambda_{m+1})}q_{s(\zeta_m)s(\eta_m)}q_{r(\zeta_m)r(\eta_m)}
$$

$$
\cdots q_{s(\zeta_1)s(\eta_1)}q_{r(\zeta_1)r(\eta_1)}q_{r(\zeta_1)r(\eta_1)}q_{s(\zeta_1)s(\eta_1)}\cdots q_{r(\zeta_m)r(\eta_m)}q_{s(\zeta_m)s(\eta_m)}
$$

Now consider the term $q_{s(\zeta_1)s(\eta_1)}q_{r(\zeta_1)r(\eta_1)}q_{s(\zeta_1)s(\eta_1)}$. As ζ_1 varies over all edges such that $\zeta_1 \zeta_2 \ldots \zeta_m$ is an *m*-length path, as in the proof of [3.5,](#page-12-0) we can replace this term in the summation with $\sum_{k,l\in V} q_{ks(\eta_1)} q_{lr(\eta_1)} q_{ks(\eta_1)}$ which is 1 as $\sum_k q_{ki} = 1$ for any *i*. Therefore, the summation reduces to

$$
\sum_{\substack{\xi,\zeta\\d(\xi)\geq d(\zeta)}} q_{s(\xi_n)s(\lambda_n)} q_{r(\xi_n)r(\lambda_n)} \cdots q_{s(\xi_{m+1})s(\lambda_{m+1})} q_{r(\xi_{m+1})r(\lambda_{m+1})}
$$

$$
\sum_{\substack{d(\xi)\geq d(\zeta)\\s(\xi)=i}} q_{s(\zeta_m)s(\eta_m)} q_{r(\zeta_m)r(\eta_m)} \cdots q_{s(\zeta_2)s(\eta_2)} q_{r(\zeta_2)r(\eta_2)} q_{s(\zeta_2)s(\eta_2)}
$$

By using the same trick repeatedly, the above summation can further be reduced to

$$
\sum_{\xi,s(\xi)=i} q_{s(\xi_n)s(\lambda_n)} q_{r(\xi_n)r(\lambda_n)} \cdots q_{s(\xi_{m+1})s(\lambda_{m+1})} q_{r(\xi_{m+1})r(\lambda_{m+1})}
$$

Now note that in the above expression, $(r(\xi_n), s(\xi_{n-1}), \cdots, s(\xi_{m+1}), r(\xi_{m+1}))$ can be replaced by $(l_n, k_{n-1}, l_{n-1}, \cdots, k_{m+1}, l_{m+1})$ where l_i 's and k_i 's vary over the vertex set. This is because if $((k_{m+1}, l_{m+1}), \cdots, (k_{n-1}, l_{n-1}), (i, l_n))$ forms a path of degree *n* − *m* starting at *i* then the corresponding coefficients already appear in the summation. Otherwise, as $\lambda_{m+1} \cdots \lambda_n$ is an $(n-m)$ -length path,

$$
q_{is(\lambda_n)}q_{l_nr(\lambda_n)}\cdots q_{l_{m+1}r(\lambda_{m+1})}=0.
$$

Therefore, the coefficient of $\chi_{[i]}$ in the summation becomes

$$
\sum_{\substack{l_n, k_{n-1}, \ldots, k_{m+1}, l_{m+1} \\ s(\xi) = s(\xi_n) = i}} q_{s(\xi_n)s(\lambda_n)} q_{l_n r(\lambda_n)} \cdots q_{k_{m+1}s(\lambda_{m+1})} q_{l_{m+1} r(\lambda_{m+1})} = q_{is(\lambda)}.
$$

Now consider the sub-case where there is no path β such that $\lambda = \eta \beta$ which means there exists $k \in \{1, 2, \dots, m\}$ such that $\lambda_k \neq \eta_k$ and hence $\pi(S_\lambda^*)\chi_{[\eta]} =$ 0. For simplicity let's assume that $\lambda_1 \neq \eta_1$. The case $k > 1$ can be proved along the same line of argument. Using the same computation as above the term $(\pi \otimes [.))\alpha(S^*_\lambda)U(\chi_{[\eta]})$ in the LHS is expanded as,

$$
\rho(\Lambda)^{-d/2} \sum_{\substack{\xi,\zeta\\d(\xi)\geq d(\zeta)}} \chi_{[s(\xi)]} \otimes q_{s(\xi_n)s(\lambda_n)} q_{r(\xi_n)r(\lambda_n)} \cdots q_{s(\xi_{m+1})s(\lambda_{m+1})} q_{r(\xi_{m+1})r(\lambda_{m+1})}
$$

$$
q_{s(\zeta_m)s(\lambda_m)} q_{r(\zeta_m)r(\lambda_m)} \cdots q_{s(\zeta_1)s(\lambda_1)} q_{r(\zeta_1)r(\lambda_1)} q_{r(\zeta_1)r(\eta_1)}
$$

$$
q_{s(\zeta_1)s(\eta_1)} \cdots q_{r(\zeta_m)r(\eta_m)} q_{s(\zeta_m)s(\eta_m)}.
$$

As $\lambda_1 \neq \eta_1$ that means either $r(\lambda_1) \neq r(\eta_1)$ or $s(\lambda_1) \neq s(\eta_1)$. Now for the case $r(\lambda_1) \neq r(\eta_1)$ the term $q_{r(\zeta_1)r(\lambda_1)}q_{r(\zeta_1)r(\eta_1)} = 0$ otherwise lets assume that $r(\lambda_1) =$ $r(\eta_1)$ and $s(\lambda_1) \neq s(\eta_1)$, now using the same argument as before, we can reduce the term $q_{s(\zeta_1)s(\lambda_1)}q_{r(\zeta_1)r(\eta_1)}q_{s(\zeta_1)s(\eta_1)}$ to $\sum_{k,l} q_{ks(\lambda_1)}q_{lr(\eta_1)}q_{ks(\eta_1)} = \sum_k$ $\sum_{k} q_{ks(\lambda_1)} q_{ks(\eta_1)} = 0$

and we are done.

Case 2: $d(\lambda) < d(\eta)$: Again as in case 1, it can be divided into two subcases: there is some β such that $\lambda \beta = \eta$ and there is no β such that $\lambda \beta = \eta$. In the first subcase, $\pi(S^*_{\lambda})(\chi_{[\eta]}) = \chi_{[\beta]}$. In the second subcase $\pi(S^*_{\lambda})(\chi_{[\eta]}) = 0$. In both the cases, using the same computational technique, it can be shown that

$$
U(\pi(S^*_\lambda)(\chi_{[\eta]}))=(\pi\otimes [.])\alpha(S^*_\lambda)U(\chi_{[\eta]}).
$$

Therefore, for any path $\eta \in \Lambda^*$, we have $(\pi \otimes [.)\alpha(S^*_{\lambda})U(\chi_{[\eta]}) = U(\pi(S_{\lambda^*}))(\chi_{[\eta]})$ and hence for any $q \in C(\text{Aut}^+(\Lambda)),$

$$
(\pi \otimes [.])\alpha(S_{\lambda}^*)\widetilde{U}(\chi_{[\eta]} \otimes q) = \widetilde{U}(\pi(S_{\lambda}^*) \otimes [1])(\chi_{\eta} \otimes q).
$$

By density of $\text{Sp}\{\chi_{[\eta]}\otimes q : \eta \in \Lambda^*, q \in C(\text{Aut}^+(\Lambda))\}\$ in the Hilbert space $\mathcal{H}\otimes$ $L^2(\text{Aut}^+(\Lambda))$ and unitarity of \widetilde{U} , we have for any $\lambda \in \Lambda^*$,

(3.5)
$$
(\pi \otimes [.])\alpha(S^*_{\lambda}) = \widetilde{U}(\pi(S^*_{\lambda}) \otimes [1])\widetilde{U}^* \in \mathcal{B}(\mathcal{H} \otimes L^2(\text{Aut}^+(\Lambda))).
$$

Using the same computational technique it can be shown that

(3.6)
$$
(\pi \otimes [.)\alpha(S_{\lambda}) = \widetilde{U}(\pi(S_{\lambda}) \otimes [1])\widetilde{U}^* \in \mathcal{B}(\mathcal{H} \otimes L^2(\text{Aut}^+(\Lambda)))
$$

for any $\lambda \in \Lambda^*$. Alternatively, one can see as before that Equation [3.6](#page-16-0) is equivalent to the following for $\lambda, \eta \in \Lambda^*$:

(3.7)
$$
(\pi \otimes [.)\alpha(S_{\lambda})U(\chi_{[\eta]}) = U(\pi(S_{\lambda}))(\chi_{[\eta]}).
$$

Again recalling the representation π of $C^*(\Lambda)$, we see that RHS of the above is equal to $\rho(\Lambda)^{d(\lambda/2)}U(\chi_{\{ \lambda n \}})$ whereas the LHS is equal to

$$
(3.8) \quad \rho(\Lambda)^{d(\mu)/2} \sum \chi_{[\mu\xi]} \otimes q_{r(\mu_1)r(\lambda_1)} q_{s(\mu_1)s(\lambda_1)} \cdots q_{r(\mu_n)r(\lambda_n)} q_{s(\mu_n)s(\lambda_n)} q_{r(\xi_1)r(\eta_1)}
$$

 $q_s(\xi_1)s(\eta_1) \cdots q_r(\xi_m)r(\eta_m)q_s(\xi_m)s(\eta_m)$

where $\mu = \mu_1 \dots \mu_n$ and $\xi = \xi_1 \dots \xi_m$;

$$
\alpha(S_{\lambda}) = \sum_{\mu} S_{\mu} \otimes q_{r(\mu_1)r(\lambda_1)} q_{s(\mu_1)s(\lambda_1)} \cdots q_{r(\mu_n)r(\lambda_n)} q_{s(\mu_n)s(\lambda_n)}
$$

and

$$
U(\chi_{[\eta]})=\sum_{\xi}\chi_{[\xi]}\otimes q_{r(\xi_1)r(\eta_1)}q_{s(\xi_1)s(\eta_1)}\cdots q_{r(\xi_m)r(\eta_m)}q_{s(\xi_m)s(\eta_m)}.
$$

A moment's reflection reveals that the coefficients of $\chi_{\lbrack\mu\xi\rbrack}$ in [3.8](#page-16-1) are the coefficients of $S_{\mu}S_{\xi}$ in the expression of $\alpha(S_{\lambda})\alpha(S_{\eta})$. Then the homomorphic property of α coupled with the fact that $d(\mu) = d(\lambda)$ for all μ proves Equation [3.7.](#page-16-2) Since both $(\pi \otimes [.)\alpha$ and $ad_{\widetilde{U}}$ are *C*^{*}-homomorphisms, we have the following from [3.5](#page-16-3) and [3.6:](#page-16-0)

$$
(\pi \otimes [.])\alpha(S_{\lambda}S_{\mu}^*) = \widetilde{U}(\pi(S_{\lambda}S_{\mu}^*) \otimes [1])\widetilde{U}^* \in \mathcal{B}(\mathcal{H} \otimes L^2(\text{Aut}^+(\Lambda))),
$$

for $\lambda, \mu \in \Lambda^*$. As $\text{Sp}\{S_\lambda S_\mu^* : \lambda, \mu \in \Lambda^*\}$ is dense in $C^*(\Lambda)$, using the continuity of the maps involved, we get

$$
(\pi \otimes [.]\alpha(a) = \widetilde{U}(\pi(a) \otimes [1])\widetilde{U}^* \in \mathcal{B}(\mathcal{H} \otimes L^2(\operatorname{Aut}^+(\Lambda))),
$$

for all $a \in C^*(\Lambda)$, proving the proposition. \square

Proof of Theorem [3.1](#page-8-1): Combining Proposition [3.6](#page-13-0) and Proposition [3.7,](#page-14-1) we see that the action α satisfies conditions (I) and (II) of Definition [2.18.](#page-8-0) This completes the proof of the main theorem.

4. A non-isometric action on the Cuntz algebra

We begin this section with an observation from the foregoing section. Note that in the proof of unitarity of \tilde{U} , the invariance of KMS-state under the given action α was crucially used (see Lemma [3.4\)](#page-12-1). So as mentioned in the introduction, it is natural to ask the question whether any object in the category of CQG's acting linearly on the graph C^* -algebra of a strongly connected graph in the sense of [\[14\]](#page-19-5) such that the KMS state is invariant act isometrically or not. In this section we give a counter example. To that end let us recall that the quantum unitary group U_n^+ acts linearly on the Cuntz algebra \mathcal{O}_n and it is the universal object in the category of CQG's acting linearly on \mathcal{O}_n such that the unique KMS-sate is invariant. Let us recall the action α : $\mathcal{O}_n \to \mathcal{O}_n \otimes C(U_n^+)$ given by (see [\[14\]](#page-19-5))

(4.1)
$$
\alpha(S_{e_i}) = \sum_{j=1}^{n} S_{e_j} \otimes q_{ji}
$$

where $\{e_1, \ldots, e_n\}$ are the loops of the graph of the Cuntz algebra ; S_{e_i} 's are the generating partial isometries; and $\{q_{ij}\}_{i,j=1,\dots,n}$ are the generators of $C(U_n^+)$. In the following we are going to retain the standard notations used in the previous section.

Theorem [4.1](#page-17-0). *The action* α *of* U_n^+ *on* \mathcal{O}_n *given* by 4.1 *is not isometric in the sense of Definition [2.18.](#page-8-0)*

Proof. To prove the theorem we have to show that there is no unitary co-representation of U_n^+ on the Hilbert space of the infinite path space (say $\mathcal H$ as before) commutting with the Dirac operator can implement the given action. To that end let us take a unitary co-representation \hat{U} that commutes with the Dirac operator D . We shall work with the corresponding \mathbb{C} -linear map $U : \mathcal{H} \to \mathcal{H} \overline{\otimes} C(U_n^+)$. Then *U* must preserve the eigen space \mathcal{R}_0 of *D* which is one-dimensional and spanned by the constant function 1 which is $\chi_{[v]}$ where *v* is the only vertex of the graph. Then $U(1) = 1 \otimes q$ for some unitary $q \in C(U_n^+)$. If possible, let *U* implement *α*. Then we have $(\pi \otimes [.)\alpha(S_{e_i})U(1) = U(\pi(S_{e_i}).1) = U(\chi_{[e_i]}).$ By expanding $\alpha(S_{e_i}),$ we get

$$
(\pi \otimes [.])\alpha(S_{e_i})U(1) = (\sum_j \pi(S_{e_j}) \otimes [q_{ji}]) (1 \otimes q)
$$

$$
= \sum_j \pi(S_{e_j}).1 \otimes q_{ji}q
$$

$$
= \sum_j \chi_{[e_j]} \otimes q_{ji}q
$$

Thus we have $U(\chi_{[e_i]}) = \sum_j \chi_{[e_j]} \otimes q_{ji}q$. But $1 = \sum_j \chi_{[e_j]}$ and therefore $U(1) =$ $U(\sum)$ $\sum_{i,j} \chi_{[e_i]}$) = $\sum_{i,j} \chi_{[e_j]} \otimes q_{ji} q$. As $U(1) = 1 \otimes q$, we get !

(4.2)
$$
\sum_{j} \chi_{[e_j]} \otimes \left(\sum_{i} q_{ji} q\right) = \sum_{j} \chi_{[e_j]} \otimes q.
$$

Now choose a point $x \in \Lambda^{\infty}$ such that $x \in [e_k]$ for a unique k. Then evaluating both sides of the Equation [4.2](#page-18-4) at the point *x*, we get $\sum_{i} q_{ki}q = q$ which, by unitarity of q, implies that $\sum_{i} q_{ki} = 1$, which is a contradiction as the generators of $C(U_n^+)$ do not satisfy these relations. \Box

Remark 4.2. (1) The quantum permutation group S_n^+ has a 'linear' action α on \mathcal{O}_n given by

$$
\alpha(S_{e_i}) = \sum_{j=1}^n S_{e_j} \otimes q_{ji},
$$

where $\{q_{ij}\}_{i,j=1,\dots,n}$ are the generators of $C(S_n^+)$. Then it can be verified that the unique KMS state of \mathcal{O}_n is invariant under the action α . Using the same computational technique and same line of arguments it can be shown that the action α is also isometric in the sense of Definition [2.18.](#page-8-0) The quantum group S_n^+ is a quantum symmetry object of the underlying graph of \mathcal{O}_n in the sense of Goswami et al. ([\[12\]](#page-19-19)) where they consider quantum symmetry of finite graphs possibly with multiple edges. It seems reasonable to expect that the quantum symmetry of the underlying graph of a graph *C* ∗ -algebra is well behaved in the context of quantum isometric actions with respect to the class of spectral triple on graph *C* ∗ -algebras constructed by Farsi et al. in [\[7\]](#page-19-10). On can explore further in this direction by considering graphs with possible multiple edges.

(2) Since the spectral triple is Θ-summable for generic choices of eigenvalues of the Dirac operator, one can consider quantum isometry in terms of a 'good Laplacian' in the sense of [\[10\]](#page-19-0) and possibly compare with our consideration.

Acknowledgement The first author would like to acknowledge the support provided by SERB MATRICS grant (MTR/2022/000515) provided by the government of India. Both the authors would like to acknowledge the support provided by the FIST grant (SR/FST/MS-II/2019/51) provided by the government of India. The first author would like to thank Debashish Goswami for some helpful inputs.

REFERENCES

- [1] Teodor Banica, *Quantum automorphism groups of homogeneous graphs*, J. Funct. Anal. **224** (2005), no. 2, 243–280, DOI 10.1016/j.jfa.2004.11.002. MR2146039
- [2] Suvrajit Bhattacharjee and Soumalya Joardar, *Equivariant C*[∗] *-correspondences and compact quantum group actions on Pimsner algebras*, Canadian Journal of Mathematics, posted on 2023, 1–40, DOI 10.4153/S0008414X23000810.
- [3] Jyotishman Bhowmick and Debashish Goswami, *Quantum group of orientationpreserving Riemannian isometries*, J. Funct. Anal. **257** (2009), no. 8, 2530–2572, DOI 10.1016/j.jfa.2009.07.006. MR2555012
- [4] Kenny De Commer, *Actions of compact quantum groups*, arXiv (2016), available at <1604.00159>.

20 JOARDAR AND SHARMA

- [5] Caterina Consani and Matilde Marcolli, *Noncommutative geometry, dynamics, and* ∞*-adic Arakelov geometry*, Selecta Math. (N.S.) **10** (2004), no. 2, 167–251, DOI 10.1007/s00029-004- 0369-3. MR2080121
- [6] Roberto Conti and Stefano Rossi, *Groups of isometries of the Cuntz algebras*, Doc. Math. **26** (2021), 1799–1815. MR4353340
- [7] Carla Farsi, Elizabeth Gillaspy, Antoine Julien, Sooran Kang, and Judith Packer, *Spectral triples for higher-rank graph C*[∗] *-algebras*, Math. Scand. **126** (2020), no. 2, 321–338, DOI 10.7146/math.scand.a-119260. MR4102567
- [8] Carla Farsi, Elizabeth Gillaspy, Sooran Kang, and Judith A. Packer, *Separable representations, KMS states, and wavelets for higher-rank graphs*, J. Math. Anal. Appl. **434** (2016), no. 1, 241–270, DOI 10.1016/j.jmaa.2015.09.003. MR3404559
- [9] Fulton, *The quantum automorphism group and directed trees*, Ph. D. thesis, Virginia (2006).
- [10] Debashish Goswami, *Quantum group of isometries in classical and noncommutative geometry*, Comm. Math. Phys. **285** (2009), no. 1, 141–160, DOI 10.1007/s00220-008-0461-1. MR2453592
- [11] Debashish Goswami and Soumalya Joardar, *Non-existence of faithful isometric action of compact quantum groups on compact, connected Riemannian manifolds*, Geom. Funct. Anal. **28** (2018), no. 1, 146–178, DOI 10.1007/s00039-018-0437-z. MR3777415
- [12] D. Goswami and Sk. Asfaq Hossain, *Quantum symmetry in multigraphs*, https://doi.org/10.48550/arXiv.2302.08726.
- [13] Astrid an Huef, Marcelo Laca, Iain Raeburn, and Aidan Sims, *KMS states on the C*∗ *-algebras of finite graphs*, J. Math. Anal. Appl. **405** (2013), no. 2, 388–399, DOI 10.1016/j.jmaa.2013.03.055. MR3061018
- [14] Soumalya Joardar and Arnab Mandal, *Quantum symmetry of graph C*∗*-algebras associated with connected graphs*, Infin. Dimens. Anal. Quantum Probab. Relat. Top. **21** (2018), no. 3, 1850019, 18, DOI 10.1142/S0219025718500194. MR3860252
- [15] _____, *Invariance of KMS states on graph C*[∗]-algebras under classical and quantum sym*metry*, Proc. of the Edin. Math. Soc. (2) **64** (2021), 762-778.
- [16] , *Quantum symmetry of graph C*[∗] *-algebras at critical inverse temperature*, Studia Math. **256** (2021), no. 1, 1-20, DOI 10.4064/sm190102-30-9. MR4159553
- [17] Ujjal Karmakar and Arnab Mandal, *Unitary Easy Quantum Groups Arising as Quantum Symmetries of Graph C*-algebras* (2024), available at <2406.13484>.
- [18] Alex Kumjian, David Pask, and Iain Raeburn, *Cuntz-Krieger algebras of directed graphs*, Pacific J. Math. **184** (1998), no. 1, 161–174, DOI 10.2140/pjm.1998.184.161. MR1626528
- [19] Efton Park, *Isometries of noncommutative metric spaces*, Proc. Amer. Math. Soc. **123** (1995), no. 1, 97–105, DOI 10.2307/2160614. MR1213868
- [20] Ann Maes and Alfons Van Daele, *Notes on compact quantum groups*, Nieuw Arch. Wisk. (4) **16** (1998), no. 1-2, 73–112. MR1645264
- [21] Sergey Neshveyev and Lars Tuset, *Compact quantum groups and their representation cat*egories, Cours Spécialisés [Specialized Courses], vol. 20, Société Mathématique de France, Paris, 2013. MR3204665
- [22] David Pask and Alex Kumjian, *Higher rank graph C*[∗] *-algebras*, New York J. Math. **6** (2020), 1-20.
- [23] Iain Raeburn, *Graph algebras*, CBMS Regional Conference Series in Mathematics, vol. 103, Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 2005. MR2135030
- [24] Simon Schmidt and Moritz Weber, *Quantum symmetries of graph C*[∗] *-algebras*, Canad. Math. Bull. **61** (2018), no. 4, 848–864, DOI 10.4153/CMB-2017-075-4. MR3846752
- [25] Shuzhou Wang, *Quantum symmetry groups of finite spaces*, Comm. Math. Phys. **195** (1998), no. 1, 195–211, DOI 10.1007/s002200050385. MR1637425
- [26] S. L. Woronowicz, *Compact quantum groups*, Symétries quantiques (Les Houches, 1995), North-Holland, Amsterdam, 1998, pp. 845–884. MR1616348

Department of Mathematics and Statistics, Indian Institute of Science Education and Research Kolkata, Mohanpur - 741246, West Bengal, India

Email address: soumalya@iiserkol.ac.in

Department of Mathematics and Statistics, Indian Institute of Science Education and Research Kolkata, Mohanpur - 741246, West Bengal, India

Email address: js20ip007@iiserkol.ac.in