ON POSITIVE JACOBI MATRICES WITH COMPACT INVERSES

PAVEL ŠŤOVÍČEK AND GRZEGORZ ŚWIDERSKI

Abstract. We consider positive Jacobi matrices J with compact inverses and consequently with purely discrete spectra. A number of properties of the corresponding sequence of orthogonal polynomials is studied including the convergence of their zeros, the vague convergence of the zero counting measures and of the Christoffel–Darboux kernels on the diagonal. Particularly, if the inverse of J belongs to some Schatten class, we identify the asymptotic behaviour of the sequence of orthogonal polynomials and express it in terms of its regularized characteristic function. In the even more special case when the inverse of J belongs to the trace class we derive various formulas for the orthogonality measure, eigenvectors of J as well as for the functions of the second kind and related objects. These general results are given a more explicit form in the case when -J is a generator of a Birth–Death process. Among others we provide a formula for the trace of the inverse of J. We illustrate our results by introducing and studying a modification of q-Laguerre polynomials corresponding to a determinate moment problem.

1. Introduction

Given two sequences $a = (a_n : n \in \mathbb{N}_0)$, $b = (b_n : n \in \mathbb{N}_0)$ of positive and real numbers, respectively, we define an infinite tridiagonal matrix by

$$\mathcal{J} = \begin{pmatrix} b_0 & a_0 & & \\ a_0 & b_1 & a_1 & \\ & a_1 & b_2 & \ddots \\ & & \ddots & \ddots \end{pmatrix}.$$

Action of \mathcal{J} on any sequence of finite support treated as a column vector is defined by the formal matrix multiplication. Let the operator J be its closure in the Hilbert space $\ell^2(\mathbb{N}_0)$. Let us recall that

$$\ell^2(\mathbb{N}_0) = \left\{ x = (x_0, x_1, \dots) : \sum_{n=0}^{\infty} |x_n|^2 < \infty \right\}, \quad \langle x, y \rangle = \sum_{n=0}^{\infty} \overline{x_n} y_n.$$

For any $n \in \mathbb{N}_0$ by e_n we shall denote the sequence with one on the *n*th position and zero elsewhere. Then $(e_n : n \in \mathbb{N}_0)$ is an orthonormal basis of $\ell^2(\mathbb{N}_0)$, the so called standard (or canonical) orthonormal basis.

The symmetric operator J has always self-adjoint extensions on $\ell^2(\mathbb{N}_0)$. Let \tilde{J} be any such extension. Define a probability measure on the Borel subsets of the real line by

$$\mu(\cdot) = \langle \boldsymbol{e}_0, E_{\tilde{I}}(\cdot)\boldsymbol{e}_0 \rangle, \tag{1.1}$$

where $E_{\tilde{J}}$ is the projection-valued spectral measure of \tilde{J} , i.e. $\tilde{J} = \int_{\mathbb{R}} x \, dE_{\tilde{J}}(x)$ in the weak sense. The support of μ coincides with the spectrum $\sigma(\tilde{J})$.

Associated with \mathcal{J} is the sequence of polynomials $(P_n:n\in\mathbb{N}_0)$ satisfying the three-term recurrence relation

$$zP_n(z) = a_{n-1}P_{n-1}(z) + b_nP_n(z) + a_nP_{n+1}(z), \quad n \ge 1$$
(1.2)

with the initial conditions

$$P_0(z) \equiv 1, \quad P_1(z) = \frac{z - b_0}{a_0}.$$
 (1.3)

1

²⁰²⁰ Mathematics Subject Classification. Primary 47B36; Secondary 42C05, 33C45.

Key words and phrases. Orthogonal polynomials; Jacobi matrix; functions of the second kind; zero counting measures; Christoffel–Darboux kernels; *q*-Laguerre polynomials.

It turns out that $(P_n : n \ge 0)$ is a sequence of orthonormal polynomials in $L^2(\mu)$, i.e.

$$\int_{\mathbb{R}} \overline{P_n(x)} P_m(x) d\mu(x) = \begin{cases} 1 & n = m, \\ 0 & \text{otherwise.} \end{cases}$$

It is well-known that the Hamburger moment problem for such an orthonormal polynomial sequence $(P_n(x) : n \ge 0)$ is determinate, i.e. the sequence is orthonormal with respect to a unique probability measure on the real line, if and only if the Jacobi operator J itself is self-adjoint (see e.g. [21, Theorem 6.10]).

For our purposes particularly interesting will be those Jacobi matrices which are related to generators of Birth-Death processes. In that case

$$a_n = \sqrt{\lambda_n \mu_{n+1}}, \quad b_n = \lambda_n + \mu_n,$$
 (1.4)

where $(\lambda_n : n \in \mathbb{N}_0)$ and $(\mu_n : n \in \mathbb{N}_0)$ are sequences such that $\lambda_n > 0$, $\mu_{n+1} > 0$ for any $n \in \mathbb{N}_0$, and $\mu_0 \ge 0$. For the motivation of Jacobi parameters (1.4) see e.g. [23, Section 5.1]. For probabilistic applications of -J see e.g. [10, Chapter 3].

In this paper we focus mainly on the case when J is positive definite and the inverse of its specific self-adjoint extension, the so-called *Friedrichs extension*, belongs to the trace class or, more generally, to a Schatten class. Some first results on this topic have been obtained in papers [25, 26]. Here these results are substantially extended and generalized in several directions.

In Sections 2 and 3 we assume that J is positive definite and J^{-1} belongs to the trace class. This situation is firstly studied, in Section 2, on a general level. Naturally, under the aforementioned assumptions the spectrum of J is pure point and also the orthogonality measure for $(P_n(x): n \ge 0)$ is purely atomic. As a first step, we derive a convenient criterion, expressed in terms of the real sequence $(P_n(0): n \ge 0)$, which implies the desired properties of J. Then we derive several formulas for the orthogonality measure and for eigenvectors of J. The characteristic function of J and its associated semi-infinite principal submatrices as well as the functions of the second kind play an important role in this study.

Then, in Section 3, we specialize these general results to Jacobi operators corresponding to generators of Birth-Death processes. This is a broad class of Jacobi operators which are widely used in various applications. For this type of Jacobi operators we derive a criterion for the trace class inverse, a formula for the characteristic function and an additional formula related to the orthogonality measure. This way we extend substantially the results obtained earlier in [26] but only for a very special class of Jacobi operators.

In Section 4 we advance our studies by considering a more general case when the inverse of the Friedrichs extension of J is supposed to be compact. In Section 4.2 we study weak convergence of the zero counting measures of P_n and in Section 4.3 the weak behaviour of the Christoffel–Darboux kernels on the diagonal. In Section 4.4, under the additional assumption that the inverse of the Friedrichs extension of J belongs to an arbitrary Schatten class, the regularized characteristic function of the Jacobi operator is defined and it is shown to be the locally uniform limit of a certain sequence of elementary analytic functions. Thus we generalize a result from [25] where the characteristic function of J has been introduced and studied under the assumption that J^{-1} is a trace class operator. Moreover, in Section 4.5 we show that for Jacobi operators which are not semibounded a similar convergence does not hold in general.

Finally as an application, in Section 5, we introduce and study an alternative q-version to the Laguerre polynomials $(L_n^{(\alpha)}(x):n\geq 0)$ which we call the modified q-Laguerre polynomials and denote $(\tilde{L}_n^{(\alpha)}(x;q):n\geq 0)$. The particular case with $\alpha=0$ has been studied in [26]. A distinguished feature of these newly introduced q-Laguerre polynomials is that the respective Hamburger moment problem is determinate. Furthermore, they can be shown to correspond to the generators of a Birth-Death process. They are related to the (standard) q-Laguerre polynomials, as introduced and studied in [20], by a construction based on the notion of quasi-orthogonal sequence [4]. The q-Laguerre polynomials are known explicitly and, owing to this kind of construction, the same is in principle true for the modified q-Laguerre polynomials. Notably, the moment sequence for $(\tilde{L}_n^{(\alpha)}(x;q):n\geq 0)$ can be derived explicitly and the orthogonality measure turns out to be supported on the roots of the (appropriately rescaled) q-Bessel function $J_{\alpha+1}^{(2)}(2\sqrt{z};q)$.

2. More on positive Jacobi operators with a trace class inverse

2.1. A criterion for the trace class. In Sections 2 and 3 we assume that J is positive definite and J^{-1} is a trace class operator. This is what has been assumed also in the earlier paper [25]. In this subsection we prove a convenient criterion that guarantees J^{-1} to exist and to belong to the trace class. It turns out that with this criterion the assumption on positive definiteness of J can be relaxed so that it is fully sufficient to assume that J is just positive-semidefinite meaning that its quadratic form is non-negative. The positive definiteness is then obtained as a straightforward consequence. This simplifies the discussion notably.

In the sequel, the orthogonality measure for the orthonormal polynomial sequence $(P_n(x) : n \ge 0)$ is called μ and $\varrho(J)$ stands for the resolvent set of J while $\sigma(J)$ denotes its spectrum.

Theorem 2.1. Assume that

$$\sum_{n=0}^{\infty} P_n(0)^2 = \infty {(2.1)}$$

and J is positive-semidefinite. Then J is self-adjoint and invertible. Moreover, J^{-1} belongs to the trace class if and only if

$$-\sum_{n=0}^{\infty} \left(\sum_{j=n}^{\infty} \frac{1}{a_j P_j(0) P_{j+1}(0)} \right) P_n(0)^2 < \infty.$$
 (2.2)

If so then $Tr(J^{-1})$ equals the sum in (2.2) (including the sign).

Proof. It follows from (2.1) that the Hamburger moment problem for J is determinate and so J is self-adjoint, and also that $0 \notin \sigma_p(J)$, i.e. J^{-1} exists. The positivity of J then implies that the support of the spectral measure of J is contained in the closed positive real half-line, and the same is true for the orthogonality measure μ . Then all roots of the polynomials $P_n(x)$, $n \ge 1$, are necessarily contained in the interior of the same interval (see e.g. [7, Theorem I.5.2]), i.e. in $(0, +\infty)$.

Moreover, recalling that $(a_n : n \ge 0)$ is a sequence of positive numbers, we have $\operatorname{sgn} P_n(0) = (-1)^n$ for any $n \ge 0$, and for any $n \ge 1$ the functions $P_n(x)^2$ are all decreasing on the interval $(-\infty, 0]$.

According to equations (28) and (7) in [25], for all $z \in \rho(J)$,

$$-\left(\sum_{j=n}^{\infty} \frac{1}{a_j P_j(z) P_{j+1}(z)}\right) P_n(z)^2 = \langle \boldsymbol{e}_n, (J - zI)^{-1} \boldsymbol{e}_n \rangle$$
$$= \int_{\mathbb{R}} \frac{P_n(x)^2}{x - z} \, \mathrm{d}\mu(x).$$

Then, since μ is supported on the positive half-line, we can apply twice the monotone convergence theorem and get

$$\int_{\mathbb{R}} \frac{P_n(x)^2}{x} d\mu(x) = \lim_{\varepsilon \to 0+} \int_{\mathbb{R}} \frac{P_n(x)^2}{x + \varepsilon} d\mu(x)$$

$$= -\lim_{\varepsilon \to 0+} \left(\sum_{j=n}^{\infty} \frac{1}{a_j P_j(-\varepsilon) P_{j+1}(-\varepsilon)} \right) P_n(-\varepsilon)^2$$

$$= -\left(\sum_{j=n}^{\infty} \frac{1}{a_j P_j(0) P_{j+1}(0)} \right) P_n(0)^2.$$

Thus, for any given $n \in \mathbb{N}_0$, $\boldsymbol{e}_n \in \mathrm{Dom}(J^{-1/2})$ if and only if

$$||J^{-1/2}\boldsymbol{e}_n||^2 = \int \frac{P_n(x)^2}{x} d\mu(x) = -\left(\sum_{j=n}^{\infty} \frac{1}{a_j P_j(0) P_{j+1}(0)}\right) P_n(0)^2 < \infty.$$
 (2.3)

If J^{-1} belongs to the trace class then $J^{-1/2}$ is a Hilbert-Schmidt operator and, in regard of (2.3), condition (2.2) holds.

Conversely, if (2.2) is true then $\forall n \in \mathbb{N}_0$, $\boldsymbol{e}_n \in \mathrm{Dom}(J^{-1/2})$, and, by a standard estimate, $J^{-1/2}$ is bounded on span{ $\boldsymbol{e}_n : n \geq 0$ }. In detail, $\forall \boldsymbol{v} \in \mathrm{span}\{\boldsymbol{e}_n : n \geq 0\}$,

$$\|J^{-1/2}v\|^2 = \sum_{n=0}^{\infty} |\langle \boldsymbol{e}_n, J^{-1/2}v \rangle|^2 = \sum_{n=0}^{\infty} |\langle J^{-1/2}\boldsymbol{e}_n, v \rangle|^2 \le \sum_{n=0}^{\infty} \|J^{-1/2}\boldsymbol{e}_n\|^2 \|v\|^2.$$

Consequently, since $J^{-1/2}$ is closed, it is necessarily an everywhere defined bounded operator. And again by (2.2), $J^{-1/2}$ is a Hilbert-Schmidt operator, J^{-1} is a trace class operator and

$$\operatorname{Tr}(J^{-1}) = \sum_{n=0}^{\infty} \langle \boldsymbol{e}_n, J^{-1} \boldsymbol{e}_n \rangle = \sum_{n=0}^{\infty} \|J^{-1/2} \boldsymbol{e}_n\|^2 = -\sum_{n=0}^{\infty} \left(\sum_{j=n}^{\infty} \frac{1}{a_j P_j(0) P_{j+1}(0)} \right) P_n(0)^2.$$

This concludes the proof.

2.2. Functions of the second kind, the orthogonality measure. Using the same notation as in Section 1, here we fully focus on the case when the Jacobi operator J is positive definite and its inverse belongs to the trace class. This also means that 0 belongs to the resolvent set of J. Moreover, then the spectrum of J is discrete and consists of simple eigenvalues of J only. The eigenvalues can be enumerated, $\sigma(J) = \{\zeta_n : n \ge 1\}$, so that (ζ_n) is a strictly increasing sequence of positive numbers, and $\lim_{n\to\infty} \zeta_n = +\infty$.

Weyl's function w(z) associated with J is by definition the Stieltjes transformation of the measure μ ,

$$w(z) = \int_{\mathbb{R}} \frac{\mathrm{d}\mu(x)}{x - z} = \langle \boldsymbol{e}_0, (J - z)^{-1} \boldsymbol{e}_0 \rangle.$$

More generally, the so called functions of the second kind are defined as

$$w_k(z) = \int_{\mathbb{R}} \frac{P_k(x)}{x - z} \, \mathrm{d}\mu(x) = \langle \boldsymbol{e}_k, (J - z)^{-1} \boldsymbol{e}_0 \rangle, \quad k \ge 0.$$
 (2.4)

In particular, $w(z) \equiv w_0(z)$. In our case, $w_k(z)$ are meromorphic functions on \mathbb{C} , its singular points are located at the eigenvalues of J, and all of them are simple poles. The measure μ can be reconstructed from Weyl's function by evaluating the residues,

$$\forall \zeta \in \sigma(J), \ \mathop{\mathrm{Res}}_{z=\zeta} w(z) = -\mu(\{\zeta\}).$$

The characteristic function of J can be naturally defined as

$$\mathfrak{F}(z) = \prod_{n=1}^{\infty} \left(1 - \frac{z}{\zeta_n} \right).$$

A more general situation will be considered in Subsection 4.4 and this definition is a particular case of (4.20) for k = 1. $\mathfrak{F}(z)$ is an entire function whose zeroes are all simple and coincide with the eigenvalues of J. It is known that

$$\mathfrak{F}(z) = 1 - z \sum_{n=0}^{\infty} w_n(0) P_n(z), \tag{2.5}$$

and the series converges locally uniformly in $z \in \mathbb{C}$, see [25, Thm. 2].

Let

$$\mathfrak{W}(z) = w(z)\mathfrak{F}(z). \tag{2.6}$$

Then, clearly, all singularities of $\mathfrak{W}(z)$ are removable which is to say that $\mathfrak{W}(z)$ is an entire function. We have, for every $\zeta \in \sigma(J)$,

$$\mu(\{\zeta\}) = -\lim_{z \to \zeta} (z - \zeta) w(z) = -\lim_{z \to \zeta} \frac{(z - \zeta) \mathfrak{W}(z)}{\mathfrak{F}(z) - \mathfrak{F}(\zeta)}.$$

Hence

$$\forall \zeta \in \sigma(J), \ \mu(\{\zeta\}) = -\frac{\mathfrak{W}(\zeta)}{\mathfrak{F}'(\zeta)}. \tag{2.7}$$

It may be worthwhile to recall, too, that

$$\forall \zeta \in \sigma(J), \ \mu(\{\zeta\}) = \left(\sum_{n=0}^{\infty} P_n(\zeta)^2\right)^{-1},$$

see e.g. [22, Corollary 2.6, p. 45] or [1, Thm. 2.5.3].

As an auxiliary step in the course of the proof of Theorem 2 in [25], the following result has been derived but not stated there as an independent proposition. Here it is recalled as a lemma.

Lemma 2.2. For $n \geq 0$,

$$w_n(0)P_n(z) = \sqrt{\kappa_n} \langle \boldsymbol{e}_n, (I - zK)^{-1} \boldsymbol{k} \rangle,$$

where

$$\kappa_n = w_n(0)P_n(0), \quad \mathbf{k} = (\sqrt{\kappa_0}, \sqrt{\kappa_1}, \sqrt{\kappa_2}, \ldots)^T,$$

and

$$K_{m,n} = \begin{cases} \frac{P_n(0)}{P_m(0)} \sqrt{\frac{\kappa_m}{\kappa_n}} (w_m(0) P_n(0) - w_n(0) P_m(0)) & \text{for } m > n \\ 0 & \text{otherwise.} \end{cases}$$
(2.8)

Hence, in view of (2.5), the characteristic function of J reads

$$\mathfrak{F}(z) = 1 - z\mathbf{k}^T \cdot (\mathrm{Id} - zK)^{-1} \cdot \mathbf{k}. \tag{2.9}$$

Remark. The inverse operator $(\operatorname{Id} - zK)^{-1}$ can be evaluated as a geometric series, and doing so the scalar product $\langle e_n, (\operatorname{Id} - zK)^{-1} k \rangle$ reduces to a finite sum.

Further we recall that the sequence of so-called *orthogonal polynomials of the second kind*, $(Q_n : n \ge 0)$, is defined by the three-term recurrence relation

$$Q_0(x) = 0, \quad Q_1(x) = \frac{1}{a_0},$$

$$a_{n-1}Q_{n-1}(x) + b_nQ_n(x) + a_nQ_{n+1}(x) = xQ_n(x), \quad n \ge 1,$$

and it holds true that

$$Q_n(x) = \int_{\mathbb{R}} \frac{P_n(x) - P_n(u)}{x - u} \, \mathrm{d}\mu(u), \quad n \ge 0.$$

In the beginning of the proof of Theorem 2 in [25], there was shown that

$$\mathfrak{F}(z) = \lim_{n \to \infty} \frac{P_n(z)}{P_n(0)} \tag{2.10}$$

locally uniformly on \mathbb{C} . For a different proof, see Corollary 4.10 below.

In [3], a generalization of Markov's Theorem has been derived which is applicable to any Jacobi matrix \mathcal{J} which is the matrix in the canonical basis of an essentially self-adjoint operator in $\ell^2(\mathbb{N}_0)$. Denote by \mathcal{Z}_n the set of roots of $P_n(x)$ and put

$$\Lambda := \bigcap_{N=1}^{\infty} \overline{\bigcup_{n=N}^{\infty} \mathcal{Z}_n}.$$

Then supp $\mu \subset \Lambda$ and

$$\lim_{n \to \infty} \frac{Q_n(z)}{P_n(z)} = -w(z) \tag{2.11}$$

locally uniformly on $\mathbb{C} \setminus \Lambda$.

As a consequence of Markov's Theorem it has been observed in [25, Eq. (28)] that

$$w_n(z) = -P_n(z) \sum_{j=n}^{\infty} \frac{1}{a_j P_j(z) P_{j+1}(z)}, \quad n \ge 0,$$
 (2.12)

holds on $\mathbb{C} \setminus \Lambda$, and the series converges locally uniformly on this set. Furthermore, combining (2.10) and (2.11) with the defining relation (2.6) we immediately get

$$\mathfrak{W}(z) = -\lim_{n \to \infty} \frac{Q_n(z)}{P_n(0)}$$

locally uniformly on $\mathbb{C} \setminus \Lambda$.

By the assumptions formulated in the beginning of this section, $J \ge \gamma$ for some $\gamma > 0$ (of course, the best choice is $\gamma = \zeta_1$).

By $J^{[k]}$, $k \ge 1$, let us denote the Jacobi matrix operator obtained from J by removing the first k rows and columns. Then $J^{[k]}$ obeys the same assumptions as J does, see [26, Prop. 4]. All objects associated with $J^{[k]}$ will be distinguished by the upper index [k]. Particularly this concerns the orthonormal polynomial sequence $(P_n^{[k]}(x):n\ge 0)$ and the functions of the second kind $(w_n^{[k]}(z):n\ge 0)$, see (2.4), and also the characteristic function $\mathfrak{F}^{[k]}(z)$. The upper index [0] corresponds to objects which are related to the original Jacobi operator J.

Some attention should be paid to the polynomials of the second kind, however. As in [25], we define them so that

$$Q_0^{[k]}(x) = Q_1^{[k]}(x) = \dots = Q_k^{[k]}(x) = 0, \ Q_{k+n}^{[k]}(x) = \frac{1}{a_k} P_{n-1}^{[k+1]}(x) \text{ for } n \ge 1.$$
 (2.13)

Let us recall that there exists a generalization of Markov's theorem (2.11) due to Van Assche, see [24, Thm. 2], which claims that, for $k \ge 0$,

$$w_k(z) = -\lim_{n \to \infty} \frac{Q_n^{[k]}(z)}{P_n(z)} \quad \text{locally uniformly in } z \in \mathbb{C} \setminus [\gamma, +\infty). \tag{2.14}$$

It is also useful to recall that

$$\forall k \ge 0, \ w_k(z) = w(z)P_k(z) + Q_k(z),$$
 (2.15)

(see [25, Eq. (13)]). This relation should be understood as an equality between two meromorphic functions on \mathbb{C} . Moreover, the functions of the second kind form a square summable sequence for all z from the resolvent set, i.e.

$$\forall z \in \rho(J), \ (w_0(z), w_1(z), w_2(z), \dots)^T \in \ell^2(\mathbb{N}_0).$$
 (2.16)

Furthermore, by [25, Eq. (26)],

$$\forall m \ge n \ge 0, \ Q_m^{[n]}(x) = Q_m(x)P_n(x) - P_m(x)Q_n(x). \tag{2.17}$$

Theorem 2.3. It holds true that

$$\forall k \ge 0, \ w_k^{[1]}(z) = -\frac{w_{k+1}(z)}{a_0 w(z)},\tag{2.18}$$

and

$$\forall k \ge 0, \ w_k(z) = \frac{w_k(0)\mathfrak{F}^{[k+1]}(z)}{\mathfrak{F}(z)}.$$
 (2.19)

These equations are both understood as equalities between meromorphic functions on \mathbb{C} .

Proof. In regard of (2.14) and (2.13),

$$w_k^{[1]}(z) = -\lim_{m \to \infty} \frac{Q_m^{[k+1]}(z)}{P_{m-1}^{[1]}(z)} = -\lim_{m \to \infty} \frac{Q_m^{[k+1]}(z)}{a_0 Q_m(z)}$$

locally uniformly on $\mathbb{C} \setminus [\gamma, +\infty)$. Using (2.17), (2.11) and (2.15) we get

$$\begin{split} w_k^{[1]}(z) &= \frac{1}{a_0} \left(-P_{k+1}(z) + Q_{k+1}(z) \lim_{m \to \infty} \frac{P_m(z)}{Q_m(z)} \right) \\ &= -\frac{1}{a_0} \left(P_{k+1}(z) + \frac{Q_{k+1}(z)}{w(z)} \right) \\ &= -\frac{w_{k+1}(z)}{a_0 w(z)}. \end{split}$$

Now it suffices to observe that, under our assumptions, all functions of the second kind are non-zero meromorphic functions on \mathbb{C} .

Similarly, taking into account also (2.10),

$$w_{k}(z) = -\lim_{n \to \infty} \frac{Q_{n}^{[k]}(z)}{P_{n}(z)}$$

$$= -\lim_{n \to \infty} \frac{P_{n-k-1}^{[k+1]}(z)}{P_{n-k-1}^{[k+1]}(0)} \cdot \frac{Q_{n}^{[k]}(0)}{P_{n}(0)} \cdot \frac{P_{n}(0)}{P_{n}(z)}$$

$$= \mathfrak{F}^{[k+1]}(z)w_{k}(0)\frac{1}{\mathfrak{F}(z)}.$$

Again, the limit is locally uniform on $\mathbb{C} \setminus [\gamma, +\infty)$. Finally notice that the involved characteristic functions are all entire functions.

Remark. One can also prove (2.19) by mathematical induction on k, with the aid of (2.18).

Corollary 2.4. It holds true that

$$w_k(0)\mathfrak{F}^{[k+1]}(z) = w(0)P_k(z)\mathfrak{F}^{[1]}(z) + Q_k(z)\mathfrak{F}(z) \quad \text{for } k \ge 0$$
 (2.20)

and all $z \in \mathbb{C}$.

Proof. In equation (2.15) we substitute for $w_k(z)$ and $w(z) \equiv w_0(z)$ from (2.19) and get (2.20).

Corollary 2.5. For every eigenvalue ζ of J, a corresponding eigenvector can be chosen as the column vector

$$(w(0)\mathfrak{F}^{[1]}(\zeta), w_1(0)\mathfrak{F}^{[2]}(\zeta), w_2(0)\mathfrak{F}^{[3]}(\zeta), \dots)^T.$$
(2.21)

Proof. For every $z \in \mathbb{C}$, the column vector

$$P(z) = (P_0(z), P_1(z), P_2(z), \dots)^T$$

is a formal eigenvector of J, i.e. a solution of the equation $(\mathcal{J} - z \operatorname{Id}) P(z) = 0$ in \mathbb{C}^{∞} , and $x \in \mathbb{R}$ is an eigenvalue of J if and only if P(x) has a finite ℓ^2 norm.

Letting $z = \zeta \in \sigma(J)$ in (2.20) we see that the column vector (2.21) is a multiple of $P(\zeta)$.

Moreover, the first entry in (2.21), i.e. $w(0)\mathfrak{F}^{[1]}(\zeta)$, is necessarily non-zero. Actually, it is known that the eigenvalues of J and $J^{[1]}$ interlace [17, Corollary 10.23]. Hence the zeroes of $\mathfrak{F}(z)$ and $\mathfrak{F}^{[1]}(z)$ interlace. But we have $\mathfrak{F}(\zeta)=0$. This is also seen from (2.19) for k=0 since Weyl's function has a singularity at $z=\zeta$ which is not removable and the zero of $\mathfrak{F}(z)$ at $z=\zeta$ is simple.

Remark. Note that the column vector (2.21) has a finite ℓ^2 norm for all $\zeta \in \mathbb{C}$ and not only for the eigenvalues of J. If ζ is not an eigenvalue of J then it follows from (2.16) and (2.19).

Theorem 2.6. We have that

$$\mathfrak{W}(z) = w(0) + z \sum_{n=1}^{\infty} w_n(0) Q_n(z), \tag{2.22}$$

and the series converges locally uniformly on \mathbb{C} .

Proof. By Theorem 2 in [25], for every Jacobi operator J fulfilling our assumptions it holds true that the series $\sum_{n=0}^{\infty} w_n(0) P_n(z)$ converges locally uniformly on $\mathbb C$ and the respective characteristic function $\mathfrak{F}(z)$ satisfies equation (2.5). In particular, all these claims are true for $J^{[1]}$ as well.

According to (2.18) and (2.13),

$$w_n(0)Q_n(z) = -w(0)w_{n-1}^{[1]}(0)P_{n-1}^{[1]}(z)$$
 for $n \ge 1$.

Hence the series $\sum_{n=1}^{\infty} w_n(0)Q_n(z)$ converges locally uniformly on $\mathbb C$ and

$$w(0) + z \sum_{n=1}^{\infty} w_n(0) Q_n(z) = w(0) \left(1 - z \sum_{n=0}^{\infty} w_n^{[1]}(0) P_n^{[1]}(z) \right)$$

$$= w(0) \mathfrak{F}^{[1]}(z)$$

$$= w(z) \mathfrak{F}(z)$$

$$= \mathfrak{W}(z).$$

In the last two equations we have used (2.19), with k=0, and the very definition of $\mathfrak{W}(z)$.

Remark 2.7. Finally a short remark concerning Green's function. It is valid for arbitrary Jacobi matrix \mathcal{J} . Recall that Green's function of \mathcal{J} , called \mathcal{G} , is the unique semi-infinite strictly lower triangular matrix which is a right inverse of \mathcal{J} , i.e. $\mathcal{J}\mathcal{G} = \operatorname{Id}$, see [12]. Then, for all $z \in \mathbb{C}$,

$$\mathbf{P}(z) = (I - z\mathbf{G})^{-1}\mathbf{P}(0)$$

(the inverse $(\operatorname{Id} - z\mathcal{G})^{-1}$ can be defined by the geometric series) [25, Eq. (30)]. Similarly,

$$Q(z) = (Q_0(z), Q_1(z), Q_2(z), \dots)^T$$

is a formal solution of the equation $(\mathcal{J} - z \operatorname{Id}) \mathbf{Q}(z) = \mathbf{e}_0$ in \mathbb{C}^{∞} . The solution is unique with the additional assumption that the first entry is zero. It holds again true that

$$\mathbf{Q}(z) = (I - z\mathbf{G})^{-1}\mathbf{Q}(0).$$

- 3. Jacobi matrices with trace resolvent related to generators of Birth-Death processes
- 3.1. A criterion for the trace class in the case of generators of Birth-Death processes. In this section we specialize the general results derived in the preceding section to Jacobi matrices corresponding to generators of Birth-Death processes, namely, with Jacobi parameters as given in (1.4). This way we extend substantially the class of Jacobi operators studied earlier in [26] which is briefly recalled in Example 3.4.

Let us define

$$\pi_n = \begin{cases} \frac{\lambda_0 \lambda_1 \dots \lambda_{n-1}}{\mu_1 \mu_2 \dots \mu_n} & n \ge 1, \\ 1 & n = 0. \end{cases}$$
(3.1)

Occasionally, apart of orthonormal polynomials we shall also need a monic version of $(P_n : n \in \mathbb{N}_0)$. They satisfy

$$p_n(z) = a_0 a_1 \dots a_{n-1} P_n(z) \tag{3.2}$$

and

$$zp_n(z) = a_{n-1}^2 p_{n-1}(z) + b_n p_n(z) + p_{n+1}(z), \quad n \ge 1$$
(3.3)

with the initial conditions

$$p_0(z) \equiv 1, \quad p_1(z) = z - b_0.$$
 (3.4)

The following result can be derived from [14, formula (2.4)]. Our proof seems to be new.

Lemma 3.1. Consider Jacobi parameters (1.4) with $\mu_0 \ge 0$. Then for any $n \ge 0$

$$p_n(0) = (-1)^n \lambda_0 \dots \lambda_{n-1} \left(1 + \mu_0 \sum_{k=0}^{n-1} \frac{1}{\lambda_k \pi_k} \right)$$
 (3.5)

$$= (-1)^{n} \lambda_{0} \dots \lambda_{n-1} \left(1 + \sum_{k=0}^{n-1} \frac{\mu_{0} \dots \mu_{k}}{\lambda_{0} \dots \lambda_{k}} \right), \tag{3.6}$$

and consequently,

$$P_n(0) = (-1)^n \sqrt{\pi_n} \left(1 + \mu_0 \sum_{k=0}^{n-1} \frac{1}{\lambda_k \pi_k} \right).$$
 (3.7)

Proof. By (3.3) and (3.4) we have

$$zp_n(z) = p_{n+1}(z) + (\mu_n + \lambda_n)p_n(z) + \lambda_{n-1}\mu_n p_{n-1}(z), \quad n \ge 1$$
(3.8)

with the initial conditions: $p_0(z) \equiv 1$, $p_1(z) = z - \lambda_0 - \mu_0$. We shall only prove that (3.6) holds, because then (3.5) and (3.7) are simple consequences of (1.4), (3.1) and (3.2).

Notice that the equation (3.8) can be rewritten in the form

$$\begin{pmatrix} p_n(z) \\ p_{n+1}(z) \end{pmatrix} = B_n(z) \begin{pmatrix} p_{n-1}(z) \\ p_n(z) \end{pmatrix}, \quad n \ge 1, \tag{3.9}$$

where

$$B_n(z) = \begin{pmatrix} 0 & 1 \\ -\lambda_{n-1}\mu_n & z - \lambda_n - \mu_n \end{pmatrix}.$$

Notice that for any $n \ge 1$,

$$B_n(0) \begin{pmatrix} 1 \\ -\lambda_{n-1} \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ -\lambda_{n-1}\mu_n & -\lambda_n - \mu_n \end{pmatrix} \begin{pmatrix} 1 \\ -\lambda_{n-1} \end{pmatrix} = -\lambda_{n-1} \begin{pmatrix} 1 \\ -\lambda_n \end{pmatrix}, \tag{3.10}$$

$$B_n(0) \begin{pmatrix} 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 1 \\ -\lambda_n - \mu_n \end{pmatrix} = \begin{pmatrix} 1 \\ -\lambda_n \end{pmatrix} - \mu_n \begin{pmatrix} 0 \\ 1 \end{pmatrix}. \tag{3.11}$$

We claim that for any $n \ge 0$,

$$\binom{p_n(0)}{p_{n+1}(0)} = (-1)^n \lambda_0 \dots \lambda_{n-1} \left(1 + \sum_{k=0}^{n-1} \frac{\mu_0 \dots \mu_k}{\lambda_0 \dots \lambda_k} \right) \binom{1}{-\lambda_n} + (-1)^{n+1} \mu_0 \dots \mu_n \binom{0}{1}.$$
 (3.12)

This claim holds true for n = 0. Now, assuming that it holds for n we compute using (3.9) and (3.12), and then (3.10) and (3.11)

$$\begin{pmatrix} p_{n+1}(0) \\ p_{n+2}(0) \end{pmatrix} = B_{n+1}(0) \begin{pmatrix} p_n(0) \\ p_{n+1}(0) \end{pmatrix}
= (-1)^n \lambda_0 \dots \lambda_{n-1} \left(1 + \sum_{k=0}^{n-1} \frac{\mu_0 \dots \mu_k}{\lambda_0 \dots \lambda_k} \right) B_{n+1}(0) \begin{pmatrix} 1 \\ -\lambda_n \end{pmatrix} + (-1)^{n+1} \mu_0 \dots \mu_n B_{n+1}(0) \begin{pmatrix} 0 \\ 1 \end{pmatrix}
= (-1)^{n+1} \lambda_0 \dots \lambda_{n-1} \lambda_n \left(1 + \sum_{k=0}^n \frac{\mu_0 \dots \mu_k}{\lambda_0 \dots \lambda_k} \right) \begin{pmatrix} 1 \\ -\lambda_{n+1} \end{pmatrix} + (-1)^{n+2} \mu_0 \dots \mu_{n+1} \begin{pmatrix} 0 \\ 1 \end{pmatrix},$$

which shows that then (3.12) holds true also for n+1. Thus, by induction the proof of (3.12) is complete. \Box

Lemma 3.2. Assume, in addition to the assumptions of Lemma 3.1, that

$$\sum_{n=0}^{\infty} \pi_n (1 + \mathfrak{s}_n)^2 = \infty, \quad \text{where} \quad \mathfrak{s}_n := \mu_0 \sum_{k=0}^{n-1} \frac{1}{\lambda_k \pi_k}.$$
 (3.13)

Then J is self-adjoint and positive-semidefinite. Set

$$\mathfrak{s}_{\infty} := \lim_{n \to \infty} \mathfrak{s}_n$$

Then, for $n \in \mathbb{N}_0$, we have

$$-\left(\sum_{j=n}^{\infty} \frac{1}{a_j P_j(0) P_{j+1}(0)}\right) P_n(0) = (-1)^n \frac{\sqrt{\pi_n}}{1 + \mathfrak{s}_{\infty}} \sum_{j=n}^{\infty} \frac{1}{\lambda_j \pi_j}$$
(3.14)

if $\mathfrak{s}_{\infty} < \infty$, and

$$-\left(\sum_{j=n}^{\infty} \frac{1}{a_j P_j(0) P_{j+1}(0)}\right) P_n(0) = (-1)^n \frac{\sqrt{\pi_n}}{\mu_0}$$
(3.15)

if $\mathfrak{s}_{\infty} = \infty$.

Proof. One can employ formula (3.7). Likewise in the proof of Theorem 2.1, (3.13) implies that J is self-adjoint. Moreover, a simple consequence of [14, Thm. 1] is that J is positive-semidefinite.

If $\mu_0 = 0$ then $P_n(0) = (-1)^n \sqrt{\pi_n}$ and $\mathfrak{s}_n = 0$ for any $n \ge 0$. Thus, $\mathfrak{s}_{\infty} = 0$ and (3.14) is immediately seen to hold. Note that

$$a_j = \lambda_j \sqrt{\frac{\pi_j}{\pi_{j+1}}}\,,$$

For $\mu_0 > 0$ we have

$$-\frac{1}{a_j P_j(0) P_{j+1}(0)} = \frac{1}{\lambda_j \pi_j (1 + \mathfrak{s}_j) (1 + \mathfrak{s}_{j+1})} = \frac{1}{\mu_0} \left(\frac{1}{1 + \mathfrak{s}_j} - \frac{1}{1 + \mathfrak{s}_{j+1}} \right).$$

Whence

$$-\left(\sum_{j=n}^{\infty} \frac{1}{a_j P_j(0) P_{j+1}(0)}\right) P_n(0) = (-1)^n \frac{\sqrt{\pi_n} (1 + \mathfrak{s}_n)}{\mu_0} \lim_{N \to \infty} \left(\frac{1}{1 + \mathfrak{s}_n} - \frac{1}{1 + \mathfrak{s}_N}\right)$$
$$= (-1)^n \frac{\sqrt{\pi_n}}{\mu_0} \left(1 - \frac{1 + \mathfrak{s}_n}{1 + \mathfrak{s}_\infty}\right).$$

Depending on the value of \mathfrak{s}_{∞} we get either (3.14) or (3.15).

Remark. If $\mu_0 > 0$, then (3.13) is in fact *necessary* for self-adjointness of J, see [10, formula (3.83)].

A straightforward combination of Theorem 2.1, Lemma 3.1 and Lemma 3.2 yields the following theorem.

Theorem 3.3. Let J be the Jacobi operator corresponding to (1.4). If

$$\sum_{n=0}^{\infty} \pi_n (1 + \mathfrak{s}_n)^2 = \infty,$$

then J is a positive-semidefinite self-adjoint operator and $0 \notin \sigma_p(J)$. Moreover, if $\mathfrak{s}_{\infty} < \infty$ then

$$\operatorname{Tr}(J^{-1}) = \frac{1}{1 + \mathfrak{s}_{\infty}} \sum_{n=0}^{\infty} \pi_n (1 + \mathfrak{s}_n) \sum_{j=n}^{\infty} \frac{1}{\lambda_j \pi_j}$$
$$= \frac{1}{1 + \mathfrak{s}_{\infty}} \sum_{j=0}^{\infty} \frac{1}{\lambda_j \pi_j} \sum_{n=0}^{j} \pi_n (1 + \mathfrak{s}_n), \tag{3.16}$$

and if $\mathfrak{s}_{\infty} = \infty$ then

$$Tr(J^{-1}) = \frac{1}{\mu_0} \sum_{n=0}^{\infty} \pi_n (1 + \mathfrak{s}_n). \tag{3.17}$$

Example 3.4. In [26] the following Jacobi parameters

$$a_n = \kappa \tilde{a}_n, \quad b_n = \tilde{a}_n + \kappa^2 \tilde{a}_{n-1}, \quad n \in \mathbb{N}_0,$$

where $\tilde{a}_{-1} := 0, \kappa \in (0, 1)$ and $(\tilde{a}_n : n \in \mathbb{N}_0)$ is a positive sequence such that

$$\sum_{n=0}^{\infty} \frac{1}{\tilde{a}_n} < \infty. \tag{3.18}$$

Notice that by setting

$$\lambda_n = \tilde{a}_n, \quad \mu_n = \kappa^2 \tilde{a}_{n-1}$$

leads to the setup (1.4) with $\mu_0 = 0$. Thus, by (3.1) we have for $n \ge 1$,

$$\pi_n = \frac{\lambda_0 \lambda_1 \dots \lambda_{n-1}}{\mu_1 \mu_2 \dots \mu_n} = \frac{1}{\kappa^{2n}}.$$

Therefore, if $0 \notin \sigma_{ess}(J)$, then the identity (3.16) reads

$$\mathrm{Tr}(J^{-1}) = \sum_{j=0}^{\infty} \frac{1}{\tilde{a}_j} \sum_{n=0}^{j} \kappa^{2(j-n)} = \sum_{j=0}^{\infty} \frac{1}{\tilde{a}_j} \frac{1-\kappa^{2j+2}}{1-\kappa^2}.$$

which, thanks to (3.18), is finite.

3.2. **The characteristic function.** In this subsection we are still focusing on Jacobi operators with the parameters as given in (1.4). To simplify the discussion we shall assume that

$$\sum_{j=0}^{\infty} \frac{1}{\lambda_j \pi_j} < \infty.$$

In addition to formula (3.7), we also have

$$w_n(0) = (-1)^n \sqrt{\pi_n} \left(1 + \mu_0 \sum_{j=0}^{\infty} \frac{1}{\lambda_j \pi_j} \right)^{-1} \sum_{j=n}^{\infty} \frac{1}{\lambda_j \pi_j}.$$
 (3.19)

This equation is obtained as a combination of (2.12) with (3.14).

With regard to applications of Lemma 2.2 to follow it is convenient to set

$$S_n = \sum_{j=n}^{\infty} \frac{1}{\lambda_j \pi_j}, \quad n \ge 0.$$

Hence $\mathfrak{s}_n = \mu_0 S_n$, see (3.13). Then we can write

$$P_n(0) = (-1)^n \sqrt{\pi_n} \left(1 + \mu_0 (S_0 - S_n) \right), \quad w_n(0) = (-1)^n \frac{\sqrt{\pi_n} S_n}{1 + \mu_0 S_0},$$

and

$$\kappa_n = w_n(0)P_n(0) = \frac{1 + \mu_0(S_0 - S_n)}{1 + \mu_0S_0} \pi_n S_n.$$

Further we put, by definition,

$$S_{-1} = \frac{1}{\mu_0} + S_0$$

if $\mu_0 > 0$, and $S_{-1} = \infty$ if $\mu_0 = 0$. Then

$$\frac{1 + \mu_0(S_0 - S_n)}{1 + \mu_0 S_0} = 1 - \frac{\mu_0 S_n}{1 + \mu_0 S_0} = 1 - \frac{S_n}{S_{-1}}.$$

Hence

$$\kappa_n = \left(1 - \frac{S_n}{S_{-1}}\right) \pi_n S_n.$$

Using this notation we have

$$K_{m,n} = -\sqrt{\frac{1 - S_n/S_{-1}}{1 - S_m/S_{-1}}} \sqrt{\pi_m S_m \pi_n S_n} \left(1 - \frac{S_m}{S_n}\right) \quad \text{for } m > n,$$

see (2.8). Thus we can rewrite (2.9) as follows,

$$\mathfrak{F}(z) = 1 + \sum_{m=1}^{\infty} (-1)^m z^m \sum_{0 \le j_1 < j_2 < \dots < j_m} \pi_{j_1} S_{j_1} \pi_{j_2} S_{j_2} \dots \pi_{j_m} S_{j_m} \left(1 - \frac{S_{j_1}}{S_{-1}} \right) \left(1 - \frac{S_{j_2}}{S_{j_1}} \right) \dots \left(1 - \frac{S_{j_m}}{S_{j_{m-1}}} \right). \tag{3.20}$$

Now consider the case when $\mu_0 = 0$. Then (see (3.7), (3.19) with $\mu_0 = 0$)

$$P_n(0) = (-1)^n \sqrt{\pi_n}, \quad w_n(0) = (-1)^n \sqrt{\pi_n} S_n, \text{ and } \kappa_n = \pi_n S_n.$$

Using Lemma 2.2 we can express the characteristic function as

$$\mathfrak{F}(z) = 1 + \sum_{m=1}^{\infty} (-1)^m z^m \sum_{0 \le j_1 < j_2 < \dots < j_m} \pi_{j_1} \pi_{j_2} \dots \pi_{j_m} (S_{j_1} - S_{j_2}) \dots (S_{j_{m-1}} - S_{j_m}) S_{j_m}$$

This can be rewritten as follows,

$$\mathfrak{F}(z) = 1 + \sum_{m=1}^{\infty} (-1)^m z^m \sum_{\substack{0 \le j_1 \le k_2 \le \dots \le k_{m-1} \le j_m \le k_m}} \frac{\pi_{j_1} \pi_{j_2} \dots \pi_{j_m}}{\lambda_{k_1} \pi_{k_1} \dots \lambda_{k_{m-1}} \pi_{k_m} \lambda_{k_m} \pi_{k_m}}.$$
 (3.21)

Denote

$$T_k = \sum_{j=0}^k \pi_j, \quad k \ge 0.$$
 (3.22)

Changing the order of summation we get

$$\mathfrak{F}(z) = 1 + \sum_{m=1}^{\infty} (-1)^m z^m \sum_{0 \le k_1 < k_2 < \dots < k_m} \frac{T_{k_1} (T_{k_2} - T_{k_1}) \dots (T_{k_m} - T_{k_{m-1}})}{\lambda_{k_1} \pi_{k_1} \dots \lambda_{k_{m-1}} \pi_{k_{m-1}} \lambda_{k_m} \pi_{k_m}}.$$
(3.23)

3.3. **The function** $\mathfrak{B}(z)$. Let us make a few comments on computing the function $\mathfrak{B}(z)$, as given in (2.6) and in (2.22), but still assuming that $\mu_0 = 0$. $J^{[1]}$ again stands for the first associated matrix to J obtained by removing the first row and first column, and $\lambda_n^{[1]}$, $\mu_n^{[1]}$ for the corresponding Birth-Death parameters. In a similar fashion we introduce the symbol $\pi_n^{[1]}$ etc. Then

$$\lambda_n^{[1]} = \lambda_{n+1}, \quad \mu_n^{[1]} = \mu_{n+1}, \quad \pi_n^{[1]} = \pi_{n+1} \frac{\mu_1}{\lambda_0}.$$

Furthermore,

$$S_n^{[1]} = \sum_{j=n}^{\infty} \frac{1}{\lambda_j^{[1]} \pi_j^{[1]}} = \frac{\lambda_0}{\mu_1} S_{n+1},$$

and also $(\pi_0 = 1)$

$$S_{-1}^{[1]} = \frac{1}{\mu_0^{[1]}} + S_0^{[1]} = \frac{1}{\mu_1} + \frac{\lambda_0}{\mu_1} S_1 = \frac{\lambda_0}{\mu_1} \left(\frac{1}{\lambda_0 \pi_0} + S_1 \right) = \frac{\lambda_0}{\mu_1} S_0.$$

Regarding the characteristic function of $J^{[1]}$, since $\mu_0^{[1]} \neq 0$, we can use formula (3.20), namely

$$\mathfrak{F}^{[1]}(z) = 1 + \sum_{m=1}^{\infty} (-1)^m z^m$$

$$\times \sum_{\substack{0 \le j_1 < j_2 < \dots < j_m \\ 0 \le j_1 \le j_2 < \dots \le j_m}} \pi_{j_1}^{[1]} S_{j_1}^{[1]} \dots \pi_{j_m}^{[1]} S_{j_m}^{[1]} \left(1 - \frac{S_{j_1}^{[1]}}{S_{j_1}^{[1]}}\right) \left(1 - \frac{S_{j_2}^{[1]}}{S_{j_1}^{[1]}}\right) \dots \left(1 - \frac{S_{j_m}^{[1]}}{S_{j_m}^{[1]}}\right).$$

Hence

$$\mathfrak{F}^{[1]}(z) = 1 + \frac{1}{S_0} \sum_{m=1}^{\infty} (-1)^m z^m \sum_{1 \le j_1 < j_2 < \dots < j_m} \pi_{j_1} \dots \pi_{j_m} (S_0 - S_{j_1}) (S_{j_1} - S_{j_2}) \dots (S_{j_{m-1}} - S_{j_m}) S_{j_m}.$$

Substituting for S_i we get (note that $S_0 = w(0)$)

$$\mathfrak{W}(z) = S_0 \mathfrak{F}^{[1]}(z) = \sum_{m=0}^{\infty} (-1)^m z^m \sum_{0 \le k_1 < j_1 \le k_2 < j_2 \dots \le k_m < j_m \le k_{m+1}} \frac{\pi_{j_1} \pi_{j_2} \dots \pi_{j_m}}{\lambda_{k_1} \pi_{k_1} \dots \lambda_{k_m} \pi_{k_m} \lambda_{k_{m+1}} \pi_{k_{m+1}}}$$

(one may wish to compare this equation to (3.21)). This equation can be also rewritten with the aid of the symbol T_k , as defined in (3.22),

$$\mathfrak{W}(z) = \sum_{m=0}^{\infty} (-1)^m z^m \sum_{0 \le k_1 < \dots < k_m < k_{m+1}} \frac{(T_{k_2} - T_{k_1}) \cdots (T_{k_{m+1}} - T_{k_m})}{\lambda_{k_1} \pi_{k_1} \dots \lambda_{k_m} \pi_{k_m} \lambda_{k_{m+1}} \pi_{k_{m+1}}}$$

(again, this may be compared to (3.23)).

4. Some results for positive Jacobi matrices with compact inverses

4.1. On the convergence of the zeros of orthogonal polynomials. Suppose that Jacobi operator J is positive. Let J_F be its $Friedrichs\ extension(^1)$ (see, e.g. [21, Chapter 8.1]). Suppose that J_F^{-1} is compact and let $\sigma(J_F) = \{\zeta_j : j \ge 1\}$ be ordered by

$$0 < \zeta_1 < \zeta_2 < \dots \tag{4.1}$$

Let $(p_n : n \ge 0)$ be the sequence of monic orthogonal polynomials associated with \mathcal{J} . Then by [7, Theorem I.5.2] we have that all the zeros of $(p_n : n \ge 1)$ are real, simple and located in (ζ_1, ∞) . In particular,

$$\forall n \ge 1 \ \forall z \in \mathbb{C} \setminus (\zeta_1, \infty), \quad p_n(z) \ne 0. \tag{4.2}$$

Thus we have

$$p_n(z) = (z - x_{1:n}) \dots (z - x_{n:n}), \tag{4.3}$$

where

$$0 < \zeta_1 < x_{1:n} < x_{2:n} < \dots < x_{n:n}. \tag{4.4}$$

The following result will be instrumental in what follows.

Theorem 4.1. Suppose that Jacobi operator J is positive and let J_F be its Friedrichs extension. Let J_F^{-1} be compact. Let $\sigma(J_F) = \{\zeta_j : j \geq 1\}$ be ordered as in (4.1) and let the monic orthogonal polynomials be written as in (4.3) and (4.4). Then

$$\zeta_j < x_{j;n}, \quad j = 1, 2, \dots, n; \ n \ge 1,$$
 (4.5)

$$\zeta_j = \lim_{n \to \infty} x_{j;n}, \quad j = 1, 2, \dots$$
 (4.6)

Proof. First of all, by [7, Theorem II.4.1] we get that(2)

$$\sharp ((x_{i:n}, x_{i+1:n}) \cap \sigma(J_F)) \ge 1, \quad i = 1, 2, \dots, n-1.$$

$$(4.7)$$

By (4.4) and (4.1) it easily implies (4.5). Now if J is not-self-adjoint, i.e. $J \neq J_F$, the formula (4.6) has been proven in [5, p. 479]. On the other hand, if J is self-adjoint, then formula (4.7) together with Theorems I.5.3 and II.4.3 from [7] imply (4.6).

¹If J is self-adjoint, then its every self-adjoint extension is equal to J. In particular, then $J = J_F$.

 $^{{}^{2}}$ By $\sharp X$ we denote the number of elements in X.

4.2. The zero counting measures. Suppose that we are working in the setup of Section 4.1. In this section we shall analyse weak convergence of the sequence of zero counting measures of P_n defined by

$$\nu_n = \sum_{j=1}^n \delta_{x_{j;n}},\tag{4.8}$$

see formulas (4.3) and (4.4).

Theorem 4.2. Suppose that a Jacobi matrix J is positive and J_F^{-1} is compact. Let $\sigma(J_F) = \{\zeta_j : j \geq 1\}$ be ordered as in (4.1). Suppose that for some continuous non-decreasing function $w : \mathbb{R} \to (0, \infty)$ we have

$$\sum_{i=1}^{\infty} \frac{1}{w(\zeta_j)} < \infty. \tag{4.9}$$

Then for any(3) $f \in C_b(\mathbb{R})$ such that $||wf||_{\infty} < \infty$ we have

$$\lim_{n \to \infty} \int_{\mathbb{R}} f \, \mathrm{d}\nu_n = \int_{\mathbb{R}} f \, \mathrm{d}\nu_\infty, \tag{4.10}$$

where v_n is defined in (4.8) and

$$\nu_{\infty} = \sum_{i=1}^{\infty} \delta_{\zeta_j}.$$
 (4.11)

Proof. In order to prove (4.10) it is enough to show that

$$\lim_{n \to \infty} \sum_{j=1}^{n} \frac{\tilde{f}(x_{j;n})}{w(x_{j;n})} = \sum_{j=1}^{\infty} \frac{\tilde{f}(\zeta_j)}{w(\zeta_j)}, \quad \tilde{f} \in C_b(\mathbb{R}).$$
(4.12)

Indeed, to obtain (4.10) it is enough to apply (4.12) for $\tilde{f}(x) = w(x)f(x)$. Notice that by (4.5) and the fact that the function w is non-decreasing we get

$$\frac{\tilde{f}(x_{j;n})}{w(x_{j;n})} \le \frac{\|\tilde{f}\|_{\infty}}{w(x_{j;n})} \le \frac{\|\tilde{f}\|_{\infty}}{w(\zeta_{j})}.$$
(4.13)

Moreover, by (4.6) and the continuity of \tilde{f} and w we get that

$$\lim_{n \to \infty} \frac{\tilde{f}(x_{j;n})}{w(x_{j:n})} = \frac{\tilde{f}(\zeta_j)}{w(\zeta_j)}.$$
(4.14)

By combining (4.14), (4.13) and (4.9), by Dominated Convergence Theorem, we get (4.13) what concludes the proof.

Remark 4.3. The hypothesis (4.9) is only important for continuous functions f with non-compact support. Indeed, if (4) $f \in C_c(\mathbb{R})$, then the convergence (4.10) holds by (4.6) as the both sides of (4.10) involves only sums with uniformly bounded number of terms.

Example 4.4. Suppose that J is positive and J_F^{-1} belongs to the Schatten class S_k for some $k \ge 1$. Let $\sigma(J_F) = \{\zeta_i : j \ge 1\}$ be ordered as in (4.1). Then since

$$\sum_{j=1}^{\infty} \frac{1}{\zeta_j^k} < \infty$$

we can take in Theorem 4.2

$$w(x) = \begin{cases} (1+x^2)^{k/2} & x > 0, \\ 1 & x \le 0. \end{cases}$$

 $^{^3}$ By $C_b(\mathbb{R})$ we denote the space of continuous bounded functions on the real line equipped with the supremum norm $\|\cdot\|_{\infty}$.

⁴The set $C_c(\mathbb{R}) \subset C_b(\mathbb{R})$ consists of compactly supported continuous functions on \mathbb{R} .

4.3. **The Christoffel–Darboux kernels.** Let us recall that given J, let \tilde{J} be a self-adjoint extension on $\ell^2(\mathbb{N}_0)$ of J. We the can define a Borel probability measure $\tilde{\mu}$ by the formula

$$\tilde{\mu}(\cdot) = \langle \boldsymbol{e}_0, E_{\tilde{\boldsymbol{J}}}(\cdot)\boldsymbol{e}_0 \rangle,$$

where $E_{\tilde{J}}$ is the projection-valued spectral measure of \tilde{J} . By Riesz's theorem (see e.g. [22, Theorem 2.14, p. 62]) the sequence $(P_n : n \ge 0)$ forms an orthonormal basis in $L^2(\tilde{\mu})$. Now, for any $n \ge 1$ let

$$K_n(x, y) = \sum_{j=0}^{n-1} p_j(x) p_j(y), \quad x, y \in \mathbb{R}$$

be the nth Christoffel-Darboux kernel. In this section we shall study the weak convergence of the sequence

$$d\eta_n = K_n(x, x) d\tilde{\mu}(x). \tag{4.15}$$

Let us start with a general result.

Theorem 4.5. Let \tilde{J} be a self-adjoint extension on $\ell^2(\mathbb{N}_0)$ of J. If $\sigma_{\text{ess}}(\tilde{J}) = \emptyset$, then

$$\lim_{n \to \infty} \int_{\mathbb{R}} f \, \mathrm{d}\eta_n = \int_{\mathbb{R}} f \, \mathrm{d}\eta_\infty \tag{4.16}$$

for any function $f \in L^1(\eta_\infty)$, where

$$\eta_{\infty} = \sum_{\zeta \in \sigma(\tilde{J})} \delta_{\zeta}. \tag{4.17}$$

Proof. Recall that $\sigma(\tilde{J}) = \operatorname{supp}(\tilde{\mu})$. Since $\sigma_{\operatorname{ess}}(\tilde{J}) = \emptyset$, it follows that $\tilde{\mu}$ is purely discrete. Let us recall that

$$K_{\infty}(\zeta,\zeta) := \lim_{n \to \infty} K_n(\zeta,\zeta) = \frac{1}{\tilde{\mu}(\{\zeta\})}, \quad \zeta \in \operatorname{supp}(\tilde{\mu}),$$

see e.g. [22, Corollary 2.6, p. 45] in case $J = \tilde{J}$ and [22, Theorem 2.13, p. 60] otherwise. Thus,

$$\int_{\mathbb{R}} f \, \mathrm{d}\eta_n = \sum_{\zeta \in \sigma(\tilde{J})} f(\zeta) \frac{K_n(\zeta, \zeta)}{K_\infty(\zeta, \zeta)}.$$

Since

$$|f(\zeta)| \frac{K_n(\zeta,\zeta)}{K_m(\zeta,\zeta)} \le |f(\zeta)|$$

and

$$\lim_{n\to\infty}\frac{K_n(\zeta,\zeta)}{K_\infty(\zeta,\zeta)}=1$$

we get (4.16) by Dominated Convergence Theorem.

Let v_n be defined in (4.8). It is well-known that the sequences of probability measures $(\frac{1}{n}v_n : n \ge 1)$ and $(\frac{1}{n}\eta_n : n \ge 1)$ are equiconvergent in the sense

$$\lim_{n\to\infty} \left| \frac{1}{n} \int_{\mathbb{R}} f \, d\nu_n - \frac{1}{n} \int_{\mathbb{R}} f \, d\eta_n \right| = 0, \quad f \in C_c(\mathbb{R}),$$

see [13, Theorem 2.1]. Notice that since (ν_n) does not depend on the self-adjoint extension of J, then the limit (if exists) of $(\frac{1}{n}\eta_n)$ also does not depend on the self-adjoint extension of J. The following remark states that in general a similar result *does not hold* for the sequences $(\nu_n : n \ge 1)$ and $(\eta_n : n \ge 1)$ themselves.

Remark 4.6. Suppose that J is positive, let J_F be its Friedrichs extension and \tilde{J} be an arbitrary self-adjoint extension of J on $\ell^2(\mathbb{N}_0)$. Suppose that J_F^{-1} is compact. Then by Remark 4.3

$$\lim_{n\to\infty}\int_{\mathbb{R}}f\,\mathrm{d}\nu_n=\int_{\mathbb{R}}f\,\mathrm{d}\nu_\infty,\quad f\in C_c(\mathbb{R}),$$

where v_{∞} is defined in (4.11). If J is self-adjoint, then $J = J_F = \tilde{J}$. So by Theorem 4.5

$$\lim_{n \to \infty} \int_{\mathbb{R}} f \, d\eta_n = \int_{\mathbb{R}} f \, d\eta_{\infty}, \quad f \in C_c(\mathbb{R}), \tag{4.18}$$

where η_{∞} is defined in (4.17), and since $\tilde{J} = J_F$ we get $\nu_{\infty} = \eta_{\infty}$. On the other hand, if J is not self-adjoint, then $\sigma_{\rm ess}(\tilde{J}) = \emptyset$, see e.g. [21, Theorem 7.7(i)]. So we still have the convergence (4.18) for η_n obtained from $\tilde{\mu}$ via the formula (4.15). Since all self-adjoint extensions on $\ell^2(\mathbb{N}_0)$ of J have disjoint spectra, see e.g. [21, Theorem 7.7(ii)], we get $\nu_{\infty} = \eta_{\infty}$ iff $\tilde{J} = J_F$.

4.4. The regularized characteristic function for Schatten classes. Let a sequence of distinct real numbers $(\zeta_j : j \ge 1)$ ordered by $0 < |\zeta_1| \le |\zeta_2| < \dots$ satisfies for some $k \in \{1, 2, \dots\}$

$$\sum_{j=1}^{\infty} \frac{1}{|\zeta_j|^k} < \infty. \tag{4.19}$$

Then one can define an entire function $\mathfrak{F}_k : \mathbb{C} \to \mathbb{C}$ by the formula

$$\mathfrak{F}_k(z) = \prod_{i=1}^{\infty} E_{k-1} \left(\frac{z}{\zeta_i} \right), \tag{4.20}$$

where *nth elementary factor* is defined by by

$$E_n(w) = (1 - w) \exp\left(\sum_{i=1}^n \frac{1}{j} w^j\right), \quad w \in \mathbb{C},$$
(4.21)

(see e.g. [9, Definition VII.5.10]). We have $\mathfrak{F}_k(z) = 0$ iff $z = \zeta_j$ for some j (see e.g. [9, Theorem VII.5.12]). Suppose now that we working in the setup of Section 4.1. Suppose additionally, that J_F^{-1} exists and belongs to some Schatten class S_k for some $k \in \{1, 2, \ldots\}$. It means that for $\sigma(J_F) = \{\zeta_j : j \ge 1\}$ ordered as in (4.1) we have (4.19). In particular, we can define the *regularized characteristic function* of J_F by the formula (4.20). Notice that we also have

$$\mathfrak{F}_k(z) = \det \left[E_{k-1} \left(z J_F^{-1} \right) \right],$$

where det is the Fredholm determinant.

Let us recall that monic orthogonal polynomials (see (3.2)) satisfy

$$p_n(z) = \det(z \operatorname{Id}_n - \mathcal{J}_n), \tag{4.22}$$

where Id_n is $n \times n$ identity matrix and for any $n \ge 1$ the truncation of \mathcal{J} , \mathcal{J}_n , is defined by

$$\mathcal{J}_n = \begin{pmatrix} b_0 & a_0 & & & & \\ a_0 & b_1 & a_1 & & & & \\ & a_1 & \ddots & \ddots & & \\ & & \ddots & \ddots & a_{n-2} \\ & & & a_{n-2} & b_{n-1} \end{pmatrix}.$$

Having fixed the notation we can formulate our main result of this section.

Theorem 4.7. Suppose that J is positive and J_F^{-1} belongs to the Schatten class S_k for some $k \in \{1, 2, ...\}$. Then

$$\lim_{n \to \infty} \det \left[E_{k-1}(z\mathcal{J}_n^{-1}) \right] = \mathfrak{F}_k(z) \tag{4.23}$$

locally uniformly with respect to $z \in \mathbb{C}$.

Proof. By (4.2) for any $n \ge 1$ the matrix \mathcal{J}_n is invertible (cf. (4.22)) and by (4.3)

$$\det \left[E_{k-1}(z\mathcal{J}_n^{-1}) \right] = \prod_{i=1}^n E_{k-1} \left(\frac{z}{x_{j;n}} \right). \tag{4.24}$$

In view of (4.20), in order to prove (4.23) we need to prove that

$$\lim_{n\to\infty} \prod_{j=1}^n E_{k-1}\left(\frac{z}{x_{j;n}}\right) = \prod_{j=1}^\infty E_{k-1}\left(\frac{z}{\zeta_j}\right)$$

locally uniformly with respect to $z \in \mathbb{C}$.

Let us take R > 0. Notice that by (4.6) for any $j \ge 1$,

$$\lim_{n \to \infty} \frac{z}{x_{j;n}} = \frac{z}{\zeta_j} \tag{4.25}$$

uniformly for $|z| \le R$. Notice that for any fixed $1 \le j_0 \le n$

$$\prod_{j=1}^{n} E_{k-1} \left(\frac{z}{x_{j;n}} \right) = \prod_{j=1}^{j_0-1} E_{k-1} \left(\frac{z}{x_{j;n}} \right) \cdot \prod_{j=j_0}^{n} E_{k-1} \left(\frac{z}{x_{j;n}} \right).$$

Thus, by continuity of E_{k-1} and (4.25) we need to only prove that for some $j_0 \ge 1$

$$\lim_{n \to \infty} \prod_{j=j_0}^{n} E_{k-1} \left(\frac{z}{x_{j;n}} \right) = \prod_{j=j_0}^{\infty} E_{k-1} \left(\frac{z}{\zeta_j} \right)$$
 (4.26)

uniformly for $|z| \le R$. Now, let $j_0 \ge 1$ be such that $\frac{R}{\lambda_j} \le \frac{1}{2}$ for any $j \ge j_0$. By (4.5) we also have $\frac{R}{x_{j;n}} \le \frac{1}{2}$ for any $j \ge j_0$ and $n \ge 1$. Since

$$|E_{k-1}(w) - 1| \le |w|^k$$
, $|w| \le \frac{1}{2}$

see e.g. [9, Lemma VII.5.11], we get by (4.5)

$$\left| E_{k-1} \left(\frac{z}{x_{j;n}} \right) - 1 \right| \le \left(\frac{R}{x_{j;n}} \right)^k \le \left(\frac{R}{\zeta_j} \right)^k.$$

Since $J^{-1} \in \mathcal{S}_k$ the last sequence is summable. Thus, in view of uniform Tannery's theorem for infinite products (for a pointwise version see e.g. [16, Theorem 7.7]) together with (4.25) we get (4.26), what concludes the proof.

Remark 4.8. It is known (see e.g. [2, Theorem 2.11]) that if J is not self-adjoint, then J_F^{-1} is a Hilbert–Schmidt operator, so in such a case in Theorem 4.7 one can take any $k \ge 2$.

Using (4.3) and (4.24) one can verify that for any $k \ge 1$,

$$\det\left[E_{k-1}(z\mathcal{J}_n^{-1})\right] = \frac{p_n(z)}{p_n(0)} \exp\left(\sum_{j=1}^{k-1} \frac{z^j}{j} \operatorname{tr}\left(\mathcal{J}_n^{-j}\right)\right). \tag{4.27}$$

In order to compute the value of (4.27) in terms of p_n we need the following proposition.

Proposition 4.9. Let $n \ge 1$. Define for $1 \le k \le n$

$$\mathfrak{p}_k = \operatorname{tr} \left(\mathcal{J}_n^{-k} \right), \quad \mathfrak{e}_k = \frac{(-1)^k}{k!} \frac{p_n^{(k)}(0)}{p_n(0)}.$$

Then we have

$$\mathfrak{p}_k = \sum_{i=1}^{k-1} (-1)^{i-1} \mathfrak{p}_{k-i} \mathfrak{e}_i + (-1)^{k-1} k \mathfrak{e}_k, \quad 1 \le k \le n.$$
 (4.28)

In particular,

$$\mathfrak{p}_1 = \mathfrak{e}_1, \quad \mathfrak{p}_2 = \mathfrak{e}_1^2 - 2\mathfrak{e}_2, \quad \mathfrak{p}_3 = \mathfrak{e}_1^3 - 3\mathfrak{e}_1\mathfrak{e}_2 + 3\mathfrak{e}_3$$
 (4.29)

Proof. Notice that by (4.22) and (4.3) we have

$$\mathfrak{p}_k = \sum_{i=1}^n \frac{1}{x_{i:n}^k}. (4.30)$$

Next, by (4.3)

$$\frac{p_n(z)}{p_n(0)} = \prod_{i=1}^n \left(1 - \frac{z}{x_{i;n}}\right).$$

By induction on $1 \le k \le n$ one can prove that

$$\left[\prod_{i=1}^{n} \left(1 - \frac{z}{x_{i;n}}\right)\right]^{(k)}(z) = \sum_{\substack{I \subset \{1, \dots, n\} \\ \sharp I = k}} \prod_{i \in I} \frac{-1}{x_{i;n}} \cdot \prod_{i \in \{1, \dots, n\} \setminus I} \left(1 - \frac{z}{x_{i;n}}\right).$$

Thus,

$$\mathbf{e}_{k} = \frac{(-1)^{k}}{k!} \frac{p_{N}^{(k)}(0)}{p_{n}(0)} = \sum_{1 \le i_{1} \le \dots \le i_{k} \le n} \prod_{j=1}^{k} \frac{1}{x_{i_{j};n}}.$$
(4.31)

Now, in view of (4.30) and (4.31), the formula (4.28) is just Newton's identity, see e.g. [19]. Then the formula (4.29) is an application of (4.29).

In view of (4.27) and Proposition 4.9 we can rewrite the argument of the exponential on left-hand side of (4.23) in terms of $p_n^{(j)}(0)/p_n(0)$. In particular, it leads to

$$\det\left[E_0(z\mathcal{J}_n^{-1})\right] = \frac{p_n(z)}{p_n(0)} \tag{4.32}$$

$$\det\left[E_1(z\mathcal{J}_n^{-1})\right] = \frac{p_n(z)}{p_n(0)} \exp\left(-z\frac{p'_n(0)}{p_n(0)}\right) \tag{4.33}$$

$$\det\left[E_2(z\mathcal{J}_n^{-1})\right] = \frac{p_n(z)}{p_n(0)} \exp\left(-z\frac{p_n'(0)}{p_n(0)} + \frac{z^2}{2}\left\{\left(\frac{p_n'(0)}{p_n(0)}\right)^2 - \frac{p_n''(0)}{p_n(0)}\right\}\right). \tag{4.34}$$

Let us recall that in [25, Section 3] the following particular case was proven.

Corollary 4.10. Suppose that J is positive and J^{-1} is a trace class operator. Then

$$\lim_{n \to \infty} \frac{p_n(z)}{p_n(0)} = \mathfrak{F}_1(z) \tag{4.35}$$

locally uniformly with respect to $z \in \mathbb{C}$.

Proof. Apply Theorem 4.7 with k = 1 and use (4.32).

Remark 4.11. Theorem 4.1 together with [6, Theorem 2] imply that if the operator J is positive with J_F^{-1} compact, then the left-hand side of (4.35) is convergent locally uniformly on $\mathbb C$ to an entire function with the zero set $\sigma(J_F)$ iff J_F^{-1} is trace class. In [6] the right-hand side of (4.35) was not identified.

4.5. **Importance of the positivity.** Suppose that J is a self-adjoint Jacobi matrix with $b_n \equiv 0$. Let $(P_n : n \geq 0)$ be the corresponding orthonormal polynomials and let μ be their unique orthonormality measure, i.e. defined in (1.1). Then it is well-known that the measure μ is symmetric around 0 and (see e.g. formulas (1.6)–(1.9) in [8]) that

$$P_{2n}(z) = \tilde{P}_n(z^2), \quad P_{2n+1}(z) = z \hat{P}_n(z^2), \quad n \ge 0,$$

where the sequences $(\tilde{P}_n : n \ge 0)$, $(\hat{P}_n : n \ge 0)$ are orthonormal with respect to the measures

$$\tilde{\mu}(X) = \mu(\{\pm \sqrt{x} : x \in X \cap [0, \infty)\}) \quad \text{and} \quad \hat{\mu}(X) = \int_X x \, \mathrm{d}\tilde{\mu}(x), \tag{4.36}$$

respectively.

Suppose that J^{-1} is a Hilbert–Schmidt operator, then

$$\sum_{j=1}^{\infty} \frac{1}{\zeta_j^2} < \infty$$

and its 2-regularized characteristic function is equal to

$$\mathfrak{F}_2(z) = \prod_{j=1}^{\infty} \left[\left(1 - \frac{z}{\zeta_j} \right) \exp\left(\frac{z}{\zeta_j} \right) \right] \left[\left(1 - \frac{z}{-\zeta_j} \right) \exp\left(\frac{z}{-\zeta_j} \right) \right] = \prod_{j=1}^{\infty} \left(1 - \frac{z^2}{\zeta_j^2} \right).$$

Note that if supp $\mu = \{\pm \zeta_j : j \ge 1\}$, where $0 < \zeta_1 < \zeta_2 < \dots$ then by (4.36) supp $\tilde{\mu} = \text{supp } \hat{\mu} = \{\zeta_j^2 : j \ge 1\}$. Consequently, their characteristic functions are equal to

$$\tilde{\mathfrak{F}}_1(z) = \hat{\mathfrak{F}}_1(z) = \prod_{j=1}^{\infty} \left(1 - \frac{z}{\zeta_j^2}\right).$$

Suppose that both $\tilde{\mu}$ and $\hat{\mu}$ are determinate (in Example 4.12 we show that it is possible). Then the corresponding Jacobi matrices \tilde{J}, \hat{J} are positive and their inverses belong to the trace class. Thus, by Corollary 4.10, for any fixed $w \in \mathbb{C}_+$

$$\frac{P_{2n}(z)}{P_{2n}(w)} = \frac{\tilde{P}_n(z^2)}{\tilde{P}_n(w^2)} \to \frac{\tilde{\mathfrak{F}}_1(z^2)}{\tilde{\mathfrak{F}}_1(w^2)} = \frac{\mathfrak{F}_2(z)}{\mathfrak{F}_2(w)}$$

and

$$\frac{P_{2n+1}(z)}{P_{2n+1}(w)} = \frac{z\hat{P}_n(z^2)}{w\hat{P}_n(w^2)} \to \frac{z}{w} \frac{\hat{\mathfrak{F}}_1(z^2)}{\hat{\mathfrak{F}}_1(w^2)} = \frac{z}{w} \frac{\mathfrak{F}_2(z)}{\mathfrak{F}_2(w)},$$

Consequently, we have

$$\lim_{n \to \infty} \frac{P_{2n}(z)}{P_{2n}(w)} = \frac{\mathfrak{F}_2(z)}{\mathfrak{F}_2(w)} \quad \text{whereas} \quad \lim_{n \to \infty} \frac{P_{2n+1}(z)}{P_{2n+1}(w)} = \frac{z}{w} \frac{\mathfrak{F}_2(z)}{\mathfrak{F}_2(w)}, \tag{4.37}$$

which are not equal when $z, w \in \mathbb{C}_+$ are such that $z \neq w$.

Example 4.12 (Symmetric Al-Salam–Carlitz II polynomials). Let a > 0 and 0 < q < 1. Let J be the Jacobi matrix with Jacobi parameters

$$a_{2n} = \sqrt{aq^{-n}}, \quad a_{2n+1} = \sqrt{q^{-n-1} - 1}, \quad b_n \equiv 0, \qquad n \ge 0.$$

According to [8, p. 342] all of the measures μ , $\tilde{\mu}$, $\hat{\mu}$ are determinate iff $aq^2 \ge 1$. Then, $\tilde{J} > 0$ and \tilde{J}^{-1} is trace class. Therefore, in view of (4.36), the operator J^{-1} is Hilbert–Schmidt. Thus, it falls into the setup of this subsection, and consequently, we have (4.37).

5. The modified q-Laguerre polynomials

5.1. **Preliminaries.** In this section we introduce and study the so called modified q-Laguerre polynomials. In contrast to the well established standard q-Laguerre polynomials (see [20]) they are distinguished by the fact that the respective Hamburger moment problem is determinate. Furthermore, they correspond to a Birth-Death process and they are derived from the standard q-Laguerre polynomials by a construction based on the notion of quasi-orthogonal sequence (see [4]).

The (standard) q-Laguerre polynomials are defined by the three-term recurrence relation (see e.g. [15, formula (14.21.5)])

$$\begin{split} (1-q^{n+1})L_{n+1}^{(\alpha)}(x;q) - \big(1-q^{n+1} + q(1-q^{n+\alpha})\big)L_n^{(\alpha)}(x;q) \\ &+ q(1-q^{n+\alpha})L_{n-1}^{(\alpha)}(x;q) = -q^{2n+\alpha+1}xL_n^{(\alpha)}(x;q), \quad n \geq 0, \end{split}$$

with $L_{-1}^{(\alpha)}(x;q)=0$, $L_{0}^{(\alpha)}(x;q)=1$, and $\alpha>-1$ being a parameter. The normalized recurrence relation for the respective monic polynomials $p_{n}^{(\alpha)}(x;q)$ then reads

$$\begin{split} p_{n+1}^{(\alpha)}(x;q) + q^{-2n-\alpha-1} \big(1 - q^{n+1} + q(1-q^{n+\alpha}) \big) p_n^{(\alpha)}(x;q) \\ &+ q^{-4n-2\alpha+1} (1-q^n) (1-q^{n+\alpha}) p_{n-1}^{(\alpha)}(x;q) = x p_n^{(\alpha)}(x;q), \end{split}$$

where

$$L_n^{(\alpha)}(x;q) = \frac{(-1)^n q^{n(n+\alpha)}}{(q;q)_n} p_n^{(\alpha)}(x;q). \tag{5.1}$$

Explicitly, see [15, Sec. 14.21],

$$L_n^{(\alpha)}(x;q) = \frac{(q^{\alpha+1};q)_n}{(q;q)_n} {}_1\phi_1 \left(\begin{matrix} q^{-n} \\ q^{\alpha+1} \end{matrix}; q, -q^{n+\alpha+1} x \right)$$

$$= \frac{(q^{\alpha+1};q)_n}{(q;q)_n} \sum_{k=0}^n \frac{(q^{-n};q)_k}{(q^{\alpha+1};q)_k (q;q)_k} q^{k(k-1)/2} (q^{n+\alpha+1} x)^k, \tag{5.2}$$

and

$$p_n^{(\alpha)}(x;q) = (-1)^n q^{-n(n+\alpha)} (q^{\alpha+1};q)_{n \ 1} \phi_1 \left(\frac{q^{-n}}{q^{\alpha+1}}; q, -q^{n+\alpha+1} x \right). \tag{5.3}$$

The following relation derived in [20] will be of importance in the sequel,

$$q^n L_n^{(\alpha)}(x;q) + L_{n-1}^{(\alpha+1)}(x;q) - L_n^{(\alpha+1)}(x;q) = 0, \quad n \geq 0.$$

In regard of (5.1), for the monic q-Laguerre polynomials this means that

$$p_n^{(\alpha)}(x;q) = p_n^{(\alpha+1)}(x;q) + q^{-2n-\alpha}(1-q^n)p_{n-1}^{(\alpha+1)}(x;q), \quad n \ge 0.$$
 (5.4)

The parameters $a_n^{(\alpha)},\,b_n^{(\alpha)}$ in the corresponding Jacobi matrix are given by

$$(a_n^{(\alpha)})^2 = q^{-4n-2\alpha-3}(1-q^{n+1})(1-q^{n+\alpha+1}), \quad b_n^{(\alpha)} = q^{-2n-\alpha-1}(1-q^{n+1}+q(1-q^{n+\alpha})),$$

 $n \ge 0$. We put, too,

$$d_n^{(\alpha)} = (a_{n-1}^{(\alpha)})^2 = q^{-4n-2\alpha+1}(1-q^n)(1-q^{n+\alpha}), \quad n \ge 1,$$

so that the normalized three-term recurrence relation reads

$$p_{n+1}^{(\alpha)}(x;q) = (x - b_n^{(\alpha)}) p_n^{(\alpha)}(x;q) - d_n^{(\alpha)} p_{n-1}^{(\alpha)}(x;q).$$

It turns out that we are dealing with a Jacobi parameters of the form (1.4), i.e.

$$\left(a_n^{(\alpha)}\right)^2 = \lambda_n^{(\alpha)} \mu_{n+1}^{(\alpha)}, \quad b_n^{(\alpha)} = \lambda_n^{(\alpha)} + \mu_n^{(\alpha)},$$

where

$$\lambda_n^{(\alpha)} = q^{-2n-\alpha-1}(1-q^{n+1}), \quad \mu_n^{(\alpha)} = q^{-2n-\alpha}(1-q^{n+\alpha}).$$

The Hamburger moment problem for the q-Laguerre polynomials is known to be indeterminate. A detailed discussion of the N-extremal orthogonality measures can be found in [20].

The notion of the so-called quasi-orthogonal polynomials of order $k \in \mathbb{N}$ turns out to be quite useful for our purposes although only the order k = 2 will be needed in the current paper. Here we recall a basic result which is a particular case of Theorem 2.1 in [4].

Theorem 5.1. Let $(p_n(x) : n \ge 0)$ be a monic orthogonal polynomial sequence obeying a three-term recurrence relation

$$p_{n+1}(x) = (x - b_n)p_n(x) - d_n p_{n-1}(x)$$
 for $n \ge 0$,

with $p_{-1}(x) = 0$ (hence, necessarily, $b_n \in \mathbb{R}$ for $n \ge 0$ and $d_n > 0$ for $n \ge 1$). Let $(f_n : n \ge 0)$ be a real sequence, with $f_0 = 0$ and $f_n \ne 0$ for $n \ge 1$. Let us define

$$\tilde{p}_n(x) = p_n(x) + f_n p_{n-1}(x)$$
 for $n \ge 0$,

and

$$\tilde{b}_n = b_n + f_n - f_{n+1}$$
 for $n \ge 0$,
 $\tilde{d}_n = d_n + f_n(b_{n-1} - b_n - f_n + f_{n+1})$ for $n \ge 1$.

Then $(\tilde{p}_n(x): n \geq 0)$ is a monic orthogonal polynomial sequence if and only if

$$f_{n+1} = f_n + b_n - b_{n-1} - \frac{d_n}{f_n} + \frac{d_{n-1}}{f_{n-1}} \quad \text{for } n \ge 2,$$
 (5.5)

and $\tilde{d}_n > 0$ for $n \ge 1$. If so then $(\tilde{p}_n(x) : n \ge 0)$ obeys the three-term recurrence relation

$$\tilde{p}_{n+1}(x) = (x - \tilde{b}_n)\tilde{p}_n(x) - \tilde{d}_n\tilde{p}_{n-1}(x)$$
 for $n \ge 0$,

with $\tilde{p}_{-1}(x) = 0$. Moreover, the coefficients \tilde{d}_n also satisfy

$$\tilde{d}_n = \frac{f_n}{f_{n-1}} d_{n-1}, \quad n \ge 2.$$

Remark 5.2. The polynomials $\tilde{p}_n(x)$, as defined in Theorem 5.1, are called *quasi-orthogonal polynomials* of order 2. This construction can be generalized to any order $k \in \mathbb{N}$, however, as explained in [4]. The case k = 1 is trivial as it means that $\tilde{p}_n(x) = p_n(x)$ for all $n \ge 0$. The particular case of order 2 has been explored already earlier in [18].

Further we recall a theorem which is a particular case of Theorem 2.2 in [4], with some details taken from its proof ibidem.

Theorem 5.3. Suppose that the quasi-orthogonal polynomial sequence $(\tilde{p}_n(x) : n \ge 0)$, as defined in Theorem 5.1, is in fact a monic orthogonal polynomial sequence.

Denote by Λ and Λ their moment functionals, and by $(M_n:n\geq 0)$ and $(\tilde{M}_n:n\geq 0)$ the moment sequences associated with the polynomial sequences $(p_n(x):n\geq 0)$ and $(\tilde{p}_n(x):n\geq 0)$, respectively. Using the notation of Theorem 5.1 let

$$h(x) = 1 - \frac{f_1 \tilde{b}_0}{\tilde{d}_1} + \frac{f_1}{\tilde{d}_1} x.$$

Then for every polynomial f,

$$\Lambda(f) = \tilde{\Lambda}(hf).$$

Consequently,

$$\forall n \ge 0, \ \tilde{M}_{n+1} = \frac{\tilde{d}_1}{f_1} M_n + \left(\tilde{b}_0 - \frac{\tilde{d}_1}{f_1} \right) \tilde{M}_n.$$
 (5.6)

5.2. **Definition.** Let

$$\gamma(u, v; q) = \frac{(q^u; q)_{\infty}}{(q^{u+v}; q)_{\infty}}.$$

In particular, for v > 0,

$$\gamma(0, v; q) = 0.$$

For $m \in \mathbb{N}_0$ we have $\gamma(u, m; q) = (q^u; q)_m$. In regard to the q-Binomial Theorem telling us that [11, Eq. (II.3)]

$${}_{1}\phi_{0}\left({a\atop -};q,z\right) = \frac{(az;q)_{\infty}}{(z;q)_{\infty}}, \quad |z| < 1, \tag{5.7}$$

we also have, for u > 0 and u + v > 0,

$$\gamma(u, v; q) = \frac{1}{1\phi_0\left(\frac{q^v}{-}; q, q^u\right)} = 1\phi_0\left(\frac{q^{-v}}{-}; q, q^{u+v}\right). \tag{5.8}$$

Furthermore, one can rewrite the function $\gamma(u, v; q)$ in terms of the q-Gamma function. Recall that ([11, Eq. (I.35)] or [15, Eq. (1.9.1)])

$$\Gamma_q(z) = \frac{(q;q)_{\infty}}{(q^z;q)_{\infty}} (1-q)^{1-z}.$$

We have

$$\gamma(u, v; q) = \frac{\Gamma_q(u+v)}{\Gamma_q(u)} (1-q)^v. \tag{5.9}$$

For later convenience, we show some identities for function $\gamma(u, v; q)$.

Lemma 5.4. For $m \in \mathbb{N}_0$ and $\beta > -1$,

$$\sum_{k=0}^{m-1} q^k \gamma(k+1, \beta; q) = \frac{\gamma(m, \beta+1; q)}{1 - q^{\beta+1}}$$

and

$$\sum_{k=0}^{\infty} q^k \gamma(k+1,\beta;q) = \frac{1}{1 - q^{\beta+1}}.$$
 (5.10)

In particular, for $\ell \in \mathbb{N}_0$,

$$\sum_{k=0}^{m-1} q^k (q^{k+1}; q)_{\ell} = \frac{(q^m; q)_{\ell+1}}{1 - q^{\ell+1}}$$
 (5.11)

and

$$\sum_{k=0}^{\infty} q^k (q^{k+1}; q)_{\ell} = \frac{1}{1 - q^{\ell+1}}.$$
 (5.12)

Proof. Letting $a = q^{\beta+1}$ and $c = q^{-n}$ in the q-Vandermonde Sum (see e.g. [11, Eq. (II.6)]) telling us that

$$_{2}\phi_{1}\begin{pmatrix} a, q^{-n} \\ c \end{pmatrix}; q, q = \frac{(c/a; q)_{n}}{(c; q)_{n}}a^{n},$$

we get

$$\sum_{k=0}^{n} q^{k} \frac{(q^{\beta+1}; q)_{k}}{(q; q)_{k}} = \frac{(q^{\beta+2}; q)_{n}}{(q; q)_{n}}, \quad n \ge 0.$$
 (5.13)

Then

$$\begin{split} \sum_{k=0}^{m-1} q^k \gamma(k+1,\beta;q) &= \frac{(q;q)_{\infty}}{(q^{\beta+1};q)_{\infty}} \sum_{k=0}^m q^k \frac{(q^{\beta+1};q)_k}{(q;q)_k} \\ &= \frac{(q;q)_{\infty}}{(q^{\beta+1};q)_{\infty}} \frac{(q^{\beta+2};q)_{m-1}}{(q;q)_{m-1}} \\ &= \frac{\gamma(m,\beta+1;q)}{1-q^{\beta+1}}. \end{split}$$

In the limit $m \to \infty$ we get (5.10). Equations (5.11) and (5.12) are particular cases for $\beta = \ell \in \mathbb{N}_0$. Let us note that to show (5.12) one can alternatively use the *q*-Binomial Theorem (5.7) while letting $a = q^{\ell+1}$ and z = q. Let

$$r_n^{(\alpha)} = \frac{1 - \gamma(n+1, \alpha+1; q)}{1 - \gamma(n, \alpha+1; q)}.$$
 (5.14)

Then

$$1 - r_n^{(\alpha)} = \frac{q^n (1 - q^{\alpha + 1}) \gamma(n + 1, \alpha; q)}{1 - \gamma(n, \alpha + 1; q)}.$$
 (5.15)

To see it, note that

$$(1 - \gamma(u, v + 1; q)) - (1 - \gamma(u + 1, v + 1; q)) = q^{u}(1 - q^{v+1})\gamma(u + 1, v; q).$$

It is known that $\lim_{q \uparrow 1} \Gamma_q(x) = \Gamma(x)$ [11, Eq. (I.36)]. Hence from (5.9) we infer that, for $u \ge 0$, v > 0,

$$\gamma(u, v; q) \approx \frac{\Gamma(u + v)}{\Gamma(u)} (1 - q)^{v}$$
 as $q \uparrow 1$.

From here and (5.14), while still assuming that $\alpha > -1$, one finds that

$$\forall n \in \mathbb{N}_0, \ \lim_{q \uparrow 1} r_n^{(\alpha)} = 1. \tag{5.16}$$

We define the modified q-Laguerre polynomials $\tilde{L}_{n}^{(\alpha)}(x;q)$ as

$$\tilde{L}_n^{(\alpha)}(x;q) = q^n r_n^{(\alpha)} L_n^{(\alpha)}(x;q) + \left(1 - r_n^{(\alpha)}\right) L_n^{(\alpha+1)}(x;q).$$

The respective monic polynomials are denoted $\tilde{p}_n^{(\alpha)}(x;q)$, and they fulfill

$$\tilde{p}_{n}^{(\alpha)}(x;q) = r_{n}^{(\alpha)} p_{n}^{(\alpha)}(x;q) + (1 - r_{n}^{(\alpha)}) p_{n}^{(\alpha+1)}(x;q). \tag{5.17}$$

Thus the monic modified q-Laguerre polynomials are defined as a convex combination of the "standard" monic q-Laguerre polynomials.

It is known that [15]

$$\lim_{q \uparrow 1} L_n^{(\alpha)} \left((1 - q)x; q \right) = L_n^{(\alpha)}(x)$$

where $L_n^{(\alpha)}(x)$ are the ordinary Laguerre polynomials. Therefore, in view of (5.16), we also have

$$\lim_{q\uparrow 1} \tilde{L}_n^{(\alpha)}\big((1-q)x;q\big) = L_n^{(\alpha)}(x).$$

5.3. Quasi-orthogonal polynomials as Birth-Death polynomials. Let

$$\tilde{\lambda}_n^{(\alpha)} = \frac{q^{-2n-\alpha-1}(1-q^{n+\alpha+1})}{r_n^{(\alpha)}}, \quad \tilde{\mu}_n^{(\alpha)} = q^{-2n-\alpha}(1-q^n)r_n^{(\alpha)}, \tag{5.18}$$

where $r_n^{(\alpha)}$ is defined in (5.14) (hence $\tilde{\mu}_0^{(\alpha)} = 0$).

The modified q-Laguerre polynomials in the monic form, $\tilde{p}_n^{(\alpha)}(x;q)$, have been defined in (5.17) as a convex combination of $p_n^{(\alpha)}(x;q)$ and $p_n^{(\alpha+1)}(x;q)$. But owing to identity (5.4) we can also write

$$\tilde{p}_{n}^{(\alpha)}(x;q) = p_{n}^{(\alpha+1)}(x;q) + \tilde{\mu}_{n}^{(\alpha)} p_{n-1}^{(\alpha+1)}(x;q). \tag{5.19}$$

Hence $(\tilde{p}_n^{(\alpha)}(x;q):n\geq 0)$ is a quasi-orthogonal polynomial sequence derived from the monic orthogonal polynomial sequence $(p_n^{(\alpha+1)}(x;q):n\geq 0)$. With this observation, Theorem 5.1 is directly applicable to the modified q-Laguerre polynomials.

Theorem 5.5. The quasi-orthogonal polynomials $(\tilde{p}_n^{(\alpha)}(x;q):n\geq 0)$ constitute a monic orthogonal polynomial sequence corresponding to the Birth-Death process with the parameters (5.18). Hence this polynomial sequence obeys a three-term recurrence relation

$$\tilde{p}_{n+1}^{(\alpha)}(x;q) = \left(x - \tilde{b}_n^{(\alpha)}\right) \tilde{p}_n^{(\alpha)}(x;q) - \tilde{d}_n^{(\alpha)} \tilde{p}_{n-1}^{(\alpha)}(x;q), \quad n \geq 0,$$

with $\tilde{p}_{-1}^{(\alpha)}(x;q)=0$, where the coefficients are given by $\tilde{b}_{n}^{(\alpha)}=\tilde{\lambda}_{n}^{(\alpha)}+\tilde{\mu}_{n}^{(\alpha)}$, $n\geq 0$, and

$$\tilde{d}_n^{(\alpha)} = \tilde{\lambda}_{n-1}^{(\alpha)} \tilde{\mu}_n^{(\alpha)} = q^{-4n-2\alpha+1} (1 - q^n) (1 - q^{n+\alpha}) \frac{r_n^{(\alpha)}}{r_{n-1}^{(\alpha)}} \quad n \ge 1.$$

Proof. Comparing to the notation used in Theorem 5.1 we have

$$b_n \equiv b_n^{(\alpha+1)}, \quad d_n \equiv d_n^{(\alpha+1)}, \quad f_n = \tilde{\mu}_n^{(\alpha)}.$$

Thus we have to show that

$$\tilde{\mu}_{n+1}^{(\alpha)} = \tilde{\mu}_n^{(\alpha)} + b_n^{(\alpha+1)} - b_{n-1}^{(\alpha+1)} - \frac{d_n^{(\alpha+1)}}{\tilde{\mu}_n^{(\alpha)}} + \frac{d_{n-1}^{(\alpha+1)}}{\tilde{\mu}_{n-1}^{(\alpha)}} \quad \text{for } n \ge 2.$$

Obviously, it suffices to show that

$$\tilde{\mu}_{n+1}^{(\alpha)} - b_n^{(\alpha+1)} + \frac{d_n^{(\alpha+1)}}{\tilde{\mu}_n^{(\alpha)}} = 0 \quad \text{for } n \ge 1.$$
 (5.20)

But this equation can be verified by a straightforward computation.

Furthermore, it is also straightforward to verify that

$$\frac{\tilde{\mu}_n^{(\alpha)}}{\tilde{\mu}_{n-1}^{(\alpha)}}d_n^{(\alpha+1)} = \tilde{d}_n^{(\alpha)} \quad \text{and} \quad b_n^{(\alpha+1)} + \tilde{\mu}_n^{(\alpha)} - \tilde{\mu}_{n+1}^{(\alpha)} = \tilde{b}_n^{(\alpha)}.$$

Finally, one observes that $\tilde{d}_n^{(\alpha)} > 0$ for $n \ge 1$.

Further we can apply Theorem 5.3 to determine the moments associated with the modified q-Laguerre polynomials. Let us recall that the moment sequence $(M_n^{(\alpha)}:n\geq 0)$ associated with the q-Laguerre polynomials is well known, see [20]. When consulting this source some caution is needed, however, because the notation in [20] slightly differs from the commonly used notation like that introduced in [15]. Namely, the variable x in the q-Laguerre polynomials is rescaled by the factor (1-q) in the former source, and the same is true for the q-Bessel functions. We have

$$\forall n \ge 0, \ M_n^{(\alpha)} = q^{-\alpha n - (n+1)n/2} (q^{\alpha+1}; q)_n. \tag{5.21}$$

Proposition 5.6. The moments $\tilde{M}_n^{(\alpha)}$ associated with the modified q-Laguerre polynomials take the values

$$\tilde{M}_{0}^{(\alpha)} = 1$$
 and $\tilde{M}_{n}^{(\alpha)} = q^{-\alpha n - (n+1)n/2} \frac{(q^{\alpha+1};q)_{n}}{r_{0}^{(\alpha)}}$ for $n \ge 1$.

Proof. Sticking to the notation of Theorem 5.1 one can readily check that (5.5) is equivalent to the equation

$$f_n - \tilde{b}_n + \frac{\tilde{d}_{n+1}}{f_{n+1}} = f_{n+1} - \tilde{b}_{n+1} + \frac{\tilde{d}_{n+2}}{f_{n+2}}, \quad n \ge 0.$$

Furthermore, one finds that

$$f_n - \tilde{b}_n + \frac{\tilde{d}_{n+1}}{f_{n+1}} = f_{n+1} - b_n + \frac{d_n}{f_n}, \quad n \ge 1.$$

Dealing with the modified monic *q*-Laguerre polynomials regarded as quasi-orthogonal polynomials one has to substitute $\tilde{\mu}_n^{(\alpha)}$ for f_n , see (5.19). In view of (5.20) we get

$$\tilde{\mu}_{n}^{(\alpha)} - \tilde{b}_{n}^{(\alpha)} + \frac{\tilde{d}_{n+1}^{(\alpha)}}{\tilde{\mu}_{n+1}^{(\alpha)}} = \tilde{\mu}_{n+1}^{(\alpha)} - \tilde{b}_{n+1}^{(\alpha)} + \frac{\tilde{d}_{n+2}^{(\alpha)}}{\tilde{\mu}_{n+2}^{(\alpha)}} = \tilde{\mu}_{n+2}^{(\alpha)} - b_{n+1}^{(\alpha+1)} + \frac{d_{n+1}^{(\alpha+1)}}{\tilde{\mu}_{n+1}^{(\alpha)}} = 0, \quad n \geq 0.$$

In particular, for n = 0, $\tilde{\mu}_n^{(\alpha)} = 0$ and

$$\tilde{b}_0^{(\alpha)} = \frac{\tilde{d}_1^{(\alpha)}}{\tilde{\mu}_1^{(\alpha)}} = \frac{q^{-\alpha - 1}(1 - q^{\alpha + 1})}{r_0^{(\alpha)}}.$$

By Theorem 5.3, equation (5.6), we get

$$\forall n \geq 0, \ \tilde{M}_{n+1}^{(\alpha)} = \frac{q^{-\alpha-1}(1-q^{\alpha+1})}{r_0^{(\alpha)}} M_n^{(\alpha+1)}.$$

In view of (5.21), the proposition readily follows.

Remark 5.7. As already noted, the moment problem for the q-Laguerre polynomials is indeterminate. On the other hand, we shall see later that the moment problem for the modified q-Laguerre polynomials is determinate. The fact that, except the first terms, both moment sequences are proportional immediately implies that Carleman's sufficient condition for the determinacy (see e.g. [21, Theorem 4.3]) is *not* applicable to the modified q-Laguerre polynomials.

5.4. **The Jacobi matrix operator.** For the Birth-Death polynomials with the parameters (5.18) we have

$$\tilde{\pi}_{n}^{(\alpha)} = \prod_{k=0}^{n-1} \frac{\tilde{\lambda}_{k}^{(\alpha)}}{\tilde{\mu}_{k+1}^{(\alpha)}} = \frac{q^{n}(q^{\alpha+1};q)_{n}(1-\gamma(1,\alpha+1;q))}{(q;q)_{n}(1-\gamma(n,\alpha+1;q))(1-\gamma(n+1,\alpha+1;q))}$$

and

$$\tilde{\lambda}_{n}^{(\alpha)}\tilde{\pi}_{n}^{(\alpha)} = \frac{q^{-n-\alpha-1}(q^{\alpha+1};q)_{n+1}(1-\gamma(1,\alpha+1;q))}{(q;q)_{n}(1-\gamma(n+1,\alpha+1;q))^{2}}.$$
(5.22)

The corresponding Jacobi parameters $\tilde{a}_n^{(\alpha)}$, $\tilde{b}_n^{(\alpha)}$, $n \ge 0$, are given by

$$\begin{split} &(\tilde{a}_{n}^{(\alpha)})^{2} = \tilde{\lambda}_{n}^{(\alpha)} \tilde{\mu}_{n+1}^{(\alpha)} = q^{-4n-2\alpha-3} (1-q^{n+1}) (1-q^{n+\alpha+1}) \frac{r_{n+1}^{(\alpha)}}{r_{n}^{(\alpha)}}, \\ &\tilde{b}_{n}^{(\alpha)} = \tilde{\lambda}_{n}^{(\alpha)} + \tilde{\mu}_{n}^{(\alpha)} = q^{-2n-\alpha-1} \bigg(\frac{(1-q^{n+\alpha+1})}{r_{n}^{(\alpha)}} + q (1-q^{n}) r_{n}^{(\alpha)} \bigg). \end{split}$$

The Jacobi matrix operator with these entries will be denoted $\tilde{J}^{(\alpha)}$.

Lemma 5.8. For $n \in \mathbb{N}_0$,

$$\sum_{j=0}^n \tilde{\pi}_j^{(\alpha)} = \frac{(q^{\alpha+2};q)_n \left(1-\gamma(1,\alpha+1;q)\right)}{(q;q)_n \left(1-\gamma(n+1,\alpha+1;q)\right)}.$$

Proof. Note that

$$\frac{1}{1 - \gamma(j+1,\alpha+1;q)} - \frac{1}{1 - \gamma(j,\alpha+1;q)} = \frac{q^{j}(1 - q^{\alpha+1})\gamma(j+1,\alpha;q)}{(1 - \gamma(j,\alpha+1;q))(1 - \gamma(j+1,\alpha+1;q))}.$$

Thus we have

$$\begin{split} \sum_{j=0}^{n} \tilde{\pi}_{j}^{(\alpha)} &= \left(1 - \gamma(1, \alpha + 1; q)\right) \sum_{j=0}^{n} \frac{q^{j}(q^{\alpha+1}; q)_{j}}{(q; q)_{j} \left(1 - \gamma(j, \alpha + 1; q)\right) \left(1 - \gamma(j + 1, \alpha + 1; q)\right)} \\ &= \left(1 - \gamma(1, \alpha + 1; q)\right) \frac{(q^{\alpha+2}; q)_{\infty}}{(q; q)_{\infty}} \sum_{j=0}^{n} \left(\frac{1}{1 - \gamma(j + 1, \alpha + 1; q)} - \frac{1}{1 - \gamma(j, \alpha + 1; q)}\right) \\ &= \left(1 - \gamma(1, \alpha + 1; q)\right) \frac{(q^{\alpha+2}; q)_{\infty}}{(q; q)_{\infty}} \left(\frac{1}{1 - \gamma(n + 1, \alpha + 1; q)} - 1\right) \\ &= \frac{(q^{\alpha+2}; q)_{n}}{(q; q)_{n}} \frac{1 - \gamma(1, \alpha + 1; q)}{1 - \gamma(n + 1, \alpha + 1; q)}. \end{split}$$

Theorem 5.9. The Hamburger moment problem for the modified q-Laguerre polynomials is determinate or, equivalently, the Jacobi operator $\tilde{J}^{(\alpha)}$ is self-adjoint. Moreover, $\tilde{J}^{(\alpha)}$ is positive, the inversion $(\tilde{J}^{(\alpha)})^{-1}$ exists and belongs to the trace class.

Proof. We claim that

$$\sum_{j=0}^{\infty} \tilde{\pi}_{j}^{(\alpha)} = \infty \quad \text{and} \quad \sum_{j=0}^{\infty} \frac{1}{\tilde{\lambda}_{j}^{(\alpha)} \tilde{\pi}_{j}^{(\alpha)}} \sum_{n=0}^{j} \tilde{\pi}_{n}^{(\alpha)} < \infty.$$

Firstly, by Lemma 5.8,

$$\sum_{j=0}^{\infty} \tilde{\pi}_j^{(\alpha)} = \lim_{n \to \infty} \frac{(q^{\alpha+2}; q)_n (1 - \gamma(1, \alpha + 1; q))}{(q; q)_n (1 - \gamma(n + 1, \alpha + 1; q))} = \infty.$$

Indeed, in virtue of (5.8),

$$\lim_{n\to\infty}\gamma(n+1,\alpha+1;q)=\lim_{n\to\infty}\frac{1}{1\phi_0\left(\begin{smallmatrix}q^{\alpha+1}\\-\end{smallmatrix};q,q^{n+1}\right)}=1.$$

Further,

$$\sum_{j=0}^{\infty} \frac{1}{\tilde{\lambda}_{j}^{(\alpha)} \tilde{\pi}_{j}^{(\alpha)}} \sum_{n=0}^{j} \tilde{\pi}_{n}^{(\alpha)} = \frac{q^{\alpha+1}}{1-q^{\alpha+1}} \sum_{j=0}^{\infty} q^{j} \left(1-\gamma(j+1,\alpha+1;q)\right) < \infty.$$

The theorem now follows straightforwardly from Theorem 3.3. Note that $\tilde{\mu}_0^{(\alpha)} = 0$ and therefore $\mathfrak{s}_n \equiv 0$.

We can compute the trace of $(\tilde{J}^{(\alpha)})^{-1}$ explicitly. By Theorem 3.3 and using (5.10) we obtain

$$\operatorname{Tr}\left((\tilde{J}^{(\alpha)})^{-1}\right) = \frac{q^{\alpha+1}}{1 - q^{\alpha+1}} \sum_{j=0}^{\infty} q^{j} \left(1 - \gamma(j+1, \alpha+1; q)\right) = \frac{q^{\alpha+2}}{(1-q)(1-q^{\alpha+2})}.$$
 (5.23)

5.5. The characteristic function, the orthogonality measure. It follows from Theorem 5.9 that the Jacobi operator $\tilde{J}^{(\alpha)}$ has a trace class resolvent, its spectrum consists of countably many simple positive eigenvalues, denoted as $\{\zeta_k^{(\alpha)}: k \ge 1\}$, and

$$\sum_{k=1}^{\infty} \frac{1}{\zeta_k^{(\alpha)}} = \operatorname{Tr}\left((\tilde{J}^{(\alpha)})^{-1}\right) < \infty.$$

The eigenvalues are supposed to be ordered increasingly, $0 < \zeta_1^{(\alpha)} < \zeta_2^{(\alpha)} < \zeta_3^{(\alpha)} < \dots$ The characteristic function of $\tilde{J}^{(\alpha)}$, as introduced in Section 4.4, reads

$$\tilde{\mathfrak{F}}^{(\alpha)}(z) = \prod_{k=1}^{\infty} \left(1 - \frac{z}{\zeta_k^{(\alpha)}}\right).$$

The function $\tilde{\mathfrak{F}}^{(\alpha)}$ is entire and, as stated in Corollary 4.10,

$$\lim_{n \to \infty} \frac{\tilde{p}_n^{(\alpha)}(z;q)}{\tilde{p}_n^{(\alpha)}(0;q)} = \tilde{\mathfrak{F}}^{(\alpha)}(z)$$

locally uniformly with respect to $z \in \mathbb{C}$.

The function $\tilde{\mathfrak{F}}^{(\alpha)}$ turns out to be expressible in terms of the q-Bessel function. Recall that the Jackson q-Bessel functions of the second kind are

$$J_{\nu}^{(2)}(x;q) := \frac{(q^{\nu+1};q)_{\infty}}{(q;q)_{\infty}} \left(\frac{x}{2}\right)^{\nu} {}_{0}\phi_{1}\left(\begin{array}{c} - \\ q^{\nu+1} \end{array};q,-\frac{q^{\nu+1}x^{2}}{4}\right).$$

Theorem 5.10. The characteristic function of the of the Jacobi operator $\tilde{J}^{(\alpha)}$ equals

$$\tilde{\mathfrak{F}}^{(\alpha)}(z) = {}_{0}\phi_{1}\left(\begin{matrix} - \\ q^{\alpha+2} \end{matrix}; q, -q^{\alpha+2}z \right) = \frac{(q;q)_{\infty}}{(q^{\alpha+2};q)_{\infty}} z^{-(\alpha+1)/2} J_{\alpha+1}^{(2)}(2\sqrt{z};q). \tag{5.24}$$

Proof. Let us write

$$p_n^{(\alpha)}(x;q) = \sum_{k=0}^n p_{n,k}^{(\alpha)} x^k, \quad \tilde{p}_n^{(\alpha)}(x;q) = \sum_{k=0}^n \tilde{p}_{n,k}^{(\alpha)} x^k.$$

The q-Laguerre polynomials are known explicitly, see (5.2) and (5.3). With regard to (5.17), we find that, for $n \ge k \ge 0$,

$$\frac{\tilde{p}_{n,k}^{(\alpha)}}{\tilde{p}_{n,0}^{(\alpha)}} = r_n^{(\alpha)} \frac{p_{n,k}^{(\alpha)}}{\tilde{p}_{n,0}} + (1 - r_n^{(\alpha)}) \frac{p_{n,k}^{(\alpha+1)}}{\tilde{p}_{n,0}} \\
= \frac{(-1)^k q^{k(k-1)+k(\alpha+1)} (q^{n-k+1}; q)_k}{(q^{\alpha+1}; q)_k (q; q)_k} \left(1 - \frac{1 - q^k}{1 - q^{k+\alpha+1}} \frac{(q^{n+1}; q)_{\infty}}{(q^{n+\alpha+2}; q)_{\infty}}\right).$$
(5.25)

Further we write

$$\tilde{\mathfrak{F}}^{(\alpha)}(z) = \sum_{k=0}^{\infty} F_k z^k,$$

and chose any C^1 positively oriented Jordan curve C with 0 in its interior. Then, owing to the locally uniform convergence, for every $k \in \mathbb{N}_0$ it holds true that

$$F_{k} = \frac{1}{2\pi i} \int_{C} \frac{\tilde{\mathfrak{F}}^{(\alpha)}(z)}{z^{k+1}} dz = \frac{1}{2\pi i} \int_{C} \lim_{n \to \infty} \frac{\tilde{p}_{n}^{(\alpha)}(z;q)}{\tilde{p}_{n}^{(\alpha)}(0;q)z^{k+1}} dz$$
$$= \frac{1}{2\pi i} \lim_{n \to \infty} \int_{C} \frac{\tilde{p}_{n}^{(\alpha)}(z;q)}{\tilde{p}_{n}^{(\alpha)}(0;q)z^{k+1}} dz = \lim_{n \to \infty} \frac{\tilde{p}_{n,k}^{(\alpha)}}{\tilde{p}_{n,0}^{(\alpha)}}.$$

In view of (5.25), we have

$$\lim_{n \to \infty} \frac{\tilde{p}_{n,k}^{(\alpha)}}{\tilde{p}_{n,0}^{(\alpha)}} = \frac{(-1)^k q^{k(k-1)+k(\alpha+1)}}{(q^{\alpha+1};q)_k(q;q)_k} \left(1 - \frac{1-q^k}{1-q^{k+\alpha+1}}\right) = \frac{(-1)^k q^{k(k+\alpha+1)}}{(q^{\alpha+2};q)_k(q;q)_k}.$$

Whence

$$\tilde{\mathfrak{F}}^{(\alpha)}(z) = \sum_{k=0}^{\infty} \frac{(-1)^k q^{k(k+\alpha+1)}}{(q^{\alpha+2};q)_k (q;q)_k} z^k = {}_{0}\phi_1 \begin{pmatrix} - \\ q^{\alpha+2} \end{pmatrix}; q, -q^{\alpha+2}z .$$

This shows (5.24).

Remark 5.11. Notice that from (5.24) one can rederive the trace formula (5.23),

$$\operatorname{Tr}\left((\tilde{J}^{(\alpha)})^{-1}\right) = -\frac{\operatorname{d}\tilde{\mathfrak{F}}^{(\alpha)}(0)}{\operatorname{d}z} = \frac{q^{\alpha+2}}{(1-q)(1-q^{\alpha+2})}.$$

The moment functional $\tilde{\Lambda}^{(\alpha)}$ for the modified q-Laguerre polynomials is described in Proposition 5.6. The corresponding orthogonality measure $\tilde{\mu}^{(\alpha)}$ is a unique probability Borel measure on \mathbb{R} such that for every polynomial f,

$$\tilde{\Lambda}^{(\alpha)}(f) = \int_{\mathbb{D}} f(x) \, \mathrm{d}\tilde{\mu}^{(\alpha)}(x).$$

As an immediate corollary of Theorem 5.10 and some general results including Eq. (2.7) we have the following description of $\tilde{\mu}^{(\alpha)}$.

Corollary 5.12. The orthogonality measure $\tilde{\mu}^{(\alpha)}$ is supported on the set of positive roots of the function $J_{\alpha+1}^{(2)}(2\sqrt{z};q)$. For every positive root ζ of $J_{\alpha+1}^{(2)}(2\sqrt{z};q)$,

$$\tilde{\mu}^{(\alpha)}(\{\zeta\}) = \left(\sum_{n=0}^{\infty} \tilde{P}_n^{(\alpha)}(\zeta)^2\right)^{-1} = -\tilde{\mathfrak{W}}^{(\alpha)}(\zeta) \left(\frac{\mathrm{d}\tilde{\mathfrak{F}}^{(\alpha)}(\zeta)}{\mathrm{d}z}\right)^{-1},$$

where

$$\tilde{\mathfrak{B}}^{(\alpha)}(z) = \tilde{w}^{(\alpha)}(z)\tilde{\mathfrak{F}}^{(\alpha)}(z)$$

and $\tilde{w}^{(\alpha)}(z)$ is the corresponding Weyl function.

5.6. The characteristic function continued. Applying (3.23) to the modified q-Laguerre polynomials we can re-derive formula (5.24) for the characteristic function.

In view of Lemma 5.8 and in agreement with definition (3.22) we put

$$\tilde{T}_{k}^{(\alpha)} = \frac{(q^{\alpha+2}; q)_{k} (1 - \gamma(1, \alpha + 1; q))}{(q; q)_{k} (1 - \gamma(k + 1, \alpha + 1; q))}, \quad k \ge 0.$$
(5.26)

Lemma 5.13. The sequences $\{\tilde{\lambda}_k^{(\alpha)}\tilde{\pi}_k^{(\alpha)}\}$ and $\{\tilde{T}_k^{(\alpha)}\}$, as given in (5.22) and (5.26), respectively, satisfy the equations

$$\sum_{k=0}^{m-1} (q^k; q^{-1})_{\ell} \frac{\tilde{T}_k^{(\alpha)} (\tilde{T}_m^{(\alpha)} - \tilde{T}_k^{(\alpha)})}{\tilde{\lambda}_k^{(\alpha)} \tilde{\pi}_k^{(\alpha)}} = \frac{q^{2\ell + \alpha + 2} (q^m; q^{-1})_{\ell+1}}{(1 - q^{\ell+1})(1 - q^{\ell + \alpha + 2})} \tilde{T}_m^{(\alpha)}, \quad m, \ell \ge 0,$$
 (5.27)

and

$$\sum_{k=0}^{\infty} (q^k; q^{-1})_{\ell} \frac{\tilde{T}_k^{(\alpha)}}{\tilde{\lambda}_k^{(\alpha)} \tilde{\pi}_k^{(\alpha)}} = \frac{q^{2\ell + \alpha + 2}}{(1 - q^{\ell + 1})(1 - q^{\ell + \alpha + 2})}, \quad \ell \ge 0.$$
 (5.28)

Proof. First let us consider the sum

$$\sum_{k=0}^{m-1} (q^k; q^{-1})_{\ell} \frac{\tilde{T}_k^{(\alpha)}}{\tilde{\lambda}_k^{(\alpha)} \tilde{\pi}_k^{(\alpha)}}, \quad m, \ell \ge 0.$$

Notice that this expression vanishes for $m \le \ell$. Assuming that $m \ge \ell + 1$, the sum equals

$$\frac{q^{\alpha+1}}{1-q^{\alpha+1}} \sum_{k=0}^{m-1} q^k (q^k; q^{-1})_{\ell} (1-\gamma(k+1, \alpha+1; q))$$

$$=\frac{q^{\ell+\alpha+1}}{1-q^{\alpha+1}}\sum_{k=0}^{m-\ell-1}q^k(q^{k+\ell};q^{-1})_\ell\bigg(1-\frac{(q^{k+\ell+1};q)_\infty}{(q^{k+\ell+\alpha+2};q)_\infty}\bigg).$$

Notice that $(q^{k+\ell}; q^{-1})_{\ell} = (q^{k+1}; q)_{\ell}$. Applying (5.11) and afterwards (5.13) we get the expression

$$\frac{q^{\ell+\alpha+1}}{1-q^{\alpha+1}} \left(\frac{(q^{m-\ell};q)_{\ell+1}}{1-q^{\ell+1}} - \frac{(q;q)_{\infty}}{(q^{\ell+\alpha+2};q)_{\infty}} \sum_{k=0}^{m-\ell-1} q^{k} \frac{(q^{\ell+\alpha+2};q)_{k}}{(q;q)_{k}} \right) \\
= \frac{q^{\ell+\alpha+1}}{1-q^{\alpha+1}} \left(\frac{(q^{m-\ell};q)_{\ell+1}}{1-q^{\ell+1}} - \frac{(q;q)_{\infty}}{(q^{\ell+\alpha+2};q)_{\infty}} \frac{(q^{\ell+\alpha+3};q)_{m-\ell-1}}{(q;q)_{m-\ell-1}} \right) \\
= \frac{q^{\ell+\alpha+1}}{1-q^{\alpha+1}} \left(\frac{(q^{m-\ell};q)_{\ell+1}}{1-q^{\ell+1}} - \frac{(q^{m-\ell};q)_{\ell+1}(q^{m+1};q)_{\infty}}{(1-q^{\ell+\alpha+2})(q^{m+\alpha+2};q)_{\infty}} \right).$$

Finally we find that

$$\sum_{k=0}^{m-1} (q^k; q^{-1})_{\ell} \frac{\tilde{T}_k^{(\alpha)}}{\tilde{\lambda}_k^{(\alpha)} \tilde{\pi}_k^{(\alpha)}} = \frac{q^{\ell+\alpha+1}(q^{m-\ell}; q)_{\ell+1}}{(1-q^{\alpha+1})(1-q^{\ell+\alpha+2})} \left(\frac{q^{\ell+1}(1-q^{\alpha+1})}{1-q^{\ell+1}} + 1 - \gamma(m+1, \alpha+1; q) \right). \tag{5.29}$$

Note that (5.29) holds for $m \le \ell$ as well since both sides vanish.

Sending $m \to \infty$ in (5.29) we get (5.28).

Next, again assuming that $m \ge \ell + 1$, we consider the sum

$$\begin{split} \sum_{k=0}^{m-1} (q^k; q^{-1})_\ell \, \frac{(\tilde{T}_k^{(\alpha)})^2}{\tilde{\lambda}_k^{(\alpha)} \tilde{\pi}_k^{(\alpha)}} &= \left(1 - \gamma(1, \alpha + 1; q)\right) \frac{q^{\alpha + 1}}{1 - q^{\alpha + 1}} \sum_{k=0}^{m-1} q^k (q^k; q^{-1})_\ell \, \frac{(q^{\alpha + 2}; q)_k}{(q; q)_k} \\ &= \left(1 - \gamma(1, \alpha + 1; q)\right) \frac{q^{\ell + \alpha + 1} (q^{\alpha + 2}; q)_\ell}{1 - q^{\alpha + 1}} \sum_{k=0}^{m-\ell-1} q^k \, \frac{(q^{\ell + \alpha + 2}; q)_k}{(q; q)_k}. \end{split}$$

Applying (5.13) we get the expression

$$\begin{split} & \left(1 - \gamma(1, \alpha + 1; q)\right) \frac{q^{\ell + \alpha + 1}(q^{\alpha + 2}; q)_{\ell}(q^{\ell + \alpha + 3}; q)_{m - \ell - 1}}{(1 - q^{\alpha + 1})(q; q)_{m - \ell - 1}} \\ &= \left(1 - \gamma(1, \alpha + 1; q)\right) \frac{q^{\ell + \alpha + 1}(q^m; q^{-1})_{\ell + 1}(q^{\alpha + 2}; q)_m}{(1 - q^{\alpha + 1})(1 - q^{\ell + \alpha + 2})(q; q)_m}. \end{split}$$

With regard to (5.26) we obtain

$$\sum_{k=0}^{m-1} (q^k; q^{-1})_{\ell} \frac{(\tilde{T}_k^{(\alpha)})^2}{\tilde{\lambda}_k^{(\alpha)} \tilde{\pi}_k^{(\alpha)}} = \frac{q^{\ell+\alpha+1} (q^m; q^{-1})_{\ell+1}}{(1-q^{\alpha+1})(1-q^{\ell+\alpha+2})} (1 - \gamma(k+1, \alpha+1; q)) \tilde{T}_m^{(\alpha)}.$$
 (5.30)

A combination of (5.29) and (5.30) gives (5.27).

Proposition 5.14. The sequences $(\tilde{\lambda}_k^{(\alpha)}\pi_k)$ and $(\tilde{T}_k^{(\alpha)})$, as given in (5.22) and (5.26), respectively, satisfy the equations, for $m \ge 1$ and $\ell \ge 0$,

$$\sum_{0 \le k_1 < k_2 < \dots < k_m} (q^{k_1}; q^{-1})_{\ell} \frac{\tilde{T}_{k_1}^{(\alpha)} (\tilde{T}_{k_2}^{(\alpha)} - \tilde{T}_{k_1}^{(\alpha)}) \dots (\tilde{T}_{k_m}^{(\alpha)} - \tilde{T}_{k_{m-1}}^{(\alpha)})}{\tilde{\lambda}_{k_1}^{(\alpha)} \tilde{\pi}_{k_1}^{(\alpha)} \dots \tilde{\lambda}_{k_{m-1}}^{(\alpha)} \tilde{\pi}_{k_m}^{(\alpha)} \tilde{\lambda}_{k_m}^{(\alpha)} \tilde{\pi}_{k_m}^{(\alpha)}} = \frac{q^{m(2\ell + m + \alpha + 1)}}{(q^{\ell + 1}; q)_m (q^{\ell + \alpha + 2}; q)_m}.$$
 (5.31)

Proof. We shall proceed by mathematical induction on m. Equation (5.31) for m = 1 coincides with (5.28). Suppose (5.31) is true for a given $m \in \mathbb{N}$. Then, in view of (5.27) and using the induction hypothesis,

$$\begin{split} & \sum_{0 \leq k_1 < k_2 < \ldots < k_m < k_{m+1}} (q^{k_1}; q^{-1})_{\ell} \frac{\tilde{T}_{k_1}^{(\alpha)} (\tilde{T}_{k_2}^{(\alpha)} - \tilde{T}_{k_1}^{(\alpha)}) (\tilde{T}_{k_3}^{(\alpha)} - \tilde{T}_{k_2}^{(\alpha)}) \ldots (\tilde{T}_{k_{m+1}}^{(\alpha)} - \tilde{T}_{k_m}^{(\alpha)})}{\tilde{\lambda}_{k_1}^{(\alpha)} \tilde{\pi}_{k_1}^{(\alpha)} \tilde{\lambda}_{k_2}^{(\alpha)} \tilde{\pi}_{k_2}^{(\alpha)} \ldots \tilde{\lambda}_{k_m}^{(\alpha)} \tilde{\pi}_{k_m}^{(\alpha)} \tilde{\lambda}_{k_{m+1}}^{(\alpha)} \tilde{\pi}_{k_{m+1}}^{(\alpha)}} \\ &= \frac{q^{2\ell + \alpha + 2}}{(1 - q^{\ell + 1}) (1 - q^{\ell + \alpha + 2})} \sum_{0 \leq k_2 < k_3 \ldots < k_m < k_{m+1}} (q^{k_2}; q^{-1})_{\ell + 1} \frac{\tilde{T}_{k_2}^{(\alpha)} (\tilde{T}_{k_3}^{(\alpha)} - \tilde{T}_{k_2}^{(\alpha)}) \ldots (\tilde{T}_{k_{m+1}}^{(\alpha)} - \tilde{T}_{k_m}^{(\alpha)})}{\tilde{\lambda}_{k_2}^{(\alpha)} \tilde{\pi}_{k_2}^{(\alpha)} \ldots \tilde{\lambda}_{k_m}^{(\alpha)} \tilde{\pi}_{k_m}^{(\alpha)} \tilde{\lambda}_{k_{m+1}}^{(\alpha)} \tilde{\pi}_{k_{m+1}}^{(\alpha)}} \\ &= \frac{q^{(m+1)} (2\ell + m + \alpha + 2)}{(q^{\ell + 1}; q)_{m+1} (q^{\ell + \alpha + 2}; q)_{m+1}}. \end{split}$$

This shows the induction step.

Another proof of formula (5.24). Letting $\ell = 0$ in (5.31) we obtain

$$\sum_{0 \le k_1 < k_2 < \dots < k_m} \frac{\tilde{T}_{k_1}^{(\alpha)} (\tilde{T}_{k_2}^{(\alpha)} - \tilde{T}_{k_1}^{(\alpha)}) \dots (\tilde{T}_{k_m}^{(\alpha)} - \tilde{T}_{k_{m-1}}^{(\alpha)})}{\tilde{\lambda}_{k_1}^{(\alpha)} \tilde{\pi}_{k_1}^{(\alpha)} \dots \tilde{\lambda}_{k_{m-1}}^{(\alpha)} \tilde{\pi}_{k_{m-1}}^{(\alpha)} \tilde{\lambda}_{k_m}^{(\alpha)} \tilde{\pi}_{k_m}^{(\alpha)}} = \frac{q^{m(m+\alpha+1)}}{(q;q)_m (q^{\alpha+2};q)_m}.$$

In view of (3.23), the characteristic function equals

$$\tilde{\mathfrak{F}}^{(\alpha)}(z) = 1 + \sum_{m=1}^{\infty} (-1)^m z^m \frac{q^{m(m+\alpha+1)}}{(q;q)_m (q^{\alpha+2};q)_m} = {}_{0}\phi_1 \begin{pmatrix} - \\ q^{\alpha+2} \end{pmatrix}; q, -q^{\alpha+2}z \end{pmatrix}.$$

REFERENCES

- [1] N.I. Akhiezer, The classical moment problem and some related questions in analysis, Hafner Publishing Co., New York, 1965.
- [2] B. Beckermann, Complex Jacobi matrices, vol. 127, 2001, pp. 17–65.
- [3] Ch. Berg, Markov's theorem revisited, J. Approx. Theory **78** (1994), no. 2, 260–275.
- [4] C.F. Bracciali, F. Marcellán, and S. Varma, *Orthogonality of quasi-orthogonal polynomials*, Filomat **32** (2018), no. 20, 6953–6977.
- [5] T.S. Chihara, On determinate Hamburger moment problems, Pacific J. Math. 27 (1968), 475–484.
- [6] ______, Convergent sequences of orthogonal polynomials, J. Math. Anal. Appl. 38 (1972), 335–347.
- [7] ______, An introduction to orthogonal polynomials, Mathematics and its Applications, Vol. 13, Gordon and Breach Science Publishers, New York-London-Paris, 1978.
- [8] ______, Indeterminate symmetric moment problems, J. Math. Anal. Appl. 85 (1982), no. 2, 331–346.
- [9] J.B. Conway, Functions of one complex variable, second ed., Graduate Texts in Mathematics, vol. 11, Springer-Verlag, New York-Berlin, 1978.
- [10] M.D. de la Iglesia, Orthogonal Polynomials in the Spectral Analysis of Markov Processes: Birth-Death Models and Diffusion, Encyclopedia of Mathematics and its Applications, Cambridge University Press, 2021.
- [11] G. Gasper and M. Rahman, *Basic hypergeometric series*, second ed., Encyclopedia of Mathematics and its Applications, vol. 96, Cambridge University Press, Cambridge, 2004.
- [12] J.S. Geronimo, On the spectra of infinite-dimensional Jacobi matrices, J. Approx. Theory 53 (1988), no. 3, 251–265.
- [13] J.S. Geronimo, E.M. Harrell II, and W. Van Assche, *On the asymptotic distribution of eigenvalues of banded matrices*, Constr. Approx. **4** (1988), no. 4, 403–417.
- [14] S. Karlin and J.L. McGregor, *The differential equations of birth-and-death processes, and the Stieltjes moment problem*, Trans. Amer. Math. Soc. **85** (1957), 489–546.
- [15] R. Koekoek, P.A. Lesky, and R.F. Swarttouw, *Hypergeometric orthogonal polynomials and their q-analogues*, Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2010.
- [16] P. Loya, Amazing and aesthetic aspects of analysis, Springer, New York, 2017.
- [17] M. Lukić, A first course in spectral theory, Graduate Studies in Mathematics, vol. 226, American Mathematical Society, Providence, RI, 2022.
- [18] P. Maroni, Sur la suite de polynômes orthogonaux associée à la forme $u = \delta_c + \lambda (x c)^{-1}L$, Period. Math. Hungar. **21** (1990), no. 3, 223–248.
- [19] D.G. Mead, Newton's identities, Amer. Math. Monthly 99 (1992), no. 8, 749–751.
- [20] D.S. Moak, The q-analogue of the Laguerre polynomials, J. Math. Anal. Appl. 81 (1981), no. 1, 20-47.
- [21] K. Schmüdgen, The moment problem, Graduate Texts in Mathematics, vol. 277, Springer, Cham, 2017.
- [22] J.A. Shohat and J.D. Tamarkin, *The Problem of Moments*, American Mathematical Society Mathematical Surveys, Vol. I, American Mathematical Society, New York, 1943.
- [23] G. Świderski, Spectral properties of unbounded Jacobi matrices with almost monotonic weights, Constr. Approx. 44 (2016), no. 1, 141–157.
- [24] W. Van Assche, *Orthogonal polynomials, associated polynomials and functions of the second kind*, J. Comput. Appl. Math. **37** (1991), no. 1-3, 237–249.
- [25] P. Šťovíček, On infinite Jacobi matrices with a trace class resolvent, J. Approx. Theory 249 (2020), 105306.
- [26] _____, On family of orthogonal polynomials corresponding to Jacobi matrices with a trace class inverse, J. Approx. Theory **279** (2022), 105769.

PAVEL ŠŤOVÍČEK, DEPARTMENT OF MATHEMATICS, FACULTY OF NUCLEAR SCIENCE, CZECH TECHNICAL UNIVERSITY IN PRAGUE, TROJANOVA 13, 120 00 PRAHA, CZECH REPUBLIC

Email address: stovicek@fjfi.cvut.cz

Grzegorz Świderski, Institute of Mathematics, Polish Academy of Sciences, ul. Śniadeckich 8, 00-696 Warsaw, Poland

Email address: grzegorz.swiderski@pwr.edu.pl