# Asymmetric GARCH modelling without moment conditions

Yuxin Tao

Department of Statistics and Data Science, Southern University of Science and Technology

and

Dong Li<sup>∗</sup>

Department of Statistics and Data Science, Tsinghua University

October 2, 2024

#### Abstract

There is a serious and long-standing restriction in the literature on heavy-tailed phenomena in that moment conditions, which are unrealistic, are almost always assumed in modelling such phenomena. Further, the issue of stability is often insufficiently addressed. To this end, we develop a comprehensive statistical inference for an asymmetric generalized autoregressive conditional heteroskedasticity model with standardized non-Gaussian symmetric stable innovation (sAGARCH) in a unified framework, covering both the stationary case and the explosive case. We consider first the maximum likelihood estimation of the model including the asymptotic properties of the estimator of the stable exponent parameter among others. We then propose a modified Kolmogorov-type test statistic for diagnostic checking, as well as those for strict stationarity and asymmetry testing. We conduct Monte Carlo simulation studies to examine the finite-sample performance of our entire statistical inference procedure. We include empirical examples of stock returns to highlight the usefulness and merits of our sAGARCH model.

Keywords: Heavy tails; Kolmogorov-type test; Maximum likelihood estimation; Nonstationarity; Stable distribution

<sup>∗</sup>Li's work is supported in part by the NSFC (no.72471127).

### 1 Introduction

#### 1.1 Motivation and related works

Heavy-tailed phenomena are ubiquitous in the real world and can be observed across a wide range of scientific fields. On such phenomena, there have been numerous research results, including, e.g., the recent monographs of [Resnick](#page-34-0) [\(2007\)](#page-34-0), [Harvey](#page-32-0) [\(2013\)](#page-32-0), [Buraczewski et](#page-30-0) [al.](#page-30-0) [\(2016\)](#page-30-0), [Ibragimov and Prokhorov](#page-32-1) [\(2017\)](#page-32-1), [Kulik and Soulier](#page-32-2) [\(2020\)](#page-32-2), [Nolan](#page-33-0) [\(2020\)](#page-33-0), [Taleb](#page-34-1) [\(2020\)](#page-34-1), [Nair et al.](#page-33-1) [\(2022\)](#page-33-1) and others. Nowadays, studies of heavy-tailed phenomena continue to generate significant interest in the fields of economics, finance, and statistics. Many heavy-tailed distributions, such as the Student's  $t$ , the Pareto, and the stable distributions, have been introduced and have a wide range of practical applications.

Financial return time series is one of the most typical examples among others. Generally, they exihibit some important stylized features, including heavy tails, volatility clustering, asymmetry, aggregational Gaussianity, quasi long range dependence [\(Rydberg,](#page-34-2) [2000\)](#page-34-2), etc. Since the seminal works of [Engle](#page-31-0) [\(1982\)](#page-31-0) and [Bollerslev](#page-30-1) [\(1986\)](#page-30-1), the generalized autoregressive conditional heteroskedasticity (GARCH) model has been used extensively in empirical work and is now regarded as the benchmark by many econometricians and financial practitioner [\(Fan and Yao,](#page-31-1) [2017\)](#page-31-1). Over the past four decades, numerous variants of GARCH models have been proposed to cater for various practical needs (see the monograph [Francq](#page-31-2) and Zakoïan  $(2019)$  for a comprehensive review).

Although theoretically GARCH models are capable of capturing several of the stylized features mentioned above, they do not always perform well as far as tail-thickness is concerned. Typically, there can still be excess kurtosis left in the standardized residuals of fitted models in practice; see, e.g., [Bai](#page-30-2) [\(2003\)](#page-30-2). Thus, the commonly imposed assumption of finite fourth or second moments for the innovations in GARCH-type models is found to be too stringent. To address this problem, the use of Student's  $t$  innovations, or generalized

Gaussian innovations have been proposed. However, these distributions still face several challenges. One issue is that they lack stability under addition. A distribution is considered stable under addition if the distribution of the sum of i.i.d. random variables is the same as that of the individual summands, up to location-scale transforms (see [Fama](#page-31-3) [\(1965\)](#page-31-3)), and we call it stable distribution. It also has an appealing property that it is the only possible limiting distributions of sums of i.i.d. random variables. This is required when the error terms are assumed to be the sum of all external effects that are not captured by the model. It is also of particular importance in portfolio and risk management, as it allows for the modelling of aggregate behavior, such as price changes of individual stocks in a portfolio. Even if the individual changes have different distributions, the cumulative change is approximately stable, thus enabling stable distribution to model the overall risk of a portfolio (see also e.g. [Fama](#page-31-3) [\(1965\)](#page-31-3), [Samuelson](#page-34-3) [\(1967\)](#page-34-3), [Calzolari et al.](#page-31-4) [\(2014\)](#page-31-4), and [Nolan](#page-33-2) [\(2014,](#page-33-2) [2020\)](#page-33-0)). Moreover, stable distribution allows for infinite mean and asymmetry, which is another property that  $t$  or generalized Gaussian distribution does not share.

To take on the challenges, we suggest the use of stable distribution<sup>[1](#page-2-0)</sup> for innovations. In fact, the statistical use of stable distribution in finance can be dated back to [Mandelbrot](#page-33-3) [\(1963\)](#page-33-3) and [Fama](#page-31-3) [\(1965\)](#page-31-3), who suggested the stable Paretian distribution as a model to fit the unconditional asset returns. However, applications of stable distribution were hampered due to the lack of closed-form density function and limited computing power. With the development of powerful modern computers and numerical techniques, it is now feasible to use stable models efficiently in practice. An occasional argument against using non-normal stable models is that they have infinite variance or even infinite mean, which may seem contradictary to the finite sample moments observed in applications. However, [Mandelbrot](#page-33-4)

<span id="page-2-0"></span><sup>&</sup>lt;sup>1</sup>The stable distribution was first studied by Lévy [\(1925\)](#page-32-3), and then developed by [Gnedenko and Kol](#page-32-4)[mogorov](#page-32-4) [\(1954\)](#page-32-4), [Fama and Roll](#page-31-5) [\(1968\)](#page-31-5), etc. Monographs of stable distributions include [Zolotarev](#page-34-4) [\(1986\)](#page-34-4), [Samorodnitsky and Taqqu](#page-34-5) [\(1994\)](#page-34-5), and [Nolan](#page-33-0) [\(2020\)](#page-33-0).

[\(1997\)](#page-33-4) (p.88-92) pointed out that the sample moments do not always converge as the sample size increases, so the finiteness of sample moments does not necessarily imply the finiteness of the underlying distribution's moment. In fact, empirical studies suggest that power-law models with divergent second or even first moments are ubiquitous in finance<sup>[2](#page-3-0)</sup>, with tail indices less than 2 or 1. The infinite-mean models sometimes fit the observed reality statistically better, and they can provide useful insights on risk management problems (see, e.g., [Silverberg and Verspagen](#page-34-6) [\(2007\)](#page-34-6), [Clauset et al.](#page-31-6) [\(2009\)](#page-31-6), and [Chen and Wang](#page-31-7) [\(2024\)](#page-31-7)).

In the context of GARCH models, the use of stable innovations was first proposed by [McCulloch](#page-33-5) [\(1985\)](#page-33-5) and studied by [Liu and Brorsen](#page-33-6) [\(1995\)](#page-33-6) within a more general context, both focusing on applications. [Panorska et al.](#page-34-7) [\(1995\)](#page-34-7) and [Mittnik et al.](#page-33-7) [\(2002\)](#page-33-7) studied the stationarity conditions of GARCH models with stable innovations. With respect to statistical estimation, [Liu and Brorsen](#page-33-6) [\(1995\)](#page-33-6) proposed the maximum likelihood estimation approach. [Calzolari et al.](#page-31-4) [\(2014\)](#page-31-4) discussed the indirect inference method by using Student's t innovations as auxiliary models. As for heavy-tailed GARCH-type models, [Hall and](#page-32-5) [Yao](#page-32-5) [\(2003\)](#page-32-5) discussed GARCH models with heavy-tailed innovations, assuming innovations with infinite fourth moments but unit variance. However, to the best of our knowledge, the asymptotic properties of the maximum likelihood estimators (MLE) of parameters of GARCH-type models with infinite-moment innovations are absent in the literature.

#### 1.2 Stable distribution and our model

Generally, stable distributions are defined by their characteristic functions since their densities have no closed forms except for three special cases (i.e., Gaussian, Cauchy, and Lévy distributions). Generally, a stable distribution  $\mathbf{S}(\alpha,\beta,\gamma,\delta)$  contains four parameters:

<span id="page-3-0"></span><sup>2</sup>Applications of stable laws can be found in a variety of fields, including finance, telecommunications and physics. See, e.g. [Adler et al.](#page-30-3) [\(1998\)](#page-30-3), [Uchaikin and Zolotarev](#page-34-8) [\(1999\)](#page-34-8), [Rachev and Mittnik](#page-34-9) [\(2000\)](#page-34-9), [Ibragimov et al.](#page-32-6) [\(2015\)](#page-32-6), [Peng and Qi](#page-33-8) [\(2017\)](#page-33-8), [Nolan](#page-33-0) [\(2020\)](#page-33-0), and [Taleb](#page-34-1) [\(2020\)](#page-34-1).

 $\alpha, \beta, \gamma$ , and  $\delta$ , representing stability exponent, skewness, scale, and location, respectively. In this paper, we consider the standardized<sup>[3](#page-4-0)</sup> symmetric stable distribution  $\mathbf{S}(\alpha,0,1,0)$ , of which the characteristic function is  $\phi(s) = \exp(-|s|^{\alpha})$ ,  $s \in \mathbb{R}$ , where  $\alpha \in (0, 2]$ . Its density is

<span id="page-4-1"></span>
$$
f_{\alpha}(x) = \frac{1}{\pi} \int_0^{\infty} \exp(-s^{\alpha}) \cos(sx) ds, \quad x \in \mathbb{R}.
$$
 (1.1)

It has no variance when  $\alpha < 2$  and no expectation when  $\alpha \leq 1$ . Particularly, when  $\alpha = 1$ , it is the standard Cauchy distribution, and when  $\alpha = 2$ , it reduces to  $\mathcal{N}(0, 2)$ . More properties on stable distribution can be found in Propositions 2.3-2.5 in the Appendix C.

Conditional asymmetry or leverage effects are also highly relevant in financial applications, in the sense that negative returns tend to have stronger impact on future volatilities than positive returns of the same magnitude. The original GARCH model fails to capture this feature, which motivates a variety of extensions including Threshold GARCH, Asymmetric GARCH and Exponential GARCH. In view of stable distributions, a natural way to describe leverage effects is the direct use of a nonzero skewness parameter  $\beta$ . However, the density of a skew stable distribution is much more complicated than that for the symmetric case, which will entail massive computation in numerical optimization. On the other hand, as  $\alpha$  approaches 2, generally all stable distributions will tend to be symmetric and  $\beta$  will become meaningless and harder to estimate accurately; see [Nolan](#page-33-0)  $(2020)(p.12)$  $(2020)(p.12)$ . Note that  $β$  is insignificant when  $α = 2$  under Gaussian laws. To avoid such problem, we propose to introduce asymmetry to the model structure rather than to the innovations.

To better capture excess kurtosis, asymmetry and volatility clustering jointly, and to conduct inference within a unified framework of both the stationary and explosive cases, we propose the first-order asymmetric GARCH model with standardized non-Gaussian

<span id="page-4-0"></span><sup>&</sup>lt;sup>3</sup>The term "standardized" stems from [Nolan](#page-33-0) [\(2020\)](#page-33-0), meaning the scale  $\gamma = 1$  and the location  $\delta = 0$ . Throughout the paper, we adopt the 0-parametrization of  $\mathbf{S}(\alpha,\beta,\gamma,\delta)$ , which has the simplest form of the characteristic function that is continuous in all parameters, as recommended in [Nolan](#page-33-0) [\(2020\)](#page-33-0) (p.5).

symmetric stable innovation (hereafter sAGARCH(1,1)) defined as

<span id="page-5-0"></span>
$$
\begin{cases}\ny_t = \sigma_t \eta_t, & t \in \mathbb{Z}_+ := \{1, 2, \ldots\}, \\
\sigma_t^2 = \omega + \phi_+ (y_{t-1}^+) ^2 + \phi_- (y_{t-1}^-)^2 + \psi \sigma_{t-1}^2,\n\end{cases} \tag{1.2}
$$

with initial values  $y_0$  and  $\sigma_0 \geq 0$ , where  $\omega > 0$ ,  $\phi_+ \geq 0$ ,  $\phi_- \geq 0$ ,  $\psi \geq 0$ ,  $x^+ = \max\{x, 0\}$ ,  $x^- = -\min\{x, 0\}$ .  $\{\eta_t\}$  is a sequence of i.i.d. standardized non-Gaussian symmetric stable random variables with the stable exponent  $\alpha \in (0, 2)$ . Heavy-tailedness is modeled through stable innovation  $\{\eta_t\}$ , and asymmetry is captured by different parameters  $\phi_+$  and  $\phi_-$ .

Following [Bougerol and Picard](#page-30-4) [\(1992a](#page-30-4)[,b\)](#page-30-5), model [\(1.2\)](#page-5-0) is strictly stationary if and only if the top Lyapunov exponent

<span id="page-5-1"></span>
$$
\gamma_{\alpha} := E \log a(\eta) < 0
$$
, where  $a(x) = \phi_+(x^+)^2 + \phi_-(x^-)^2 + \psi$ , (1.3)

and  $\eta$  is a generic random variable with the same distribution as  $\eta_t$  and independent of  $\{\eta_t\}$ . Particularly, when  $\alpha = 1$ ,  $\eta$  follows the standard Cauchy distribution and  $\gamma_1 =$  $\log[(\sqrt{\phi_+} +$  $\sqrt{\psi}$ )( $\sqrt{\phi_-}$ + √  $[\overline{\psi}]$ . Fig. [1\(](#page-6-0)a) plots the densities  $f_{\alpha}(x)$  for different values of  $\alpha$ , and Fig. [1\(](#page-6-0)b)-(d) plots the surface of  $\gamma_{\alpha} = 0$  in [\(1.3\)](#page-5-1) in terms of  $(\phi_+, \phi_-, \psi)$  within a finite region  $(0, 1)^3$ . The strict stationarity region of model  $(1.2)$  is the closed one encompassed by the surface of  $\gamma_{\alpha} = 0$  and the plains  $\phi_{+} = 0$ ,  $\phi_{-} = 0$ , and  $\psi = 0$ . From the subfigures, we can see that: (i) the larger  $\alpha$  is, the bigger the strictly stationary region becomes; (ii) in the strictly stationary region,  $\psi$  is strictly no more than 1, while  $\phi_+$  or  $\phi_-$  could be greater than 1, provided that the other two parameters are small enough.

#### <span id="page-5-2"></span>1.3 Contributions and outline

The first contribution of this work is the proposal of an asymmetric GARCH model with stable innovations, which allows for no moment conditions. The model is capable of effectively addressing the excess kurtosis commonly encountered in financial time series, in addition to other key features such as volatility clustering and leverage effects. We study



<span id="page-6-0"></span>Figure 1: (a) The densities  $f_{\alpha}(x)$ ; (b)-(d) The surfaces determined by  $\{(\phi_+, \phi_-, \psi) : \gamma_{\alpha} = 0\}$ in [\(1.3\)](#page-5-1) for different fixed values of  $\alpha = 1.5, 1$ , and 0.5. The vertical axis represents  $\psi$ , and the other two represent  $\phi_+$  and  $\phi_-\$ . In (d), the black curve represents the boundary of the surface in the  $\psi = 0$  plane. Here the surfaces are plotted within a finite region  $(0, 1)^3$  for the sake of clarity, while the whole surface can stretch very far.

the MLE of model parameters in  $(1.2)$  and investigate their asymptotic properties within a unified framework, that encompasses both the stationary and explosive cases, thus filling a theoretical gap in the statistical inference for GARCH models within heavy-tail settings. It is shown that the MLE is strongly consistent and asymptotically normal with a convergence rate of  $n^{1/2}$  (except for  $\omega$  in the explosive case). This result differs from the findings of

[Hall and Yao](#page-32-5) [\(2003\)](#page-32-5), who showed that for heavy-tailed errors with infinite fourth moment but unit variance, the asymptotic distributions of the quasi-MLE in GARCH models follow multivariate stable laws, which are typically *non-normal*. Although model  $(1.2)$  specifies the distribution of innovations, it can accommodate cases with much heavier tails, and the asymptotic properties of MLE are more standard and easier to obtain for statistical inference. We provide two estimation approaches for the Fisher information matrix in statistical inference. A universal estimator of the asymptotic variance of MLE is also proposed.

We would like to emphasize the technical contributions of our paper. Although the model setup  $(1.2)$  is a variant of those in Francq and Zakoïan  $(2012, 2013)$  $(2012, 2013)$  $(2012, 2013)$ , the key difference is that the innovation in model  $(1.2)$  has no moment condition, while the latter ones require a unit variance assumption in the innovation terms. Thus, to obtain the asymptotic properties of MLE in model  $(1.2)$ , we need new techniques to get rid of moment restrictions. The main technical contributions are summarized as follows: (i) The asymptotic behavior of partial derivatives of log-stable densities, as summarized in Proposition 2.5 in the Supplementary Material. This provides more precise results compared to the existing ones and will be useful in studying the inference of statistical models with stable innovations; (ii) The identifiability of the stability parameter  $\alpha$  in view of the properties of stable distribution; (iii) A substantial improvement in an exponential convergence rate for the process  $\sup_{\theta \in \Theta_0^*} |\sigma_t^2(\theta)/\sigma_t^2(\theta_0) - v_t(\theta)|$ . This is crucial for controlling the infinite moments of stable innovations.

Second, although theoretically the MLE of the intercept  $\omega$  is consistent and asymptotically normal for the stationary case, its finite-sample performance is not satisfactory, especially with heavy-tailed innovations. This feature has been well recognized [\(Francq](#page-31-8) and Zakoïan, [2012\)](#page-31-8), although no plausible explanation has been given so far. Now, we give an intuitive explanation as follows. Our model reveals that this phenomenon is probably related to the extreme values of the observations. An intuitive reason is that the intercept term  $\omega$  can be viewed as a scale parameter of  $y_t$ , as model  $(1.2)$  is equivalent to

$$
y_t/\sqrt{\omega} = (\sigma_t/\sqrt{\omega})\eta_t, \quad (\sigma_t/\sqrt{\omega})^2 = 1 + \phi_+(y_{t-1}^+/\sqrt{\omega})^2 + \phi_-(y_{t-1}^-/\sqrt{\omega})^2 + \psi(\sigma_{t-1}/\sqrt{\omega})^2.
$$

This is also an sAGARCH model for the scaled series with intercept 1 and other parameters remaining the same. As  $\alpha$  decreases, the tail of innovation becomes heavier, leading to extreme values of  $y_t$ . So the estimator of scale parameter  $\omega$  is likely to be affected by the scale of  $y_t$  and be overestimated in some cases. Moreover, random number generators sometimes work poorly for heavy-tailed distributions, as outliers are often generated and may violate the theoretical distribution. These reasons all account for the poor performance of the estimator of  $\omega$ . Thus, through our model with heavy-tailed innovations, we have provided a deeper understanding of the finite-sample performance of the intercept estimator.

Third, we develop a powerful modified Kolmogorov-type test statistic for model diagnostic checking within a unified framework, using the transformation method inspired by [Bai](#page-30-2) [\(2003\)](#page-30-2). We also provide statistics for testing strict stationarity and asymmetry of the model. Last, Monte Carlo simulation studies are conducted to confirm our theoretical findings and assess the finite-sample performance of the MLE and test statistics. Empirical examples of stock returns are analyzed and results are compared with existing ones, so as to illustrate the efficacy of the sAGARCH model. Considering the complex nature of stable densities, we also provide efficient algorithms that leverage advanced integration techniques for the practical implementation and estimation of our model.

The rest of the article is organized as follows. Section [2](#page-9-0) studies the MLE of sAGARCH model and establishes its asymptotics, together with a universal estimator of its asymptotic variance. Section [3](#page-13-0) considers the estimation of Fisher information matrices and computational issues. Section [4](#page-15-0) develops a Kolmogorov-type test statistic for diagnostic checking, as well as tests for strict stationarity and asymmetry of the model. Section [5](#page-19-0) reports Monte Carlo simulation studies on the performance of the MLE and test statistics. Section [6](#page-24-0) analyzes real stock return series. Section [7](#page-28-0) concludes the paper. All technical proofs are postponed to the Supplementary Material.

### <span id="page-9-0"></span>2 Maximum Likelihood Estimation

Let  $\theta = (\omega, \vartheta')' = (\omega, \phi_+, \phi_-, \psi, \alpha)' \in \mathbb{R}_+^4 \times (0, 2)$ , where  $\mathbb{R}_+ = (0, \infty)$ . Suppose that the observations  $\{y_0, y_1, ..., y_n\}$  are from model [\(1.2\)](#page-5-0) with true parameter  $\theta_0 = (\omega_0, \vartheta'_0)' =$  $(\omega_0, \phi_{0+}, \phi_{0-}, \psi_0, \alpha_0)'$ . Then the (conditional) log-likelihood function is defined as

$$
\widetilde{L}_n(\theta) = \sum_{t=1}^n \widetilde{\ell}_t(\theta), \quad \widetilde{\ell}_t(\theta) = -\log \widetilde{\sigma}_t(\theta) + \log f_\alpha\left(\frac{y_t}{\widetilde{\sigma}_t(\theta)}\right),\tag{2.1}
$$

where  $f_{\alpha}(x)$  is defined in [\(1.1\)](#page-4-1), and

$$
\widetilde{\sigma}_t^2(\theta) = \omega + \phi_+(y_{t-1}^+)^2 + \phi_-(y_{t-1}^-)^2 + \psi \widetilde{\sigma}_{t-1}^2(\theta), \quad t \ge 1,
$$
\n(2.2)

with initial values  $(y_0, \tilde{\sigma}_0(\theta)) \equiv (0, 0)$ . The MLE of  $\theta_0$  is

<span id="page-9-1"></span>
$$
\widehat{\theta}_n := \arg \max_{\theta \in \Theta} \widetilde{L}_n(\theta),
$$

where the parameter space  $\Theta$  is a subset of  $\mathbb{R}^4_+ \times (0,2)$ .

To study the asymptotic properties of  $\widehat{\theta}_n$  for the stationary case, define the theoretical (conditional) log-likelihood function as follows:

$$
L_n(\theta) = \sum_{t=1}^n \ell_t(\theta), \quad \ell_t(\theta) = -\log \sigma_t(\theta) + \log f_\alpha\left(\frac{y_t}{\sigma_t(\theta)}\right),
$$

where

$$
\sigma_t^2(\theta) = \omega + \phi_+(y_{t-1}^+)^2 + \phi_-(y_{t-1}^-)^2 + \psi \sigma_{t-1}^2(\theta), \quad t \in \mathbb{Z}.
$$
 (2.3)

Thus, when  $0 \leq \psi < 1$ , we have

$$
\sigma_t^2(\theta) = \sum_{j=0}^{\infty} \psi^j \{ \omega + \phi_+(y_{t-1-j}^+)^2 + \phi_-(y_{t-1-j}^-)^2 \}, \quad t \in \mathbb{Z}.
$$
 (2.4)

If  $\{y_t : t \in \mathbb{Z}\}\)$  are from model  $(1.2)$  with true parameter  $\theta_0$ , then  $\{y_t\}$  is strictly stationary and ergodic. Since  $\sigma_t^2(\theta)$  is a measurable function of  $\{y_t\}$ ,  $\{\sigma_t^2(\theta) : t \in \mathbb{Z}\}$  is also an ergodic strictly stationary sequence.

#### 2.1 Consistency and asymptotic normality

To study the asymptotics of  $\widehat{\theta}_n$ , the following assumptions are needed. These assumptions are standard in the literature on volatility models.

<span id="page-10-0"></span>**Assumption 1**  $\{\eta_t\}$  is a sequence of i.i.d. standardized non-Gaussian symmetric stable random variables with the density function  $f_{\alpha_0}(x)$ .

<span id="page-10-1"></span>**Assumption 2** The parameter space  $\Theta$  is compact and  $\theta_0 \in \Theta$ .

<span id="page-10-2"></span>**Assumption 3** The true parameter  $\theta_0$  is an interior point of  $\Theta$ .

Now, we are ready to state our main results as follows.

<span id="page-10-3"></span>**Theorem 2.[1](#page-10-0)** Suppose that Assumptions  $1-\frac{2}{3}$  hold.

- (i). If  $\gamma_{\alpha_0} < 0$ , for  $\Theta$  such that  $\forall \theta \in \Theta$ ,  $\psi < 1$ , then  $\widehat{\theta}_n \rightarrow_{a.s.} \theta_0$  as  $n \rightarrow \infty$ .
- (ii). If  $\gamma_{\alpha_0} > 0$ , then  $\widehat{\vartheta}_n \to_{a.s.} \vartheta_0$  as  $n \to \infty$ .

<span id="page-10-4"></span>**Theorem 2.2** Suppose that Assumptions  $1-\frac{3}{5}$  $1-\frac{3}{5}$  hold.

(i). If  $\gamma_{\alpha_0} < 0$ , for  $\Theta$  such that  $\forall \theta \in \Theta$ ,  $\psi < 1$ , then  $\sqrt{n}(\widehat{\theta}_n - \theta_0) \rightarrow_d \mathcal{N}(0, \Sigma^{-1})$  as  $n \to \infty$ , where  $\hookrightarrow_d$  stands for convergence in distribution, and

$$
\Sigma = E\left\{\frac{\partial \ell_t(\theta_0)}{\partial \theta} \frac{\partial \ell_t(\theta_0)}{\partial \theta'}\right\} = \left(\frac{\Sigma_{\tilde{\theta}\tilde{\theta}'} \left| \Sigma'_{\tilde{\theta}\alpha} \right|}{\Sigma_{\tilde{\theta}\alpha} \left| \Sigma_{\alpha\alpha} \right|}\right),
$$

with  $\tilde{\theta} = (\omega, \phi_+, \phi_-, \psi)'$  and

$$
\Sigma_{\tilde{\theta}\tilde{\theta}'} = \frac{1}{4} E_{\infty} \Big\{ \frac{1}{\sigma_t^4(\theta_0)} \frac{\partial \sigma_t^2(\theta_0)}{\partial \tilde{\theta}} \frac{\partial \sigma_t^2(\theta_0)}{\partial \tilde{\theta}'} \Big\} E \Big\{ 1 + \frac{\partial \log f_{\alpha_0}(\eta_t)}{\partial x} \eta_t \Big\}^2,
$$
\n
$$
\Sigma_{\tilde{\theta}\alpha} = -\frac{1}{2} E_{\infty} \Big\{ \frac{1}{\sigma_t^2(\theta_0)} \frac{\partial \sigma_t^2(\theta_0)}{\partial \tilde{\theta}} \Big\} E \Big\{ \frac{\partial \log f_{\alpha_0}(\eta_t)}{\partial x} \frac{\partial \log f_{\alpha_0}(\eta_t)}{\partial \alpha} \eta_t \Big\}, \ \Sigma_{\alpha\alpha} = E \Big\{ \frac{\partial \log f_{\alpha_0}(\eta_t)}{\partial \alpha} \Big\}^2.
$$
\n(ii). If  $\gamma_{\alpha_0} > 0$ , then  $\sqrt{n} (\hat{\theta}_n - \theta_0)' \to_d \mathcal{N}(0, \Upsilon^{-1})$  as  $n \to \infty$ , where  $\Upsilon = \begin{pmatrix} \Upsilon_{\tilde{\theta}\tilde{\theta}'} & \Upsilon_{\tilde{\theta}\alpha} \\ \Upsilon_{\tilde{\theta}\alpha} & \Upsilon_{\alpha\alpha} \end{pmatrix}$ 

with  $\tilde{\vartheta} = (\phi_+, \phi_-, \psi)'$  and

$$
\begin{split}\n\Upsilon_{\widetilde{\vartheta}\widetilde{\vartheta}'} &= \frac{1}{4} E(d_t d_t') E\Big\{ 1 + \frac{\partial \log f_{\alpha_0}(\eta_t)}{\partial x} \eta_t \Big\}^2, \\
\Upsilon_{\widetilde{\vartheta}\alpha} &= -\frac{1}{2} E(d_t) E\Big\{ \frac{\partial \log f_{\alpha_0}(\eta_t)}{\partial x} \frac{\partial \log f_{\alpha_0}(\eta_t)}{\partial \alpha} \eta_t \Big\}, \qquad \Upsilon_{\alpha\alpha} = E\Big\{ \frac{\partial \log f_{\alpha_0}(\eta_t)}{\partial \alpha} \Big\}^2.\n\end{split}
$$

The explicit forms of  $E(d_t d'_t)$  and  $E(d_t)$  are provided in (D.2)-(D.3) in the Appendix D of the Supplementary Material.

**Remark 1** It is interesting to note that, when  $\alpha_0 = 1$ , i.e.,  $\eta$  follows the standard Cauchy distribution, the following values (also given in [Li, et.al.](#page-32-7)  $(2023)$ ) are available:

$$
E\left\{\frac{\partial \log f_{\alpha_0}(\eta)}{\partial x}\right\}^2 = E\left\{1 + \frac{\partial \log f_{\alpha_0}(\eta)}{\partial x}\eta\right\}^2 = \frac{1}{2},
$$
  
\n
$$
E\left\{\frac{\partial \log f_{\alpha_0}(\eta)}{\partial x}\frac{\partial \log f_{\alpha_0}(\eta)}{\partial \alpha}\eta\right\} = \frac{\mathcal{C} - 1 + \log 2}{2},
$$
  
\n
$$
E\left\{\frac{\partial \log f_{\alpha_0}(\eta)}{\partial \alpha}\right\}^2 = \frac{(\mathcal{C} - 1 + \log 2)^2}{2} + \frac{\pi^2}{12},
$$

where  $C = 0.577215664 \cdots$  is the Euler-Mascheroni constant. Based on these, the exact value of Fisher matrix  $\Upsilon$  can be calculated, which is given in the Appendix D.

**Remark 2** Theorems [2.1\(](#page-10-3)ii) and [2.2\(](#page-10-4)ii) show that the intercept  $\omega_0$  is not estimable for the explosive case, due to the non-identifiability of  $\omega$  in the limit of  $L_n(\theta)/n$ . Such an unidentifiable phenomenon of partial parameters in nonstationary time series analysis was first observed by [Jensen and Rahbek](#page-32-8)  $(2004a,b)$  $(2004a,b)$  who studied the QMLE of a nonstationary  $ARCH(1)$  model, and then by Francq and Zakoïan [\(2012,](#page-31-8) [2013\)](#page-31-9) who studied nonstationary GARCH models.

**Remark 3** Note that Theorems [2.1](#page-10-3)[-2.2](#page-10-4) exclude the critical case  $\gamma_{\alpha_0} = 0$ . As far as we know, for this critical case, the behavior of  $y_t$  remains essentially open in the literature. For example, although Francq and Zakoïan [\(2012\)](#page-31-8) claimed to have resolved this issue, their additional Assumption A is difficult to check within the setting of stable innovations. Our

extensive simulations suggest that their assumption does not hold. We can certainly impose more straightforward assumptions on  $y_t$  in order to obtain consistency and asymptotic normality of  $\hat{\vartheta}_n$ , such as  $|y_t|/\rho^t \to_{a.s.} \infty$ ,  $\rho > 1$ , as  $t \to \infty$ , while such assumptions are normally unverifiable in practice. Thus, we leave it as future research.

# <span id="page-12-1"></span>2.2 A universal estimator of the asymptotic variance of  $\widehat{\vartheta}_n$

This subsection discusses a universal form of the asymptotic variance of  $\widehat{\vartheta}_n$  without any stationarity condition.

From Theorem [2.2\(](#page-10-4)i), in the stationary case  $\gamma_{\alpha_0} > 0$ , the asymptotic distribution of  $\widehat{\vartheta}_n$ is  $\sqrt{n}(\widehat{\vartheta}_n - \vartheta_0) \rightarrow_d \mathcal{N}(0, \Upsilon^{-1}_*)$ , where

$$
\Upsilon_* = \Sigma_{\vartheta\vartheta} - \Sigma_{\vartheta\omega} \Sigma_{\omega\omega}^{-1} \Sigma_{\vartheta\omega}' = \left(\frac{\Sigma_{\widetilde{\vartheta}\widetilde{\vartheta}} \left| \Sigma_{\widetilde{\vartheta}\alpha} \right|}{\Sigma_{\widetilde{\vartheta}\alpha} \left| \Sigma_{\alpha\alpha} \right|} \right) - \Sigma_{\omega\omega}^{-1} \left(\Sigma_{\widetilde{\vartheta}\omega} \left| \Sigma_{\alpha\omega} \right| \Sigma_{\alpha\omega}\right).
$$

Define their estimators of six block submatrices contained in  $\Upsilon_*$  as

<span id="page-12-0"></span>
$$
\begin{split}\n\widehat{\Sigma}_{\widetilde{\partial}\widetilde{\theta}} &= \frac{1}{4} \Big\{ \frac{1}{n} \sum_{t=1}^{n} \frac{1}{\widetilde{\sigma}_{t}^{4}(\widehat{\theta}_{n})} \frac{\partial \widetilde{\sigma}_{t}^{2}(\widehat{\theta}_{n})}{\partial \widetilde{\theta}} \frac{\partial \widetilde{\sigma}_{t}^{2}(\widehat{\theta}_{n})}{\partial \widetilde{\theta}^{y}} \Big\} \frac{1}{n} \sum_{t=1}^{n} \Big\{ 1 + \frac{\partial \log f_{\widehat{\alpha}_{n}}(\widehat{\eta}_{t})}{\partial x} \widehat{\eta}_{t} \Big\}^{2}, \\
\widehat{\Sigma}_{\widetilde{\theta}\alpha} &= -\frac{1}{2} \Big\{ \frac{1}{n} \sum_{t=1}^{n} \frac{1}{\widetilde{\sigma}_{t}^{2}(\widehat{\theta}_{n})} \frac{\partial \widetilde{\sigma}_{t}^{2}(\widehat{\theta}_{n})}{\partial \widetilde{\theta}} \Big\} \Big\{ \frac{1}{n} \sum_{t=1}^{n} \frac{\partial \log f_{\widehat{\alpha}_{n}}(\widehat{\eta}_{t})}{\partial x} \frac{\partial \log f_{\widehat{\alpha}_{n}}(\widehat{\eta}_{t})}{\partial \alpha} \widehat{\eta}_{t} \Big\}, \\
\widehat{\Sigma}_{\alpha\alpha} &= \frac{1}{n} \sum_{t=1}^{n} \Big\{ \frac{\partial \log f_{\widehat{\alpha}_{n}}(\widehat{\eta}_{t})}{\partial \alpha} \Big\}^{2}, \\
\widehat{\Sigma}_{\omega\omega} &= \frac{1}{4} \Big\{ \frac{1}{n} \sum_{t=1}^{n} \frac{1}{\widetilde{\sigma}_{t}^{4}(\widehat{\theta}_{n})} \frac{\partial \widetilde{\sigma}_{t}^{2}(\widehat{\theta}_{n})}{\partial \omega} \frac{\partial \widetilde{\sigma}_{t}^{2}(\widehat{\theta}_{n})}{\partial \omega} \Big\} \Big\{ \frac{1}{n} \sum_{t=1}^{n} \Big( 1 + \frac{\partial \log f_{\widehat{\alpha}_{n}}(\widehat{\eta}_{t})}{\partial x} \widehat{\eta}_{t} \Big)^{2} \Big\}, \\
\widehat{\Sigma}_{\widetilde{\theta}\omega} &= \frac{1}{4} \Big\{ \frac{1}{n} \sum_{t=1}^{n
$$

where the residual  $\hat{\eta}_t$  is calculated by  $\hat{\eta}_t = y_t/\tilde{\sigma}_t(\hat{\theta}_n)$ . Let  $\hat{\Upsilon}_* = \hat{\Sigma}_{\vartheta\vartheta} - \hat{\Sigma}_{\vartheta\omega}\hat{\Sigma}_{\omega\omega}^{-1}\hat{\Sigma}_{\vartheta\omega}'$ . It can be shown that  $\hat{\Upsilon}_*$  is a consistent estimator of  $\Upsilon_*$  in the stationary case. The following theorem shows that such an estimator also provides a consistent estimator of the asymptotic variance of  $\widehat{\vartheta}_n$  in the explosive case  $\gamma_{\alpha_0} > 0$ .

<span id="page-13-1"></span>**Theorem 2.3** Suppose that Assumptions  $1-\frac{2}{3}$  $1-\frac{2}{3}$  hold.

- (i). If  $\gamma_{\alpha_0} < 0$ , then  $\hat{\Upsilon}_* \rightarrow_{a.s.} \Upsilon_*$ , as  $n \rightarrow \infty$ .
- (ii). If  $\gamma_{\alpha_0} > 0$ , then  $\hat{\Upsilon}_* \to_{a.s.} \Upsilon$ , as  $n \to \infty$ .

In each case,  $\hat{\Upsilon}_*^{-1}$  is a strongly consistent estimator of the asymptotic variance of  $\hat{\theta}_n$ . Thus, we can conduct asymptotically valid inference for  $\vartheta_0$  within this unified framework of both stationary and explosive cases, without prior stationarity test. The asymmetry test in Corollary [4.1](#page-17-0) hereinafter is an immediate application of this theorem.

### <span id="page-13-0"></span>3 Estimation of  $\Sigma$  and  $\Upsilon$

For statistical inference on  $\theta_0$  in practice, we need to estimate the Fisher informaiton matrix  $\Sigma$  or  $\Upsilon$  in Theorem [2.2.](#page-10-4) First, notice that each entry of  $\Sigma$  contains two factors. Taking the entry  $\Sigma_{\psi\psi}$  as an example, we can write  $\Sigma_{\psi\psi} = \Sigma_{\sigma} \Sigma_{\eta}$ , where

$$
\Sigma_{\sigma} = \frac{1}{4} E \Big\{ \frac{1}{\sigma_t^4(\theta_0)} \frac{\partial \sigma_t^2(\theta_0)}{\partial \psi} \frac{\partial \sigma_t^2(\theta_0)}{\partial \psi} \Big\} \quad \text{and} \quad \Sigma_{\eta} = E \Big\{ 1 + \frac{\partial \log f_{\alpha_0}(\eta)}{\partial x} \eta \Big\}^2.
$$

With observations  $\{y_0, y_1, ..., y_n\}$ , similar to  $(2.5)$ , it is not hard to estimate  $\Sigma_{\sigma}$  by

$$
\widehat{\Sigma}_{\sigma} = \frac{1}{4n} \sum_{t=1}^{n} \frac{1}{\widetilde{\sigma}_{t}^{4}(\widehat{\theta}_{n})} \frac{\partial \widetilde{\sigma}_{t}^{2}(\widehat{\theta}_{n})}{\partial \psi} \frac{\partial \widetilde{\sigma}_{t}^{2}(\widehat{\theta}_{n})}{\partial \psi}.
$$

As for the factor  $\Sigma_{\eta}$ , here we provide two estimation approaches. One is based on the integral expression of  $\Sigma_{\eta}$ , namely

$$
\widehat{\Sigma}_{\eta}^{\text{int}} = \int_{\mathbb{R}} \left\{ 1 + \frac{\partial \log f_{\widehat{\alpha}_n}(u)}{\partial x} u \right\}^2 f_{\widehat{\alpha}_n}(u) du.
$$

The other is based on the residuals  $\{\hat{\eta}_t\}$ , which is the same as that in [\(2.5\)](#page-12-0), namely

$$
\widehat{\Sigma}_{\eta}^{\text{res}} = \frac{1}{n} \sum_{t=1}^{n} \left\{ 1 + \frac{\partial \log f_{\widehat{\alpha}_n}(\widehat{\eta}_t)}{\partial x} \widehat{\eta}_t \right\}^2.
$$

Combining  $\hat{\Sigma}_{\sigma}$  with these two estimators of  $\Sigma_{\eta}$ , we can thus obtain two estimators of  $\Sigma_{\psi\psi}$ , say,  $\hat{\Sigma}_{\psi\psi}^{\text{int}}$  and  $\hat{\Sigma}_{\psi\psi}^{\text{res}}$ . It can be shown that they are consistent in probability by the mean value theorem, Lebesgue's dominated convergence theorem, and Theorems [2.1-](#page-10-3)[2.2.](#page-10-4) The other entries can be dealt with analogously, and we obtain two estimators  $\hat{\Sigma}^{\text{int}}$  and  $\hat{\Sigma}^{\text{res}}$ .

The estimation of  $\Upsilon$  in the explosive case can be dealt with similarly. Note that in  $\Upsilon$ ,  $E(d_t d'_t)$  and  $E(d_t)$  only involves the innovation  $\eta$ . Take  $E(d_t^{\psi})$  $(t_t^{\psi})^2$  as an example.  $E(d_t^{\psi})$  $\binom{w}{t}^2 =$  $\nu_2(1+\nu_1)/\{\psi_0^2(1-\nu_2)(1-\nu_1)\}\,$ , where

$$
\nu_i = E\{\psi_0/a_0(\eta_t)\}^i, \quad \text{with } a_0(x) = \phi_{0+}(x^+)^2 + \phi_{0-}(x^-)^2 + \psi_0, \quad i = 1, 2. \tag{3.1}
$$

Similar to the preceding approach, for  $\nu_i$ , we can construct two estimators of  $\nu_i$  as

<span id="page-14-2"></span>
$$
\widehat{\nu}_{i}^{\text{int}} = \int_{\mathbb{R}} \left\{ \frac{\widehat{\psi}_{n}}{\widehat{\phi}_{n+}(u^{+})^{2} + \widehat{\phi}_{n-}(u^{-})^{2} + \widehat{\psi}_{n}} \right\}^{i} f_{\widehat{\alpha}_{n}}(u) du,
$$
\n
$$
\widehat{\nu}_{i}^{\text{res}} = \frac{1}{n} \sum_{t=1}^{n} \left\{ \frac{\widehat{\psi}_{n}}{\widehat{\phi}_{n+}(\widehat{\eta}_{t}^{+})^{2} + \widehat{\phi}_{n-}(\widehat{\eta}_{t}^{-})^{2} + \widehat{\psi}_{n}} \right\}^{i}, \quad i = 1, 2.
$$

The rest part of  $\Upsilon_{\psi\psi}$  can be dealt with similarly, and we derive two estimators of  $\Upsilon_{\psi\psi}$ , say,  $\hat{\Upsilon}_{\psi\psi}^{\text{int}}$  and  $\hat{\Upsilon}_{\psi\psi}^{\text{res}}$ , and in turn get two estimators  $\hat{\Upsilon}^{\text{int}}$  and  $\hat{\Upsilon}^{\text{res}}$  of  $\Upsilon$ , respectively.

Finally, we provide two estimators for the Lyapunov exponent  $\gamma_{\alpha_0}$  in [\(1.3\)](#page-5-1):

<span id="page-14-1"></span>
$$
\widehat{\gamma}_n^{\text{int}} = \int_{\mathbb{R}} \log \left[ \widehat{\phi}_{n+}(u^+)^2 + \widehat{\phi}_{n-}(u^-)^2 + \widehat{\psi}_n \right] f_{\widehat{\alpha}_n}(u) du,
$$
\n
$$
\widehat{\gamma}_n^{\text{res}} = \frac{1}{n} \sum_{t=1}^n \log \left[ \widehat{\phi}_{n+}(\widehat{\eta}_t^+)^2 + \widehat{\phi}_{n-}(\widehat{\eta}_t^-)^2 + \widehat{\psi}_n \right].
$$
\n(3.2)

As for computation, since the density  $f_{\alpha}(x)$  involves improper integrals of oscillating functions and has no explicit formula, numerical integration techniques are necessary. Some transformation techniques (see, e.g., [Nolan](#page-33-9) [\(1997\)](#page-33-9), Chapter 4 in [Uchaikin and Zolotarev](#page-34-8) [\(1999\)](#page-34-8), and [Matsui and Takemura](#page-33-10) [\(2006\)](#page-33-10)) are recommended for all numerical calculations that involve stable density  $f_{\alpha}(x)^{4}$  $f_{\alpha}(x)^{4}$  $f_{\alpha}(x)^{4}$ .

Numerical studies in Section [5](#page-19-0) show that two types of estimators (i.e.,  $\hat{\Sigma}^{\text{int}}$  and  $\hat{\Sigma}^{\text{res}}, \hat{\Upsilon}^{\text{int}}$ and  $\hat{\Upsilon}^{\text{res}}$ ) have good performance and each has its own pros and cons. We here provide an empirical guidance for the choices of estimators: the estimator based on the integration (i.e.

<span id="page-14-0"></span><sup>&</sup>lt;sup>4</sup>The R code for calculation of stable densities and implementation of MLE is available upon request.

 $\widehat{\Sigma}^{\text{int}}$  or  $\widehat{\Upsilon}^{\text{int}}$ ) is recommended if  $\widehat{\alpha}_n \in (0, 1]$ , while the estimator based on the residuals (i.e.  $\hat{\Sigma}^{\text{res}}$  or  $\hat{\Upsilon}^{\text{res}}$ ) is recommended if  $\hat{\alpha}_n \in (1, 2)$  in practice. As an estimator of  $\gamma_{\alpha_0}$ , the empirical mean of  $\hat{\gamma}_n^{\text{res}}$  is closer to the true value than that of  $\hat{\gamma}_n^{\text{int}}$ , although the empirical standard deviation of  $\hat{\gamma}_n^{\text{res}}$  is slightly larger. From the theoretical view, obtaining the asymptotics of  $\hat{\gamma}_n^{\text{res}}$  is relatively easier than that of  $\hat{\gamma}_n^{\text{int}}$ . Thus we recommend the estimator  $\hat{\gamma}_n^{\text{res}}$ .

### <span id="page-15-0"></span>4 Testing

In this section we discuss three types of hypothesis tests, which are stationarity testing, asymmetry testing, and a Kolmogorov-type test for model diagnostic checking.

#### 4.1 Strict stationarity testing

Consider the following two test problems:

(i) For the strict stationarity testing,

<span id="page-15-2"></span>
$$
H_0: \gamma_{\alpha_0} < 0 \quad \text{vs} \quad H_1: \gamma_{\alpha_0} \ge 0 \tag{4.1}
$$

and (ii) for the explosivity testing,

<span id="page-15-3"></span>
$$
H_0: \gamma_{\alpha_0} > 0 \quad \text{vs} \quad H_1: \gamma_{\alpha_0} \le 0. \tag{4.2}
$$

Let  $\gamma_n(\theta) = \frac{1}{n} \sum_{t=1}^n \log[\phi_+\{q_t^+(\theta)\}^2 + \phi_-\{q_t^-(\theta)\}^2 + \psi], \ \theta \in \Theta$ , where  $q_t(\theta) = y_t/\tilde{\sigma}_t(\theta)$ and  $\tilde{\sigma}_t(\theta)$  is defined in [\(2.2\)](#page-9-1). Then  $\hat{\gamma}_n := \gamma_n(\hat{\theta}_n)$  is an estimator of  $\gamma_0$ , which is the estimator  $\hat{\gamma}_n^{\text{res}}$  defined in [\(3.2\)](#page-14-1). We first give the asymptotic distribution of  $\hat{\gamma}_n$ .

<span id="page-15-1"></span>**Theorem 4.1** Let  $u_t = \log a_0(\eta_t) - \gamma_{\alpha_0}$  and  $\sigma_u^2 = Eu_t^2 < \infty$ , where  $a_0(x)$  is defined in  $(3.1)$ . Then, under the assumptions of Theorem [2.2,](#page-10-4) it follows that

$$
\sqrt{n}(\widehat{\gamma}_n - \gamma_0) \to_d \mathcal{N}(0, \sigma_\gamma^2) \quad \text{as } n \to \infty,
$$

where

$$
\sigma_{\gamma}^{2} = \begin{cases}\n\sigma_{u}^{2} + \{a_{1}'\Sigma^{-1}a_{2} - 4(1 - \nu_{1})^{2}/c_{1}\}, & \text{if } \gamma_{\alpha_{0}} < 0, \\
\sigma_{u}^{2}, & \text{if } \gamma_{\alpha_{0}} > 0,\n\end{cases}
$$
\n
$$
a_{1} = (0, -\tilde{\nu}_{1+}, -\tilde{\nu}_{1-}, -\nu_{1}/\psi_{0}, 2c_{2}(1 - \nu_{1} + c^{*})/c_{1} - 2\tilde{c}',
$$
\n
$$
a_{2} = (0, -\tilde{\nu}_{1+}, -\tilde{\nu}_{1-}, -\nu_{1}/\psi_{0}, 2c_{2}(1 - \nu_{1})/c_{1}),
$$
\n
$$
c^{*} = E\left[u_{t}\left(1 + \frac{\partial \log f_{\alpha_{0}}(\eta_{t})}{\partial x}\eta_{t}\right)\right], \quad \tilde{c} = E\left(u_{t}\frac{\partial \log f_{\alpha_{0}}(\eta_{t})}{\partial \alpha}\right),
$$
\n
$$
c_{1} = E\left(1 + \frac{\partial \log f_{\alpha_{0}}(\eta_{t})}{\partial x}\eta_{t}\right)^{2}, \quad c_{2} = E\left(\frac{\partial \log f_{\alpha_{0}}(\eta_{t})}{\partial x}\frac{\partial \log f_{\alpha_{0}}(\eta_{t})}{\partial \alpha}\eta_{t}\right),
$$
\n
$$
\tilde{\nu}_{1+} = E\left\{\frac{(\eta_{1}^{+})^{2}}{a_{0}(\eta_{1})}\right\}, \quad \tilde{\nu}_{1-} = E\left\{\frac{(\eta_{1}^{-})^{2}}{a_{0}(\eta_{1})}\right\}, \quad \nu_{1} = E\left\{\frac{\psi_{0}}{a_{0}(\eta_{1})}\right\}.
$$

**Remark 4** Compared with Theorem 4.1 in Francq and Zakoïan [\(2013\)](#page-31-9), the difference is that the asymptotic variance of  $\hat{\gamma}_n$  has a much complicated form. It is due to different innovation assumptions, so the asymptotic behavior of  $\widehat{\gamma}_n$  involves the estimation effect of stability parameter  $\alpha$ . Also,  $\sigma_u^2 < \infty$  can be guaranteed by  $E(|\eta_t|^{\alpha_0/2}) < \infty$ .

Denote

$$
\widehat{\sigma}_u^2 = \frac{1}{n} \sum_{t=1}^n \left\{ \log \left[ \widehat{\phi}_{n+}(\widehat{\eta}_t^+)^2 + \widehat{\phi}_{n-}(\widehat{\eta}_t^-)^2 + \widehat{\psi}_n \right] \right\}^2 - \left\{ \frac{1}{n} \sum_{t=1}^n \log \left[ \widehat{\phi}_{n+}(\widehat{\eta}_t^+)^2 + \widehat{\phi}_{n-}(\widehat{\eta}_t^-)^2 + \widehat{\psi}_n \right] \right\}^2.
$$

Under the assumptions of Theorem [4.1,](#page-15-1) it is not hard to show that  $\hat{\sigma}_u^2 \to_p \sigma_u^2$  whenever  $\gamma_{\alpha_0}$  < 0 or  $\gamma_{\alpha_0}$  > 0. Similar to Francq and Zakoïan [\(2013\)](#page-31-9), we construct the test statistic

$$
\mathrm{T}_n:=\sqrt{n}\widehat{\gamma}_n/\widehat{\sigma}_u
$$

Thus, for the testing problem  $(4.1)$  [resp.,  $(4.2)$ ], the test defined by the stationary [resp., explosive] critical region

$$
C^{ST} = \{T_n > \Phi^{-1}(1-\underline{\alpha})\} \quad \text{[resp.,} \quad C^{ET} = \{T_n < \Phi^{-1}(\underline{\alpha})\}\text{]},
$$

has its asymptotic significance level bounded by  $\underline{\alpha} \in (0,1)$  and is consistent.

#### 4.2 Asymmetry testing

It is particularly interesting to test the existence of a leverage effect in financial asset returns. Benefit from the framework of our sAGARCH model [\(1.2\)](#page-5-0), the asymmetry testing problem has a simple form of

<span id="page-17-1"></span>
$$
H_0: \phi_{0+} = \phi_{0-} \quad \text{vs} \quad H_1: \phi_{0+} \neq \phi_{0-}.
$$
 (4.3)

Similar to Francq and Zakoïan  $(2013)$ , we consider the following test statistic for symmetry

$$
\mathcal{T}_n^{\mathcal{S}} = \frac{\sqrt{n}(\widehat{\phi}_{n+} - \widehat{\phi}_{n-})}{\widehat{\sigma}_s}, \quad \text{with} \quad \widehat{\sigma}_s = \sqrt{\mathbf{e}^{\prime} \widehat{\mathcal{T}}_*^{-1} \mathbf{e}} \quad \text{and} \quad \mathbf{e} = (1, -1, 0, 0)^{\prime}.
$$

Note that the symmetry test does not require any stationarity assumption, as we take advantage of the universal estimator  $\hat{\Upsilon}_*$ . By Theorem [2.3,](#page-13-1) the following corollary holds.

<span id="page-17-0"></span>Corollary 4.1 Under the assumptions of Theorem [2.2,](#page-10-4) for the symmetry testing problem  $(4.3)$ , the test defined by the critical region

$$
C^{S} = \left\{ |T_{n}^{S}| > \Phi^{-1}(1 - \underline{\alpha}/2) \right\}
$$

has the asymptotic significance level  $\underline{\alpha} \in (0,1)$  and is consistent.

#### 4.3 Diagnostic checking

Diagnostic checking is important for time series modeling. However, the most commonly used portmanteau test based on the autocorrelation of the residuals or the squared residuals does not work for assessing the adequacy of model  $(1.2)$ , since the innovation is specified to follow a certain stable distribution. Here, we construct a Kolmogorov-type test for diagnostic checking. Consider the null hypothesis as follows

<span id="page-17-2"></span>
$$
H_0: \eta_t \sim f_{\alpha_*}(x) \quad \text{(i.e., } \alpha_0 = \alpha_*\text{)},\tag{4.4}
$$

where  $\alpha_* \in (0, 2)$  is a fixed constant. In practice,  $\alpha_*$  can be chosen to be  $\hat{\alpha}_n$  or an approximate value of  $\widehat{\alpha}_n$ .

Let  $F_{\alpha}(x)$  be the cumulative distribution functions of the standardized symmetric stable random variable. Define

$$
U_t = F_{\alpha_*}(\eta_t)
$$
 and  $V_n(r) = \frac{1}{\sqrt{n}} \sum_{t=1}^n [I(U_t \le r) - r]$ , for  $r \in [0, 1]$ .

Clearly, if  $H_0$  in [\(4.4\)](#page-17-2) holds, then  $\{U_t\}$  are i.i.d.  $U[0, 1]$  random variables, and

<span id="page-18-0"></span>
$$
\sup_{0 \le r \le 1} |V_n(r)| \to_d \sup_{0 \le r \le 1} |\mathbb{B}(r) - r \mathbb{B}(1)| \quad \text{as } n \to \infty,
$$
\n(4.5)

where  $\mathbb{B}(\cdot)$  is a standard Brownian motion, by the Donsker Theorem and continuous mapping theorem. Since  $\eta_t$  is not observable in practice, we replace it by the residual  $\tilde{\eta}_t$ , where  $\widetilde{\eta}_t = y_t/\sigma_t(\widetilde{\theta}_n), t = 1, ..., n$ , and  $\widetilde{\theta}_n = (\widehat{\omega}_n, \widehat{\phi}_{n+}, \widehat{\phi}_{n-}, \widehat{\psi}_n)'$  is the restricted MLE of  $\widetilde{\theta}_0$  under  $H_0$ . Accordingly, we replace  $U_t$  and  $V_n(r)$  by  $\widehat{U}_t$  and  $\widehat{V}_n(r)$  respectively, where

$$
\widehat{U}_t = F_{\alpha_*}(\widetilde{\eta}_t)
$$
 and  $\widehat{V}_n(r) = \frac{1}{\sqrt{n}} \sum_{t=1}^n [I(\widehat{U}_t \le r) - r],$  for  $r \in [0, 1].$ 

Unfortunately as opposed to  $(4.5)$ ,  $\sup_{0 \le r \le 1} |V_n(r)|$  is no longer asymptotically distributionfree, since it involves the effect of parameter estimation. To obtain an asymptotically distribution-free test statistic, we adopt the martingale transformation of [Khmaladze](#page-32-10) [\(1981\)](#page-32-10) inspired by [Bai](#page-30-2) [\(2003\)](#page-30-2). Specifically, let  $g(r) = (g_1(r), g_2(r))' = (r, f_{\alpha_*}(F_{\alpha_*}^{-1}(r))F_{\alpha_*}^{-1}(r))'$ with its derivative  $\dot{g}(r) = (1, \dot{g}_2(r))'$ , where

$$
\dot{g}_2(r) = 1 + \frac{\dot{f}_{\alpha_*}(F_{\alpha_*}^{-1}(r))}{f_{\alpha_*}(F_{\alpha_*}^{-1}(r))} F_{\alpha_*}^{-1}(r)
$$
 and  $\dot{f}_{\alpha_*}(x) = \frac{\partial f_{\alpha_*}(x)}{\partial x}$ .

Define Khmaladze's transformation

$$
\widehat{W}_n(r) = \widehat{V}_n(r) - \int_0^r \left( \dot{g}(s)' C^{-1}(s) \int_s^1 \dot{g}(u) d\widehat{V}_n(u) \right) ds,
$$

where  $C(r) = \int_r^1 \dot{g}(s)\dot{g}(s)'ds$ . Then, the Kolmogorov-type test statistic is defined as

$$
\mathcal{T}_n^{\mathcal{D}} = \sup_{0 \le r \le 1} |\widehat{W}_n(r)|.
$$

**Theorem 4.2** Suppose model [\(1.2\)](#page-5-0) is well specified with  $\alpha_0 = \alpha_*$  and the conditions in Theorem [2.2](#page-10-4) hold. Then for both  $\gamma_{\alpha_*}$  < 0 and  $\gamma_{\alpha_*}$  > 0 cases, it follows that

$$
T_n^D \to_d \sup_{0 \le r \le 1} |\mathbb{B}(r)| \quad \text{as } n \to \infty.
$$

Thus, for the testing problem [\(4.4\)](#page-17-2), the critical region is defined as

$$
C^D = \left\{T^D_n > cv^D_{\underline{\alpha}}\right\}
$$

at the asymptotic significance level  $\underline{\alpha} \in (0,1)$ , where  $cv_{\underline{\alpha}}^D$  is the  $\underline{\alpha}$  upper percentile of the limiting distribution of  $T_n^D$ . The critical values at the significance levels of 10%, 5%, and 1% are 1.9600, 2.2414, and 2.8070, respectively, via simulation. Numerical simulations in Section [5](#page-19-0) show that  $T_n^D$  has a satisfactory power in all cases, even for small samples.

**Remark 5** In application,  $T_n^D$  can be approximately computed with

$$
\max_{1 \le j \le n} \sqrt{n} \left| \frac{j}{n} - \frac{1}{n} \sum_{k=1}^{j} \dot{g}(v_k)' C_k^{-1} D_k(v_k - v_{k-1}) \right|,
$$

where  $D_k = \sum_{i=k}^n \dot{g}(v_i)$  and  $C_k = \sum_{i=k}^n \dot{g}(v_i) \dot{g}(v_i)'(v_{i+1} - v_i)$ , and  $v_1, ..., v_n$  are ordered values of  $\widehat{U}_1, ..., \widehat{U}_n$  with the convention  $v_0 = 0$  and  $v_{n+1} = 1$ .

### <span id="page-19-0"></span>5 Simulation studies

#### <span id="page-19-1"></span>5.1 Performance of the MLE

To assess the finite-sample performance of the MLE of  $\theta_0$  in the sAGARCH model [\(1.2\)](#page-5-0) in the stationary and explosive cases, we choose the number of observations  $n = 200, 500$ , 1000, each with 1000 replications, with three different values of  $\alpha_0$  being 1.5, 1.0, and 0.5.

First, for the stationary case, true parameters are set to be  $\theta_0 = (\omega_0, \phi_{0+}, \phi_{0-}, \psi_0, \alpha_0)' =$  $(0.2, 0.1, 0.2, 0.5, 1.5)$ <sup>'</sup>,  $(0.1, 0.1, 0.2, 0.3, 1)$ <sup>'</sup>, and  $(0.05, 0.02, 0.05, 0.1, 0.5)$ <sup>'</sup>, respectively. The true top Lyapunov exponents  $\gamma_{\alpha_0}$  are calculated and summarized in Table A.1.



Table 1: Simulation results for the MLE  $\hat{\theta}$  for sAGABCH(1.1) under stationary cases  $\theta_n$  for sAGARCH(1,1) under stationary cases.

Table [5.1](#page-19-1) summarizes the empirical biases (Bias), empirical standard deviations (ESD), asymptotic standard deviations (ASD) of the MLE  $\widehat{\theta}_n$ , together with two estimators  $\widehat{\text{ASD}}^{\text{int}}$ and  $\widehat{ASD}^{\text{res}}$  of ASDs from  $\widehat{\Sigma}^{\text{int}}$  and  $\widehat{\Sigma}^{\text{res}}$  in Section [3.](#page-13-0) The ASDs are calculated from the asymptotic covariance matrix in Theorem  $2.2$  (i). From Table  $5.1$ , we can find that (i) for the MLE  $\widehat{\vartheta}_n$ , the biases are small in each case and the values of the ESD and ASD are close to each other, especially for large n; (ii) The two estimators of the ASD of  $\{\hat{\phi}_{n+}, \hat{\phi}_{n-}, \hat{\psi}_n\}$ perform similarly well; for  $\widehat{\alpha}_n$ ,  $\widehat{ASD}^{int}$  tends to outperform  $\widehat{ASD}^{res}$  when  $\alpha_0 < 1$ , while  $\widehat{ASD}^{\text{res}}$  outperforms  $\widehat{ASD}^{\text{int}}$  when  $\alpha_0 \in [1,2)$  on the whole; their differences diminish as n increases; (iii) The MLE of the intercept  $\omega$  performs relatively poorly, especially for small  $\alpha$  and small n; the ESDs are large compared with the corresponding ASDs, and the estimated ASD is quite large. Its performance becomes better as  $n$  increases to a certain extent. This is a very interesting phenomenon, which has been recognized in the existing literature but has not been studied heretofore. As discussed in Section [1.3,](#page-5-2) we find that  $\omega$  can be interpreted as the scale parameter of the model, so its estimation is likely to be affected by the scale of observations and be overestimated sometimes in finite samples. Thus, our model provides a deeper understanding of the finite-sample performance of the intercept estimator. Fig. [2](#page-22-0) plots the histograms of  $\sqrt{n}(\hat{\theta}_n - \theta_0)$  with  $n = 1000$  and  $\theta_0 = (0.2, 0.1, 0.2, 0.5, 1.5)'$  $\theta_0 = (0.2, 0.1, 0.2, 0.5, 1.5)'$  $\theta_0 = (0.2, 0.1, 0.2, 0.5, 1.5)'$ . From Fig. 2, we can see that the overall performance of  $\hat{\theta}_n$  is good and conforms to the asymptotic normal distribution.

Due to space limitations, the simulation results for the performance of MLE in the explosive cases and the estimation of the Lyapunov exponents are presented in the Appendix A.1. The finite-sample performance also validates the theoretical findings.

#### 5.2 Performance of the test statistics

In this subsection, we examine the finite-sample performance of the Kolmogorov-type test statistic  $T_n^D$  in Section [4.](#page-15-0) The simulation results for the stationarity and asymmetry tests



<span id="page-22-0"></span>Figure 2: The histograms of  $\sqrt{n}(\hat{\theta}_n - \theta_0)$  with  $n = 1000$  and  $\theta_0 = (0.2, 0.1, 0.2, 0.5, 1.5)'$ .

are presented in Appendix A.2 of the Supplementary Material due to the space limit.

For the stationary cases, two scenarios are considered:  $(I)$  sAGARCH $(1,1)$  model with  $\theta_0 = (0.2, 0.1, 0.2, 0.5, 1.5)'$ . The null hypothesis is that  $H_0^D : \eta_t \sim f_{1.5}$ , which is based on the empirical example in Section [6,](#page-24-0) while the alternative one  $H_1: \eta_t \sim f_{\alpha_*}$  with  $\alpha_* \neq 1.5$ . (II) sAGARCH(1,1) model with  $\theta_0 = (0.1, 0.1, 0.2, 0.3, 1)'$ . The null hypothesis  $H_0: \eta_t \sim f_1$ (i.e.,  $\eta_t \sim$  standard Cauchy), with alternative  $\eta_t \sim$  Student's  $t_{\nu}$ -distribution with degree of freedom  $\nu \in [0.5, 5]$ . For both two scenarios, the model is strictly stationary and ergodic under  $H_0$ . Fig. [3](#page-23-0) plots the size and the power of  $T_n^D$  at the significance level 5%. Here, *n* is 500, 1000, and 2000 for (I), and 200, 500,1000 for (II), each with 1000 replications. Fig. [3](#page-23-0) demonstrates that the size of  $T_n^D$  is close to the nominal level (except for  $n = 500$  under scenario (I)), and its power increases as the stable exponent  $\alpha$  or the degree of freedom  $\nu$ deviates from the null hypothesis. The size and power improve with increasing  $n$ .

Further, we examine the finite-sample performance of  $T_n^D$  for explosive cases. Similarly, we consider two testing scenarios for sAGARCH $(1,1)$  model with  $\theta_0 = (0.1, 0.1, 0.2, 0.5, 1)'$ : (I) The null hypothesis is  $H_0$ :  $\eta_t \sim f_1$ , while the alternative  $H_1 : \eta_t \sim f_{\alpha_*}$  with  $\alpha_* \neq 1$ . (II) The null hypothesis is  $H_0$ :  $\eta_t \sim f_1$ , while the alternative  $H_1$ :  $\eta_t \sim t_{\nu}$ -distribution



<span id="page-23-0"></span>Figure 3: Finite-sample performance of diagnostic test statistic  $T_n^D$  in the stationary cases. (a) The alternative is  $\eta_t \sim f_\alpha(x)$  with  $\alpha \in [1,2]$ . The size corresponds to  $\alpha = 1.5$ . (b) The alternative is  $\eta_t \sim$  Student's  $t_{\nu}$ -distribution with  $\nu \in [0.5, 5]$ . The size corresponds to  $\nu = 1$ (i.e.,  $\eta_t \sim f_1$ ). The horizontal dotted line at the bottom denotes the 5% significance level.

with  $\nu \in [0.5, 5]$ . Fig. [4](#page-24-1) plots the size and the power of  $T_n^D$  at the significance level 5%, when  $n = 500$ , 1000, 2000 for (I), and  $n = 200$ , 500, 1000 for (II), each with 1000 replications. From Fig. [4,](#page-24-1) we can find that for explosive cases  $T_n^D$  has a more satisfactory size and power as the stable exponent  $\alpha$  or the degree of freedom  $\nu$  deviates from the null hypothesis, compared with the stationary cases. Particularly, in scenario (II), the test statistics can effectively capture the differences between the true  $t_{\nu}$  innovation and the misspecified stable innovation. These facts indicate the usefulness of the modified Kolmogorov-type test statistic  $T_n^D$  for diagnostic checking and the identification of model misspecification even in finite-sample settings.



<span id="page-24-1"></span>Figure 4: The size and power of  $T_n^D$  for explosive cases. (a)  $H_1: \eta_t \sim f_\alpha(x)$ , with  $\alpha \in$ [0.5, 1.5]; (b)  $H_1: \eta_t \sim t_{\nu}$ -distribution, with  $\nu \in [0.5, 5]$ . The horizontal dotted line denotes the 5% significance level.

### <span id="page-24-0"></span>6 Empirical examples

To showcase the merits of model [\(1.2\)](#page-5-0), we analyze four individual stock return series, which are the same as those in Francq and Zakoïan  $(2012, 2013)$  $(2012, 2013)$  $(2012, 2013)$  for ease of comparison. They are the daily series of Icagen (NasdaqGM: ICGN, May 31, 2007–Feb.7, 2011), Monarch Community Bancorp (NasdaqCM: MCBF, Aug. 28, 2007 – Feb. 7, 2011), KV Pharmaceutical (NYSE: KV-A, Mar. 31, 2006 – Feb. 7, 2011), and China MediaExpress (NasdaqGS: CCME, Mar. 31,  $2009 - \text{Feb. } 7$ ,  $2011$ <sup>[5](#page-26-0)</sup>. Fig. 5 plots the four stock return series.

We fit the  $\text{sAGARCH}(1,1)$  model  $(1.2)$  to the data and the results are summarized in Table [2,](#page-25-0) together with the explosivity test statistics  $T_n$ , the asymmetry test statistics  $T_n^S$ , and the diagnostic checking statistics  $T_n^D$ . The estimated ASDs in parentheses are calculated by the universal estimator  $\hat{\Upsilon}_*$  in Section [2.2.](#page-12-1) The estimation results of an asymmetric  $GARCH(1,1)$  model by the QMLE in Francq and Zakoïan  $(2013)$  are also reported, as well

<span id="page-24-2"></span> $5$ The result of the Community Bankers Trust (AMEX: BTC) is similar to that in Francq and Zakoïan  $(2013)$ , so we omit it here.

<span id="page-25-0"></span>Table 2: The upper rows display the model-fitting results of  $sAGARCH(1,1)$  model, explosivity test statistics  $T_n$  with p-value, asymmetry test statistics  $T_n^S$  with p-value, diagnostic checking statistics  $T_n^D$ , the log-likelihood and the value of AIC for four stock returns, respectively. The lower four rows display the previous results of  $AGARCH(1,1)$  model in [Francq](#page-31-9) and Zakoïan [\(2013\)](#page-31-9), i.e., the p-value of the nonstationarity test "p-value-NS", p-value of the asymmetry test "p-value-S", log-likelihood and AIC, for comparison. The differences in the testing results for the two models are marked in bold.

Model		<b>ICGN</b>	<b>MCBF</b>	KV-A	CCME
sAGARCH (1,1)	$\boldsymbol{n}$	928	868	1221	488
	$\widehat{\omega}_{n+}$	2.688	0.044	0.070	0.002
	$\widehat{\phi}_{n+}$	0.098(0.026)	0.022(0.008)	0.033(0.008)	0.084(0.022)
	$\widehat{\phi}_{n-}$	0.164(0.035)	0.029(0.008)	0.039(0.008)	0.090(0.027)
	$\widehat{\psi}_{n}$	0.419(0.056)	0.884(0.022)	0.835(0.018)	0.766(0.034)
	$\widehat{\alpha}_n$	1.556(0.057)	1.369(0.065)	1.587(0.045)	1.527(0.084)
	$\mathrm{T}_n$	$-14.974$	$-1.585$	$-4.614$	$-0.569$
	p-value	0.000	$\bf 0.056$	0.000	0.285
	$T_n^S$	$-1.609$	$-0.625$	$-0.632$	$-0.190$
	p-value	0.108	0.532	0.527	0.849
	$T_n^{\rm D}$	2.697	4.837	1.043	2.276
	$log-lik$	$-3000.7$	$-2488.6$	$-3134.1$	$-1087.7$
	<b>AIC</b>	6011.4	4987.2	6278.1	2185.4
	p-value-NS	0.008	0.515	0.708	0.611
<b>AGARCH</b>	p-value-S	0.037	0.850	0.052	0.503
(1,1)	$log-lik$	$-3211.2$	$-2633.8$	$-3693.8$	$-1132.1$
	<b>AIC</b>	6430.5	5275.5	7395.7	2272.1

as the values of the log-likelihood and of the AIC. The significant differences in the testing results for these two models are highlighted in bold.

Table [2](#page-25-0) reveals some interesting new findings:

(i) We can see that all the estimated parameters are significant. The  $\text{sAGARCH}(1,1)$ model has a better fit than the  $AGARCH(1,1)$  one, as evidenced by the values of the AIC.

(ii) The stationarity test results of both the ICGN and the CCME accord with those in Francq and Zakoïan  $(2013)$ , while for the KV-A, the null hypothesis of explosivity is rejected at significance level 5%, and the MCBF is tested to be stationary at the 10% level. It reveals that the series which appears to be explosive under the framework of GARCH model with normal innovations turns out to be stationary in the heavy-tail setting.

(iii) For the stock ICGN, the null hypothesis of symmetry cannot be rejected at the 5% level using our model, while it is tested to be asymmetric using the  $AGARCH(1,1)$  model in Francq and Zakoïan [\(2013\)](#page-31-9). It shows that the existence of leverage effect may depend on the heavy-tail assumption of the model. The seemingly asymmetric volatility of the series is probably caused by a few extreme innovation terms instead of the leverage effect. As a matter of fact, after removing one obvious outlier around September 22, 2010 in ICGN,



<span id="page-26-0"></span>Figure 5: The graphs of four stock return series (%): ICGN, MCBF, KV-A, and CCME.

the series is tested to be symmetric even using  $AGARCH(1,1)$  model.

(iv) The values of  $T_n^D$  for the ICGN, KV-A, and CCME are smaller than critical values of  $\sup_{0 \leq r \leq 1} |\mathbb{B}(r)|$  at the significance levels of either 1% or 5%, while for the MCBF, the values of  $T_n^D$  are relatively large. The latter phenomenon is probably due to the seeming presence of change points in the time series of the MCBF, as seen from Fig [5,](#page-26-0) which also contribute to the nonstationarity of the series. The volatility increases sharply around the middle of the year 2008, which may be related to the financial crisis at that time. Thus, testing the existence of change points is an interesting topic for future studies.

We further examine the model-fitting performance of  $\text{AGARCH}(1,1)$  by examining residuals in Fig. [6,](#page-28-1) and simulated paths in Fig. [7.](#page-28-2) Fig. [6](#page-28-1) plots the cumulative distribution function (CDF) of stable distribution with estimated stability exponent  $\hat{\alpha}_n$  and the empirical CDF (ECDF) of the residual. It is shown that for ICGN, KV-A and CCME, the empirical CDF generally coincides with the true one, which illustrates the goodness-of-fit of sAGARCH(1,1) model on these datasets. As for MCBF, there exists certain difference between two CDFs, e.g., for values around 0. It accords with the relatively large test statistics  $T_n^D = 4.837$ , and implies that the model could be improved to better fit this dataset, for example by taking the change points into consideration. Nevertheless, our  $\text{sAGARCH}(1,1)$ model outperforms the original  $AGARCH(1,1)$  model in Francq and Zakoïan [\(2013\)](#page-31-9).

We also analyze several popular portfolio returns and compare the performance of model [\(1.2\)](#page-5-0) and AGARCH model with t-innovation. The results illustrate the merits of our model for modelling aggregate behavior and characterizing risk of portfolios. See Appendix A.3 in the Supplementary Material for more details.



<span id="page-28-1"></span>Figure 6: The ECDFs of residuals of four stock return series.



<span id="page-28-2"></span>Figure 7: Simulated paths (in red) of stock return series(%), with their true values in black.

# <span id="page-28-0"></span>7 Concluding remarks

In this paper, we have succeeded in delivering asymmetric GARCH modelling without moment conditions. The success is achieved by proposing a first-order asymmetric GARCH

(or sAGARCH) model with standardized non-Gaussian symmetric stable innovation. The sAGARCH model is shown to be effective in addressing particularly excess kurtosis, as well as other important stylized features observed in financial return series, namely volatility clustering and leverage effects. We have also developed a comprehensive statistical inference theory for this model within a unified framework that covers both the stationary and the explosive cases, thus filling the significant gap in the statistical inference for heavy-tailed GARCH-type models. Further, via extensive Monte Carlo simulations, we have found an intriguing phenomenon that the estimate of the intercept of the heavy-tailed GARCH-type model may exhibit unsatisfactory finite-sample performance, even for a large sample size, in the stationary cases. Such a phenomenon brings with it potential risks when forecasting financial returns. So far, this point has been overlooked in the existing literature.

To conclude, let us mention several possible extensions of our work. First, it would be interesting to extend our first-order sAGARCH model to a higher-order one. Like [Chan](#page-30-6) [and Ng](#page-30-6) [\(2009\)](#page-30-6), the corresponding statistical inference theory can be developed although many complicated technical difficulties would be involved in the proof. Second, practical issues on inference in a general  $\text{sAGARCH}(p, q)$  model include over-parametrization and over-identification. When the true parameters are on the boundary of the parameter space, statistical inference will become nonstandard and much complicated. Third, change points may exist in the real return series, so change point detection or anomaly detection becomes relevant for  $sAGARCH(p, q)$  model. Last but not least, the asymptotical behavior of sAGARCH(1,1) model for the critical case  $\gamma_{\alpha_0} = 0$  should be further investigated. We leave these topics for future research.

#### SUPPLEMENTARY MATERIAL

The Supplementary Material contains additional simulation results, empirical results and proofs of all theorems. Appendix A provides additional simulation results of MLE,

stationarity and asymmetry testings, and empirical results on portfolio returns. Appendix B gives proofs of all theoretical results. Appendix C provides useful properties of stable densities and lemmas with proofs. Appendix D gives the explicit expressions of partial derivatives of  $\ell_t(\theta)$  and  $\Upsilon$ .

## References

- <span id="page-30-3"></span>Adler, R.J., Feldman, R.E. and Taqqu, M.S. (1998). A Practical Guide to Heavy Tails: Statistical Techniques and Applications. Birkhäuser, Boston.
- Andrews, B., Calder, M. and Davis, R.A. (2009). Maximum likelihood estimation for  $\alpha$ stable autoregressive processes. Ann. Statist. 37(4), 1946–1982.
- <span id="page-30-2"></span>Bai, J. (2003). Testing parametric conditional distributions of dynamic models. Review of Economics and Statistics 85(3), 531–549
- <span id="page-30-1"></span>Bollerslev, T. (1986). Generalized autoregressive conditional heteroskedasticity. J. Econometrics 31(3), 307–327.
- <span id="page-30-4"></span>Bougerol, P. and Picard, N. (1992a) Stationarity of GARCH processes and of some nonnegative time series. J. Econometrics  $52(1-2)$ ,  $115-127$ .
- <span id="page-30-5"></span>Bougerol, P. and Picard, N. (1992b) Strict Stationarity of generalized autoregressive processes. Ann. Probability  $20(4)$ , 1714–1730.
- <span id="page-30-0"></span>Buraczewski, D., Damek, E. and Mikosch, T. (2016). Stochastic Models with Power-Law Tails: The Equation  $X = AX + B$ . Springer, Cham.
- <span id="page-30-6"></span>Chan, N.H. and Ng, C.T. (2009). Statistical inference for non-stationary  $GARCH(p,q)$ models. Electron. J. Stat. 3, 956–992.
- <span id="page-31-4"></span>Calzolari, G., Halbleib, R. and Parrini, A. (2014). Estimating GARCH-type models with symmetric stable innovations: Indirect inference versus maximum likelihood. Comput. Statist. and Data Analysis 76, 158–171.
- <span id="page-31-6"></span>Clauset, A., Shalizi, C. R. and Newman, M. E. (2009). Power-law distributions in empirical data. SIAM Review, 51(4):661–703.
- <span id="page-31-7"></span>Chen, Y. and Wang, R. (2024). Infinite-mean models in risk management: Discussions and recent advances. arXiv:2408.08678v1.
- <span id="page-31-0"></span>Engle, R.F. (1982). Autoregressive conditional heteroscedasticity with estimates of the variance of United Kingdom inflation. Econometrica 50(4), 987–1007.
- <span id="page-31-5"></span><span id="page-31-3"></span>Fama, E. (1965). The behavior of stock market prices. J. Bus.  $38(1)$ ,  $34-105$ .
- Fama, E.F. and Roll, R. (1968). Some properties of symmetric stable distributions. J. Amer. Statist. Assoc. 63(323), 817–836.
- <span id="page-31-1"></span>Fan, J. and Yao, Q. (2017). The Elements of Financial Econometrics. Cambridge University Press, Cambridge.
- Francq, C. and Zakoïan, J.-M. (2004). Maximum likelihood estimation of pure GARCH and ARMA-GARCH processes. Bernoulli 10(4), 605–637.
- <span id="page-31-8"></span>Francq, C. and Zakoïan, J.-M. (2012). Strict stationarity testing and estimation of explosive and stationary generalized autoregressive conditional heteroscedasticity models. *Econometrica* **80**(2), 821–861.
- <span id="page-31-9"></span>Francq, C. and Zakoïan, J.-M. (2013). Inference in nonstationary asymmetric GARCH models. Ann. Stat. 41(4), 1970-1998.
- <span id="page-31-2"></span>Francq, C. and Zakoïan, J.-M. (2019). GARCH Models: Structure, Statistical Inference and Financial Applications, 2nd Edn. John Wiley.
- <span id="page-32-4"></span>Gnedenko, B.V. and Kolmogorov, A.N. (1954). Limit Distributions for Sums of Independent Random Variables. Addison-Wesley.
- <span id="page-32-5"></span>Hall, P. and Yao, Q. (2003). Inference in ARCH and GARCH models with heavy-tailed errors. *Econometrica*  $71(1)$ , 285–317.
- <span id="page-32-0"></span>Harvey, A.C. (2013). Dynamic Models for Volatility and Heavy Tails: With Applications to Financial and Economic Time Series. Cambridge University Press, Cambridge.
- Hörmann, S. (2008). Augmented GARCH sequences: Dependence structure and asymptotics. *Bernoulli*  $14(2)$ , 543–561.
- <span id="page-32-6"></span>Ibragimov, M., Ibragimov, R. and Walden, J. (2015). Heavy-Tailed Distributions and Robustness in Economics and Finance. Springer, Cham.
- <span id="page-32-1"></span>Ibragimov, R. and Prokhorov, A. (2017). Heavy Tails and Copulas: Topics in Dependence Modelling in Economics and Finance. World Scientific, Hackensack, NJ.
- <span id="page-32-8"></span>Jensen, S. T. and Rahbek, A. (2004a). Asymptotic normality of the QMLE estimator of ARCH in the nonstationary case. *Econometrica*  $72(2)$ , 641–646.
- <span id="page-32-9"></span>Jensen, S. T. and Rahbek, A. (2004b). Asymptotic inference for nonstationary GARCH. *Econometric Theory* **20**(6), 1203–1226.
- <span id="page-32-10"></span>Khmaladze, E.V. (1981). Martingale approach in the theory of goodness-of-fit tests. Theory of Probability and its Applications  $26(2)$ ,  $240-257$ .
- <span id="page-32-3"></span><span id="page-32-2"></span>Kulik, R. and Soulier, P. (2020). Heavy-Tailed Time Series. Springer, New York.
- <span id="page-32-7"></span>Lévy, P. (1925). *Calcul des Probabilités*. Gauthier-Villars, Paris.
- Li, D., Tao, Y., Yang, Y. and Zhang, R. (2023). Maximum likelihood estimation for  $\alpha$ -stable double autoregressive models. J. Econometrics 236(1), 105471.
- <span id="page-33-6"></span>Liu, S.M. and Brorsen, B.W. (1995). Maximum likelihood estimation of a GARCH-stable model. J. Applied Econometrics 10(3), 273–285.
- <span id="page-33-3"></span>Mandelbrot, B.  $(1963)$ . The variation of certain speculative prices. J. Bus. **36**(4), 394–419.
- <span id="page-33-4"></span>Mandelbrot, B. (1997). Fractals and Scaling in Finance: Discontinuity, Concentration, Risk. Springer, New York.
- <span id="page-33-10"></span>Matsui, M. and Takemura, A. (2006). Some improvements in numerical evaluation of symmetric stable density and its derivatives. *Comm. Statist. Theory Methods* **35**(1), 149–172.
- <span id="page-33-5"></span>McCulloch, J. (1985). Interest-risk sensitive deposit insurance premia: Stable ACH estimates. Journal of Banking and Finance  $9(1)$ , 137–156.
- <span id="page-33-7"></span>Mittnik, S., Paolella, M. and Rachev, S. (2002). Stationarity of stable power-GARCH processes. J. Econometrics  $106(1)$ , 97-107.
- <span id="page-33-1"></span>Nair, J., Wierman, A. and Zwart, B. (2022). The Fundamentals of Heavy Tails: Properties, Emergence, and Estimation. Cambridge University Press, Cambridge.
- <span id="page-33-9"></span>Nolan, J.P. (1997). Numerical calculation of stable densities and distribution functions. Comm. Statist. Stachastic Models. 13(4), 759–774.
- <span id="page-33-2"></span>Nolan, J.P. (2014). Financial modeling with heavy-tailed stable distributions. WIREs Comput. Stat.  $6(1)$ , 45–55.
- <span id="page-33-0"></span>Nolan, J.P. (2020). Univariate Stable Distributions: Models for Heavy Tailed Data. Springer, Cham.
- <span id="page-33-8"></span>Peng, L. and Qi, Y. (2017). Inference for Heavy-Tailed Data: Applications in Insurance and Finance. Academic Press, London.
- <span id="page-34-7"></span>Panorska, A., Mittnik, S. and Rachev, S. (1995). Stable GARCH models for financial time series. Applied Mathematics Letters 8(5), 33–37.
- <span id="page-34-9"></span>Rachev, S.T. and Mittnik, S. (2000). Stable Paretian Models in Finance. Wiley, New York.
- <span id="page-34-0"></span>Resnick, S.I. (2007). Heavy-Tail Phenomena: Probabilistic and Statistical Modeling. Springer, New York.
- <span id="page-34-2"></span>Rydberg, T. (2000). Realistic statistical modelling of financial data. Int. Stat. Rev. 68(3), 233–258.
- <span id="page-34-5"></span>Samorodnitsky, G. and Taqqu, M.S. (1994). Stable Non-Gaussian Random Processes: Stochastic Models with Infinite Variance. Chapman & Hall, New York.
- <span id="page-34-3"></span>Samuelson, P.A. (1967). Efficient portfolio selection for Pareto-Lévy investments. The Journal of Financial and Quantitative Analysis 2(2), 107–122.
- <span id="page-34-6"></span>Silverberg, G. and Verspagen, B. (2007). The size distribution of innovations revisited: An application of extreme value statistics to citation and value measures of patent significance. Journal of Econometrics, 139(2), 318–339.
- <span id="page-34-1"></span>Taleb, N.N. (2020). Statistical Consequences of Fat Tails: Real World Preasymptotics, Epistemology, and Applications. STEM Academic Press.
- <span id="page-34-8"></span>Uchaikin, V.V. and Zolotarev, V.M. (1999). Chance and Stability: Stable Distributions and their Applications. VSP, Utrecht.
- <span id="page-34-4"></span>Zolotarev, V. M. (1986). One-Dimensional Stable Distributions. Amer. Math. Soc., Providence, RI.