INTEGRAL POINTS ON SINGULAR TORIC VARIETIES AND CYCLIC NORMAL POLYNOMIALS

ANDREW O'DESKY

Abstract. We establish a formula for the height zeta function for integral points on a class of projective toric varieties. Our method builds on the harmonic analysis approach of Batyrev–Tschinkel for rational points and is applicable even when the toric variety has cyclic quotient singularities. As an application, we determine the leading term in the asymptotic number of monic integral polynomials of bounded height with linearly independent roots and a given cyclic Galois group.

CONTENTS

1. INTRODUCTION

Let Y be a projective normal toric variety defined over $\mathbb Q$ with at worst cyclic quotient singularities. Let $T \subset Y$ be the open torus of Y. The problem of computing the asymptotic number of rational points on T of bounded height was taken up in $[3]$. Their strategy was to extend the height function to the space of adelic points $T(A)$ and exploit the spectral decomposition of the $T(\mathbb{Q})$ -invariant function

$$
g\mapsto \sum_{t\in T(\mathbb{Q})}\frac{1}{H(gt,s)}
$$

using the Poisson summation formula. The quantity s denotes an element of the complex vector space $Pic(Y) \otimes \mathbb{C}$. If the real part of s is sufficiently positive with respect to the effective cone then the sum converges absolutely, and the Poisson formula implies that

$$
\sum_{t \in T(\mathbb{Q})} \frac{1}{H(gt, s)} = \int_{(T(\mathbb{A})/T(\mathbb{Q}))^{\vee}} \widehat{H}(\chi, -s) \chi(g) d\chi
$$

where $H(\chi, -s)$ is the Fourier transform of $H(\cdot, -s) = H(\cdot, s)^{-1} \in L^2(T(\mathbb{A}))$. When $g = 1$ and s is proportional to the class $[L]$ of a big line bundle, this multiple Dirichlet series

Date: October 2, 2024. Research supported by NSF grant DMS-2103361.

specializes to the single Dirichlet series (height zeta function)

$$
Z(s) = \sum_{t \in T(\mathbb{Q})} \frac{1}{H(t, s)}
$$

for rational points on T of bounded L-height. When $T(\mathbb{A})/T(\mathbb{Q})$ is compact and Y is smooth—as assumed in [\[3\]](#page-42-1)—the terms on the spectral side of the Poisson formula are easily isolated and expressed using Hecke L-functions. From the principal terms one can deduce the singularities of the height zeta function using analytic properties of Dedekind zeta functions.

Difficulties arise when $T(\mathbb{A})/T(\mathbb{Q})$ is not compact or Y is not smooth. In the non-compact case, the spectral side of the Poisson formula involves an integral over the continuous part of the automorphic spectrum of T, and it is difficult to determine where $Z(s)$ has singularities from the Fourier transform $\hat{H}(\chi, -s)$. In their follow-up papers [\[4\]](#page-43-0), [\[5\]](#page-43-1) treating the smooth non-compact case, this integral over the continuous spectrum was approximated by relating $\hat{H}(\chi, -s)$ to simpler functions constructed using certain cones ("X-functions"). By their method they inferred the abscissa of convergence for $Z(s)$, however their method has not been successfully applied to compute either secondary terms for the distribution of rational points, or leading terms for the distribution of integral points.

Let D be a T-stable divisor of $Y - T$ and let U be an integral model of $Y - D$. A rational point of T is called D-integral if it extends to an integral point of U. The present paper is concerned with the height zeta function in the integral singular non-compact setting:

$$
Z(s) = \sum_{\substack{t \in T(\mathbb{Q}) \\ D\text{-integral}}} \frac{1}{H(t,s)}.
$$

The Poisson strategy was applied in the integral smooth non-compact setting to study $Z(s)$ in [\[8\]](#page-43-2). Unfortunately their application of the Poisson strategy is flawed and has a published counterexample [\[22\]](#page-43-3) proven by a different method. According to [\[22,](#page-43-3) Remark 3.2.2], there is a gap in the proof of [\[8,](#page-43-2) Lemma 3.11.4] where it is assumed that certain \mathcal{X} -functions are non-vanishing.

In this paper we do not use X-functions and thus do not follow the strategy of [\[4\]](#page-43-0), [\[5\]](#page-43-1), [\[8\]](#page-43-2) in treating the integral over the continuous spectrum of $T(\mathbb{A})/T(\mathbb{Q})$. The form to be integrated is $\widehat{H}(\chi, -s) d\chi$ which has singularities along affine hyperplanes depending on the fan of Y. This is a strong constraint, and our strategy is to use the multidimensional residue formula from [\[14\]](#page-43-4) which applies to such integrals. Poles of meromorphic forms with hyperplane singularities are indexed by *flags*, and our residue formula shows that only a small subset of flags contribute a residue to the integral which does not cancel identically with residues from other flags. Since this residue formula does not involve approximation, we obtain an exact formula for $Z(s)$ in terms of simpler multiple Dirichlet series and rational functions that can be computed from the fan of Y . Our residue formula requires that the fan of Y satisfies a certain compatibility condition (see §[3.2\)](#page-16-1).

Theorem 1. Assume the residue method is applicable for the fan Σ , i.e. each Σ -convex singular flag is compatible with the polyhedron Π_{σ} for every maximal cone $\sigma \in \Sigma_D(r)$. Then the height zeta function $Z(s)$ for D-integral rational points on the torus is equal to

$$
Z(s) = \frac{|\mu|}{|C||R|P} \sum_{\xi \in R^{\vee}} \sum_{\psi \in C^{\vee}} \sum_{\tau \in \Sigma_D} \sum_{\gamma \in Z_{\sigma_{\tau}}} L(s + 2\pi i m_{\gamma\xi}, (\xi \psi)^{-1}, -\tau^{\circ}, D) R_{\gamma}(s + 2\pi i m_{\xi})
$$

in the region of $s \in Pic(Y) \otimes \mathbb{C}$ where both sides converge.

The arithmetic and combinatorial invariants appearing in the formula can be described as follows:

- $-$ μ is the torsion subgroup of $T(\mathbb{Q}) \cap (T(\mathbb{R}) \times \prod_{v \text{ finite}} K_v)$ where K_v is the open compact subgroup of $T(\mathbb{Q}_v)$ stabilizing $U(\mathbb{Z}_v)$ (§[2.3.2\)](#page-11-0),
- |R| is the volume of the level K regulator group R of the torus T, i.e. the identity component of $T(\mathbb{Q})\backslash T(\mathbb{A})^1/K$ where $T(\mathbb{A})^1$ is the norm-one subgroup of $T(\mathbb{A})$ and $K = K_{\infty}^{\max} \times \prod_{v \text{ finite}} K_v \text{ (§2.3.1)},$ $K = K_{\infty}^{\max} \times \prod_{v \text{ finite}} K_v \text{ (§2.3.1)},$ $K = K_{\infty}^{\max} \times \prod_{v \text{ finite}} K_v \text{ (§2.3.1)},$
- C is the level K class group of the torus T, i.e. the quotient of $T(\mathbb{Q})\backslash T(\mathbb{A})^1/K$ by R $(§2.3.1),$ $(§2.3.1),$ $(§2.3.1),$
- (−) [∨] denotes Pontryagin dual,
- Σ_D is a subfan of Σ_{Γ} determined by T, D and Σ (§[3.4\)](#page-25-0) where
- $-\Sigma_{\Gamma}$ is the restricted fan of Σ with respect to the action of the Galois group Γ of the splitting field of T (§[2.2.3\)](#page-9-0),
- $-\sigma_{\tau}$ is any maximal cone of Σ_{D} containing τ ,
- Z_{σ} is the subset of flags terminating in Π_{σ} cut out by a collection of Π_{σ} -stable singular hyperplanes of the meromorphic form $\widetilde{H_{\infty}}(\chi_m, -s)$ dm for a cone σ in $\Sigma_{\Gamma}(r)$ (§[3.2\)](#page-16-1) where
- $-\Pi_{\sigma} = \mathfrak{t}^{\vee} + i\sigma^{\vee} \subset \mathfrak{t}^{\vee} \otimes \mathbb{C}$ is a polyhedron with boundary \mathfrak{t}^{\vee} (§[3.2.2\)](#page-17-0) where
- $-$ t^{\vee} is the Galois invariant subspace of the real vector space $\mathfrak{t}_E^{\vee} = \mathbb{R} \otimes N_E$ where E is the splitting field of T and N_E is the cocharacter lattice of T_E (§[2.1\)](#page-6-1),
- $-m_{\gamma\xi}$ is the terminal point of the flag γ of $\widehat{H_{\infty}}(\chi_m, -(s+2\pi im_{\xi})) dm$ (§[3.2.4\)](#page-18-0) and m_{ξ} is the local exponent of ξ ,
- $-\tau^{\circ} = \tau \cup_{\tau' \subsetneq \tau} \tau'$ is the interior of a cone τ ,
- $\Sigma(1)$ is the set of generators of Σ ,
- $-L(s, \xi, \sigma, D)$ is a certain multiple Dirichlet series in $s \in \mathbb{C}^{\Sigma(1)}$ (§[3.3.4\)](#page-25-1)
- $-R_{\gamma}(s)$ is a certain rational function on Pic(Y) defined by Definition [8](#page-19-0) and given explicitly by (9) $(\S3.3)$ $(\S3.3)$,
- and P is a period determined by the fan given by

$$
P = \sum_{\gamma \in Z_{\sigma_1}} \int_{\mathfrak{t}_0^{\vee}} R_{\gamma}(s_{AC} + 2\pi im) \, dm
$$

where s_{AC} is the element of $\mathbb{Z}^{\Sigma(1)}$ corresponding to the sum of the toric divisors of Y, σ_1 is an arbitrary maximal cone of Σ_D , and t_0 is the trace-zero subspace of t (§[2.3.2\)](#page-11-0).

Our work is incomplete in the sense that we do not make contact with Peyre's formula for the leading constant in the rational case, nor Batyrev–Manin's conjecture on the shape of the leading term. Both conjectures have recently been formulated in the integral and smooth setting with a proof using universal torsors for toric varieties in a recent preprint [\[18\]](#page-43-5). One would like to verify that the leading term predicted by our formula in the case that Y is smooth agrees with [\[18\]](#page-43-5) though we have yet to carry out this important verification.

1.0.1. Explicit formulas for summatory functions. In favorable cases, applying Perron's formula to our formula for $Z(s)$ results in explicit formulas for the sum of Dirichlet coefficients in terms of automorphic data. These formulas are quite complicated but can sometimes be analyzed to extract main terms and secondary terms. The idea is that one expects the "conical" multiple Dirichlet series $L(s + 2\pi i m_{\gamma\xi}, (\xi\psi)^{-1}, -\tau^{\circ}, D)$ in the formula for $Z(s)$ to

4 ANDREW O'DESKY

be nearly automorphic and the rational functions R_{γ} to be nonvanishing. (We have been able to prove that R_{γ} is nonvanishing under a strong convexity assumption (Proposition [4\)](#page-21-0) though we expect it to hold generally.)

We illustrate our formula with the real quadratic Severi–Brauer surface associated to a real quadratic field E^1 E^1 . See Figure [4](#page-24-0) for the fan. For an appropriate boundary divisor, the integral Diophantine problem has the nonzero algebraic integers in E as its solutions, and the height zeta function is, by definition,

$$
Z(s) = \sum_{\substack{a \in O_E \\ a \neq 0}} \frac{1}{\max(|a|, |a'|)^s}
$$

where a' is the conjugate of a. The regulator group R is $\mathbb{R}/\mathbb{Z}\log u$ where u is a positive generator of O_E^{\times} . The regulator characters are ξ^k for integers k where $\xi(y) = |y|^{2\pi i/\log u}$. The fan Σ_D consists of a single cone σ , and Z_σ consists of a single flag. Our formula is

$$
Z(s) = \frac{4}{hR} \sum_{k \in \mathbb{Z}} \sum_{\psi \in C^{\vee}} L(\frac{s}{2}, \psi \xi^k) \frac{s}{s^2 + (\frac{2\pi k}{R})^2}
$$

where h denotes the class number, R the regulator, and $L(s, \chi) = \sum_{I \subset O_E} \chi(I) N(I)^{-s} =$ $\sum_{N} b_N(\chi) N^{-s}$ is the Hecke L-function of χ . Since Y has Picard number one, the conical Dirichlet series reduce to one-variable Dirichlet series. This is a useful formula since we have related the distribution of integral points of bounded height—a global problem—to the distribution of ideals of bounded norm, and the latter can be described with Hecke characters. Now we apply Perron's formula to obtain an explicit formula for the number of algebraic integers in E of bounded height in terms of automorphic data:

$$
\# \{ a \in O_E : 0 < |a|, |a'| < X \} = \frac{4}{hR} \sum_{\psi \in C^{\vee}} \sum_{N < X^2} b_N(\psi) \log(X/\sqrt{N}) + \frac{8\pi}{hR^2} \sum_{\substack{k \in \mathbb{Z} \\ k \neq 0}} \sum_{\psi \in C^{\vee}} \sum_{N < X^2} k b_N(\psi \xi^k) \sin\left(\frac{2\pi k}{R} \log(\sqrt{N}/X)\right).
$$

With the class number formula we can verify that the first term is the principal term:

$$
\frac{4}{hR} \sum_{\psi \in C^{\vee}} \sum_{N < X^2} b_N(\psi) \log(X/\sqrt{N}) = \frac{2}{hR} \operatorname{res}[\zeta_E] \cdot X^2 + O(X^{2-\delta}) = \frac{4}{\sqrt{d_E}} \cdot X^2 + O(X^{2-\delta}),
$$

which is the expected number of points of a lattice with covolume $\sqrt{d_E}$ in a box of sidelength 2X. The contribution of the rational functions $R_{\gamma} = \frac{s}{\gamma^2 + s^2}$ $s^2+\left(\frac{2\pi k}{R}\right)$ $\frac{1}{2}$ to this formula are the $\log(X/\sqrt{N})$ and $\sin\left(\frac{2\pi k}{R}\right)$ $\frac{\pi k}{R} \log(\sqrt{N}/X)$ terms (obtained from well-known inverse Mellin transforms).

¹For imaginary quadratic Severi–Brauer surfaces the height zeta function is a specialization of a real analytic Eisenstein series for GL_2 and the rational functions R_γ all have degree zero, so it is "not interesting" as far as illustrating our formula. For quasi-split tori the function R_{γ} is the reciprocal of a polynomial whose degree is the rank of the unit group, and real quadratic fields are the first interesting examples.

Other explicit formulas related to cubic abelian fields have been derived in [\[6\]](#page-43-6), [\[13\]](#page-43-7) by this method. In [\[6\]](#page-43-6) this was used to obtain leading and secondary terms for the number of integral monic trace-one cubic abelian polynomials of bounded root height.[2](#page-4-0)

1.1. Cyclic normal polynomials. We apply our formula to a problem in arithmetic statistics: counting monic integral polynomials with a given cyclic Galois group and linearly independent roots in their splitting field. We call an irreducible polynomial *normal* if its roots are Q-linearly independent in the splitting field. Normal polynomials with a given Galois group admit an orbit parametrization $[15]$. Let G be a finite group. The orbit parametrization from [\[15\]](#page-43-8) is similar to a prehomogeneous space in that it is a projective variety $Y = Y_G$ with an action of a reductive group $\mathcal G$ (the unit group of the group algebra of G) and containing a dense open G-orbit $T \subset Y$, however the orbit T is not generally birational to projective space. The rational points of T parametrize data $(K/\mathbb{Q}, x)$ where K/\mathbb{Q} is an étale \mathbb{Q} -algebra with Galois group G and x is a normal element of K. (An element of K is normal if its G-conjugates are $\mathbb Q$ -linearly independent.) We recall this construction in §[4.](#page-29-0) When G is abelian this orbit parametrization Y has a natural toric structure, and we may apply the methods of this paper to study the distribution of such polynomials.

Theorem 2. Let $n \geq 5$. Let $N(C_n, H)$ be the number of normal monic degree n integral polynomials with Galois group C_n and root height less than H. Then

$$
N(C_n, H) = \varphi(n)^{-1} \frac{|\mu||C \cap \ker \gamma^*|}{|C||R|P} \kappa_0 \left(\sum_{d|n} \kappa_d\right) \frac{1}{n} H^n(1 + o(1))
$$

where

$$
\kappa_0 = \left(\prod_{d|n} \text{res}[\zeta_{\mathbb{Q}(\zeta_d)}]\right) \prod_p Q_p(p^{-\frac{1}{e_p}s_{AC}})
$$

and

$$
\kappa_d = \left(\frac{\varphi(d)}{2n}\right)^{\frac{1}{2}(1-\varphi(n))} \sum_{\tau \in \Sigma_{\Gamma_{\infty}}(r_{\infty})} \left(\frac{\tau}{\sigma_{(d)}}\right) \frac{[N_{\infty} : \mathbb{Z}\tau(1)]}{[N : \mathbb{Z}\sigma_{(d)}(1)]}.
$$

The factors of the leading constant may be interpreted as follows:

- $-\varphi(n)$ is Euler's totient function, and also the number of twists of a rational point $(K/\mathbb{Q}, x)$ for a C_n -field K by an outer automorphism of C_n ; there are $\varphi(n)$ such rational points of T giving rise to any C_n -polynomial so we divide by this factor (cf. Corollary [2](#page-32-1) or the examples in [\[6,](#page-43-6) §2]),
- the open torus $T \subset Y$ is the quotient torus $T = \mathcal{G}/C_n$ where $\mathcal G$ is the group of units in the group algebra of C_n ,
- $\frac{|\mu|}{|C||E}$ $\frac{|\mu|}{|C||R|P}$ comes from our formula for $Z(s)$,
- $-\gamma$: $P = \mathbb{G}_m \times \mathcal{G} \to \mathcal{G} \to T$ where $\mathcal{G} \to T$ is the natural quotient homomorphism and $\gamma^*: T(\mathbb{A})^\vee \to P(\mathbb{A})^\vee$ is the pullback homomorphism on characters (§[2.5\)](#page-15-0),
- $-|C \cap \ker \gamma^*|$ is the number of Hecke characters on $R \times C$ which contribute to the principal terms,
- κ_0 is part of the leading constant of the principal conical Dirichlet series,

²The root height of a polynomial $f = t^n + a_1 t^{n-1} + \cdots + a_n \in \mathbb{Z}[t]$ is $H(f) = \max(|a_1|, |a_2|^{1/2}, \ldots, |a_n|^{1/n})$.

6 ANDREW O'DESKY

- Q_p is a polynomial determined by the fan which measures the singularities of Y; it only depends on the decomposition group at p in the splitting field E/\mathbb{Q} (Proposition [9\)](#page-35-0),
- $-\sum_{d} \kappa_d$ is the sum of the values of the rational functions R_γ at the abscissa of convergence,
- $N = N_{E}^{\Gamma}$ and $N_{\infty} = N_{E}^{\Gamma_{\infty}}$ where Γ_{∞} is the decomposition group at infinity,
- $-r_{\infty}=\frac{1}{2}$ $\frac{1}{2}(1+\sum_{d|n}\varphi(d))$ is the rank of $\prod_{d|n}\mathbb{Z}[\zeta_d]^{\times}$,
- Σ is the fan for Y constructed in §[5,](#page-32-0)
- $-\sigma_{(d)}$ is the cone of Σ_D corresponding to the unique Galois orbit in $C_n^{\vee} \subset \Sigma(1)$ of size $\varphi(d)$ (§[6.1\)](#page-33-1),
- $-\int_{a}^{T}$ σ is the number of unordered r-subsets $E \subset \tau(1)$ such that $\pi E = \sigma(1)$ where $\pi \colon N_E \to N$ denotes restriction to M_E^{Γ} and $M_E = N_E^{\vee}$.

Our proofs do not make use of any known cases of Malle's conjecture nor class field theory, though we do make use of known zero-free regions in the critical strip for L-functions of Hecke characters on cyclotomic fields. We expect that the same asymptotic formula holds when counting all C_n -polynomials (see Remark [20\)](#page-42-2).

1.1.1. *Malle's conjecture*. Our theorem on abelian polynomials may be compared with another asymptotic formula for $N(C_n, H)$ implied by Malle's conjecture. Let G be a regular permutation group of order $n \geq 4$, and let $D_G(s)$ be the "discriminant series"

$$
D_G(s) = \sum_{K/\mathbb{Q} \text{ } G \text{-field}} |d_K|^{-s}.
$$

Malle's conjecture predicts there is a positive constant $a < \frac{1}{2}$ such that the number $F(G, H)$ of G-fields K/\mathbb{Q} with $|d_K| < X$ satisfies $X^{a-\varepsilon} \ll_{\varepsilon} F(G, H) \ll_{\varepsilon} X^{a+\varepsilon}$ for any $\varepsilon > 0$ as X goes to infinity; in particular, $D_G(1/2)$ is finite. Let $f_x \in \mathbb{Z}[t]$ denote the characteristic polynomial of an element $x \in K$. As H goes to infinity, we have that

$$
N(G,H) = \sum_{K/\mathbb{Q}} \frac{1}{G} \# \{ x \in O_K : K = \mathbb{Q}(x), H(f_x) < H \} = c_G D_G(1/2) H^n(1 + o(1))
$$

where c_G is the scaling ratio between root height and Minkowski norm. It would be interesting to compare the formulas for $D_{C_p}(1/2)$ when p is prime provided by our theorem and [\[9\]](#page-43-9).

We are very grateful to Anshul Adve, Tim Browning, Vesselin Dimitrov, Peter Sarnak, Victor Wang, and Shou-Wu Zhang for helpful discussions. In particular, we would like to thank Manjul Bhargava for many inspiring discussions and also for showing us the argument in §[1.1.1.](#page-5-0)

Notation. The places of a number field E are denoted P_E and the completion of E at a place w is denoted E_w . For a finite place w of E we set $|x|_w := q_w^{\text{ord}_w(x)}$ where $\text{ord}_w : E^\times \to \mathbb{Z}$ is the normalized additive valuation associated to w and q_w is the size of the residue field of the ring of integers of E_w . The norm $|x|_w$ is also the multiplier for the additive Haar measure on E_w under $y \mapsto yx$. For an infinite place w corresponding to an embedding $i: E \hookrightarrow \mathbb{C}$, let $|x|_w := |ix|_{\mathbb{C}}^{d_w}$ where $d_w = 1$ (resp. $d_w = 2$) if w is real (resp. if w is complex), and we set ord $_{w}(x) = -\frac{\log |x|_{w}}{\log q_{w}}$ $\frac{\log |x|_w}{\log q_w}$ where $q_w \coloneqq e^{d_w}$. With these norms the product formula takes the form $\prod_{w\in P_{\mathcal{E}}}|x|_w=1$ for any nonzero x. If E is Galois over Q and v is a place of Q we let e_v (resp. f_v) denote the ramification (resp. inertia) index of any place of E lying over v; by

convention, $e_v = 1$ if $v = \infty$. We set $e(x) = e^{2\pi ix}$. For a fan Σ we let $|\Sigma|$ denote the union of its cones in the ambient space t. A subset of t is called Σ -convex if it is contained in a cone of Σ.

2. Integral points on toric varieties

2.1. Preliminaries on algebraic tori. Let T be an *n*-dimensional algebraic torus defined over Q. Let E/\mathbb{Q} be the minimal Galois extension which splits T and let $\Gamma = G(E/\mathbb{Q})$ be its Galois group. For a field F containing Q write $T_F = T \otimes F$ for the base-change of T to F. Let $X^*T = \text{Hom}(T, \mathbb{G}_{m, \mathbb{Q}})$ be the set of rational characters of T and let $X_*T = \text{Hom}(\mathbb{G}_{m, \mathbb{Q}}, T)$ be the set of rational cocharacters of T; we will also need $X^*T_E = \text{Hom}(T, \mathbb{G}_{m,E})$ and $X_*T_E = \text{Hom}(\mathbb{G}_{m,E}, T_E)$. These sets are finite free abelian groups and there is a canonical pairing $X^*T \times X_*T \to \mathbb{Z}$ given by $(m, n) \mapsto \deg m(n)$. While basic examples show this pairing is generally not perfect, it does identify X_*T with a finite index sublattice of the dual lattice to X^*T so this pairing induces a canonical isomorphism on the level of real vector spaces:

$$
\mathbb{R} \otimes X_*T \cong \mathbb{R} \otimes X^*T^{\vee}.
$$

The character lattice X^*T_E has a natural Galois action by post-composition and we equip X_*T_E with the corresponding dual representation. It is however more convenient to use additive notation for these lattices. We denote these with M and N :

$$
M_E \cong X^*T_E: u \leftrightarrow \chi^u, \quad M = (M_E)^{\Gamma} = X^*T, \quad N_E = X_*T_E, \quad N = N_E^{\Gamma} = X_*T.
$$

The associated real vector spaces will be denoted

$$
\mathfrak{t}_E^{\vee} = \mathbb{R} \otimes M_E, \quad \mathfrak{t}^{\vee} = (\mathfrak{t}_E^{\vee})^{\Gamma}, \quad \mathfrak{t}_E = \mathbb{R} \otimes N_E = \text{LieAlg}(T_E), \quad \mathfrak{t} = \mathfrak{t}_E^{\Gamma}
$$

Definition 1. Let w be a place of E. For any $t \in T(E_w)$ the function $\chi \mapsto \text{ord}_w(\chi(t))$ on characters $\chi \in X^*T_{E_w}$ determines an element of $\mathbb{R} \otimes X_*T_{E_w}$. Let

$$
\ell_w(t) \in \mathfrak{t}_E
$$

be the corresponding cocharacter under the canonical isomorphism $\mathbb{R} \otimes X_* T_{E_w} \cong \mathbb{R} \otimes X_* T_E$ induced by base change of the split torus T_E along the inclusion $E \to E_w$.

Let w be a place of E and let v be its restriction to \mathbb{Q} . Let

$$
N_w = N_E^{\Gamma_w}, \quad \mathfrak{t}_w = \mathfrak{t}_E^{\Gamma_w}
$$

where Γ_w is the decomposition group at w. Let $K_v^{\max} \subset T(\mathbb{Q}_v)$ be the maximal compact subgroup.

Proposition 1 ([\[10,](#page-43-10) (1.3), p. 449]). Let w be a place of E and let v be its restriction to \mathbb{Q} . There is an exact sequence

$$
1 \longrightarrow K_v^{\max} \longrightarrow T(\mathbb{Q}_v) \xrightarrow{\ell_w} \mathfrak{t}_w.
$$

If w is infinite then ℓ_w is surjective, if w is finite and unramified then the image of ℓ_w is the lattice N_w , and if v is ramified then the image of ℓ_w is a finite index subgroup of N_w .

We recall the definition of the toric height on Y following [\[3\]](#page-42-1).

Definition 2. Let $\varphi: \mathfrak{t}_E \to \mathbb{C}$ be a function. For $t = (t_w)_w \in T(\mathbb{A}_E)$ let

$$
H(t,\varphi) = \prod_{w \in P_E} \left(q_w^{-\varphi(\ell_w(t_w))} \right)^{\frac{1}{[E:\mathbb{Q}]}}
$$

where $\varphi(\ell_w(t_w))$ is evaluated using the canonical isomorphism $X_*T_E \cong X_*T_{E_w}$.

The reader may easily verify the following basic properties:

- (1) $H(-, \varphi)$ is invariant under the action of the maximal compact subgroup of $T(\mathbb{A}_E)$.
- (2) $H(t, \varphi) = 1$ if $t \in T(\mathbb{Q})$ and φ is linear.
- (3) If $t \in T(\mathbb{Q}_v)$ where w lies over v and φ is Γ-invariant, then $\varphi(\ell_w(t))$ is independent of w, even though $\ell_w(t)$ depends on the choice of w over v in general.
- (4) If $t = (t_w)_w \in T(\mathbb{A})$ embedded diagonally in $T(\mathbb{A}_E)$ and φ is Γ-invariant, then

$$
H(t,\varphi) = \prod_{v \in P_{\mathbb{Q}}} q_v^{-\frac{1}{e_v} \varphi(\ell_w(t_w))}.
$$
\n(1)

Remark 1. The toric height defined in [\[3\]](#page-42-1) differs slightly from ours. We have introduced normalization factors that are needed for (2) to hold, which is itself needed in order for the height zeta function to descend to a function on Pic Y and not just $Pic^T Y$.

2.2. Preliminaries on toric varieties. Let Σ denote the fan of Y_E in \mathfrak{t}_E . We do not assume Y is smooth and permit cyclic quotient singularities. This means the fan Σ is simplicial: all the cones are simplices. Let $Div_T Y$ denote the group of T-Cartier divisors on Y. See [\[12\]](#page-43-11) for more details. We let $\Sigma(k)$ denote the set of k-dimensional cones in Σ and identify onedimensional cones with their minimal lattice generators. The next proposition is well-known $[12, §3.4]$ $[12, §3.4]$.

Proposition 2. There is a commutative diagram with exact rows:

The two vertical inclusions become isomorphisms after tensoring with \mathbb{Q} . The canonical Γaction on N_E preserves the fan of Y_E . The group $\text{Div}_T Y_E$ is naturally isomorphic with the group of Σ -linear^{[3](#page-7-0)} functions $\varphi: \mathfrak{t}_E \to \mathbb{R}$ which are \mathbb{Z} -valued on N_E . The Σ -linear function φ corresponding to the line bundle $\mathcal{O}(\sum_{e \in \Sigma(1)} s_e D_e)$ satisfies $\varphi(e) = -s_e$ for each $e \in \Sigma(1)$. A T-Cartier divisor on Y_E descends to a T-Cartier divisor on Y if and only if its associated Σ-linear function is Galois invariant.

To each $s \in \mathbb{C}^{\Sigma(1)}$ we associate the unique Σ -linear function φ satisfying

$$
\varphi(e) = -s_e
$$

for all $e \in \Sigma(1)$, and set

$$
H(t,s) \coloneqq H(t,\varphi).
$$

Remark 2. Our sign convention on φ follows [\[12\]](#page-43-11) and not [\[3\]](#page-42-1). In particular, the linear function $\varphi_{\text{div}(m)}: \mathfrak{t}_E \to \mathbb{R}$ is equal to $-m$ and not m.

³i.e. continuous functions $\varphi: \mathfrak{t}_E \to \mathbb{R}$ such that the restriction of φ to each cone $\sigma \in \Sigma$ is equal to the restriction of some linear form $u(\sigma) \in M_E$.

2.2.1. Integral points. Let D be an effective T-stable Weil divisor of Y ("boundary divisor") and let U be an integral model for $Y - D$. An adelic point $t = (t_v)_v$ of T is D-integral if $t_v \in U(\mathbb{Z}_v)$ for all finite places v. The indicator function on D-integral points in $T(\mathbb{A})$ is a locally constant factorizable function

$$
\mathbf{1}^D = \prod_{v \in P_{\mathbb{Q}}} \mathbf{1}_v^D
$$

where 1^D_{∞} is identically one on $T(\mathbb{R})$. For finite places v let $K_v^D \subset T(\mathbb{Q}_v)$ be the (open compact) subgroup under which 1_D^D is invariant, and let K^D_{∞} denote the maximal compact subgroup of $T(\mathbb{R})$. Set $K^D = \prod_v \dot{K}_v^D$.

Warning 1. The toric height $t \mapsto H(t, s)$ is invariant under translation in x by elements of the maximal compact subgroup K^{\max} of $T(\mathbb{A})$, but to count D-integral points we must use $H(t, s)1^D(t)$ instead of $H(t, s)$ and the indicator function 1_D is generally not invariant under the maximal compact subgroup K^{max}. (E.g. the subgroup $K^{\overline{D}} \subset K$ has index 6 for the Diophantine problem in [\[6\]](#page-43-6), see [\[6,](#page-43-6) Lemma 2].) This means that in the Poisson-style approach to counting integral points, the integral [\(7\)](#page-16-2) over the automorphic spectrum of the torus T is complicated by characters with non-maximal level.

Remark 3 (Cartier divisors). When the boundary divisor D is Cartier there is a canonical integral model for $Y - D$ defined as follows. Let φ be the Σ -linear function corresponding to the T-Cartier divisor D. On each cone $\sigma \in \Sigma$ the restriction of φ to σ is equal to the restriction of some linear form $u(\sigma) \in M_E$. Let $U_{\sigma} = \text{Spec } \mathbb{Q}[\sigma^{\vee} \cap M_E] \subset Y_E$ denote the affine open subvariety corresponding to a cone σ . The ideal $I \subset \mathcal{O}(U_{\sigma})$ corresponding to $D \cap U_{\sigma}$ is generated by the character corresponding to $u(\sigma)$, i.e.

$$
I = \mathbb{Q}[\sigma^{\vee} \cap M_E] \chi^{u(\sigma)}.
$$

From this description it is clear that $U_{\sigma} - D \cap U_{\sigma}$ has the integral model

$$
\text{Spec } \mathbb{Z}[\sigma^{\vee} \cap M_E][\chi^{-u(\sigma)}].
$$

The affine opens $\{U_{\sigma}\}\$ form an open cover of Y_E and glue to an integral model U for $Y - D$. We have the following integrality criterion:

$$
t \in T(\mathbb{Q}_v) \cap U(\mathbb{Z}_v) \iff \chi^{u(\sigma)}(t) \in \mathbb{Z}_v^{\times}
$$
 whenever $t \in U_{\sigma}$.

Nonetheless, in applications it is important to consider integral points with respect to non-Cartier divisors, e.g. the boundary divisor in [\[6\]](#page-43-6) is not Cartier and its notion of integrality corresponds to the natural condition of being monic for integral cubic abelian polynomials.

2.2.2. The height zeta function. Let $\Phi = \prod_{v \in P_{\mathbb{Q}}} \Phi_v : T(\mathbb{A}) \to \mathbb{C}$ be a locally constant factorizable function for which Φ_{∞} is identically one on $T(\mathbb{R})$.

Definition 3. Define the multiple Dirichlet series

$$
Z(s, \Phi) = \sum_{t \in T(\mathbb{Q})} \frac{\Phi(t)}{H(t, s)}.
$$

The height zeta function (for D-integral points in T) is

$$
Z(s) = Z(s, 1_D) = \sum_{\substack{t \in T(\mathbb{Q}) \\ D\text{-integral}}} \frac{1}{H(t, s)}.
$$

This converges to a holomorphic function of several complex variables on the region of $Pic(Y) \otimes \mathbb{C}$ whose real part is sufficiently positive with respect to the effective cone.

2.2.3. Restricted fans. We fix once and for all a choice of place w of E lying over each place v of Q. If $\Gamma_w \subset \Gamma$ is the decomposition group at the chosen place w of E over v then we write Γ_v for Γ_w .

Definition 4 (restricted fan). For any subgroup Γ_v of Γ let Σ_{Γ_v} denote the fan in $\mathfrak{t}_E^{\Gamma_v}$ E consisting of the cones $\sigma \cap \mathfrak{t}_E^{\Gamma_v}$ for each cone σ in Σ .

The next lemma describes the restricted fan and its generators for a simplicial fan Σ which is not necessarily regular; cf. [\[3,](#page-42-1) Prop. 1.3.2] for the regular case.

Lemma 1.

- (1) For each $\sigma \in \Sigma_{\Gamma}$, there is a unique cone $\tau_{\sigma} \in \Sigma$ which is minimal among cones in Σ containing σ. The cone τ_{σ} will be fixed by Γ (i.e. $\gamma(\tau_{\sigma}) = \tau_{\sigma}$ for all $\gamma \in \Gamma$) and satisfies $\tau_{\sigma} \cap \mathfrak{t} = \sigma$. The map $\sigma \to \tau_{\sigma}$ determines a bijection between Σ_{Γ} and the subset of cones of Σ which are fixed by Γ .
- (2) Let $e_0 \in \Sigma_{\Gamma}(1)$ be a lattice generator for a ray σ in Σ_{Γ} . Then Γ acts transitively on $\tau_{\sigma}(1)$ and

$$
e_0 = \frac{1}{k} \sum_{e \in \tau_\sigma(1)} e \tag{2}
$$

for a positive integer determined by σ . If τ_{σ} is regular, then $k = 1$. The bijection $\sigma \leftrightarrow \tau_{\sigma}$ determines a bijection between $\Sigma_{\Gamma}(1)$ and the set of Γ -orbits in $\Sigma(1)$ equal to $\tau(1)$ for some cone $\tau \in \Sigma$.

(3) The fan Σ_{Γ} is simplicial. If $\tau \in \Sigma$ is regular then $\tau \cap \mathfrak{t} \in \Sigma_{\Gamma}$ is regular.

Proof. Since Σ is closed under intersection, τ_{σ} is simply the intersection of all cones in Σ containing σ ; if τ_{σ} were not fixed by Γ, then it would not be minimal. By definition of Σ_{Γ} there is a cone $\tau \in \Sigma$ for which $\tau \cap \mathfrak{t} = \sigma$. Since τ contains τ_{σ} we see that $\tau_{\sigma} \cap \mathfrak{t} = \sigma$. To see this is a bijection, we will show that any Γ-fixed cone τ of Σ satisfies $\tau = \tau_{\sigma}$ with $\sigma = \tau \cap \mathfrak{t}$. Clearly τ contains τ_{σ} , so it suffices to show that if $\tau' \subset \tau$ is another Γ-fixed cone satisfying $\tau' \cap \mathfrak{t} = \sigma$, then $\tau' = \tau$. To the contrary, if $\tau' \neq \tau$ then there is some $e \in \tau(1) \setminus \tau'(1)$. Since τ' is Γ-fixed, $\tau'(1)$ is a proper Γ-stable subset of $\tau(1)$, so the Γ-orbit of e has no intersection with $\tau'(1)$. This shows that $\sum_{\gamma} \gamma(e) \notin \tau'$, for otherwise we would have a nontrivial relation among the linearly independent generators of τ ; thus $\tau' \cap \mathfrak{t} \neq \sigma$ since it fails to contain the point $\sum_{\gamma} \gamma(e) \in \sigma$.

Let $e_0 \in \Sigma_\Gamma(1)$ be a lattice generator for a ray σ in Σ_Γ . Then $\sum_{e \in \tau_\sigma(1)} e \in \sigma \cap N$ so it is equal to ke₀ for some integer $k \in \underline{\mathbb{Z}}_{\geq 1}$. Since τ_{σ} is minimal the group Γ must act transitively on $\tau_{\sigma}(1)$. If τ_{σ} is regular, then $\frac{1}{k} \sum_{e \in \tau_{\sigma}(1)}^{\infty} e$ can only be a lattice point if $k = 1$.

Let $\tau \in \Sigma$ be Γ-fixed and let $\sigma = \tau \cap \mathfrak{t}$. Each generator of σ corresponds via [\(2\)](#page-9-1) to a Γ-orbit in $\tau(1)$. Since $\tau(1)$ extends to a Q-basis of QN_E, the generators of σ given by [\(2\)](#page-9-1) extend to a Q-basis of QN. For the implication that τ regular $\implies \tau \cap \tau$ regular see [\[3,](#page-42-1) Prop. 1.3.2]. Prop. 1.3.2].

Remark 4. Example [3](#page-20-0) gives a regular cone $\sigma \in \Sigma_{\Gamma}$ for which $\tau_{\sigma} \in \Sigma$ is not regular.

2.3. Automorphic characters on the torus. Now we describe automorphic characters of the torus. We fix an arbitrary open compact subgroup K_{fin} of $T(\mathbb{A}_{fin})$ and set $K =$ $K_{\text{fin}} \times K_{\infty}^{\text{max}}$.

2.3.1. The norm-one subgroup of $T(\mathbb{A})$. We recall a logarithm map from the construction of Langlands–Eisenstein series, cf. [\[19,](#page-43-12) §6]: the homomorphism $\ell \colon T(\mathbb{A}) \to \mathfrak{t}$ defined by the formula

 $\ell(t)(m) = \log |\chi^m(t)|_A^{-1} \quad (m \in M).$

The norm-one subgroup $T(\mathbb{A})^1$ is the kernel of ℓ . A construction from [\[17\]](#page-43-13) shows that ℓ admits a splitting.

Lemma 2 ([\[17,](#page-43-13) Lemma 3.8.1, p. 132]). The logarithm map ℓ admits a continuous splitting s: $t \to T(\mathbb{A})$ which takes values in $T(\mathbb{R})$. In particular, ℓ is surjective.

Fix such a splitting $s: \mathfrak{t} \to T(\mathbb{A})$. This determines isomorphisms in the usual manner

$$
T(\mathbb{A}) \cong_s \mathfrak{t} \times T(\mathbb{A})^1 \colon t \mapsto (\ell(t), ts(\ell(t)))^{-1}), \qquad T(\mathbb{Q}) \backslash T(\mathbb{A}) / K \cong_s \mathfrak{t} \times T(\mathbb{Q}) \backslash T(\mathbb{A})^1 / K. \tag{3}
$$

In particular, any character on $T(\mathbb{Q})\backslash T(\mathbb{A})^1/K$ extends to a character on $T(\mathbb{Q})\backslash T(\mathbb{A})/K$.

Example 1 (restriction tori). Take $T = R_{\mathbb{Q}}^E \mathbb{G}_m$ where E is a finite Galois extension of Q. There are *n* characters χ_1, \ldots, χ_n generating X^*T_E which realize the *n* embeddings $T(\mathbb{Q}) = E^{\times} \to \mathbb{C}$ on the Q-points of T. The rational character lattice $M \cong X^*T$ is generated by the norm character $m_N \leftrightarrow N = \prod \chi_i$ where $N: T(\mathbb{A}) = \mathbb{A}_E^\times \to \mathbb{G}_m(\mathbb{A}) = \mathbb{A}^\times$. If $(e_i)_i \subset N_E$ is the dual basis to χ_1, \ldots, χ_n and $e_\infty := -\sum e_i$, then

$$
\ell(x) = \frac{1}{n} \log |N(x)|_{\mathbb{A}} e_{\infty}.
$$
\n(4)

A natural choice of splitting is given by

$$
s: \mathfrak{t} = \mathbb{R}e_{\infty} \to T(\mathbb{A}) = T(\mathbb{R}) \times T(\mathbb{A}_{fin})
$$

$$
e_{\infty} \mapsto s(e_{\infty}) = (q_{\infty}1_{\infty}; 1_{fin})
$$

where 1_{∞} (resp. 1_{fin}) is the identity of the group $T(\mathbb{R}) = E_{\mathbb{R}}^{\times}$ (resp. $T(\mathbb{A}_{\text{fin}})$) and $q_{\infty} = e^{d_{\infty}} = e$ (resp. e^2) if E is totally real (resp. not totally real).

Next we turn to characters of $T(\mathbb{Q})\backslash T(\mathbb{A})^1/K$. We will use non-standard but equivalent definitions for the regulator and class groups that are more convenient for the Poisson formula.

Definition 5. Let R denote the identity component of $T(\mathbb{Q})\setminus T(\mathbb{A})^1/K$ and let C denote the quotient of $T(\mathbb{Q})\backslash T(\mathbb{A})^1/K$ by R. We call R the *(level K) regulator group* and C the (level K) class group.

It is a fundamental fact $[17,$ Theorem 3.1.1 that R is compact and C is finite. The terminology reflects the fact that if $T = R_{\mathbb{Q}}^E \mathbb{G}_m$ and K is maximal, then R is the quotient in the trace-zero hyperplane of $E_{\mathbb{R}}$ by the lattice of logarithms of units and C is the ideal class group of E .

Lemma 3. Assume that $K_{\infty} = K_{\infty}^{\max}$. The class group C is isomorphic to $T(\mathbb{Q})\setminus T(\mathbb{A})/(T(\mathbb{R})K)$ (so our definition agrees with the standard definition of the class group of an algebraic group). If $\psi \in (T(\mathbb{Q}) \setminus T(\mathbb{A})^1 / K)^\vee$ factors through C, then the local archimedean component ψ_{∞} of ψ , given by $T(\mathbb{R}) \to T(\mathbb{Q}) \backslash T(\mathbb{A}) \to_s T(\mathbb{Q}) \backslash T(\mathbb{A})^1/K \to C \to \mathbb{C}^\times$, is trivial.

Proof. Let $C' = T(\mathbb{Q}) \setminus T(\mathbb{A}) / (T(\mathbb{R})K)$. The kernel of $T(\mathbb{Q}) \setminus T(\mathbb{A}) / K \to C'$ is $T(\mathbb{Q}) \setminus (T(\mathbb{Q})T(\mathbb{R})K) / K$. Observe that $T(\mathbb{R})K/K = T(\mathbb{R})/K_{\infty}$ is connected (Proposition [1\)](#page-6-2), and since the natural map

$$
T(\mathbb{R})K/K \to T(\mathbb{Q})\backslash (T(\mathbb{Q})T(\mathbb{R})K)/K
$$

12 ANDREW O'DESKY

is clearly surjective, we see that $T(\mathbb{Q})\setminus (T(\mathbb{Q})T(\mathbb{R})K)/K$ is also connected. It is also closed and open since it is the preimage of the identity in C' , so the above kernel is the identity component of $T(\mathbb{Q})\backslash T(\mathbb{A})/K$. The isomorphism [\(3\)](#page-10-1) identifies the identity component of $T(\mathbb{Q})\backslash T(\mathbb{A})/K$ with $\mathfrak{t}\times R$ so we have the exact sequence

$$
1 \longrightarrow \mathfrak{t} \times R \longrightarrow \mathfrak{t} \times T(\mathbb{Q}) \backslash T(\mathbb{A})^1 / K \longrightarrow_s C' \longrightarrow 1.
$$

Restricting this short exact sequence to the subgroup $\{0\} \times T(\mathbb{Q}) \setminus T(\mathbb{A})^1/K$ shows that $C \cong_s C'.$

Let w be an infinite place of E. The local component ψ_{∞} factors through the map

$$
T(\mathbb{R}) \to T(\mathbb{Q}) \backslash T(\mathbb{A}) \to_s T(\mathbb{Q}) \backslash T(\mathbb{A})^1 / K \to C
$$

which itself factors through $T(\mathbb{R}) \xrightarrow{\ell_w} \mathfrak{t}_w$ (Proposition [1\)](#page-6-2), but every homomorphism $\mathfrak{t}_w \to C$ is constant so ψ_{∞} is trivial.

2.3.2. The regulator group of level K. Let r_{∞} be the rank of N_w for any infinite place w of E and let r be the rank of N .

Definition 6. The subspace $t \subset t_w$ of Γ-invariants has a canonical subspace complement

$$
\mathfrak{t}_0=\bigoplus_{\psi\neq 1}((\mathfrak{t}_E)_\psi)^{\Gamma_w}
$$

coming from the Γ-isotypic decomposition of \mathfrak{t}_E . We call \mathfrak{t}_0 the *trace-zero subspace* of \mathfrak{t}_w .

The abelian subgroup

$$
T(\mathbb{Q}) \cap \left(T(\mathbb{R}) \times \prod_{v \text{ finite}} K_v\right)
$$

of $T(A)$ is finitely generated with rank $r_{\infty} - r$ [\[16,](#page-43-14) Thm. 4, p. 285] and canonically splits as

$$
\Upsilon \times \mu = T(\mathbb{Q}) \cap \left(T(\mathbb{R}) \times \prod_{v \text{ finite}} K_v \right)
$$

where $\mu = T(\mathbb{Q}) \cap K$ is its torsion subgroup. It is well-known that the natural map

$$
\Upsilon \to T(\mathbb{R}) \xrightarrow{\ell w} \mathfrak{t}_w
$$

embeds Υ as a (full-rank) lattice of the trace-zero subspace t_0 .

Remark 5. If D is Cartier and the canonical integral model is used to define D-integrality (cf. Remark [3\)](#page-8-0), then μ is independent of D and equal to $T(\mathbb{Q}) \cap K^{\max}$.

2.4. Fourier transform of the height zeta function. If $t \in T(\mathbb{Q}_v)$ and $s \in (\mathbb{C}^{\Sigma(1)})^{\Gamma}$, then we may write

$$
H_v(t,s) = q_v^{-\frac{1}{e_v}\varphi(\ell_w(t))}
$$

where w is any place of E lying over v (cf. [\(1\)](#page-7-1)); although $\ell_w(t)$ may depend on the choice of w over v, the quantity $H_v(t, s)$ only depends on v.

2.4.1. Haar measures. For each place v of $\mathbb Q$ let a place w of E lying over v be chosen and fixed. For each finite place v let $d\mu_v$ be the Haar measure on $T(\mathbb{Q}_v)$ for which the maximal compact subgroup has measure one. At the infinite place choose the Haar measure $d\mu_{\infty}$ on $T(\mathbb{R})$ for which $N_w \subset \mathfrak{t}_w$ is a unimodular lattice for the pushforward of $d\mu_{\infty}$ along $\ell_w: T(\mathbb{R}) \to \mathfrak{t}_w$. The product measure $d\mu = \prod_v d\mu_v$ is a Haar measure on the restricted product $T(A)$ and the Fourier transform of any factorizable integrable function $f = \otimes_v f_v \in$ $L^1(T(\mathbb{A}))$ is defined by

$$
\widehat{f}(\chi) = \int_{T(\mathbb{A})} f(x) \chi(x)^{-1} d\mu(x) = \prod_{v} \int_{T(\mathbb{Q}_v)} f_v(x) \chi_v(x)^{-1} d\mu_v(x).
$$

2.4.2. Local Fourier transforms: nonarchimedean case. Let v be a finite place of \mathbb{Q} . Write

$$
H_v(t, s, D) = H_v(t, s)1_v^D(t).
$$

Let K_v^D denote the subgroup of K_v^{\max} fixing the indicator function 1_v^D .

Lemma 4. Let $s \in (\mathbb{C}^{\Sigma(1)})^{\Gamma}$ and let $\varphi: \mathfrak{t} \to \mathbb{C}$ be the corresponding Σ -linear function (Propo-sition [2\)](#page-7-2). Let w be any fixed place of E lying over v and set $\Sigma_v = \Sigma_{\Gamma_w}$. For any character $\chi \in T(\mathbb{Q}_v)^\vee$ the integral

$$
\widehat{H_v}(\chi, -s, D) = \int_{T(\mathbb{Q}_v)} H_v(x, -s, D) \chi(x)^{-1} d\mu_v(x)
$$

converges absolutely to a holomorphic function of s in the region

$$
\{s \in (\mathbb{C}^{\Sigma(1)})^{\Gamma} : \text{Re}(s_e) > 0 \text{ for every } e \in \Sigma(1)\}.
$$

This holomorphic function admits a meromorphic continuation to a rational function in the variables $\{q_v^{\varphi(e)} : e \in \Sigma_v(1)\}$. Furthermore $\widehat{H}_v(\chi, -s, D)$ is identically zero if χ is nontrivial on K_v^D , and otherwise is given by

$$
\widehat{H_v}(\chi, -s, D) = \sum_{n \in T(\mathbb{Q}_v)/K_v^D} \chi(n)^{-1} 1_v^D(n) q_v^{\frac{1}{e_v} \varphi(\ell_w(n))}.
$$

Remark 6. The local Fourier transforms must be computed before restricting to the line in $(\text{Pic}^T Y) \otimes \mathbb{C}$ spanned by the line bundle determining the height of interest since $x \mapsto$ $H_v(x, -s, D)$ will generally not be integrable once some of the coordinates of s vanish, no matter how large and positive $\text{Re}(s_e)$ is for the other coordinates.

Remark 7. One aspect where the Poisson-style analysis of integral points differs from the analysis of rational points [\[3\]](#page-42-1) is that 1_v^D and thus $H_v(\chi, -s, D)$ may have compact support, cf. [\[6,](#page-43-6) Prop. 4].

Proof. If χ is nontrivial on K_v^D then the Fourier transform of $H_v(x, s)^{-1}1_v^D(x)$ vanishes by Schur's lemma. Now suppose $\chi|_{K_v^D} = 1$. Then the integrand is K_v^D -invariant and

$$
\int_{T(\mathbb{Q}_v)} H_v(x, -s, D)\chi(x)^{-1} d\mu_v(x) = \sum_{n \in T(\mathbb{Q}_v)/K_v^D} q_v^{\frac{1}{e_v} \varphi(\ell_w(n))} 1_v^D(n) \chi(n)^{-1}.
$$

The number of n in the sum which are equivalent modulo K_v^{max} is bounded by the index of K_v^D in K_v^{max} which is finite, so the sum is absolutely convergent since $\text{Re}(s_e) > 0$ for all $e.$ 2.4.3. Toric ideals. We amplify the last computation by considering all finite places at once. For this purpose we introduce some notation. Let K_{fin} be an arbitrary open compact subgroup of $T(\mathbb{A}_{fin})$ and set $K = K_{fin} \times K_{\infty}^{\max}$.

Definition 7 (toric ideals). The group of fractional ideals of T (of level K) is

$$
\mathcal{Q} = T(\mathbb{A}_{fin})/K_{fin}.
$$

Now assume K contains K^D . The set of D-integral ideals of T (of level K) is

$$
\mathcal{N} = \{ n \in \mathcal{Q} : 1^D(n) = 1 \}.
$$

A D-integral ideal $n \in \mathcal{N}$ is *principal* if $\psi(n) = 1$ for any class group character $\psi \in C^{\vee}$. The set of D-integral principal ideals is denoted

$$
\mathscr{P} = \{ n \in \mathcal{N} : \psi(n) = 1 \text{ for all } \psi \in C^{\vee} \}.
$$

For any $n \in \mathcal{Q}$ and Γ -invariant Σ -linear function $\varphi \leftrightarrow s \in (\mathbb{C}^{\Sigma(1)})^{\Gamma}$ we set

$$
n^{-s} := H(n, -s) = \prod_{p \in P_{\mathbb{Q}}^{\text{fin}}} p^{\frac{1}{e_p} \varphi(\ell_w(n_p))} \in \mathbb{C}^{\times}.
$$

Corollary 1. Let $\chi \in (T(\mathbb{A})/K^D)^{\vee}$ be a character. The finite part $\hat{H}_{fin}(\chi, -s, D)$ of \hat{H} is equal to

$$
\widehat{H}_{\text{fin}}(\chi, -s, D) = \sum_{n \in \mathcal{N}} \chi(n)^{-1} n^{-s} \tag{5}
$$

where N is the set of D-integral ideals of T of level K^D . There is a constant $\sigma > 0$ such that the right-hand side converges absolutely to a holomorphic function of s in the region $\{s \in (\mathbb{C}^{\Sigma(1)})^{\Gamma} : \text{Re}(s_e) > \sigma \text{ for every } e \in \Sigma(1)\}.$

Proof. For each place v choose a place w of E lying over v. The finite part of the Fourier transform of the height is an infinite product

$$
\prod_{p\in P^{\text{fin}}_{\mathbb{Q}}} \sum_{n\in T(\mathbb{Q}_p)/K_p^D} \chi_p(n)^{-1} 1_v^D(n) p^{\frac{1}{e_p}\varphi(\ell_w(n))}.
$$

For all but finitely many p we have $K_p^D = K_p^{\text{max}}$, $e_p = 1$, and p-factor (Proposition [1\)](#page-6-2)

$$
\sum_{n \in N_w} \chi_p(n)^{-1} 1_p^D(n) p^{\varphi(n)}.
$$

Let $|\cdot|$ be any norm on \mathfrak{t}_E . Then

$$
\left|\chi_p(n)^{-1}1_p^D(n)p^{\varphi(n)}\right| \le p^{-\rho t|n|}
$$

where $t = \min\{\text{Re}(s_e) : e \in \Sigma(1)\}\$ and $\rho > 0$ is some constant independent of p and n. Then

$$
\sum_{n \in N_w} \chi_p(n)^{-1} 1_p^D(n) p^{\varphi(n)} = 1 + O(p^{-\rho t |n_0|})
$$

for a minimally chosen n_0 and an implied constant independent of p. This shows that the infinite product defining $H_{fin}(\chi, -s, D)$ converges to a holomorphic function of s and may
be multiplied out into an absolutely convergent infinite sum, for some constant $\sigma > 0$. be multiplied out into an absolutely convergent infinite sum, for some constant $\sigma > 0$.

Remark 8. It is tempting to think of n^{-s} as $e^{\varphi(\ell(n))}$ but in general these are different quantities due to nonlinearity of φ . However if φ is linear on the cone generated by every element

 $\ell_w(n_p)$ log q_w

where $n_p \in T(\mathbb{Q}_p)$ is any element such that $\ell_w(n_p)$ is D-integral and $w \in P_E^{\text{fin}}$ lies over v , then $n^{-s} = e^{\varphi(\ell(n))}$. This occurs notably when φ is linear or when the D-integral region of \mathfrak{t}_E is Σ -convex.

Example 2 (toric ideals for restriction tori). We verify that our notions on toric ideals agree with the usual ones for restriction tori. Take $T = R_{\mathbb{Q}}^E \mathbb{G}_m$ where E is a finite Galois extension of Q. There is a short exact sequence with finite kernel

$$
1 \longrightarrow K^{\max}/(\mu \cdot K^D) \longrightarrow \mathcal{Q} \longrightarrow \{\text{fractional ideals of } E\} \longrightarrow 1.
$$

We may compactify T according to the usual fan of projective space \mathbb{P}_n to obtain a Severi– Brauer compactification^{[4](#page-14-0)} of T. Now take the boundary divisor to be $D = D_{\infty}$ corresponding to the unique Galois-fixed generator e_{∞} of the fan. This divisor is Cartier and we use the canonical integral model for the complement of D_{∞} (Remark [3\)](#page-8-0). Then $K^D = K^{\text{max}}$ and the D_{∞} -integral ideals of T of level $K = K^{\max}$ are precisely the (nonzero) ideals of O_E . An element $n \in \mathcal{N}$ is principal by our definition if $\psi(n) = 1$ for every class group character ψ . This is equivalent to requiring that the image [n] of n under $\mathscr{N} \to T(\mathbb{A}_{fin}) \to T(\mathbb{A}) \to_s$ $T(\mathbb{A})^1 \to T(\mathbb{Q}) \backslash T(\mathbb{A})^1 / K$ is in the identity component R. With the help of Example [1](#page-10-2) one finds that

$$
[n] = [N(I)^{\frac{d_{\infty}}{[E:\mathbb{Q}]}}\mathbb{1}_{\infty}; t_{\text{fin}}]
$$

where I is the ideal of O_E corresponding to n and $t_{fin} \in T(\mathbb{A}_{fin})$ is any element mapping to n. Regarding [n] as an element of $T(\mathbb{Q})\backslash T(\mathbb{A})/K$ via the natural inclusion $T(\mathbb{A})^1 \subset T(\mathbb{A})$ and using the description of the identity component of $T(\mathbb{Q})\backslash T(\mathbb{A})/K$ from the proof of Lemma [3](#page-10-3) shows that

$$
[N(I)^{\frac{d_{\infty}}{[E:\mathbb{Q}]} } 1_{\infty}; t_{\text{fin}}] \in T(\mathbb{Q}) \setminus (T(\mathbb{Q})T(\mathbb{R})K)/K \subset T(\mathbb{Q}) \setminus T(\mathbb{A})/K
$$

which is equivalent to $[N(I)^{\frac{d_{\infty}}{[E:\mathbb{Q}]}1_{\infty}}; t_{fin}] \in T(\mathbb{Q})T(\mathbb{R})K$. This shows that $t_{fin} \in T(\mathbb{Q})_{fin}K_{fin}$ which is the condition that I is principal in the usual sense. We may also evaluate n^{-s} for $n \in \mathcal{N}$. The D-integral region is Σ -convex so $n^{-s} = e^{\varphi(\ell(n))}$ (Remark [8\)](#page-14-1). Any $s \in (\mathbb{C}^{\Sigma(1)})^{\Gamma}$ satisfies $s_1 = \cdots = s_{[E:\mathbb{Q}]}$ and so $n^{-s} = N(I)^{-s_1}$ by integrality of I and [\(4\)](#page-10-4). The general quasi-split case is similar: if $T = R_{\mathbb{Q}}^{E_1} \mathbb{G}_m \times \cdots \times R_{\mathbb{Q}}^{E_r} \mathbb{G}_m$ then the elements n of N correspond to ideals $I_1 \times \cdots \times I_r$ of $O_{E_1} \times \cdots \times O_{E_r}$ and $n^{-s} = N(I_1)^{-s_{O_1}} \cdots N(I_r)^{-s_{O_r}}$ where s_{O_j} is the common value of s_e on the *j*th Galois orbit O_j of $\Sigma(1)$ corresponding to E_j .

2.4.4. Local Fourier transforms: archimedean case. In the nonarchimedean setting, the local Fourier transform has contributions from cones of every dimension whereas in the archimedean setting, the image of ℓ_w is \mathfrak{t}_w and only cones of maximal dimension have positive measure. Let w be an infinite place of E. Let r_{∞} be the rank of t_w . Write Σ_{∞} for the restricted fan $\Sigma_{\Gamma_w}.$

Proposition 3. Let $s \in (\mathbb{C}^{\Sigma(1)})^{\Gamma_w}$ and let $\varphi: \mathfrak{t}_w \to \mathbb{C}$ be the corresponding Σ -linear function (Proposition [2\)](#page-7-2). For any $e \in \Sigma_{\infty}(1)$ let $s_e = -\varphi(e)$. Assume $\text{Re}(s_e) > 0$ for all $e \in \Sigma_{\infty}(1)$. Let $\chi \in T(\mathbb{R})^{\vee}$ be a character. If χ is K_{∞} -ramified then $H_{\infty}(\chi, -s, D)$ is identically zero.

⁴This compactification is isomorphic to \mathbb{P}_n over \mathbb{Q} as a variety but not as a toric variety: any such isomorphism will not be equivariant for the toric actions.

If χ is K_{∞} -unramified, then there is a unique $m \in \mathfrak{t}_{w}^{\vee}$ such that $\chi(t) = e(\langle \ell_{w}(t), m \rangle)$ for all $t \in T(\mathbb{R})$, and we have that

$$
\widehat{H_{\infty}}(\chi, -s, D) = \left(\frac{1}{2\pi i}\right)^{r_{\infty}} \sum_{\tau \in \Sigma_{\infty}(r_{\infty})} [N_w : \mathbb{Z}\tau(1)] \prod_{e \in \tau(1)} \left(m(e) + \frac{s_e}{2\pi i}\right)^{-1}.
$$

Proof. Note that 1^D_∞ is identically one on $T(\mathbb{R})$ and H_∞ is $K_\infty = K_\infty^{\max}$ -invariant. If χ is ramified, then $\widetilde{H_{\infty}}(\chi, -s, D)$ vanishes by Schur's lemma for χ on K_{∞}^{\max} . If χ is unramified, then

$$
\widehat{H_{\infty}}(\chi, -s, D) = \int_{\mathfrak{t}_w} H_{\infty}(y, -s)e(-\langle y, m \rangle) d\mu_{\infty}(y) = \int_{\mathfrak{t}_w} e^{\varphi(y)} e(-\langle y, m \rangle) d\mu_{\infty}(y) \tag{6}
$$

Since Y is projective, the restricted fan Σ_{∞} is complete so

$$
\widehat{H_{\infty}}(\chi,-s,D)=\sum_{\tau \in \Sigma_{\infty}(r_{\infty})}\int_{\tau}e^{\varphi(y)}e(-\langle y,m\rangle)\,d\mu_{\infty}(y).
$$

Since Σ_{∞} is simplicial, the generators $\tau(1) = \{e_{j_1}, \ldots, e_{j_{r_{\infty}}}\}\$ of τ form a basis of \mathfrak{t}_w . Write $y = x_1 e_{j_1} + \cdots + x_{r_\infty} e_{j_{r_\infty}}$. Under the isomorphism $\mathbb{R}^{r_\infty} \to \mathfrak{t}_w$ determined by $\tau(1)$, the measure μ_{∞} pulls back to $[N_w : \mathbb{Z} \tau(1)] \mu_{\text{Lebesgue}}$. Then

$$
\int_{\tau} e^{\varphi(y)} e(-\langle y, m \rangle) d\mu_{\infty}(y) = [N_w : \mathbb{Z}\tau(1)] \int_{\mathbb{R}_{\geq 0}^{\tau_{\infty}}} e^{-x_1 s_{j_1} - \dots - x_{r_{\infty}} s_{j_{r_{\infty}}}} e(-\langle y, m \rangle) dx_1 \wedge \dots \wedge dx_{r_{\infty}}
$$

\n
$$
= [N_w : \mathbb{Z}\tau(1)] \prod_{e \in \tau(1)} \int_0^{\infty} e^{-x(s_e + 2\pi i \langle e, m \rangle)} dx = [N_w : \mathbb{Z}\tau(1)] \prod_{e \in \tau(1)} (s_e + 2\pi i \langle e, m \rangle)^{-1}.
$$

The integral converges if and only if $\text{Re}(s_e) > 0$ for all $e \in \Sigma_{\infty}(1)$.

2.5. Characters contributing to the expected principal term. Let P denote the quasisplit torus whose character lattice is $Pic^{T}(Y_{E}) \cong \mathbb{Z}^{\Sigma(1)}$. The natural map $M_{E} \to \mathbb{Z}^{\Sigma(1)}$ (Proposition [2\)](#page-7-2) induces an isogeny

 $\gamma\colon P\to T$

of algebraic tori. The summands of our formula for $Z(s)$ are indexed by automorphic characters of T on $R \times C$, and the summands which contribute to the expected principal term are those characters which become trivial after pulling back by γ^* to P. The following essentially trivial lemma will ensure that every such character is trivial on the regulator group R.

Lemma 5. Let $\gamma: T' \to T$ be a surjective homomorphism of algebraic tori and let $\gamma^*: T(\mathbb{A})^\vee \to T(\mathbb{A})$ $T'(\mathbb{A})^\vee$ denote the pullback homomorphism. Then there is a positive integer $C(\gamma)$ annihilating the kernel of γ^* .

Proof. Recall the category of diagonalizable algebraic groups which split over E/\mathbb{Q} is equivalent to the category of finite-type $\mathbb{Z}\Gamma$ -modules. Thus the cokernel of γ is representable as a finite group scheme. If χ is a unitary character of $T(\mathbb{A})$ for which $\chi \circ \gamma = 1$, then χ factors through the cokernel. Thus χ is annihilated by $C(\gamma) = |\text{coker}(\gamma)(\overline{\mathbb{Q}})|$.

Remark 9. [\[3,](#page-42-1) Theorem 3.1.1], [\[2,](#page-42-3) Proposition 6.4], and [\[4,](#page-43-0) p. 19] assert that ker γ^* is dual to the obstruction group to weak approximation^{[5](#page-15-1)} of T and cite [\[10\]](#page-43-10). In the later paper [\[8\]](#page-43-2) this assertion appears as Lemma 3.8.1, and their explanation cites the proof of [\[21,](#page-43-15)

 5 See [\[21,](#page-43-15) Theorem 6] for a definition of this group.

Theorem 6]. We were unable to see how their assertion follows from either [\[10\]](#page-43-10) or the proof of [\[21,](#page-43-15) Theorem 6], so we will not make use of this description of ker γ^* . For our purposes, it is enough to know that ker γ^* is annihilated by $C(\gamma)$.

3. Iterated residues of multiple Dirichlet series

3.1. The Poisson summation formula. Let B be a locally compact abelian group with Haar measure db. Let $f \in L^1(B)$. The Fourier transform of f given by

$$
\widehat{f}(\chi) = \int_B f(b) \chi(b)^{-1} \, db
$$

converges and defines a continuous function on B^{\vee} .

Now let A be a closed subgroup of B and let $A^{\perp} \subset B^{\vee}$ denote the subgroup of characters on B that are trivial on A . The general Poisson summation formula — whose proof is essentially the classical proof for $\mathbb{Z} \subset \mathbb{R}$ — says that if $\widehat{f}|_{A^{\perp}} \in L^1(A^{\perp})$ then

$$
\int_{A} f(ab) da = \int_{A^{\perp}} \widehat{f}(\chi) \chi(b) d\chi \tag{7}
$$

for a suitably normalized Haar measure on A^{\perp} and all $b \in B$ away from a subset of measure zero [\[11,](#page-43-16) Theorem 4.4.2, p. 105].

We will apply the Poisson formula for the discrete subgroup of rational points in the adelic space of the torus:

$$
A = T(\mathbb{Q}) \subset B = T(\mathbb{A}).
$$

This leads to a difficult integral over the automorphic spectrum $T(\mathbb{Q})^{\perp} = (T(\mathbb{A})/T(\mathbb{Q}))^{\vee}$ of T that was first studied in general in $[5]$. Whenever the torus T admits nonconstant characters defined over Q, the domain of integration is noncompact; this has an interesting effect on the integral which is not limited to expected difficulties concerning convergence. We will apply the method in [\[14\]](#page-43-4) to obtain an exact expression for this integral in terms of higher-dimensional residues. This improves on the analysis in [\[5\]](#page-43-1) which only approximated this integral.

Remark 10. This use of Poisson summation is different from Tate's thesis where Poisson summation is applied for the subgroup $E \subset A_E$ in the *additive* adeles; unlike the additive group A_E , the multiplicative group $T(A)$ is not self-dual. On the spectral side, the Poisson formula in the multiplicative setting involves an integral over Hecke characters on T of general level. It is better regarded as a special case of the Arthur–Selberg trace formula.

3.2. The method of iterated residues. The Poisson formula will lead to the following integral:

$$
\int_{\mathfrak{t}^\vee} \chi_m(n)^{-1} \widehat{H_\infty}(\xi \chi_m, -s) \, dm
$$

where $n \in \mathcal{N}$ is a toric ideal, $\chi_m : \mathfrak{t}^\vee \to \mathbb{C}^\times$ is the unitary character $\chi_m(n) = e(m(n)),$ and $\xi \in (T(\mathbb{Q}) \setminus T(\mathbb{A})/K)^\vee$ is an automorphic character of level K. We will evaluate this integral using the residue formula from [\[14,](#page-43-4) Theorem 1]. This formula proceeds by choosing a polyhedron Π and iteratively applying Cauchy's residue formula by deforming along the rays of the imaginary part of Π one at a time. This expresses the integral as a sum over *iterated residues* which are indexed by *affine flags* cut out by the affine hyperplanes where $H_{\infty}(\chi_m, -s)$ is singular, and the problem is to determine which flags contribute a residue.

This is answered in [\[14,](#page-43-4) Theorem 1] using sign conditions on the minors of the defining matrices for each flag.

3.2.1. Minors. For an $r \times r$ real matrix $J = (a_{ij})$, we consider the kth leading principal minor

$$
p_k = \det \begin{bmatrix} a_{11} & \cdots & a_{1k} \\ \vdots & \ddots & \vdots \\ a_{k1} & \cdots & a_{kk} \end{bmatrix} \qquad (k \in \{1, \ldots, r\}),
$$

the $k \times k$ minor

$$
q_{k\ell} = \det \begin{bmatrix} a_{11} & \cdots & a_{1,k-1} & a_{1\ell} \\ \vdots & \ddots & \vdots & \vdots \\ a_{k1} & \cdots & a_{k,k-1} & a_{k\ell} \end{bmatrix} \qquad (\ell \in \{k+1, \ldots, r\}),
$$

and the $(k-1) \times (k-1)$ minor

$$
r_{jk} = \det \begin{bmatrix} a_{11} & \cdots & a_{1,k-1} \\ \vdots & \ddots & \vdots \\ a_{j-1,1} & \cdots & a_{j-1,k-1} \\ a_{j+1,1} & \cdots & a_{j+1,k-1} \\ \vdots & \ddots & \vdots \\ a_{k1} & \cdots & a_{k,k-1} \end{bmatrix} \qquad (j \in \{1, \ldots, k-1\}).
$$

Let J be a $k \times r$ real matrix. We say J is stable if

 $p_1, \ldots, p_k > 0$ and $(-1)^{\ell-j} r_{j\ell} \ge 0$ for all $1 \le j < \ell \le k$.

We say J is *compatible* if either it is not stable or

$$
q_{j\ell} \le 0 \text{ for all } 1 \le j < \ell \le k.
$$

3.2.2. Polyhedra. By a polyhedron (with boundary \mathfrak{t}^{\vee}) we mean a closed subset $\Pi \subset \mathfrak{t}^{\vee}_{\mathbb{C}}$ and a linear isomorphism $\mathfrak{t}^\vee \stackrel{\sim}{\to} \mathbb{R}^r$ such that Π is identified with the product of the closed upper half-planes in \mathbb{C}^r under the induced linear isomorphism $\mathfrak{t}_{\mathbb{C}}^{\vee} \stackrel{\sim}{\to} \mathbb{C}^r$. We consider the defining map $\mathfrak{t}^{\vee} \xrightarrow{\sim} \mathbb{R}^r$ to be part of the data of Π . Let $\Pi = \mathfrak{t}^{\vee} + i\Theta$ be a polyhedron. Its imaginary cone $\Theta = \mathbb{R}_{\geq 0} \langle v_1, \ldots, v_r \rangle$ is generated by the distinguished basis (v_1, \ldots, v_r) of \mathfrak{t}^{\vee} determined by the defining map. Let z be the dual basis to (v_1, \ldots, v_r) . Let

$$
\gamma : \mathfrak{t}_{\mathbb{C}}^{\vee} \supset H_1 \supset H_1 \cap H_2 \supset \cdots \supset H_1 \cap \cdots \cap H_r
$$

be a flag determined by a collection $H = (H_1, \ldots, H_r)$ of linearly independent singular hyperplanes with respective defining forms $e_1, \ldots, e_r \in E$. Let $J = J_{H,\Pi}$ be the Jacobian matrix

$$
J = \frac{\partial e}{\partial z} = \begin{bmatrix} e_1(v_1) & \cdots & e_1(v_r) \\ \vdots & \ddots & \vdots \\ e_r(v_1) & \cdots & e_r(v_r) \end{bmatrix}.
$$

We say H is Π -stable (resp. Π -compatible) if J is stable (resp. compatible).

3.2.3. The residue formula. Let $\gamma: \mathfrak{t}_\mathbb{C}^{\vee} \supset H_1 \supset H_1 \cap H_2 \supset \cdots \supset H_1 \cap \cdots \cap H_k$ be a flag. We write itres_z[ω , γ] for the iterated residue of the form ω along the poles prescribed by the flag $γ$, cf. [\[14,](#page-43-4) §1]. This is a meromorphic top-degree form on $γ(k)$ defined as follows. We write

$$
res_{z_1}[g\,dz_1,\gamma(1)] = res_{z_1}[g\,dz_1,\gamma(1)](z_2,\ldots,z_r)
$$

for the residue of $g dz_1$ with respect to z_1 if there exists $z_1^* \in \mathbb{C}$ such that the vector $v \in V_{\mathbb{C}}$ with coordinates $z(v) = (z_1^*, z_2(v), \ldots, z_r(v))$ is in $\gamma(1)$. If no such z_1 exists, then we set this equal to zero. This recursively defines the *iterated residue of* ω along γ :

$$
itres_z[\omega, \gamma](z_{k+1},\ldots,z_r) = \operatorname{res}_{z_k}[\cdots \operatorname{res}_{z_2}[\operatorname{res}_{z_1}[g\,dz_1,\gamma(1)]\,dz_2,\gamma(2)]\cdots dz_k,\gamma(k)],
$$

Let ω be a meromorphic r-form on $\mathfrak{t}_\mathbb{C}^\vee$ which is regular on the complement of an affine hyperplane arrangement $\mathfrak{t}_{\mathbb{C}}^{\vee} - \cup_{e \in E} H_e$ where $E \subset \mathfrak{t}^{\vee}$ is a finite set of nonzero real linear forms and $H_e = \{m \in \mathfrak{t}_{\mathbb{C}}^{\vee} : e(m) + \frac{s_e}{2\pi i} = 0\} \subset \mathfrak{t}_{\mathbb{C}}^{\vee}$ for some parameters s_e with positive real part.

Theorem 3 ([\[14,](#page-43-4) Theorem 1]). If ω admits an iterated residue expansion along the polyhedron Π and every singular flag of ω is Π -compatible, then

$$
\frac{1}{(2\pi i)^r} \int_V \frac{h(z) dz}{g_1(z) \cdots g_R(z)} = \sum_{\gamma \in Z_*} \text{itres}_z[\omega, \gamma]
$$

where $Z_* \subset \{ \gamma \in Z : \gamma(r) \in \Pi \}$ is the subset of flags cut out by a collection of Π -stable singular hyperplanes.

3.2.4. Transfer maps and poles. Let $\pi: \mathfrak{t}_E \to \mathfrak{t}$ denote the linear projection map given by restriction to \mathfrak{t}^{\vee} . Recall that \mathfrak{t}_{w} has a canonical subspace decomposition

$$
\mathfrak{t}_w=\mathfrak{t}\oplus \mathfrak{t}_0.
$$

If $E \subset \mathfrak{t}_w$ is an r-subset whose projection πE is a basis of \mathfrak{t} , then it can be used to transfer linear forms on \mathfrak{t}_w to linear forms on \mathfrak{t} . More precisely, for such a subset E we define a linear map

$$
\mu_E\colon \mathfrak t_w^\vee \to \mathfrak t^\vee
$$

by requiring that

$$
(\mu_E m)(\pi e) = m(e) \text{ for all } e \in E.
$$

Equivalently, the projection π restricts to a bijection of cones

$$
\mathfrak{t}_w \supset \mathbb{R}\langle E \rangle \xrightarrow{\pi} \mathbb{R}\langle \pi E \rangle \subset \mathfrak{t},
$$

and $\mu_E m$ is the unique linear map on t corresponding to $m|_{\mathbb{R}(E)}$ under this identification.

Remark 11. The transfer map μ_E equals the natural restriction map $\mathfrak{t}^\vee_w \to \mathfrak{t}^\vee$ if and only if $E \subset \mathfrak{t}.$

Transfer maps help with analyzing the poles of H_{∞} . Let $s \in (\mathbb{C}^{\Sigma(1)})^{\Gamma}$, and let $\xi \in R^{\vee}$. Let φ be the Σ -linear map corresponding to s. Let the exponent $m_{\xi} \in \mathfrak{t}_0^{\vee}$ be determined by the identity $\xi(t_{\infty}) = e(\langle m_{\xi}, \ell_{w}(t_{\infty}) \rangle)$ for $t_{\infty} \in T(\mathbb{R}) \subset T(\mathbb{A})$. The next lemma determines the poles and their variation with ξ . We omit the easy proof.

Lemma 6. Let $E \subset \Sigma_w(1)$ be any Σ -convex r-subset whose projection πE is a basis of t. Say $\tau \in \Sigma_w(r_w)$ is a cone containing E. The form $\widehat{H_\infty}(\xi \chi_m, -s)$ dm is singular along the hyperplanes $H_e(\xi) = \{m \in \mathfrak{t}_{\mathbb{C}}^{\vee} : e(m) + e(m_{\xi}) + \frac{s_e}{2\pi i} = 0\}$ for $e \in E$ and we let $m_{E\xi} =$ $\bigcap_{e \in E} H_e(\xi)$ denote their intersection. Then $m_{E_1} \in \mathfrak{t}_\mathbb{C}^{\vee}$ is equal to

$$
m_{E1} = (2\pi i)^{-1} \mu_E(\varphi|_{\tau})
$$

and

$$
m_{E\xi}=m_1-\mu_E(m_\xi)
$$

where $\varphi|_{\tau} \in \mathfrak{t}_{w}^{\vee} \otimes \mathbb{C}$ is the unique linear map agreeing with φ on τ .

3.3. Residual functions. Fix an archimedean place w of E . Let dm be a generator of $\wedge^r {\mathfrak t}^\vee$ for which the lattice M is unimodular. Let $z = (z_1, \ldots, z_r)$ be any basis of t. For each singular flag $\gamma : H_1 \supset H_1 \cap H_2 \supset \cdots \supset H_1 \cap \cdots \cap H_r$ with defining equations $H_k = \{m \in$ $\mathfrak{t}_{\mathbb{C}}^{\vee} : e_k(m) + \frac{s_k}{2\pi i} = 0$, let $J_{\gamma} = \partial e/\partial z$ be the Jacobian of the map $(e_1, \ldots, e_r) : \mathfrak{t}^{\vee} \to \mathbb{R}^r$ with respect to z. Let $B \subset GL_r(\mathbb{C})$ denote the subgroup of upper-triangular matrices. Let $\mathbb{C}_+^{\Sigma(1)} \subset \mathbb{C}^{\Sigma(1)}$ denote the half-space with positive real parts.

Definition 8. Let $s \in (\mathbb{C}_{+}^{\Sigma(1)})^{\Gamma_w}$. Let $\omega(m) = \widehat{H_{\infty}}(\chi_m, -s) dm$, a meromorphic r-form on \mathfrak{t}^{\vee} . For any singular flag γ whose Jacobian J_{γ} is in $B^{T}B$ we set

$$
R_{\gamma}(s) = (2\pi i)^{r} \text{ itres}_{z}[\omega, \gamma].
$$

The condition that $J_{\gamma} \in B^{T}B$ ensures that R_{γ} is a rational function of $s \in (\mathbb{C}^{\Sigma(1)})^{\Gamma_{w}}$ which is independent of z (cf. [\[14,](#page-43-4) Prop. 4]). Note that whether or not J_{γ} is in $B^T B$ does not depend on s nor the particular choice of defining equations for γ .

Remark 12. It is important to allow for $s \in (\mathbb{C}^{\Sigma(1)})^{\Gamma_w}$ and not just $s \in (\mathbb{C}^{\Sigma(1)})^{\Gamma}$ even though in the end we are really only concerned with the values of the height zeta function on $(\mathbb{C}^{\Sigma(1)})^{\Gamma}$. In the Poisson formula each residual function is summed over the lattice of local exponents m_{ξ} of regulator characters $\xi \in R^{\vee}$ in $\mathfrak{t}_0^{\vee} \subset (\mathbb{C}^{\Sigma(1)})^{\Gamma_w}$ which are not in $(\mathbb{C}^{\Sigma(1)})^{\Gamma}$ if $\xi \neq 1$.

Fix an $s \in (\mathbb{C}^{\Sigma(1)})^{\Gamma_w}$. Let z be the terminal point of a singular flag of $H_{\infty}(\chi_m, -s)$ dm. If there are distinct r-subsets $E, E' \subset \Sigma_w(1)$ such that $\bigcap_{e \in E} H_e = \{z\} = \bigcap_{e' \in E'} H_{e'}$ then z is ambiguous, otherwise z is simple.

Lemma 7. There is a dense open subset $U \subset (\mathbb{C}_{+}^{\Sigma(1)})^{\Gamma_w}$ such that for any fixed $s \in U$, the terminal point of each singular flag of $\widehat{H_{\infty}}(\chi_m, -s)$ dm is simple. Thus, for any s in this open set, each singular flag γ in $\mathfrak{t} \vee \otimes \mathbb{C}$ is cut out by a unique ordered r-subset $E_{\gamma} \subset \Sigma_w(1)$. Let z_{γ} denote the terminal point of γ and set $s_{\gamma} = -2\pi i z_{\gamma}$. The residual function R_{γ} for such a flag γ is given by

$$
R_{\gamma}(s) = \sum_{\substack{\tau \in \Sigma_w(r_{\infty}) \\ E_{\gamma} \subset \tau(1)}} \frac{[N_w : \mathbb{Z}\tau(1)]}{[N : \mathbb{Z}\pi E_{\gamma}]} \prod_{\substack{e \in \tau(1) \\ e \notin E_{\gamma}}} \frac{1}{(s - s_{\gamma})_e}.
$$
 (8)

Now consider an arbitrary parameter $s \in (\mathbb{C}_+^{\Sigma(1)})^{\Gamma_w}$ and a singular flag γ of $\widehat{H_\infty}(\chi_m, -s)$ dm. Let F be the set of singular flags of $H_{\infty}(\chi_m, -s')$ dm which converge to γ when $s' \to s$ through values in U. Then

$$
R_{\gamma}(s) = \lim_{\substack{s' \to s \\ s' \in U}} \sum_{\gamma' \in F} R_{\gamma'}(s'). \tag{9}
$$

In the last expression it suffices to sum over those $\gamma' \in F$ whose defining subset $E_{\gamma'}$ is Σ-convex.

Proof. Each singular hyperplane $H_e = \{e(m) + \frac{s_e}{2\pi i} = 0\}$ is associated to an independent parameter s_e ($e \in \Sigma_w(1)$) as s varies in $(\mathbb{C}^{\Sigma(1)})^{\Gamma_w}$ so the first assertion is clear. Now fix $s \in U$ and let γ be a singular flag of $H_{\infty}(\chi_m, -s) dm$. Since the pole z_{γ} is simple, each step of the iterated residue (starting with $\overline{H_{\infty}}(\chi_m, -s) dm$) can be computed by eliminating the unique singular factor and restricting the remaining quantities to the next step of γ . This process results in the first formula. The general case follows from continuity of iterated residues in the parameters s (cf. [\[20,](#page-43-17) $\S II.5.4$] and [\[14,](#page-43-4) Prop. 4]). For the last statement, if $E_{\gamma'}$ is not Σ -convex then $R_{\gamma'} \equiv 0$ since the indexing set for the sum in [\(8\)](#page-19-3) is empty.

Lemma 8. Each residual function is invariant under translation by $s \mapsto s + 2\pi i m$ for any linear form $m \in \mathfrak{t}^{\vee} \otimes \mathbb{C}$. Thus, each residual function factors through the quotient $(\mathbb{C}^{\Sigma(1)})^{\Gamma} \to (\mathbb{C}^{\Sigma(1)})^{\Gamma}/(\mathfrak{t}^{\vee} \otimes \mathbb{C}) = (\text{Pic } Y) \otimes \mathbb{C} \dashrightarrow \mathbb{C}$ to a rational function on the Picard group.

Proof. We claim $s - s_\gamma$ is invariant under the given translation. Let $\pi: \mathfrak{t}_E \to \mathfrak{t}$ denote the linear projection map given by restriction to \mathfrak{t}^{\vee} . The pole z_{γ} is the intersection of the hyperplanes $H_e \subset \mathfrak{t}_\mathbb{C}^{\vee}$ for $e \in E_\gamma$ so it is the unique linear map on $\mathfrak{t}_\mathbb{C}$ satisfying $z_\gamma(\pi e) =$ $e(z_{\gamma}) = -\frac{s_e}{2\pi i}$ for each $e \in E_{\gamma}$. From this one easily finds that, if z'_{γ} denotes the terminal point of the translation of γ , then $z'_{\gamma} = z_{\gamma} - m$ and $s'_{\gamma} = s_{\gamma} + 2\pi im$. This proves invariance of R_{γ} .

Now pass to Γ-invariants of the first row from the diagram in Proposition [2](#page-7-2) to obtain the long exact sequence $0 \longrightarrow M \longrightarrow \text{Div}_T Y \longrightarrow \text{Pic} Y \longrightarrow H^1(\Gamma, M_E) \longrightarrow \cdots$. Since $H^1(\Gamma, M_E)$ is finite the complexified Picard group is isomorphic to $(\mathbb{C}^{\Sigma(1)})^{\Gamma}/(\mathfrak{t}^{\vee}\otimes\mathbb{C})$. \Box

3.3.1. Nonvanishing of residual forms. We next consider a class of \mathbb{P}_n -like toric varieties whose Galois action satisfies a convexity condition.

Definition 9. The Galois action on the fan Σ is *convex* if the Γ-orbit Γe is Σ-convex for every ray generator $e \in \Sigma_w(1)$. Consider a Σ -convex r-subset $E \subset \Sigma_w(1)$ whose restriction to \mathfrak{t}^\vee is linearly independent. If the Galois action on Σ is convex and the set $\pi E = {\pi e : e \in E}$ is Σ -convex for every such E then we say the Galois action on Σ is *strongly convex*.

Remark 13. Twists of projective space, and more generally their quotients by finite subgroups of the torus, have fans with strongly convex Galois action.

Example 3. The depicted fan with Galois action determines a complete toric surface over Q with cyclic quotient singularities whose open torus splits over a quadratic extension. The Galois action is not convex.

Proposition 4. Assume the Galois action is strongly convex. Let $\sigma \in \Sigma(r)$ and let $\gamma \in Z_{\sigma}$. Then

$$
R_{\gamma}(s) > 0
$$

for any $s \in \text{Pic } Y \otimes \mathbb{R}$ in the interior of the cone of effective divisors. In particular, if L is a big line bundle on Y, then $R_{\gamma}(t[L]) > 0$ for any $t > 0$.

Proof. The rational function $R_{\gamma}(s)$ is regular for any $s \in (\mathbb{C}_{+}^{\Sigma(1)})^{\Gamma_w}$ since the iterated residue is always well-defined. The iterated residue does not vanish identically since γ may be defined by a Π_{σ} -stable Jacobian. In view of [\(9\)](#page-19-1) it suffices to consider the case that the terminal point of γ is simple. It is clear from [\(8\)](#page-19-3) that the sign of the real number $R_{\gamma}(s)$ can only change at a singularity of R_{γ} , so it suffices to show that $R_{\gamma}(s) > 0$ for a single s in the interior of the effective cone. Since Y is projective, it admits an ample line bundle L. Replacing L with a positive multiple of itself, we may assume it admits a T -linearization since $H^1(\Gamma, M_E)$ is finite. Let s be the element of $(\mathbb{Z}^{\Sigma(1)})^{\Gamma} \subset (\mathbb{C}^{\Sigma(1)})^{\Gamma}$ corresponding to L with its T-linearization (Proposition [2\)](#page-7-2). Let φ be the corresponding Σ -linear function. Since L is ample it is globally generated (not true in general for proper varieties but true for proper toric varieties [\[12,](#page-43-11) p. 70]). It is well-known that L is globally generated if and only if φ is upper convex $[12, p. 68]$, so

$$
(\varphi - \varphi|_{\tau})(v) \le 0 \tag{10}
$$

for any cone $\tau \in \Sigma$ and $v \notin \tau$ where $\varphi|_{\tau}$ is a linear map on \mathfrak{t}_E agreeing with φ . The terminal point of $\gamma = \gamma(s)$ is $z_{\gamma} = (2\pi i)^{-1} \mu_{E_{\gamma}}(\varphi_{|\tau})$ (Lemma [6\)](#page-18-1). Since the Galois action is convex we have $\mu_{E_{\gamma}}(\varphi|_{\tau}) = \varphi|_{\pi E_{\gamma}}$ where $\varphi|_{\pi E_{\gamma}} \in \mathfrak{t}^{\vee} \otimes \mathbb{C}$ is the unique linear map agreeing with φ on the basis πE_γ . Since the Galois action is strongly convex, $\varphi|_{\pi E_\gamma} = \varphi|_{\sigma}$ where $\sigma = \langle \pi E_\gamma \rangle \in \Sigma_{\Gamma}(r)$. In particular, $s_{\sigma} = s_{\gamma}$ only depends on σ . Now let $\tau \in \Sigma_w(r_w)$ be a cone containing E_{γ} and let $e \in \tau(1) - E_{\gamma}$. Since the Galois action is convex, the ray $\tau_e = \mathbb{R}_{\geq 0} \pi e$ is a cone in Σ_{Γ} by Lemma [1.](#page-9-2) Suppose the ray $\mathbb{R}_{\geq 0}\pi e$ were contained in σ . Then it would be an edge of σ , and so $\pi e = \lambda \pi e'$ for some $e' \in E_{\gamma}$ and $\lambda > 0$. This would imply e and e' are in the same Galois orbit. Since s is Γ-invariant, this would mean that $e + \frac{s_e}{2\pi i}$ and $e' + \frac{s_{e'}}{2\pi i}$ define the same hyperplane in $\mathfrak{t} \otimes \mathbb{C}$, which contradicts simplicity of γ . We conclude that πe is not contained in σ by simplicity of γ . Now the desired positivity

$$
(s - s_{\gamma})_e = -(\varphi - \varphi|_{\sigma})(e) = -(\varphi - \varphi|_{\sigma})(\pi e) > 0
$$

for e follows from [\(10\)](#page-21-1) and regularity of R_{γ} .

3.3.2. Estimates on residual forms. In this section we show that the form $\omega = \widehat{H_{\infty}}(\chi_m, -s)$ dm admits an iterated residue expansion for any fixed $s \in (\mathbb{C}^{\Sigma(1)})^{\Gamma_w}$ with positive real parts.

Lemma 9. Let $\gamma: \mathfrak{t}_\mathbb{C}^{\vee} \supset H_1 \supset \cdots \supset H_1 \cap \cdots \cap H_k$ be a k-step singular flag in $\mathfrak{t}_\mathbb{C}^{\vee}$. Then

$$
|\mathsf{itres}_z[\omega,\gamma]| \ll \frac{1}{|m|} \sum_{\tau \in \Sigma_{\infty}(r_{\infty})} \prod_{\substack{e \in \tau(1) \\ H_e \notin \{H_1, \dots, H_k\}}} \left| m(e) + \frac{s_e}{2\pi i} \right|^{-1} \qquad (m \in \mathfrak{t}_{\mathbb{C}}^{\vee}).
$$

Proof. We use part of the proof of [\[3,](#page-42-1) Prop. 2.3.2]. Recall the earlier formula [\(6\)](#page-15-2) for $\widehat{H_{\infty}}$:

$$
\widehat{H_{\infty}}(\chi_m, -s) = \int_{\mathfrak{t}_w} e^{\varphi(y)} e(-\langle y, m \rangle) d\mu_{\infty}(y).
$$

$$
\Box
$$

Let (v_1, \ldots, v_{r_w}) be any basis for \mathfrak{t}_w and write a general element of \mathfrak{t}_w as $y = y_1v_1 + \cdots + y_{r_w}v_{r_w}$. Integrating by parts shows that for any $j \in \{1, \ldots, r_w\}$ and $m \notin \text{ker } v_j$,

$$
\widehat{H_{\infty}}(\chi_m, -s) = \int_{\mathfrak{t}_w} e^{\varphi(y)} e(-\langle y, m \rangle) d\mu_{\infty}(y) = \frac{1}{-2\pi i v_j(m)} \int_{\mathfrak{t}_w} \frac{\partial}{\partial y_j} (e^{\varphi(y)}) e(-\langle y, m \rangle) d\mu_{\infty}(y).
$$

The function $\varphi(y)$ is linear on each cone so the formula $\frac{\partial}{\partial y_j}(e^{\varphi(y)}) = \frac{\partial \varphi}{\partial y_j}e^{\varphi(y)}$ is valid and well-defined almost everywhere. Splitting up the integral over cones obtains

$$
\widehat{H_{\infty}}(\chi_m, -s, D) = \frac{1}{-2\pi i v_j(m)} \sum_{\tau \in \Sigma_{\infty}(r_{\infty})} \frac{\partial \varphi|_{\tau}}{\partial y_j} \int_{\tau} e^{\varphi(y)} e(-\langle y, m \rangle) d\mu_{\infty}(y)
$$

=
$$
\frac{1}{-2\pi i v_j(m)} \left(\frac{1}{2\pi i}\right)^{r_w} \sum_{\tau \in \Sigma_{\infty}(r_{\infty})} [N_w : \mathbb{Z}\tau(1)] \frac{\partial \varphi|_{\tau}}{\partial y_j} \prod_{e \in \tau(1)} \left(m(e) + \frac{s_e}{2\pi i}\right)^{-1}.
$$

The nonzero form v_j cannot vanish identically on any of the affine hyperplanes H_1, \ldots, H_k so we may evaluate the residual form with this expression. Thus itres_z[ω , γ] is a linear combination of forms (with coefficients depending only on Σ)

$$
\frac{1}{v_j(m)} \prod_{\substack{e \in \tau(1) \\ H_e \notin \{H_1, \dots, H_k\}}} \left(m(e) + \frac{s_e}{2\pi i} \right)^{-1} dm,
$$

and

$$
|\mathsf{itres}_z[\omega,\gamma]| \ll \frac{1}{|v_j(m)|} \sum_{\tau \in \Sigma_{\infty}(r_{\infty})} \prod_{\substack{e \in \tau(1) \\ H_e \notin \{H_1,\ldots,H_k\}}} \left| m(e) + \frac{s_e}{2\pi i} \right|^{-1}
$$

This holds for any $j \in \{1, ..., r_w\}$ which proves the claimed formula.

Proposition 5. The r-form $\omega = \widehat{H_{\infty}}(\chi_m, -s)$ dm admits an iterated residue expansion with respect to any polyhedron Π with boundary \mathfrak{t}^{\vee} . In particular, $\chi_m(n)^{-1}\omega$ admits an iterated residue expansion with respect to Π if $\chi_m(n)^{-1}$ is bounded on Π .

Note that $\chi_m(n)^{-1}$ is bounded on Π if and only if $-\ell(n) \in \Theta^{\vee}$ where $\Theta = \text{Im}(\Pi)$.

Proof. Let $\gamma : \mathfrak{t}^{\vee} \supset H_1 \supset \cdots \supset H_1 \cap \cdots \cap H_k$ $(1 \leq k < r)$ be a k-step singular flag. Let S denote the subset of generators $e \in \Sigma_w(1)$ whose corresponding hyperplane H_e is not in $\{H_1,\ldots,H_k\}$. For any $\delta > 0$ let $T_\delta \subset \gamma(k)$ denote the open subset of elements $m \in \gamma(k)$ satisfying

$$
\left| e(m) + \frac{s_e}{2\pi i} \right| > \delta \tag{11}
$$

for all $e \in S$.

We claim that

$$
itres_z[\omega, \gamma] = O_{\delta,s}(|m|^{-2})
$$

as $m \in T_\delta \subset \gamma(k)$ tends to infinity. The convergence of the iterated residue expansion follows immediately from this estimate. Let $\pi_k : \mathfrak{t}_{\mathbb{C}} \to \mathfrak{t}_{\mathbb{C}}/\langle e_1, \ldots, e_k \rangle$ be the natural quotient map. Suppose a cone τ in $\Sigma_{\infty}(r_{\infty})$ has a summand in H_{∞} which contributes to itres_z[ω , γ]. The generators in $\tau(1)$ which are not in S are in the subspace $\langle e_1, \ldots, e_k \rangle$. Since τ is simplicial, the image of $\tau(1) \cap S$ under π_k will span the quotient space $\mathfrak{t}_\mathbb{C}/\langle e_1, \ldots, e_k \rangle$. Thus

$$
\max_{e \in \tau(1) \cap S} \left| e(m) + \frac{s_e}{2\pi i} \right| \gg_s |m| \tag{12}
$$

.

for $m \in \gamma(k)$. With the help of [\(11\)](#page-22-0) and [\(12\)](#page-22-1) we have

$$
\prod_{e \in \tau(1) \cap S} \left| e(m) + \frac{s_e}{2\pi i} \right|^{-1} \ll_{\delta, s} |m|^{-1} \qquad (m \in T_\delta \subset \gamma(k)).
$$

Now use Lemma [9.](#page-21-2) It is easy to verify $\chi_m(n)^{-1}$ is bounded on Π if and only if $-\ell(n) \in \Theta^{\vee}$. \Box

Remark 14. The following estimate is proven in [\[3,](#page-42-1) Prop. 2.3.2] for fixed $s \in (\mathbb{C}^{\Sigma(1)})^{\Gamma_w}$ and $m \in \mathfrak{t}_w^{\vee}$ tending to infinity:

$$
\widehat{H_{\infty}}(\chi_m, -s) \ll_s \max_{\tau \in \Sigma_w(r_w)} \prod_{e \in \tau(1)} \left(\frac{1}{1+|m(e)|} \right)^{1+1/r_w}.
$$
\n(13)

.

In a later paper [\[2,](#page-42-3) Lemma 4.5] this estimate is cited to justify the weaker estimate

$$
\widehat{H_{\infty}}(\chi_m, -s) \ll_s \left(\frac{1}{1+|m|}\right)^{1+1/r_w}
$$

Unlike [\(13\)](#page-23-0) this does not suffice to prove that the integral $\int_{\mathfrak{t}^\vee} \widehat{H_\infty}(\chi_m, -s) dm$ converges, which is what is claimed in [\[2,](#page-42-3) Theorem 4.3] and needed for the Poisson formula. In the subsequent paper $[4,$ Prop. 4.12 this is apparently corrected and (13) is now cited to justify the stronger estimate

$$
\widehat{H_{\infty}}(\chi_m, -s) \ll_s \left(\frac{1}{1+|m|}\right)^{r_w+1}
$$

however this is false in general. Already for the projective plane over $\mathbb Q$ it is possible for $\widehat{H_{\infty}}(\chi_m, -s)$ to be as large as $\gg_s |m|^{-2}$ along certain lines in \mathfrak{t}_w^{\vee} : the Fourier transform $\widehat{H_{\infty}}$ is given by

$$
\widehat{H_{\infty}}(\chi_m, -s) = \left(\frac{1}{2\pi i}\right)^2 \frac{(s_0 + s_1 + s_2)/2\pi i}{(x + \frac{s_1}{2\pi i})(y + \frac{s_2}{2\pi i})(-x - y + \frac{s_0}{2\pi i})}
$$
(14)

which is $\gg |m|^{-2}$ when m tends to infinity along $x + y = 0$.

3.3.3. Periods. Let γ be a singular flag. We will need to evaluate the following constants:

$$
\rho_{\gamma} = \lim_{\substack{s = (t,\ldots,t) \\ t \to \infty}} \sum_{\xi \in R^{\vee}} R_{\gamma}(s + 2\pi i m_{\xi}).
$$

We will see that this equals the product of the level K regulator of T with a period obtained by integrating R_{γ} . Recall that R_{γ} factors through the quotient $(\mathbb{C}^{\Sigma(1)})^{\Gamma} \twoheadrightarrow (\mathbb{C}^{\Sigma(1)})^{\Gamma}/(\mathfrak{t}^{\vee} \otimes$ \mathbb{C} = (Pic Y) $\otimes \mathbb{C}$ -- $\rightarrow \mathbb{C}$ to a rational function on the Picard group. In particular, it defines a rational function on the subspace $\mathfrak{t}^\vee_w/\mathfrak{t}^\vee \subset (\text{Pic } Y) \otimes \mathbb{C}$; this subspace is isomorphic to the dual \mathfrak{t}_0^{\vee} of the trace-zero subspace.

Lemma 10. The constant ρ_{γ} is equal to

$$
|R_K| \int_{\mathfrak{t}_0^{\vee}} R_{\gamma}(s_{AC} + 2\pi im) \, dm
$$

where $|R_K|$ is the volume of the level K regulator group of T.

Proof. Let Υ denote the free part of the finitely generated abelian group

$$
T(\mathbb{Q}) \cap \left(T(\mathbb{R}) \times \prod_{v \text{ finite}} K_v\right).
$$

The group Υ has rank $r_w - r$ [\[16,](#page-43-14) Thm. 4, p. 285]. It is well-known that the natural map

$$
\Upsilon \to T(\mathbb{R}) \xrightarrow{\ell w} \mathfrak{t}_w
$$

embeds Υ as a full-rank lattice of the trace-zero subspace \mathfrak{t}_0 , and the dual group R^{\vee} is canonically isomorphic to the dual lattice to $\ell_w \Upsilon$ in \mathfrak{t}_0^{\vee} . By the definition of the Riemann integral,

$$
\lim_{t \to \infty} \sum_{\xi \in R^{\vee}} R_{\gamma}(ts_{AC} + 2\pi im_{\xi}) = \lim_{t \to \infty} t^{-(r_w - r)} \sum_{\xi \in R^{\vee}} R_{\gamma}(s_{AC} + 2\pi im_{\xi}/t) = V^{-1} \int_{t_0^{\vee}} R_{\gamma}(s_{AC} + 2\pi im) dm
$$

where V is the volume of a fundamental domain for the dual lattice to $\ell_w \Upsilon$ in \mathfrak{t}_0^{\vee} . Since this is isomorphic to R^{\vee} we have $V^{-1} = |R_K|$.

Example 4 (real quadratic Severi–Brauer surfaces). The depicted fan determines a complete smooth toric surface Y over Q whose open torus $T = R_{\mathbb{Q}}^E \mathbb{G}_m$ splits over a real quadratic extension E with Galois group $\Gamma = \langle \gamma \rangle$. (This surface is isomorphic to \mathbb{P}_2 over $\mathbb Q$ as a variety but not as a toric variety.) The boundary divisor D is Cartier and we use the canonical

FIGURE 1. The fan for a quadratic Severi–Brauer surface.

integral model for $Y - D$ (Remark [3\)](#page-8-0). The D-integral region for the divisor D corresponding to the ray e_0 is shaded in the figure. The D-integral rational points of T are $[1 : a : \gamma a]$ for $a \in O_E \setminus \{0\}$. With the help of [\(14\)](#page-23-1), one finds the residual function for the flag γ determined by the hyperplane H_0 is (with respect to the coordinate $z = -(x + y)$ on t)

$$
R_{\gamma}(s) = (2\pi i) \text{ itres}_{z}[\widehat{H_{\infty}}(\chi_{m}, -s) dm, \gamma] = \frac{s_{0} + s_{1} + s_{2}}{(\frac{s_{0}}{2} + s_{1})(\frac{s_{0}}{2} + s_{2})}.
$$

The regulator group $R \subset E^{\times} \backslash A_E^{\times,1}/K$ is isomorphic to $\mathbb{R}/(\mathbb{Z} \log u)$ where u is a positive generator of O_E^{\times} . (Indeed, if $E^{\times}xK$ is in R then its image in the class group of E is trivial, which means $x_{fin}K_{fin} \in E^{\times}K_{fin}$. Thus we have an isomorphism $R \to \mathbb{R}/(\mathbb{Z} \log u) : E^{\times}xK \mapsto$ log y where we have translated by $E^{\times}K$ to ensure that $x = x_{fin}x_{\infty} = (1, \ldots, 1; y, 1/y)$ and y > 0.) A generating Hecke character is given by $\xi(E^{\times}xK) = |y|^{\frac{2\pi i}{\log u}}$. This character

has local exponent $m_{\xi} = \frac{-1}{2 \log u}(x - y)$. Thus for any Γ-invariant s we have $s + 2\pi i m_{\xi} =$ $\left(s_0, s_1 - \frac{\pi i}{\log}\right)$ $\frac{\pi i}{\log u}, s_1 + \frac{\pi i}{\log u}$ $\log u$). The constant ρ_{γ} is $\rho_{\gamma} = \lim_{\Delta \to 0}$ $\lim_{s=(t,t,t)\to\infty} R_{\gamma}(s+2\pi i m_{\xi}) = \lim_{t\to\infty} \sum_{k\in\mathbb{Z}}$ $k\in\mathbb{Z}$ 4t $t^2+\left(\frac{2\pi k}{\log n}\right)$ $\log u$ $\overline{\lambda^2} = \lim_{t \to \infty}$ $2\log u \coth\left(\frac{1}{2}\right)$ $\frac{1}{2}t\log u$ = $2\log u$.

Or using the lemma, we check that

$$
\rho_{\gamma} = |R_K| \int_{-\infty}^{\infty} \frac{4}{1 + (2\pi m)^2} dm = 2 \log u.
$$

3.3.4. Conical multiple Dirichlet series. Let K_{fin} be an arbitrary open compact subgroup of $T(\mathbb{A}_{fin})$ and set $K = K_{fin} \times K_{\infty}^{\max}$. Assume K contains K^D . Let $\Phi = \prod_{v \in P_{\mathbb{Q}}} \Phi_v : T(\mathbb{A}) \to \mathbb{C}$ be a locally constant factorizable K-invariant function (e.g. $\Phi = \xi \in (T(\mathbb{Q}) \setminus T(\mathbb{A})/K^D)^{\vee}$ may be an automorphic character of level K^D). Let $\sigma \subset \mathfrak{t}$ be a subset (e.g. an open or closed cone). The multiple Dirichlet series of Φ over D-integral ideals in σ is

$$
L(s, \Phi, \sigma, D) = \sum_{\substack{n \in \mathcal{N} \\ \ell(n) \in \sigma}} \Phi(n) n^{-s}.
$$

The multiple Dirichlet series of Φ over principal D-integral ideals in σ is

$$
L_1(s, \Phi, \sigma, D) = \sum_{\substack{n \in \mathscr{P} \\ \ell(n) \in \sigma}} \Phi(n) n^{-s} = \frac{1}{|C|} \sum_{\psi \in C^{\vee}} L(s, \Phi \psi, \sigma, D).
$$

The difference between these conical series and the multiple Dirichlet series $\widehat{H}_{fin}(\chi, -s, D)$ from Corollary [1](#page-13-0) is that these series impose global conditions. Accordingly, $Z(s, \xi, \sigma, D)$ and $Z_1(s, \xi, \sigma, D)$ do not admit Euler products in general even though $\widehat{H}_{fin}(\chi, -s, D)$ always does, and this is where the failure of local-to-global properties begins for the associated integral Diophantine problem. More precisely, the condition $n \in \mathscr{P}$ is a global condition (at least when $C \neq 1$), and the condition $\ell(n) \in \sigma$ is a global condition (when σ does not contain $\ell(\mathcal{N})$.

Example 5 (Comparison with Hecke L-series for restriction tori). Continue the context of Example [2](#page-14-2) with $T = R_{\mathbb{Q}}^E$ and $K^D = K^{\max}$. The formula $n^{-s} = N(I)^{-s_1}$ shows that if σ is the positive $(D_{\infty}$ -integral) cone of t, then

$$
L(s_1, \xi) = L(s, \xi, \sigma, D_{\infty})
$$

where $L(s_1, \xi)$ is the Hecke L-series of ξ . This conical series does admit an Euler product, and this is because σ contains $\ell(\mathcal{N})$.

3.4. Proof of the main theorem on height zeta functions. A flag in Z is called Σ *convex* if it is cut out by singular hyperplanes H_1, \ldots, H_r whose generators e_1, \ldots, e_r form a Σ-convex set. For each cone σ in $\Sigma_{\Gamma}(r)$ let

$$
\Pi_{\sigma} = \mathfrak{t}^{\vee} + i\sigma^{\vee} \subset \mathfrak{t}^{\vee} \otimes \mathbb{C},
$$

a polyhedron with boundary \mathfrak{t}^{\vee} and defining isomorphism determined by $\sigma(1)$, and let Z_{σ} denote the subset of Σ -convex singular flags in Z cut out by a Π_{σ} -stable collection of singular hyperplanes. Let Σ_D be the smallest subfan of Σ_{Γ} whose support $|\Sigma_D|$ contains $-\ell(\mathcal{N})$. For each $\tau \in \Sigma_D$ choose any maximal dimensional cone $\sigma_{\tau} \in \Sigma_D(r)$ containing τ . Let $\tau^{\circ} = \tau - \bigcup_{\tau' \subsetneq \tau} \tau'$ denote the interior of a cone τ .

Theorem 4. Assume that each Σ -convex singular flag is compatible with the polyhedron Π_{σ} for every $\sigma \in \Sigma_D(r)$. Then the height zeta function for D-integral rational points on the torus

$$
Z(s) = \sum_{\substack{t \in T(\mathbb{Q}) \\ D\text{-integral}}} \frac{1}{H(t, s)}
$$

is equal to

$$
Z(s) = \frac{|\mu|}{|C||R|P} \sum_{\xi \in R^{\vee}} \sum_{\psi \in C^{\vee}} \sum_{\tau \in \Sigma_D} \sum_{\gamma \in Z_{\sigma_{\tau}}} L(s + 2\pi i m_{\gamma\xi}, (\xi \psi)^{-1}, -\tau^{\circ}, D) R_{\gamma}(s + 2\pi i m_{\xi})
$$

where

$$
P = \sum_{\gamma \in Z_{\sigma_1}} \int_{\mathfrak{t}_0^{\vee}} R_{\gamma}(s_{AC} + 2\pi im) \, dm.
$$

in the region of $s \in (\mathbb{C}^{\Sigma(1)})^{\Gamma}$ where both sides converge.

Remark 15. As is well-known, the left-hand side converges absolutely in a half-space of $(\mathbb{C}^{\Sigma(1)})^{\Gamma}$ where $\text{Re}(s_e) \gg_{\Sigma,T} 0$ for every $e \in \Sigma(1)$. Meanwhile the right-hand side converges absolutely for any $s \in (\mathbb{C}^{\Sigma(1)})^{\Gamma}$ for which

$$
\sum_{\xi \in R^{\vee}} R_{\gamma}(s + 2\pi i m_{\xi}) \tag{15}
$$

is absolutely convergent for every cone $\sigma \in \Sigma_D(r)$ and singular flag $\gamma \in Z_{\sigma}$.

Proof. The formula for Z will be obtained by applying the Poisson formula to the subgroup $T(\mathbb{Q}) \subset T(\mathbb{A})$ and the function $x \mapsto H(x, -s, D)$, then evaluating the integral using the residue formula [\[14,](#page-43-4) Theorem 1]. We must check that $H(x, -s, D)$ is in $L^1(T(\mathbb{A}))$ and the restriction of $\hat{H}(\chi, -s, D)$ is in $L^1(T(\mathbb{Q})^{\perp})$. The integral to be computed is

$$
Z(s) = \int_{(T(\mathbb{Q}) \backslash T(\mathbb{A}))^{\vee}} \widehat{H}(\chi, -s, D) d\chi.
$$

First we verify that $x \mapsto f(x) = H(x, -s, D)$ is in $L^1(T(\mathbb{A}))$. The chain of inequalities

$$
\int_{T(\mathbb{A})} f(x) d\mu(x) = \lim_{C \text{ compact}} \int_C f(x) d\mu(x) \le \lim_{S \text{ finite}} \int_{T(\mathbb{A}_S)} f(x) d\mu(x) \le \int_{T(\mathbb{A})} f(x) d\mu(x)
$$

in the limits of larger C and S shows that

$$
\int_{T(\mathbb{A})} f(x) d\mu(x) = \lim_{S} \int_{T(\mathbb{A}_S)} f(x) d\mu(x) \le \lim_{S} \prod_{v \in S} \int_{T(\mathbb{Q}_v)} f_v(x) d\mu_v(x).
$$

The convergence of the right-hand side follows from Proposition [3](#page-14-3) and Corollary [1.](#page-13-0)

Next we check that the restriction of $H(\chi, -s, D)$ to $T(\mathbb{Q})^{\perp} \cong (T(\mathbb{A})/T(\mathbb{Q}))^{\vee}$ is integrable. Let $K = K^D$ denote the subgroup of the maximal compact subgroup fixing 1^D . We have the exact sequence

$$
1 \longrightarrow C^{\vee} \longrightarrow (T(\mathbb{Q}) \backslash T(\mathbb{A})^1 / K)^{\vee} \longrightarrow R^{\vee} \longrightarrow 1.
$$

28 ANDREW O'DESKY

Since R^{\vee} is a free abelian group this sequence splits, so any character of R lifts to a character of $T(\mathbb{Q})\backslash T(\mathbb{A})^1/K$. Furthermore, any character of $T(\mathbb{Q})\backslash T(\mathbb{A})^1/K$ lifts to a character of $T(\mathbb{Q})\backslash T(\mathbb{A})/K$ (see [\(3\)](#page-10-1)). Thus, any character of $T(\mathbb{Q})\backslash T(\mathbb{A})^1/K$ may be factored uniquely as a product of three characters

$$
\xi \psi \chi
$$

where (identifying characters with their lifts to $T(\mathbb{Q})\setminus T(\mathbb{A})/K$) $\xi \in R^{\vee}$, $\psi \in C^{\vee}$ and $\chi = \chi_m$ is the character given by $\chi_m(t) = e(\langle \ell(t), m \rangle)$ for $m \in \mathfrak{t}^\vee$. Recall that $T(\mathbb{Q}) \setminus T(\mathbb{A})^1$ is compact [\[17,](#page-43-13) Theorem 3.1.1] so $(T(\mathbb{Q})\setminus T(\mathbb{A})^1)^\vee$ is discrete and equipped with the counting measure, so the subset $(T(\mathbb{Q})\setminus T(\mathbb{A})^1/K)^\vee$ is also equipped with the counting measure. Since \hat{H} is supported on $(T(\mathbb{Q})\backslash T(\mathbb{A})/K)^\vee$, the height zeta function is proportional to

$$
\sum_{\xi \in R^{\vee}} \sum_{\psi \in C^{\vee}} \int_{m \in \mathfrak{t}^{\vee}} \widehat{H}(\xi \psi \chi_m, -s, D) \, dm
$$

The infinite component $\widehat{H}_{\infty}(\xi\psi\chi_m, -s, D)$ is independent of ψ by the last assertion of Lemma [3.](#page-10-3) Recall the notation $\mathscr{Q} = T(\mathbb{A}_{fin})/K_{fin}^D$ of §[2.4.3,](#page-13-1) where for each $n \in \mathscr{Q}$ we have set

$$
n^{-s} := H(n, -s) = \prod_{p \in P_{\mathbb{Q}}^{\text{fin}}} p^{\frac{1}{e_p} \varphi(\ell_w(n_p))} \in \mathbb{C}^{\times}.
$$

Expand \hat{H}_{fin} of \hat{H} as an absolutely convergent multiple Dirichlet series (Corollary [1\)](#page-13-0) and then substitute it into $Z(s)$ to obtain

$$
Z(s) = \kappa \sum_{\xi \in R^{\vee}} \sum_{\psi \in C^{\vee}} \int_{m \in \mathfrak{t}^{\vee}} \sum_{n \in \mathcal{D}} (\xi \psi \chi_m)^{-1}(n) 1^{D}(n) n^{-s} \widehat{H}_{\infty}(\xi \chi_m, -s) dm
$$

=
$$
\kappa \sum_{\xi \in R^{\vee}} \sum_{\psi \in C^{\vee}} \sum_{n \in \mathcal{N}} (\xi \psi)^{-1}(n) n^{-s} \int_{m \in \mathfrak{t}^{\vee}} \chi_m(n)^{-1} \widehat{H}_{\infty}(\xi \chi_m, -s) dm
$$

for some yet to be determined positive constant κ .

Now we evaluate the inner integral using the residue formula from [\[14\]](#page-43-4). For each $n \in$ N let σ_n denote any cone in $\Sigma_D(r)$ containing $-\ell(n)$. By Proposition [5](#page-22-2) the integral $\int_{m\in\mathcal{N}} \chi_m(n)^{-1} \widehat{H_\infty}(\xi\chi_m, -s) dm$ admits an iterated residue expansion with respect to the polyhedron $\Pi_{\sigma_n} = \mathfrak{t}^\vee + i\sigma_n^\vee$. Let $z = \sigma(1)$ be the dual basis to $\sigma(1)^\vee$. The residue formula is

$$
\int_{m\in\mathfrak{t}^\vee} \chi_m(n)^{-1} \widehat{H_\infty}(\xi\chi_m,-s) dm = (2\pi i)^r \sum_{\gamma \in Z_{\sigma_n}} \text{itres}_z \left[\chi_m(n)^{-1} \widehat{H_\infty}(\xi\chi_m,-s) dm, \gamma \right].
$$

By definition of R_{γ} we have that

$$
(2\pi i)^r \operatorname{itres}_z \left[\chi_m(n)^{-1} \widehat{H_{\infty}}(\chi_m, -s) \, dm, \gamma \right] = \chi_{m_{\gamma}}(n)^{-1} R_{\gamma}(s)
$$

for any $s \in (\mathbb{C}^{\Sigma(1)})^{\Gamma_w}$ where $m_\gamma = m_\gamma(s)$ is the terminal point of γ . With our convention that m is associated to the element $(m(e))_e$ in $\mathbb{C}^{\Sigma(1)}$ one easily checks that $\chi_m(n) = n^{2\pi im}$ (Remark [8\)](#page-14-1) and $\widehat{H_{\infty}}(\xi \chi_m, -s) = \widehat{H_{\infty}}(\chi_m, -(s+2\pi i m_{\xi}))$ for any $\xi \in R^{\vee}$ where $m_{\xi} \in \mathfrak{t}_0^{\vee}$ is determined by the identity $\xi(t_{\infty}) = e(\langle m_{\xi}, \ell_{w}(t_{\infty}) \rangle)$ for $t_{\infty} \in T(\mathbb{R}) \subset T(\mathbb{A})$. Write

$$
m_{\gamma\xi} := m_{\gamma}(s + 2\pi i m_{\xi}) \in \mathfrak{t}_{\mathbb{C}}^{\vee}.
$$
 Then

$$
\int_{m\in\mathfrak{t}^{\vee}} \chi_m(n)^{-1} \widehat{H_{\infty}}(\xi \chi_m, -s) dm = \sum_{\gamma \in Z_{\sigma_n}} \chi_{m_{\gamma\xi}}(n)^{-1} R_{\gamma}(s + 2\pi i m_{\xi}) = \sum_{\gamma \in Z_{\sigma_n}} n^{-2\pi i m_{\gamma\xi}} R_{\gamma}(s + 2\pi i m_{\xi}).
$$

Putting this back into the earlier expression for Z obtains

$$
Z(s) = \kappa \sum_{\xi \in R^{\vee}} \sum_{\psi \in C^{\vee}} \sum_{n \in \mathcal{N}} (\xi \psi)^{-1}(n) n^{-s} \sum_{\gamma \in Z_{\sigma_n}} n^{-2\pi i m_{\gamma\xi}} R_{\gamma}(s + 2\pi i m_{\xi}). \tag{16}
$$

Collecting $n \in \mathcal{N}$ into summands according to whichever cone $\tau \in \Sigma_D$ contains $-\ell(n)$ rearranges to the claimed formula up to the constant κ .

To determine κ , specialize $s = (t, \ldots, t) = ts_{AC}$ and let $t \to +\infty$. By the defining series for $Z(s)$, we see it approaches the number of D-integral rational points of T with height one. This is also the order of the torsion subgroup μ of

$$
T(\mathbb{Q}) \cap \left(T(\mathbb{R}) \times \prod_{v \text{ finite}} K_v^D\right).
$$

On the other hand $Z(s)$ approaches

$$
\kappa \sum_{\xi \in R^{\vee}} \sum_{\psi \in C^{\vee}} \sum_{\substack{n \in \mathcal{N} \\ n^{-s} = 1}} (\xi \psi)^{-1}(n) \sum_{\gamma \in Z_{\sigma_1}} R_{\gamma}(ts_{AC} + 2\pi i m_{\xi})
$$

The subset of $n \in \mathcal{N}$ satisfying $n^{-s} = 1$ for all s is K_{fin}^{\max}/K_{fin}^D . Recall that $(\xi\psi)^{-1}(n)$ is evaluated using the maps

$$
K_{\text{fin}}^{\text{max}}/K_{\text{fin}}^D \hookrightarrow T(\mathbb{Q})\backslash T(\mathbb{A})/K^D \xrightarrow{\sim} T(\mathbb{Q})\backslash T(\mathbb{A})^1/K^D \times \mathfrak{t}^{\vee} \xrightarrow{\sim} R \times C \times \mathfrak{t}^{\vee}.
$$

The identity component of $T(\mathbb{Q})\backslash T(\mathbb{A})/K^D$ is $T(\mathbb{Q})\backslash (T(\mathbb{Q})T(\mathbb{R})K^D)/K^D$ (see the proof of Lemma [3\)](#page-10-3) which shows that

$$
(K_{\mathrm{fin}}^{\mathrm{max}}/K_{\mathrm{fin}}^D) \cap (R \times 1 \times \mathfrak{t}^{\vee}) = 1.
$$

Thus $(K_{\text{fin}}^{\text{max}}/K_{\text{fin}}^D) \to C$ is injective, so by orthogonality of characters $Z(s)$ approaches

$$
\kappa|C|\sum_{\xi\in R^\vee}\sum_{\gamma\in Z_{\sigma_1}}R_\gamma(ts_{AC}+2\pi im_\xi).
$$

With the help of Lemma [10](#page-23-2) this approaches

$$
\kappa|C|\sum_{\gamma\in Z_{\sigma_1}}\rho_\gamma=\kappa|C||R_{K^D}|\sum_{\gamma\in Z_{\sigma_1}}\int_{\mathfrak{t}_0^\vee}R_\gamma(s_{AC}+2\pi im)\,dm=|\mu|.
$$

Now assume [\(15\)](#page-26-0) is absolutely convergent for every cone $\sigma \in \Sigma_D(r)$ and singular flag $\gamma \in Z_{\sigma}$. Then [\(16\)](#page-28-0) is absolutely convergent, i.e. $\hat{H}(\chi, -s, D) \in L^{1}((T(\mathbb{Q})\setminus T(\mathbb{A}))^{\vee})$. It follows from the Poisson formula (cf. §[3.1\)](#page-16-3) that

$$
\sum_{x \in T(\mathbb{Q})} H(xy, -s, D) = \int_{(T(\mathbb{Q}) \backslash T(\mathbb{A}))^{\vee}} \widehat{H}(\chi, -s, D) \chi(y) \, d\chi \tag{17}
$$

holds for almost every $y \in T(\mathbb{A})$. Since $H(\chi, -s, D) \in L^1((T(\mathbb{Q})\setminus T(\mathbb{A}))^{\vee})$ the right-hand side is continuous in y. The equality thus holds for every y if $x \mapsto H(xy, -s, D)$ is in $L^1(T(\mathbb{Q}))$ for any $y \in T(\mathbb{A})$, as this implies continuity of the left-hand side. Since y only affects the height at finitely many places in a manner independent of x , it suffices to consider $y = 1$. The left-hand side can be bounded from above by the same sum where s is replaced by a Z-linear combination of toric divisors representing a T-Cartier divisor and satisfying $\text{Re}(s_e) > 0$ for all $e \in \Sigma(1)$. Such an s corresponds to a big T-linearized line bundle L. Since L is big, the number of rational points of L-height $\leq H$ not in the base locus of L is bounded by a polynomial in H . The base locus of L is disjoint from T , and this proves the left-hand side is absolutely convergent in a half-space of $(\mathbb{C}^{\Sigma(1)})^{\Gamma}$ where $\text{Re}(s_e) \gg_{\Sigma,T} 0$ for every $e \in \Sigma(1)$. In particular, the formula holds when $y = 1$.

Remark 16. The proof shows the twisted Poisson formula [\(17\)](#page-28-1) is valid for any parameter $y \in T(\mathbb{A})$ whenever both sides converge. This can be useful for applications [\[13\]](#page-43-7).

4. An orbit parametrization for polynomials with given Galois group

Let G be a finite group. Consider the projective space of lines

$$
\mathbb{P}(\mathbb{A}^1 \oplus \mathbb{A}^n) = \text{Proj } \mathbb{Z}[Y, X_1, \ldots, X_n]
$$

in the G-representation formed from the trivial representation $\mathbb{A}^1 = \text{Spec } \mathbb{Z}[Y]$ and the regular representation $\mathbb{A}^n = \text{Spec } \mathbb{Z}[X_1, \ldots, X_n]$ where X_1, \ldots, X_n is the dual basis to $G =$ ${g_1 = e, \ldots, g_n}$. In this section we show that the quotient $\mathbb{P}(\mathbb{A}^1 \oplus \mathbb{A}^n)/G$ can be used to parametrize polynomials with Galois group G . This is a modification of the construction from [\[15\]](#page-43-8) where a trace condition was imposed. We will later see that $\mathbb{P}(\mathbb{A}^1 \oplus \mathbb{A}^n)/G$ is a toric variety when G is abelian.

4.1. Normal bases for torsors of finite groups. Let G act by S-automorphisms on a finite S-scheme $f: T \to S$. If $(f_*\mathcal{O}_T)^G = \mathcal{O}_S$ and the induced action of G on the set $T(k)$ is free for any S-field k, then we say T/S is Galois with Galois group G, or that T/S is a G-torsor. A G-stable ordered list (x_1, \ldots, x_n) of regular functions on a G-torsor T/S is called a *normal basis* if it forms a basis of $\mathcal{O}(T)$ as an $\mathcal{O}(S)$ -module and is isomorphic as a G-set to G itself with its left regular action. Let $\mathcal L$ be a G-line bundle over T. Let $x \mapsto {}^gx$ denote the action of G on sections given by ${}^gx(t) = g(x(g^{-1}t))$ for $t \in T$, where the outer g on the right-hand side denotes the action of g on the fibers $(g^{-1}t)^*\mathcal{L} \to t^*\mathcal{L}$ coming from the structure of $\mathcal L$ as a G-line bundle.

We will equip G-torsors T/S with extra data:

- (1) a G-line bundle $\mathcal L$ over T, and
- (2) global sections y and x of $\mathcal L$

subject to three conditions:

- (1) $Gx \cup \{y\}$ generates \mathcal{L} ,
- (2) $^{g}x(t) \neq^{h}x(t)$ if $g \neq h$ and $t \in T$, and
- (3) y is *G*-invariant.

We regard data $(T/S, \mathcal{L}, x, y)$ and $(T'/S, \mathcal{L}', x', y')$ as equivalent if there is a G-equivariant isomorphism $\phi \colon T \to T'$ over S and an isomorphism $\psi \colon \phi^* \mathcal{L}' \to \mathcal{L}$ of G-line bundles such that $(\psi \phi^* x', \psi \phi^* y') = (x, y)$. We call $(T/S, \mathcal{L}, x, y)$ monic if y is nowhere vanishing. If $(T/S, \mathcal{L}, x, y)$ is monic then by rescaling x and y we may take $\mathcal{L} = \mathcal{O}(T)$, and by rescaling x by $\mathcal{O}(T)^{\times}$ we may take $y = 1$; thus monic points may be written more compactly as $(T/S, x)$. Let $A \subset \mathbb{A}^n$ denote the complement of the union of the hyperplane divisors $\{X_i/Y = X_j/Y\}$ for all $1 \leq i < j \leq n$. Let $\mathcal{G} \subset A$ denote the open subscheme where the group determinant $\Delta_G = \det(X_{gh}/Y)_{g,h \in G}$ is invertible. The group algebra structure on \mathbb{A}^n gives $\mathcal G$ the structure of a group scheme.

Proposition 6. The S-points of the affine variety A/G are in bijection with equivalence classes of $(T/S, \mathcal{L}, x, y)$, functorially in S. Any S-point of \mathcal{G}/G is monic. Let $(T/S, x)$ be a monic S-point of A/G and write (x_1, \ldots, x_n) for the regular functions $({}^{g_1}x, \ldots, {}^{g_n}x)$ on T. Then $(T/S, x)$ is contained in \mathcal{G}/G if and only if $(x_1(t), \ldots, x_n(t))$ is a normal basis for $\mathcal{O}(T) \otimes k(f(t))$ over $k(f(t))$ for every $t \in T$.

Proof. First observe that A is contained in the open locus of \mathbb{P} where G has no isotropy, and therefore G acts freely on $A(k)$ for any S-field k. It follows that the scheme A/G represents the stack theoretic quotient $[A/G]$. This means that the S-points of A/G are in bijection with equivalence classes of pairs $(T/S, \phi)$ where T/S is a G-torsor and ϕ is a G-equivariant morphism $\phi: T \to A$. Thus ϕ is determined by the line bundle $\mathcal{L} = \phi^* \mathcal{O}(1)|_A$ and the global sections $y = \phi^* Y, x = x_1 = \phi^* X_1, \dots, x_n = \phi^* X_n$. Since ϕ is G-equivariant, these global sections satisfy the required conditions: they clearly generate \mathcal{L} , and if $^gx(t) = ^hx(t)$ then $\phi(t) \in \{X_q/Y = X_h/Y\}$ which implies $g = h$. Meanwhile $\mathcal L$ has a canonical G-line bundle structure inherited from $\mathcal{O}(1)$ for which y is invariant since $gy = g\phi^*Y = \phi^*gY = \phi^*Y = y$. In the other direction, from a datum $(T/S, \mathcal{L}, x, y)$, we get a morphism $\phi \colon T \to A$ from y and the sections $x_g := {}^gx$ since $Gx \cup \{y\}$ is generating. This morphism will be G-equivariant since

$$
\begin{aligned} \phi(gt) &= [y(gt) : (x_h(gt))_h] = [g^{(g^{-1}}y)(t) : (g^{(g^{-1}}x_h)(t))_h] \\ &= [gy(t) : (gx_{g^{-1}h}(t))_h] \\ &= g\phi(t). \end{aligned}
$$

An S-point of \mathcal{G}/G is automatically monic since $\mathcal G$ is contained in the open locus of $\mathbb P$ where Y is invertible. A monic S-point $(T/S, x)$ will be contained in \mathcal{G}/G if and only if ϕ factors through G, which occurs if and only if $\phi^* \Delta_G$ is invertible on T. If $t \in T$, then $\phi^* \Delta_G(t) = \det(x_{gh}(t))_{g,h}$ is invertible if and only if $(x_1(t), \ldots, x_n(t))$ forms a basis for $\mathcal{O}(T) \otimes k(f(t))$ over $k(f(t))$.

4.2. Characteristic maps for torsors of finite groups. The root height of a polynomial $f = t^{n} + a_1 t^{n-1} + \cdots + a_n \in \mathbb{Z}[t]$ is

$$
H(f) = \max(|a_1|, |a_2|^{1/2}, \ldots, |a_n|^{1/n}).
$$

We now explain how to express the root height in terms of a Weil height on \mathbb{P}/G in the usual sense, i.e. the height function obtained by pulling back the standard height on projective space via some morphism from \mathbb{P}/G to projective space. To do this, we consider some rational maps from P/G to projective space built from the characteristic polynomial. The coefficients of the characteristic polynomial will be valued in a certain line bundle on \mathbb{P}/G . Write $\mathbb{P} = \mathbb{P}(\mathbb{A}^1 \oplus \mathbb{A}^n)$ and let $\pi \colon \mathbb{P} \to \mathbb{P}/G$ denote the quotient morphism. Let $l = \text{lcm}(1, \ldots, n)$. Let $e_k(X) \in \mathcal{O}(k)$ denote the degree k elementary symmetric polynomial.

Lemma 11. The coherent sheaf $\mathcal{L}_G := (\pi_* \mathcal{O}(l))^G$ is an ample line bundle on \mathbb{P}/G . The lth power of the root height is a Weil height on \mathbb{P}/G for the linear system of \mathcal{L}_G spanned by the generating sections Y^l and $e_k^{l/k} \in H^0(\mathbb{P}/G, \mathcal{L}_G)$ for $1 \leq k \leq n$.

Proof. By [\[15,](#page-43-8) Prop. 3.2] the G-line bundle $\mathcal{O}(l)$ on \mathbb{P} descends to a line bundle on \mathbb{P}/G (passing to the lth power of $\mathcal{O}(1)$ trivializes any action of isotropy subgroups on fibers over fixed points). Sections of this descended line bundle on an open subset $U \subset \mathbb{P}/G$ are canonically identified via pullback along π with G-invariant sections of $\mathcal{O}(l)$ on $\pi^{-1}(U)$. Note \mathcal{L}_G is ample since π is finite and its pullback $\pi^* \mathcal{L}_G = \mathcal{O}(l)$ is ample.

Consider the projective morphism Ch: $\mathbb{P}/G \to \mathbb{P}_n$ defined by the generating sections Y^l and $e_k^{l/k} \in H^0(\mathbb{P}/G, \mathcal{L}_G)$ for $1 \leq k \leq n$. On monic S-points of A/G it is given by

$$
Ch(T/S, \mathcal{O}_T, x, y) = [y^l : e_1(Gx)^l : e_2(Gx)^{l/2} : \cdots : e_n(Gx)^{l/n}].
$$

These sections are globally generating for \mathcal{L}_G since $e_1(X) = e_2(X) = \cdots = e_n(X) = 0$ if and only if $X = 0$, so the base locus of this linear system is empty. only if $X = 0$, so the base locus of this linear system is empty.

Remark 17 (coefficient height). By contrast, the coefficient height $\max(|a_1|, |a_2|, \ldots, |a_n|)$ is not equivalent to a power of a Weil height on P/G . The proof of the lemma also shows that $(\pi_*\mathcal{O}(n))^G$ is a line bundle. There is a closely related map with coordinates $Y^{n-k}e_k \in \mathbb{R}^n$ $H^0(\mathbb{P}/G, (\pi_*\mathcal{O}(n))^G)$ for $0 \le k \le n$. On monic S-points of A/G this map is given by

$$
(T/S, \mathcal{O}_T, x, y) \dashrightarrow [y^n : y^{n-1}e_1(Gx) : y^{n-2}e_2(Gx) : \cdots : e_n(Gx)].
$$

The "problem" with this map is that it has a base locus $B = \{Y = e_n(X) = 0\}$ which is not G-stable. The line bundle associated to this height is not on \mathbb{P}/G but on the blowup of \mathbb{P}/G along B , and when G is abelian the blowup no longer has a canonical toric variety structure and the methods of this paper are not applicable. The "correct" compactification of A/G to consider the coefficient height is $(\mathbb{P}_1 \times \cdots \times \mathbb{P}_1)/G$ where G permutes the G copies of the projective line, however this alternative quotient does not have a natural toric structure.

Remark 18 (anticanonical height). Let $D = \{Y = 0\}$. Then $\mathcal{L}_G = \mathcal{O}(lD)$. The divisor D is toric (T-stable) and corresponds to the ray e_0 in the fan of Y. Let $s_D \in (\mathbb{Z}^{\Sigma(1)})^{\Gamma}$ be the corresponding element (1 in the e_0 coordinate and 0 elsewhere). One computes that $s_{AC} \equiv (n+1)s_D \pmod{M}$, so the root-height is equivalent to $H_{AC}^{\frac{1}{n+1}}$.

4.2.1. Fibers of the characteristic map. Any G-torsor over $\mathbb Q$ has the form $T = \text{Spec } K \sqcup$ $\cdots \sqcup$ Spec K for some field K with Galois group $D \leq G$. We call D the monodromy group of T.

Lemma 12. Let $D \leq G$ and let $(K/\mathbb{Q}, x)$ be a rational point of \mathcal{G}/G with monodromy group D. The number of elements of $(\mathcal{G}/G)(\mathbb{Q})$ which have the same characteristic polynomial as $(K/\mathbb{Q},x)$ is equal to $[N_{S_n}(D):N_G(D)].$

Proof. Suppose two rational points $(K/\mathbb{Q}, x)$, $(K'/\mathbb{Q}, x')$ have the same characteristic polynomial. Since G acts transitively on the roots, we may assume that $x = x'$. Now we are in the situation that K and K' are isomorphic as \mathbb{Q} -algebras but not necessarily as G-algebras, since an isomorphism of G-algebras must be G-equivariant. A G-equivariant isomorphism $K \to K'$ that fixes x must be the identity, so we are computing the number of non-isomorphic G-algebra structures on a given G-algebra with monodromy group D. A G-algebra (ring of functions on a G-torsor) with monodromy group D is the same thing as a cocycle of $H^1(\mathbb{Q}, G)$ whose image is G-conjugate to $D \leq G$, while an étale Q-algebra of degree *n* with monodromy group *D* is the same thing as a cocycle of $H^1(\mathbb{Q}, S_n)$ whose image is S_n -conjugate to D. We must compute the number of cocycles in $H^1(\mathbb{Q}, G)$ lying over a given cocycle $[K/\mathbb{Q}] \in H^1(\mathbb{Q}, S_n)$ whose image is S_n -conjugate to D under the natural homomorphism $H^1(\mathbb{Q}, G) \to H^1(\mathbb{Q}, S_n)$. This is equal to the given quantity.

One can check that $N_{S_n}(C_n) = n\varphi(n)$ which proves the following corollary.

Corollary 2. Suppose $G = C_n$. The number of rational points of $T = \mathcal{G}/C_n$ with the same irreducible characteristic polynomial is $\varphi(n) = |\text{Out}(C_n)|$ where φ is Euler's totient function.

5. THE TORIC STRUCTURE ON P/G

Assume G is abelian. Let G denote the group of units in the group algebra of G, an algebraic torus over $\mathbb Q$ of rank n. The group G is a constant subgroup of $\mathcal G$ and $T = \mathcal G/G$ is again an algebraic torus of rank n. The quotient variety \mathbb{P}/G inherits a T-action from the G-action on $\mathbb{Q}G$ making it into a (normal) toric variety. We now describe its fan.

Let $G^{\vee} = \text{Hom}(G, \mathbb{C}^{\times})$ denote the set of characters of G. Let E be the field over Q generated by the values of every character $\chi \in G^{\vee}$. The elementary idempotents v_{χ} of CG are in bijection with characters $\chi \in G^{\vee}$ where $v_{\chi} = \frac{1}{n}$ $\frac{1}{n} \sum_{g \in G} \chi(g^{-1})g$. We define

$$
N'_E = \mathbb{Z}\langle v_\chi : \chi \in G^\vee \rangle \subset \mathbb{C}G, \qquad \mathfrak{t}_E = N'_E \otimes \mathbb{R}.
$$

Let $v_0 = -\sum_{\chi} v_{\chi}$. Let Σ be the set of cones σ_I where $I \subset \{v_0\} \cup \{v_{\chi} : \chi \in G^{\vee}\}\$ is any proper subset and $\sigma_I \subset \mathfrak{t}_E$ is the convex span of I. For a root of unity ζ , let $\ell(\zeta) \in [0,1) \cap \mathbb{Q}$ be the unique rational number satisfying $e(\ell(\zeta)) = \zeta$.

Proposition 7. Let $G = \langle c_1 \rangle \times \cdots \times \langle c_K \rangle$ be a decomposition of G into cyclic groups. Define the elements

$$
\omega_j = \sum_{\chi \in \langle c_j \rangle^\vee} \ell(\chi(c_j^{-1})) v_\chi \in N'_E \otimes \mathbb{Q}
$$

and the lattice

$$
N_E = N'_E + \sum_{j=1}^K \mathbb{Z} \omega_j.
$$

The toric variety over Q associated to the fan Σ and the lattice N_E with its canonical Galois action by $\Gamma = \text{Gal}(E/\mathbb{Q})$ is the quotient variety \mathbb{P}/G , and its open torus is T.

Proof. Observe that the group G acts by left translation on $N'_E \otimes \mathbb{C}$, and the dual action on $M'_E \otimes \mathbb{C}$ is given by

$$
g \cdot v_{\chi}^{\vee} = \chi(g^{-1})v_{\chi}^{\vee}
$$

where $\{v_\chi^\vee\}$ is the dual basis to $\{v_\chi\}$. This induces a G-action on $\mathbb{C}[M_E'] = \mathbb{C}[\{x_\chi^{\pm 1}\}]$ where $x_{\chi} = \chi^{v_{\chi}^{\vee}}.$

Now observe that the quotient group N_E/N'_E acts on $\mathbb{C}[M'_E]$ by

$$
v \cdot \chi^u = e(\langle u, v \rangle) \chi^u
$$

where $\langle \cdot, \cdot \rangle: M'_E/(N'_E) \times N_E/N'_E \to \mathbb{Q}/\mathbb{Z}$ is the canonical perfect pairing. We have the short exact sequence

$$
0 \longrightarrow N'_E \longrightarrow N_E \longrightarrow G \longrightarrow 0
$$

where the quotient map takes ω_{C_i} to the fixed generator of C_i . For any character χ ,

$$
\omega_C \cdot x_\chi = e(\langle v_\chi^\vee, \omega_C \rangle) x_\chi = e(\ell(\chi(c^{-1}))) x_\chi = \chi(c^{-1}) x_\chi = c \cdot x_\chi.
$$

Thus the isomorphism $N_E/N'_E = G$ is compatible with the action on $\mathcal{G}_{\mathbb{C}} = \text{Spec } \mathbb{C}[M'_E]$. As a G-representation, $\mathbb{C}[M'_E]$ is a direct sum over the weight vectors χ^u for all $u \in M'_E$ which implies that the subspace $\mathbb{C}[M'_E]^G$ of G-invariants is a direct sum over the weight vectors χ^u with trivial G-action. In view of the isomorphism $N_E/N'_E = G$, we see χ^u is G-invariant if and only if $u \in N_E^{\vee}$. Thus $\mathbb{C}[M_E']^G = \mathbb{C}[N_E^{\vee}]$, and more generally

$$
\mathbb{C}[\sigma^{\vee} \cap M'_E]^G = \mathbb{C}[\sigma^{\vee} \cap N_E^{\vee}]
$$

for any cone $\sigma \in \Sigma$. The toric variety defined by Σ and N'_E is \mathbb{P} , which is covered by the set of affine toric varieties $U_{\sigma} = \text{Spec } \mathbb{C}[\sigma^{\vee} \cap M'_{E}]$ for $\sigma \in \Sigma$. Each U_{σ} is \mathcal{G} -stable hence also G-stable, so

$$
(\mathbb{P}/G)_{\mathbb{C}} = \bigcup_{\sigma \in \Sigma} U_{\sigma}/G = \bigcup_{\sigma \in \Sigma} \text{Spec } \mathbb{C}[\sigma^{\vee} \cap N_{E}^{\vee}].
$$

 \Box

6. Monic cyclic polynomials of bounded height

In this section we prove Theorem [2](#page-4-1) by applying Theorem [4](#page-26-1) to an integral Diophantine problem for the toric variety constructed in §[5.](#page-32-0) To define the integrality condition, we use the divisor $D = \{Y = 0\} \subset \mathbb{P}/G$. This is a Q-Cartier toric divisor which is not Cartier, however the multiple lD is Cartier where $l = \text{lcm}(1,\ldots,n)$ and corresponds to the line bundle $\mathcal{L}_G = (\pi_* \mathcal{O}(l))^G$ from §[4.2.](#page-30-0) In fact nD is already Cartier, but we use lD so that the coefficients $Y^{\ell}, e_k^{\ell/k} \in H^0(Y, \mathcal{L}_G)$ of the characteristic polynomial are generators of the line bundle. The quotient $\mathbb{P}/G = \text{Proj }\mathbb{Z}[Y, X_1, \ldots, X_n]^G$ is defined over \mathbb{Z} , as is the boundary divisor $D = \{Y = 0\}$, so for our integral model we simply take U to be the complement of D. For this integral model, the indicator function on D-integral points in $T(\mathbb{A})$

$$
1^D = \prod_{v \in P_{\mathbb{Q}}} 1^D_v
$$

is invariant under the maximal compact subgroup K^{\max} of $T(\mathbb{A})$. The preimage of U in \mathbb{P} is $\mathbb{P} - D = \mathbb{A}^n = \text{Spec } \mathbb{Z}[X_1/Y, \ldots, X_n/Y].$ The group algebra structure on \mathbb{A}^n is defined over \mathbb{Z} , so the action of the unit group \mathcal{G} on $\mathbb{P} - D$ and therefore the action of the quotient group $T = \mathcal{G}/G$ on $U = \mathbb{A}^n/G$ are defined over \mathbb{Z} . For finite places v the maximal compact subgroup is $K_v^{\max} = T(\mathbb{Z}_v) \subset T(\mathbb{Q}_v)$, which shows that if $t \in U(\mathbb{Z}_v)$ and $k \in K_v^{\max}$ then $tk \in U(\mathbb{Z}_v)$. Thus $K_v^D = K_v^{\max}$.

Remark 19. For the integral Diophantine problem in [\[6\]](#page-43-6), the indicator function on D-integral points was not invariant under the maximal compact subgroup, cf. [\[6,](#page-43-6) Prop. 4]. This occured because of the trace-one condition that was imposed on polynomials. This condition corresponds to splitting off the factor of \mathbb{G}_m from the unit group $\mathcal{G} = \mathbb{G}_m \times \mathcal{G}_1$ corresponding to the trivial representation of C_3 , and using the torus $T_1 = \mathcal{G}_1/C_3$ instead of \mathcal{G}/C_3 . However the splitting $\mathcal{G} = \mathbb{G}_m \times \mathcal{G}_1$ is only defined over $\mathbb{Z}[\frac{1}{3}]$ $\frac{1}{3}$ (the trivial idempotent is $\frac{1}{3}(1+\sigma+\sigma^2)$), so the action of T_1 on the integral model is not defined over $\mathbb Z$ and the argument given here does not apply.

6.1. Describing the subfan Σ_D . We describe the smallest subfan $\Sigma_D \subset \Sigma$ whose support $|\Sigma_D|$ contains $-\ell(\mathcal{N})$. The ray generators in Σ_{Γ} correspond to Galois orbits in $\Sigma(1)$ = ${e_0} \sqcup {e_\chi}_{\chi \in G^\vee}$ (Lemma [1\)](#page-9-2). The Galois action on $\Sigma(1)$ is identified with the cyclotomic action on triv⊔G^{\vee} and the \mathbb{Z}^{\times} -orbits of G^{\vee} are naturally in bijection with the cyclic subgroups of G^{\vee} . The cones in Σ_{Γ} containing e_0 are in bijection with proper subsets of the set $G^{\vee}/\mathbb{Z}^{\times}$

of \mathbb{Z}^{\times} -orbits on G^{\vee} . Let O_1, \ldots, O_r be the Galois orbits in $\{e_{\chi}\}_{\chi \in G^{\vee}}$. We may assume that $O_1 = \{e_1\}$. Set $f_i = \sum_{e \in O_i} e$. Then (f_1, \ldots, f_r) is a basis of **t**. Let

$$
\Lambda_j = -\mathbb{R}_{\geq 0} \langle e_0, f_i : i \neq j \rangle = \{ b_1 f_1 + \dots + b_r f_r : b_j \geq 0 \text{ and } b_j \geq b_i \text{ for all } i \} \subset \mathfrak{t}.
$$

Any element of $\ell(\mathcal{N})$ may be expressed as $b_1f_1 + \cdots + b_rf_r$ for $b_i \geq 0$, and it is contained in each region Λ_j where j is an index for which b_j is largest.

6.2. Compatibility of the polyhedra. Here we verify that the residue method is applicable for the fan Σ to make use of Theorem [4.](#page-26-1) Let Π be a polyhedron and let H be a collection of r affine hyperplanes in $\mathfrak{t}_\mathbb{C}^{\vee}$ defined by equations $f_1 - is_1, \ldots, f_r - is_r$ for real linear maps $f_1, \ldots, f_r : \mathfrak{t}^{\vee} \to \mathbb{R}$ and constants s_1, \ldots, s_r with positive real parts. The associated Jacobian is $f(x) = f(x)$ $f(x)$

$$
J(H, \Pi) = \begin{bmatrix} f_1(v_1) & f_1(v_2) & \cdots & f_1(v_r) \\ f_2(v_1) & f_2(v_2) & \cdots & f_2(v_r) \\ \vdots & \vdots & \ddots & \vdots \\ f_r(v_1) & f_r(v_2) & \cdots & f_r(v_r) \end{bmatrix}
$$

where (v_1, \ldots, v_r) is the defining basis of Π . We verify that every every collection H_1, \ldots, H_r of linearly independent singular hyperplanes of $\omega = H_{\infty}(\chi, -s) dm$ is compatible with the polyhedron Π_{σ} for every $\sigma \in \Sigma_D(r)$. Such a collection corresponds to a choice $E =$ ${e_1, \ldots, e_r}$ of r forms in $\Sigma_w(1)$ whose restrictions πE to \mathfrak{t}^\vee are linearly independent.

The maximal cones in $\Sigma_D(r)$ are $\sigma_{(j)} = -\Lambda_j$ for $1 \leq j \leq r$, and the polyhedra to be integrated along are of the form $\Pi_j = \tilde{\mathfrak{t}}^{\vee} + i\sigma_{(j)}^{\vee}$ for $1 \leq j \leq r$. Any proper subset of $\Sigma(1)$ is Σ-convex, so any such collection E will project to the set $πE = σ(1)$ of generators for some $\sigma = \sigma_{(i)} \in \Sigma_{\Gamma}(r)$ (Lemma [1\)](#page-9-2). One can check that the Jacobian $J(\sigma, \Pi_i)$ is compatible for every $\sigma \in \Sigma_{\Gamma}(r)$ and $1 \leq j \leq r$. They also satisfy the conjecture of [\[14,](#page-43-4) Remark 3]. The Jacobians $J(\sigma, \Pi_i)$ follow the following pattern, e.g. for $r = 4$:

$$
J(\sigma_{(1)}, \Pi_1) = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \qquad J(\sigma_{(1)}, \Pi_2) = \begin{bmatrix} 1 & 0 & 0 & 0 \\ -1 & -1 & -1 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}
$$

$$
J(\sigma_{(1)}, \Pi_3) = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ -1 & -1 & -1 & -1 \\ 0 & 0 & 0 & 1 \end{bmatrix} \qquad J(\sigma_{(1)}, \Pi_4) = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -1 & -1 & -1 & -1 \end{bmatrix}
$$

$$
J(\sigma_{(2)}, \Pi_1) = \begin{bmatrix} 1 & 0 & 0 & 0 \\ -1 & -1 & -1 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \qquad J(\sigma_{(2)}, \Pi_2) = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}
$$

$$
J(\sigma_{(2)}, \Pi_3) = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ -1 & -1 & -1 & -1 \\ 0 & 0 & 0 & 1 \end{bmatrix} \qquad J(\sigma_{(2)}, \Pi_4) = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ -1 & -1 & -1 & -1 \end{bmatrix}
$$

36 ANDREW O'DESKY

$$
J(\sigma_{(\infty)}, \Pi_1) = \begin{bmatrix} -1 & -1 & -1 & -1 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \qquad J(\sigma_{(\infty)}, \Pi_2) = \begin{bmatrix} 0 & 1 & 0 & 0 \\ -1 & -1 & -1 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}
$$

$$
J(\sigma_{(\infty)}, \Pi_3) = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ -1 & -1 & -1 & -1 \\ 0 & 0 & 0 & 1 \end{bmatrix} \qquad J(\sigma_{(\infty)}, \Pi_4) = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -1 & -1 & -1 & -1 \end{bmatrix}.
$$

6.3. Finding the poles. Having covered $-\ell(\mathcal{N})$ with rational cones in Σ_D , we still need to compute $n^{-(s-s_{\gamma})}$ for each $\gamma \in Z_*$ in order to determine the poles of the multiple Dirichlet series appearing in the formula for $Z(s)$.

Proposition 8. Let $s \in (\mathbb{C}^{\Sigma(1)})^{\Gamma}$ be any element with corresponding Σ -linear map $\varphi: \mathfrak{t}_E \to$ $\mathbb C$. The flags in $Z_{-\Lambda_j}$ have a common terminal point which is the unique linear map $m_{-\Lambda_j}\in \mathfrak t_\mathbb C^\vee$ agreeing with $(2\pi i)^{-1}\varphi$ on $-\Lambda_j$. Each Γ -invariant Σ -linear function on \mathfrak{t}_E is determined by its values on e_0, f_1, \ldots, f_r . The function $\varphi - 2\pi i m_{-\Lambda_j}$ (corresponding to $s - s_\gamma$) takes zero on each of these elements except for f_i where it has the value $-(s_0 + \cdots + s_n)$.

Proof. Since $\varphi - 2\pi i m_{-\Lambda_j}$ vanishes identically on $-\Lambda_j$, it follows that $\varphi - 2\pi i m_{-\Lambda_j}$ is zero on e_0 and every f_k with $k \neq j$. Meanwhile

$$
2\pi i m_{-\Lambda_j}(f_j) = 2\pi i m_{-\Lambda_j}\left(-e_0 - \sum_{k \neq j} f_k\right) = -\varphi(e_0) - \sum_{\substack{e \notin O_j \\ e \neq e_0}} \varphi(e)
$$

and $\varphi(f_j) = \sum_{e \in O_j} \varphi(e)$ by Σ -linearity and convexity of the Galois action on Σ . Thus

$$
(\varphi - 2\pi i m_{-\Lambda_j})(f_j) = \varphi(f_j) - 2\pi i m_{-\Lambda_i}(f_j) = -(s_0 + \cdots + s_n).
$$

Recall the formula [\(5\)](#page-13-2) for the finite part of the Fourier transform of the height function:

$$
\widehat{H}_{\text{fin}}(\chi, -s, D) = \prod_{p \in P_{\mathbb{Q}}^{\text{fin}}} \sum_{n \in T(\mathbb{Q}_p)/K_p^D} \chi_p(n)^{-1} 1_v^D(n) p^{\frac{1}{ep}\varphi(\ell_w(n))} = \sum_{n \in \mathcal{N}} \chi(n)^{-1} n^{-s}.
$$

To deduce the location of the right-most pole of $Z(s)$ from Theorem [4,](#page-26-1) we first need a more precise description of the Euler factors of \widehat{H}_{fin} . For regular fans this is accomplished by [\[3,](#page-42-1) Prop. 2.2.3], but for our application to polynomials we must consider simplicial fans. When a cone in the fan is not regular, the local Euler factors are modified by an additional factor Q_{Δ} which generally contributes a pole to the Dirichlet series. However, we will show such poles occur to the left of the right-most pole. Let $\Delta = {\sum_{\chi \in G^{\vee}} a_{\chi} e_{\chi} : 0 \le a_{\chi} < 1}$. For each Galois orbit $O_j \subset \Sigma(1) - \{e_\infty\}$, let $T_j = R_{\mathbb{Q}}^{E_j} \mathbb{G}_m$ be the torus over \mathbb{Q} determined by the Galois permutation action on O_j . There are natural homomorphisms $T_j \to T$ of algebraic tori induced by the Galois equivariant lattice maps $\mathbb{Z}O_j \to N_E$.

Proposition 9. Let p be a finite place and let w be a place of E lying over p. Let $s \in (\mathbb{C}_{+}^{\Sigma(1)})^{\Gamma}$ and let χ be a character of $T(\mathbb{Q}_p)$ which is trivial on K_p^D . Let $O_1^w, \ldots, O_{r_w}^w$ be the Γ_w -orbits in $\{e_x\}_{x \in G}$. Then

$$
\widehat{H_p}(\chi, -s, D) = Q_p(\chi, s) \prod_{j=1}^{r_w} \left(1 - \chi(O_j^w)^{-1} p^{\frac{1}{e_p} \varphi(O_j^w)}\right)^{-1} = Q_p(\chi, s) \prod_{j=1}^r L_p(\chi_j, s_{O_j})
$$

where $\varphi(O_j^w) = \sum_{e \in O_j^w} \varphi(e), \ \chi(O_j^w) = \prod_{e \in O_j^w} \chi(e), \ s_{O_j} = -\varphi(e)$ for any $e \in O_j$, χ_j is the pullback of χ along $T_j(\mathbb{Q}_p) \to T(\mathbb{Q}_p)$, L_p is the Euler factor at p of the Hecke L-function of $\chi_j,\ and$

$$
Q_p(\chi, s) = \sum_{n \in \text{im } \ell_w \cap \Delta} \chi(n)^{-1} p^{\frac{1}{e_p} \varphi(n)}.
$$

In particular, the multiple Dirichlet series $\widehat{H}_{fin}(\chi, -s, D)$ for an automorphic character χ of level K^D is equal to

$$
Q(\chi, s) \prod_{j=1}^r L(\chi_j, s_{O_j})
$$

where $Q(\chi, s) = \prod_{p \in P_{\mathbb{Q}}^{\text{fin}}} Q_p(\chi, s)$. Set $o_k = |O_k|$. The multiple Dirichlet series $Q(\chi, s)$ converges absolutely on the half-space $\{\frac{1}{2}\text{Re}(o_2s_{O_2}+\cdots+o_r s_{O_r})>1\}\subset (\mathbb{C}_+^{\Sigma(1)})^{\Gamma}$, and the multiple Dirichlet series $\widehat{H}_{fin}(\chi, -s, D)$ converges absolutely on the polyhedral region

$$
\left\{\frac{1}{2}\text{Re}(o_2s_{O_2}+\cdots+o_r s_{O_r})>1\right\}\cap\left\{\text{Re}(s_{O_2})>1\right\}\cap\cdots\cap\left\{\text{Re}(s_{O_r})>1\right\}.
$$

If $\chi = \xi \psi \chi_m$ and $\xi \psi \neq 1$ then the multiple Dirichlet series $Q(\chi, s)$ converges conditionally on the half-space $\{ \text{Re}(o_2s_{O_2} + \cdots + o_r s_{O_r}) > 0 \} \subset (\mathbb{C}_+^{\Sigma(1)})^{\Gamma}$, and the multiple Dirichlet series $\widehat{H}_{fin}(\chi, -s, D)$ converges conditionally on the polyhedral region

$$
\{ \text{Re}(s_{O_2}) > 0 \} \cap \cdots \cap \{ \text{Re}(s_{O_r}) > 0 \}.
$$

Finally, there is a constant $c > 1$ such that for any $u \in \mathbb{C}$ with positive real part

$$
Q_p(1, us_{AC}) = \begin{cases} 1 + \phi(n)p^{-\frac{n-1}{2}u} + O(p^{-\frac{c(n-1)}{2}u}) & if p \text{ split in } E/\mathbb{Q}, \\ 1 + O(p^{-\frac{c(n-1)}{2ep}u}) & otherwise \end{cases}
$$
(18)

where ϕ is the Euler totient function, $\omega = \frac{1}{n}$ $\frac{1}{n}(e_2 + 2e_3 + \cdots + (n-1)e_n)$, and the implied constant is independent of p. If G has prime order, then this simplifies to

$$
Q_p(1,s) = \begin{cases} 1 + (n-1)p^{-\frac{n-1}{2}u} & \text{if } p \text{ split in } E/\mathbb{Q}, \\ 1 & \text{otherwise.} \end{cases}
$$

Proof. The first formula is obtained by applying a known formula for the exponential gen-erating function of integer points in polyhedra [\[1,](#page-42-4) p. 110]. The series $Q(\chi, s)$ is dominated by

$$
\prod_{p} \sum_{n \in N_E \cap \Delta} p^{\frac{1}{e_p} \varphi(n)}.
$$

The convergence of this series is determined by those nonzero elements of $N_E \cap \Delta$ for which $\text{Re}(\varphi(n))$ is smallest; these minima occur when n is in the Galois orbit of ω . We have that

$$
\pi \omega = \frac{1}{|\Gamma|} \sum_{\chi \in G^{\vee}} \ell(\chi(c^{-1})) \sum_{\gamma} \gamma e_{\chi}
$$

$$
= \sum_{j=2}^{b} \frac{1}{o_j} f_j \sum_{e_{\chi} \in O_j} \ell(\chi(c^{-1}))
$$

$$
= \sum_{j=2}^{b} \frac{1}{o_j} f_j \frac{o_j}{2} \qquad (n \ge 3)
$$

$$
= \frac{1}{2} (f_2 + \dots + f_r).
$$

Since the Galois action on Σ is convex we know that $\varphi(\omega') = \varphi(\pi \omega)$ for any ω' in the Galois orbit of ω , and thus $\varphi(\omega') = \frac{1}{2}(\varphi(Q_2) + \cdots + \varphi(Q_r)) = -\frac{1}{2}$ $\frac{1}{2}(o_2s_{O_2}+\cdots+o_r s_{O_r})$. The claimed regions of convergence for Q and \hat{H}_{fin} now follow from the well-known regions of convergence for Hecke L-functions.

Now we prove the last claim. The formula for Q_p follows from summing over $n \in N_E \cap \Delta$ ordered by $\text{Re}(\varphi(n))$. The minimal value of $\frac{1}{2}\text{Re}(o_2s_{O_2} + \cdots + o_r s_{O_r})$ is $\frac{n-1}{2}\text{Re}(u)$ which is achieved on the Galois orbit of ω which has size $[E : \mathbb{Q}] = \phi(n)$. The set im $\ell_w \cap \Delta$ is contained in $N_E \cap \Delta$ which has $[N_E : N'_E] = n$ elements where $N'_E = \mathbb{Z}\langle e_1, \ldots, e_n \rangle$. If G has prime order then the orbit of ω under $\overline{\Gamma} = \text{Gal}(\mathbb{Q}(\zeta_n)/\mathbb{Q})$ has size $n-1$. Thus if p is split then im $\ell_w \cap \Delta$ entirely consists of $0 \in N_E$ and the Galois orbit of ω . As ω is moved by any nontrivial element of Γ , the set im $\ell_w \cap \Delta$ is $\{0\}$ if p is not split. nontrivial element of Γ, the set im $\ell_w \cap \Delta$ is $\{0\}$ if p is not split.

6.4. **Parametrizing toric ideals.** We will use the above description of \widehat{H}_p to describe toric ideals in terms of ideals of number fields. To ease the analysis, we will restrict attention to those toric ideals which are needed to determine the asymptotic rate of growth of the sum of Dirichlet coefficients of the height zeta function $Z(s)$. Let Sp_E denote the set of square-free positive integers which are only divisible by primes that are split in E/\mathbb{Q} . By the description of the Euler factor [\(18\)](#page-36-0), we see that primes which are not split in E/\mathbb{Q} may be ignored in the sense that

$$
\frac{\prod_{p \in Sp_E} (1 + \phi(n)p^{-\frac{n-1}{2}u})}{Q(1,us_{AC})}
$$

is regular and nonvanishing at $u = (\frac{n-1}{2})^{-1}$. We set

$$
Q_1(u) = \prod_{p \in Sp_E} (1 + \phi(n)p^{-\frac{n-1}{2}u}) = \sum_{M \in Sp_E} \phi(n)^{\omega(M)} M^{-\frac{1}{2}(n-1)u}
$$

where $\omega(M)$ is the number of distinct prime factors of M. Let $\mathcal{N}_1 \subset \mathcal{N}$ denote the subset of toric ideals in bijection with $Sp_E \times \text{Ideals}(O_{E_1}) \times \cdots \times \text{Ideals}(O_{E_r})$. We will regard elements of \mathcal{N}_1 as weighted by the factor $\phi(n)^{\omega(M)}$. For any $n \in \mathcal{N}_1$ we have that

$$
n^{-s} = M^{-\frac{1}{2}(o_2 s_{O_2} + \dots + o_r s_{O_r})} N(I_1)^{-s_{O_1}} \dots N(I_r)^{-s_{O_r}}
$$
(19)

and

$$
\sum_{n \in \mathcal{N}_1} n^{-s} = \sum_{M, I_1, \dots, I_r} \phi(n)^{\omega(M)} M^{-\frac{1}{2}(\log_{\mathcal{O}_2} + \dots + \log_{\mathcal{O}_r})} N(I_1)^{-s_{O_1}} \cdots N(I_r)^{-s_{O_r}}
$$

where $I_1 \times \cdots \times I_r$ is an ideal of $O_{E_1} \times \cdots \times O_{E_r}$. (The formula for n^{-s} in terms of norms of ideals is explained at the end of Example [2.](#page-14-2))

6.5. Proof of the main theorem on polynomials. We begin by restricting $Z(s)$ to the complex line spanned by the anticanonical line bundle $[-K] \in Pic(Y)$. Let $s_{AC} \in (\mathbb{Z}^{\Sigma(1)})^{\Gamma}$ denote the element with all coordinates equal to 1. It is well-known that s_{AC} maps to $[-K]$ under the canonical map $(\mathbb{Z}^{\Sigma(1)})^{\Gamma} \to Pic(Y)$. Let $s = us_{AC}$ for $u \in \mathbb{C}$ with sufficiently large real part. We write $Z(us_{AC})=\sum_{N}a_{N}N^{-(n+1)u}$ and let $A(X)=\sum_{N.$

By the Poisson formula,

$$
Z(us_{AC}) = \frac{|\mu|}{|C||R|P} \sum_{\xi \in R^{\vee}} \sum_{\psi \in C^{\vee}} \int_{\mathfrak{t}^{\vee}} \widehat{H}_{\text{fin}}(\xi \psi \chi_m, -s, D) \widehat{H}_{\infty}(\xi \chi_m, -s) \, dm = \sum_{\xi, \psi} Z_{\xi \psi}(u)
$$

where

$$
Z_{\xi\psi}(u) = \frac{|\mu|}{|C||R|P} \int_{\mathfrak{t}^{\vee}} \widehat{H}_{\text{fin}}(\xi\psi\chi_m, -s, D)\widehat{H}_{\infty}(\xi\chi_m, -s) \, dm.
$$

We evaluate $Z_{\xi\psi}$ by distributing the integral over \mathfrak{t}^\vee into the sum over toric ideals and then applying the residue formula (Theorem [3\)](#page-18-2). (The hypotheses for the residue formula were verified in §[6.2.](#page-34-0)) This results in the $\xi\psi$ -summand of the formula for $Z(s)$ in Theorem [4,](#page-26-1) which is

$$
Z_{\xi\psi}(u) = \frac{|\mu|}{|C||R|P} \sum_{\tau \in \Sigma_D} \sum_{\gamma \in Z_{\sigma_{\tau}}} L(s + 2\pi i m_{\gamma\xi}, (\xi\psi)^{-1}, -\tau^{\circ}, D) R_{\gamma}(us_{AC} + 2\pi i m_{\xi}). \tag{20}
$$

To determine the main term of $A(X)$ it suffices to consider maximal cones σ in the sum over cones $\tau \in \Sigma_D$. Since the difference between a cone σ and σ° is formed from lower-dimensional cones, to compute the main term we may use either σ or σ° . So we turn to the Dirichlet series $L(s+2\pi im_{\gamma\xi},(\xi\psi)^{-1},-\sigma,D)$ where σ is a maximal cone of Σ_D . The quantity $s+2\pi im_{\gamma\xi}$ may be evaluated with the help of Lemma [6](#page-18-1) which shows that $2\pi i m_{\gamma\xi} = \mu_{E_{\gamma}}(\varphi|_{\tau}) - 2\pi i \mu_{E_{\gamma}} m_{\xi}$ where $E_{\gamma} \subset \Sigma_w(1)$ is any defining set of generators for γ . Since the Galois action is convex we have $\mu_{E_{\gamma}}(\varphi|_{\tau}) = \varphi|_{\pi E_{\gamma}}$ where $\varphi|_{\pi E_{\gamma}} \in \mathfrak{t}^{\vee} \otimes \mathbb{C}$ is the unique linear map agreeing with φ on the basis πE_γ . Since the Galois action is strongly convex, $\varphi|_{\pi E_\gamma} = \varphi|_{\sigma}$ where $\sigma = \langle \pi E_\gamma \rangle$ is a maximal cone of Σ_D . Let $s_\gamma \in (\mathbb{C}^{\Sigma(1)})^{\Gamma}$ correspond to $\varphi|_{\sigma}$ so $s + 2\pi i m_\gamma = s - s_\gamma$. We conclude that $s + 2\pi i m_{\gamma\xi} = s - s_{\sigma} - 2\pi i \mu_{E_{\gamma}} m_{\xi}$ where $s_{\sigma} = s_{\gamma}$ only depends on σ .

The maximal cones of Σ_D were described in §[6.1:](#page-33-1) they have the form $\sigma_{(j)} = -\Lambda_j$ for $1 \leq j \leq r$ where Λ_j consists of those elements $b_1f_1 + \cdots + b_rf_r \in \mathfrak{t}$ with $b_i \geq 0$ and b_i maximal. We may deduce the main term up to a constant from the part of $L(s$ $s_{\sigma} - 2\pi i \mu_{E_{\gamma}} m_{\xi}, (\xi \psi)^{-1}, -\sigma, D$ summed over the subset $\mathcal{N}_1 \subset \mathcal{N}$. Namely, if c is the constant of the main term of the summatory function of the Dirichlet series $L(s - s_{\sigma} 2\pi i\mu_{E_\gamma}m_{\xi}$, $({\xi\psi})^{-1}, -\sigma, D$ summed over the subset $\mathscr{N}_1 \subset \mathscr{N}$, then the constant of the main term for the summatory function of $L(s - s_{\sigma} - 2\pi i\mu_{E_{\gamma}}m_{\xi}, (\xi\psi)^{-1}, -\sigma, D)$ without restricting to $\mathcal{N}_1 \subset \mathcal{N}$ may be obtained by multiplying c by the value of $Q(1, u s_{AC})/Q_1(u)$ at the abscissa of convergence (cf. §[6.4\)](#page-37-0). We saw in §[6.4](#page-37-0) that each toric ideal $n \in \mathcal{N}_1$ corresponds to a tuple (M, I_1, \ldots, I_r) where $M \in Sp_E$ and $I_1 \times \cdots \times I_r$ is an ideal of $O_{E_1} \times \cdots \times O_{E_r}$. The formulas from §[6.4](#page-37-0) imply that

$$
\ell(n) = f_1 \log N(I_1) + \frac{1}{o_2} f_2 \left(\log N(I_2) + \frac{1}{2} o_2 \log M \right) + \dots + \frac{1}{o_r} f_r \left(\log N(I_r) + \frac{1}{2} o_r \log M \right).
$$
\n(21)

First we consider Λ_1 . From [\(21\)](#page-38-0) we see that $\ell(n) \in \Lambda_1$ if and only if

$$
N(I_k)\sqrt{M}^{o_k} \le N(I_1)^{o_k} \quad \text{for all } k \neq 1.
$$

Similarly $\ell(n) \in \Lambda_{\ell}$ for $2 \leq \ell \leq r$ if and only if

$$
N(I_1)^{o_\ell} \le N(I_\ell) \sqrt{M}^{o_\ell}, \quad N(I_k)^{\frac{o_\ell}{o_k}} \le N(I_\ell) \quad \text{for all } k \ne 1.
$$

We sum over tuples (M, I_1, \ldots, I_r) satisfying one of the above systems of inequalities to determine the Dirichlet coefficients. First note that

$$
L(s - s_{\sigma} - 2\pi i\mu_{E_{\gamma}} m_{\xi}, (\xi\psi)^{-1}, -\sigma, D) = L(s - s_{\sigma}, (\xi\psi\chi)^{-1}, -\sigma, D)
$$

where $\chi(n) = n^{-2\pi i \mu_{E_\gamma} m_{\xi}}$ is a Hecke character of level K^D . For the maximal cone $\sigma = -\Lambda_1$ we use [\(19\)](#page-37-1) and Proposition [8](#page-35-1) to see that

$$
n^{-(s-s_{\sigma})} = N^{-t}, \quad N = N(I_1), \quad t = s_0 + \cdots + s_n = (n+1)u.
$$

From the inequalities defining Λ_1 we find that the Dirichlet coefficient of N^{-t} is

$$
\sum_{M=1}^{N^2} 1_{Sp_E}(M)\phi(n)^{\omega(M)} \prod_{k=2}^r \left(\sum_{\substack{I_k \subset O_{E_k} \\ 1 \le N(I_k) \le (N/\sqrt{M})^{o_k}}} (\xi \psi \chi)_k (I_k)^{-1} \right).
$$
 (22)

Let $\delta_{(\xi\psi)_k} = 1$ if $(\xi\psi)_k = 1$ and otherwise 0. We also set $\beta_k := 1 - \frac{2}{1+\epsilon}$ $\frac{2}{1+\mathfrak{o}_k}$. By known zero-free regions of Hecke L-functions this is

$$
\sum_{M=1}^{N^2} 1_{Sp_E}(M) \phi(n)^{\omega(M)} \prod_{k=2}^r (\text{res}[\zeta_{E_k}] \delta_{(\xi \psi)_k} (N/\sqrt{M})^{o_k} + O((N/\sqrt{M})^{o_k \beta_k})).
$$

This is largest when $(\xi \psi)_k = 1$ for all k, which occurs if and only if $\xi \psi \in \ker \gamma^*$ where γ is the surjective homomorphism $P \to T$ from §[2.5.](#page-15-0) Since ker γ^* is torsion and R^{\vee} is free we must have $\xi = 1$ if $\xi \psi \in \text{ker } \gamma^*$. For any such $\xi \psi = \psi \in \text{ker } \gamma^*$ the Dirichlet coefficient is

$$
\text{res}[\zeta_{E_2}] \cdots \text{res}[\zeta_{E_r}] \sum_{M=1}^{N^2} 1_{Sp_E}(M) \phi(n)^{\omega(M)} (N/\sqrt{M})^{n-1} ((1+O((N/\sqrt{M})^{\sum_k o_k \beta_k - (n-1)})).
$$
 (23)

Observe that

 $\overline{1}$

$$
\sum_{M=1}^{N^2} 1_{Sp_E}(M)\phi(n)^{\omega(M)}(N/\sqrt{M})^{n-1} = N^{n-1} \prod_{p \in Sp_E} \left(1 + \phi(n)p^{-\frac{n-1}{2}}\right)
$$

and

$$
\sum_{M=1}^{N^2} 1_{Sp_E}(M) \phi(n)^{\omega(M)} (N/\sqrt{M})^{n-1} O((N/\sqrt{M})^{\sum_k o_k \beta_k - (n-1)})
$$

= $N^{\sum_k o_k \beta_k} \sum_{M=1}^{N^2} 1_{Sp_E}(M) \phi(n)^{\omega(M)} O((1/\sqrt{M})^{\sum_k o_k \beta_k}).$

We have that $\sum_{k} o_k \beta_k = n - 1 - 2 \sum_{k|n,k>1} (1 + \phi(k))^{-1} > n - 1 - 2 \sum_{k|n,k>1} 1/k \ge 2$, so this secondary term is bounded by $O_{\varepsilon}(N^{\sum_k o_k \beta_k + \varepsilon}) = O_{\delta}(N^{n-1-\delta})$ for some $\delta > 0$. In conclusion, the coefficient of N^{-t} in the multiple Dirichlet series $L(s - s_{\sigma}, (\xi \psi \chi)^{-1}, -\sigma, D)$ restricted to \mathcal{N}_1 if $\xi = 1$ and $\psi \in \ker \gamma^*$ is

$$
\operatorname{res}[\zeta_{E_2}]\cdots\operatorname{res}[\zeta_{E_r}]\prod_{p\in Sp_E}\left(1+\phi(n)p^{-\frac{n-1}{2}}\right)N^{n-1}(1+O_{\delta}(N^{-1-\delta})),
$$

the coefficient of N^{-t} for $L(s - s_{\sigma}, (\xi \psi \chi)^{-1}, -\sigma, D)$ if $\xi = 1$ and $\psi \in \ker \gamma^*$ is

$$
\operatorname{res}[\zeta_{E_2}] \cdots \operatorname{res}[\zeta_{E_r}] \left(\prod_p \sum_{n \in \operatorname{im} \ell_w \cap \Delta} p^{\frac{1}{e_p} \varphi_{AC}(n)} \right) N^{n-1} (1 + O_\delta(N^{-1-\delta})), \tag{24}
$$

while the coefficient for $L(s - s_{\sigma}, (\xi \psi \chi)^{-1}, -\sigma, D)$ if $\xi \psi \notin \ker \gamma^*$ is

$$
\mathrm{res}[\zeta_{E_2}]\cdots\mathrm{res}[\zeta_{E_r}]\left(\prod_p\sum_{n\in\mathrm{im}\,\ell_w\cap\Delta}p^{\frac{1}{e_p}\varphi_{AC}(n)}\right)N^{n-1}O_\delta(N^{-1-\delta}).
$$

The other regions may be handled similarly. For $\sigma = -\Lambda_{\ell}$, $2 \leq \ell \leq r$, we find

$$
n^{-(s-s_{\sigma})} = N^{-t}, \quad N = N(I_{\ell})^{1/o_{\ell}} \sqrt{M}.
$$

Once again the coefficient of N^{-t} is largest if and only if $\xi \psi \in \ker \gamma^*$, in which case the coefficient is equal to

$$
\sum_{M=1}^{N^2} 1_{Sp_E}(M) \phi(n)^{\omega(M)} \left(\sum_{\substack{I_1 \subset O_{E_1}\\1\leq N(I_1)\leq N}}1\right) \left(\sum_{\substack{I_\ell \subset O_{E_\ell}\\N(I_\ell)=(N/\sqrt{M})^{o_\ell}}1}\right) \prod_{\substack{2\leq k\leq r\\k\neq \ell}}\left(\sum_{\substack{I_k \subset O_{E_k}\\1\leq N(I_k)\leq (N/\sqrt{M})^{o_k}}1}\right).
$$

This is bounded by

$$
\sum_{M=1}^{N^2} 1_{Sp_E}(M)\phi(n)^{\omega(M)}N^{1+\varepsilon} \prod_{\substack{2\leq k\leq r\\k\neq\ell}} ((N/\sqrt{M})^{o_k} + O((N/\sqrt{M})^{o_k \beta_k}))
$$
\n
$$
= \sum_{M=1}^{N^2} 1_{Sp_E}(M)\phi(n)^{\omega(M)}N^{1+\varepsilon}(N/\sqrt{M})^{n-1-o_\ell}(1+O((N/\sqrt{M})^{\sum_{k\neq\ell,1}o_k\beta_k-(n-1-o_\ell)}))
$$
\n
$$
= N^{n-o_\ell+\varepsilon} \sum_{M=1}^{N^2} 1_{Sp_E}(M)\phi(n)^{\omega(M)}(1/\sqrt{M})^{n-1-o_\ell}(1+O((N/\sqrt{M})^{\sum_{k\neq\ell,1}o_k\beta_k-(n-1-o_\ell)})).
$$

We have that

$$
\sum_{M=1}^{N^2} 1_{Sp_E}(M)\phi(n)^{\omega(M)}(1/\sqrt{M})^{n-1-o_\ell} = O_\varepsilon(N^{2+\varepsilon})
$$

for any $\varepsilon > 0$ and

$$
O(N^{\sum_{k \neq \ell, 1} o_k \beta_k - (n-1-o_{\ell})}) \sum_{M=1}^{N^2} 1_{Sp_E}(M) \phi(n)^{\omega(M)} = O_{\delta}(N^{2-\delta})
$$

for some $\delta > 0$. Thus the coefficient of N^{-t} in the multiple Dirichlet series for Λ_{ℓ} with $\ell \neq 1$ is $O_{\delta}(N^{4-\delta})$ for some $\delta > 0$.

42 ANDREW O'DESKY

Now we determine the main term of $A(X)$. The summands of [\(20\)](#page-38-1) for non-maximal cones are bounded by $O(N^{n-1})$ which is smaller than the secondary terms $O_{\delta}(N^{n-\delta})$ of the summands for maximal cones. When $\xi \psi \in \ker \gamma^*$, the estimate [\(24\)](#page-40-0) shows the abscissa of absolute convergence of the Dirichlet series $L(s - s_{\sigma}, (\xi \psi \chi)^{-1}, -\sigma, D)$ is at $u = \frac{n}{n+1}$. Since $R_{\gamma}(us_{AC})$ is regular and nonvanishing (Proposition [4\)](#page-21-0) at $u = \frac{n}{n+1}$, the leading coefficient of $Z(u)$ in the Laurent expansion at $u = \frac{n}{n+1}$ is also the leading coefficient of

$$
\frac{|\mu|}{|C||R|P} \sum_{\sigma \in \Sigma_D(r)} \left(\sum_{\gamma \in Z_{\sigma}} R_{\gamma}(\frac{n}{n+1} s_{AC}) \right) \sum_{\psi \in \ker \gamma^*} L(s - s_{\sigma}, \psi^{-1}, -\sigma, D) \tag{25}
$$

in the asymptotic expansion as u tends to $\frac{n}{n+1}$ from above. For maximal cones $\sigma \in \Sigma_D(r)$ and $\tau \in \sum_{w} (r_w)$ let $\begin{pmatrix} \tau \\ \sigma \end{pmatrix}$ $(\tau) \in \mathbb{Z}_{\geq 0}$ denote the number of ordered r-subsets $E \subset \tau(1)$ such that $\pi E = \sigma(1)$ (as unordered sets) and the flag γ defined by E is in Z_{σ} .^{[6](#page-41-0)} Say $\sigma = \sigma_{(j)}$. With the help of Lemma [7](#page-19-4) and Proposition [8](#page-35-1) one finds that

$$
\sum_{\gamma \in Z_{\sigma}} R_{\gamma}(\frac{n}{n+1} s_{AC}) = \left(\frac{o_j}{2n}\right)^{r_w - r} \sum_{\tau \in \Sigma_w(r_w)} \left(\frac{\tau}{\sigma}\right) \frac{[N_w : \mathbb{Z}\tau(1)]}{[N : \mathbb{Z}\sigma(1)]}.
$$

The ratio of the Dirichlet series [\(25\)](#page-41-1) and $Z(u)$ tends to 1 as u tends to $\frac{n}{n+1}$ from above, so their summatory functions have the same main terms. Set

$$
\kappa_0 = \text{res}[\zeta_{E_2}] \cdots \text{res}[\zeta_{E_r}] \prod_p \sum_{n \in \text{im } \ell_w \cap \Delta} p^{\frac{1}{e_p} \varphi_{AC}(n)} \quad \text{and} \quad \kappa_\sigma = \left(\frac{o_j}{2n}\right)^{r_w - r} \sum_{\tau \in \Sigma_w(r_w)} \binom{\tau}{\sigma} \frac{[N_w : \mathbb{Z}\tau(1)]}{[N : \mathbb{Z}\sigma(1)]}.
$$

From the asymptotic formula [\(24\)](#page-40-0) for the coefficients of $L(s - s_{\sigma}, \psi^{-1}, -\sigma, D)$ we conclude that the summatory function $A(X)$ of $Z(us_{AC})$ satisfies

$$
A(X) = \frac{|\mu||C \cap \ker \gamma^*|}{|C||R|P} \kappa_0 \left(\sum_{\sigma \in \Sigma_D(r)} \kappa_\sigma \right) \frac{1}{n} X^{\frac{n}{n+1}} (1 + o(1)).
$$

To complete the proof, we need to relate $A(X)$ to the count $N(C_n, X)$ of normal C_n polynomials. Recall the rational points of T are in bijection with equivalence classes of pairs $(K/\mathbb{Q}, x)$ where $K = \mathcal{O}(T)$ is the ring of functions on a G-torsor over Spec \mathbb{Q} and $x \in K$ is a normal element (Proposition [6\)](#page-30-1). The number of twists of $(K/\mathbb{Q}, x)$ for a C_n -field K by an outer automorphism of C_n is equal to $\varphi(n)$, so there are $\varphi(n)$ such rational points of T giving rise to any C_n -polynomial (Corollary [2\)](#page-32-1).

We bound the number of reducible rational points on T . Recall $\mathcal G$ is a product of simple tori indexed by the Q-irreducible representations of C_n . In particular, $\mathcal{G} = \mathbb{G}_m \times \mathcal{G}_1$ where \mathbb{G}_m corresponds to the trivial representation. The monodromy group of a rational point $(K/\mathbb{Q}, x)$ is contained in a given subgroup $D \subsetneq C_n$ if and only if $(K/\mathbb{Q}, x)$ is the image of a rational point on \mathcal{G}/D via the natural quotient map $\mathcal{G}/D \to \mathcal{G}/G = T$. Let $\Upsilon =$ $\Upsilon_D \subset T(\mathbb{Q})_{D\text{-integral}} = T(\mathbb{Q}) \cap U(\mathbb{Z})$ be the subset of such points which are D-integral. By expressing this quotient map as $\mathbb{G}_m \times \mathcal{G}_1/D \to \mathbb{G}_m \times T_1$ where $T_1 = \mathcal{G}_1/G$ we see that the projection of Υ to the T_1 factor of T is itself a thin subset $\Upsilon_1 \subset T_1$ of type II. We make use of ideas from $|7|$ to show Υ has vanishing density. It suffices to show that it has a vanishing

⁶Since the Jacobians of Σ satisfy the conjecture of [\[14,](#page-43-4) Remark 3] (cf. §[6.2\)](#page-34-0), the integer $\binom{\tau}{\sigma}$ can more simply be defined as the number of unordered r-subsets $E \subset \tau(1)$ such that $\pi E = \sigma(1)$.

contribution to the main term [\(23\)](#page-39-0) of the Dirichlet coefficients of $Z(us_{AC})$. For a finite place v let $\Phi_v: T(\mathbb{Q}_v)$ integral $\to \{0,1\}$ be the characteristic function defined by

$$
\Phi_v(x_v) = \begin{cases} 1 & x_v \in \Upsilon \pmod{p_v}, \\ 0 & \text{otherwise.} \end{cases}
$$

Set $\Phi = \Phi_{\infty} \times \prod_{v \text{ finite}} \Phi_v$ where Φ_{∞} is identically one on $T(\mathbb{R})$. The same computation we carried out for $Z(us_{AC}) = Z(us_{AC}, 1_D$ goes through for $Z(us_{AC}, \Phi 1_D) = \sum_{t \in T(\mathbb{Q})}$ $\Phi(t)1_D(t)$ $H(t,s)$ since Φ is also K^D-invariant, so the Poisson formula for $Z(us_{AC},\Phi 1_D)$ involves the same integral over level K^D automorphic characters. From [\(22\)](#page-39-1) we see the contribution to the leading term in the Dirichlet coefficients of $Z(us_{AC})$ is

$$
\sum_{M=1}^{N^2} 1_{Sp_E}(M) \phi(n)^{\omega(M)} \prod_{k=2}^r \left(\sum_{\substack{I_k \subset O_{E_k} \\ 1 \leq N(I_k) \leq (N/\sqrt{M})^{o_k}}} \Phi(I_k)\right)
$$

where $\Phi(I_k)$ is evaluated using the bijection between $n \in \mathcal{N}_1$ and $(M, I_1, \ldots, I_r) \in Sp_E \times$ Ideals $(O_{E_1}) \times \cdots \times \text{Ideals}(O_{E_r})$. Note that [\(22\)](#page-39-1) only depends on the projection $\pi(n)$, so we are really counting points in the thin set Υ_1 in T_1 . We use the strategy from the proof of [\[7,](#page-43-18) Theorem 1.2. There is a constant $0 < c < 1$ and an extension L/\mathbb{Q} only depending on Υ_1 such that for all primes p which split in L the number of points in $\Upsilon_1 \pmod{p}$ is at most $cp^{\dim T_1} = cp^{n-1}$ by [\[7,](#page-43-18) Lemma 3.8]. Thus [\(22\)](#page-39-1) is bounded by

$$
\sum_{M=1}^{N^2} 1_{Sp_E}(M) \phi(n)^{\omega(M)} c^{\omega_L(N(I_2)\cdots N(I_r))} (N/\sqrt{M})^{n-1}
$$

where $\omega_L(N)$ is the number of distinct prime divisors of N which are split in L. Since $0 < c < 1$ this is $o(N^{n-1})$ which shows these points have vanishing density.

Remark 20. We expect the asymptotic formula of Theorem [2](#page-4-1) is also the correct formula for the count of all (monic integral) C_n -polynomials since non-normal polynomials form a thin set of \mathbb{P}/C_n . Non-normal C_n -polynomials such as t^3-3t-1 are the images of rational points on the boundary locus $\mathbb{P}/C_n - D$. This closed set is a union of toric divisors defined over Q so the distribution of its rational points is amenable to toric methods in principle, however these toric divisors may be geometrically reducible so they are not themselves toric varieties. The geometrically irreducible toric divisors in $\mathbb{P}/C_n - D$ are toric varieties but defined over cyclotomic fields. We expect the methods of this paper, suitably extended to more general ground fields, would verify that these thin sets have vanishing density. We similarly expect that an analysis of the height zeta function for the toric orbit parametrization from §[5](#page-32-0) would produce an asymptotic formula for G-polynomials of bounded root height for any finite abelian group G.

REFERENCES

- [1] A. Barvinok. *Integer points in polyhedra*. Zur. Lect. Adv. Math. Zürich: European Mathematical Society (EMS), 2008.
- [2] V. V. Batyrev and Y. Tschinkel. Manin's conjecture for toric varieties, 1995.
- [3] V. V. Batyrev and Y. Tschinkel. Rational points of bounded height on compactifications of anisotropic tori. Internat. Math. Res. Notices, (12):591–635, 1995.

44 ANDREW O'DESKY

- [4] V. V. Batyrev and Y. Tschinkel. Height zeta functions of toric varieties. *https: // math. nyu. edu/ ~ [tschinke/ papers/ yuri/ 96march/ march96. pdf](https://math.nyu.edu/~tschinke/papers/yuri/96march/march96.pdf)* , 1996.
- [5] V. V. Batyrev and Y. Tschinkel. Manin's conjecture for toric varieties. J. Algebr. Geom., 7(1):15–53, 1998.
- [6] S. Bhattacharya and A. O'Desky. On monic abelian trace-one cubic polynomials, 2023.
- [7] T. Browning and D. Loughran. Sieving rational points on varieties. Trans. Amer. Math. Soc., 371(8):5757–5785, 2019.
- [8] A. Chambert-Loir and Y. Tschinkel. Integral points of bounded height on toric varieties, 2010.
- [9] H. Cohen, F. Diaz y Diaz, and M. Olivier. On the density of discriminants of cyclic extensions of prime degree. J. Reine Angew. Math., 550:169–209, 2002.
- [10] P. K. J. Draxl. L-Funktionen algebraischer Tori. J. Number Theory, 3:444–467, 1971.
- [11] G. B. Folland. A course in abstract harmonic analysis. Studies in Advanced Mathematics. CRC Press, Boca Raton, FL, 1995.
- [12] W. Fulton. Introduction to toric varieties, volume 131 of Annals of Mathematics Studies. Princeton University Press, Princeton, NJ, 1993. The William H. Roever Lectures in Geometry.
- [13] A. O'Desky. On trace-one generators of abelian cubic fields, 2024.
- [14] A. O'Desky. A residue formula for integrals with hyperplane singularities, 2024.
- [15] A. O'Desky and J. Rosen. The moduli space of G-algebras. Advances in Mathematics, 431:109240, 2023.
- [16] T. Ono. On some arithmetic properties of linear algebraic groups. Ann. of Math. (2), 70:266–290, 1959.
- [17] T. Ono. Arithmetic of algebraic tori. Ann. of Math. (2), 74:101–139, 1961.
- [18] T. Santens. Manin's conjecture for integral points on toric varieties, 2023.
- [19] F. Shahidi. Eisenstein series and automorphic L-functions, volume 58 of Colloq. Publ., Am. Math. Soc. Providence, RI: American Mathematical Society (AMS), 2010.
- [20] A. K. Tsikh. Multidimensional residues and their applications. Transl. from the Russian by E. J. F. Primrose. Transl. edited by S. Gelfand, volume 103 of Transl. Math. Monogr. Providence, RI: American Mathematical Society, 1992.
- [21] V. E. Voskresenskii. Birational properties of linear algebraic groups. Math. USSR, Izv., 4:1–17, 1971.
- [22] F. Wilsch. Integral points of bounded height on a certain toric variety. Trans. Am. Math. Soc., Ser. B, 11:567–599, 2024.

Department of Mathematics PRINCETON UNIVERSITY Princeton, NJ 08544-1000, USA

Email address: andy.odesky@gmail.com