THE LAPLACIAN WITH COMPLEX MAGNETIC FIELDS

DAVID KREJČIŘÍK, THO NGUYEN DUC, AND NICOLAS RAYMOND

ABSTRACT. We study the two-dimensional magnetic Laplacian when the magnetic field is allowed to be complex-valued. Under the assumption that the imaginary part of the magnetic potential is relatively form-bounded with respect to the real part of the magnetic Laplacian, we introduce the operator as an m-sectorial operator. In two dimensions, sufficient conditions are established to guarantee that the resolvent is compact. In the case of non-critical complex magnetic fields, a WKB approach is used to construct semiclassical pseudomodes, which do not exist when the magnetic field is real-valued.

1. INTRODUCTION

Since the appearance of the highly influential paper [1] at the turn of the millennium, there has been a growing interest in Schrödinger operators with complex-valued electric potentials. This is explained not only by diverse physical motivations, but notably due to a new concept of quantum mechanics where observables can be represented by non-self-adjoint operators [26, 21]. Moreover, the mathematical studies lead to unprecedented spectral properties such as the existence of pseudomodes [9, 10, 20, 3] or the existence of eigenvalues accumulating at non-zero points of the essential spectrum [4, 8, 5].

It is about time that someone would address the case of Schrödinger operators where the magnetic potential is complex-valued now. This is the subject of the present paper. Our motivation is not merely our mathematical curiosity, but also strong physical motivations. Among these, let us mention superconductors [13, 14], quantum statistical physics [24, 2], speculations about a novel type of magnetic monopoles [25], stability of black holes in general relativity [16, 17] and the concept of quasi-self-adjointness again [19]. What is more, complexifying the magnetic field is mathematically much more challenging than its electric counterpart. That is probably why rigorous results do not exist in the literature and the objective of this paper is to fill in this gap.

1.1. The problem. We are concerned with the differential expression

$$(-ih\nabla - \mathbf{A})^2 = \sum_{j=1}^d \left(-ih\partial_{x_j} - A_j\right)^2, \qquad (1.1)$$

where $d \ge 1$ is the dimension, $\mathbf{A} = (A_1, \ldots, A_d) : \mathbb{R}^d \to \mathbb{C}^d$ is the complexified (magnetic) vector potential and h is a small positive (semiclassical) parameter. If the imaginary part of \mathbf{A} is non-zero, it is not even clear that (1.1) leads to a well defined operator in $L^2(\mathbb{R}^d)$. More specifically, our first concern is

(I) to identify the right subspace of $L^2(\mathbb{R}^d)$ as a domain on which (1.1) is realised as a closed operator with non-empty resolvent set.

Our next curiosity is about the realisation of complexified "magnetic bottles", *i.e.*,

(II) to find conditions on **A** which guarantee that the operator has compact resolvent. Since it is the pseudospectrum which describes non-self-adjoint phenomena, our last task is

(III) to construct pseudomodes in the semiclassical limit $h \to 0$.

Of course, other problems could be raised, but already these three tasks lead to considerable mathematical challenges.

1.2. Main results.

1.2.1. The magnetic Laplacian as an m-sectorial operator. In this paper, we deal with task (I) by assuming that the imaginary part of the vector potential is relatively formbounded with respect to the real part of the magnetic Laplacian. More specifically, we always make the following hypothesis.

Assumption 1.1. Let $\mathbf{A} \in L^2_{loc}(\mathbb{R}^d, \mathbb{C}^d)$ and assume that there exist two constants $a \in (0, 1)$ and $b \ge 0$ such that

$$\int_{\mathbb{R}^d} |(\operatorname{Im} \mathbf{A})u|^2 \, \mathrm{d}x \leqslant a \int_{\mathbb{R}^d} |(-ih\nabla - \operatorname{Re} \mathbf{A})u|^2 \, \mathrm{d}x + b \int_{\mathbb{R}^d} |u|^2 \, \mathrm{d}x \,, \tag{1.2}$$

for all $u \in C_c^{\infty}(\mathbb{R}^d)$.

We define the sesquilinear form naturally associated with (1.1), *i.e.*,

$$Q_{h,\mathbf{A}}(u,v) := \int_{\mathbb{R}^d} (-ih\nabla - \mathbf{A})u \cdot \overline{(-ih\nabla - \mathbf{A})v} \, \mathrm{d}x \,, \tag{1.3}$$
$$\mathrm{Dom}(Q_{h,\mathbf{A}}) := \left\{ u \in L^2(\mathbb{R}^d) : (-ih\nabla - \mathrm{Re}\,\mathbf{A})u \in L^2(\mathbb{R}^d, \mathbb{C}^d), \, (\mathrm{Im}\,\mathbf{A})u \in L^2(\mathbb{R}^d, \mathbb{C}^d) \right\} \,.$$

The objective of task (I) is fulfilled by the following theorem.

Theorem 1.2. Under Assumption 1.1, $Q_{h,A}$ is densely defined, closed and sectorial. Then, the operator defined by

$$Dom(\mathscr{L}_{h,\mathbf{A}}) \coloneqq \left\{ u \in Dom(Q_{h,\mathbf{A}}) : \exists f \in L^2(\mathbb{R}^d), \forall v \in Dom(Q_{h,\mathbf{A}}), Q_{h,\mathbf{A}}(u,v) = \langle f, v \rangle \right\},$$
$$\mathscr{L}_{h,\mathbf{A}} \coloneqq f,$$

is m-sectorial.

While the closed representative $\mathscr{L}_{h,\mathbf{A}}$ of (1.1) is easily introduced in all dimensions, it is well known that its spectral analysis becomes cumbersome in higher dimensions even in the self-adjoint case. Therefore, we modestly restrict to dimension d = 2 in the sequel. In this case, the magnetic field

$$\mathbf{B} \coloneqq \operatorname{curl} \mathbf{A} = \partial_{x_1} A_2 - \partial_{x_2} A_1$$

is a scalar function. Then pointwise sufficient conditions which guarantee Assumption 1.1 are contained in the following proposition.

Proposition 1.3. Let $\mathbf{A} \in L^2_{loc}(\mathbb{R}^2, \mathbb{C}^2)$ and $\mathbf{B} \in L^1_{loc}(\mathbb{R}^2)$. Assume one of the following conditions:

(C1): there exist $\varepsilon_1 \in (0,1)$, $C_1 \in \mathbb{R}$ such that

$$|\operatorname{Im} \boldsymbol{A}(x)|^2 \leqslant \pm \varepsilon_1 h \operatorname{Re} \boldsymbol{B}(x) + C_1, \qquad \forall x \in \mathbb{R}^2;$$

(C2): there exist $\varepsilon_2 \in (0, \frac{1}{2}), C_2 \in \mathbb{R}$ such that

$$|\operatorname{Im} \boldsymbol{A}(x)|^2 \leqslant \pm \varepsilon_2 h \operatorname{Im} \boldsymbol{B}(x) + C_2, \qquad \forall x \in \mathbb{R}^2.$$

Then Assumption 1.1 holds.

Of course, the conditions are automatically satisfied if $\text{Im } \mathbf{A} = 0$ and if $\text{Re } \mathbf{B}$ is bounded from below or from above. If $\text{Im } \mathbf{B} = 0$, note that one can choose a real vector potential \mathbf{A} , but that one can also choose a complex gauge (leading to non unitarily equivalent operators).

In practice, when **A** and **B** are continuous on \mathbb{R}^2 , it suffices to verify (**C1**) or (**C2**) for large values of |x|. Furthermore, if Im **A** is bounded and $\operatorname{Re} \mathbf{B}(x)$ (respectively, Im $\mathbf{B}(x)$) does not change sign for large |x|, then (**C1**) (respectively, (**C2**)) holds.

Example 1 (Homogeneous magnetic field). Unfortunately, Assumption 1.1 excludes the important case of constant magnetic field $\mathbf{B} = c \in \mathbb{C}$ corresponding to

$$\mathbf{A}(x) := c \frac{1}{2} (-x_2, x_1)$$
 or $\mathbf{A}(x) := c (0, x_1)$,

unless Im c = 0. In fact, implementing task (I) in this case seems particularly non-trivial.

Example 2 (Complexified Miller–Simon's potential). Consider

$$\mathbf{A}(x) := \left(-\frac{cx_2}{(1+|x|)^{\alpha}}, \frac{cx_1}{(1+|x|)^{\alpha}}\right),\,$$

where $c := c_1 + ic_2 \in \mathbb{C}$ with $c_1, c_2 \in \mathbb{R}$ and $\alpha > 0$. When c is real, the magnetic Laplacian with this type of magnetic potential was considered in [23]. When $c_2 \neq 0$ and if we assume further that $\alpha \ge 1$ then it can be checked that both condition (C1) and (C2) are satisfied. Indeed, since $|\mathbf{A}|$ is bounded and that

$$\mathbf{B}(x) = c \left(\frac{2}{(1+|x|)^{\alpha}} - \frac{\alpha |x|}{(1+|x|)^{\alpha+1}} \right)$$

is also bounded on \mathbb{R}^2 , we can therefore find sufficient large constants C_1 and C_2 such that (C1) and (C2) hold. With more effort, we can see that $\text{Dom}(Q_{h,\mathbf{A}}) = H^1(\mathbb{R}^2)$ and $\text{Dom}(\mathscr{L}_{h,\mathbf{A}}) = H^2(\mathbb{R}^2)$. In particular, when $c_1 = 0$, we have an example of purely imaginary magnetic fields such that our magnetic Laplacian is well defined.

Example 3. (Purely imaginary exponential magnetic potentials) Now we give an example of an unbounded purely imaginary \mathbf{A} for which (C1) fails but (C2) still holds. Consider

$$\mathbf{A}(x) \coloneqq ice^{|x|^2}(-x_2, x_1), \qquad (1.4)$$

where $c \in (-h, h)$ and h > 0. It can be checked that

$$|\text{Im } \mathbf{A}(x)|^2 = c^2 |x|^2 e^{|x|^2}, \qquad \text{Im } \mathbf{B}(x) = 2c(|x|^2 + 1)e^{|x|^2}.$$

Hence, we have

$$|\operatorname{Im} \mathbf{A}(x)|^2 \leq \frac{c}{2} \operatorname{Im} \mathbf{B}(x), \qquad \forall x \in \mathbb{R}^2.$$

By choosing $C_2 = 0$ and $\varepsilon_2 \in \left(\frac{|c|}{2h}, \frac{1}{2}\right)$, (C2) is verified.

1.2.2. *Compactness of the resolvent*. Let us now consider task (II) about the compactness of the resolvent (which implies that the spectrum is purely discrete).

Theorem 1.4. Let A satisfy Assumption 1.1 and $\operatorname{Re} A \in L^{\infty}_{\operatorname{loc}}(\mathbb{R}^2, \mathbb{R}^2)$. Assume one of the following conditions:

(H1): Re
$$\boldsymbol{B} \in C^{0}(\mathbb{R}^{2})$$
 and $\lim_{|x| \to +\infty} |\operatorname{Re} \boldsymbol{B}(x)| = +\infty$,
(H2): Im $\boldsymbol{B} \in C^{0}(\mathbb{R}^{2})$ and $\lim_{|x| \to +\infty} |\operatorname{Im} \boldsymbol{B}(x)| = +\infty$,
(H3): Im $\boldsymbol{A} \in C^{0}(\mathbb{R}^{2}, \mathbb{R}^{2})$ and $\lim_{|x| \to +\infty} |\operatorname{Im} \boldsymbol{A}(x)| = +\infty$.

Then $\mathscr{L}_{h,A}$ has compact resolvent.

The first condition of the theorem extends the usual magnetic bottle realisation [15] to the non-self-adjoint setting. More interestingly, it follows that the magnetic Laplacian of Example 3 (with $c \neq 0$) has compact resolvent despite Re $\mathbf{B} = 0$.

1.2.3. Semiclassical pseudomodes. Finally, let us present our implementation of task (III) about the construction of semiclassical pseudomodes. To this purpose, let us restrict to magnetic potentials $\mathbf{A} \in C^{\infty}(\mathbb{R}^2, \mathbb{C}^2)$ satisfying Assumption 1.1. In the plane \mathbb{R}^2 , we define the subset

$$\Gamma := \left\{ x \in \mathbb{R}^2 \, \middle| \begin{array}{c} \operatorname{Im} \mathbf{A}(x) = 0, \ \mathbf{B}(x) \neq 0, \ \partial_{\overline{z}} \mathbf{B}(x) \neq 0, \\ Q_1(x) > 0, \ Q_1(x) Q_3(x) - Q_2^2(x) > 0 \end{array} \right\} \,,$$

where

$$Q_{1}(x) := \frac{1}{4} \operatorname{Re} \left[\mathbf{B}(x) \left(1 + \frac{\partial_{z} \mathbf{B}}{\partial_{\overline{z}} \mathbf{B}}(x) \right) \right] + \frac{1}{2} \partial_{x_{1}} \operatorname{Im} A_{1}(x) ,$$

$$Q_{2}(x) := \frac{1}{4} \operatorname{Im} \left[\mathbf{B}(x) \frac{\partial_{z} \mathbf{B}}{\partial_{\overline{z}} \mathbf{B}}(x) \right] + \frac{1}{4} \left(\partial_{x_{1}} \operatorname{Im} A_{2} + \partial_{x_{2}} \operatorname{Im} A_{1}(x) \right) , \qquad (1.5)$$

$$Q_{3}(x) := \frac{1}{4} \operatorname{Re} \left[\mathbf{B}(x) \left(1 - \frac{\partial_{z} \mathbf{B}}{\partial_{\overline{z}} \mathbf{B}}(x) \right) \right] + \frac{1}{2} \partial_{x_{2}} \operatorname{Im} A_{2}(x) ,$$

and $\partial_z := \frac{1}{2} (\partial_{x_1} - i \partial_{x_2})$ and $\partial_{\overline{z}} := \frac{1}{2} (\partial_{x_1} + i \partial_{x_2})$ stand for the usual Wirtinger derivatives. Our main result is the following theorem.

Theorem 1.5. Let $x^0 \in \Gamma$ and assume that **B** is real-analytic at x^0 . Then there exist constants $C, h_0 > 0$ and a family of functions $(u_h)_{0 < h \leq h_0} \subset C_c^{\infty}(\mathbb{R}^2)$ such that, for all $h \in (0, h_0)$,

$$\left\| \left(\mathscr{L}_{h,\boldsymbol{A}} - h\boldsymbol{B}(x^0) \right) u_h \right\| \leq \exp\left(-\frac{C}{h^{1/7}}\right) \|u_h\|.$$
(1.6)

Note that $\Gamma = \emptyset$ if Im $\mathbf{A} = 0$, because $Q_1Q_3 - Q_2^2 = 0$ in this case. It follows that the existence of the pseudomode $(u_h)_{0 < h \leq h_0}$ is possible only in the present non-self-adjoint setting. Below we provide examples of magnetic potentials for which $\Gamma \neq \emptyset$; these include polynomial and oscillating complex magnetic fields.

In order to prove Theorem 1.5, we use the ideas of the magnetic WKB strategy as developed in [6, 12] in the self-adjoint case to find approximations of eigenfunctions. In our non-self-adjoint development, it is remarkable that the magnetic potential \mathbf{A} does not need to be analytic for this result to hold. This technical outcome is due to the use of a formal gauge, which allows for the local transformation to an appropriate analytic potential at x^0 , see Section 3.1.

We stress that Theorem 1.5 is not covered by [10]. Indeed, the Weyl symbol of $\mathscr{L}_{h,\mathbf{A}}$ is

$$p(x,\xi) = |\xi - \operatorname{Re} \mathbf{A}|^2 - |\operatorname{Im} \mathbf{A}|^2 - 2i\langle\xi - \operatorname{Re} \mathbf{A}, \operatorname{Im} \mathbf{A}\rangle.$$

Note that $p(x,\xi) = 0$ is equivalent to $\xi - \operatorname{Re} \mathbf{A} = \pm (\operatorname{Im} \mathbf{A})^{\perp}$ and that

$$\{\operatorname{Re} p, \operatorname{Im} p\}(x, \xi) = -4(\xi - \operatorname{Re} \mathbf{A}) \cdot \partial_x \langle \xi - \operatorname{Re} \mathbf{A}, \operatorname{Im} \mathbf{A} \rangle - 2\partial_x (|\xi - \operatorname{Re} \mathbf{A}|^2 - |\operatorname{Im} \mathbf{A}|^2) \cdot \operatorname{Im} \mathbf{A},$$

where $\{\cdot, \cdot\}$ is the Poisson bracket. Given $x^0 \in \Gamma$, then for any point $(x^0, \xi^0) \in \mathbb{R}^4$ such that $p(x^0, \xi^0) = 0$, we have $\{\operatorname{Re} p, \operatorname{Im} p\}(x^0, \xi^0) = 0$ (since $\operatorname{Im} \mathbf{A}(x^0) = 0$), so [10, Thm. 1.2] does not apply.

1.3. Structure of the paper. In Section 2 we deal with tasks (I) and (II); in particular, we introduce the operator $\mathscr{L}_{h,\mathbf{A}}$ and prove Proposition 1.3 and Theorem 1.4. Task (III) is considered in Section 3; namely, we establish Theorem 1.5 and provide the specific examples which the theorem applies to.

2. Sectorial magnetic Laplacians

The main purpose of this section is to realise the differential expression (1.1) as an msectorial operator in $L^2(\mathbb{R}^d)$ via a sesquilinear form and study its spectral property compact resolvent.

Assume $\mathbf{A} \in L^2_{\text{loc}}(\mathbb{R}^d, \mathbb{C}^d)$. Expanding $(-ih\nabla - \mathbf{A})^2$, the action of this operator should be

$$(-ih\nabla - \operatorname{Re} \mathbf{A})^2 - (\operatorname{Im} \mathbf{A})^2 - i(\operatorname{Im} \mathbf{A} \cdot (-ih\nabla - \operatorname{Re} \mathbf{A}) + (-ih\nabla - \operatorname{Re} \mathbf{A}) \cdot \operatorname{Im} \mathbf{A}) .$$

Aiming at the variational definition, this suggests considering the form domain

$$\mathcal{V}_{h,\mathbf{A}} := \left\{ u \in L^2(\mathbb{R}^d) : (-ih\nabla - \operatorname{Re}\mathbf{A})u \in L^2(\mathbb{R}^d, \mathbb{C}^d), \, (\operatorname{Im}\mathbf{A})u \in L^2(\mathbb{R}^d, \mathbb{C}^d) \right\}$$
(2.1)

equipped with the natural inner product

$$\langle u, v \rangle_{\mathcal{V}_{h,\mathbf{A}}} := \langle u, v \rangle + \langle (-ih\nabla - \operatorname{Re}\mathbf{A})u, (-ih\nabla - \operatorname{Re}\mathbf{A})v \rangle + \langle (\operatorname{Im}\mathbf{A})u, (\operatorname{Im}\mathbf{A})v \rangle, \quad (2.2)$$

where $\langle \cdot, \cdot \rangle$ denotes the inner product of $L^2(\mathbb{R}^d)$, linear in the first component.

Remark 2.1. Since $\mathbf{A} \in L^2_{\text{loc}}(\mathbb{R}^d, \mathbb{C}^d)$, it follows that for all $u \in L^2(\mathbb{R}^d)$,

 $(\operatorname{Re} \mathbf{A})u \in L^1_{\operatorname{loc}}(\mathbb{R}^d, \mathbb{C}^d), \quad (\operatorname{Im} \mathbf{A})u \in L^1_{\operatorname{loc}}(\mathbb{R}^d, \mathbb{C}^d).$

In particular, $(-ih\nabla - \operatorname{Re} \mathbf{A})u$ and $(\operatorname{Im} \mathbf{A})u$ can be understood in the sense of distributions in (2.1).

Just as in the self-adjoint case, we have the following properties of the form domain.

Proposition 2.2. Let h > 0 and $\mathbf{A} \in L^2_{loc}(\mathbb{R}^d, \mathbb{C}^d)$, the following holds.

- (a) $(\mathcal{V}_{h,A}, \langle \cdot, \cdot \rangle_{\mathcal{V}_{h,A}})$ is a Hilbert space.
- (b) $C_c^{\infty}(\mathbb{R}^d)$ is dense in $\mathcal{V}_{h,A}$.
- (c) Under the assumption 1.1, $\mathcal{V}_{h,A} = \mathcal{V}_{h,\text{Re}A}$, and two norms $\|\cdot\|_{V_{h,\text{Re}A}}$ and $\|\cdot\|_{V_{h,A}}$ are equivalent.

Proof. The proof of (a) is standard, and the proof of the density result in (b) follows the same steps as in [22, Theorem 7.22], so we omit the details here. Using (b), we extend the assumption (1.2) to hold on the space $\mathcal{V}_{h,\mathbf{A}}$. Consequently, statements in (c) are established.

After exploring several key properties of the space $\mathcal{V}_{h,\mathbf{A}}$, we are now ready to prove our first main theorem.

Proof of Theorem 1.2. We recall the sesquilinear form $Q_{h,\mathbf{A}} : \mathcal{V}_{h,\mathbf{A}} \to \mathbb{C}$ defined in (1.3). It is evident that $Q_{h,\mathbf{A}}$ is densely defined. Writing $\mathbf{A} = \operatorname{Re} \mathbf{A} + i \operatorname{Im} \mathbf{A}$, we expand $Q_{h,\mathbf{A}}$ as follows

$$Q_{h,\mathbf{A}}(u,v) = \int_{\mathbb{R}^d} (-ih\nabla - \operatorname{Re}\mathbf{A} - i\operatorname{Im}\mathbf{A})u \cdot \overline{(-ih\nabla - \operatorname{Re}\mathbf{A} + i\operatorname{Im}\mathbf{A})v} \, \mathrm{d}x$$
$$= \int_{\mathbb{R}^d} \left(|(-ih\nabla - \operatorname{Re}\mathbf{A})u|^2 - |\operatorname{Im}\mathbf{A}|^2 |u|^2 \right) \, \mathrm{d}x - 2i\operatorname{Re}\left\langle \operatorname{Im}\mathbf{A}u, (-ih\nabla - \operatorname{Re}\mathbf{A})u \right\rangle.$$

Under the assumption (1.2) (which extends to $\mathcal{V}_{h,\mathbf{A}}$ by the density of $C_c^{\infty}(\mathbb{R}^d)$ in $\mathcal{V}_{h,\mathbf{A}}$), we obtain that for all $u \in \mathcal{V}_{h,\mathbf{A}}$,

$$\operatorname{Re} Q_{h,\mathbf{A}}(u,u) \ge (1-a) \int_{\mathbb{R}^d} |(-ih\nabla - \operatorname{Re} \mathbf{A})u|^2 \mathrm{d}x - b||u||^2, \qquad (2.3)$$

where $a \in (0, 1)$ and $b \ge 0$ are constants from (1.2).

Next, we estimate the imaginary part of $Q_{h,\mathbf{A}}(u,u)$:

$$|\operatorname{Im} Q_{h,\mathbf{A}}(u,u)| \leq 2 \|\operatorname{Im} \mathbf{A} u\| \| (-ih\nabla - \operatorname{Re} \mathbf{A})u\|$$

$$\leq \|\operatorname{Im} \mathbf{A} u\|^{2} + \| (-ih\nabla - \mathbf{A})u\|^{2}$$

$$\leq (1+a)\| (-ih\nabla - \mathbf{A})u\|^{2} + b\|u\|^{2}.$$

Combining these inequalities, we deduce

$$|\operatorname{Im} Q_{h,\mathbf{A}}(u,u)| \leq \frac{1+a}{1-a} \operatorname{Re} Q_{h,\mathbf{A}}(u,u) + \frac{2b}{1-a} ||u||^2.$$

This shows that the form $Q_{h,\mathbf{A}}$ is sectorial.

We now verify the closedness of the form. Consider the form

$$t(u,v) \coloneqq \langle (-i\nabla - \operatorname{Re} \mathbf{A})u, (-i\nabla - \operatorname{Re} \mathbf{A})v \rangle,$$

which is sectorial and closed on $\mathcal{V}_{h,\text{Re}\mathbf{A}}$. By applying [18, Thm. VI.1.33] and noting that the form $\langle \text{Im} \mathbf{A} \cdot, \text{Im} \mathbf{A} \cdot \rangle$ is *t*-bounded, we conclude that the form $\text{Re} Q_{h,\mathbf{A}}$ is closed. By a remark in [18, Sec. VI.1.3], the sectorial form $Q_{h,\mathbf{A}}$ is also closed.

Consequently, $Q_{h,\mathbf{A}}$ gives rise to an m-sectorial operator $\mathscr{L}_{h,\mathbf{A}}$ via the standard representation theorem [18, Thm. VI.2.1].

From now on, we restrict to dimension d = 2.

Lemma 2.3. Let
$$\mathbf{A} \in L^2_{\text{loc}}(\mathbb{R}^2, \mathbb{C}^2)$$
 and $\mathbf{B} \in L^1_{\text{loc}}(\mathbb{R}^2)$. Then, for all $u \in C^{\infty}_c(\mathbb{R}^2)$, we have

$$\left| \int_{\mathbb{R}^2} h \operatorname{Re} \boldsymbol{B} |u|^2 \, \mathrm{d}x \right| \leq \int_{\mathbb{R}^2} \left| (-ih\nabla - \operatorname{Re} \boldsymbol{A}) u \right|^2 \, \mathrm{d}x \,,$$
$$\left| \int_{\mathbb{R}^2} h \operatorname{Im} \boldsymbol{B} |u|^2 \, \mathrm{d}x \right| \leq \int_{\mathbb{R}^2} \left| (-ih\nabla - \operatorname{Re} \boldsymbol{A}) u \right|^2 \, \mathrm{d}x + \int_{\mathbb{R}^2} \left| (\operatorname{Im} \boldsymbol{A}) u \right|^2 \, \mathrm{d}x \,.$$

Proof. Since $\mathbf{A} \in L^2_{loc}(\mathbb{R}^2, \mathbb{C}^2)$ and $\mathbf{B} \in L^1_{loc}(\mathbb{R}^2)$, all the integrals appearing in the statements of this lemma are well defined. Let $u \in C^{\infty}_c(\mathbb{R}^2)$. By observing that

$$\left[(-ih\partial_{x_1} - A_1), \, (-ih\partial_{x_2} - A_2)\right] = ih\mathbf{B}\,,$$

we have

$$ih\mathbf{B}|u|^2 = \overline{u}(-ih\partial_{x_1} - A_1)(-ih\partial_{x_2} - A_2)u - \overline{u}(-ih\partial_{x_2} - A_2)(-ih\partial_{x_1} - A_1)u.$$

Integrating by parts, we get

$$\int_{\mathbb{R}^2} ih\mathbf{B}|u|^2 \,\mathrm{d}x = \langle (-ih\partial_{x_2} - A_2)u, (-ih\partial_{x_1} - \overline{A_1})u \rangle - \langle (-ih\partial_{x_1} - A_1)u, (-ih\partial_{x_2} - \overline{A_2})u \rangle.$$

By writing out the real and imaginary parts of A_1 and A_2 , we find that

$$\int_{\mathbb{R}^2} ih \mathbf{B} |u|^2 \, \mathrm{d}x = 2\mathrm{Im} \left\langle (\mathrm{Im} \, A_2)u, (-ih\partial_{x_1} - \mathrm{Re} \, A_1)u \right\rangle - 2\mathrm{Im} \left\langle (\mathrm{Im} \, A_1)u, (-ih\partial_{x_2} - \mathrm{Re} \, A_2)u \right\rangle + i2\mathrm{Im} \left\langle (-ih\partial_{x_2} - \mathrm{Re} \, A_2)u, (-ih\partial_{x_1} - \mathrm{Re} \, A_1)u \right\rangle.$$

In other words, we have

$$\int_{\mathbb{R}^2} h \operatorname{Re} \mathbf{B} |u|^2 \, \mathrm{d}x = 2 \operatorname{Im} \left\langle (-ih\partial_{x_2} - \operatorname{Re} A_2)u, (-ih\partial_{x_1} - \operatorname{Re} A_1)u \right\rangle,$$

and

$$\int_{\mathbb{R}^2} h \operatorname{Im} \mathbf{B} |u|^2 \, \mathrm{d}x = 2 \operatorname{Im} \left\langle (\operatorname{Im} A_1) u, (-ih\partial_{x_2} - \operatorname{Re} A_2) u \right\rangle - 2 \operatorname{Im} \left\langle (\operatorname{Im} A_2) u, (-ih\partial_{x_1} - \operatorname{Re} A_1) u \right\rangle.$$

The conclusion follows from the Cauchy–Schwarz inequality.

Now we are in a position to establish Proposition 1.3.

Proof of Proposition 1.3. When Im A and Re B satisfy (C1), then Assumption 1.1 is a direct consequence of the first inequality in Lemma 2.3.

Now, we assume that $\operatorname{Im} \mathbf{A}$ and $\operatorname{Im} \mathbf{B}$ satisfy (C2) with the plus sign (for instance). From the second inequality in Lemma 2.3,

$$\int_{\mathbb{R}^2} |(\operatorname{Im} \mathbf{A})u|^2 \, \mathrm{d}x \leqslant \varepsilon_2 h \int_{\mathbb{R}^2} \operatorname{Im} \mathbf{B}|u|^2 \, \mathrm{d}x + C_2 ||u||^2$$
$$\leqslant \varepsilon_2 \int_{\mathbb{R}^2} |(-ih\nabla - \operatorname{Re} \mathbf{A})u|^2 \, \mathrm{d}x + \varepsilon_2 \int_{\mathbb{R}^2} |(\operatorname{Im} \mathbf{A})u|^2 \, \mathrm{d}x + C_2 ||u||^2.$$

This implies that Assumption 1.1 holds with $a = \frac{\varepsilon_2}{1-\varepsilon_2} \in (0,1)$ and $b = \frac{C_2}{1-\varepsilon_2}$.

Now we turn to task (II) about the compactness of the resolvent of $\mathscr{L}_{h,\mathbf{A}}$.

Proof of Theorem 1.4. Thanks to [7, Prop. 4.24], it is equivalent to proving that the injection

$$\left(\operatorname{Dom}(\mathscr{L}_{h,\mathbf{A}}), \|\cdot\|_{\mathscr{L}_{h,\mathbf{A}}}\right) \hookrightarrow \left(L^2(\mathbb{R}^2), \|\cdot\|\right)$$

is compact, where $\|\cdot\|_{\mathscr{L}_{h,\mathbf{A}}} \coloneqq \|\mathscr{L}_{h,\mathbf{A}}\cdot\| + \|\cdot\|$ is the graph norm. From (2.3) and Proposition 2.2(c), there exist $\gamma > 0$ and $\mu > 0$ such that for all $u \in \text{Dom}(\mathscr{L}_{\mathbf{A}})$, we have

$$\gamma \|u\|_{\mathcal{V}_{h,\mathbf{A}}}^{2} \leq |Q_{h,\mathbf{A}}(u,u) + \mu\|u\|^{2}| \leq \|(\mathscr{L}_{h,\mathbf{A}} + \mu)u\|\|u\| \leq \left(\frac{1}{2} + \mu\right) \|u\|_{\mathscr{L}_{h,\mathbf{A}}}^{2}$$

which shows that the injection

$$\left(\mathrm{Dom}(\mathscr{L}_{h,\mathbf{A}}), \|\cdot\|_{\mathscr{L}_{h,\mathbf{A}}}\right) \hookrightarrow \left(\mathcal{V}_{h,\mathbf{A}}, \|\cdot\|_{\mathcal{V}_{h,\mathbf{A}}}\right)$$

is continuous. Since the space of compact operators forms an ideal within the space of bounded operator, it remains to explain why $(\mathcal{V}_{h,\mathbf{A}}, \|\cdot\|_{\mathcal{V}_{h,\mathbf{A}}})$ is compactly embedded in $L^2(\mathbb{R}^2).$

Let us consider $D \coloneqq \{u \in \mathcal{V}_{h,\mathbf{A}} : \|u\|_{\mathcal{V}_{h,\mathbf{A}}} \leq 1\}$ and prove its precompactness in $L^2(\mathbb{R}^2)$ by means of the Kolmogorov–Riesz theorem (see [7, Thm. 4.14]). We only need to check the following:

- (i) For all ε > 0, there exists ω ⊂⊂ ℝ² such that ∫_{ℝ²\ω} |u|² dx ≤ ε², for all u ∈ D.
 (ii) For all ε > 0 and for all ω ⊂⊂ ℝ², there exists δ > 0 such that

$$\int_{\omega} |u(x+s) - u(x)|^2 \, \mathrm{d}x \leqslant \varepsilon^2$$

for all $s \in \mathbb{R}^2$ with $|s| \leq \delta$ and for all $u \in D$.

The equi-integrability condition (i) is satisfied as long as at least one of the assumptions (H1), (H2), or (H3) holds. This follows from Assumption 1.1, Lemma 2.3, Proposition $2.2(\mathbf{b})$, and the continuity of the functions considered in assumptions (H1), (H2), and (H3). Below, we provide a detailed proof of (i) under the assumption (H1). Since the proof is identical under the other assumptions, we will omit those cases.

Assume that $\operatorname{Re} \mathbf{B}$ satisfies (H1). Since $\operatorname{Re} \mathbf{B}$ is continuous, by the multivariate intermediate value theorem [11, Thm. 1.9.5] and the connectedness of the punctured disk in \mathbb{R}^2 , the condition $\lim_{|x|\to+\infty} |\operatorname{Re} \mathbf{B}(x)| = +\infty$ implies that

$$\lim_{x \to +\infty} \operatorname{Re} \mathbf{B}(x) = +\infty \quad \text{or} \quad \lim_{|x| \to +\infty} -\operatorname{Re} \mathbf{B}(x) = +\infty.$$

Without loss of generality, we assume the first possibility. By adding a constant C > 0such that $\operatorname{Re} \mathbf{B} + C \ge 0$ on \mathbb{R}^2 (to apply Fatou's lemma in the following step), and using the first inequality in Lemma 2.3, we obtain

$$(C+1)\|u\|_{\mathcal{V}_{h,\mathbf{A}}}^2 \geqslant \int_{\mathbb{R}^2} (\operatorname{Re}\mathbf{B} + C)|u|^2 \,\mathrm{d}x\,, \qquad \forall u \in C_c^\infty(\mathbb{R}^2)\,.$$

To extend this inequality to the space $\mathcal{V}_{h,\mathbf{A}}$, we utilise the density of $C_c^{\infty}(\mathbb{R}^2)$ in $\mathcal{V}_{h,\mathbf{A}}$. More precisely, let $u \in \mathcal{V}_{h,\mathbf{A}}$, there exists a sequence $(u_n)_{n\in\mathbb{N}} \subset C_c^{\infty}(\mathbb{R}^2)$ such that $u_n \xrightarrow{n\to+\infty} u$ in $\mathcal{V}_{h,\mathbf{A}}$. By considering a subsequence, still denoted by u_n , such that $u_n(x) \xrightarrow{n\to+\infty} u(x)$ for almost every $x \in \mathbb{R}^2$, Fatou's lemma yields that

$$\begin{aligned} (C+1)\|u\|_{\mathcal{V}_{h,\mathbf{A}}}^2 &= \lim_{n \to +\infty} (C+1)\|u_n\|_{\mathcal{V}_{h,\mathbf{A}}} \geqslant \liminf_{n \to +\infty} \int_{\mathbb{R}^2} (\operatorname{Re} \mathbf{B} + C)|u_n|^2 \, \mathrm{d}x \\ &\geqslant \int_{\mathbb{R}^2} (\operatorname{Re} \mathbf{B} + C)|u|^2 \, \mathrm{d}x \,, \qquad \forall u \in \mathcal{V}_{\mathbf{A}} \,. \end{aligned}$$

Given $\varepsilon > 0$, using this estimate and the unboundedness of Re B at infinity, there exists a constant R > 0 such that

$$\int_{|x|>R} |u|^2 \,\mathrm{d}x < \varepsilon \,, \qquad \forall u \in D$$

Let us now consider (ii). Let $\varepsilon > 0$ and $\omega \subset \mathbb{R}^2$, we consider a function $\chi \in C_c^{\infty}(\mathbb{R}^2)$ and $\chi = 1$ in some neighbourhood of $\overline{\omega}$. For all $u \in D$,

$$-ih\nabla(\chi u) = \operatorname{Re} \mathbf{A}\chi u + (-ih\nabla\chi)u + \chi(-ih\nabla - \operatorname{Re} \mathbf{A})u \in L^2(\mathbb{R}^2),$$

where we used the fact that $\operatorname{Re} \mathbf{A} \in L^{\infty}_{\operatorname{loc}}(\mathbb{R}^2, \mathbb{R}^2)$. There exists C > 0 such that, for all $u \in D$,

$$\|\chi u\|_{H^1(\mathbb{R}^2)} \leqslant C \,.$$

Then, we notice that, for |s| small enough such that $\chi(x+s) = 1$ on ω , we have, for all $u \in D$,

$$\int_{\omega} |u(x+s) - u(x)|^2 dx = \int_{\omega} |(\chi u)(x+s) - (\chi u)(x)|^2 dx$$
$$\leq \int_{\mathbb{R}^2} |(\chi u)(x+s) - (\chi u)(x)|^2 dx \leq C^2 |s|^2,$$

where we used [7, Prop. 2.94].

3. WKB CONSTRUCTION OF PSEUDOMODES

This section is concerned with task (III) about the construction of semiclassical pseudomodes. In particular, we establish Theorem 1.5. Throughout this section, we assume that $\mathbf{B} \in C^{\infty}(\mathbb{R}^2)$.

Definition 3.1. We say that **B** is real-analytic at x^0 , when, in the neighbourhood of x^0 , **B** is the sum of a converging series:

$$\mathbf{B}(x_1, x_2) = \sum_{m,n \ge 0} b_{mn} (x_1 - x_1^0)^m (x_2 - x_2^0)^n, \qquad b_{mn} \in \mathbb{C}.$$

The main technical result of this section is the following theorem.

Theorem 3.2. Let $x^0 \in \Gamma$ and assume that **B** is real-analytic at x^0 . Then, there exist

- i) a neighbourhood \mathcal{U} of x^0 in \mathbb{R}^2 ;
- ii) a real-analytic function P on U satisfying

$$\operatorname{Re} P(x) = Q(x_1 - x_1^0, x_2 - x_2^0) + \mathcal{O}\left(|x - x^0|^3\right), \qquad (3.1)$$

where $Q(u,v) := Q_1(x^0)u^2 - 2Q_2(x^0)uv + Q_3(x^0)v^2$ is a positive definite quadratic form on \mathbb{R}^2 with Q_1, Q_2, Q_3 defined as in (1.5);

iii) a sequence of real-analytic functions $(a_j)_{j\in\mathbb{N}}$ on \mathfrak{U} with $a_0(x^0) = 1$ and $a_j(x^0) = 0$ for $j \ge 1$;

such that, for all $N \in \mathbb{N}$,

$$e^{P/h}\left(\mathscr{L}_{h,\boldsymbol{A}} - h\boldsymbol{B}(x^0)\right)\left(e^{-P/h}\sum_{j=0}^N h^j a_j\right) = \mathcal{O}\left(h^{N+2}\right)$$
(3.2)

locally uniformly on \mathcal{U} .

After a translation, we can assume that $x^0 = 0$.

Notation 3.3 (Complexification of a real-analytic function). Assume that a is real-analytic near the point $0 \in \mathbb{R}^2$. We denote by \tilde{a} the function defined near $0 \in \mathbb{C}^2$ by

$$\widetilde{a}(z,w) \coloneqq a\left(\frac{z+w}{2}, \frac{z-w}{2i}\right)$$

Note that

$$\widetilde{a}(z,\overline{z}) = a(\operatorname{Re} z,\operatorname{Im} z), \qquad \partial_z \widetilde{a} = \widetilde{\partial_z a}, \qquad \partial_w \widetilde{a} = \widetilde{\partial_{\overline{z}} a}.$$
(3.3)

3.1. A choice of the magnetic potential.

Lemma 3.4. There exists a real-analytic and complex-valued function φ in a neighbourhood Ω of 0 such that

$$\Delta \varphi = \boldsymbol{B}, \qquad \varphi(x_1, x_2) = \frac{\boldsymbol{B}(0)}{4} \left(x_1^2 + x_2^2 \right) + \mathcal{O}\left(|x|^3 \right)$$

Proof. By considering the complexification of \mathbf{B} in the neighbourhood of 0

$$\widetilde{\mathbf{B}}(z,w) = \sum_{(\alpha,\beta) \in \mathbb{N}^2} a_{\alpha,\beta} z^{\alpha} w^{\beta} \,,$$

with $a_{0,0} = \mathbf{B}(0)$, we introduce the power series

$$\widetilde{\varphi}(z,w) = \frac{1}{4} \sum_{(\alpha,\beta) \in \mathbb{N}^2} \frac{a_{\alpha,\beta}}{(\alpha+1)(\beta+1)} z^{\alpha+1} w^{\beta+1}$$

Then we get

$$4\partial_z \partial_w \widetilde{\varphi}(z, w) = \widetilde{\mathbf{B}}(z, w)$$

The function $z \mapsto \widetilde{\varphi}(z, \overline{z})$ satisfies the required properties.

Let φ be the function given by Lemma 3.4, and define

$$\mathcal{M} := (-\partial_{x_2}\varphi, \partial_{x_1}\varphi) \; .$$

Then \mathcal{M} satisfies

$$\frac{\partial \mathcal{M}_2}{\partial x_1} - \frac{\partial \mathcal{M}_1}{\partial x_2} = \mathbf{B} \qquad \text{on } \Omega.$$

By the Poincaré lemma, there exists a function $\theta \in C^{\infty}(\Omega, \mathbb{C})$ such that

$$\mathcal{M} = \mathbf{A} + \nabla \theta \qquad \text{on } \Omega \,. \tag{3.4}$$

From this, we obtain

$$\mathscr{L}_{h,\mathcal{M}} = e^{i\theta/h} \mathscr{L}_{h,\mathbf{A}} e^{-i\theta/h} \qquad \text{on } \Omega \,. \tag{3.5}$$

3.2. WKB analysis. In this section, we construct a pseudomode for the operator $\mathscr{L}_{h,\mathcal{M}}$. More precisely, for $N \in \mathbb{N}$, we look for a pseudomode in the form

$$u_h(x) = e^{-S(x)/h} \sum_{j=0}^N a_j(x)h^j$$
,

attached to a quasi-eigenvalue $\lambda(h) = h\mu$. Here, S and a_j are real-analytic functions defined in the neighbourhood of $0 \in \mathbb{R}^2$, and $\mu \in \mathbb{C}$.

Let us consider the formal conjugated operator acting locally:

$$\begin{aligned} \mathscr{L}_{h,\mathcal{M}}^{S} &:= e^{S/h} \mathscr{L}_{h,\mathcal{M}} e^{-S/h} \\ &= (-ih\partial_{1} - \mathcal{M}_{1} + i\partial_{x_{1}}S)^{2} + (-ih\partial_{2} - \mathcal{M}_{2} + i\partial_{x_{2}}S)^{2} \\ &= E_{0} + hE_{1} + h^{2}E_{2} \,, \end{aligned}$$

where the differential expressions E_0 , E_1 and E_2 are given by

$$E_0 := (-\mathcal{M}_1 + i\partial_{x_1}S)^2 + (-\mathcal{M}_2 + i\partial_{x_2}S)^2$$

$$E_1 := \Delta S + 2(\nabla S + i\mathcal{M}) \cdot \nabla,$$

$$E_2 := -\Delta.$$

Then, we have

$$e^{S/h} \left(\mathscr{L}_{h,\mathcal{M}} - \lambda(h)\right) u_h(x) = \left[E_0 + h(E_1 - \mu) + h^2 E_2\right] \sum_{j=0}^N a_j(x) h^j = \sum_{j=0}^{N+2} \phi_j(x) h^j,$$

where the functions ϕ_j are explicitly given by

and the last function ϕ_{N+2} is

$$h^{N+2}: E_2 a_N =: \phi_{N+2}.$$

3.2.1. The eikonal equation. Let us find S such that $E_0 = 0$, *i.e.*,

$$(-\mathcal{M}_1 + i\partial_{x_1}S)^2 + (-\mathcal{M}_2 + i\partial_{x_2}S)^2 = 0.$$

It is equivalent to the equation

$$(i\partial_{x_1}S - \partial_{x_2}S - \mathcal{M}_1 - i\mathcal{M}_2)(i\partial_{x_1}S + \partial_{x_2}S - \mathcal{M}_1 + i\mathcal{M}_2) = 0$$

Let us choose S such that

$$i\partial_{x_1}S - \partial_{x_2}S - \mathcal{M}_1 - i\mathcal{M}_2 = 0$$

Then we have

$$2\partial_{\overline{z}}S = \mathcal{M}_2 - i\mathcal{M}_1 = 2\partial_{\overline{z}}\varphi$$

In particular, $S - \varphi$ is holomorphic. By using Notation 3.3, this suggests taking

$$\widetilde{S}(z,w) = \widetilde{\varphi}(z,w) + f(z) \,,$$

where f(z) is a holomorphic function (in a neighbourhood of $0 \in \mathbb{R}^2$) to be determined later in the neighbourhood of $0 \in \mathbb{C}$. Note that $\Delta S = \mathbf{B}$. 3.2.2. Towards the transport equations. Now, we consider the operator E_1 . By using the expression of \mathcal{M} and the choice of \widetilde{S} , we have

$$(\nabla S + i\mathcal{M}) \cdot \nabla = (\partial_{x_1}S + i\mathcal{M}_1) \,\partial_{x_1} + (\partial_{x_2}S + i\mathcal{M}_2) \,\partial_{x_2}$$

= $(\partial_{x_1}\varphi - i\partial_{x_2}\varphi + f'(z)) \,\partial_{x_1} + (\partial_{x_2}\varphi + i\partial_{x_1}\varphi + if'(z)) \,\partial_{x_2}$
= $2 \left(2\partial_z\varphi + f'(z)\right) \,\partial_{\overline{z}}$.

Thus, for any real-analytic function a near the point x^0 , we have

$$\widetilde{E_1 a} = \left[4 \left(2\partial_z \widetilde{\varphi} + f'(z) \right) \partial_w + \widetilde{\mathbf{B}} \right] \widetilde{a} \,.$$

Note also that

$$\widetilde{E_2 a} = -4\partial_z \partial_w \widetilde{a}$$

From (3.6), we are led to the system of the transport equations

$$h^{1}: \qquad \left[4\left(2\partial_{z}\widetilde{\varphi}+f'(z)\right)\partial_{w}+\widetilde{\mathbf{B}}-\mu\right]\widetilde{a}_{0}=0,$$

$$h^{2}: \qquad \left[4\left(2\partial_{z}\widetilde{\varphi}+f'(z)\right)\partial_{w}+\widetilde{\mathbf{B}}-\mu\right]\widetilde{a}_{1}=4\partial_{z}\partial_{w}\widetilde{a}_{0},$$

$$\vdots$$

$$h^{N+1}: \qquad \left[4\left(2\partial_{z}\widetilde{\varphi}+f'(z)\right)\partial_{w}+\widetilde{\mathbf{B}}-\mu\right]\widetilde{a}_{N}=4\partial_{z}\partial_{w}\widetilde{a}_{N-1}$$

3.2.3. Choosing f and determining S. Since $0 \in \Gamma$, we have $\partial_{\overline{z}} \mathbf{B}(0) \neq 0$ and thus

$$\partial_w \widetilde{\mathbf{B}}(0) = \widetilde{\partial_{\overline{z}} \mathbf{B}}(0) = \partial_{\overline{z}} \mathbf{B}(0) \neq 0.$$

In virtue of the holomorphic implicit function theorem, there exists a unique holomorphic function w, in the neighborhood of $0 \in \mathbb{C}$, such that

$$w(0) = 0, \qquad \mathbf{B}(z, w(z)) = \mathbf{B}(0).$$
 (3.7)

In particular,

$$w'(0) = -\frac{\partial_z \mathbf{B}}{\partial_w \mathbf{B}}(0) = -\frac{\partial_z \mathbf{B}}{\partial_{\overline{z}} \mathbf{B}}(0).$$
(3.8)

Remark 3.5. Note that the function w solves the same effective equation as in [6], but that we are *not* in the case of a magnetic well.

In order to solve the above transport equations, we will choose a function f(z) such that

$$2\partial_z \widetilde{\varphi}(z, w(z)) + f'(z) = 0$$

Lemma 3.6. Let φ be given in Lemma 3.4, w be the holomorphic function given in (3.7) and θ given in (3.4). Letting

$$f(z) := -\int_{[0,z]} 2\partial_z \widetilde{\varphi}(\zeta, w(\zeta)) \,\mathrm{d}\zeta + \operatorname{Im} \theta(0) \,, \tag{3.9}$$

we have

$$f(0) = \operatorname{Im} \theta(0), \qquad f'(0) = 0, \qquad f''(0) = \frac{B(0)}{2} \frac{\partial_z B}{\partial_{\overline{z}} B}(0).$$

In particular,

$$\operatorname{Re} S(x) = \operatorname{Im} \theta(0) + \tilde{Q}_1 x_1^2 - 2\tilde{Q}_2 x_1 x_2 + \tilde{Q}_3 x_2^2 + \mathcal{O}(|x|^3) ,$$

where

$$\tilde{Q}_{1} = \frac{1}{4} \operatorname{Re} \left[\boldsymbol{B}(0) \left(1 + \frac{\partial_{z} \boldsymbol{B}}{\partial_{\overline{z}} \boldsymbol{B}}(0) \right) \right], \qquad \tilde{Q}_{2} = \frac{1}{4} \operatorname{Im} \left[\boldsymbol{B}(0) \frac{\partial_{z} \boldsymbol{B}}{\partial_{\overline{z}} \boldsymbol{B}}(0) \right],$$
$$\tilde{Q}_{3} = \frac{1}{4} \operatorname{Re} \left[\boldsymbol{B}(0) \left(1 - \frac{\partial_{z} \boldsymbol{B}}{\partial_{\overline{z}} \boldsymbol{B}}(0) \right) \right].$$

Proof. A straightforward computation gives

$$f''(0) = -2\partial_{zz}^2 \widetilde{\varphi}(0) - 2w'(0)\partial_{zw}^2 \widetilde{\varphi}(0) \,.$$

Due to Lemma 3.4, we have

$$\partial_{zz}^2 \widetilde{\varphi}(0) = 0, \quad \partial_{zw}^2 \widetilde{\varphi}(0) = \frac{\mathbf{B}(0)}{4},$$

and we get the value of f''(0) by using (3.8).

Since $S(x_1, x_2) = \varphi(x_1, x_2) + f(x_1 + ix_2)$, we can write

$$S(x_1, x_2) = \operatorname{Im} \theta(0) + \frac{\mathbf{B}(0)}{4} \left[x_1^2 + x_2^2 + \frac{\partial_z \mathbf{B}(0)}{\partial_{\overline{z}} \mathbf{B}(0)} (x_1 + ix_2)^2 \right] + \mathcal{O}(|x|^3),$$

so that

$$\operatorname{Re} S(x) = \operatorname{Im} \theta(0) + \frac{1}{4} \operatorname{Re} \left[\mathbf{B}(0) \left(1 + \frac{\partial_z \mathbf{B}(0)}{\partial_{\overline{z}} \mathbf{B}(0)} \right) \right] x_1^2 + \frac{1}{4} \operatorname{Re} \left[\mathbf{B}(0) \left(1 - \frac{\partial_z \mathbf{B}(0)}{\partial_{\overline{z}} \mathbf{B}(0)} \right) \right] x_2^2 - \frac{1}{2} \operatorname{Im} \left[\mathbf{B}(0) \frac{\partial_z \mathbf{B}(0)}{\partial_{\overline{z}} \mathbf{B}(0)} \right] x_1 x_2 + \mathcal{O}(|x|^3) = \operatorname{Im} \theta(0) + \tilde{Q}_1 x_1^2 - 2 \tilde{Q}_2 x_1 x_2 + \tilde{Q}_3 x_2^2 + \mathcal{O}(|x|^3).$$

3.2.4. Solving the first transport equation. With the choice (3.9), we can write

$$8\left[\partial_z \widetilde{\varphi}(z,w) - \partial_z \widetilde{\varphi}(z,w(z))\right] \partial_w \widetilde{a}_0(z,w) + \left(\widetilde{\mathbf{B}}(z,w) - \mu\right) \widetilde{a}_0(z,w) = 0.$$
(3.10)

Taking w = w(z) and using (3.7), we see that (3.10) has a holomorphic solution \tilde{a}_0 such that $\tilde{a}_0(0) \neq 0$ if and only if

$$\mu = \mathbf{B}(0) \,.$$

Let us explain this. The Taylor formula gives

$$\partial_z \widetilde{\varphi}(z, w) - \partial_z \widetilde{\varphi}(z, w(z)) = (w - w(z))V(z, w) ,$$

$$\widetilde{\mathbf{B}}(z, w) - \widetilde{\mathbf{B}}(z, w(z)) = (w - w(z))F(z, w) ,$$

where

$$V(z,w) = \int_0^1 \widetilde{\mathbf{B}}(z,w(z) + t(w - w(z)) \, \mathrm{d}t, \quad F(z,w) = \int_0^1 \partial_w \widetilde{\mathbf{B}}(z,w(z) + t(w - w(z)) \, \mathrm{d}t.$$

Notice that $V(0) = \mathbf{B}(0) \neq 0$, so that near $0 \in \mathbb{C}^2$, we have $V(z, w) \neq 0$. By using $\mu = \mathbf{B}(0)$, the first transport equation becomes

$$\partial_w \widetilde{a}_0(z,w) + \frac{1}{8} \frac{F(z,w)}{V(z,w)} \widetilde{a}_0(z,w) = 0$$

Therefore, we have

$$\widetilde{a}_0(z,w) = \mathcal{A}_0(z)J(z,w), \quad J(z,w) \coloneqq \exp\left(-\int_{[w(z),w]} \frac{1}{8} \frac{F(z,u)}{V(z,u)} \,\mathrm{d}u\right)$$

where $\mathcal{A}_0(z)$ is a holomorphic function to be determined such that $\mathcal{A}_0(0) = 1$.

13

3.2.5. Solving the second transport equation. Let us now consider the second transport equation

$$(w - w(z)) \left[8V(z, w)\partial_w + F(z, w)\right] \tilde{a}_1(z, w) = 4\partial_z \partial_w \tilde{a}_0(z, w).$$
(3.11)

Letting w = w(z), we necessarily get that

$$4\partial_z \partial_w \widetilde{a}_0(z, w(z)) = 0 \,,$$

which means that

$$\partial_w J(z, w(z)) \mathcal{A}'_0(z) + \partial_z \partial_w J(z, w(z)) \mathcal{A}_0(z) = 0$$

This allows us to determine \mathcal{A}_0 of the previous step. From the definition of J, we have

$$\partial_w J(z, w(z)) = -\frac{1}{8} \frac{F(z, w(z))}{V(z, w(z))},$$

and thus

$$\partial_w J(0, w(0)) = -\frac{1}{8} \frac{\partial_w \widetilde{\mathbf{B}}}{\widetilde{\mathbf{B}}}(0) = -\frac{1}{8} \frac{\partial_{\overline{z}} \mathbf{B}}{\mathbf{B}}(0) \neq 0$$

This leads to the choice

$$\mathcal{A}_0(z) = \exp\left(-\int_{[0,z]} \frac{\partial_z \partial_w J}{\partial_w J}(u, w(u)) \,\mathrm{d}u\right) \,,$$

which is not allowed in the case of a magnetic well in the self-adjoint case. In particular,

$$\widetilde{a}_0(z,w) = \exp\left(-\int_{[0,z]} \frac{\partial_z \partial_w J}{\partial_w J}(u,w(u)) \,\mathrm{d}u\right) \exp\left(-\int_{[w(z),w]} \frac{1}{8} \frac{F(z,u)}{V(z,u)} \,\mathrm{d}u\right)$$

With this choice, (3.11) becomes

$$\partial_w \tilde{a}_1(z, w) + \frac{1}{8} \frac{F(z, w)}{V(z, w)} \tilde{a}_1(z, w) = \frac{1}{2} \frac{T_0(z, w)}{V(z, w)},$$

with

$$T_0(z,w) = \int_0^1 \partial_w^2 \partial_z \widetilde{a}_0(z,w(z) + t(w - w(z))) \,\mathrm{d}t \,.$$

We have

$$\widetilde{a}_1(z,w) = J(z,w) \int_{[w(z),w]} \frac{T_0(z,u)}{2J(z,u)V(z,u)} \,\mathrm{d}u + \mathcal{A}_1(z)J(z,w),$$

where $\mathcal{A}_1(z)$ is a holomorphic function to be determined with $\mathcal{A}_1(0) = 0$.

3.2.6. Induction. By induction, the solution of the j+1-th transport equation can be written as

$$\widetilde{a}_{j+1}(z,w) = J(z,w) \int_{[w(z),w]} \frac{T_j(z,u)}{2J(z,u)V(z,u)} \,\mathrm{d}u + \mathcal{A}_{j+1}(z)J(z,w) \,,$$

with

$$T_j(z,w) = \int_0^1 \partial_w^2 \partial_z \widetilde{a}_j(z,w(z) + t(w - w(z))) \,\mathrm{d}t$$

and \mathcal{A}_{j+1} is determined by the constraint (coming from the j + 2-th equation):

$$4\partial_z \partial_w \widetilde{a}_{j+1}(z, w(z)) = 0.$$
(3.12)

We have

$$\partial_w \tilde{a}_{j+1}(z, w) = \partial_w J(z, w) \int_{[w(z), w]} \frac{T_j(z, u)}{2J(z, u)V(z, u)} \, \mathrm{d}u + \frac{T_j(z, w)}{2V(z, w)} + \mathcal{A}_{j+1}(z) \partial_w J(z, w) \,,$$

so that

$$\partial_z \partial_w \widetilde{a}_{j+1}(z, w(z)) = \left[-\partial_w J w'(z) \frac{T_j}{2V} + \partial_z J \frac{T_j}{2V} + \partial_z \left(\frac{T_j}{2V} \right) + \mathcal{A}'_{j+1} \partial_w J + \mathcal{A}_{j+1} \partial_{zw}^2 J \right] (z, w(z)) \,.$$

Since J(z, w(z)) = 1, we have $\partial_z J(z, w(z)) = -w'(z)\partial_w J(z, w(z))$. Therefore, \mathcal{A}_{j+1} satisfies the equation

$$\partial_w J(z, w(z)) \mathcal{A}'_{j+1}(z) + \partial^2_{zw} J(z, w(z)) \mathcal{A}_{j+1}(z) = G_j(z) ,$$

where

$$G_j(z) := -\left[\partial_z J \frac{T_j}{V} + \partial_z \left(\frac{T_j}{2V}\right)\right](z, w(z))$$

Thus,

$$\mathcal{A}_{j+1}(z) = \mathcal{A}_0(z) \int_{[0,z]} \frac{G_j(u)}{\mathcal{A}_0(u)} \,\mathrm{d}u \,.$$

Therefore, we have, for all $j \ge 0$,

$$\widetilde{a}_{j+1}(z,w) = J(z,w) \int_{[w(z),w]} \frac{T_j(z,u)}{2J(z,u)V(z,u)} \,\mathrm{d}u + \widetilde{a}_0(z,w) \int_{[0,z]} \frac{G_j(u)}{\mathcal{A}_0(u)} \,\mathrm{d}u \,, \tag{3.13}$$

where

$$T_j(z,w) = \int_0^1 \partial_w^2 \partial_z \widetilde{a}_j(z,w(z) + t(w - w(z))) \,\mathrm{d}t \,,$$

and

$$G_j(u) = \left[\left(\frac{\partial_z J}{\mathbf{B}(0)} + \frac{1}{2} \partial_z \left(\frac{1}{V} \right) \right) \partial_w^2 \partial_z \widetilde{a}_j + \frac{1}{2\mathbf{B}(0)} \partial_w^2 \partial_z^2 \widetilde{a}_j + \frac{w'(u)}{4\mathbf{B}(0)} \partial_w^3 \partial_z \widetilde{a}_j \right] (u, w(u)) \,.$$

3.3. **Proof of Theorem 3.2.** We have constructed holomorphic functions $\widetilde{S}(z, w)$ and $(\widetilde{a}_j(z, w))_{j \in \mathbb{N}_0}$ which are defined in a neighbourhood of $0 \in \mathbb{C}^2$. Now we take $w = \overline{z}$ and recall Notation 3.3. For all $N \ge 0$, we have

$$e^{S/h} \left(\mathscr{L}_{h,\mathcal{M}} - h\mu \right) \left(e^{-S/h} \sum_{j=0}^{N} h^j a_j \right) = (-\Delta a_N) h^{N+2}$$

From (3.5), we have

$$e^{P/h} \left(\mathscr{L}_{h,\mathbf{A}} - h\mu \right) \left(e^{-P/h} \sum_{j=0}^{N} h^j a_j \right) = (-\Delta a_N) h^{N+2}, \quad P := S + i\theta.$$
 (3.14)

From the definitions of θ and \mathcal{M} , we have $\nabla \theta(0) = \mathcal{M}(0) - \mathbf{A}(0) = -\mathbf{A}(0)$ and thus $\nabla \operatorname{Im} \theta(0) = -\operatorname{Im} \mathbf{A}(0) = 0$. Notice that

$$\operatorname{Im} \partial_1^2 \theta(0) = \operatorname{Im} \left(-\partial_{12}^2 \varphi(0) - \partial_1 A_1(0) \right) = -\partial_1 \operatorname{Im} A_1(0) ,$$

$$\operatorname{Im} \partial_2^2 \theta(0) = \operatorname{Im} \left(\partial_{12}^2 \varphi(0) - \partial_2 A_2(0) \right) = -\partial_2 \operatorname{Im} A_2(0) ,$$

$$\operatorname{Im} \partial_{12}^2 \theta(0) = \operatorname{Im} \left(\partial_{11}^2 \varphi(0) - \partial_1 A_2(0) \right)$$

$$= \frac{\operatorname{Im} \mathbf{B}(0)}{2} - \partial_1 \operatorname{Im} A_2(0)$$

$$= -\frac{1}{2} \left(\partial_1 \operatorname{Im} A_2(0) + \partial_2 \operatorname{Im} A_1(0) \right) .$$

From Lemma 3.6 and using Taylor expansion for $\text{Im}\,\theta$, we deduce that

$$\operatorname{Re} P(x) = Q_1(0)x_1^2 - 2Q_2(0)x_1x_2 + Q_3(0)x_2^2 + \mathcal{O}(|x|^3).$$

This concludes the proof of Theorem 3.2.

3.4. Proof of Theorem 1.5.

3.4.1. Preliminaries.

Notation 3.7. We will use the following notation for a polydisc

$$P(0; R_1, R_2) \coloneqq \{(z, w) \in \mathbb{C}^2 : |z| < R_1 \text{ and } |w| < R_2\}.$$

For $K \subset \mathbb{C}^2$, we write

$$\|\widetilde{a}(z,w)\|_{K} \coloneqq \sup_{(z,w)\in K} |\widetilde{a}(z,w)|$$

When $K = P(0; R_1, R_2)$, we simply write $\|\cdot\|_K = \|\cdot\|_{R_1, R_2}$. For $m, n \in \mathbb{N}$, we denote $[[m, n]] := [m, n] \cap \mathbb{Z}$.

Lemma 3.8. For a holomorphic function $\tilde{a}(z, w)$ defined in a neighbourhood of $P(0; R_1, R_2)$, we have, for all $(z, w) \in P(0; R_1, R_2)$,

$$\left|\partial_{w}^{2}\partial_{z}\widetilde{a}(z,w)\right| \leq \frac{2R_{1}R_{2}\|\widetilde{a}\|_{R_{1},R_{2}}}{\left(R_{1}-|z|\right)^{2}\left(R_{2}-|w|\right)^{3}}$$

and

$$\left|\partial_{w}^{2}\partial_{z}^{2}\widetilde{a}(z,w)\right| \leqslant \frac{4R_{1}R_{2}\|\widetilde{a}\|_{R_{1},R_{2}}}{(R_{1}-|z|)^{3}\left(R_{2}-|w|\right)^{3}}, \quad \left|\partial_{w}^{3}\partial_{z}\widetilde{a}(z,w)\right| \leqslant \frac{6R_{1}R_{2}\|\widetilde{a}\|_{R_{1},R_{2}}}{(R_{1}-|z|)^{2}\left(R_{2}-|w|\right)^{4}}$$

Proof. For example, the Cauchy formula gives

$$\begin{aligned} \left|\partial_{w}^{2}\partial_{z}\widetilde{a}(z,w)\right| &= \left|\frac{2}{(2\pi i)^{2}}\int_{|z|=R_{1}}\int_{|w|=R_{2}}\frac{\widetilde{a}(\xi_{1},\xi_{2})}{(\xi_{1}-z)^{2}(\xi_{2}-w)^{3}}\,\mathrm{d}\xi_{1}\mathrm{d}\xi_{2}\right| \\ &\leqslant \frac{2R_{1}R_{2}\|\widetilde{a}\|_{R_{1},R_{2}}}{(R_{1}-|z|)^{2}\left(R_{2}-|w|\right)^{3}}\,. \end{aligned}$$

Lemma 3.9. Let $(a_j)_{j \in \mathbb{N}}$ be the real-analytic sequence given in Theorem 3.2, then there exist constants m > 0, $\delta_0 > 0$ such that, for all $j \ge 0$ and for all $x \in D(x^0, \delta_0)$,

 $|a_j(x)| \leq m^{j+1} j^{7j}, \qquad |\nabla a_j(x)| \leq m^{j+1} j^{7j}, \qquad |\Delta a_j(x)| \leq m^{j+1} j^{7j}.$ (3.15)

Proof. Let us prove that there exists m > 0 such that, for all $j \ge 0$ and all (z, w) in a neighbourhood of $0 \in \mathbb{C}^2$,

$$|\tilde{a}_j(z,w)| \leqslant m^{j+1} j^{7j} \,. \tag{3.16}$$

The key to get the growth control is the recursion relation (3.13): we see that \tilde{a}_{j+1} is related to derivatives $\partial_w^2 \partial_z \tilde{a}_j$, $\partial_w^2 \partial_z^2 \tilde{a}_j$, and $\partial_w^3 \partial_z \tilde{a}_j$.

Consider $C, R_1, R_2 > 0$ such that, for all $z \in D(0, R_1)$,

$$|w(z)| \leqslant C|z| < R_2. \tag{3.17}$$

For all $(z, w) \in P(0; R_1, R_2)$ and for all $t \in [0, 1]$, we have

$$w(z) + t(w - w(z))| < \max(C|z|, |w|) < R_2.$$
(3.18)

Then, Lemma 3.8 yields that, for all $(z, w) \in P(0; R_1, R_2)$,

$$|T_{j}(z,w)| \leq \int_{0}^{1} |\partial_{w}^{2} \partial_{z} \tilde{a}_{j}(z,w(z) + t(w - w(z))| dt$$

$$\leq \int_{0}^{1} \frac{2R_{1}R_{2} \|\tilde{a}_{j}\|_{R_{1},R_{2}}}{(R_{1} - |z|)^{2} (R_{2} - |w(z) + t(w - w(z))|)^{3}} dt$$

$$\leq \frac{2R_{1}R_{2} \|\tilde{a}_{j}\|_{R_{1},R_{2}}}{(R_{1} - |z|)^{2} (R_{2} - \max\{C|z|, |w|\})^{3}}.$$

Then, the first term in (3.13) can be controlled, for all $(z, w) \in P(0; R_1, R_2)$,

$$\begin{aligned} \left| J(z,w) \int_{[w(z),w]} \frac{T_j(z,u)}{2J(z,u)V(z,u)} \, \mathrm{d}u \right| \\ &= \left| J(z,w) \int_0^1 \frac{T_j(z,w(z) + t(w - w(z)))}{2(JV)(z,w(z) + t(w - w(z)))} (w - w(z)) \, \mathrm{d}t \right| \\ &\leqslant \|J\|_{R_1,R_2} \left\| \frac{1}{JV} \right\|_{R_1,R_2} \frac{R_1R_2 \|\widetilde{a}_j\|_{R_1,R_2}}{(R_1 - |z|)^2 (R_2 - \max\{C|z|, |w|\})^3} |w - w(z)| \\ &\leqslant \|J\|_{R_1,R_2} \left\| \frac{1}{JV} \right\|_{R_1,R_2} \frac{2R_1R_2^2 \|\widetilde{a}_j\|_{R_1,R_2}}{(R_1 - |z|)^2 (R_2 - \max\{C|z|, |w|\})^3} \right|. \end{aligned}$$

For the second term in (3.13), we have, for all $(z, w) \in P(0; R_1, R_2)$,

$$\begin{aligned} \left| \widetilde{a}_{0}(z,w) \int_{[0,z]} \frac{G_{j}(w)}{\mathcal{A}_{0}(w)} \, \mathrm{d}u \right| \\ \leqslant |z| \left\| \widetilde{a}_{0} \right\|_{R_{1},R_{2}} \left\| \frac{1}{\mathcal{A}_{0}} \right\|_{R_{1}} \\ \times \int_{0}^{1} \left| \left[\left(\frac{\partial_{z}J}{\mathbf{B}(0)} + \frac{1}{2} \partial_{z} \left(\frac{1}{V} \right) \right) \partial_{w}^{2} \partial_{z} \widetilde{a}_{j} + \frac{1}{2\mathbf{B}(0)} \partial_{w}^{2} \partial_{z}^{2} \widetilde{a}_{j} + \frac{w'(tz)}{4\mathbf{B}(0)} \partial_{w}^{3} \partial_{z} \widetilde{a}_{j} \right] (tz,w(tz)) \right| \, \mathrm{d}t \\ \leqslant |z| \left\| \widetilde{a}_{0} \right\|_{R_{1},R_{2}} \left\| \frac{1}{\mathcal{A}_{0}} \right\|_{R_{1}} \left[\left(\frac{\left\| \partial_{z}J \right\|_{R_{1},R_{2}}}{|\mathbf{B}(0)|} + \frac{1}{2} \left\| \partial_{z} \left(\frac{1}{V} \right) \right\|_{R_{1},R_{2}} \right) \int_{0}^{1} \left| \partial_{w}^{2} \partial_{z} \widetilde{a}_{j}(tz,w(tz)) \right| \, \mathrm{d}t \\ + \frac{1}{2|\mathbf{B}(0)|} \int_{0}^{1} \left| \partial_{w}^{2} \partial_{z}^{2} \widetilde{a}_{j}(tz,w(tz)) \right| \, \mathrm{d}t + \frac{\|w'\|_{R_{1}}}{4|\mathbf{B}(0)|} \int_{0}^{1} \left| \partial_{w}^{3} \partial_{z} \widetilde{a}_{j}(tz,w(tz)) \right| \, \mathrm{d}t \right] \, . \end{aligned}$$

From Lemma 3.8, we have, for all $(z, w) \in P(0; R_1, R_2)$,

$$\int_{0}^{1} \left| \partial_{w}^{2} \partial_{z} \tilde{a}_{j}(tz, w(tz)) \right| |z| dt \leq \frac{2R_{1}^{2}R_{2} \left\| \widetilde{a}_{j} \right\|_{R_{1}, R_{2}}}{(R_{1} - |z|)^{2} (R_{2} - C|z|)^{2}},$$

$$\int_{0}^{1} \left| \partial_{w}^{2} \partial_{z}^{2} \tilde{a}_{j}(tz, w(tz)) \right| |z| dt \leq \frac{4R_{1}^{2}R_{2} \left\| \widetilde{a}_{j} \right\|_{R_{1}, R_{2}}}{(R_{1} - |z|)^{3} (R_{2} - C|z|)^{3}},$$

and

$$\int_{0}^{1} \left| \partial_{w}^{3} \partial_{z} \tilde{a}_{j}(tz, w(tz)) \right| |z| \, \mathrm{d}t \leqslant \frac{6R_{1}^{2}R_{2} \left\| \widetilde{a}_{j} \right\|_{R_{1}, R_{2}}}{(R_{1} - |z|)^{2} \left(R_{2} - C|z|\right)^{4}}$$

Let us consider $R_1^0, R_2^0, C_1, C > 0$ such that, for all $R_1 \leq 2R_1^0, R_2 \leq 2R_2^0$, $\|J\|_{R_1, R_2} + \left\|\frac{1}{JV}\right\|_{R_1, R_2} + \|\widetilde{a}_0\|_{R_1, R_2} + \left\|\frac{1}{\mathcal{A}_0}\right\|_{R_1} + \|\partial_z J\|_{R_1, R_2} + \left\|\partial_z \left(\frac{1}{V}\right)\right\|_{R_1, R_2} + \|w'\|_{R_1} \leq C_1$, and, for all $z \in D(0, 2R_1^0)$,

$$|w(z)| \le C|z| < 2R_2^0.$$
(3.19)

From (3.13), there exists a constant $C_2 > 0$ (depending only on R_1^0, R_2^0) such that, for all $\ell \ge 0$ and for all $(z, w) \in P(0; R_1, R_2)$,

$$|\tilde{a}_{\ell+1}(z,w)| \leq C_2 \frac{\|\tilde{a}_{\ell}\|_{R_1,R_2}}{(R_1 - |z|)^3 (R_2 - \max\{C|z|,w\})^4}.$$
(3.20)

We fix $j \in \mathbb{N}$ and let, for all $k \in [[0, j-1]]$,

$$r_{1,k} \coloneqq \left(2 - \frac{k}{j}\right) R_1^0, \qquad r_{2,k} \coloneqq \left(2 - \frac{k}{j}\right) R_2^0$$

Thanks to (3.19), we see that (3.17) holds for $R_1 = r_{1,k}$ and $R_2 = r_{2,k}$ with a constant C defined in (3.19). Thus, the estimate (3.20) also holds for all $k \in [0, j-1]$, *i.e.*, for all $(z,w) \in P(0;r_{1,k},r_{2,k}),$

$$|\widetilde{a}_{k+1}(z,w)| \leq C_2 \frac{\|\widetilde{a}_k\|_{r_{1,k},r_{2,k}}}{(r_{1,k}-|z|)^3 (r_{2,k}-\max\{C|z|,w\})^4}.$$
(3.21)

This holds in particular for all $(z, w) \in P(0; r_{1,k+1}, r_{2,k+1})$. From the second inequality in $(3.19), CR_1^0 < R_2^0$, and thus, for $|z| < r_{1,k+1}$, we have

$$C|z| < Cr_{1,k+1} = C\left(2 - \frac{k+1}{j}\right)R_1^0 < \left(2 - \frac{k+1}{j}\right)R_2^0 = r_{2,k+1}$$

This shows that, for all $(z, w) \in P(0; r_{1,k+1}, r_{2,k+1})$,

$$\frac{1}{(r_{1,k} - |z|)^3 (r_{2,k} - \max\{C|z|, w\})^4} \leq \frac{1}{(r_{1,k} - r_{1,k+1})^3 (r_{2,k} - r_{2,k+1})^4} = \frac{j^7}{(R_1^0)^3 (R_2^0)^4}$$

m (3.21) (with $\ell = k$), there exists $C_2 > 0$ such that, for all $k \in [0, j-1]$

From (3.21) (with $\ell = k$), there exists $C_3 > 0$ such that, for all $k \in [0, j]$ ⊥∥,

$$\|\widetilde{a}_{k+1}\|_{r_{1,k+1},r_{2,k+1}} \leqslant C_3 j^7 \|\widetilde{a}_k\|_{r_{1,k},r_{2,k}}.$$

Multiplying these estimates, we get

$$\|\widetilde{a}_{j}\|_{R_{1}^{0},R_{2}^{0}} \leqslant (C_{3}j^{7})^{j}\|\widetilde{a}_{0}\|_{2R_{1}^{0},2R_{2}^{0}}.$$
(3.22)

Then, the estimate (3.16) follows with $m = \max\{C_3, \|\tilde{a}_0\|_{2R_{2}^{0}, 2R_{2}^{0}}\}$. Finally, the estimates on the derivatives of \tilde{a}_i are elementary consequences of (3.22). Indeed,

$$\begin{split} \widetilde{\partial_{x_1}a_j}(z,w) &= \left(\partial_z \widetilde{a}_j + \partial_w \widetilde{a}_j\right)(z,w) \,, \qquad \widetilde{\partial_{x_2}a_j}(z,w) = i \left(\partial_z \widetilde{a}_j - \partial_w \widetilde{a}_j\right)(z,w) \,, \\ \widetilde{\Delta a_j}(z,w) &= 4 \partial_z \partial_w \widetilde{a}_j(z,w) \,, \end{split}$$

so that Cauchy estimates give, for all $(z, w) \in P(0; R_1^0, R_2^0)$,

$$\begin{split} \left| \widetilde{\partial_{x_1} a_j}(z, w) \right| &\leqslant \frac{R_1^0 R_2^0 (R_1^0 + R_2^0) \|\widetilde{a}_j\|_{R_1^0, R_2^0}}{(R_1^0 - |z|)^2 (R_2^0 - |w|)^2}, \qquad \left| \widetilde{\partial_{x_2} a_j}(z, w) \right| \leqslant \frac{R_1^0 R_2^0 (R_1^0 + R_2^0) \|\widetilde{a}_j\|_{R_1^0, R_2^0}}{(R_1^0 - |z|)^2 (R_2^0 - |w|)^2}, \\ \left| \widetilde{\Delta a_j}(z, w) \right| \leqslant \frac{4R_1^0 R_2^0 \|\widetilde{a}_j\|_{R_1^0, R_2^0}}{(R_1^0 - |z|)^2 (R_2^0 - |w|)^2}. \end{split}$$

Then, by using theses estimates (3.22) for all $(z,w) \in P\left(0; \frac{R_1^0}{2}, \frac{R_2^0}{2}\right)$, this concludes the proof.

3.4.2. Proof of Theorem 1.5. Let us recall (3.1). Since Q is positive, there exist $\delta > 0$ and $M_1, M_2 > 0$ such that

$$M_1|x|^2 \leq \operatorname{Re} P(x) \leq M_2|x|^2$$
, for all $x \in D(0, \delta)$. (3.23)

By considering δ sufficiently small, we can assume that (3.15) holds.

Let $\chi \in C_c^{\infty}(\mathbb{R}^2)$ be a smooth cut-off function which is equal to 1 near 0 and has support in a compact subset of $D(0, \delta)$. We define our pseudomode as

$$u_h(x) \coloneqq \chi(x) e^{-P(x)/h} \left[a_0(x) + \sum_{j=1}^{N(h)} h^j a_j(x) \right], \qquad N(h) \coloneqq \lfloor (emh)^{-1/7} \rfloor.$$

Here, m is a constant appearing in (3.15) and $|\cdot|$ denotes the floor function.

By construction in Theorem 3.2, for all $j \ge 1$, $a_j(0) = 0$. Thus, for all $x \in D(0, \delta)$,

$$|a_j(x)| = \left| \int_0^1 \frac{\mathrm{d}}{\mathrm{d}t} a_j(tx) \,\mathrm{d}t \right| = \left| \int_0^1 x \cdot \nabla a_j(tx) \,\mathrm{d}t \right| \leqslant m^{j+1} j^{7j} |x|.$$

Then, by using (3.15) and the definition of N(h), we have, for some $C_1 > 0$,

$$\sum_{j=1}^{N(h)} h^j |a_j(x)| \leqslant \sum_{j=1}^{N(h)} h^j m^{j+1} j^{7j} |x| \leqslant \sum_{j=1}^{N(h)} m e^{-j} |x| \leqslant C_1 |x|.$$
(3.24)

Since $a_0(0) = 1$, there exists $C_2 > 0$ (independent of h) such that, in a neighbourhood of 0,

$$\left| a_0(x) + \sum_{j=1}^{N(h)} h^j a_j(x) \right| \ge |a_0(x)| - C_1 |x| \ge C_2$$

By using (3.23), we get

$$\int_{\mathbb{R}^2} |u_h(x)|^2 \,\mathrm{d}x \gtrsim h^{\frac{1}{2}} \,.$$

Let us now prove that for each $\varepsilon \in (0,1)$, $e^{\varepsilon P/h} (\mathscr{L}_{h,\mathbf{A}} - h\mu) u_h(x)$ is exponentially small. We write

$$e^{\varepsilon P/h} \left(\mathscr{L}_{h,\mathbf{A}} - h\mu\right) u_h(x)$$

$$= \underbrace{e^{\varepsilon P/h} \left[\mathscr{L}_{h,\mathbf{A}}, \chi\right] \left(e^{-P/h} \sum_{j=0}^N h^j a_j\right)}_{I_h} + \underbrace{\chi e^{\varepsilon P/h} \left(\mathscr{L}_{h,\mathbf{A}} - h\lambda\right) \left(e^{-P/h} \sum_{j=0}^N h^j a_j\right)}_{J_h},$$

where

$$[\mathscr{L}_{h,\mathbf{A}},\chi] = -2h^2 \,\nabla\chi \cdot \nabla - h^2 \,\Delta\chi + 2h \,i\mathbf{A} \cdot \nabla\chi\,.$$

The term I_h is a function supported on $\operatorname{supp}(\nabla \chi)$:

$$I_h = e^{-\frac{(1-\varepsilon)P}{h}} \left[-2h^2 \nabla \chi \cdot \sum_{j=0}^{N(h)} h^j \nabla a_j + \left(-h^2 \Delta \chi + 2h \nabla P \cdot \nabla \chi + 2h \, i \mathbf{A} \cdot \nabla \chi \right) \left(\sum_{j=0}^{N(h)} h^j a_j \right) \right].$$

Thanks to (3.24), we notice that $\sum_{j=0}^{N(h)} h^j a_j(x)$ is bounded uniformly in $x \in D(0, \delta)$ and in

h. Similarly, $\sum_{j=0}^{N(h)} h^j \nabla a_j(x)$ is also uniformly bounded.

Then, there exist $C, \tilde{C} > 0$ such that, for all $h \in (0, h_0)$ and for all $x \in D(0, \delta)$,

$$|I_h(x)| \leqslant C e^{-(1-\varepsilon)M_1|x|^2/h} \mathbf{1}_{\operatorname{supp} \nabla \chi}(x) \leqslant C e^{-(1-\varepsilon)\tilde{C}/h}$$

For J_h , we can notice that, by construction and by using Lemma 3.9,

$$|J_h(x)| = |\chi(x)e^{-(1-\varepsilon)P(x)/h}h^{N(h)+2}\Delta a_{N(h)}| \lesssim (hmN(h)^7)^{N(h)} \lesssim e^{-N(h)} \lesssim e^{-C/h^{1/7}},$$

where, in the last estimate, we used the fact that $N(h) > \frac{1}{(emh)^{1/7}} - 1$.

This concludes the proof of Theorem 1.5.

3.5. Examples. Of course, Theorem 1.5 is only interesting if one can ensure that Γ is not empty. Let us discuss conditions on **A** and **B** to ensure that $0 \in \Gamma$.

First, since $\partial_{\overline{z}} \mathbf{B}(0) \neq 0$, we observe that 0 cannot be a critical point of **B** at 0. Then, also since Im $\mathbf{A}(0) = 0$, we write

$$\mathbf{B}(x_1, x_2) = a + bx_1 + cx_2 + \mathcal{O}\left(|x|^2\right) ,$$

Im $A_1(x_1, x_2) = dx_1 + ex_2 + \mathcal{O}\left(|x|^2\right) ,$
Im $A_2(x_1, x_2) = fx_1 + gx_2 + \mathcal{O}\left(|x|^2\right) ,$

in a neighbourhood of 0, where $a, b, c \in \mathbb{C}$ and $d, e, f, g \in \mathbb{R}$. Below, we will denote a_1, b_1, c_1 (respectively, a_2, b_2, c_2) as the real parts (respectively, imaginary parts) of a, b, c. Let us find conditions on these coefficients so that $0 \in \Gamma$. We have

$$\mathbf{B}(0) = a_1 + ia_2, \quad \partial_z \mathbf{B}(0) = \frac{1}{2} [b_1 + c_2 + i(b_2 - c_1)], \quad \partial_{\overline{z}} \mathbf{B}(0) = \frac{1}{2} [b_1 - c_2 + i(b_2 + c_1)], \\ \partial_{x_1} \operatorname{Im} A_1(0) = d, \quad \partial_{x_2} \operatorname{Im} A_1(0) = e, \quad \partial_{x_1} \operatorname{Im} A_2(0) = f, \quad \partial_{x_2} \operatorname{Im} A_2(0) = g.$$

Then, we obtain

$$\frac{\partial_z \mathbf{B}}{\partial_{\overline{z}} \mathbf{B}}(0) = \frac{b_1^2 + b_2^2 - c_1^2 - c_2^2 - i2(b_1c_1 + b_2c_2)}{(b_1 - c_2)^2 + (b_2 + c_1)^2},$$

Re $\left[B(0)\frac{\partial_z \mathbf{B}}{\partial_{\overline{z}} \mathbf{B}}(0)\right] = \frac{a_1\left(b_1^2 + b_2^2 - c_1^2 - c_2^2\right) + 2a_2(b_1c_1 + b_2c_2)}{(b_1 - c_2)^2 + (b_2 + c_1)^2},$
Im $\left[B(0)\frac{\partial_z \mathbf{B}}{\partial_{\overline{z}} \mathbf{B}}(0)\right] = \frac{a_2\left(b_1^2 + b_2^2 - c_1^2 - c_2^2\right) - 2a_1(b_1c_1 + b_2c_2)}{(b_1 - c_2)^2 + (b_2 + c_1)^2},$

and thus,

$$Q_{1}(0) = \frac{1}{2} \left[\frac{a_{1} (b_{1}^{2} + b_{2}^{2} - b_{1}c_{2} + b_{2}c_{1}) + a_{2}(b_{1}c_{1} + b_{2}c_{2})}{(b_{1} - c_{2})^{2} + (b_{2} + c_{1})^{2}} + d \right],$$

$$Q_{2}(0) = \frac{1}{2} \left[\frac{a_{2} (b_{1}^{2} + b_{2}^{2} - b_{1}c_{2} + b_{2}c_{1}) - a_{1}(b_{1}c_{1} + b_{2}c_{2})}{(b_{1} - c_{2})^{2} + (b_{2} + c_{1})^{2}} + e \right],$$

$$Q_{3}(0) = \frac{1}{2} \left[\frac{a_{1} (c_{1}^{2} + c_{2}^{2} - b_{1}c_{2} + b_{2}c_{1}) - a_{2}(b_{1}c_{1} + b_{2}c_{2})}{(b_{1} - c_{2})^{2} + (b_{2} + c_{1})^{2}} + g \right].$$

Here, in $Q_2(0)$, we have used $f - e = a_2$ deduced from $\partial_{x_1} \operatorname{Im} A_2(0) - \partial_{x_2} \operatorname{Im} A_1(0) = \operatorname{Im} \mathbf{B}(0)$. Then, $0 \in \Gamma$ if and only if

$$a_1 \neq 0 \text{ or } a_2 \neq 0,$$
 (3.25)

$$b_1 \neq c_2 \text{ or } b_2 \neq -c_1 ,$$
 (3.26)

$$Q_1(0) > 0, (3.27)$$

$$Q_1(0)Q_3(0) - Q_2^2(0) > 0. (3.28)$$

Example 4 (Polynomial complex magnetic fields). In this example, we want to provide a class of polynomial **B** such that $\mathscr{L}_{h,\mathbf{A}}$ is well defined and $0 \in \Gamma$. We consider

$$\mathbf{B}(x_1, x_2) = a + bx_1 + cx_2 + R(x_1, x_2),$$

where

- $a, b, c \in \mathbb{C}$ such that $a_1 > 0, a_2 = 0$ and $b_2c_1 b_1c_2 > 0$, $R : \mathbb{R}^2 \to \mathbb{R}$ is a polynomial such that

$$R(0) = \partial_{x_1} R(0) = \partial_{x_2} R(0) = 0$$
 and $|x|^4 = o(R(x))$ as $|x| \to +\infty$.

Let us choose the following magnetic potential

$$A_1(x_1, x_2) = 0$$
, $A_2(x_1, x_2) = \int_0^{x_1} \mathbf{B}(s, x_2) \, \mathrm{d}s$.

Since R is real-valued, we have

Re **B**
$$(x_1, x_2) = a_1 + b_1 x_1 + c_1 x_2 + R(x_1, x_2)$$
, Im $A_2(x_1, x_2) = \frac{b_2}{2} x_1^2 + c_2 x_1 x_2$

From the asymptotic behaviour of R, we deduce that

$$|\operatorname{Im} \mathbf{A}(x)|^2 = o(\operatorname{Re} \mathbf{B}(x)), \quad \text{as } |x| \to +\infty.$$

Then, (C1) is satisfied (and thus Assumption 1.1 holds). This allows us to define $\mathscr{L}_{h,\mathbf{A}}$ by Theorem 1.2. Furthermore, since $|\operatorname{Re} \mathbf{B}(x)| \to +\infty$ as $|x| \to +\infty$, the operator has discrete eigenvalues by Theorem 1.4.

Since $a_1 > 0$ and $b_2c_1 - b_1c_2 > 0$, (3.25) and (3.26) hold. Note that d = e = g = 0 and $a_2 = 0$, we have

$$Q_1(0) = \frac{1}{2} \frac{a_1 (b_1^2 + b_2^2 + b_2 c_1 - b_1 c_2)}{(b_1 - c_2)^2 + (b_2 + c_1)^2},$$

$$Q_2(0) = -\frac{1}{2} \frac{a_1 (b_1 c_1 + b_2 c_2)}{(b_1 - c_2)^2 + (b_2 + c_1)^2},$$

$$Q_3(0) = \frac{1}{2} \frac{a_1 (c_1^2 + c_2^2 + b_2 c_1 - b_1 c_2)}{(b_1 - c_2)^2 + (b_2 + c_1)^2}.$$

Obviously, $Q_1(0) > 0$. Let us compute

$$Q_1(0)Q_3(0) - Q_2(0)^2 = \frac{a_1^2 \left[(b_1^2 + b_2^2 + b_2c_1 - b_1c_2) (c_1^2 + c_2^2 + b_2c_1 - b_1c_2) - (b_1c_1 + b_2c_2)^2 \right]}{4 \left[(b_1 - c_2)^2 + (b_2 + c_1)^2 \right]^2}$$

Notice that

$$(b_1^2 + b_2^2 + b_2c_1 - b_1c_2) (c_1^2 + c_2^2 + b_2c_1 - b_1c_2) - (b_1c_1 + b_2c_2)^2$$

= $(b_2c_1 - b_1c_2) [(b_1 - c_2)^2 + (b_2 + c_1)^2].$

This yields that

$$Q_1(0)Q_3(0) - Q_2(0)^2 = \frac{a_1^2 (b_2 c_1 - b_1 c_2)}{4(b_1 - c_2)^2 + (b_2 + c_1)^2} > 0$$

Thus, we have $0 \in \Gamma$. By Theorem 1.5, there exist a pseudomode $u(\cdot, h)$ such that

$$\left\| \left(\mathscr{L}_{h,\mathbf{A}} - ha_1 \right) u_h(x) \right\| \leqslant \exp\left(-\frac{C}{h^{1/7}}\right) \left\| u_h(x) \right\|, \quad \text{as } h \to 0.$$
(3.29)

Example 5 (Bounded oscillating magnetic fields). Consider the magnetic potential

$$A_1(x_1, x_2) = -\sin(x_1)x_2 + i\cos(x_2), \qquad A_2(x_1, x_2) = i\cos(x_2)$$

We have

 $\mathbf{B}(x_1, x_2) = \sin(x_1) + i \sin(x_2).$

Since Im **A** is bounded, both conditions (C1) and (C2) hold, therefore, the magnetic Laplacian $\mathscr{L}_{h,\mathbf{A}}$ is well-defined by Theorem 1.2. Let us find the set Γ in this explicit example. Take $x^0 = (x_1, x_2) \in \Gamma$, we have

$$\operatorname{Im} \mathbf{A}(x^0) = 0 \Longleftrightarrow \cos(x_2) = 0.$$

Then, it implies that $\mathbf{B}(x^0) \neq 0$ and

$$\partial_{\overline{z}} \mathbf{B}(x^0) = \partial_z \mathbf{B}(x^0) = \frac{1}{2} \cos(x_1).$$

Therefore, it is obvious that

$$\partial_{\overline{z}} \mathbf{B}(x^0) \neq 0 \iff \cos(x_1) \neq 0$$
.

By straightforward calculation, we obtain

$$Q_1(x^0) = \frac{1}{2}\sin(x_1), \qquad Q_2(x^0) = 0, \qquad Q_3(x^0) = -\frac{1}{2}\sin(x_2).$$

Therefore, it yields that

$$Q_1(x^0) > 0 \iff \sin(x_1) > 0,$$

 $Q_1(x^0)Q_3(x^0) - Q_2^2(x^0) > 0 \iff \sin(x_2) = -1.$

The above analysis leads to

$$\Gamma = \{ (x_1, x_2) \in \mathbb{R}^2 : \sin(x_1) > 0, \ \cos(x_1) \neq 0, \ \cos(x_2) = 0, \ \sin(x_2) = -1 \} \\ = \left\{ (0, \pi) \setminus \left\{ \frac{\pi}{2} \right\} + 2k\pi : k \in \mathbb{Z} \right\} \times \left\{ -\frac{\pi}{2} + 2\pi n : n \in \mathbb{Z} \right\} .$$

Then, Theorem 1.5 states that: for each $x_1^0 \in (0,\pi) \setminus \left\{\frac{\pi}{2}\right\} + 2k\pi$ for some $k \in \mathbb{Z}$, we can construct a pseudomode $u(\cdot, h)$ such that

$$\left\| \left(\mathscr{L}_{h,\mathbf{A}} - h(\sin(x_1^0) - i) \right) u_h(x) \right\| \leq \exp\left(-\frac{C}{h^{1/7}}\right) \|u_h(x)\|, \quad \text{as } h \to 0.$$
(3.30)

Acknowledgements

T.N.D. would like to express his heartfelt gratitude to his late mother, Thi Quy Nguyen, whose love and support have been a constant source of inspiration. This work is dedicated to her memory. D.K. and T.N.D. were supported by the EXPRO grant number 20-17749X of the Czech Science Foundation (GAČR). N.R. and T.N.D. are grateful to CIRM where this work was started.

References

- A. A. Abramov, A. Aslanyan, and E. B. Davies, Bounds on complex eigenvalues and resonances, J. Phys. A: Math. Gen. 34 (2001), 57–72.
- 2. N. Ananikian and R. Kenna, Imaginary magnetic fields in the real world, Physics 8 (2015), 2.
- A. Arnal and P. Siegl, Resolvent estimates for one-dimensional Schrödinger operators with complex potentials, J. Funct. Anal. 284 (2023), 109856.
- S. Bögli, Schrödinger operator with non-zero accumulation points of complex eigenvalues, Comm. Math. Phys. 352 (2017), 629–639.
- S. Bögli and J.-C. Cuenin, Counterexample to the Laptev-Safronov conjecture, Comm. Math. Phys. 398 (2023), 1349–1370.
- Y. Bonthonneau and N. Raymond, WKB constructions in bidimensional magnetic wells, Math. Res. Lett. 27 (2020), 647–664.
- C. Cheverry and N. Raymond, A guide to spectral theory: Applications and exercises, Birkhäuser Advanced Texts: Basler Lehrbücher, Birkhäuser/Springer, Cham, 2021.
- J.-C. Cuenin, Schrödinger operators with complex sparse potentials, Comm. Math. Phys. 392 (2022), 951–992.
- E. B. Davies, Semi-classical states for non-self-adjoint Schrödinger operators, Comm. Math. Phys. 200 (1999), 35–41.
- N. Dencker, J. Sjöstrand, and M. Zworski, Pseudospectra of semiclassical (pseudo-) differential operators, Comm. Pure Appl. Math. 57 (2004), 384–415.
- J. J. Duistermaat and J. A. C. Kolk, *Multidimensional real analysis. I. Differentiation*, Cambridge Studies in Advanced Mathematics, vol. 86, Cambridge University Press, Cambridge, 2004.
- Y. Guedes Bonthonneau, T. Nguyen Duc, N. Raymond, and S. Vũ Ngọc, Magnetic WKB constructions on surfaces, Rev. Math. Phys. 33 (2021), 2150022.
- N. Hatano and D. R. Nelson, Localization transitions in non-Hermitian quantum mechanics, Phys. Rev. Lett. 77 (1996), 570–573.
- 14. _____, Non-Hermitian delocalization and eigenfunctions, Phys. Rev. B 58 (1998), 8384–8390.
- A. Iwatsuka, Magnetic Schrödinger operators with compact resolvent, J. Math. Kyoto Univ. 26 (1986), 357–374.
- J. L. Jaramillo, Black hole horizons and quantum charged particles, Class. Quantum Grav. 32 (2015), 132001.
- 17. _____, A perspective on black hole horizons from the quantum charged particle, J. Phys. Conf. Ser. 600 (2015), 012037.
- T. Kato, Perturbation theory for linear operators, Classics in Mathematics, Springer-Verlag, Berlin, 1995, Reprint of the 1980 edition.
- D. Krejčiřík, Complex magnetic fields: an improved Hardy-Laptev-Weidl inequality and quasi-selfadjointness, SIAM J. Math. Anal. 51 (2019), 790–807.
- D. Krejčiřík and P. Siegl, Pseudomodes for Schrödinger operators with complex potentials, J. Funct. Anal. 276 (2019), 2856–2900.

- D. Krejčiřík, P. Siegl, M. Tater, and J. Viola, Pseudospectra in non-Hermitian quantum mechanics, J. Math. Phys. 56 (2015), 103513, 32.
- 22. E. H. Lieb and M. Loss, *Analysis*, second ed., Graduate Studies in Mathematics, vol. 14, American Mathematical Society, Providence, RI, 2001.
- K. Miller and B. Simon, Quantum magnetic Hamiltonians with remarkable spectral properties, Phys. Rev. Lett. 44 (1980), 1706–1707.
- 24. X. Peng, H. Zhou, B.-B. Wei, J. Cui, J. Du, and R.-B. Liu, *Experimental observation of Lee-Yang zeros*, Phys. Rev. Lett. **114** (2015), 010601.
- J. A. Reggia, Generalizing Maxwell's equations to complex-valued electromagnetic fields, Physica Scripta 99 (2023), 015513.
- F. G. Scholtz, H. B. Geyer, and F. J. W. Hahne, Quasi-Hermitian operators in quantum mechanics and the variational principle, Ann. Phys 213 (1992), 74–101.

(David Krejčiřík) DEPARTMENT OF MATHEMATICS, FACULTY OF NUCLEAR SCIENCES AND PHYSICAL ENGINEERING, CZECH TECHNICAL UNIVERSITY IN PRAGUE, TROJANOVA 13, 12000 PRAGUE, CZECHIA. *Email address*: david.krejcirik@fjfi.cvut.cz

(Tho Nguyen Duc) DEPARTMENT OF MATHEMATICS, FACULTY OF NUCLEAR SCIENCES AND PHYSICAL ENGINEERING, CZECH TECHNICAL UNIVERSITY IN PRAGUE, TROJANOVA 13, 12000 PRAGUE, CZECHIA. *Email address:* nguyed16@fjfi.cvut.cz

(Nicolas Raymond) UNIV ANGERS, CNRS, LAREMA, INSTITUT UNIVERSITAIRE DE FRANCE, SFR MATHSTIC, F-49000 ANGERS, FRANCE

Email address: nicolas.raymond@univ-angers.fr