# A FEW LAST WORDS ON POINTWISE MULTIPLIERS OF CALDERÓN–LOZANOVSKIĬ SPACES

TOMASZ KIWERSKI AND JAKUB TOMASZEWSKI

ABSTRACT. We will provide a complete description of the space  $M(X_F, X_G)$  of pointwise multipliers between two Calderón–Lozanovskiĭ spaces  $X_F$  and  $X_G$  built upon a rearrangement invariant space  $X$  and two Young functions  $F$  and  $G$ . Meeting natural expectations, the space  $M(X_F, X_G)$  turns out to be another Calderón–Lozanovskiĭ space  $X_{G\ominus F}$  with  $G\ominus F$  being the appropriately understood generalized Young conjugate of G with respect to F. Nevertheless, our argument is not a mere transplantation of existing techniques and requires a rather delicate analysis of the interplay between the space X and functions  $F$  and  $G$ . Furthermore, as an example to illustrate applications, we will solve the factorization problem for Calderón–Lozanovski $\tilde{a}$  spaces. All this not only complements and improves earlier results (basically giving them the final touch), but also confirms the conjecture formulated by Kolwicz, Leśnik and Maligranda in [Pointwise multipliers of Calderón–Lozanovski $\check{i}$  spaces, Math. Nachr. 286 (2012), no. 8-9, 876– 907]. We will close this work by formulating a number of open questions that outline a promising panorama for future research.

#### CONTENTS



1

#### 1. Introduction

<span id="page-1-0"></span>1.a. **Goals.** The main goal of this paper is to provide a description of the space  $M(X_F, X_G)$ of pointwise multipliers between two Calderón–Lozanovkiĭ spaces  $X_F$  and  $X_G$ . By this we simply mean that we are looking for conditions on the function  $f$  guaranteeing that the induced multiplication operator  $M_f : g \mapsto fg$  is bounded when acting from  $X_F$  into  $X_G$ . Here, and hereinafter, X is a rearrangement invariant space, while F and G are two Young functions. (For all unexplained concepts that have already appeared, or will appear, we refer to Sections [3,](#page-4-1) [4.a](#page-9-2) and [5.a.](#page-18-1)) However, before presenting our results in more detail, let us first try to explain what the sources of this problem are, what was our motivation and why all this may be even important.

1.b. Back to Orlicz spaces. From a historical perspective, the *fons et origo* of the problem of describing the space of pointwise multipliers  $M(X, Y)$  between two function spaces X and Y can most likely be traced back to O'Neil's work from 1965. In  $[O'Ne65, Prob$ lem 6.2], he asks

**Problem.** (O'Neil, 1965). How, given two Young functions  $F$  and  $G$ , to choose the third Young function, say H, so that the space  $M(L_F, L_G)$  of pointwise multipliers between two Orlicz spaces  $L_F$  and  $L_G$  is another Orlicz space  $L_H$ ?

Many authors over nearly half a century have obtained mainly partial<sup>\*</sup> results in this direction (see, for example, [\[And60\]](#page-25-1), [\[O'Ne65\]](#page-27-0), [\[Mau74\]](#page-27-1), [\[MP89\]](#page-26-0), [\[MN10\]](#page-26-1) and [\[ZR67\]](#page-27-2)). Putting some technical details aside, the generic result of this type looks more-or-less like this: If three given Young functions satisfy the relation  $F^{-1}G^{-1} \approx H^{-1}$ , then  $M(L_F, L_G) =$  $L_H$ . Unfortunately, such a results do not lead to any explicit formula for the Young function H generating the space of pointwise multipliers. Even worse, there are pairs of Young functions, say  $F$  and  $G$ , for which there is no third Young function  $H$  satisfying the relation  $F^{-1}G^{-1} \approx H^{-1}$  even though the space  $M(L_F, L_G)$  is a non-trivial Orlicz space. There are probably two noted examples of this kind in the existing literature, namely, [\[KLM12,](#page-26-2) Example 7.8] and [\[KT22,](#page-26-3) Example A.2]. The former one concerns Orlicz spaces defined on  $\mathbb{I} = [0, 1]$  and gives  $M(L_F, L_G) = L_{\infty}[0, 1]$ , while the second one

Date: Thursday 3<sup>rd</sup> October, 2024.

<sup>2020</sup> Mathematics Subject Classification. Primary: 46E30; Secondary: 46B03, 46B20, 46B42.

Key words and phrases. Calderón–Lozanovskiĭ spaces; generalized Young conjugate; pointwise multipliers; pointwise products; factorization.

<span id="page-1-1"></span><sup>∗</sup>Note that some satisfactory results crowning these struggles have only been obtained much more recently in [\[DR00\]](#page-25-2) and [\[LT17\]](#page-26-4).

covers both remaining cases, that is,  $\mathbb{I} = [0, \infty)^*$  and  $\mathbb{I} = \mathbb{N}$ , but is also somehow more sophisticated, because  $M(L_F, L_G) \neq L_{\infty}$ .

For the reasons outlined above, our work can naturally be placed within the framework of O'Neil's problem in which the classical Orlicz spaces  $L_F$  and  $L_G$  are replaced by a much more general<sup>[†](#page-2-1)</sup> constructions of the Calderón–Lozanovskiı̆ spaces  $X_F$  and  $X_G$ , respectively.

1.c. One conjecture. Apart from these historical marginalia, our work starts where [\[KLM12\]](#page-26-2) and [\[KLM14\]](#page-26-5) ends. In fact, our initial motivation comes form the following conjecture formulated at the end of [\[KLM12\]](#page-26-2)

Conjecture. (Kolwicz, Leśnik and Maligranda, 2012). Let X be a rearrangement invariant function space defined on  $\mathbb{I} = [0, 1]$ . Suppose that  $X \neq L_{\infty}[0, 1]$ . Then we have the equality  $M(X_F, X_G) = X_{G \ominus F}$ .

While working on this problem we realize, to our pleasant surprise, that we were not only able to confirm the above conjecture, but even provide a necessary and sufficient conditions on the triple  $(X, F, G)$  for equality  $M(X_F, X_G) = X_{G\ominus F}$  to holds (this, we hope, justifies the slightly funeral character of the title of our work).

Notabene, we were also able to resolve the case when the space  $X$  is defined on  $\mathbb{I} = [0, \infty)$ . This is also a bit of surprise, because the structure of rearrangement invariant spaces defined on  $\mathbb{I} = [0, \infty)$  is much more complicated that on  $\mathbb{I} = [0, 1]$ . For instance, there is a whole menagerie of spaces that resemble  $L_{\infty}[0,\infty)$  in the sense that they have no order continuous functions (they are, so to speak, "extremely" non-separable). In fact, just take any rearrangement invariant space defined on  $\mathbb{I} = [0, \infty)$ , say X, and consider the intersection space  $X \cap L_{\infty}$  defined via the norm  $||f||_{X \cap L_{\infty}} = \max{||f||_X, ||f||_{L_{\infty}}}$ . On the other hand, up to the equivalence of norms, there is only one rearrangement invariant space defined on  $\mathbb{I} = [0, 1]$  devoid of order continuous functions, namely,  $L_{\infty}[0, 1]$  itself. This is most likely the reason why the above conjecture was formulated only in the case when the space X is defined on  $\mathbb{I} = [0, 1]$ .

1.d. Overview. Let us take a closer look at our results. The first one concerns the description of the space of pointwise multipliers.

<span id="page-2-2"></span>Theorem A (Theorem [4.b.3](#page-11-0) and Theorem [4.b.5\)](#page-14-0).

- (1)  $M(X_F, X_G) = X_{G\ominus F}$  if, and only if, the triple  $(X, F, G)$  is nice.
- (2)  $M(X_F, X_G) = X_{G\ominus_1 F}$  if, and only if, either the triple  $(X, F, G)$  fails to be nice or the space  $X$  is a sequence space.

In particular, since triples  $(X, F, G)$ , where X is a rearrangement invariant space on  $\mathbb{I} = [0, 1]$  such that  $X \neq L_{\infty}[0, 1]$ , are always nice, so (1) corresponds exactly with Kolwicz, Leśnik and Maligranda's conjecture.

<span id="page-2-0"></span><sup>\*</sup>Actually, what is written in [\[KT22,](#page-26-3) Example A.2] does not cover the case  $\mathbb{I} = [0, \infty)$  at all, but with some modicum of solid effort can be modified to work then too.

<span id="page-2-1"></span><sup>&</sup>lt;sup>†</sup>After all, the class of Orlicz spaces  $L_F$  is just a particular case of Calderón–Lozanovskiı's construction  $X_F$ , where the space X is chosen to be  $L_1$ .

<span id="page-3-0"></span>Inspired by [\[KLM14\]](#page-26-5), we will use these results to provide a final picture on the factorization of Calderón–Lozanovski $\tilde{a}$  spaces. The overall result in this direction is as follows.

**Theorem B** (Theorem [5.c.1\)](#page-22-1). Let X be a rearrangement invariant space.

- (1) Suppose that  $X \neq L_{\infty}$ . Then the factorization  $X_F \odot M(X_F, X_G) = X_G$  holds if, and only if,  $F^{-1}(G \ominus F)^{-1} \approx G^{-1}$  (this equivalence should be understood in the appropriate way).
- (2) Suppose that  $X = L_{\infty}$ . Then  $X_F \odot M(X_F, X_G) = X_G$  regardless of the functions F and G.

Theorems [A](#page-2-2) and [B](#page-3-0) extends and completes many earlier results obtained, for example, by Ando [\[And60\]](#page-25-1); Dankert [\[Dan74\]](#page-25-3); Djakov and Ramanujan [\[DR00\]](#page-25-2); Kolwicz, Leśnik and Maligranda [\[KLM12\]](#page-26-2), [\[KLM14\]](#page-26-5); Leśnik and Tomaszewski [\[LT17\]](#page-26-4), [\[LT21\]](#page-26-6); Maligranda and Nakai [\[MN10\]](#page-26-1); Maligranda and Persson [\[MP89\]](#page-26-0); O'Neil [\[O'Ne65\]](#page-27-0); Zabreĭko and Rutickiĭ  $|ZR67|$ .

It seems to us that there is one more, slightly obscure, thing about Theorems [A](#page-2-2) and [B](#page-3-0) that encompasses aesthetic imperative. Their formulations are simply transparent, which quite ruthlessly corresponds with previous results.

1.e. Key ideas. A few things that could be considered the most innovative in our approach should probably be clearly highlighted. Let's do this now.

 $\star$  Contrary to the results so far, we are not trying to find some conditions on the Young functions F and G that guarantee that the space  $M(X_F, X_G)$  is again the Calderón– Lozanovskiĭ space. Instead, we simply want to describe the space  $M(X_F, X_G)$ . Anyhow, the fact that  $M(X_F, X_G)$  is indeed another Calderón–Lozanovskiĭ space can be considered a fortunate coincidence.

**★** We heavily use the machinery of generalized Young's conjugate functions. This idea is deeply rooted in the recent work of Leśnik and the second-named author [\[LT17\]](#page-26-4) and [\[LT21\]](#page-26-6), which in turn draw inspiration from the pioneering paper of Djakov and Ramanujan [\[DR00\]](#page-25-2).

**★** While working on factorization, we are not interested in the factorization of the form  $X_F \odot X_H = X_G$ . Instead, we want to know when  $X_F \odot M(X_F, X_G) = X_G$ . This is only apparently weaker. In fact, if  $X_F \odot X_H = X_G$  then  $X_H \hookrightarrow M(X_F, X_G)$ . This simple observation, combined with the knowledge that  $M(X_F, X_G) = X_{G\ominus F}$ , leads to a factorization problem of the form  $X_F \odot X_{G\ominus F} = X_G$ . This is somehow a little easier than the initial problem.

 $\star$  We don't need any separability assumptions imposed on the space X. This is probably quite remarkable, because this assumption has been present, in one form or anther, in almost all previous results. This also clearly distinguishes our case from the case of Orlicz spaces (after all,  $L_F = (L_1)_F$  and  $L_1$  is separable).

 $\star$  The formalism we propose supports all three separable measure spaces, namely,  $\mathbb{I} = [0, 1]$  or  $\mathbb{I} = [0, \infty)$  with the Lebesgue measure m and  $\mathbb{I} = \mathbb{N}$  with the counting measure  $\#$ , simultaneously. From the perspective of Luxemburg's representation theorem, that's all one could ever expect in the realm of rearrangement invariant spaces.

1.f. Outline. Let us now briefly describe the organization of this work. Overall, the paper is divided into six sections.

The first section after Introduction, that is, Section [3,](#page-4-1) is of a preliminary nature. Here we will recollect the needful background and provide a handful of useful facts. We will pay special attention to the Calderón–Lozanovski˘ı construction (Section [3.b\)](#page-6-0), without forgetting about the construction of the space of pointwise multipliers (Section [3.c\)](#page-8-0) and pointwise products (Section [3.d\)](#page-9-0).

Next part, that is, Section [4,](#page-9-1) should be considered as the main part of this work. Apart from a brief discussion about the generalized Young conjugate (Section [4.a\)](#page-9-2), it contains essentially two results, which are Theorem [4.b.3](#page-11-0) and Theorem [4.b.5.](#page-14-0) They provide necessary and sufficient condition for the equality  $M(X_F, X_G) = X_{G\ominus F}$  to hold. We will conclude this section with some specific examples to illustrate our general results (see Examples [4.b.6,](#page-16-0) [4.b.7,](#page-16-1) [4.b.8](#page-17-0) and [4.b.9\)](#page-17-1).

Section [5](#page-18-0) is devoted to applications. After a short introduction to the factorization problem (Section [5.a\)](#page-18-1), we present several technical lemmas that will prove necessary later (Section [5.b\)](#page-18-2). The culmination is Section [5.c,](#page-22-0) where we prove factorization theorem for Calderón–Lozanovskiĭ spaces (precisely, see Theorem [5.c.1\)](#page-22-1).

Lastly, in Section [6,](#page-23-0) we have collected several open problems whose potential solution would significantly extend and complement the results obtained here.

1.g. Acknowledgments. The results presented here confirm certain beliefs that the second-named author acquired after completing his doctoral dissertation [\[Tom21\]](#page-26-7) (written under supervision of Karol Leśnik and Ryszard Płuciennik).

#### 2. Statements and Declarations

<span id="page-4-0"></span>Funding. Our research was carried out at the Poznań University of Technology (grant number 0213/SBAD/0116).

Conflict of Interest. The authors declare that they have no known competing financial interest or personal relationship that could have appeared to influence the work reported in this paper.

<span id="page-4-1"></span>Data Availibility. All data generated or analysed during this study are included in this article.

### 3. Toolbox

Our notation and terminology is rather standard and is in-line with what can be found in the classical monographs by Bennett and Sharpley [\[BS88\]](#page-25-4), and by Lindenstrauss and Tzafriri [\[LT79\]](#page-26-8). Below, we provide in detail the most important definitions, terminology and some essential facts which we will use hereinafter. This is dictated not only by the reader's convenience, but also by the need to organize some of the material and adapt it to our purposes. We have also made some effort to ensure that this work is largely self-contained. Other concepts omitted here will be introduced where necessary.

<span id="page-5-0"></span>3.a. **Banach ideal spaces.** For a complete and  $\sigma$ -finite measure space  $(\Omega, \Sigma, \mu)$ , let  $L_0(\Omega)$ , briefly just  $L_0$  if the context leaves no ambiguity, be the set consisting of all equivalence classes, by the Kolmogorov quotient modulo equality almost everywhere, of real- or complex-valued measurable functions defined on  $\Sigma$ . As usual, we consider the space  $L_0$  with the topology of convergence in measure on sets of finite measure, that is, the topology of local convergence in measure. This makes  $L_0$  an  $F$ -space.

A Banach space X is called a **Banach ideal space** (using another common nomenclature, a Köthe space or a Banach function lattice) if the following two conditions hold:

- (1) X is a linear subspace of  $L_0$ ;
- (2) if  $|f(\omega)| \leq |g(\omega)|$  almost everywhere on  $\Omega$ , with f measurable and  $g \in X$ , then also  $f \in X$  and  $||f|| \le ||g||$  (the so-called **ideal property**).

Further, recall that a Banach ideal space  $X$  is said to be rearrangement invariant (or symmetric) if additionally:

- (3) for any two measurable functions, say f and g, with  $\mu(\{\omega \in \Omega : |f(\omega)| > \lambda\}) =$  $\mu(\{\omega \in \Omega : |g(\omega)| > \lambda\})$  for all  $\lambda \geq 0$  and  $f \in X$ , it follows that  $g \in X$  and  $||f|| = ||q||;$
- (4) for any increasing sequence  $\{f_n\}_{n=1}^{\infty}$  of non-negative functions from X converging almost everywhere to f such that  $\sup\{\|f_n\| : n \in \mathbb{N}\}\$ is finite, it follows that f belongs to X and  $||f|| = \sup{||f_n|| : n \in \mathbb{N} \}$  (this is the so-called **Fatou prop**erty[∗](#page-5-1) ).

Using the language of the interpolation theory, rearrangement invariant spaces defined as above are exactly interpolation spaces with respect to the copule  $(L_{\infty}, L_1)$ . Evidently, condition (3) ensures that  $||f|| = ||f^*||$ , where  $f^*$  is the **non-increasing rearrangement** of f, that is,  $f^*(t) := \inf \{ \lambda > 0 : \mu (\{ \omega \in \Omega : |f(\omega)| > \lambda \}) \leq t \}$  for  $t \geq 0$ . Moreover, thanks to Luxemburg's representation theorem (see [\[BS88,](#page-25-4) Theorem 4.10, p. 62] and [\[LT79,](#page-26-8) pp. 114–115]), it is enough to consider rearrangement invariant spaces defined on one of the following three separable[†](#page-5-2) measure spaces:

- $\star$  the set of positive integers  $\mathbb{I} = \mathbb{N}$  with the counting measure # (in this situation we will talk about **sequence spaces**);
- $\star$  the unit interval  $\mathbb{I} = [0, 1]$  or the half-line  $\mathbb{I} = [0, \infty)$  with the usual Lebesgue measure (with the convention that we will call them **function spaces**).

Note that the most prolific classes of function spaces such as Lebesgue spaces, Orlicz spaces and Lorentz spaces are indeed rearrangement invariant.

A function f from a Banach ideal space X is said to be **order continuous**<sup>[‡](#page-5-3)</sup> if, for any sequence  $\{f_n\}_{n=1}^{\infty}$  of positive functions that is bounded above by  $|f|$  and converges almost everywhere to zero, it follows that  $\{f_n\}_{n=1}^{\infty}$  is norm null sequence. By  $X_o$  we denote a

<span id="page-5-1"></span><sup>\*</sup>Roughly speaking, this means that the closed unit ball of  $X$  is also closed with respect to the topology of local convergence in measure.

<span id="page-5-2"></span><sup>&</sup>lt;sup>†</sup>It is well-known that the measure space  $(\Omega, \Sigma, \mu)$  is separable if, and only if,  $L_1(\Omega)$  is separable as a Banach space (see [\[Zaa67,](#page-27-3) p. 137]).

<span id="page-5-3"></span><sup>&</sup>lt;sup>‡</sup>Plainly, any function f from  $L_1(\Omega)$  is order continuous. Thus, this definition is nothing else, but Lebesgue's dominated convergence theorem under more abstract clothes.

closed subspace of all order continuous functions from X. We will say that the space X is order continuous if  $X = X_o$ . Note that the a Banach ideal space X is order continuous if, and only if, it is separable (see [\[BS88,](#page-25-4) Theorem 5.5, p. 27]). Therefore, hereinafter, we will use these terms interchangeably.

By the Köthe dual (or, the associated space)  $X^{\times}$  of a given Banach ideal space X we understand here a vector space all measurable functions  $f$  such that  $fg$  is integrable for all  $g \in X$  equipped with the norm  $||f||_{X^{\times}} := \sup{||fg||}_{L_1} : ||g||_X \leq 1$ . Recall that  $X \equiv X^{\times}$  if, and only if, the norm in X has the Fatou property. Moreover, for any given order continuous Banach ideal space X, its Köthe dual  $X^*$  can be naturally identified with the topological dual  $X^*$ , that is, the space of all continuous linear forms on X (see [\[BS88,](#page-25-4) Corollary 4.3, p. 23]). Basically for this reason, a Banach ideal space X is reflexive if, and only if, both X and  $X^{\times}$  are order continuous (see [\[BS88,](#page-25-4) Corollary 4.4., p. 23]).

Recall that the **fundamental function**  $\psi_X : \mathbb{I} \to [0, \infty)$  of a rearrangement invariant space X is defined by the formula  $\psi_X(t) := ||1_{[0,t)}||_X$  for  $t \in \mathbb{I}$ . We will say that the fundamental function  $\psi_X$  does not vanish at zero if  $\psi_X(0^+) := \lim_{t \to 0^+} \psi_X(t) > 0$ . It is straightforward to see that  $\psi_X$  does not vanish at zero if, and only if, X is a subspace of  $L_{\infty}$  if, and only if, the ideal  $X_o$  is trivial (see, for example, [\[KT17,](#page-26-9) Theorem B]).

We ought to mention that the embedding  $X \subset Y$  between two Banach ideal spaces is always continuous, that is,  $\|\text{id}: X \to Y\| = \sup\{\|f\|_Y : \|f\|_X = 1\}$  is finite. To duly emphasize this fact, we shall rather write  $X \hookrightarrow Y$ . Moreover, the symbol  $X = Y$  indicate that the spaces  $X$  and  $Y$  are the same as vector spaces and their norms are equivalent. Plainly,  $X = Y$  if, and only if,  $X \hookrightarrow Y$  and  $Y \hookrightarrow X$ . Occasionally, we will write  $X \equiv Y$ , understanding that  $X = Y$ , but this time both norms are even equal.

We refer to the books by Bennett and Sharpley [\[BS88\]](#page-25-4), Brudnyĭ and Krugljak [\[BK91\]](#page-25-5), and Lindenstrauss and Tzafriri [\[LT79\]](#page-26-8) for a comprehensive information about the theory of Banach ideal spaces and, in particular, rearrangement invariant spaces. An inexhaustible source of information about order continuity is Wnuk's monograph [\[Wnu99\]](#page-27-4). To place the mentioned structures within the general framework of abstract Banach lattices and Riesz spaces we recommend taking a look at [\[M-N91\]](#page-27-5).

<span id="page-6-0"></span>3.b. Calderón–Lozanovski $\mathbf{v}$  construction. Let us denote by  $\mathcal{U}$  the set of all nonnegative, concave and positively homogeneous functions  $\varrho: [0, \infty) \times [0, \infty) \to [0, \infty]$  which vanish only at  $(0, 0)$ . For a function  $\rho$  from U and two Banach ideal spaces X and Y, both defined on the same measure space, by the **Calderón–Lozanovski**<sup>†</sup> construction  $\rho(X, Y)$  (or just the **Calderón–Lozanovski** space) we understand a vector space consisting of all measurable functions, say  $f$ , such that

$$
|f(t)| \leq \lambda \varrho(|g(t)|, |h(t)|)
$$

for some  $\lambda > 0$  with  $g \in \text{Ball}(X)$  and  $h \in \text{Ball}(Y)$ ; the norm  $||f||_{\varrho(X,Y)}$  of a function f from  $\rho(X, Y)$  is defined as the infimum over all  $\lambda > 0$  for which the above inequality holds. It is straightforward to see that

$$
||f||_{\varrho(X,Y)} = \inf \max_{7} \{ ||g||_X, ||h||_Y \},
$$

where the infimum is taken over all  $g \in X$  and  $h \in Y$  with  $|f(t)| \leq \varrho(|g(t)|, |h(t)|)$ . This construction was introduced in the mid 60's by Alberto Calderón [\[Cal64\]](#page-25-6) and later systematically developed by Grigorii Lozanovskiı̆ in a series of papers (see, for example, [\[Loz71\]](#page-26-10), [\[Loz73\]](#page-26-11) and [\[Loz78\]](#page-26-12); cf. [\[KL10\]](#page-26-13) and [\[Mal04,](#page-26-14) Section 15]). It is inextricably linked to the interpolation theory, because  $\varrho(\cdot,\cdot)$  is an interpolation functor for positive<sup>\*</sup> linear operators. Although, we will be not interested in this general construction per se, its unifying character is hard to overestimate.

More precisely, when  $\varrho(s,t) = tF^{-1}(s/t)$  for  $s,t > 0$ , where  $F^{-1}$  is the right-continuous inverse<sup>[†](#page-7-1)</sup> of a Young function F, the corresponding Calderón–Lozanovski<sub>v</sub> space  $\varrho(X, L_{\infty})$ is usually denoted by  $X_F$  and is sometimes called the **generalized Orlicz space** (see [\[Mal04,](#page-26-14) Example 2, p. 178]). It is only a matter of simple calculations to be convinced that the space  $X_F$  consists of those measurable functions f such that  $F(|f(\cdot)|/\lambda)$  belongs to X for some  $\lambda > 0$  (see, for example, [\[Mal04,](#page-26-14) Example 2, p. 178]). Moreover,

<span id="page-7-2"></span>
$$
||f||_{\varrho(X,L_{\infty})} = ||f||_{X_F} := \inf \{ \lambda > 0 \colon \mathcal{M}_F(f/\lambda) \leq 1 \},
$$

where  $\mathcal{M}_F(f) \coloneqq \left\| F\left(|f| \right) \right\|_X$  is the **modular** (pedantically speaking,  $\mathcal{M}_F$  should be called the convex and left-continuous semi-modular). Here, we follow the convention that if  $F(|f|) \notin X$ , then  $\mathcal{M}_F(f) = \infty$ . Fortunately, for our purposes, the voluminous theory of modular spaces boils down to the following three simple relations:

(3.1) 
$$
\mathcal{M}_F(f) = 1 \implies ||f||_{X_F} = 1;
$$

<span id="page-7-5"></span>(3.2) 
$$
||f||_{X_F} \le 1 \iff \mathcal{M}_F(f) \le 1
$$
 (see [Mal04, Theorem 1.4(b), p. 9]);

<span id="page-7-6"></span>(3.3)  $||f||_{X_F} < 1 \implies \mathcal{M}_F(f) \le ||f||_{X_F}$  (see the proof of Theorem 3 in [\[DR00\]](#page-25-2)).

In general, none of the above implications can be reversed (for example, this can be done in [\(3.1\)](#page-7-2) if, and only if, the space  $X_F$  satisfies the so-called norm-modular condition; see [\[KL10,](#page-26-13) Remark 26] and references given there). It is known that the construction  $X_F$  inherits many<sup>[‡](#page-7-3)</sup> properties of the space X. For example, if the space X has the Fatou property or is rearrangement invariant then the same can be said about the space  $X_F$ . Note also that

<span id="page-7-4"></span>(3.4) 
$$
\psi_{X_F}(t) = \frac{1}{F^{-1}(1/\psi_X(t))}
$$

for  $t \in \mathbb{I}$  (cf. [\[Mal04,](#page-26-14) Corollary 4, p. 58]).

In particular, the space  $\varrho(L_1, L_\infty) = (L_1)_F$  coincides, up to the equality of norms, with the familiar **Orlicz space**  $L_F$ , whilst the space  $\varrho(\Lambda(w), L_\infty) = (\Lambda(w))_F$  coincides with the **Orlicz–Lorentz space**  $\Lambda_F(w)$  (see Section [6.b](#page-23-2) for more details). Evidently, if the

<span id="page-7-0"></span><sup>∗</sup>We can move from positive to arbitrary linear operators by paying the price of assuming the Fatou property.

<span id="page-7-1"></span><sup>&</sup>lt;sup>†</sup>Recall that for a given Young function F its right-continuous inverse  $F^{-1}$ :  $[0,\infty] \to [0,\infty]$  is defined as follows  $F^{-1}(s) := \inf\{t \geq 0 : F(t) > s\}$  for  $0 \leq s < \infty$  and  $F^{-1}(\infty) := \lim_{s \to \infty} F^{-1}(s)$  (with the convention that inf $\{\emptyset\} = \infty$ ).

<span id="page-7-3"></span><sup>&</sup>lt;sup>‡</sup>However, this is not always that obvious. For example, the space  $X_F$  is separable provided the space X is separable and F satisfies the appropriately understood  $\Delta_2$ -condition. Let alone some geometric properties like rotundity or uniform convexity (much work in this direction was done between 1990 and 2010; see [\[KL10\]](#page-26-13) and their references).

Young function F is just a power function, that is,  $F(t) = t^p$  for some  $1 \leq p < \infty$ , then the Orlicz space  $L_F$  is nothing else but the classical Lebesgue space  $L_p$ .

There are two more constructions that should be mentioned. In the case of power functions, that is, when  $\varrho(s,t) = s^{1-\theta}t^{\theta}$  for some  $0 \le \theta \le 1$ , the space  $\varrho(X,Y)$  coincide with the so-called **Calderón product**  $X^{1-\theta}Y^{\theta}$  (see [\[Cal64\]](#page-25-6) and [\[Mal89,](#page-26-15) p. 176]). In particular, for  $1 < p < \infty$ , the *p*-convexification  $X^{(p)}$  of X, is defined as

$$
X^{1/p}(L_{\infty})^{1-1/p} \equiv X^{(p)} \coloneqq \{f \in L_0 \colon |f|^p \in X\}
$$

with  $||f||_{X^{(p)}} \coloneqq |||f|^p||_X^{1/p}$  $X<sup>1/P</sup>$  (for more about *p*-convexification of Banach ideal spaces see, for example, [\[MP89\]](#page-26-0)). Let us add that the above construction  $X^{(p)}$  makes sense for  $0 < p < 1$ , but then it is usually referred to as the  $p$ -concavification.

Much more information about Calderón–Lozanovski˘ı spaces (but also modular spaces and, in particular, Orlicz spaces), their Banach space structure and connections with interpolation theory can be found in Maligranda's book [\[Mal89\]](#page-26-15) (see also [\[BM05\]](#page-25-7), [\[Mal04\]](#page-26-14), [\[KL10\]](#page-26-13) and references therein).

<span id="page-8-0"></span>3.c. Pointwise multipliers. By the space of pointwise multipliers  $M(X, Y)$  between two Banach ideal spaces X and Y we understand a vector space

$$
M(X,Y) \coloneqq \{ f \in L_0 \colon fg \in Y \text{ for all } g \in X \}
$$

furnished with the natural<sup>[∗](#page-8-1)</sup> operator norm  $||f||_{M(X,Y)} := \sup_{||g||_X=1} ||fg||_Y$ . Cooked in this fashion, the space  $M(X, Y)$  becomes a Banach ideal space itself (see [\[MP89,](#page-26-0) Proposition 2.). Note that the space  $M(X,Y)$  is non-trivial if, and only if,  $X \stackrel{locally}{\hookrightarrow} Y$ , that is, for any  $f \in X$ , and any set A with positive but finite measure, it follows that  $||f1_A||_Y \leq C ||f1_A||_X$  for some constant  $C > 0$  dependent only(!) on A (see [\[KLM12,](#page-26-2) Proposition 2.3]). Moreover, the space  $M(X, Y)$  is a rearrangement invariant provided X and Y are rearrangement invariant too (see [\[KLM12,](#page-26-2) Theorem 2.2] and [\[KT22,](#page-26-3) Lemma 4.3]). To put this construction on familiar ground, let us observe that the space  $M(X, L_1)$  coincide, up to the equality of norms, with the Köthe dual  $X^*$  of X. This leads, among other things, to a general variant of Hölder–Rogers inequality

(3.5) 
$$
||fg||_Y \le ||f||_X ||g||_{M(X,Y)}
$$

It is also straightforward to see that  $M(X, X) \equiv L_{\infty}$  (see [\[MP89,](#page-26-0) Theorem 1]). Informally, the space  $M(X, Y)$  may be regarded as a "pointwise quotient" of the space Y by X.

<span id="page-8-2"></span>.

Using this construction one can, for example, provide a characterization of compact and weakly compact multiplication operators acting between two Banach ideal spaces or describe compact Fourier multipliers acting on Banach spaces of analytic functions (see [\[KT22\]](#page-26-3) for more).

It is also worth mentioning that a lot has been said about pointwise multipliers acting on function spaces with some smoothness, like Besov–Sobolev–Triebel–Lizorkin' type spaces (see, for example, [\[Sic99\]](#page-27-6) and [\[Tri03\]](#page-27-7); classical book on this topic is [\[MS85\]](#page-27-8)).

There is an extensive literature devoted to this topic. We refer to [\[And60\]](#page-25-1), [\[Ben96\]](#page-25-8), [\[Ber23\]](#page-25-9), [\[BL93\]](#page-25-10), [\[CDSP08\]](#page-25-11), [\[Cro69\]](#page-25-12), [\[DSP10\]](#page-25-13), [\[DR00\]](#page-25-2), [\[KT22\]](#page-26-3), [\[KLM12\]](#page-26-2), [\[LT17\]](#page-26-4), [\[LT21\]](#page-26-6),

<span id="page-8-1"></span><sup>\*</sup>After all, every function  $f \in M(X, Y)$  induces the multiplication operator  $M_f: X \to Y$  given as  $M_f: g \leadsto fg$  and, moreover, there holds  $||f||_{M(X,Y)} = ||M_f: X \to Y||.$ 

[\[MN10\]](#page-26-1), [\[MP89\]](#page-26-0), [\[Nak95\]](#page-27-9), [\[Nak16\]](#page-27-10), [\[O'Ne65\]](#page-27-0), [\[OT72\]](#page-27-11), [\[Ray92\]](#page-27-12) and [\[Sch10\]](#page-27-13) (see also Nakai's survey [\[Nak17\]](#page-27-14) and references given there).

<span id="page-9-0"></span>3.d. **Pointwise products.** For two Banach ideal spaces  $X$  and  $Y$ , the **pointwise prod**uct space  $X \odot Y$  of X and Y is defined as

$$
X \odot Y \coloneqq \{gh \colon g \in X \text{ and } h \in Y\},\
$$

and endowed with the quasi<sup>\*</sup>-norm(!)

<span id="page-9-5"></span>
$$
||f||_{X \odot Y} := \inf \{ ||g||_X ||h||_Y : f = gh, g \in X \text{ and } h \in Y \}.
$$

The fact that  $X \odot Y$  is a vector space is not entirely obvious but follows, in one way or another, from the ideal property of  $X$  and  $Y$ . It seems noteworthy that if both spaces  $X$ and Y have the Fatou property, then the space  $X \odot Y$  has the Fatou property as well. Similarly, the space  $X \odot Y$  is rearrangement invariant as long as both spaces X and Y are rearrangement invariant. All this is essentially due to the fact that the product space  $X\odot Y$ is isometric to the  $\frac{1}{2}$ -concavification of the Calderón product  $X^{1/2}Y^{1/2}$ . Furthermore, for any two given rearrangement invariant spaces, say  $X$  and  $Y$ , the following nice formula holds

(3.6) 
$$
\psi_{X \odot Y}(t) = \psi_X(t)\psi_Y(t)
$$

for  $t \in \mathbb{I}$ . In other words, the fundamental function  $\psi_{X \odot Y}$  of the product space  $X \odot Y$  is just a product of fundamental function  $\psi_X$  of X and  $\psi_Y$  of Y (see [\[KLM12,](#page-26-2) Theorem 2]).

Plainly, taking pointwise product  $X \odot Y$  seems somehow opposite to taking "pointwise" quotient"  $M(X, Y)$ . For this reason, the problem whether  $X \odot M(X, Y)$  is the same as Y is not without significance. A generic example of this kind is **Lozanovski**<sup>'</sup>'s factorization **theorem**, which says that  $X \odot M(X, L_1) = L_1$  (we will come back to this problem in Section [5.c](#page-22-0) and say more about factorization).

More about pointwise products and factorization can be found in  $\left| \underline{\text{Ben96}} \right|$ ,  $\left| \underline{\text{Bun87}} \right|$ , [\[CS14\]](#page-25-15), [\[CS17\]](#page-25-16), [\[Gil81\]](#page-25-17), [\[JR76\]](#page-25-18), [\[KT24\]](#page-26-16), [\[KLM14\]](#page-26-5), [\[KLM19\]](#page-26-17), [\[LT17\]](#page-26-4), [\[LT21\]](#page-26-6), [\[LT-J80\]](#page-27-15), [\[Mau74\]](#page-27-1), [\[Nil85\]](#page-27-16), [\[Rei81\]](#page-27-17) and [\[Sch10\]](#page-27-13).

#### 4. Main results

<span id="page-9-2"></span><span id="page-9-1"></span>4.a. Generalized Young conjugate. This section essentially can be seen as a spin-off from [\[LT17\]](#page-26-4) (see also [\[And60\]](#page-25-1), [\[KLM12,](#page-26-2) Section 7], [\[Mal04,](#page-26-14) pp. 77–78], [\[MP89,](#page-26-0) pp. 334– 335] and [\[O'Ne65\]](#page-27-0)).

In what follows it will be convenient to write that a Young function  $F$  jumps to **infinity** if  $b_F := \sup\{t \ge 0: M(t) < \infty\}$  is finite. Otherwise, that is, when  $b_M = \infty$ , we will say that  $M$  is finite.

<span id="page-9-4"></span>Definition 4.a.1 (Generalized Young conjugate). For a given two Young functions, say F and G, we define the **generalized Young conjugate**  $G \ominus F$  of G with respect to F

<span id="page-9-3"></span><sup>∗</sup>The quasi-norm is, to put it briefly, just the norm in which the △-inequality holds but with a constant greater than 1 (see, for example, [\[Kal03\]](#page-26-18) and references therein for much more about quasi-norms and quasi-Banach spaces).

(in that order!), in the following way

<span id="page-10-1"></span>(4.1) 
$$
(G \ominus F)(t) := \sup_{0 \le s < b_F} \{G(st) - F(s)\}
$$

for  $t \geq 0$ . Moreover, for  $0 < a < b_F$ , we define the function  $G \ominus_a F$ , which can be seen as a truncated version of  $G \ominus F$ , as follows

<span id="page-10-2"></span>(4.2) 
$$
(G \ominus_a F)(t) := \sup_{0 \le s \le a} \{G(st) - F(s)\}
$$

for  $t \geqslant 0$ .

One can show that both functions  $G \ominus F$  and  $G \ominus_a F$  are again Young functions (see, for example,  $[And60, Theorem 4]$  and  $[LT17, Lemma 2]$ . Moreover, for  $t \ge 0$ , there holds

<span id="page-10-6"></span>(4.3) 
$$
\lim_{a \to b_F^-} (G \ominus_a F)(t) = (G \ominus F)(t).
$$

It is also clear that the so-called generalized Young inequality holds, that is, for any two Young functions  $F$  and  $G$ ,

(4.4) 
$$
G(st) \leqslant (G \ominus_a F)(t) + F(s),
$$

where  $t \geqslant 0$  and  $0 \leqslant s \leqslant a$ .

The idea behind [\(4.1\)](#page-10-1) and [\(4.2\)](#page-10-2) goes back to the work of Maurey [\[Mau74\]](#page-27-1). Roughly speaking, both constructions [\(4.1\)](#page-10-1) and [\(4.2\)](#page-10-2) are intended to properly generalize the Köthe duality theory of Orlicz spaces. To see this, recall that the Köthe dual  $(L_M)^{\times}$  of the Orlicz space  $L_F$  coincide, up to the equivalence of norms, with another Orlicz space  $L_{F^*}$ , where the function  $F^*$  is defined as  $F^*(t) := \sup_{s>0} \{st - F(s)\}\$ for  $t \geq 0$ , and is customarily called the Young conjugate of M (all of this is classic; see [\[Mal89,](#page-26-15) Chapters 8 and 9] and [\[O'Ne65,](#page-27-0) Definition 1.5]). In other words, we have

<span id="page-10-5"></span><span id="page-10-3"></span>
$$
(L_F)^{\times} = M(L_F, L_1) = L_{F^*} = L_{\text{id} \oplus F}.
$$

Thus, in general, it is perfectly natural to suspect that

$$
(4.5) \t\t M(L_F, L_G) = L_{G \ominus F}.
$$

In fact, after many partial results, this conjecture was finally confirmed in full generality by Karol Leśnik and the second-named author in [\[LT17\]](#page-26-4) (see also [\[LT21\]](#page-26-6) for further generalization to the setting of Musielak–Orlicz spaces).

Note also that  $(4.5)$  in its most rudimentary form, that is, when both Young functions M and N are just a power functions, looks as follows: For  $1 \leqslant q < p < \infty$ , we have  $M(L_p, L_q) = L_r$  with  $1/r = 1/q - 1/p$ .

<span id="page-10-0"></span>4.b. Pointwise multipliers of Calderón–Lozanovskiĭ spaces. Before we go any further, let us introduce some notation that will make our live a little more bearable.

<span id="page-10-4"></span>**Notation 4.b.1** (Nice triple). Let X be a rearrangement invariant function space. Further, let F and G be two Young functions. We will say that the triple  $(X, F, G)$  is nice if the following three conditions:

- (1) the fundamental function  $\psi_X$  of the space X does not vanish at zero;
- (2) the function  $F$  is finite;
- (3) the function  $G$  jump to infinity,

does not(!) meet simultaneously.

Remark 4.b.2. Unwinding the above definition, it is straightforward to see that the triple  $(X, F, G)$ , where X is a rearrangement invariant space X defined on  $\mathbb{I} = [0, 1]$  is nice if, and only if,  $X \neq L_{\infty}[0, 1].$ 

<span id="page-11-0"></span>After this modest preparation, we are finally ready to prove the following

**Theorem 4.b.3** (Pointwise multipliers of Calderón–Lozanovskiĭ spaces). Let X be a rearrangement invariant function space. Further, let F and G be two Young functions. Then  $M(X_F, X_G) = X_{G \ominus F}$  if, and only if, the triple  $(X, F, G)$  is "nice" (see Notation [4.b.1](#page-10-4) for clarification).

*Proof.* Proof of necessity. Suppose that the triple  $(X, F, G)$  fails to be nice. We have to prove that

<span id="page-11-1"></span>
$$
(A) \t M(X_F, X_G) \neq X_{G \ominus F}.
$$

Actually, a simple plan to justify  $(\clubsuit)$  is to show that the space  $X_{G\ominus F}$  is trivial and  $M(X_F, X_G)$  is not. The former is easy. Indeed, since the function G jump to infinity and F is finite, so the Young conjugate function  $G \oplus F$  is identically equal to infinity outside zero. This means that the space  $X_{G\ominus F}$  can only contain one function, namely, the one that is equal to zero almost everywhere. With this in mind, let us make one more observation. Since the fundamental function  $\psi_X$  of the space X does not vanish at zero, so it follows that  $X \hookrightarrow L_{\infty}$ . Thus, also  $X_F \hookrightarrow (L_{\infty})_F = L_{\infty}$ . We have

$$
||1_{[0,1]}||_{M(X_F, X_G)} = \sup\{||f1_{[0,1]}||_{X_G} : f \in \text{Ball}(X_F)\}\
$$
  
\$\leqslant ||X\_F \hookrightarrow L\_{\infty}|| \sup\{||f1\_{[0,1]}||\_{X\_G} : f \in \text{Ball}(L\_{\infty})\}\\$  
\$\leqslant ||X\_F \hookrightarrow L\_{\infty}|| ||1\_{[0,1]}||\_{X\_G}\$  
= ||X\_F \hookrightarrow L\_{\infty}|| [G^{-1} (1/\psi\_X(1))]^{-1},

where the last equality is due to [\(3.4\)](#page-7-4). This means, of course, that the space  $M(X_F, X_G)$ is non-trivial and  $\left(\bullet\right)$  follows.

**Proof of sufficiency.** Suppose that the triple  $(X, F, G)$  is nice. This is, unfortunately, the moment where things start to get a little more cumbersome, because formally we have five different situations to consider. However, be not of faint heart, we can divide the whole argument into just two cases, one of which is almost obvious.

**★** The case when  $b_F = \infty$ ,  $b_G < \infty$  and  $\psi_X(0) = 0$ . From what we said above, we already know that in this situation the space  $X_{G\ominus F}$  is trivial. Moreover, due to our assumptions,  $X_F \nleftrightarrow L_\infty$  and  $X_G \hookrightarrow L_\infty$ . But this means that the space  $M(X_F, X_G)$ is trivial as well. (Otherwise, we would have that  $||f1_{[0,1]}||_{L_{\infty}} \leq C ||f1_{[0,1]}||_{X_F}$ , which is clearly impossible, because the space  $X_F$  contains unbounded functions.) There is no doubt that  $0 = 0$ .

 $\star$  The remaining case. We can assume right away that  $(G \ominus F)(t)$  is finite for some  $t > 0$ , because the opposite is only possible when  $b_G < \infty$  and  $b_F = \infty$ . Therefore, the space  $X_{G\ominus F}$  is non-trivial. Moreover, without any loss of generality we can assume that

 $b_F \geq 1$ . Let us start with the embedding

<span id="page-12-0"></span>
$$
X_{G\ominus F}\hookrightarrow M(X_F,X_G).
$$

Take  $f \in X_{G \ominus F}$  with  $||f||_{X_{G \ominus F}} \leq 1/2$  and  $g \in X_F$  with  $||g||_{X_F} \leq 1/2$ . We have

$$
\mathcal{M}_G(fg) = ||G(|fg|)||_X
$$
  
\n
$$
\leq ||(G \ominus F)(|f|) + F(|g|)||_X \quad \text{(using (4.4))}
$$
  
\n
$$
\leq ||(G \ominus F)(|f|)||_X + ||F(|g|)||_X \quad \text{(by $\triangle$-inequality$)}
$$
  
\n
$$
\leq 1.
$$

Consequently, using [\(3.2\)](#page-7-5),  $fg \in X_G$  with  $||fg||_{X_G} \le 1$  and ( $\heartsuit$ ) follows. Now, let us focus on the opposite embedding

<span id="page-12-1"></span>
$$
(\diamondsuit) \t\t M(X_F, X_G) \hookrightarrow X_{G\ominus F}.
$$

Note, that in order to show  $(\diamondsuit)$ , it is enough to prove something a little easier, namely that there is a constant  $C > 0$  such that for all positive simple functions, say f, the following inequality

<span id="page-12-2"></span>(4.6) 
$$
||f||_{X_{G \ominus F}} \leqslant C ||f||_{M(X_F, X_G)}
$$

holds. Indeed, a straightforward argument based on the Fatou property of the space X will do the job. Observe, however, that we can make one more reduction and instead of proving [\(4.6\)](#page-12-2), we can only show that for any  $1 < a < b_F$  the following modular inequality

(4.7) MG⊖a<sup>F</sup> (f) 6 1 2

holds. To see this, let f be a positive simple function, that is,  $f = \sum_{n=1}^{N} a_n 1_{A_n}$ , where  $a_n$ 's are positive reals and  $A_n$ 's are sets of positive but finite measure. Without the loss of generality we can assume that

$$
(4.8) \t\t\t\t||f||_{M(X_F,X_G)} \leqslant \frac{1}{2\delta},
$$

where  $\delta = b_F$  provided F jumps to infinity or  $\delta = 1$  otherwise. Remembering about [\(4.3\)](#page-10-6) and using  $(4.7)$ , we have

<span id="page-12-6"></span><span id="page-12-5"></span><span id="page-12-4"></span><span id="page-12-3"></span>
$$
\mathcal{M}_{G \ominus F}(f) = ||(G \ominus F)(f)||_{X} = \liminf_{a \to b_{F}^{-}} ||(G \ominus_{a} F)(f)||_{X} \leq \frac{1}{2}.
$$

Thus, due to [\(3.2\)](#page-7-5), we have  $||f||_{X_{G\ominus F}} \le 1$  and [\(4.6\)](#page-12-2) follows with  $C = 2\delta$ . To recap, form this point on, we can focus all our efforts on showing  $(4.7)$ . Fix  $1 < a < b_F$ . It is easy to see that for any  $1 \leq n \leq N$  there is a positive real number  $b_n$  with

(4.9) 
$$
G(a_n b_n) = (G \ominus_a F)(a_n) + F(b_n).
$$

Set  $g := \sum_{n=1}^{N} b_n 1_{A_n}$ . Then, in view of [\(4.9\)](#page-12-4), (4.10)  $G(fq) = (G \ominus_a F)(f) + F(q).$ 

We claim that

<span id="page-12-7"></span>(4.11)  $||g||_{X_F} \leq 1.$ 

We will justify this in three steps.

**Step 1:** Divide and conquer. We are going to divide a support  $\mathbb{I}$  of the space X into a sequence  $\{\Delta_n\}_{n=1}^{\infty}$  of pairwise disjoint sets with positive but finite measure. The division algorithm will depend on two factors, namely, whether  $F$  is finite or not and whether  $\psi_X(0^+)$  is equal to or greater than zero. We will explain how to do this in the simplest situation, that is, when both F and G are finite and  $\psi_X(0^+) = 0$ . Later we will show how to modify this construction to work in other cases as well. Since  $\psi_{X_F}(0^+) = 0$ , so simple functions are order continuous in X. In consequence, there is  $\lambda > 0$  with  $||1_{\Delta}||_{X_F} \leq 1/a$ for all  $\Delta \subset \mathbb{I}$  with  $m(\Delta) \leq \lambda$ . Plainly, since the space X is rearrangement invariant, so  $\mathbb{I} = \bigcup_{n=1}^{\infty} \Delta_n$ , where  $\Delta_n \coloneqq [(n-1)\lambda, n\lambda] \cap \mathbb{I}$  for  $n \in \mathbb{N}$ . Moreover,  $||1_{\Delta_n}||_{X_F} \leq 1/a$  for all  $n \in \mathbb{N}$ . Now, suppose that  $b_F < \infty$  and  $\psi_X(0^+) \geq 0$ . Without any loss of the generality, we can assume that  $b_F > 1$  and  $\psi_X(0^+) \leq 1$ . Then  $\psi_{X_F}(0^+) \leq 1/b_F < 1$ . Thus, again, we can find  $\lambda > 0$  with  $||1_\Delta||_{X_F} \leq 1$  for all  $\Delta \subset \mathbb{I}$  with  $m(\Delta) \leq \lambda$ . The rest is already known. The algorithm is finished. Next, since  $|g(t)| \le a < b_F$ , so remembering about [\(4.8\)](#page-12-5), we get

<span id="page-13-0"></span>
$$
(4.12) \t\t\t||fg1_{\Delta_n}||_{X_G} \leq ||f||_{M(X_F,X_G)} ||g1_{\Delta_n}||_{X_F} \leq \frac{a}{2\delta} ||1_{\Delta_n}||_{X_F} \leq \frac{1}{2}.
$$

Thus, using  $(3.3)$  along with  $(4.10)$ ,

(4.13) 
$$
\mathcal{M}_G(g1_{\Delta_n}) \leq \mathcal{M}_G(fg1_{\Delta_n}) \leq \|fg1_{\Delta_n}\|_{X_G} \leq \frac{1}{2},
$$

where the last inequality follows from [\(4.12\)](#page-13-0). Finally, for  $n \in \mathbb{N}$ , set

<span id="page-13-3"></span><span id="page-13-1"></span>
$$
g_n := \sum_{i=1}^n g1_{\Delta_i}.
$$

## End of Step 1.

Step 2: Inductive argument. Now, we will show that

(4.14) M<sup>F</sup> (gn) 6 1 2

for all  $n \in \mathbb{N}$ . For  $n = 1$  this follows from [\(4.13\)](#page-13-1). Thus, let  $n \geq 2$  and assume that  $\mathcal{M}_F(g_{n-1}) \leqslant 1/2$ . Then

<span id="page-13-2"></span>
$$
\mathcal{M}_F(g_n) = \mathcal{M}_F(g_{n-1}) + \mathcal{M}_F(g_1 \Delta_n) \leq 1.
$$

In consequence, due to [\(3.2\)](#page-7-5),

$$
(4.15) \t\t\t\t \|g_n\|_{X_F} \leq 1.
$$

We have

$$
\mathcal{M}_F(g_n) \leq \mathcal{M}_G(fg_n) \qquad \text{(since, due to (4.10), } F(g) \leq M(fg))
$$
  
\n
$$
\leq \|fg_n\|_{X_G} \qquad \text{(in view of (3.3), (4.8) and (4.15))}
$$
  
\n
$$
\leq \|f\|_{M(X_F, X_G)} \|g_n\|_{X_F} \qquad \text{(by the Hölder-Rogers inequality (3.5))}
$$
  
\n
$$
\leq \frac{1}{2} \qquad \text{(by (4.8) together with (4.15))}.
$$

In other words,  $(4.14)$  follows. **End of Step 2.** 

**Step 3: Limit argument.** We know, thanks to  $(4.14)$ , that  $\mathcal{M}_F(g_n) \leq 1/2$  for all  $n \in \mathbb{N}$ . Thus, invoking the Fatou property, we get

(4.16) 
$$
||g||_{X_F} = \sup{||g_n||_{X_F} : n \in \mathbb{N}} \le 1.
$$

But this is exactly what we wanted: [\(4.11\)](#page-12-7) follows. End of Step 3.

Having  $(4.11)$  in hand, we can finally finish the whole proof by showing  $(4.7)$ . This is how it goes

<span id="page-14-1"></span>
$$
\mathcal{M}_{G\ominus_a F}(f) \leq \mathcal{M}_G(fg) \qquad \text{(since } (G\ominus_a F)(f) \leq G(fg))
$$
  
\n
$$
\leq \|fg\|_{X_G} \qquad \text{(using (3.3), (4.8) and (4.16))}
$$
  
\n
$$
\leq \|f\|_{M(X_F, X_G)} \|g\|_{X_F} \qquad \text{(by the Hölder-Rogers inequality (3.5))}
$$
  
\n
$$
\leq \frac{1}{2} \qquad \text{(using (4.8) and (4.11))}.
$$

<span id="page-14-2"></span>The proof has been completed.

**Remark 4.b.4** (What if the triple  $(X, F, G)$  is not nice?). Note that it follows from the proof of Theorem [4.b.3](#page-11-0) that if the triple  $(X, F, G)$  fails to be nice, then although  $M(X_F, X_G) \neq X_{G \ominus F}$ , the space  $M(X_F, X_G)$  is nevertheless non-trivial. The question that certainly looms on the horizon is: How to describe the space  $M(X_F, X_G)$ ? In the case when X is defined on the unit interval  $\mathbb{I} = [0, 1]$ , things are rather straightforward. In fact, since the fundamental function  $\psi_X$  does not vanish at zero, so  $X = L_\infty[0, 1]$  and

$$
M(X_F, X_G) \equiv M((L_{\infty}[0, 1])_F, (L_{\infty}[0, 1])_G)
$$
  
=  $M(L_{\infty}[0, 1], L_{\infty}[0, 1])$   
 $\equiv L_{\infty}[0, 1].$ 

However, the remaining case, when X is defined on the half-line  $\mathbb{I} = [0, \infty)$ , is definitely less obvious.

<span id="page-14-0"></span>The next result clarifies this situation.

**Theorem 4.b.5.** Let X be a rearrangement invariant space. Further, let F and G be two Young functions. Suppose that either the triple  $(X, F, G)$  is not "nice" (see Notation [4.b.1](#page-10-4)) for clarification) or the space X is a sequence space. Then  $M(X_F, X_G) = X_{G \ominus_1 F}$ .

Proof. To facilitate the impending maneuvers, without any loss of generality, we may assume that

- $F(1) = G(1) = 1;$
- $b_G > 1$  and  $\psi_X(0) = 1$  provided X is a function space;
- $\psi_X(1) = 1$  provided X is a sequence space.

Moreover, by Remark [4.b.4,](#page-14-2) we can ignore the case when the space X is defined on  $\mathbb{I} = [0, 1]$ . Now, we will consider both embeddings separately.

Embedding  $X_{G\ominus_1F} \hookrightarrow M(X_F, X_G)$ . Take  $f \in X_{G\ominus_1F}$  with  $||f||_{X_{G\ominus_1F}} \leq 1/2$  and  $g \in X_F$  with  $||g||_{X_F} \leq 1/2$ . Our plan is to show that the product  $fg$  belongs to  $X_G$  with  $||fg||_{X_G} \leq 1$ . Recall the following variant of Young's inequality

<span id="page-14-3"></span>
$$
(4.17) \tG(st) \leq (G \ominus_1 F)(s) + F(t),
$$

where  $0 \le s, t \le 1$ , follows directly from the definition of the function  $G \ominus_1 F$  (see Definition [4.a.1](#page-9-4) and [\(4.4\)](#page-10-5)). Moreover, in view of our assumptions,  $||X \hookrightarrow L_{\infty}|| = 1$ , so  $||f||_{L_{\infty}} \leq 1$  and  $||g||_{L_{\infty}} \leq 1$ . We have

$$
\mathcal{M}_G(fg) = ||G(|fg|)||_X
$$
  
\n
$$
\leq ||(G \ominus_1 F)(|f|) + F(|g|)||_X \quad \text{(using (4.17))}
$$
  
\n
$$
\leq ||(G \ominus_1 F)(|f|)||_X + ||F(|g|)||_X \quad \text{(via $\triangle$-inequality)}
$$
  
\n
$$
\leq \frac{1}{2} + \frac{1}{2}.
$$

Here, the last inequality is due to the well-known relation between the norm and the modular [\(3.3\)](#page-7-6). This means that indeed  $fg \in X_G$  with  $||fg||_{X_G} \leq 1$ . That's all we wanted for now.

**Embedding**  $M(X_F, X_G) \hookrightarrow X_{G\ominus_1 F}$ . Note that we only need to show that the following inequality

<span id="page-15-1"></span>
$$
\left(\spadesuit\right) \qquad \qquad \left\|f\right\|_{X_{G\ominus_1 F}} \leqslant C \left\|f\right\|_{M(X_F, X_G)}
$$

holds for some constant  $C > 0$  and all positive, simple functions f. Then, using the Fatou property, we can lift this inequality to the whole space  $M(X_F, X_G)$ . Keeping this in mind, let f be a positive, simple function with  $||f||_{M(X_F,X_G)} \leq \frac{1}{2}$  $\frac{1}{2}$ . Plainly, there is a sequence  ${a_n}_{n=1}^N$  of positive reals together with a sequence  ${A_n}_{n=1}^\infty$  of sets with positive but finite measure, such that  $f = \sum_{n=1}^{N} a_n 1_{A_n}$ . Since  $M(X_F, L_\infty) \equiv L_\infty$  and, due to our assumptions,  $||M(X_F, X_G) \hookrightarrow M(X_F, L_\infty)|| \leq 1$ , so  $||f||_{L_\infty} \leq 1$ . Moreover, using the standard compactness argument, for every  $n \in \mathbb{N}$ , we can find  $0 \leq b_k \leq 1$  with

<span id="page-15-0"></span>
$$
G(a_n b_n) = (G \ominus_1 F)(a_n) + F(b_n).
$$

Set  $g \coloneqq \sum_{n=1}^{\infty} b_n 1_{A_n}$ . We claim that

$$
(4.18) \t\t\t ||g||_{X_F} \leq 1.
$$

To see this, we will divide the measure space underlying X into a sequence  $\{\Delta_n\}_{n=1}^{\infty}$  of pairwise disjoint sets with positive and finite measure. Let us explain how to do this.

 $\star$  The situation in which X is a sequence space is the most straightforward. Just take  $\Delta_n$  as the n<sup>th</sup> atom, that is,  $\Delta_n = \{n\}$  for  $n \in \mathbb{N}$ . Then, making use of [\(3.4\)](#page-7-4) along with the fact that the space  $X$  is rearrangement invariant, we infer that

$$
||1_{\{n\}}||_{X_F} = ||1_{\{1\}}||_{X_F} = [F^{-1}(1/\psi_X(1))]^{-1} = [F^{-1}(1)]^{-1} \leq 1.
$$

**★** Next, suppose that X is a function space. It is enough to take  $\Delta_n = [n-1, n]$  for  $n \in \mathbb{N}$ . Then, exactly as above,

$$
||1_{[n-1,n]}||_{X_F} = ||1_{[0,1]}||_{X_F} = [F^{-1}(1/\psi_X(1))]^{-1} \leq [F^{-1}(1)]^{-1} \leq 1.
$$

All this is, of course, very similar to the proof of Theorem [4.b.3](#page-11-0) (see "Step 1"). For this reason, to obtain [\(4.18\)](#page-15-0), it is enough to mimic the inductive argument in "Step 2" contained therein. However, since repeating all this here may seem rather boring, we leave easy-to-fill details for the inquisitive reader.

Going ahead, we have

$$
\mathcal{M}_{G\ominus_{1}F}(f) \leq \mathcal{M}_{G}(fg) \qquad \text{(since } (G\ominus_{1}F)(f) \leq G(fg))
$$
  
\n
$$
\leq \|fg\|_{X_{G}} \qquad \text{(using (3.3))}
$$
  
\n
$$
\leq \|f\|_{M(X_{F},X_{G})}\|g\|_{X_{F}} \qquad \text{(by the Hölder-Rogers inequality (3.5))}
$$
  
\n
$$
\leq \frac{1}{2} \qquad \text{(in view of (4.18))}.
$$

But this means that  $(\bullet)$  holds. In consequence, the proof has been completed.

It seems quite instructive to support the above results with some concrete calculations. This is by no means difficult, but rather tedious. Recall that the  $p^{\text{th}}$ -power function  $F_p: [0, \infty) \to [0, \infty)$  with  $1 \leqslant p < \infty$  is defined in the following way

$$
F_p(t) \coloneqq \frac{1}{p} t^p.
$$

<span id="page-16-0"></span>Example 4.b.6. Let  $1 \leq p, q < \infty$  with  $1 \leq p < q < \infty$  and  $1/r = 1/p - 1/q$ . Our goal is to compute

$$
(F_p \ominus F_q)(t) \coloneqq \sup_{s>0} \left\{ \frac{1}{p} (st)^p - \frac{1}{q} s^q \right\}.
$$

Some elementary calculations show us that for fixed  $t > 0$  the extreme value of the function

$$
f(s,t) = \frac{1}{p}(st)^p - \frac{1}{q}s^q
$$

is attained at the point  $s_{\text{ext}} := t^{p/(q-p)}$ . In consequence, for  $t \geq 0$ , we have

$$
(F_p \ominus F_q)(t) = \frac{1}{p}(s_{ext}t)^p - \frac{1}{q}s_{ext}^q
$$
  
=  $\frac{1}{p}(t^{p/(q-p)}t)^p - \frac{1}{q}(t^{p/(q-p)})^q$   
=  $\frac{1}{p}(t^{(p+q-p)/(q-p)})^p - \frac{1}{q}(t^{pq/(q-p)})$   
=  $\frac{1}{p}t^r - \frac{1}{q}t^r$   
=  $\frac{1}{r}t^r$   
=  $F_r(t)$ .

<span id="page-16-1"></span>**Example 4.b.7.** Let  $0 < b < \infty$  and  $1 \leqslant p, q < \infty$ . Now, let us modify the function  $F_p$ in the following way

$$
F_{p,b}(t) := \begin{cases} F_p(t) & \text{if } 0 \leqslant t \leqslant b \\ \infty & \text{if } t > b. \end{cases}
$$

It is easy to see that  $(F_{p,b} \ominus F_q)(t) = \infty$  for all  $t > 0$ . Thus, as Theorem [4.b.5](#page-14-0) teaches us, to obtain a non-trivial Young's function we should instead consider the function  $F_{p,b} \ominus_1 F_q$ .

Let us start with the case  $1 \leqslant p < q \leqslant \infty$ . The same calculations as in Example [4.b.6](#page-16-0) shows that for  $0 \leq t \leq \min\{1, b\}$  and  $1/r = 1/p - 1/q$ , we have

$$
(F_{p,b} \ominus_1 F_q)(t) = F_r(t).
$$

Furthermore,  $(F_{p,b} \ominus_1 F_q)(t) = \infty$  for  $t > b$ . On the other hand, for  $1 < t \leq b$ , we have

$$
(F_{p,b} \ominus_1 F_q)(t) = \sup_{0 \le s \le 1} \left\{ \frac{1}{p} (st)^p - \frac{1}{q} s^q \right\}.
$$

Again, due to Example [4.b.6,](#page-16-0) we already know that the extreme value of the function  $f(s,t) = (st)^p/p - s^q/q$  with fixed  $t > 0$  is attained in the point  $s_{\text{ext}} = t^{p/(q-p)} > 1$ , that is, outside the range of the parameter s. Since the function  $s \mapsto f(s,t)$  is increasing, so the supremum is attained at the end of the interval [0, 1]. Consequently,

$$
(F_{p,b} \ominus_1 F_q)(t) = \frac{1}{p}t^p - \frac{1}{q}.
$$

In summary,

$$
(F_{p,b} \ominus_1 F_q)(t) = \begin{cases} F_r(t) & \text{for} \quad 0 \leq t \leq \min\{1, b\} \\ \frac{1}{p}t^p - \frac{1}{q} & \text{for} \quad 1 < t \leq b \\ \infty & \text{for} \quad t > b. \end{cases}
$$

Finally, let us consider one more situation when  $1 \leqslant q \leqslant p < \infty$ . In this case  $(st)^p/p \leqslant$  $s^q/q$  for  $t > 0$  and  $0 \le s \le 1$ , so

<span id="page-17-2"></span>
$$
(F_{p,b} \ominus_1 F_q)(t) = \begin{cases} 0 & \text{for } & 0 \leq t \leq b \\ \infty & \text{for } & t > b. \end{cases}
$$

One of the immediate conclusions from the above result is the following example, first noted by Maligranda and Persson in [\[MP89,](#page-26-0) Corollary 2].

<span id="page-17-0"></span>**Example 4.b.8** (Maligranda and Persson, 1989). Let X be a rearrangement invariant space. Further, let  $1 \leq q < p < \infty$  with  $1/r = 1/q - 1/p$ . We claim that

(4.19) 
$$
M(X^{(p)}, X^{(q)}) \equiv X^{(r)}.
$$

To see this, note that if  $F_p(t) = t^p/p$  and  $F_q(t) = t^q/q$ , then the Calderón–Lozanovskii constructions  $X_{F_p}$  and  $X_{F_q}$  coincide with the p-convexification  $X^{(p)}$  of X and the qconvexification  $X^{(q)}$  of X, respectively. Moreover, it is crystal clear that the triple  $(X, F_p, F_q)$  is nice. Thus, after realizing that  $(F_q \ominus F_p)(t) = t^r/r$  (just in case, Example [4.b.6](#page-16-0) may be helpful), it is enough to call Theorem [4.b.3](#page-11-0) on stage. In particular, if  $X = L_1$ , the formula [\(4.19\)](#page-17-2) reduces to the well-known fact that  $M(L_p, L_q) \equiv L_r$ .

<span id="page-17-1"></span>**Example 4.b.9.** Let X be a rearrangement invariant space with  $X \hookrightarrow L_{\infty}$ . Further, let  $1 \leqslant q < p < \infty$  with  $1/r = 1/q - 1/p$  and  $b = 1$ . It follows from Theorem [4.b.5](#page-14-0) and Example [4.b.7](#page-16-1) that

$$
M(X_{F_{p,b}}, X_{F_q}) = X_{F_{r,b}}.
$$

In particular, for  $X = L_1 \cap L_\infty$ , we get

$$
M(L_p \cap L_{\infty}, L_q \cap L_{\infty}) = L_r \cap L_{\infty}.
$$

#### 5. Applications

<span id="page-18-0"></span>In this section we will show how, form the description of the space of pointwise multipliers between Calderón–Lozanovski˘ı spaces (see Theorem [4.b.3\)](#page-11-0), one can deduce the factorization of these spaces.

<span id="page-18-1"></span>5.a. A bird's eye view of factorization. Recall that classical Lozanovskii's factorization [\[Loz69,](#page-26-19) Theorem 6] (see also [\[Gil81\]](#page-25-17), [\[JR76\]](#page-25-18), [\[Mal04,](#page-26-14) Example 6, p. 185] and [\[Rei81\]](#page-27-17)) teaches us that for any  $\varepsilon > 0$  each function f from  $L_1$  can be written as a pointwise product of two functions, say g and h, one from X and the other from the Köthe dual  $X^*$  of X, in such a way that

$$
||f||_{L_1} \leq ||g||_X ||h||_{X^{\times}} \leq (1+\varepsilon) ||f||_{L_1}.
$$

Furthermore, knowing that the space X posses the Fatou property, we can set  $\varepsilon = 0$ . In other words,  $L_1$  can be factorized through X, that is,

$$
X \odot X^{\times} = X \odot M(X, L_1) = L_1.
$$

Clearly, one can replace  $L_1$  from Lozanovskii's factorization by an arbitrary Banach ideal space Y and ask whether Y can be factorized through  $X$ ? It turns out that an answer to this question is in general very difficult and the equality  $X \odot M(X, Y) = Y$ does not holds without some extra assumptions on  $X$  and  $Y$  (see, for example, [\[KLM14,](#page-26-5) Example 2 and  $\vert$ KT22, Example A.2..

The topic of factorization of Banach ideal spaces seems to be very much in vouge recently; see, for example, the papers [\[DR00\]](#page-25-2) and [\[LT17\]](#page-26-4) for Orlicz spaces; [\[CS17\]](#page-25-16), [\[KLM14\]](#page-26-5) and  $[Rei81]$  for Lorentz and Marcinkiewicz spaces;  $|KLM12|$  for Calderón–Lozanovski spaces; [\[KT22\]](#page-26-3), [\[KLM19\]](#page-26-17) and [\[Sch10\]](#page-27-13) for Cesáro and Tandori spaces; and [\[LT21\]](#page-26-6) for Musielak–Orlicz spaces.

Nevertheless, factorization has much more to offer that just a simple analogy with the Lozanovski˘ı theorem. For example, using some factorization techniques, Nilsson was able to gave another proof of Pisier's result (see [\[Nil85,](#page-27-16) Theorem 2.4]). Moreover, Odell and Schlumprecht's proof that  $\ell_2$  is arbitrarily distortable makes a use of Lozanovskii's factorization (see [\[OS94,](#page-27-18) p. 261]). There is also a beautiful connection between complex interpolation, Lozanovski<sup>†</sup>'s factorization and the construction of twisted sums of Banach spaces (see [\[CS14\]](#page-25-15), [\[Cor22\]](#page-25-19), [\[CS17\]](#page-25-16), [\[CGF17\]](#page-25-20), [\[Kal92\]](#page-26-20) and references therein).

There is, however, a lot of life outside of the "ideal" world of Banach ideal spaces. In the realm of harmonic analysis, for example, (weak) factorization is a domesticated and powerful technique. Some remarkable factorization results for Bergman spaces, Hardy spaces and tent spaces, can be found in [\[CRW76\]](#page-25-21), [\[CV00\]](#page-25-22), [\[Hor77\]](#page-25-23), [\[JR76\]](#page-25-18) and [\[PZ15\]](#page-27-19).

<span id="page-18-2"></span>5.b. Products of Calderón–Lozanovskii spaces. The next step, en route to our factorization results, goes through a series of rather technical lemmas. But first, some inevitable notation.

<span id="page-18-3"></span>**Notation 5.b.1** (Vinogradov's notation). For two given quantities, say A and B, depending (maybe) on certain parameters, we will write  $A \preccurlyeq B$  understanding that there exists an absolute constant  $C > 0$  (that is, independent of all involved parameters) such that  $A \leqslant CB$ . Moreover, we will write  $A \approx B$  meaning that  $A \preccurlyeq B$  and  $B \preccurlyeq A$ .

**Notation 5.b.2** (Small/Large/All arguments). Let  $F, G: [0, \infty) \to [0, \infty)$  be two realvalued function. Further, let the symbol  $\Box$  denote either the relation  $\preccurlyeq$  or  $\approx$  (see Notation [5.b.1\)](#page-18-3). We will say that

- F $\Box G$  holds for small arguments if there is  $T > 0$  such that  $F(t) \Box G(t)$  for all  $0 \leqslant t \leqslant T$ ;
- $F \Box G$  holds for **large arguments** if there is  $T > 0$  such that  $F(t) \Box G(t)$  for all  $t \geqslant T$ ;
- $F \Box G$  holds for all arguments if  $F(t) \Box G(t)$  for all  $t \geq 0$ .

**Remark 5.b.3.** Plainly, if both F and G are Young functions then  $F \square G$  holds for all arguments if, and only if,  $F\square G$  holds for small and large arguments.

Let us get to the point. The first of the lemmas we need is well-known. (However, many partial results in this direction are scattered throughout the literature; see, for example, [\[And60,](#page-25-1) Theorem 1], [\[Dan74,](#page-25-3) pp. 63–68], [\[KR61,](#page-26-21) Theorems 13.7 and 13.8], [\[Mal04,](#page-26-14) pp. 69– 75], [\[O'Ne65,](#page-27-0) Section VI] and [\[ZR67,](#page-27-2) Theorem 8].)

<span id="page-19-3"></span>**Lemma 5.b.4.** Let X be a Banach ideal space with the Fatou property. Further, let  $F$ ,  $G$ and H be three Young functions. Suppose that one of the following three conditions holds:

- (1)  $F^{-1}G^{-1} \approx H^{-1}$  for all arguments and neither  $L_{\infty} \hookrightarrow X$  nor  $X \hookrightarrow L_{\infty}$ ;
- (2)  $F^{-1}G^{-1} \approx H^{-1}$  for large arguments and  $L_{\infty} \hookrightarrow X$ ;
- (3)  $F^{-1}G^{-1} \approx H^{-1}$  for small arguments and  $X \hookrightarrow L_{\infty}$ .

Then  $X_F \odot X_G = X_H$ .

*Proof.* This is just Theorems 4.1, 4.2 and 4.5 from [\[KLM12\]](#page-26-2) in tandem with Theorem  $5(a)$ from [\[KLM14\]](#page-26-5) (see also [\[KLM14,](#page-26-5) Theorem A(a)]; cf. [\[Mal04,](#page-26-14) Theorem 10.1, p. 69]).  $\blacksquare$ 

<span id="page-19-2"></span>The *crux* of the entire route is proof of the following

**Lemma 5.b.5.** Let  $X$  be a rearrangement invariant function space. Further, let  $F$ ,  $G$ and H be three Young functions. Suppose that  $X_H \hookrightarrow X_F \odot X_G$ . Then

- (1)  $H^{-1} \preccurlyeq F^{-1}G^{-1}$  for small arguments provided  $L_{\infty} \not\hookrightarrow X$ ;
- (2)  $H^{-1} \preccurlyeq F^{-1}G^{-1}$  for large arguments provided  $X \not\hookrightarrow L_{\infty}$ ;
- (3)  $H^{-1} \preccurlyeq F^{-1}G^{-1}$  for all arguments provided neither  $L_{\infty} \hookrightarrow X$  nor  $X \hookrightarrow L_{\infty}$ .

*Proof.* We will only show  $(1)$  in details. Once this is done, the proof of  $(2)$  is completely analogous. Moreover, the proof of (3) is just a simple combination of (1) and (2).

Suppose that  $L_{\infty} \nleftrightarrow X$  and the condition  $H^{-1} \preccurlyeq F^{-1}G^{-1}$  does not holds for small arguments. This means that there is a decreasing null sequence  $\{a_n\}_{n=1}^{\infty}$  of reals with

<span id="page-19-0"></span>(5.1) 
$$
0 < a_n < \min\left\{1, (F^{-1})^{-1}(b_F)\right\}
$$

for all  $n \in \mathbb{N}$ , where  $(F^{-1})^{-1}(b_F) = \inf\{t > 0 : F^{-1}(t) = b_F\}$ , and

<span id="page-19-1"></span>(5.2) 
$$
2^{n} F^{-1}(a_{n}) G^{-1}(a_{n}) \leq H^{-1}(a_{n})
$$

for all  $n \in \mathbb{N}$ . Keeping this in mind, we claim that there is a sequence  $\{A_n\}_{n=1}^{\infty}$  of sets of positive but finite measure such that  $||a_n1_{A_n}||_X = 1$ . To see this, just note that the function  $\psi_X$  is continuous and unbounded. (Otherwise,  $L_{\infty} \hookrightarrow X$ , which is obviously not the case.) Thus, thanks to the Darboux property of continuous functions, we can effortlessly find  $A_n$ 's with  $||a_n1_{A_n}||_X = 1$  for  $n \in \mathbb{N}$ . Our claim follows. Knowing this, for  $n \in \mathbb{N}$ , we set

$$
f_n := F^{-1}(a_n)1_{A_n}, \quad g_n := G^{-1}(a_n)1_{A_n} \quad \text{and} \quad h_n := f_n g_n.
$$

Now, for  $0 < \lambda < 1$ , we have

$$
\mathcal{M}_F(f_n/\lambda) = ||F[\lambda^{-1}F^{-1}(a_n)]1_{A_n}||_X
$$
  
\n
$$
\geq \lambda^{-1} ||(F \circ F^{-1})(a_n)1_{A_n}||_X \qquad \text{(due to the convexity of } F)
$$
  
\n
$$
= \lambda^{-1} ||a_n 1_{A_n}||_X \qquad \text{(thanks to (5.1), } (F \circ F^{-1})(a_n) = a_n)
$$
  
\n
$$
> 1.
$$

This means that  $||f_n||_{X_F} \geq 1$ . By reasoning in exactly the same way, we can show that also  $||g_n||_{X_F} \geq 1$ . In consequence, we have

$$
||h_n||_{X_F \odot X_G} = F^{-1}(a_n) G^{-1}(a_n) \psi_{X_F \odot X_G}(m(A_n))
$$
  
=  $F^{-1}(a_n) \psi_{X_F}(m(A_n)) G^{-1}(a_n) \psi_{X_G}(m(A_n))$  (by (3.6))  
=  $||f_n||_{X_F} ||g_n||_{X_F} \ge 1$ .

Thus  $||h_n||_{X_F \odot X_G}$  is "big". Now we just need to show that  $||h_n||_{X_H}$  can be arbitrarily "small". We have

$$
\mathcal{M}_H(2^n h_n) = \|H\left[2^n F^{-1}(a_n) G^{-1}(a_n)\right] 1_{A_n}\|_X
$$
  
\n
$$
\leq \| (H \circ H^{-1})(a_n) 1_{A_n}\|_X \qquad \text{(in view of (5.2))}
$$
  
\n
$$
\leq \|a_n 1_{A_n}\|_X \qquad \text{(see [O'Ne65, Property 1.3])}
$$
  
\n
$$
\leq 1.
$$

In consequence,  $||h_n||_{X_H} \leq 1/2^n$ . Putting this two facts together,

$$
2^n\left\|h_n\right\|_{X_H}\leqslant \|h_n\|_{X_F\odot X_G}
$$

for  $n \in \mathbb{N}$ , so  $X_H \not\hookrightarrow X_F \odot X_G$ .

The last lemma is quite interesting. This is almost literally [\[KLM14,](#page-26-5) Theorem 5(d)]. However, a very slight change in the proof resulted in drastic improvement<sup>\*</sup> of the result.

**Lemma 5.b.6.** Let  $X$  be a rearrangement invariant sequence space. Further, let  $F$ ,  $G$ and H be three Young functions. Suppose that  $\ell_{\infty} \nleftrightarrow X$  and  $X_H \hookrightarrow X_F \odot X_G$ . Then  $H^{-1} \preccurlyeq F^{-1}G^{-1}$  for small arguments.

*Proof.* Suppose that  $\ell_{\infty} \nleftrightarrow X$  and  $H^{-1} \preccurlyeq F^{-1}G^{-1}$  does not hold for small arguments. This means that there is a null sequence  $\{a_n\}_{n=1}^{\infty}$  of positive integers such that

$$
2^{n} F^{-1}(a_{n}) G^{-1}(a_{n}) \leqslant H^{-1}(u_{n})
$$

<span id="page-20-0"></span><sup>\*</sup>Strictly speaking, instead of assuming that the space  $X$  is separable, which rules out a lot of spaces like, for example, Marcinkiewicz sequence spaces  $m_{\psi}$  or Orlicz sequence spaces  $\ell_F$  without the  $\Delta_{2}$ condition, we merely assume that X is not  $\ell_{\infty}$ .

for all  $n \in \mathbb{N}$ . Since  $\ell_{\infty} \not\hookrightarrow X$ , so  $\lim_{n\to\infty} ||\sum_{i=1}^{n} e_n||_X = \infty$ . (Note also that the Fatou property is important here, because otherwise  $c_0$  is a simple counter-example.) In consequence, for every  $n \in \mathbb{N}$ , there exists  $M(n) \in \mathbb{N}$  such that

(5.3) 
$$
a_n \left\| \sum_{i=1}^{M(n)} e_i \right\|_X \leq 1 < a_n \left\| \sum_{i=1}^{M(n)+1} e_i \right\|_X.
$$

Since any fundamental function is quasi-concave, so  $\psi_X$  is increasing, while

<span id="page-21-1"></span><span id="page-21-0"></span>
$$
n \mapsto \frac{\psi_X(n)}{n} = \frac{\left\| \sum_{i=1}^n e_i \right\|_X}{n}
$$

is non-increasing (see [\[BS88,](#page-25-4) Corollary 5.3, p. 67]). In consequence, for  $n \in \mathbb{N}$ ,

(5.4) 
$$
1 \geqslant \frac{\psi_X(n)}{\psi_X(n+1)} \geqslant \frac{n}{n+1} \geqslant \frac{1}{2}.
$$

Thus, combining  $(5.3)$  with  $(5.4)$ , we see that

(5.5) 
$$
\frac{1}{2} \leq a_n \left\| \sum_{i=1}^{M(n)} e_i \right\|_X \leq 1.
$$

From this point on, the rest of the argument is the same to the letter. However, instead of referring to Lemma  $5.b.5$  (or [\[KLM14,](#page-26-5) Theorem  $5(d)$ ]), let us finish what we started. Set

<span id="page-21-2"></span>
$$
x_n := F^{-1}(a_n) \sum_{i=1}^{M(n)} e_i
$$
,  $y_n := G^{-1}(a_n) \sum_{i=1}^{M(n)} e_i$  and  $z_n := x_n y_n$ .

Using  $(5.5)$ , we have

$$
\mathcal{M}_F(x_n) \leq a_n \left\| \sum_{i=1}^{M(n)} e_i \right\|_X \leq 1
$$

and

$$
\mathcal{M}_G(3x_n) = F\left[3F^{-1}(a_n)\right] \left\| \sum_{i=1}^{M(n)} e_i \right\|_X \ge 3a_n \left\| \sum_{i=1}^{M(n)} e_i \right\|_X > 1.
$$

In other words,  $1/3 \leq ||x_n||_{X_F} \leq 1$  for  $n \in \mathbb{N}$ . Repeating the same argument for  $y_n$ 's in place of  $x_n$ 's we get  $1/3 \leqslant ||y_n||_{X_G} \leqslant 1$  for  $n \in \mathbb{N}$ . Now, it is easy to see that  $||z_n||_{X_F \odot X_G} \geq 1/9$  and  $||z_n||_{X_H} \leq 1/2^n$  for  $n \in \mathbb{N}$ . Thus, for  $n \in \mathbb{N}$ ,

$$
2^n\left\|z_n\right\|_{X_H}\leqslant 9\left\|z_n\right\|_{X_F\odot X_G}
$$

and  $X_H \nleftrightarrow X_F \odot X_G$ .

<span id="page-22-0"></span>5.c. Factorization of Calderón–Lozanovskiĭ spaces. Before the *grand finale*, we need one more definition.

For a given two Banach ideal spaces X and Y, we will say that X is Y-perfect if  $M(M(X, Y), Y) = X$ . Incidentally, L<sub>1</sub>-perfectness of X is exactly the Fatou property under disguise. To see this, it is enough to compare the equality

$$
M(M(X, L_1), L_1) = M(X^{\times}, L_1) = X^{\times \times}
$$

with the well-known fact that  $X^{\times} = X$  if, and only if, the space X has the Fatou property. Moreover, note that if  $X$  has the Fatou property and  $Y$  factorizes through X then X is Y-perfect. Indeed, using the so-called "cancellation" property (see, for example, [\[KLM14,](#page-26-5) Theorem 4]), we infer that

$$
M(M(X,Y),Y) = M(L_{\infty} \odot M(X,Y), X \odot M(X,Y)) = M(L_{\infty},X) = X.
$$

Alas, in general, there is no hope for the reverse implication to hold (in fact, there is a three-dimensional counter-example by Bolobás and Brightwell; see [\[Sch10,](#page-27-13) Example 3.6] for a detailed presentation).

<span id="page-22-1"></span>Now, we are ready to show the following

**Theorem 5.c.1** (Factorization of Calderón–Lozanovskiı̆ spaces). Let X be a rearrangement invariant space such that  $X \neq L_{\infty}$ . Further, let F and G be two Young functions. Then the space  $X_G$  can be factorize through  $X_F$ , that is,  $X_F \odot M(X_F, X_G) = X_G$  if, and only if, one the following four conditions holds:

- (1)  $F^{-1}(G \ominus F)^{-1} \approx G^{-1}$  for all arguments and neither  $L_{\infty} \hookrightarrow X$  nor  $X \hookrightarrow L_{\infty}$ ;
- (2)  $F^{-1}(G \ominus F)^{-1} \approx G^{-1}$  for large arguments and  $L_{\infty} \hookrightarrow X$ ;
- (3)  $F^{-1}(G \ominus F)^{-1} \approx G^{-1}$  for small arguments,  $X \hookrightarrow L_{\infty}$  and the triple  $(X, F, G)$  is nice;
- (4)  $F^{-1}(G \ominus_1 F)^{-1} \approx G^{-1}$  for small arguments and either the triple  $(X, F, G)$  fails to be nice or  $X$  is a sequence space.

In particular, in this situation, the space  $X_F$  is  $X_G$ -perfect.

*Proof.* Let X, F and G be as above. We will only explain how to prove  $(1)$ , because the rest is essentially the same.

Suppose that  $X_F \odot M(X_F, X_G) = X_G$ . Since we are only interested in (1) anyway, so we can assume that neither  $L_{\infty} \hookrightarrow X$  nor  $X \hookrightarrow L_{\infty}$ . Then, due to our assumption that  $X \neq L_{\infty}$ , so Lemma [5.b.5](#page-19-2) teaches us that  $G^{-1} \preccurlyeq F^{-1}(G \ominus F)^{-1}$  for all arguments. Therefore, it only remains to explain that also  $F^{-1}(G \ominus F)^{-1} \preccurlyeq G^{-1}$  for all arguments. To see this, let us recall that the conjugate function  $G \ominus F$  always satisfy generalized Young's inequality, that is,  $G(st) \leqslant (G \ominus F)(t) + F(s)$  for  $s, t \geqslant 0$  (see [\(4.a.1\)](#page-9-4)). Now, it is enough to invoke Theorem 6.1 from [\[O'Ne65\]](#page-27-0) (see also [\[KLM12,](#page-26-2) pp. 892–893] and [\[KLM14,](#page-26-5) Remark 6]).

In order to obtain the reverse implication, suppose that  $F^{-1}(G \ominus F)^{-1} \approx G^{-1}$  for all arguments and neither  $L_{\infty} \hookrightarrow X$  nor  $X \hookrightarrow L_{\infty}$ . Then, it follows from Lemma [5.b.4](#page-19-3) that  $X_F \odot X_{G\ominus F} = X_G$ . However, we also know from Theorem [4.b.3](#page-11-0) that the space  $X_{G\ominus F}$ coincide with  $M(X_F, X_G)$ . In consequence, the factorization  $X_F \odot M(X_F, X_G) = X_G$ holds.

<span id="page-23-0"></span>Finally, the last part about  $X_G$ -perfectness of the space  $X_F$  follows directly from the discussion preceding the theorem.

# 6. What's next?

<span id="page-23-1"></span>In this closing section, let us gather and shortly discuss some problems which naturally arise from this work.

6.a. How badly can factorization fail? The following problem (in a slightly different version) was proposed to authors by Professor Mieczysław Mastyło.

**Problem 6.a.1.** Suppose that F, G and  $G \ominus F$  are N-functions<sup>\*</sup>. Is it then true that  $F^{-1}(G \ominus F)^{-1} \approx G^{-1}$ ?

This is a very intriguing question, because in every existing in the literature example showing that  $F^{-1}(G \ominus F)^{-1} \not\approx G^{-1}$ , at least on of the functions F or G is not an  $\mathcal N$ -function. A positive answer to the aforementioned question would indicate that factorization may fail only in somewhat pathological situations. Conversely, a negative outcome would yield new intriguing examples of Orlicz functions.

<span id="page-23-2"></span>6.b. Orlicz–Lorentz spaces. A function  $w \in L_0$  is called the weight whenever it is nonnegative and decreasing. For a given weight w, by the **weighted Orlicz space**  $L_F(w)$ we understand the Orlicz space associated to the Young function  $F$  and the measure  $wdt$ , that is,

$$
L_F(w) \coloneqq \left\{ f \in L_0 \colon \int F(\lambda |f(t)|) w(t) dt < \infty \text{ for some } \lambda = \lambda(f) > 0 \right\}.
$$

Recall that the **Orlicz–Lorentz space**  $\Lambda_F(w)$  is defined as a *symmetrization* of the corresponding weighted Orlicz space  $L_F(w)$ , that is, the space  $\Lambda_F(w)$  consists of all  $f \in L_0$ such that  $f^* \in L_F(w)$ . In particular, if F is just a power function, that is,  $F(t) = t^p$ for some  $p \geq 1$ , the space  $\Lambda_F(w)$  is usually denoted by  $\Lambda_p(w)$  and sometimes called the Lorentz–Sharpley space.

In the early 90's, Yves Raynaud showed that the space  $M(\Lambda_p(w), \Lambda_q(v))$  of pointwise multipliers between two Lorentz–Sharpley spaces  $\Lambda_p(w)$  and  $\Lambda_q(v)$  coincide, up to the equivalence of norms, with anther Lorentz–Sharpley space  $\Lambda_r(u)$ . Here,  $1/r = 1/q - 1/p$ and the weight u verify the relation  $u^{1/r}v^{1/q} \approx w^{1/p}$  (see [\[Ray92,](#page-27-12) Proposition 25]).

<span id="page-23-4"></span>**Problem 6.b.1.** Provide a representation of the space  $M(\Lambda_M(w), \Lambda_N(v))$  and deduce the factorization of Orlicz–Lorentz spaces.

Note that Theorem [4.b.3](#page-11-0) gives an answer to the above problem only in the case when both weights are the same. To see this, it is enough to observe that the Orlicz–Lorentz space  $\Lambda_F(w)$  coincide with the Calderón–Lozanovskiı̆ construction  $(\Lambda(w))_F$ , where the space  $\Lambda(w)$  is defined via the norm  $||f||_{\Lambda(w)} := \int f^*(t)w(t)dt$ . Then, plainly,

$$
M(\Lambda_F(w), \Lambda_G(w)) = M((\Lambda(w))_F, (\Lambda(w))_G) = (\Lambda(w))_{G \ominus F} = \Lambda_{G \ominus F}(w).
$$

<span id="page-23-3"></span>\*Recall that an Orlicz function F is called the N-function if  $\lim_{t\to 0^+} \frac{F(t)}{t} = 0$  and  $\lim_{t\to\infty} \frac{F(t)}{t} = \infty$ (cf. [\[Mal04,](#page-26-14) p 47]).

<span id="page-24-0"></span>6.c. Musielak–Orlicz setting. Another, much more general, way to look at the situation presented in this work is to consider a variant of Calderón–Lozanovskiı's construction  $X_F$  in which the Young function F is replaced by the so-called Musielak–Orlicz function Φ (see [\[Mal04,](#page-26-14) p. 11] for details). For lack of a better idea, and only for purposes of this paragraph, let us call them the **generalized Calderón–Lozanovski** $\check{\mathbf{x}}_0$  spaces  $X_{\Phi}$ . In the light of the recent results obtained in [\[LT21\]](#page-26-6), the following problem seems tempting.

<span id="page-24-3"></span>**Problem 6.c.1.** Lift the results of this paper to the setting of generalized Calderón– Lozanovskiĭ spaces.

We firmly believe that an answer to the above question is within reach, but we have not verified all the details. However we have some thoughts and tips for the committed reader. To approach this problem, we need to generalise Lemmas 3 and 4 from [\[LT21\]](#page-26-6) to the setting of generalized Calderón–Lozanovskiĭ spaces. Then we can follow the proof of Theorem [4.b.3](#page-11-0) (replacing Young conjugates with the version appropriate for Musielak– Orlicz functions), with the main change in "Step 1" where we need the above lemmas to proceed. The rest is just checking the details. (Parenthetically speaking, we have deliberately refrained from providing any exact formulas or definitions here because they are quite ugly and technical. Anyway, everything - modulo references to literature - can be found in [\[LT21\]](#page-26-6).) Note also that an answer to Problem [6.c.1](#page-24-3) will immediately yield an answer to Problem [6.b.1.](#page-23-4) To see this, just consider Musielak–Orlicz functions of the following form  $\Phi(t, s) = F(t)w(s)$ , where F is a Young function and w is a weight.

Actually, there is one more thing.

**Problem 6.c.2.** Solve the generalized version of Problem [6.b.1](#page-23-4) in which the class of Orlicz–Lorentz spaces  $\Lambda_F(w)$  is replaced by the class of symmetrizations of the generalized Calderón–Lozanovski $\check{\alpha}$  spaces  $X_{\Phi}$ .

<span id="page-24-1"></span>For now, however, this problem looks like the "ultimate horror".

6.d. Non-symmetric variant. The most straightforward way to generalize our results is to give up the assumption about symmetry.

**Problem 6.d.1.** Prove Theorems [4.b.3](#page-11-0) and [4.b.5](#page-14-0) without assuming that the space X is rearrangement invariant.

It seems that this problem is essentially similar to the transition from the case of Orlicz spaces to Musielak–Orlicz spaces (as done in [\[LT21\]](#page-26-6)), Most likely, its proof will rely on the "localization" of the arguments used to show Theorems [4.b.3](#page-11-0) and [4.b.5.](#page-14-0) As before, the main obstacle will be "Step 1" in the proof of Theorem [4.b.3.](#page-11-0) While solving this problem, however, one should not only consider the behaviour of the Young functions  $F$  and  $G$ , but also the rate of decay (or lack thereof) of the norm of the indicator functions.

<span id="page-24-2"></span>6.e. Non-commutative affairs. There are many papers (like, for example, those by Han [\[Han15\]](#page-25-24), Han, Shao and Yan [\[HSY21\]](#page-25-25) or de Jager and Labuschagne [\[JL19\]](#page-25-26)) contemplating non-commutative analogues of the results obtained in [\[KLM12\]](#page-26-2) and [\[KLM14\]](#page-26-5). The prospect of enhancing them with the technology invented here seems very promising. All this encourages us to pose the following

**Problem 6.e.1.** Provide non-commutative variants of the main results obtained here.

# <span id="page-25-0"></span>**REFERENCES**

<span id="page-25-20"></span><span id="page-25-16"></span><span id="page-25-15"></span><span id="page-25-14"></span><span id="page-25-11"></span><span id="page-25-10"></span><span id="page-25-9"></span><span id="page-25-8"></span><span id="page-25-7"></span><span id="page-25-6"></span><span id="page-25-5"></span><span id="page-25-4"></span><span id="page-25-1"></span>

<span id="page-25-26"></span><span id="page-25-25"></span><span id="page-25-24"></span><span id="page-25-23"></span><span id="page-25-22"></span><span id="page-25-21"></span><span id="page-25-19"></span><span id="page-25-18"></span><span id="page-25-17"></span><span id="page-25-13"></span><span id="page-25-12"></span><span id="page-25-3"></span><span id="page-25-2"></span>[JR76] R. E. Jamison and W. H. Ruckle, Factoring absolutely convergent series, Math. Ann. 244 (1976), no. 2, 143–148.

- <span id="page-26-20"></span>[Kal92] N. J. Kalton, Differentials of complex interpolation processes for Köthe function spaces, Trans. Amer. Math. Soc. 333 (1992), 479–529.
- <span id="page-26-18"></span>[Kal03] N. J. Kalton, *Quasi-Banach spaces*, in: Handbook of the Geometry of Banach Spaces, Vol. 2, pp. 1099–1130, North-Holland, Amsterdam 2003.
- [KN20] R. Kawasumi and E. Nakai, Pointwise multipliers on weak Orlicz spaces, Hiroshima Math. J. 50 (2020), no. 2, 169–184.
- <span id="page-26-9"></span>[KT17] T. Kiwerski and J. Tomaszewski, Local approach to order continuity in Cesàro function spaces, J. Math. Anal. Appl. 455 (2017), no. 2, 1636–1654.
- <span id="page-26-3"></span>[KT22] T. Kiwerski and J. Tomaszewski, Essential norms of pointwise multipliers in the nonalgebraic setting, preprint available on [arxiv.org/abs/2212.06723](https://arxiv.org/abs/2212.06723), December 2022.
- <span id="page-26-16"></span>[KT24] T. Kiwerski and J. Tomaszewski, Arithmetic, interpolation and factorization of amalgams, preprint available on [arxiv.org/abs/2401.05526](https://arxiv.org/abs/2401.05526), January 2024.
- <span id="page-26-13"></span>[KL10] P. Kolwicz and K. Leśnik, Topological and geometrical structure of Calderón–Lozanovski $\tilde{i}$ construction, Math. Ineq. Appl. 13 (2010), no. 1, 175–196.
- <span id="page-26-2"></span>[KLM12] P. Kolwicz, K. Leśnik and L. Maligranda, Pointwise multipliers of Calderón–Lozanovski $\tilde{u}$ spaces, Math. Nachr. 286 (2012), no. 8-9, 876–907.
- <span id="page-26-5"></span>[KLM14] P. Kolwicz, K. Leśnik and L. Maligranda, Pointwise products of some Banach function spaces and factorization, J. Funct. Anal. 266 (2014), no. 2, 616–659.
- <span id="page-26-17"></span>[KLM19] P. Kolwicz, K. Leśnik and L. Maligranda, Symmetrization, factorization and arithmetic of quasi-Banach function spaces, J. Math. Anal. Appl. 470 (2019), 1136–1166.
- <span id="page-26-21"></span>[KR61] M. A. Krasnoselskiĭ and Ja. B. Rutickiĭ, Convex Functions and Orlicz Spaces, Gosudarstv. Izdat. Fiz.-Mat. Lit., Moscow, 1958 (in Russian); English translation in: Noordhoff, Groningen, 1961.
- <span id="page-26-4"></span>[LT17] K. Leśnik and J. Tomaszewski, Pointwise multipliers of Orlicz function spaces and factorization, Positivity 21 (2017), no. 4, 1563–1573.
- <span id="page-26-6"></span>[LT21] K. Leśnik and J. Tomaszewski, Pointwise multipliers of Musielak–Orlicz spaces and factorization, Rev. Math. Complut. 34 (2021), 489–509.
- <span id="page-26-7"></span>[Tom21] J. Tomaszewski, Mnożniki punktowe pomiędzy przestrzeniami Orlicza, PhD thesis (2021), 81 pp. (in Polish).
- <span id="page-26-8"></span>[LT79] J. Lindenstrauss and L. Tzafriri, Classical Banach Spaces II. Function Spaces, Springer-Verlag, Berlin-New York 1979.
- <span id="page-26-19"></span>[Loz69] G. Ja. Lozanovskiĭ, On some Banach lattices, Sibirsk. Math. Zh. 10 (1969), 584–599 (in Russian); English translation in: Siberian Math. J. 10 (1969), no. 3, 419–431.
- <span id="page-26-10"></span>[Loz71] G. Ja. Lozanovskiĭ, The Banach lattices and concave functions, Dokl. Akad. Nauk SSSR 199 (1971), 536–539 (in Russian); English translation in: Soviet Math. Dokl. 12 (1971), 1114–1117.
- <span id="page-26-11"></span>[Loz73] G. Ja. Lozanovski˘ı, Certain Banach lattices. IV, Sibirsk. Mat. Zh. 14 (1973), 140–155 (in Russian); English translation in: Siberian Math. J. 14 (1973), 97–108.
- <span id="page-26-12"></span>[Loz78] G. Ja. Lozanovskiĭ, On the conjugate space of a Banach lattice, Theor. Funktsiĭ Funktsional. Anal. i Prilozhen. 30 (1978), 85–90 (in Russian).
- <span id="page-26-15"></span>[Mal89] L. Maligranda, Orlicz Spaces and Interpolation, Sem. Mat. 5, Universidade Estadual de Campinas, Dep. Mat., Campinas 1989.
- <span id="page-26-14"></span>[Mal04] L. Maligranda, *Calderón–Lozanovskiĭ* construction for mixed norm spaces, Acta Math. Hungar. 103 (2004), no. 4, 279–302.
- <span id="page-26-1"></span>[MN10] L. Maligranda and E. Nakai, *Pointwise multipliers of Orlicz spaces*, Arch. Math. **95** (2010), no. 3, 251–256.
- <span id="page-26-0"></span>[MP89] L. Maligranda and L. E. Persson, Generalized duality of some Banach function spaces, Indag. Math. 51 (1989), no. 3, 323–338.
- [MW91] L. Maligranda and W. Wnuk, Landau type theorem for Orlicz spaces, Math. Z. 208 (1991), 57–64.

<span id="page-27-18"></span><span id="page-27-16"></span><span id="page-27-15"></span><span id="page-27-14"></span><span id="page-27-11"></span><span id="page-27-10"></span><span id="page-27-9"></span><span id="page-27-8"></span><span id="page-27-5"></span><span id="page-27-1"></span><span id="page-27-0"></span>

<span id="page-27-19"></span><span id="page-27-17"></span><span id="page-27-13"></span><span id="page-27-12"></span><span id="page-27-7"></span><span id="page-27-6"></span><span id="page-27-4"></span><span id="page-27-3"></span><span id="page-27-2"></span>(Tomasz Kiwerski) Poznań University of Technology, Institute of Mathematics, Piotrowo 3A, 60-965 Poznań, Poland

Email address: [tomasz.kiwerski@gmail.com](mailto:tomasz.kiwerski@gmail.com)

(Jakub Tomaszewski) Poznań University of Technology, Institute of Mathematics, Piotrowo 3A, 60-965 Poznań, Poland

Email address: [tomaszewskijakub@protonmail.com](mailto:tomaszewskijakub@protonmail.com)